

Contents

1 Preliminaries	1
1.1 Function spaces	1
1.2 Chemotaxis	8
2 Critical and supercritical spaces	13
3 Mapping properties of Keller–Segel nonlinearities	19
3.1 Preliminaries	19
3.2 The inhomogeneous parabolic-elliptic model	21
3.3 The homogeneous parabolic-elliptic model	27
3.4 The parabolic-parabolic model, I	32
3.5 The parabolic-parabolic model, II	35
4 Equations of Keller–Segel type	39
4.1 Preparations	39
4.2 The inhomogeneous parabolic-elliptic model	41
4.3 The homogeneous parabolic-elliptic model	44
4.4 The parabolic-parabolic model, I	45
4.5 The parabolic-parabolic model, II	47
4.6 Stability and well-posedness	49
4.7 Decay properties	52
4.8 Positivity	57
4.9 Conservation properties	60
4.10 A discussion: Damping constants and negative chemotaxis	62
5 Further PDE models for chemotaxis	65
5.1 Fokker–Planck equations	65
5.2 Models with logistic terms	69
5.3 Models for competing species	76
5.4 Density-dependent sensitivity models	77
5.5 Decay, positivity, and well-posedness	85
5.6 Peaks, troughs, stripes, spots, and Faber bases	85
6 Navier–Stokes equations	95
6.1 Preliminaries	95
6.2 Critical and supercritical spaces	98
6.3 Main assertions	100
6.4 Stability and well-posedness	102

7 Chemotaxis Navier–Stokes equations	107
7.1 Introduction	107
7.2 Critical and supercritical spaces	111
7.3 Chemotaxis Navier–Stokes nonlinearities	113
7.4 Main assertions	115
7.5 Stability and well-posedness	117
7.6 Double chemotaxis Navier–Stokes equations	118
7.7 Decay, positivity, and well-posedness	119
Bibliography	121
Symbols	127
Index	129