

# Contents

<b>1</b>	<b>Introduction</b>	1
1.1	Main result and historical context	1
1.2	Statement of the main results	7
1.3	Basic notation	16
<b>2</b>	<b>KAM for PDEs and strategy of proof</b>	19
2.1	The Newton–Nash–Moser algorithm	19
2.2	The reducibility approach to KAM for PDEs	22
2.2.1	Transformation laws and reducibility	22
2.2.2	Perturbative reducibility	23
2.3	Reducibility results	27
2.3.1	KAM for 1-dimensional NLW and NLS equations with Dirichlet boundary conditions	27
2.3.2	KAM for 1-dimensional NLW and NLS equations with periodic boundary conditions	30
2.3.3	Space multidimensional PDEs	32
2.3.4	1-dimensional quasi- and fully nonlinear PDEs, water waves	34
2.4	The multiscale approach to KAM for PDEs	41
2.4.1	Time-periodic case	44
2.4.2	Quasiperiodic case	47
2.4.3	The multiscale analysis of Chapter 5	51
2.5	Outline of the proof of Theorem 1.2.1	57
<b>3</b>	<b>Hamiltonian formulation</b>	69
3.1	Hamiltonian form of NLW equation	69
3.2	Action-angle and normal variables	71
3.3	Admissible Diophantine directions $\bar{\omega}_\varepsilon$	73
<b>4</b>	<b>Functional setting</b>	77
4.1	Phase space and basis	77
4.2	Linear operators and matrix representation	79
4.3	Decay norms	85
4.4	Off-diagonal decay of $\sqrt{-\Delta + V(x)}$	96
4.5	Interpolation inequalities	105
<b>5</b>	<b>Multiscale Analysis</b>	113
5.1	Multiscale proposition	113
5.2	Matrix representation	117
5.3	Multiscale step	120
5.4	Separation properties of bad sites	123

5.5	Definition of the sets $\Lambda(\varepsilon; \eta, X_{r,\mu})$	134
5.6	Right inverse of $[\mathcal{L}_{r,\mu}]_N^{2N}$ for $\bar{N} \leq N < N_0^2$	136
5.7	Inverse of $\mathcal{L}_{r,\mu,N}$ for $N \geq N_0^2$	146
5.7.1	The set $\Lambda(\varepsilon; 1, X_{r,\mu})$ is good at any scale	146
5.7.2	Inverse of $\mathcal{L}_{r,\mu,N}$ for $N \geq N_0^2$	151
5.8	Measure estimates	153
5.8.1	Preliminaries	153
5.8.2	Measure estimate of $\Lambda \setminus \tilde{\mathcal{G}}$	155
5.8.3	Measure estimate of $\tilde{\Lambda} \setminus \mathcal{G}_{N,\frac{1}{2}}^0$ for $N \geq N_0^2$	156
5.8.4	Measure estimate of $\tilde{\Lambda} \setminus \mathcal{G}_{N_k,\frac{1}{2}}^0$ for $k \geq 1$	157
5.8.5	Stability of the $L^2$ good parameters under variation of $X_{r,\mu}$	161
5.8.6	Conclusion: proof of (5.1.17) and (5.1.18)	162
<b>6</b>	<b>Nash–Moser theorem</b>	165
6.1	Statement	165
6.2	Shifted tangential frequencies up to $O(\varepsilon^4)$	171
6.3	First approximate solution	173
<b>7</b>	<b>Linearized operator at an approximate solution</b>	175
7.1	Symplectic approximate decoupling	175
7.2	Proof of Proposition 7.1.1	179
7.3	Proof of Lemma 7.1.2	185
<b>8</b>	<b>Splitting of low-high normal subspaces up to <math>O(\varepsilon^4)</math></b>	189
8.1	Choice of $\mathbb{M}$	189
8.2	Homological equations	191
8.3	Averaging step	195
<b>9</b>	<b>Approximate right inverse in normal directions</b>	199
9.1	Split admissible operators	199
9.2	Approximate right inverse	201
<b>10</b>	<b>Splitting between low-high normal subspaces</b>	205
10.1	Splitting step and corollary	205
10.2	The linearized homological equation	218
10.3	Solution of homological equations: proof of Lemma 10.2.2	220
10.4	Splitting step: proof of Proposition 10.1.1	239
<b>11</b>	<b>Construction of approximate right inverse</b>	247
11.1	Splitting of low-high normal subspaces	247
11.2	Approximate right inverse of $\mathcal{L}_D$	248
11.3	Approximate right inverse of $\mathcal{L} = \mathcal{L}_D + \varrho$	260
11.4	Approximate right inverse of $\tilde{\omega}_\varepsilon \cdot \partial_\varphi - J(A_0 + \rho)$	263

**12 Proof of the Nash–Moser Theorem** . . . . . 271

    12.1 Approximate right inverse of  $\mathcal{L}_\omega$  . . . . . 271

    12.2 Nash–Moser iteration . . . . . 275

    12.3  $C^\infty$  solutions . . . . . 291

**13 Genericity of the assumptions** . . . . . 295

    13.1 Genericity of nonresonance and nondegeneracy conditions . . . . . 295

**A Hamiltonian and reversible PDEs** . . . . . 311

    A.1 Hamiltonian and reversible vector fields . . . . . 311

    A.2 Nonlinear wave and Klein–Gordon equations . . . . . 312

    A.3 Nonlinear Schrödinger equation . . . . . 315

    A.4 Perturbed KdV equations . . . . . 316

**B Multiscale Step** . . . . . 319

    B.1 Matrices with off-diagonal decay . . . . . 319

    B.2 Multiscale step proposition . . . . . 326

**C Normal form close to an isotropic torus** . . . . . 337

    C.1 Symplectic coordinates near an invariant torus . . . . . 337

    C.2 Symplectic coordinates near an approximately invariant torus . . . . . 343

**Bibliography** . . . . . 349

**Index** . . . . . 357