

WHEN IS AN EQUIVALENCE RELATION CLASSIFIABLE?

GREG HJORTH

ABSTRACT. One finds in certain branches of analysis the idea that a classifiable equivalence relation is one for which we can assign points in a very concrete space as a complete invariant. Results by Effros, Glimm, and Mackey, and then later Harrington, Kechris, and Louveau, have given a thorough analysis of when such a classification is possible. In the last few years a similar analysis has been undertaken by descriptive set theorists regarding when an equivalence relation is classifiable by *countable structures considered up to isomorphism*. There is a kind of parallel theory of which equivalence relations can be assigned countable structures as complete invariants.

1991 Mathematics Subject Classification: 04A15

Keywords and Phrases: Equivalence relations, effective cardinality, classification, Polish group actions.

§0 ONE ANSWER The question posed in the title of this talk is admittedly a vague one. Not only is the question itself vague, but moreover any answer to this question will necessarily be subjective, since a classification theorem will only be satisfactory if it is judged as such for some specific purposes.

Nevertheless, in certain branches of mathematics, especially those influenced by the works of George Mackey, one finds the idea that a *classifiable* equivalence relation is one for which points in some very concrete spaces – such as \mathbb{R} , \mathbb{C} , \mathbb{T} , $C([0, 1])$ – can be assigned in some reasonably ‘nice’, preferably Borel, manner. Ultimately I will discuss some alternative notions of *classifiable* and present motivating examples for this line of research. Before continuing we should understand the following definition.

0.1 DEFINITION Let E be an equivalence relation on a Polish space X . E is *smooth* or *tame* if there is Polish space Y and a Borel function

$$\theta : X \rightarrow Y$$

such that for all $x, y \in X$

$$xEy \Leftrightarrow \theta(x) = \theta(y).$$

Just so there are no confusions about the definitions, a *Polish space* is a separable topological space that admits a complete compatible metric – and so the class of Polish spaces includes objects like the reals, the complex numbers,

Hilbert space, and so on. A function between Polish spaces is said to be *Borel* if the pullback of any open set is Borel.

It is also customary in this context to refer to a Polish space stripped down to its Borel structure as a *standard Borel space*; that is to say, (Y, \mathcal{B}) is a standard Borel space if there is a Polish topology τ on Y with respect to which \mathcal{B} is the σ -algebra generated by the τ -open sets.

In definition 0.1 we could just as well insist that Y be \mathbb{R} , since any Polish space allows a Borel injection into the reals.

It may then be helpful to think of the function

$$\theta : X \rightarrow \mathbb{R}$$

from 0.1 as lifting to an injection

$$\hat{\theta} : X/E \rightarrow \mathbb{R},$$

and that in this sense the *Borel cardinality* of X/E is less than or equal to the Borel cardinality of \mathbb{R} . Indeed this is an important theme in this branch of descriptive set theory: Determine the *effective cardinality* of quotients of the form X/E .

Another equivalent formulation of smoothness is that the space of equivalence classes, $X/E = \{[x]_E : x \in X\}$, be a subspace of a *standard Borel space* in the quotient Borel structure – that is to say, if we let \mathcal{B}_E be the collection of subsets of X/E of the form $\{[x]_E : x \in A\}$ for $A \subset X$ an E -invariant (any $x \in A$ has $[x]_E \subset A$) Borel set, then there is some standard Borel space (Y, \mathcal{B}) with $Y \supset X/E$ and $\mathcal{B}_E = \{A \cap X/E : A \in \mathcal{B}\}$. Finally, E is smooth if and only if there is a countable sequence $(A_n)_{n \in \mathbb{N}}$ of E -invariant such that for all $x, y \in X$

$$xEy \Leftrightarrow \forall n(x \in A_n \Leftrightarrow y \in A_n).$$

I suppose that for a mathematician approaching this from another area the restriction to the Borel category may seem rather arbitrary. It turns out that many mathematical objects can be naturally realized as either points in some Polish space or as equivalence classes in some Polish space, and in fact the context of these problems is far wider than it may initially appear. The theorems stated below in §4 for Borel functions all pass to much more general classes of *reasonably definable* functions.

Historically the notion of smoothness as classifiability is extremely important. Not only does one find the notion in papers such as [2], [3], and [5], and perhaps [15]. These papers suggest a wider project to determine which equivalence relations are smooth and which classification problems are no harder than that of the equality relation on \mathbb{R} .

§1 EXAMPLES: SMOOTH

1.1 EXAMPLE: COMPACT RIEMANN SURFACES A very natural classification problem is that of compact Riemann surfaces considered up to conformal equivalence. In this case there exists a reduction to the equality relation on the reals. The classical theory, as at say [11], obtains points in some standard Borel space as a complete invariant.

Of course one can ask how this sits with the original definition at 0.1. Here it is routine (but see [13] for details) to obtain a standard Borel space parameterizing (separable) complex manifolds in some natural manner. In this context one has that the set of points parameterizing the *compact complex surfaces* is Borel and the equivalence relation of conformal equivalence restricted to this Borel set is indeed smooth in exactly the sense of 0.1.

1.2 EXAMPLE: BERNOULLI SHIFTS Let $S = \{s_1, \dots, s_n\}$ be a finite alphabet, $\sigma : S^{\mathbb{Z}} \rightarrow S^{\mathbb{Z}}$ be the shift map, and for p_1, p_2, \dots, p_n a finite sequence of positive numbers summing to 1 let μ the product measure resulting from giving s_i the weight p_i . We may choose to think of two such systems as being equivalent if there is an invertible measure preserving map that conjugates them: that is, set $(S_1, \sigma_1, \mu_1) \sim (S_2, \sigma_2, \mu_2)$ if there is a measurable preserving bijection

$$\pi : (S_1)^{\mathbb{Z}} \rightarrow (S_2)^{\mathbb{Z}}$$

such that

$$\begin{aligned} \sigma_1 &= \pi^{-1} \circ \sigma_2 \circ \pi \\ \forall A \subset (S_2)^{\mathbb{Z}} (\mu_2(A) &= \mu_1(\pi^{-1}(A))). \end{aligned}$$

Ornstein in [16] shows a single real number, the *entropy* of the system (S, σ, μ) , provides a complete invariant. Moreover in a suitable standard Borel structure, this invariant *can* be calculated in a Borel fashion. Here as a suitable Borel structure one may represent the shift by the sequence $p_1, p_2, \dots, p_n \in \mathbb{R}^n$ for various n ; the point is that a countable union of standard Borel spaces, such as $\bigcup_n \mathbb{R}^n$ is again standard Borel.

1.3 EXAMPLE: GROUP REPRESENTATIONS Consider the irreducible representations of the group \mathbb{Z} . Given a complex Hilbert space H with associated unitary group U of all inner product respecting transformations, we can let $\text{Irr}(\mathbb{Z}, H)$ be the space of homomorphisms

$$\tau : \mathbb{Z} \rightarrow U$$

where U has no non-trivial invariant subspaces under $\tau[\mathbb{Z}]$. It is natural to think of τ_1 and τ_2 as somehow presenting equivalent representations if there is some $T \in U$ with

$$\tau_1(g) = T \circ \tau_2(g) \circ T^{-1}$$

for all $g \in \mathbb{Z}$.

The space of *all* representations may be naturally identified with a closed subspace of $H^{\mathbb{Z}}$, and hence it is a Polish space. Furthermore the equivalence relation of interest here is induced by the continuous action of the group H .

Here $\text{Irr}(\mathbb{Z}, H)$ is non-empty if and only if H is one dimensional. Moreover we may identify the elements of $\text{Irr}(\mathbb{Z}, H)$ with characters, and thus a complete classification of these objects may be given by points in \mathbb{T} , and hence \mathbb{R} .

On the other hand if G is finite the space $\text{Irr}(G, H)$ will be non-empty only when H is finite dimensional. Then the above equivalence relation will be induced by the a continuous action of the now compact group U on the Polish space

$\text{Irr}(G, H)$. In general such orbit equivalence relations are always classifiable by points in \mathbb{R} .

§2 EXAMPLES: NON-SMOOTH

2.1 EXAMPLE: GENERAL COMPLEX DOMAINS One can view Becker, Henson, and Rubel in [1] as obtaining non-classifiability by a process tantamount to embedding the equivalence relation E_0 of eventual agreement on infinite sequences of 0's and 1's into conformal equivalence on complex domains – so that for $f, g : \mathbb{N} \rightarrow \{0, 1\}$ we have fE_0g if there is some $N \in \mathbb{N}$ such that

$$\forall n > N (f(n) = g(n)).$$

Here E_0 is an F_σ equivalence relation on $\{0, 1\}^{\mathbb{N}}$, the space of all infinite binary sequences in the product topology, and is in some ways (compare [9]) the canonical example of a non-smooth equivalence relation.

In fact if we assign \mathcal{D} , the space of open subsets of \mathbb{C} , with the *Effros standard Borel structure* – under which it does have a natural Borel structure – then their argument can be seen as showing that there is a Borel function

$$\theta : \{0, 1\}^{\mathbb{N}} \rightarrow \mathcal{D}$$

such that fE_0g if and only if $\theta(f)$ and $\theta(g)$ are biholomorphic. Since E_0 is non-smooth we obtain non-smoothness of conformal equivalence on arbitrary complex surfaces, even with respect to the Borel structure articulated in [13].

2.2 EXAMPLE: ARBITRARY MEASURE PRESERVING TRANSFORMATIONS Consider M_∞ the group of all invertible measure preserving transformations of the unit interval. In the topology it inherits from its action on $L^2([0, 1])$ this is a topological group that is Polish as a space – that is to say, it is a *Polish group*. For instance, if (U_n) enumerates the basic open subsets of $[0, 1]$ we obtain a complete metric with

$$d(\pi_1, \pi_2) = \sum_{n \in \mathbb{N}} 2^{-n} (\lambda(\pi_1(U_n) \Delta \pi_2(U_n)) + \lambda(\pi_1^{-1}(U_n) \Delta \pi_2^{-1}(U_n))).$$

The obvious classification problem is for the conjugacy equivalence relation – it is natural to say that $\pi_1, \pi_2 : [0, 1] \rightarrow [0, 1]$ are *equivalent* if they are conjugate, in the sense of their being some $\sigma \in M_\infty$ such that

$$\sigma \circ \pi_1 = \pi_2 \circ \sigma \text{ a.e.}$$

This equivalence relation was observed by Feldman [5] to be non-smooth. As with 2.1 the proof rested on embedding E_0 .

2.3 EXAMPLE: GROUP REPRESENTATIONS AGAIN Let G be a countable discrete group that is *not* abelian-by-finite. Let H_∞ be a separable infinite dimensional Hilbert space and U_∞ the unitary group on H_∞ . Again take $\text{Irr}(G, H_\infty)$ to be space of irreducible representations $\tau : G \rightarrow U_\infty$ with the equivalence relation of conjugacy –

$$\tau_1 \approx \tau_2 \Leftrightarrow \exists A \in U_\infty \forall g \in G (\tau_1(g) = A^{-1} \circ \tau_2(g) \circ A).$$

It is known from [17] and [8] that $\text{Irr}(G, H_\infty)$ is non-empty and \approx is not smooth: there is no Borel assignment of reals as complete invariants to $\text{Irr}(G, H_\infty)/\approx$.

§3 MORE EXAMPLES: PUZZLING CASES The above were deliberately chosen with the view to supporting the intuition that *classifiable* means *smooth*. In the cases where there is a proof of smoothness, it is generally accepted as a classification theorem. In the cases where the equivalence relation does not admit points in \mathbb{R} as a complete invariant, the authors seemed to take *that* as a proof of at least some manner of non-classifiability. Consequently I hope the position that takes classifiable to mean smooth will seem an initially attractive one.

This much said, let us consider some examples where there is a more generous notion of classifiability implicit; these in turn have motivated the search for new tools in the study of Borel and analytic equivalence relations.

3.1 QUESTION: COMPLEX SURFACES Becker, Henson, and Rubel in [1] explicitly ask: is there some reasonably non-pathological way to assign to every domain $D \subset \mathbb{C}$ some countable set of complex numbers S_D such that

$$D \cong D'$$

if and only if

$$S_D = S_{D'}?$$

3.2 EXAMPLE: DISCRETE SPECTRUM MPT'S Halmos and von Neumann in [10] showed that for *discrete spectrum* elements of M_∞ , we may assign a countable collection $\{c_i(\pi) : i \in \mathbb{N}\}$ of complex numbers that completely describe the equivalence class of π . While conjugacy on discrete measure preserving transformations is not smooth, the Halmos-von Neumann theorem would seem to constitute some sort of weaker notion of classification, and it certainly appears to be accepted as such.

3.3 EXAMPLE: C^* -ALGEBRAS AND TOPOLOGICAL DYNAMICS (This is not quite analogous to examples 1.3 and 2.3, but derives from roughly the same area.) Giordano, Putnam, and Skau in [6] consider the problem of classifying *minimal Cantor systems up to orbit equivalence*. Two continuous

$$\varphi_1 : X_1 \rightarrow X_1,$$

$$\varphi_2 : X_2 \rightarrow X_2$$

which are *minimal* in the sense of having no non-trivial closed invariant sets and are *Cantor* in the sense of X_1, X_2 being compact, uncountable and completely disconnected metric spaces, are said to be *orbit equivalent* if there is a homeomorphism $F : X_1 \rightarrow X_2$ which respects the orbit structure set wise, in that for all x

$$\{\varphi_2^i(F(x)) : i \in \mathbb{Z}\} = F[\{\varphi_1^i(x) : i \in \mathbb{Z}\}].$$

This problem is in turn equivalent to classifying a certain class of C^* -algebras.

Here they produce countable ordered abelian groups as complete invariants. One similarly finds discussion of the classification of certain C^* by countable discrete structures considered up to isomorphism in papers such as [4].

It is important to note a link between the kind of classification one finds in 3.1-2 and 3.3: Any equivalence relation that can be classified by a countable set of reals can be classified by countable structures considered up to isomorphism. For instance, if we let (q_n) enumerate the rationals, then to a countable unordered set $A \subset \mathbb{R}$ we can associate the model $\mathcal{M}_A = \{x_a : a \in A\}$ with unary predicates (P_n) governed by the rule that

$$\mathcal{M}_A \models P_n(x_a)$$

if and only if $q_n < a$. Trivially then reduction to the equality relation on \mathbb{R} implies classification by countable sets of reals, and hence classification by countable structures.

Thus we may be led to formulate a more generous notion of classifiability.

3.4 QUESTION For which E can we provide some kind of countable structure considered up to isomorphism as a complete invariant?

Letting \mathcal{L} be a countable language, we form $\text{Mod}(\mathcal{L})$, the space of all \mathcal{L} -structures on \mathbb{N} with the topology generated by quantifier free formulas. This is a Polish space, and therefore there is a precise version of the question.

For which equivalence relations E on Polish X can we find a Borel $\theta : X \rightarrow \text{Mod}(\mathcal{L})$ such that for all $x, y \in X$

$$xEy \Leftrightarrow \theta(x) \cong \theta(y)?$$

In very general terms these examples may illustrate the kinds of concerns driving the descriptive set theory of equivalence relations, as well as the particular problem of classification by countable structures. I should add to these general remarks that the isomorphism relation on countable structures is historically important in logic, and that for someone in my area it seems intriguing to ask which classification problems may be simply reduced to that of countable models considered up to isomorphism.

§4 SOME THEOREMS I will begin with two sufficient conditions for classifiability, the first of which is trivial.

4.1 THEOREM(folklore) Let G be a compact metrizable group acting continuously on a Polish space X with induced orbit equivalence relation E_G . Then E_G is smooth.

4.2 THEOREM (Kechris [14]) Let G be a locally compact Polish group acting continuously on a Polish space X . Then there is a countable sequence of Borel functions $(f_i)_{i \in \mathbb{N}}$ such that for all $x, y \in X$

$$xE_Gy \Leftrightarrow \{f_i(x) : i \in \mathbb{N}\} = \{f_i(y) : i \in \mathbb{N}\}.$$

In other words, we may classify by *countable unordered sets* of reals.

And two sufficient conditions for non-classifiability:

4.3 THEOREM (folklore) Let G be a Polish group and X a Polish space. Suppose that

- (i) some orbit is dense;
- (ii) every orbit is meager (its complement includes the intersection of countably many open dense sets). Then E_G is not smooth.

4.4 THEOREM (Hjorth [12]) Let G be a Polish group and X a Polish space. Suppose that

- (i) some orbit is dense;
- (ii) every orbit is meager (its complement includes the intersection of countably many open dense sets);
- (iii) for some $x \in X$, the *local orbits of x* are all somewhere dense; that is to say, if V is an open neighborhood of 1_G , U is an open set containing x , and if $O(x, U, V)$ is the set of all $\hat{x} \in [x]_G$ such that there is a finite sequence $(x_i)_{i \leq k} \subset U$ such that $x_0 = x$, $x_k = \hat{x}$, and each $x_{i+1} \in V \cdot x_i$, then the closure of $O(x, U, V)$ contains an open set.

Then there is no Borel (or even Baire measurable) $\theta : X \rightarrow \text{Mod}(\mathcal{L})$ such that for all $x, y \in X$

$$xE_Gy \Leftrightarrow \theta(x) \cong \theta(y).$$

Consequently there is no sequence $(f_i)_{i \in \mathbb{N}}$ of Borel (or even *reasonably definable*) functions

$$f_i : X \rightarrow \mathbb{R}$$

such that

$$xE_Gy \Leftrightarrow \{f_i(x) : i \in \mathbb{N}\} = \{f_i(y) : i \in \mathbb{N}\}.$$

A Polish group action satisfying 4.4(i)-(iii) is called *generically turbulent*.

Again I will return to the motivation and examples in the next and final section. These examples on their own may suggest that 4.4 is the *right theorem* for showing this kind of non-classifiability.

However there are also results in [12] reinforcing this view. The presence of a generically turbulent action is necessary for non-classifiability in the sense that if E_G arises from the continuous action of Polish G on Polish X then either E_G is reducible to isomorphism on countable structures (using say *universally Baire measurable* functions) or there is a generically turbulent Polish G -space Y which admits a continuous G -embedding into X . (Here a function θ is said to be *universally Baire measurable* if for any Borel function ρ we have that $\theta \circ \rho$ is Baire measurable – in the sense of pulling back open sets to sets with the Baire property.)

§5 EXAMPLES AGAIN

5.1 EXAMPLE: COMPLEX MANIFOLDS AGAIN By the uniformization theorem, conformal equivalence on complex surfaces may be reduced to an appropriately chosen locally compact group action.

THEOREM (Hjorth-Kechris [13]) Let \mathcal{D} be the space of all complex domains. Then there is a *definable* assignment

$$M \mapsto S_M$$

of countable sets of reals to domains such that for all $M, N \in \mathcal{D}$

$$M \cong N \Leftrightarrow S_M = S_N.$$

Moreover it is Borel in the sense of there existing a countable sequence (f_n) of Borel functions from \mathcal{D} to \mathbb{R} such that S_M always equals the (unordered) set $\{f_n(M) : n \in \mathbb{N}\}$.

But in higher dimensions one may embed a generically turbulent orbit equivalence relation and obtain:

THEOREM (Hjorth-Kechris [13]) Let \mathcal{M}^2 be the space of two dimensional complex manifolds. Then there is no Borel assignment of countable structures up to isomorphism as complete invariants. Consistently with ZFC there is no *definable* assignment.

5.2 EXAMPLE: MEASURE PRESERVING TRANSFORMATIONS AGAIN

THEOREM (Hjorth) Let M_∞ be the space of invertible measure preserving transformations on the unit interval. Consider the conjugacy equivalence relation \sim : $\pi_1 \sim \pi_2$ if there is $\sigma \in M_\infty$ such that

$$\sigma \circ \pi_1 = \pi_2 \circ \sigma \text{ a.e.}$$

Then there is no sequence $(f_i)_{i \in \mathbb{N}}$ of Borel functions

$$f_i : M_\infty \rightarrow \mathbb{R}$$

such that

$$\pi_1 \sim \pi_2 \Leftrightarrow \{f_i(\pi_1) : i \in \mathbb{N}\} = \{f_i(\pi_2) : i \in \mathbb{N}\}.$$

In fact, \sim is strictly more complicated than isomorphism on countable models: there is a Borel $\theta : \text{Mod}(\mathcal{L}) \rightarrow M_\infty$ such that for all $M, N \in \text{Mod}$

$$M \cong N \Leftrightarrow \theta(M) \sim \theta(N),$$

but (for any choice of \mathcal{L}) there is no Borel (or even universally Baire measurable) $\rho : M_\infty \rightarrow \text{Mod}(\mathcal{L})$ such that for all $\pi_1, \pi_2 \in M_\infty$

$$\pi_1 \sim \pi_2 \Leftrightarrow \rho(\pi_1) \cong \rho(\pi_2).$$

5.3 EXAMPLE: DISCRETE GROUP REPRESENTATIONS AGAIN

THEOREM (Hjorth) Let G be a countable group that is *not* abelian-by-finite. Let H_∞ be an separable infinite dimensional Hilbert space, let $\text{Irr}(G, H_\infty)$ be the space of irreducible representations of G in H_∞ . Then there is no sequence $(f_i)_{i \in \mathbb{N}}$ of Borel functions

$$f_i : \text{Irr}(G, H_\infty) \rightarrow \mathbb{R}$$

such that

$$\tau_1 \approx \tau_2 \Leftrightarrow \{f_i(\tau_1) : i \in \mathbb{N}\} = \{f_i(\tau_2) : i \in \mathbb{N}\}.$$

In fact there is no *reasonably definable* assignment of countable models considered up to isomorphism as complete invariants.

REFERENCES

- [1] J. Becker, C.W. Henson, L. Rubel, *First order conformal invariants*, ANNALS OF MATHEMATICS, vol. 102(1980), pp. 124-178.
- [2] E. Effros, *Transformation groups and C^* -algebras*, ANNALS OF MATHEMATICS, ser 2, vol. 81(1975), pp. 38-55.
- [3] E. Effros, *Polish transformation groups and classification problems*, GENERAL TOPOLOGY AND MODERN ANALYSIS (Proc. Conf., Univ. California, Riverside, Calif., 1980), pp.217-227, Academic Press, New York-London, 1981.
- [4] G.A. Elliott, *On the classification of inductive limits of sequences of semisimple finite-dimensional algebras*, JOURNAL OF ALGEBRA, vol. 38(1976), pp. 29-44.
- [5] J. Feldman, *Borel structure and invariants for measurable transformations*, PROCEEDINGS OF THE AMERICAN MATHEMATICAL SOCIETY, vol. 46(1974), pp. 383-394.
- [6] T. Giordano, I.F. Putnam, C.F. Skau, *Topological orbit equivalence and C^* -crossed products*, JOURNAL REINE UND ANGEWANDT MATHEMATISCHE, vol. 469(1995), pp. 51-111.
- [7] J. Glimm, *Locally compact transformations groups*, TRANSACTIONS OF THE AMERICAN MATHEMATICAL SOCIETY, vol. 101(1961), pp. 124-138.
- [8] J. Glimm, *Type I C^* -algebras*, ANNALS OF MATHEMATICS, 1961, pp. 572-612.
- [9] L.A. Harrington, A.S. Kechris, A. Louveau, *A Glimm-Effros dichotomy for Borel equivalence relations*, JOURNAL OF THE AMERICAN MATHEMATICAL SOCIETY, vol. 3(1990), pp. 903-928.
- [10] P.R. Halmos, J. von Neumann, *Operator methods in classical mechanics, II*, ANNALS OF MATHEMATICS, vol. 43(1942), pp. 332-50.
- [11] Y. Imayoshi, M. Taniguchi, AN INTRODUCTION TO TEICHMÜLLER SPACES, Springer-Verlag, Berlin, 1992.
- [12] G. Hjorth, CLASSIFICATION AND ORBIT EQUIVALENCE RELATIONS, manuscript (available at www.math.ucla.edu/~greg/research.html).
- [13] G. Hjorth, A.S. Kechris, THE COMPLEXITY OF THE CLASSIFICATION OF RIEMANN SURFACES BY COMPLEX MANIFOLDS, preprint, UCLA, 1998.
- [14] A.S. Kechris, *Countable sections for locally compact group actions*, JOURNAL OF ERGODIC THEORY AND DYNAMICAL SYSTEMS, vol. 12(1992), pp. 283-295.
- [15] G.W. Mackey, *Infinite-dimensional group representations*, BULLETIN OF THE AMERICAN MATHEMATICAL SOCIETY, vol. 69(1963), pp.628-686.

- [16] D.S. Ornstein, *Bernoulli shifts with the same entropy are isomorphic*, ADVANCES IN MATHEMATICS, vol. 4(1970), pp. 337-52.
- [17] E. Thoma, *Eine Charakterisierung diskreter Gruppen vom type I*, INVENTIONES MATHEMATIQUE, vol. 6(1968), pp. 190-196.

Greg Hjorth
Mathematics
UCLA
CA90095-1555
USA
greg@math.ucla.edu