

WEDDERBURN'S THEOREM FOR REGULAR LOCAL RINGS

MANUEL OJANGUREN

Received: July 24, 2014

ABSTRACT. Wedderburn's theorem is extended to Azumaya algebras over certain regular local rings.

2000 Mathematics Subject Classification: 16H05

Keywords and Phrases: Division ring, Azumaya algebra, regular local ring.

In [Pa] Ivan Panin proved the following theorem.

THEOREM 1. *Let R be a regular local ring, K its field of fractions and (V, Φ) a quadratic space over R . Suppose R contains a field of characteristic zero. If $(V, \Phi) \otimes_R K$ is isotropic over K , then (V, Φ) is isotropic over R .*

The proof rests on a series of lemmas which can be summarized in a single one:

LEMMA 2. *Let k be a field of characteristic zero, u a closed point of a smooth k -variety and $R = \mathcal{O}_{U,u}$ the local ring of U at u . Let further \mathcal{X} be a projective R -scheme, smooth over R . Let K be the field of fractions of R and suppose that \mathcal{X} has a K -point. Then, for every prime number p there exist an integral R -étale algebra S of degree prime to p and an S -point of \mathcal{X} .*

Proof. See [Pa], Lemma 3, Lemma 4 and proof of Theorem 1.

I want to show that the argument used for proving Theorem 1 also yields the following extension of Wedderburn's theorem to a large class of regular local rings.

THEOREM 3. *Let R be a regular local ring, K its field of fractions and A an Azumaya algebra over R . Suppose R contains a field k of characteristic zero. If $A \otimes_R K$ is isomorphic to $M_n(D)$ where D is a central division algebra over K , then A is isomorphic to $M_n(\Delta)$ where Δ is a maximal (unramified) R -order of D . In other words, every class of the Brauer group of R is represented by an Azumaya algebra Δ such that $\Delta \otimes_R K$ is a division K -algebra.*

Proof. Let d^2 be the dimension of D over K . It suffices to show that A contains a right ideal I such that A/I is free of rank $(n^2 - n)d^2$ over R . In fact, since

any A -module is projective over A if and only if it is projective over R , the projection $A \rightarrow A/I$ splits, I is a direct factor of the right A -module A , and $\Delta := \text{End}_A(I)$ is an Azumaya algebra equivalent to A . Clearly $\Delta \otimes_R K = D$ and by Morita theory

$$A = \text{End}_\Delta(\text{Hom}_A(I, A)) = M_n(\Delta).$$

In order to find a right ideal I of the right rank we consider the set \mathcal{I} of all such ideals or, more precisely, we consider the functor \mathcal{I} that associates to any R -algebra S the set of such ideals in $A \otimes_R S$.

LEMMA 4. \mathcal{I} is a smooth closed subscheme of the Grassmannian scheme \mathcal{G} consisting of all the free R -submodules of A which are direct factors of A and have rank nd^2 .

Proof. We denote by m the maximal ideal of R . To show that \mathcal{I} is closed we first remark that A , as an R -module, is generated by the set A^* of all invertible elements of A . In fact for any $a \in A$ and any $\lambda \in k$ the reduced norm of $\lambda + a$ is a polynomial

$$P(\lambda) = \lambda^{nd} + c_1 \lambda^{n-1} + \cdots + c_{nd}$$

whose coefficients are in R and only depend on a . Choosing λ in k^* such that $P(\lambda)$ is not 0 in R/m insures that $\lambda + a$ is invertible and allows to write $a = (\lambda + a) - \lambda$. So an R -submodule M of A is an ideal if $aM = M$ for every unit a . In other words, we must show that the set of fixed points of \mathcal{G} under the action of A^* is closed. This is well-known.

The second point is the smoothness of \mathcal{I} . This means that for any R -algebra S and any ideal I of S , any S/I -point of \mathcal{X} can be lifted to an S/I^2 -point. But points correspond to right ideals generated by an idempotent and it is well-known that idempotents can be lifted.

Note that it suffices to treat the case when A is of prime power order in the Brauer group $Br(R)$ of R . In fact the class of A is a product of classes $[A_i]$ of order $p_i^{e_i}$ for some distinct primes p_1, \dots, p_r . If each of them is represented by an order Δ_i in $D_i = \Delta_i \otimes_R K$ then A is Brauer equivalent to $\Delta_1 \otimes_R \cdots \otimes_R \Delta_r$ which is an order in $D = D_1 \otimes_K \cdots \otimes_K D_r$ and we know that D is a division algebra.

We now assume that R is of geometric type, in other words R is the local ring of a closed point u of a smooth k -variety. The general case then follows from this special case by a standard application of Dorin Popescu's theorem, saying that a regular ring containing a field is an inductive limit of smooth algebras. A self-contained proof of Popescu's theorem in the form needed here has been given by R. Swan [Sw]. For the original articles by Popescu see the references in [Sw].

Suppose now that A is of prime power exponent in $Br(R)$ and that the degree of D is p^e for some prime number p . Since $A \otimes_R K = M_n(D)$ the scheme

\mathcal{I} has a K -point and according to Lemma 2 it also has an S -point, where S is an integral étale algebra whose degree d is prime to p . This means that $A \otimes_R S = M_n(B)$ for some maximal order B in $D \otimes_K L$, L being the field of fractions of S . Note that $D \otimes_K L$ remains a division algebra because the degree of L over K is prime to p . So the Brauer class $[A]_S$ of $A \otimes_R S$ in $Br(S)$ is represented by a degree p^e algebra. In [Ga] (see also [AdJ], Proposition 2.6.1) Gabber proved that any class $\alpha \in Br(R)$ which is represented by a degree m algebra when extended to a finite faithfully flat R -algebra S of degree d can be represented by an R -algebra of degree dm . We can thus find an Azumaya algebra A_1 of degree dp^e in the same class as A . On the other hand, we may also use Ferrand's [Fe] norm functor $N_{S/R}$ from S -algebras to R -algebras. Applying it to B we find that $N_{S/R}(B) = A_2$ is an Azumaya R -algebra equivalent to $A^{\otimes d}$ ([Fe], section 7.3), of degree p^{ed} ([Fe], Théorème 4.3.4). If the integer c is an inverse of d modulo p^e , the algebra $A_3 = A_2^{\otimes c}$ is Brauer equivalent to A and its degree is p^{cde} . Recall now that DeMeyer [DM] proved that every class in $Br(R)$ is represented by a unique "minimal" Azumaya algebra Δ with the property that every algebra in the same class is isomorphic to some matrix algebra over Δ . What is the degree m of this Δ in our case? We must have $A_1 \simeq M_{s_1}(\Delta)$ and $A_3 \simeq M_{s_3}(\Delta)$, hence $s_1 m = dp^e$ and $s_3 m = p^{cde}$. Since d is prime to p , this implies that m divides p^e and extending the scalars to K shows that $m = p^e$. The theorem is proved.

Easy and well-known examples (the simplest one being the usual quaternion algebra extended to $\mathbb{R}[x, y, z]/(x^2 + y^2 + z^2)$ localized at the origin) show that we cannot replace regularity by, say, normality.

Remark. As the referee pointed out, the proof of Theorem 3 could be extended to the case of a semi-local regular ring containing a field k of characteristic zero, although I do not see how to proceed if k has positive characteristic. Fortunately, since the time this article was written, new and stronger results have appeared. In [AB2] Benjamin Antieau and Ben Williams have generalized Theorem 3 to arbitrary semi-local regular rings. In [AB1] they have shown that Theorem 3 fails for arbitrary regular rings, in particular for certain smooth complex affine algebras of dimension 6.

REFERENCES

- [AB1] Benjamin Antieau and Ben Williams, *Unramified division algebras do not always contain Azumaya maximal orders*, *Invent. math.* **197** (2014), 47–56.
- [AB2] ———, *Topology and purity of torsors*, preprint, <http://arxiv.org/abs/1311.5273> (2013).
- [AdJ] Michel Artin and Aise Johan de Jong, *Stable orders over surfaces*, www.math.lsa.umich.edu/courses/711/.
- [DM] Frank DeMeyer, *Projective modules over central separable algebras*, *Canad. J. Math.* **21** (1969), 39–43.

- [Fe] Daniel Ferrand, *Un foncteur norme*, Bull. Soc. Math. France **126** (1998), 1–49.
- [Ga] Ofer Gabber, *Some theorems on Azumaya algebras*, The Brauer group, Lecture Notes in Math, vol. 844, Springer, Berlin-New York, 1981, pp. 129–209.
- [Pa] Ivan Panin, *Rationally isotropic quadratic spaces are locally isotropic*, Invent. Math. **176** (2009), 397–403.
- [Sw] Richard G. Swan, *Néron-Popescu desingularization*, Algebra and geometry (Taipei, 1995), Lect. Algebra, vol. 2, Int. Press, Cambridge, MA, 1998, pp. 135–192.

Chemin de la Raye 11, CH-
1024 Écublens