

# Contents

<b>1</b>	<b>Introduction</b>	1
<b>2</b>	<b>Basic frameworks and key constants</b>	13
2.1	Framework	13
2.2	Conductance constant	21
2.3	Combinatorial modulus	24
2.4	Neighbor disparity constant	28
<b>3</b>	<b>Conductive homogeneity and its consequences</b>	35
3.1	Construction of $p$ -energy: $p > \dim_{AR}(K, d)$	35
3.2	Construction of $p$ -energy: $p \leq \dim_{AR}(K, d)$	49
3.3	Conductive homogeneity	51
<b>4</b>	<b>Conductive homogeneity of self-similar sets</b>	57
4.1	Self-similar sets and self-similarity of energy	57
4.2	Conductive homogeneity of self-similar sets	62
4.3	Subsystems of (hyper)cubic tiling	64
4.4	Examples: subsystems of (hyper)cubic tiling	76
4.5	Rationally ramified Sierpiński crosses	81
4.6	Nested fractals	88
<b>5</b>	<b>Knight move implies conductive homogeneity</b>	95
5.1	Conductance and Poincaré constants	95
5.2	Relations of constants	99
5.3	Proof of Theorem 3.33	104
<b>6</b>	<b>Miscellanea</b>	107
6.1	Uniformity of constants	107
6.2	Modification of the structure of a graph	111
6.3	Open problems	112
<b>Appendices</b>	115	
A	Basic inequalities	115
B	Basic facts on $p$ -energy	115
C	Useful facts on combinatorial modulus	117
D	An Arzelà–Ascoli theorem for discontinuous functions	119

E Geometric properties of strongly symmetric self-similar sets . . . . .	120
F List of definitions and notations . . . . .	125
<b>References</b> . . . . .	127