

PERFECT BASES IN REPRESENTATION THEORY: THREE MOUNTAINS AND THEIR SPRINGS

JOEL KAMNITZER

ABSTRACT

In order to give a combinatorial descriptions of tensor product multiplicities for semisimple groups, it is useful to find bases for representations which are compatible with the actions of Chevalley generators of the Lie algebra. There are three known examples of such bases, each of which flows from geometric or algebraic mountain. Remarkably, each mountain gives the same combinatorial shadow: the crystal $B(\infty)$ and the Mirković–Vilonen polytopes. In order to distinguish between the three bases, we introduce measures supported on these polytopes. We also report on the interaction of these bases with the cluster structure on the coordinate ring of the maximal unipotent subgroup.

MATHEMATICS SUBJECT CLASSIFICATION 2020

Primary 22E46; Secondary 14M15, 16G20, 13F60, 05E10

KEYWORDS

Canonical basis, crystals, affine Grassmannian, quiver varieties

1. REPRESENTATIONS AND THEIR BASES

1.1. Semisimple Lie algebras and their representations

Let G be a complex semisimple group. The representation theory of G is very well understood. The irreducible G -representations are labeled by dominant weights, and every representation is a direct sum of these irreducible representations. For $\lambda \in P_+$, the irreducible representation $V(\lambda)$ admits a decomposition into eigenspaces $V(\lambda)_\mu$ for the action of T . These eigenspaces are called weight spaces and their dimensions are called *weight multiplicities*.

The tensor product of two irreducible representations decomposes into a direct sum of irreducible representations with *tensor product multiplicities* $c_{\lambda\mu}^v$,

$$V(\lambda) \otimes V(\mu) \cong \bigoplus_{v \in P_+} V(v)^{\oplus c_{\lambda\mu}^v}.$$

Problem 1.1. Determine combinatorial formulae for weight multiplicities and tensor product multiplicities.

Weight and tensor product multiplicities are closely related by the following construction. Let $C_{\lambda\mu}^v = \text{Hom}(V(v), V(\lambda) \otimes V(\mu))$, a vector space whose dimension is $c_{\lambda\mu}^v$.

Proposition 1.2. *There is an injective map $C_{\lambda\mu}^v \rightarrow V(\lambda)_{v-\mu}$ with image $\bigcap_{i \in I} \ker e_i^{\alpha_i^\vee(\lambda)+1}$.*

Here we use the Chevalley presentation of \mathfrak{g} , with generators e_i, f_i, α_i^\vee , for $i \in I$.

1.2. Good and perfect bases

Problem 1.1 was first solved by Littelmann [34] and Berenstein–Zelevinsky [9], following an approach first proposed by Gel'fand–Zelevinsky [21]. They suggested finding weight bases for each $V(\lambda)$ which restrict to bases of tensor product multiplicity spaces.

Let V be a G -representation. A *weight basis* for V is a basis consisting of weight vectors. A weight basis B for $V(\lambda)$ is called *good*, if for each $i \in I$, it is compatible with the filtration of $V(\lambda)$ given by the kernels of powers of e_i . From Proposition 1.2, it follows that a good basis restricts to a basis of each tensor product multiplicity space.

A slight strengthening of the notion of good basis was proposed by Berenstein–Kazhdan [8]. One might imagine that we could find a basis for a representation such that each e_i takes each basis vector to another basis vector (or 0). However, this is not always possible (see Example 1.4). So instead we will demand that each e_i permutes the basis up to lower order terms.

To formulate this, we define a map $\varepsilon_i : V \rightarrow \mathbb{N}$ giving the nilpotence degree of e_i on a vector $v \in V$; more precisely, $\varepsilon_i(v) = \max\{n \in \mathbb{N} : e_i^n v \neq 0\}$.

A good basis B of V is called *perfect*, if for each $i \in I$, and $b \in B$, either $e_i b = 0$ or there exists $\tilde{e}_i(b) \in B$ such that

$$e_i b = \varepsilon_i(b) \tilde{e}_i(b) + v \quad \text{for some } v \in \ker e_i^{\varepsilon_i(b)-1}.$$

In other words (up to a predictable scalar) $e_i b$ equals $\tilde{e}_i(b)$ modulo a vector with lower nilpotence degree. Note that this definition only requires V to be a representation of the Borel subalgebra \mathfrak{b} .

Example 1.3. To understand these scalars and gain some intuition, it is instructive to consider the case of $\mathfrak{g} = \mathfrak{sl}_2$. In this case, $P_+ = \mathbb{N}$ and $V(n) = \mathbb{C}[x, y]_n$, the space of homogeneous polynomials of degree n . The Chevalley generator e acts by $y\partial_x$ (on both the left and right) and the unique perfect basis (up to a scalar) is $\{x^n, x^{n-1}y, \dots, y^n\}$.

Note that $y\partial_x(x^k y^{n-k}) = kx^{k-1}y^{n-k+1}$ and $\varepsilon(x^k y^{n-k}) = k$. In this case there is no lower order term.

Example 1.4. The simplest irreducible representation where lower order terms occur is the adjoint representation of \mathfrak{sl}_3 . In this representation, $V = \mathfrak{sl}_3$ with the action given by matrix commutator. If we assume that B contains the highest weight vector

$$\begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix},$$

then it is easy to see that the perfect basis condition forces B to contain

$$\begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix}.$$

The choice of basis for the diagonal matrices is more interesting. The requirement that B be compatible with the kernels of e_1, e_2 forces B to contain matrices of the form

$$\begin{pmatrix} -a & 0 & 0 \\ 0 & -a & 0 \\ 0 & 0 & 2a \end{pmatrix}, \begin{pmatrix} 2b & 0 & 0 \\ 0 & -b & 0 \\ 0 & 0 & -b \end{pmatrix}$$

for some nonzero a, b . We are forced to take $b = 1/3$ and similarly $a = 1/3$, since

$$e_1 \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix} = 2 \begin{pmatrix} 2/3 & 0 & 0 \\ 0 & -1/3 & 0 \\ 0 & 0 & -1/3 \end{pmatrix} + \begin{pmatrix} -1/3 & 0 & 0 \\ 0 & -1/3 & 0 \\ 0 & 0 & 2/3 \end{pmatrix},$$

where the second term is of lower nilpotence degree (since it lies in the kernel of e_1).

1.3. Perfect bases and crystals

Any perfect basis gives rise to a combinatorial structure called a crystal. Crystals were first introduced by Kashiwara [30] as the $q = 0$ limit of a basis for a representation of a quantum group. However, we prefer to view them as recording the leading order behavior of e_i acting on a perfect basis.

A crystal is a finite set B , along with a map $\text{wt} : B \rightarrow P$, and for each $i \in I$, a partially defined map $\tilde{e}_i : B \rightarrow B$. If B is a perfect basis, then it automatically acquires a crystal structure. The following result of Berenstein–Kazhdan [8, THEOREM 5.37] shows that this combinatorial structure depends only on the representation.

Theorem 1.5. *Let V be a representation and let B, B' be two perfect bases. Then there exists a bijection $B \cong B'$ which is an isomorphism of crystals.*

Because of this theorem, we may speak of *the* crystal of a representation. In particular, the crystal of $V(\lambda)$ is denoted $B(\lambda)$. Many different explicit combinatorial realizations of $B(\lambda)$ are possible. In this talk, we will focus on MV polytopes Section 2.3.

1.4. Biperfect bases

Rather than looking at each irreducible representation individually, we can study them all at once, using the following trick. The maximal unipotent subgroup N has left and right actions of N and thus the coordinate ring $\mathbb{C}[N]$ has left and right actions of \mathfrak{n} by differential operators. For each $i \in I$, we write $e_i : \mathbb{C}[N] \rightarrow \mathbb{C}[N]$ for the left action and $e_i^* : \mathbb{C}[N] \rightarrow \mathbb{C}[N]$ for the right action.

For each $\lambda \in P_+$, choose a highest weight vector $v_\lambda \in V(\lambda)$ and let $v_\lambda^* : V(\lambda) \rightarrow \mathbb{C}$ be a dual linear form. We define an N -equivariant map

$$\Psi_\lambda : V(\lambda) \rightarrow \mathbb{C}[N], \quad \Psi_\lambda(v)(g) = v_\lambda^*(gv).$$

This linear map is injective and its image is

$$\text{im } \Psi_\lambda = \bigcap_{i \in I} \ker(e_i^*)^{\alpha_i^\vee(\lambda)+1} \subset \mathbb{C}[N]. \tag{1.1}$$

Thus a basis for $\mathbb{C}[N]$ compatible with the kernels of all powers of e_i^* gives a basis for each $V(\lambda)$. Conversely, a collection of bases for each $V(\lambda)$ can sometimes glue together to give a basis for $\mathbb{C}[N]$.

A basis B of $\mathbb{C}[N]$ is called *biperfect* if it contains 1, and it is perfect with respect to both the left and right actions of \mathfrak{n} . Thus, B will have two families of crystal operators, written $\tilde{e}_i, \tilde{e}_i^*$ and two families of maps $\varepsilon_i, \varepsilon_i^* : B \rightarrow \mathbb{N}$ (but only one weight map).

From Proposition 1.2 and (1.1), we immediately deduce the following corollary, which can be regarded as a generalization (from the canonical basis to arbitrary biperfect bases) of [9, COROLLARY 3.4].

Corollary 1.6. *Let B be a biperfect basis of $\mathbb{C}[N]$.*

(1) *For any $\lambda \in P_+$, the set $\{b \in B : \varepsilon_i^*(b) \leq \alpha_i^\vee(\lambda)\}$ restricts via Ψ_λ to a perfect basis for $V(\lambda)$.*

(2) *For any $\lambda, \mu, \nu \in P_+$, the set*

$$\{b \in B : \text{wt}(b) = \nu - \mu - \lambda \text{ and } \forall i \in I, \varepsilon_i(b) \leq \alpha_i^\vee(\mu), \varepsilon_i^*(b) \leq \alpha_i^\vee(\lambda)\}$$

restricts to a basis for $C_{\lambda\mu}^\nu$.

Thus we can solve Problem 1.1 by understanding well the bicrystal structure on B .

1.5. The bicrystal $B(\infty)$


The Berenstein–Kazhdan result (Theorem 1.5) generalizes to biperfect bases.

Theorem 1.7 ([5, THEOREM 2.4]). *Let B and B' be two biperfect bases of $\mathbb{C}[N]$. Then there is a unique bijection $B \cong B'$ that respects the bicrystal structure.*

The abstract combinatorial crystal underlying any biperfect basis is denoted $B(\infty)$. On $B(\infty)$ we have the Kashiwara involution $*$: $B(\infty) \rightarrow B(\infty)$ exchanging \tilde{e}_i and \tilde{e}_i^* .

Remark 1.8. The algebra $\mathbb{C}[N]$ has an involutive automorphism $*$ (coming from the inverse map on N) which exchanges the left and right actions of \mathfrak{n} . A $*$ -invariant perfect basis is a perfect basis which is invariant under $*$. Every example of a biperfect basis that we know is $*$ -invariant. The Kashiwara involution $*$ on $B(\infty)$ is the combinatorial manifestation of the involution $*$ on $\mathbb{C}[N]$.

Example 1.9. When $G = \mathbf{SL}_3$, $B(\infty)$ can be drawn in the following way. The action of e_1 is given by right-pointing diagonal arrows and the action of e_2 is given by the left-pointing ones. Each horizontal group of dots have the same weight and the Kashiwara involution flips each such group. We would like to thank Mark Haiman for showing us this drawing many years ago.



1.6. Biperfect bases in small rank

For small rank groups, it is easy to show the existence and uniqueness of biperfect bases of $\mathbb{C}[N]$ by elementary means.

Theorem 1.10. *For $G = \mathbf{SL}_2, \mathbf{SL}_3, \mathbf{SL}_4$, $\mathbb{C}[N]$ has a unique biperfect basis.*

Example 1.11. Suppose $G = \mathbf{SL}_2$, then $\mathbb{C}[N] = \mathbb{C}[x]$ and $\Psi_n : \mathbb{C}[x, y]_n \rightarrow \mathbb{C}[x]$ is the map sending y to 1. The left and right actions of $e \in \mathfrak{n}$ on $\mathbb{C}[x]$ agree and are given by $e = \partial_x$. The unique biperfect basis of $\mathbb{C}[x]$ is $\{1, x, x^2, \dots\}$.

Example 1.12. Suppose $G = \mathbf{SL}_3$, with the standard choice for B , T and N . Then $\mathbb{C}[N] = \mathbb{C}[x, y, z]$ where x , y and z are the three matrix entries of an upper unitriangular matrix

$$\begin{pmatrix} 1 & x & z \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix} \in N.$$

The unique bipерfect basis of $\mathbb{C}[N]$ is

$$B = \{x^a z^b (xy - z)^c : (a, b, c) \in \mathbb{N}^3\} \cup \{y^a z^b (xy - z)^c : (a, b, c) \in \mathbb{N}^3\}.$$

1.7. Three different bipерfect bases

For general G , bipерfect bases are not unique, nor is it very easy to show their existence.

The first example of a bipерfect basis was Lusztig's *dual canonical basis* which is also known as Kashiwara's upper global basis [31, 37]. This is actually a basis for the corresponding quantum deformation of $\mathbb{C}[N]$, but it can be specialized at $q = 1$ to give a bipерfect basis.

Another example, when \mathfrak{g} is simply-laced, is Lusztig's *dual semicanonical basis* [38], which is constructed by means of the representation theory of the preprojective algebra.

A third example is the *Mirković–Vilonen basis* [39] coming from the geometry of the affine Grassmannian.

This trichotomy of bases will be the focus of this paper. Each of these bases comes from a complicated algebraic or geometric source. Following Arun Ram, we can imagine three high mountains whose springs give these three bases.

These bases are all different, combining [5, THM. 1.7], [4, PROP. 2.7], and [18, (3)].

Theorem 1.13. *For $G = \mathbf{SL}_6$ in the weight space $2\alpha_1 + 4\alpha_2 + 4\alpha_3 + 4\alpha_4 + 2\alpha_5$, and for $G = \mathbf{SO}_8$ in the weight space $2\alpha_1 + 4\alpha_2 + 2\alpha_3 + 2\alpha_4$, there is a point of $B(\infty)$ whose corresponding dual canonical, dual semicanonical, and MV bases are all different.*

Moreover, in both these examples, we have the following specific situation:

$$d = b + v, \quad c = b + 2v, \tag{1.2}$$

where b, c, d denote the MV, dual semicanonical, and dual canonical basis vectors, all of which define the same point in $B(\infty)$, and v denotes a vector common to all three bases.

Question 1.14. What can we say about the set of all bipерfect bases of $\mathbb{C}[N]$ for a fixed G ?

2. MIRKOVIĆ–VILONEN BASIS

2.1. MV cycles

Mirković–Vilonen [39] used the geometric Satake correspondence to define the MV basis for irreducible representations of G . This basis is indexed by certain subvarieties in the affine Grassmannian, known as MV cycles.

Let G^\vee be the Langlands dual group and let $\text{Gr} = G^\vee((t))/G^\vee[[t]]$ denote the affine Grassmannian of this group. By definition, the coweight lattice of G^\vee coincides with the weight lattice P of G . For each coweight $\mu \in P$, we get a point of $G^\vee((t))$ and hence a point L_μ in Gr . Let $S_\pm^\mu := N_\pm^\vee((t))L_\mu$ denote semiinfinite orbits in Gr , where N_\pm^\vee denote opposite unipotent subgroups in G^\vee .

For $\lambda \in P_+$, let $\text{Gr}^\lambda := \overline{G^\vee[[t]]L_\lambda}$ be a *spherical Schubert variety*. This is a finite-dimensional singular projective variety whose geometry is closely related to the irreducible representation $V(\lambda)$. Let $P(\text{Gr})$ denote the category of perverse sheaves on Gr which are constructible with respect to the stratification by $G^\vee[[t]]$ orbits. This is a semisimple category whose simple objects are the intersection cohomology sheaves IC_λ of the spherical Schubert varieties. There is a monoidal structure on $P(\text{Gr})$ by convolution.

The following geometric Satake correspondence was established by Mirković–Vilonen [39], following earlier work by Lusztig [35] and Ginzburg [23].

Theorem 2.1. (1) *There is an equivalence of monoidal categories, $P(\text{Gr}) \cong \text{Rep } G$.*

(2) *Under this equivalence, for each $\lambda \in P_+$, IC_λ is sent to $V(\lambda)$.*

(3) *Under this equivalence, for each $\mu \in P$, the hyperbolic stalk functor $H_{S_\pm^\mu}^\bullet(-)$ matches the functor of taking the μ -weight space.*

Combining these statements, we conclude $H_{\text{top}}(\text{Gr}^\lambda \cap S_\pm^\mu) \cong V(\lambda)_\mu$. The irreducible components of $\text{Gr}^\lambda \cap S_\pm^\mu$ are called *MV cycles*. Via this theorem, they provide a basis for each $V(\lambda)_\mu$.

2.2. Stable MV cycles

For bases of $\mathbb{C}[N]$, we will be concerned with the intersection of opposite semiinfinite orbits. For any $\nu \in Q_+$, the positive root cone, the irreducible components of $S_+^\nu \cap S_-^0$ are called *stable MV cycles*.

Given an MV cycle $Z \subset \text{Gr}^\lambda \cap S_-^\mu$, we can translate by $t^{-\mu}$ to produce a stable MV cycle $t^{-\mu}Z$. This process is the geometric analog of the map $\Psi_\lambda : V(\lambda) \rightarrow \mathbb{C}[N]$.

In [5], we combined work of Ginzburg [23] and Mirković–Vilonen [39] to prove the following result, which had been conjectured by Anderson [1].

Theorem 2.2. (1) *The MV bases for each $V(\lambda)$ can be collected together to form a bipermut basis for $\mathbb{C}[N]$, which is indexed by stable MV cycles.*

(2) *For each i , the action of e_i on an MV basis vector b_Z is given by the intersection of the stable MV cycle Z with a hyperplane section.*

(3) *Given two MV cycles Z_1, Z_2 , the product $b_{Z_1}b_{Z_2}$ in $\mathbb{C}[N]$ is given by the Beilinson–Drinfeld degeneration of $Z_1 \times Z_2$.*

In particular, the structure constants for the action of e_i and for the multiplication are nonnegative integers.

2.3. MV polytopes

For each stable MV cycle Z , we define $\text{Pol}(Z)$ to be its moment map image (for a real Hamiltonian torus action). Equivalently, we have

$$\text{Pol}(Z) = \text{Conv}(\mu : L_\mu \in Z).$$

The polytopes produced this way are called MV polytopes. In [27], we proved the following result.

Theorem 2.3. *The map $Z \mapsto \text{Pol}(Z)$ gives a bijection between the stable MV cycles and the MV polytopes. The MV polytopes are precisely those lattice polytopes whose dual fan is a coarsening of the Weyl fan and whose 2-faces are MV polygons for the appropriate rank 2 groups (which can be described explicitly).*


This theorem was reinterpreted by Goncharov–Shen [24] as the following statement.

Corollary 2.4. *The MV polytopes are in natural bijection with $(G^\vee / B^\vee)(\mathbb{Z}_{\text{trop}})_{\geq}$, the non-negative tropical points of the flag variety.*

Following the historical order, we have described MV polytopes as the moment map images of MV cycles. However, we emphasize that Theorem 2.3 shows that they are purely combinatorial objects. We will see in the next two sections that these same polytopes are naturally obtained from general preprojective algebras modules and simple KLR modules. They are the common shadows from all three mountains.

In [26], we gave an explicit description of the crystal structure on the set of MV polytopes. This provides a convenient combinatorial framework for describing the crystal $B(\infty)$ and is easily connected to many other combinatorial models. In particular, for each reduced word $s_{i_1} \cdots s_{i_m} = w_0$ for the longest element of the Weyl group, Lusztig [36] constructed a bijection $B(\infty) \rightarrow \mathbb{N}^m$ using the relation between PBW monomials and the canonical basis. In [26], we showed that the Lusztig datum of $b \in B(\infty)$ is the list of lengths along a path following the edges of the MV polytope $\text{Pol}(b)$, in root directions determined by the reduced word.

Example 2.5. Take $G = \text{SL}_3$. In this case, an MV polytope is a hexagon with all 120° angles, whose “width” A is equal to the maximum of its two “heights” B, C .



For this polytope, the two Lusztig data are $(3, 2, 1)$ and $(2, 1, 4)$.

3. DUAL SEMICANONICAL BASIS

3.1. Preprojective algebra

Assume for this section that \mathfrak{g} is simply-laced. Let H denote the set of oriented edges of the Dynkin diagram of \mathfrak{g} . If $h = (i, j)$, write $\bar{h} = (j, i)$. Fix a map $\tau : H \rightarrow \{1, -1\}$ such that for each h , $\tau(h) + \tau(\bar{h}) = 0$ (such a τ corresponds to an orientation of each edge of the Dynkin diagram).

The *preprojective algebra* Λ is the quotient of the path algebra of (I, H) by the relation $\sum_{h \in H} \tau(h) h \bar{h} = 0$. So a Λ -module M consists of vector spaces M_i , for $i \in I$, and linear maps $M_h : M_i \rightarrow M_j$ for each $h = (i, j) \in H$, such that

$$\sum_{h \in H} \tau(h) M_h M_{\bar{h}} = 0. \quad (3.1)$$

Given a Λ -module M , we define its dimension vector by

$$\underline{\dim} M = \sum_{i \in I} (\dim M_i) \alpha_i.$$

We write S_i for the simple module at vertex i , the unique module with $\underline{\dim} S_i = \alpha_i$.

For each $v = \sum_{i \in I} v_i \alpha_i \in Q_+$, we consider the affine variety of Λ -module structures on $\bigoplus_{i \in I} \mathbb{C}^{v_i}$. More precisely, we define

$$\Lambda(v) \subset \bigoplus_{(i,j) \in H} \text{Hom}(\mathbb{C}^{v_i}, \mathbb{C}^{v_j})$$

to be the subvariety defined by equation (3.1).

3.2. The dual semicanonical basis

Let M be a Λ -module. Following Lusztig [38] and Geiss–Leclerc–Schröer [18, §5], we define an element $\xi_M \in \mathbb{C}[N]$ as follows. First, for each $\underline{i} \in I^p$, we define the projective variety of composition series of type \underline{i} ,

$$F_{\underline{i}}(M) = \{0 = M^0 \subset M^1 \subset \cdots \subset M^p = M : M^k / M^{k-1} \cong S_{i_k} \text{ for all } k\}$$

and then we define $\xi_M \in \mathbb{C}[N]$ by requiring that

$$\langle e_{i_1} \cdots e_{i_p}, \xi_M \rangle = \chi(F_{\underline{i}}(M))$$

for any $\underline{i} \in I^p$, where χ denotes topological Euler characteristic, and where $\langle \cdot, \cdot \rangle$ denotes the pairing between $U\mathfrak{n}$ and $\mathbb{C}[N]$.

This map $M \mapsto \xi_M$ is constructible and so for any irreducible component $Y \subset \Lambda(v)$, we can define $c_Y \in \mathbb{C}[N]_v$ by setting $c_Y = \xi_M$, for M a general point in Y .

The following result is due to Lusztig [38].

- Theorem 3.1.**
- (1) For each $v \in Q_+$, $\{c_Y \mid Y \in \text{Irr } \Lambda(v)\}$ is a basis for $\mathbb{C}[N]_v$.
 - (2) Together they form a biperfect basis of $\mathbb{C}[N]$, called the dual semicanonical basis.

3.3. Polytopes from preprojective algebra modules

In the resulting bicrystal structure on the set $\sqcup_v \text{Irr } \Lambda(v)$, we have

$$\varepsilon_i(Y) = \dim \text{Hom}_\Lambda(M, S_i), \quad \varepsilon_i^*(Y) = \dim \text{Hom}_\Lambda(S_i, M),$$

where M is a general point of Y .

Remark 3.2. Fix $\lambda \in P_+$. From Corollary 1.6(1), those components Y which satisfy $\varepsilon_i^*(Y) \leq \alpha_i^*(\lambda)$ index a basis for $V(\lambda)$. These same irreducible components form the core of the corresponding Nakajima quiver variety via the correspondence explained in [43, SECTION 4.6].

The bicrystal $\sqcup \text{Irr } \Lambda(v)$ is isomorphic to $B(\infty)$ by Theorem 1.7. Thus, an MV polytope is canonically associated to component Y . We can describe this polytope using the module structure on a general point of Y .

Theorem 3.3 ([6, §1.3]). *Let Y be a component of $\Lambda(v)$ and M be a general point of Y . The MV polytope of the basis vector c_Y is given by the Harder–Narasimhan polytope of M ,*

$$\text{Pol}(M) := \text{Conv}(\underline{\dim} N : N \subseteq M \text{ is a submodule}).$$

Example 3.4. Take $\mathfrak{g} = \mathfrak{sl}_3$, and consider $v = \alpha_1 + \alpha_2$.

Then $\Lambda(v) = \{(a, b) \in \mathbb{C}^2 : ab = 0\}$ where (a, b) corresponds to the Λ -module

$$\mathbb{C} \begin{array}{c} b \\ \xleftrightarrow{\quad} \\ a \end{array} \mathbb{C}.$$

Further, $\Lambda(v)$ has two components in this case. If Y is the component given by $b = 0$, then for general $M \in Y$, we have submodules in M of dimension 0, α_1 , $\alpha_1 + \alpha_2$ and so $\text{Pol}(M)$ is the triangle with these vertices.

4. DUAL CANONICAL BASES

4.1. KLR algebras

Let \mathfrak{g} be an arbitrary semisimple Lie algebra. For each $v \in Q_+$, Khovanov–Lauda–Rouquier defined an algebra R_v . It can either be defined by generators and relations using a modification of the presentation of a Hecke algebra [42], or as an algebra of decorated string diagrams as in [32]. This algebra can also be realized as an Ext algebra of certain perverse sheaves constructed by Lusztig (see [45]).

In this section, we work over \mathbb{C} , though it is possible to work over fields of positive characteristic; this produces different bases, called the dual p -canonical bases.

The algebra R_v contains idempotents $e_{\underline{i}}$ indexed by sequences $(i_1, \dots, i_p) \in I^p$ such that $\alpha_{i_1} + \dots + \alpha_{i_p} = v$. We write $K(R_v)_{\mathbb{C}}$ for the complexified Grothendieck group of finite-dimensional R_v modules. The following result is due independently to Rouquier and Khovanov–Lauda.

Theorem 4.1. *For each $v \in Q_+$, there is an isomorphism $K(R_v)_{\mathbb{C}} \cong \mathbb{C}[N]_v$, written $[L] \mapsto d_L$ such that $\langle e_{\underline{i}}, d_L \rangle = \dim e_{\underline{i}}L$, for any module L and \underline{i} as above.*

The vector space $K(R_\nu)_\mathbb{C}$ has a basis given by the simple finite-dimensional R_ν -modules. By [45], the resulting basis for $\mathbb{C}[N]$ coincides with Lusztig’s dual canonical basis.

4.2. Polytopes from KLR modules

The bicrystal structure on the set of simple KLR modules was carefully studied by Lauda–Vazirani [33]. By Theorem 1.7, this bicrystal is isomorphic to $B(\infty)$, and thus an MV polytope is canonically associated to each simple KLR module.

On the other hand, these algebras come with nonunital morphisms

$$R_\mu \otimes R_{\nu-\mu} \rightarrow R_\nu.$$

We write $e_{\mu,\nu-\mu}$ for the image of the identity under this map. Tingley–Webster [44] used these morphisms to define a polytope associated to each KLR module.

Theorem 4.2. *The MV polytope of a simple R_ν -module L is the character polytope*

$$\text{Pol}(L) := \text{Conv}(\mu : e_{\mu,\nu-\mu}L \neq 0).$$

4.3. Generalizations to affine and Kac–Moody cases

Unlike the MV basis, the dual semicanonical basis and dual canonical basis admit straightforward generalizations to the setting where \mathfrak{g} is a symmetric (resp. symmetrizable) Kac–Moody Lie algebra.

The polytopes $\text{Pol}(M)$ and $\text{Pol}(L)$ associated to a general Λ -module or a simple KLR module admit obvious generalizations in this setting. However, due to higher root multiplicities, the polytope is not enough to characterize the point in $B(\infty)$. Thus, we must enhance the polytope with some extra information. This was carried out in [6] (using Λ -modules) and in [44] (using KLR modules).

In the affine case, this extra information consists of partitions associated to vertical edges of the polytope (vertical edges are those pointing in the imaginary root direction). Moreover, these decorated polytopes are characterized by their 2-faces (as in the finite case, Theorem 2.3) and the new relevant polygons were described combinatorially in [3].

- Question 4.3.**
- (1) Is it possible to give a “tropical” description of affine MV polytopes, similar to Corollary 2.4?
 - (2) Outside of the affine type, it is possible to give a combinatorial description to the extra information carried on the MV polytope?
 - (3) Though more complicated, the theory of MV cycles exists for affine Kac–Moody Lie algebras. How can we relate these MV cycles to the affine MV polytopes? In particular, what information about the cycles is encoded in the partitions along vertical edges?

5. COMPARING BIPERFECT BASES

5.1. Change of basis matrix

Let B, B' be two biperfect bases for $\mathbb{C}[N]$. By Theorem 1.7, we obtain bijections $B \rightarrow B(\infty) \leftarrow B'$ and thus a bijection between B and B' . Thus, it makes sense to speak of the change of basis matrix between B and B' . In [2], Baumann proved that this matrix is upper unitriangular with respect to a partial order on $B(\infty)$, defined combinatorially using the crystal structure. Many elements of $B(\infty)$ are incomparable using this order. Thus, many off-diagonal elements of the change of basis matrix must vanish. In low rank, the order becomes trivial and gives the proof of Theorem 1.10.

5.2. Measures

We think of an MV polytope as the shadow of a biperfect basis vector. Unfortunately, this shadow is not precise enough to distinguish between different biperfect basis vectors which represent the same element of $B(\infty)$. For this reason, we now introduce a measure supported on the MV polytope.

Consider the vector space $\text{Dist}(\mathfrak{t}_{\mathbb{R}}^*)$ of \mathbb{C} -valued compactly supported distributions on $\mathfrak{t}_{\mathbb{R}}^*$. It forms an algebra under convolution, using the addition map $\mathfrak{t}_{\mathbb{R}}^* \times \mathfrak{t}_{\mathbb{R}}^* \xrightarrow{+} \mathfrak{t}_{\mathbb{R}}^*$.

Let $\Delta^p := \{(c_0, \dots, c_p) \in \mathbb{R}^{p+1} : \text{each } c_i \geq 0, c_0 + \dots + c_p = 1\}$ be the standard p -simplex. For $\underline{i} \in I^p$, we define the linear map $\pi_{\underline{i}} : \mathbb{R}^{p+1} \rightarrow \mathfrak{t}_{\mathbb{R}}^*$ by

$$\pi_{\underline{i}}(c_0, \dots, c_p) = \sum_{k=0}^p c_k (\alpha_{i_1} + \dots + \alpha_{i_k}).$$

We define the measure $D_{\underline{i}}$ on $\mathfrak{t}_{\mathbb{R}}^*$ by $D_{\underline{i}} := (\pi_{\underline{i}})_*(\delta_{\Delta^p})$, the push-forward of Lebesgue measure on the p -simplex. The measures $D_{\underline{i}}$ satisfy the shuffle identity.

Lemma 5.1 ([5, LEMMA 8.5]). *For $\underline{j} \in I^p, \underline{k} \in I^q$,*

$$D_{\underline{j}} * D_{\underline{k}} = \sum_{\underline{i} \in \underline{j} \sqcup \underline{k}} D_{\underline{i}},$$

where $\underline{j} \sqcup \underline{k}$ is the set of all sequences obtained by shuffling \underline{j} and \underline{k} .

Elementary considerations involving the coproduct structure on $U\mathfrak{n}$ show that the shuffle identity implies that there is an algebra morphism $D : \mathbb{C}[N] \rightarrow \text{Dist}(\mathfrak{t}_{\mathbb{R}}^*)$ defined by

$$D(f) = \sum_{\underline{i}} \langle e_{\underline{i}}, f \rangle D_{\underline{i}}.$$

5.3. Fourier transform

For each weight $\beta \in P$, we define e^{β} to be the function $x \mapsto e^{\langle \beta, x \rangle}$ on $\mathfrak{t}_{\mathbb{C}}$. Given a measure $D(f)$ as above, we can consider its Fourier transform $\text{FT}(D(f))$ which lies in the space of meromorphic functions on $\mathfrak{t}_{\mathbb{C}}$, spanned by these exponentials over the field $\mathbb{C}(t)$ of rational functions. In this way, we obtain the following result.

Proposition 5.2. *The composition $\text{FT} \circ D$ defines an algebra morphism*

$$\mathbb{C}[N] \rightarrow \mathbb{C}(t) \otimes \mathbb{C}[T].$$

This algebra morphism defines a rational map $\mathfrak{t} \times T \rightarrow N$. This map is actually regular on $\mathfrak{t}^{\text{reg}} \times T$, where $\mathfrak{t}^{\text{reg}}$ is the complement of the root hyperplanes in \mathfrak{t} .

Theorem 5.3 ([5, THEOREM 8.11]). (1) For all $x \in \mathfrak{t}^{\text{reg}}$, there exists a unique $n_x \in N$ such that $\text{Ad}_{n_x}(x) = x + e$.

(2) The rational map from Proposition 5.2 is given by $(x, t) \mapsto t^{-1}n_x t n_x^{-1}$.

We now study a simpler invariant \overline{D} . For a sequence $\underline{i} = (i_1, \dots, i_p)$, we define

$$\overline{D}_{\underline{i}} = \prod_{k=1}^p \frac{1}{\alpha_{i_k} + \dots + \alpha_{i_p}} \in \mathbb{C}(\mathfrak{t}).$$

Proposition 5.4 ([5, PROPOSITION 8.4 AND LEMMA 8.7]). (1) These rational functions satisfy the shuffle identity from Lemma 5.1 and thus define an algebra morphism $\overline{D} : \mathbb{C}[N] \rightarrow \mathbb{C}(\mathfrak{t})$ by $\overline{D}(f) = \sum_{\underline{i}} \langle e_{\underline{i}}, f \rangle \overline{D}_{\underline{i}}$.

(2) For any $f \in \mathbb{C}[N]$, $\overline{D}(f)$ is the coefficient of e^0 in $\text{FT}(D(f))$.

(3) The algebra morphism from (1) comes from the morphism of varieties $\mathfrak{t}^{\text{reg}} \rightarrow N$ given by $x \mapsto n_x$.

5.4. Duistermaat–Heckman measure

In symplectic geometry, the Duistermaat–Heckman (DH) measure of a symplectic manifold with a Hamiltonian torus action is defined to be the push-forward of the Liouville measure under the moment map. Brion–Procesi [14] reformulated this notion in algebraic geometry by considering the asymptotics of sections of equivariant line bundles.

Fix a W -invariant bilinear form on \mathfrak{t} (normalized so that short roots have length 1). This leads to a central extension of $G^\vee(\mathfrak{t})$ and thus an equivariant line bundle $\mathcal{O}(1)$ on Gr .

Let $Z \subset \text{Gr}$ be an MV cycle. The torus T^\vee acts on the space of sections $\Gamma(Z, \mathcal{O}(n))$. We consider $[\Gamma(Z, \mathcal{O}(n))]$ as a distribution on $\mathfrak{t}_{\mathbb{R}}^*$ by

$$[\Gamma(Z, \mathcal{O}(n))] = \sum_{\mu \in P} \dim \Gamma(Z, \mathcal{O}(n))_{\mu} \delta_{\mu}.$$

(Implicitly, we use the bilinear form to identify \mathfrak{t} and \mathfrak{t}^* .)

Let $\tau_n : \mathfrak{t}_{\mathbb{R}}^* \rightarrow \mathfrak{t}_{\mathbb{R}}^*$ be the automorphism given by scaling by $\frac{1}{n}$. The Duistermaat–Heckman measure of Z is defined to be the limit

$$\text{DH}(Z) := \lim_{n \rightarrow \infty} \frac{1}{n^{\dim Z}} (\tau_n)_* [\Gamma(Z, \mathcal{O}(n))]$$

within the space of distributions on $\mathfrak{t}_{\mathbb{R}}^*$. Note that each $(\tau_n)_* [\Gamma(Z, \mathcal{O}(n))]$ is supported on $\text{Pol}(Z)$, and hence so is $\text{DH}(Z)$.

Via the Fourier transform, $\text{DH}(Z)$ is closely related to the class of Z in the equivariant homology of the affine Grassmannian. The following ideas are not specific to the affine Grassmannian: they apply to any (ind-)projective variety equipped with a torus action having isolated fixed points. First, we have the localization theorem in equivariant homology (see, for example, [13]).

Theorem 5.5. *The inclusion $\mathrm{Gr}^{T^\vee} \rightarrow \mathrm{Gr}$ induces an isomorphism*

$$H_\bullet^{T^\vee}(\mathrm{Gr}^{T^\vee}) \otimes_{\mathbb{C}[\mathfrak{t}]} \mathbb{C}(\mathfrak{t}) \xrightarrow{\sim} H_\bullet^{T^\vee}(\mathrm{Gr}) \otimes_{\mathbb{C}[\mathfrak{t}]} \mathbb{C}(\mathfrak{t}).$$

Because of this theorem and using $\mathrm{Gr}^{T^\vee} = \{L_\mu : \mu \in P\}$, we can write

$$[Z] = \sum_{\mu \in P} m_\mu(Z) [\{L_\mu\}]$$

for unique $m_\mu(Z) \in \mathbb{C}(\mathfrak{t})$. Following Brion [13], we call $m_\mu(Z)$ the *equivariant multiplicity* of Z at L_μ . In [5], we proved the following result (again in the general context of projective varieties with torus actions), following ideas of Brion.

Theorem 5.6. *For any MV cycle Z ,*

$$\mathrm{FT}(\mathrm{DH}(Z)) = \sum_{\mu \in P} m_\mu(Z) e^\mu.$$

5.5. DH measures and measures from $\mathbb{C}[N]$

The T^\vee -equivariant homology of Gr was computed by Yun–Zhu [46]. They defined a commutative convolution algebra structure on $H_\bullet^{T^\vee}(\mathrm{Gr})$ and described this algebra using the geometric Satake correspondence.

Let $e = \sum e_i$ be a regular nilpotent element. We define the *universal centralizer space* to be

$$C := \{(x, b) \in \mathfrak{t} \times B : \mathrm{Ad}_b(x + e) = x + e\}.$$

Remark 5.7. For any $x \in \mathfrak{t}$, $x + e$ is regular and has centralizer contained in B . Thus our space C is the base change over $\mathfrak{t} \rightarrow \mathfrak{t}/W$ of the usual universal centralizer (often denoted J), as defined in, for example [10, §2.2].

From the definition, we have a map $C \rightarrow \mathfrak{t} \times T$ given by $(x, tn) \mapsto (x, t)$. The dual algebra map fits into the following diagram.

Theorem 5.8 ([46, PROP 3.3 AND 5.7]). *There is an isomorphism of algebras $\theta : \mathbb{C}[C] \rightarrow H_\bullet^{T^\vee}(\mathrm{Gr})$ making the following diagram commute:*

$$\begin{array}{ccc} \mathbb{C}[\mathfrak{t}] \otimes \mathbb{C}[T] & \xrightarrow{\sim} & H_\bullet^{T^\vee}(\mathrm{Gr}^{T^\vee}) \\ \downarrow & & \downarrow \text{Theorem 5.5} \\ \mathbb{C}[C] & \xrightarrow{\theta} & H_\bullet^{T^\vee}(\mathrm{Gr}) \end{array}$$

Recall the map $D : \mathbb{C}[N] \rightarrow \mathrm{Dist}(\mathfrak{t}_{\mathbb{R}}^*)$ defined in §5.2 and the map $\overline{D} : \mathbb{C}[N] \rightarrow \mathbb{C}(\mathfrak{t})$ defined in Section 5.3. Combining Theorems 5.3, 5.6, and 5.8, we proved the following in [5].

Corollary 5.9. *For any stable MV cycle Z ,*

$$D(b_Z) = \mathrm{DH}(Z), \quad \overline{D}(b_Z) = m_0(Z).$$

This corollary is very useful since the equivariant multiplicity $m_0(Z)$ is easily computed using computer algebra programs. In the appendix of [5] (written with C. Morton-Ferguson and A. Dranowski), we used this approach to establish part of Theorem 1.13.

5.6. Measures from preprojective algebra modules

Let M be a Λ -module of dimension vector ν . By the definition of ξ_M and the map D , we have that

$$D(\xi_M) = \sum_{\underline{i}} \chi(F_{\underline{i}}(M)) D_{\underline{i}}.$$

In the previous section (Corollary 5.9), we saw that the measure $D(b_Z)$ of an MV basis vector equals the asymptotics of sections of line bundles on Z . In a similar fashion, we will now explain that $D(\xi_M)$ can also be regarded as an asymptotic.

Consider the algebra $\Lambda[t] := \Lambda \otimes_{\mathbb{C}} \mathbb{C}[t]$. We define

$$\mathbb{G}_{\mu}(M[t]/t^n) = \{N \subset M \otimes \mathbb{C}[t]/t^n : N \text{ is a } \Lambda[t]\text{-submodule, } \underline{\dim} N = \mu\}.$$

We will record the information of the Euler characteristics of these varieties as an element of $\text{Dist}(t_{\mathbb{R}}^*)$ by

$$[H^{\bullet}(\mathbb{G}(M[t]/t^n))] = \sum_{\mu \in Q_+} \chi(\mathbb{G}_{\mu}(M[t]/t^n)) \delta_{\mu}.$$

Theorem 5.10 ([5, THEOREM 11.4 AND LEMMA 12.3]). *For any Λ -module M , we have*

$$D(\xi_M) = \lim_{n \rightarrow \infty} \frac{1}{n^{\dim M}} (\tau_n)_* [H^{\bullet}(\mathbb{G}(M[t]/t^n))].$$

5.7. A conjecture and symplectic duality

Suppose that $Y \in \text{Irr } \Lambda(\nu)$, with general point M , and Z is a stable MV cycle, such that $c_Y = b_Z$ (there are many such pairs, conjecturally). Then $D(c_Y) = D(b_Z)$. Via Theorem 5.10 and Corollary 5.9, both sides are the asymptotics of T^{\vee} -representations. So it is natural to expect equality before taking the limit. If we further assume that the odd cohomology of $\mathbb{G}_{\mu}(M[t]/t^n)$ vanishes, this implies that there is an isomorphism of T^{\vee} -representations,

$$\Gamma(Z, \mathcal{L}^{\otimes n}) \cong H^{\bullet}(\mathbb{G}(M[t]/t^n)), \quad \text{for all } n \in \mathbb{N}, \quad (5.1)$$

where T^{\vee} acts on the right-hand side through the decomposition $\mathbb{G}(M[t]/t^n) = \sqcup \mathbb{G}_{\mu}(M[t]/t^n)$.

The left-hand sides of (5.1) form the components of a graded algebra, so it is natural to search for a similar structure on the right-hand sides. After studying this question for some time, we are pessimistic about finding this algebra structure. On the other hand, $\mathbb{C}[Z] := \bigoplus_n \Gamma(Z, \mathcal{L}^{\otimes n})$ is also a module over $\mathbb{C}[\overline{S_+^{\nu}} \cap \overline{S_-^0}]$. We believe that such a module structure naturally exists for the direct sums of the right-hand side of (5.1).

Conjecture 5.11. (1) *For any preprojective algebra module M of dimension vector ν , $\bigoplus_{n \in \mathbb{N}} H^{\bullet}(\mathbb{G}(M[t]/t^n))$ carries the structure of a $\mathbb{C}[\overline{S_+^{\nu}} \cap \overline{S_-^0}]$ -module.*

(2) *When $b_Z = c_Y$ and M is a general point of Y , then there is an isomorphism (5.1) of $\mathbb{C}[\overline{S_+^{\nu}} \cap \overline{S_-^0}]$ -modules.*

This conjecture should be a manifestation of the symplectic duality between generalized affine Grassmannian slices and Nakajima quiver varieties. In particular, Braverman–Finkelberg–Nakajima [12] proved that a generalized affine Grassmannian slice is the Coulomb branch associated to the quiver gauge theory defining the corresponding Nakajima quiver variety. In [25], with Hilburn and Weekes, we developed a Springer theory for Coulomb branch algebras and proved a weak form of Conjecture 5.11 for those modules M which come from a representation of the undoubled quiver.

The symplectic singularity viewpoint is also a useful framework for thinking about our three bases. In particular, following the philosophy of Braden–Licata–Proudfoot–Webster [11], the MV cycles and quiver variety components can be categorified using category \mathcal{O} for quantizations of affine Grassmannian slices and quiver varieties, respectively. Moreover, these categories are closely related to categories of modules for KLR algebras [28]. From this perspective, the failures of these bases to agree with the dual canonical basis in Theorem 1.13 can be attributed to the nonirreducibility of the character varieties of simple objects in these categories.

6. CLUSTER STRUCTURES

6.1. Cluster structures on $\mathbb{C}[N]$

Cluster algebras were defined by Fomin–Zelevinsky in order to understand the dual canonical basis of $\mathbb{C}[N]$. A cluster algebra is a commutative algebra A with a distinguished collection of “clusters.” Each *cluster* consists of an algebraically independent subset $T = \{x_1, \dots, x_n\} \subset A$, such that $A \subset \mathbb{C}(x_1, \dots, x_n)$. We pass from one cluster to another using an “exchange procedure” which removes one of the x_i and replaces it with a certain rational function. A *cluster monomial* is a monomial in the variables in one cluster.

Berenstein–Fomin–Zelevinsky [7] proved that $\mathbb{C}[N]$ carries a cluster algebra structure. Every reduced word $s_{i_1} \cdots s_{i_m} = w_0$ for the longest element of the Weyl group gives us a cluster $T(i)$ (though these are not all the clusters).

Geiss–Leclerc–Schröer [19] established the following beautiful result explaining how the theory of preprojective algebras provides an additive categorification of the cluster algebra structure on $\mathbb{C}[N]$.

Theorem 6.1. *Assume that \mathfrak{g} is simply-laced.*

- (1) *A maximal rigid Λ -module T gives a cluster with cluster variables $x_1 = \xi_{T_1}, \dots, x_n = \xi_{T_n}$, where T_i are the distinct indecomposable summands of T .*
- (2) *Every cluster is of this form and all cluster monomials lie in the dual semicanonical basis.*
- (3) *The exchange relations in $\mathbb{C}[N]$ comes from short exact sequences in Λ -mod.*

On the other hand, Kang–Kashiwara–Kim–Oh [29] proved that the categories of R_ν -modules provide a monoidal categorification of the cluster algebra $\mathbb{C}[N]$. In particular, they proved the following result, which was obtained at around the same time by Qin [40].

Theorem 6.2. *Every cluster monomial in $\mathbb{C}[N]$ lies in the dual canonical basis.*

Together, Theorems 6.1 and 6.2 imply that the dual semicanonical and canonical bases contain many common elements, since they both contain all cluster monomials.

6.2. g -vectors

Fix a cluster $T = \{x_1, \dots, x_n\}$ in a cluster algebra A . Let $u \in A$ be a cluster monomial. Fomin–Zelevinsky [17] defined combinatorially the g -vector $g_T(u) \in \mathbb{Z}^n$ of u using a mutation procedure. In [15], Derksen–Weyman–Zelevinsky proved that this g -vector encodes the “leading monomial” appearing in u . In this case, we say u is g -pointed. As we vary the cluster T , the data of these $g_T(u)$ defines a tropical point in the Langlands dual cluster X variety, as studied by Fock–Goncharov [16].

The following observation is due to Genz–Koshevoy–Schumann [22, SECTION 6].

Proposition 6.3. *Let \underline{i} be a reduced word for w_0 , giving a cluster $T(\underline{i})$. Let u be a cluster monomial. Then $g_{T(\underline{i})}(u)$ agrees with the \underline{i} -Lusztig data of u , up to a simple linear change of coordinates.*

In this way, we see explicitly how the information of all these g -vectors is the same information as the MV polytope (this is also closely related to Corollary 2.4).

In the setting of the dual semicanonical basis, this can be generalized as follows. Let T be a maximal rigid Λ -module and let $M \in \Lambda\text{-mod}$. Geiss–Leclerc–Schröer [20] defined $g_T(M) \in \mathbb{Z}^n$ using homological algebra, extending the above notion of g -vector. Moreover, they proved that ξ_M is g -pointed in each cluster.

6.3. Theta basis

A cluster algebra that contains finitely many clusters is called *finite type*. The cluster algebra $\mathbb{C}[N]$ is of finite type only when $G = \mathbf{SL}_2, \mathbf{SL}_3, \mathbf{SL}_4, \mathbf{SL}_5$. When a cluster algebra A is not finite type, then the cluster monomials do not span A . It was a longstanding open problem to extend the set of cluster monomials to a basis for A . This problem was solved by the following remarkable theorem of Gross–Hacking–Keel–Kontsevich.

Theorem 6.4. *Let A be a cluster algebra, satisfying some hypotheses (which hold for $\mathbb{C}[N]$). There is a natural basis for A , called the theta basis, extending the set of cluster monomials. This theta basis is parametrized by the set of tropical points of the Langlands dual cluster X variety. Moreover, each theta basis element is g -pointed in each cluster.*

Combining Theorem 6.4 with Proposition 6.3 or Corollary 2.4, we get a natural parametrization of the theta basis of $\mathbb{C}[N]$ by $B(\infty)$. In Section 1.5, we saw that biperfect bases are parametrized by $B(\infty)$, so it is natural to ask the following.

Question 6.5. Is the theta basis for $\mathbb{C}[N]$ a biperfect basis?

6.4. Cluster structure on the MV basis

From Theorems 6.1 and 6.2 the dual semicanonical and dual canonical bases for $\mathbb{C}[N]$ contain all cluster monomials. In another direction, Qin [41] studied bases which are g -pointed in each cluster and gave a description of the set of all such bases. In particular, he showed that all such bases contain all the cluster monomials.

This motivates the following conjecture.

Conjecture 6.6. *The MV basis for $\mathbb{C}[N]$ contains all cluster monomials. Moreover, its elements are g -pointed in each cluster.*

Note that the conjecture would imply that the MV basis and dual semicanonical basis agree for \mathbf{SL}_5 which is not known.

As evidence for this conjecture, let us mention that the first counterexamples (in both \mathbf{SO}_8 and \mathbf{SL}_6) to the equality of the MV and dual semicanonical bases (see Theorem 1.13) occur for the square of the simplest basis element which is not a cluster monomial.

Baumann–Gaussent–Littelmann proved this conjecture for certain clusters.

Theorem 6.7 ([4, PROP 7.2]). *If a reduced word \underline{i} for w_0 satisfies a certain condition (which holds for all reduced words in small rank), then all cluster monomials in the cluster $T(\underline{i})$ lie in the MV basis.*

At the moment, we are far from Conjecture 6.6, but thinking about this conjecture motivates the following questions.

Question 6.8. (1) Is there a refinement of the notion of biperfect basis which would imply that such a basis contains all cluster monomials?

(2) What do cluster exchange relations correspond to geometrically? Which collections of MV cycles form clusters?

Finally, we close with the following wild conjecture.

Conjecture 6.9. *The MV and theta bases for $\mathbb{C}[N]$ coincide.*

We have three pieces of weak evidence for this conjecture. First, the way in which the MV, dual canonical, and dual semicanonical bases differ in $\mathbb{C}[N]$ for \mathbf{SL}_6 in (1.2) is very reminiscent of the way in which the theta, dual canonical, and generic bases differ for rank 2, affine type cluster algebras. Second, the construction of the MV and theta bases are both related to the geometry of loop spaces. Finally, the trichotomy of bases studied here seems to match the trichotomy of bases for cluster algebras, see [41].

ACKNOWLEDGMENTS

I would like to thank Pierre Baumann and Peter Tingley for many years of fruitful collaboration and friendship. I also thank Alexander Braverman, Michel Brion, Elie Casbi, Anne

Dranowski, Tom Dunlap, Michael Finkelberg, Stephane Gaussent, Justin Hilburn, Allen Knutson, Bernard Leclerc, Peter Littelmann, Calder Morton-Ferguson, Dinakar Muthiah, Ben Webster, Alex Weekes, and Harold Williams for collaboration and useful conversations. Finally, I would like to thank Arun Ram his three mountains analogy.

FUNDING

This work was partially supported by an NSERC Discovery Grant.

REFERENCES

- [1] J. E. Anderson, A polytope calculus for semisimple groups. *Duke Math. J.* **116** (2003), 567–588.
- [2] P. Baumann, The canonical basis and the quantum Frobenius morphism. 2012, arXiv:1201.0303.
- [3] P. Baumann, T. Dunlap, J. Kamnitzer, and P. Tingley, Rank 2 affine MV polytopes. *Represent. Theory* **17** (2013), 442–468.
- [4] P. Baumann, S. Gaussent, and P. Littelmann, Bases of tensor products and geometric Satake correspondence. 2020, arXiv:2009.00042.
- [5] P. Baumann, J. Kamnitzer, and A. Knutson, The Mirković–Vilonen basis and Duistermaat–Heckman measures. *Acta Math.* **227** (2021), 1–101.
- [6] P. Baumann, J. Kamnitzer, and P. Tingley, Affine Mirković–Vilonen polytopes. *Publ. Math. Inst. Hautes Études Sci.* **120** (2014), 113–205.
- [7] A. Berenstein, S. Fomin, and A. Zelevinsky, Cluster algebras III: upper bounds and double Bruhat cells. *Duke Math. J.* **126** (2005), 1–52.
- [8] A. Berenstein and D. Kazhdan, Geometric and unipotent crystals. II. From unipotent bicrystals to crystal bases. In *Quantum groups (Israel Math. Conf. Proc.)*, pp. 13–88, Contemp. Math. 433, Amer. Math. Soc., 2007.
- [9] A. Berenstein and A. Zelevinsky, Tensor product multiplicities, canonical bases and totally positive varieties. *Invent. Math.* **143** (2001), 77–128.
- [10] R. Bezrukavnikov, M. Finkelberg, and I. Mirković, Equivariant homology and K-theory of affine Grassmannians and Toda lattices. *Compos. Math.* **141** (2005), 746–768.
- [11] T. Braden, T. Licata, N. Proudfoot, and B. Webster, Quantizations of conical symplectic resolutions II: category O and symplectic duality. *Astérisque* **384** (2016), 75–179.
- [12] A. Braverman, M. Finkelberg, and H. Nakajima, Coulomb branches of $3d \mathcal{N} = 4$ quiver gauge theories and slices in the affine Grassmannian. *Adv. Theor. Math. Phys.* **23** (2019), no. 1, 75–166.
- [13] M. Brion, Equivariant Chow groups for torus actions. *Transform. Groups* **3** (1997), 225–267.

- [14] M. Brion and C. Procesi, Action d'un tore dans une variété projective. In *Operator algebras, unitary representations, enveloping algebras, and invariant theory (Paris, 1989)*, pp. 509–539, Progr. Math. 92, Birkhauser, Boston, 1990.
- [15] H. Derksen, J. Weyman, and A. Zelevinsky, Quivers with potentials and their representations II: applications to cluster algebras. *J. Amer. Math. Soc.* **23** (2010), no. 3, 749–790.
- [16] V. Fock and A. Goncharov, Cluster ensembles, quantization and the dilogarithm. *Ann. Sci. Éc. Norm. Supér.* **42** (2009), no. 6, 865–930.
- [17] S. Fomin and A. Zelevinsky, Cluster algebras IV: coefficients. *Compos. Math.* **143** (2007), 112–164.
- [18] C. Geiss, B. Leclerc, and J. Schröer, Semicanonical bases and preprojective algebras. *Ann. Sci. Éc. Norm. Supér.* **38** (2005), 193–253.
- [19] C. Geiss, B. Leclerc, and J. Schröer, Rigid modules over preprojective algebras. *Invent. Math.* **165** (2006), 589–632.
- [20] C. Geiss, B. Leclerc, and J. Schröer, Generic bases for cluster algebras and the Chamber ansatz. *J. Amer. Math. Soc.* **25** (2012), 21–76.
- [21] I. Gelfand and A. Zelevinsky, Polytopes in the pattern space and canonical bases for the irreducible representations of gl_3 . *Funct. Anal. Appl.* **19** (1985), 72–75.
- [22] V. Genz, G. Koshevoy, and B. Schumann, Polyhedral parametrizations of canonical bases and cluster duality. *Adv. Math.* **369** (2020).
- [23] V. Ginzburg, Perverse sheaves on a loop group and Langlands duality. 2000, arXiv:alg-geom/9511007.
- [24] A. Goncharov and L. Shen, Geometry of canonical bases and mirror symmetry. *Invent. Math.* **202** (2015), no. 2, 487–633.
- [25] J. Hilburn, J. Kamnitzer, and A. Weekes, BFN Springer theory. 2020, arXiv:2004.14998.
- [26] J. Kamnitzer, The crystal structure on the set of Mirković–Vilonen polytopes. *Adv. Math.* **215** (2007), 66–93.
- [27] J. Kamnitzer, Mirković–Vilonen cycles and polytopes. *Ann. of Math.* **171** (2010), 245–294.
- [28] J. Kamnitzer, P. Tingley, B. Webster, A. Weekes, and O. Yacobi, On category \mathcal{O} for affine Grassmannian slices and categorified tensor products. *Proc. Lond. Math. Soc.* **119** (2019), no. 5, 1179–1233.
- [29] S.-J. Kang, M. Kashiwara, M. Kim, and S.-j. Oh, Monoidal categorification of cluster algebras. *J. Amer. Math. Soc.* **31** (2018), 349–426.
- [30] M. Kashiwara, Crystalizing the q -analogue of universal enveloping algebras. *Comm. Math. Phys.* **133** (1990), no. 2, 249–260.
- [31] M. Kashiwara, Global crystal bases of quantum groups. *Duke Math. J.* **69** (1993), 455–485.
- [32] M. Khovanov and A. Lauda, A diagrammatic approach to categorification of quantum groups I. *Represent. Theory* **13** (2009), 309–347.

- [33] A. Lauda and M. Vazirani, Crystals from categorified quantum groups. *Adv. Math.* **228** (2011), 803–861.
- [34] P. Littelmann, The path model for representations of symmetrizable Kac–Moody algebras. In *Proc. ICM, Zürich, 1994*. Birkhäuser, Basel, 1995.
- [35] G. Lusztig, Singularities, character formulas and a q -analog of weight multiplicities. *Astérisque* **101–102** (1983), 208–229.
- [36] G. Lusztig, Canonical bases arising from quantized enveloping algebras. *J. Amer. Math. Soc.* **3** (1990), no. 2, 447–498.
- [37] G. Lusztig, Canonical bases arising from quantized enveloping algebras II. *Progr. Theoret. Phys. Suppl.* **102** (1990), 175–201.
- [38] G. Lusztig, Semicanonical bases arising from enveloping algebras. *Adv. Math.* **151** (2000), 129–139.
- [39] I. Mirković and K. Vilonen, Geometric Langlands duality and representations of algebraic groups over commutative rings. *Ann. of Math.* **166** (2007), 95–143.
- [40] F. Qin, Triangular bases in quantum cluster algebras and monoidal categorification conjectures. *Duke Math. J.* **166** (2017), no. 12, 2337–2442.
- [41] F. Qin, Bases for upper cluster algebras and tropical points. *J. Eur. Math. Soc. (JEMS)* (to appear).
- [42] R. Rouquier, 2-Kac–Moody Lie algebras. 2008, arXiv:0812.5023.
- [43] Y. Saito, Crystal bases and quiver varieties. *Math. Ann.* **324** (2002), 675–688.
- [44] P. Tingley and B. Webster, Mirković–Vilonen polytopes and Khovanov–Lauda–Rouquier algebras. *Compos. Math.* **152** (2016), 1648–1696.
- [45] M. Varagnolo and E. Vasserot, Canonical bases and KLR algebras. *J. Reine Angew. Math.* **659** (2011), 67–100.
- [46] Z. Yun and X. Zhu, Integral homology of loop groups via Langlands dual groups. *Represent. Theory* **15** (2011), 347–369.

JOEL KAMNITZER

Department of Mathematics, University of Toronto, Toronto, ON, Canada,
jkamnitz@math.toronto.edu

