

References

- [1] S. Albeverio and S. Kusuoka, [The invariant measure and the flow associated to the \$\Phi_3^4\$ -quantum field model](#). *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **20** (2020), no. 4, 1359–1427
- [2] H. Bahouri, J.-Y. Chemin, and R. Danchin, [Fourier analysis and nonlinear partial differential equations](#). Grundlehren Math. Wiss. 343, Springer, Heidelberg, 2011
- [3] N. Barashkov and M. Gubinelli, [A variational method for \$\Phi_3^4\$](#) . *Duke Math. J.* **169** (2020), no. 17, 3339–3415
- [4] N. Barashkov and M. Gubinelli, [The \$\Phi_3^4\$ measure via Girsanov’s theorem](#). *Electron. J. Probab.* **26** (2021), article no. 81
- [5] Á. Bényi, T. Oh, and O. Pocovnicu, [On the probabilistic Cauchy theory of the cubic nonlinear Schrödinger equation on \$\mathbb{R}^d\$, \$d \geq 3\$](#) . *Trans. Amer. Math. Soc. Ser. B* **2** (2015), 1–50
- [6] Á. Bényi, T. Oh, and T. Zhao, [Fractional Leibniz rule on the torus](#). *Proc. Amer. Math. Soc.* **153** (2025), no. 1, 207–221
- [7] J.-M. Bony, [Calcul symbolique et propagation des singularités pour les équations aux dérivées partielles non linéaires](#). *Ann. Sci. École Norm. Sup. (4)* **14** (1981), no. 2, 209–246
- [8] M. Boué and P. Dupuis, [A variational representation for certain functionals of Brownian motion](#). *Ann. Probab.* **26** (1998), no. 4, 1641–1659
- [9] J. Bourgain, [Periodic nonlinear Schrödinger equation and invariant measures](#). *Comm. Math. Phys.* **166** (1994), no. 1, 1–26
- [10] J. Bourgain, [Invariant measures for the 2D-defocusing nonlinear Schrödinger equation](#). *Comm. Math. Phys.* **176** (1996), no. 2, 421–445
- [11] J. Bourgain, [Invariant measures for the Gross–Piatevskii equation](#). *J. Math. Pures Appl. (9)* **76** (1997), no. 8, 649–702
- [12] J. Bourgain, [Nonlinear Schrödinger equations](#). In *Hyperbolic equations and frequency interactions (Park City, UT, 1995)*, pp. 3–157, IAS/Park City Math. Ser. 5, American Mathematical Society, Providence, RI, 1999
- [13] B. Bringmann, [Invariant Gibbs measures for the three-dimensional wave equation with a Hartree nonlinearity I: Measures](#). *Stoch. Partial Differ. Equ. Anal. Comput.* **10** (2022), no. 1, 1–89
- [14] B. Bringmann, [Invariant Gibbs measures for the three-dimensional wave equation with a Hartree nonlinearity II: Dynamics](#). *J. Eur. Math. Soc. (JEMS)* **26** (2024), no. 6, 1933–2089
- [15] B. Bringmann, Y. Deng, A. R. Nahmod, and H. Yue, [Invariant Gibbs measures for the three dimensional cubic nonlinear wave equation](#). *Invent. Math.* **236** (2024), no. 3, 1133–1411
- [16] D. C. Brydges and G. Slade, [Statistical mechanics of the 2-dimensional focusing nonlinear Schrödinger equation](#). *Comm. Math. Phys.* **182** (1996), no. 2, 485–504

- [17] E. A. Carlen, J. Fröhlich, and J. Lebowitz, **Exponential relaxation to equilibrium for a one-dimensional focusing non-linear Schrödinger equation with noise.** *Comm. Math. Phys.* **342** (2016), no. 1, 303–332
- [18] R. Catellier and K. Chouk, **Paracontrolled distributions and the 3-dimensional stochastic quantization equation.** *Ann. Probab.* **46** (2018), no. 5, 2621–2679
- [19] J. Colliander and T. Oh, **Almost sure well-posedness of the cubic nonlinear Schrödinger equation below $L^2(\mathbb{T})$.** *Duke Math. J.* **161** (2012), no. 3, 367–414
- [20] G. Da Prato and A. Debussche, **Strong solutions to the stochastic quantization equations.** *Ann. Probab.* **31** (2003), no. 4, 1900–1916
- [21] G. Da Prato and J. Zabczyk, ***Stochastic equations in infinite dimensions.*** 2nd edn., Encyclopedia Math. Appl. 152, Cambridge University Press, Cambridge, 2014
- [22] Y. Deng, A. R. Nahmod, and H. Yue, **Invariant Gibbs measure and global strong solutions for the Hartree NLS equation in dimension three.** *J. Math. Phys.* **62** (2021), no. 3, article no. 031514
- [23] A. Deya, **A nonlinear wave equation with fractional perturbation.** *Ann. Probab.* **47** (2019), no. 3, 1775–1810
- [24] A. Deya, **On a non-linear 2D fractional wave equation.** *Ann. Inst. Henri Poincaré Probab. Stat.* **56** (2020), no. 1, 477–501
- [25] W. E, A. Jentzen, and H. Shen, **Renormalized powers of Ornstein–Uhlenbeck processes and well-posedness of stochastic Ginzburg–Landau equations.** *Nonlinear Anal.* **142** (2016), 152–193
- [26] D. Fan and S. Sato, **Transference on certain multilinear multiplier operators.** *J. Aust. Math. Soc.* **70** (2001), no. 1, 37–55
- [27] C. Fefferman, **The multiplier problem for the ball.** *Ann. of Math. (2)* **94** (1971), 330–336
- [28] J. Forlano and L. Tolomeo, **On the unique ergodicity for a class of 2 dimensional stochastic wave equations.** *Trans. Amer. Math. Soc.* **377** (2024), no. 1, 345–394
- [29] J. Ginibre, Y. Tsutsumi, and G. Velo, **On the Cauchy problem for the Zakharov system.** *J. Funct. Anal.* **151** (1997), no. 2, 384–436
- [30] J. Ginibre and G. Velo, **Generalized Strichartz inequalities for the wave equation.** *J. Funct. Anal.* **133** (1995), no. 1, 50–68
- [31] L. Grafakos, ***Modern Fourier analysis.*** 3rd edn., Grad. Texts in Math. 250, Springer, New York, 2014
- [32] M. Gubinelli and M. Hofmanová, **Global solutions to elliptic and parabolic Φ^4 models in Euclidean space.** *Comm. Math. Phys.* **368** (2019), no. 3, 1201–1266
- [33] M. Gubinelli and M. Hofmanová, **A PDE construction of the Euclidean ϕ_3^4 quantum field theory.** *Comm. Math. Phys.* **384** (2021), no. 1, 1–75
- [34] M. Gubinelli, P. Imkeller, and N. Perkowski, **Paracontrolled distributions and singular PDEs.** *Forum Math. Pi* **3** (2015), article no. e6
- [35] M. Gubinelli, H. Koch, and T. Oh, **Renormalization of the two-dimensional stochastic nonlinear wave equations.** *Trans. Amer. Math. Soc.* **370** (2018), no. 10, 7335–7359

- [36] M. Gubinelli, H. Koch, and T. Oh, [Paracontrolled approach to the three-dimensional stochastic nonlinear wave equation with quadratic nonlinearity](#). *J. Eur. Math. Soc. (JEMS)* **26** (2024), no. 3, 817–874
- [37] M. Gubinelli, H. Koch, T. Oh, and L. Tolomeo, [Global dynamics for the two-dimensional stochastic nonlinear wave equations](#). *Int. Math. Res. Not. IMRN* **2022** (2022), no. 21, 16954–16999
- [38] T. Gunaratnam, T. Oh, N. Tzvetkov, and H. Weber, [Quasi-invariant Gaussian measures for the nonlinear wave equation in three dimensions](#). *Probab. Math. Phys.* **3** (2022), no. 2, 343–379
- [39] M. Hairer, [A theory of regularity structures](#). *Invent. Math.* **198** (2014), no. 2, 269–504
- [40] M. Hairer and K. Matetski, [Discretisations of rough stochastic PDEs](#). *Ann. Probab.* **46** (2018), no. 3, 1651–1709
- [41] M. Keel and T. Tao, [Endpoint Strichartz estimates](#). *Amer. J. Math.* **120** (1998), no. 5, 955–980
- [42] R. Killip, B. Stovall, and M. Visan, [Blowup behaviour for the nonlinear Klein–Gordon equation](#). *Math. Ann.* **358** (2014), no. 1-2, 289–350
- [43] A. Kupiainen, [Renormalization group and stochastic PDEs](#). *Ann. Henri Poincaré* **17** (2016), no. 3, 497–535
- [44] J. L. Lebowitz, H. A. Rose, and E. R. Speer, [Statistical mechanics of the nonlinear Schrödinger equation](#). *J. Statist. Phys.* **50** (1988), no. 3-4, 657–687
- [45] P. G. Lemarié-Rieusset, [Recent developments in the Navier–Stokes problem](#). Chapman & Hall/CRC Res. Notes Math. 431, Chapman & Hall/CRC, Boca Raton, FL, 2002
- [46] H. Lindblad and C. D. Sogge, [On existence and scattering with minimal regularity for semilinear wave equations](#). *J. Funct. Anal.* **130** (1995), no. 2, 357–426
- [47] J.-C. Mourrat and H. Weber, [The dynamic \$\Phi_3^4\$ model comes down from infinity](#). *Comm. Math. Phys.* **356** (2017), no. 3, 673–753
- [48] J.-C. Mourrat and H. Weber, [Global well-posedness of the dynamic \$\Phi^4\$ model in the plane](#). *Ann. Probab.* **45** (2017), no. 4, 2398–2476
- [49] J.-C. Mourrat, H. Weber, and W. Xu, [Construction of \$\Phi_3^4\$ diagrams for pedestrians](#). In *From particle systems to partial differential equations*, pp. 1–46, Springer Proc. Math. Stat. 209, Springer, Cham, 2017
- [50] E. Nelson, A quartic interaction in two dimensions. In *Mathematical Theory of Elementary Particles (Proc. Conf., Dedham, Mass., 1965)*, pp. 69–73, MIT Press, Cambridge, Massachusetts-London, 1966
- [51] D. Nualart, [The Malliavin calculus and related topics](#). 2nd edn., Probab. Appl. (N. Y.), Springer, Berlin, 2006
- [52] T. Oh and M. Okamoto, [Comparing the stochastic nonlinear wave and heat equations: A case study](#). *Electron. J. Probab.* **26** (2021), article no. 9
- [53] T. Oh, M. Okamoto, and T. Robert, [A remark on triviality for the two-dimensional stochastic nonlinear wave equation](#). *Stochastic Process. Appl.* **130** (2020), no. 9, 5838–5864

- [54] T. Oh, M. Okamoto, and L. Tolomeo, [Focusing \$\Phi_2^4\$ -model with a Hartree-type nonlinearity](#). *Mem. Am. Math. Soc.* **304** (2024), no. 1529
- [55] T. Oh, M. Okamoto, and N. Tzvetkov, [Uniqueness and non-uniqueness of the Gaussian free field evolution under the two-dimensional Wick ordered cubic wave equation](#). *Ann. Inst. Henri Poincaré Probab. Stat.* **60** (2024), no. 3, 1684–1728
- [56] T. Oh, O. Pocovnicu, and N. Tzvetkov, [Probabilistic local Cauchy theory of the cubic nonlinear wave equation in negative Sobolev spaces](#). *Ann. Inst. Fourier (Grenoble)* **72** (2022), no. 2, 771–830
- [57] T. Oh, T. Robert, P. Sosoe, and Y. Wang, [Invariant Gibbs dynamics for the dynamical sine-Gordon model](#). *Proc. Roy. Soc. Edinburgh Sect. A* **151** (2021), no. 5, 1450–1466
- [58] T. Oh, T. Robert, P. Sosoe, and Y. Wang, [On the two-dimensional hyperbolic stochastic sine-Gordon equation](#). *Stoch. Partial Differ. Equ. Anal. Comput.* **9** (2021), no. 1, 1–32
- [59] T. Oh, T. Robert, and N. Tzvetkov, [Stochastic nonlinear wave dynamics on compact surfaces](#). *Ann. H. Lebesgue* **6** (2023), 161–223
- [60] T. Oh, T. Robert, and Y. Wang, [On the parabolic and hyperbolic Liouville equations](#). *Comm. Math. Phys.* **387** (2021), no. 3, 1281–1351
- [61] T. Oh, K. Seong, and L. Tolomeo, [A remark on Gibbs measures with log-correlated Gaussian fields](#). *Forum Math. Sigma* **12** (2024), article no. e50
- [62] T. Oh, P. Sosoe, and L. Tolomeo, [Optimal integrability threshold for Gibbs measures associated with focusing NLS on the torus](#). *Invent. Math.* **227** (2022), no. 3, 1323–1429
- [63] T. Oh and L. Thomann, [A pedestrian approach to the invariant Gibbs measures for the 2-d defocusing nonlinear Schrödinger equations](#). *Stoch. Partial Differ. Equ. Anal. Comput.* **6** (2018), no. 3, 397–445
- [64] T. Oh and L. Thomann, [Invariant Gibbs measures for the 2-d defocusing nonlinear wave equations](#). *Ann. Fac. Sci. Toulouse Math. (6)* **29** (2020), no. 1, 1–26
- [65] T. Oh, Y. Wang, and Y. Zine, [Three-dimensional stochastic cubic nonlinear wave equation with almost space-time white noise](#). *Stoch. Partial Differ. Equ. Anal. Comput.* **10** (2022), no. 3, 898–963
- [66] G. Parisi and Y. S. Wu, [Perturbation theory without gauge fixing](#). *Sci. Sinica* **24** (1981), no. 4, 483–496
- [67] B. C. Rider, [On the \$\infty\$ -volume limit of the focusing cubic Schrödinger equation](#). *Comm. Pure Appl. Math.* **55** (2002), no. 10, 1231–1248
- [68] S. Ryang, T. Saito, and K. Shigemoto, [Canonical stochastic quantization](#). *Progr. Theoret. Phys.* **73** (1985), no. 5, 1295–1298
- [69] I. Shigekawa, [*Stochastic analysis*](#). Translated from the 1998 Japanese original by the author. Iwanami Series in Modern Mathematics. Transl. Math. Monogr. 224, American Mathematical Society, Providence, RI, 2004
- [70] B. Simon, [*The \$P\(\phi\)_2\$ Euclidean \(quantum\) field theory*](#). Princeton Ser. Phys., Princeton University Press, Princeton, NJ, 1974
- [71] L. Thomann and N. Tzvetkov, [Gibbs measure for the periodic derivative nonlinear Schrödinger equation](#). *Nonlinearity* **23** (2010), no. 11, 2771–2791

- [72] L. Tolomeo, Unique ergodicity for a class of stochastic hyperbolic equations with additive space-time white noise. *Comm. Math. Phys.* **377** (2020), no. 2, 1311–1347
- [73] L. Tolomeo, Global well posedness of the two-dimensional stochastic nonlinear wave equation on an unbounded domain. *Ann. Probab.* **49** (2021), no. 3, 1402–1426
- [74] L. Tolomeo, Ergodicity for the hyperbolic $P(\Phi)_2$ -model. 2023, arXiv:[2310.02190v1](https://arxiv.org/abs/2310.02190v1)
- [75] L. Tolomeo and H. Weber, Phase transition for invariant measures of the focusing Schrödinger equation. [v1] 2023, [v2] 2024, arXiv:[2306.07697v2](https://arxiv.org/abs/2306.07697v2)
- [76] H. Triebel, *Tempered homogeneous function spaces*. EMS Ser. Lect. Math., European Mathematical Society (EMS), Zürich, 2015
- [77] A. S. Üstünel, Variational calculation of Laplace transforms via entropy on Wiener space and applications. *J. Funct. Anal.* **267** (2014), no. 8, 3058–3083
- [78] C. Villani, *Topics in optimal transportation*. Grad. Stud. Math. 58, American Mathematical Society, Providence, RI, 2003