Références

[1] J. Arthur, A trace formula for reductive groups. II. Applications of a truncation operator. *Compos. Math.* **40** (1980), n° 1, 87–121

[2] J. Arthur, On a family of distributions obtained from Eisenstein series. II. Explicit formulas. *Amer. J. Math.* **104** (1982), n° 6, 1289–1336

[3] J. Arthur, On the inner product of truncated Eisenstein series. *Duke Math. J.* **49** (1982), n° 1, 35–70

[4] J. Arthur, A measure on the unipotent variety. *Canad. J. Math.* **37** (1985), n° 6, 1237–1274

[5] J. Arthur, On a family of distributions obtained from orbits. *Canad. J. Math.* **38** (1986), n° 1, 179–214

[6] J. Arthur, The invariant trace formula. I. Local theory. *J. Amer. Math. Soc.* **1** (1988), n° 2, 323–383

[7] J. Arthur, The invariant trace formula. II. Global theory. *J. Amer. Math. Soc.* **1** (1988), n° 3, 501–554

[8] J. Arthur, A local trace formula. *Publ. Math. Inst. Hautes Études Sci.* (1991), n° 73, 5–96

[9] J. Arthur, A stable trace formula. II. Global descent. *Invent. Math.* **143** (2001), n° 1, 157–220

[10] J. Arthur, A stable trace formula. I. General expansions. *J. Inst. Math. Jussieu* **1** (2002), n° 2, 175–277

[11] J. Arthur, A stable trace formula. III. Proof of the main theorems. *Ann. of Math. (2)* **158** (2003), n° 3, 769–873

[12] J. Arthur, *The endoscopic classification of representations*. Amer. Math. Soc. Colloq. Publ. 61, American Mathematical Society, Providence, RI, 2013

[13] J. G. Arthur, A trace formula for reductive groups. I. Terms associated to classes in $G(\mathbf{Q})$. *Duke Math. J.* **45** (1978), n° 4, 911–952

[14] A. Borel, *Introduction aux groupes arithmétiques*. Publications de l'Institut de Mathématique de l'Université de Strasbourg, XV. Actualités Scientifiques et Industrielles 1341, Hermann, Paris, 1969

[15] A. Borel, *Linear algebraic groups*. 2e éd., Grad. Texts in Math. 126, Springer, New York, 1991

[16] A. Borel and J. Tits, Groupes réductifs. *Publ. Math. Inst. Hautes Études Sci.* (1965), n° 27, 55–150

[17] V. G. Drinfel′d, Cohomology of compactified moduli varieties of F-sheaves of rank 2 (in Russian). *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)* **162** (1987), 107–158, 189. English translation: *Journal of Soviet Mathematics* **46** (1989), 1789–1821

[18] V. G. Drinfel′d, Varieties of modules of F-sheaves (in Russian). *Funktsional. Anal. i Prilozhen.* **21** (1987), n° 2, 23–41. English translation: *Functional Analysis and Its Applications* **21** (1987), 107–122

[19] G. Harder, Halbeinfache Gruppenschemata über vollständigen Kurven. *Invent. Math.* **6** (1968), 107–149

[20] G. Harder, Minkowskische Reduktionstheorie über Funktionenkörpern. *Invent. Math.* **7** (1969), 33–54

[21] G. Harder, Chevalley groups over function fields and automorphic forms. *Ann. of Math. (2)* **100** (1974), 249–306

[22] H. Jacquet and R. P. Langlands, *Automorphic forms on* GL(2). Lecture Notes in Math. 114, Springer, Berlin-New York, 1970

[23] J.-P. Labesse, L-indistinguishable representations and trace formula for SL(2). In *Lie groups and their representations (Proc. Summer School, Bolyai János Math. Soc., Budapest, 1971)*, pp. 331–338, Halsted Press, New York-Toronto, Ont., 1975

[24] J.-P. Labesse and R. P. Langlands, L-indistinguishability for SL(2). *Canad. J. Math.* **31** (1979), n° 4, 726–785

[25] J.-P. Labesse and J.-L. Waldspurger, *La formule des traces tordue d'après le* Friday Morning Seminar. CRM Monogr. Ser. 31, American Mathematical Society, Providence, RI, 2013

[26] L. Lafforgue, *Chtoucas de Drinfeld et conjecture de Ramanujan-Petersson*. Astérisque 243, Soc. math. de France, 1997

[27] L. Lafforgue, Chtoucas de Drinfeld et correspondance de Langlands. *Invent. Math.* **147** (2002), n° 1, 1–241

[28] R. P. Langlands, *On the functional equations satisfied by Eisenstein series*. Lecture Notes in Math. 544, Springer, Berlin-New York, 1976

[29] R. P. Langlands, *Base change for* GL(2). Ann. of Math. Stud. 96, Princeton University Press, Princeton, NJ ; University of Tokyo Press, Tokyo, 1980

[30] G. Laumon, *Cohomology of Drinfeld modular varieties. Part II*. Cambridge Stud. Adv. Math. 56, Cambridge University Press, Cambridge, 1997

[31] B. Lemaire and G. Henniart, *Représentations des espaces tordus sur un groupe réductif connexe* $\mathfrak{p}$*-adique*. Astérisque 386, Soc. math. de France, 2017

[32] C. Mœglin and J.-L. Waldspurger, *Décomposition spectrale et séries d'Eisenstein*. Progr. Math. 113, Birkhäuser, Basel, 1994

[33] C. Mœglin and J.-L. Waldspurger, *Stabilisation de la formule des traces tordue. Vol. 1*. Progr. Math. 316, Birkhäuser/Springer, Cham, 2016

[34] C. Mœglin and J.-L. Waldspurger, *La formule des traces locale tordue*. Mem. Amer. Math. Soc. 251, Amer. Math. Soc., 2018

[35] L. E. Morris, Eisenstein series for reductive groups over global function fields. I. The cusp form case. *Canad. J. Math.* **34** (1982), n° 1, 91–168

[36] L. E. Morris, Eisenstein series for reductive groups over global function fields. II. The general case. *Canad. J. Math.* **34** (1982), n° 5, 1112–1182

[37] T. Ngô Dac, Sur le développement spectral de la formule des traces d'Arthur-Selberg sur les corps de fonctions. *Bull. Soc. Math. France* **137** (2009), n° 4, 545–586

[38] A. Selberg, Harmonic analysis and discontinuous groups in weakly symmetric Riemannian spaces with applications to Dirichlet series. *J. Indian Math. Soc. (N.S.)* **20** (1956), 47–87

[39] T. A. Springer, Reduction theory over global fields. *Proc. Indian Acad. Sci. Math. Sci.* **104** (1994), n° 1, 207–216