

References

- [1] T. Arakawa, [Vector-valued Siegel’s modular forms of degree two and the associated Andrianov \$L\$ -functions](#). *Manuscripta Math.* **44** (1983), no. 1-3, 155–185
- [2] A. Ash, D. Mumford, M. Rapoport, and Y.-S. Tai, [Smooth compactifications of locally symmetric varieties](#). 2nd edn., Cambridge Math. Lib., Cambridge University Press, Cambridge, 2010
- [3] W. L. Baily, Jr. and A. Borel, [Compactification of arithmetic quotients of bounded symmetric domains](#). *Ann. of Math. (2)* **84** (1966), 442–528
- [4] N. Bergeron, Z. Li, J. Millson, and C. Moeglin, [The Noether–Lefschetz conjecture and generalizations](#). *Invent. Math.* **208** (2017), no. 2, 501–552
- [5] R. E. Borcherds, [Automorphic forms on \$O_{s+2,2}\(\mathbf{R}\)\$ and infinite products](#). *Invent. Math.* **120** (1995), no. 1, 161–213
- [6] R. E. Borcherds, L. Katzarkov, T. Pantev, and N. I. Shepherd-Barron, Families of $K3$ surfaces. *J. Algebraic Geom.* **7** (1998), no. 1, 183–193
- [7] A. Borel, [Density and maximality of arithmetic subgroups](#). *J. Reine Angew. Math.* **224** (1966), 78–89
- [8] J. H. Bruinier, [Borcherds products on \$O\(2, 1\)\$ and Chern classes of Heegner divisors](#). Lecture Notes in Math. 1780, Springer, Berlin, 2002
- [9] F. Cléry and G. van der Geer, [Generators for modules of vector-valued Picard modular forms](#). *Nagoya Math. J.* **212** (2013), 19–57
- [10] F. Cléry and G. van der Geer, [Constructing vector-valued Siegel modular forms from scalar-valued Siegel modular forms](#). *Pure Appl. Math. Q.* **11** (2015), no. 1, 21–47
- [11] A. Collino, Indecomposable motivic cohomology classes on quartic surfaces and on cubic fourfolds. In *Algebraic K-theory and its applications (Trieste, 1997)*, pp. 370–402, World Scientific, River Edge, NJ, 1999
- [12] T. Enright, R. Howe, and N. Wallach, [A classification of unitary highest weight modules](#). In *Representation theory of reductive groups (Park City, Utah, 1982)*, pp. 97–143, Progr. Math. 40, Birkhäuser, Boston, MA, 1983
- [13] T. J. Enright and R. Parthasarathy, [A proof of a conjecture of Kashiwara and Vergne](#). In *Noncommutative harmonic analysis and Lie groups (Marseille, 1980)*, pp. 74–90, Lecture Notes in Math. 880, Springer, Berlin, 1981
- [14] G. Faltings and C.-L. Chai, [Degeneration of abelian varieties](#). *Ergeb. Math. Grenzgeb. (3)* **22**, Springer, Berlin, 1990
- [15] E. Freitag, [Ein Verschwindungssatz für automorphe Formen zur Siegelschen Modulgruppe](#). *Math. Z.* **165** (1979), no. 1, 11–18
- [16] E. Freitag and R. Salvati Manni, [Vector-valued modular forms on a three-dimensional ball](#). *Trans. Amer. Math. Soc.* **371** (2019), no. 8, 5293–5308
- [17] E. Freitag and R. S. Salvati Manni, [Vector-valued Hermitian and quaternionic modular forms](#). *Kyoto J. Math.* **55** (2015), no. 4, 819–836

- [18] W. Fulton and J. Harris, *Representation theory*. Grad. Texts in Math. 129, Springer, New York, 1991
- [19] R. Goodman and N. R. Wallach, *Symmetry, representations, and invariants*. Grad. Texts in Math. 255, Springer, Dordrecht, 2009
- [20] V. Gritsenko, Modular forms and moduli spaces of abelian and $K3$ surfaces (in Russian). *Algebra i Analiz* **6** (1994), no. 6, 65–102. English translation. *St. Petersburg Math. J.* **6** (1995), no. 6, 1179–1208
- [21] V. A. Gritsenko, K. Hulek, and G. K. Sankaran, *The Kodaira dimension of the moduli of $K3$ surfaces*. *Invent. Math.* **169** (2007), no. 3, 519–567
- [22] V. A. Gritsenko, K. Hulek, and G. K. Sankaran, Moduli of $K3$ surfaces and irreducible symplectic manifolds. In *Handbook of moduli. Vol. I*, pp. 459–526, Adv. Lect. Math. (ALM) 24, International Press, Somerville, MA, 2013
- [23] M. Harris, *Special values of zeta functions attached to Siegel modular forms*. *Ann. Sci. École Norm. Sup. (4)* **14** (1981), no. 1, 77–120
- [24] J. E. Humphreys, *Representations of semisimple Lie algebras in the BGG category $\mathcal{O}$* . Grad. Stud. Math. 94, American Mathematical Society, Providence, RI, 2008
- [25] T. Ibukiyama, Dimension formulas of Siegel modular forms of weight 3 and supersingular abelian varieties. In *Proceedings of the 4-th Spring Conference on modular forms and related topics (Hamana lake)*, pp. 39–60, 2007
- [26] T. Ibukiyama, Vector valued Siegel modular forms of symmetric tensor weight of small degrees. *Comment. Math. Univ. St. Pauli* **61** (2012), no. 1, 51–75
- [27] T. Ibukiyama and R. Kyomura, *A generalization of vector valued Jacobi forms*. *Osaka J. Math.* **48** (2011), no. 3, 783–808
- [28] H. P. Jakobsen, *Hermitian symmetric spaces and their unitary highest weight modules*. *J. Funct. Anal.* **52** (1983), no. 3, 385–412
- [29] R. C. King, *Modification rules and products of irreducible representations of the unitary, orthogonal, and symplectic groups*. *J. Mathematical Phys.* **12** (1971), no. 8, 1588–1598
- [30] R. C. King, *Branching rules for classical Lie groups using tensor and spinor methods*. *J. Phys. A* **8** (1975), 429–449
- [31] K. Koike and I. Terada, *Young-diagrammatic methods for the representation theory of the classical groups of type B_n , C_n , D_n* . *J. Algebra* **107** (1987), no. 2, 466–511
- [32] K. Koike and I. Terada, *Young diagrammatic methods for the restriction of representations of complex classical Lie groups to reductive subgroups of maximal rank*. *Adv. Math.* **79** (1990), no. 1, 104–135
- [33] E. Looijenga, *Moduli spaces and locally symmetric varieties*. In *Development of moduli theory—Kyoto 2013*, pp. 33–75, Adv. Stud. Pure Math. 69, The Mathematical Society of Japan, Tokyo, 2016
- [34] S. Ma, *Special holomorphic tensors on orthogonal modular varieties and applications to the Lang conjecture*. *Math. Ann.* **386** (2023), no. 1–2, 1127–1150
- [35] S. Ma, *Irregular cusps of orthogonal modular varieties*. *Canad. J. Math.* **77** (2025), no. 3, 739–769

- [36] D. Mumford, [Hirzebruch's proportionality theorem in the noncompact case](#). *Invent. Math.* **42** (1977), 239–272
- [37] D. Mumford and T. Oda, *Algebraic geometry. II*. Texts Read. Math. 73, Hindustan Book Agency, New Delhi, 2015
- [38] S. Okada, *Representation theory of the classical groups and combinatorics* (in Japanese). Baifukan, 2006
- [39] K. Pommerening, [Die Fortsetzbarkeit von Differentialformen auf arithmetischen Quotienten von hermiteschen symmetrischen Räumen](#). *J. Reine Angew. Math.* **356** (1985), 194–220
- [40] I. I. Pyateskii-Shapiro, *Automorphic functions and the geometry of classical domains*. Math. Appl. 8, Gordon and Breach Science, New York, 1969
- [41] M. S. Raghunathan, *Discrete subgroups of Lie groups*. *Ergeb. Math. Grenzgeb.* 68, Springer, New York, 1972
- [42] T. Satoh, [On certain vector valued Siegel modular forms of degree two](#). *Math. Ann.* **274** (1986), no. 2, 335–352
- [43] N.-P. Skoruppa, [Jacobi forms of critical weight and Weil representations](#). In *Modular forms on Schiermonnikoog*, pp. 239–266, Cambridge University Press, Cambridge, 2008
- [44] G. van der Geer, [Siegel modular forms and their applications](#). In *The 1-2-3 of modular forms*, pp. 181–245, Universitext, Springer, Berlin, 2008
- [45] C. Voisin, [Variations of Hodge structure and algebraic cycles](#). In *Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Zürich, 1994)*, pp. 706–715, Birkhäuser, Basel, 1995
- [46] C. Voisin, [Hodge theory and complex algebraic geometry. I](#). Cambridge Stud. Adv. Math. 76, Cambridge University Press, Cambridge, 2002
- [47] R. Weissauer, [Vektorwertige Siegelsche Modulformen kleinen Gewichtes](#). *J. Reine Angew. Math.* **343** (1983), 184–202
- [48] E. Witt, [Eine Identität zwischen Modulformen zweiten Grades](#). *Abh. Math. Sem. Univ. Hamburg* **14** (1941), 323–337