

Graph product structure: Theory and applications

Vida Dujmović and David R. Wood

Abstract. Graph product structure theory describes graphs in complicated graph classes as subgraphs of products of graphs in simpler graph classes. This essay selectively surveys recent results and applications in this field.

1. Introduction

One of the aims of the field of graph structure theory is to describe the global structure of graphs in a particular class $\mathcal{G}$ in terms of simpler graphs. Graph product structure theory achieves this goal by describing graphs in $\mathcal{G}$ as subgraphs of products of graphs in simpler graph classes, typically with bounded treewidth. The *treewidth* of a graph G , denoted by $\text{tw}(G)$, is the standard measure of how similar G is to a tree. Graph classes with bounded treewidth are considered to be simple; for example, numerous NP-complete problems are efficiently solvable on graphs with bounded treewidth. See [5, 39, 52] for background on treewidth, including the definition.

The following planar graph product structure theorem by Dujmović, Joret, Micek, Morin, Ueckerdt, and Wood [29] is the classical example of a graph product structure theorem. Here, a graph H is *contained* in a graph G if H is isomorphic to a subgraph of G , written $H \lesssim G$. As illustrated in Figure 1, the *strong product* of graphs A and B , denoted by $A \boxtimes B$, is the graph with vertex set $V(A) \times V(B)$, where distinct vertices $(v, x), (w, y)$ are adjacent if

- $v = w$ and $xy \in E(B)$, or
- $x = y$ and $vw \in E(A)$, or
- $vw \in E(A)$ and $xy \in E(B)$.

Theorem 1.1 ([29]). *For every planar graph G , there is a graph H and a path P such that $\text{tw}(H) \leq 8$ and $G \lesssim H \boxtimes P$.*




Figure 1. Example of a strong product.

Theorem 1.1 implies and generalises many results from the literature on the global structure of planar graphs, including the Lipton–Tarjan balanced separator theorem [49], the Baker decomposition and bounded local treewidth [2, 12, 37], and layered treewidth [32].

Theorem 1.1 has been the key tool for resolving several open problems regarding queue layouts [29], nonrepetitive colourings [24], centred colourings [11], adjacency labelling schemes [6, 23, 38], twin-width [7, 47], infinite graphs [45], clustered colouring [25, 26], and comparable box dimension [35]. Now we give one example.

Motivated by measuring the relative power of stacks and queues as data structures, Heath, Leighton, and Rosenberg [40, 41] defined the *stack number* $\text{sn}(G)$ and *queue number* $\text{qn}(G)$ of a graph G . They asked whether planar graphs have bounded queue number, which became the most important open problem in the field and a key case to consider in the comparison of the relative power of stacks and queues.

Prior to [29] the best known upper bound on the queue number of n -vertex planar graphs was $O(\log n)$ [14, 22]. With Theorem 1.1 in hand, Dujmović et al. [29] showed that an $O(1)$ bound is easily proved using already established tools. In particular, Dujmović, Morin, and Wood [34] showed in 2005 that $\text{qn}(G) \leq f(\text{tw}(G))$ for some function f . The best known bound is $\text{qn}(G) \leq 2^{\text{tw}(G)} - 1$ due to Wiechert [56]. Wood [57] showed in 2005 that $\text{qn}(H \boxtimes P) \leq 3 \text{qn}(H) + 1$ for any graph H and any path P . Now consider any planar graph G . By Theorem 1.1, $G \subseteq H \boxtimes P$, where $\text{tw}(H) \leq 8$ and P is a path. Since queue number is subgraph monotone, the above results together imply that

$$\text{qn}(G) \leq \text{qn}(H \boxtimes P) \leq 3 \text{qn}(H) + 1 \leq 3(2^{\text{tw}(H)} - 1) + 1 \leq 3(2^8 - 1) + 1 = 766.$$

This shows that planar graphs have bounded queue number, resolving the 28-year-old open problem of Heath, Leighton, and Rosenberg [40, 41].

The key tool in this proof is the planar graph product structure theorem. We emphasise that the authors of [29] were not trying to prove Theorem 1.1. They first solved the queue-number problem and then realised that their proof contained

a structural description of planar graphs that could be written as a product, as in Theorem 1.1. It turns out that, back in 2008, Wood [58] had conjectured a version of Theorem 1.1, where the subgraph relation was replaced by the weaker notion of shallow minor. Theorem 1.1 implies this conjecture.

Theorem 1.1 has been extended in various ways. The following result is the state of the art for the product structure of planar graphs.

Theorem 1.2. *Every planar graph G is contained in:*

- (a) $H \boxtimes P$, where $\text{tw}(H) \leq 6$ and P is a path [55],
- (b) $H \boxtimes P \boxtimes K_2$, where $\text{tw}(H) \leq 4$ and P is a path [55],
- (c) $H \boxtimes P \boxtimes K_3$, where $\text{tw}(H) \leq 3$ and P is a path [29].

Theorem 1.2 (c) is often the most useful version of the planar graph product structure theorem since the dependence on $\text{tw}(H)$ in applications is often exponential, as in the queue-number example above. Indeed, using Theorem 1.2 (c), Dujmović et al. [29] showed that planar graphs have queue number at most 49, which was improved to 42 by Bekos, Gronemann, and Raftopoulou [3]. The treewidth 3 bound in Theorem 1.2 (c) is the best possible for any result showing that planar graphs are contained in $H \boxtimes P \boxtimes K_c$ for any constant c (see [29]). Improved bounds are known for special classes of planar graphs, such as squaregraphs [42]. Note that some extra properties hold in Theorems 1.1 and 1.2. Notably, H is a minor of the given graph G , and thus H is planar. On the other hand, Dujmović, Joret, Micek, Morin, and Wood [30] proved that in Theorems 1.1 and 1.2, H must have unbounded maximum degree, even for planar graphs G with maximum degree 5.

The proofs of Theorems 1.1 and 1.2 were inspired by an elegant result of Pilipczuk and Siebertz [51] which uses Sperner's lemma at its heart. The method is constructive and leads to an efficient algorithm for computing the mapping into the product graph. Indeed, Bose, Morin, and Odak [9] gave a linear-time algorithm for doing so. Note that Illingworth, Scott, and Wood [46] gave a purely combinatorial proof of Theorem 1.2 (c) that avoids planar embeddings and avoids the use of Sperner's lemma. Roughly speaking, their result describes how any tree decomposition of a graph excluding a fixed minor can be converted into a product-like structure.

Motivated by Theorem 1.1, Bose, Dujmović, Javarsineh, Morin, and Wood [8] introduced the following definition. The *row treewidth* of a graph G , denoted by $\text{rtw}(G)$, is the minimum $c \in \mathbb{N}$ such that $G \subsetneq H \boxtimes P$ for some path P and graph H with $\text{tw}(H) \leq c$. Theorem 1.2 (a) says that planar graphs have row treewidth at most 6. It is open whether planar graphs have row treewidth at most 3. Bose et al. [8] studied the relationship between row treewidth and layered treewidth $\text{ltw}(G)$. While it is easily seen that $\text{ltw}(G) \leq \text{rtw}(G)$, Bose et al. [8] described a class of graphs G

with $\text{ltw}(G) = 1$ and unbounded row treewidth. Thus, bounded row treewidth is a strictly stronger setting than bounded layered treewidth.

Other classes with bounded row treewidth include graphs embeddable on any fixed surface [29]. The best bounds are due to Distel, Hickingbotham, Huynh, and Wood [18].

Theorem 1.3 ([18]). *Every graph G of Euler genus g is contained in:*

- (a) $H \boxtimes P$, where $\text{tw}(H) \leq 2g + 6$ and P is a path,
- (b) $H \boxtimes P \boxtimes K_{\max\{2g, 3\}}$, where $\text{tw}(H) \leq 3$ and P is a path.

Dujmović et al. [29] established the following characterisation of minor-closed classes with bounded row treewidth. A graph X is *apex* if $X - a$ is planar for some vertex $a \in V(X)$.

Theorem 1.4 ([29]). *A minor-closed class $\mathcal{G}$ has bounded row treewidth if and only if some apex graph is not in $\mathcal{G}$.*

The product structure of graphs excluding an apex minor has been further studied in [27, 46]. For an apex graph X , let k_X be the minimum integer such that, for some integer c , every X -minor-free graph is contained in $H \boxtimes P \boxtimes K_c$, where H is a graph with $\text{tw}(H) \leq k_X$ and P is a path. Illingworth et al. [46] showed that k_X is at most the vertex cover number of X . Tightening this result, Dujmović et al. [27] showed that $\text{td}(X) - 2 \leq k_X \leq 2^{\text{tw}(X)-1}$, where $\text{td}(X)$ is the tree depth of X . This shows that the tree depth of X determines the minimum treewidth in such a product structure theorem.

While graphs excluding an apex minor are the largest class with bounded row treewidth, graph products can be used to describe the structure of graphs excluding any fixed minor. Dujmović et al. [25] showed that graphs excluding a fixed minor and with bounded maximum degree have bounded row treewidth. Dujmović et al. [29] showed the following ‘graph minor product structure theorem’, where $+K_a$ means the complete join with a complete graph on a vertices.

Theorem 1.5 ([29]). *For any graph X , there exist integers k, a such that every X -minor-free graph can be obtained from clique sums of subgraphs of graphs of the form $(H \boxtimes P) + K_a$, where $\text{tw}(H) \leq k$.*

This result has enabled many of the above applications, including bounded queue number, to be extended to the setting of any proper minor-closed class. As another application, Dujmović et al. [29] showed that Theorem 1.5 implies the following classical theorem of DeVos, Ding, Oporowski, Sanders, Reed, Seymour, and Vertigan [13]: for any graph X and integer $k \geq 2$, every X -minor-free graph has an

(improper) vertex k -colouring such that the union of any $k - 1$ colour classes has bounded treewidth. Another application relates to Hadwiger's conjecture. Using Theorem 1.5 as a starting point, Dujmović et al. [26] showed that graphs excluding a fixed K_t -minor are $(t - 1)$ -colourable with monochromatic components of bounded size. This resolves the so-called clustered Hadwiger conjecture.

Graph product structure theorems have also been established for various non-minor-closed graph classes [4, 19, 33, 43]. Here, Theorems 1.1 and 1.2 are used as building blocks for establishing product structure theorems for more general classes, as illustrated by the following example. A graph is (g, k) -planar if it has a drawing in a surface of Euler genus g such that each edge is in at most k crossings (and no three edges cross at the same point).

Theorem 1.6. *Every (g, k) -planar graph G is contained in:*

- (a) $H \boxtimes P \boxtimes K_c$, where $\text{tw}(H) \leq O(k^3)$, P is a path, and $c \in O(gk^2)$ [33],
- (b) $H \boxtimes P \boxtimes K_c$, where $\text{tw}(H) \leq 10^{10}$, P is a path, and $c = c(g, k)$ [19].

We now briefly summarise some of the recent directions in which graph product structure theory is moving.

- Huynh et al. [45] extended all of the above results to the setting of countably infinite graphs. This is interesting, since in the infinite setting, the graph $H \boxtimes P$ is universal, in the sense that it contains every countable planar graph.
- Several authors have established results which demonstrate that graphs of given treewidth are contained in products of simpler graphs [10, 15, 16, 20, 21, 50]. Indeed, this direction predates Theorem 1.1 through the classical work of Ding and Oporowski [15, 16]. The characterisation of graphs with polynomial growth due to Krauthgamer and Lee [48] is another early example of a graph product structure theorem.
- Hliněný and Jedelský [44] introduced graph product structure theory for dense graph classes and the induced subgraph relation.
- A number of classical theorems about graph minors have recently been revisited in light of graph product structure theory. Dujmović, Hickingbotham, Joret, Micek, Morin, and Wood [28] showed that for every tree T of radius h , there is an integer c such that every T -minor-free graph is contained in $H \boxtimes K_c$ for some graph H with pathwidth at most $2h - 1$. This is a qualitative strengthening of the excluded tree minor theorem of Robertson and Seymour [53]. They also showed that radius is the right parameter to consider in this setting, and $2h - 1$ is the best possible bound. Dujmović et al. [27] proved analogous strengthenings of the grid minor theorem of Robertson and Seymour [54]. Here the right parameter of the excluded minor is tree depth.

- Several recent results show that for various graph classes $\mathcal{G}$, every n -vertex graph in $\mathcal{G}$ is contained in $H \boxtimes K_m$, where H has bounded treewidth and $m \in O(\sqrt{n})$. This direction was initiated by Illingworth et al. [46]. For planar graphs, Distel, Dujmović, Eppstein, Hickingbotham, Joret, Micek, Morin, Seweryn, and Wood [17] showed this result with $\text{tw}(H) \leq 2$ which is best possible. For K_t -minor-free graphs, they showed this result with $\text{tw}(H) \leq 4$. These two results are qualitative strengthenings of classical balanced separator results of Lipton and Tarjan [49] and Alon, Seymour, and Thomas [1]. Dvořák and Wood [36] studied analogous questions for any graph class admitting strongly sublinear separators. Recently, Dujmović, Joret, Micek, Morin, and Wood [31] showed that every n -vertex planar graph (and indeed any graph with bounded row treewidth) is contained in $H \boxtimes K_m$, where H is a fan graph (which has pathwidth 2) and $m \in O(\sqrt{n} \log^2 n)$.

We finish with two open problems. The first asks how far these product structure theorems can be generalised. For example, is there a product structure theorem (using other ingredients in addition to products) for any monotone graph class admitting strongly sublinear separators? For example, the Koebe disc packing theorem says every planar graph can be represented as the touching graph of discs in $\mathbb{R}^2$. What can be said about the product structure of touching graphs of spheres in $\mathbb{R}^3$? Our second open problem concerns applications. Does every monotone graph class admitting strongly sublinear separators have bounded queue number? We expect that a successful answer to this question will unlock the right structure theorem to answer the first question. The interplay between theory and applications is a feature of this field of research.

Acknowledgements. We thank all our collaborators on this topic: Édouard Bonnet, Prosenjit Bose, Rutger Campbell, Katie Clinch, Marc Distel, David Eppstein, Louis Esperet, Cyril Gavoille, Pascal Gollin, Kevin Hendrey, Jędrzej Hodor, Robert Hickingbotham, Tony Huynh, Freddie Illingworth, Mehrnoosh Javarsineh, Gwenaél Joret, Paul Jungeblut, O-joung Kwon, Hoang La, Laura Merker, Bojan Mohar, Pat Morin, Piotr Micek, Clément Rambaud, Robert Šámal, Alex Scott, Michał Seweryn, Youri Tamitegama, Jane Tan, Carsten Thomassen, Torsten Ueckerdt, Bartosz Walczak, Wendy Yi.

Funding. Research by Dujmović is supported by NSERC and a University of Ottawa Research Chair. Research by Wood is supported by the Australian Research Council and by NSERC.

References

- [1] N. Alon, P. Seymour, and R. Thomas, [A separator theorem for nonplanar graphs](#). *J. Amer. Math. Soc.* **3** (1990), no. 4, 801–808 Zbl [0747.05051](#) MR [1065053](#)
- [2] B. S. Baker, [Approximation algorithms for NP-complete problems on planar graphs](#). *J. ACM* **41** (1994), no. 1, 153–180 Zbl [0807.68067](#) MR [1369197](#)
- [3] M. Bekos, M. Gronemann, and C. N. Raftopoulou, [An improved upper bound on the queue number of planar graphs](#). *Algorithmica* **85** (2023), no. 2, 544–562 Zbl [1515.68223](#) MR [4541066](#)
- [4] M. A. Bekos, G. Da Lozzo, P. Hliněný, and M. Kaufmann, [Graph product structure for \$h\$ -framed graphs](#). In *Proceedings of the 33rd international symposium on algorithms and computation (ISAAC 2022)*, LIPIcs - Leibniz Int. Proc. Inform. 248, Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, 2022, article no. 23, 15 pp. Zbl [1570.05113](#) MR [4526676](#)
- [5] H. L. Bodlaender, [A partial \$k\$ -arboretum of graphs with bounded treewidth](#). *Theoret. Comput. Sci.* **209** (1998), no. 1–2, 1–45 Zbl [0912.68148](#) MR [1647486](#)
- [6] M. Bonamy, C. Gavaille, and M. Pilipczuk, [Shorter labeling schemes for planar graphs](#). *SIAM J. Discrete Math.* **36** (2022), no. 3, 2082–2099 Zbl [1548.68148](#) MR [4474794](#)
- [7] É. Bonnet, O.-j. Kwon, and D. R. Wood, [Reduced bandwidth: a qualitative strengthening of twin-width in minor-closed classes \(and beyond\)](#). *J. Combin. Theory Ser. B* **178** (2026), 27–66 MR [4998097](#)
- [8] P. Bose, V. Dujmović, M. Javarsineh, P. Morin, and D. R. Wood, [Separating layered treewidth and row treewidth](#). *Discrete Math. Theor. Comput. Sci.* **24** (2022), no. 1, article no. 18, 10 pp. Zbl [1520.05028](#) MR [4425386](#)
- [9] P. Bose, P. Morin, and S. Odak, [An optimal algorithm for product structure in planar graphs](#). In *Proceedings of the 18th Scandinavian symposium and workshops on algorithm theory (SWAT 2022)*, LIPIcs - Leibniz Int. Proc. Inform. 227, Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, 2022, article no. 19, 14 pp. Zbl [07853691](#) MR [4473458](#)
- [10] R. Campbell, K. Clinch, M. Distel, J. P. Gollin, K. Hendrey, R. Hickingbotham, T. Huynh, F. Illingworth, Y. Tamitegama, J. Tan, and D. R. Wood, [Product structure of graph classes with bounded treewidth](#). *Combin. Probab. Comput.* **33** (2024), no. 3, 351–376 Zbl [1551.05333](#) MR [4730906](#)
- [11] M. Dębski, S. Felsner, P. Micek, and F. Schröder, [Improved bounds for centered colorings](#). *Adv. Comb.* (2021), article no. 8, 28 pp. Zbl [1478.05050](#) MR [4309118](#)
- [12] E. D. Demaine and M. T. Hajiaghayi, [Diameter and treewidth in minor-closed graph families, revisited](#). *Algorithmica* **40** (2004), no. 3, 211–215 Zbl [1082.05086](#) MR [2080518](#)
- [13] M. DeVos, G. Ding, B. Oporowski, D. P. Sanders, B. Reed, P. Seymour, and D. Vertigan, [Excluding any graph as a minor allows a low tree-width 2-coloring](#). *J. Combin. Theory Ser. B* **91** (2004), no. 1, 25–41 Zbl [1042.05036](#) MR [2047529](#)

- [14] G. Di Battista, F. Frati, and J. Pach, [On the queue number of planar graphs](#). *SIAM J. Comput.* **42** (2013), no. 6, 2243–2285 Zbl [1300.05075](#) MR [3141759](#)
- [15] G. Ding and B. Oporowski, [Some results on tree decomposition of graphs](#). *J. Graph Theory* **20** (1995), no. 4, 481–499 Zbl [0837.05044](#) MR [1358539](#)
- [16] G. Ding and B. Oporowski, [On tree-partitions of graphs](#). *Discrete Math.* **149** (1996), no. 1–3, 45–58 Zbl [0844.05078](#) MR [1375097](#)
- [17] M. Distel, V. Dujmović, D. Eppstein, R. Hickingbotham, G. Joret, P. Micek, P. Morin, M. T. Seweryn, and D. R. Wood, [Product structure extension of the Alon–Seymour–Thomas theorem](#). *SIAM J. Discrete Math.* **38** (2024), no. 3, 2095–2107 Zbl [1543.05172](#) MR [4770382](#)
- [18] M. Distel, R. Hickingbotham, T. Huynh, and D. R. Wood, [Improved product structure for graphs on surfaces](#). *Discrete Math. Theor. Comput. Sci.* **24** (2022), no. 2, article no. 6, 10 pp. Zbl [1516.05145](#) MR [4504777](#)
- [19] M. Distel, R. Hickingbotham, M. Seweryn, and D. R. Wood, [Powers of planar graphs, product structure, and blocking partitions](#). *Innov. Graph Theory* **1** (2024), 39–86 Zbl [1556.05032](#) MR [4882662](#)
- [20] M. Distel and D. R. Wood, Tree-partitions with small bounded degree trees. [v1] 2022, [v2] 2023, arXiv:[2210.12577v2](#)
- [21] M. Distel and D. R. Wood, [Tree-partitions with bounded degree trees](#). In D. R. Wood, J. de Gier, and C. E. Praeger (Eds.), *2021–2022 MATRIX annals*, pp. 203–212, MATRIX Book Ser. 5, Springer, Cham, 2024 Zbl [1548.05085](#) MR [4807276](#)
- [22] V. Dujmović, [Graph layouts via layered separators](#). *J. Combin. Theory Ser. B* **110** (2015), 79–89 Zbl [1303.05036](#) MR [3279388](#)
- [23] V. Dujmović, L. Esperet, C. Gavaille, G. Joret, P. Micek, and P. Morin, [Adjacency labelling for planar graphs \(and beyond\)](#). *J. ACM* **68** (2021), no. 6, article no. 42, 33 pp. Zbl [1499.05165](#) MR [4402353](#)
- [24] V. Dujmović, L. Esperet, G. Joret, B. Walczak, and D. R. Wood, [Planar graphs have bounded nonrepetitive chromatic number](#). *Adv. Comb.* (2020), article no. 5, 11 pp. Zbl [1450.05012](#) MR [4125346](#)
- [25] V. Dujmović, L. Esperet, P. Morin, B. Walczak, and D. R. Wood, [Clustered 3-colouring graphs of bounded degree](#). *Combin. Probab. Comput.* **31** (2022), no. 1, 123–135 Zbl [1510.05071](#) MR [4356460](#)
- [26] V. Dujmović, L. Esperet, P. Morin, and D. R. Wood, [Proof of the clustered Hadwiger conjecture](#). In *Proceedings of the 2023 IEEE 64th annual symposium on foundations of computer science (FOCS 2023)*, pp. 1921–1930, IEEE Computer Society, 2023 Zbl [08080132](#) MR [4720358](#)
- [27] V. Dujmović, R. Hickingbotham, J. Hodor, G. Joret, H. La, P. Micek, P. Morin, C. Rambaud, and D. R. Wood, [The grid-minor theorem revisited](#). In *Proceedings of the 2024 annual ACM-SIAM symposium on discrete algorithms (SODA 2024)*, pp. 1241–1245, SIAM, 2024 Zbl [07951121](#) MR [4699293](#)

- [28] V. Dujmović, R. Hickingbotham, G. Joret, P. Micek, P. Morin, and D. R. Wood, [The excluded tree minor theorem revisited](#). *Combin. Probab. Comput.* **33** (2024), no. 1, 85–90 Zbl 1551.05381 MR 4680491
- [29] V. Dujmović, G. Joret, P. Micek, P. Morin, T. Ueckerdt, and D. R. Wood, [Planar graphs have bounded queue-number](#). *J. ACM* **67** (2020), no. 4, article no. 22, 38 pp. Zbl 1466.05047 MR 4148600
- [30] V. Dujmović, G. Joret, P. Micek, P. Morin, and D. R. Wood, [Bounded-degree planar graphs do not have bounded-degree product structure](#). *Electron. J. Combin.* **31** (2024), no. 2, article no. P2.51, 15 pp. Zbl 1548.05281 MR 4758205
- [31] V. Dujmović, G. Joret, P. Micek, P. Morin, and D. R. Wood, [Planar graphs in blowups of fans](#). In *Proceedings of the 2025 annual ACM-SIAM symposium on discrete algorithms (SODA 2025)*, pp. 3382–3391, SIAM, 2025 MR 4863525
- [32] V. Dujmović, P. Morin, and D. R. Wood, [Layered separators in minor-closed graph classes with applications](#). *J. Combin. Theory Ser. B* **127** (2017), 111–147 Zbl 1371.05282 MR 3704658
- [33] V. Dujmović, P. Morin, and D. R. Wood, [Graph product structure for non-minor-closed classes](#). *J. Combin. Theory Ser. B* **162** (2023), 34–67 Zbl 1519.05211 MR 4586042
- [34] V. Dujmović, P. Morin, and D. R. Wood, [Layout of graphs with bounded tree-width](#). *SIAM J. Comput.* **34** (2005), no. 3, 553–579 Zbl 1069.05055 MR 2137079
- [35] Z. Dvořák, D. Gonçalves, A. Lahiri, J. Tan, and T. Ueckerdt, [On comparable box dimension](#). In *Proceedings of the 38th international symposium on computational geometry (SoCG 2022)*, LIPIcs - Leibniz Int. Proc. Inform. 224, Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, 2022, article no. 38, 14 pp. Zbl 1558.68235 MR 4470917
- [36] Z. Dvořák and D. R. Wood, [Product structure of graph classes with strongly sublinear separators](#). *Innov. Graph Theory* **2** (2025), 191–222 Zbl 08099014 MR 4960622
- [37] D. Eppstein, [Diameter and treewidth in minor-closed graph families](#). *Algorithmica* **27** (2000), no. 3–4, 275–291 Zbl 0963.05128 MR 1759751
- [38] L. Esperet, G. Joret, and P. Morin, [Sparse universal graphs for planarity](#). *J. Lond. Math. Soc. (2)* **108** (2023), no. 4, 1333–1357 Zbl 1527.05041 MR 4655266
- [39] D. J. Harvey and D. R. Wood, [Parameters tied to treewidth](#). *J. Graph Theory* **84** (2017), no. 4, 364–385 Zbl 1359.05030 MR 3623383
- [40] L. S. Heath, F. T. Leighton, and A. L. Rosenberg, [Comparing queues and stacks as mechanisms for laying out graphs](#). *SIAM J. Discrete Math.* **5** (1992), no. 3, 398–412 Zbl 0764.05093 MR 1172748
- [41] L. S. Heath and A. L. Rosenberg, [Laying out graphs using queues](#). *SIAM J. Comput.* **21** (1992), no. 5, 927–958 Zbl 0778.05078 MR 1181408
- [42] R. Hickingbotham, P. Jungeblut, L. Merker, and D. R. Wood, [The product structure of squaregraphs](#). *J. Graph Theory* **105** (2024), no. 2, 179–191 Zbl 1531.05057 MR 4684232

- [43] R. Hickingbotham and D. R. Wood, [Shallow minors, graph products, and beyond-planar graphs](#). *SIAM J. Discrete Math.* **38** (2024), no. 1, 1057–1089 Zbl 1535.05076 MR 4716720
- [44] P. Hliněný and J. Jedelský, [\$\mathcal{H}\$ -clique-width and a hereditary analogue of product structure](#). In *Proceedings of the 49th international symposium on mathematical foundations of computer science (MFCS 2024)*, LIPIcs - Leibniz Int. Proc. Inform. 306, Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, 2024, article no. 61, 16 pp. MR 4802177
- [45] T. Huynh, B. Mohar, R. Šámal, C. Thomassen, and D. R. Wood, [Universality in minor-closed graph classes](#). 2021, arXiv:2109.00327v1
- [46] F. Illingworth, A. Scott, and D. R. Wood, [Product structure of graphs with an excluded minor](#). *Trans. Amer. Math. Soc. Ser. B* **11** (2024), 1233–1248 Zbl 1554.05157 MR 4817445
- [47] H. Jacob and M. Pilipczuk, [Bounding twin-width for bounded-treewidth graphs, planar graphs, and bipartite graphs](#). In *Proceedings of the 48th international workshop on graph-theoretic concepts in computer science (WG 2022)*, pp. 287–299, Lecture Notes in Comput. Sci. 13453, Springer, 2022 Zbl 1570.05030 MR 4606560
- [48] R. Krauthgamer and J. R. Lee, [The intrinsic dimensionality of graphs](#). *Combinatorica* **27** (2007), no. 5, 551–585 Zbl 1164.05050 MR 2375717
- [49] R. J. Lipton and R. E. Tarjan, [A separator theorem for planar graphs](#). *SIAM J. Appl. Math.* **36** (1979), no. 2, 177–189 Zbl 0432.05022 MR 0524495
- [50] C.-H. Liu and D. R. Wood, [Quasi-tree-partitions of graphs with an excluded subgraph](#). [v1] 2024, [v2] 2025, arXiv:2408.00983v2
- [51] M. Pilipczuk and S. Siebertz, [Polynomial bounds for centered colorings on proper minor-closed graph classes](#). *J. Combin. Theory Ser. B* **151** (2021), 111–147 Zbl 1490.05082 MR 4278114
- [52] B. A. Reed, [Tree width and tangles: a new connectivity measure and some applications](#). In R. A. Bailey (Ed.), *Surveys in combinatorics*, pp. 87–162, London Math. Soc. Lecture Note Ser. 241, Cambridge University Press, Cambridge, 1997 Zbl 0895.05034 MR 1477746
- [53] N. Robertson and P. D. Seymour, [Graph minors. I. Excluding a forest](#). *J. Combin. Theory Ser. B* **35** (1983), no. 1, 39–61 Zbl 0521.05062 MR 0723569
- [54] N. Robertson and P. D. Seymour, [Graph minors. V. Excluding a planar graph](#). *J. Combin. Theory Ser. B* **41** (1986), no. 1, 92–114 Zbl 0598.05055 MR 0854606
- [55] T. Ueckerdt, D. R. Wood, and W. Yi, [An improved planar graph product structure theorem](#). *Electron. J. Combin.* **29** (2022), no. 2, article no. P2.51, 12 pp. Zbl 1527.05046 MR 4441087
- [56] V. Wiechert, [On the queue-number of graphs with bounded tree-width](#). *Electron. J. Combin.* **24** (2017), no. 1, article no. P1.65, 18 pp. Zbl 1358.05060 MR 3651947
- [57] D. R. Wood, [Queue layouts of graph products and powers](#). *Discrete Math. Theor. Comput. Sci.* **7** (2005), no. 1, 255–268 Zbl 1153.05325 MR 2183176

- [58] D. R. Wood, The structure of Cartesian products. In *Banff workshop on graph minors*, 2008, <https://www.birs.ca/workshops/2008/08w5079/report08w5079.pdf> visited on 2 February 2026

Vida Dujmović

School of Computer Science and Electrical Engineering, University of Ottawa, Ottawa, Canada; vida.dujmovic@uottawa.ca

David R. Wood

School of Mathematics, Monash University, Melbourne, Australia; david.wood@monash.edu