

An abstract Nash–Moser theorem with parameters and applications to PDEs

M. Berti ^{a,*}, P. Bolle ^b, M. Procesi ^{a,1}

^a *Dipartimento di Matematica e Applicazioni “R. Caccioppoli”, Università degli Studi Napoli Federico II, Via Cintia, Monte S. Angelo, I-80126 Napoli, Italy*

^b *Université d’Avignon et des Pays de Vaucluse, Laboratoire d’Analyse non Linéaire et Géométrie (EA 2151), F-84018 Avignon, France*

Received 26 February 2009; received in revised form 8 November 2009; accepted 8 November 2009

Available online 3 December 2009

Abstract

We prove an abstract Nash–Moser implicit function theorem with parameters which covers the applications to the existence of finite dimensional, differentiable, invariant tori of Hamiltonian PDEs with merely differentiable nonlinearities. The main new feature of the abstract iterative scheme is that the linearized operators, in a neighborhood of the expected solution, are invertible, and satisfy the “tame” estimates, only for proper subsets of the parameters. As an application we show the existence of periodic solutions of nonlinear wave equations on Riemannian Zoll manifolds. A point of interest is that, in presence of possibly very large “clusters of small divisors”, due to resonance phenomena, it is more natural to expect solutions with only Sobolev regularity.

© 2009 L’Association Publications de l’Institut Henri Poincaré. Published by Elsevier B.V. All rights reserved.

1. Introduction

1.1. Small divisors problems in Hamiltonian PDEs

Bifurcation problems of periodic and quasi-periodic solutions for Hamiltonian PDEs are naturally affected by small divisors difficulties: the standard implicit function theorem cannot be applied because the linearized operators have an unbounded inverse, due to arbitrarily “small divisors” in their Fourier series expansions. This problem has been handled for PDEs with analytic nonlinearities via KAM methods, see e.g. Kuksin [22,23], Wayne [29], Pöschel [27], Eliasson and Kuksin [15], or via Newton-type iterative schemes as developed in Craig and Wayne [14] and Bourgain [7–10].

The pioneering KAM results in [22,29,27] were limited to 1-dimensional PDEs, with Dirichlet boundary conditions, because they required the eigenvalues of the Laplacian to be *simple* (the square roots of the eigenvalues are the normal mode frequencies of small oscillations). In this case one can impose the so-called “second order Melnikov” non-resonance conditions between the “tangential” and the “normal” frequencies of the expected KAM torus to solve

* Corresponding author.

E-mail addresses: m.berti@unina.it (M. Berti), philippe.bolle@univ-avignon.fr (P. Bolle), michela.procesi@dma.unina.it (M. Procesi).

¹ Supported by the European Research Council under FP7 “New Connections between dynamical systems and Hamiltonian PDEs with small divisors phenomena”.

the homological equations which arise at each step of the KAM iteration. Such equations are linear PDEs with constant coefficients and can be solved simply using Fourier series. Unfortunately, yet for periodic boundary conditions, where two consecutive eigenvalues are possibly equal, the second order Melnikov non-resonance conditions are violated. For wave equations this case has been later handled via KAM method by Chierchia and You in [11].

On the other hand, the Lyapunov–Schmidt decomposition approach, combined with the Newton method developed in [14,7–10], has the advantage to require only the “minimal” non-resonance conditions, which, for example, are fulfilled in higher dimensional PDE applications (we refer to [15] for the KAM approach in higher dimension). As a drawback, its main difficulty relies on the inversion of the linearized operators in a neighborhood of the expected solution, and in obtaining estimates of their inverses in analytic (or Gevrey) norms. Indeed these operators come from linear PDEs with non-constant coefficients and are small perturbations of a diagonal operator having arbitrarily small eigenvalues. Their spectrum depends very sensitively on the parameters, whence they are invertible only over complicated Cantor-like set of parameters with possibly positive measure.

We also mention that, more recently, the Lindstedt series renormalization method has been developed by Gentile, Mastropietro and Procesi to prove the existence of periodic solutions for analytic PDEs, one-dimensional in [16,17] and also higher dimensional in [18].

In all the mentioned results analyticity is deeply exploited, either for the convergence proof of the iterative scheme, or in obtaining suitable estimates for inverse linearized operators.

The existence of periodic solutions of Hamiltonian PDEs with merely differentiable nonlinearities has been recently proved in [3,4] along the lines of the Craig–Wayne–Bourgain approach. The iterative scheme is combined with a smoothing procedure and interpolation estimates to ensure convergence in spaces of functions with only Sobolev regularity. The key step in [3,4] is to prove the “tame” estimates of the inverse operators in high Sobolev norms.

The aim of this paper is to generalize the previous approach in an abstract functional analytic setting, proving a Nash–Moser theorem “ready for applications” (Theorems 1–2), in particular, to prove the existence of lower dimensional, differentiable, invariant tori of PDEs with only differentiable nonlinearities.

The problem consists in solving a nonlinear equation

$$F(\varepsilon, \lambda, u) = 0 \tag{1}$$

where $\varepsilon \in [0, \varepsilon_0]$ is small, $\lambda \in \Lambda \subset \mathbb{R}^q$ with Λ open and bounded, and u belongs to some Banach space. Assuming that $F(0, \lambda, 0) = 0$, $\forall \lambda \in \Lambda$ (see hypothesis (F1) in the next subsection), the aim is to find, for ε_0 small enough, a function $u(\varepsilon, \lambda)$ with $u(0, \lambda) = 0$, which solves (1) for all (ε, λ) in a positive measure “Cantor-like” subset of $[0, \varepsilon_0] \times \Lambda$.

In typical applications the parameters λ can be “frequencies” – as in Section 3 – or vectors of a “resonant space” – when solving the “range equation” (called “P-equation”) obtained after a Lyapunov–Schmidt reduction, see e.g. [2,13].

We assume that the nonlinear map F satisfies abstract “tame” properties, see (F2)–(F4), which in applications are easily verified by differential and composition operators in scales of Sobolev functions, see e.g. [20] and Section 3.

The Nash–Moser theory has been well developed till now, see e.g. [20,21] and references therein. The main difference between the present Theorems 1–2 and the “standard” Nash–Moser theory is the abstract assumption (L) (or $(L_{\mathcal{K}})$) in Section 1.2: the “tame” estimates for the inverse operators hold only for *proper* subsets of the parameters. On the contrary, the standard Nash–Moser theory requires the invertibility of the derivative $\partial_u F$ in a *full* neighborhood, albeit in a tame sense. Indeed, in typical small divisors problems – as the search of finite dimensional tori for PDEs considered in this paper – an unbounded inverse $(\partial_u F)^{-1}$ does not exist for all the values of the parameters, but only for a “Cantor like” subset.

In [30] Zehnder observed that, for the convergence of the Nash–Moser iterative scheme, it is sufficient to assume only the existence of an “approximate inverse” of $\partial_u F$, which is required to be an exact inverse only at the solutions. This weaker property has been proved in [30], in a full neighborhood of the expected solution, for many conjugacy problems, thanks to algebraic features of the problem. In particular these theorems were sufficient to prove the existence of invariant Lagrangian tori for finite dimensional Hamiltonian systems. On the other hand, we remark that for lower dimensional tori – as for finite dimensional tori of PDEs – one cannot expect, by general arguments, the existence of an approximate inverse.

The existence of an inverse of the linearized operator for a Cantor set of parameters, together with estimates in scales of analytic functions, is the core of the Craig–Wayne–Bourgain method for analytic PDEs. With respect to these estimates, the main novelty of our assumptions is to require only tame estimates for the inverse, see (4).

By assumption (L), we ensure the invertibility of the linearized operators, at each step of the Nash–Moser iteration, only on smaller and smaller open sets of “non-resonant” parameters. A task of the iteration is to prove that, at the end of the recurrence, we have obtained a positive measure “Cantor-like” set of parameters where the solution is defined. This is the common scenario in these type of problems, see [2–5,7–12,14,19]. Such a property is implied by the abstract measure theoretical assumptions (7)–(8) in (L) and the rapid convergence of the iterative scheme, see the proof of Theorem 1. This abstract framework highlights specific constructions which were implicitly used in all the previous works. By means of a cut-off procedure at each step of the iteration, our solution can be smoothly extended in the whole space of parameters.

The abstract assumptions (F1)–(F4) and, in particular, hypothesis (L), make transparent the iterative procedures that, in specific contexts, have been performed in previous papers. In order to separate clearly the inductive argument and the measure estimates obtained in Theorem 1 (Section 2.5), we prove first the iterative Theorem 3 (Sections 2.2–2.4), where we do not assume hypothesis (L). We introduce an improvement with respect to the iterative scheme of [3,4] in order to prove the “ C^∞ -result” of Theorem 2 (Section 2.6).

Returning to PDE applications, a point of interest in developing a Nash–Moser theory for solutions with only Sobolev regularity is that, in presence of possibly very large clusters of small divisors (typical for higher dimensional PDEs), it is more natural to expect solutions with only Sobolev regularity, instead of analytic or Gevrey ones. An intuitive reason is that huge clusters of eigenvalues can produce strong resonance effects, having a consequence on the regularity of the solutions.

In Section 3 we present an application of Theorems 1–3 to the existence of periodic solutions of Klein–Gordon equations on a Zoll manifold \mathcal{M} , e.g. spheres, recently considered in [1], see Theorem 4. Other applications are given in [5]. The main issue for proving Theorem 4 is to verify the abstract assumption (L). For that, we exploit that the eigenvalues of $(-\Delta + V(x))^{1/2}$ on \mathcal{M} are contained in disjoint intervals, growing linearly to infinity, see Lemma 3.1. The corresponding geometry of the small divisors, see Lemma 3.6, suggests to look for solutions which are more regular in the time variable t than in the spatial variable x . Actually, a key idea is to look for solutions in the Sobolev scale (54) of time-periodic functions with values in a fixed Sobolev space $H^{s_1}(\mathcal{M})$, see Remark 3.2. Interestingly, many tools in our proof are reminiscent of those used in the normal form result in [1].

A final comment is in order: in [25] Moser introduced the related technique of analytic smoothing to approach the differentiable case. The idea is to first approximate, in a very accurate way, the differentiable Hamiltonian by analytic ones. Then one constructs, using an analytic KAM theorem, a sequence of analytic approximate invariant tori which actually converge to a differentiable torus of the original system. This powerful approach has been efficiently developed by Pöschel [26] and Salamon and Zehnder [28], to prove, for finite dimensional systems, the existence of invariant Lagrangian tori under the optimal finite regularity assumptions on the Hamiltonian. We think that this technique cannot, in general, be directly implemented in PDE applications when, for the presence of large clusters of small divisors, the resonance effects are so strong that the existence of analytic tori is doubtful. This is the main reason why, in this paper, we develop a Nash–Moser iterative procedure that is in spirit more similar to the original one in [24].

1.2. Functional setting and abstract Nash–Moser theorems

We consider a scale of Banach spaces $(X_s, \| \cdot \|_s)_{s \geq 0}$ such that

$$\forall s \leq s', \quad X_{s'} \subseteq X_s, \quad \|u\|_s \leq \|u\|_{s'}, \quad \forall u \in X_{s'},$$

and we define

$$X := \bigcap_{s \geq 0} X_s.$$

We assume that there are an increasing family $(E^{(N)})_{N \geq 0}$ of closed subspaces of X such that $\bigcup_{N \geq 0} E^{(N)}$ is dense in X_s for every $s \geq 0$, and that there are projectors

$$\Pi^{(N)} : X_0 \rightarrow E^{(N)} \text{ of range } E^{(N)}$$

satisfying, $\forall s \geq 0, \forall d \geq 0$,

- (S1) $\|\Pi^{(N)}u\|_{s+d} \leq C(s, d)N^d \|u\|_s, \forall u \in X_s$;
- (S2) $\|(I - \Pi^{(N)})u\|_s \leq C(s, d)N^{-d} \|u\|_{s+d}, \forall u \in X_{s+d}$

where $C(s, d)$ are positive constants. The projectors $\Pi^{(N)}$ can be seen as smoothing operators.

Note that by (S1) the norms $\|\cdot\|_s$ restricted to each $E^{(N)}$ are all equivalent. Moreover, by the density of $\bigcup_{N \geq 0} E^{(N)}$ in X_s , for $u \in X_s, \|u - \Pi^{(N)}u\|_s \rightarrow 0$ as $N \rightarrow \infty$.

Example (Sobolev scale). If X_s is the Sobolev space $H^s(\mathbb{T}^d), s \geq 0, \mathbb{T}^d := \mathbb{R}^d/2\pi\mathbb{Z}^d$, then $X = C^\infty(\mathbb{T}^d)$ and we can choose $E^{(N)} := \text{Span}\{e^{ik \cdot y}, k \in \mathbb{Z}^d, |k| \leq N\}$ and $\Pi^{(N)}$ the L^2 -orthogonal projector on $E^{(N)}$.

In every Banach scale with smoothing operators satisfying (S1)–(S2) as above, the following interpolation inequality holds.

Lemma 1.1 (Interpolation). $\forall 0 < s_1 < s_2$ there is $K(s_1, s_2) > 0$ such that, $\forall t \in [0, 1]$,

$$\|u\|_{ts_1+(1-t)s_2} \leq K(s_1, s_2) \|u\|_{s_1}^t \|u\|_{s_2}^{1-t}, \quad \forall u \in X_{s_2}.$$

Proof. Suppose $u \neq 0$. Setting $s := ts_1 + (1 - t)s_2$, we have, $\forall N \geq 1$,

$$\|u\|_s \leq \|\Pi^{(N)}u\|_s + \|u - \Pi^{(N)}u\|_s \stackrel{(S1),(S2)}{\leq} C(s_1, s_2) (N^{s-s_1} \|u\|_{s_1} + N^{s-s_2} \|u\|_{s_2})$$

and the result follows taking $N \geq 1$ as the integer part of $(\|u\|_{s_2}/\|u\|_{s_1})^{1/(s_2-s_1)}$. \square

We consider a C^2 map

$$F : [0, \varepsilon_0) \times \Lambda \times X_{s_0+\nu} \rightarrow X_{s_0} \tag{2}$$

where $s_0 \geq 0, \nu > 0, \varepsilon_0 > 0$ and Λ is a bounded open domain of \mathbb{R}^q . We assume

- (F1) $F(0, \lambda, 0) = 0, \forall \lambda \in \Lambda$,

and the ‘‘tame’’ properties:

$\exists S \in (s_0, \infty]$ such that $\forall s \in [s_0, S), \forall u \in X_{s+\nu}$ with $\|u\|_{s_0} \leq 2, \forall (\varepsilon, \lambda) \in [0, \varepsilon_0) \times \Lambda$,

- (F2)² $\|\partial_{(\varepsilon, \lambda)} F(\varepsilon, \lambda, u)\|_s \leq C(s)(1 + \|u\|_{s+\nu}), \|D_u F(\varepsilon, \lambda, 0)[h]\|_s \leq C(s)\|h\|_{s+\nu}$;
- (F3) $\|D_u^2 F(\varepsilon, \lambda, u)[h, v]\|_s \leq C(s)(\|u\|_{s+\nu}\|h\|_{s_0}\|v\|_{s_0} + \|v\|_{s+\nu}\|h\|_{s_0} + \|h\|_{s+\nu}\|v\|_{s_0})$;
- (F4) $\|\partial_{(\varepsilon, \lambda)} D_u F(\varepsilon, \lambda, u)[h]\|_s \leq C(s)(\|h\|_{s+\nu} + \|u\|_{s+\nu}\|h\|_{s_0})$.

From (F1)–(F4) we can deduce tame properties also for $F(\varepsilon, \lambda, u)$ and $(D_u F)(\varepsilon, \lambda, u)$, see Section 2.1.

The main assumption concerns the invertibility of the linear operators

$$L^{(N)}(\varepsilon, \lambda, u) := \Pi^{(N)} D_u F(\varepsilon, \lambda, u)|_{E^{(N)}}.$$

We consider two parameters $\mu \geq 0, \sigma \geq 0$, such that

$$\sigma > 4(\mu + \nu), \quad \bar{s} := s_0 + 4(\mu + \nu + 1) + 2\sigma < S. \tag{3}$$

For all $\gamma > 0$, we define appropriate subsets

² The symbol $\partial_{(\varepsilon, \lambda)}$ denotes either the partial derivative ∂_ε , or $\partial_{\lambda_i}, i = 1, \dots, q$.

$$J_{\gamma,\mu}^{(N)} \subseteq \{ (\varepsilon, \lambda, u) \in [0, \varepsilon_0] \times \Lambda \times E^{(N)} \mid L^{(N)}(\varepsilon, \lambda, u) \text{ is invertible and } \forall s \in \{s_0, \bar{s}\}, \\ \|L^{(N)}(\varepsilon, \lambda, u)^{-1}[h]\|_s \leq \frac{N^\mu}{\gamma} (\|h\|_s + \|u\|_s \|h\|_{s_0}), \forall h \in E^{(N)} \}. \tag{4}$$

Given $\mathbb{K} > 0$, we define

$$\mathcal{U}_{\mathbb{K}}^{(N)} := \{u \in C^1([0, \varepsilon_0] \times \Lambda, E^{(N)}) \mid \|u\|_{s_0} \leq 1, \|\partial_{(\varepsilon,\lambda)} u\|_{s_0} \leq \mathbb{K}\} \tag{5}$$

and, for all $u \in \mathcal{U}_{\mathbb{K}}^{(N)}$, we set

$$G_{\gamma,\mu}^{(N)}(u) := \{(\varepsilon, \lambda) \in [0, \varepsilon_0] \times \Lambda \mid (\varepsilon, \lambda, u(\varepsilon, \lambda)) \in J_{\gamma,\mu}^{(N)}\}. \tag{6}$$

We assume that

- (L) There exist $\sigma \geq 0, \mu \geq 0$ satisfying (3), $\bar{\gamma} > 0, M \in \mathbb{N}, C > 0$, such that:

$$(i) \quad \forall \gamma \in (0, \bar{\gamma}], \forall \varepsilon \in (0, \varepsilon_0], \quad |(G_{\gamma,\mu}^{(M)}(0))^c \cap ([0, \varepsilon] \times \Lambda)| \leq C\gamma\varepsilon. \tag{7}$$

- (ii) $\forall \gamma \in (0, \bar{\gamma}], \bar{\mathbb{K}} > 0, \exists \bar{\varepsilon} := \bar{\varepsilon}(\gamma, \bar{\mathbb{K}}) \in (0, \varepsilon_0]$ such that, $\forall \varepsilon \in (0, \bar{\varepsilon}], N' \geq N \geq M, u_1 \in \mathcal{U}_{\bar{\mathbb{K}}}^{(N)}, u_2 \in \mathcal{U}_{\bar{\mathbb{K}}}^{(N')}$ with $\|u_2 - u_1\|_{s_0} \leq N^{-\sigma}$,

$$|(G_{\gamma,\mu}^{(N')}(u_2))^c \setminus (G_{\gamma,\mu}^{(N)}(u_1))^c \cap ([0, \varepsilon] \times \Lambda)| \leq C \frac{\gamma\varepsilon}{N}. \tag{8}$$

Condition (7) says that $L^{(M)}(\varepsilon, \lambda, 0)$ is invertible for most parameters in $[0, \varepsilon] \times \Lambda$ and condition (8) says that the sets of “good” parameters $G_{\gamma,\mu}^{(N')}(u_2), G_{\gamma,\mu}^{(N)}(u_1)$ do not change too much for u_1, u_2 close enough in “low” Sobolev norm.

In applications, the verification of (L) strongly depends on the PDE. If in definition (4) we consider only $s = s_0$, then, by eigenvalue variation arguments, we can verify, for many PDEs, properties (i)–(ii). The main difficulty is to pass from informations on the eigenvalues of $L^{(N)}(\varepsilon, \lambda, u)$ to the interpolation estimates (4) in the high Sobolev norm $\|\cdot\|_{\bar{s}}$. Typically this requires “separation” properties on the small divisors of the PDE, see e.g. [10,13] and, in Section 3, Proposition 3.1 and Lemmas 3.4, 3.5.

Theorem 1. Assume (F1)–(F4), (L), (3). Then there is $C > 0$ and, $\forall \gamma \in (0, \bar{\gamma})$, there exists $\varepsilon_3 := \varepsilon_3(\gamma) \in (0, \varepsilon_0]$ and a C^1 map

$$u : [0, \varepsilon_3] \times \Lambda \rightarrow X_{s_0+v} \tag{9}$$

such that $u(0, \lambda) = 0$ and $F(\varepsilon, \lambda, u(\varepsilon, \lambda)) = 0$ except in a set \mathcal{C}_γ of Lebesgue measure $|\mathcal{C}_\gamma| \leq C\gamma\varepsilon_3$. Moreover, for all $\varepsilon \in (0, \varepsilon_3), |\mathcal{C}_\gamma \cap ([0, \varepsilon] \times \Lambda)| \leq C\gamma\varepsilon$.

Remark 1.1. As a consequence, if the freely chosen parameter $\gamma \rightarrow 0$, then³

$$\frac{|\mathcal{C}_\gamma \cap ([0, \varepsilon_3(\gamma)) \times \Lambda|}{|[0, \varepsilon_3(\gamma)) \times \Lambda|} \rightarrow 0,$$

namely the “bad” set \mathcal{C}_γ of parameters has asymptotically zero measure.

Remark 1.2. If u_1, u_2 are the maps in (9) associated respectively to γ_1, γ_2 , with $\gamma_1 < \gamma_2$, then $\mathcal{C}_{\gamma_1} \subset \mathcal{C}_{\gamma_2}$ and, for $\varepsilon \leq \min(\varepsilon_3(\gamma_1), \varepsilon_3(\gamma_2)), u_1$ and u_2 coincide outside \mathcal{C}_{γ_2} . This is easily seen from the construction of u in Section 2.

Remark 1.3. In the applications to PDEs with small divisors, the “good” parameters (ε, λ) such that $u(\varepsilon, \lambda)$ is a solution of $F(\varepsilon, \lambda, u) = 0$ form typically a Cantor-like set. The property that the solution can be extended to a C^1 function $u(\cdot, \cdot)$ defined on all the space of parameters could be seen as a Whitney extension theorem. However, here, it

³ Also $\varepsilon_3(\gamma) \rightarrow 0$.

is just a consequence of the use of a smooth cut-off function at each step of the Nash–Moser iteration. Such a property has been first proved in Pöschel [26] for KAM tori, and for PDEs it appeared in Kuksin [22,23] (actually it is sufficient to consider only Lipschitz extensions).

The conclusions of Theorem 1 can be strengthened under slightly stronger assumptions. Given a non-decreasing function $\mathcal{K} : [0, \infty) \rightarrow [1, \infty)$, we define the subsets

$$J_{\gamma,\mu,\mathcal{K}}^{(N)} \subseteq \{ (\varepsilon, \lambda, u) \in [0, \varepsilon_0] \times \Lambda \times E^{(N)} \mid L^{(N)}(\varepsilon, \lambda, u) \text{ is invertible and } \forall s \geq s_0, \\ \|L^{(N)}(\varepsilon, \lambda, u)^{-1}[h]\|_s \leq \mathcal{K}(s) \frac{N^\mu}{\gamma} (\|h\|_s + \|u\|_s \|h\|_{s_0}), \forall h \in E^{(N)} \}, \tag{10}$$

and the corresponding set $G_{\gamma,\mu,\mathcal{K}}^{(N)}(u)$ as in (6). We consider the stronger hypothesis

- $(L_{\mathcal{K}})$ The analogue to (L), with the sets $G_{\gamma,\mu}^{(N)}(\cdot)$ instead of $G_{\gamma,\mu,\mathcal{K}}^{(N)}(\cdot)$ in (7)–(8).

We remark that in typical PDEs applications, see Section 3, assumption $(L_{\mathcal{K}})$ is proved to hold for some \mathcal{K} with just slightly more effort than (L).

Theorem 2 (Regularity). Assume (F1)–(F4) with $S = \infty$ and $(L_{\mathcal{K}})$. Then the conclusion of Theorem 1 holds with $u \in C^1([0, \varepsilon_3(\gamma)] \times \Lambda; X)$ where $X := \bigcap_{s \geq 0} X_s$.

The proofs of Theorems 1 and 2 are based on an iterative Nash–Moser scheme that we describe in the next section.

2. The Nash–Moser iterative scheme

We shall deduce Theorem 1 from the following iterative result, where

$$N_n := N_0^{2^n}, \tag{11}$$

$N_0 \in \mathbb{N}$ will be chosen large enough (depending on γ), and $E_n, \Pi_n, J_{\gamma,\mu}^n$ are abbreviations for $E^{(N_n)}, \Pi^{(N_n)}, J_{\gamma,\mu}^{(N_n)}$ respectively. Given a set A and $\eta > 0$ we denote by $\mathcal{N}(A, \eta)$ the open neighborhood of A of width η (which is empty if A is empty).

Theorem 3. Assume (F1)–(F4) and (3). Then, for all $\gamma > 0$ there are $N_0 := N_0(\gamma), K_0(\gamma) > 0, \varepsilon_2 := \varepsilon_2(\gamma) \in (0, \varepsilon_0]$ and a sequence $(u_n)_{n \geq 0}$ of C^1 maps $u_n : [0, \varepsilon_2) \times \Lambda \rightarrow X_{s_0+\nu}$ with the following properties:

- $(P1)_n$ $u_n(\varepsilon, \lambda) \in E_n, u_n(0, \lambda) = 0, \|u_n\|_{s_0} \leq 1, \|\partial_{(\varepsilon,\lambda)} u_n\|_{s_0} \leq K_0(\gamma) N_0^{\sigma/2}.$
- $(P2)_n$ For $1 \leq k \leq n, \|u_k - u_{k-1}\|_{s_0} \leq N_k^{-\sigma-1}, \|\partial_{(\varepsilon,\lambda)}(u_k - u_{k-1})\|_{s_0} \leq N_k^{-1-\nu}.$
- $(P3)_n$ Let $A_n := \bigcap_{k=0}^n G_{\gamma,\mu}^{(N_k)}(u_{k-1})$ with $u_{-1} := 0.$ If $(\varepsilon, \lambda) \in \mathcal{N}(A_n, \gamma N_n^{-\sigma/2})$ then $u_n(\varepsilon, \lambda)$ solves the equation

$$(\mathcal{F}_n) \quad \Pi_n F(\varepsilon, \lambda, u) = 0.$$

- $(P4)_n$ $B_n := 1 + \|u_n\|_{\bar{s}}, B'_n := 1 + \|\partial_{(\varepsilon,\lambda)} u_n\|_{\bar{s}}$ (where \bar{s} is defined in (3)) satisfy

$$(i) \quad B_n \leq 2N_{n+1}^{\mu+\nu}, \quad (ii) \quad B'_n \leq 2N_{n+1}^{\mu+\nu+\sigma/2}.$$

The sequence $(u_n)_{n \geq 0}$ converges uniformly in $C^1([0, \varepsilon_2] \times \Lambda, X_{s_0+\nu})$ (endowed with the sup-norm of the map and its partial derivatives) to u with $u(0, \lambda) = 0$ and

$$(\varepsilon, \lambda) \in A_\infty := \bigcap_{n \geq 0} A_n \quad \Rightarrow \quad F(\varepsilon, \lambda, u(\varepsilon, \lambda)) = 0.$$

Note that in Theorem 3 we do *not* use any hypothesis on the linearized operators $L^{(N)}(\varepsilon, \lambda, u)$, in particular we do not assume (L). Then it could happen that $A_{n_0} = \emptyset$ for some n_0 . In such a case $u_n = u_{n_0}, \forall n \geq n_0$, and $A_\infty = \emptyset$. This is certainly the case if γ is chosen too large or μ too small.

Then, in Section 2.5, we show, assuming also (L), that the Lebesgue measure of the set A_∞ is large, deducing Theorem 1.

Remark 2.1. A minor difference between Theorem 3 and most Nash–Moser iterative schemes is that we solve exactly, at each step, the Galerkin approximate equations (\mathcal{F}_n) . It could be possible also to solve it only approximately, following a standard Newton iteration plus smoothing. This different procedure accounts for the classical quadratic convergence of our scheme where $N_n := N_0^{2^n}$ (see (11)) whereas a rapid convergence scheme in tame setting usually requires $N_n := e^{\alpha\chi^n}$ with $1 < \chi < 2$.

Let us give an outline of the convergence proof of Theorem 3. The sequence of approximate solutions u_n is constructed in Sections 2.2 and 2.3 solving the Galerkin approximate equations (\mathcal{F}_n) . First, in Section 2.2, we find u_0 as a fixed point of the nonlinear operator \mathcal{G}_0 , defined in (18). We prove that \mathcal{G}_0 is a contraction on a ball of $(E_0, \|\cdot\|_{s_0})$, taking ε sufficiently small. Then, in Section 2.3, by induction, we construct $u_{n+1} = u_n + h_{n+1}$ from u_n , finding h_{n+1} as a fixed point of \mathcal{G}_{n+1} defined in (29), see Lemma 2.4.

At the origin of the convergence of this Nash–Moser iteration, is the fact that L_{n+1}^{-1} satisfies the “tame” estimates (27), that the “remainder” term r_n is supported on the “high Fourier modes”, and that $R_n(h)$ is “quadratic” in h , see (22) for the definition of $L_{n+1}, r_n, R_n(h)$. Then r_n has a very small low norm $\|\cdot\|_{s_0}$ thanks to the smoothing estimates (S2), the tame estimate (F5), and the controlled growth of the high norms $\|u_n\|_{\bar{s}}$ of the approximate solutions given in $(P4)_n$ (see the proof of Lemma 2.4). Actually, the main point is to prove that $\|u_n\|_s$ does not grow, as $n \rightarrow \infty$, faster than some power of N_n independent of s , see Lemmas 2.5, 2.8, and Section 2.6.

We remark that the term r_n does not appear in a purely quadratic Newton scheme because it is a consequence of the smoothing procedure (projections). In the PDEs applications considered in [14,7–10] a term like r_n is proved to be small by decreasing the analyticity width at each step.

Finally, in Section 2.4, we conclude the convergence proof of Theorem 3. The proof of Theorem 1 is completed in Section 2.5 and, in Section 2.6, using the stronger assumption $(L_{\mathcal{K}})$ and the interpolation Lemma 1.1, we prove Theorem 2.

2.1. Preliminaries

From (F1)–(F3) we deduce, using Taylor formula, the tame properties: for $s \in [s_0, S)$, there is $C(s) > 0$ such that $\forall \|u\|_{s_0} \leq 2, \|h\|_{s_0} \leq 1$,

- (F5) $\|F(\varepsilon, \lambda, u)\|_s \leq C(s)(\varepsilon + \|u\|_{s+\nu})$;
- (F6) $\|(D_u F)(\varepsilon, \lambda, u)[h]\|_s \leq C(s)(\|u\|_{s+\nu}\|h\|_{s_0} + \|h\|_{s+\nu})$;
- (F7) $\|F(\varepsilon, \lambda, u+h) - F(\varepsilon, \lambda, u) - D_u F(\varepsilon, \lambda, u)[h]\|_s \leq C(s)(\|u\|_{s+\nu}\|h\|_{s_0}^2 + \|h\|_{s+\nu}\|h\|_{s_0})$.

We have the following perturbation lemmas:

Lemma 2.1. *Let A, R be linear operators in $E^{(N)}$ (A being possibly unbounded). Assume that A is invertible and that the following bounds hold for some $s > s_0$ and some $\alpha, \beta, \rho, \delta \geq 0$:*

$$\|A^{-1}v\|_{s_0} \leq \alpha\|v\|_{s_0}, \quad \|A^{-1}v\|_s \leq \alpha\|v\|_s + \beta\|v\|_{s_0}, \tag{12}$$

$$\|Rk\|_{s_0} \leq \delta\|k\|_{s_0}, \quad \|Rk\|_s \leq \delta\|k\|_s + \rho\|k\|_{s_0}. \tag{13}$$

If $\alpha\delta \leq 1/2$ then $A + R$ is invertible and

$$\|(A + R)^{-1}v\|_{s_0} \leq 2\alpha\|v\|_{s_0}, \quad \|(A + R)^{-1}v\|_s \leq 2\alpha\|v\|_s + 4(\beta + \alpha^2\rho)\|v\|_{s_0}. \tag{14}$$

Proof. The fact that $A + R$ is invertible and the first bound in (14) are standard: it is enough to write $A + R = (I + RA^{-1})A$ and to notice that $I + RA^{-1}$ is invertible because $\|RA^{-1}\|_{s_0} \leq 1/2$ and $E^{(N)}$ is a Banach space.

For the second bound, let $k := (A + R)^{-1}v$. We have $k = A^{-1}(v - Rk)$ and so

$$\|k\|_s \stackrel{(12)}{\leq} \alpha \|v - Rk\|_s + \beta \|v - Rk\|_{s_0} \stackrel{(13)}{\leq} \alpha \|v\|_s + \alpha \delta \|k\|_s + \alpha \rho \|k\|_{s_0} + \beta \|v\|_{s_0} + \beta \delta \|k\|_{s_0}.$$

Hence, since $\alpha \delta \leq 1/2$ and $\|k\|_{s_0} = \|(A + R)^{-1}v\|_{s_0} \leq 2\alpha \|v\|_{s_0}$, we obtain

$$\|k\|_s \leq 2(\alpha \|v\|_s + (2\alpha^2 \rho + \beta + 2\beta \delta \alpha) \|v\|_{s_0}) \leq 2\alpha \|v\|_s + 4(\beta + \alpha^2 \rho) \|v\|_{s_0}$$

proving the second inequality in (14). \square

Lemma 2.2. Let $(\varepsilon, \lambda, u) \in J_{\gamma, \mu}^{(N)}$ and $\|u\|_{s_0} \leq 1$. There is $c_0 := c_0(\bar{s}) > 0$ such that, if $|(\varepsilon', \lambda') - (\varepsilon, \lambda)| + \|h\|_{s_0} \leq c_0 \gamma N^{-(\mu+\nu)}$, $h \in E^{(N)}$, then $L^{(N)}(\varepsilon', \lambda', u + h)$ is invertible and $\forall v \in E^{(N)}$

$$\|L^{(N)}(\varepsilon', \lambda', u + h)^{-1}[v]\|_{s_0} \leq 4 \frac{N^\mu}{\gamma} \|v\|_{s_0}, \tag{15}$$

$$\|L^{(N)}(\varepsilon', \lambda', u + h)^{-1}[v]\|_{\bar{s}} \leq 4 \frac{N^\mu}{\gamma} \|v\|_{\bar{s}} + K \frac{N^{2\mu+\nu}}{\gamma^2} (\|u\|_{\bar{s}} + \|h\|_{\bar{s}}) \|v\|_{s_0}. \tag{16}$$

Proof. For brevity we set $z := (\varepsilon, \lambda)$, $z' := (\varepsilon', \lambda')$ and we apply Lemma 2.1 with $A = L^{(N)}(z, u)$ and $R = L^{(N)}(z', u + h) - L^{(N)}(z, u)$. Since $\|u\|_{s_0} \leq 1$, the bounds in (12) hold by (4) with $\alpha = 2\gamma^{-1}N^\mu$ and $\beta = \gamma^{-1}N^\mu \|u\|_{\bar{s}}$. By (F3) and (F4) we have, for $s = s_0$ or $s = \bar{s}$,

$$\begin{aligned} \|Rk\|_s &\leq |z' - z| C(s) (\|k\|_{s+\nu} + (\|u\|_{s+\nu} + \|h\|_{s+\nu}) \|k\|_{s_0}) \\ &\quad + C(s) ((\|u\|_{s+\nu} + \|h\|_{s+\nu}) \|h\|_{s_0} \|k\|_{s_0} + \|h\|_{s+\nu} \|k\|_{s_0} + \|h\|_{s_0} \|k\|_{s+\nu}) \\ &\leq C(s) N^\nu (|z' - z| + \|h\|_{s_0}) \|k\|_s + C(s) N^\nu (|z' - z| + \|h\|_{s_0}) (\|u\|_s + \|h\|_s) + \|h\|_s \|k\|_{s_0}. \end{aligned}$$

Hence, the bounds in (13) are satisfied with $\delta = C(\bar{s}, s_0) N^\nu (|z' - z| + \|h\|_{s_0})$ and $\rho = C(\bar{s}) (\|u\|_{\bar{s}} + 2\|h\|_{\bar{s}}) N^\nu$, for suitable positive constants $C(\bar{s}, s_0)$, $C(\bar{s})$. Then

$$\alpha \delta \leq 2\gamma^{-1} N^\mu C(\bar{s}, s_0) N^\nu c_0 \gamma N^{-\mu-\nu} = \frac{1}{2}, \quad \text{for } c_0 := \frac{1}{4C(\bar{s}, s_0)},$$

and Lemma 2.1 can be applied. Then we deduce (15)–(16) by (14). \square

The two following subsections are devoted to the construction of the sequence (u_n) of Theorem 3. Throughout this construction we shall take $N_0 := N_0(\gamma)$ large enough.

2.2. Initialization in the iterative Nash–Moser scheme

Let $A_0 := G_{\gamma, \mu}^{(N_0)}(0)$. By the definition (6), the parameters (ε, λ) are in A_0 if and only if $(\varepsilon, \lambda, 0) \in J_{\gamma, \mu}^{(N_0)}$. Then, by Lemma 2.2, if N_0 is large enough, $\forall (\varepsilon, \lambda) \in \mathcal{N}(A_0, 2\gamma N_0^{-\sigma/2})$, the operator $L^{(N_0)}(\varepsilon, \lambda, 0)$ is invertible and

$$\|L^{(N_0)}(\varepsilon, \lambda, 0)^{-1}\|_{s_0} \leq 4N_0^\mu \gamma^{-1}, \quad \|L^{(N_0)}(\varepsilon, \lambda, 0)^{-1}\|_{\bar{s}} \leq 4N_0^\mu \gamma^{-1} \tag{17}$$

(recall that $\sigma > 4(\mu + \nu)$ by (3)). Let us introduce the notations $L_0 := L^{(N_0)}(\varepsilon, \lambda, 0)$, $r_{-1} := \Pi_0 F(\varepsilon, \lambda, 0)$, and

$$R_{-1}(u) := \Pi_0 (F(\varepsilon, \lambda, u) - F(\varepsilon, \lambda, 0) - D_u F(\varepsilon, \lambda, 0)[u]).$$

A fixed point of

$$\mathcal{G}_0 : E_0 \rightarrow E_0, \quad \mathcal{G}_0(u) := -L_0^{-1}(r_{-1} + R_{-1}(u)), \tag{18}$$

is a solution of equation (\mathcal{F}_0) . If $0 \leq \varepsilon \leq \varepsilon_2(N_0, \gamma)$ is sufficiently small, \mathcal{G}_0 maps

$$\mathcal{B}_0 := \{u \in E_0 \mid \|u\|_{s_0} \leq \rho_0 := C_0 N_0^\mu \varepsilon \gamma^{-1}\}$$

into itself for some $C_0 := C_0(s_0)$. Indeed, by (17), (F5)–(F7), (S1), $\forall u \in \mathcal{B}_0$, $\|u\|_{s_0} \leq \rho_0$,

$$\begin{aligned} \|\mathcal{G}_0(u)\|_{s_0} &\leq 4N_0^\mu \gamma^{-1} (\|r_{-1}\|_{s_0} + \|R_{-1}(u)\|_{s_0}) \leq 4N_0^\mu \gamma^{-1} C(s_0) (\varepsilon + N_0^\nu \|u\|_{s_0}^2) \\ &\leq 4C(s_0) N_0^\mu \varepsilon \gamma^{-1} + 4N_0^{\mu+\nu} \gamma^{-1} C(s_0) \rho_0^2 \leq \rho_0 := C_0 N_0^\mu \varepsilon \gamma^{-1}, \end{aligned} \tag{19}$$

taking $C_0 := 8C(s_0)$ and ε so small that

$$4N_0^{\mu+\nu} \gamma^{-1} C(s_0) \rho_0 = 4N_0^{2\mu+\nu} \gamma^{-2} C(s_0) C_0 \varepsilon \leq \frac{1}{2}. \tag{20}$$

In the same way, if ε is small enough, we have by (F3), $\forall u \in \mathcal{B}_0$, $\|D\mathcal{G}_0(u)[h]\|_{s_0} \leq \|h\|_{s_0}/2$. Hence \mathcal{G}_0 is a contraction on $(\mathcal{B}_0, \|\cdot\|_{s_0})$ and it has a unique fixed point in this set.

Remark 2.2. The only difference between the proofs in this first step and those of Section 2.3 (and that is why this section is rather concise) is that the term r_{-1} is small thanks to the smallness of ε .

Let $\tilde{u}_0(\varepsilon, \lambda)$ denote the unique solution in \mathcal{B}_0 of (\mathcal{F}_0) , defined for all $(\varepsilon, \lambda) \in \mathcal{N}(A_0, 2\gamma N_0^{-\sigma/2})$. By (F1), if $(0, \lambda) \in \mathcal{N}(A_0, 2\gamma N_0^{-\sigma/2})$ then $\tilde{u}_0(0, \lambda) = 0$. Moreover, by the implicit function theorem, $\tilde{u}_0 \in C^1(\mathcal{N}(A_0, 2\gamma N_0^{-\sigma/2}); \mathcal{B}_0)$ and $\partial_{(\varepsilon, \lambda)} \tilde{u}_0 = -L^{(N_0)}(\varepsilon, \lambda, \tilde{u}_0)^{-1} [\Pi_0 \partial_{(\varepsilon, \lambda)} F(\varepsilon, \lambda, \tilde{u}_0)]$. By (F2), (15) and (20) we have $\|\partial_{(\varepsilon, \lambda)} \tilde{u}_0\|_{s_0} \leq K N_0^\mu \gamma^{-1}$.

Then we define the C^1 map $u_0 := \psi_0 \tilde{u}_0 : [0, \varepsilon_2] \times \Lambda \rightarrow E_0$ where the C^1 cut-off function $\psi_0 : [0, \varepsilon_2] \times \Lambda \rightarrow [0, 1]$ takes the values 1 on $\mathcal{N}(A_0, \gamma N_0^{-\sigma/2})$ and 0 outside $\mathcal{N}(A_0, 2\gamma N_0^{-\sigma/2})$, and $|\partial_{(\varepsilon, \lambda)} \psi_0| \leq C N_0^{\sigma/2} \gamma^{-1}$. The map u_0 satisfies property $(P3)_0$.

Moreover, $u_0(0, \lambda) = 0$, and, by the previous estimates, property $(P1)_0$ holds:

$$\|u_0\|_{s_0} \leq \frac{1}{2}, \quad \|\partial_{(\varepsilon, \lambda)} u_0\|_{s_0} \leq (C N_0^{\sigma/2} + K N_0^\mu) \gamma^{-1} \leq \frac{K_0(\gamma)}{2} N_0^{\sigma/2} \tag{21}$$

for some constant $K_0(\gamma)$. It remains to show $(P4)_0$. By (17), proceeding as in (19), provided that $4N_0^{\mu+\nu} \gamma^{-1} C(\bar{s}) \rho_0 \leq 1/2$, we have $\|\tilde{u}_0\|_{\bar{s}} \leq K(\gamma) N_0^\mu \varepsilon$, and, similarly,

$$\|\partial_{(\varepsilon, \lambda)} \tilde{u}_0\|_{\bar{s}} \stackrel{(16)}{\leq} 4 \frac{N_0^\mu}{\gamma} \|\partial_{(\varepsilon, \lambda)} F(\varepsilon, \lambda, \tilde{u}_0)\|_{\bar{s}} + K \frac{N_0^{2\mu+\nu}}{\gamma^2} \|\tilde{u}_0\|_{\bar{s}} \|\partial_{(\varepsilon, \lambda)} F(\varepsilon, \lambda, \tilde{u}_0)\|_{s_0} \leq K(\gamma) N_0^\mu.$$

Hence

$$\|\tilde{u}_0\|_{\bar{s}} \leq 2N_1^{\mu+\nu} \quad \text{and} \quad \|\partial_{(\varepsilon, \lambda)} \tilde{u}_0\|_{\bar{s}} \leq 2N_1^{\mu+\nu+(\sigma/2)}$$

for $N_0(\gamma)$ large enough (since $N_1 \geq N_0^2/2$ by (11)).

2.3. Iteration in the Nash–Moser scheme

In the previous subsection, we have proved that there is u_0 that satisfies $(P1)_0$ (more precisely (21)), $(P3)_0$ and $(P4)_0$. Note that $(P2)_0$ is automatically satisfied.

By induction, now suppose that we have already defined $u_n \in C^1([0, \varepsilon_2] \times \Lambda, E_n)$ satisfying the properties $(P1)_n$ – $(P4)_n$. We define the next approximation term u_{n+1} via the following modified Nash–Moser scheme.

For $h \in E_{n+1}$ we write

$$\Pi_{n+1} F(\varepsilon, \lambda, u_n(\varepsilon, \lambda) + h) = r_n + L_{n+1}[h] + R_n(h)$$

where

$$\begin{aligned} r_n &:= \Pi_{n+1} F(\varepsilon, \lambda, u_n), \quad L_{n+1} := L_{n+1}(\varepsilon, \lambda) := L^{(N_{n+1})}(\varepsilon, \lambda, u_n(\varepsilon, \lambda)), \\ R_n(h) &:= \Pi_{n+1} (F(\varepsilon, \lambda, u_n + h) - F(\varepsilon, \lambda, u_n) - D_u F(\varepsilon, \lambda, u_n)[h]). \end{aligned} \tag{22}$$

The “quadratic” term $R_n(h)$ is estimated, by (F7), as

$$\|R_n(h)\|_s \leq C(s)(\|u_n\|_{s+\nu}\|h\|_{s_0}^2 + \|h\|_{s+\nu}\|h\|_{s_0}). \tag{23}$$

By (P3)_n, if $(\varepsilon, \lambda) \in \mathcal{N}(A_n; \gamma N_n^{-\sigma/2})$ then u_n solves equation (\mathcal{F}_n) and so

$$r_n = \Pi_{n+1}F(\varepsilon, \lambda, u_n) - \Pi_n F(\varepsilon, \lambda, u_n) = \Pi_{n+1}(I - \Pi_n)F(\varepsilon, \lambda, u_n). \tag{24}$$

By (6) and (4), the operator $L_{n+1}(\varepsilon, \lambda)$ is invertible on the set $A_{n+1} = A_n \cap G_{\gamma, \mu}^{(N_{n+1})}(u_n)$. If $A_{n+1} = \emptyset$ we define $u_k := u_n, \forall k > n$. Otherwise we continue the iteration.

Note that, by (11), for N_0 large enough, we have the inclusion

$$\mathcal{N}(A_{n+1}, 2\gamma N_{n+1}^{-\sigma/2}) \subset \mathcal{N}(A_n, \gamma N_n^{-\sigma/2}). \tag{25}$$

Lemma 2.3. *For all $(\varepsilon, \lambda) \in \mathcal{N}(A_{n+1}, 2\gamma N_{n+1}^{-\sigma/2})$ the operator $L_{n+1}(\varepsilon, \lambda)$ is invertible,*

$$\|L_{n+1}^{-1}[v]\|_{s_0} \leq 4 \frac{N_{n+1}^\mu}{\gamma} \|v\|_{s_0}, \quad \forall v \in E_{n+1}, \tag{26}$$

and

$$\|L_{n+1}^{-1}[v]\|_{\bar{s}} \leq K(\gamma)N_{n+1}^\mu (\|v\|_{\bar{s}} + N_{n+1}^{2(\mu+\nu)}\|v\|_{s_0}), \quad \forall v \in E_{n+1}. \tag{27}$$

Proof. We apply Lemma 2.2. In fact, if $z := (\varepsilon, \lambda) \in \mathcal{N}(A_{n+1}, 2\gamma N_{n+1}^{-\sigma/2})$, there is $z' := (\varepsilon', \lambda') \in A_{n+1}$ (i.e. $(z', u_n(z')) \in J_{\gamma, \mu}^{(N_{n+1})}$) such that $|z - z'| \leq 2\gamma N_{n+1}^{-\sigma/2}$, and then

$$|z - z'| + \|u_n(z) - u_n(z')\|_{s_0} \stackrel{(P1)_n}{\leq} 2\gamma N_{n+1}^{-\sigma/2} (1 + K_0(\gamma)N_0^{\sigma/2}) \leq c_0\gamma N_{n+1}^{-(\mu+\nu)}$$

for $N_0 := N_0(\gamma)$ large enough, using (3) and (11). Thus (15) gives (26) and (16), together with the bound $\|u_n(z') - u_n(z)\|_s \leq \|u_n(z')\|_s + \|u_n(z)\|_s \leq 2B_n$, provides

$$\|L_{n+1}^{-1}[v]\|_{\bar{s}} \leq \frac{K'}{\gamma} N_{n+1}^\mu \left(\|v\|_{\bar{s}} + \frac{N_{n+1}^{\mu+\nu}}{\gamma} B_n \|v\|_{s_0} \right) \tag{28}$$

which implies (27) by (P4)_n. \square

Defining for $(\varepsilon, \lambda) \in \mathcal{N}(A_{n+1}, 2\gamma N_{n+1}^{-\sigma/2})$ the map

$$\mathcal{G}_{n+1} : E_{n+1} \rightarrow E_{n+1}, \quad \mathcal{G}_{n+1}(h) := -L_{n+1}^{-1}[r_n + R_n(h)], \tag{29}$$

the equation (\mathcal{F}_{n+1}) is equivalent to the fixed point problem $h = \mathcal{G}_{n+1}(h)$.

Lemma 2.4 (Contraction). *Let $(\varepsilon, \lambda) \in \mathcal{N}(A_{n+1}, 2\gamma N_{n+1}^{-\sigma/2})$. For $N_0(\gamma)$ large enough \mathcal{G}_{n+1} is a contraction in $\mathcal{B}_{n+1} := \{h \in E_{n+1} \mid \|h\|_{s_0} \leq \rho_{n+1} := N_{n+1}^{-\sigma-1}\}$ endowed with the norm $\|\cdot\|_{s_0}$.*

Proof. For all $(\varepsilon, \lambda) \in \mathcal{N}(A_{n+1}, 2\gamma N_{n+1}^{-\sigma/2})$, by (26) and (29), we have

$$\|\mathcal{G}_{n+1}(h)\|_{s_0} \leq 4N_{n+1}^\mu \gamma^{-1} (\|r_n\|_{s_0} + \|R_n(h)\|_{s_0}) \tag{30}$$

and r_n has the form (24) because of (25). Now, if $\|h\|_{s_0} \leq \rho_{n+1} := N_{n+1}^{-\sigma-1}$ then

$$\begin{aligned} \|r_n\|_{s_0} + \|R_n(h)\|_{s_0} &\stackrel{(S2), (23)}{\leq} K(N_n^{-(\bar{s}-s_0)}\|F(\varepsilon, \lambda, u_n)\|_{\bar{s}} + \|u_n\|_{s_0+\nu}\|h\|_{s_0}^2 + \|h\|_{s_0}\|h\|_{s_0+\nu}) \\ &\stackrel{(F5), (S1), (11)}{\leq} K'(N_{n+1}^{-(\bar{s}-s_0)/2}N_n^\nu B_n + N_{n+1}^\nu\|h\|_{s_0}^2) \\ &\stackrel{(P4)_n, (3)}{\leq} K_1(N_{n+1}^{-\mu-\sigma-2} + N_{n+1}^\nu\rho_{n+1}^2) \\ &\leq K_1\rho_{n+1}(N_{n+1}^{-\mu-1} + N_{n+1}^{\nu-\sigma-1}) \stackrel{(3)}{\leq} K_2\rho_{n+1}N_{n+1}^{-\mu-1}. \end{aligned}$$

As a consequence, for $N_0 := N_0(\gamma)$ large enough, we have

$$\|h\|_{s_0} \leq \rho_{n+1} \quad \Rightarrow \quad \|r_n\|_{s_0} + \|R_n(h)\|_{s_0} \leq \rho_{n+1} N_{n+1}^{-\mu} \gamma / 4. \tag{31}$$

Hence by (30), $\mathcal{G}_{n+1}(\mathcal{B}_{n+1}) \subset \mathcal{B}_{n+1}$.

Next, differentiating (29) with respect to h and using (22), we get, $\forall h \in \mathcal{B}_{n+1}$,

$$D_h \mathcal{G}_{n+1}(h)[v] = -L_{n+1}^{-1} \Pi_{n+1} (D_u F(\varepsilon, \lambda, u_n + h)[v] - D_u F(\varepsilon, \lambda, u_n)[v])$$

and

$$\|D_h \mathcal{G}_{n+1}(h)[v]\|_{s_0} \stackrel{(26),(F3),(P1)_n}{\leq} \frac{K}{\gamma} N_{n+1}^{\mu+\nu} \rho_{n+1} \|v\|_{s_0} \stackrel{(3)}{\leq} \frac{K}{\gamma} N_{n+1}^{-1} \|v\|_{s_0} \leq \frac{\|v\|_{s_0}}{2}$$

for N_0 large enough. Hence \mathcal{G}_{n+1} is a contraction in \mathcal{B}_{n+1} . \square

Let $\tilde{h}_{n+1} := \tilde{h}_{n+1}(\varepsilon, \lambda) \in E_{n+1}$ be the unique fixed point of \mathcal{G}_{n+1} , for $(\varepsilon, \lambda) \in \mathcal{N}(A_{n+1}, 2\gamma N_{n+1}^{-\sigma/2})$. Since \tilde{h}_{n+1} solves

$$U_{n+1}(\varepsilon, \lambda, h) := \Pi_{n+1} F(\varepsilon, \lambda, u_n(\varepsilon, \lambda) + h) = 0 \tag{32}$$

and $u_n(0, \lambda) \stackrel{(P1)_n}{=} 0$, we deduce, by (F1) and the uniqueness of the fixed point, that

$$(0, \lambda) \in \mathcal{N}(A_{n+1}, 2\gamma N_{n+1}^{-\sigma/2}) \quad \Rightarrow \quad \tilde{h}_{n+1}(0, \lambda) = 0. \tag{33}$$

Lemma 2.5 (Estimate in high norm). $\forall (\varepsilon, \lambda) \in \mathcal{N}(A_{n+1}, 2\gamma N_{n+1}^{-\sigma/2})$ we have

$$\|\tilde{h}_{n+1}\|_{\bar{s}} \leq N_{n+1}^{2(\mu+\nu)}. \tag{34}$$

Proof. By $\tilde{h}_{n+1} = \mathcal{G}_{n+1}(\tilde{h}_{n+1})$ we estimate

$$\|\tilde{h}_{n+1}\|_{\bar{s}} \stackrel{(27)}{\leq} K(\gamma) N_{n+1}^{\mu} (\|r_n\|_{\bar{s}} + \|R_n(\tilde{h}_{n+1})\|_{\bar{s}} + N_{n+1}^{2(\mu+\nu)} (\|r_n\|_{s_0} + \|R_n(\tilde{h}_{n+1})\|_{s_0})). \tag{35}$$

By (22) and (F5),

$$\|r_n\|_{\bar{s}} \leq K(\varepsilon + \|u_n\|_{\bar{s}+\nu}) \stackrel{(S1)}{\leq} K' N_n^{\nu} B_n \stackrel{(P4)_n, (11)}{\leq} K'' N_{n+1}^{\mu+\frac{3}{2}\nu}. \tag{36}$$

By (23) and (S1)

$$\begin{aligned} \|R_n(\tilde{h}_{n+1})\|_{\bar{s}} &\leq K(N_n^{\nu} B_n \|\tilde{h}_{n+1}\|_{s_0}^2 + N_{n+1}^{\nu} \|\tilde{h}_{n+1}\|_{s_0} \|\tilde{h}_{n+1}\|_{\bar{s}}) \\ &\leq N_{n+1}^{-\sigma-1} + K N_{n+1}^{\nu-\sigma-1} \|\tilde{h}_{n+1}\|_{\bar{s}}, \end{aligned} \tag{37}$$

using (P4)_n, $\|\tilde{h}_{n+1}\|_{s_0} \leq \rho_{n+1} := N_{n+1}^{-\sigma-1}$ (Lemma 2.4) and $\sigma > 4(\mu + \nu)$. Inserting in (35) the estimates (36)–(37) and (31) we get, for $N_0 := N_0(\gamma)$ large enough,

$$\|\tilde{h}_{n+1}\|_{\bar{s}} \leq \frac{1}{2} N_{n+1}^{2(\mu+\nu)} + K'(\gamma) N_{n+1}^{\mu+\nu-\sigma-1} \|\tilde{h}_{n+1}\|_{\bar{s}} \leq \frac{1}{2} N_{n+1}^{2(\mu+\nu)} + \frac{1}{2} \|\tilde{h}_{n+1}\|_{\bar{s}}$$

and (34) follows. \square

Lemma 2.6 (Estimates of the derivatives). The map \tilde{h}_{n+1} is in $C^1(\mathcal{N}(A_{n+1}, 2\gamma N_{n+1}^{-\sigma/2}); \mathcal{B}_{n+1})$ and

$$(i) \quad \|\partial_{(\varepsilon, \lambda)} \tilde{h}_{n+1}\|_{s_0} \leq \frac{1}{2} N_{n+1}^{-1-\nu}, \quad (ii) \quad \|\partial_{(\varepsilon, \lambda)} \tilde{h}_{n+1}\|_{\bar{s}} \leq N_{n+1}^{2(\mu+\nu)+\sigma}. \tag{38}$$

Proof. We set for brevity $z := (\varepsilon, \lambda)$. Recall that $U_{n+1}(z, \tilde{h}_{n+1}(z)) = 0$, see (32). The partial derivative $D_h U_{n+1}(z, \tilde{h}_{n+1}) = L^{(N_{n+1})}(z, u_n(z) + \tilde{h}_{n+1})$ is invertible by Lemma 2.2. Actually, arguing as in the proof of Lemma 2.3, since $\|\tilde{h}_{n+1}\|_{s_0} \leq N_{n+1}^{-\sigma-1} \ll c_0 \gamma N_{n+1}^{-(\mu+\nu)}$ for N_0 large, the estimates (15)–(16) imply

$$\|(D_h U_{n+1}(z, \tilde{h}_{n+1}))^{-1}[v]\|_{s_0} \leq 4\gamma^{-1} N_{n+1}^\mu \|v\|_{s_0}, \quad \forall v \in E_{n+1}, \tag{39}$$

$$\begin{aligned} \|(D_h U_{n+1}(z, \tilde{h}_{n+1}))^{-1}[v]\|_{\bar{s}} &\stackrel{(P4)_n}{\leq} K(\gamma) N_{n+1}^\mu (\|v\|_{\bar{s}} + N_{n+1}^{\mu+\nu} (N_{n+1}^{\mu+\nu} + \|\tilde{h}_{n+1}\|_{\bar{s}}) \|v\|_{s_0}) \\ &\stackrel{(34)}{\leq} K'(\gamma) N_{n+1}^\mu (\|v\|_{\bar{s}} + N_{n+1}^{3(\mu+\nu)} \|v\|_{s_0}). \end{aligned} \tag{40}$$

Then, by the implicit function theorem, $\tilde{h}_{n+1} \in C^1(\mathcal{N}(A_{n+1}, 2\gamma N_{n+1}^{-\sigma/2}); \mathcal{B}_{n+1})$ and

$$\partial_z \tilde{h}_{n+1} = -((D_h U_{n+1})(z, \tilde{h}_{n+1}))^{-1}(\partial_z U_{n+1})(z, \tilde{h}_{n+1}). \tag{41}$$

Now, using that $u_n(z)$ solves (\mathcal{F}_n) for $z \in \mathcal{N}(A_n, \gamma N_n^{-\sigma/2})$, we get by (25)

$$\partial_z U_{n+1}(z, h) = \Pi_{n+1}(\partial_z F(z, u_n + h) + D_u F(z, u_n + h)[\partial_z u_n]) \tag{42}$$

$$\begin{aligned} &= \Pi_{n+1}(\partial_z F)(z, u_n + h) - \Pi_n(\partial_z F)(z, u_n) \\ &\quad + \Pi_{n+1}(D_u F)(z, u_n + h)[\partial_z u_n] - \Pi_n(D_u F)(z, u_n)[\partial_z u_n] \\ &= \Pi_{n+1}((\partial_z F)(z, u_n + h) - (\partial_z F)(z, u_n)) \end{aligned} \tag{43}$$

$$+ \Pi_{n+1}((D_u F)(z, u_n + h) - (D_u F)(z, u_n))[\partial_z u_n] \tag{44}$$

$$+ \Pi_{n+1}(I - \Pi_n)(\partial_z F(z, u_n) + D_u F(z, u_n)[\partial_z u_n]). \tag{45}$$

Using (F4), (F3), $(P1)_n$, (S1), we get

$$\|(43)\|_{s_0} + \|(44)\|_{s_0} \leq K(\gamma) N_{n+1}^\nu \|\tilde{h}_{n+1}\|_{s_0} \leq K(\gamma) N_{n+1}^{\nu-\sigma-1} \tag{46}$$

by Lemma 2.4. By the smoothing estimate (S2), and (F2), (F3), (F6), $(P1)_n$,

$$\begin{aligned} \|(45)\|_{s_0} &\leq K(\gamma) N_n^{-(\bar{s}-s_0)} (1 + \|u_n\|_{\bar{s}+\nu} + \|\partial_z u_n\|_{\bar{s}+\nu}) \\ &\stackrel{(S1), (P4)_n}{\leq} K'(\gamma) N_n^{-(\bar{s}-s_0)} N_n^\nu N_{n+1}^{\nu+\mu+\frac{\sigma}{2}} \stackrel{(3)}{\leq} K'(\gamma) N_{n+1}^{-\frac{1}{2}(\mu+\nu+\sigma+4)}. \end{aligned} \tag{47}$$

From (41), (39), (46)–(47) we deduce estimate (38)(i) for $N_0(\gamma)$ large enough. To prove (38)(ii) we use (41) and estimate (40), whence

$$\begin{aligned} \|\partial_z \tilde{h}_{n+1}\|_{\bar{s}} &\leq K'(\gamma) N_{n+1}^\mu (\|\partial_z U_{n+1}(z, \tilde{h}_{n+1})\|_{\bar{s}} + N_{n+1}^{3(\mu+\nu)} \|\partial_z U_{n+1}(z, \tilde{h}_{n+1})\|_{s_0}) \\ &\leq \tilde{K}(\gamma) N_{n+1}^\mu (\|u_n\|_{\bar{s}+\nu} + \|\tilde{h}_{n+1}\|_{\bar{s}+\nu} + \|\partial_z u_n\|_{\bar{s}+\nu} + N_{n+1}^{2(\mu+\nu)}) \\ &\stackrel{(P4)_n, (34)}{\leq} K''(\gamma) N_{n+1}^{\mu+\nu} (N_{n+1}^{\mu+\nu+\sigma/2} + N_{n+1}^{2(\mu+\nu)}) \leq N_{n+1}^{2(\mu+\nu)+\sigma} \end{aligned} \tag{48}$$

for $N_0 := N_0(\gamma)$ large enough. To obtain (48) we have used (F2), (F6) and $(P1)_n$ in (42) to bound $\|\partial_z U_{n+1}(z, \tilde{h}_{n+1})\|_{\bar{s}}$ and (46)–(47) to bound $\|\partial_z U_{n+1}(z, \tilde{h}_{n+1})\|_{s_0}$. \square

We now define a C^1 -extension of $(\tilde{h}_{n+1})|_{A_{n+1}}$ onto the whole $[0, \varepsilon_2) \times \Lambda$.

Lemma 2.7 (Extension). *There is $h_{n+1} \in C^1([0, \varepsilon_2) \times \Lambda, \mathcal{B}_{n+1})$ satisfying*

$$h_{n+1}(0, \lambda) = 0, \quad \|h_{n+1}\|_{s_0} \leq N_{n+1}^{-\sigma-1}, \quad \|\partial_{(\varepsilon, \lambda)} h_{n+1}\|_{s_0} \leq N_{n+1}^{-\nu-1}$$

and that is equal to \tilde{h}_{n+1} on $\mathcal{N}(A_{n+1}, \gamma N_{n+1}^{-\sigma/2})$.

Proof. Let

$$h_{n+1}(\varepsilon, \lambda) := \begin{cases} \psi_{n+1}(\varepsilon, \lambda)\tilde{h}_{n+1}(\varepsilon, \lambda) & \text{if } (\varepsilon, \lambda) \in \mathcal{N}(A_{n+1}, 2\gamma N_{n+1}^{-\sigma/2}), \\ 0 & \text{if } (\varepsilon, \lambda) \notin \mathcal{N}(A_{n+1}, 2\gamma N_{n+1}^{-\sigma/2}) \end{cases} \quad (50)$$

where ψ_{n+1} is a C^1 cut-off function satisfying $0 \leq \psi_{n+1} \leq 1$, $\psi_{n+1} = 1$ on $\mathcal{N}(A_{n+1}, \gamma N_{n+1}^{-\sigma/2})$, $\psi_{n+1} = 0$ outside $\mathcal{N}(A_{n+1}, 2\gamma N_{n+1}^{-\sigma/2})$, and $|\partial_{(\varepsilon, \lambda)}\psi_{n+1}| \leq C\gamma^{-1}N_{n+1}^{\sigma/2}$.

By (33) and the definition of ψ_{n+1} we get $h_{n+1}(0, \lambda) = 0, \forall \lambda \in \Lambda$.

By the definition, $\|h_{n+1}\|_{s_0} \leq \|\tilde{h}_{n+1}\|_{s_0} \leq \rho_{n+1} = N_{n+1}^{-\sigma-1}$ by Lemma 2.4, and

$$\|\partial_{(\varepsilon, \lambda)}h_{n+1}\|_{s_0} \leq |\partial_{(\varepsilon, \lambda)}\psi_{n+1}|\|\tilde{h}_{n+1}\|_{s_0} + \|\partial_{(\varepsilon, \lambda)}\tilde{h}_{n+1}\|_{s_0} \leq N_{n+1}^{-\nu-1}$$

for $N_0(\gamma)$ large enough, by the previous bound on $|\partial_{(\varepsilon, \lambda)}\psi_{n+1}|$ and Lemma 2.6. \square

Finally we define $u_{n+1} \in C^1([0, \varepsilon_2) \times \Lambda, E_{n+1})$ as $u_{n+1} := u_n + h_{n+1}$. By Lemma 2.7, on $\mathcal{N}(A_{n+1}, \gamma N_n^{-\sigma/2})$ we have $h_{n+1} = \tilde{h}_{n+1}$ that solves Eq. (32) and so u_{n+1} solves Eq. (\mathcal{F}_{n+1}) . Hence property $(P3)_{n+1}$ holds. By Lemma 2.7, property $(P2)_{n+1}$ holds. By (21) and $(P2)_{n+1}$, for $N_0(\gamma)$ large enough,

$$\|u_{n+1}\|_{s_0} \leq \frac{1}{2} + \sum_{k=0}^n \|h_{k+1}\|_{s_0} \leq 1,$$

$$\|\partial_{(\varepsilon, \lambda)}u_{n+1}\|_{s_0} \leq \frac{K_0(\gamma)}{2}N_0^{\sigma/2} + \sum_{k=0}^n \|\partial_{(\varepsilon, \lambda)}h_{k+1}\|_{s_0} \leq K_0(\gamma)N_0^{\sigma/2}.$$

Moreover, still by Lemma 2.7 we have $u_{n+1}(0, \lambda) = 0, \forall \lambda \in \Lambda$, and also property $(P1)_{n+1}$ is verified. The induction of Theorem 3 is concluded in the following lemma.

Lemma 2.8. For $N_0 := N_0(\gamma)$ large, property $(P4)_{n+1}$ holds.

Proof. By the definition (50) and (34) we have $\|h_{n+1}\|_{\bar{s}} \leq N_{n+1}^{2(\mu+\nu)}$ and, by $(P4)_n$,

$$B_{n+1} \leq B_n + \|h_{n+1}\|_{\bar{s}} \leq 2N_{n+1}^{\mu+\nu} + N_{n+1}^{2(\mu+\nu)} \leq 2N_{n+2}^{\mu+\nu}$$

for $N_0 := N_0(\gamma)$ large enough. The second inequality follows similarly by

$$\begin{aligned} \|\partial_{(\varepsilon, \lambda)}h_{n+1}\|_{\bar{s}} &\leq |\partial_{(\varepsilon, \lambda)}\psi_{n+1}|\|\tilde{h}_{n+1}\|_{\bar{s}} + \|\partial_{(\varepsilon, \lambda)}\tilde{h}_{n+1}\|_{\bar{s}} \\ &\stackrel{(34), (38)}{\leq} \frac{C}{\gamma}N_{n+1}^{(\sigma/2)+2(\mu+\nu)} + N_{n+1}^{2(\mu+\nu)+\sigma} \leq \frac{3}{2}N_{n+2}^{\mu+\nu+\sigma/2} \end{aligned}$$

for $N_0 := N_0(\gamma)$ large enough. \square

2.4. Proof of Theorem 3 completed

The sequence of maps $u_n \in C^1([0, \varepsilon_2) \times \Lambda, E_n)$ converges in $C^1([0, \varepsilon_2) \times \Lambda, X_{s_0+\nu})$ to u , because $X_{s_0+\nu}$ is a Banach space and

$$\sum_{n \geq 0} \|u_n - u_{n-1}\|_{s_0+\nu} \stackrel{(S1)}{\leq} K \sum_{n \geq 0} N_n^\nu \|u_n - u_{n-1}\|_{s_0} \stackrel{(P2)_n}{\leq} K \sum_{n \geq 0} N_n^{\nu-\sigma-1} \leq \sum_{n \geq 0} N_n^{-1} < \infty$$

and, similarly, $\sum_{n \geq 0} \|\partial_{(\varepsilon, \lambda)}u_n - \partial_{(\varepsilon, \lambda)}u_{n-1}\|_{s_0+\nu} \leq K' \sum_{n \geq 0} N_n^{-1} < \infty$.

Finally, if $(\varepsilon, \lambda) \in A_\infty := \bigcap_{n \geq 0} A_n$ then $F(\varepsilon, \lambda, u) = 0$ because

$$F(\varepsilon, \lambda, u) = \Pi_n(F(\varepsilon, \lambda, u) - F(\varepsilon, \lambda, u_n)) + (I - \Pi_n)F(\varepsilon, \lambda, u) \xrightarrow{\|\cdot\|_{s_0}} 0$$

for $n \rightarrow \infty$.

2.5. Proof of Theorem 1

In order to deduce Theorem 1 from Theorem 3 it is sufficient to prove that assumption (L) implies $|A_\infty^c \cap ([0, \varepsilon) \times \Lambda)| \leq C\gamma\varepsilon, \forall \varepsilon \in (0, \varepsilon_3)$ for some $\varepsilon_3 \leq \varepsilon_2$.

Setting $G_n := G_{\gamma, \mu}^{(N_n)}(u_{n-1})$ for $n \geq 1$, and $G_0 := G_{\gamma, \mu}^{(N_0)}(0)$ we have $A_\infty = \bigcap_{n=0}^\infty G_n$. Its complementary set in $[0, \varepsilon) \times \Lambda$ is (here the apex c denotes the complementary in $[0, \varepsilon) \times \Lambda$)

$$A_\infty^c = \bigcup_{n=0}^\infty G_n^c \subset H^c \cup (G_0^c \setminus H^c) \cup \bigcup_{n=1}^\infty (G_n^c \setminus G_{n-1}^c)$$

where $H := G_{\gamma, \mu}^{(M)}(0)$, and $N_0 \geq M$. This implies, by (7)–(8), the measure estimate

$$|A_\infty^c| \leq |H^c| + |G_0^c \setminus H^c| + \sum_{n=1}^\infty |G_n^c \setminus G_{n-1}^c| \leq C\gamma\varepsilon(1 + M^{-1}) + \sum_{n=1}^\infty C\gamma\varepsilon N_{n-1}^{-1} \leq 2C\gamma\varepsilon$$

where we can apply (8) for

$$0 < \varepsilon \leq \varepsilon_3(\gamma) := \min(\varepsilon_1(\gamma, \bar{K}), \varepsilon_2(\gamma)) \quad \text{with } \bar{K} = K_0(\gamma)N_0^{\sigma/2}(\gamma)$$

because, by $(P1)_n$, we have $u_n \in \mathcal{U}_{\bar{K}}^{(N_n)}$ and $\|u_n - u_{n-1}\|_{s_0} \leq N_n^{-\sigma-1}$ by $(P2)_n$ for all n .

2.6. Proof of Theorem 2

Under $(L_{\mathcal{K}})$ we can apply Theorem 3 with $A_n = \bigcap_{k=0}^n G_{\gamma, \mu, \mathcal{K}}^{(N_k)}(u_{k-1})$, and the conclusion of Theorem 1 holds. We have to check that u is in $C^1([0, \varepsilon_3) \times \Lambda; X_{s'})$ for all $s' > 0$. For this, the main point is property $(P4)'_n$ below whose proof requires only small changes in the arguments used in Lemmas 2.5 and 2.6.

Lemma 2.9. For any $s > \bar{s}$, $B_n(s) := 1 + \|u_n\|_s, B'_n(s) := 1 + \|\partial_{(\varepsilon, \lambda)} u_n\|_s$ satisfy

$$(P4)'_n \quad B_n(s) \leq C(s)N_{n+1}^{\mu+\nu}, \quad B'_n(s) \leq C(s)N_{n+1}^{\mu+\nu+\sigma/2}.$$

This implies $\|h_n\|_s \leq 2C(s)N_{n+1}^{\mu+\nu}$.

Proof. First consider the map \tilde{h}_{n+1} defined on $\mathcal{N}(A_{n+1}, 2\gamma N_{n+1}^{-\sigma/2})$ after Lemma 2.4. Applying Lemma 2.2 with \bar{s} replaced by s , we get, for all $n \geq n_0(s)$ large enough,

$$\|L_{n+1}^{-1}[v]\|_s \leq K(\gamma, s)N_{n+1}^\mu (\|v\|_s + N_{n+1}^{\mu+\nu} B_n(s)\|v\|_{s_0}),$$

similarly to (28) in the proof of Lemma 2.3. Then, from $\tilde{h}_{n+1} = \mathcal{G}_{n+1}(\tilde{h}_{n+1})$ we derive (using (31))

$$\|\tilde{h}_{n+1}\|_s \leq K(\gamma, s)N_{n+1}^\mu (\|r_n\|_s + \|R_n(\tilde{h}_{n+1})\|_s) + K(\gamma, s)N_{n+1}^{\mu+\nu} B_n(s)\rho_{n+1}.$$

As in (36)–(37), we get

$$\|r_n\|_s \leq C(s)N_n^\nu B_n(s), \quad \|R_n(\tilde{h}_{n+1})\|_s \leq C(s)(N_n^\nu B_n(s)\rho_{n+1}^2 + N_{n+1}^\nu \|\tilde{h}_{n+1}\|_s \rho_{n+1}).$$

Since $\rho_{n+1} = N_{n+1}^{-\sigma-1}$, for $n \geq n_0(s)$ large enough, $K(\gamma, s)C(s)N_{n+1}^{\mu+\nu} \rho_{n+1} \leq 1/2$, and we derive from the previous inequalities, using (3) again, that

$$\|\tilde{h}_{n+1}\|_s \leq K'(\gamma, s)N_{n+1}^\mu N_n^\nu B_n(s) \leq N_{n+1}^{\mu+\nu} B_n(s).$$

Hence, as in Lemma 2.7, $\|h_{n+1}\|_s \leq N_{n+1}^{\mu+\nu} B_n(s)$ and

$$B_{n+1}(s) \leq (1 + N_{n+1}^{\mu+\nu}) B_n(s)$$

for $n \geq n_0(s)$, which implies that the sequence $(B_n(s) N_{n+1}^{-\mu-\nu})_n$ is bounded. This proves the first bound in $(P4)'_n$. With similar changes in Lemma 2.6 we obtain the second bound in $(P4)'_n$. \square

Now, consider any $s > s' > s_0$. By Lemma 1.1, writing $s' := (1 - t)s_0 + ts$, $t \in (0, 1)$,

$$\|h_n\|_{s'} \leq K(s_0, s) \|h_n\|_{s_0}^{1-t} \|h_n\|_s^t \leq K'(s) N_n^{-(\sigma+1)(1-t)} N_n^{2(\mu+\nu)t} = K'(s) N_n^{-1}$$

using $\|h_n\|_{s_0} \leq N_n^{-\sigma-1}$ (Lemma 2.4), $\|h_n\|_s \leq 2C(s) N_n^{2(\mu+\nu)}$ (Lemma 2.9), and choosing s large such that

$$t = \frac{s' - s_0}{s - s_0} = \frac{\sigma}{2(\mu + \nu) + \sigma + 1}.$$

Hence $\sum \|h_n\|_{s'} < \infty$ and, since $X_{s'}$ is a Banach space, $u \in X_{s'}$. We prove exactly in the same way that $\|\partial_{(\varepsilon, \lambda)} h_n\|_{s'} \leq C(s) N_n^{-1}$ and we derive that u is C^1 to $X_{s'}$. Since $s' \geq s_0$ is arbitrary we conclude that u is in $C^1([0, \varepsilon_3] \times \Lambda, X)$ where $X := \bigcap_{s \geq s_0} X_s$.

3. An application to PDEs

We present here an application of Theorems 1–2 to the search of periodic solutions of nonlinear wave equations

$$u_{tt} - \Delta u + V(x)u = \varepsilon f(\omega t, x, u), \quad x \in \mathcal{M}, \tag{51}$$

where \mathcal{M} is a d -dimensional connected, compact, Riemannian C^∞ -manifold without boundary, of Zoll type, namely the geodesic flow on the unit tangent bundle is periodic of minimal period $T > 0$. Classical examples of Zoll manifolds are the spheres and the symmetric compact spaces of rank 1 endowed with the canonical Riemannian structure. By results of Zoll, Funk, Guillemin and Weinstein, there exist many different metrics on the spheres, besides the standard one, whose geodesics are all simple closed curves of equal length, see e.g. [6].

In (51), Δ denotes the Laplace–Beltrami operator and we assume that the potential satisfies

$$V(x) \geq 0, \quad V \in C^p(\mathcal{M}) \text{ for some } p > \max\{2, d/2\}, \quad V \neq 0 \tag{52}$$

the forcing term f is differentiable only finitely many times, and $f(\omega t, x, u)$ is $(2\pi/\omega)$ -periodic in time, i.e. $f(\cdot, x, u)$ is 2π -periodic.

Remark 3.1. Wave equations on Zoll manifolds have been recently studied in [1] for time independent C^∞ -nonlinearities. The present techniques, written in the forced case for simplicity, apply also to such autonomous PDEs.

For $\varepsilon = 0$ the equilibrium $u = 0$ is a solution of (51). If $\varepsilon \neq 0$ and $f(t, x, 0) \neq 0$ then $u = 0$ is no more a solution. After a rescaling in time, we look for periodic solutions of

$$\omega^2 u_{tt} - \Delta u + V(x)u - \varepsilon f(t, x, u) = 0 \tag{53}$$

for $\varepsilon \neq 0$ small enough, in the Sobolev scale

$$H^s := H^s(\mathbb{T}, H^{s_1}(\mathcal{M}, \mathbb{R})), \quad s \geq 0, \tag{54}$$

of real, 2π -periodic in time functions with values in the Sobolev space $H^{s_1}(\mathcal{M}, \mathbb{R})$, where $s_1 \in (\max\{2, d/2\}, p]$. For $s_1 > d/2$, the Sobolev space $H^{s_1}(\mathcal{M}) \subset L^\infty(\mathcal{M})$ is a Banach algebra. Thanks to this property, for $s > 1/2$, each H^s is a Banach algebra too, see e.g. [2].

We define the closed subspaces of H^0

$$E^{(N)} := \left\{ u = \sum_{|l| \leq N} e^{ilt} u_l(x), \quad u_l \in H^{s_1}(\mathcal{M}, \mathbb{C}), \quad \bar{u}_l(x) = u_{-l}(x) \right\}$$

and the corresponding L^2 -orthogonal projectors $\Pi^{(N)}$. The smoothing properties (S1)–(S2) hold. Moreover

$$E^{(N)} \subset \bigcap_{s \geq 0} H^s = C^\infty(\mathbb{T}, H^{s_1}(\mathcal{M}, \mathbb{R})).$$

We need informations on the eigenvalues of the unbounded, self-adjoint operator

$$P := \sqrt{-\Delta + V(x)}$$

densely defined on $L^2(\mathcal{M}) := L^2(\mathcal{M}, \mathbb{C})$. The eigenvalues of P are the normal mode frequencies of the membrane. The spectrum $\sigma(P)$ of P is discrete, real and every $\lambda \in \sigma(P)$ is an eigenvalue of P of finite multiplicity. The following lemma, due to Colin de Verdière [12] and taken from [1], describes the asymptotic distribution of the eigenvalues of P when \mathcal{M} is a Zoll manifold.

Lemma 3.1. *If \mathcal{M} is a Zoll manifold, there are constants $\alpha \in \mathbb{R}$, $c_0 > 0$, $\delta \in (0, 1)$, $C_0 > 0$, and disjoint compact intervals $(I_j)_{j \geq 1}$ with I_1 at the left of I_2 , and*

$$I_j := \left[\frac{2\pi}{T} j + \alpha - \frac{c_0}{j^\delta}, \frac{2\pi}{T} j + \alpha + \frac{c_0}{j^\delta} \right], \quad j \geq 2, \tag{55}$$

such that the spectrum of P satisfies

$$\sigma(P) \subset \bigcup_{j \geq 1} I_j \quad \text{and} \quad \text{cardinality}(\sigma(P) \cap I_j) \leq C_0 j^{d-1} \tag{56}$$

(counted with multiplicity).

We call $\omega_{j,k}$, $1 \leq k \leq d_j$, $d_j \leq C_0 j^{d-1}$, the eigenvalues of P in each I_j . There is an orthonormal basis of $L^2(\mathcal{M})$ composed of corresponding eigenvectors $\varphi_{j,k}$. Since the manifold \mathcal{M} has no boundary, the Sobolev norms in $H^{s_1}(\mathcal{M}) := H^{s_1}(\mathcal{M}, \mathbb{C})$ can be defined as

$$\left\| \sum_{1 \leq j, 1 \leq k \leq d_j} v_{j,k} \varphi_{j,k} \right\|_{H^{s_1}(\mathcal{M})}^2 = \sum_{1 \leq j, 1 \leq k \leq d_j} (1 + \omega_{j,k}^2)^{s_1} |v_{j,k}|^2.$$

We consider forcing frequencies ω that are not in resonance with the normal mode frequencies $\omega_{j,k}$ of the membrane. More precisely, fixed some $\tau > d - 1$, we restrict to ω such that

$$|\omega^2 l^2 - \omega_{j,k}^2| \geq \frac{\gamma}{1 + |l|^\tau}, \quad \forall l \in \mathbb{Z}, j \in \mathbb{N}, k \in [1, d_j], \tag{57}$$

for some $\gamma \in (0, 1)$. By standard arguments, and taking into account (56), the non-resonance condition (57) is satisfied $\forall \omega \in (\omega_1, \omega_2)$ but a subset of measure $O(\gamma)$.

Theorem 4. *Let \mathcal{M} be a Zoll manifold and assume (52). Fix $0 < \omega_1 < \omega_2$ and $s_1 \in (\max\{2, d/2\}, p]$. Then*

(i) *Existence. There exists $s^* > 1/2$, $k^* \in \mathbb{N}$ such that: $\forall f \in C^{k^*}(\mathbb{T} \times \mathcal{M} \times \mathbb{R})$, $\forall \gamma \in (0, 1)$, there is $\varepsilon_0 := \varepsilon_0(\gamma) > 0$, a map*

$$u \in C^1([0, \varepsilon_0] \times (\omega_1, \omega_2), H^{s^*}) \quad \text{with } u(0, \omega) = 0,$$

such that $u(\varepsilon, \omega)$ is a solution of (53) for all $(\varepsilon, \omega) \in [0, \varepsilon_0] \times (\omega_1, \omega_2)$ except in a set \mathcal{C}_γ of Lebesgue measure $O(\gamma \varepsilon_0)$. Moreover, $\forall 0 < \varepsilon \leq \varepsilon_0(\gamma)$, $|\mathcal{C}_\gamma \cap ([0, \varepsilon] \times (\omega_1, \omega_2))| = O(\gamma \varepsilon)$.

(ii) *Regularity. If $f \in C^\infty(\mathbb{T} \times \mathcal{M} \times \mathbb{R})$ then*

$$u \in C^1([0, \varepsilon_0] \times (\omega_1, \omega_2), C^\infty(\mathbb{T}, H^{s_1}(\mathcal{M}, \mathbb{R}))).$$

The proof Theorem 4 is an application of Theorems 1 and 2.

Applying the linear operator $Q := (-\Delta + V(x) + I)^{-1}$ in (53), we look for zeros of

$$F(\varepsilon, \omega, u) := \omega^2 Qu_{tt} + u - Qu - \varepsilon Qf(t, x, u) \tag{58}$$

in the Sobolev scale $(H^s)_{s \geq 0}$.

By classical elliptic estimates the operator Q is regularizing of order 2 in the spatial variables: more precisely, we have

$$\|(-\Delta + V(x) + I)^{-1} u\|_{s, s'_1} \leq \|u\|_{s, \max(0, s'_1 - 2)}, \quad \forall u \in H^{s, s'_1}, \tag{59}$$

where $H^{s, s'_1} := H^s(\mathbb{T}, H^{s'_1}(\mathcal{M}, \mathbb{R}))$, $s'_1 \geq 0$, with Hilbert norms

$$\|u\|_{s, s'_1}^2 = \sum_{l \in \mathbb{Z}} \langle l \rangle^{2s} \|u_l\|_{H^{s'_1}(\mathcal{M})}^2, \quad \langle l \rangle := \max(1, |l|). \tag{60}$$

When $s'_1 = s_1$ we shall more simply denote $\| \cdot \|_{s, s'_1} = \| \cdot \|_{s, s_1} = \| \cdot \|_s$ the norm in H^s . Finally, given a linear operator L in H^{s, s'_1} , $\|L\|_{s, s'_1}$ denotes the associated operatorial norm.

Lemma 3.2. *If $f \in C^k(\mathbb{T} \times \mathcal{M} \times \mathbb{R})$ with $S := k - s_1 - 2 > s_0 > 1/2$, the map F satisfies (2), with $v = 2$, $\Lambda = (\omega_1, \omega_2) \subset \mathbb{R}$, and (F1) holds. Moreover F is C^2 and the tame properties (F2)–(F4) hold for all $s \in [s_0, S]$.*

Proof. Use standard properties for the composition operators in Sobolev spaces, see e.g. [3]. \square

There remains to verify properties (L) and $(L_{\mathcal{K}})$ concerning the linearized operators

$$L^{(N)}(u)[v] = Q\mathcal{L}^{(N)}(u)[v] = \omega^2 Qv_{tt} + v - Qv - \varepsilon \Pi^{(N)} Q(b(t, x)v), \quad v \in E^{(N)}, \tag{61}$$

where $b(t, x) := (\partial_u f)(t, x, u(t, x))$ and

$$\mathcal{L}^{(N)}(u)[v] := \mathcal{L}^{(N)}(\varepsilon, \omega, u)[v] := \omega^2 v_{tt} - \Delta v + V(x)v - \varepsilon \Pi^{(N)}(b(t, x)v).$$

We shall prove in detail property $(L_{\mathcal{K}})$, assuming that f is in C^∞ . The proof of (L) is similar.

Proposition 3.1. *For all $\tau > 0$, $\tau_0 > 1$, there exist constants $\mu_0 \geq 0$, $\tilde{s} > 1/2$, a non-decreasing function $\mathcal{K} : \mathbb{R}_+ \rightarrow [1, \infty)$ and, $\forall \gamma > 0$, a constant $\eta(\gamma) > 0$ such that: if $\varepsilon(\|b\|_{\tilde{s}} + 1) \leq \eta(\gamma)$,*

$$\left| \omega l - \frac{2\pi}{T} p \right| \geq \frac{\gamma}{(1 + |l|)^{\tau_0}}, \quad \forall (l, p) \in \mathbb{Z}^2 \setminus \{(0, 0)\}, \tag{62}$$

and

$$\forall 1 \leq K \leq N, \quad \|(\mathcal{L}^{(K)}(u))^{-1}\|_{0,0} \leq 4 \frac{K^\tau}{\gamma}, \tag{63}$$

then, $\forall s \geq \tilde{s}$,

$$\|(\mathcal{L}^{(N)}(u))^{-1} h\|_{s,0} \leq \frac{\mathcal{K}(s)}{\gamma} N^{\mu_0} (\|h\|_{s,0} + \|b\|_s \|h\|_{\tilde{s},0}), \quad \forall h \in E^{(N)}. \tag{64}$$

Postponing the proof of Proposition 3.1 to the end of the section, we complete the proof of property $(L_{\mathcal{K}})$. By a bootstrap type argument, (64) implies a similar estimate for $\|(L^{(N)}(u))^{-1} h\|_s$.

Lemma 3.3. *Under the assumptions of Proposition 3.1, $\forall s \geq \tilde{s}$,*

$$\|(L^{(N)}(u))^{-1} h\|_s \leq \frac{\mathcal{K}(s)}{\gamma} N^\mu (\|h\|_s + \|u\|_s \|h\|_{\tilde{s}}), \quad \forall h \in E^{(N)},$$

where $\mu := \mu_0 + s_1 + 2$, taking, if necessary, $\mathcal{K}(s)$ larger.

Proof. Setting $h := (L^{(N)}(u))[v] = v + Q(\omega^2 v_{tt} - v - \varepsilon \Pi^{(N)}(bv))$ in (61), we estimate

$$\begin{aligned} \|v\|_s &= \|Q(-\omega^2 v_{tt} + v + \varepsilon \Pi^{(N)}(bv)) + h\|_s \\ &\stackrel{(59)}{\leq} \|- \omega^2 v_{tt} + v + \varepsilon \Pi^{(N)}(bv)\|_{s,s_1-2} + \|h\|_s \leq CN^2 \|v\|_{s,s_1-2} + \varepsilon C(s) \|b\|_s \|v\|_{\tilde{s},s_1-2} + \|h\|_s \end{aligned}$$

by interpolation inequality (82). Using $\|v\|_{\tilde{s},s_1-2} \leq CN^2 \|v\|_{\tilde{s},\max(0,s_1-4)} + \|h\|_{\tilde{s}}$, and iterating, we obtain

$$\|v\|_s \leq C(s)N^{s_1+2} (\|v\|_{s,0} + \|h\|_s + \|b\|_s \|v\|_{\tilde{s},0} + \|b\|_s \|h\|_{\tilde{s}}). \tag{65}$$

Since $v = (\mathcal{L}^{(N)}(u))^{-1}(-\Delta + V(x) + I)h$,

$$\|v\|_{s,0} \stackrel{(64)}{\leq} \frac{\mathcal{K}(s)}{\gamma} N^{\mu_0} (\|h\|_{s,2} + \|b\|_s \|h\|_{\tilde{s},2}) \leq \frac{\mathcal{K}'(s)}{\gamma} N^{\mu_0} (\|h\|_s + \|u\|_s \|h\|_{\tilde{s}}), \tag{66}$$

using $s_1 \geq 2$ and $\|b\|_s = \|(\partial_u f)(t, x, u)\|_s \leq C(s)(1 + \|u\|_s)$. By (65) and (66) the lemma follows. \square

To conclude the proof of property $(L_{\mathcal{K}})$ we have to define $J_{\gamma,\mu,\mathcal{K}}^{(N)}$ and show the measure estimates (7) and (8). Fix $\tau \geq d + 2$ (the exponent in (57) and in (63)), $\tau_0 > 1$ (the exponent in (62)) and define

$$G := \{(\varepsilon, \omega) \in [0, \varepsilon_0] \times (\omega_1, \omega_2) \mid \omega \text{ satisfies (57) and (62)}\}.$$

By standard arguments $|G^c \cap ([0, \varepsilon] \times (\omega_1, \omega_2))| = O(\gamma\varepsilon)$. We also define

$$J_{\gamma,\mu,\mathcal{K}}^{(N)} := \{(\varepsilon, \omega, u) \in [0, \varepsilon_0] \times (\omega_1, \omega_2) \times E^{(N)} \mid (\varepsilon, \omega) \in G, \|u\|_{s_0} \leq 1, \text{ and (63) holds}\}.$$

By Proposition 3.1 and Lemma 3.3, for $\varepsilon_0 > 0$ small enough, the inclusion (10) is satisfied, with

$$\mu := \mu_0 + s_1 + 2 \quad \text{and} \quad s_0 > \max\{1/2, \tilde{s}\}.$$

Next, given a function $u \in \mathcal{U}_{\mathbb{K}}^{(N)}$ (see (5)), $\mathbb{K} > 0$, the set $G_{\gamma,\mu,\mathcal{K}}^{(N)}(u)$ defined as in (6) can be written as

$$G_{\gamma,\mu,\mathcal{K}}^{(N)}(u) = \bigcap_{1 \leq K \leq N} B_K(u) \cap G \tag{67}$$

where

$$B_K(u) := \left\{ (\varepsilon, \omega) \in [0, \varepsilon_0] \times (\omega_1, \omega_2) \mid \|(\mathcal{L}^{(K)}(u))^{-1}\|_{0,0} \leq 4 \frac{K^\tau}{\gamma} \right\}.$$

Lemma 3.4. *If $\varepsilon_0 \gamma^{-1} M^\tau \leq c$ is small enough, then $G_{\gamma,\mu,\mathcal{K}}^{(M)}(0) = G$. Hence (7) holds.*

Proof. We have $\mathcal{L}^{(K)}(u) = D^{(K)} + T^{(K)}$ with

$$D^{(K)}h := \omega^2 h_{tt} - \Delta h + V(x)h \quad \text{and} \quad T^{(K)}h := -\varepsilon \Pi^{(K)}(bh). \tag{68}$$

If ω satisfies (57) then $\|(D^{(K)})^{-1}\|_{0,0} \leq 2K^\tau \gamma^{-1}$. Moreover $\|T^{(K)}\|_{0,0} \leq C\varepsilon \|b\|_{\tilde{s}}$. By Lemma 2.1, if $2M^\tau \gamma^{-1} C\varepsilon \|b\|_{\tilde{s}} < 1/2$, then, $\forall 1 \leq K \leq M$, $\mathcal{L}^{(K)}(u)$ is invertible in $H^{0,0}$ and $\|(\mathcal{L}^{(K)}(u))^{-1}\|_{0,0} \leq 4K^\tau \gamma^{-1}$. \square

We fix $\sigma > \max\{4(\mu + 2), d + 2\}$ (the first condition is (3) with $\nu = 2$).

Lemma 3.5. *The measure estimate (8) holds.*

Proof. Fix $\tilde{\varepsilon} \in (0, \varepsilon_0]$. As in the proof of Lemma 3.4, for all $N, N' \leq N_{\tilde{\varepsilon}} := (c\gamma/\tilde{\varepsilon})^{1/\tau}$, for all $u_1 \in \mathcal{U}_{\mathbb{K}}^{(N')}$, $u_2 \in \mathcal{U}_{\mathbb{K}}^{(N')}$, it results $G_{\gamma,\mu,\mathcal{K}}^{(N)}(u_1) = G_{\gamma,\mu,\mathcal{K}}^{(N')}(u_2) = G$ and thus (8) is trivially satisfied in such cases. Given a set $A \subset (0, \varepsilon_0] \times [\omega_1, \omega_2]$ let A^c represent the complementary in $(0, \tilde{\varepsilon}] \times [\omega_1, \omega_2]$. For $N' \geq N$,

$$\begin{aligned} (G_{\gamma,\mu,\mathcal{K}}^{(N')})^c \setminus (G_{\gamma,\mu,\mathcal{K}}^{(N)})^c &= (G_{\gamma,\mu,\mathcal{K}}^{(N')})^c(u_2) \cap G_{\gamma,\mu,\mathcal{K}}^{(N)}(u_1) \\ &\subset \left[\bigcup_{K \leq N} (B_K^c(u_2) \cap B_K(u_1) \cap G) \right] \cup \left[\bigcup_{K > N} B_K^c(u_2) \cap G \right]. \end{aligned}$$

As we have just seen, if $K \leq N_{\tilde{\varepsilon}}$ then $B_K^c(u_2) \cap G = \emptyset$. Hence it is enough to prove that, if $\|u_1 - u_2\|_{s_0} \leq N^{-\sigma}$, then

$$\mathcal{B} := \sum_{K \leq N} |B_K^c(u_2) \cap B_K(u_1)| + \sum_{K > \max\{N, N_{\tilde{\varepsilon}}\}} |B_K^c(u_2)| \leq \bar{C} \frac{\gamma \tilde{\varepsilon}}{N}. \tag{69}$$

Since $\mathcal{L}^{(K)}(u)$ is selfadjoint in $H^{0,0}$ and $(CI + \mathcal{L}^{(K)}(u))^{-1}$ is compact for some large C depending on K , $H^{0,0}$ has an orthonormal basis of eigenvectors of $\mathcal{L}^{(K)}(u)$, and $\|(\mathcal{L}^{(K)}(u))^{-1}\|_{0,0}$ is the inverse of the eigenvalue of smallest modulus.

Since $\|\mathcal{L}^{(K)}(u_2) - \mathcal{L}^{(K)}(u_1)\|_{0,0} = O(\varepsilon\|u_2 - u_1\|_{s_0}) = O(\varepsilon N^{-\sigma})$, if one of the eigenvalues of $\mathcal{L}^{(K)}(u_2)$ is in $[-4\gamma K^{-\tau}, 4\gamma K^{-\tau}]$ then, by the variational characterization of the eigenvalues of $\mathcal{L}^{(K)}(u)$, one of the eigenvalues of $\mathcal{L}^{(K)}(u_1)$ is in $[-4\gamma K^{-\tau} - C\varepsilon N^{-\sigma}, 4\gamma K^{-\tau} + C\varepsilon N^{-\sigma}]$. As a result

$$\begin{aligned} B_K^c(u_2) \cap B_K(u_1) &\subset \{(\varepsilon, \omega) \mid \exists \text{ at least one eigenvalue of } \mathcal{L}^{(K)}(\varepsilon, \omega, u_1) \\ &\text{with modulus in } [4\gamma K^{-\tau}, 4\gamma K^{-\tau} + C\varepsilon N^{-\sigma}]\}. \end{aligned}$$

By a simple eigenvalue variation argument, as is Lemma 3.2 of [4], we have that: if ε is small enough (depending on \bar{K}), if I is a compact interval in $[-\gamma, \gamma]$ of length $|I|$, then

$$|\{\omega \in [\omega_1, \omega_2] \text{ s.t. at least one eigenvalue of } \mathcal{L}^{(K)}(\varepsilon, \omega, u_1) \text{ belongs to } I\}| \leq C \frac{K^d |I|}{\omega_1}. \tag{70}$$

As a consequence $|\{\omega \mid (\varepsilon, \omega) \in B_K^c(u_2) \cap B_K(u_1)\}| \leq C\varepsilon N^{-\sigma} K^d / \omega_1$ for each $\varepsilon \in (0, \tilde{\varepsilon}]$, whence $|B_K^c(u_2) \cap B_K(u_1)| \leq C'\tilde{\varepsilon}^2 K^d N^{-\sigma}$. Moreover, still by (70), $|B_K^c(u_2)| \leq C\tilde{\varepsilon} K^d \gamma K^{-\tau} / \omega_1 \leq C'\tilde{\varepsilon} \gamma K^{d-\tau}$. Hence \mathcal{B} defined in (69) satisfies

$$\begin{aligned} \mathcal{B} &\leq C\tilde{\varepsilon}^2 \left(\sum_{K \leq N} K^d \right) N^{-\sigma} + C\tilde{\varepsilon} \gamma \left(\sum_{K > \max\{N, N_{\tilde{\varepsilon}}\}} K^{d-\tau} \right) \\ &\leq C\tilde{\varepsilon}^2 N^{d+1-\sigma} + C'\tilde{\varepsilon} \gamma (\max\{N, N_{\tilde{\varepsilon}}\})^{d+1-\tau} \leq \bar{C} \gamma \tilde{\varepsilon} N^{-1}, \end{aligned}$$

for $\sigma, \tau \geq d + 2$. This proves the measure estimate (8). \square

We have verified all the assumptions of Theorems 1–2 whence Theorem 4 follows.

Proof of Proposition 3.1. Fixed $\rho > 0$, we consider the “singular” \mathcal{S} and “regular” \mathcal{R} sites

$$\mathcal{S} := \{l \in \mathbb{Z} \cap [-N, N] \mid \|D_l(\omega)^{-1}\|_{\mathcal{L}(L^2(\mathcal{M}))} > \rho^{-1}\}, \quad \mathcal{R} := \mathcal{S}^c,$$

where $D_l(\omega) := -\omega^2 l^2 - \Delta + V(x)$ are self-adjoint, unbounded operators, densely defined in $L^2(\mathcal{M})$. \square

The singular sites \mathcal{S} are “separated” like in the 1-dimensional wave equations.

Lemma 3.6. Assume the diophantine condition (62). Then $\exists c(\gamma) > 0$, $\delta_0 := \delta_0(\tau_0, \delta) \in (0, 1)$, such that $\forall l, l' \in \mathcal{S}$ with $l \neq l'$, we have $|l - l'| \geq c(\gamma)(|l| + |l'|)^{\delta_0}$.

Proof. Suppose that $l_1, l_2 > 0$; if $l_1, l_2 \in \mathcal{S}$ then there are $j_1, k_1 \in [1, d_{j_1}]$, $j_2, k_2 \in [1, d_{j_2}]$ such that

$$|\omega l_1 - \omega_{j_1, k_1}| \leq C \frac{\rho}{|l_1|}, \quad |\omega l_2 - \omega_{j_2, k_2}| \leq C \frac{\rho}{|l_2|}.$$

Using the spectral asymptotics in (55), and the diophantine condition, we get, if $l_1 \neq l_2$,

$$\frac{\gamma}{(1 + |l_1 - l_2|)^{\tau_0}} \leq \left| \omega(l_1 - l_2) - \frac{2\pi}{T}(j_1 - j_2) \right| \leq \frac{c}{|l_1|^\delta} + \frac{c}{|l_2|^\delta}$$

and the thesis follows, using $|l_1| + |l_2| \leq 2 \min(|l_1|, |l_2|) + |l_1 - l_2|$. \square

Remark 3.2. According to the definitions in [13,14,7] the singular sites are the integers (l, j, k) such that $|\omega^2 l^2 + \omega_{j,k}^2| < \rho$, where $\omega_{j,k}^2$ are the eigenvalues of $-\Delta + V(x)$. Due to the multiplicity of such eigenvalues they may form very large clusters. However, the previous lemma shows good separation properties for their projection in time-Fourier indices. This is the main motivation for working with the spaces H^s defined in (54). This setting enables to proceed similarly to the 1-dimensional wave equation; the only difference is that, after decomposing in time Fourier series, we get matrices of spatial operators.

Now, we shall follow closely the procedure in [4], which is here much simpler because the singular sites are singletons (in time-Fourier indices), see Lemma 3.6. A difference is that, in order to prove the C^∞ -result, Theorem 4(ii), we need to assume $\varepsilon(\|b\|_{\tilde{s}} + 1)$ small (independently of s).

According to the orthogonal decomposition $E^{(N)} := E_R \oplus E_S$, where

$$E_R := \left\{ u = \sum_{l \in R} e^{ilt} u_l(x) \in E^{(N)} \right\} \quad \text{and} \quad E_S := \left\{ u = \sum_{l \in S} e^{ilt} u_l(x) \in E^{(N)} \right\},$$

for $(\varepsilon, \omega) \in G$, we represent $\mathcal{L}^{(N)} := \mathcal{L}^{(N)}(u)$ as the self-adjoint block matrix (of spatial operators)

$$\mathcal{L}^{(N)} = \begin{pmatrix} \Pi_R \mathcal{L}|_{E_R} & \Pi_R \mathcal{L}|_{E_S} \\ \Pi_S \mathcal{L}|_{E_R} & \Pi_S \mathcal{L}|_{E_S} \end{pmatrix} = \begin{pmatrix} \mathcal{L}_R & \mathcal{L}_R^S \\ \mathcal{L}_S^R & \mathcal{L}_S \end{pmatrix}$$

where $\Pi_S : E^{(N)} \rightarrow E_S, \Pi_R : E^{(N)} \rightarrow E_R$ denote the corresponding orthogonal projectors. It results that $\mathcal{L}_R^S = (\mathcal{L}_S^R)^\dagger, \mathcal{L}_R^\dagger := \mathcal{L}_R, \mathcal{L}_S^\dagger = \mathcal{L}_S$. We fix

$$\tilde{s} := 1 + (\tau + 2)\delta_0^{-1} \tag{71}$$

where δ_0 is given by Lemma 3.6.

Lemma 3.7. For $\varepsilon \|b\|_{\tilde{s}}$ small enough, \mathcal{L}_R is invertible and, $\forall s \geq \tilde{s}$,

$$\|\mathcal{L}_R^{-1} h\|_{s,0} \leq 2\rho^{-1} \|h\|_{s,0} + \rho^{-2} \varepsilon C(s) \|b\|_s \|h\|_{\tilde{s},0}, \quad \forall h \in E^{(N)}. \tag{72}$$

Proof. We have $\mathcal{L}_R = D_R^{(N)} + T_R^{(N)}$ as in (68). By the definition of $R, \forall s \geq 0, \|(D_R^{(N)})^{-1}\|_{s,0} \leq \rho^{-1}$, and, by Lemma 4.1,

$$\|T_R^{(N)} h\|_{s,0} \leq \varepsilon C_0(\tilde{s}) \|b\|_{\tilde{s}} \|h\|_{s,0} + \varepsilon C_1(s, \tilde{s}) \|b\|_s \|h\|_{\tilde{s},0}.$$

Hence, by Lemma 2.1, if $\rho^{-1} \varepsilon \|b\|_{\tilde{s}}$ is small enough, then \mathcal{L}_R is invertible and (72) follows with $C(s) := 4C_1(s, \tilde{s})$. \square

The invertibility of $\mathcal{L}^{(N)}$ is then reduced to the invertibility of the self-adjoint operator

$$U := (U_l^2)_{l_1, l_2 \in S} := \mathcal{L}_S - \mathcal{L}_S^R \mathcal{L}_R^{-1} \mathcal{L}_R^S : E_S \rightarrow E_S \tag{73}$$

by the “resolvent” identity

$$(\mathcal{L}^{(N)})^{-1} = \begin{pmatrix} I & -\mathcal{L}_R^{-1} \mathcal{L}_R^S \\ 0 & I \end{pmatrix} \begin{pmatrix} \mathcal{L}_R^{-1} & 0 \\ 0 & U^{-1} \end{pmatrix} \begin{pmatrix} I & 0 \\ -\mathcal{L}_S^R \mathcal{L}_R^{-1} & I \end{pmatrix}.$$

Then (64), and so Proposition 3.1, is a consequence of the following lemma.

Lemma 3.8. If (62)–(63) are satisfied and $\varepsilon(\|b\|_{\tilde{s}} + 1) \leq \eta(\gamma)$ is small enough, then, $\forall s \geq \tilde{s}$,

$$\|U^{-1} h\|_{s,0} \leq \frac{K(s)}{\gamma} N^{\mu_0} (\|h\|_{s,0} + \|b\|_s \|h\|_{0,0}), \quad \forall h \in H_S, \tag{74}$$

with $\mu_0 := 2\tau + 2$.

Proof. To prove (74) we use that, for all $l_1, l_2 \in \mathcal{S}$,

$$(i) \quad \|(U_{l_1}^{l_1})^{-1}\|_{\mathcal{L}(L^2(\mathcal{M}))} \leq C \frac{|l_1|^\tau}{\gamma}, \quad (ii) \quad \|U_{l_1}^{l_2}\|_{\mathcal{L}(L^2(\mathcal{M}))} \leq \frac{C(s)\varepsilon\|b\|_s}{|l_2 - l_1|^{s-1/2}}, \quad l_1 \neq l_2. \quad (75)$$

Estimate (75)(ii) is a consequence of the decay of the Fourier coefficients $\|b_l\|_{H^{s_1}(\mathcal{M})}$, as in Lemma 3.12 of [4]. Moreover it can be proved that, by the separation of the singular sites, assumption (63) can be translated to estimate (75)(i), like in Lemma 3.13 of [4]. To prove (74) we write

$$U = \mathcal{D}(I + \mathcal{D}^{-1}\mathcal{R}), \quad \mathcal{D} := \text{diag}(U_l^l)_{l \in \mathcal{S}}, \quad \mathcal{R} := U - \mathcal{D}. \quad (76)$$

Given $L_1 \in \mathbb{N}_+$, we estimate

$$\begin{aligned} \|(I - \Pi^{(L_1)})\mathcal{D}^{-1}\mathcal{R}h\|_{s,0} &\leq \sum_{l_1 \in \mathcal{S}, |l_1| > L_1} |l_1|^s \left\| (U_{l_1}^{l_1})^{-1} \sum_{l_2 \in \mathcal{S}, l_2 \neq l_1} U_{l_1}^{l_2} h_{l_2} \right\|_{L^2(\mathcal{M})} \\ &\stackrel{(75)(i)}{\leq} C \sum_{l_1 \in \mathcal{S}, |l_1| > L_1} \frac{|l_1|^{s+\tau}}{\gamma} \left(\sum_{l_2 \in \mathcal{S}, l_2 \neq l_1} \|U_{l_1}^{l_2}\|_{\mathcal{L}(L^2(\mathcal{M}))} \|h_{l_2}\|_{L^2(\mathcal{M})} \right) \\ &= (P1) + (P2) \end{aligned} \quad (77)$$

where in (P1), resp. (P2), the sum is restricted to the indices $L_1 \leq |l_1| \leq 2|l_2|$, resp. $|l_1| > 2|l_2|$. By (75)(ii), Lemma 3.6, Schwarz inequality, and since $\delta_0 \in (0, 1)$, we deduce

$$\begin{aligned} (P1) &\leq C(\gamma) \sum_{l_1 \in \mathcal{S}, |l_1| > L_1} \varepsilon |l_1|^{s+\tau} \|b\|_{\tilde{s}} \left(\sum_{|l_2| \geq |l_1|/2} \frac{|l_2|^s \|h_{l_2}\|_{L^2(\mathcal{M})}}{|l_2|^{s+\delta_0(\tilde{s}-1/2)}} \right) \\ &\leq C(\gamma) \sum_{l_1 \in \mathcal{S}, |l_1| > L_1} \varepsilon |l_1|^{s+\tau} \|b\|_{\tilde{s}} \|h\|_{s,0} \left(\sum_{|l_2| \geq |l_1|/2} |l_2|^{-2s-\delta_0(2\tilde{s}-1)} \right)^{1/2} \\ &\leq \varepsilon C(\gamma) C(s) \|b\|_{\tilde{s}} \|h\|_{s,0} \sum_{l_1 \in \mathcal{S}, |l_1| > L_1} |l_1|^{\tau+1-\delta_0\tilde{s}} \leq \varepsilon C(\gamma) C(s) \|b\|_{\tilde{s}} \|h\|_{s,0} L_1^{-\alpha} \end{aligned} \quad (78)$$

where $\alpha := \delta_0\tilde{s} - \tau - 2 > 0$ by the definition of \tilde{s} in (71). By (75)(ii) and, since in (P2) we have $|l_1 - l_2| \geq |l_1| - |l_2| \geq |l_1| - (|l_1|/2) = |l_1|/2$, we deduce that

$$\begin{aligned} (P2) &\leq \frac{C(s)}{\gamma} \sum_{l_1 \in \mathcal{S}} |l_1|^{s+\tau} \frac{\varepsilon \|b\|_s}{|l_1|^{s-1/2}} \left(\sum_{|l_2| < |l_1|/2} \|h_{l_2}\|_{L^2(\mathcal{M})} \right) \\ &\leq \frac{C(s)}{\gamma} \varepsilon \|b\|_s \sum_{l_1 \in \mathcal{S}} |l_1|^{\tau+1} \|h\|_{0,0} \leq \frac{C(s)}{\gamma} \varepsilon \|b\|_s N^{\mu_1} \|h\|_{0,0} \end{aligned} \quad (79)$$

where $\mu_1 := \tau + 2$. Similarly one obtains

$$\|\mathcal{D}^{-1}\mathcal{R}h\|_{0,0} \leq C(\gamma)\varepsilon\|b\|_{\tilde{s}}\|h\|_{0,0} \quad (80)$$

and then

$$\|\Pi^{(L_1)}\mathcal{D}^{-1}\mathcal{R}h\|_{s,0} \stackrel{(S1)}{\leq} L_1^s \|\mathcal{D}^{-1}\mathcal{R}h\|_{0,0} \leq L_1^s C(\gamma)\varepsilon\|b\|_{\tilde{s}}\|h\|_{0,0}. \quad (81)$$

We choose $L_1 := L_1(s)$ large enough so that in estimate (78) it results $C(s)L_1^{-\alpha} \leq 1$. Then we deduce from (77)–(81) that there is $\eta(\gamma) > 0$ such that, for $\varepsilon(\|b\|_{\tilde{s}} + 1) \leq \eta(\gamma)$,

$$\|\mathcal{D}^{-1}\mathcal{R}h\|_{s,0} \leq \frac{1}{2} \|h\|_{s,0} + C'(s)\|b\|_s N^{\mu_1} \|h\|_{0,0}, \quad \|\mathcal{D}^{-1}\mathcal{R}h\|_{0,0} \leq \frac{1}{2} \|h\|_{0,0}.$$

Hence, by Lemma 2.1, for $\varepsilon(\|b\|_{\tilde{s}} + 1) \leq \eta(\gamma)$, $I + \mathcal{D}^{-1}\mathcal{R}$ is invertible in $H^{0,0}$ and

$$\|(I + \mathcal{D}^{-1}\mathcal{R})^{-1}h\|_{s,0} \leq 2\|h\|_{s,0} + 4C'(s)\|b\|_s N^{\mu_1} \|h\|_{0,0}.$$

Finally, (74) follows by (75)(i), with $\mu_0 := \mu_1 + \tau = 2\tau + 2$. \square

4. Appendix

Lemma 4.1. Fix $\tilde{s} > 1/2$, $s_1 > d/2$. For all $s \geq \tilde{s}$, $s'_1 \in [0, s_1]$ there exist constants $C_0(\tilde{s})$, $C_1(\tilde{s}, s) > 0$ such that, $\forall b \in H^s$, $u \in H^{s, s'_1}$, we have

$$\|bu\|_{s, s'_1} \leq C_0(\tilde{s}) \|b\|_{\tilde{s}} \|u\|_{s, s'_1} + C_1(\tilde{s}, s) \|b\|_s \|u\|_{\tilde{s}, s'_1}. \tag{82}$$

Proof. We estimate

$$\|bu\|_{s, s'_1}^2 \stackrel{(60)}{:=} \sum_{m \in \mathbb{Z}} \langle m \rangle^{2s} \left\| \sum_{l \in \mathbb{Z}} b_l u_{m-l} \right\|_{H^{s'_1}(\mathcal{M})}^2 \leq \sum_{m \in \mathbb{Z}} \langle m \rangle^{2s} \left(\sum_{l \in \mathbb{Z}} \|b_l u_{m-l}\|_{H^{s'_1}(\mathcal{M})} \right)^2 \tag{83}$$

$$\leq C(s_1) \sum_{m \in \mathbb{Z}} \langle m \rangle^{2s} \left(\sum_{l \in \mathbb{Z}} \|b_l\|_{H^{s_1}(\mathcal{M})} \|u_{m-l}\|_{H^{s'_1}(\mathcal{M})} \right)^2 \leq 2C(s_1)((P1) + (P2)) \tag{84}$$

where in (P1) the sum is restricted to the indices such that

$$\frac{\langle m \rangle}{\langle m-l \rangle} \leq 1 + \eta(s) \quad \text{with } \eta(s) := 2^{1/s} - 1 > 0, \tag{85}$$

and in (P2) on the complementary set of indices. In passing from (83) to (84) we use that the multiplication operator T_b for $b \in H^{s_1}(\mathcal{M}) \subset L^\infty(\mathcal{M})$, $s_1 > d/2$, satisfies

$$\|T_b\|_{\mathcal{L}(L^2(\mathcal{M}))} \leq \|b\|_{L^\infty(\mathcal{M})} \leq C(s_1) \|b\|_{H^{s_1}(\mathcal{M})}, \quad \|T_b\|_{\mathcal{L}(H^{s_1}(\mathcal{M}))} \leq C(s_1) \|b\|_{H^{s_1}(\mathcal{M})},$$

and so, by interpolation theory (see [23, Chapter 1], and references therein), $\forall 0 \leq s'_1 \leq s_1$, we have

$$\|T_b\|_{\mathcal{L}(H^{s'_1}(\mathcal{M}), H^{s'_1}(\mathcal{M}))} \leq C(s_1) \|b\|_{H^{s_1}(\mathcal{M})}.$$

Using Cauchy–Schwartz inequality (for brevity $\| \cdot \|_{H^{s_1}} := \| \cdot \|_{H^{s_1}(\mathcal{M})}$)

$$\begin{aligned} (P1) &:= \sum_{m \in \mathbb{Z}} \left(\sum_{l \text{ s.t. (85) holds}} \|b_l\|_{H^{s_1}} \langle l \rangle^{\tilde{s}} \|u_{m-l}\|_{H^{s'_1}} \langle m-l \rangle^s \frac{\langle m \rangle^s}{\langle l \rangle^{\tilde{s}} \langle m-l \rangle^s} \right)^2 \\ &\stackrel{(85)}{\leq} \sum_{m \in \mathbb{Z}} \left(\sum_{l \in \mathbb{Z}} \|b_l\|_{H^{s_1}}^2 \langle l \rangle^{2\tilde{s}} \|u_{m-l}\|_{H^{s'_1}}^2 \langle m-l \rangle^{2s} \right) \left(\sum_{l \in \mathbb{Z}} \frac{2}{\langle l \rangle^{2\tilde{s}}} \right) = C(\tilde{s}) \|b\|_{\tilde{s}}^2 \|u\|_{s, s'_1}^2. \end{aligned} \tag{86}$$

Next, in the sum (P2) we have $\langle l \rangle > \langle m \rangle - \frac{\langle m \rangle}{1+\eta(s)} = \langle m \rangle \eta(s) (1 + \eta(s))^{-1}$ and, arguing as in (86),

$$(P2) \leq \|b\|_{\tilde{s}}^2 \|u\|_{s, s'_1}^2 C(s, \tilde{s}). \tag{87}$$

By (84), (86) and (87) we deduce (82). \square

References

- [1] D. Bambusi, J.M. Delort, B. Grebert, J. Szeftel, Almost global existence for Hamiltonian semilinear Klein–Gordon equations with small Cauchy data on Zoll manifolds, *Comm. Pure Appl. Math.* 60 (11) (2007) 1665–1690.
- [2] M. Berti, *Nonlinear Oscillations in Hamiltonian PDEs*, *Progr. Nonlinear Differential Equations Appl.*, vol. 74, Birkhäuser, Boston, 2007.
- [3] M. Berti, P. Bolle, Cantor families of periodic solutions of wave equations with C^k nonlinearities, *NoDEA* 15 (2008) 247–276.
- [4] M. Berti, P. Bolle, Sobolev periodic solutions of nonlinear wave equations in higher spatial dimensions, *Archive for Rational Mechanics and Analysis*, published on line 21-1-2009.
- [5] M. Berti, M. Procesi, *Nonlinear wave and Schrödinger equations on compact Lie groups and homogeneous spaces*, preprint, 2009.
- [6] A. Besse, *Manifolds all of whose Geodesics are Closed*, with appendices by D.B.A. Epstein, J.-P. Bourguignon, L. Bérard-Bergery, M. Berger and J.L. Kazdan, in: *Results in Mathematics and Related Areas*, vol. 93, Springer-Verlag, Berlin/New York, 1978.
- [7] J. Bourgain, Construction of quasi-periodic solutions for Hamiltonian perturbations of linear equations and applications to nonlinear PDE, *Int. Math. Res. Not.* 11 (1994) 475–497.
- [8] J. Bourgain, Construction of periodic solutions of nonlinear wave equations in higher dimension, *Geom. Funct. Anal.* 5 (1995) 629–639.
- [9] J. Bourgain, Quasi-periodic solutions of Hamiltonian perturbations of 2D linear Schrödinger equations, *Ann. of Math.* 148 (1998) 363–439.

- [10] J. Bourgain, Green's Function Estimates for Lattice Schrödinger Operators and Applications, *Ann. of Math. Stud.*, vol. 158, Princeton University Press, Princeton, 2005.
- [11] L. Chierchia, J. You, KAM tori for 1D nonlinear wave equations with periodic boundary conditions, *Comm. Math. Phys.* 211 (2000) 497–525.
- [12] Y. Colin de Verdière, Sur le spectre des opérateurs elliptiques à bicaractéristiques toutes périodiques, *Comment. Math. Helv.* 54 (1979) 508–522.
- [13] W. Craig, Problèmes de petits diviseurs dans les équations aux dérivées partielles, *Panor. Synthèses*, vol. 9, Société Mathématique de France, Paris, 2000.
- [14] W. Craig, C.E. Wayne, Newton's method and periodic solutions of nonlinear wave equation, *Comm. Pure Appl. Math.* 4 (1993) 1409–1498.
- [15] L.H. Eliasson, S. Kuksin, KAM for the nonlinear Schrödinger equation, *Annals of Math.*, in press.
- [16] G. Gentile, V. Mastropietro, Construction of periodic solutions of nonlinear wave equations with Dirichlet boundary conditions by the Lindstedt series method, *J. Math. Pures Appl.* (9) 83 (8) (2004) 1019–1065.
- [17] G. Gentile, V. Mastropietro, M. Procesi, Periodic solutions for completely resonant nonlinear wave equations, *Comm. Math. Phys.* 256 (2) (2005) 437–490.
- [18] G. Gentile, M. Procesi, Periodic solutions for a class of nonlinear partial differential equations in higher dimension, *Comm. Math. Phys.* 289 (3) (2009) 863–906.
- [19] G. Iooss, P. Plotnikov, J. Toland, Standing waves on an infinitely deep perfect fluid under gravity, *Arch. Ration. Mech. Anal.* 177 (3) (2005) 367–478.
- [20] R.S. Hamilton, The inverse function theorem of Nash and Moser, *Bull. Amer. Math. Soc. (N.S.)* 7 (1982) 65–222.
- [21] L. Hörmander, On the Nash Moser implicit function theorem, *Ann. Acad. Sci. Fenn. Ser. A I Math.* 10 (1985) 255–259.
- [22] S. Kuksin, Hamiltonian perturbations of infinite-dimensional linear systems with imaginary spectrum, *Funktsional. Anal. i Prilozhen.* 21 (1987) 22–37.
- [23] S. Kuksin, *Analysis of Hamiltonian PDEs*, Oxford Lecture Ser. Math. Appl., vol. 19, Oxford University Press, 2000.
- [24] J. Moser, A new technique for the construction of solutions of nonlinear differential equations, *Proc. Natl. Acad. Sci.* 47 (1961) 1824–1831.
- [25] J. Moser, A rapidly convergent iteration method and non-linear partial differential equations I & II, *Ann. Sc. Norm. Super. Pisa* (3) 20 (1966) 265–315, 499–535.
- [26] J. Pöschel, Integrability of Hamiltonian systems on Cantor sets, *Comm. Pure Appl. Math.* 35 (1982) 653–695.
- [27] J. Pöschel, A KAM-theorem for some nonlinear PDEs, *Ann. Sc. Norm. Super. Pisa Cl. Sci.* 23 (1996) 119–148.
- [28] D. Salamon, E. Zehnder, KAM theory in configuration space, *Comment. Math. Helv.* 64 (1989) 84–132.
- [29] E. Wayne, Periodic and quasi-periodic solutions of nonlinear wave equations via KAM theory, *Comm. Math. Phys.* 127 (1990) 479–528.
- [30] E. Zehnder, Generalized implicit function theorems with applications to some small divisors problems I–II, *Comm. Pure Appl. Math.* 28 (1975) 91–140, *Comm. Pure Appl. Math.* 29 (1976) 49–113.