

Asymptotic analysis for surfaces with large constant mean curvature and free boundaries

Paul Laurain

UMPA, 46 allée d'Italie, Lyon, France

Received 9 February 2011; received in revised form 27 July 2011; accepted 16 September 2011

Available online 29 September 2011

Abstract

We prove that simply connected H -surfaces with bounded area and free boundary in a domain necessarily concentrate at a critical point of the mean curvature of the boundary of this domain.

© 2011 L'Association Publications de l'Institut Henri Poincaré. Published by Elsevier B.V. All rights reserved.

Introduction

The aim of this article is to understand the asymptotic behaviour of sequences of surfaces with large constant mean curvature and free boundaries. These surfaces arise naturally in the partitioning problem which consists in dividing a domain into two parts of prescribed volumes by a surface of minimal area. The existence of solutions of this problem is given by the geometric measure theory (see for instance Morgan [18]). However we get no information about the topology of such surfaces, except in the case of strictly convex domains, where we know that such a surface is connected and we get some bounds on the number of components of its boundary as well as its genus, see Ros and Vergasta [20]. Moreover it is conjectured that, in this case, the surface is homeomorphic to a disk, see Ritoré and Ros [19].

In the following, we let Ω be a smooth domain of \mathbb{R}^3 and we will consider H -surface as a map $u \in C^2(\overline{\mathbb{D}}, \mathbb{R}^3)$ where

$$\mathbb{D} = \{z \in \mathbb{R}^2 \text{ s.t. } |z| < 1\}$$

which is an immersion and which satisfies

$$\begin{cases} \Delta u = -2Hu_x \wedge u_y, \\ \langle u_x, u_y \rangle = |u_x|^2 - |u_y|^2 = 0, \\ u(z) \in \partial\Omega \quad \text{for all } z \in \partial\mathbb{D}, \\ \partial_\nu u(z) \perp T_{u(z)}\partial\Omega \quad \text{for all } z \in \partial\mathbb{D}, \end{cases} \quad (1)$$

where $\Delta = -\frac{\partial^2}{\partial x^2} - \frac{\partial^2}{\partial y^2}$.

E-mail address: paul.laurain@umpa.ens-lyon.fr.

Then $u(\overline{D})$ is a regular surface of constant mean curvature H with boundary contained in $\partial\Omega$ and which meets $\partial\Omega$ orthogonally.

The first result of existence of solutions of (1) is due to Struwe [21], which finds solutions in domains diffeomorphic to a ball using a parabolic version of our equation. Another idea to find solutions of (1) in a general domain is to look for solutions with large mean curvature (i.e. with a small diameter). In fact in this case the topology of the domain play no role and the geometry is under control.

This intuition was confirmed by Fall [9]. However, the existence of such solutions is subject to a local condition on the curvature of the boundary of Ω . Fall proved in [9] the following: given any smooth domain $\Omega \subset \mathbb{R}^3$ and $p \in \partial\Omega$ be a non-degenerate critical point of the mean curvature of $\partial\Omega$. There exists a family of solutions $u^\varepsilon \in C^2(\overline{D}, \mathbb{R}^3)$ of (1) for $H = \frac{1}{\varepsilon}$ such that $u^\varepsilon(\overline{D})$ is embedded and $\|u^\varepsilon - p\|_\infty \rightarrow 0$ when $\varepsilon \rightarrow 0$. Moreover $\frac{1}{\varepsilon}u^\varepsilon$, correctly translated, converges to a hemisphere of radius 1.

This result is similar to the result of Ye [25] concerning the existence of closed surfaces with constant mean curvature in a curved manifold. Indeed they are similar in their statement but also in the method of proof which takes a solution of the limit equation, here one hemisphere, and tries to perturb it via the implicit functions theorem. This remark done, the question of the necessity of the condition that p is a critical point of the mean curvature of $\partial\Omega$ comes naturally. A first answer in this direction is provided by Fall [10]. Indeed he shows that the solutions to the problem of partitioning, which are equivalent to the isoperimetric problem solutions in this context, converge to a point of maximal mean curvature when their volume tends to zero.

This theorem is similar to the result of Druet [7] concerning the location of small isoperimetric domains in a curved manifold. Indeed Druet proved that these domains are near global maxima of the scalar curvature. In [17], we proved under suitable assumptions that surfaces of large constant mean curvature and small diameter in a 3-dimensional manifold are necessarily located near a critical point of the scalar curvature. Here we show under reasonable assumptions that surfaces of large constant mean curvature with boundary included in $\partial\Omega$ and meeting $\partial\Omega$ orthogonally are necessarily located near a critical point of the mean curvature of $\partial\Omega$. First, we assume that the diameter is controlled in order to avoid solutions that collapse along some geodesics (such examples were constructed by Mahmoudi and Fall [11]). Second, we assume that the area is controlled to avoid an infinity of bubbles. Then we prove the following theorem.

Theorem 0.1. *Let Ω be a smooth domain of \mathbb{R}^3 and a sequence of embedded surfaces Σ^ε in Ω satisfying the following assumptions:*

- (i) $\partial\Sigma^\varepsilon \subset \partial\Omega$ and Σ^ε and $\partial\Omega$ meet orthogonally,
- (ii) Σ^ε has constant mean curvature equal to $\frac{1}{\varepsilon}$,
- (iii) the diameter and the area of Σ^ε are respectively an $O(\varepsilon)$ and an $O(\varepsilon^2)$.

Then, up to a subsequence, Σ^ε converges to $p \in \partial\Omega$ which is a critical point of the mean curvature of $\partial\Omega$.

This theorem can be explained in the following way: given $\Omega \subset \mathbb{R}^3$ a smooth domain, for any $\delta > 0$, any $C > 0$, there exists $\varepsilon_0 > 0$ such that any embedded surface orthogonal to the boundary Σ of constant mean curvature $\frac{1}{\varepsilon}$ with $\varepsilon < \varepsilon_0$, $\text{diameter}(\Sigma) \leq C\varepsilon$, $\text{Area}(\Sigma) \leq C\varepsilon^2$, satisfies that $\Sigma \subset B(p, \delta)$ for some critical point $p \in \partial\Omega$ of the mean curvature of $\partial\Omega$.

Note that the bound on the diameter and the area are scale invariant with respect to the mean curvature.

This article is organized as follows. In the first section we remind some useful results about regularity of constant mean curvature surfaces with free boundaries. In the second section we remind the classification of the solution of the constant mean curvature equation on the whole plane and we extend it to domain like disk or half-plane. Finally in a third section we give a proof of the theorem, dividing it in three parts; first we perform a blow-up analysis decomposing our sequence in a sum of spheres and hemispheres; then we insure the existence of at least one hemisphere in the decomposition using notably the Aleksandrov reflexion principle, finally we achieve the proof applying the balancing formula. The main difficulty is to understand precisely the asymptotic behaviour of our sequence of surfaces Σ^ε on the boundary of Ω . Some technical lemmas are postponed to Appendix A.

1. Regularity and *a priori* estimates on constant mean curvature surfaces with free boundaries

In this section we give a general result on the regularity of constant mean curvature surfaces with free boundaries, the reader will find all the details in Chapter 7 of [6].

Theorem 1.1. *Let Ω be a $C^{m,\alpha}$ domain of \mathbb{R}^3 with $m \geq 3$ and $\alpha > 0$, then every solution of (1) is $C^{m,\alpha}$.*

The solutions inherit of the regularity of the Ω provided it is sufficiently smooth.

The proof of this result is divided into three steps. The first shows, using the isoperimetric inequality for surfaces, that the solutions are $C^{0,\eta}$ up to the boundary. Then, using *a priori* estimates in the spaces $H^{k,p}$, we deduce the $C^{1,\frac{1}{2}}$ regularity up to the boundary. Finally, using a classical argument of bootstrap, we obtain that the solutions are smooth inside and inherit of the regularity of the domain up to the boundary as soon as it is at least $C^{3,\alpha}$.

We give here the *a priori* estimate which is the keystone of the second step and that will be used later.

Theorem 1.2. *Let Ω be a smooth domain, whose metric of the boundary will be denoted by g , and u be a solution of (1). We assume that u belongs to $C^{0,\eta}(\overline{\mathbb{D}})$. Then, for every open set U of $\overline{\mathbb{D}}$ and every $2 < p < +\infty$, there exists a constant c depending only on $\|g\|_3$, U , p , $\int_U |\nabla u|^2 dz$ and the modulus of continuity of u such that*

$$\int_U |\nabla u|^p dz < c.$$

This estimate and the standard elliptic theory lead to uniform bounds of the type

$$\|u\|_{2+\eta,U} < c,$$

where c depends only on $\|g\|_3$, U , p , $\int_U |\nabla u|^2 dz$ and the modulus of continuity of u .

Remark 1.1. In particular, we note that from any sequence of solutions whose gradient is uniformly bounded on an open set U of $\overline{\mathbb{D}}$, we can extract a subsequence which converges uniformly in $C^2(U)$.

2. Classification of solution of the limit equation

We start by remind a crucial result of Brezis and Coron [2] which states that the only solutions of

$$\Delta u = -2u_x \wedge u_y \quad \text{on } \mathbb{R}^2$$

with bounded energy are exactly, up to a conformal reparametrization, the inverse of the stereographic projection. This result can be seen as a variant of the Hopf’s theorem where the hypothesis of conformality is replaced by a bound on the area.

Lemma 2.1 (Lemma A.1 of [2]). *Let $\omega \in L^1_{loc}(\mathbb{R}^2, \mathbb{R}^3)$ which satisfies*

$$\begin{aligned} \Delta \omega &= -2\omega_x \wedge \omega_y, \\ \int_{\mathbb{R}^2} |\nabla \omega|^2 dz &< +\infty. \end{aligned} \tag{2}$$

Then ω has precisely the form

$$\omega(z) = \pi_N^{-1} \left(\frac{P(z)}{Q(z)} \right) + C,$$

where $N \in S^2$, P and Q are polynomial, C is a constant and π_N is the stereographic projection from the north pole N . In addition

$$\int_{\mathbb{R}^2} |\nabla \omega|^2 = 8\pi k \quad \text{with } k = \max\{\deg P, \deg Q\},$$

provided that $\frac{P}{Q}$ is irreducible.

It could be useful to remark that the gradient of such an ω satisfies the following formula

$$|\nabla \omega| = \frac{2\sqrt{2}|P'Q - Q'P|}{|P|^2 + |Q|^2}.$$

Then we define a special class of solutions which will be important in what follows: the spheres which are parametrized only once.

Definition 2.1. A solution ω of (2) is said to be simple if

$$\omega(z) = \pi_N^{-1}\left(\frac{P(z)}{Q(z)}\right) + C,$$

with $\frac{P}{Q}$ is irreducible and $\max\{\deg P, \deg Q\} = 1$.

In particular, if ω is a simple solution of (2), then we have

$$|\nabla \omega^\varepsilon(x)| = O\left(\frac{\lambda^\varepsilon}{|x - a^\varepsilon|^2 + (\lambda^\varepsilon)^2}\right), \tag{3}$$

where $\omega^\varepsilon = \omega\left(\frac{\cdot - a^\varepsilon}{\lambda^\varepsilon}\right)$, a^ε and λ^ε are respectively a sequence of points in \mathbb{R}^2 and a sequence of positive numbers.

Finally we give a generalization of the result of Brezis and Coron for solutions defined on the disk or the half-plane with appropriate boundary conditions.

Lemma 2.2. Let $\Omega = \mathbb{H}$ or \mathbb{D} and $u : \Omega \rightarrow \mathbb{R}^2 \times \mathbb{R}_+$ be such that

$$\begin{aligned} \Delta u &= -2u_x \wedge u_y, \\ \langle u_x, u_y \rangle &= |u_x| - |u_y| = 0, \\ \|\nabla u\|_2 &< +\infty, \\ u|_{\partial\Omega} &\subset \mathbb{R}^2 \times \{0\} \end{aligned}$$

and such that the angle between $u(\Omega)$ and $\mathbb{R}^2 \times \{0\}$ when it is defined is right. Then

$$u = C + \pi^{-1}\left(\frac{P}{Q}\right)$$

where π is the stereographic projection and P and Q are two polynoms of $\mathbb{C}[z]$. Moreover, $u|_{\partial\Omega}$ describes a circle of radius one.

Proof of Lemma 2.2. First of all we can assume that $\Omega = \mathbb{D}$. Indeed let $\phi : \mathbb{D} \setminus \{1\} \rightarrow \overline{\mathbb{H}}$ defined by

$$\phi(z) = -i \frac{z + 1}{z - 1}.$$

It is well known that this application is a conformal isomorphism. Hence if $u : \mathbb{H} \rightarrow \mathbb{R}^2 \times \mathbb{R}_+$ satisfies the hypothesis of the lemma, it is the same for $\tilde{u} : \mathbb{D} \setminus \{1\} \rightarrow \mathbb{R}^2 \times \mathbb{R}_+$ defined by $\tilde{u} = u \circ \phi$. But, since $\|\nabla \tilde{u}\|_2 = \|\nabla u\|_2 < +\infty$, thanks to the regularity theory \tilde{u} can be extended smoothly at 1.

Then we can extend u to the whole plane, setting

$$u(z) = -\begin{pmatrix} u^1 \\ u^2 \\ -u^3 \end{pmatrix} \begin{pmatrix} 1 \\ z \end{pmatrix} \quad \text{for all } z \in \mathbb{R}^2 \setminus \mathbb{D}.$$




Fig. 1. A surface with boundary.

This extension is C^1 and moreover satisfies the hypothesis of Lemma 2.1, since the energy is simply doubled by this extension. This proves the first part of the theorem. Finally we easily remark that $u|_{\partial\Omega}$ describes a circle of 1, remarking that u is equal, up to a sign change, to its Gauss map. But our hypothesis forces to the Gauss map to be contained in a great circle on the boundary, which achieves the proof of the theorem. \square

3. Proof of theorem

The main idea is to apply the balancing formula to the boundary of our sequences of surfaces in order to detect the geometry of $\partial\Omega$. We remind that the balancing formula is an identity discovered by Kusner [16] which concerns the shape of the boundary of a surface with constant mean curvature. Let S be a surface with constant mean curvature. Then

$$\int_{\partial S} \vec{\eta} ds = 2H_0 \int_{\Sigma} \vec{v} d\sigma, \tag{4}$$

where Σ is a smooth surface having the same boundary as S , \vec{v} is the normal of Σ and $\vec{\eta}$ is the conormal of ∂S , see Fig. 1. The reader will find a proof of (4) at Chapter 7 of [15].

In order to exploit this formula, we need a precise description of the behaviour of our surfaces. In order to obtain it, we start by proving that the sequence decomposes asymptotically as a sum of spheres and hemispheres. Then we shall prove that this decomposition contains at least one hemisphere, that is to say that the boundaries of our sequence of surfaces do not collapse to a point.

Now we consider a smooth domain Ω of \mathbb{R}^3 and a sequence of embedded disks Σ^ε in Ω which satisfy the following assumptions

- (i) $\partial \Sigma^\varepsilon \subset \partial \Omega$ and $\partial \Sigma^\varepsilon$ and $\partial \Omega$ meet orthogonally,
- (ii) Σ^ε has constant mean curvature equal to $\frac{1}{\varepsilon}$,
- (iii) the diameter and the area of Σ^ε are respectively an $O(\varepsilon)$ and an $O(\varepsilon^2)$.

Up to translate Ω and to extract a subsequence of Σ^ε , we can assume that Σ^ε goes to 0 and that $0 \in \partial \Sigma^\varepsilon$. Then we rescale the space by a factor $\frac{1}{\varepsilon}$ and we choose a conformal parametrization for our sequence of surfaces, that is to say a sequence of $u^\varepsilon : \mathbb{D} \rightarrow \mathbb{R}^3$ such that

$$\begin{cases} \Delta u^\varepsilon = -2u_x^\varepsilon \wedge u_y^\varepsilon, \\ \langle u_x^\varepsilon, u_y^\varepsilon \rangle = \|u_x^\varepsilon\|^2 - \|u_y^\varepsilon\|^2 = 0, \end{cases} \quad \text{on } \mathbb{D},$$

$$\|u^\varepsilon\|_\infty = O(1) \quad \text{and} \quad \|\nabla u^\varepsilon\|_2 = O(1),$$

$$u^\varepsilon(\partial\mathbb{D}) \subset \partial\Omega^\varepsilon \quad \text{and} \quad \langle u_x^\varepsilon \wedge u_y^\varepsilon, N^\varepsilon \rangle = 0 \quad \text{on } \partial\mathbb{D}, \tag{5}$$

where $\Omega^\varepsilon = \frac{1}{\varepsilon}\Omega$ and N^ε is the exterior normal of $\partial\Omega^\varepsilon$. The regularity of such a sequence of functions depends on the regularity of the surface where its free boundary lives. Here, since $\partial\Omega$ is smooth, our sequence is smooth up to the boundary.

3.1. Decomposition of u^ε as a sum of spheres and hemispheres

We start performing a decomposition of our surfaces as a sum of spheres and hemispheres in the spirit of what has been done by Brezis and Coron in [2]. However there are two big changes. On the one hand, there are two limit solutions (sphere and hemisphere). On the other hand, we must obtain an L^∞ -estimate rather than estimates on the gradient, while the equation lends itself much better to obtain estimates on the gradient.

Theorem 3.1. *Let u^ε be a sequence of maps in $C^2(\overline{\mathbb{D}})$ which are non-constant solutions of (5), then either u^ε converges uniformly to 0 or there exist $p \in \mathbb{N}$ and*

- (i) $\omega^1, \dots, \omega^p$ non-constant solutions of (2),
- (ii) $a_1^\varepsilon, \dots, a_p^\varepsilon$ sequences of $\overline{\mathbb{D}}$, and
- (iii) $\lambda_1^\varepsilon, \dots, \lambda_p^\varepsilon$ sequences of positive real numbers such that $\lim_{\varepsilon \rightarrow 0} \lambda_i^\varepsilon < +\infty$,

such that, for a subsequence u^ε (still denote u^ε), we get

$$u_i^\varepsilon \rightarrow \omega^i \quad \text{in } C_{loc}^2(\Omega_i \setminus S_i) \text{ as } \varepsilon \rightarrow 0 \text{ for all } 1 \leq i \leq p, \tag{A}$$

where $u_i^\varepsilon = u^\varepsilon(\lambda_i^\varepsilon \cdot + a_i^\varepsilon)$, $\Omega_i = \lim_{\varepsilon \rightarrow 0} \{z \in \mathbb{R}^2 \text{ s.t. } \lambda_i^\varepsilon \cdot + a_i^\varepsilon \in \mathbb{D}\}$ and $S_i = \lim_{\varepsilon \rightarrow 0} \{\frac{a_j^\varepsilon - a_i^\varepsilon}{\lambda_i^\varepsilon} \text{ s.t. } j \in \{1, \dots, p\} \setminus \{i\}\}$.

$$\lim_{\varepsilon \rightarrow 0} \frac{d_i^\varepsilon(a_j^\varepsilon)}{\lambda_j^\varepsilon} + \frac{d_j^\varepsilon(a_i^\varepsilon)}{\lambda_i^\varepsilon} = +\infty \quad \text{for all } i \neq j, \tag{B}$$

where $d_i^\varepsilon(x) = \sqrt{(\lambda_i^\varepsilon)^2 + |a_i^\varepsilon - x|^2}$,

$$d\left(u^\varepsilon(\mathbb{D}), \bigcup_{i=1}^p B_i\right) \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0, \tag{C}$$

where B_i is the limit set of $\omega_i^\varepsilon(\mathbb{D})$ as ε goes to zero, with $\omega_i^\varepsilon = \omega_i(\frac{\cdot - a_i^\varepsilon}{\lambda_i^\varepsilon})$, that is to say some spheres and hemispheres.

Proof. We are going to extract the bubble by induction, the process will stop thanks to our uniform estimate on the energy of u^ε . In fact, such an extraction will be done until a “weak estimate” is not satisfied on the reminder, which is by now an almost classical technics since the work of Druet, Hebey and Robert about strong estimate for sequences of solution of Yamabe-type equation, see [8]. The advantage of this method is to insure a C_{loc}^2 -converge rather than an H^1 -converge.

Let $k \geq 1$, we say that u^ε satisfies the property (P_k) if there exist

- (i) $\omega^1, \dots, \omega^k$ non-constant solution of (2),
- (ii) $a_1^\varepsilon, \dots, a_k^\varepsilon$ sequences of $\overline{\mathbb{D}}$ and
- (iii) $\lambda_1^\varepsilon, \dots, \lambda_k^\varepsilon$ sequences of positive real numbers such that $\lim_{\varepsilon \rightarrow 0} \lambda_i^\varepsilon < +\infty$,

such that, for a subsequence u^ε (still denoted u^ε), we get

$$u_i^\varepsilon \rightarrow \omega_i \quad \text{in } C_{loc}^2(\Omega_i \setminus S_i) \text{ as } \varepsilon \rightarrow 0 \text{ for all } 1 \leq i \leq k, \tag{A_k}$$

where $u_i^\varepsilon = u^\varepsilon(\lambda_i^\varepsilon \cdot + a_i^\varepsilon)$, $\Omega_i = \lim_{\varepsilon \rightarrow 0} \{z \in \mathbb{R}^2 \text{ s.t. } \lambda_i^\varepsilon \cdot + a_i^\varepsilon \in \mathbb{D}\}$ and $S_i = \lim_{\varepsilon \rightarrow 0} \{\frac{a_j^\varepsilon - a_i^\varepsilon}{\lambda_i^\varepsilon} \text{ s.t. } j \in \{1, \dots, p\} \setminus \{i\}\}$.

$$\frac{d_i^\varepsilon(a_j^\varepsilon)}{\lambda_j^\varepsilon} + \frac{d_j^\varepsilon(a_i^\varepsilon)}{\lambda_i^\varepsilon} \rightarrow +\infty \quad \text{as } \varepsilon \rightarrow 0 \text{ for all } i \neq j, \tag{B_k}$$

where $d_i^\varepsilon(x) = \sqrt{(\lambda_i^\varepsilon)^2 + |a_i^\varepsilon - x|^2}$. Moreover, when $\Omega_i \neq \mathbb{R}^2$, $\omega_i^\varepsilon|_{\partial\Omega_i}$ describes (perhaps several times) a circle of radius 1.

Claim 1. If (P_k) holds for some $k \geq 1$, then either (P_{k+1}) holds or

$$\limsup_{\varepsilon \rightarrow 0} \sup_{z \in \mathbb{D}} \left(\min_{1 \leq i \leq k} d_i^\varepsilon(z) \right) \left| \nabla \left(u^\varepsilon - \sum_{i=1}^k \omega_i^\varepsilon \right) (z) \right| = 0, \tag{6}$$

where $\omega_i^\varepsilon = \omega_i \left(\frac{\cdot - a_i^\varepsilon}{\lambda_i^\varepsilon} \right)$.

Proof. Assume that (P_k) holds and that there exist $\gamma_0 > 0$ and a subsequence u^ε (still denoted u^ε) such that

$$\sup_{z \in \mathbb{D}} \left(\min_{1 \leq i \leq k} d_i^\varepsilon(z) \right) \left| \nabla \left(u^\varepsilon - \sum_{i=1}^k \omega_i^\varepsilon \right) (z) \right| \geq \gamma_0. \tag{7}$$

Let $a_{k+1}^\varepsilon \in \overline{\mathbb{D}}$ be such that

$$\left(\min_{1 \leq i \leq k} d_i^\varepsilon(a_{k+1}^\varepsilon) \right) \left| \nabla \left(u^\varepsilon - \sum_{i=1}^k \omega_i^\varepsilon \right) (a_{k+1}^\varepsilon) \right| = \sup_{z \in \mathbb{D}} \left(\min_{1 \leq i \leq k} d_i^\varepsilon(z) \right) \left| \nabla \left(u^\varepsilon - \sum_{i=1}^k \omega_i^\varepsilon \right) (z) \right|.$$

We define $\lambda_{k+1}^\varepsilon$ by

$$\left| \nabla \left(u^\varepsilon - \sum_{i=1}^k \omega_i^\varepsilon \right) (a_{k+1}^\varepsilon) \right| = \frac{1}{\lambda_{k+1}^\varepsilon}.$$

Remarking that $\min_{1 \leq i \leq k} d_i^\varepsilon(a_{k+1}^\varepsilon)$ is bounded, it is clear that

$$\lim_{\varepsilon \rightarrow 0} \lambda_{k+1}^\varepsilon < +\infty. \tag{8}$$

There are now two cases to consider.

First case:

$$\lim_{\varepsilon \rightarrow 0} \frac{\min_{1 \leq i \leq k} d_i^\varepsilon(a_{k+1}^\varepsilon)}{\lambda_{k+1}^\varepsilon} = +\infty. \tag{9}$$

In that case, (B_{k+1}) is automatically satisfied. We set

$$u_{k+1}^\varepsilon(z) = u^\varepsilon(\lambda_{k+1}^\varepsilon z + a_{k+1}^\varepsilon) \quad \text{for all } z \in \overline{\Omega_{k+1}^\varepsilon}$$

where $\Omega_{k+1}^\varepsilon = \{z \in \mathbb{R}^2 \text{ s.t. } \lambda_{k+1}^\varepsilon z + a_{k+1}^\varepsilon \in \mathbb{D}\}$. Let $z \in \Omega_{k+1}^\varepsilon$, we get

$$\begin{aligned} \left| \nabla u_{k+1}^\varepsilon(z) \right| &= \lambda_{k+1}^\varepsilon \left| \nabla u^\varepsilon(\lambda_{k+1}^\varepsilon z + a_{k+1}^\varepsilon) \right| \\ &\leq \lambda_{k+1}^\varepsilon \left| \nabla \left(u^\varepsilon - \sum_{i=1}^k \omega_i^\varepsilon \right) (\lambda_{k+1}^\varepsilon z + a_{k+1}^\varepsilon) \right| + \lambda_{k+1}^\varepsilon \left| \nabla \left(\sum_{i=1}^k \omega_i^\varepsilon \right) (\lambda_{k+1}^\varepsilon z + a_{k+1}^\varepsilon) \right|. \end{aligned} \tag{10}$$

Thanks to (A_k) and (9), we easily get that

$$\lambda_{k+1}^\varepsilon \left| \nabla \left(\sum_{i=1}^k \omega_i^\varepsilon \right) (\lambda_{k+1}^\varepsilon z + a_{k+1}^\varepsilon) \right| = o(1). \tag{11}$$

Then, using the definition of a_{k+1}^ε , (9), (10) and (11), we get

$$\left| \nabla u_{k+1}^\varepsilon(z) \right| \leq \frac{\min_{1 \leq i \leq k} d_i^\varepsilon(a_{k+1}^\varepsilon)}{\min_{1 \leq i \leq k} d_i^\varepsilon(\lambda_{k+1}^\varepsilon z + a_{k+1}^\varepsilon)} + o(1) = 1 + o(1). \tag{12}$$

Then $|\nabla u_{k+1}^\varepsilon|$ is bounded on every compact subset of $\overline{\Omega_{k+1}^\varepsilon}$. Moreover, thanks to conformal invariance of our equation, u_{k+1}^ε still satisfies (5). Hence, using standard elliptic theory, see Section 1 and [13], we see that there exist a subsequence of u^ε (still denoted u^ε) and $\omega^{k+1} \in C^2(\overline{\Omega_{k+1}})$ such that

$$u_{k+1}^\varepsilon \rightarrow \omega^{k+1} \quad \text{in } C_{loc}^2(\overline{\Omega_{k+1}})$$

and

$$\Delta \omega^{k+1} = -2\omega_x^{k+1} \wedge \omega_y^{k+1} \quad \text{on } \Omega_{k+1},$$

where $\Omega_{k+1} = \lim_{\varepsilon \rightarrow 0} \Omega_{k+1}^\varepsilon$. Here there are again two cases: either Ω_{k+1} is the whole plane or this is a disk or an half-plane (which is conformally equivalent). In the case of the disk or the half-plane, the boundary condition passes to the limit, that is to say, up to rotation,

$$\omega^{k+1}(\partial \Omega_{k+1}) \subset \mathbb{R}^2 \times \{0\}$$

and

$$\langle \omega_x^{k+1} \wedge \omega_y^{k+1}, N \rangle = 0 \quad \text{on } \partial \Omega_{k+1},$$

where $N = (0, 0, 1)$. Moreover, thanks to conformal invariance of $\|\nabla \cdot\|_2$, up to extract a subsequence, we get

$$u_{k+1}^\varepsilon \rightharpoonup \omega^{k+1} \quad \text{in } L^2(\mathbb{R}^2)$$

and

$$\|\nabla \omega^{k+1}\|_2 \leq \liminf_{\varepsilon \rightarrow 0} \|\nabla u_{k+1}^\varepsilon\|_2 = \liminf_{\varepsilon \rightarrow 0} \|\nabla u^\varepsilon\|_2 < +\infty.$$

Finally, thanks to Lemmas 2.1 and 2.2, ω^{k+1} has the desired shape, moreover ω^{k+1} is non-constant since $|\nabla \omega^{k+1}(0)| = 1$. This achieves the proof of (P_{k+1}) in the first case.

Second case:

$$\lim_{\varepsilon \rightarrow 0} \frac{\min_{1 \leq i \leq k} d_i^\varepsilon(a_{k+1}^\varepsilon)}{\lambda_{k+1}^\varepsilon} = \gamma > 0. \tag{13}$$

First of all, we need to show that (B_{k+1}) holds. We assume by contradiction that (B_{k+1}) does not hold, then, up to extract a subsequence, there exists $1 \leq i_0 \leq k$ such that

$$d_{k+1}^\varepsilon(a_{i_0}^\varepsilon) = O(\lambda_{i_0}^\varepsilon) \quad \text{and} \quad d_{i_0}^\varepsilon(a_{k+1}^\varepsilon) = O(\lambda_{k+1}^\varepsilon). \tag{14}$$

On the one hand, (14) gives

$$\lim_{\varepsilon \rightarrow 0} \frac{\lambda_{k+1}^\varepsilon}{\lambda_{i_0}^\varepsilon} = c \quad \text{and} \quad |a_{i_0}^\varepsilon - a_{k+1}^\varepsilon| = O(\lambda_{i_0}^\varepsilon), \tag{15}$$

where c is a positive constant. On the other hand, thanks to (A_k) and (B_k) , we get

$$\nabla \left(\left(u^\varepsilon - \sum_{i=1}^k \omega_i^\varepsilon \right) (\lambda_{i_0}^\varepsilon + a_{i_0}^\varepsilon) \right) \rightarrow 0 \quad \text{in } C_{loc}^2(\Omega_{i_0} \setminus S_{i_0}). \tag{16}$$

Hence, thanks to (13) and (15), we necessarily get

$$d \left(\frac{a_{k+1}^\varepsilon - a_{i_0}^\varepsilon}{\lambda_{i_0}^\varepsilon}, S_{i_0} \right) = o(1).$$

Let $j \in \{1, \dots, k\} \setminus \{i_0\}$ be such that

$$\left| \frac{a_{k+1}^\varepsilon - a_j^\varepsilon}{\lambda_{i_0}^\varepsilon} \right| = o(1).$$

Using (13) and (15), we remark that for ε small enough,

$$\frac{\lambda_j^\varepsilon}{\lambda_{k+1}^\varepsilon} \geq \frac{\gamma}{2},$$

and, using again (15), we remark that, for ε small enough,

$$\frac{\lambda_j^\varepsilon}{\lambda_{i_0}^\varepsilon} \geq \frac{\gamma}{4c}.$$

Since $\frac{a_{i_0}^\varepsilon - a_j^\varepsilon}{\lambda_{i_0}^\varepsilon} = O(1)$ and i_0 and j satisfy (B_k) , we necessarily get

$$\lambda_{i_0}^\varepsilon = o(\lambda_j^\varepsilon).$$

Hence, for all j such that $\frac{a_{k+1}^\varepsilon - a_j^\varepsilon}{\lambda_{i_0}^\varepsilon} = o(1)$, we get

$$\lambda_{i_0}^\varepsilon = o(\lambda_j^\varepsilon).$$

In particular, thanks to (A_k) , there exists $\delta > 0$ such that for all $z \in B(0, \delta)$, we have

$$\lambda_{i_0}^\varepsilon |\nabla \omega_i^\varepsilon(a_{k+1}^\varepsilon + z\lambda_{i_0}^\varepsilon)| = o(1) \quad \text{for all } i \neq i_0.$$

We easily see that

$$\lambda_{i_0}^\varepsilon |\nabla u^\varepsilon| = O(1) \quad \text{on } B(a_{k+1}^\varepsilon, \delta\lambda_{i_0}^\varepsilon).$$

Using standard elliptic theory, up to extract a subsequence, we get that $u_{i_0}^\varepsilon$ converges to ω^{i_0} in $C_{loc}^2(B(c_{k+1}, \frac{\delta}{2}))$ where $c_{k+1} = \lim_{\varepsilon \rightarrow 0} \frac{a_{k+1}^\varepsilon - a_{i_0}^\varepsilon}{\lambda_{i_0}^\varepsilon}$. Then we deduce that

$$|\nabla(u_{i_0}^\varepsilon - \omega^{i_0})(a_{k+1}^\varepsilon)| \rightarrow 0,$$

which leads to

$$\lambda_{i_0}^\varepsilon \left| \nabla \left(\left(u^\varepsilon - \sum_{i=1}^k \omega_i^\varepsilon \right) (a_{k+1}^\varepsilon) \right) \right| \rightarrow 0,$$

which, thanks to (16), is a contradiction with (13) and proves (B_{k+1}) .

Now, we set

$$u_{k+1}^\varepsilon = u^\varepsilon(\lambda_{k+1}^\varepsilon \cdot + a_{k+1}^\varepsilon) \quad \text{for all } z \in \Omega_{k+1}^\varepsilon,$$

where $\Omega_{k+1}^\varepsilon = \{z \in \mathbb{R}^2 \text{ s.t. } \lambda_{k+1}^\varepsilon z + a_{k+1}^\varepsilon \in \mathbb{D}\}$. Let $z \in \overline{\Omega_{k+1}^\varepsilon} \setminus \{S_{k+1}\}$, we have

$$\begin{aligned} |\nabla u_{k+1}^\varepsilon(z)| &= \lambda_{k+1}^\varepsilon |\nabla u^\varepsilon(\lambda_{k+1}^\varepsilon z + a_{k+1}^\varepsilon)| \\ &\leq \lambda_{k+1}^\varepsilon \left| \nabla \left(u^\varepsilon - \sum_{i=1}^k \omega_i^\varepsilon \right) (\lambda_{k+1}^\varepsilon z + a_{k+1}^\varepsilon) \right| + \lambda_{k+1}^\varepsilon \left| \nabla \left(\sum_{i=1}^k \omega_i^\varepsilon \right) (\lambda_{k+1}^\varepsilon z + a_{k+1}^\varepsilon) \right|. \end{aligned} \tag{17}$$

Thanks to (A_k) and (13), we obtain

$$\lambda_{k+1}^\varepsilon \left| \nabla \left(\sum_{i=1}^k \omega_i^\varepsilon \right) (\lambda_{k+1}^\varepsilon \cdot + a_{k+1}^\varepsilon) \right| = O\left(\frac{1}{d(z, S_{k+1})} + |z| + 1\right). \tag{18}$$

With the consention that $d(z, \emptyset) = +\infty$.

Then using the definition of a_{k+1}^ε , (17) and (18), we get

$$\begin{aligned} |\nabla u_{k+1}^\varepsilon(z)| &\leq \frac{\min_{1 \leq i \leq k} d_i^\varepsilon(a_{k+1}^\varepsilon)}{\min_{1 \leq i \leq k} d_i^\varepsilon(\lambda_{k+1}^\varepsilon z + a_{k+1}^\varepsilon)} + O\left(\frac{1}{d(z, S_{k+1})} + |z| + 1\right) \\ &= O\left(\frac{1}{d(z, S_{k+1})} + |z| + 1\right). \end{aligned} \tag{19}$$

Then $|\nabla u_{k+1}^\varepsilon|$ is bounded on every compact subset of $\overline{\Omega_{k+1}^\varepsilon} \setminus S_{k+1}$. Moreover, thanks to the conformal invariance of our equation, u_{k+1}^ε still satisfies (5). Hence, thanks to the standard elliptic theory, see Section 1 and [13], there exist a subsequence of u^ε (still denoted u^ε) and $\omega^{k+1} \in C^2(\overline{\Omega_{k+1}} \setminus S_{k+1})$ such that

$$u_{k+1}^\varepsilon \rightarrow \omega^{k+1} \quad \text{in } C_{loc}^1(\overline{\Omega_{k+1}} \setminus S_{k+1}),$$

and

$$\Delta \omega^{k+1} = -2\omega_x^{k+1} \wedge \omega_y^{k+1} \quad \text{on } \Omega_{k+1} \setminus S_{k+1}$$

where $\Omega_{k+1} = \lim_{\varepsilon \rightarrow 0} \Omega_{k+1}^\varepsilon$. As before, there are two possibilities; either Ω_{k+1} is the whole plane or this is a disk or a half-plane (which is conformally equivalent). If it is a disk or a half-plane, the boundary condition passes to the limit, that is to say, up to rotation,

$$\omega^{k+1}(\partial \Omega_{k+1}) \subset \mathbb{R}^2 \times \{0\}$$

and

$$\langle \omega_x^{k+1} \wedge \omega_y^{k+1}, N \rangle = 0 \quad \text{on } \partial \Omega_{k+1},$$

where $N = (0, 0, 1)$.

Moreover, thanks the conformal invariance of $\|\nabla \cdot\|_2$, up to extract a subsequence, we get

$$u_{k+1}^\varepsilon \rightharpoonup \omega^{k+1} \quad \text{in } L^2(\mathbb{R}^2)$$

and

$$\|\nabla \omega^{k+1}\|_2 \leq \liminf_{\varepsilon \rightarrow 0} \|\nabla u_{k+1}^\varepsilon\|_2 = \liminf_{\varepsilon \rightarrow 0} \|\nabla u^\varepsilon\|_2 < +\infty.$$

Then, ω^{k+1} is a solution of (2) on Ω_{k+1} , and ω^{k+1} has the desired shape. Finally, we need to show that ω^{k+1} is non-constant. This is trivial if $0 \notin S_{k+1}$, since in that case $|\nabla \omega^{k+1}(0)| = 1$. Else, for all i_0 such that

$$\frac{|a_{i_0}^\varepsilon - a_{k+1}^\varepsilon|}{\lambda_{k+1}^\varepsilon} = o(1),$$

thanks to (13) and (B_{k+1}) , we get

$$\lambda_{i_0}^\varepsilon = o(\lambda_{k+1}^\varepsilon).$$

Then mimicking the argument of the proof of (B_{k+1}) we show that

$$\nabla u_{k+1}^\varepsilon \rightarrow \nabla \omega^{k+1} \quad \text{on } B(0, \delta),$$

where $\delta > 0$. This leads in every case to $|\nabla \omega^{k+1}(0)| = 1$ and then ω^{k+1} is non-constant. This proves (P_{k+1}) in this second case. The study of these two cases ends the proof of Claim 1. \square

Then, we need to prove a claim about the energy of a sum of bubbles. In fact, using (B_k) , we show that the bubbles do not interact in a weak sense and that each one provides at least the energy of a simple hemisphere, that is to say 4π .

Claim 2. Let $k \in \mathbb{N}^*$ and

- (i) $\omega^1, \dots, \omega^k$ non-constant solutions of (2),
- (ii) $a_1^\varepsilon, \dots, a_k^\varepsilon$ sequences of $\overline{\mathbb{D}}$, and
- (iii) $\lambda_1^\varepsilon, \dots, \lambda_k^\varepsilon$ sequences of positive real numbers such that $\lim_{\varepsilon \rightarrow 0} \lambda_i^\varepsilon = 0$,

such that, with u^ε , they satisfy (P_k) . Then

$$\liminf_{\varepsilon \rightarrow 0} \|\nabla u^\varepsilon\|_2^2 \geq \sum_{i=1}^k \|\nabla \omega^i\|_2^2 \geq 4\pi k.$$

Proof. Let R be a positive real number, thanks to (B_k) , for ε small enough, we get

$$\int_{\mathbb{D}} |\nabla u^\varepsilon|^2 dz \geq \sum_{i=1}^k \int_{\mathbb{D} \cap B(a_i^\varepsilon, R\lambda_i^\varepsilon) \setminus \Omega_i^\varepsilon(R)} |\nabla u^\varepsilon|^2 dz,$$

where

$$\Omega_i^\varepsilon(R) = \left\{ z \in B(a_j^\varepsilon, R\lambda_j^\varepsilon) \text{ where } j \text{ is such that } \lim_{\varepsilon \rightarrow 0} \frac{\lambda_j^\varepsilon}{\lambda_i^\varepsilon} = 0 \right\}.$$

Then, thanks to (A_k) , we get

$$\begin{aligned} \int_{\mathbb{R}^2} |\nabla u^\varepsilon|^2 dz &\geq \sum_{i=1}^k \int_{(B(-\frac{a_i^\varepsilon}{\lambda_i^\varepsilon}, \frac{1}{\lambda_i^\varepsilon}) \cap B(0, R)) \setminus \Omega_i(R)} |\nabla \omega_i|^2 dz + \delta_{\varepsilon, R} \\ &\geq 4\pi k + \delta_{\varepsilon, R} \end{aligned} \tag{20}$$

where $\Omega_i(R) = \bigcup_{x \in S_i} B(x, \frac{1}{R})$ and $\lim_{R \rightarrow +\infty} \lim_{\varepsilon \rightarrow 0} \delta_{\varepsilon, R} = 0$. Here we bound the energy of a solution by the smallest possible, that is to say the energy of a hemisphere. \square

Proof of the theorem. We start setting $a_1^\varepsilon \in \overline{\mathbb{D}}$ and λ_1^ε as

$$|\nabla u^\varepsilon(a_1^\varepsilon)| = \sup_{z \in \mathbb{D}} |\nabla u^\varepsilon(z)|$$

and

$$|\nabla u^\varepsilon(a_1^\varepsilon)| = \frac{1}{\lambda_1^\varepsilon}.$$

Either λ_1^ε goes to infinity and then u^ε converges uniformly to 0 which proves the theorem. Or we set

$$u_1^\varepsilon(z) = u^\varepsilon(a_1^\varepsilon + \lambda_1^\varepsilon z) \quad \text{for all } z \in \Omega_1^\varepsilon$$

where $\Omega_1^\varepsilon = \{z \in \mathbb{R}^2 \text{ s.t. } a_1^\varepsilon + \lambda_1^\varepsilon z \in \mathbb{D}\}$.

It is clear that $|\nabla u_1^\varepsilon|$ is bounded on every compact subset of $\overline{\Omega_1^\varepsilon}$. Moreover thanks to conformal invariance of our equation, u_1^ε still satisfies (5). Hence, applying standard elliptic theory, see Section 1 and [13]. We see that there exists a subsequence of u_1^ε (still denoted u_1^ε) and $\omega^1 \in C^2(\Omega_1)$ such that

$$u_1^\varepsilon \rightarrow \omega^1 \quad \text{in } C_{loc}^2(\overline{\Omega_1})$$

and

$$\Delta \omega^1 = -2\omega_x^1 \wedge \omega_y^1 \quad \text{on } \Omega_1,$$

where $\Omega_1 = \lim_{\varepsilon \rightarrow 0} \Omega_1^\varepsilon$. There are two possibilities for Ω_1 ; it is either the whole plane or disk or a half-plane (which is conformally equivalent). In the last case, the boundary condition passes to the limit, that is to say, up to rotation,

$$\omega^1(\partial\Omega_1) \subset \mathbb{R}^2 \times \{0\}$$

and

$$\langle \omega_x^1 \wedge \omega_y^1, N \rangle = 0 \quad \text{on } \partial\Omega_k,$$

where $N = (0, 0, 1)$. Moreover, thanks to conformal invariance of $\|\nabla \cdot\|_2$, up to extract a subsequence, we get

$$u_1^\varepsilon \rightharpoonup \omega^1 \quad \text{in } L^2(\mathbb{R}^2)$$

and

$$\|\nabla\omega^1\|_2 \leq \liminf_{\varepsilon \rightarrow 0} \|\nabla u_1^\varepsilon\|_2 = \liminf_{\varepsilon \rightarrow 0} \|\nabla u^\varepsilon\|_2 < +\infty.$$

Then, thanks to Lemmas 2.1 and 2.2, ω^1 has the desired shape. Finally ω^1 is non-constant since $|\nabla\omega^1(0)| = 1$.

Now we can start our induction. Indeed, thanks to Claims 1 and 2 and the fact that the energy is uniformly bounded, there exists $k \in \mathbb{N}^*$ such that (P_k) is satisfied and

$$\limsup_{\varepsilon \rightarrow 0} \left(\min_{z \in \mathbb{D}} \left(\min_{0 \leq i \leq k} d_i^\varepsilon(z) \right) \left| \nabla \left(u^\varepsilon - \sum_{i=0}^k \omega_i^\varepsilon \right) (z) \right| \right) = 0, \tag{21}$$

where $\omega_i^\varepsilon = \omega_i(\frac{\cdot - a_i^\varepsilon}{\lambda_i^\varepsilon})$. This proves (A) and (B).

It suffices to show (C) to conclude. We start with the following claim.

Claim 3.

$$\left\| \nabla \left(u^\varepsilon - \sum_{i=1}^k \omega_i^\varepsilon \right) \right\|_2 \rightarrow 0 \quad \text{when } \varepsilon \rightarrow 0. \tag{22}$$

Proof. We set

$$R^\varepsilon = u^\varepsilon - \sum_{i=1}^k \omega_i^\varepsilon$$

and we assume that there exists $\delta > 0$ such that

$$\|\nabla R^\varepsilon\|_2 \geq \delta.$$

With those assumptions, we are going to prove the existence of a new bubble which will contradict (21). In order to find this bubble, we follow the method developed in [2].

First of all, we introduce the concentration function

$$C^\varepsilon(t) = \sup_{z \in \mathbb{D}} \int_{B(z,t)} |\nabla R^\varepsilon|^2 dz.$$

It is clear that C^ε is continuous, increasing with respect to t and that $C^\varepsilon(0) = 0$. We fix ν such that

$$0 < \nu < \min \left\{ \frac{1}{2C_0}, \frac{\delta}{2} \right\},$$

where C_0 is the constant in the Wente inequality given by Lemma A.1. Hence there exists $a^\varepsilon \in \overline{\mathbb{D}}$ and $\lambda^\varepsilon > 0$ such that

$$C^\varepsilon(\lambda^\varepsilon) = \int_{B(a^\varepsilon, \lambda^\varepsilon)} |\nabla R^\varepsilon|^2 dz = \nu.$$

Then we rescale around a^ε , setting $\tilde{f} = f(\lambda^\varepsilon \cdot + a^\varepsilon)$ for all $z \in \Omega^\varepsilon = \{z \in \mathbb{R}^2 \text{ s.t. } \lambda^\varepsilon z + a^\varepsilon \in \mathbb{D}\}$, and we get

$$\int_{\Omega^\varepsilon} |\nabla \tilde{R}^\varepsilon|^2 dz = \|\nabla R^\varepsilon\|_2^2 \leq C,$$

and

$$\|\tilde{R}^\varepsilon\|_\infty \leq C,$$

where C is a positive real. Moreover, using (5), we remark that \tilde{R}^ε satisfies

$$\Delta \tilde{R}^\varepsilon = -2\tilde{R}_x^\varepsilon \wedge \tilde{R}_y^\varepsilon + O \left(\sum_{i=0}^k |\nabla \tilde{\omega}_i^\varepsilon| \left(\sum_{j \neq i} |\nabla \tilde{\omega}_j^\varepsilon| + |\nabla \tilde{R}^\varepsilon| \right) \right).$$

However, thanks to (B), we get

$$|\nabla \tilde{\omega}_i^\varepsilon| |\nabla \tilde{\omega}_j^\varepsilon| \rightarrow 0 \quad \text{in } L^1_{loc}(\overline{\Omega^0}) \text{ for } i \neq j$$

and, thanks to (21), we get

$$|\nabla \tilde{\omega}_i^\varepsilon| |\nabla \tilde{R}^\varepsilon| \rightarrow 0 \quad \text{in } L^1_{loc}(\overline{\Omega^0}) \text{ for all } i,$$

with $\Omega^0 = \lim_{\varepsilon \rightarrow 0} \Omega^\varepsilon$. Finally

$$\Delta \tilde{R}^\varepsilon = -2\tilde{R}_x^\varepsilon \wedge \tilde{R}_y^\varepsilon + h^\varepsilon,$$

where $h^\varepsilon \rightarrow 0$ in $L^1_{loc}(\overline{\Omega^0})$ when $\varepsilon \rightarrow 0$. Then, up to extract a subsequence, we get

$$\tilde{R}^\varepsilon \rightarrow R \quad \text{p.p. on } \Omega^0$$

and

$$\nabla \tilde{R}^\varepsilon \rightharpoonup \nabla R \quad \text{weakly in } L^2(\Omega^0).$$

Moreover R is a weak solution of

$$\Delta R = -2R_x \wedge R_y \quad \text{on } \Omega^0.$$

Now, thanks to our choice of v , we are going to show that the weak convergence is in fact strong. Let $v^\varepsilon = \tilde{R}^\varepsilon - R$, then v^ε satisfy

$$\Delta v^\varepsilon = -2v_x^\varepsilon \wedge v_y^\varepsilon - 2(v_x^\varepsilon \wedge R_y + R_x \wedge v_y^\varepsilon) + h^\varepsilon.$$

Moreover, thanks to Corollary A.1, there exists $\psi_\varepsilon \in H_0^1(\Omega^0)$ a solution of

$$\Delta \psi^\varepsilon = -2(v_x^\varepsilon \wedge R_y + R_x \wedge v_y^\varepsilon)$$

satisfying

$$\|\nabla \psi^\varepsilon\|_2 + \|\psi^\varepsilon\|_\infty \leq \|\nabla v^\varepsilon\|_2 \|\nabla R\|_2. \tag{23}$$

However,

$$\int_{\Omega^0} |\nabla \psi^\varepsilon|^2 dz = -2 \int_{\Omega^0} \langle \psi^\varepsilon, v_x^\varepsilon \wedge R_y + R_x \wedge v_y^\varepsilon \rangle dz.$$

Then using (23), we get that $\psi^\varepsilon \wedge R_x$ and $\psi^\varepsilon \wedge R_y$ are bounded in $L^2(\Omega^0)$. Hence, since $\nabla v^\varepsilon \rightarrow 0$ weakly in $L^2(\Omega^0)$, we get that

$$\int_{\Omega^0} |\nabla \psi^\varepsilon|^2 dz \rightarrow 0.$$

Then we deduce that

$$\Delta v^\varepsilon = -2v_x^\varepsilon \wedge v_y^\varepsilon + g^\varepsilon,$$

where $g^\varepsilon \rightarrow 0$ in $D'(\overline{\Omega^0})$.

Finally, let $\phi \in C_c^\infty(\Omega^0)$ be such that $\text{supp}(\phi)$ is contained in a ball of radius 1, using Lemma A.1, we get

$$\begin{aligned} \int_{\Omega^0} |\nabla(\phi v^\varepsilon)|^2 dz &= -2 \int_{\Omega^0} \langle v^\varepsilon, \phi v_x^\varepsilon \wedge \phi v_y^\varepsilon \rangle dz + o(1), \\ &\leq 2(C_0 \|\nabla v^\varepsilon|_{\text{supp}(\phi)}\|_2) \|\nabla(\phi v^\varepsilon)\|_2^2 + o(1). \end{aligned}$$

Thanks to our choice of λ^ε , we get $C_0 \|\nabla v^\varepsilon|_{\text{supp}(\phi)}\|_2 \leq \frac{1}{2}$, which gives finally

$$\int_{\Omega^0} |\nabla(\phi v^\varepsilon)|^2 dz = o(1)$$

which proves

$$\nabla \tilde{R}^\varepsilon \rightarrow \nabla R \quad \text{strongly in } L^2_{loc}(\overline{\Omega^0}).$$

Indeed, we can remark that R isn't constant since $\|\nabla R\|_2 = \nu > 0$. But, thanks to (21), we have, for all $z \in \mathbb{R}^2$, that there exists i such that

$$|\nabla \tilde{R}^\varepsilon(z)| = o\left(\frac{1}{\sqrt{(\frac{\lambda_i^\varepsilon}{\lambda^\varepsilon})^2 + |z + \frac{a^\varepsilon - a_i^\varepsilon}{\lambda^\varepsilon}|^2}}\right),$$

which is a contradiction and proves (22). \square

In order to conclude, we have to transform this H^1 -estimate in an L^∞ -estimate. An idea could be to use the Wente inequality as it is done by Brezis and Coron in [2] in order to get L^∞ -estimate. But contrary to Brezis and Coron, here we don't control what happens on the boundary. In order to overpass this difficulty we are going to extend our surfaces.

Usually extend a surface across its boundary smoothly is not an easy fact, but here, thanks to the fact that our surfaces and the boundary of our domains meet orthogonally, this will be possible without perturbing too much the condition to be with constant mean curvature.

The idea is to reflect our surface through $\partial\Omega^\varepsilon$ which is almost a plane so that our transformation will be almost an isometry (in fact a symmetry) and will almost conserve the mean curvature. Moreover the new surfaces will be at least $C^{1,1}$ thanks to the fact that our surfaces meet $\partial\Omega^\varepsilon$ orthogonally.

Since $\partial\Omega^\varepsilon$ converges uniformly to a plane, there exists a diffeomorphism $\psi^\varepsilon : B(0, 2R) \rightarrow \mathbb{R}^3$, where R is chosen such that $u^\varepsilon(\mathbb{D}) \subset B(0, R)$, which sends $\partial\Omega^\varepsilon \cap B(0, 2)$ in $\mathbb{R}^2 \times \{0\}$ and which preserves the orthogonality on $\partial\Omega^\varepsilon$. In fact it suffices to straighten up the local foliation of the normal bundle of $\partial\Omega^\varepsilon$ to $\mathbb{R}^2 \times \mathbb{R}$. Then now we get new surfaces which have almost constant mean curvature equal to 1. Then we extend our map to S^2 . Here S^2 will be identified with the Riemann sphere $\hat{\mathbb{C}}$. We set

$$v^\varepsilon(z) = s\left(v^\varepsilon\left(\frac{1}{z}\right)\right) \quad \text{for all } z \in \hat{\mathbb{C}} \setminus \mathbb{D},$$

where s is the symmetry through $\mathbb{R}^2 \times \{0\}$ and $v^\varepsilon = \psi^\varepsilon \circ u^\varepsilon$. Using the fact that $v^\varepsilon(\overline{\mathbb{D}})$ and $\mathbb{R}^2 \times \{0\}$ meet orthogonally we easily show that v^ε is $C^{1,1}$. Then we set $\tilde{u}^\varepsilon = \psi^{-1} \circ v^\varepsilon$ which is also $C^{1,1}$ and its mean curvature uniformly converges to 1.

Then we are in position to prove our theorem. Assume by contradiction that

$$d\left(\tilde{\Sigma}^\varepsilon, \bigcup_{i=1}^k \tilde{B}_i\right) \rightarrow 0,$$

where $\tilde{\Sigma}^\varepsilon = \tilde{u}^\varepsilon(S^2)$ and \tilde{B}_i is the union of B_i and its symmetry through $T_0\partial\Omega^\varepsilon$.

Then there exists $y^\varepsilon \in \tilde{\Sigma}^\varepsilon$ such that

$$d\left(y^\varepsilon, \bigcup_{i=1}^k \tilde{B}_i\right) \rightarrow 0.$$

Let $z^\varepsilon \in \hat{\mathbb{C}}$ be such that $\tilde{u}^\varepsilon(z^\varepsilon) = y^\varepsilon$. We are going to prove that there is some area in a neighbourhood of z^ε . The idea is that if a surface has bounded mean curvature, passes through the center of a ball, and has no boundary inside the ball, then it has to use a certain amount of area to leave the ball. Since the mean curvature is bounded, then the Gaussian curvature of our surface is uniformly bounded from above by a constant K_0 . Let $r_0 > 0$ be such that

$$B(y^\varepsilon, r_0) \cap \left(\bigcup_{i=1}^k \tilde{B}_i\right) = \emptyset.$$

Then using a Bishop comparison, like Theorem III.4.2 of [4], we see that

$$\text{Vol}\left(B\left(y^\varepsilon, \frac{r_0}{2}\right) \cap \tilde{\Sigma}^\varepsilon\right) \geq \text{Vol}\left(B_{\tilde{\Sigma}^\varepsilon}\left(y^\varepsilon, \frac{r_0}{2}\right)\right) \geq \text{Vol}\left(B_{M_{K_0}}\left(y^\varepsilon, \frac{r_0}{2}\right)\right) \geq C_0 r_0^2,$$

where M_{K_0} is the space of constant curvature K^0 and C^0 a positive constant. Hence we see that u^ε necessarily get some area in a neighbourhood of z^ε whose image is far from the bubbles, which is a contradiction with Claim 3 since all the area of u^ε is devoted to cover the bubbles. This proves (C) and achieves the proof of the theorem. \square

3.2. There is at least one hemisphere in the decomposition

In order to show our result, we have to eliminate the sequence of surfaces whose boundaries collapse. It suffices to show that in Theorem 3.1, there is at least one bubble whose domain of definition is not the whole plane, that is to say there is at least one hemisphere.

Since our surfaces are embedded, we can assume that our bubbles are simple. In fact, we just need to prove that $\max\{\deg P_i, \deg Q_i\} = 1$ for all $1 \leq i \leq p$, with $\omega_i = \pi_{P_i}\left(\frac{P_i}{Q_i}\right)$ where $\frac{P_i}{Q_i}$ is irreducible. But this is an easy consequence of the fact that our surfaces are embedded, (A) and the following lemma.

Lemma 3.1. *Let $u^\varepsilon : B(0, 1) \rightarrow \mathbb{R}^3$ a sequence of smooth embedding such that there exist $u^0 \in C^1(B(0, 1), \mathbb{R}^3)$ and*

$$u^\varepsilon \rightarrow u^0 \quad \text{in } C_{loc}^2(B(0, 1) \setminus \{0\}).$$

Then u^0 can't be a multiple parametrization, that is to say there is no embedded $U_0 \in C^1(B(0, 1), \mathbb{R}^3)$, $\Phi \in \mathcal{O}(B(0, 1), \mathbb{C})$ a holomorphic function and an integer $k \geq 2$ such that

$$u^0 = U^0 \circ \Phi$$

and

$$\Phi(z) = z^k + o(|z|^k) \quad \text{as } z \rightarrow 0.$$

Proof. First of all, up to a diffeomorphism of a neighbourhood of 0, we can assume that

$$u^\varepsilon \rightarrow U^0(z^l) \quad \text{in } C_{loc}^2(B(0, \delta) \setminus \{0\}),$$

where $l \geq 2$ and $\delta > 0$. Let $A_\delta = B(0, \frac{\delta}{2}) \setminus B(0, \frac{\delta}{3})$ and C_r be the cylinder of center $U^0(0)$, radius r and orthogonal to $T_{U^0(0)}U^0(B(0, 1))$, the tangent plane to the image of U^0 at $U^0(0)$. Let $\delta > 0$ and $r > 0$ be small enough such that $C_r \cap U_0(A_\delta)$ is a simple curve. Then, for ε small enough, we easily see that the intersection of $u^\varepsilon(A_\delta)$ and C_r turns l times around the cylinder, hence $u^\varepsilon(A_\delta)$ necessarily intersect, which is a contradiction and proves the lemma. \square

Claim. *Let u^ε be a sequence of C^2 -solutions of (5). We note p the number of bubbles given by the decomposition 3.1, this number splits into k spheres and l hemispheres, such that $p = k + l$. Then we necessarily have that $l \geq 1$.*

Proof. We assume by contradiction that $l = 0$. We show first that necessarily $k \leq 1$, that is to say there is no neck as in Fig. 2.

We assume for contradiction that $k \geq 2$ and we consider the highest bubble, that is to say the one which corresponds to the smallest λ_i^ε . Up to reorder, we assume that $i = 1$. Thanks to (B), there is no bubble closed to this one, that is to say

$$u_1^\varepsilon \rightarrow \omega^1 \quad \text{in } C_{loc}^2(\mathbb{R}^2)$$

where $u_1^\varepsilon = u^\varepsilon(a_1^\varepsilon + \lambda_1^\varepsilon \cdot)$.

We claim that the highest bubble is over another bubble. More precisely, there exist $i > 1$ and $R_0 > 0$ such that for all $R > 0$ one gets $B(a_1^\varepsilon, R\lambda_1^\varepsilon) \subset B(a_i^\varepsilon, R_0\lambda_i^\varepsilon)$ for ε small enough.

Else this bubble would be isolated and will become tangent to $\partial\Omega$ at 0. Indeed, there exists $z^0 \in \partial\mathbb{D}$ such that for all $R > 0$ and for all $z^\varepsilon \in \partial B(a_1^\varepsilon, R\lambda_1^\varepsilon)$, there exists a curve Γ of $\bar{\mathbb{D}}$ joining z^ε to z^0 staying far from the other bubble.




Fig. 2. Two bubbles joined by a neck.




Fig. 3. $\tilde{\Sigma}^\varepsilon$ range by u^ε of a neighbourhood of the highest bubble.

Hence thanks to the estimate (21), for ε small enough, we see that the bubble ω_1^ε would be almost tangent to $\partial\Omega^\varepsilon$. This makes impossible the existence of an other isolated bubble, since it would be also almost tangent to $\partial\Omega^\varepsilon$ at the same point, since we can take the same z^0 for the two bubbles. This would contradict the fact that the bubble is embedded and in the interior of Ω^ε . Hence the second bubble should be over the first one and so higher, which is a contradiction.

There exist i_0 and $R_0 > 0$ such that for all $R > 0$ we get $B(a_1^\varepsilon, R\lambda_1^\varepsilon) \subset B(a_{i_0}^\varepsilon, R_0\lambda_{i_0}^\varepsilon)$. Then we choose the minimal $\lambda_{i_0}^\varepsilon$ satisfying this property. In this case we consider a neighbourhood of a_1^ε , $B(a_1^\varepsilon, r\lambda_{i_0}^\varepsilon)$, where $r > 0$ is chosen such that this neighbourhood contains no other bubble. For ε small enough, the range of this neighbourhood by u^ε , which will be noted by $\tilde{\Sigma}^\varepsilon$, seems like a sphere glued on a spherical cap, see Fig. 3.

Now we are in position to apply the Aleksandrov reflexion principle, as described in Chapter VII of [14]. Let P^ε be the tangent plane to $\tilde{\Sigma}^\varepsilon$ at $u^\varepsilon(a_1^\varepsilon)$ and ν^ε the exterior normal at this point. We set $P_t^\varepsilon = P^\varepsilon + t\nu^\varepsilon$, $\tilde{\Sigma}_t^{\varepsilon,+} = \tilde{\Sigma}^\varepsilon \cap \{P^\varepsilon + u\nu^\varepsilon \text{ s.t. } u \geq t\}$ and $\tilde{\Sigma}_t^{\varepsilon,-}$ the reflexion of $\tilde{\Sigma}_t^{\varepsilon,+}$ with respect to P_t^ε . We consider the first negative time t such that $\tilde{\Sigma}_t^{\varepsilon,-}$ meets $\tilde{\Sigma}^\varepsilon$. For ε small enough, the contact point of $\tilde{\Sigma}_t^{\varepsilon,-}$ and $\tilde{\Sigma}^\varepsilon$ can't belong to the boundary thanks to the presence of the neck. Moreover at the contact point the surfaces get the same orientation since $\tilde{\Sigma}_t^{\varepsilon,-}$ comes from the interior of the bubble. Hence applying the Aleksandrov principle, we get $\tilde{\Sigma}_t^{\varepsilon,-} = \tilde{\Sigma}^\varepsilon \setminus \tilde{\Sigma}_t^{\varepsilon,+}$, which is clearly a contradiction with the fact that the surface is embedded and with a boundary. Which proves that $k \leq 1$.

Now we have to exclude $k = 0$ and $k = 1$. If $k = 0$, then $p = 0$ and in that case the surface collapses. In fact his area will go to 0, see the proof of Theorem 3.1. In that case, we rescale our space in order to get a new surface, denoted by $\hat{\Sigma}^\varepsilon$, whose area is equal to 1. This imposes to the mean curvature of our new surface to go to 0. Then our new sequence of surfaces $\hat{\Sigma}^\varepsilon$ goes to a minimal surface which bounds a plane curve. Indeed to insure the convergence it suffices to prove that $|\nabla\hat{u}^\varepsilon|$ is uniformly bounded, where \hat{u}^ε is a conformal parametrization of $\hat{\Sigma}^\varepsilon$. The regularity theory given in Section 1, will give the convergence in $C^2(\mathbb{D})$.

Let us assume by contradiction that $\sup_{\mathbb{D}} |\nabla\hat{u}^\varepsilon| \rightarrow +\infty$ when $\varepsilon \rightarrow 0$. Then we set $a_1^\varepsilon \in \mathbb{D}$ and λ_1^ε such that

$$|\nabla\hat{u}^\varepsilon(a_1^\varepsilon)| = \sup_{z \in \mathbb{D}} |\nabla\hat{u}^\varepsilon(z)|$$

and

$$|\nabla\hat{u}^\varepsilon(a_1^\varepsilon)| = \frac{1}{\lambda_1^\varepsilon}.$$

Then we set

$$\hat{u}_1^\varepsilon(z) = \hat{u}^\varepsilon(a_1^\varepsilon + \lambda_1^\varepsilon z) \quad \text{for all } z \in \Omega_1^\varepsilon$$

where $\Omega_1^\varepsilon = \{z \in \mathbb{R}^2 \text{ s.t. } a_1^\varepsilon + \lambda_1^\varepsilon z \in \mathbb{D}\}$.

It is clear that $|\nabla\hat{u}_1^\varepsilon|$ is bounded on every compact subset of $\overline{\Omega_1^\varepsilon}$. Moreover, thanks to conformal invariance of our equation, \hat{u}_1^ε satisfies

$$\Delta \hat{u}_1^\varepsilon = o((\hat{u}_1^\varepsilon)_x \wedge (\hat{u}_1^\varepsilon)_y)$$

and

$$\langle (\hat{u}_1^\varepsilon)_x, (\hat{u}_1^\varepsilon)_y \rangle = |(\hat{u}_1^\varepsilon)_x| - |(\hat{u}_1^\varepsilon)_y| = 0.$$

Hence, applying standard elliptic theory, see Section 1 and [13], we see that there exists a subsequence of \hat{u}_1^ε (still denoted by \hat{u}_1^ε) and $\beta^1 \in C^2(\Omega_1)$ such that

$$\hat{u}_1^\varepsilon \rightarrow \beta^1 \quad \text{in } C_{loc}^2(\overline{\Omega_1})$$

and

$$\begin{aligned} \Delta \beta_1 &= 0, \\ \langle (\beta_1)_x, (\beta_1)_y \rangle &= |(\beta_1)_x| - |(\beta_1)_y| = 0, \end{aligned}$$

where $\Omega_1 = \lim_{\varepsilon \rightarrow 0} \Omega_1^\varepsilon$. Then there are two possibilities for Ω_1 , either it is the whole plane or it is a disk or a half-plane (which is conformally equivalent). In this last case the boundary condition passes to the limit, that is to say, up to a rotation,

$$\beta^1(\partial\Omega_1) \subset \mathbb{R}^2 \times \{0\}$$

and

$$\langle \beta_x^1 \wedge \beta_y^1, N \rangle = 0 \quad \text{on } \partial\Omega_k,$$

where $N = (0, 0, 1)$. In that case β_1 can be extended by symmetry in a C^1 function defined on the whole plane.

Moreover, thanks to the conformal invariance of $\|\nabla \cdot\|_2$, we get

$$\|\nabla \beta^1\|_2 \leq 2 \liminf_{\varepsilon \rightarrow 0} \|\nabla \hat{u}_1^\varepsilon\|_2 = 2 \liminf_{\varepsilon \rightarrow 0} \|\nabla \hat{u}^\varepsilon\|_2 = 2.$$

Thanks to the Liouville theorem, we necessarily get $\nabla \beta_1 \equiv 0$, which is a contradiction with the fact that $|\nabla \beta_1(0)| = 1$. This proves that $|\nabla \hat{u}^\varepsilon|$ is uniformly bounded and also the convergence of the sequence of surfaces $\hat{\Sigma}^\varepsilon$.

With this convergence, the boundary condition passes to the limit, that is to say the minimal surface which is obtained meets the plane which contains its boundary orthogonally. But thanks to classical theory of minimal surfaces, see [5], these surfaces should be flat, which contradicts the fact it must meet orthogonally the plane which contains its boundary.




Fig. 4. A bubble meeting $\partial\Omega$.

Finally, the last possibility is $k = 1$, that is to say there is only one bubble as in Fig. 4.

But we can apply once more the Aleksandrov reflexion principle with respect to the tangent plane Σ^ε at the furthest point to $\partial\Omega^\varepsilon$. Then the contact point between the surface and the reflected part is necessarily at the boundary, else the surface should be closed without boundary. Indeed if we have a contact in the interior, the local equality given by the reflexion principle would be global thanks to connexity, which is impossible since the upper part is simply connected and the lower is not. Hence the contact is done at the boundary $\partial\Sigma^\varepsilon$, but the tangent plane to Σ^ε at the furthest point to $\partial\Omega^\varepsilon$ becomes parallel to the one of $\partial\Omega$ at 0, which forces the angle between $\partial\Omega^\varepsilon$ and Σ^ε at the contact point to go to zero when ε goes to 0, which is a contradiction and achieves the proof of the claim. \square




Fig. 5. ω^ε which is bounded by $\partial\Sigma^\varepsilon$.

3.3. Proof of Theorem 0.1

Thanks to the previous section, $u^\varepsilon(S^1)$ converges uniformly to a union of circles with radius 1 centered at points (c_i) of $T_0\partial\Omega$ (see Fig. 5). Now we are in position to prove Theorem 0.1. In order to do it we apply the balancing formula (4) to the sequence Σ^ε . This gives

$$2 \int_{\omega^\varepsilon} \vec{N}^\varepsilon dv - \int_{\partial\Sigma^\varepsilon} \vec{v}^\varepsilon d\sigma = 0 \tag{24}$$

where \vec{v}^ε is the conormal.

The fact that Σ^ε and Ω^ε meet orthogonally imposes $\vec{v}^\varepsilon = \vec{N}^\varepsilon$. We make a Taylor expansion of \vec{N}^ε . Since $\partial\Omega^\varepsilon$ is a graph above its tangent plane, we make the expansion in those coordinates. In fact, we are not going to do the expansion with respect to 0 but with respect to $c^\varepsilon \in \partial\Omega^\varepsilon$ a point closed to 0 which will be fixed later.

$$\vec{N}^\varepsilon(z) = \vec{N}^\varepsilon(c^\varepsilon) + \varepsilon d\vec{N}_{c^\varepsilon}(z - c^\varepsilon) + \varepsilon^2 d^2\vec{N}_{c^\varepsilon}(z - c^\varepsilon)(z - c^\varepsilon) + o(\varepsilon^2).$$

At the first order, the left-hand side (24) gives

$$(2|\omega^\varepsilon| - |\partial\Sigma^\varepsilon|)\vec{N}(c^\varepsilon). \tag{25}$$

We can remark here that, thanks to Theorem 3.1, we have,

$$|\omega^\varepsilon| \rightarrow l\pi,$$

and

$$\liminf_{\varepsilon \rightarrow 0} |\partial\Sigma^\varepsilon| \geq 12\pi.$$

In the first equality, we use the L^∞ -convergence while in the second, we use the C^2_{loc} -convergence. Then, thanks to (25), we get

$$\lim_{\varepsilon \rightarrow 0} |\partial\Sigma^\varepsilon| = 12\pi,$$

which proves that $\partial\Sigma^\varepsilon$ converges to a union of l circles as a current which justifies the fact that we will pass to the limit in the integral defined over this set.

In order to eliminate (25) we project the left-hand side term of (24) orthogonally to $\vec{N}^\varepsilon(c^\varepsilon)$, which gives to the second order

$$\varepsilon\pi^\varepsilon \left(2 \int_{\omega^\varepsilon} d\vec{N}_{c^\varepsilon}(z - c^\varepsilon) dv - \int_{\partial\Sigma^\varepsilon} d\vec{N}_{c^\varepsilon}(z - c^\varepsilon) d\sigma \right) \tag{26}$$

where π^ε is the orthogonal projection parallel to $\vec{N}^\varepsilon(c^\varepsilon)$.

Then we remark that there exists c^ε such that (26) vanishes. Indeed

$$2 \int_{\omega^\varepsilon} (z - c^\varepsilon) dv - \int_{\partial\Sigma^\varepsilon} (z - c^\varepsilon) d\sigma$$

is the weighted barycenter of $(\omega^\varepsilon, 2)$ and $(\partial\Sigma^\varepsilon, -1)$, then it suffices to choose c^ε as the corresponding barycenter to vanish (26).

Then it remains the order two terms, in which we pass to the limit after dividing them by ε^2 , which gives

$$\pi^0 \left(2 \int_{\omega^0} d^2 \vec{N}_0(z - c^0)(z - c^0) dv - \int_{\partial \Sigma^0} d^2 \vec{N}_{c^0}(z - c^0)(z - c^0) d\sigma \right) = 0 \tag{27}$$

where π^0 is the orthogonal projection parallel to $\vec{N}(0)$, c^0 , ω^0 and $\partial \Sigma^0$ are respectively the limit of c^ε , ω^ε and $\partial \Sigma^\varepsilon$. As already remarked at the beginning of this section, ω^0 and $\partial \Sigma^0$ are respectively a union of disks with radius 1 and union of circles with radius 1 centered at some points c_i . Then we decompose the integral on this subset, which gives

$$\begin{aligned} & \sum_{i=1}^l \pi^0 \left(2 \int_{D(c_i, 1)} d^2 \vec{N}_0(z - c^0)(z - c^0) dv - \int_{\partial D(c_i, 1)} d^2 \vec{N}_{c^0}(z - c^0)(z - c^0) d\sigma \right) \\ &= \sum_{i=1}^l \pi^0 \left(2 \int_{D(c_i, 1)} d^2 \vec{N}_0(z - c^i)(z - c^i) dv - \int_{\partial D(c_i, 1)} d^2 \vec{N}_{c^0}(z - c^i)(z - c^i) d\sigma \right). \end{aligned}$$

Here we use the fact that the integral vanishes if it contains an odd number of $z - c^i$ and the fact that $2|D(c_i, 1)| = |\partial D(c_i, 1)|$. Now we integrate, which traces $d^2 \vec{N}$ and gives

$$l \left(2 \int_{\mathbb{D}} |z|^2 dz - 1 \right) \pi^0(\Delta \vec{N}(0)) = 0.$$

But the general equation of a Gauss map of an immersion X is given by

$$\Delta \vec{N} = |\nabla \vec{N}|^2 \vec{N} - 2|\nabla X|^2 \nabla H(u), \tag{28}$$

hence we get

$$\frac{l}{2} \nabla H(0) = 0,$$

which achieves the proof of the theorem. \square

Acknowledgements

I would to thank Frank Pacard for submitted me this question and the work of Fall. It is also my pleasure to express my deep thanks to my thesis advisor O. Druet for his encouragement and helpful comments during the preparation of this work.

Appendix A. Wente’s inequality and application

The aim of this section is to remind some Wente’s inequalities, originally proved in [23].

Theorem A.1. *Let Ω be a bounded open set of \mathbb{R}^2 and $a, b \in H^1(\Omega)$. Let $u \in W_0^{1,1}(\Omega)$ be the solution of*

$$\Delta u = a_x b_y - a_y b_x \quad \text{on } \Omega,$$

then

$$\|u\|_\infty \leq \frac{1}{2\pi} \|\nabla a\|_2 \|\nabla b\|_2,$$

and

$$\|\nabla u\|_2 \leq \sqrt{\frac{3}{16\pi}} \|\nabla a\|_2 \|\nabla b\|_2.$$

Moreover the constant are optimal.

Which is remarkable here is that the constant is independent of Ω . We will find the proof in [22] and [12], see also [24] and [1]. These inequalities have been extend to function defined on surfaces. In particular, we have the following theorem.

Theorem A.2. *Let Σ a compact Riemannian surface without boundary and $v \in H^1(\Sigma, \mathbb{R}^2)$. Then if $u \in W^{1,1}(\Sigma, \mathbb{R})$ be the solution of*

$$\Delta u = \det(\nabla v) \quad \text{on } \Sigma,$$

then

$$\text{osc}(u) + \|\nabla u\|_2 \leq \left(\frac{1}{4\pi} + \sqrt{\frac{3}{128\pi}} \right) \|\nabla v\|_2^2,$$

where $\text{osc}(u) = \sup_{x,y \in \Sigma} |u(x) - u(y)|$.

Then, assuming that $u \in H^1$, we extend such an equality to $\Omega = \mathbb{R}^2$.

Corollary A.1. *Let $v \in H^1(\mathbb{R}^2, \mathbb{R}^3)$ and $u \in H^1(\mathbb{R}^2, \mathbb{R}^3)$ be a solution of*

$$\Delta u = -2v_x \wedge v_y \quad \text{on } \mathbb{R}^2$$

then

$$\text{osc}(u) + \|\nabla u\|_2 \leq \left(\frac{1}{\pi} + \sqrt{\frac{3}{8\pi}} \right) \|\nabla v\|_2^2.$$

Here the constant is *a priori* not optimal.

Proof. Let π the standard stereographic projection from S^2 to \mathbb{R}^2 . Thanks to the conformal invariance of the equation, $u \circ \pi^{-1}$ and $v \circ \pi^{-1}$ satisfy the hypothesis of Theorem A.2 when $\Sigma = S^2$, hence we get that

$$\text{osc}(u^1) + \|\nabla u^1\|_2 \leq \left(\frac{1}{2\pi} + \sqrt{\frac{3}{32\pi}} \right) (\|\nabla v^2\|_2^2 + \|\nabla v^3\|_2^2),$$

$$\text{osc}(u^2) + \|\nabla u^2\|_2 \leq \left(\frac{1}{2\pi} + \sqrt{\frac{3}{32\pi}} \right) (\|\nabla v^1\|_2^2 + \|\nabla v^3\|_2^2)$$

and

$$\text{osc}(u^3) + \|\nabla u^3\|_2 \leq \left(\frac{1}{2\pi} + \sqrt{\frac{3}{32\pi}} \right) (\|\nabla v^1\|_2^2 + \|\nabla v^2\|_2^2).$$

Then summing these inequalities, we get the desired inequality. \square

To conclude this section we remind a useful Wente's type inequality, see [3] for example.

Lemma A.1. *Let $u \in H^1(\mathbb{D}) \cap L^\infty(\mathbb{D})$ and $v \in H_0^1(\mathbb{D})$, then there exists C , independent of u and v , such that*

$$\left| \int_{\Omega} \langle u, v_x \wedge v_y \rangle \right| \leq C \|\nabla u\|_2 \|\nabla v\|_2^2.$$

References

- [1] Sami Baraket, Estimations of the best constant involving the L^∞ norm in Wente’s inequality, *Ann. Fac. Sci. Toulouse Math.* (6) 5 (3) (1996) 373–385.
- [2] H. Brezis, J.-M. Coron, Convergence of solutions of H -systems or how to blow bubbles, *Arch. Ration. Mech. Anal.* 89 (1) (1985) 21–56.
- [3] Haïm Brezis, Jean-Michel Coron, Multiple solutions of H -systems and Rellich’s conjecture, *Comm. Pure Appl. Math.* 37 (2) (1984) 149–187.
- [4] Isaac Chavel, *Riemannian Geometry: A Modern Introduction*, second edition, Cambridge Studies in Advanced Mathematics, vol. 98, Cambridge University Press, Cambridge, 2006.
- [5] Tobias H. Colding, William P. Minicozzi II, *Minimal Surfaces*, Courant Lecture Notes in Mathematics, vol. 4, New York University, Courant Institute of Mathematical Sciences, New York, 1999.
- [6] Ulrich Dierkes, Stefan Hildebrandt, Albrecht Küster, Ortwin Wohlrab, *Minimal Surfaces. II: Boundary Regularity*, Grundlehren der Mathematischen Wissenschaften (Fundamental Principles of Mathematical Sciences), vol. 296, Springer-Verlag, Berlin, 1992.
- [7] Olivier Druet, Sharp local isoperimetric inequalities involving the scalar curvature, *Proc. Amer. Math. Soc.* 130 (8) (2002) 2351–2361 (electronic).
- [8] Olivier Druet, Emmanuel Hebey, Frédéric Robert, *Blow-up Theory for Elliptic PDEs in Riemannian Geometry*, Mathematical Notes, vol. 45, Princeton University Press, Princeton, NJ, 2004.
- [9] Mouhamed Moustapha Fall, Embedded disc-type surfaces with large constant mean curvature and free boundaries, preprint SISSA, 2007.
- [10] Mouhamed Moustapha Fall, Area-minimizing regions with small volume in Riemannian manifolds with boundary, *Pacific J. Math.* 244 (2) (2010) 235–260.
- [11] Mouhamed Moustapha Fall, Fethi Mahmoudi, Hypersurfaces with free boundary and large constant mean curvature: Concentration along submanifolds, *Ann. Sc. Norm. Super. Pisa Cl. Sci.* (5) 7 (3) (2008) 407–446.
- [12] Yuxin Ge, Estimations of the best constant involving the L^2 norm in Wente’s inequality and compact H -surfaces in Euclidean space, *ESAIM Control Optim. Calc. Var.* 3 (1998) 263–300 (electronic).
- [13] David Gilbarg, Neil S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Classics in Mathematics, Springer-Verlag, Berlin, 2001 (reprint of the 1998 edition).
- [14] Heinz Hopf, *Differential Geometry in the Large*, Lecture Notes in Mathematics, vol. 1000, Springer-Verlag, Berlin, 1983 (notes taken by Peter Lax and John Gray, with a preface by S.S. Chern).
- [15] Katsuei Kenmotsu, *Surfaces with Constant Mean Curvature*, Translations of Mathematical Monographs, vol. 221, American Mathematical Society, Providence, RI, 2003 (translated from the 2000 Japanese original by Katsuhiko Moriya and revised by the author).
- [16] Nicholas J. Korevaar, Rob Kusner, Bruce Solomon, The structure of complete embedded surfaces with constant mean curvature, *J. Differential Geom.* 30 (2) (1989) 465–503.
- [17] Paul Laurain, Concentration of CMC surfaces in a Riemannian manifold, I.M.R.N., in press.
- [18] Frank Morgan, *Geometric Measure Theory: A Beginner’s Guide*, fourth edition, Elsevier/Academic Press, Amsterdam, 2009.
- [19] Manuel Ritoré, Antonio Ros, Some updates on isoperimetric problems, *Math. Intelligencer* 24 (3) (2002) 9–14.
- [20] Antonio Ros, Enaldo Vergasta, Stability for hypersurfaces of constant mean curvature with free boundary, *Geom. Dedicata* 56 (1) (1995) 19–33.
- [21] Michael Struwe, *Plateau’s Problem and the Calculus of Variations*, Mathematical Notes, vol. 35, Princeton University Press, Princeton, NJ, 1988.
- [22] Peter Topping, The optimal constant in Wente’s L^∞ estimate, *Comment. Math. Helv.* 72 (2) (1997) 316–328.
- [23] Henry C. Wente, An existence theorem for surfaces of constant mean curvature, *J. Math. Anal. Appl.* 26 (1969) 318–344.
- [24] Henry C. Wente, Large solutions to the volume constrained Plateau problem, *Arch. Ration. Mech. Anal.* 75 (1) (1980/81) 59–77.
- [25] Ye Rugang, Foliation by constant mean curvature spheres, *Pacific J. Math.* 147 (2) (1991) 381–396.