

Asymptotic stability of solitary waves in generalized Gross–Neveu model

Andrew Comech^{a,b,*}, Tuoc Van Phan^{c,1}, Atanas Stefanov^{d,2}

^a Texas A&M University, College Station, TX 77843, USA

^b IITP, Moscow 101447, Russia

^c Department of Mathematics, University of Tennessee, Knoxville, TN 37996, USA

^d Department of Mathematics, University of Kansas, Lawrence, KS 66045, USA

Received 5 July 2014; received in revised form 28 April 2015; accepted 4 November 2015

Available online 2 December 2015

Abstract

For the nonlinear Dirac equation in $(1 + 1)D$ with scalar self-interaction (Gross–Neveu model), with quintic and higher order nonlinearities (and within certain range of the parameters), we prove that solitary wave solutions are asymptotically stable in the “even” subspace of perturbations (to ignore translations and eigenvalues $\pm 2\omega i$). The asymptotic stability is proved for initial data in H^1 . The approach is based on the spectral information about the linearization at solitary waves which we justify by numerical simulations. For the proof, we develop the spectral theory for the linearized operators and obtain appropriate estimates in mixed Lebesgue spaces, with and without weights.

© 2015 L'Association Publications de l'Institut Henri Poincaré. Published by Elsevier B.V. All rights reserved.

Keywords: Gross–Neveu model; Nonlinear Dirac equation; Solitary waves; Asymptotic stability; Weighted spaces

1. Introduction

Models of self-interacting spinor fields have been appearing in particle physics for many years [19,16,15,17]. The most common examples of nonlinear Dirac equation are the massive Thirring model [35] (vector self-interaction) and the Soler model [30] (scalar self-interaction). The $(1 + 1)D$ analogue of the latter model is widely known as the massive Gross–Neveu model [23]. In the present paper, we address the asymptotic stability of solitary waves in this model. We require that the nonlinearity in the equation vanishes of order at least five; the common case of cubic nonlinearity seems out of reach with the current technology; there is a similar situation with other popular dispersive models in one spatial dimension, such as the Schrödinger and Klein–Gordon equations (see [6,7,24,13,20] and the references therein).

* Corresponding author.

E-mail address: comech@math.tamu.edu (A. Comech).

¹ T.V. Phan is supported by Simons Foundation, grant number # 354889.

² A. Stefanov is supported in part by NSF-DMS under contract # 1313107.

We only consider perturbations in the class of “even” spinors (same parity as the solitary waves under consideration). The restriction to this subspace allows us to ignore spatial translations and the $\pm 2\omega i$ eigenvalues which are present in the spectrum of the linearization at solitary waves [12]. This paper therefore may be considered as the extension of [29] to the translation-invariant systems (in that paper, the potential was needed to obtain the desired spectrum of linearization at small amplitude solitary waves).

A similar result – asymptotic stability of solitary waves in the translation-invariant nonlinear Dirac equation in three spatial dimensions – is obtained in [3]. Authors base their highly technical approach on a series of assumptions about the spectrum of the linearizations at solitary waves; these assumptions cannot be verified yet for a particular model. The authors also restrict the perturbations to a certain subspace to avoid spatial translations and issues caused by the presence of $\pm 2\omega i$ eigenvalues [12] and only consider the solitary waves with $\omega > m/3$. Contrary to [3], our results are obtained for models for which the spectrum is known (albeit numerically); our technical restriction is $|\omega| < m/3$.

We briefly review the related research on stability of solitary waves in nonlinear Dirac equation. There have been numerous approaches to this question based on considering the energy minimization at particular families of perturbations, but the scientific relevance of these conclusions has never been justified; see the review and references in e.g. [2,31]. The linear (spectral) stability of the nonlinear Dirac equation is still being settled. According to [2,11], one expects that the linear stability properties of solitary waves in the nonrelativistic limit of the nonlinear Dirac equation (solitary waves with $\omega \lesssim m$) are similar to linear stability of nonlinear Schrödinger equation; in particular, the stability of the *ground states* (no-node solutions) is described by the Vakhitov–Kolokolov stability criterion [36], $\partial_\omega Q(\omega) < 0$, with $Q(\omega) = \|\phi_\omega\|_{L^2}^2$ the charge of a solitary wave. Away from the nonrelativistic limit, the border of the instability region can be indicated by the conditions $\partial_\omega Q(\omega) = 0$ or $E(\omega) = 0$ (the value of the energy functional at a solitary wave), see [4]. The instability could also develop from the bifurcation of the quadruple of complex eigenvalues from the embedded thresholds $\pm i(m + |\omega|)$ as in [8], which in particular can take place at the collision of thresholds at $\lambda = \pm im$ when $\omega = 0$ as in [22]. We do not have a good criterion when such bifurcation takes place.

Let us mention that our results are at odds with the numerical simulations in [31] which are interpreted as instability of the cubic Gross–Neveu model ($k = 1$) for $\omega \leq \omega_c \approx 0.56$, of the quintic model ($k = 2$) for $\omega \leq \omega_c \approx 0.92$, and of the $k = 3$ model for all $\omega < m$. We expect that the observed instability is related to the boundary effects, when certain harmonics, instead of being dispersed, are reflected into the bulk of the solution, where the nonlinearity creates higher harmonics; this process keeps repeating, and eventually the space–time discretization becomes insufficient. This explanation is corroborated by the fact that the characteristic instability times *grow almost proportionally with the size of the domain* (see the instability times for the one-humped solitary wave with $k = 1$, $\omega = 0.5$ in [31, Table II]), suggesting the link not to the linear instability but to the boundary contribution. Our numerics show no complex eigenvalues away from the union of real and imaginary axes in the Gross–Neveu model with $1 \leq k \leq 9$. The presence of real eigenvalues (as on Fig. 2) agrees with the Vakhitov–Kolokolov stability criterion, $dQ(\omega)/d\omega > 0$.

The approach in our paper is standard, being based on modulation equations, dispersive wave decay estimates, and the Strichartz inequalities. Instead of explaining our approach, we provide a detailed outline of the paper, which will elucidate the main steps and ideas involved in the proof. In Section 2, we state the Gross–Neveu model, describe the standing wave solutions and their properties, and formulate our main result on the asymptotic stability. We provide numerics which suggest that, at least for certain range of the parameters, we have a favorable for us spectral picture: that is, *the absence of unstable spectrum, as well as the absence of marginally stable point spectrum, except at zero*. In Section 3, we obtain the form of the linearized operator around the solitary wave for the corresponding nonlinear evolution and obtain the modulation equations. Section 4 is the most challenging from a technical point of view. Therein, we develop the spectral theory for the linearized operator. We use the four linearly independent Jost solutions to construct the resolvent explicitly. This allows us to obtain (among other things) a limiting absorption principle for the linearized operator (Proposition 4.14), which is crucial for the types of estimates required to establish asymptotic stability. (Let us mention a related result [21] on local energy decay for the Dirac equation on one dimension, which we will also need.) In Section 5, we use the spectral theory developed in the previous section to establish various dispersive estimates for the linearized Dirac evolution semigroup. Namely, we establish weighted decay estimates, which in turn imply Strichartz estimates. We also state and prove estimates between Strichartz spaces and weighted $L_t^\infty L^2$ spaces – in all this, we have been greatly helped by the Christ–Kiselev lemma and Born expansions. In Section 6, we present the fixed point argument in the appropriate spaces, which finally shows well-posedness for small data for the equation of the residuals.

2. The model and the main results

2.1. Generalized Gross–Neveu model

The generalized Soler model (classical fermionic field with scalar self-interaction) corresponds to the Lagrangian density

$$\mathcal{L} = \bar{\psi}(i\gamma^\mu \partial_\mu - m)\psi + F(\bar{\psi}\psi), \quad \psi(x, t) \in \mathbb{C}^N, \quad x \in \mathbb{R}^n, \tag{2.1}$$

where $F \in C^\infty(\mathbb{R})$, $F(0) = 0$, $\bar{\psi}$ is a common notation from the Quantum Field Theory,

$$\bar{\psi} = \psi^* \gamma^0, \tag{2.2}$$

with ψ^* being the Hermitian conjugate, and γ^μ , $0 \leq \mu \leq n$, are the Dirac gamma-matrices:

$$\gamma^\mu \gamma^\nu + \gamma^\nu \gamma^\mu = 2h^{\mu\nu} I_n, \quad 0 \leq \mu, \nu \leq n,$$

with $h^{\mu\nu} = \text{diag}[1, -1, \dots, -1]$ (the inverse of) the Minkowski metric tensor and I_n the identity matrix. The one-dimensional analogue of (2.1) is called the Gross–Neveu model; from now on, we set

$$n = 1, \quad N = 2.$$

The equation of motion corresponding to the Lagrangian (2.1) is then given by the following nonlinear Dirac equation:

$$i\partial_t \psi = D_m \psi - f(\psi^* \beta \psi) \beta \psi, \quad \psi(x, t) \in \mathbb{C}^2, \quad x \in \mathbb{R}, \tag{2.3}$$

where $f = F' \in C^\infty(\mathbb{R})$, $\alpha = \gamma^0 \gamma^1$, $\beta = \gamma^1$, and $D_m = -i\alpha \frac{\partial}{\partial x} + \beta m$ is the Dirac operator, with α , β the self-adjoint Dirac matrices satisfying

$$\alpha^2 = \beta^2 = I_2, \quad \alpha\beta + \beta\alpha = 0.$$

A particular choice of the Dirac matrices is irrelevant (this is known as the Dirac–Pauli theorem, see e.g. [34, Lemma 2.25]). For definiteness, we take

$$\alpha = -\sigma_2 = \begin{bmatrix} 0 & i \\ -i & 0 \end{bmatrix}, \quad \beta = \sigma_3 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}.$$

Without loss of generality, we will also assume that the mass is equal to $m = 1$. Then one has

$$D_m = -i\alpha \frac{\partial}{\partial x} + m\beta = \begin{bmatrix} 1 & \partial_x \\ -\partial_x & -1 \end{bmatrix}. \tag{2.4}$$

The Hamiltonian density derived from the Lagrangian density (2.1) is given by

$$\mathcal{E}(\psi, \dot{\psi}) = \frac{\partial \mathcal{L}}{\partial \dot{\psi}} \dot{\psi} - \mathcal{L}. \tag{2.5}$$

The value of the energy functional

$$E(\psi) = \int_{\mathbb{R}} \mathcal{E}(\psi, \dot{\psi}) dx \tag{2.6}$$

is (formally) conserved for the solutions to (2.3). Due to the U(1)-invariance of the Lagrangian (2.1), the total charge of the solutions to (2.3),

$$Q(\psi) = \int_{\mathbb{R}} \psi^*(x, t) \psi(x, t) dx, \tag{2.7}$$

is also (formally) conserved.

We mention recent results on local and global well-posedness of nonlinear Dirac equation on one dimension: [33, 25,9,28,18].

2.2. Existence and properties of solitary waves

Equation (2.3) can be written explicitly as

$$\begin{cases} i\partial_t\psi_1 = \partial_x\psi_2 + \psi_1 - f(|\psi_1|^2 - |\psi_2|^2)\psi_1, \\ i\partial_t\psi_2 = -\partial_x\psi_1 - \psi_2 + f(|\psi_1|^2 - |\psi_2|^2)\psi_2. \end{cases} \tag{2.8}$$

In the abstract form, we write (2.3) as

$$i\partial_t\psi = D_m\psi + \mathbf{N}(\psi), \tag{2.9}$$

with the Dirac operator

$$D_m = -i\alpha\partial_x + \beta = \begin{bmatrix} 1 & \partial_x \\ -\partial_x & -1 \end{bmatrix}$$

and the nonlinearity

$$\mathbf{N}(\psi) = \begin{bmatrix} -f(|\psi_1|^2 - |\psi_2|^2) & 0 \\ 0 & f(|\psi_1|^2 - |\psi_2|^2) \end{bmatrix} \psi. \tag{2.10}$$

Definition 2.1. Solitary waves are solutions of the form

$$\psi_\omega(x, t) = \phi_\omega(x)e^{-i\omega t}, \quad \phi_\omega \in H^1(\mathbb{R}, \mathbb{C}^2), \quad \omega \in \mathbb{R}. \tag{2.11}$$

Substituting this Ansatz into (2.9), we see that ϕ_ω solves

$$\omega\phi_\omega = D_m\phi_\omega + \mathbf{N}(\phi_\omega). \tag{2.12}$$

The existence of solitary waves follows from [14,1]:

Proposition 2.2. Let F be the antiderivative of $f \in C^\infty(\mathbb{R})$ such that $F(0) = 0$. Assume that for given $\omega \in \mathbb{R}$, $0 < \omega < 1$, there exists $\Gamma_\omega > 0$ such that

$$\omega\Gamma_\omega = \Gamma_\omega - F(\Gamma_\omega), \quad \omega \neq 1 - f(\Gamma_\omega), \quad \omega s < s - F(s), \quad \text{for } s \in (0, \Gamma_\omega).$$

Then there is a solitary wave solution $\psi_\omega(x, t) = \phi_\omega(x)e^{-i\omega t}$ to (2.3), with

$$\phi_\omega(x) = \begin{bmatrix} v(x, \omega) \\ u(x, \omega) \end{bmatrix}, \quad v(\cdot, \omega), u(\cdot, \omega) \in H^1(\mathbb{R}). \tag{2.13}$$

This solution is unique if we require that v, u are real-valued, v even and positive, and u odd. Both v and u are exponentially decaying as $|x| \rightarrow \infty$ and satisfy $|u(x, \omega)| < |v(x, \omega)|$, $x \in \mathbb{R}$.

Moreover, there is $c_\omega < \infty$ such that

$$|\phi_\omega(x)| \leq c_\omega e^{-\delta_\omega|x|}, \quad x \in \mathbb{R}, \tag{2.14}$$

where

$$\delta_\omega = \sqrt{1 - \omega^2}. \tag{2.15}$$

Similarly, there is $c_\omega < \infty$ such that

$$|\partial_\omega\phi_\omega(x)| \leq c_\omega(x)e^{-\delta_\omega|x|}, \quad |\partial_\omega^2\phi_\omega(x)| \leq c_\omega(x)^2e^{-\delta_\omega|x|}, \quad x \in \mathbb{R}, \tag{2.16}$$

$$|\partial_x\partial_\omega\phi_\omega(x)| \leq c_\omega(x)e^{-\delta_\omega|x|}, \quad |\partial_x\partial_\omega^2\phi_\omega(x)| \leq c_\omega(x)^2e^{-\delta_\omega|x|}, \quad x \in \mathbb{R}. \tag{2.17}$$

Proof. The proof is given in e.g. [1, Lemma 3.2]. The sharp rate of decay (2.14) can be proved as in e.g. [10, Appendix A]. The bounds (2.16) on $|\partial_\omega\phi_\omega(x)|$ and $|\partial_\omega^2\phi_\omega(x)|$ follow from differentiating (2.12) with respect to ω and applying similar techniques. Applying ∂_ω -derivatives to (2.12) and using the bounds (2.16), one obtains (2.17). \square

Remark 2.3. According to Proposition 2.2, for $f(s) = \alpha s^k$ with any $\alpha > 0$ and $k \in \mathbb{N}$, there are solitary wave solutions for $\omega \in (0, 1)$.

2.3. Main result

Let

$$X = \{ \phi \in L^2(\mathbb{R}, \mathbb{C}^2); \quad \phi_1(x) = \phi_1(-x), \quad \phi_2(x) = -\phi_2(-x) \}. \tag{2.18}$$

Assumption 2.4. Assume that $f \in C^\infty(\mathbb{R})$ is such that $f(s) = \mathcal{O}(s^k)$ for $|s| \leq 1$, with $k \geq 2$, and that there is an open interval Ω ,

$$\Omega \subset \left(-\frac{1}{3}, \frac{1}{3} \right),$$

such that the following takes place:

- (i) For each $\omega \in \Omega$, there are solitary wave solutions $\psi_\omega(x, t) = \phi_\omega(x)e^{-i\omega t}$, $\phi_\omega \in H^1(\mathbb{R}, \mathbb{C}^2)$, to (2.3), with the map $\Omega \rightarrow H^1$, $\omega \mapsto \phi_\omega$ being C^2 .
- (ii) Non-degeneracy:

$$\partial_\omega Q(\omega) \neq 0, \quad \omega \in \Omega.$$

Here $Q(\omega)$ is the value of the charge functional (2.7) evaluated at the solitary wave $\phi_\omega(x)e^{-i\omega t}$.

- (iii) The linearization of (2.3) at a solitary wave with $\omega \in \Omega$ has no eigenvalues with nonzero real part and no purely imaginary eigenvalues $\lambda \in i\mathbb{R}$ with eigenfunctions from X (of the same parity as ϕ_ω), and no resonances at $\lambda = 1 \pm |\omega|$ with generalized L^∞ eigenfunctions of the same parity as ϕ_ω .
- (iv) For $\omega \in \Omega$, the Evans function $E(\lambda, \omega)$ (cf. Definition 4.5) does not vanish at $\lambda \in i\mathbb{R}$ with $|\lambda| \geq 1 - |\omega|$.

Above, by the generalized L^∞ eigenfunction of a differential operator A we mean the nontrivial solution to $(A - \lambda)u = 0$ which has finite L^∞ -norm. Also, the Evans function is defined Definition 4.5 below.

The following theorem is the main result of our paper.

Theorem 2.5 (Asymptotic stability of solitary waves in nonlinear Dirac equation). *Let Assumption 2.4 hold. Let $\omega_0 \in \Omega$ and $\phi_{\omega_0}(x)e^{-i\omega_0 t}$ be the corresponding solitary wave with $\phi_{\omega_0} \in X \cap H^1(\mathbb{R}, \mathbb{C}^2)$. There exist $\epsilon_0 > 0$ and $C < \infty$ such that if $\psi_0 \in X$ satisfies*

$$\inf_{\gamma \in [0, 2\pi]} \left\| \psi_0 - e^{i\gamma} \phi_{\omega_0} \right\|_{H^1} \leq \epsilon^2, \quad \text{with some } \epsilon \in (0, \epsilon_0),$$

then the solution ψ of (2.3) with $\psi|_{t=0} = \psi_0$ exists globally in time and there are

$$\omega, \gamma \in C^1(\mathbb{R}_+, \mathbb{R}), \quad \varphi \in X \cap L^\infty(\mathbb{R}_+, H^1(\mathbb{R}, \mathbb{C}^2)) \cap L^4(\mathbb{R}_+, L^\infty(\mathbb{R}, \mathbb{C}^2))$$

with $\omega(t) \xrightarrow{t \rightarrow \infty} \omega_\infty \in \Omega$ and $\lim_{t \rightarrow \infty} \|\varphi\|_{L_x^\infty} = 0$, such that

$$\psi(x, t) = (\phi_{\omega(t)}(x) + \varphi(x, t))e^{-i \int_0^t \omega(s) ds - i\gamma(t)}$$

and

$$|\omega_0 - \omega_\infty| \leq \epsilon, \quad \|\dot{\gamma}\|_{L^1(0, \infty)} + \|\dot{\gamma}\|_{L^1(0, \infty)} \leq \epsilon, \quad \|\varphi\|_{L_t^\infty H_x^1} + \|\varphi\|_{L_t^4 L_x^\infty} \leq C\epsilon.$$

Several remarks are in order.

Remark 2.6. The precise structure of the nonlinearity of the Gross–Neveu model, $f(\psi^* \beta \psi)\beta \psi$, does not play any particular role in our considerations. Yet, we choose to base our consideration on this model: being a relativistically invariant nonlinear Dirac equation with minimal coupling, it is a ubiquitous model in Physics.

Remark 2.7. The solitary waves to classical Gross–Neveu model ($k = 1$, cubic nonlinearity) are known to be linearly stable [1], but our argument does not apply to this situation. The assumption $k \geq 2$ allows us to close the argument

in Section 5.2 using the Strichartz estimates, making the argument sufficiently compact. Similar requirements on the order of vanishing of the nonlinearity being sufficiently high are common in the research on asymptotic stability of solitons in nonlinear Schrödinger equation, starting with the seminal papers [6,7].

Remark 2.8. In the Assumption 2.4, we require that $\Omega \subset (-1/3, 1/3)$ to avoid the situation when the eigenvalues $\pm 2\omega i$ become embedded into the essential spectrum ($\lambda \in i\mathbb{R}$, $|\lambda| \geq 1 - |\omega|$). In that case, our construction of the resolvent in Section 4.2 does not allow us to obtain the necessary estimates (see Remark 3.4 and estimate (4.17) below). Yet, the restriction to $|\omega| < 1/3$ seems to be merely technical; we still expect that for $1/3 \leq |\omega| < 1$, the resolvent of the linearized operator restricted to X has the same properties as stated in Proposition 4.14 even in the vicinity of the embedded eigenvalues $\pm 2\omega i$ and that the asymptotic stability could be proved.

Remark 2.9. By [12,4], the assumptions $E(\omega) \neq 0$ and $\partial_\omega Q(\omega) \neq 0$ guarantee that the generalized null space of the linearization at a solitary wave is (exactly) four-dimensional. (Above, $E(\omega)$ and $Q(\omega)$ are the values of the energy and charge functionals (2.6), (2.7) at the solitary wave $\phi_\omega e^{-i\omega t}$.) We do not need to impose the condition $E(\omega) \neq 0$ since although the vanishing of $E(\omega)$ leads to the increase of the Jordan block of the linearization at a solitary wave, this increase is absent when we restrict the operator to the subspace X . More details are in Section 3.1 (see in particular Remark 3.4).

Remark 2.10. The Vakhitov–Kolokolov stability condition [36] states that the ground state solution to the NLS ($\phi(x)e^{-i\omega t}$ with $\phi > 0$ and monotonically decreasing as $|x| \rightarrow \infty$) are linearly stable (positive eigenvalues of the linearization at a solitary wave are absent) as long as $\partial_\omega Q(\omega) < 0$. In the NLD context, it could be shown that a similar conclusion can be drawn in the nonrelativistic limit, $\omega \lesssim m$. In the L^2 -critical case, while the charge in the NLS does not depend on ω , this is “resolved” in the NLD: one now has $\partial_\omega Q(\omega) < 0$ and consequently the eigenvalue $\lambda = 0$ does not have additional degeneracy; instead, there are point eigenvalues in the spectrum (see Fig. 1). For $k > 2$, the positive eigenvalues of the linearized operator are only present in the spectrum for ω near m but disappear below certain value of ω . This is accompanied by the change in sign of $\partial_\omega Q(\omega)$ (see Fig. 2).

Remark 2.11. We numerically verified Assumption 2.4-(iii) in the following cases:

- (i) For the Gross–Neveu model with $f(s) = s^2$ and $\Omega = (0.23, 0.33)$ (see Fig. 1);
- (ii) For the Gross–Neveu model with $f(s) = s^3$ and $\Omega = (0.14, 0.33)$ (see Fig. 2).

We also mention that in the Gross–Neveu model with $k = 1, 2, \dots, 9$, we found no complex eigenvalues for the linearizations at solitary waves with $\omega = 0.1, 0.2, \dots, 0.9$ in the domain $0.0008 < |\operatorname{Re} \lambda| < 0.59$, $|\operatorname{Im} \lambda| < 2.5$. Moreover, according to [2], the bifurcations of point eigenvalues off the imaginary axis could result only from the collision of purely imaginary eigenvalues or from eigenvalues embedded into the continuous spectrum, and also from resonances at the embedded thresholds, $\lambda = \pm i(m + |\omega|)$ (in one-dimensional case, the resonances correspond to the generalized L^∞ eigenfunctions). Our numerics show that there are no resonances at the embedded thresholds in the Gross–Neveu model with $k = 2$ and $k = 3$ for all $\omega \in (0, m)$, justifying the observed absence of complex eigenvalues away from $\mathbb{R} \cup i\mathbb{R}$.

We expect that the Evans function never has zeros at $\lambda \in i\mathbb{R}$, $|\operatorname{Im} \lambda| \geq 1 + |\omega|$, but could not prove this. Instead, we check this assumption numerically; all the zeros of the Evans function which we found are plotted as solid curves on Figs. 1 and 2 (these zeros correspond to the point eigenvalues of the linearized operator). The absence of zeros of the Evans function for $\lambda \rightarrow \pm i\infty$ follows from Lemma 4.10.

Let us summarize that most of our assumptions are technical; the only essential assumption is that the spectrum of the linearized operator has no eigenvalues in the right half-plane and that the Jordan block of $\lambda = 0$ is (exactly) four-dimensional. We expect that the presence of purely imaginary eigenvalues does not lead to instability unless these eigenvalues are of higher algebraic multiplicity. More generally, we expect that, similarly to the case of the nonlinear Schrödinger equation and similar systems, the (dynamic) instability takes place when either there is a linear instability or when the eigenvalues on the imaginary axis are of higher algebraic multiplicities (when we are at the threshold of




Fig. 1. Gross–Neveu model, $k = 2$ (the quintic case). Linearization at a solitary wave. Horizontal axis: $\omega \in (0, 1)$. Vertical axis: spectrum on the upper half of the imaginary axis. Solid vertical (green) lines: part of the continuous spectrum between the threshold $i(1 - |\omega|)$ and the embedded threshold $i(1 + |\omega|)$. Solid red curves: eigenvalues with eigenfunctions from \mathbf{X} (of the same parity as ϕ_ω ; see (3.16)), which we cannot ignore; our result holds in the regions where such eigenvalues are absent. Solid blue curve (near $\omega = 0$ and $\lambda = i$) and the line $\lambda = 2\omega i$ denote eigenvalues with eigenfunctions from \mathbf{X}^\perp (see (3.17)), which remain orthogonal to our perturbation. Dotted red and blue curves: antibound states of different parity (from \mathbf{X} and \mathbf{X}^\perp); we do not mention them in the argument. Antibound states correspond to zeros of Evans functions on the “wrong” Riemann sheet, which corresponds to generalized eigenfunctions with exponential growth at infinity. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)




Fig. 2. Gross–Neveu model, $k = 3$. Hollow red diamonds (on bottom right) denote positive eigenvalues (thus linear instability) present in the spectrum for $\omega \in (0.85, 1)$. These eigenvalues are superimposed on the imaginary axis. **Theorem 2.5** on asymptotic stability applies for solitary waves with ω such that there are neither hollow red diamonds (linear instability) nor solid red curves (purely imaginary eigenvalues with eigenfunctions from \mathbf{X}) in the spectrum. Note that the dotted kink indicates collision of antibound states at $\omega_b \approx 0.1$ on the imaginary axis and their bifurcation off the imaginary axis for $\omega < \omega_b$. (Location of these values of λ off the imaginary axis does not lead to instability since the corresponding antibound states have infinite L^2 -norm.) (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

linear instability). It should be pointed out, though, that the presence of additional point eigenvalues in the spectral gap of the linearization at a solitary wave is very likely to considerably increase the difficulty of the analysis, just like in the case of the NLS.

3. Linearization at a solitary wave and modulation equations

3.1. Linearization at a solitary wave

To study stability of a solitary wave $\phi_\omega(x)e^{-i\omega t}$, with $\phi_\omega(x) = \begin{bmatrix} v(x,\omega) \\ u(x,\omega) \end{bmatrix} \in \mathbb{R}^2$, we consider the solution in the form

$$\psi(x, t) = (\phi_\omega(x) + \rho(x, t))e^{-i\omega t}, \quad \rho(x, t) \in \mathbb{C}^2.$$

Substituting this Ansatz into (2.9), we obtain:

$$i\partial_t \rho = (D_m - \omega I_2)\rho + \mathbf{N}(\phi_\omega + \rho) - \mathbf{N}(\phi_\omega). \quad (3.1)$$

Thus, the linearization at a solitary wave (the linearized equation on ρ derived from (3.1)) can be written as follows:

$$\dot{R} = \mathbf{J}\mathbf{L}R, \quad R = \begin{bmatrix} \operatorname{Re} \rho \\ \operatorname{Im} \rho \end{bmatrix} \in \mathbb{R}^4, \quad (3.2)$$

where

$$\mathbf{J} = \begin{bmatrix} 0 & I_2 \\ -I_2 & 0 \end{bmatrix}, \quad \mathbf{L}(\omega) = \mathbf{D}_m - \omega I_4 + \mathbf{W}(x, \omega), \quad (3.3)$$

with

$$\mathbf{W}(x, \omega) = \begin{bmatrix} W_1(x, \omega) & 0 \\ 0 & W_0(x, \omega) \end{bmatrix}, \quad W_0(x, \omega) = \begin{bmatrix} -f(v^2 - u^2) & 0 \\ 0 & f(v^2 - u^2) \end{bmatrix}, \quad W_1(x, \omega) = W_0(x, \omega) - 2f'(v^2 - u^2) \begin{bmatrix} v^2 & -vu \\ -vu & u^2 \end{bmatrix}. \quad (3.4)$$

The free Dirac operator takes the form

$$\mathbf{D}_m = \mathbf{J}\alpha\partial_x + \beta, \quad (3.5)$$

with

$$\alpha = \begin{bmatrix} \operatorname{Re} \alpha & -\operatorname{Im} \alpha \\ \operatorname{Im} \alpha & \operatorname{Re} \alpha \end{bmatrix} = \begin{bmatrix} 0 & \operatorname{Im} \sigma_2 \\ -\operatorname{Im} \sigma_2 & 0 \end{bmatrix}, \quad \beta = \begin{bmatrix} \operatorname{Re} \beta & -\operatorname{Im} \beta \\ \operatorname{Im} \beta & \operatorname{Re} \beta \end{bmatrix} = \begin{bmatrix} \sigma_3 & 0 \\ 0 & \sigma_3 \end{bmatrix}; \quad (3.6)$$

\mathbf{J} , α , and β represent $-i$, α , and β when acting on $\begin{bmatrix} \operatorname{Re} \psi \\ \operatorname{Im} \psi \end{bmatrix}$, with $\psi \in \mathbb{C}^2$. We then have

$$\mathbf{D}_m = \begin{bmatrix} D_m & 0 \\ 0 & D_m \end{bmatrix}, \quad \text{where } D_m = \begin{bmatrix} 1 & \partial_x \\ -\partial_x & -1 \end{bmatrix}. \quad (3.7)$$

Note that since v , u both depend on ω , the potentials W_1 , W_0 also depend on it. We will often omit this dependence in our notations.

Lemma 3.1. *There is $C_\omega < \infty$ such that the matrix-valued potential \mathbf{W} satisfies*

$$\|\mathbf{W}(x, \omega)\|_{\mathbb{C}^4 \rightarrow \mathbb{C}^4} \leq C_\omega e^{-2k|x|\delta_\omega}, \quad x \in \mathbb{R}. \quad (3.8)$$

Proof. This bound is an immediate consequence of the exponential decay of ϕ_ω in Proposition 2.2 (see (2.14)), the assumption $f(s) = \mathcal{O}(s^k)$ for $|s| \leq 1$, and (3.4). \square

Lemma 3.2.

$$\sigma_{\text{ess}}(\mathbf{J}\mathbf{L}) = i\mathbb{R} \setminus (-i(1 - |\omega|), i(1 - |\omega|)).$$

Proof. This is an immediate consequence of Weyl’s theorem on the essential spectrum. \square

Denote

$$\phi(x) = \phi_\omega(x) = \begin{bmatrix} \operatorname{Re} \phi_\omega(x) \\ \operatorname{Im} \phi_\omega(x) \end{bmatrix} = \begin{bmatrix} \phi_\omega(x) \\ 0 \end{bmatrix}. \tag{3.9}$$

Thanks to the invariance of (2.12) with respect to the phase rotation and the translation, we have

$$\mathbf{JL}\mathbf{J}\phi = 0, \quad \mathbf{JL}\partial_x\phi = 0.$$

The null space of \mathbf{L} is given by

$$N(\mathbf{L}) = (\mathbf{J}\phi, \partial_x\phi). \tag{3.10}$$

Remark 3.3. This could be readily justified by analyzing the Jost solutions of

$$L_1(\omega) = D_m - \omega I_2 + W_1, \quad L_0(\omega) = D_m - \omega I_2 + W_0. \tag{3.11}$$

Namely, for each of L_1 and L_0 , there are two Jost solutions corresponding to $\lambda = 0$: one decreasing and one increasing. More details on the Jost solutions are in Section 4.1.

Moreover,

$$\mathbf{JL}\partial_\omega\phi = \mathbf{J}\phi, \tag{3.12}$$

$$\mathbf{JL}\left(\omega x\mathbf{J}\phi - \frac{1}{2}\alpha\phi\right) = \partial_x\phi, \tag{3.13}$$

where

$$\omega x\mathbf{J}\phi - \frac{1}{2}\alpha\phi = \begin{bmatrix} 0 \\ i\frac{1}{2}\alpha\phi - \omega x\phi \end{bmatrix} = \begin{bmatrix} 0 \\ -i\frac{1}{2}\sigma_2\phi - \omega x\phi \end{bmatrix}.$$

Therefore,

$$\left\{ \mathbf{J}\phi, \partial_x\phi, \partial_\omega\phi, \omega x\mathbf{J}\phi - \frac{1}{2}\alpha\phi \right\} \subset N_g(\mathbf{JL}). \tag{3.14}$$

By [4], if $\partial_\omega Q(\omega) \neq 0$ and $E(\omega) \neq 0$, then the above vectors form a basis in the generalized null space $N_g(\mathbf{JL})$:

$$N_g(\mathbf{JL}) = \operatorname{Span}\left(\mathbf{J}\phi, \partial_x\phi, \partial_\omega\phi, \omega x\mathbf{J}\phi - \frac{1}{2}\alpha\phi\right). \tag{3.15}$$

Following the definition (2.18), we define

$$\mathbf{X} = \left\{ \psi \in L^2(\mathbb{R}, \mathbb{C}^4); \psi_k(x) = \psi_k(-x), \quad k = 1, 3; \quad \psi_k(x) = -\psi_k(-x), \quad k = 2, 4 \right\}; \tag{3.16}$$

$$\mathbf{X}^\perp = \left\{ \psi \in L^2(\mathbb{R}, \mathbb{C}^4); \psi_k(x) = \psi_k(-x), \quad k = 2, 4; \quad \psi_k(x) = -\psi_k(-x), \quad k = 1, 3 \right\}. \tag{3.17}$$

From now on, we shall restrict $\mathbf{JL}(\omega)$ to \mathbf{X} . Noting that $\mathbf{J}\phi$ and $\partial_\omega\phi$ (as well as ϕ) belong to the space \mathbf{X} , while $\partial_x\phi$, $x\mathbf{J}\phi$, and $\alpha\phi$ belong to \mathbf{X}^\perp , we conclude that

$$N(\mathbf{JL}|_{\mathbf{X}}) = \operatorname{Span}(\mathbf{J}\phi), \quad N_g(\mathbf{JL}|_{\mathbf{X}}) = \operatorname{Span}(\mathbf{J}\phi, \partial_\omega\phi). \tag{3.18}$$

The operator \mathbf{JL} acts invariantly in \mathbf{X} and in \mathbf{X}^\perp .

Remark 3.4. The restriction of $\mathbf{JL}(\omega)$ onto \mathbf{X} allows one to exclude certain eigenvalue directions, significantly simplifying the problem. In particular, by [12], one has

$$\mathbf{JL} \begin{bmatrix} \sigma_1\phi \\ i\sigma_1\phi \end{bmatrix} = 2i\omega \begin{bmatrix} \sigma_1\phi \\ i\sigma_1\phi \end{bmatrix}, \quad \mathbf{JL} \begin{bmatrix} \sigma_1\phi \\ -i\sigma_1\phi \end{bmatrix} = -2i\omega \begin{bmatrix} \sigma_1\phi \\ -i\sigma_1\phi \end{bmatrix}, \tag{3.19}$$

where σ_1 is the Pauli matrix; this shows that $\pm 2\omega i \in \sigma_p(\mathbf{JL}(\omega))$. On the other hand, the restriction of \mathbf{JL} to \mathbf{X} satisfies $\pm 2\omega i \notin \sigma_d(\mathbf{JL}|_{\mathbf{X}})$.

Let us also mention that while the dimension of the generalized space $N_g(\mathbf{JL})$ would grow if either $\partial_\omega Q(\omega)$ or $E(\omega)$ vanish [4], the dimension of $N_g(\mathbf{JL}|_{\mathbf{X}})$ would grow only if $\partial_\omega Q(\omega)$ vanishes; we no longer have to worry whether $E(\omega)$ vanishes or not.

Since $(\mathbf{JL})^* = -\mathbf{LJ}$, it follows from (3.12), (3.13) that the corresponding generalized kernel for the adjoint is

$$\mathbf{X}_g((\mathbf{JL})^*) = N_g((\mathbf{JL})^*) \cap \mathbf{X} = \{ \mathbf{J}\partial_\omega\phi, \phi \}.$$

We decompose the space \mathbf{X} as follows:

$$\mathbf{X} = \mathbf{X}_g(\mathbf{JL}) \oplus \mathbf{X}_c(\mathbf{JL}), \quad \text{where} \quad \mathbf{X}_c(\mathbf{JL}) = \mathbf{X}_g((\mathbf{JL})^*)^\perp. \tag{3.20}$$

The subspaces $\mathbf{X}_g(\mathbf{JL})$ and $\mathbf{X}_c(\mathbf{JL})$ are invariant under the action of \mathbf{JL} , and any $R_1 \in \mathbf{X}_g(\mathbf{JL})$, $R_2 \in \mathbf{X}_c(\mathbf{JL})$ satisfy the following symplectic orthogonality condition:

$$\langle \mathbf{J}R_1, R_2 \rangle = 0.$$

It then follows that any $R \in \mathbf{X}$ can be uniquely decomposed into

$$R = 2 \frac{\langle \phi, R \rangle}{Q'(\omega)} \partial_\omega\phi + 2 \frac{\langle \mathbf{J}\partial_\omega\phi, R \rangle}{Q'(\omega)} \mathbf{J}\phi + U, \quad U \in \mathbf{X}_c(\mathbf{JL}), \tag{3.21}$$

where $Q(\omega)$ is the charge functional (2.7) evaluated at $\phi_\omega e^{-i\omega t}$. Thus, a vector function $U \in \mathbf{X}_c(\mathbf{JL})$ satisfies the following two symplectic orthogonality conditions:

$$\langle \phi, U \rangle = 0, \quad \langle \mathbf{J}\partial_\omega\phi, U \rangle = 0. \tag{3.22}$$

Remark 3.5. Note that $Q'(\omega) \neq 0$ by Assumption 2.4.

Let $P_d(\omega)$ denote the symplectically orthogonal projection onto the generalized null space of $\mathbf{JL}(\omega)$ restricted onto the space \mathbf{X} from (3.16). By (3.21),

$$P_d(\omega)R = 2 \frac{\langle \phi, R \rangle}{Q'(\omega)} \partial_\omega\phi + 2 \frac{\langle \mathbf{J}\partial_\omega\phi, R \rangle}{Q'(\omega)} \mathbf{J}\phi, \tag{3.23}$$

while the projection onto \mathbf{X}_c is

$$P_c(\omega) = 1 - P_d(\omega). \tag{3.24}$$

3.2. Modulation equations

We consider the solution ψ of equation (2.9) in the form

$$\psi(x, t) = (\phi_{\omega(t)}(x) + \rho(x, t))e^{-i\theta(t)}, \quad \text{with} \quad \theta(t) = \int_0^t \omega(s) ds + \gamma(t), \quad x, t \in \mathbb{R}. \tag{3.25}$$

Substituting this Ansatz into (2.9), we get

$$i\partial_t \rho = (D_m - I_2\omega - \dot{\gamma}I_2)\rho - \dot{\gamma}\phi - i\dot{\omega}\partial_\omega\phi + \mathbf{N}(\phi + \rho) - \mathbf{N}(\phi), \tag{3.26}$$

with \mathbf{N} defined in (2.10). As in (3.2), (3.9), we use the notations

$$R = \begin{bmatrix} \text{Re } \rho \\ \text{Im } \rho \end{bmatrix}, \quad \phi_\omega = \begin{bmatrix} \text{Re } \phi_\omega \\ \text{Im } \phi_\omega \end{bmatrix} = \begin{bmatrix} \phi_\omega \\ 0 \end{bmatrix}.$$

Then equation (3.26) takes the form

$$\partial_t R = \mathbf{JL}R - \dot{\gamma}\mathbf{J}R - \dot{\gamma}\mathbf{J}\phi - \dot{\omega}\partial_\omega\phi + \mathbf{JN}_1, \tag{3.27}$$

where

$$\mathbf{N}_1(R, \omega) = \begin{bmatrix} \operatorname{Re}(\mathbf{N}(\phi + \rho) - \mathbf{N}(\phi)) \\ \operatorname{Im}(\mathbf{N}(\phi + \rho) - \mathbf{N}(\phi)) \end{bmatrix} - \mathbf{W}R, \tag{3.28}$$

with \mathbf{W} from (3.4).

Remark 3.6. Let us point out that since we take the initial data of certain parity, $\psi|_{t=0} \in X$, then we also have $\psi \in X$ for all $t \geq 0$, so that $\rho \in X$; therefore, $R \in \mathbf{X}$ and $\mathbf{J}\mathbf{N}_1 \in \mathbf{X}$ (see Definitions 2.18, 3.16). Moreover, the operators $\mathbf{J}\mathbf{L}(\omega)$, $P_d(\omega)$, and $P_c(\omega)$ act invariantly in \mathbf{X} .

We impose the requirement $R(t) \in \mathbf{X}_c(\omega(t))$. Together with the symplectic orthogonality condition (3.22), this requirement implies that

$$\langle \phi, R \rangle = \langle \mathbf{J}\partial_\omega\phi, R \rangle = 0. \tag{3.29}$$

Taking the time derivative of the relations (3.29), we get

$$\langle \phi, \dot{R} \rangle = -\dot{\omega}\langle \partial_\omega\phi, R \rangle = -\dot{\omega}\operatorname{Re}\langle \varphi, \rho \rangle, \quad \langle \mathbf{J}\partial_\omega\phi, \dot{R} \rangle = -\dot{\omega}\langle \mathbf{J}\partial_\omega^2\phi, R \rangle = \dot{\omega}\operatorname{Im}\langle \partial_\omega\varphi, \rho \rangle, \tag{3.30}$$

where

$$\varphi_\omega = \partial_\omega\phi_\omega.$$

Coupling (3.27) with ϕ and with $\mathbf{J}\partial_\omega\phi$ and using the symplectic relations (3.22) and the relations (3.30), we obtain

$$\mathcal{A}(t) \begin{bmatrix} \dot{\omega} \\ \dot{\gamma} \end{bmatrix} = \begin{bmatrix} \langle \phi, \mathbf{J}\mathbf{N}_1 \rangle \\ \langle \mathbf{J}\partial_\omega\phi, \mathbf{N}_1 \rangle \end{bmatrix}, \tag{3.31}$$

where

$$\mathcal{A}(t) = \begin{bmatrix} \langle \phi, \partial_\omega\phi \rangle - \langle \partial_\omega\phi, R \rangle & \langle \phi, \mathbf{J}R \rangle \\ -\langle \mathbf{J}\partial_\omega^2\phi, R \rangle & \langle \mathbf{J}\partial_\omega\phi, \mathbf{J}\phi \rangle + \langle \mathbf{J}\partial_\omega\phi, \mathbf{J}R \rangle \end{bmatrix}, \tag{3.32}$$

where ω and R are evaluated at the moment t .

Define

$$\mu(x) := e^{-\delta_\Omega(x)/(4k)}, \quad \delta_\Omega := \inf_{\omega \in \Omega} \sqrt{1 - \omega^2} > 0; \tag{3.33}$$

by (2.14) and (2.16), there is $C < \infty$ such that for any $\omega \in \Omega$,

$$|\phi_\omega(x)| + |\partial_\omega\phi_\omega(x)| + |\partial_\omega^2\phi_\omega(x)| \leq C\mu(x)^{2k}, \quad x \in \mathbb{R}, \quad \omega \in \Omega. \tag{3.34}$$

Lemma 3.7. *There is $\varepsilon > 0$ such that if $\langle \mu, |R(t)| \rangle < \varepsilon$, then*

$$\|\mathcal{A}(t)^{-1}\| < 2 \left(\inf_{\omega \in \Omega} \langle \phi_\omega, \partial_\omega\phi_\omega \rangle \right)^{-1} < \infty.$$

Proof. From (3.32) and (3.34), we have

$$\mathcal{A}(t) = \begin{bmatrix} \langle \phi_\omega, \partial_\omega\phi_\omega \rangle & 0 \\ 0 & \langle \phi_\omega, \partial_\omega\phi_\omega \rangle \end{bmatrix} + \mathcal{O}(\langle \mu^{2k}, |R| \rangle),$$

where $\omega = \omega(t)$; we took into account the bounds (2.14) and (2.16). By Assumption 2.4, one has $2\langle \phi_\omega, \partial_\omega\phi_\omega \rangle = Q'(\omega)$ with $\inf_{\omega \in \Omega} |Q'(\omega)| > 0$; therefore, one can choose $\varepsilon > 0$ so small that $\mathcal{A}(t)$ is invertible and satisfies the conclusion of the lemma. \square

To control ρ (or equivalently R), let us define

$$Z(t) = P_c(\omega_0)R(t), \tag{3.35}$$

so that

$$Z(0) = P_c(\omega_0)R(0) = P_c(\omega_0) \begin{bmatrix} \operatorname{Re}(\Psi_0 - \phi_0) \\ \operatorname{Im}(\Psi_0 - \phi_0) \end{bmatrix}. \tag{3.36}$$

Since $Z = P_c(\omega_0)R$ and $R = P_c(\omega)R$, and by (3.23), we have

$$Z - R = P_c(\omega_0)R - P_c(\omega)R = (P_d(\omega) - P_d(\omega_0))R = \mathcal{O}(\omega - \omega_0)\langle \mu^{2k}, |R| \rangle \mu^{2k}. \tag{3.37}$$

Therefore, if $|\omega - \omega_0|$ is sufficiently small, to control R , it suffices to control Z ; in particular, it follows from (3.37) that if either Z or R is from H^1 in x , then so is the other function, and moreover

$$\|Z - R\|_{H^1_x} \leq C|\omega - \omega_0|\langle \mu^{2k}, |R| \rangle, \tag{3.38}$$

with some constant $C < \infty$ which depends only on Ω and on the nonlinearity f in (2.3). The weight $\mu(x)^{2k} = e^{-\delta_\Omega(x)/2}$ (cf. (3.33)) comes from the bounds (3.34) on the eigenfunctions that span the generalized null space (3.18) of the operator $\mathbf{JL}(\omega)$ and from the explicit form (3.23) of the projector $P_d(\omega)$.

4. Spectral theory for the linearized operator

In this section, we consider dispersive estimates for the complexification of the linearized equation (3.2),

$$\partial_t R = \mathbf{JL}R, \quad R \in \mathbb{C}^4. \tag{4.1}$$

More precisely, we will show that similarly to the free Dirac evolution, the linear evolution of (4.1) projected onto the continuous spectrum of \mathbf{JL} scatters the initial data. This phenomena in the related Schrödinger equation context manifests itself in a variety of useful estimates; see for example the work of Mizumachi [24].

Before proceeding to specific estimates for the solution of (3.2), let us take a moment to properly define $e^{\mathbf{JL}t} P_c$. Since

$$\sigma_{\text{ess}}(\mathbf{JL}(\omega)) = (-i\infty, -i(1 - |\omega|]) \cup [i(1 - |\omega|), i\infty),$$

we define $e^{\mathbf{JL}(\omega)t} P_c(\omega)$ by the following Cauchy formula:

$$\begin{aligned} e^{\mathbf{JL}t} P_c(\omega)f &= -\frac{1}{2\pi i} \oint_{\Gamma} R_{\mathbf{JL}}(\lambda) f \, d\lambda \\ &= -\frac{1}{2\pi i} \left(\int_{-i\infty}^{-i(1-|\omega|)} + \int_{i(1-|\omega|)}^{i\infty} \right) e^{\lambda t} \left([R_{\mathbf{JL}}^+(\lambda) - R_{\mathbf{JL}}^-(\lambda)] f \right) d\lambda \\ &= -\frac{1}{2\pi} \left(\int_{-\infty}^{|\omega|-1} + \int_{1-|\omega|}^{+\infty} \right) e^{i\Lambda t} \left([R_{\mathbf{JL}}^+(i\Lambda) - R_{\mathbf{JL}}^-(i\Lambda)] f \right) d\Lambda, \end{aligned} \tag{4.2}$$

where Γ is a positively-oriented contour around the essential spectrum of \mathbf{JL} . For $\lambda \in i\mathbb{R}$ the operators

$$R_{\mathbf{JL}}^\pm(\lambda) := \lim_{\varepsilon \rightarrow 0^+} (\mathbf{JL} - (\lambda \pm \varepsilon))^{-1}$$

are to be interpreted in a certain appropriate sense (for example, as operators from $L^2_\alpha \rightarrow L^2_{-\alpha}$, for certain $\alpha > 0$, by the limiting absorption principle).

4.1. The Jost solutions and the Evans function of the operator \mathbf{JL}

The eigenvalue problem for the operator $\mathbf{JL}(\omega)$ (cf. (3.3), (3.4)),

$$\mathbf{J}(\mathbf{D}_m - \omega + \mathbf{W}(x, \omega))\psi = \mathbf{J}(\mathbf{J}\alpha\partial_x + \beta - \omega + \mathbf{W}(x, \omega))\psi = \lambda\psi,$$

can be rewritten as

$$(\partial_x - \alpha\mathbf{J}\beta + \omega\alpha\mathbf{J} + \alpha\lambda - \alpha\mathbf{JW}(x, \omega))\psi = 0. \tag{4.3}$$

The construction of Jost solutions is based on considering solutions to the constant coefficient equation

$$(\partial_x - \mathcal{M}_0(\lambda, \omega))\psi = 0, \quad \mathcal{M}_0(\lambda, \omega) := \alpha\mathbf{J}\beta - \omega\alpha\mathbf{J} - \alpha\lambda, \tag{4.4}$$

and using the Duhamel representation to construct solutions to equation (4.3) with variable coefficients, written in the form

$$(\partial_x - \mathcal{M}(x, \lambda, \omega))\psi = 0, \quad \mathcal{M}(x, \lambda, \omega) := \mathcal{M}_0(\lambda, \omega) + \alpha\mathbf{J}\mathbf{W}(x, \omega). \tag{4.5}$$

Lemma 4.1. *Let $\omega \in [-1, 1]$, $\lambda \in \mathbb{C}$. Then:*

(i)

$$\sigma(\mathcal{M}_0(\lambda, \omega)) = \{\pm\sqrt{1 - (\omega \pm i\lambda)^2}\};$$

(ii)

$$\sup_{\lambda \in i\mathbb{R}} \{|\operatorname{Re} \zeta|; \zeta \in \sigma(\mathcal{M}_0(\lambda, \omega))\} = 1;$$

(iii)

$$\sup_{\lambda \in \sigma_{\text{ess}}(\mathbf{J}\mathbf{L}(\omega))} \{|\operatorname{Re} \zeta|; \zeta \in \sigma(\mathcal{M}_0(\lambda, \omega))\} = 2\sqrt{\omega - \omega^2}.$$

Proof. We need to find all $z \in \mathbb{C}$ such that

$$\mathcal{M}_0(\lambda, \omega) - z = \alpha\mathbf{J}\beta - \omega\alpha\mathbf{J} - \alpha\lambda - z = -\alpha\lambda - \omega\alpha\mathbf{J} - z + \alpha\mathbf{J}\beta$$

is degenerate. Multiplying the above matrix in the right-hand side by $-\alpha\mathbf{J}$, we need to find out when the matrix

$$\mathbf{J}\lambda - \omega + \alpha\mathbf{J}z + \beta$$

is degenerate. Since $\alpha\beta$ anticommutes with both α and β , while $\det \alpha = \det \beta = 1$, one has:

$$\begin{aligned} \det(\mathbf{J}\lambda - \omega + \alpha\mathbf{J}z + \beta)^2 &= \det((\mathbf{J}\lambda - \omega + \alpha\mathbf{J}z + \beta)\alpha\beta(\mathbf{J}\lambda - \omega + \alpha\mathbf{J}z + \beta)\beta\alpha) \\ &= \det((\mathbf{J}\lambda - \omega + \alpha\mathbf{J}z + \beta)(\mathbf{J}\lambda - \omega - \alpha\mathbf{J}z - \beta)) = \det((\mathbf{J}\lambda - \omega)^2 - (-z^2 + 1)). \end{aligned}$$

Since $\sigma(\mathbf{J}) = \{\pm i\}$, we conclude that the above determinant vanishes (hence $z \in \sigma(\mathcal{M}_0(\lambda, \omega))$) if and only if

$$z^2 - 1 + (\pm i\lambda - \omega)^2 = 0.$$

The conclusion about the spectrum of \mathcal{M}_0 follows.

Other statements are checked by direct computation. \square

Due to the symmetry of the potential \mathbf{W} (see (4.6) below), we have the following results.

Lemma 4.2. *If $\psi(x)$ solves (4.5) for $\lambda \in \mathbb{C}$, then $\theta(x) = \beta\psi(-x)$ also solves (4.5) for the same $\lambda \in \mathbb{C}$.*

Proof. Since v is even and u is odd, and since $\beta = \sigma_3$ anticommutes with σ_1 , there are the relations

$$W_0(x)\beta = \beta W_0(-x), \quad W_1(x)\beta = \beta W_1(-x), \tag{4.6}$$

for W_0, W_1 from the Gross–Neveu model (3.4). (It is convenient to notice that for each of these models, W_0 and W_1 can be written as linear combinations of the form $w_a(x)\sigma_1 + w_b(x)\sigma_3 + w_c(x)I_2$, with scalar-valued functions w_b and w_c symmetric in x and w_a skew-symmetric.) The conclusion follows. \square

Lemma 4.3. *For any $x \in \mathbb{R}$, $\omega \in \Omega$, and $\lambda \in \mathbb{C}$, the matrix \mathcal{M} from (4.5) satisfies $\operatorname{tr} \mathcal{M}(x, \lambda, \omega) = 0$.*

Proof. The statement is immediate for all the terms from \mathcal{M}_0 (cf. (4.4)). The remaining relation $\text{tr} \alpha \mathbf{J} \mathbf{W} = 0$ is checked with the explicit expressions (3.4). \square

We now turn to the construction of the Jost solutions, which are defined as eigenfunctions of $\mathbf{J} \mathbf{L}(\omega)$ with the same asymptotic behavior as eigenfunctions of $\mathbf{J}(\mathbf{D}_m - \omega)$. To do this, for $\lambda \in \mathbb{C}$, we first define

$$\xi_1(\lambda, \omega) = \sqrt{(\omega - i\lambda)^2 - 1}, \quad \xi_2(\lambda, \omega) = \sqrt{(\omega + i\lambda)^2 - 1}, \tag{4.7}$$

so that $\sigma(\mathcal{M}_0(\lambda, \omega)) = \{\pm \xi_1(\lambda, \omega), \pm \xi_2(\lambda, \omega)\}$ (cf. Lemma 4.1). Without loss of generality, we will only consider the case

$$\omega \geq 0, \quad \text{Re } \lambda \leq 0, \quad \text{Im } \lambda \geq 0; \tag{4.8}$$

in each of the two square roots in (4.7), we choose the branch that is positive for $\lambda \in i\mathbb{R}, \text{Im } \lambda \gg 1$.

We define

$$\Xi_1(\lambda, \omega) = \frac{1}{c_1(\lambda, \omega)} \begin{bmatrix} i\xi_1 \\ -i\lambda - 1 + \omega \\ -\xi_1 \\ \lambda - i(1 - \omega) \end{bmatrix}, \quad \mathbf{H}_1(\lambda, \omega) = \frac{1}{c_1(\lambda, \omega)} \begin{bmatrix} i\xi_1 \\ i\lambda + 1 - \omega \\ -\xi_1 \\ -\lambda + i(1 - \omega) \end{bmatrix}, \tag{4.9}$$

$$\Xi_2(\lambda, \omega) = \frac{1}{c_2(\lambda, \omega)} \begin{bmatrix} i\xi_2 \\ i\lambda - 1 + \omega \\ \xi_2 \\ \lambda + i(1 - \omega) \end{bmatrix}, \quad \mathbf{H}_2(\lambda, \omega) = \frac{1}{c_2(\lambda, \omega)} \begin{bmatrix} i\xi_2 \\ -i\lambda + 1 - \omega \\ \xi_2 \\ -\lambda - i(1 - \omega) \end{bmatrix}, \tag{4.10}$$

with the constants

$$c_1(\lambda, \omega) > 0, \quad c_2(\lambda, \omega) > 0 \tag{4.11}$$

chosen so that $|\Xi_j| = |\mathbf{H}_j| = 1, j = 1, 2$. Note that $\mathbf{H}_j = \beta \Xi_j; j = 1, 2$. The plane waves

$$\Xi_1(\lambda, \omega) e^{i\xi_1(\lambda, \omega)x}, \quad \Xi_2(\lambda, \omega) e^{i\xi_2(\lambda, \omega)x}, \quad \mathbf{H}_1(\lambda, \omega) e^{-i\xi_1(\lambda, \omega)x}, \quad \mathbf{H}_2(\lambda, \omega) e^{-i\xi_2(\lambda, \omega)x}$$

satisfy the equation $(\mathbf{J}(\mathbf{D}_m - \omega) - \lambda)\psi(x) = 0$ (and thus (4.4)).

By (4.8), we see that

$$\xi_1 > \xi_2 \geq 0 \quad \text{for } \lambda \in i\mathbb{R}, \quad |\lambda| \geq 1 + |\omega|; \quad |\xi_2| > |\xi_1| \quad \text{for } \lambda \in i\mathbb{R}, \quad |\lambda| \leq 1 - |\omega|.$$

We denote

$$\kappa_1 = |\text{Im } \xi_1|, \quad \kappa_2 = |\text{Im } \xi_2|; \tag{4.12}$$

then one has

$$\kappa_2 > \kappa_1 \geq 0 \quad \text{for } \lambda \in i\mathbb{R}, \quad |\lambda| \leq 1 - |\omega|.$$

Now we introduce the Jost solutions of the operator $\mathbf{J} \mathbf{L}$ defined in (3.3), (3.4).

Proposition 4.4. *Let $f \in C^\infty$ be such that for $|s| \leq 1$ $f(s) = \mathcal{O}(s^k)$ for $|s| \leq 1$ with $k \geq 2$. Let $\phi_\omega \in H^1(\mathbb{R}, \mathbb{C}^2)$, $\omega \in (-1, 1)$ be a solitary wave profile, and let the operator $\mathbf{J} \mathbf{L}$ be defined in (3.3), (3.4). There are the Jost solutions $\mathbf{f}_j(x, \lambda, \omega), \mathbf{F}_j(x, \lambda, \omega), j = 1, 2$, which satisfy the equation $(\mathbf{J} \mathbf{L}(\omega) - \lambda)u = 0$ and have the following properties. There is $c(\omega) < \infty$ such that*

- For $\lambda \in i\mathbb{R}, |\lambda| \leq 1 - |\omega|$,

$$|e^{\kappa_j x} \mathbf{f}_j(x, \lambda, \omega) - \Xi_j(\lambda, \omega)| + |e^{-\kappa_j x} \mathbf{F}_j(x, \lambda, \omega) - \mathbf{H}_j(\lambda, \omega)| \leq c(\omega) e^{-2k\delta_\omega x}, \quad x \geq 0, \quad j = 1, 2. \tag{4.13}$$

- For $\lambda \in i\mathbb{R}$, $1 - |\omega| \leq |\lambda| \leq 1 + |\omega|$,

$$\begin{aligned} |e^{-i\xi_1 x} \mathbf{f}_1(x, \lambda, \omega) - \Xi_1(\lambda, \omega)| + |e^{i\xi_1 x} \mathbf{F}_1(x, \lambda, \omega) - \mathbf{H}_1(\lambda, \omega)| &\leq c(\omega)e^{-2k\delta_\omega x}, & x \geq 0, \\ |e^{\kappa_2 x} \mathbf{f}_2(x, \lambda, \omega) - \Xi_2(\lambda, \omega)| + |e^{-\kappa_2 x} \mathbf{F}_2(x, \lambda, \omega) - \mathbf{H}_2(\lambda, \omega)| &\leq c(\omega)e^{-2k\delta_\omega x}, & x \geq 0. \end{aligned}$$

- For $\lambda \in i\mathbb{R}$, $|\lambda| \geq 1 + |\omega|$,

$$|e^{-i\xi_j x} \mathbf{f}_j(x, \lambda, \omega) - \Xi_j(\lambda, \omega)| + |e^{i\xi_j x} \mathbf{F}_j(x, \lambda, \omega) - \mathbf{H}_j(\lambda, \omega)| \leq c(\omega)e^{-2k\delta_\omega x}, \quad x \geq 0, \quad j = 1, 2.$$

- For $\lambda \in i\mathbb{R}$,

$$|\mathbf{f}_j(x, \lambda, \omega)| + |\mathbf{F}_j(x, \lambda, \omega)| \leq c(\omega)(\langle x \rangle + e^{\kappa_2|x|}), \quad x \leq 0, \quad j = 1, 2.$$

- For $\lambda \in i\mathbb{R}$, $|\lambda| \geq 3$,

$$|\mathbf{f}_j(x, \lambda, \omega)| + |\mathbf{F}_j(x, \lambda, \omega)| \leq c(\omega), \quad x \in \mathbb{R}, \quad j = 1, 2. \tag{4.14}$$

Above, $\delta_\omega = \sqrt{1 - \omega^2}$ (cf. (2.15)).

Using the above proposition, we also define the Jost solutions with appropriate asymptotics as $x \rightarrow -\infty$ by

$$\mathbf{g}_j(x) = \beta \mathbf{f}_j(-x), \quad \mathbf{G}_j(x) = \beta \mathbf{F}_j(-x), \quad x \in \mathbb{R}, \quad j = 1, 2. \tag{4.15}$$

Proof. The proof is quite standard. However, since the decay rate of the potential \mathbf{W} depends on ω and k (cf. Assumption 2.4), we choose to provide the details. Given $\varkappa \in \sigma(\mathcal{M}_0(\lambda, \omega))$, with $\omega \in \Omega$ and $\lambda \in i\mathbb{R}$, let $\Xi \in \mathbb{C}^4$ be a corresponding eigenvector, with $|\Xi| = 1$. To find a solution $\psi(x) \sim \Xi e^{\varkappa x}$, $x \rightarrow +\infty$ of (4.4), we define $\xi(x)$ by

$$\psi(x) = e^{\varkappa x} \xi(x), \quad \text{so that} \quad \xi|_{x=+\infty} = \Xi;$$

then $\partial_x \xi = (\mathcal{M}_0 - \varkappa)\xi + \alpha \mathbf{JW}\xi$, and hence we can write

$$\partial_x (e^{-(\mathcal{M}_0 - \varkappa)x} \xi) = e^{-(\mathcal{M}_0 - \varkappa)x} \alpha \mathbf{JW}\xi.$$

We construct $\xi(x)$ in the form of the power series $\xi(x) = \sum_{n=0}^{\infty} \xi_n(x)$, where $\xi_0 = \Xi$ and

$$\partial_x (e^{-(\mathcal{M}_0 - \varkappa)x} \xi_n(x)) = e^{-(\mathcal{M}_0 - \varkappa)x} \alpha \mathbf{JW}(x, \omega) \xi_{n-1}(x), \quad \xi_n(+\infty) = 0, \quad n \geq 1;$$

hence

$$\xi_n(x) = - \int_x^{+\infty} e^{(\mathcal{M}_0 - \varkappa)(x-y)} \alpha \mathbf{JW}(y, \omega) \xi_{n-1}(y) dy, \quad n \geq 1.$$

Let P_ζ denote the Riesz projector onto the eigenspace corresponding to $\zeta \in \sigma(\mathcal{M}_0)$. Then, for $x \geq 0$,

$$\begin{aligned} |P_\zeta \xi_n(x)| &\leq \int_x^{+\infty} \left\| P_\zeta e^{(\mathcal{M}_0 - \varkappa)(x-y)} \right\| \|\mathbf{W}(y, \omega)\|_{\text{End}(\mathbb{C}^4)} |\xi_{n-1}(y)| dy \\ &\leq \sup_{y \geq x} |\xi_{n-1}(y)| \int_x^{+\infty} a e^{(x-y) \text{Re}(\zeta - \varkappa)} \langle x - y \rangle K e^{-2k\delta_\omega y} dy \leq c e^{-2k\delta_\omega x} \sup_{y \geq x} |\xi_{n-1}(y)|, \end{aligned} \tag{4.16}$$

for some $c = c(\omega, K) < \infty$. Above, we used the bound $\|P_\zeta e^{(\mathcal{M}_0 - \varkappa)x}\| \leq a e^{x \text{Re}(\zeta - \varkappa)} \langle x \rangle$, with some $a < \infty$ (which depends on ω but does not depend on ζ), with the factor $\langle x \rangle$ due to the possibility of the Jordan block of \mathcal{M}_0 (when $\zeta = 0$). For the convergence of the integration in y , we used the bound (3.8) and the inequalities

$$|\text{Re } \zeta| \leq k\delta_\omega, \quad |\text{Re } \varkappa| \leq k\delta_\omega, \tag{4.17}$$

which are trivially satisfied under conditions of Assumption 2.4: one has $k \geq 2$, $|\omega| < 1/3$, hence $k\delta_\omega \geq 2\sqrt{8/9}$, while $\zeta \in \sigma(\mathcal{M}_0(\lambda, \omega))$ for any $\lambda \in i\mathbb{R}$, $\omega \in (-1, 1)$, satisfy $|\text{Re } \zeta| \leq 1$ (cf. Lemma 4.1 (ii)).

Then the integration in y in (4.16) can be estimated as follows:

$$\int_x^{+\infty} e^{(x-y) \operatorname{Re}(\zeta-x)} \langle x-y \rangle e^{-2k\delta_\omega y} dy = e^{-2k\delta_\omega x} \int_0^{+\infty} e^{-z \operatorname{Re}(\zeta-x)} \langle z \rangle e^{-2k\delta_\omega z} dz$$

$$\leq e^{-2k\delta_\omega x} \left(\frac{1}{2(k-1)\delta_\omega} + \frac{1}{(2(k-1)\delta_\omega)^2} \right).$$

We conclude that

$$\sup_{y \geq x} |\xi_n(y)| = \sup_{y \geq x} \left| \sum_{\zeta \in \sigma(\mathcal{M}_0)} P_\zeta \xi_n(y) \right| \leq \sum_{\zeta \in \sigma(\mathcal{M}_0)} \sup_{y \geq x} |P_\zeta \xi_n(y)| \leq 4ce^{-2k\delta_\omega x} \sup_{y \geq x} |\xi_{n-1}(y)|, \quad x \geq 0.$$

Therefore, there is $C < \infty$ such that $\sum_{n=1}^\infty |\xi_n(x)| \leq Ce^{-2k\delta_\omega x}$, for all $x \geq 0$, hence

$$|\xi(x) - \Xi| = Ce^{-2k\delta_\omega x}, \quad x \geq 0.$$

Let us prove the uniform bounds (4.14). Let us write (4.5) in the form

$$(\partial_x - \mathcal{M}_0(\lambda, \omega))\psi = \alpha \mathbf{JW}(x, \omega)\psi. \tag{4.18}$$

Using the Green function for the operator $\partial_x - \mathcal{M}_0(\lambda, \omega)$, which is given by

$$\mathcal{G}(x, y, \lambda, \omega) = \left(\Xi_1 \otimes \theta_1^* e^{i(x-y)\xi_1} + \Xi_2 \otimes \theta_2^* e^{i(x-y)\xi_2} + \mathbf{H}_1 \otimes \eta_1^* e^{-i(x-y)\xi_1} + \mathbf{H}_2 \otimes \eta_2^* e^{-i(x-y)\xi_2} \right) \Theta(x-y),$$

where Θ is the Heaviside step-function and $\theta_j, \eta_j \in \mathbb{C}^4, j = 1, 2$, is the basis dual to $\Xi_j, \mathbf{H}_j \in \mathbb{C}^4$, one can construct the solutions $\mathbf{f}_j(x, \lambda, \omega), \mathbf{F}_j(x, \lambda, \omega)$, in the form of the power series

$$\psi = \sum_{n=0}^\infty \psi_n, \tag{4.19}$$

with $\psi_0(x) = \Xi_j e^{i\xi_j x}$ or $\psi_0(x) = \mathbf{H}_j e^{-i\xi_j x}$ (according to (4.15), these are asymptotics of $\mathbf{f}_j(x), \mathbf{F}_j(x)$ for $x \gg 1$), and with $\psi_n(x), n \geq 1$ solving

$$(\partial_x - \mathcal{M}_0(\lambda, \omega))\psi_n(x) = \alpha \mathbf{JW}(x, \omega)\psi_{n-1}(x).$$

For definiteness, we will consider $\mathbf{f}_1(x)$ only (other functions are considered in the same way). For any $x \in \mathbb{R}$, the series (4.19) converges due to the estimate

$$\begin{aligned} |\psi_n(x)| &\leq \int_x^\infty \|\mathbf{W}(x_1)\|_{\operatorname{End}(\mathbb{C}^4)} |\psi_{n-1}(x_1)| dx_1 \\ &\leq \int_x^\infty \int_{x_1}^\infty \|\mathbf{W}(x_1)\|_{\operatorname{End}(\mathbb{C}^4)} \|\mathbf{W}(x_2)\|_{\operatorname{End}(\mathbb{C}^4)} |\psi_{n-2}(x_2)| dx_1 dx_2 \leq \dots \\ &\leq \int_{x < x_1 < \dots < x_n < \infty} \left(\prod_{l=1}^n \|\mathbf{W}(x_l)\|_{\operatorname{End}(\mathbb{C}^4)} \right) |\psi_0(x_n)| dx_1 \dots dx_n \\ &\leq \frac{1}{n!} \int_{x_1 > x} \dots \int_{x_n > x} \left(\prod_{l=1}^n \|\mathbf{W}(x_l)\|_{\operatorname{End}(\mathbb{C}^4)} \right) |\psi_0(x_n)| dx_1 \dots dx_n \leq \frac{(\int_x^\infty \|\mathbf{W}(y)\|_{\operatorname{End}(\mathbb{C}^4)} dy)^n}{n!}, \end{aligned}$$

where we represented the integration over the simplex $x < x_1 < \dots < x_n < \infty$ in \mathbb{R}^n as a fraction of the integration over the quadrant $x_l > x, 1 \leq l \leq n$, and substituted $|\psi_0(x_n)| = |\Xi_1| = 1$. Therefore, $|\psi(x)| \leq \sum_{n \geq 0} |\psi_n(x)| \leq \exp(\int_{\mathbb{R}} \|\mathbf{W}(y)\|_{\operatorname{End}(\mathbb{C}^4)} dy)$, for any $x \in \mathbb{R}$. This proves (4.14).

Finally, Lemma 4.2 allows us to use (4.15) to obtain the Jost solutions with required asymptotic behavior at $-\infty$. \square

Definition 4.5. We define the Evans function by

$$E(\lambda, \omega) = \det [\mathbf{f}_1(x, \lambda, \omega), \mathbf{f}_2(x, \lambda, \omega), \mathbf{g}_1(x, \lambda, \omega), \mathbf{g}_2(x, \lambda, \omega)]. \tag{4.20}$$

We note that, by Liouville’s formula and by Lemma 4.3,

$$\frac{\partial}{\partial x} \ln (\det [\mathbf{f}_1, \mathbf{f}_2, \mathbf{g}_1, \mathbf{g}_2]) = \text{tr } \mathcal{M} = 0,$$

hence the right-hand side of (4.20) does not depend on $x \in \mathbb{R}$.

The following lemma gives the relation between the eigenvalues and the zeros of the Evans function.

Lemma 4.6. Fix $\omega \in \Omega$.

- (i) Let $\lambda \in i\mathbb{R}$, $|\lambda| \in (1 - |\omega|, 1 + |\omega|)$. Then $E(\lambda, \omega) = 0$ at some $\lambda \in i\mathbb{R}$, $|\lambda| \in (1 - |\omega|, 1 + |\omega|)$, if and only if λ is an L^2 eigenvalue of \mathbf{JL} .
- (ii) At $\lambda = \pm i(1 + |\omega|)$, one has $E(\lambda, \omega) = 0$ if and only if there is a generalized L^∞ -eigenfunction corresponding to λ , which has the asymptotics $\psi \sim a\Xi_2$ as $x \rightarrow +\infty$, $\psi \sim b\mathbf{H}_2$ as $x \rightarrow -\infty$.

Remark 4.7. The statement of the lemma at the thresholds is non-trivial since at the threshold points the solution to $(\mathbf{JL} - \lambda)\psi = 0$ which is bounded for $x \rightarrow +\infty$ could be linearly growing as $x \rightarrow -\infty$.

Proof. Let us start with the easy “if” statements. For the “if” statement of Part 1, we consider the case $\pm\lambda \in i(1 - |\omega|, 1 + |\omega|)$. If ψ is an eigenfunction (L^2 solution to (4.5)), then, due to the asymptotics of the Jost solutions (Proposition 4.4), $\psi = C\mathbf{f}_2 = C'\mathbf{g}_2$ for some nonzero $C, C' \in \mathbb{C}$, hence \mathbf{f}_2 and \mathbf{g}_2 are linearly dependent, resulting in $E(\lambda, \omega) = 0$. The proof of the “if” part of Part 2 is similar. Let $\lambda = \pm i(1 + |\omega|)$. If ψ is a generalized L^∞ eigenfunction of (4.5) with the asymptotics as assumed in Part 2 of the lemma, due to the asymptotics of the Jost solutions (\mathbf{f}_1 and $\mathbf{F}_j, 1 \leq j \leq 2$, are not small when $x \rightarrow +\infty$; \mathbf{g}_1 and $\mathbf{G}_j, 1 \leq j \leq 2$, is not small when $x \rightarrow -\infty$), one again has $\psi = C\mathbf{f}_2 = C'\mathbf{g}_2$ for some nonzero $C, C' \in \mathbb{C}$, hence \mathbf{f}_2 and \mathbf{g}_2 are linearly dependent, thus $E(\lambda, \omega) = 0$.

Let us prove the “only if” statement of Part 1. If $\det[\mathbf{f}_1, \mathbf{f}_2, \mathbf{g}_1, \mathbf{g}_2] = 0$ for some $\lambda \in i\mathbb{R}$, then there are $a_1, a_2, b_1, b_2 \in \mathbb{C}$, not all of them equal to zero, one has

$$\Phi(x) := \sum_{j=1}^2 a_j \mathbf{f}_j(x, \lambda, \omega) = \sum_{j=1}^2 b_j \mathbf{g}_j(x, \lambda, \omega), \quad x \in \mathbb{R}. \tag{4.21}$$

Since \mathbf{f}_j are linearly independent, and so are \mathbf{g}_j , the function Φ thus defined is not identically zero.

Define

$$\Sigma = i\mathbf{J} = \begin{bmatrix} 0 & iI_2 \\ -iI_2 & 0 \end{bmatrix}.$$

Let us consider the auxiliary Dirac equation

$$i \Sigma \partial_t \Psi = \mathbf{L}\Psi, \quad \Psi(x, t) \in \mathbb{C}^4, \quad x \in \mathbb{R}, \tag{4.22}$$

where $\mathbf{L} = \mathbf{J}\alpha\partial_x + \beta + \mathbf{W} - \omega$. This is a Hamiltonian system with the Hamiltonian density

$$\mathfrak{h} = \Psi^* \mathbf{L}\Psi = \Psi^* (\mathbf{J}\alpha\partial_x + \beta + \mathbf{W} - \omega)\Psi$$

and the Lagrangian density

$$\mathfrak{l} = \Psi^* (i \Sigma \partial_t - \mathbf{L})\Psi.$$

If $\Phi \in C^1(\mathbb{R}, \mathbb{C}^4)$ satisfies $\lambda\Phi = \mathbf{JL}\Phi$, which we write as $(i\lambda)(i\mathbf{J})\Phi = -\lambda\mathbf{J}\Phi = \mathbf{L}\Phi$, then we have

$$\Omega \Sigma \Phi = \mathbf{L}\Phi, \quad \Omega := i\lambda \in \mathbb{R}.$$

Thus, $\Psi(x, t) = \Phi(x)e^{-i\Omega t}$ is a “solitary wave solution” to (4.22), except that Φ is not necessarily in L^2 .

Equation (4.22) conserves the Krein charge; its density is

$$\sigma(x, t) = \Psi^* \Sigma \Psi = \Phi^* \Sigma \Phi, \tag{4.23}$$

while the density of the corresponding current is

$$j(x, t) = \Psi^* \Sigma \alpha \Psi = \Phi^* \Sigma \alpha \Phi. \tag{4.24}$$

Remark 4.8. We call the quantity $\langle \Psi, \Sigma \Psi \rangle$ the “Krein charge” in view of its relation to the Krein index considerations. Namely, the relation $\mathbf{JL}\Phi = \lambda\Phi$ implies that $\langle \Phi, \mathbf{L}\Phi \rangle = -\lambda \langle \Phi, \mathbf{J}\Phi \rangle$, with $\langle \Phi, \mathbf{L}\Phi \rangle$ real and $\langle \Phi, \mathbf{J}\Phi \rangle$ purely imaginary; hence $\text{Re } \lambda \neq 0$ leads to $\langle \Phi, \mathbf{J}\Phi \rangle = 0$, $\langle \Phi, \mathbf{L}\Phi \rangle = 0$. Thus, the Krein signature is zero (\mathbf{L} is not sign-definite on the corresponding eigenspace) for any eigenvalue away from the imaginary axis. (The above could also be interpreted as follows. We could say that if $\Psi = \Phi(x)e^{-i\Omega t}$ (with $\Omega = i\lambda$) is a solitary wave solution to (4.22) and $\text{Im } \Omega = \text{Re } \lambda \neq 0$, then the conservation of the “Krein charge” $\langle \Psi(t), \Sigma \Psi(t) \rangle = \langle \Phi, \Sigma \Phi \rangle e^{2 \text{Im } \Omega t}$ requires that this charge is zero, $\langle \Phi, \Sigma \Phi \rangle = 0$.) It follows that purely imaginary eigenvalues $\lambda \in i\mathbb{R} \setminus 0$ with nonzero Krein signature, $\langle \Phi, i\mathbf{J}\Phi \rangle \neq 0$, cannot bifurcate off the imaginary axis into the complex plane.

Since the Krein charge density does not depend on time, the local conservation of the Krein charge in the system (4.22) leads to the equality of the Krein current (4.24) evaluated at the endpoints of the interval $(-l, l)$, $l > 0$. Therefore, taking into account that

$$\Sigma \Xi_1 = -\Xi_1, \quad \Sigma \Xi_2 = \Xi_2, \quad \Sigma H_1 = -H_1, \quad \Sigma H_2 = H_2,$$

under the assumption that $E(\lambda, \omega) = 0$, we compute for Φ from (4.21):

$$\begin{aligned} 0 &= \lim_{l \rightarrow +\infty} \Phi^* \Sigma \alpha \Phi \Big|_{-l}^l = \lim_{l \rightarrow +\infty} \left((a_1 \Xi_1 e^{i\xi_1 x} + a_2 \Xi_2 e^{-\kappa_2 x})^* \alpha (-a_1 \Xi_1 e^{i\xi_1 x} + a_2 \Xi_2 e^{-\kappa_2 x}) \right) \Big|_{x=l} \\ &\quad - \lim_{l \rightarrow +\infty} \left((b_1 H_1 e^{-i\xi_1 x} + b_2 H_2 e^{-\kappa_2 |x|})^* \alpha (-b_1 H_1 e^{-i\xi_1 x} + b_2 H_2 e^{-\kappa_2 |x|}) \right) \Big|_{x=-l} \\ &= \lim_{l \rightarrow +\infty} \left((a_1 \Xi_1 e^{i\xi_1 x} + a_2 \Xi_2 e^{-\kappa_2 x})^* \alpha (-a_1 \Xi_1 e^{i\xi_1 x} + a_2 \Xi_2 e^{-\kappa_2 x}) \right) \Big|_{x=l} \\ &\quad - \lim_{l \rightarrow +\infty} \left((b_1 \Xi_1 e^{-i\xi_1 x} + b_2 \Xi_2 e^{-\kappa_2 |x|})^* \alpha (b_1 \Xi_1 e^{-i\xi_1 x} - b_2 \Xi_2 e^{-\kappa_2 |x|}) \right) \Big|_{x=-l}. \end{aligned} \tag{4.25}$$

In the last relation, we took into account that $H_j = \beta \Xi_j$ and that β anticommutes with α . Taking into account that $\Xi_1^* \alpha \Xi_2 = \Xi_2^* \alpha \Xi_1 = 0$, we rewrite the above as

$$0 = (|a_1|^2 + |b_1|^2) \Xi_1^* \alpha \Xi_1 + (|a_2|^2 + |b_2|^2) \Xi_2^* \alpha \Xi_2 \lim_{l \rightarrow +\infty} e^{-2\kappa_2 l}. \tag{4.26}$$

For Part 1, when $\lambda \in i\mathbb{R}$ and $1 - |\omega| < |\lambda| < 1 + |\omega|$, one has $\kappa_2 > 0$, hence the second term in the right-hand side of (4.26) vanishes. On the other hand,

$$\Xi_1^* \alpha \Xi_1 = \frac{4i(\lambda - i(1 - \omega))\xi_1}{c_1^2} > 0 \quad \text{for } \lambda \in i\mathbb{R}, \quad \text{Im } \lambda > 1 - |\omega|.$$

Then it follows from (4.26) that $a_1 = b_1 = 0$, and we conclude that Φ is exponentially decaying for $x \rightarrow \pm\infty$, so that λ is an L^2 eigenvalue. This finishes the proof of Part 1 of the lemma.

Finally, let us prove the “only if” statement of Part 2. When $\lambda = \pm(1 + |\omega|)i$, one has $\kappa_2 = 0 = \xi_2$, and, using (4.10), one computes

$$\Xi_2^* \alpha \Xi_2 = \frac{4i(\lambda + i(1 - \omega))\xi_2}{c_2^2} = 0;$$

therefore, the assumption $E(\lambda, \omega) = 0$ which leads to (4.26) results in $a_1 = b_1 = 0$. \square

For $\lambda \in i\mathbb{R}$, $|\lambda| > 1 + |\omega|$, the solutions $\mathbf{f}_1(\lambda, \omega)$, $\mathbf{f}_2(\lambda, \omega)$, $\mathbf{F}_1(\lambda, \omega)$, $\mathbf{F}_2(\lambda, \omega)$ are linearly independent, constituting a fundamental set of solutions to (4.5); hence, there are $A(\lambda, \omega)$, $B(\lambda, \omega) \in \mathbb{C}^{4 \times 4}$, locally bounded in λ, ω , such that

$$\mathbf{g}_j(x, \lambda, \omega) = \sum_{k=1}^2 \mathbf{f}_k(x, \lambda, \omega) A_{kj}(\lambda, \omega) + \sum_{k=1}^2 \mathbf{F}_k(x, \lambda, \omega) B_{kj}(\lambda, \omega), \quad j = 1, 2. \tag{4.27}$$

We note that, by (4.15), applying β to (4.27) and flipping x , we also have

$$\mathbf{f}_j(x, \lambda, \omega) = \sum_{k=1}^2 \mathbf{g}_k(x, \lambda, \omega) A_{kj}(\lambda, \omega) + \sum_{k=1}^2 \mathbf{G}_k(x, \lambda, \omega) B_{kj}(\lambda, \omega), \quad j = 1, 2. \tag{4.28}$$

Lemma 4.9. For each $\omega \in \Omega$, the matrices $A(\lambda, \omega)$, $B(\lambda, \omega)$ from (4.27) satisfy

$$\lim_{\lambda \rightarrow \pm i\infty} \|A(\lambda, \omega)\|_{\text{End}(\mathbb{C}^4)} = 0, \quad \lim_{\lambda \rightarrow \pm i\infty} B(\lambda, \omega) = B_\infty(\omega), \tag{4.29}$$

with $\|B_\infty(\omega)\|_{\text{End}(\mathbb{C}^N)} < \infty$. Moreover,

$$\det B_\infty(\omega) = 1. \tag{4.30}$$

Proof. The bound (4.14) (which is also valid for \mathbf{g}_j , \mathbf{G}_j in view of (4.15)), together with (4.27) and with the asymptotic behavior of \mathbf{f} , \mathbf{F} for $x \gg 1$ (Proposition 4.4) and linear independence of Ξ_j , H_j , $1 \leq j \leq 2$, leads to

$$\lim_{\lambda \rightarrow \pm i\infty} (\|A(\lambda, \omega)\|_{\text{End}(\mathbb{C}^2)} + \|B(\lambda, \omega)\|_{\text{End}(\mathbb{C}^2)}) < \infty.$$

Following the proof of (4.14) from Proposition 4.4 and using the stationary phase method, which yields

$$\int_x^\infty \Xi_j \otimes \theta_k^* e^{i(x-y)\xi_j} \mathbf{W}(y) H_l e^{-iy\xi_l} dy = \mathcal{O}\left(\frac{1}{\xi_j}\right) \rightarrow 0 \quad \text{as } \lambda \rightarrow \pm i\infty,$$

one shows that $\|A(\lambda, \omega)\|_{\text{End}(\mathbb{C}^4)} \rightarrow 0$ as $\lambda \rightarrow \pm i\infty$.

Let us show that $\det B_\infty(\omega) = 1$. First, we note from (4.9), (4.10) that

$$\lim_{\lambda \rightarrow \pm i\infty} \Xi_1(\lambda, \omega) = M \lim_{\lambda \rightarrow \pm i\infty} H_2(\lambda, \omega), \quad \lim_{\lambda \rightarrow \pm i\infty} \Xi_2(\lambda, \omega) = M \lim_{\lambda \rightarrow \pm i\infty} H_1(\lambda, \omega),$$

where $M = \begin{bmatrix} I_2 & 0 \\ 0 & -I_2 \end{bmatrix}$. Therefore, taking into account that for each $x \in \mathbb{R}$ one has $\lim_{\lambda \rightarrow \pm i\infty} |x| |\xi_1 - \xi_2| \rightarrow 0$, we have $\lim_{\lambda \rightarrow \pm i\infty} \|(\mathbf{f}_1, \mathbf{f}_2) - M(\mathbf{F}_2, \mathbf{F}_1)e^{2i\xi_1 x}\|_{\mathbb{C}^4 \times \mathbb{C}^4} \rightarrow 0$, for each fixed $x \gg 1$, and hence (due to continuous dependence of solutions to (4.18) on the initial data) for each fixed $x \in \mathbb{R}$:

$$\lim_{\lambda \rightarrow \pm i\infty} \|(\mathbf{f}_1(x, \lambda, \omega), \mathbf{f}_2(x, \lambda, \omega)) - M(\mathbf{F}_2(x, \lambda, \omega), \mathbf{F}_1(x, \lambda, \omega))e^{2i\xi_1 x}\|_{\mathbb{C}^4 \times \mathbb{C}^4} = 0, \quad x \in \mathbb{R}.$$

Similarly, comparing asymptotics for $x \ll -1$, we conclude that

$$\lim_{\lambda \rightarrow \pm i\infty} \|(\mathbf{G}_1(x, \lambda, \omega), \mathbf{G}_2(x, \lambda, \omega)) - M(\mathbf{g}_2(x, \lambda, \omega), \mathbf{g}_1(x, \lambda, \omega))e^{2i\xi_1 x}\|_{\mathbb{C}^4 \times \mathbb{C}^4} = 0, \quad x \in \mathbb{R}.$$

Therefore, besides (4.27), which yields $\lim_{\lambda \rightarrow \pm i\infty} \|(\mathbf{g}_1, \mathbf{g}_2) - (\mathbf{F}_1, \mathbf{F}_2)B\|_{L^\infty} = 0$ (due to (4.29)), we also have

$$\lim_{\lambda \rightarrow \pm i\infty} \|(\mathbf{G}_1(x, \lambda, \omega), \mathbf{G}_2(x, \lambda, \omega)) - (\mathbf{f}_1(x, \lambda, \omega), \mathbf{f}_2(x, \lambda, \omega))B(\lambda, \omega)\|_{\mathbb{C}^4 \times \mathbb{C}^4} = 0, \quad x \in \mathbb{R}.$$

On the other hand, from (4.28), taking into account (4.29), we also have

$$\lim_{\lambda \rightarrow \pm i\infty} \|(\mathbf{f}_1(x, \lambda, \omega), \mathbf{f}_2(x, \lambda, \omega)) - (\mathbf{G}_1(x, \lambda, \omega), \mathbf{G}_2(x, \lambda, \omega))B(\lambda, \omega)\|_{\mathbb{C}^4 \times \mathbb{C}^4} = 0, \quad x \in \mathbb{R}. \tag{4.31}$$

It follows that $\lim_{\lambda \rightarrow \pm i\infty} B(\lambda, \omega)^2 = I_2$, hence $\lim_{\lambda \rightarrow \pm i\infty} \det B(\lambda, \omega) = \pm 1$. The conclusion $\lim_{\lambda \rightarrow \pm i\infty} \det B(\lambda, \omega) = 1$ can be made by substituting the “interaction term” \mathbf{W} with $s\mathbf{W}$, $s \in [0, 1]$, and using the continuity argument when changing s from 0 to 1. \square

Lemma 4.10. For each $\omega \in \Omega$, one has $\lim_{\lambda \rightarrow \pm i\infty} |E(\lambda, \omega)| = 1$.

Proof. Using (4.9) and (4.10), we compute:

$$\det[\Xi_1, \Xi_2, H_1, H_2] = 1 + \mathcal{O}(|\lambda|^{-1}), \quad \lambda \rightarrow \pm i\infty. \tag{4.32}$$

On the other hand, by (4.27),

$$\begin{aligned} E(\lambda, \omega) &= \det[\mathbf{f}_1, \mathbf{f}_2, \mathbf{g}_1, \mathbf{g}_2] = \det[\mathbf{f}_1, \mathbf{f}_2, \sum_{j=1}^2 \mathbf{F}_j B_{j1}, \sum_{j=1}^2 \mathbf{F}_j B_{j2}] = \det B(\lambda, \omega) \det[\mathbf{f}_1, \mathbf{f}_2, \mathbf{F}_1, \mathbf{F}_2] \\ &= \det B(\lambda, \omega) \lim_{x \rightarrow +\infty} \det[\mathbf{f}_1, \mathbf{f}_2, \mathbf{F}_1, \mathbf{F}_2] = \det B(\lambda, \omega) \det[\Xi_1, \Xi_2, H_1, H_2], \end{aligned}$$

where we used the asymptotics of $\mathbf{f}_j, \mathbf{F}_j$ from Proposition 4.4. Therefore, by Lemma 4.9 and (4.32),

$$\lim_{\lambda \rightarrow \pm i\infty} E(\lambda, \omega) = \lim_{\lambda \rightarrow \pm i\infty} \det B(\lambda, \omega) \lim_{\lambda \rightarrow \pm i\infty} \det[\Xi_1, \Xi_2, H_1, H_2] = 1.$$

This finishes the proof. \square

Remark 4.11. For $\omega \in \Omega, \lambda \in i\mathbb{R}$ with $|\lambda| > 1 + |\omega|$, the Jost solutions $\mathbf{f}_j, \mathbf{F}_j, j = 1, 2$ (and similarly $\mathbf{g}_j, \mathbf{G}_j, j = 1, 2$) are linearly independent (since so are the vectors $\Xi_j, H_j, j = 1, 2$ from (4.9), (4.10)); hence there is a “scattering matrix” $S(\lambda, \omega) \in \mathbb{C}^{4 \times 4}$ such that

$$\begin{aligned} &(\mathbf{g}_1(x, \lambda, \omega), \mathbf{g}_2(x, \lambda, \omega), \mathbf{G}_1(x, \lambda, \omega), \mathbf{G}_2(x, \lambda, \omega)) \\ &= (\mathbf{f}_1(x, \lambda, \omega), \mathbf{f}_2(x, \lambda, \omega), \mathbf{F}_1(x, \lambda, \omega), \mathbf{F}_2(x, \lambda, \omega))S(\lambda, \omega). \end{aligned}$$

Taking into account the relations (4.15) between \mathbf{f}_j and \mathbf{g}_j and between \mathbf{F}_j and \mathbf{G}_j , we conclude that one also has

$$(\mathbf{f}_1, \mathbf{f}_2, \mathbf{F}_1, \mathbf{F}_2) = (\mathbf{g}_1, \mathbf{g}_2, \mathbf{G}_1, \mathbf{G}_2)S,$$

hence $S^2 = I, \det S = \pm 1$. Taking into account that $S \rightarrow \begin{bmatrix} 0 & I_2 \\ I_2 & 0 \end{bmatrix}$ in the limit of zero interaction (when $\mathbf{W}(x, \omega)$ in (4.3) is substituted by zero), we conclude that $\det S = 1$.

4.2. Explicit construction of the resolvent of the operator \mathbf{JL}

In this section, we will not restrict \mathbf{JL} onto \mathbf{X} and give a general construction of the resolvent in the case when $E(\lambda, \omega) \neq 0$.

Remark 4.12. Although for applications to asymptotic stability we will only need the resolvent of $\mathbf{JL}(\omega)$ for λ in the essential spectrum, we will make our construction for all $\lambda \in i\mathbb{R}$.

Definition 4.13. For $1 \leq p \leq \infty$ and $s \in \mathbb{R}$, we will use the weighted L^p spaces with polynomial weights:

$$\|f\|_{L^p_s} := \|\langle \cdot \rangle^s f\|_{L^p}.$$

For $f(x, t)$, we will denote

$$\|f\|_{(L^p_s)_x} := \|f(\cdot, t)\|_{L^p_s} = \|\langle \cdot \rangle^s f(\cdot, t)\|_{L^p}.$$

Proposition 4.14. Fix $\omega \in \Omega$. Assume that $\lambda \in i\mathbb{R}, |\lambda| \geq 1 - |\omega|$, is such that $E(\lambda, \omega) \neq 0$.

- There are resolvents $G^\pm(x, y, \lambda, \omega)$ of the operator $\mathbf{JL} = -\alpha(\partial_x - \mathcal{M}(x, \lambda, \omega))$ which satisfy

$$-\alpha(\partial_x - \mathcal{M}(x, \lambda \pm 0, \omega))G^\pm(x, y, \lambda \pm 0, \omega) = \delta(x - y)I_4, \tag{4.33}$$

and for some $C(\lambda, \omega) < \infty$ (locally bounded in λ and ω) one has

$$|G^\pm(x, y, \lambda, \omega)| \leq C(\lambda \pm 0, \omega) \min(\langle x \rangle, \langle y \rangle)\langle y \rangle, \quad (x, y) \in \mathbb{R}^2. \tag{4.34}$$

- For each $\omega \in \Omega$, there is $C(\omega) < \infty$ such that

$$\limsup_{\Lambda \rightarrow \pm\infty} \|G^\pm(x, y, i\Lambda \pm 0, \omega)\|_{\text{End}(\mathbb{C}^N)} \leq C(\omega), \quad (x, y) \in \mathbb{R}^2. \tag{4.35}$$

- For every $s > 3$ and $K > 0$, there is a constant $C_{s,K,\omega} < \infty$ such that for all $\lambda \in i\mathbb{R}$ with $|\lambda| < K$ one has

$$\sup_{\lambda \in i\mathbb{R} \pm 0, |\lambda| < K} \|(\mathbf{JL}(\omega) - \lambda)^{-1}\|_{L^2_s \rightarrow L^2_{-s}} \leq C_{s,K,\omega}. \tag{4.36}$$

- There is a constant $C_\omega < \infty$ such that

$$\limsup_{\lambda \in i\mathbb{R} \pm 0, |\lambda| \rightarrow \infty} \|(\mathbf{JL}(\omega) - \lambda)^{-1}\|_{L^2_1 \rightarrow L^2_{-1}} \leq C_\omega. \tag{4.37}$$

Proof. We will only provide a construction of G^- ; see Remark 4.19 below.

Recall that $\mathbf{f}_1, \mathbf{f}_2$ are Jost solutions decaying (or oscillating) for $x \rightarrow +\infty$, while $\mathbf{F}_1, \mathbf{F}_2$ are the growing ones (or oscillating ones). $\mathbf{f}_1, \mathbf{F}_1$ have κ_1 as the rate of decay and growth, respectively; \mathbf{f}_2 and \mathbf{F}_2 have the rate κ_2 , with $\kappa_2 > \kappa_1 \geq 0$ (cf. (4.12)). Similarly with $\mathbf{g}_1, \mathbf{g}_2, \mathbf{G}_1, \mathbf{G}_2$, for $x \rightarrow -\infty$.

Recall that if $\xi_j = 0$, then $\mathbf{F}_j = \mathbf{f}_j$, hence the set $\{\mathbf{f}_1, \mathbf{f}_2, \mathbf{F}_1, \mathbf{F}_2\}$ is no longer linearly independent. To overcome this issue, let us modify \mathbf{F}_j . For $\xi_j \neq 0$, denote

$$\tilde{\mathbf{F}}_j(x, \lambda, \omega) = \mathbf{F}_j(x, \lambda, \omega) + \frac{\mathbf{f}_j(x, \lambda, \omega) - \mathbf{F}_j(x, \lambda, \omega)}{2i\xi_j}, \quad j = 1, 2; \tag{4.38}$$

$$\tilde{\mathbf{G}}_j(x, \lambda, \omega) = \mathbf{G}_j(x, \lambda, \omega) + \frac{\mathbf{g}_j(x, \lambda, \omega) - \mathbf{G}_j(x, \lambda, \omega)}{2i\xi_j}. \quad j = 1, 2. \tag{4.39}$$

Note that by (4.15) one has $\tilde{\mathbf{G}}(x, \lambda, \omega) = \beta \tilde{\mathbf{F}}(-x, \lambda, \omega)$.

For $\lambda \in i\mathbb{R}$ such that $\xi_j(\lambda, \omega) = 0$, we define $\mathbf{F}_j(x, \lambda, \omega)$ by the pointwise limit:

$$\tilde{\mathbf{F}}_j(x, \lambda, \omega) = \mathbf{F}_j(x, \lambda, \omega) + \lim_{\lambda' \rightarrow \lambda; \xi_j(\lambda') > 0} \left\{ \frac{\mathbf{f}_j(x, \lambda', \omega) - \mathbf{F}_j(x, \lambda', \omega)}{2i\xi_j(\lambda')} \right\},$$

and similarly for $\tilde{\mathbf{G}}_j$; then one has $\tilde{\mathbf{F}}_j(x, \lambda, \omega) \sim \Xi_j \langle x \rangle$ for $x \gg 1$ and $\tilde{\mathbf{G}}_j(x, \lambda, \omega) \sim \mathbf{H}_j \langle x \rangle$ for $x \ll -1$.

By Proposition 4.4, we have the following asymptotics for $\tilde{\mathbf{F}}_j, \tilde{\mathbf{G}}_j$:

Lemma 4.15. For each $\omega \in \Omega, \lambda \in i\mathbb{R}$, one has:

$$\begin{aligned} |\tilde{\mathbf{F}}_j(x, \lambda, \omega)| &\leq C(\omega) \langle x \rangle e^{\kappa_j x}, & x \geq 0, & \quad j = 1, 2, \\ |\tilde{\mathbf{G}}_j(x, \lambda, \omega)| &\leq C(\omega) \langle x \rangle e^{\kappa_j |x|}, & x \leq 0, & \quad j = 1, 2, \end{aligned}$$

where $C(\omega)$ is locally bounded in ω .

Remark 4.16. In Lemma 4.15, the estimates remain true when λ is above the corresponding threshold, so that $\xi_j > 0$ while $\kappa_j = 0$ (cf. definition (4.12)).

Proof. This follows from Proposition 4.4 and definitions (4.38), (4.39). \square

Abusing the notations (cf. (4.27)), we assume that $A(\lambda, \omega), B(\lambda, \omega) \in \mathbb{C}^{4 \times 4}$ are such that

$$\mathbf{g}_k(x, \lambda, \omega) = \sum_{j=1}^2 \mathbf{f}_j(x, \lambda, \omega) A_{jk}(\lambda, \omega) + \sum_{j=1}^2 \tilde{\mathbf{F}}_j(x, \lambda, \omega) B_{jk}(\lambda, \omega), \quad k = 1, 2,$$

which we write as

$$(\mathbf{g}_1, \mathbf{g}_2) = (\mathbf{f}_1, \mathbf{f}_2)A + (\tilde{\mathbf{F}}_1, \tilde{\mathbf{F}}_2)B. \tag{4.40}$$

Multiplying (4.40) by β , flipping the sign of x , and using (4.15), we arrive at

$$(\mathbf{f}_1, \mathbf{f}_2) = (\mathbf{g}_1, \mathbf{g}_2)A + (\tilde{\mathbf{G}}_1, \tilde{\mathbf{G}}_2)B, \tag{4.41}$$

with the same A, B as in (4.40).

Lemma 4.17. *If $E(\lambda, \omega) \neq 0$, then the matrix $B(\lambda, \omega)$ is non-degenerate.*

Proof. By (4.40), $E(\lambda, \omega) = \det[\mathbf{f}_1, \mathbf{f}_2, \mathbf{g}_1, \mathbf{g}_2] = \det[\mathbf{f}_1, \mathbf{f}_2, (\mathbf{f}_1, \mathbf{f}_2)A + (\tilde{\mathbf{F}}_1, \tilde{\mathbf{F}}_2)B] = \det[\mathbf{f}_1, \mathbf{f}_2, \tilde{\mathbf{F}}_1, \tilde{\mathbf{F}}_2] \det B$. \square

If λ is neither an eigenvalue nor a resonance, so that the Jost solutions

$$\{\mathbf{f}_1(x, \lambda, \omega), \mathbf{f}_2(x, \lambda, \omega), \mathbf{g}_1(x, \lambda, \omega), \mathbf{g}_2(x, \lambda, \omega)\} \tag{4.42}$$

are linearly independent, we define:

$$G(x, y, \lambda, \omega) = -\alpha \sum_{j,k=1}^2 \left[\Theta(x-y)\mathbf{f}_j(x)\Gamma_{jk}(\lambda, \omega) \otimes \mathbf{g}_k^*(y) + \Theta(y-x)\mathbf{g}_j(x)\Gamma_{jk}(\lambda, \omega) \otimes \mathbf{f}_k^*(y) \right] \Delta(y, \lambda, \omega)^{-1}, \tag{4.43}$$

where Θ is the Heaviside step-function, the Jost solutions also depend on (λ, ω) (this is not explicitly indicated), the matrix $\Gamma(\lambda, \omega)$ is defined by

$$\Gamma(\lambda, \omega) = \frac{1}{\sqrt{|B_{21}|^2 + |B_{22}|^2}} \begin{bmatrix} |B_{22}| & |B_{21}|e^{-is} \\ |B_{21}|e^{is} & -|B_{22}| \end{bmatrix}, \tag{4.44}$$

so that $\det \Gamma = -1$; here $s \in \mathbb{R}$ is chosen so that

$$\sum_{j=1}^2 B_{2j}\Gamma_{j1} = B_{21}\Gamma_{11} + B_{22}\Gamma_{21} = \frac{B_{21}|B_{22}| + B_{22}|B_{21}|e^{is}}{\sqrt{|B_{21}|^2 + |B_{22}|^2}} = 0. \tag{4.45}$$

(This choice of Γ is justified later by the need to have appropriate estimates on $G(x, y, \lambda, \omega)$.) The matrix $\Delta(y, \lambda, \omega)$ in (4.43) is defined by

$$\Delta(y, \lambda, \omega) = \mathbf{f}_j(y, \lambda, \omega)\Gamma_{jk}(\lambda, \omega) \otimes \mathbf{g}_k^*(y, \lambda, \omega) - \mathbf{g}_j(y, \lambda, \omega)\Gamma_{jk}(\lambda, \omega) \otimes \mathbf{f}_k^*(y, \lambda, \omega). \tag{4.46}$$

Since $\det \Gamma \neq 0$, the matrix (4.46) is invertible as long as $\{\mathbf{f}_1, \mathbf{f}_2, \mathbf{g}_1, \mathbf{g}_2\}$ are linearly independent. Moreover,

$$\det \Delta(y, \lambda, \omega) = |E(\lambda, \omega)|^2. \tag{4.47}$$

The relation (4.47) follows from the following identity:

Lemma 4.18. *For any $u_j, v_j \in \mathbb{C}^N, 1 \leq j \leq N, A \in \mathbb{C}^{N \times N}$, one has*

$$\det \left(\sum_{j,k=1}^N u_j A_{jk} \otimes v_k^* \right) = \det A \det[u_1, \dots, u_N] \overline{\det[v_1, \dots, v_N]}. \tag{4.48}$$

Proof. If v_j are linearly dependent, the rank of the matrix in the left-hand side is smaller than N , and both sides in (4.48) vanish. Otherwise, the proof follows from computing the determinants of both sides of the identity

$$\left(\sum_{j,k=1}^N u_j A_{jk} \otimes v_k^* \right) ([v_1, \dots, v_N]^*)^{-1} = \left[\sum_{j=1}^N u_j A_{j1}, \dots, \sum_{j=1}^N u_j A_{jN} \right]. \quad \square$$

Applying Lemma 4.18 to (4.46), thus setting $[u_1, \dots, u_4] = [v_1, \dots, v_4] = [\mathbf{f}_1, \mathbf{f}_2, \mathbf{g}_1, \mathbf{g}_2]$ and $A = \begin{bmatrix} 0 & \Gamma \\ -\Gamma & 0 \end{bmatrix}$, one derives:

$$\det \Delta(y, \lambda, \omega) = (\det \Gamma(\lambda, \omega))^2 \left| \det[\mathbf{f}_1(y, \lambda, \omega), \mathbf{f}_2(y, \lambda, \omega), \mathbf{g}_1(y, \lambda, \omega), \mathbf{g}_2(y, \lambda, \omega)] \right|^2,$$

arriving at (4.47).

As follows from the definition, one has

$$-\alpha(\partial_x - \mathcal{M}(x, \lambda, \omega))G(x, y, \lambda, \omega) = \delta(x - y)I_4.$$

Remark 4.19. At this point, we need to recall that the limit of the Green function is not uniquely defined at the essential spectrum. Since the expression (4.43) has the asymptotics $\sim e^{i\xi x}$, $\xi \approx -i\lambda$ for $\lambda \in i\mathbb{R}$, $\text{Im } \lambda \gg 1$ (cf. (4.7) and our convention that ξ_1, ξ_2 are positive for $\lambda \in i\mathbb{R}$, $\text{Im } \lambda \gg 1$), we conclude that (4.43) will remain bounded for λ near $i\mathbb{R}$ with $\text{Re } \lambda < 0$; thus, (4.43) corresponds to the limit $G^-(x, y, \lambda, \omega) := G(x, y, \lambda - 0, \omega)$ of the Green function to the left of the upper branch of the essential spectrum (this is consistent with (4.8)). To define the limit on the right of the essential spectrum, one would need to interchange in the above considerations $\mathbf{f}_j \sim e^{i\xi_j x}$ and $\mathbf{F}_j \sim e^{-i\xi_j x}$, as well as \mathbf{g}_j and \mathbf{G}_j (this is assuming that $\text{Im } \lambda$ is large enough so that $\xi_j > 0$, hence $\mathbf{f}_j, \mathbf{F}_j$ with particular j oscillate as $x \rightarrow +\infty$).

Let us now find the bounds on $G(x, y, \lambda, \omega)$. Our goal is to show that (4.43) does not grow exponentially when x and or y go to infinity. For example, when $y \rightarrow +\infty$, the fastest growing term is $\tilde{\mathbf{F}}_2(y)$. We need to show that when (4.43) is written solely in terms of $\mathbf{f}_j, \tilde{\mathbf{F}}_k$, then in the combinations $\mathbf{f}_j(x) \otimes \tilde{\mathbf{F}}_k^*(y)$ one always has $x \geq y$, and moreover the coefficient at the term $\mathbf{f}_1(x) \otimes \tilde{\mathbf{F}}_2^*(y)$ vanishes (this is the only problematic term, when the decay of $\mathbf{f}_j(x)$ with $x \geq y$, $x \gg 1$, $y \gg 11$, does not compensate for the growth of $\tilde{\mathbf{F}}_k(y)$). We claim that the choice of Γ in (4.44) specifically guarantees this.

For $x \geq y$, we only need to consider the first term from (4.43):

$$\sum_{j,k} \mathbf{f}_j(x) \Gamma_{jk} \otimes \mathbf{g}_k^*(y), \quad x \geq y. \tag{4.49}$$

It is enough to consider the following two (intersecting) cases: (1) $x \geq y, y \leq 0$ and (2) $x \geq y, x \geq 0$. (In the intersection, one has $x \geq 0, y \leq 0$, hence (4.49) is uniformly bounded.)

Let us consider the case $x \geq y, x \geq 0$. By (4.40), the factor at $\mathbf{f}_1(x)$ in (4.49) is given by

$$\sum_k \Gamma_{1k} \mathbf{g}_k^*(y) = \sum_{j,k} (\mathbf{f}_j(y) A_{jk} \bar{\Gamma}_{1k} + \tilde{\mathbf{F}}_j(y) B_{jk} \bar{\Gamma}_{1k})^* = \sum_{j,k} (\mathbf{f}_j(y) A_{jk} \bar{\Gamma}_{1k})^* + \sum_k (\tilde{\mathbf{F}}_1(y) B_{1k} \bar{\Gamma}_{1k})^*; \tag{4.50}$$

in the last equality, we took into account (4.44) and (4.45):

$$\sum_k B_{2k} \bar{\Gamma}_{1k} = B_{21} \bar{\Gamma}_{11} + B_{22} \bar{\Gamma}_{12} = B_{21} \Gamma_{11} + B_{22} \Gamma_{21} = 0.$$

It follows that when we rewrite (4.49) in terms of $\mathbf{f}, \tilde{\mathbf{F}}$ only, then the only term which can become exponentially large for $x \geq y, x \geq 0$, namely $\mathbf{f}_1(x) \otimes \tilde{\mathbf{F}}_2^*(y)^*$, drops out! Hence, (4.49) is bounded by $C(\lambda, \omega) \langle y \rangle$ for $x \geq 0, x \geq y$. The linear growth in y may come from $\mathbf{f}_j(x) \otimes \tilde{\mathbf{F}}_j^*(y)^*$ when $0 \ll y \leq x$, whenever $\lambda \in i\mathbb{R}$ is near $i(1 \pm |\omega|)$, so that $\xi_j \approx 0$.

Let $x \geq y, y \leq 0$. By (4.41), the factor at $\mathbf{g}_1^*(y, \lambda, \omega)$ in (4.49) is given by

$$\sum_j \mathbf{f}_j(x) \Gamma_{j1} = \sum_{j,k} (\mathbf{g}_k(x) A_{kj} + \tilde{\mathbf{G}}_k(x) B_{kj}) \Gamma_{j1} = \sum_{j,k} \mathbf{g}_k(x) A_{kj} \Gamma_{j1} + \sum_j \tilde{\mathbf{G}}_1(x) B_{1j} \Gamma_{j1}; \tag{4.51}$$

in the last equality, we took into account that the coefficient at $\tilde{\mathbf{G}}_2(x) \otimes \mathbf{g}_1^*(y)$ is given by $B_{21} \Gamma_{11} + B_{22} \Gamma_{21} = 0$, by (4.45). Thus, when we rewrite (4.49) in terms of \mathbf{g} and $\tilde{\mathbf{G}}$, the coefficient at the term $\tilde{\mathbf{G}}_2(x) \otimes \mathbf{g}_1^*(y)$, the only one out of $\tilde{\mathbf{G}}_j(x) \otimes \mathbf{g}_k^*(y)$ which can be exponentially large for $x \geq y, y \rightarrow -\infty$, drops out. It follows that (4.49) is bounded by $C(\omega) \langle x \rangle$ for $y \leq 0, x \geq y$. The linear growth in x may come from $\tilde{\mathbf{G}}_j(x) \otimes \mathbf{g}_j(y)$ for $y \leq x \ll 0$ (when writing (4.49) as a linear combination of $\mathbf{g}_j \otimes \mathbf{g}_k^*, \tilde{\mathbf{G}}_j \otimes \mathbf{g}_k^*$, via the substitution (4.41)), whenever λ is near $i(1 \pm |\omega|)$ so that the corresponding ξ_j is near zero. By (4.14), as $|\lambda| \rightarrow \infty$, $\|\mathbf{f}_j(\cdot, \lambda, \omega)\|_{L^\infty}$ and $\|\mathbf{g}_j(\cdot, \lambda, \omega)\|_{L^\infty}$ are bounded by $c(\omega) < \infty$.

We summarize the cases $x \geq y, y \leq 0$ and $x \geq y, x \geq 0$: Thus, for some $c(\lambda, \omega) < \infty$,

$$\left\| \sum_{j,k} \Gamma_{jk} \mathbf{f}_j(x) \otimes \mathbf{g}_k^*(y) \right\|_{\text{End}(\mathbb{C}^4)} \leq c(\lambda, \omega) \min(\langle x \rangle, \langle y \rangle), \quad x \geq y. \tag{4.52}$$

The case $x \leq y$ follows from the above once we notice that $\Delta(-y, \lambda, \omega) = \beta \Delta(y, \lambda, \omega) \beta$ and then

$$G(-x, -y, \lambda, \omega) = -\beta G(x, y, \lambda, \omega) \beta;$$

we arrive at the same bound but now for $x \leq y$:

$$\left\| \sum_{j,k} \Gamma_{jk} \mathbf{g}_j(x) \otimes \mathbf{f}_k^*(y) \right\|_{\text{End}(\mathbb{C}^4)} \leq c(\lambda, \omega) \min(\langle x \rangle, \langle y \rangle), \quad x \leq y. \tag{4.53}$$

Let us study the contribution of the matrix $\Delta(y, \lambda, \omega)$ defined in (4.46). By (4.52) and (4.53), $\Delta(y, \lambda, \omega)$ satisfies

$$\|\Delta(y, \lambda, \omega)\|_{\text{End}(\mathbb{C}^4)} \leq c(\lambda, \omega) \langle y \rangle, \tag{4.54}$$

with the linear growth only for $x \approx \pm i(1 \pm \omega)$.

By (4.47) and (4.54), there is $C(\lambda, \omega) < \infty$ such that

$$\left\| \Delta(y, \lambda, \omega)^{-1} \right\|_{\text{End}(\mathbb{C}^4)} \leq C(\lambda, \omega) \langle y \rangle. \tag{4.55}$$

(Here, we need to argue that the minors of Δ cannot grow faster than $\langle y \rangle$; at most one of $\tilde{\mathbf{G}}_j(y) \otimes \mathbf{g}_j(y)^*$, $j = 1, 2$ can grow linearly at a given value of λ , hence, in the appropriate basis, only one element of Δ grows linearly while others are bounded uniformly in $y \in \mathbb{R}$.) Combining (4.52) and (4.53) with (4.55), we arrive at the bound (4.34).

Let us now study the behavior of $G(x, y, \lambda, \omega)$ for $\lambda \in i\mathbb{R}$, $|\lambda| \rightarrow \infty$. By Proposition 4.4, the Jost solutions $\mathbf{f}_j, \tilde{\mathbf{F}}_j, \mathbf{g}_j, \tilde{\mathbf{G}}_j$ are bounded uniformly in x as long as $|\lambda|$ is sufficiently large. By Lemma 4.10 and (4.47), for $\lambda \in i\mathbb{R}$, $|\lambda| \rightarrow \infty$, one has $|\det \Delta(y, \lambda, \omega)| \rightarrow 1$, while the components of $\Delta(y, \lambda, \omega)$ are uniformly bounded for $\lambda \rightarrow \pm i\infty$. It follows that the components of the matrix $G(x, y, \lambda, \omega)$ defined in (4.43) are bounded uniformly in x and y as long as $|\lambda|$ is sufficiently large.

Finally, the bounds (4.36) and (4.37) follow from the pointwise estimates (4.34) and (4.35) for Green’s function. This concludes the proof of Proposition 4.14. \square

5. Dispersive estimates for the semigroup

In this section, we develop set of dispersive estimates, which will be useful in the sequel for controlling the radiation portion of the perturbation.

5.1. Weighted decay estimates

We start with an estimate, which is typical in this context, namely $e^{t\mathbf{JL}(\omega)} P_c(\omega) : L^2 \rightarrow L^\infty(\langle x \rangle^{-3} L_t^2)$.

Proposition 5.1. *Let $\omega \in \Omega$. Then there exists $C < \infty$ such that for all $t > 0$, the following estimates hold:*

$$\begin{aligned} \sup_x \langle x \rangle^{-3} \left\| [e^{t\mathbf{JL}(\omega)} P_c(\omega) f](x) \right\|_{L_t^2} &\leq C \|f\|_{L_x^2}, \\ \left\| \int_{-\infty}^{\infty} e^{t\mathbf{JL}(\omega)} P_c(\omega) F(t, \cdot) dt \right\|_{L_x^2} &\leq C \|F\|_{(L_x^1)_x L_t^2}. \end{aligned}$$

Remark 5.2.

(i) The estimates in Proposition 5.1 can be upgraded to include derivatives. For example,

$$\sup_x \langle x \rangle^{-3} \left\| \partial_x [e^{t\mathbf{JL}} P_c(\omega) f](x) \right\|_{L_t^2} \leq C \|f\|_{H_x^1}.$$

Note that the last estimate presents a challenge, since $\partial_x e^{t\mathbf{JL}} \neq e^{t\mathbf{JL}} \partial_x$. Nevertheless, since

$$\mathbf{L}(\omega) = \mathbf{D}_m - \omega I_4 + \mathbf{W}(x, \omega),$$

with \mathbf{D}_m from (3.5), we may essentially commute the derivative with $e^{t\mathbf{JL}}$ modulo low order error terms, whence the result generalizes to include derivatives.

(ii) Interestingly, Proposition 5.1 fails for the free Dirac operator. This is due to an essentially resonant behavior of the free Dirac operator close to the edges of the continuous spectrum. We have an alternative estimate for the free Dirac operator in Lemma 5.4.

Proof of Proposition 5.1. Clearly, the two estimates in the claim of Proposition 5.1 are dual to each other, so it suffices to establish the first one.

Pick an even function $\chi \in C_{\text{comp}}^\infty(\mathbb{R})$ such that

$$\text{supp } \chi \subset [-4, 4], \quad \chi(\Lambda) = 1 \quad \text{for } |\Lambda| \leq 3. \tag{5.1}$$

Decompose the evolution into two pieces:

$$e^{t\mathbf{JL}} P_c(\omega) f = \chi(i\mathbf{JL}) e^{t\mathbf{JL}} P_c(\omega) f + (1 - \chi(i\mathbf{JL})) e^{t\mathbf{JL}} P_c(\omega) f,$$

with the terms in the right-hand side defined in view of (4.2) as the inverse Fourier transforms in time of the functions

$$-\chi(\Lambda) ([R_{\mathbf{JL}}^+(i\Lambda) - R_{\mathbf{JL}}^-(i\Lambda)]f)(x), \quad -(1 - \chi(\Lambda)) ([R_{\mathbf{JL}}^+(i\Lambda) - R_{\mathbf{JL}}^-(i\Lambda)]f)(x). \tag{5.2}$$

The required estimate will follow from

$$\sup_x \|(1 - \chi(i\mathbf{JL})) e^{t\mathbf{JL}} P_c(\omega) f\|_{L_t^2} \leq C \|f\|_{L_x^2}, \tag{5.3}$$

$$\sup_x \langle x \rangle^{-3} \|\chi(i\mathbf{JL}) e^{t\mathbf{JL}} P_c(\omega) f\|_{L_t^2} \leq C \|f\|_{L_x^2}. \tag{5.4}$$

Using the Fourier transforms (5.2), we see that (5.3) will follow from

$$\sup_x \|(1 - \chi(\Lambda)) R_{\mathbf{JL}}^\pm(i\Lambda) f(x)\|_{L_\Lambda^2} \leq C \|f\|_{L_x^2}. \tag{5.5}$$

Similarly, (5.4) will follow from

$$\sup_x \langle x \rangle^{-3} \|\chi(\Lambda) R_{\mathbf{JL}}^\pm(i\Lambda) f(x)\|_{L_\Lambda^2} \leq C \|f\|_{L_x^2}. \tag{5.6}$$

We now prove (5.5) and (5.6).

Proof of (5.5). For brevity, we denote

$$R_{\mathbf{W}}(\Lambda) := (\mathbf{D}_m - \omega I_4 - \Lambda \mathbf{J}^{-1} + \mathbf{W})^{-1}.$$

From the resolvent identity, we have $R_{\mathbf{W}} = R_0 - R_{\mathbf{W}} \mathbf{W} R_0 = R_0 - R_0 \mathbf{W} R_{\mathbf{W}}$, whence the following Born expansion holds:

$$R_{\mathbf{W}} = R_0 - R_0 \mathbf{W} R_0 + R_0 \mathbf{W} R_{\mathbf{W}} \mathbf{W} R_0. \tag{5.7}$$

Observe that $R_0 = \begin{bmatrix} (D_m - (\omega + \Lambda) I_2)^{-1} & 0 \\ 0 & (D_m - (\omega - \Lambda) I_2)^{-1} \end{bmatrix}$. The restrictions imposed by the cut-off $(1 - \chi)$ (cf. (5.1)) implies that $|\omega \pm \Lambda| > 3$. It follows that it is enough to show that

$$\sup_x \int_3^\infty |(D_m - \mu I_2)^{-1} f(x)|^2 d\mu \leq C \|f\|_{L_x^2}^2; \tag{5.8}$$

$$\sup_x \int_3^\infty |(D_m - \mu I_2)^{-1} W_v (D_m - \mu I_2)^{-1} f(x)|^2 d\mu \leq C \|\mathbf{W}\|_{L_x^1}^2 \|f\|_{L_x^2}^2; \tag{5.9}$$

$$\sup_x \int_3^\infty |(D_m - \mu I_2)^{-1} W_v R_{\mathbf{W}} W_v (D_m - \mu I_2)^{-1} f(x)|^2 d\mu \leq C \|\langle x \rangle^\alpha \mathbf{W}\|_{L_x^2}^2 \|f\|_{L_x^2}^2. \tag{5.10}$$

Above, $\alpha > 3/2$ and W_v is either of the potentials W_1, W_0 . Similar estimates were shown in [29, Section VIII], but we provide the details here for completeness. Note that

$$(D_m - \mu I_2)^{-1} = (1 - \partial_x^2 - \mu^2)^{-1} \begin{pmatrix} 1 + \mu & \partial_x \\ -\partial_x & \mu - 1 \end{pmatrix}.$$

Thus, setting $\mu = \sqrt{k^2 + 1}$, the operator $(D_m - \mu I_2)^{-1}$ is represented as a linear combination of operators with the following kernels:

$$e^{\pm ik|x|} \operatorname{sgn}(x), \quad \frac{e^{\pm ik|x|}}{k}, \quad \frac{e^{\pm ik|x|} \sqrt{k^2 + 1}}{k}.$$

Clearly, for the purposes of showing (5.8), (5.9), (5.10), it is enough to consider the operator with kernel $e^{\pm ik|x|} \operatorname{sgn}(x)$. For the proof of (5.8), we have by Plancherel’s

$$\begin{aligned} & \sup_{x \in \mathbb{R}} \int_{\sqrt{8}}^{\infty} \left| \int e^{\pm ik|x-y|} \operatorname{sgn}(x-y) f(y) dy \right|^2 \frac{k dk}{\sqrt{k^2 + 1}} \\ & \leq 2 \sup_{x \in \mathbb{R}} \int_{\sqrt{8}}^{\infty} \left\{ \left| \int_{-\infty}^x e^{\mp iky} f(y) dy \right|^2 + \left| \int_x^{\infty} e^{\pm iky} f(y) dy \right|^2 \right\} dk \leq 4 \|f\|_{L^2}^2. \end{aligned}$$

Similarly, for (5.9) we have (by Minkowski’s)

$$\begin{aligned} & \sup_x \int_{\sqrt{8}}^{\infty} \left| \int e^{\pm ik|x-y|} \operatorname{sgn}(x-y) \mathbf{W}(y) [R_0(\sqrt{1+k^2})f](y) dy \right|^2 \frac{k dk}{\sqrt{k^2 + 1}} \\ & \leq \int_{\sqrt{8}}^{\infty} \left| \int |\mathbf{W}(y)| \left| [R_0(\sqrt{1+k^2})f](y) \right| dy \right|^2 dk \\ & \leq \left(\int |\mathbf{W}(y)| \left(\int_{\sqrt{8}}^{\infty} |[R_0(\sqrt{1+k^2})f](y)|^2 dk \right)^{1/2} dy \right)^2 \\ & \leq \|\mathbf{W}\|_{L^1}^2 \sup_y \int_{\sqrt{8}}^{\infty} |[R_0(\sqrt{1+k^2})f](y)|^2 dk \leq C \|\mathbf{W}\|_{L^1}^2 \|f\|_{L^2}^2. \end{aligned}$$

This shows (5.9). Finally, for (5.10), we estimate

$$\begin{aligned} & \sup_x \int_{\sqrt{8}}^{\infty} \left| \int e^{\pm ik|x-y|} \operatorname{sgn}(x-y) \mathbf{W}(y) R_{\mathbf{W}}[\mathbf{W}R_0 f](y) dy \right|^2 dk \\ & \leq \|\langle y \rangle^3 \mathbf{W}(y)\|_{L^2}^2 \int_{\sqrt{8}}^{\infty} \|\langle y \rangle^{-3} R_{\mathbf{W}}(y)^{-3} [\langle y \rangle^3 \mathbf{W}(y) [R_0(\sqrt{1+k^2})f](y)]\|_{L^2_y}^2 dk \\ & \leq \|\langle y \rangle^3 \mathbf{W}(y)\|_{L^2}^4 \|R_{\mathbf{W}}\|_{(L^2_3)_x \rightarrow (L^2_{-3})_x}^2 \sup_y \int_{\sqrt{8}}^{\infty} \left| R_0(\sqrt{1+k^2})f(y) \right|^2 dk \\ & \leq C \|\langle y \rangle^3 \mathbf{W}(y)\|_{L^2}^4 \|f\|_{L^2_x}^2. \end{aligned}$$

In the last estimate, we have used the estimates from Proposition 4.14 which are uniform for large Λ (for large values of the spectral parameter $\Lambda > \sqrt{8}$), $R_{\mathbf{W}} : L^2_3(\mathbb{R}) \rightarrow L^2_{-3}(\mathbb{R})$.

Proof of (5.6). The statement for low frequencies follows from the following result:

Lemma 5.3. Let $C_0(\mathbb{R})$ be the space of continuous, compactly supported functions. Define $A : C_0(\mathbb{R}) \rightarrow C(\mathbb{R} \times \mathbb{R})$ by

$$u \mapsto Au(x, \Lambda) = \chi(\Lambda) \int_{\mathbb{R}} G(x, y, i\Lambda, \omega)u(y) dy. \tag{5.11}$$

Then A extends to a continuous operator $L^2(\mathbb{R}) \rightarrow L^\infty_{\text{loc}}(\mathbb{R}, L^2_\Lambda(\mathbb{R}))$, and moreover there is $C < \infty$ such that

$$\sup_x \langle x \rangle^{-3} \|Au(x, \cdot)\|_{L^2_\Lambda} \leq C \|u\|_{L^2}. \tag{5.12}$$

Proof. Let $u \in L^2(\mathbb{R}, \mathbb{C}^4)$. Without loss of generality, we assume that $\text{supp } u \subset \mathbb{R}_+$, so that in (4.43) we have $y \geq 0$.

The case $x \geq 0$. We use the expression (4.43) for $G(x, y, i\Lambda, \omega)$; expressing in (4.43) the Jost solutions \mathbf{g}_j in terms of \mathbf{f}_j and $\tilde{\mathbf{F}}_j$, we see that it suffices to check that the expressions

$$\int_0^\infty \Theta(\pm(x-y))\mathbf{f}_j(x)\mathbf{f}_k^*(y)u(y) dy, \quad \int_0^\infty \Theta(y-x)\tilde{\mathbf{F}}_j(x)\mathbf{f}_k^*(y)u(y) dy, \quad \int_0^\infty \Theta(x-y)\mathbf{f}_j(x)\tilde{\mathbf{F}}_k^*(y)u(y) dy, \tag{5.13}$$

with $j, k = 1, 2$, are bounded in L^2 as functions of Λ , with an appropriate bound on the growth with x . Above, we omitted the weight $\chi(\Lambda)$ present in (5.11); this weight will become important when we will integrate by parts.

In (5.13) and in the rest of the proof, the Jost solutions are evaluated at $\lambda = i\Lambda$ and ω , which we usually do not indicate explicitly to shorten the notations. The first two terms in (5.13) are analyzed similarly; the more difficult being the second one, so we focus on it.

- Assume that $\mathbf{f}_k(y, i\Lambda, \omega)$ is exponentially decaying, so that

$$\mathbf{f}_k(y, i\Lambda, \omega) \sim e^{-\kappa_k y}, \quad y \gg 1,$$

with $\kappa_k > 0$ (cf. (4.12)).

When $\tilde{\mathbf{F}}_j(x, i\Lambda, \omega)$ remains bounded or grows linearly in x for $x \gg 1$,

$$\left| \int_0^\infty \tilde{\mathbf{F}}_j(x)\mathbf{f}_k^*(y)u(y) dy \right| \leq C \langle x \rangle \int_0^\infty |\mathbf{f}_k(y)||u(y)| dy \leq C \langle x \rangle \|\Theta(\cdot)\mathbf{f}_k\| \|u\| \leq \frac{C \langle x \rangle}{\sqrt{\kappa_k}} \|u\|.$$

Note that $\kappa_k^{-1/2}$ is L^2 in Λ near the thresholds $\Lambda = \pm(1 \pm \omega)$.

When $\tilde{\mathbf{F}}_j(x, i\Lambda, \omega)$ is exponentially growing, by Lemma 4.15, we have $|\tilde{\mathbf{F}}_j(x)| \leq C(\Lambda, \omega)\langle x \rangle e^{\kappa_j x}$ for $x \geq 0$, and moreover we only need to consider terms with $\kappa_j \leq \kappa_k$ due to our construction of G in Proposition 4.14 (the term $\tilde{\mathbf{F}}_2(x)\mathbf{f}_1^*(y)$ is absent in the expansion of $G(x, y)$ over $\mathbf{f}_j(x)\mathbf{f}_k^*(y)$, $\tilde{\mathbf{F}}_j(x)\mathbf{f}_k^*(y)$, and $\mathbf{f}_j(x)\tilde{\mathbf{F}}_k^*(y)$), and with $C(\Lambda, \omega)$ locally bounded in Λ and ω , with $\limsup_{\Lambda \rightarrow \pm\infty} C(\Lambda, \omega) \leq C(\omega) < \infty$. Then, again,

$$\left| \int_0^\infty \Theta(y-x)\tilde{\mathbf{F}}_j(x)\mathbf{f}_k^*(y)u(y) dy \right| \leq C \langle x \rangle \int_x^\infty e^{\kappa_j x} e^{-\kappa_k y} |u(y)| dy \leq \frac{C \langle x \rangle}{\sqrt{\kappa_k}} \|u\|.$$

- Assume that $\mathbf{f}_k(y, i\Lambda, \omega) \sim e^{i\xi_k y}$ is oscillating:

$$|\Lambda \pm \omega| > 1, \quad \xi_k(i\Lambda, \omega) = \sqrt{(\Lambda \pm \omega)^2 - 1} > 0. \tag{5.14}$$

(According to the construction of the Green function, since \mathbf{f}_k is oscillating, we only need to consider the terms in (5.13) with $\tilde{\mathbf{F}}_j(x)$ also oscillating: $\xi_j > 0$.) In this case, the integration in spatial variables becomes possible after integrating by parts with the aid of the operator $L_\Lambda = \frac{1}{i(y-z)}\partial_\Lambda$; we only give a sketch, substituting the Jost solutions by their asymptotic behavior $\mathbf{f}_k(x) \sim e^{i\xi_k x}$ and $\tilde{\mathbf{F}}_j(x) \sim e^{-i\xi_j x} + \frac{e^{i\xi_j x} - e^{-i\xi_j x}}{2i\xi_j}$ (cf. (4.38)). Then the integration by parts yields

$$\begin{aligned}
 & \left| \int_{\mathbb{R}} \chi(\Lambda) d\Lambda \int_{\mathbb{R} \times \mathbb{R}} |\tilde{\mathbf{F}}_j(x)|^2 \overline{\mathbf{f}_k^*(z)u(z)} \mathbf{f}_k^*(y)u(y) dy dz \right| \\
 &= \left| \int_{\mathbb{R}} \chi(\Lambda) d\Lambda \int_{\mathbb{R} \times \mathbb{R}} |\tilde{\mathbf{F}}_j(x)|^2 L_\Lambda^2 \left(\overline{\mathbf{f}_k^*(z)u(z)} \mathbf{f}_k^*(y)u(y) \right) dy dz \right| \\
 &\leq \langle x \rangle^2 \int_{\mathbb{R}} \chi(\Lambda) d\Lambda \int_{\mathbb{R} \times \mathbb{R}} \frac{C|u(y)||u(z)| dy dz}{1 + |\mu_j(x, \Lambda, \omega)^{-1}(y - z)\partial_\Lambda \xi_k|^2} \leq C \langle x \rangle^2 \int_{\mathbb{R}} \chi(\Lambda) d\Lambda \frac{\mu_j(x, \Lambda, \omega) \|u\|^2}{|\partial_\Lambda \xi_k|}. \tag{5.15}
 \end{aligned}$$

Above,

$$\mu_j(x, \Lambda, \omega) := C \max \left(1, |x| |\partial_\Lambda \xi_j|, \frac{|\partial_\Lambda^2 \xi_j|}{|\partial_\Lambda \xi_j|} \right) \tag{5.16}$$

is the bound on the contribution of ∂_Λ during the integration by parts (the last term in (5.16) is the contribution from the derivative falling onto $\partial_\Lambda \xi_j$ during the second integration by parts). In the last inequality in (5.15), we used the Schur test. Due to (5.14), one has

$$\partial_\Lambda \xi_j = \frac{\Lambda \pm \omega}{\xi_j}, \quad |\partial_\Lambda^2 \xi_j| \leq \frac{C \langle \Lambda \rangle^2}{\xi_j^3};$$

hence, (5.16) can be continued as follows:

$$\mu_j(x, \Lambda, \omega) = C \max \left(1, |x| |\partial_\Lambda \xi_j|, \frac{|\partial_\Lambda^2 \xi_j|}{|\partial_\Lambda \xi_j|^2} \right) \leq C \max \left(1, \frac{\langle x \rangle}{\xi_j} \right).$$

It follows that

$$\frac{\mu_j(x, \Lambda, \omega)}{|\partial_\Lambda \xi_k|} \leq \frac{C \langle x \rangle}{\xi_k(i \Lambda, \omega)}$$

is locally integrable in $\Lambda \in \text{supp } \chi$ (and such that $|\Lambda \pm \omega| > 1$), and moreover $\langle x \rangle^{-3} \int \langle x \rangle^2 \frac{\mu_j(x, \Lambda, \omega)}{|\partial_\Lambda \xi_k|} \chi(\Lambda) d\Lambda$ is bounded uniformly in x . The factor $\langle x \rangle^2$ under the integral comes from the bound $|\tilde{\mathbf{F}}_j(x, \lambda, \omega)| \leq C \langle x \rangle$ which remains valid uniformly in $\xi_j > 0$ when $\xi_j \rightarrow 0+$ (cf. Lemma 4.15). This leads to (5.12).

Let us analyze the last term in (5.13). When $\tilde{\mathbf{F}}_k(y)$ is oscillating, we use the same consideration as above, in the case when $\mathbf{f}_k(y)$ was oscillating. Let us consider the situation when $\tilde{\mathbf{F}}_k(y)$ is exponentially growing as $y \rightarrow +\infty$. Since this growth is compensated by the decay of $\Theta(x - y)\mathbf{f}_j(x)$ due to the choice of $B_{jk}(\lambda, \omega)$ in (4.44) (as we mentioned above, the construction of G is such that we only need to treat terms with $\kappa_k \leq \kappa_j$), it suffices to consider the terms $\Theta(x - y)\mathbf{f}_j(x)\tilde{\mathbf{F}}_k^*(y)$ which are bounded by $\Theta(x - y)\langle x \rangle e^{-\kappa_j|x|} e^{\kappa_k|y|}$, with $\kappa_j \geq \kappa_k$. We have:

$$\langle x \rangle \int \Theta(x - y) e^{-\kappa_j|x|} e^{\kappa_k|y|} |u(y)| dy \leq C \langle x \rangle \int_0^x |u(y)| dy \leq C \langle x \rangle^{3/2} \|u\|,$$

which immediately leads to (5.12).

The case $x \leq 0$. This case is in fact much simpler. In this case, from (4.43), we only need to consider the contribution from $\sum_{j,k=1}^2 \mathbf{g}_j(x)\Gamma_{jk}\mathbf{f}_k^*(y)$; we need to prove that the expressions

$$\int_{\mathbb{R}_+} \mathbf{g}_j(x)\Gamma_{jk}\mathbf{f}_k^*(y)u(y) dy,$$

with $j, k = 1, 2$, are L^2 -bounded in Λ , for $\Lambda \in \text{supp } \chi$. Since $\mathbf{g}_j(x)$ are bounded for $x \leq 0$, the proof follows the lines of our argument for the case $x \geq 0$, except that we do not need to worry whether the decay of $\mathbf{f}_k(x)$ compensates the growth $\mathbf{g}_j(x)$ since the latter terms are bounded for $x \leq 0$. This finishes the proof. \square

This completes the proof of Proposition 5.1. \square

Next, we state and prove the estimate for the “free” Dirac operator, which is reminiscent of Proposition 5.1. As we have pointed out before, Proposition 5.1 does not hold for \mathbf{D}_m , unless one adds a derivative correction that takes care of the low frequency component of f .

Lemma 5.4. *We have the following estimate for the evolution of the “free” Dirac operator:*

$$\sup_x \|e^{t\mathbf{JL}^0} f\|_{L_t^2} \leq \|Mf\|_{L_x^2}, \tag{5.17}$$

$$\| \int e^{t\mathbf{JL}^0} F(t, \cdot) dt \| \leq C \|MF\|_{L_x^1 L_t^2}, \tag{5.18}$$

where $M = \sqrt{\langle \nabla \rangle} / |\nabla|$ or more precisely $\widehat{Mg}(\xi) = \frac{(1+\xi^2)^{1/4}}{|\xi|^{1/2}} \hat{g}(\xi)$. In addition, by a simple duality argument, there is also

$$\| \int e^{t\mathbf{JL}^0} F(t, \cdot) dt \|_{L_x^2} \leq C \|MF\|_{L_x^1 L_t^2}. \tag{5.19}$$

Proof. Clearly, (5.18) is just a dual to (5.17), so we concentrate on (5.17). Due to the block-diagonal structure of \mathbf{D}_m , the problem $iu_t = \mathbf{D}_m u$ reduces to the following linear system:

$$i \partial_t h = D_m h, \quad h|_{t=0} = h^0,$$

which in the components of $h \in \mathbb{C}^2$ takes the following form:

$$\begin{cases} i \partial_t h_1 = h_1 + \partial_x h_2, \\ i \partial_t h_2 = -\partial_x h_1 - h_2, \\ h_1(0) = h_1^0, \quad h_2(0) = h_2^0. \end{cases}$$

It follows that h_1, h_2 both satisfy the Klein–Gordon equation $\partial_{tt} h_{1,2} - \partial_{xx} h_{1,2} + h_{1,2} = 0$ with the corresponding initial data. Thus, (5.17) reduces to

$$\sup_x \|e^{it\langle \nabla \rangle} f\|_{L_t^2} \leq C \|Mf\|_{L^2},$$

where $\widehat{\langle \nabla \rangle} g(\xi) = \sqrt{1 + \xi^2} \hat{g}(\xi)$. Changing the variables $\kappa = \text{sgn}(\xi) \sqrt{1 + \xi^2}$ and using Plancherel’s theorem, we have:

$$\begin{aligned} \|e^{it\langle \nabla \rangle} f\|_{L_t^2}^2 &= \int \left| \int e^{it\sqrt{1+\xi^2}} \hat{f}(\xi) e^{i\xi x} d\xi \right|^2 dt \\ &= \int \left| \int_{|\kappa|>1} e^{it\kappa} \hat{f}(\sqrt{\kappa^2-1}) e^{ix\sqrt{\kappa^2-1}} \frac{\kappa d\kappa}{\sqrt{\kappa^2-1}} \right|^2 dt \\ &= \int_{|\kappa|>1} \frac{|\hat{f}(\sqrt{\kappa^2-1})|^2 \kappa^2 d\kappa}{\kappa^2-1} = \int \frac{|\hat{f}(\xi)|^2 \sqrt{1+\xi^2}}{|\xi|} d\xi = \|Mf\|_{L^2}^2. \quad \square \end{aligned}$$

Note that it becomes clear in the proof that the term $\|e^{it\langle \nabla \rangle} f\|_{L_t^2}$ is actually a constant in x . Thus, adding weights (as in Proposition 5.1) would not have salvaged a statement in Proposition 5.1, if we insist on having $\|f\|_{L_x^2}$ on the right hand side. One indeed needs to have instead $\|Mf\|_{L^2}$ as we have established above.

Next, we present an estimate for the retarded term in the Duhamel representation, in the spirit of Proposition 5.1.

Lemma 5.5. *Let $\omega \in \Omega$. There exists $C < \infty$ so that*

$$\sup_x \langle x \rangle^{-3} \left\| \int_0^t e^{-(t-\tau)\mathbf{JL}} P_c(\omega) F(\tau, \cdot) d\tau \right\|_{L_t^2} \leq C \|F\|_{(L^{\frac{1}{3}})_x L_t^2}. \tag{5.20}$$

Proof. It is well-known that these type of estimates are essentially dual estimates to the one presented in Proposition 5.1. In fact, recall that from Proposition 5.1,

$$\left\| \int_0^\infty e^{\tau \mathbf{JL}} P_c(\omega) F(\tau, \cdot) d\tau \right\|_{L_x^2} \leq C \|F\|_{(L^1_x) L_t^2}.$$

Thus, if one deals with the related quantity $\int_0^\infty e^{-(t-\tau)\mathbf{JL}} P_c(\omega) F(\tau, \cdot) d\tau$, we have, by virtue of Proposition 5.1 and its dual estimate,

$$\begin{aligned} \left\| \langle x \rangle^{-3} \int_0^\infty e^{-(t-\tau)\mathbf{JL}} P_c(\omega) F(\tau, \cdot) d\tau \right\|_{L_x^\infty L_t^2} &= \left\| \langle x \rangle^{-3} e^{-t\mathbf{JL}} \int_0^\infty e^{\tau \mathbf{JL}} P_c(\omega) F(\tau, \cdot) d\tau \right\|_{L_x^\infty L_t^2} \\ &\leq C \left\| \int_0^\infty e^{\tau \mathbf{JL}} P_c(\omega) F(\tau, \cdot) d\tau \right\|_{L_x^2} \leq C \|F\|_{(L^1_x) L_t^2}. \end{aligned}$$

However, as one observes quickly, we have to deal with \int_0^t in the retarded term in the Duhamel representation, instead of \int_0^∞ in our previous consideration. This is a non-trivial issue, which has been resolved in the literature, see [24, Lemma 11] and [29, Lemma 2]. In short, these results allows one to write for $F(t, x) = g_1(t)g_2(x)$,

$$\begin{aligned} U(t, \cdot) &= 2 \int_0^t e^{(t-\tau)\mathbf{JL}} P_c(\omega) F(\tau, \cdot) d\tau + \left(\int_{-\infty}^0 - \int_0^\infty \right) e^{(t-\tau)\mathbf{JL}} P_c(\omega) F(\tau, \cdot) d\tau, \\ U(t, x) &= \frac{i}{\sqrt{2\pi}} \int_{-\infty}^\infty e^{-it\Lambda} \check{g}_1(\Lambda) \left([R_{\mathbf{JL}}^+(i\Lambda) + R_{\mathbf{JL}}^-(i\Lambda)] g_2 \right) (x) d\Lambda. \end{aligned}$$

Since we have already shown the estimates for the term $\int_0^\infty \dots$ (and the estimates for $\int_{-\infty}^0 \dots$ are similar), it remains to show the appropriate estimates for U . By the Plancherel theorem in the t -variable,

$$\begin{aligned} \|U(t, \cdot)\|_{(L_{-3}^\infty)_x L_t^2} &= \left\| \langle x \rangle^{-3} \check{g}_1(\Lambda) [R_{\mathbf{JL}}^+(i\Lambda) + R_{\mathbf{JL}}^-(i\Lambda)] g_2 \right\|_{L_\Lambda^2} \Big\|_{L_x^\infty} \\ &\leq C \check{g}_1 \Big\|_{L_\Lambda^2} \sup_{\Lambda \in \mathbb{R}} \|R_{\mathbf{JL}}^\pm(i\Lambda)\|_{L^1_{-3} \rightarrow L_{-3}^\infty} \|g_2\|_{(L^1_x)} \leq C \|g_1\|_{L_t^2} \|g_2\|_{(L^1_x)}. \end{aligned}$$

All in all, we have shown the required estimate (5.20) for the case $F = g_1(t)g_2(x)$. Note however that the domain space $(L^1_x) L_t^2$ may be embedded in the bigger space $(\mathcal{M}_3)_x L_t^2$, where \mathcal{M}_3 is the space of Borel measures with the weight $\langle x \rangle^3$. By the Krein–Milman theorem, elements of this space may be represented as weak* limits of linear combinations of Dirac masses of the form $\delta(x - a)g(t)$. Thus, to show bounds of the form $T : (\mathcal{M}_3)_x L_t^2 \rightarrow Y$ for any linear operator T and Banach space Y , it suffices to prove such an estimate for elements $F = g_2(x)g_1(t)$, with $g_2 \in \mathcal{M}_3, g_1 \in L^2$ as we have done above. \square

5.2. Further linear estimates for $e^{t\mathbf{JL}}$

We will now state and derive the Strichartz estimates.

Definition 5.6. We say that a pair (q, r) is Strichartz-admissible (for the Dirac equation in one spatial dimension), if

$$q \geq 2, \quad r \geq 2, \quad \frac{2}{q} + \frac{1}{r} \leq \frac{1}{2}.$$

Equivalently, the admissible set is a triangle in the $(\frac{1}{q}, \frac{1}{r})$ plane, with endpoints corresponding to $(q, r) = (4, \infty)$ and $(q, r) = (\infty, 2)$.

In view of the representation of the Strichartz-admissible set as a triangle in the $(\frac{1}{q}, \frac{1}{r})$ coordinates, we will state the estimates only at the vertices, with the estimates in the interior of the triangle obtained by interpolation.

Next, before we can state our Strichartz type estimates, we need a variant of the well-known Christ–Kiselev lemma, an abstract result which allows one to pass between estimates for dual operators and retarded terms in the Duhamel representation. We state a version which is due to Smith and Sogge [32].

Lemma 5.7. *Let X, Y be Banach spaces and $\mathcal{K} : L^p(\mathbb{R}; X) \rightarrow L^q(\mathbb{R}, Y)$ be a bounded linear operator such that $\mathcal{K}f(t) = \int_{-\infty}^{\infty} K(t, s)f(s) ds$. Then the operator*

$$\tilde{\mathcal{K}}f(t) = \int_0^t K(t, s)f(s) ds \tag{5.21}$$

is bounded from $L^p(\mathbb{R}; X)$ to $L^q(\mathbb{R}, Y)$, provided that $p < q$. Moreover, there is $C_{p,q} > 0$ such that

$$\|\tilde{\mathcal{K}}\|_{L^p(\mathbb{R}; X) \rightarrow L^q(\mathbb{R}, Y)} \leq C_{p,q} \|\mathcal{K}\|_{L^p(\mathbb{R}; X) \rightarrow L^q(\mathbb{R}, Y)}.$$

Lemma 5.8. *Let (q, r) be a Strichartz-admissible pair. Then, for any $\epsilon > 0$ and $s \geq 0$, there is $C_\epsilon < \infty$ so that*

$$\|e^{t\mathbf{JL}}P_c(\omega)f\|_{L_t^q L_x^\infty} \leq C\|f\|_{H^{3/4+\epsilon}}, \tag{5.22}$$

$$\|e^{t\mathbf{JL}}P_c(\omega)f\|_{L_t^\infty H_x^s} \leq C\|f\|_{H^s}, \tag{5.23}$$

$$\left\| \int_{-\infty}^{\infty} e^{\tau\mathbf{JL}}P_c(\omega)F(\tau, \cdot) \right\|_{L_t^\infty H_x^1 \cap L_t^q L_x^r} \leq \|F\|_{L_t^1 H_x^1}, \tag{5.24}$$

$$\left\| \int_0^t e^{(t-\tau)\mathbf{JL}}P_c(\omega)F(\tau, \cdot) \right\|_{L_t^\infty H_x^1 \cap L_t^q L_x^r} \leq \|F\|_{L_t^1 H_x^1}. \tag{5.25}$$

Proof. We start with the estimates (5.22) and (5.23). Let us note that we can easily upgrade (5.22) to add derivatives on the evolution. An interpolation between these two estimates then yields (cf. (5.27) below for the free Dirac case):

$$\|e^{t\mathbf{JL}}P_c(\omega)f\|_{L_t^q W_x^{s,r}} \leq C_\epsilon \|f\|_{H^{s+\frac{1}{2}+\frac{1}{q}-\frac{1}{r}+\epsilon}}, \tag{5.26}$$

for $s \geq 0$ and for all Strichartz-admissible pairs (q, r) .

The proof of (5.25) is based on an application of the dual to (5.26) and Lemma 5.7. Thus, it remains to show (5.22) and (5.23). The approach follows what has become standard in recent years: we employ the available results for the “free” Dirac operator, in addition to the weighted decay estimates that we have proved in the previous section, namely Proposition 5.1 and Lemma 5.5. In fact, we follow closely the approach in [29, Lemma 4].

Let us recall first the estimates for the free Dirac operator. Let us prove the Strichartz estimates for e^{itD_m} in the form (5.22), (5.23), (5.25). The corresponding linear equations

$$i\partial_t h_1 = h_1 + \partial_x h_2, \quad i\partial_t h_2 = -\partial_x h_1 - h_2$$

reduce to the Klein–Gordon equation for each component h_1, h_2 , as we have shown in the proof of Lemma 5.4. Thus, the “free” Dirac estimates follow from the respective estimates for the Klein–Gordon equation, which can be found in the recent work of Nakamura–Ozawa, [26, Lemma 2.1] (where one takes $\theta = 1, \Lambda = 3/2, n = 1$). These estimates read as follows: for every $\epsilon > 0$,

$$\|e^{-itD_m}f\|_{L_t^q W_x^{s,r}} \leq C_\epsilon \|f\|_{H^{s+\frac{1}{2}+\frac{1}{q}-\frac{1}{r}+\epsilon}}. \tag{5.27}$$

These are of course the variants of the estimates (5.22) and (5.23); the estimate (5.25) holds in a similar manner for the free Dirac case. One important improvement of (5.27), which is implicit in [26],³ concerns the low frequency component of f . Namely, for the particular case $q = 4, r = \infty$, we have:

³ This is the estimate (2.15) in [26], which holds with the homogeneous Besov spaces version.

$$\|e^{-itD_m} f\|_{L_t^4 L_x^\infty} \leq C_\epsilon \|\partial_x\|^{3/4} f\|_{H^\epsilon}. \tag{5.28}$$

Let us now consider $\mathbf{JL} = \mathbf{J}(\mathbf{L}_0 + \mathbf{W})$, with a potential \mathbf{W} of Schwartz class. We may write the perturbed evolution in terms of the free evolution as follows:

$$e^{t\mathbf{JL}} f = e^{t\mathbf{JL}_0} f + \int_0^t e^{(t-s)\mathbf{JL}_0} \mathbf{JW} e^{s\mathbf{JL}} f ds.$$

We now have to deal with the two endpoint cases of Strichartz pairs: $(q, r) = (4, \infty)$ and $(q, r) = (\infty, 2)$. We only present the first case, the second being similar. To that end, let $\mathbf{W}(x) = V_1(x)V_2(x)$, with $V_1(x) = e^{-\delta_\Omega(x)}$ and $V_2(x) = e^{\delta_\Omega(x)}\mathbf{W}(x)$, with

$$\delta_\Omega = \inf_{\omega \in \Omega} \delta_\omega = \inf_{\omega \in \Omega} \sqrt{1 - \omega^2} > 0 \tag{5.29}$$

so that $V_2(x)$ is also exponentially decaying (cf. (3.8)). For $f \in H^{\frac{3}{4}+\epsilon}$, we have:

$$\begin{aligned} \|e^{t\mathbf{JL}} P_c(\omega) f\|_{L_t^4 L_x^\infty} &\leq \|e^{t\mathbf{JL}_0} f\|_{L_t^4 L_x^\infty} + \left\| \int_0^t e^{(t-s)\mathbf{JL}_0} \mathbf{J}V_1 V_2 e^{s\mathbf{JL}} P_c(\omega) f ds \right\|_{L_t^4 L_x^\infty} \\ &\leq C_\epsilon \|f\|_{H_x^{\frac{3}{4}+\epsilon}} + \left\| \int_0^t e^{(t-s)\mathbf{JL}_0} \mathbf{J}V_1 V_2 e^{s\mathbf{JL}} P_c(\omega) f ds \right\|_{L_t^4 L_x^\infty}. \end{aligned}$$

We now use the Christ–Kiselev lemma (Lemma 5.7) with $K(t, s) = e^{(t-s)\mathbf{JL}_0} \mathbf{J}V_1 : L_t^2 H^{\frac{3}{4}+\epsilon} \rightarrow L_t^4 L_x^\infty$. Following (5.21),

$$\tilde{\mathcal{K}}[V_2 e^{s\mathbf{JL}} P_c(\omega) f] = \int_0^t e^{(t-s)\mathbf{JL}_0} \mathbf{J}V_1 V_2 e^{s\mathbf{JL}} P_c(\omega) f ds.$$

According to Lemma 5.7, we have

$$\begin{aligned} \left\| \int_0^t e^{(t-s)\mathbf{JL}_0} \mathbf{J}V_1 V_2 e^{s\mathbf{JL}} P_c(\omega) f ds \right\|_{L_t^4 L_x^\infty} &= \|\tilde{\mathcal{K}}[V_2 e^{s\mathbf{JL}} P_c(\omega) f]\|_{L_t^4 L_x^\infty} \\ &\leq C \|\mathcal{K}\|_{L_t^2 H^{\frac{3}{4}+\epsilon} \rightarrow L_t^4 L_x^\infty} \|V_2 e^{t\mathbf{JL}} P_c(\omega) f\|_{L_t^2 H^{\frac{3}{4}+\epsilon}}. \end{aligned}$$

From the interpolation between the cases $s = 0$ and $s = 1$, the decay and smoothness properties of V_2 and the weighted decay estimate from Proposition 5.1, we conclude that $\|V_2 e^{t\mathbf{JL}} P_c(\omega) f\|_{L_t^2 H_x^s} \leq C \|f\|_{H^s}$, and we arrive at the estimate $\|V_2 e^{t\mathbf{JL}} P_c(\omega) f\|_{L_t^2 H^{\frac{3}{4}+\epsilon}} \leq C \|f\|_{H^{\frac{3}{4}+\epsilon}}$.

It remains to obtain the appropriate estimate for $\|\mathcal{K}\|_{L_t^2 H^{\frac{3}{4}+\epsilon} \rightarrow L_t^4 L_x^\infty}$. We have again by the Strichartz estimates for the free Dirac evolution (more precisely, the version of (5.28)):

$$\begin{aligned} \left\| \int_{-\infty}^\infty e^{(t-s)\mathbf{JL}_0} \mathbf{J}V_1 G(s, \cdot) ds \right\|_{L_t^4 L_x^\infty} &= \|e^{t\mathbf{JL}_0} \int_{-\infty}^\infty e^{-s\mathbf{JL}_0} \mathbf{J}V_1 G(s, \cdot) ds\|_{L_t^4 L_x^\infty} \\ &\leq C \|\partial_x\|^{3/4} \int_{-\infty}^\infty e^{-s\mathbf{JL}_0} \mathbf{J}V_1 G(s, \cdot) ds\|_{H^\epsilon}. \end{aligned}$$

From Lemma 5.4 (and more precisely from (5.18)), we have

$$\| |\partial_x|^{3/4} \int_{-\infty}^{\infty} e^{-s\mathbf{JL}_0} \mathbf{J}V_1 G(s, \cdot) ds \|_{H^\epsilon} \leq C \| |\partial_x|^{3/4} M[\mathbf{J}V_1 G(s)] \|_{L_x^1 H^\epsilon}.$$

Note that in the low frequencies, $|\partial_x|^{3/4} M \sim |\partial_x|^{1/4}$ is not singular anymore, while in the high frequencies one has $|\partial_x|^{3/4} M \sim |\partial_x|^{3/4}$. Thus, with V_1 in the Besov space $B_2^{1,1}$, we have

$$\| |\partial_x|^{3/4} M[\mathbf{J}V_1 G(s)] \|_{L_x^1 H_x^\epsilon} \leq C \| V_1 \|_{B_2^{1,1}} \| G \|_{L_x^2 H^{3/4+\epsilon}}.$$

With that, Lemma 5.8 is proved in full. \square

Our next lemma is another essential component of the fixed point arguments to be presented in Section 6. Namely, it connects the Strichartz estimates to the weighted decay estimates.

Lemma 5.9. *There is $C < \infty$ such that*

$$\| \int_0^t e^{(t-\tau)\mathbf{JL}} P_c(\omega) F(\tau, \cdot) d\tau \|_{L_t^\infty H_x^1 \cap L_t^4 L_x^\infty} \leq C [\| F \|_{(L^1)_x L_t^2} + \| \partial_x F \|_{(L^1)_x L_t^2}], \tag{5.30}$$

$$\sup_x \langle x \rangle^{-3} \| \int_0^t e^{(t-\tau)\mathbf{JL}} P_c(\omega) F(\tau, \cdot) d\tau \|_{L_t^2} \leq C \| F \|_{L_t^1 L_x^2}. \tag{5.31}$$

Proof. For the proof of (5.30), by Lemma 5.7, we may consider the Duhamel’s operator in the form $\int_{-\infty}^{\infty} \dots$, instead of the retarded term with $\int_0^t \dots$ in the Duhamel representation. By (5.22) and (5.23),

$$\begin{aligned} \left\| \int_{-\infty}^{\infty} e^{(t-\tau)\mathbf{JL}} P_c(\omega) F(\tau, \cdot) d\tau \right\|_{L_t^\infty H_x^1 \cap L_t^4 L_x^\infty} &= \left\| e^{t\mathbf{JL}} P_c(\omega) \int_{-\infty}^{\infty} e^{-\tau\mathbf{JL}} F(\tau, \cdot) d\tau \right\|_{L_t^\infty H_x^1 \cap L_t^4 L_x^\infty} \\ &\leq \left\| \int_{-\infty}^{\infty} e^{-\tau\mathbf{JL}} P_c(\omega) F(\tau, \cdot) d\tau \right\|_{H_x^1}. \end{aligned}$$

To prove (5.30), we need to estimate two terms: one with a derivative and one without a derivative. The term without a derivative is dealt with by Proposition 5.1:

$$\left\| \int_{-\infty}^{\infty} e^{-\tau\mathbf{JL}} P_c(\omega) F(\tau, \cdot) d\tau \right\|_{L_x^2} \leq C \| F \|_{(L^1)_x L_t^2}. \tag{5.32}$$

For the term $\| \int_{-\infty}^{\infty} \partial_x [e^{-\tau\mathbf{JL}} P_c(\omega) F(\tau, \cdot)] d\tau \|_{L_x^2}$, we are facing a difficulty since $\partial_x e^{-\tau\mathbf{JL}} \neq e^{-\tau\mathbf{JL}} \partial_x$. Nevertheless, due to the fact that $\mathbf{L} = \mathbf{D}_m - \omega I_4 + \mathbf{W}$, we use the L_x^2 estimate (5.32) to derive

$$\begin{aligned} \left\| \int_{-\infty}^{\infty} \partial_x [e^{-\tau\mathbf{JL}} P_c(\omega) F(\tau, \cdot)] d\tau \right\|_{L_x^2} &\leq \left\| \int_{-\infty}^{\infty} (\mathbf{L} - \beta + \omega I_4 - \mathbf{W}) [e^{-\tau\mathbf{JL}} P_c(\omega) F(\tau, \cdot)] d\tau \right\|_{L_x^2} \\ &\leq C \left\{ \| \mathbf{L} F \|_{(L^1)_x L_t^2} + (1 + |\omega| + \| \mathbf{W} \|_{L_x^\infty}) \| F \|_{(L^1)_x L_t^2} \right\}. \end{aligned} \tag{5.33}$$

Taking into account the specific form of \mathbf{JL} , it follows from (5.32) and (5.33) that

$$\left\| \int_{-\infty}^{\infty} e^{-\tau\mathbf{JL}} g_1(\tau) P_c(\omega) g_2 d\tau \right\|_{H_x^1} \leq C [\| F \|_{(L^1)_x L_t^2} + \| \partial_x F \|_{(L^1)_x L_t^2}].$$

We now turn to proving (5.31) Because of the weak* density of linear combinations $\{\delta(t - \tau_0)G(x) : \tau_0 \in \mathbb{R}^1, G \in L_x^2(\mathbb{R}^1)\}$ in $L_t^1 L_x^2$, it suffices to prove (5.31) for $F(x) = \delta(t - \tau_0)G(x)$. By Proposition 5.1,

$$\begin{aligned} \sup_x \langle x \rangle^{-3} \left\| \int_0^t e^{(t-\tau)\mathbf{JL}} P_c(\omega) \delta(\tau - \tau_0) G(x) d\tau \right\|_{L_t^2} &= \sup_x \langle x \rangle^{-3} \|e^{(t-\tau_0)\mathbf{JL}} P_c(\omega) G(x)\|_{L_t^2} \\ &= \sup_x \langle x \rangle^{-3} \|e^{t\mathbf{JL}} P_c(\omega) G(x)\|_{L_t^2} \leq C \|G\|_{L_x^2}. \quad \square \end{aligned}$$

6. Proof of the main theorem

In this section, the constants C may change from one instance to another; they all depend only on Ω and on the nonlinearity f in (2.3). We analyze the modulation equations (3.31) and the PDE (6.6). Let $\epsilon > 0$ be sufficiently small and

$$\psi_0 e^{i\theta_0} = \phi_{\omega_0} + \rho_0, \quad \rho_0 \in \mathbf{X}_c(\omega_0), \quad \theta_0 \in \mathbb{R}, \quad \|\rho_0\|_{H^1} \leq \epsilon^2.$$

Without loss of generality, we assume that $\theta_0 = 0$.

Definition 6.1. For fixed $N > 10$ and $T > 0$, let

$$\begin{aligned} \|Z\|_{\mathcal{X}_T} &= \|Z\|_{L_t^1 L_x^\infty} + \|Z\|_{L_t^\infty H_x^1} + \left\| \langle x \rangle^{-N} Z \right\|_{L_x^\infty L_t^2} + \left\| \langle x \rangle^{-N} \partial_x Z \right\|_{L_x^\infty L_t^2}, \\ \|F\|_{\mathcal{Y}_T} &= \inf_{F=A+B} \left[\|A\|_{L_t^1 H_x^1} + \left\| \langle x \rangle^N B \right\|_{L_x^1 L_t^2} + \left\| \langle x \rangle^N \partial_x B \right\|_{L_x^1 L_t^2} \right], \end{aligned}$$

where $L_t^\alpha = L^\alpha[0, T]$ and $L_x^\alpha = L^\alpha(\mathbb{R})$.

Lemma 6.2. *There is $C < \infty$ such that for each $\omega_0 \in \Omega$ there is $\epsilon_1 \in (0, \text{dist}(\omega_0, \partial\Omega))$ such that if ω and $Z \in H^1(\mathbb{R}, \mathbb{C}^4)$ satisfy $|\omega - \omega_0| < \epsilon_1$, $\|\langle x \rangle^{-N} Z\|_{H_x^1} \leq \epsilon_1$ with $N > 10$ from Definition 6.1, then*

$$|\langle \phi, \mathbf{N}_1(R, \omega) \rangle| + |\langle \mathbf{J} \partial_\omega \phi, \mathbf{N}_1(R, \omega) \rangle| \leq C \langle \mu, |Z|^2 \rangle,$$

where $\mathbf{N}_1(R, \omega)$ is from (3.28), $R = P_c(\omega_0)R$, and $Z = P_c(\omega_0)R$.

Proof. From (3.28), Taylor’s expansion, and Young’s inequality, we see that

$$\mathbf{N}_1 = \mathbf{N}(\phi + \rho) - \mathbf{N}(\phi) - \mathbf{W}R = \mathcal{O}(|\phi|^{2k-1}|R|^2 + |R|^{2k+1}). \tag{6.1}$$

Note that the above makes sense pointwise in $x \in \mathbb{R}$ since $Z \in H^1(\mathbb{R}, \mathbb{C}^4)$, and by (3.38) one also has $R \in H^1(\mathbb{R}, \mathbb{C}^4)$.

By (3.34), this leads to

$$\langle \phi, |\mathbf{N}_1| \rangle \leq C \langle \mu, |R|^2 \rangle (1 + \|\mu R\|_{L_x^\infty}^{2k-1}) \leq C \langle \mu, |Z|^2 \rangle (1 + \|\mu Z\|_{H_x^1}^{2k-1}). \tag{6.2}$$

Let us explain the last inequality. By (3.37) and the triangle inequality,

$$||Z| - |R|| \leq |(P_d(\omega) - P_d(\omega_0))R|, \quad x \in \mathbb{R};$$

multiplying the above by $|R| + |Z|$ and coupling the result with μ , we have

$$|\langle \mu, |Z|^2 \rangle - \langle \mu, |R|^2 \rangle| \leq C |\omega - \omega_0| \langle \mu, |R|(|R| + |Z|) \rangle \leq 2C |\omega - \omega_0| \langle \mu, |R|^2 + |Z|^2 \rangle,$$

with some $C > 0$. It follows that if $|\omega - \omega_0|$ is sufficiently small, then

$$\frac{1}{2} \langle \mu, |Z|^2 \rangle \leq \langle \mu, |R|^2 \rangle \leq 2 \langle \mu, |Z|^2 \rangle.$$

Since $\|\langle x \rangle^{-N} Z\|_{H_x^1} \leq \epsilon_1$, we have $\|\mu Z\|_{H_x^1} \leq C$; therefore, the inequality (6.2) finishes the proof. \square

Applying the projection $P_c(\omega_0)$ to equation (3.27), we obtain:

$$\begin{aligned} \partial_t Z - \mathbf{JL}(\omega_0)Z + (\dot{\gamma}(t) + \omega(t) - \omega_0)P_c(\omega_0)\mathbf{J}Z \\ = P_c(\omega_0)(\mathbf{J}(\mathbf{W}(\omega) - \mathbf{W}(\omega_0))R - \dot{\gamma}\mathbf{J}\Phi - \dot{\omega}\partial_\omega\phi_\omega + \mathbf{JN}_1). \end{aligned} \tag{6.3}$$

We denote

$$\alpha(t) = \dot{\gamma}(t) + \omega(t) - \omega_0 \tag{6.4}$$

and

$$F_0(t) = \mathbf{J}(\mathbf{W}(\omega) - \mathbf{W}(\omega_0))R - \dot{\gamma}\mathbf{J}\Phi - \dot{\omega}\partial_\omega\phi_\omega + \mathbf{JN}_1; \tag{6.5}$$

then (6.3) takes the form

$$\partial_t Z - \mathbf{JL}(\omega_0)Z + \alpha(t)\mathbf{J}Z = P_c(\omega_0)F_0 + \alpha(t)[\mathbf{J}, P_c(\omega_0)]Z. \tag{6.6}$$

We assume that there exist $T > 0$ and $C_0 > 1$ depending on ω_0 such that the solution $(\omega(t), \gamma(t), Z(t))$ to the modulation equations (3.31) and the PDE (6.6) exists on $[0, T]$ and

$$\|\dot{\omega}\|_{L^1[0,T]} + \|\dot{\gamma}\|_{L^1[0,T]} \leq C_0\epsilon, \quad \|Z\|_{\mathcal{X}_T} \leq C_0\epsilon. \tag{6.7}$$

Lemma 6.3. *Assume that (6.7) holds. If $\epsilon > 0$ is sufficiently small, then the estimates (6.7) can be improved as follows:*

$$\|\dot{\omega}\|_{L^1[0,T]} + \|\dot{\gamma}\|_{L^1[0,T]} \leq \epsilon, \quad \|Z\|_{\mathcal{X}_T} \leq \epsilon. \tag{6.8}$$

Proof. By (3.31), the invertibility of $\mathcal{A}(t)$ (cf. Lemma 3.7), and the bounds from Lemma 6.2, we conclude that

$$|\dot{\gamma}| + |\dot{\omega}| \leq C\langle \mu, |Z(t)|^2 \rangle,$$

hence, for small enough $\epsilon > 0$,

$$\|\dot{\omega}\|_{L^1[0,T]} + \|\dot{\gamma}\|_{L^1[0,T]} \leq C \int_0^T \langle \mu, |Z(t)|^2 \rangle dt \leq C \left\| \mu^{1/3} Z \right\|_{L_x^\infty L_t^2}^2 \leq C \|Z\|_{\mathcal{X}_T}^2 \leq CC_0^2\epsilon^2 \leq \epsilon; \tag{6.9}$$

we used the bound on $\|Z\|_{\mathcal{X}_T}$ from (6.7). This proves the first estimate in (6.8).

With (6.9), we also have

$$\|\omega - \omega_0\|_{L^\infty[0,T]} \leq \|\dot{\omega}\|_{L^1[0,T]} \leq C \|Z\|_{\mathcal{X}_T}^2 \leq CC_0^2\epsilon^2 \leq \epsilon. \tag{6.10}$$

It follows from (6.6) with the initial data (3.36), (6.10), and from Lemma A.1 below that if $\epsilon > 0$ is sufficiently small, then

$$\|Z\|_{\mathcal{X}_T} \leq C \left[\|Z(0)\|_{H^1} + \|F\|_{\mathcal{Y}_T} \right], \quad F(t) := P_c(\omega_0)F_0(t) + \alpha(t)[\mathbf{J}, P_c(\omega_0)]Z(t). \tag{6.11}$$

From the definition (6.5) of F_0 , we see that

$$\|F_0 - \mathbf{JN}_1\|_{\mathcal{Y}_T} \leq C \left[\|\omega - \omega_0\|_{L_t^\infty[0,T]} \|Z\|_{\mathcal{X}_T} + \|\dot{\gamma}\|_{L_t^\infty[0,T]} + \|\dot{\omega}\|_{L_t^\infty[0,T]} \right]. \tag{6.12}$$

We used the bound $\|R\|_{\mathcal{X}_T} \leq C \|Z\|_{\mathcal{X}_T}$ which follows from (3.38). Noting that $[\mathbf{J}, P_c(\omega_0)]$ is localized in space and recalling that $\alpha(t) = \dot{\gamma}(t) + \omega(t) - \omega_0$, we also have

$$\begin{aligned} \|P_c(\omega_0)(F_0 - \mathbf{JN}_1) + \alpha(t)[\mathbf{J}, P_c(\omega_0)]Z\|_{\mathcal{Y}_T} \\ \leq C \left[(\|\dot{\gamma}\|_{L_t^\infty[0,T]} + \|\omega - \omega_0\|_{L_t^\infty[0,T]}) \|Z\|_{\mathcal{X}_T} + \|\dot{\gamma}\|_{L_t^\infty[0,T]} + \|\dot{\omega}\|_{L_t^\infty[0,T]} \right]. \end{aligned} \tag{6.13}$$

By Lemma 6.2,

$$|\dot{\omega}(t)| + |\dot{\gamma}(t)| \leq C \|Z(t)\|_{L_x^2}^2 \leq C \|Z\|_{\mathcal{X}_T}^2 \leq CC_0^2\epsilon^2 \leq \epsilon, \quad 0 \leq t \leq T, \tag{6.14}$$

as long as $\epsilon > 0$ is sufficiently small. Applying (6.14) and (6.10) in (6.13), we conclude that there is $C < \infty$ such that

$$\|P_c(\omega_0)(F_0 - \mathbf{JN}_1) + \alpha(t)[\mathbf{J}, P_c(\omega_0)]Z\|_{\mathcal{Y}_T} \leq C \|Z\|_{\mathcal{X}_T}^2. \tag{6.15}$$

By (3.38) and (6.1), using Young’s inequality, we see that

$$\mathbf{N}_1 = \mathcal{O}(|\phi|^{2k-1}|R|^2 + |R|^{2k+1}) = \mathcal{O}(|\phi|^{2k-1}|Z|^2 + |Z|^{2k+1} + \mu|\omega - \omega_0|^2 \langle \mu^{2k}, |R| \rangle^2).$$

Then, it follows from (3.37) and (6.10) that

$$\|\mathbf{N}_1\|_{\mathcal{Y}_T} \leq C \left(\left\| |\phi|^{2k-1}|Z|^2 \right\|_{\mathcal{Y}_T} + \left\| |Z|^{2k+1} \right\|_{\mathcal{Y}_T} + C \|Z\|_{\mathcal{X}_T}^4 \right). \tag{6.16}$$

On the other hand, from the definitions of $\|\cdot\|_{\mathcal{X}_T}$, $\|\cdot\|_{\mathcal{Y}_T}$ (cf. Definition 6.1), we observe that

$$\left\| |\phi|^{2k-1}|Z|^2 \right\|_{\mathcal{Y}_T} \leq C \left\| \langle x \rangle^n |\phi|^{2k-1}|Z|^2 \right\|_{L_x^1 L_t^2} + C \left\| \langle x \rangle^N \partial_x [|\phi|^{2k-1}|Z|^2] \right\| \leq C \|Z\|_{\mathcal{X}_T}^2 \leq C \|Z\|_{\mathcal{X}_T}^2. \tag{6.17}$$

Similarly, we have

$$\left\| |Z|^{2k+1} \right\|_{\mathcal{Y}_T} \leq C \left\| |Z|^{2k+1} \right\|_{L_t^1 H_x^1} \leq C \left\| (|Z| + |\partial_x Z|)|Z|^{2k} \right\|_{L_t^1 L_x^2} \leq C \|Z\|_{L_t^\infty H_x^1} \|Z\|_{L_t^{2k} L_x^\infty}^{2k}.$$

We note that $\|Z\|_{L_t^\infty H_x^1} \leq \|Z\|_{\mathcal{X}_T}$; since $k \geq 2$, we arrive at

$$\|Z\|_{L_t^{2k} L_x^\infty} \leq \|Z\|_{L_t^4 L_x^\infty}^{2/k} \|Z\|_{L_t^\infty L_x^\infty}^{1-2/k} \leq C \|Z\|_{\mathcal{X}_T}.$$

Therefore,

$$\left\| |Z|^{2k+1} \right\|_{\mathcal{Y}_T} \leq C \|Z\|_{\mathcal{X}_T}^{2k+1}. \tag{6.18}$$

In summary, it follows from (6.15), (6.16), (6.17), and (6.18) that there is $C < \infty$ such that

$$\|P_c(\omega_0)F_0 + \alpha(t)[\mathbf{J}, P_c(\omega_0)]Z\|_{\mathcal{Y}_T} \leq C \|Z\|_{\mathcal{X}_T}^2.$$

From this and (6.11), we infer that if $\epsilon > 0$ is sufficiently small, then we have

$$\|Z\|_{\mathcal{X}_T} \leq C[\|Z(\cdot, 0)\|_{H^1} + \|Z\|_{\mathcal{X}_T}^2] \leq C[1 + C_0]\epsilon^2 \leq \epsilon. \tag{6.19}$$

This proves the second estimate in (6.8), completing the proof of the lemma. \square

From Lemma 6.3 and the local existence theory [28], it follows that there exists unique global solution to equation (2.9),

$$\psi(x, t) = (\phi_{\omega(t)}(x) + \rho(x, t))e^{-i\left(\int_0^t \omega(s) ds + \gamma(t)\right)}, \quad t \geq 0,$$

with ω, γ , and $Z = P_c(\omega_0) \left(\begin{bmatrix} \operatorname{Re} \rho \\ \operatorname{Im} \rho \end{bmatrix} \right)$ satisfying the estimates

$$\|\dot{\omega}\|_{L^1(\mathbb{R}_+)} + \|\dot{\gamma}\|_{L^1(\mathbb{R}_+)} \leq \epsilon, \quad \|Z\|_{\mathcal{X}_\infty} \leq \epsilon.$$

From this, we infer that there exist $\omega_\infty, \gamma_\infty \in \mathbb{R}$ such that

$$\lim_{t \rightarrow \infty} \omega(t) = \omega_\infty, \quad \lim_{t \rightarrow \infty} \gamma(t) = \gamma_\infty, \quad \lim_{t \rightarrow \infty} \|Z(t)\|_{L_x^\infty} = 0.$$

The last relation is due to $\|Z\|_{L_t^4 L_x^\infty} \leq \|Z\|_{\mathcal{X}_\infty}$. Due to (3.38) and (6.10), assuming that $\epsilon > 0$ is sufficiently small, we also have

$$\lim_{t \rightarrow \infty} \|\rho(t)\|_{L_x^\infty} = 0.$$

This completes the proof of the main theorem.

Conflict of interest statement

There is no conflict of interests.

Appendix A. Estimates for the linear perturbed equation

This subsection proves the estimate (6.11) on Z . We recall that $\mathbf{X}_c(\omega_0)$ is defined in (3.20), \mathbf{J} , and \mathbf{L} are defined in (3.3). Also, we denote $\mathcal{X} = \mathcal{X}_T$, $\mathcal{Y} = \mathcal{Y}_T$, where \mathcal{X}_T and \mathcal{Y}_T are defined in Definition 6.1 for some $T \in (0, \infty]$. The main result is the following lemma:

Lemma A.1. Fix $\omega_0 \in \Omega$ and assume that the Assumption 2.4 holds. Let $Z(t) \in \mathbf{X}_c(\omega_0)$ be a solution to the equation

$$\begin{cases} \partial_t Z - \mathbf{JL}(\omega_0)Z + \alpha(t)\mathbf{J}Z = F, & t \in (0, T), \\ Z(0) = Z_0 \in \mathbf{X}_c(\omega_0). \end{cases}$$

Then there exist $c_0 > 0$ and $C < \infty$ independent on T such that if $\|\alpha\|_{L^\infty((0,T))} \leq c_0$, we have

$$\|Z\|_{\mathcal{X}} \leq C \left[\|Z(0)\|_{H^1} + \|F\|_{\mathcal{Y}} \right].$$

Proof. It follows from our linear estimates in Section 5 that Lemma A.1 holds when $\alpha = 0$. The proof therefore is a perturbative argument. We base our argument on [27, Appendix B], which originates in [5]. In the perturbation argument, instead of using the free operator as in [5,27], we shall make use of the operator

$$\mathbf{L}_\nu = \begin{bmatrix} H_\nu & 0 \\ 0 & H_\nu \end{bmatrix}, \quad \text{with } H_\nu := D_m - \omega_0 + V_\nu,$$

where V_ν is a fixed matrix-valued potential which is sufficiently small and decays exponentially, and such that the point spectrum $\sigma_d(H_\nu)$ of H_ν is empty and there is no resonance at thresholds $\Lambda = \pm m - \omega_0$. The advantage of using \mathbf{L}_ν is that it has stronger decay estimates (A.6) which essentially follow from [21, Theorem 3.7].

We now denote $\mathbf{W}_\nu = \mathbf{L}(\omega_0) - \mathbf{L}_\nu$, the exponentially decaying matrix potential; thus,

$$\mathbf{L}(\omega_0) = \mathbf{L}_\nu + \mathbf{W}_\nu.$$

For fixed $\kappa > 0$ and for $P_d(\omega_0) := \text{Id} - P_c(\omega_0)$, we consider the auxiliary equation

$$\partial_t \Psi - \mathbf{JL}(\omega_0)P_c(\omega_0)\Psi + \kappa P_d(\omega_0)\Psi + \alpha\mathbf{J}P_c(\omega_0)\Psi = F, \quad \Psi(0) = Z(0). \tag{A.1}$$

We note that $Z = P_c(\omega_0)\Psi$, therefore it suffices to prove the estimate for Ψ . Let us denote

$$\beta(t) = \int_0^t \alpha(s) ds, \quad U(t) = e^{\beta(t)\mathbf{J}}, \quad \Psi(t) = U(t)\Phi.$$

Then it follows from (A.1) that

$$\partial_t \Phi + U^{-1}(-\mathbf{JL}(\omega_0) + \kappa P_d(\omega_0))U\Phi = G, \quad G := U^{-1}F + \alpha(t)U^{-1}\mathbf{J}P_d(\omega_0)U\Phi.$$

Since \mathbf{J} commutes with \mathbf{L}_ν , we obtain

$$\partial_t \Phi - \mathbf{JL}_\nu\Phi = -U^{-1}(\mathbf{W} - \mathbf{JL}(\omega_0)P_d(\omega_0) + \kappa P_d(\omega_0))U\Phi + G. \tag{A.2}$$

Now, we choose V_2 a smooth, exponentially decaying, invertible matrix potential such that the matrix

$$V_1 = (\mathbf{W} - \mathbf{JL}P_d + \kappa P_d)V_2^{-1}$$

is also smooth and exponentially decaying. Then, note that $\Phi(0) = \Psi(0)$, and \mathbf{L}_ν commutes with \mathbf{J} . Therefore, applying $U(t)$ to both sides of equation (A.2), we infer that

$$\begin{aligned} \Psi(t) &= U(t)e^{-t\mathbf{JL}_\nu}\Psi(0) + \int_0^t e^{-(t-s)\mathbf{JL}_\nu} \left[U(t)U^{-1}(s)V_1V_2\Psi(s) - U(t)G(s) \right] ds \\ &= U(t)e^{-t\mathbf{JL}_\nu}\Psi(0) + \int_0^t e^{-(t-s)\mathbf{JL}_\nu} U(t)U^{-1}(s) \left[(V_1 - \alpha(s)\mathbf{J}P_d(\omega_0)V_2^{-1})V_2\Psi - F(s) \right] ds. \end{aligned} \tag{A.3}$$

On the other hand, it follows from [29, Section VIII] that

$$\|\Psi\|_{\mathcal{X}} \leq C \left[\|\Psi(0)\|_{H^1} + \|F\|_{\mathcal{Y}} + \|V_2\Psi\|_{L_t^2 H_x^1} + \|\alpha\|_{L^\infty} \|P_d\Psi\|_{\mathcal{Y}} \right].$$

Note that $\|P_d\Psi\|_{\mathcal{Y}} \leq C \|\Psi\|_{\mathcal{X}}$. Therefore, if $\|\alpha\|_{L^\infty}$ is sufficiently small, we obtain

$$\|\Psi\|_{\mathcal{X}} \leq C \left[\|\Psi(0)\|_{H^1} + \|F\|_{\mathcal{Y}} + \|V_2\Psi\|_{L_t^2 H_x^1} \right]. \tag{A.4}$$

Next, we need to control $\|V_2\Psi\|_{L_t^2 H_x^1}$. We denote

$$T_0 f(t) = V_2 \int_0^t e^{-(t-s)\mathbf{J}\mathbf{L}_v} U(t)U^{-1}(s)V_1 f(\cdot, s) ds.$$

From Lemma A.2 below, we see that the mapping $I - T_0 : L_t^2 H_x^1 \rightarrow L_t^2 H_x^1$ is invertible and there exists $C < \infty$ such that $\|(I - T_0)^{-1}\|_{L_t^2 H_x^1 \rightarrow L_t^2 H_x^1} \leq C$. By (A.3), we see that

$$(I - T_0)V_2\Psi = V_2 U(t)e^{-t\mathbf{J}\mathbf{L}_v}\Psi(0) - V_2 \int_0^t e^{-(t-s)\mathbf{J}\mathbf{L}_v} U(t)U^{-1}(s)[F(s) + \alpha(s)\mathbf{J}P_d\Psi(s)] ds. \tag{A.5}$$

Therefore, using again the linear estimates from [29], we obtain:

$$\begin{aligned} \|V_2\Psi\|_{L_t^2 H_x^1} &\leq \left\| V_2 U(t)e^{-t\mathbf{J}\mathbf{L}_v}\Psi(0) \right\|_{L_t^2 H_x^1} + \left\| V_2 \int_0^t e^{-(t-s)\mathbf{J}\mathbf{L}_v} U(t)U^{-1}(s)[F(s) + \alpha(s)\mathbf{J}P_d\Psi(s)] ds \right\|_{L_t^2 H_x^1} \\ &\leq C \left[\|\Psi(0)\|_{H_x^1} + \|F\|_{\mathcal{Y}} + \|\alpha\|_{L^\infty} \|\Psi\|_{\mathcal{X}} \right]. \end{aligned}$$

From this and (A.4), we see that there is $c_0 > 0$ sufficiently small such that if $\|\alpha\|_{L^\infty} \leq c_0$, then one has $\|\Psi\|_{\mathcal{X}} \leq C \left[\|\Psi(0)\|_{H^1} + \|F\|_{\mathcal{Y}} \right]$. Since $Z = P_c(\omega_0)\Psi$, this completes the proof of the lemma. \square

Lemma A.2. For $k = 0, 1$, the map $I - T_0 : L_t^2 H_x^k \mapsto L_t^2 H_x^k$ is invertible and therefore there exists $C < \infty$ such that

$$\|(I - T_0)^{-1}\|_{L_t^2 H_x^k \rightarrow L_t^2 H_x^k} \leq C.$$

Proof. First, note that it follows from the linear estimates in [29, Section VIII] that T_0 is well-defined as an operator from $L_t^2 H_x^k$ to $L_t^2 H_x^k$, with $k = 0, 1$. We now let

$$T_1 f(t) = V_2 \int_0^t e^{-(t-s)\mathbf{J}\mathbf{L}_v} V_1 f(\cdot, s) ds.$$

It follows from our linear estimates in Section 5 that T_1 is also well-defined from $L_t^2 H_x^1$ to $L_t^2 H_x^1$. Also, note that

$$(T_1 - T_0)f = V_2 \int_0^t e^{-(t-s)\mathbf{J}\mathbf{L}_v} \left(e^{\mathbf{J} \int_t^s \alpha(\tau) d\tau} - 1 \right) V_1 f(\cdot, s) ds.$$

By [21, Theorem 3.7], we have

$$\|e^{-t\mathbf{J}\mathbf{L}_v}\|_{L_\sigma^2 \rightarrow L_{-\sigma}^2} \leq C_\sigma \langle t \rangle^{-3/2}, \quad \sigma > 5/2.$$

From this, we further infer that $\|\mathbf{L}_v e^{-t\mathbf{J}\mathbf{L}_v} f\|_{L_{-\sigma}^2} \leq C_\sigma \langle t \rangle^{-3/2} \|\mathbf{L}_v f\|_{L_\sigma^2}$. Since $\|f\|_{H_x^1} \sim \|f\|_{L^2} + \|\mathbf{L}_v f\|_{L^2}$, we see that

$$\left\| e^{-t\mathbf{J}\mathbf{L}_v} \right\|_{H_\sigma^k \rightarrow H_{-\sigma}^k} \leq C_\sigma \langle t \rangle^{-3/2}, \quad \sigma > 5/2, \quad k = 0, 1. \quad (\text{A.6})$$

Using (A.6) and the fact that

$$\left| e^{\mathbf{J} \int_t^s \alpha(\tau) d\tau} - 1 \right| \leq \min(1, \|\alpha\|_{L^\infty}(t-s)),$$

we obtain:

$$\left\| V_2 e^{-(t-s)\mathbf{J}\mathbf{L}_v} \left[e^{\mathbf{J} \int_t^s \alpha(\tau) d\tau} - 1 \right] V_1 f(\cdot, s) ds \right\|_{H^k} \leq C \|\alpha\|_{L^\infty}^{1/4} \langle t-s \rangle^{-5/4} \|f(\cdot, s)\|_{H^k}.$$

Thus, if $\|\alpha\|_{L^\infty}$ is sufficiently small, we see that

$$\|T_1 - T_0\|_{L_t^2 H_x^k \rightarrow L_t^2 H_x^k} \leq C \|\alpha\|_{L^\infty}^{1/4} < 1.$$

Therefore, it suffices to prove that $I - T_1$ is invertible. The lemma then follows exactly as in [27, Lemma B.2] by using the linear estimates on $e^{-t\mathbf{J}\mathbf{L}(\omega_0)}$ from Section 5. \square

References

- [1] G. Berkolaiko, A. Comech, On spectral stability of solitary waves of nonlinear Dirac equation in 1D, *Math. Model. Nat. Phenom.* 7 (2012) 13–31.
- [2] N. Boussaïd, A. Comech, On spectral stability of nonlinear Dirac equation, 2012, ArXiv e-prints.
- [3] N. Boussaïd, S. Cuccagna, On stability of standing waves of nonlinear Dirac equations, *Commun. Partial Differ. Equ.* 37 (2012) 1001–1056.
- [4] G. Berkolaiko, A. Comech, A. Sukhtayev, Vakhitov–Kolokolov and energy vanishing conditions for linear instability of solitary waves in models of classical self-interacting spinor fields, *Nonlinearity* 28 (2015) 577–592.
- [5] M. Beceanu, New estimates for a time-dependent Schrödinger equation, *Duke Math. J.* 159 (2011) 417–477.
- [6] V.S. Buslaev, G.S. Perel'man, On nonlinear scattering of states which are close to a soliton, in: *Méthodes semi-classiques*, vol. 2, Nantes, 1991, *Astérisque* 210 (1992) 49–63.
- [7] V.S. Buslaev, G.S. Perel'man, Scattering for the nonlinear Schrödinger equation: states that are close to a soliton, *Algebra Anal.* 4 (1992) 63–102.
- [8] I.V. Barashenkov, D.E. Pelinovsky, E.V. Zemlyanaya, Vibrations and oscillatory instabilities of gap solitons, *Phys. Rev. Lett.* 80 (1998) 5117–5120.
- [9] T. Candy, Global existence for an L^2 critical nonlinear Dirac equation in one dimension, *Adv. Differ. Equ.* 16 (2011) 643–666.
- [10] A. Comech, S. Cuccagna, D.E. Pelinovsky, Nonlinear instability of a critical traveling wave in the generalized Korteweg–de Vries equation, *SIAM J. Math. Anal.* 39 (2007) 1–33.
- [11] A. Comech, M. Guan, S. Gustafson, On linear instability of solitary waves for the nonlinear Dirac equation, *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* 31 (2014) 639–654.
- [12] A. Comech, On the meaning of the Vakhitov–Kolokolov stability criterion for the nonlinear Dirac equation, 2011, ArXiv e-prints.
- [13] S. Cuccagna, On asymptotic stability in energy space of ground states of NLS in 1D, *J. Differ. Equ.* 245 (2008) 653–691.
- [14] T. Cazenave, L. Vázquez, Existence of localized solutions for a classical nonlinear Dirac field, *Commun. Math. Phys.* 105 (1986) 35–47.
- [15] R. Finkelstein, C. Fronsdal, P. Kaus, Nonlinear spinor field, *Phys. Rev.* 103 (1956) 1571–1579.
- [16] R. Finkelstein, R. LeLevier, M. Ruderman, Nonlinear spinor fields, *Phys. Rev.* 83 (1951) 326–332.
- [17] W. Heisenberg, Quantum theory of fields and elementary particles, *Rev. Mod. Phys.* 29 (1957) 269–278.
- [18] H. Huh, Global solutions to Gross–Neveu equation, *Lett. Math. Phys.* 103 (2013) 927–931.
- [19] D.D. Ivanenko, Notes to the theory of interaction via particles, *Zh. Eksp. Teor. Fiz.* 8 (1938) 260–266.
- [20] J. Krieger, K. Nakanishi, W. Schlag, Global dynamics above the ground state energy for the one-dimensional NLKG equation, *Math. Z.* 272 (2012) 297–316.
- [21] E.A. Kopylova, Weighted energy decay for 1D Dirac equation, *Dyn. Partial Differ. Equ.* 8 (2011) 113–125.
- [22] T. Kapitula, B. Sandstede, Edge bifurcations for near integrable systems via Evans function techniques, *SIAM J. Math. Anal.* 33 (2002) 1117–1143.
- [23] S.Y. Lee, A. Gavrielides, Quantization of the localized solutions in two-dimensional field theories of massive fermions, *Phys. Rev. D* 12 (1975) 3880–3886.
- [24] T. Mizumachi, Asymptotic stability of small solitary waves to 1D nonlinear Schrödinger equations with potential, *J. Math. Kyoto Univ.* 48 (2008) 471–497.
- [25] S. Machihara, K. Nakanishi, K. Tsugawa, Well-posedness for nonlinear Dirac equations in one dimension, *Kyoto J. Math.* 50 (2010) 403–451.
- [26] M. Nakamura, T. Ozawa, The Cauchy problem for nonlinear Klein–Gordon equations in the Sobolev spaces, *Publ. Res. Inst. Math. Sci.* 37 (2001) 255–293.
- [27] K. Nakanishi, W. Schlag, Global dynamics above the ground state energy for the cubic NLS equation in 3D, *Calc. Var. Partial Differ. Equ.* 44 (2012) 1–45.
- [28] D. Pelinovsky, Survey on global existence in the nonlinear Dirac equations in one spatial dimension, in: *Harmonic Analysis and Nonlinear Partial Differential Equations*, in: *RIMS Kôkyûroku Bessatsu*, vol. B26, Res. Inst. Math. Sci. (RIMS), Kyoto, 2011, pp. 37–50.

- [29] D.E. Pelinovsky, A. Stefanov, Asymptotic stability of small gap solitons in nonlinear Dirac equations, *J. Math. Phys.* 53 (2012) 073705.
- [30] M. Soler, Classical, stable, nonlinear spinor field with positive rest energy, *Phys. Rev. D* 1 (1970) 2766–2769.
- [31] S. Shao, N.R. Quintero, F.G. Mertens, F. Cooper, A. Khare, A. Saxena, Stability of solitary waves in the nonlinear Dirac equation with arbitrary nonlinearity, 2014, ArXiv e-prints.
- [32] H.F. Smith, C.D. Sogge, Global Strichartz estimates for nontrapping perturbations of the Laplacian, *Commun. Partial Differ. Equ.* 25 (2000) 2171–2183.
- [33] S. Selberg, A. Tesfahun, Low regularity well-posedness for some nonlinear Dirac equations in one space dimension, *Differ. Integral Equ.* 23 (2010) 265–278.
- [34] B. Thaller, *The Dirac Equation*, Texts and Monographs in Physics, Springer-Verlag, Berlin, 1992.
- [35] W.E. Thirring, A soluble relativistic field theory, *Ann. Phys.* 3 (1958) 91–112.
- [36] N.G. Vakhitov, A.A. Kolokolov, Stationary solutions of the wave equation in the medium with nonlinearity saturation, *Radiophys. Quantum Electron.* 16 (1973) 783–789.