

A stochastic Lagrangian proof of global existence of the Navier–Stokes equations for flows with small Reynolds number

Gautam Iyer¹

Department of Mathematics, Stanford University, USA

Received 13 February 2007; received in revised form 23 September 2007; accepted 11 October 2007

Available online 1 November 2007

Abstract

We consider the incompressible Navier–Stokes equations with spatially periodic boundary conditions. If the Reynolds number is small enough we provide an elementary short proof of the existence of global in time Hölder continuous solutions. Our proof uses a stochastic representation formula to obtain a decay estimate for heat flows in Hölder spaces, and a stochastic Lagrangian formulation of the Navier–Stokes equations.

© 2007 L'Association Publications de l'Institut Henri Poincaré. Published by Elsevier B.V. All rights reserved.

MSC: primary 76D03; secondary 76D05, 60K40

Keywords: Stochastic Lagrangian; Incompressible Navier–Stokes; Global existence

1. Introduction

The Navier–Stokes equations

$$\partial_t u + (u \cdot \nabla)u - \nu \Delta u + \nabla p = 0, \quad (1.1)$$

$$\nabla \cdot u = 0 \quad (1.2)$$

describe the evolution of the velocity field of an incompressible fluid with kinematic viscosity $\nu > 0$. One of the (still open) million dollar problems posed by the Clay Institute [10] is to show that given a smooth initial data u_0 the solution to (1.1)–(1.2) in three dimensions remains smooth for all time.

In two dimensions, the long time existence of (1.1)–(1.2) is well known [3]. In three or higher dimensions, long time existence is known provided a smallness condition is imposed on the initial data. In 1962, Kato and Fujita [8] showed existence of global solutions to (1.1)–(1.2) when the initial data is small in L^3 . Global existence when the initial data is small in H^1 is standard and can be found in books. Possibly the most general result of this type (see Koch and Tataru [17]) shows global existence of (1.1)–(1.2) assuming that the initial data is small in BMO^{-1} . We also remark that recently Chemin and Gallagher [13] found a (non-linear) criterion on the initial data that guarantees global existence of (1.1)–(1.2), and does not reduce to a smallness criterion in BMO^{-1} .

E-mail address: gautam@math.stanford.edu.

¹ The author was partly supported by NSF grant DMS-0707920.

In this paper we prove global existence of (1.1)–(1.2) provided our initial data has small Hölder norm. Though global existence under our assumptions can be deduced from the Koch–Tataru result, the proof we present here (Section 5) is short, ‘elementary’ and essentially relies only on the decay of heat flows (Section 4), and a stochastic representation of the Navier–Stokes equations using particle trajectories (Section 3, see also [5,15]).

2. Notational conventions and description of results

In this section we describe the notational convention we use, and state the main result we prove. Let $L > 0$ be a fixed length scale, and $\mathcal{I} = [0, L]$. We define the Hölder norms and semi-norms on \mathcal{I}^d by

$$\begin{aligned} |u|_\alpha &= \sup_{x,y \in \mathcal{I}^d} L^\alpha \frac{|u(x) - u(y)|}{|x - y|^\alpha}, \\ \|u\|_{C^k} &= \sum_{|m| \leq k} L^{|m|} \sup_{\mathcal{I}^d} |D^m u|, \\ \|u\|_{k,\alpha} &= \|u\|_{C^k} + \sum_{|m|=k} L^k |D^m u|_\alpha, \end{aligned}$$

where D^m denotes the derivative with respect to the multi index m . We let C^k denote the space of all k -times continuously differentiable spatially periodic functions on \mathcal{I} , and $C^{k,\alpha}$ denote the space of all spatially periodic $k + \alpha$ Hölder continuous functions. The spaces C^k and $C^{k,\alpha}$ are endowed with the norms $\|\cdot\|_{C^k}$ and $\|\cdot\|_{k,\alpha}$ respectively.

We use I to denote the identity function on \mathbb{R}^d (or on \mathcal{I}^d depending on the context), and use \mathbb{I} to denote the identity matrix. The main theorem we prove in this paper is

Theorem 2.1. *Let $k \geq 1$, $\alpha \in (0, 1)$ and $u_0 \in C^{k+1,\alpha}(\mathcal{I}^d)$ be spatially periodic, divergence free and have mean 0. Let $R = \frac{L}{\nu} \|u_0\|_{k+1,\alpha}$ be the Reynolds number of the flow. Then $\exists T = T(k, \alpha, d, \frac{1}{L} \|u_0\|_{k+1,\alpha})$ and $R_0 = R_0(k, \alpha, d)$ such that for all $0 < R < R_0$ the solution u of (1.1)–(1.2) with Reynolds number R , initial data u_0 and periodic boundary conditions is in $C^{k+1,\alpha}$ for time T , and satisfies*

$$\|u_T\|_{k+1,\alpha} \leq \|u_0\|_{k+1,\alpha}. \tag{2.1}$$

We prove Theorem 2.1 in Section 5. A few remarks are in order.

Remark 2.2. Local existence (Theorem A.5) combined with the Theorem 2.1 immediately show that for given initial data, we can choose ν large enough so that (1.1)–(1.2) have time global $C^{k+1,\alpha}$ solutions. Alternately for fixed viscosity, if $\|u_0\|_{k+1,\alpha}$ is small enough, Theorems 2.1 and A.5 again give time global $C^{k+1,\alpha}$ existence of (1.1)–(1.2).

Remark 2.3. The assumption that u_0 has mean 0 is not restrictive. First note that our boundary conditions imply that $\int u_t$ is conserved in time. Set $\bar{u} = \frac{1}{L^d} \int u_0$ to be the mean velocity. Now if we change to coordinates moving with the mean velocity by letting $u'(x, t) = u(x + \bar{u}t, t) - \bar{u}$, then u' solves (1.1)–(1.2) with mean 0 initial data $u_0 - \bar{u}$. Thus the smallness assumption in Remark 2.2 is really smallness assumptions on the deviation from the mean velocity.

Remark 2.4. Theorem 2.1 shows that for some time T , Eq. (2.1) holds. Unfortunately our proof does not show that $\|u_t\|_{k+1,\alpha}$ is decreasing in time.

3. The stochastic Lagrangian formulation

The Kolmogorov forward equation (or Feynman–Kac formula) [12,16] have been extensively used to represent solutions of linear parabolic PDE’s as the average of a stochastic process. In this section we briefly describe here a different approach used in [5,15,6] (see also [19,21]), which we use to provide a representation of the Navier–Stokes equations based on noisy particle paths.

Let $u : \mathbb{R}^d \times [0, \infty) \rightarrow \mathbb{R}^d$ be some given (time dependent) vector field, and θ a solution to the heat equation

$$\partial_t \theta + (u \cdot \nabla) \theta - \nu \Delta \theta = 0 \tag{3.1}$$

with initial data θ_0 . We impose either periodic or decay at infinity boundary conditions on θ .

We express θ as the expected value of a stochastic process as follows: Let W be a d dimensional Wiener process, and let $X : \mathbb{R}^d \rightarrow \mathbb{R}^d$ be a solution to the SDE

$$dX = u dt + \sqrt{2\nu} dW$$

with initial data $X_0(a) = a$. Standard theory² [20] shows that the flow X is a homeomorphism, and as spatially differentiable as u . We let A_t denote the spatial inverse of the flow map X_t .

Proposition 3.1. *If $u \in C^1$, $\theta_0 \in C^2$ then the unique solution θ of (3.1) with initial data θ_0 and either periodic or decay at infinity boundary conditions is given by*

$$\theta_t = \mathbf{E}\theta_0(A_t), \tag{3.2}$$

where \mathbf{E} denotes the expected value with respect to the Wiener measure.

Note that if $\nu = 0$, then Proposition 3.1 is nothing but the method of characteristics. If $\nu > 0$, this can be interpreted as solving along random characteristics, and then averaging. Notice also that the Wiener process $\sqrt{2\nu}W_t$ is the natural one to consider here, as it's generator is $\nu\Delta$.

The reason we use the representation (3.2) and not the Kolmogorov forward equation is because the Kolmogorov forward equation in it's natural setting involves final conditions, and not initial conditions. Thus the standard method employed by probabilists is to make a $t = T - s$ substitution [11]. The process obtained in this manner will have the same one dimensional distribution as the process A_t above, however spatial covariances and gradients of the two processes will in general be different. The stochastic representation of the Navier–Stokes equations we describe below involves spatial gradients of the flow map A , and for this reason our representation will not be valid if we use the Feynman Kac formula.

We now use Proposition 3.1 to represent the solution to the Navier–Stokes equations as the expected value of a system that is nonlinear in the sense of McKean. The essential idea is to find a representation of the Euler equations involving particle trajectories [4], and then add noise and average as in Proposition 3.1 (as opposed to attempting to use the Kolmogorov forward equation).

Theorem 3.2. *Let $\nu > 0$, W be an n -dimensional Wiener process, $k \geq 1$ and $u_0 \in C^{k+1,\alpha}$ be a given deterministic divergence free vector field. Let the pair u, X satisfy the stochastic system*

$$dX_t = u_t dt + \sqrt{2\nu} dW_t, \tag{3.3}$$

$$A_t = X_t^{-1}, \tag{3.4}$$

$$u_t = \mathbf{EP}[(\nabla^* A_t)(u_0 \circ A_t)] \tag{3.5}$$

with initial data

$$X(a, 0) = a. \tag{3.6}$$

We impose boundary conditions by requiring u and $X - I$ are either spatially periodic, or decay at infinity. Then u satisfies the incompressible Navier–Stokes equations (1.1)–(1.2) with initial data u_0 .

Here \mathbf{P} in equation (3.5) denotes the Leray–Hodge projection onto divergence free vector fields [2] and $\nabla^* A_t$ denotes the transpose of the Jacobian of A_t . We remark that (3.5) is algebraically equivalent to

$$\omega_t = \mathbf{E}[(\nabla X_t)u_0] \circ A_t, \tag{3.7}$$

$$u_t = -\Delta^{-1}\nabla \times \omega \tag{3.8}$$

and (3.5) can be replaced with (3.7)–(3.8) in Theorem 3.2. Note that (3.8) is exactly the Biot–Savart law. When $\nu = 0$, Eq. (3.7) reduces to the well known vorticity transport for the Euler equations [2], and in this case (3.3)–(3.6) (or equivalently the system (3.3), (3.4), (3.6)–(3.8)) are exactly a Lagrangian formulation of the Euler equations [4].

² See also [5,15,6] for an elementary proof for flows of the type we consider here.

We do not prove Proposition 3.1 or Theorem 3.2 here, and we refer the reader to [5,15] instead. A generalization of Proposition 3.1 where the diffusion matrix is not spatially constant can be found in [15,6,19,21].

4. Decay of heat flows

In this section we prove a decay estimate for solutions to the heat equation with an incompressible drift. Our first estimate is an $L^\infty \rightarrow L^\infty$ estimate that is independent of the drift. A more general $L^1 \rightarrow L^\infty$ version of this estimate appeared for example in [7] and [9]. We provide a proof that follows the proof in [7] and keeps track of the dependence of the constants on viscosity and our length scale L .

Lemma 4.1. *Let $u \in C^1([0, T], \mathcal{I}^d)$ be divergence free, and θ be a solution to Eq. (3.1) with initial data θ_0 . If θ_0 is spatially periodic, mean 0, and the dimension $d \geq 3$, then there exists an constant $c = c(d)$ such that*

$$\|\theta_t\|_\infty \leq \frac{cL^d}{(vt)^{d/2}} \|\theta_0\|_\infty.$$

Proof. Let φ be mean zero and periodic, $p \geq \frac{d+2}{4}$ and $c = c(d, p)$ be a constant that changes from line to line. Then the Hölder, Poincaré and Sobolev in inequalities give

$$\int \varphi^2 = \int \varphi^{1/p} \varphi^{(2p-1)/p} \leq \|\varphi\|_{L^1}^{1/p} \|\varphi\|_{L^{(2p-1)/(p-1)}}^{(2p-1)/p} \leq cL^{(4p-d-2)/2p} \|\varphi\|_{L^1}^{1/p} \|\nabla \varphi\|_{L^2}^{(2p-1)/p}.$$

If we set $q = \frac{2}{2p-1}$ this gives

$$\|\nabla \varphi\|_{L^2}^2 \geq cL^{(qd-4)/2} \|\varphi\|_{L^2}^{2+q} \|\varphi\|_{L^1}^{-q}.$$

Now let θ' and θ'' to be solutions of (3.1) with initial conditions θ_0^- and θ_0^+ respectively. Integrating (3.1) immediately shows that $\int \theta'$ and $\int \theta''$ are conserved. Since θ' and θ'' are of constant sign, this means that $\|\theta'\|_{L^1}$ and $\|\theta''\|_{L^1}$ are conserved in time. Finally, the maximum principle implies that $\theta' \leq \theta \leq \theta''$, and hence $\|\theta\|_{L^1}$ is nondecreasing in time.

Thus multiplying (3.1) by θ and integrating over \mathcal{I}^d gives

$$\partial_t \|\theta\|_{L^2}^2 = -2v \|\nabla \theta\|_{L^2}^2 \leq -cvL^{(qd-4)/2} \|\theta\|_{L^1}^{-q} \|\theta\|_{L^2}^{2+q} \leq -cvL^{(qd-4)/2} \|\theta_0\|_{L^1}^{-q} \|\theta\|_{L^2}^{2+q}.$$

Dividing by $\|\theta\|_{L^2}^{2+q}$ and integrating in time gives

$$\|\theta\|_{L^2} \leq c \frac{L^{2/q-d/2}}{(vt)^{1/q}} \|\theta_0\|_{L^1}.$$

Let $\mathcal{P}_t(u)$ be the solution operator of (3.1). The above estimate shows

$$\|\mathcal{P}_t(u)\|_{L^1 \rightarrow L^2} \leq c \frac{L^{2/q-d/2}}{(vt)^{1/q}}.$$

Since u is divergence free the dual operator $\mathcal{P}^*(u) = \mathcal{P}(-u)$, and hence satisfies the same bound. Thus

$$\begin{aligned} \|\mathcal{P}_{2t}\|_{L^1 \rightarrow L^\infty} &\leq \|\mathcal{P}_t\|_{L^1 \rightarrow L^2} \|\mathcal{P}_t\|_{L^2 \rightarrow L^\infty} = \|\mathcal{P}_t\|_{L^1 \rightarrow L^2} \|\mathcal{P}_t^*\|_{(L^\infty)^* \rightarrow L^2} \\ &\leq \|\mathcal{P}_t\|_{L^1 \rightarrow L^2} \|\mathcal{P}_t^*\|_{L^1 \rightarrow L^2} \leq c \frac{L^{4/q-d}}{(vt)^{2/q}}. \end{aligned}$$

Hence

$$\|\theta\|_{L^\infty} \leq c \frac{L^{4/q-d}}{(vt)^{2/q}} \|\theta_0\|_{L^1} \leq c \frac{L^{4/q}}{(vt)^{2/q}} \|\theta_0\|_{L^\infty}.$$

Finally, $p \geq \frac{d+2}{4}$ is the same as $q \leq \frac{4}{d}$, and choosing $q = \frac{4}{d}$ concludes the proof. \square

Remark 4.2. When $d = 2$, $p \geq \frac{d+2}{4}$ needs to be replaced with $p > \frac{d+2}{4}$, and hence the above proof will show that for any $\varepsilon > 0$,

$$\|\theta_t\|_\infty \leq \frac{c_\varepsilon L^{d+\varepsilon}}{(\nu t)^{(d+\varepsilon)/2}} \|\theta_0\|_\infty.$$

Lemma 4.1 cannot directly be used in our proof of Theorem 2.1. This is because the relation between the velocity field u and the inverse flow map (Eq. (3.5)) involves the Leray–Hodge projection \mathbf{P} . The Leray–Hodge projection is a Calderón–Zygmund type singular integral operator and is known to be unbounded on C^k (see [22] for instance). However, singular integral operators, and in particular the Leray–Hodge projection is known to be bounded on Hölder spaces, and thus we now proceed to extend Lemma 4.1 to Hölder spaces.

Using the stochastic flows from [5,15] we obtain the Hölder space estimate we need in an elementary manner. We remark that the usual PDE methods [18] will only provide Hölder estimates that grow exponentially in time. The estimate we provide here will in general also grow exponentially in time, however decays in time when the viscosity is large, or drift U is small.

Lemma 4.3. Let $d \geq 3$ and $u \in C^{k+1,\alpha}([0, T], \mathcal{I}^d)$ be divergence free and define U by

$$U = \sup_{t \in [0, T]} \|u_t\|_{k+1,\alpha}. \tag{4.1}$$

Let $\theta_0 \in C(\mathcal{I}^d)$ have mean 0, and θ satisfy Eq. (3.1) with initial data θ_0 . Then there exists $T' = T'(\frac{U}{L}, d, k, \alpha)$ and a constant $c = c(\frac{U}{L}, d, k, \alpha)$ such that

$$\|\theta_t\|_{k+1,\alpha} \leq c \left(\frac{L^d}{(\nu t)^{d/2}} + \left(\frac{U t}{L} \right)^\alpha \right) \|\theta_0\|_{k+1,\alpha}$$

holds for all $t \in [0, T']$. If $d = 2$, the above estimate is still true if we replace $\frac{L^d}{(\nu t)^{d/2}}$ with $\frac{c_\varepsilon L^{d+\varepsilon}}{(\nu t)^{(d+\varepsilon)/2}}$ for any $\varepsilon > 0$.

Remark 4.4. Note that the growth term is independent of the viscosity, and the decay term is independent of the drift u .

Proof. We present the proof for $d \geq 3$. The $d = 2$ case will then follow by replacing d with $d + \varepsilon$. Define X, A by Eqs. (3.3) and (3.4) respectively. From [5,15] and uniqueness of strong solutions to (3.1) we know

$$\theta_t = \mathbf{E}\theta_0 \circ A_t.$$

Let $\ell = A - I$ be the Lagrangian displacement. First notice that if $f \in C^{0,\alpha}$ then Lemma A.4 shows

$$|f \circ A_t|_\alpha \leq c |f|_\alpha \left(\frac{U t}{L} \right)^\alpha \quad \text{a.s.} \tag{4.2}$$

Now, let m a multi index with $1 \leq |m| \leq k$. We note that $D^m \theta_t$ is a sum of terms of the form

$$D^n \theta_0|_{A_t} \prod_{1 \leq i \leq |n|} D^{n_i} \ell_t \quad \text{and} \quad D^n \theta_0|_{A_t}, \tag{4.3}$$

where n_i 's are multi indices with $|n_i| \geq 1$ and $|n| + \sum_i |n_i| = |m|$. By Proposition 3.1 we know that $\mathbf{E} D^n \theta_0|_{A_t}$ satisfies (3.1) with initial data $D^n \theta_0$, and hence by Lemma 4.1 we know

$$\|\mathbf{E}[D^n \theta_0] \circ A_t\|_{L^\infty} \leq \frac{c L^d}{(\nu t)^{d/2}} \|D^n \theta_0\|_{L^\infty}.$$

Thus using Lemma A.4, inequality (4.2) we have

$$\|\mathbf{E}[D^n \theta_0] \circ A_t\|_{0,\alpha} \leq c \left(\frac{L^d}{(\nu t)^{d/2}} + \left(\frac{U t}{L} \right)^\alpha \right) \|D^n \theta_0\|_{0,\alpha}. \tag{4.4}$$

Using (4.4) and Lemma A.4, we bound the remaining terms of (4.3), concluding the proof. \square

5. Global existence

In this section we prove Theorem 2.1. We start with a lemma involving bounds for the Leray–Hodge projection.

Lemma 5.1. *Let $k \geq 1$, and $A, v \in C^{k+1,\alpha}$ be such that $\nabla A, v$ are spatially periodic. There exists a constant $c = c(d, \alpha)$ such that*

$$\|\mathbf{P}[(\nabla^* A)v]\|_{k+1,\alpha} \leq c \|\nabla A\|_{k,\alpha} \|v\|_{k+1,\alpha}.$$

Proof. Since \mathbf{P} vanishes on gradients, we can ‘integrate by parts’ to avoid the loss of derivatives. Note

$$\mathbf{P}[(\nabla^* u)v] = \mathbf{P}[\nabla(u \cdot v) - (\nabla^* v)u] = -\mathbf{P}[(\nabla^* v)u]$$

for any $u, v \in C^1$. Thus we have

$$\partial_i \mathbf{P}[(\nabla^* A)v] = \mathbf{P}[(\nabla^* A)\partial_i v] - \mathbf{P}[(\nabla^* v)\partial_i A].$$

Since \mathbf{P} is Calderón–Zygmund singular integral operator, it is bounded on Hölder spaces [22,1]. Finally note that the right hand side only depends on first derivatives of A and v , and the lemma follows by taking Hölder norms. \square

We now prove Theorem 2.1. We restate it here for the readers convenience.

Theorem 2.1. *Let $k \geq 1, \alpha \in (0, 1)$ and $u_0 \in C^{k+1,\alpha}(\mathcal{I}^d)$ be spatially periodic, divergence free and have mean 0. Let $R = \frac{L}{\nu} \|u_0\|_{k+1,\alpha}$ be the Reynolds number of the flow. Then $\exists T = T(k, \alpha, d, \frac{1}{L} \|u_0\|_{k+1,\alpha})$ and $R_0 = R_0(k, \alpha, d)$ such that for all $0 < R < R_0$ the solution u of (1.1)–(1.2) with Reynolds number R , initial data u_0 and periodic boundary conditions is in $C^{k+1,\alpha}$ for time T , and satisfies*

$$\|u_T\|_{k+1,\alpha} \leq \|u_0\|_{k+1,\alpha}. \tag{2.1}$$

Proof. We assume that $d \geq 3$. The $d = 2$ case follows similarly by replacing d with $d + \varepsilon$. Let C, δ be the dimensional constants in Theorem A.5. We let $U = C \|u_0\|_{k+1,\alpha}$, and choose T such that $\frac{UT}{L} < \delta$. By Theorem A.5, there exist a pair of $C^{k+1,\alpha}$ functions $X, u : [0, T] \rightarrow \mathcal{I}^d$ which are the unique (strong) solution to (3.3)–(3.6). Recall that $\|u_t\|_{k+1,\alpha} \leq U$ for all $t \in [0, T]$.

From equation (3.5) we see

$$u_t = \mathbf{E} \mathbf{P} u_0 \circ A_t + \mathbf{E} \mathbf{P} (\nabla^* \ell) u_0 \circ A_t.$$

Let $c = c(k, \alpha, d)$ be a constant that changes from line to line. Applying Lemmas 5.1 and A.4 to the second term we have

$$\|u_t\|_{k+1,\alpha} \leq c \|\mathbf{E} u_0 \circ A_t\|_{k+1,\alpha} + c \mathbf{E} \|\nabla^* \ell_t\|_{k,\alpha} \|u_0 \circ A_t\|_{k+1,\alpha}$$

and hence by Lemma 4.3 we have

$$\|u_t\|_{k+1,\alpha} \leq c \left(\frac{L^d}{(\nu t)^{d/2}} + \left(\frac{U t}{L} \right)^\alpha \right) \|u_0\|_{k+1,\alpha}.$$

Minimizing $\frac{L^d}{(\nu t)^{d/2}} + \left(\frac{U t}{L} \right)^\alpha$ in time shows that the minimum value is attained at $t_0 = \frac{cL}{U} R^{d/(2\alpha+d)}$, and the minimum value is $c R^{\alpha d/(2\alpha+d)}$. Thus we can choose R small enough to ensure $t_0 < T$ and equation (2.1) is satisfied. \square

Acknowledgement

I would like to thank Lenya Ryzhik for pointing out [7] which provides the elegant proof of Lemma 4.1 reproduced here.

Appendix A. Bounds for the Lagrangian displacement

In this section, we prove bounds on $\|\nabla X - \mathbb{I}\|_{k,\alpha}$. The estimates proved here are elementary, and are taken directly from [14,15]. We reproduce them here for completeness and the readers convenience.

We remark that the estimates provided here were used in [14,15] to prove local existence for the system (3.3)–(3.6). As the local existence proof is a little lengthier, we do not reproduce it here.

Lemma A.1. *Let X be a Banach algebra. If $x \in X$ is such that $\|x\| \leq \rho < 1$ then $1+x$ is invertible and $\|(1+x)^{-1}\| \leq \frac{1}{1-\rho}$. Further if in addition $\|y\| \leq \rho$ then*

$$\|(1+x)^{-1} - (1+y)^{-1}\| \leq \frac{1}{(1-\rho)^2} \|x-y\|.$$

Proof. The first part of the lemma follows immediately from the identity $(1+x)^{-1} = \sum (-x)^n$. The second part follows from the first part and the identity

$$(1+x)^{-1} - (1+y)^{-1} = (1+x)^{-1}(y-x)(1+y)^{-1}. \quad \square$$

Lemma A.2. *If $k \geq 1$, then there exists a constant $c = c(k, \alpha)$ such that*

$$\|f \circ g\|_{k,\alpha} \leq c \|f\|_{k,\alpha} (1 + \|\nabla g\|_{k-1,\alpha})^{k+\alpha}$$

and

$$\|f \circ g_1 - f \circ g_2\|_{k,\alpha} \leq c \|\nabla f\|_{k,\alpha} (1 + \|\nabla g_1\|_{k-1,\alpha} + \|\nabla g_2\|_{k-1,\alpha})^{k+1} \cdot \|g_1 - g_2\|_{k,\alpha}.$$

The proof of Lemma A.2 is elementary and not presented here.

Lemma A.3. *Let $X_1, X_2 \in C^{k+1,\alpha}$ be such that*

$$\|\nabla X_1 - \mathbb{I}\|_{k,\alpha} \leq d < 1 \quad \text{and} \quad \|\nabla X_2 - \mathbb{I}\|_{k,\alpha} \leq d < 1.$$

Let A_1 and A_2 be the inverse of X_1 and X_2 , respectively. Then there exists a constant $c = c(k, \alpha, d)$ such that

$$\|A_1 - A_2\|_{k,\alpha} \leq c \|X_1 - X_2\|_{k,\alpha}.$$

Proof. Let $c = c(k, \alpha, d)$ be a constant that changes from line to line (we use this convention implicitly throughout this paper). Note first $\nabla A = (\nabla X)^{-1} \circ A$, and hence by Lemma A.1

$$\|\nabla A\|_{C^0} \leq \|(\nabla X)^{-1}\|_{C^0} \leq c.$$

Now using Lemma A.1 to bound $\|(\nabla X)^{-1}\|_{0,\alpha}$ we have

$$\|\nabla A\|_{0,\alpha} = \|(\nabla X)^{-1} \circ A\|_{0,\alpha} \leq \|(\nabla X)^{-1}\|_{0,\alpha} (1 + \|\nabla A\|_{C^0}) \leq c.$$

When $k \geq 1$, we again bound $\|(\nabla X)^{-1}\|_{k,\alpha}$ by Lemma A.1. Taking the $C^{k,\alpha}$ norm of $(\nabla X)^{-1} \circ A$ we have

$$\|\nabla A\|_{k,\alpha} \leq \|(\nabla X)^{-1}\|_{k,\alpha} (1 + \|\nabla A\|_{k-1,\alpha})^k.$$

So by induction we can bound $\|\nabla A\|_{k,\alpha}$ by a constant $c = c(k, \alpha, d)$. The lemma now follows by applying Lemma A.2 to the identity

$$A_1 - A_2 = (A_1 \circ X_2 - I) \circ A_2 = (A_1 \circ X_2 - A_1 \circ X_1) \circ A_2. \quad \square$$

Lemma A.4. *Let $u \in C([0, T], C^{k+1,\alpha})$ and X satisfy the SDE (3.3) with initial data (3.6). Let $\lambda = X - I$ and $U = \sup_t \|u(t)\|_{k+1,\alpha}$. Then there exists $T = T(k, \alpha, \frac{U}{L})$ and $c = c(k, \alpha, \frac{U}{L})$ such that for $t \leq T$*

$$\|\nabla \lambda(t)\|_{k,\alpha} \leq \frac{cUt}{L} \quad \text{and} \quad \|\nabla \ell(t)\|_{k,\alpha} \leq \frac{cUt}{L}$$

hold almost surely.

Proof. From Eq. (3.3) we have

$$X(x, t) = x + \int_0^t u(X(x, s), s) ds + \sqrt{2\nu} W_t \Rightarrow \nabla X(t) = I + \int_0^t (\nabla u) \circ X \cdot \nabla X. \quad (\text{A.1})$$

Taking the C^0 norm of Eq. (A.1) and using Gronwall's Lemma we have

$$\|\nabla \lambda(t)\|_{C^0} = \|\nabla X(t) - I\|_{C^0} \leq e^{Ut/L} - 1.$$

Now taking the $C^{k,\alpha}$ norm in Eq. (A.1) we have

$$\|\nabla \lambda(t)\|_{k,\alpha} \leq c \int_0^t \|\nabla u\|_{k,\alpha} (1 + \|\nabla \lambda\|_{k-1,\alpha})^{k+\alpha} (1 + \|\nabla \lambda\|_{k,\alpha}).$$

The bound for $\|\nabla \lambda\|_{k,\alpha}$ now follows from the previous two inequalities, induction and Gronwall's Lemma. The bound for $\|\nabla \ell\|_{k,\alpha}$ then follows from Lemma A.3.

We draw attention to the fact that the above argument can only bound $\nabla \lambda$, and not λ . Fortunately, our results only rely on a bound of $\nabla \lambda$. \square

We conclude this appendix by stating a slightly modified version theorem which appeared in [14]. The only modification we make is that we trace the dependence of the constants in [14] to dimension less quantities, instead of absolute ones. The proof that appeared in [14] goes through verbatim.

Theorem A.5. *Let $k \geq 1$ and $u_0 \in C^{k+1,\alpha}$ be divergence free. There exists absolute constants $\delta = \delta(k, \alpha, d)$ and $C = C(k, \alpha, d)$ such that for $U = C\|u_0\|_{k+1,\alpha}$, and any T such that $\frac{UT}{L} < \delta$ there exist a pair of functions a pair of functions $\lambda, u \in C([0, T], C^{k+1,\alpha})$ such that u and $X = I + \lambda$ satisfy the system (3.3)–(3.6). Further for all $t \in [0, T]$ we have $\|u_t\|_{k+1,\alpha} \leq U$.*

References

- [1] A.P. Calderón, A. Zygmund, Singular integrals and periodic functions, *Studia Math.* 14 (1954) 249–271 (1955).
- [2] A.J. Chorin, J.E. Marsden, *A Mathematical Introduction to Fluid Mechanics*, third ed., Texts in Applied Mathematics, vol. 4, Springer-Verlag, New York, 1993.
- [3] P. Constantin, C. Foias, *Navier–Stokes equations*, Chicago Lectures in Mathematics, University of Chicago Press, Chicago, IL, 1988.
- [4] P. Constantin, An Eulerian–Lagrangian approach for incompressible fluids: local theory, *J. Amer. Math. Soc.* 14 (2) (2001) 263–278 (electronic).
- [5] P. Constantin, G. Iyer, A stochastic Lagrangian representation of the 3-dimensional incompressible Navier–Stokes equations, *Comm. Pure Appl. Math.* (2006), in press, available at arXiv:math.PR/0511067.
- [6] P. Constantin, G. Iyer, Stochastic Lagrangian transport and generalized relative entropies, *Commun. Math. Sci.* 4 (4) (2006) 767–777.
- [7] P. Constantin, A. Kiselev, L. Ryzhik, A. Zlatoš, Diffusion and mixing in fluid flow, *Ann. Math.* (2006), in press, available at arXiv: math.AP/0509663.
- [8] T. Kato, H. Fujita, On the nonstationary Navier–Stokes system, *Rend. Sem. Mat. Univ. Padova* 32 (1962) 243–260.
- [9] A. Fannjiang, A. Kiselev, L. Ryzhik, Quenching of reaction by cellular flows, *Geom. Funct. Anal.* 16 (1) (2006) 40–69.
- [10] C.L. Fefferman, Existence and smoothness of the Navier–Stokes equation, in: *The Millennium Prize Problems*, Clay Math. Inst., Cambridge, MA, 2006, pp. 57–67.
- [11] M. Freidlin, *Functional Integration and Partial Differential Equations*, Annals of Mathematics Studies, vol. 109, Princeton University Press, Princeton, NJ, 1985.
- [12] A. Friedman, *Stochastic Differential Equations and Applications*. Vol. 1, Probability and Mathematical Statistics, vol. 28, Academic Press Harcourt Brace Jovanovich Publishers, New York, 1975.
- [13] I. Gallagher, J.-Y. Chemin, On the global wellposedness of the 3-D Navier–Stokes equations with large initial data, Preprint, available at arXiv: math.AP/0508374, 2005.
- [14] G. Iyer, A stochastic perturbation of inviscid flows, *Commun. Math. Phys.* 266 (3) (2006) 631–645, available at arXiv: math.AP/0505066.
- [15] G. Iyer, A stochastic Lagrangian formulation of the Navier–Stokes and related transport equations, Ph.D. Thesis, University of Chicago, 2006.
- [16] I. Karatzas, S.E. Shreve, *Brownian Motion and Stochastic Calculus*, second ed., Graduate Texts in Mathematics, vol. 113, Springer-Verlag, New York, 1991.
- [17] H. Koch, D. Tataru, Well-posedness for the Navier–Stokes equations, *Adv. Math.* 157 (1) (2001) 22–35.

- [18] N.V. Krylov, *Lectures on Elliptic and Parabolic Equations in Hölder Spaces*, Graduate Studies in Mathematics, vol. 12, American Mathematical Society, Providence, RI, 1996.
- [19] N.V. Krylov, B.L. Rozovskiĭ, Stochastic partial differential equations and diffusion processes, *Uspekhi Mat. Nauk* 37 (6(228)) (1982) 75–95 (in Russian).
- [20] H. Kunita, *Stochastic Flows and Stochastic Differential Equations*, Cambridge Studies in Advanced Mathematics, vol. 24, Cambridge University Press, Cambridge, 1997. Reprint of the 1990 original.
- [21] B.L. Rozovskiĭ, *Stochastic Evolution Systems: Linear Theory and Applications to Nonlinear Filtering*, Mathematics and its Applications (Soviet Series), vol. 35, Kluwer Academic Publishers Group, Dordrecht, 1990. Translated from the Russian by A. Yarkho.
- [22] E.M. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton Mathematical Series, vol. 30, Princeton University Press, Princeton, NJ, 1970.