

Nonlinear diffusion from a delocalized source: affine self-similarity, time reversal, & nonradial focusing geometries [☆]

Jochen Denzler ^a, Robert J. McCann ^{b,*}

^a *Department of Mathematics, University of Tennessee at Knoxville, TN 37996-1300, USA*

^b *Department of Mathematics, University of Toronto, Ontario, Canada M5S 2E4*

Received 3 August 2006; accepted 19 May 2007

Available online 1 September 2007

Abstract

A family of explicit solutions is described, to the porous medium equation in its full range of nonlinearities (plus some analogous fourth-order diffusions), in which the pressure is given by a quadratic function of space at each instant in time. These include spreading solutions whose source is concentrated on any conic region of dimension lower than the ambient space, and solutions which focus at conic regions. The singular limiting distributions are affine projections of Barenblatt type solutions (with arbitrary signature) onto lower dimensional subspaces. All affine images of Barenblatt solutions form an invariant space on which the dynamics can be integrated explicitly. A time-reversal symmetry is revealed for the pressure equation which transforms spreading solutions to focusing solutions, and vice-versa. This yields new information about the long and short time asymptotics of finite-mass solutions, about the instability of focusing, and about singularity geometry.

© 2007 L'Association Publications de l'Institut Henri Poincaré. Published by Elsevier B.V. All rights reserved.

Résumé

On décrit une famille de solutions explicites pour l'équation des milieux poreux pour toute l'échelle des nonlinéarités (ainsi que pour d'autres équations de diffusion analogues de quatrième degré). Pour ces solutions, la pression à chaque instant est une fonction quadratique en x . Certaines de ces solutions sont diffusées à partir d'une source concentrée sur un domaine conique arbitraire de dimension inférieure à celle de l'espace ambiant ; d'autres se focalisent sur des régions coniques. Pour les solutions focalisantes, on obtient à la limite une distribution singulière qui est une projection affine d'une solution de type Barenblatt (à signature arbitraire) sur un espace de dimension inférieure. L'ensemble des images affines des solutions de type Barenblatt est un espace invariant, sur lequel la dynamique peut s'intégrer explicitement. Par un changement de variables qui inverse le sens du temps, on arrive à transformer une solution diffusive en une solution focalisante. Ceci donne de nouvelles informations sur l'asymptotique en temps court et long pour des solutions de masse finie, sur l'instabilité des solutions focalisantes, ainsi que sur la géométrie des singularités.

© 2007 L'Association Publications de l'Institut Henri Poincaré. Published by Elsevier B.V. All rights reserved.

[☆] This research was supported in part by Natural Sciences and Engineering Research Council of Canada Grant 217006-03 and United States National Science Foundation Grant DMS 0354729. RJM is pleased to acknowledge the hospitality of the Mathematical Sciences Research Institute (Berkeley, CA), the Institut Henri Poincaré (Paris), and the Université Paul Sabatier (Toulouse) where parts of this work were performed. The authors thank Professors Grigory Barenblatt, Richard Hamilton, and Herbert Koch for fruitful discussions, and Dragos Iftimie for help with the résumé français; they are especially grateful to Juan Luis Vázquez for bringing the work of Luc Tartar [59] to their attention during the Banff International Research Station workshop (April 15–20, 2006) at which our initial results were announced, and to the referee for further useful literature suggestions which led us to [18,47] and eventually [37,56,57,69].

* Corresponding author.

E-mail addresses: denzler@math.utk.edu (J. Denzler), mccann@math.toronto.edu (R.J. McCann).

Keywords: Porous medium equation; Nonlinear diffusion; Invariant subspaces; Fourth order diffusion; Hole-filling; Asymptotic dynamics; Exact nonradial solutions; Affinely self-similar dynamics; Ellipsoidal level sets

1. Introduction

Since the 1950s, the nonlinear diffusion equation

$$\frac{\partial \rho}{\partial t} = \frac{1}{m} \Delta(\rho^m) \quad (1)$$

has been a central model for strongly viscous spreading phenomena. With various boundary conditions and different interpretations for the evolving density $\rho(\mathbf{x}, t) \geq 0$ on $\mathbf{R}^n \times [0, \infty[$, this *porous medium* equation has been used to model population spreading, fluid seeping through soil ($m = 2$), creeping thin films ($m = 4$), heat propagation in plasmas ($m = 6$), avalanches in sandpiles ($m = 1 - \frac{n}{2}$), and geometrical curvature flows ($m = \frac{n-2}{n+2}$). An understanding of this equation is now well-developed; with $m > 1$, its phenomenology includes finite speed propagation of fluid into dry regions (or heat into cold regions), contrasting sharply with the linear diffusion $m = 1$. The radial self-similar solution to these equations discovered long ago [8,68] enjoys a prominent role in this theory: it models the eventual behavior of finite-mass spreading solutions, whether or not they enjoy spherical symmetry initially [36]. See Barenblatt [9] and Vázquez [63,64] for reviews.

More recently, (1) has emerged as a model for focusing phenomena, like the filling in of a dry spot by thin lava-like films in the viscous gravity current experiments of Diez et al. [35], or the eventual disappearance of groundwater from soil into fissures in the filtration-absorption model of Barenblatt et al. [10]. For viscous gravity currents, careful experiments supported by detailed numerics [35,5] leave little doubt that typical holes become more and more elongated as they shrink, eventually disappearing nonselfsimilarly. However, few explicit solutions have been known with this form, so one has only a formal asymptotic construction called the *closing-eye solution* by Angenent et al. [5] as a model. (Self-similar solutions with k -fold axial symmetry were shown to exist for $k \geq 3$ and $m > m_k > 1$ sufficiently large by Angenent and Aronson [4]; they are not explicit, and they capture only the bifurcation from roundness [7] and not the generic disappearances observed, which correspond rather to a $k = 2$ instability for all $m > 1$.)

The present manuscript describes closed-form solutions to (1) which are not radially symmetric, but enjoy a property we call *affine* self-similarity, meaning the density at each instant is a sheared image of its initial profile. These solutions were discovered independently some twenty years ago by Luc Tartar, who described them in an unpublished manuscript [59], which we learned of only after this work had been largely completed; his solutions are also documented and developed in the forthcoming book of Vázquez [65]. Unbeknownst to Tartar and to us, for two space dimensions these solutions had already appeared in the work of Titov and Ustinov [69]; they were rediscovered independently not only by us but by King [47], who worked out the details primarily for two space dimensions, by Rudykh and Semenov [57] and by Pukhnachev [56], apparently as unaware as we were of the work of Tartar. In this respect, as in others, Titov, Ustinov and Tartar were ahead of their time; many of the applications we discuss, which enhance the relevance of these solutions, had not yet been conceived. Like Tartar's, our solutions are obtained by quadrature; in suitable coordinates they remain reflection symmetric with respect to each spatial variable, and form an $\frac{n(n+1)}{2}$ parameter family whose evolution is completely encoded in the finite-dimensional dynamics of the shearing maps. Their level sets remain conic sections (ellipsoids or hyperboloids) for as long as the solutions exist. The spatial dimensions of these conics evolve at different rates, given by a coupled nonlinear system of first order differential equations (8), until one or more of these dimensions degenerates to zero. The number and signature of the dimensions which collapse, and their relative rates, determine the geometry of the focusing singularity if it occurs. These rates are not exact power laws, except in the classical case of spherical focusing or spreading solutions, but they are asymptotic to power laws at the time of collapse, with scaling exponents that we calculate. In suitable variables the affine dynamics can actually be integrated independently of m , whose effect is therefore reduced to prescribing how the affine orbits are parameterized.

Unfortunately, these affinely self-symmetric solutions have no nontrivial continuation beyond the time of the singularity, so do not replace the solutions of Gravelleau [42,6] and Angenent et al. [5] as models for hole-filling in the experiments of Diez et al. [35]. However, they provide appropriate solutions to the filtration-absorption problem of Barenblatt et al. [10], since our solutions remain zero after the last of the groundwater drains into the fissures. Moreover, our solutions predict that any perturbation from initial roundness will be amplified as the fluid drains, leading to

a last drop of fluid which becomes more and more elongated as it is absorbed, eventually disappearing in the form of a flattened ellipsoid, with one axis collapsed and the others having generically different positive lengths. This reflects the fact that the smallest dimension collapses faster than the others, since surface tension is absent from the model. This is in marked contrast to the *linearized* stability result for circular drainage asserted by Chertock [28].

These affinely self-similar solutions also provide new information concerning the long-time asymptotics of the spreading solution, a subject of considerable interest; see for example the works [2,19,25,22,24,32,31,45,50,54,60] of Angenent, Cáceres, Carrillo, Del Pino, Di Francesco, Dolbeault, Jüngel, Kim, Markowich, Otto, Slepčev, Toscani, Unterreiter and Vázquez. In the porous medium regime $m > 1$, they suggest that normalized initial profiles with compact support and the same center of mass evolve toward each other at rate $t^{-2/(2+(m-1)n)}$ in $L^1(\mathbf{R}^n)$ —the square of the rate for uncentered solutions [23,31,54]—and certainly no faster than that in dimensions $n > 1$. This rate is consistent with the extrapolation of the linearized dynamics we computed in [33] for the fast diffusion equation $1 - 2/n < m < 1$, but due to an eigenvalue crossing at $m = 1$ it is slower than the $1/t$ rate established at both extremes [46,50] of that regime. It also explains Angenent and Aronson’s observation, that for nonradial solutions with $m \geq 1$ [3], dimensions $n \geq 2$ are distinguished from one-dimension [67,62,2], in that no choice of delay t_0 improves the rate of decay beyond that obtained by centering the data.

A final section discusses analogous solutions to certain fourth-order diffusion equations (22) and (27) which parallel the porous medium family. These equations arise as gradient flows of the generalized Fisher information (28) with respect to the 2-Wasserstein distance. In particular, the special case $m = 3/2$ is the thin film equation, for which the $n = 2$ dimensional solutions were found independently by Betelú and King [18], while the case $m = 1$ is a quantum drift diffusion model studied by Jüngel and Pinnau in one-dimension (see [44] for references); it is also the generalization to higher dimensions of a model proposed by Derrida, Lebowitz, Speer and Spohn for the statistical mechanical fluctuations around a pinned interface [34].

2. Solution concept

The evolution (1) can also be written as a conservation law

$$\frac{\partial \rho}{\partial t} - \nabla \cdot (\rho \nabla \pi) = 0, \tag{2}$$

where $\pi = \gamma \rho^{1/\gamma}$ is the *pressure* and $\gamma = 1/(m - 1)$ is the adiabatic exponent of the gas (also called the polytropic index, but not consistently through the literature). The pressure satisfies

$$\frac{\partial \pi}{\partial t} = \frac{1}{\gamma} \pi \Delta \pi + |\nabla \pi|^2 \tag{3}$$

whenever the density $\rho(\mathbf{x}, t)$ satisfies (1). The unusual normalization factor γ in $\pi = \gamma \rho^{1/\gamma}$ is convenient both from the following mathematical and physical points of view: It allows us to avoid case distinctions between the porous medium ($m > 1 \Leftrightarrow \gamma > 0$) and fast diffusion ($m < 1 \Leftrightarrow \gamma < 0$) regimes, while guaranteeing that matter moves from higher to lower pressures independently of the sign of the pressure.

We argue that the truncation $\pi_* = \gamma(\pi/\gamma)_+ := \gamma \max\{\pi/\gamma, 0\}$ of any smooth solution $\pi(\mathbf{x}, t)$ also represents a physically meaningful notion of solution for (3), provided $\nabla \pi$ does not vanish on the zero set of π , and hence leads to a nonnegative solution ρ for (1): Indeed, if π solves (3), then the zero set $\mathcal{Z}_t := \{\mathbf{x} \in \mathbf{R}^n \mid \pi(\mathbf{x}, t) = 0\}$, which is a manifold if $\nabla \pi \neq 0$, moves in normal direction with velocity $-\pi_t \nabla \pi / |\nabla \pi|^2 = -\nabla \pi$, because $\pi \Delta \pi = 0$ on the interface. Moreover, it is exactly consistent with Darcy’s law as expressed in (2), so the support changes exactly as it should under Darcy’s law.

We are not purporting any general theory for (3), which is parabolic in forward time only where $\pi/\gamma > 0$, and parabolic in backwards direction only where $\pi/\gamma < 0$ and therefore ill-posed in the absence of a sign condition on the data. Rather, we construct nonlocal source solutions for (1) in the physically meaningful range $\pi/\gamma > 0$ from explicit (quadratic) solutions to (3) by means of truncation of a sporadic classical solution to (3). Or equivalently, by time reversal $-\pi(\mathbf{x}, -t) =: \tilde{\pi}(\mathbf{x}, t)$, solutions with $\pi/\gamma < 0$ which evolve backwards in time.

Example 1 (*Focusing by time-reversed Barenblatt*). The $\gamma > 0$ self-similar solution of Barenblatt [8], Pattle [55], Zel’dovich and Kompaneets [68] corresponds to (the positive truncation of) a ‘downwardly oriented’ quadratic pressure

$$\pi_b(\mathbf{x}, t) = \frac{b\gamma}{|t|^{n/(n+2\gamma)}} - \frac{\gamma}{(n+2\gamma)} \frac{|\mathbf{x}|^2}{|2t|}, \quad t > 0. \tag{4}$$

The mass of $\rho = (\pi_b/\gamma)_+^\gamma$ is determined by $b > 0$; its zero set spreads at rate $|\mathbf{x}| \sim t^{\gamma/(n+2\gamma)}$. Its negated time-reversal

$$\bar{\pi}_b(\mathbf{x}, t) = -\frac{b\gamma}{|t|^{n/(n+2\gamma)}} + \frac{\gamma}{(n+2\gamma)} \frac{|\mathbf{x}|^2}{|2t|}, \quad t < 0$$

is an ‘upward’ quadratic pressure, whose positive truncation vanishes in a ball shrinking to zero like $|t|^{\gamma/(n+2\gamma)}$ at the moment $t \rightarrow 0$ of the focusing singularity.

Note that the time-reversed solution described above has no nontrivial continuation beyond the singularity, so is a poor model for hole filling. Nevertheless, it is important to note that the focusing geometry is unstable precisely because it is the backwards in time flow of the spreading solution, which was shown by Friedman and Kamin [36] and Vázquez [64] to be a global attractor. In particular this implies that perturbations from sphericity will increase.

The partial differential equation $\partial u/\partial t = u\Delta u + \gamma|\nabla u|^2$, which is simply (3) with $u = \pi/\gamma$, has been studied in the fast diffusion range $\gamma \leq 0$ by Bertsch, Dal Passo and Ughi [15,17,16], using methods from Dal Passo and Luckhaus [30], and earlier in one dimension in [61]. (Their γ is our $-\gamma$.) They define a nonnegative function u as a weak solution for initial value $0 \leq u_0 \in L^\infty_{\text{loc}} \cap C(\mathbf{R}^n)$ if

$$u \in L^\infty_{\text{loc}}([0, \infty[\times\mathbf{R}^n) \cap L^2_{\text{loc}}([0, \infty[\rightarrow W^{1,2}_{\text{loc}}(\mathbf{R}^n)) \quad \text{and} \tag{5}$$

$$\int u_0 \psi(0, \cdot) \, d\mathbf{x} + \iint \left(u \frac{\partial \psi}{\partial t} - u \nabla u \cdot \nabla \psi + (\gamma - 1)|\nabla u|^2 \psi \right) \, d\mathbf{x} \, dt = 0$$

for all $\psi \in C^{1,1}([0, \infty[\times\mathbf{R}^n)$ with compact support in $[0, \infty[\times\mathbf{R}^n$. This definition makes sense for either sign of γ . We prove in an appendix that the solutions we find are weak solutions in the sense of (5). Bertsch, Dal Passo and Ughi actually require $u \in L^\infty([0, \infty[\times\mathbf{R}^n)$, but the distinction between L^∞ and L^∞_{loc} is not essential. Since only some, but not all, of our solutions are globally bounded, we prefer the L^∞_{loc} formulation (5).

Even for bounded pressures, the Cauchy problem features vast nonuniqueness of weak solutions [16] for fast diffusion ($\gamma \leq 0$): Roughly speaking, any smoothly shrinking support for u can be prescribed. The unique viscosity solution constructed in [17] has constant support in time. This will not be true for the fast diffusions we construct which have bounded pressure, so they are not viscosity solutions. We also construct solutions with nonvanishing unbounded pressures which, analogously to (4), lead to classical solutions of the fast diffusion equation in the whole space. Finally, for $\gamma < 0$, we construct signed solutions $\pi(\mathbf{x}, t)$ which are neither bounded above nor below; these correspond to unbounded densities $\rho = (\pi/\gamma)_+^\gamma \geq 0$ which are infinite (singular) on the set $\pi > 0$. Solutions of this kind have singular sets which expand, and were studied by Chasseigne and Vázquez in [27, §8].

The results [16] of Bertsch, Dal Passo and Ughi (namely solutions with shrinking support) are only intended for the fast diffusion case $\gamma \leq 0$, and in the density formulation they translate to unbounded focusing solutions. We construct such solutions (unbounded, but this time locally bounded) for the porous medium case ($\gamma > 0$). But we also construct bounded pressure solutions with spreading support, analogous (and forward asymptotic) to the Barenblatt solutions (4). Translated back into density variables, they give weak solutions to the porous medium equation with spreading support. Recall a weak solution [64] to the porous medium equation (1) with $m > 1$ is a function $\rho \geq 0$ such that $\rho^m, \nabla(\rho^m) \in L^1_{\text{loc}}(\mathbf{R}^n \times [0, \infty[)$ and

$$\int \varphi(\cdot, 0) \rho_0 \, d\mathbf{x} + \iint \left(\rho \frac{\partial \varphi}{\partial t} - \frac{1}{m} \nabla \rho^m \cdot \nabla \varphi \right) \, d\mathbf{x} \, dt = 0 \tag{6}$$

for all $\varphi \in C^1([0, \infty[\times\mathbf{R}^n)$ with compact support in $[0, \infty[\times\mathbf{R}^n$. Lemma 11 of the appendix below asserts that weak solutions $u(\mathbf{x}, t) \geq 0$ of the pressure evolution (5) give rise to weak solutions $\rho = u^\gamma$ of the density evolution (6). Thus our solutions $\pi(\mathbf{x}, t)$, after truncation, give weak porous medium flows $\rho = (\pi/\gamma)_+^\gamma$.

3. Results

We now show that any quadratic initial pressure $\pi(\mathbf{x}, 0)$ leads to a family of classical solutions

$$\pi(\mathbf{x}, t) = \frac{1}{2} \langle \mathbf{x}, \mathbb{P}(t)\mathbf{x} \rangle - \frac{1}{2} p_0(t) \tag{7}$$

which are quadratic in space at each instant in time; here $\mathbb{P}(t)$ is a symmetric $n \times n$ matrix. It costs no generality to exclude a linear term in \mathbf{x} from (7), which corresponds to centering the solution, nor to choose coordinates on \mathbf{R}^n which diagonalize the initial matrix $\mathbb{P}(0)$. The truncation π_* of (7) with the same sign as γ yields the pressure corresponding to the affinely self-similar solutions we are interested in.

The quadratic pressure ansatz (7) reduces the partial differential equation (3) to an ordinary differential equation

$$\frac{d\mathbb{P}}{dt} = \frac{\text{tr } \mathbb{P}}{\gamma} \mathbb{P} + 2\mathbb{P}^2 \tag{8}$$

for the matrix $\mathbb{P}(t)$, plus a condition determining the constant

$$\frac{dp_0}{dt} = p_0 \frac{\text{tr } \mathbb{P}}{\gamma}. \tag{9}$$

For functional calculus purposes below, note from (8) that $d\mathbb{P}/dt$ commutes with \mathbb{P} . The standard theory of differential equations guarantees that this system admits a smooth solution for short times; the only question is under what conditions the quadratic right-hand side causes some entries in the matrix to blow-up in a finite time. Remarkably, as Tartar also discovered [59], a nonlinear change of variables reduces this system to a single quadrature.

The symmetric matrix $t \mapsto \mathbb{P}(t)$ can be diagonalized in a flow-invariant manner, and we assume therefore, with no loss of generality, that $\mathbb{P}(t)$ is a diagonal matrix. Let its diagonal elements be called p_1, p_2, \dots, p_n . The hyperplanes $p_i = p_j$ are clearly invariant, and so are the hyperplanes $p_i = 0$. Therefore any ordering of the p_i and the constant 0 in the initial values will be preserved under the flow within the maximal interval of existence. If any p_j vanishes, then it vanishes identically and can be dropped from the ODE, reducing the dimension and not affecting the other coordinates. We will therefore assume, with no loss of generality, that none of the p_j vanishes.

Under this hypothesis, we first deduce invariance of the quantity (10). This expresses the height $p_0(t)$ of the pressure plateau in terms of the evolution of the matrix $\mathbb{P}(t)$. Note the *moment index* $\mu := n + 2\gamma$ used in our previous works [32,33,45,46,50] (where it was denoted by $-p = n + 2\gamma$) enters naturally as a scaling exponent. As Lemma 2 indicates, the conserved ratio (10) indexes the total mass $\|\rho_0\|_{L^1(\mathbf{R}^n)} \leq \infty$ of the density $\rho = (\pi/\gamma)_+^\gamma$ in (1).

Lemma 1 (*Evolution of pressure plateau*). *Fix $\gamma \neq 0$ and let $\pi(\mathbf{x}, t) = \frac{1}{2}(\langle \mathbf{x} \mathbb{P}(t) \mathbf{x} \rangle - p_0(t))$ satisfy (8)–(9) on an interval $]t_\alpha, t_\omega[\subset \mathbf{R}$. If $\det \mathbb{P}(t) \neq 0$ then*

$$z^2 = |p_0(t)| / |\det \mathbb{P}(t)|^{1/(n+2\gamma)} = \text{const}. \tag{10}$$

Proof. There is no problem solving (9) on any interval of time where (8) has a smooth solution. Reformulating these dynamics as

$$\frac{d}{dt} \log |\mathbb{P}| = \frac{\text{tr } \mathbb{P}}{\gamma} \mathbb{I} + 2\mathbb{P}, \tag{11}$$

the invariance of (10) follows by comparing

$$\frac{d}{dt} \log |p_0| = \frac{\text{tr } \mathbb{P}}{\gamma},$$

with the trace of the first equation:

$$\frac{d}{dt} \log \det |\mathbb{P}| = (n + 2\gamma) \frac{\text{tr } \mathbb{P}}{\gamma}. \quad \square$$

An equally elementary theorem holds the key to describing the dynamics. Before proving this theorem, observe that if $\mathbb{P}(t) \geq 0$ and $p_0(t) > 0$, the zero set \mathcal{Z}_t of the pressure (7) is given by $|\mathbb{P}(t)^{1/2} \mathbf{x}| = p_0(t)^{1/2}$. So $|\mathbb{P}(t)/p_0(t)|^{-1/2}$ scales as a length. More generally, the invariance of (10), which corresponds to conservation of mass, implies $S(t) := -|\det \mathbb{P}(t)|^{1/(n+2\gamma)} \mathbb{P}(t)^{-1}$ scales as a length squared. The matrix S will turn out to trivialize the dynamics. To relate this matrix directly to the evolving profile $\rho = (\pi/\gamma)_+^\gamma$, order the eigenvalues of $\mathbb{P}(t)$ and $S(t)$ so that $p_1(t) \geq p_2(t) \geq \dots \geq p_n(t)$ and hence

$$0 > s_1(t) \geq s_2(t) \geq \dots \geq s_j(t) > -\infty; \quad +\infty > s_{j+1}(t) \geq \dots \geq s_n(t) > 0, \tag{12}$$

where j counts the number of positive eigenvalues of $\mathbb{P}(t)$. Define the function $f_{(n,j)}$ (encoding the signature of \mathbb{P}) by

$$2\gamma f_{(n,j)}(\mathbf{x}) := -\frac{p_0(t)}{|p_0(t)|} + \sum_{i=1}^j x_i^2 - \sum_{i=j+1}^n x_i^2. \tag{13}$$

Think of $2\gamma(f_{(n,j)})_+^\gamma$ as the density of a Borel measure. For any affine matrix \mathbb{A} , let $\mathbb{A}_\#$ be the push-forward operator for measures with respect to the map $\mathbf{x} \mapsto \mathbb{A}\mathbf{x}$. We use $\rho_0 = \mathbb{A}_\#(f_{(n,j)})_+^\gamma$. Echoing the definition of the push-forward, ρ_0 is the unique Borel measure satisfying

$$\int_{\mathbf{R}^n} \xi d\rho_0 = \int \xi(\mathbb{A}\mathbf{x})(f_{(n,j)}(\mathbf{x}))_+^\gamma d\mathbf{x} \tag{14}$$

for all continuous test functions $0 \leq \xi \in C_c(\mathbf{R}^n)$ of compact support. A straightforward application of the change of variable formula shows that ρ_0 has the density $(f_{(n,j)})_+^\gamma(\mathbb{A}^{-1}\mathbf{y})/|\det \mathbb{A}|$.

Setting $\mathbb{A}(t) = |\mathbb{S}(t)|^{1/2}$ as above, the evolving density $\rho(t) = \mathbb{A}(t)_\#(f_{(n,j)})_+^\gamma$ gives the pressure profile corresponding to (7), where we choose the constant in (10) to be $z = 1$ without loss of generality hereafter. More explicitly:

Lemma 2 (Density evolution). *For $0 \neq \gamma \neq -n/2$ let $\pi(\mathbf{x}, t) = \frac{1}{2}(\langle \mathbf{x}, \mathbb{P}(t)\mathbf{x} \rangle - p_0(t))$ solve (8)–(9) on $]t_\alpha, t_\omega[\subset \mathbf{R}$. Let j be the number of positive eigenvalues of $\mathbb{P}(t)$ and define $z > 0$ and $f_{(n,j)}(\mathbf{x})$ via (10) and (13). If $p_0(t) \det \mathbb{P}(t) \neq 0$ then $\rho(t) = (\pi/\gamma)_+^\gamma = z^{n+2\gamma} (z\mathbb{A}(t))_\#(f_{(n,j)})_+^\gamma$ where $\mathbb{A}(t)^{-2} = |\mathbb{P}(t)|/|\det \mathbb{P}(t)|^{1/(n+2\gamma)}$.*

Proof. The definition of z gives $|p_0(t)| = z^2 |\det \mathbb{P}(t)|^{1/(n+2\gamma)} = z^2 |\det \mathbb{A}(t)|^{-1/\gamma}$, so

$$\begin{aligned} \pi(\mathbf{y}, t) &= \frac{|p_0(t)|}{2} \left(\left\langle \mathbf{y}, \frac{\mathbb{P}(t)}{z^2 |\det \mathbb{P}(t)|^{1/(n+2\gamma)}} \mathbf{y} \right\rangle - \frac{p_0(t)}{|p_0(t)|} \right) \\ &= \gamma z^2 |\det \mathbb{A}(t)|^{-1/\gamma} f_{(n,j)}(\mathbb{A}(t)^{-1}\mathbf{y}/z). \end{aligned}$$

Integrating $\xi(\mathbf{y})$ against $\rho = (\pi/\gamma)_+^\gamma$ yields

$$\begin{aligned} \int_{\mathbf{R}^n} \xi(\mathbf{y})\rho(\mathbf{y}, t) d\mathbf{y} &= z^{2\gamma+n} \int_{\mathbf{R}^n} \xi(\mathbf{y})(f_{(n,j)}(\mathbb{A}(t)^{-1}\mathbf{y}/z))_+^\gamma \det(z\mathbb{A}(t))^{-1} d\mathbf{y} \\ &= z^{2\gamma+n} \int_{\mathbf{R}^n} \xi(z\mathbb{A}(t)\mathbf{x})(f_{(n,j)}(\mathbf{x}))_+^\gamma d\mathbf{x}. \end{aligned}$$

The change of variables $\mathbf{y} = z\mathbb{A}(t)\mathbf{x}$ is invertible since $p_0(t) \det \mathbb{P}(t) \neq 0$. \square

Theorem 3 (Integrability). *Fix $0 \neq \gamma \neq -n/2$, and let $\mathbb{P}(t)$ be a diagonal invertible matrix solving (8) on a maximal interval $]t_\alpha, t_\omega[\subset \mathbf{R}$. Then the eigenvalues s_i of $\mathbb{S}(t) = -\mathbb{P}^{-1}(t)|\det \mathbb{P}(t)|^{1/(n+2\gamma)}$ satisfy*

$$\frac{ds_i}{dt} = 2|\det \mathbb{S}(t)|^{-\frac{1}{2\gamma}} \quad \text{for } 1 \leq i \leq n. \tag{15}$$

Proof. Differentiating $\log |s_i(t)|$ yields

$$\begin{aligned} \frac{1}{s_i} \frac{ds_i}{dt} &= -\frac{d \log |p_i|}{dt} + \frac{1}{n+2\gamma} \sum_{k=1}^n \frac{d \log |p_k|}{dt} \\ &= -\frac{\text{tr } \mathbb{P}}{\gamma} - 2p_i + \frac{1}{n+2\gamma} \left(\frac{n}{\gamma} \text{tr } \mathbb{P} + 2 \text{tr } \mathbb{P} \right) \\ &= -2p_i \\ &= +\frac{2}{s_i} |\det \mathbb{P}|^{1/(n+2\gamma)} \\ &= \frac{2}{s_i} |\det \mathbb{S}|^{-1/(2\gamma)} \end{aligned}$$

where the second identity utilized (11). \square

Corollary 4 (Fate of quadratic pressures). *The orbits of the dynamical system (15) consist of all lines $S(t) = S(t_0) + (\tau(t) - \tau(t_0))\mathbb{I}$ which terminate at the coordinate hyperplanes $\det S(t) = 0$; see Fig. 1(a). The traceless part B of $S(t)$ is a constant of motion. Away from the coordinate hyperplanes, $d\tau/dt = 2|\det S|^{-1/(2\gamma)} > 0$ is smooth and positive, and $\tau(t) = \frac{1}{n} \operatorname{tr} S(t)$.*

Proof. The corollary follows immediately from (15). \square

It may seem surprising that the orbits of $S(t)$ are completely independent of the nonlinearity γ ; the precise value γ affects only the rate at which these orbits are parameterized. As a partial explanation, observe that the evolution (1) formally preserves the ρ -expectation of every harmonic function. Since the solutions $\rho = (\pi/\gamma)_+^\gamma$ that we study can be determined, within their family, by their second moments, i.e., by testing them with quadratic polynomials, and since the harmonic polynomials are codimension-1 in this space of test functions, it follows that all but one degree of freedom of our special solutions should be fixed, independent of γ .

Let us record the asymptotics of the solutions $S(t)$ as $t \nearrow t_\omega$, which are easily read off of (15). Since $-S(-t)$ solves the same equation, the asymptotics as $t \searrow t_\alpha$ follow immediately. In the porous medium case $\gamma > 0$, the vector field (15) diverges on each of the coordinate hyperplanes $s_i = 0$, so the dynamics accelerate towards and away from this singular set, blowing up at a finite time $t_\omega < \infty$. When $\gamma < 0$, the vector field $|\det S(t)|^{-1/2\gamma}(1, \dots, 1)$ vanishes continuously along these hyperplanes, so the dynamics decelerates towards (and away from) this singular set. Whether or not the singularity is reached in finite time is then determined by how quickly the vector field vanishes; i.e., by the modulus α of Hölder continuity along its integral curves (relative to 1). Where the orbit simultaneously intersects n_1 of the coordinate hyperplanes, this Hölder exponent is given by $\alpha = -n_1/2\gamma$. Borrowing the classification philosophy for solutions of Ricci flow equation, e.g. [29], this gives a convenient criterion for deciding when a quadratic pressure solution will be *eternal* (i.e. both *ancient* ($t_\alpha = -\infty$) and *immortal* ($t_\omega = \infty$)). To be eternal requires the multiplicities n_1 and n_n of the largest negative and smallest positive eigenvalues (12) of $S(t)$ to satisfy $n_1 + 2\gamma > 0$ and $n_n + 2\gamma > 0$ with $\gamma < 0$. Thus for $\gamma \in [-1/2, 0]$, each $P(t)$ of indefinite sign leads to an eternal solution, but for $\gamma \notin [-n/4, 0]$ no quadratic pressure is eternal save the constant $P(t) = 0$. Note however, that for $\gamma < -n/2$ the orbits $S(t)$ of the *finite-dimensional* dynamics can be extended continuously (but not smoothly) to the entire time domain $t \in \mathbf{R}$, simply by electing to spend (say) as little time on the singular set as possible. This choice specifies the extension uniquely, but it does not seem to have a meaningful interpretation at the level of the partial differential equation.

Corollary 5 (Asymptotics). *Fix $0 \neq \gamma \neq -n/2$, and let $P(t)$ be a diagonal invertible matrix solving (8) on a maximal interval $]t_\alpha, t_\omega[\subset \mathbf{R}^n$. Set $\tau(t) = \operatorname{tr} S(t)/n$ and denote the eigenvalues of $S(t) = -P^{-1}(t)|\det P(t)|^{1/(n+2\gamma)}$ by $s_i(t)$ and those of $B = S(t) - \tau(t)\mathbb{I}$ by b_i . Then there exist constants $c = |\frac{\gamma}{n_1+2\gamma}| \prod_{i>n_1}^n |s_i(t_\omega)|^{1/(2\gamma)}$ specified below and t_1 such that*

(a) **Spreading:** *if $S(t) > 0$ then $\tau(t) \rightarrow \infty$ as $t \rightarrow t_\omega \leq +\infty$ and*

$$\tau(t) = \begin{cases} [\frac{n+2\gamma}{\gamma}(t - t_1)]^{2\gamma/(n+2\gamma)} + O_*((t - t_1)^{-2\gamma/(n+2\gamma)}) & \text{or} \\ [\frac{n+2\gamma}{|\gamma|}(t_\omega - t)]^{2\gamma/(n+2\gamma)} + O((t_\omega - t)^{-2\gamma/(n+2\gamma)}) \end{cases} \tag{16}$$

where the first formula holds for $(n + 2\gamma)/\gamma > 0$, hence $t_\omega = \infty$, and the second formula holds for $(n + 2\gamma)/\gamma < 0$, hence t_ω finite. The symbol O_* is the same as O , except when $\gamma = n/2$: In that case an extra factor $\log t$ enters in the O term.

(b) **Focusing:** *Otherwise, suppose $s_1(t)$ has the smallest absolute value among negative eigenvalues (12) of $S(t)$, and let n_1 denote its multiplicity. Then $S(t)$ converges to a rank $(n - n_1)$ finite limit matrix $S(t_\omega) = B + \tau_\omega\mathbb{I}$ as $t \rightarrow t_\omega$, with $s_1(t_\omega) = 0$ and $\tau_\omega := \tau(t_\omega)$, while*

$$\tau_\omega - \tau(t) = \begin{cases} [(t_\omega - t)/c]^{2\gamma/(n_1+2\gamma)} + O((t_\omega - t)^{4\gamma/(n_1+2\gamma)}) & \text{or} \\ [(t - t_1)/c]^{2\gamma/(n_1+2\gamma)} + O_*((t - t_1)^{4\gamma/(n_1+2\gamma)}) \end{cases} \tag{17}$$

where the first formula holds for $(n_1 + 2\gamma)/\gamma > 0$, hence $t_\omega < \infty$. The second formula holds for $(n_1 + 2\gamma)/\gamma < 0$, hence $t_\omega = \infty$. The symbol O_* denotes an extra logarithmic factor in the case $\gamma = -n_1/4$.

Proof. Recall from Corollary 4 that

$$\frac{d\tau}{dt} = 2 \prod_{i=1}^n |b_i + \tau(t)|^{-1/2\gamma}, \tag{18}$$

If $S > 0$, then clearly $\tau(t) \rightarrow \infty$ as $t \rightarrow t_\omega$ by the previous corollary and the continuation theorem. The absolute values in (18) are redundant in this case; for the moment assume $\gamma > 0$ or $\gamma < -n/2$, hence $1 + \frac{n}{2\gamma} > 0$. Separation of variables and integration (using $\text{tr } B = 0$) yields

$$\frac{\gamma}{n + 2\gamma} \tau^{1 + \frac{n}{2\gamma}} (1 + O_*(\tau^{-2})) = t - t_1 \tag{19}$$

with a yet undetermined constant of integration t_1 . The $*$ in $O_*(\tau^{-2})$ is only relevant for $\gamma = n/2$ and in this case refers to an extra factor $\ln \tau$ in the $O(\cdot)$ term. From (19) and $(n + 2\gamma)/\gamma > 0$, one gets $t_\omega = \infty$, and also

$$\tau (1 + O_*(\tau^{-2})) = \left[\frac{n + 2\gamma}{\gamma} (t - t_1) \right]^{2\gamma/(n+2\gamma)}.$$

Hence the claim (16). Essentially the same calculation can also be used for $-\frac{n}{2} < \gamma < 0$. However in this case, $t - t_1 \rightarrow 0$ as $\tau \rightarrow \infty$; so we have t_ω finite, equal to t_1 . We then get the corresponding second formula in (16).

Now in the case where $s_1 < 0$, the hyperplane $s_1 = 0$ (actually, the space $s_1 = \dots = s_{n_1} = 0$) is reached for a finite value of $t_\omega := \tau(t_\omega)$. The remaining factors $\prod_{i=n_1+1}^n \dots$ are analytic near t_ω . Separation of variables in (18) gives, for $1 + n_1/2\gamma > 0$,

$$c(\tau_\omega - \tau)^{1+n_1/2\gamma} (1 + O(\tau_\omega - \tau)) = t_\omega - t$$

with $t_\omega < \infty$ and $c = \frac{\gamma}{n_1+2\gamma} \prod_{i=n_1+1}^n |b_i + \tau_\omega|^{1/2\gamma}$. In contrast, for $1 + n_1/2\gamma < 0$ we get

$$c(\tau_\omega - \tau)^{1+n_1/2\gamma} (1 + O_*(\tau_\omega - \tau)) = t - t_1$$








with some constant of integration t_1 , with $c = \frac{|\gamma|}{n_1+2\gamma} \prod_{i=n_1+1}^n |b_i + \tau_\omega|^{1/2\gamma}$ and $t_\omega = \infty$; the O_* refers to a logarithmic term instead of $(\tau_\omega - \tau)^0$, i.e., if $\gamma = -n_1/4$. The claim follows immediately. \square

Note that in the immortal case (16a), for $\gamma > 0$ a time shift may or may not be absorbed into the error term, depending on the size of $\gamma > 0$ relative to $n/2$. This suggests phase transitions in the long-time porous medium asymptotics, similar to those observed [24,32] in the fast diffusion regime $\gamma < 0$. Higher asymptotics would follow from a closer study of the ordinary differential equation (18) with $\sum b_i = \text{tr } B = 0$. Let us rather examine the solutions described. Table 1 gives an overview.

Since eigenvalues do not change sign, the corollary asserts that a quadratic pressure $\pi(x, t)$ represents a spreading solution if and only if its quadratic part $P = -S^{-1} |\det S|^{-1/2\gamma}$ is negative definite. In this case the density $\rho = (\pi/\gamma)_+^\gamma$ is compactly supported if $\gamma > 0$, and otherwise has tails that decay like $|x|^{2\gamma}$. Thus $\rho \in L^1(\mathbf{R}^n)$ if and only if $\gamma \notin [-n/2, 0]$. We call this range the *mass-preserving* regime; it coincides with the range of γ for which spreading solutions are immortal ($t_\omega = +\infty$), and all other solutions focus in finite time. As the solution $\rho(t) = A(t) \# (f_{(n,0)})_+^\gamma$ spreads out, its isobars (= level sets of the pressure) are ellipsoids which become rounder and rounder as $t \rightarrow t_\omega$, since $A(t) = |\tau(t)I + B|^{1/2} \sim \tau(t)^{1/2} (I + \frac{1}{2\tau(t)} B)$ as τ grows. This is in accordance with the attractivity of the Barenblatt solution proved in the mass-preserving regime by Friedman and Kamin [36]; see also Vázquez [64]. Comparing different eigenvalues $a_i(t)$ of $A(t)$, for $\gamma > 0$ the departure from sphericity $2(a_i - a_k)/\tau(t)^{1/2} \sim (b_i - b_k)/|\frac{n+2\gamma}{\gamma} t|^{2\gamma/(n+2\gamma)}$ disappears more slowly than the discrepancy $(a_i(t + t_0) - a_i(t))/\tau(t)^{1/2} \sim |\frac{\gamma}{n+2\gamma} \frac{t_0}{t}$ between a solution and its time delay. These precise decay exponents are also predicted by the linearization we carried out in the mass-preserving fast diffusion $\gamma < -n/2$ range [33]. Note the constants $b_i \neq b_k$ are distinct unless the eigenvalues $a_i(t) = a_k(t)$ coincide for all $t \in]t_\alpha, t_\omega[$.

Turning now to the focusing case, we observe that when $S(t)$ is not positive definite, its orbit terminates on the i -th coordinate hyperplane, where $s_i(t)$ is the smallest in absolute value among the negative eigenvalues (12) of $S(t)$; $i = 1$ in our numeration (12). This eigenvalue decays to zero at a rate $|t_\omega - t|^{2\gamma/(n_1+2\gamma)}$ determined by its multiplicity n_1 ; (we tacitly replace $t_\omega - t$ by $t - t_1$ in this and similar formulae if $-n_1/2 < \gamma < 0$, since the solution in that case

Table 1
Overview of solutions

π Figure	$\gamma > 0$ Porous medium	$\gamma < -n/2$ Conservative fast diffusion
$P < 0, -p_0 > 0$ 	 spreading and getting rounder; increasing elliptical support. Affine image of Barenblatt	$P \sim \frac{-\gamma/(2\gamma+n)}{t} \mathbb{1} + O(t^{-1-2\gamma/(2\gamma+n)});$ $p_0 \sim \frac{\text{const}}{t^{n/(2\gamma+n)}};$ elliptical singular support (infinite density), spreads out and gets rounder as $t \nearrow \infty$.
$P < 0, -p_0 < 0$ 	N/A	Affine image of a Barenblatt solution
P indef., $-p_0 > 0$	 density supported in shaded area	support of P spreading as $t \nearrow t_\omega$: $\theta \sim (t_\omega - t)^{\gamma/(2\gamma+n_1)}$; ‘waistline’ \rightarrow const. As $t \searrow t_\alpha$, support shrinks to horizontal axis. density infinity in shaded area, finite in unshaded area
P indef., $-p_0 < 0$	 density supported in shaded area	support of P spreading as $t \nearrow t_\omega$: $\theta \sim (t_\omega - t)^{\gamma/(2\gamma+n_1)}$; ‘gap’ $\rightarrow 0$. As $t \searrow t_\alpha$, support shrinks to two rays with finite gap density infinity in shaded area, finite in unshaded area
$P > 0, -p_0 > 0$ 	unbounded, locally finite solution supported in whole space	N/A
$P > 0, -p_0 < 0$ 	unbounded, locally finite solution supported in the complement of an elliptical hole. As $t \nearrow t_\omega$, this hole shrinks to a slit or elliptical disc depending on dimension.	singular support (infinite density) outside ellipse/ellipsoid; finite density inside shrinks to a slit or elliptical disc.

is immortal, meaning the focusing takes infinitely long). Since the other distinct eigenvalues of $S(t)$ remain at fixed distances to s_1 , they converge to nonzero values at the same rate. Thus n_1 spatial dimensions $a_1 = \dots = a_{n_1} = |s_i|^{1/2}$ of the solution $\rho(t) = A(t)_{\#} (f_{(n,j)})_+^{\gamma}$ collapse as $t \rightarrow t_\omega$, the others tending to as many distinct nonzero values as their multiplicities permit. In the filtration-absorption model of Barenblatt, Bertsch, Chertock and Prostokishin [10] ($j = n$), this suggests that the last drop of fluid drains not as a round drop, but rather as a flattening ellipsoid with generically different axes, the shortest disappearing at a rate $x_1 \sim (t_\omega - t)^{\gamma/(2\gamma+n_1)}$, and the others tending to distinct nonzero limits as $t \rightarrow t_\omega$. Since the physical relevance of quadratic focusing solutions may be debatable, let us observe via time-reversal that the future asymptotics $t \rightarrow t_\omega$ for the $S(t) < 0$ solutions above (which are ancient in the mass-preserving regime), also describe the past asymptotics $t \rightarrow \bar{t}_\alpha = -t_\omega$ of spreading solutions at their moment of origin. Thus our spreading solutions with $S(t) > 0$ originate not at a point source (unless they are perfectly round), but rather as a characteristic density on a finite size rod, disc, or flattened ellipsoid. The next lemma shows this characteristic density is the sheared image of a Barenblatt profile in dimension $k = n - n_n$, but associated to a different nonlinearity $m' \neq m$ so that its moment index $n + 2\gamma =: \mu = \mu' := k + 2/(m' - 1)$ is the same. The shape of this profile can

also be realized by projecting the mass of $\rho_0 = (f_{(n,0)})_+^\gamma$ from \mathbf{R}^n to \mathbf{R}^k using a rank k matrix $A(t_\alpha)$. Corollary 4 makes it clear that each projection $A(t_\alpha)_\# \rho_0$ with $\det A(t_\alpha) = 0$ is actually realized as the $t \searrow t_\alpha$ limit of a (unique) orbit composed of densities diffusing with quadratic pressure, and that affine images of the Barenblatt solution form an invariant subspace for the porous medium / fast diffusion dynamics (1). We remind the reader that j continues to denote the number of positive eigenvalues in P . However, for the backwards-in-time limit $t \rightarrow t_\alpha$, it is not $p_1 > 0$ but $p_n < 0$ that blows up, and we will assume its multiplicity to be $n - k$, so that $p_k \neq p_{k+1} = \dots = p_n < 0$.

Lemma 6 (Singular initial profiles preserve moment index). *Suppose $A(t) > 0$ is a positive definite diagonal $n \times n$ matrix for each $t \in]t_\alpha, t_\omega[$, and converges to a rank k (with $k < n$) limit matrix $A(t_\alpha) = \text{diag}(a_1, \dots, a_k, 0, \dots, 0)$ as $t \rightarrow t_\alpha$. Given $j \leq k$ and $\gamma \notin [(k - n)/2, 0]$ define $\rho(t) = A(t)_\#(f_{(n,j)})_+^\gamma$ via (13)–(14). If $\gamma < 0$, assume $j = 0$ and $p_0(t) > 0$ to ensure $(f_{(n,j)})_+^\gamma$ is a Radon measure. Up to a normalization constant $z' > 0$,*

$$\lim_{t \rightarrow t_\alpha} \int_{\mathbf{R}^n} \xi(\mathbf{x}) \rho(\mathbf{x}, t) \, d\mathbf{x} = \int_{\mathbf{R}^n} \xi(A(t_\alpha)\mathbf{x}) (f_{(n,j)}(\mathbf{x}))_+^\gamma \, d\mathbf{x} = z' \int_{\mathbf{R}^n} \xi \, d\sigma, \tag{20}$$

for all continuous test functions $\xi \in C_c(\mathbf{R}^n)$ of compact support. Here $\sigma = A(t_\alpha)_\#(f_{(n,j)})_+^\gamma$ is a measure supported on the subspace $x_{k+1} = \dots = x_n = 0$; its density with respect to Hausdorff k -dimensional measure \mathcal{H}^k is given by

$$\frac{d\sigma}{d\mathcal{H}^k}(x_1, \dots, x_k, 0, \dots, 0) = \left(f_{(k,j)} \left(\frac{x_1}{a_1}, \dots, \frac{x_k}{a_k} \right) \right)_+^{\gamma + (n-k)/2}. \tag{21}$$

Proof. The definition $\rho(t) := A(t)_\#(f_{(n,j)})_+^\gamma$ gives

$$\int_{\mathbf{R}^n} \xi(\mathbf{y}) \rho(\mathbf{y}, t) \, d\mathbf{y} = \int_{\mathbf{R}^n} \xi(A(t)\mathbf{x}) (f_{(n,j)}(\mathbf{x}))_+^\gamma \, d\mathbf{x}.$$

The $t \rightarrow t_\alpha$ limit (20) will be derived using the dominated convergence theorem. For $\gamma > 0$ a constant function on a compact set supporting all integrands is a majorant: $(f_{(n,j)})_+^\gamma$ is continuous, hence $L^1_{\text{loc}}(\mathbf{R}^n)$, and the support of $\xi(A(t)\mathbf{x}) \in C_c(\mathbf{R}^n)$ is eventually contained in a cylinder $K \times \mathbf{R}^{n-k}$ with compact cross-section $K \subset \mathbf{R}^k$. The set $\{\mathbf{x} \mid f_{(n,j)}(\mathbf{x}) > 0\}$ is asymptotic to the cone $\sum_{i=j+1}^n x_i^2 \leq \sum_{i=1}^j x_i^2$, which has compact intersection with $K \times \mathbf{R}^{n-k}$ because $j \leq k$. For $\gamma < (k - n)/2 < 0$, our hypotheses $j = 0$ and $p_0(t) > 0$ ensure that $(f_{(n,j)}(\mathbf{x}))_+^\gamma$ is globally positive and integrable on $K \times \mathbf{R}^{n-k}$, so $\|\xi\|_{L^\infty(\mathbf{R}^n)}$ times this function is a majorant.

Finally, $\xi(A(t_\alpha)\mathbf{x}) = \xi(a_1 x_1, \dots, a_k x_k, 0, \dots, 0)$ implies σ is supported on the subspace $x_{k+1} = \dots = x_n = 0$. We can obtain its Lebesgue density there by integrating out x_i inductively for $n \geq i \geq k + 1$: Since $f_{(i,j)} := f_{(i,j)}(x_1, \dots, x_i) = f_{(i-1,j)} - x_i^2/\gamma$, we know for $\gamma > 0$ that $f_{(i-1,j)} > 0$ holds wherever $f_{(i,j)} > 0$, while for $\gamma < 0$ then $f_{(i,j)} > 0$ for each $i \leq n$ by our choices $j = 0$ and $p_0(t) > 0$. We integrate

$$\begin{aligned} \int_{-\infty}^{\infty} (f_{(i,j)})_+^\beta \, dx_i &= (f_{(i-1,j)})_+^{\beta+1/2} \sqrt{|\gamma|} \int_{-\infty}^{\infty} \left(1 - \frac{x_i^2}{\gamma f_{(i-1,j)}} \right)_+^\beta \frac{dx_i}{\sqrt{|\gamma| f_{(i-1,j)}}} \\ &= (f_{(i-1,j)})_+^{\beta+1/2} \sqrt{|\gamma|} \int_{-\infty}^{\infty} (1 - |\gamma| r^2/\gamma)_+^\beta \, dr \end{aligned}$$

for $i = n, n - 1, \dots, k + 1$, and $\beta = \gamma, \gamma + \frac{1}{2}, \dots, \gamma + \frac{n-k-1}{2}$. The dr integral is finite (if $\gamma > 0$, then $\beta > 0$; if $\gamma < (k - n)/2$, then $\beta < -\frac{1}{2}$) and could be evaluated in terms of the Gamma function, but we simply absorb its value into the normalization constant z' . Absorbing the Jacobian $\prod_{i=1}^k a_i$ into z' also, the invertible change of variables $A(t_\alpha): \mathbf{R}^k \rightarrow \mathbf{R}^k$ yields (21). \square

This lemma appears natural, and is divorced from the dynamics. It is therefore interesting to see the hypothesis $\gamma \notin [(k - n)/2, 0]$ required to guarantee local finiteness of the limiting measure σ is the same which permits nonlinear diffusion from this measure to occur in a finite time $t_\alpha > -\infty$. The forward focusing limit $t \rightarrow t_\omega < \infty$ would be more

awkward to address since, as in the case $\gamma < 0$ and $\mathbb{P}(t)$ of indefinite sign, infinities make (20) diverge for all non-vanishing $0 \leq \xi \in C(\mathbf{R}^n)$.

Finally, the solutions with $\mathbb{P}(t)$ of indefinite sign have more complicated focusing geometries (and a finite lifetime $]t_\alpha, t_\omega[$ in the mass-preserving regime). Some of these are included in the lemma above, and illustrated in Table 1. Assuming the eigenvalues $p_1(t) > \dots > p_n(t)$ are distinct and nonzero, the isobars will be hyperbolas/hyperboloids (in two and three-dimensions) asymptotic to a double-cone emitted by the plane $x_n = 0$ at $t = t_\alpha$, and collapsing anisotropically onto the plane $x_1 = 0$ as $t \rightarrow t_\omega$. In two-dimensions, the initial region of positive pressure can be (i) two opposite rays along $x_2 = 0$ or (ii) the entire line $x_2 = 0$. At the focusing time $t = t_\omega$ it will finally occupy all but (i) the line $x_1 = 0$ or (ii) two opposite rays along it. In three-dimensions the asymptotic cone encircles either (a) the x_3 -axis or (b) the x_1 -axis, depending on the sign of p_2 : the positive pressure region then begins as (a1) the entire plane, (a2) the complement of an ellipse, or the region (b1) between or (b2) outside the two sheets of a hyperbola in the plane $x_3 = 0$; it evolves to fill (and diverge on) all but a region (b2), (b1), (a2) or (a1) in the plane $x_1 = 0$ by time $t = t_\omega$. When $n = 3$, various degenerate limits are possible if two eigenvalues are equal. Higher dimensions allow for more exotic geometries and singularities.

4. Fourth order diffusions

4.1. A naive logarithmic diffusion in several dimensions

In this section, we observe that the same strategy can be used to expose and to analyze affinely self-similar solutions for certain fourth-order equations of recent interest governing diffusion of a nonnegative scalar $\rho(\mathbf{x}, t) \geq 0$ on $\mathbf{R}^n \times]0, \infty[$. The first of these is the naive generalization

$$\frac{\partial \rho}{\partial t} = -\Delta(\rho \Delta \log \rho) \tag{22}$$

to higher dimensions of an evolution proposed by Derrida, Lebowitz, Speer and Spohn [34] as a model governing the size x of interfacial fluctuations in a statistical mechanical spin system, as a function of the distance t to a point where the interface is pinned by the boundary conditions. The exponential change of variables $\pi(\mathbf{x}, t) = \log \rho(\mathbf{x}, t)$ converts this model to a form [44]

$$-\frac{\partial \pi}{\partial t} = \Delta^2 \pi + 2 \nabla \pi \cdot \nabla \Delta \pi + (\Delta \pi)^2 + \Delta \pi |\nabla \pi|^2$$

which makes clear that quadratic solutions $2\pi(\mathbf{x}, t) = \langle \mathbf{x} \mathbb{P}(t) \mathbf{x} \rangle - p_0(t)$ exist for any symmetric initial matrix $\mathbb{P}(t_0)$ and constant $p_0(t_0)$. The resulting differential equations

$$\frac{d\mathbb{P}(t)}{dt} = -2\mathbb{P}(t)^2 \operatorname{tr} \mathbb{P}(t), \tag{23}$$

$$\frac{dp_0(t)}{dt} = 2(\operatorname{tr} \mathbb{P}(t))^2 \tag{24}$$

are smooth, fix the harmonic polynomials $\operatorname{tr} \mathbb{P} = 0$, and can be integrated on some maximal time interval $]t_\alpha, t_\omega[\subset \mathbf{R}$ to give invariance of $e^{p_0(t)} \det \mathbb{P}(t) = e^{p_0(t_0)} \det \mathbb{P}(t_0)$ and orbits $\mathbb{P}(t)^{-1} = \mathbb{P}(t_0)^{-1} - (\tau(t) - \tau(t_0))\mathbb{I}$ parallel to the principal diagonal, as in Lemma 1 and Corollary 4; see Fig. 1(b), which we now justify. The parametrization $\tau(t)$ is governed by

$$\frac{d\tau}{dt} = 2 \sum_{i=1}^n \frac{1}{b_i + \tau(t)},$$

where b_i denote the eigenvalues of the traceless part \mathbb{B} of $\mathbb{S}(t) := -\mathbb{P}(t)^{-1}$. The dynamics governing $\mathbb{S}(t) = \mathbb{B} + \tau(t)\mathbb{I}$ is smooth away from the set $\det \mathbb{S} = 0$. It is clear from (23) that all orbits are immortal: $\operatorname{tr} \mathbb{P}$ cannot change sign because $\{\operatorname{tr} \mathbb{P} = 0\}$ is invariant; so if $\operatorname{tr} \mathbb{P} > 0$, all p_i decrease, but none can go to $-\infty$ because the trace is bounded below; an analogous argument applies if $\operatorname{tr} \mathbb{P} < 0$. The bound then gives global existence. All orbits apart from the equilibria $\operatorname{tr} \mathbb{P} = 0$ fail to be ancient: If $\operatorname{tr} \mathbb{P} > 0$, then this trace increases in backwards time, and also some $p_i > 0$. So $dp_i/d(-t) \geq cp_i^2$ implies finite time blow up. Similarly for $\operatorname{tr} \mathbb{P} < 0$. So the orbits originate at $\det \mathbb{S} = 0$ as




Fig. 1. Comparison of quadratic dynamics for three diffusion equations in 2D: In each case, the axes denote the eigenvalues of S . Gray zones denote singularities, as do white dots, to stress that the orbits run into singularities. Black dots denote equilibria.

$t \searrow t_\alpha$, departing from the intersection of n_k coordinate hyperplanes in a finite time $\tau(t) \sim \tau(t_\alpha) + 2\sqrt{n_k(t - t_\alpha)}$. If $S(t_0) > 0$ or $S(t_0) < 0$ has a definite sign, then the orbit $S(t) = B + \tau(t)\mathbb{I}$ grows like $\tau(t) \sim 2\sqrt{nt}$ as $t \rightarrow \infty$; otherwise $S(t) = S_\omega + O(e^{-\kappa t})$ converges exponentially to the surface $\text{tr } S^{-1} = 0$ as $t \rightarrow \infty$, with exponent $\kappa = 2 \text{tr}(S_\omega^{-2})$. The corresponding densities $\rho = e^\pi > 0$ are classical solutions to Eq. (22), given as the image $\rho(t) = A(t)_\# \exp(\gamma f_{(n,j)})$ under $A(t) = |S(t)|^{1/2}$ of a Gaussian (13), j counting the number of negative eigenvalues of $S(t)$. When $j = 0$, this evolves from a Gaussian measure concentrated on the subspace $A(\tau(t_\alpha))(\mathbf{R}^n)$, toward the self-similar solution $(4t)^{1/4}_\# \exp(\gamma f_{(n,0)})$ found by Derrida et al. [34] in the one-dimensional case. Since the evolution (22) preserves center of mass $\mathbf{x}_0 \in \mathbf{R}^n$, from $A(t) \sim \tau(t)^{1/2}(\mathbb{I} + \frac{1}{2\tau(t)}B)$ as $t \rightarrow \infty$ we deduce the relative rates of convergence of translations in space $\mathbf{x}_0/\tau(t)^{1/2} \sim \mathbf{x}_0/(4t)^{1/4}$, and in time $(a_i(t + t_0) - a_i(t))/\tau(t)^{1/2} \sim t_0/(4t)$, with the deviation from sphericity $2(a_i - a_k)/\tau(t)^{1/2} \sim (b_i - b_k)/(4t)^{1/2}$. For $j = n$ the solution instead grows exponentially as $\mathbf{x} \rightarrow \infty$ and $t \rightarrow \infty$, while for $1 < j < n$ it is a logarithmically quadratic function which becomes log-harmonic exponentially fast in the long time limit.

4.2. A diffusion that is the 2-Wasserstein gradient flow of the Fisher information

Another diffusion coinciding with (22) in one-dimension is the evolution

$$\frac{\partial \rho}{\partial t} = - \sum_{i,j=1}^n \partial_{x_i x_j}^2 (\rho \partial_{x_i x_j}^2 \log \rho) \tag{25}$$

explored by Jüngel and Pinnau as a model for transport phenomena in quantum systems [44] (damped by a magnetic field, for example). The same exponential change of variables converts this equation to the form

$$-\frac{\partial \pi}{\partial t} = \Delta^2 \pi + 2\langle \nabla \pi, \nabla \Delta \pi \rangle + \text{tr}(D^2 \pi D^2 \pi) + \langle \nabla \pi, D^2 \pi \nabla \pi \rangle \tag{26}$$

where $D^2 \pi$ denotes the Hessian matrix of π . The quadratic ansatz $2\pi(\mathbf{x}, t) = \langle \mathbf{x}, P(t)\mathbf{x} \rangle - p_0(t)$ proves self-consistent, reducing the dynamics to a system of ordinary differential equations $dP/dt = -2P(t)^3$ and $dp_0/dt = 2 \text{tr}(P^2)$ which can be integrated directly to yield $P(t)^{-2} = P(t_0)^{-2} + 4(t - t_0)\mathbb{I} = 4B + 4t\mathbb{I}$ and $e^{p_0(t)} \det P(t) = e^{p_0(t_0)} \det P(t_0)$ on a maximal time interval $[t_\alpha, \infty[$ with $t_\alpha > -\infty$ (unless $P \equiv 0$). The ordering among the eigenvalues $a_i(t)$ of $A(t) = |P(t)|^{-1/2}$ is preserved; all are increasing functions of time, whose asymptotics can be deduced from $A(t) = (4t)^{1/4}(\mathbb{I} + B/t)^{1/4}$. In this case, focusing is not consistent with our quadratic ansatz; all solutions spread forwards in time (and focus backwards in time). The corresponding densities $\rho = A(t)_\# \exp(\gamma f_{(n,j)})$ are classical solutions to the evolution (25) which represent Gaussians (if $P(t) < 0$ so $j = 0$); they begin evolving from a lower dimensional Gaussian at $t = t_\alpha$, and spread at rate $a_i(t) \sim (4t)^{1/4}$ so that $a_i/a_{i+1} \rightarrow 1$ as $t \rightarrow \infty$. The finer asymptotics $(a_i(t + t_0) - a_i(t))/(4t)^{1/4} \sim t_0/(4t)$ and $(a_i - a_k)/(4t)^{1/4} \sim (b_i - b_k)/(4t)$ parallel the heat equation, though translations converge at the slower rate $1/|4t|^{1/4}$. Fig. 1 gives a quick comparison of the dynamics of quadratic solutions for the different types of diffusion considered here.

Recently, the global well-posedness of (25) was announced by Gianazza, Savaré and Toscani, who exploited the fact that (25) is the gradient flow of the Fisher information

$$I(u) := \frac{1}{2} \int \frac{|\nabla u|^2}{u} dx = \frac{1}{2} \int |\nabla \ln u|^2 u dx = 2 \int |\nabla \sqrt{u}|^2 dx,$$

with respect to the 2-Wasserstein distance. This is explained in Section 1.5 of [41] and [1, Chapter XI]. For a general background, the reader may be reminded that the 2-Wasserstein distance between two measures of the same total mass is the square root of the minimal cost of transport from one measure to the other, provided the cost functional is the square of the distance.

4.3. A fourth order analog for the porous medium family

The Wasserstein gradient flow structure of (25) makes it clear that this equation represents the case $m = 1$ in a family of fourth-order diffusion equations analogous to the porous medium family discussed above. In conservation law form, these equations

$$\frac{\partial \rho}{\partial t} = -\frac{1}{m-1/2} \nabla \cdot [\rho \nabla (\rho^{m-3/2} \Delta (\rho^{m-1/2}))], \tag{27}$$

govern the evolution of a nonnegative scalar $\rho(x, t) \geq 0$ on $\mathbf{R}^n \times]0, \infty[$, and depend on a parameter $m \in \mathbf{R}$. Formally, this dynamics arises as the gradient flow of the *generalized Fisher information*

$$I_m(\rho) := \frac{1}{2} \frac{1}{(m-1/2)^2} \int_{\mathbf{R}^n} |\nabla (\rho^{m-1/2}(x, t))|^2 dx, \tag{28}$$

with respect to 2-Wasserstein distance, in the same way that the porous medium flow (1) arises as the 2-Wasserstein gradient flow of the *entropy*

$$E_m(\rho) := \frac{1}{m(m-1)} \int_{\mathbf{R}^n} (\rho(x, t)^m - \rho(x, t)) dx \tag{29}$$

in the work of Otto [54]; see Ambrosio, Gigli and Savaré [1] and Carrillo, McCann and Villani [26] for related developments.

The case $m = \frac{3}{2}$ of (27), $\rho_t = -\nabla \cdot (\rho \nabla (\Delta \rho))$, actually coincides with the case $n = 1$ of the thin film equation family $\rho_t = -\nabla \cdot (\rho^n \nabla (\Delta \rho))$. This family has been studied by many authors, especially in the 1-dimensional case. For instance, see Myers [51] and Oron, Davis and Bankoff [52] for review articles about the equation, Bernis and Friedman [11] for existence of weak solutions, Otto [53], Giacomelli and Knüpfer [38] and Giacomelli and Otto [39,40] for some recent work on the thin-film equation, and also Bertozzi, Laugesen, Pugh and Slepčev in various combinations [13,48,58] for recent work involving the competition between fourth-order stabilizing and second-order destabilizing effects, due—for example—to a competition of thin film surface tension with gravity. Bertsch et al. [14] study the thin film equation in higher space dimensions. A representative review of such work would be beyond the scope of this paper. Betelú and King [18] found quadratic solutions to the thin-film equation $u_t = -\text{div}(u \nabla (\Delta u))$. Inasmuch as this special case can be embedded into the family we are studying here, our solutions generalize theirs (but not within the thin-film hierarchy). The unifying theme is that the nonlinearities are quadratic, as was elaborated on the basis of different examples already by Galaktionov [37].

Remarkably, the dynamics (27) can also be formulated in terms of a pressure equation (30) which possesses quadratic solutions. The limit $m \rightarrow 1$ motivates nomenclature, while the fact that $dE_m(\rho)/dt = -2I_m(\rho) < 0$ under the porous medium dynamics (1) explains some of the parallels between these two families. To see directly the equivalence of (25) with the $m = 1$ instance of (27), i.e., to show

$$2\partial_i (\rho \partial_i (\rho^{-1/2} \partial_j \partial_j \rho^{1/2})) = \partial_i \partial_j (\rho \partial_i \partial_j \log \rho)$$

evaluate the rightmost derivative on either side; after canceling the common ‘factor’ ∂_i , it suffices to show the identity $\rho \partial_i \rho^{-1/2} \partial_j \rho^{-1/2} \rho_j = \partial_j \rho \partial_i \rho^{-1} \rho_j$. The left-side equals $\rho \partial_i \partial_j \rho^{-1} \rho_j - \rho \partial_i (-\frac{1}{2} \rho^{-2} \rho_j \rho_j)$. The right-hand side equals $\rho \partial_j \partial_i \rho^{-1} \rho_j + \rho_j \partial_i \rho^{-1} \rho_j$. Equality can now be checked easily. For $m \leq 1$, some quadratic solutions discussed below

will be classical; but for $m > 1$, the discussion is purely formal, since we do not attempt to identify a notion of weak solution.

As in the porous medium case, applying the identity

$$\frac{\pi^{1-\gamma/2}}{1 + \gamma/2} \Delta(\pi^{1+\gamma/2}) = \pi \Delta \pi + \gamma |\nabla \pi|^2 / 2$$

to $\gamma = 1/(m - 1)$ and $\pi = \gamma \rho^{1/\gamma}$ reexpresses the evolution (27) in the simpler form

$$-\frac{\partial \pi}{\partial t} = \frac{\pi}{\gamma} \Delta \left(\frac{\pi}{\gamma} \Delta \pi + \frac{1}{2} |\nabla \pi|^2 \right) + \nabla \pi \cdot \nabla \left(\frac{\pi}{\gamma} \Delta \pi + \frac{1}{2} |\nabla \pi|^2 \right). \tag{30}$$

Unlike the pressure version (3) of the porous medium equation, this equation is invariant under the negation $\pi(\mathbf{x}, t) \leftrightarrow -\pi(\mathbf{x}, t)$. It degenerates only where $\pi = 0$ or $\pi = \pm\infty$. The quadratic ansatz $\pi(\mathbf{x}, t) = \frac{1}{2}(\langle \mathbf{x}, \mathbb{P}(t)\mathbf{x} \rangle - p_0(t))$ proves self-consistent, and yields a system of ordinary differential equations

$$-\frac{d\mathbb{P}(t)}{dt} = 2\mathbb{P}^3 + 2\mathbb{P}^2 \frac{\text{tr } \mathbb{P}}{\gamma} + \mathbb{P} \frac{\text{tr}(\mathbb{P}^2)}{\gamma} + \mathbb{P} \frac{(\text{tr } \mathbb{P})^2}{\gamma^2} \tag{31}$$

$$-\frac{1}{p_0} \frac{dp_0}{dt} = \frac{\text{tr}(\mathbb{P}^2)}{\gamma} + \frac{(\text{tr } \mathbb{P})^2}{\gamma^2} \tag{32}$$

which clearly admit local solutions. Any initially diagonal matrix $\mathbb{P}(0)$ remains diagonal under the evolution, and any eigenvalue $p_i(0) = 0$ which vanishes initially remains zero at all subsequent times. We can thus discard any zero eigenvalues from the system (31), instead studying the analogous evolution for a smaller, nondegenerate matrix. We therefore assume $\det \mathbb{P}(t) \neq 0$ hereafter without loss of generality.

Since distinct eigenvalues remain distinct under the flow (31), and since zero eigenvalues can be ignored, we can define the effective dimension \check{n} to be the number of distinct nonzero eigenvalues of \mathbb{P} .

As in (10), it is straightforward to deduce the invariance of

$$z^2 = |p_0(t)| / |\det \mathbb{P}(t)|^{1/(n+2\gamma)} = \text{const} \tag{33}$$

from (31)–(32). Reexpressed in terms of

$$S(t) := -\mathbb{P}^{-1} |\det \mathbb{P}(t)|^{1/(n+2\gamma)}, \quad |\det S| = |\det \mathbb{P}|^{-2\gamma/(n+2\gamma)}, \tag{34}$$

a calculation similar to the proof of Theorem 3 reduces the dynamics (31) to the form

$$\frac{dS}{dt} = 2 |\det S(t)|^{-1/\gamma} \left(S^{-1} + \frac{\text{tr } S^{-1}}{\gamma} \mathbb{I} \right). \tag{35}$$

Notice that Lemma 2 now asserts the density $\rho = z^{n+2\gamma} (z\mathbb{A}(t))_{\#} (f_{(n,j)})_+^{\gamma}$ is a formal solution to our fourth order evolution (27), where $\mathbb{A}(t) = |S(t)|^{1/2}$, $f_{(n,j)}$ is defined by (13) and j counts the number of negative eigenvalues of $S(t)$. It becomes a classical solution if $j = 0 > \gamma$.


As it turns out, (31) and (35) can be reduced to quadratures in 2 dimensions (or if the effective dimension \check{n} is 2), but apparently not otherwise. When the effective dimension is larger, a qualitative analysis leads to Theorem 7. But first we study the case $\check{n} = 2$, both for the S equation (35) in rescaled time $dS/d\tau = S^{-1} + (\text{tr } S^{-1})\mathbb{I}/\gamma$, and for the \mathbb{P} equation (31). We omit the calculations for S , because they are easier than the ones for \mathbb{P} , which we do explain subsequently. The translation between (31) and (35) involves case distinctions from the value of the exponent in $|\det S| = |\det \mathbb{P}|^{-2\gamma/(n+2\gamma)}$, and both equations are useful for extracting different dynamical information. For example, the $t \searrow t_{\alpha}$ asymptotics are more easily accessed via S , whereas \mathbb{P} is more convenient for the $t \nearrow t_{\omega}$ limit.

Assume S has eigenvalues s_1 and s_2 with respective multiplicity n_1 and n_2 , so $n = n_1 + n_2$. The axes $s_1 s_2 = 0$ are excluded from the phase space, which is therefore disconnected. The orbits satisfy the homogeneous ODE

$$\frac{ds_2}{ds_1} = \frac{(\gamma + n_2)s_1 + n_1 s_2}{(\gamma + n_1)s_2 + n_2 s_1},$$

which has the invariant directions $s_2/s_1 = -(n_2 + \gamma)/(n_1 + \gamma)$ and $s_2/s_1 = 1$ unless $\gamma = -n_1 = -n_2 = -n/2$; its solutions satisfy

$$|s_2 - s_1|^{\gamma+n} |(\gamma + n_1)s_2 + (\gamma + n_2)s_1|^{\gamma} = \text{const}.$$



top: S dynamics (35) – horizontal axis: s_1 ; vertical axis: s_2
 bottom: P dynamics (31) – horizontal axis: $x = p_1$; vertical axis: $y = p_2$




Fig. 2. Dynamics of Eqs. (31) and (35) in the case $n_1 = n_2$ ($\tilde{n} = 2$). The graphs are qualitative and represent only topology and tangency, but not symmetries. Note that $\gamma < -n$ and $\gamma > 0$ have homeomorphic phase portraits, hence the cyclic arrangement of graphs.

For $\gamma = -n_1 = -n_2 = -n/2$, the solutions are $s_2/s_1 = const$, but for $\gamma = -n/2$, the P and S equations are not related anyway, because the transformation connecting them becomes singular. Phase portraits for both S and P in the case $n_1 = n_2 = n/2$ are collected in Fig. 2. Portraits for the case $n_1 = 3, n_2 = 1$, are collected in Figs. 3 and 4.

Returning to the P equation, let us assume that P has entry $p_1 =: x$ with multiplicity n_1 , and $p_2 =: y$ with multiplicity n_2 , with $n = n_1 + n_2$. Then orbits are described by the homogeneous ODE, which follows immediately from (31):

$$\frac{dy}{dx} = \frac{2y^3 + 2(n_1x + n_2y)y^2/\gamma + (n_1x^2 + n_2y^2)y/\gamma + (n_1x + n_2y)^2y/\gamma^2}{2x^3 + 2(n_1x + n_2y)x^2/\gamma + (n_1x^2 + n_2y^2)x/\gamma + (n_1x + n_2y)^2x/\gamma^2}$$

(or, of course, the reciprocal of this equation, to capture orbits where $dx/dy = 0$). It is easy to check for radial lines that are flow invariant and determine the stability or instability of the origin for the flow restricted to these lines: Equating the huge fraction with y/x gives a fourth degree polynomial equation of which three solutions $x = 0, y = 0$ and $y = x$ are obvious and allow a fourth solution to be found. The stability of the origin under the reduced dynamics on the radial direction is determined by going back to (31). The results of these routine calculations are listed in the




Fig. 3. Some cases of the dynamics of (35) and (31) for $\check{n} = 2, n_1 = 3, n_2 = 1, n = 4$. A variety of behavior in the range $-n < \gamma \leq -n/2$ already.

following table, in which the origin is stable if the given sign is negative, and unstable if it is positive. The ray consists of fixed points if the sign is 0:

Invariant direction	Stability of origin given by sign of
$y = 0$	$-(n_1 + \gamma)(n_1 + 2\gamma)$
$x = 0$	$-(n_2 + \gamma)(n_2 + 2\gamma)$
$y = x$	$-(n + \gamma)(n + 2\gamma)$
$\frac{y}{x} = -\frac{n_1 + \gamma}{n_2 + \gamma}$	$-(n_1 + \gamma)(n_2 + \gamma)(n + 2\gamma)/\gamma$

(36)

An explicit solution by quadratures can be found when $\check{n} = 2$, using the usual substitution $v = y/x$:

$$\frac{dx}{x} = \left(-\frac{n_1 + 2\gamma}{2\gamma} \frac{1}{v} + \frac{n + \gamma}{2\gamma} \frac{1}{v - 1} + \frac{1}{2} \frac{1}{v + (n_1 + \gamma)/(n_2 + \gamma)} \right) dv$$

which implies, after integration, the equation for the orbits:

$$|x|^{n_2 + 2\gamma} |y|^{n_1 + 2\gamma} |y - x|^{-n - \gamma} |(n_2 + \gamma)y + (n_1 + \gamma)x|^{-\gamma} = C. \tag{37}$$


As can be seen from table (36), a wide variety of dynamics occurs over the range of nonlinearities already in two dimensions. We give a graphical sketch for the case where $n_1 = n_2 = n/2$ in Fig. 2. For general dimension, we prove the following theorem.

Theorem 7 (Asymptotics of fourth order diffusion family).

- (a) For any $\gamma \neq 0$, the distance between distinct eigenvalues s_i of the same sign decreases as a function of time under the evolution (35).
- (b) If $\gamma > 0$ or $\gamma < -n$, then all maximal orbits $\mathbb{P}(t)$ of (31) converge to 0 as $t \rightarrow t_\omega = +\infty$, whereas $\|\mathbb{P}\| \rightarrow \infty$ as $t \rightarrow t_\alpha > -\infty$ (unless $\mathbb{P} \equiv 0$). If \mathbb{P} has a definite sign, then a more precise asymptotic is

$$\mathbb{P}(t) = \pm \frac{\gamma \mathbb{I}}{(2(n + \gamma)(n + 2\gamma)t)^{1/2}} (1 + o(1)) + O(t^{-(n+3\gamma)/2(n+\gamma)}) \quad \text{as } t \rightarrow \infty, \tag{38}$$

where the $O(t^{-(n+3\gamma)/2(n+\gamma)})$ term is traceless. As $t \rightarrow t_\alpha$, the eigenvalue of \mathbb{P} with largest absolute value blows up (the corresponding eigenvalue of \mathbb{S} with smallest absolute value goes to 0). All the other distinct eigenvalues of \mathbb{S} go to distinct finite nonzero limits. The leading order of the smallest (in absolute value) eigenvalue s_1 is $s_1 \sim c(t - t_\alpha)^{\gamma/(n_1 + 2\gamma)}$ as $t \rightarrow t_\alpha$.



top: S dynamics (35) – horizontal axis: s_1 ; vertical axis: s_2

bottom: P dynamics (31) – horizontal axis: $x = p_1$; vertical axis: $y = p_2$




Fig. 4. More cases of the dynamics of (35) and (31) for $\check{n} = 2, n_1 = 3, n_2 = 1, n = 4$.

- (c) If $\gamma = -n$, then still $t_\omega = \infty$, and each orbit converges to some multiple of the identity matrix: nonzero, if P has a definite sign, 0 otherwise. Unless $P = cI$, the same claims about t_α and its asymptotics hold as in (b).
- (d) If P has a specific sign and $-n < \gamma < -n/2$, then $\det P \rightarrow \infty$ as $t \rightarrow t_\omega$.

Proof. Claim (a) is immediate from $d(s_i - s_j)/d\tau = s_i^{-1} - s_j^{-1}$.

Write (31) as

$$-\frac{d}{dt} p_i = p_i \left[p_i^2 + \frac{1}{\gamma} \operatorname{tr} P^2 + \left(p_i + \frac{1}{\gamma} \operatorname{tr} P \right)^2 \right]. \tag{39}$$

Let $\|P\|_\infty := \max_i |p_i| =: |p_j|$. For $\gamma > 0$, it obviously follows that $\frac{d}{dt} \|P\|_\infty \leq -\|P\|_\infty^3$. A similar conclusion for $\gamma \leq -n$ follows from $\operatorname{tr} P^2 \leq n p_j^2$, namely $\frac{d}{dt} \|P\|_\infty \leq -(1 - \frac{n}{|\gamma|}) \|P\|_\infty^3$. These estimates give $t_\omega = \infty$ for $\gamma \leq -n$, as well as $t_\alpha > -\infty$ for $\gamma \notin [-n, 0]$, and hence the qualitative part of claim (b), and the first claim of (c).

If P has a definite sign, we may assume $P > 0$. (If $P < 0$, consider the same argument for $-P$.) The function $1 - (\operatorname{tr} P)^2 / (n \operatorname{tr} P^2)$ describes the sine square of the angular distance of a point from the diagonal $\mathbf{R}I$, and it turns out to be a Lyapunov function in the positive cone (for the dynamics projected on the unit sphere): Indeed, a straightforward calculation gives

$$\frac{d}{dt} \left(1 - \frac{(\operatorname{tr} P)^2}{n \operatorname{tr} P^2} \right) = \frac{4 \operatorname{tr} P}{n (\operatorname{tr} P^2)^2} \left(\operatorname{tr} P^3 \operatorname{tr} P^2 - \operatorname{tr} P^4 \operatorname{tr} P + \frac{1}{\gamma} (\operatorname{tr} P^2)^2 \operatorname{tr} P - \frac{1}{\gamma} \operatorname{tr} P^3 (\operatorname{tr} P)^2 \right). \tag{40}$$

Now the usual Hölder style interpolation of ℓ^p norms in \mathbf{R}^n gives

$$\operatorname{tr} P^3 \leq (\operatorname{tr} P^4)^{1/2} (\operatorname{tr} P^2)^{1/2} \quad \text{hence} \quad \operatorname{tr} P^4 \operatorname{tr} P \geq \frac{\operatorname{tr} P^3}{\operatorname{tr} P^2} \times (\operatorname{tr} P^3 \operatorname{tr} P).$$

With this estimate used on the second term in the parenthesis of (40), we conclude

$$\frac{d}{dt} \left(1 - \frac{(\text{tr } P)^2}{n \text{tr } P^2} \right) \leq \frac{-4 \text{tr } P}{n(\text{tr } P^2)^2} \left(\frac{\text{tr } P^3}{\text{tr } P^2} + \frac{1}{\gamma} \text{tr } P \right) \times (\text{tr } P^3 \text{tr } P - (\text{tr } P^2)^2). \tag{41}$$

The first parenthesis on the right is trivially positive if $\gamma > 0$, and positive by Hölder’s inequality if $\gamma < -n$. If $\gamma = -n$, it is still positive, unless $P \in \mathbf{RI}$. The second factor on the right is nonnegative by Hölder style interpolation again.

The orbits of (31), projected on the unit sphere, are orbits of a dynamical system there. By homogeneity of (31) different orbits whose initial points project onto the same point on the sphere, will, as a whole, have the same projection onto the sphere for all time, even though the time parametrization will depend on the radial coordinate of the initial point. We have shown that on the orthant of the unit sphere that comes from the intersection with the positive cone, $1 - (\text{tr } P)^2/n \text{tr } P^2$ is a strict Lyapunov function minimized at the diagonal point $\mathbb{I}/\|\mathbb{I}\|_2$. So this point is the $t \rightarrow t_\omega$ limit of the projected dynamics.

We have therefore shown that $P(t) = \sigma(t)\mathbb{I} + T(t)$ as $t \rightarrow t_\omega = \infty$, where the error term $T(t)$ is $o(\sigma(t))$; with no loss of generality, by a small modification of σ we may assume $T(t)$ to be traceless. The function σ is yet unspecified, except that we know $\sigma(t) \rightarrow 0$ as $t \rightarrow \infty$. Using this decomposition in (31), we conclude

$$-\frac{d\sigma}{dt} = \left(2 + \frac{n}{\gamma} \right) \left(1 + \frac{n}{\gamma} \right) (1 + o(1)) \sigma^3, \quad -\frac{dT}{dt} = \left(2 + \frac{n}{\gamma} \right) \left(3 + \frac{n}{\gamma} \right) (1 + o(1)) \sigma^2 T,$$

which can be integrated to obtain (38): first the asymptotic for σ , then for T . The $o(1)$ -term in σ is actually $O(T^2/\sigma^2)$ and could subsequently be improved from the T asymptotics if so desired. This proves the t_ω asymptotics of (b).

Still in (b), from $d \text{tr } S/d(-\tau) = -(1 + n/\gamma) \text{tr } S^{-1}$ it is clear that all $s_i > 0$ remain bounded above as $t \rightarrow t_\alpha$. Denoting the smallest s_i as s_1 , we find, in view of (a), that $s_1 = \frac{1}{n}[\text{tr } S - \sum (s_i - s_1)]$ decreases as $t \searrow t_\alpha$. So $\lim s_1$ exists. Now from (a) again, all $\lim s_i$ exist by monotonicity. If s_1 were positive, then all s_i would be bounded away from both ∞ and 0, and then the same would follow for the p_i , in contradiction to $\|P\|_\infty \rightarrow \infty$ as $t \searrow t_\alpha$. We therefore conclude $s_1 \rightarrow 0$. All eigenvalues distinct from s_1 must have a positive limit, due to (a). Absorbing all $\lim s_i$ other than 0 in a constant, Eq. (35) becomes $\frac{d}{dt} s_1 = \tilde{c} s_1^{-1-n_1/\gamma} (1 + O(s_1))$ and hence $s_1 \sim c(t - t_\alpha)^{\gamma/(n_1+2\gamma)}$ as $t \rightarrow t_\alpha$.

Concerning (c), we have already obtained the boundedness of the orbits and $t_\omega = \infty$ from the previous calculation $\frac{d}{dt} \|P\|_\infty \leq 0$. By LaSalle’s invariance principle (see, e.g., [66, VII, §30]), the ω -limit set consisting of all forward accumulation points of a given orbit must be a connected invariant subset of the set on which $\frac{d}{dt} \|P\|_\infty$ vanishes. Vanishing of this derivative implies that P is a multiple of \mathbb{I} , so the subset in question could only be an interval on the diagonal \mathbf{RI} . Each P on the diagonal is an equilibrium when $\gamma = -n$. If P is not (positive or negative) definite, this (together with the fact that the evolution preserves the signs of the p_i) only leaves the possibility $P \rightarrow 0$.

Now assume P (and hence S , too) has a definite sign; we still want to show that the compact interval on the diagonal must be a singleton. Use (35) with a rescaled time τ , namely $\frac{d}{d\tau} S = S^{-1} + \frac{1}{\gamma} \text{tr } S^{-1}$, hence $\frac{1}{2} \frac{d}{d\tau} \text{tr } S^2 = n + \frac{1}{\gamma} \text{tr } S \text{tr } S^{-1} \leq n - n^2/|\gamma|$, which equals 0 for $\gamma = -n$. By monotonicity, $\lim_{\tau \rightarrow \tau_\omega} \text{tr } S^2$ exists. This, together with the previous conclusion about the ω -limit set, forces the ω -limit set to be a singleton, i.e., $\lim_{t \rightarrow \infty} P(t) = c\mathbb{I}$ for some c . Now $c \neq 0$ because else $\text{tr } S^2 \rightarrow \infty$. This proves the t_ω claims of (c).

For the t_α limit, assume $P \neq c\mathbb{I}$. Then we still get from (41) that $\frac{d}{dt} (1 - (\text{tr } P)^2/(n \text{tr } P^2)) < 0$. So there exists an $\varepsilon > 0$ such that $(\text{tr } P)^2/(n \text{tr } P^2) < (1 - \varepsilon)$ as $t \rightarrow t_\alpha$. Then by an immediate indirect proof, there exists also some $\delta > 0$ such that $\text{tr } P < n(1 - \delta)p_j$. This, with (39), gives $-\frac{d}{dt} \|P\|_\infty \geq \delta^2 \|P\|^3$, hence $\|P\|_\infty \rightarrow \infty$ as $t \rightarrow t_\alpha > -\infty$. The arguments from (b) for the quantitative asymptotics carry over literally now.

For $-n < \gamma$, we even have $d \text{tr } S^2/d\tau \leq -c < 0$ (and hence $\tau_\omega < \infty$). We cannot conclude $S \rightarrow 0$ by Lyapunov’s method, due to the singularity on the hyperplanes $\det S = 0$. But as either Lyapunov’s conclusion applies, or else a singularity is hit, we can conclude $\det S \rightarrow 0$ as $\tau \rightarrow \tau_\omega$ either way. For $\gamma < -n/2$, this implies $\det P \rightarrow \infty$, thus proving (d). \square

A study of asymmetric multiplicities $n_1 \neq n_2$ in effective dimension $\check{n} = 2$ reveals that the exact rate of blowup of one or several eigenvalues may depend further on the initial data. Depending on γ and initial data, only some or all s_i may go to 0; see Fig. 3. In the few cases studied, the relative asymptotics observed implies $t_\omega < \infty$, but deciding validity of this conjecture in general remains a task for some further research. Moreover, for $\gamma = -n/2$, there arise

affine images of Barenblatt with $t_\alpha = -\infty$ whose initial profile as $t \searrow t_\alpha$ is not a singular measure, but a regular density. It is remarkable that in the cases $\gamma = -n$ (or equivalently $m = 1 - 1/n$) and $\gamma = -n/2$ ($m = 1 - 2/n$), the Barenblatt profile is a stationary point of the dynamics (27). According to Fig. 3, we see that this profile is unstable for $\gamma = -n/2$, but is stable at least within the invariant subspace of quadratic pressure solutions for $\gamma = -n$, due to Theorem 7(c). Whether it remains globally attractive for fourth order diffusions with $\gamma \notin [-n, 0]$ and a wide class of initial data, as in the second order case, remains an open question. For the special case $(n, m) = (1, 3/2)$ it was addressed by Bernoff and Witelski via linearization [12], nonlinearly by Carrillo and Toscani [21], and nonlinearly in all dimensions but with $m = 1$ by Gianazza, Savaré and Toscani [41]. The analogous question in a periodic domain— attractivity of the constant solution—was addressed for $(n, m) = (1, 1)$ by Cáceres, Carrillo and Toscani [20], and in higher dimensions by Jüngel and Mattes [43].

Let us conclude our study of fourth order nonlinear diffusions by showing that every degenerate matrix $\det S_0 = 0$ is either the initial or final point of at least one orbit of the dynamical system (35), and complement Theorem 7(b) by giving the asymptotics of the orbit at that limit. To do so, it is convenient to replace $S(t)$ by an effective matrix $Q(t)$ whose eigenvalues $q_i(t)$ are distinct, but coincide with those of $S(t)$ otherwise; they come with multiplicities $n_1, \dots, n_{\check{n}}$ in $S(t)$. Obviously $Q(t)$ and the integer matrix N of multiplicities are $\check{n} \times \check{n}$ diagonal matrices satisfy the evolution

$$\begin{aligned} \frac{dq_i}{d\tau} &:= \frac{1}{2|\det S(t)|^{-1/\gamma}} \frac{dq_i}{dt} \\ &= q_i^{-1} + \frac{1}{\gamma} \operatorname{tr} N Q^{-1}. \end{aligned} \tag{42}$$

Theorem 8 (Affinely self-similar fourth-order dynamics from sources concentrated on subspaces). *Let Q_0 and N be diagonal $\check{n} \times \check{n}$ matrices, with eigenvalues $\bar{q}_1, \dots, \bar{q}_{\check{n}} \in \mathbf{R}$ and $n_1, \dots, n_{\check{n}} \in \mathbf{N}$. Suppose $\bar{q}_1 = 0$ and $0 \neq \gamma \neq -n_1$. If the eigenvalues of Q_0 are distinct, then precisely two maximal smooth orbits $\det Q(t) \neq 0$ of (42) include Q_0 in their closures. These two orbits $Q(t)$ have unit tangent vectors which parallel $\pm(1 + \frac{\gamma}{n_1}, 1, \dots, 1)$ at Q_0 . If $\gamma \notin [-n_1, 0]$ then $Q_0 = \lim_{t \rightarrow t_\alpha} Q(t)$ and $t_\alpha \in \mathbf{R}$; otherwise $Q_0 = \lim_{t \rightarrow t_\omega} Q(t)$ with $t_\omega \in \mathbf{R}$ if $\gamma \in]-n_1, -n_1/2[$, and $t_\omega = +\infty$ if $\gamma \in [-n_1/2, 0[$, as in (46)–(47).*

Proof. The orbits $Q(t) = (q_1(t), \dots, q_{\check{n}}(t))$ of the dynamical system (42) are tangent to the field of unit vectors whose directions are given by

$$\frac{dq_k}{dq_1} = \frac{q_1(q_k^{-1} + \operatorname{tr} N Q^{-1}/\gamma)}{1 + q_1 \operatorname{tr} N Q^{-1}/\gamma} \tag{43}$$

for $\det Q \neq 0$ and $k > 1$. Since the eigenvalues of Q_0 are distinct, we see $q_1 \operatorname{tr} N Q^{-1}$ is a smooth function of $Q = (q_1, \dots, q_{\check{n}})$ in a neighborhood of $Q_0 = \operatorname{diag}(0, \bar{q}_2, \dots, \bar{q}_3)$, with limit $q_1 \operatorname{tr} N Q^{-1} \rightarrow n_1$ as $q_1 \rightarrow 0$. Thus the direction field (43) of unit tangent vectors extends smoothly to a neighborhood of Q_0 , with

$$\left. \frac{dq_k}{dq_1} \right|_{Q_0} = \frac{n_1/\gamma}{1 + n_1/\gamma}. \tag{44}$$

The theory of ordinary differential equations asserts this smooth direction field has a (unique) integral curve passing through Q_0 ; it intersects the plane $q_1 = 0$ transversally since (44) is nonvanishing. Since orbits of the dynamical system (42) lie on integral curves of (43), at most two such orbits $\det Q(t) \neq 0$ approach Q_0 . On the other hand, for $Q(t)$ near Q_0 , the sign of

$$q_1 \frac{dq_1}{d\tau} = 1 + \frac{n_1}{\gamma} + O(q_1) \quad \text{as } q_1 \rightarrow 0, \tag{45}$$

shows Q_0 is the initial point $\tau \searrow \tau_\alpha > -\infty$ of both orbits if $\gamma \notin [-n_1, 0]$, and the final limit $\tau \nearrow \tau_\omega < +\infty$ of both orbits $Q(t)$ otherwise. This shows Q_0 is either a fixed point of the dynamics, as for $\gamma = -n_1/2$, or a singular point. Either way, the lemma is proved. Integrating (45) yields

$$\frac{q_1(\tau)^2}{2} = \begin{cases} -(1 + \frac{n_1}{\gamma})(\tau_\omega - \tau) + O(|\tau_\omega - \tau|^{3/2}) & \text{if } \gamma \in]-n_1, 0[, \\ (1 + \frac{n_1}{\gamma})(\tau - \tau_\alpha) + O(|\tau - \tau_\alpha|^{3/2}) & \text{if } \gamma \notin [-n_1, 0]. \end{cases} \tag{46}$$

Thus $q_i(\tau) - \bar{q}_i = O(|\tau - \tau_0|^{1/2})$ as $\tau \rightarrow \tau_0$ for $i = 1, \dots, \check{n}$, where $\tau_0 = \tau_\alpha$ or τ_ω and the limit is taken from above or below according to the sign of $1 + n_1/\gamma$. Using this to integrate

$$\begin{aligned} \frac{dt}{d\tau} &= \frac{1}{2} \prod_{k=1}^{\check{n}} |q_k(\tau)|^{n_k/\gamma} \\ &= \left| 2 \left(1 + \frac{n_1}{\gamma} \right) (\tau - \tau_0) \right|^{n_1/2\gamma} \left(\frac{1}{2} + O(|\tau - \tau_0|^{1/2}) \right) \prod_{k=2}^{\check{n}} |\bar{q}_k|^{n_k/\gamma} \end{aligned}$$

yields

$$\begin{aligned} t - t_0 &\sim \frac{\gamma c}{n_1 + 2\gamma} |\tau - \tau_0|^{(n_1+2\gamma)/2\gamma} && \text{with } t_0 \in \mathbf{R} \quad \text{if } \gamma \notin [-n_1/2, 0], \\ t &\sim -\frac{c}{2} \log |\tau_\omega - \tau| && \text{with } t_\omega = +\infty \text{ if } \gamma = -n_1/2, \\ t &\sim -\frac{\gamma c}{n_1+2\gamma} |\tau_\omega - \tau|^{(n_1+2\gamma)/2\gamma} && \text{with } t_\omega = +\infty \text{ if } \gamma \in]-n_1/2, 0], \end{aligned} \tag{47}$$

as $\tau \rightarrow \tau_0$, where $c = |2(1 + \frac{n_1}{\gamma})|^{n_1/2\gamma} \prod_{k=2}^{\check{n}} |\bar{q}_k|^{n_k/\gamma} > 0$. \square

Notice the limiting tangent direction $\pm(1 + \frac{\gamma}{n_1}, 1, \dots, 1)$ of orbits in this result always lies in a plane, but rotates 180 degrees over the full range of $\gamma \in \mathbf{R}$, which helps to explain the diversity of dynamics represented in Figs. 2, 3, and 4.

5. Outlook

It is interesting to consider a combination of the porous medium model and the fourth order family (27), in the spirit of the quoted work by Pugh and co-authors [13,48,58] on the thin film equation. The 2-Wasserstein gradient flow of the combined energy $I_m(u) - \mu E_m(u)$ (with μ a constant that could actually be scaled away, except for its sign) leads to the equation

$$\frac{\partial \rho}{\partial t} = -\frac{1}{m-1/2} \nabla \cdot [\rho \nabla (\rho^{m-3/2} \Delta(\rho^{m-1/2}))] - \frac{\mu}{m} \Delta(\rho^m) \tag{48}$$

and with the same pressure substitution and quadratic pressure ansatz, we get the equations

$$-\frac{dP(t)}{dt} = 2P^3 + 2P^2 \frac{\text{tr } P}{\gamma} + P \frac{\text{tr}(P^2)}{\gamma} + P \frac{(\text{tr } P)^2}{\gamma^2} + \mu \frac{\text{tr } P}{\gamma} P + 2\mu P^2, \tag{49}$$

$$-\frac{1}{p_0} \frac{dp_0}{dt} = \frac{\text{tr}(P^2)}{\gamma} + \frac{(\text{tr } P)^2}{\gamma^2} + \mu \frac{\text{tr } P}{\gamma}. \tag{50}$$

Due to the different powers in P on the right-hand side, one may ask the question on whether there are nontrivial stationary solutions to this equation. It turns out that this is indeed the case:

Theorem 9. *Assume $m \neq 0$ and $0 \neq \gamma \neq -\frac{n}{2}$. Among invertible matrices P , the stationary solutions to (49)–(50) are exactly $P = -\frac{\gamma\mu}{n+\gamma} \mathbb{I}$.*

Proof. We are looking for solutions to $\frac{d}{dt} P = 0, \frac{d}{dt} p_0 = 0$. Canceling P from the right-hand side of (49) and taking the trace, we get $n + 2\gamma$ times the right-hand side of (50). Incidentally, this shows that (50) is not an independent condition for an equilibrium solution; this is of course to be expected, from mass preservation of the partial differential equation described by this ODE system. Subtracting \mathbb{I} times (50) from P^{-1} times (49), we can cancel another factor P and get, as a necessary condition for stationary solutions, that $P + \frac{1}{\gamma} (\text{tr } P) \mathbb{I} + \mu \mathbb{I} = 0$, i.e., P is a multiple of \mathbb{I} . Plugging the ansatz $P = \alpha \mathbb{I}$ back into the equation determines α uniquely and shows sufficiency. \square

Is this radially symmetric, stationary profile stable? Linearization of (49) about the equilibrium (with \dot{P} the linearized variable) gives

$$-\frac{d\dot{P}(t)}{dt} = \frac{2\gamma^2\mu^2}{(n+\gamma)^2}\dot{P} + \frac{\mu^2(n+3\gamma)}{(n+\gamma)^2}(\text{tr } \dot{P})\mathbb{I} =: \frac{\mu^2}{(n+\gamma)^2}(a\dot{P} + b(\text{tr } \dot{P})\mathbb{I}).$$

Again, with no loss of generality, \dot{P} is diagonal. Linearized stability *within* the reduced dynamics of the quadratic family is therefore determined by the eigenvalues of the matrix $a\mathbb{I} + b[1, \dots, 1]^T[1, \dots, 1]$, where $a = 2\gamma^2$ and $b = n + 3\gamma$. This matrix has the eigenvalues $a + nb = (n + \gamma)(n + 2\gamma)$ (simple, with eigenvector $[1, \dots, 1]^T$) and a (with eigenspace the orthocomplement of $[1, \dots, 1]^T$). This gives stability of the radially symmetric quadratic profile within the reduced dynamics provided $\gamma \notin [-n, -\frac{n}{2}]$.

The stability of this ‘parabolic soliton’ under the full dynamics is an interesting question, currently under investigation.

6. Appendix: Weak pressure and density evolutions

This appendix is devoted to showing the positive truncation $\rho = (\pi/\gamma)_+^\gamma$ of our quadratic pressure (7) yields a weak solution to the porous medium equation in the pressure form (5) with $u = \rho^{1/\gamma}$ and in the density form (6).

Lemma 10 (*Non-negative solutions from signed solutions*). *If $u \in C^2(\mathbf{R}^n \times [0, \infty[)$ is a classical solution to $\partial u/\partial t = u\Delta u + \gamma|\nabla u|^2$, then $u_+ := \max\{u, 0\}$ is a weak solution to this same equation in the sense of (5).*

Proof. Since $u_+ \in W_{\text{loc}}^{1,2}$, the term $\int u_+ \psi_t dt$ in (5) can be integrated by parts with $(u_+)_t = u_t \chi_{\{u>0\}}$, as in e.g. Lieb and Loss [49, §6]. Likewise $\nabla u_+ = (\nabla u) \chi_{\{u>0\}}$. To show that for each t ,

$$\int_{\{u(t, \cdot) > 0\}} ((-u\Delta u - \gamma|\nabla u|^2)\psi - u\nabla u \cdot \nabla \psi + (\gamma - 1)|\nabla u|^2\psi) dx = 0$$

we note that the left-hand side reduces to $-\int_{\{u>0\}} \text{div}(u\psi\nabla u) dx$, which vanishes without further geometric hypotheses on the boundary of $\Omega := \{x \in \mathbf{R}^n \mid u(x, t) > 0\}$, because the C^1 vector field $v := u\psi\nabla u$ vanishes on the boundary of the open set $\Omega \subset \mathbf{R}^n$. To prove this, fix a compact set $\Omega_\varepsilon \subset \Omega$ whose boundary is in an ε -neighborhood of $\partial\Omega$, and a smooth cutoff function $\eta_\varepsilon \in C_c^\infty(\Omega; [0, 1])$, compactly supported in Ω and with $\eta_\varepsilon = 1$ on Ω_ε . Specifically, we can mollify $\hat{\eta}_\varepsilon := \min\{1, \frac{3}{\varepsilon}(\text{dist}(x, \partial\Omega) - \frac{\varepsilon}{3})_+\}$. Then we have $\text{div } v = O(1)$ on $\Omega \setminus \Omega_\varepsilon$, and the measure of $\Omega \setminus \Omega_\varepsilon$ goes to 0 as $\varepsilon \rightarrow 0$. Therefore

$$\begin{aligned} \int_{\Omega} \text{div } v &= \int_{\Omega_\varepsilon} \text{div}(v\eta_\varepsilon) + \int_{\Omega \setminus \Omega_\varepsilon} \text{div } v \\ &= \int_{\Omega} \text{div}(v\eta_\varepsilon) - \int_{\Omega \setminus \Omega_\varepsilon} ((\text{div } v)\eta_\varepsilon + v \cdot \nabla \eta_\varepsilon) + o(1) \quad \text{as } \varepsilon \rightarrow 0. \end{aligned}$$

The first integral vanishes, since Ω can be deformed inwards to a smooth domain on which the divergence theorem applies, still avoiding the compact support of η_ε . The remaining terms are small, since $\|v\|_\infty = O(\varepsilon)$ balances $\|\nabla \eta_\varepsilon\|_\infty = O(1/\varepsilon)$ as $\varepsilon \rightarrow 0$, while $\text{div } v$ and $\|\eta_\varepsilon\|_\infty$ are bounded and the domain of integration has measure going to 0. \square

A final lemma asserts that the weak solutions (5) we described to the pressure evolution (3) also give rise to weak solutions of the density evolution (1).

Lemma 11 (*Weak pressure evolutions yield weak density evolutions*). *Fix $\gamma > 0$. Assume a weak solution u to $u_t = u\Delta u + \gamma|\nabla u|^2$, in the sense of (5) is Lipschitz (uniformly on compact subsets of $[0, \infty[\times\mathbf{R}^n$). Moreover, if $\gamma < 1$, assume $\nabla u^\gamma \in L_{\text{loc}}^1([0, \infty[\times\mathbf{R}^n)$, or $\nabla u^{(1+\gamma)/2} \in L_{\text{loc}}^2([0, \infty[\times\mathbf{R}^n)$. Then $\rho := u^\gamma$ is a weak solution of the porous medium equation in the sense of (6).*

Proof. For a Lipschitz function u to be a weak solution, the expression

$$V := \int u_0 \psi_0 \, dx + \iint (\psi_t u + (\gamma - 1) |\nabla u|^2 \psi - u \nabla u \cdot \nabla \psi) \, dx \, dt$$

vanishes for all ψ Lipschitz with compact support. Guided by the calculation with classical solutions, we'd like to choose $\psi := \gamma u^{\gamma-1} \varphi$. We can choose $\psi_\varepsilon := \gamma(u + \varepsilon)^{\gamma-1} \varphi$ for $\varphi \in C^1$ with compact support. Then

$$\begin{aligned} V/\gamma &= \int u_0 (u_0 + \varepsilon)^{\gamma-1} \varphi_0 \, dx + \iint (\gamma - 1) (u + \varepsilon)^{\gamma-2} u u_t \varphi \, dx \, dt \\ &\quad + \iint (u + \varepsilon)^{\gamma-1} u \varphi_t \, dx \, dt + (\gamma - 1) \iint |\nabla u|^2 (u + \varepsilon)^{\gamma-1} \varphi \, dx \, dt \\ &\quad - \iint u \nabla u (\gamma - 1) (u + \varepsilon)^{\gamma-2} \nabla u \varphi \, dx \, dt - \iint u \nabla u (u + \varepsilon)^{\gamma-1} \nabla \varphi \, dx \, dt. \end{aligned}$$

The first, third and sixth terms converge to their formal limits by the dominated convergence theorem. Terms 4 and 5 can be merged; term 2 gets split into an easy part and one with an explicit factor ε :

$$\begin{aligned} V/\gamma &= \int u_0^\gamma \varphi_0 \, dx + \iint u^\gamma \varphi_t \, dx \, dt - \iint u^\gamma \nabla u \nabla \varphi \, dx \, dt + (\gamma - 1) \iint (u + \varepsilon)^{\gamma-1} u_t \varphi \, dx \, dt \\ &\quad - (\gamma - 1) \varepsilon \iint (u + \varepsilon)^{\gamma-2} (u_t - |\nabla u|^2) \varphi \, dx \, dt + o(1) \quad \text{as } \varepsilon \rightarrow 0. \end{aligned}$$

Writing $(u + \varepsilon)^{\gamma-1} u_t = \gamma^{-1} \partial_t (u + \varepsilon)^\gamma$ and $(\gamma - 1) (u + \varepsilon)^{\gamma-2} u_t = \partial_t (u + \varepsilon)^{\gamma-1}$, we integrate the u_t terms by parts, set $u = \rho^{m-1} = \rho^{1/\gamma}$, and use dominated convergence again to get

$$\begin{aligned} V/\gamma &= \gamma^{-1} \int \rho_0 \varphi_0 \, dx + \gamma^{-1} \iint \rho \varphi_t \, dx \, dt - \frac{1}{\gamma + 1} \iint \nabla \rho^{(\gamma+1)/\gamma} \nabla \varphi \, dx \, dt \\ &\quad + (\gamma - 1) \varepsilon \iint (u + \varepsilon)^{\gamma-2} |\nabla u|^2 \varphi \, dx \, dt + o(1) \end{aligned}$$

as $\varepsilon \rightarrow 0$. (This uses the estimate $\varepsilon(u + \varepsilon)^{\gamma-1} = O(\varepsilon^{\min(1, \gamma)})$ on the support of φ .) We have to show that the last integral disappears as $\varepsilon \rightarrow 0$. For $\gamma \geq 2$ the dominated convergence theorem completes the proof. If $1 < \gamma < 2$, then $\varepsilon(u + \varepsilon)^{\gamma-2} \leq \varepsilon^{\gamma-1}$ implies this term is $O(\varepsilon^{\gamma-1})$. If $\gamma = 1$, its coefficient is zero anyways. Finally, for $0 < \gamma < 1$, we have $\varepsilon(u + \varepsilon)^{\gamma-2} \leq (u + \varepsilon)^{\gamma-1} \leq u^{\gamma-1}$. Since ∇u and φ are bounded on the support of φ , dominated convergence again applies in view of our hypotheses on ∇u^γ or $\nabla u^{(1+\gamma)/2}$. In the limit we retrieve (6). \square

References

- [1] L.A. Ambrosio, N. Gigli, G. Savaré, Gradient Flows in Metric Spaces and in the Space of Probability Measures, Lecture Notes in Mathematics ETH Zürich, Birkhäuser Verlag, Basel, 2005.
- [2] S.B. Angenent, Large time asymptotics for the porous medium equation, in: Nonlinear Diffusion Equations and their Equilibrium States I, in: Math. Sci. Res. Inst. Publ., vol. 12, Springer, New York, 1988, pp. 21–34.
- [3] S.B. Angenent, D.G. Aronson, Optimal asymptotics for solutions to the initial value problem for the porous medium equation, in: T.S. Angell, et al. (Eds.), Nonlinear Problems in Applied Mathematics: in honor of Professor Ivar Stakgold on his 70th birthday, Society for Industrial and Applied Mathematics, Philadelphia, 1996, pp. 10–19.
- [4] S.B. Angenent, D.G. Aronson, Non-axial self-similar hole filling for the porous medium equation, J. Amer. Math. Soc. 14 (2001) 737–782.
- [5] S.B. Angenent, D.G. Aronson, S.I. Betelú, J.S. Lowengrub, Focusing of an elongated hole in porous medium flow, Physica D 151 (2001) 228–252.
- [6] D.G. Aronson, J. Graveleau, A self-similar solution to the focusing problem for the porous medium equation, Eur. J. Appl. Math. 4 (1993) 65–81.
- [7] D.G. Aronson, J. Bouwe van den Berg, J. Hulshof, Parametric dependence of exponents and eigenvalues in focusing porous media flows, Eur. J. Appl. Math. 14 (2003) 485–512.
- [8] G.I. Barenblatt, On some unsteady motions of a liquid or gas in a porous medium, Akad. Nauk SSSR Prikl. Mat. Mekh. 16 (1952) 67–78.
- [9] G.I. Barenblatt, Scaling, Cambridge University Press, Cambridge, 2003.
- [10] G.I. Barenblatt, M. Bertsch, A.E. Chertock, V.M. Prostokishin, Self-similar intermediate asymptotics for a degenerate parabolic filtration absorption equation, Proc. Natl. Acad. Sci. USA 97 (2000) 9844–9848.
- [11] F. Bernis, A. Friedman, Higher order nonlinear degenerate parabolic equations, J. Differential Equations 83 (1990) 179–206.
- [12] A.J. Bernoff, T.P. Witelski, Linear stability of source-type similarity solutions of the thin film equation, Appl. Math. Lett. 15 (2002) 599–606.

- [13] A.L. Bertozzi, M.C. Pugh, Finite-time blow-up of solutions of some long-wave unstable thin film equations, *Indiana Univ. Math. J.* 49 (2000) 1323–1366.
- [14] M. Bertsch, R. Dal Passo, H. Garcke, G. Grün, The thin viscous flow equation in higher space dimensions, *Adv. Differential Equations* 3 (1998) 417–440.
- [15] M. Bertsch, R. Dal Passo, M. Ughi, Discontinuous ‘viscosity’ solutions of a degenerate parabolic equation, *Trans. Amer. Math. Soc.* 320 (1990) 779–798.
- [16] M. Bertsch, R. Dal Passo, M. Ughi, Nonuniqueness of solutions of a degenerate parabolic equation, *Ann. Mat. Pura Appl.* 161 (4) (1992) 57–81.
- [17] M. Bertsch, M. Ughi, Positivity properties of viscosity solutions of a degenerate parabolic equation, *Nonlinear Anal.* 14 (1990) 571–592.
- [18] S. Betelú, J.R. King, Explicit solutions of a two-dimensional fourth order nonlinear diffusion equation, *Math. Comput. Modelling* 37 (2003) 395–403.
- [19] M.J. Cáceres, G. Toscani, Kinetic approach to long time behavior of linearized fast diffusion equations, *J. Statist. Phys.* 128 (2007) 883–925.
- [20] M.J. Cáceres, J.A. Carrillo, G. Toscani, Long-time behavior for a nonlinear fourth-order parabolic equation, *Trans. Amer. Math. Soc.* 357 (2005) 1161–1175.
- [21] J.A. Carrillo, G. Toscani, Long-time asymptotics for strong solutions of the thin film equation, *Comm. Math. Phys.* 225 (2002) 551–571.
- [22] J.A. Carrillo, A. Jüngel, P.A. Markowich, G. Toscani, A. Unterreiter, Entropy dissipation methods for degenerate parabolic problems and generalized Sobolev inequalities, *Monatsh. Math.* 133 (2001) 1–82.
- [23] J.A. Carrillo, G. Toscani, Asymptotic L^1 -decay of solutions of the porous medium equation to self-similarity, *Indiana Univ. Math. J.* 49 (2000) 113–141.
- [24] J.A. Carrillo, J.L. Vázquez, Fine asymptotics for fast diffusion equations, *Comm. Partial Differential Equations* 28 (2003) 1023–1056.
- [25] J.A. Carrillo, M. Di Francesco, G. Toscani, Strict contractivity of the 2-Wasserstein distance for the porous medium equation by mass-centering, *Proc. Amer. Math. Soc.* 135 (2007) 353–363.
- [26] J.A. Carrillo, R.J. McCann, C. Villani, Contractions in the 2-Wasserstein length space and thermalization of granular media, *Arch. Rational Mech. Anal.* 179 (2006) 217–263.
- [27] E. Chasseigne, J.L. Vázquez, Theory of extended solutions for fast-diffusion equations in optimal classes of data. Radiation from singularities, *Arch. Rational Mech. Anal.* 164 (2002) 133–187.
- [28] A. Chertock, On the stability of a class of self-similar solutions to the filtration-absorption equation, *Eur. J. Appl. Math.* 13 (2002) 179–194.
- [29] B. Chow, D. Knopf, *The Ricci Flow: An Introduction*, Mathematical Surveys and Monographs, vol. 110, American Mathematical Society, Providence, RI, 2004.
- [30] R. Dal Passo, S. Luckhaus, A degenerate diffusion problem not in divergence form, *J. Differential Equations* 69 (1987) 1–14.
- [31] M. Del Pino, J. Dolbeault, Best constants for Gagliardo–Nirenberg inequalities and applications to nonlinear diffusions, *J. Math. Pures Appl.* 81 (2002) 847–875.
- [32] J. Denzler, R.J. McCann, Phase transitions and symmetry breaking in singular diffusion, *Proc. Natl. Acad. Sci. USA* 100 (2003) 6922–6925.
- [33] J. Denzler, R.J. McCann, Fast diffusion to self-similarity: complete spectrum, long time asymptotics, and numerology, *Arch. Rational Mech. Anal.* 175 (2005) 301–342.
- [34] B. Derrida, J.L. Lebowitz, E. Speer, H. Spohn, Fluctuations of a stationary nonequilibrium interface, *Phys. Rev. Lett.* 67 (1991) 165–168.
- [35] J.A. Diez, L.P. Thomas, S. Betelú, R. Gratton, B. Marino, J. Gratton, D.G. Aronson, S.B. Angenent, Noncircular converging flows in viscous gravity currents, *Phys. Rev. E* 58 (1998) 6182–6187.
- [36] A. Friedman, S. Kamin, The asymptotic behaviour of a gas in an n -dimensional porous medium, *Trans. Amer. Math. Soc.* 262 (1980) 551–563.
- [37] V.A. Galaktionov, Invariant subspaces and new explicit solutions to evolution equations with quadratic nonlinearities, *Proc. Roy. Soc. Edinburgh Sect. A* 125 (1993) 225–246.
- [38] L. Giacomelli, H. Knüpfer, Flat data solutions to the thin film equation do not rupture, Preprint SFB 611, n. 283, 2006.
- [39] L. Giacomelli, F. Otto, Variational formulation for the lubrication approximation of the Hele–Shaw flow, *Calc. Var. Partial Differential Equations* 13 (2001) 377–404.
- [40] L. Giacomelli, F. Otto, Rigorous lubrication approximation, *Interfaces Free Bound* 5 (2003) 483–529.
- [41] U. Gianazza, G. Savaré, G. Toscani, The Wasserstein gradient flow of the Fisher information and the quantum drift-diffusion equation, Preprint at <http://www.imati.cnr.it/~savare/publicazioni/preprints.html>, 2006.
- [42] J. Graveleau, Quelques solutions auto-semblables pour l’équation de la chaleur non-linéaire, Rapport Interne C.E.A., 1972.
- [43] A. Jüngel, D. Matthes, The Derrida–Lebowitz–Speer–Spohn equation: existence, non-uniqueness, and decay rates of the solutions, *SIAM J. Math. Anal.* (2007), in press.
- [44] A. Jüngel, R. Pinnau, Global nonnegative solutions of a nonlinear fourth-order parabolic equation for quantum systems, *SIAM J. Math. Anal.* 32 (2000) 760–777.
- [45] Y.J. Kim, R.J. McCann, Sharp decay rates for the fastest conservative diffusions, *C. R. Acad. Sci. Paris, Ser. I* 341 (2005) 157–162.
- [46] Y.J. Kim, R.J. McCann, Potential theory and optimal convergence rates in fast nonlinear diffusion, *J. Math Pures Appl.* 86 (2006) 42–67.
- [47] J.R. King, Exact multidimensional solutions to some nonlinear diffusion equations, *Quart. J. Mech. Appl. Math.* 46 (1993) 419–436.
- [48] R.S. Laugesen, M.C. Pugh, Heteroclinic orbits, mobility parameters and stability for thin film type equations, *Electron. J. Differential Equations* 95 (2002) 1–29.
- [49] E.H. Lieb, M. Loss, *Analysis*, American Mathematical Society, Providence, RI, 1997.
- [50] R.J. McCann, D. Slepčev, Second-order asymptotics for the fast-diffusion equation, *Int. Math. Res. Not.* 24947 (2006) 1–22.
- [51] T.G. Myers, Thin films with high surface tension, *SIAM Rev.* 40 (1998) 441–462.
- [52] A. Oron, S.H. Davis, S.G. Bankoff, Long-scale evolution of thin liquid films, *Rev. Modern Phys.* 69 (1997) 931–980.
- [53] F. Otto, Lubrication approximation with prescribed nonzero contact angle, *Comm. Partial Differential Equations* 23 (1998) 2077–2164.

- [54] F. Otto, The geometry of dissipative evolution equations: the porous medium equation, *Comm. Partial Differential Equations* 26 (2001) 101–174.
- [55] R.E. Pattle, Diffusion from an instantaneous point source with concentration dependent coefficient, *Quart. J. Mech. Appl. Math.* 12 (1959) 407–409.
- [56] V.V. Pukhnachev, Exact multidimensional solutions of the nonlinear diffusion equation, *J. Appl. Mech. Tech. Phys.* 36 (1995) 169–176.
- [57] G.A. Rudykh, E.I. Semenov, The construction of exact solutions of the multi-dimensional quasilinear heat-conduction equation, *Comp. Math. Math. Phys.* 33 (1993) 1087–1097.
- [58] D. Slepčev, M.C. Pugh, Selfsimilar blowup of unstable thin-film equations, *Indiana Univ. Math. J.* 54 (2005) 1697–1738.
- [59] L. Tartar, Solutions particulières de $U_t = \Delta U^m$ et comportement asymptotique, Unpublished, 1986.
- [60] G. Toscani, A central limit theorem for solutions of the porous medium equation, *J. Evol. Equ.* 5 (2005) 185–203.
- [61] M. Ughi, A degenerate parabolic equation modelling the spread of an epidemic, *Ann. Mat. Pura Appl.* 143 (4) (1986) 385–400.
- [62] J.L. Vázquez, Asymptotic behaviour and propagation of the one-dimensional flow of gas in a porous medium, *Trans. Amer. Math. Soc.* 277 (1983) 507–527.
- [63] J.L. Vázquez, An introduction to the mathematical theory of the porous medium equation, in: *Shape Optimization and Free Boundaries*, Montreal, PQ, 1990, in: NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., vol. 380, Kluwer Acad. Publ., 1992, pp. 347–389.
- [64] J.L. Vázquez, Asymptotic behaviour for the porous medium equation posed in the whole space, *J. Evol. Equ.* 3 (2003) 67–118.
- [65] J.L. Vázquez, *The Porous Medium Equation. Mathematical Theory*, Oxford University Press, Oxford, 2007.
- [66] W. Walter, *Ordinary Differential Equations*, Translated from the Sixth German (1996) Edition by Russell Thompson, Springer-Verlag, New York, 1998.
- [67] Ya.B. Zel'dovich, G.I. Barenblatt, The asymptotic properties of self-modelling solutions of the nonstationary gas filtration equations, *Sov. Phys. Doklady* 3 (1989) 44–47.
- [68] Ya.B. Zel'dovich, A.S. Kompaneets, Theory of heat transfer with temperature dependent thermal conductivity, in: *Collection in Honour of the 70th Birthday of Academician A.F. Ioffe*, Izdat. Akad. Nauk. SSSR, Moscow, 1950, pp. 61–71.
- [69] S.S. Titov, V.A. Ustinov, Investigation of polynomial solutions of the two-dimensional Leibenzon filtration equation with an integer adiabatic exponent, in: A.F. Sidorov, S.V. Vershinin (Eds.), *Approximate Methods for Solving Boundary Value Problems of Continuum Mechanics*, vol. 91, Akad. Nauk SSSR, Ural. Nauchn. Tsentr, Sverdlovsk, 1985, pp. 64–70 (in Russian).