

## Two-dimensional almost-Riemannian structures with tangency points <sup>☆</sup>

A.A. Agrachev <sup>a</sup>, U. Boscain <sup>b</sup>, G. Charlot <sup>c</sup>, R. Ghezzi <sup>a</sup>, M. Sigalotti <sup>d,e,\*</sup>

<sup>a</sup> *SISSA, via Beirut 2-4, 34014 Trieste, Italy*

<sup>b</sup> *Centre de Mathématiques Appliquées, École Polytechnique Route de Saclay, 91128 Palaiseau Cedex, France*

<sup>c</sup> *Institut Fourier, UMR 5582, CNRS/Université Grenoble 1, 100 rue des Maths, BP 74, 38402 St Martin d'Hères, France*

<sup>d</sup> *INRIA Nancy – Grand Est, Équipe-projet CORIDA, France*

<sup>e</sup> *Institut Élie Cartan, UMR CNRS/INRIA/Nancy Université, BP 239, 54506 Vandœuvre-lès-Nancy, France*

Received 17 August 2009; received in revised form 22 November 2009; accepted 23 November 2009

Available online 3 December 2009

### Abstract

Two-dimensional almost-Riemannian structures are generalized Riemannian structures on surfaces for which a local orthonormal frame is given by a Lie bracket generating pair of vector fields that can become collinear. We study the relation between the topological invariants of an almost-Riemannian structure on a compact oriented surface and the rank-two vector bundle over the surface which defines the structure. We analyse the generic case including the presence of tangency points, i.e. points where two generators of the distribution and their Lie bracket are linearly dependent. The main result of the paper provides a classification of oriented almost-Riemannian structures on compact oriented surfaces in terms of the Euler number of the vector bundle corresponding to the structure. Moreover, we present a Gauss–Bonnet formula for almost-Riemannian structures with tangency points.

© 2019 L'Association Publications de l'Institut Henri Poincaré. Published by Elsevier B.V. All rights reserved.

### 1. Introduction

Let  $M$  be a two-dimensional smooth manifold. A Riemannian distance on  $M$  can be seen as the minimum-time function of an optimal control problem where admissible velocities are vectors of norm one. The control problem can be written locally as

$$\dot{q} = uX(q) + vY(q), \quad u^2 + v^2 \leq 1, \quad (1)$$

by fixing an orthonormal frame  $(X, Y)$ .

Almost-Riemannian structures generalize Riemannian ones by allowing  $X$  and  $Y$  to be collinear at some points. If the pair  $(X, Y)$  is Lie bracket generating, i.e., if

<sup>☆</sup> This work was partially supported by the ANR “GCM” program “Blanc – CSD 5”.

\* Corresponding author at: Institut Élie Cartan, UMR CNRS/INRIA/Nancy Université, BP 239, 54506 Vandœuvre-lès-Nancy, France.

*E-mail addresses:* [agrachev@sissa.it](mailto:agrachev@sissa.it) (A.A. Agrachev), [ugo.boscain@cmap.polytechnique.fr](mailto:ugo.boscain@cmap.polytechnique.fr) (U. Boscain), [Gregoire.Charlot@ujf-grenoble.fr](mailto:Gregoire.Charlot@ujf-grenoble.fr) (G. Charlot), [ghezzi@sissa.it](mailto:ghezzi@sissa.it) (R. Ghezzi), [Mario.Sigalotti@inria.fr](mailto:Mario.Sigalotti@inria.fr) (M. Sigalotti).

$$\text{span}\{X(q), Y(q), [X, Y](q), [X, [X, Y]](q), \dots\} = T_q M$$

at every  $q \in M$ , then (1) is completely controllable and the minimum-time function still defines a continuous distance  $d$  on  $M$ . Notice that a Riemannian distance can be globally defined on  $M$  by a control problem as in (1) only if the Riemannian structure admits a global orthonormal frame, implying that  $M$  is parallelizable. More in general,  $X$  and  $Y$  are parallel on a set  $\mathcal{Z} \subset M$  (called *singular locus*), which is generically a one-dimensional embedded submanifold of  $M$  (possibly disconnected).

Metric structures that can be defined *locally* by a pair of vector fields  $(X, Y)$  through (1) are called *almost-Riemannian structures*. An almost-Riemannian structure  $\mathcal{S}$  can be seen as an Euclidean bundle  $E$  of rank two over  $M$  (i.e. a vector bundle whose fibre is equipped with a smoothly-varying scalar product  $\langle \cdot, \cdot \rangle_q$ ) and a morphism of vector bundles  $f : E \rightarrow TM$  such that the evaluation at  $q$  of the Lie algebra generated by

$$\{f \circ \sigma \mid \sigma \text{ section of } E\}$$

is equal to  $T_q M$  for every  $q \in M$ , see Section 2 and also [17].

If  $E$  is orientable, we say that  $\mathcal{S}$  is *orientable*. If  $E$  is isomorphic to the trivial bundle  $M \times \mathbb{R}^2$ , we say that the almost-Riemannian structure is *trivializable*. The singular locus  $\mathcal{Z}$  is the set of points  $q$  of  $M$  at which  $f(E_q)$  is one-dimensional. An almost-Riemannian structure is Riemannian if and only if  $\mathcal{Z} = \emptyset$ , i.e.  $f$  is an isomorphism of vector bundles.

The first example of genuinely almost-Riemannian structure is provided by the Grushin plane, which is the almost-Riemannian structure on  $M = \mathbb{R}^2$  with  $E = \mathbb{R}^2 \times \mathbb{R}^2$ ,  $f((x, y), (a, b)) = ((x, y), (a, bx))$  and  $\langle \cdot, \cdot \rangle$  the canonical Euclidean structure on  $\mathbb{R}^2$ . The model was originally introduced in the context of hypoelliptic operator theory [10, 11] (see also [4, 8]). Notice that the singular locus is indeed nonempty, being equal to the  $y$ -axis. Another example of (trivializable) almost-Riemannian structure appeared in problems of control of quantum mechanical systems (see [6, 7]).

The notion of almost-Riemannian structure was introduced in [1]. In that paper, an almost-Riemannian structure is defined as a locally finitely generated Lie bracket generating  $C^\infty(M)$ -submodule  $\Delta$  of  $\text{Vec}(M)$ , the space of smooth vector fields on  $M$ , endowed with a bilinear, symmetric map  $G : \Delta \times \Delta \rightarrow C^\infty(M)$  which is positive definite (in a suitable sense). This definition is equivalent to the one given above in terms of Euclidean bundles of rank two over  $M$ . A pair of vector fields  $(X, Y)$  in  $\Delta$  is said to be an *orthonormal frame for  $G$*  on some open set  $\Omega$  if  $G(X, Y)(q) = 0$  and  $G(X, X)(q) = G(Y, Y)(q) = 1$  for every  $q \in \Omega$ . Equivalently, there exists a local orthonormal frame  $(\sigma, \rho)$  for  $\langle \cdot, \cdot \rangle$  such that  $X = f \circ \sigma$ ,  $Y = f \circ \rho$ .

Almost-Riemannian structures present very interesting phenomena. For instance, even in the case where the Gaussian curvature is everywhere negative (where it is defined, i.e., on  $M \setminus \mathcal{Z}$ ), geodesics may have conjugate points. This happens for instance on the Grushin plane (see [1] and also [5] in the case of surfaces of revolution). Moreover it is possible to define non-orientable almost-Riemannian structures on orientable manifolds and orientable almost-Riemannian structures on non-orientable manifolds (see [1]).

This paper is a continuation of [1], where we provided a characterization of generic almost-Riemannian structures by means of local normal forms, and we proved a generalization of the Gauss–Bonnet formula. (For generalizations of Gauss–Bonnet formula in related contexts, see [2, 16].) Let us briefly recall these results.

The *flag* of a submodule  $\Delta$  of  $\text{Vec}(M)$  is the sequence of submodules  $\Delta = \Delta_1 \subset \Delta_2 \subset \dots \subset \Delta_m \subset \dots$  defined through the recursive formula

$$\Delta_{k+1} = \Delta_k + [\Delta, \Delta_k].$$

Denote by  $\Delta_m(q)$  the set  $\{V(q) \mid V \in \Delta_m\}$ . Under generic assumptions, the singular locus  $\mathcal{Z}$  has the following properties: (i)  $\mathcal{Z}$  is an embedded one-dimensional submanifold of  $M$ ; (ii) the points  $q \in M$  at which  $\Delta_2(q)$  is one-dimensional are isolated; (iii)  $\Delta_3(q) = T_q M$  for every  $q \in M$ . We say that  $\mathcal{S}$  satisfies (H0) if properties (i), (ii), (iii) hold true. If this is the case, a point  $q$  of  $M$  is called *ordinary* if  $\Delta(q) = T_q M$ , *Grushin point* if  $\Delta(q)$  is one-dimensional and  $\Delta_2(q) = T_q M$ , i.e. the distribution is transversal to  $\mathcal{Z}$ , and *tangency point* if  $\Delta_2(q)$  is one-dimensional, i.e. the distribution is tangent to  $\mathcal{Z}$ . If  $(\Omega, X, Y)$  is a local generator of  $\Delta$  such that  $\Omega \setminus \mathcal{Z}$  has exactly two components, then  $(X, Y)$  has different orientations on each of them. Normal forms for local generators of  $\Delta$  around ordinary, Grushin and tangency points are recalled in Section 2.1.

The main result of [1] is an extension of the Gauss–Bonnet theorem for orientable almost-Riemannian structures on orientable manifolds, under the hypothesis that there are not tangency points. More precisely, denote by  $K : M \setminus \mathcal{Z} \rightarrow$

$\mathbb{R}$  the Gaussian curvature and by  $\omega$  a volume form for the Euclidean structure on  $E$ . Let  $dA_s$  be the two-form on  $M \setminus \mathcal{Z}$  given by the pushforward of  $\omega$  along  $f$ . Fix an orientation  $\mathcal{E}$  on  $M$  and let  $M^+$  (respectively,  $M^-$ ) be the subset of  $M \setminus \mathcal{Z}$  where  $dA_s$  is a positive (respectively, negative) multiple of  $\mathcal{E}$ . The main goal of [1] was to prove the existence and to assign a value to the limit

$$\lim_{\varepsilon \searrow 0} \int_{M_\varepsilon} K dA_s, \tag{2}$$

where  $M_\varepsilon = \{q \in M \mid d(q, \mathcal{Z}) > \varepsilon\}$  and  $d(\cdot, \cdot)$  is the distance globally defined by the almost-Riemannian structure on  $M$ . If  $M$  has no tangency point, then the limit in (2) can be shown to exist and to be equal to  $2\pi(\chi(M^+) - \chi(M^-))$ , where  $\chi$  denotes the Euler characteristic. When the almost-Riemannian structure is trivialisable, we proved that  $\chi(M^+) = \chi(M^-)$  whence the limit in (2) is equal to zero. Once applied to the special subclass of Riemannian structures, such a result simply states that the integral of the curvature of a parallelizable compact oriented surface (i.e., the torus) is equal to zero. In a sense, in the standard Riemannian construction the topology of the surface gives a constraint on the total curvature through the Gauss–Bonnet formula, whereas for an almost-Riemannian structure induced by a single pair of vector fields the total curvature is equal to zero and the topology of the manifold constrains the metric to be singular on a suitable set.

The main objective of this paper is to complete the analysis in the more complicated case in which  $S$  has tangency points.

The following result provides a classification of almost-Riemannian structures in terms of the Euler number of the vector bundle associated with it (for a definition of the Euler number see Section 2).

**Theorem 1.** *Let  $M$  be a compact oriented two-dimensional manifold endowed with an oriented almost-Riemannian structure  $S = (E, f, \langle \cdot, \cdot \rangle)$  satisfying the generic hypothesis (H0). Then  $\chi(M^+) - \chi(M^-) + \tau(S) = e(E)$ , where  $e(E)$  denotes the Euler number of  $E$  and  $\tau(S)$  is the number of revolutions of  $\Delta$  on  $\mathcal{Z}$  computed with respect to the orientation induced by  $M^+$  on  $\mathcal{Z}$ .*

For a detailed definition of  $\tau(S)$  see Section 3. Notice that the Euler number  $e(E)$  measures how far the vector bundle  $E$  is from the trivial one. Indeed,  $E$  is trivial if and only if  $e(E) = 0$ . As a direct consequence of Theorem 1 we get that  $S$  is trivialisable if and only if  $\chi(M^+) - \chi(M^-) + \tau(S) = 0$ .

For what concerns the notion of integrability of the curvature with respect to the Riemannian density on  $M \setminus \mathcal{Z}$ , it turns out that the hypothesis made in [1] about the absence of tangency points is not just technical. Indeed, in Section 5.1 we present some numerical simulations strongly hinting that the limit in (2) diverges, in general, if tangency points are present. One possible explanation of this fact is the interaction between different orders in the asymptotic expansion of the almost-Riemannian distance. In this paper, to avoid this interference, we define a 3-scale integral of the curvature. Let  $\mathcal{T}$  be the set of tangency points of  $S$ . Associate with every  $q \in \mathcal{T}$ , a two parameters “rectangular” neighborhood  $B_{\delta_1, \delta_2}^q$  ( $\delta_1$  and  $\delta_2$  playing the role of lengths of the sides of the rectangle) built as follows. We consider a smooth curve  $(-1, 1) \ni s \mapsto w(s)$  passing through the tangency point  $w(0) = q$  and transversal to the distribution at  $q$ . We then consider, for each  $s \in (-1, 1)$ , the geodesic  $\gamma_s$  (parameterized by arclength) such that  $\gamma_s(0) = w(s)$  and minimizing locally the distance from  $\{w(s) \mid s \in (-1, 1)\}$  (such geodesic exists and is unique due to the transversality of the curve  $w$ ; for details see [1]). For  $\delta_1, \delta_2$  sufficiently small, the rectangle  $B_{\delta_1, \delta_2}^q$  is the subset of  $M$  containing the tangency point  $q$  and having as boundary

$$\gamma_{\delta_2}([- \delta_1, \delta_1]) \cup \gamma_{[- \delta_2, \delta_2]}(\delta_1) \cup \gamma_{- \delta_2}([- \delta_1, \delta_1]) \cup \gamma_{[- \delta_2, \delta_2]}(- \delta_1)$$

(see Fig. 1). Let  $M_{\varepsilon, \delta_1, \delta_2} = M_\varepsilon \setminus \bigcup_{q \in \mathcal{T}} B_{\delta_1, \delta_2}^q$ . We say that  $K$  is 3-scale  $S$ -integrable if

$$\lim_{\delta_1 \rightarrow 0} \lim_{\delta_2 \rightarrow 0} \lim_{\varepsilon \rightarrow 0} \int_{M_{\varepsilon, \delta_1, \delta_2}} K dA_s \tag{3}$$

exists and is finite. In this case we denote such a limit by  $\int_M K dA_s$ . The following result, proved in Section 5.3, is a generalization of the classical Gauss–Bonnet formula for Riemannian structures to generic oriented two-dimensional almost-Riemannian structures. It concludes the analysis started in [1] including the presence of tangency points.




Fig. 1. The rectangular box  $B_{\delta_1, \delta_2}^q$ .

**Theorem 2.** *Let  $M$  be a compact oriented two-dimensional manifold. If an oriented almost-Riemannian structure  $\mathcal{S}$  on  $M$  satisfies the hypothesis (H0), then  $K$  is 3-scale  $\mathcal{S}$ -integrable and*

$$\oint_M K dA_s = 2\pi e(E).$$

Notice that the construction of  $M_{\varepsilon, \delta_1, \delta_2}$  is not intrinsic since it depends on the choice of a manifold transversal to  $\mathcal{Z}$  at each tangency point and on its parameterization. The result, however, is. An interesting question is whether a canonical way of choosing these manifolds and their parameterizations exists. This is related to the problem of finding intrinsic local normal forms for almost-Riemannian structures.

As a consequence of Theorems 1 and 2 we get the following corollary.

**Corollary 1.** *Let  $M$  be a compact oriented two-dimensional manifold. For an oriented almost-Riemannian structure  $\mathcal{S}$  on  $M$  satisfying the generic hypothesis (H0) we have  $\oint_M K dA_s = 0$  if and only if  $\mathcal{S}$  is trivializable. In particular, if  $\mathcal{S}$  has not tangency points then  $\int_M K dA_s = 0$  if and only if  $\mathcal{S}$  is trivializable.*

These results complete the analysis of the relation between the integral of the curvature and the topology of the manifold for two-dimensional almost-Riemannian structures.

The structure of the paper is the following. In Section 2 we introduce the notion of  $n$ -dimensional rank-varying sub-Riemannian structure and recall some definitions and results given in [1]. In Section 3 we define the number of revolutions of a one-dimensional distribution along an oriented closed curve. In Section 4 we prove Theorem 1. First we construct a section of the sphere bundle on a tubular neighborhood of the singular locus having each tangency point as singularity. Then, we extend the section to the complement in the manifold of a finite set and show the sum of the indexes at all the singularities to be equal to  $\chi(M^+) - \chi(M^-) + \tau(\mathcal{S})$ . In Section 5, the relation between tangency points and integrability of the curvature with respect to the area form associated with an almost-Riemannian structure is discussed. In particular, we provide in Section 5.1 some numerical simulations strongly hinting that in presence of tangency points the integral  $\int_{M_\varepsilon} K dA_s$  does not converge as  $\varepsilon$  tends to zero. This leads us to introduce in Section 5.2 the notion of 3-scale  $\mathcal{S}$ -integrability. Thanks to a Gauss–Bonnet formula for almost-Riemannian surfaces with boundary given in [9], we compute in Section 5.3 the total curvature of a generic two-dimensional almost-Riemannian structure with tangency points, proving Theorem 2.

## 2. Basic definitions

Let  $M$  be an  $n$ -dimensional manifold. Throughout the paper, unless specified, manifolds are smooth (i.e.  $C^\infty$ ) and without boundary; vector fields and differential forms are smooth. Given a vector bundle  $E$  over  $M$ , the set of smooth sections of  $E$ , denoted by  $\Gamma(E)$ , is naturally endowed with the structure of  $C^\infty(M)$ -module. In the case  $E = TM$  we denote  $\Gamma(E)$  by  $\text{Vec}(M)$ .

Given an oriented vector bundle of rank  $n$  over a compact connected oriented  $n$ -manifold  $M$ , the Euler number of  $E$ , denoted by  $e(E)$ , is the self-intersection number of  $M$  in  $E$ , where  $M$  is identified with the zero section. To compute  $e(E)$ , consider a smooth section  $\sigma : M \rightarrow E$  transverse to the zero section. Then, by definition,

$$e(E) = \sum_{p|\sigma(p)=0} i(p, \sigma),$$

where  $i(p, \sigma) = 1$ , respectively  $-1$ , if  $d_p\sigma : T_pM \rightarrow T_{\sigma(p)}E$  preserves, respectively reverses, the orientation. Notice that if we reverse the orientation on  $M$  or on  $E$  then  $e(E)$  changes sign. Hence, the Euler number of an orientable vector bundle  $E$  is defined up to a sign, depending on the orientations of both  $E$  and  $M$ . Since reversing the orientation on  $M$  also reverses the orientation of  $TM$ , the Euler number of  $TM$  is defined unambiguously and is equal to  $\chi(M)$ , the Euler characteristic of  $M$ . We refer the reader to [12] for a more detailed discussion of this subject.

**Remark 1.** Assume that  $\sigma \in \Gamma(E)$  has only isolated zeros, i.e. the set  $\{p \mid \sigma(p) = 0\}$  is finite. If  $E$  is endowed with a smooth scalar product  $\langle \cdot, \cdot \rangle$ , we define  $\tilde{\sigma} : M \setminus \{p \mid \sigma(p) = 0\} \rightarrow SE$  by  $\tilde{\sigma}(q) = \frac{\sigma(q)}{\sqrt{\langle \sigma(q), \sigma(q) \rangle}}$ . Then if  $\sigma(p) = 0$ ,  $i(p, \tilde{\sigma}) = i(p, \sigma)$  is equal to the degree of the map  $\partial B \rightarrow S^{n-1}$  that associate with each  $q \in \partial B$  the value  $\tilde{\sigma}(q)$ , where  $B$  is a neighborhood of  $p$  diffeomorphic to an open ball in  $\mathbb{R}^n$  that does not contain any other zero of  $\sigma$ .

Notice that if  $i(p, \sigma) \neq 0$ , the limit  $\lim_{q \rightarrow p} \tilde{\sigma}(q)$  does not exist and, in this case, we say that  $\tilde{\sigma}$  has a singularity at  $p$ .

**Definition 1.** An  $(n, k)$ -rank-varying distribution on an  $n$ -dimensional manifold  $M$  is a pair  $(E, f)$  where  $E$  is a vector bundle of rank  $k$  over  $M$  and  $f : E \rightarrow TM$  is a morphism of vector bundles, i.e., (i) if  $\pi : TM \rightarrow M$  and  $\pi_E : E \rightarrow M$  denote the canonical projections, then the diagram

$$\begin{array}{ccc} E & \xrightarrow{f} & TM \\ & \searrow \pi_E & \downarrow \pi \\ & & M \end{array}$$

commutes and (ii)  $f$  is linear on fibers.

Moreover, we require the map  $\sigma \mapsto f \circ \sigma$  from  $\Gamma(E)$  to  $\text{Vec}(M)$  to be injective.

**Remark 2.** Definition 1 is equivalent to the definition of  $(n, k)$ -rank-varying distribution given in [1], namely a submodule  $\Delta$  of  $\text{Vec}(M)$  locally generated by  $k$  (and not less than  $k$ ) vector fields. Given an  $(n, k)$ -rank-varying distribution  $(E, f)$ , we define  $\Delta = \{f \circ \sigma \mid \sigma \in \Gamma(E)\}$ . Thanks to (i)  $\Delta$  is a submodule of  $\text{Vec}(M)$ . The injectivity assumption on the map  $\sigma \mapsto f \circ \sigma$  implies that  $\Delta$  cannot be generated by less than  $k$  vector fields. In the following, we use either definition, depending on our convenience.

Let  $(E, f)$  be an  $(n, k)$ -rank-varying distribution,  $\Delta = \{f \circ \sigma \mid \sigma \in \Gamma(E)\}$  be its associated submodule and denote by  $\Delta(q)$  the linear subspace  $\{V(q) \mid V \in \Delta\} = f(E_q) \subseteq T_qM$ . Let  $\text{Lie}(\Delta)$  be the smallest Lie subalgebra of  $\text{Vec}(M)$  containing  $\Delta$  and  $\text{Lie}_q(\Delta) = \{V(q) \mid V \in \text{Lie}(\Delta)\}$  for every  $q \in M$ . We say that  $(E, f)$  satisfies the *Lie bracket generating condition* if  $\text{Lie}_q(\Delta) = T_qM$  for every  $q \in M$ .

A property  $(P)$  defined for  $(n, k)$ -rank-varying distributions is said to be *generic* if for every vector bundle  $E$  of rank  $k$  over  $M$ ,  $(P)$  holds for every  $f$  in an open and dense subset of the set of morphisms of vector bundles from  $E$  to  $TM$ , endowed with the  $C^\infty$ -Whitney topology. E.g., generically, an  $(n, k)$ -rank-varying distribution is Lie bracket generating, provided that  $k > 1$ .

We say that an  $(n, k)$ -rank-varying distribution  $(E, f)$  is *orientable* if  $E$  is orientable as a vector bundle. Similarly,  $(E, f)$  is *trivializable* if  $E$  is isomorphic to the trivial bundle  $M \times \mathbb{R}^k$ . In this case, by definition, we have  $e(E) = 0$ .

A rank-varying sub-Riemannian structure is defined by requiring that  $E$  is an Euclidean bundle.

**Definition 2.** An  $(n, k)$ -rank-varying sub-Riemannian structure is a triple  $\mathcal{S} = (E, f, \langle \cdot, \cdot \rangle)$  where  $(E, f)$  is a Lie bracket generating  $(n, k)$ -rank-varying distribution on a manifold  $M$  and  $\langle \cdot, \cdot \rangle_q$  is a scalar product on  $E_q$  smoothly depending on  $q$ .

**Remark 3.** Definition 2 is equivalent to Definition 4 in [1], with  $G : \Delta \times \Delta \rightarrow \mathcal{C}^\infty(M)$  defined by  $G(V, W) = \langle \sigma_V, \sigma_W \rangle$  where  $\sigma_V, \sigma_W$  are the unique sections of  $E$  satisfying  $f \circ \sigma_V = V, f \circ \sigma_W = W$ .

Definition 2 generalizes several classical structures. First of all, a Riemannian structure  $(M, g)$  is an  $(n, n)$ -rank-varying sub-Riemannian structure with  $E = TM, f = 1_{TM}$  and  $\langle \cdot, \cdot \rangle = g(\cdot, \cdot)$ . Classical sub-Riemannian structures (see [3,15]) are  $(n, k)$ -rank-varying sub-Riemannian structures such that  $E$  is a proper Euclidean subbundle of  $TM$  and  $f$  is the inclusion. Finally, we define  $n$ -dimensional almost-Riemannian structures ( $n$ -ARSs for short) as  $(n, n)$ -rank-varying sub-Riemannian structures.

Let  $\mathcal{S} = (E, f, \langle \cdot, \cdot \rangle)$  be an  $(n, k)$ -rank-varying sub-Riemannian structure. For every  $q \in M$  and every  $v \in \Delta(q)$  define

$$\mathbf{G}_q(v) = \inf\{\langle u, u \rangle_q \mid u \in E_q, f(u) = v\}.$$

If  $\sigma_1, \dots, \sigma_k$  is an orthonormal frame for  $\langle \cdot, \cdot \rangle$  on an open subset  $\Omega$  of  $M$ , an *orthonormal frame for  $G$*  on  $\Omega$  is given by  $f \circ \sigma_1, \dots, f \circ \sigma_k$ . One easily proves that orthonormal frames are systems of local generators of  $\Delta$ . Notice that  $\mathcal{S}$  is trivialisable if and only if there exists a global orthonormal frame for  $\langle \cdot, \cdot \rangle$ . Hence  $\mathcal{S}$  is trivialisable if and only if there exists a global orthonormal frame for  $G$ .

A curve  $\gamma : [0, T] \rightarrow M$  is said to be *admissible* for  $\mathcal{S}$  if it is Lipschitz continuous (with respect to the differential structure on  $M$ ) and there exists a measurable essentially bounded function

$$[0, T] \ni t \mapsto u(t) \in E_{\gamma(t)}$$

called *control function*, such that  $\dot{\gamma}(t) = f(u(t))$  for almost every  $t \in [0, T]$ . Given an admissible curve  $\gamma : [0, T] \rightarrow M$ , the *length of  $\gamma$*  is

$$\ell(\gamma) = \int_0^T \sqrt{\mathbf{G}_{\gamma(t)}(\dot{\gamma}(t))} dt.$$

The function  $\ell(\gamma)$  is invariant under reparameterization of the curve  $\gamma$ . Moreover, if an admissible curve  $\gamma$  minimizes the *energy functional*  $J(\gamma) = \int_0^T \mathbf{G}_{\gamma(t)}(\dot{\gamma}(t)) dt$  (with fixed  $T$  and fixed endpoints) then  $v = \sqrt{\mathbf{G}_{\gamma(t)}(\dot{\gamma}(t))}$  is constant and  $\gamma$  is also a minimizer of  $\ell(\cdot)$ . On the other hand a minimizer  $\gamma$  of  $\ell(\cdot)$  such that  $v$  is constant is a minimizer of  $J(\cdot)$  with  $T = \ell(\gamma)/v$ . A *geodesic* for  $\mathcal{S}$  is an admissible curve  $\gamma : [0, T] \rightarrow M$  such that for every sufficiently small interval  $[t_1, t_2] \subset [0, T]$ ,  $\gamma|_{[t_1, t_2]}$  is a minimizer of  $J(\cdot)$ . A geodesic for which  $\mathbf{G}_{\gamma(t)}(\dot{\gamma}(t))$  is (constantly) equal to one is said to be parameterized by arclength.

The distance induced by  $\mathcal{S}$  on  $M$  is defined as

$$d(q_0, q_1) = \inf\{\ell(\gamma) \mid \gamma(0) = q_0, \gamma(T) = q_1, \gamma \text{ admissible}\}. \quad (4)$$

The finiteness and the continuity of  $d(\cdot, \cdot)$  with respect to the topology of  $M$  are guaranteed by the Lie bracket generating assumption on the rank-varying sub-Riemannian structure (see [13]). The distance  $d(\cdot, \cdot)$  endows  $M$  with the structure of metric space compatible with the topology of  $M$  as smooth manifold.

### 2.1. Normal forms for generic 2-ARSs

Given a 2-ARS  $\mathcal{S}$ , its *singular locus* is the set of points  $q$  where  $\Delta(q) = f(E_q)$  has not maximal rank, that is,

$$\mathcal{Z} = \{q \in M \mid \dim(\Delta(q)) = 1\}.$$

Notice that  $\dim(\Delta(q)) > 0$  for every point  $q \in M$ , because  $\Delta$  is a bracket generating distribution.

We say that  $\mathcal{S}$  satisfies condition (H0) if the following properties hold: (i)  $\mathcal{Z}$  is an embedded one-dimensional submanifold of  $M$ ; (ii) the points  $q \in M$  at which  $\Delta_2(q)$  is one-dimensional are isolated; (iii)  $\Delta_3(q) = T_q M$  for every  $q \in M$ , where  $\Delta_1 = \Delta$  and  $\Delta_{k+1} = \Delta_k + [\Delta, \Delta_k]$ .

**Proposition 1.** (See [1].) *Property (H0) is generic for 2-ARSSs.*

ARSSs satisfying hypothesis (H0) admit the following local normal forms.

**Theorem 3.** (See [1].) *Given a 2-ARS  $\mathcal{S}$  satisfying (H0), for every point  $q \in M$  there exist a neighborhood  $U$  of  $q$  and an orthonormal frame  $(X, Y)$  for  $G$  on  $U$ , such that up to a smooth change of coordinates defined on  $U$ ,  $q = (0, 0)$  and  $(X, Y)$  has one of the forms*

$$(F1) \quad X(x, y) = (1, 0), \quad Y(x, y) = (0, e^{\phi(x,y)}),$$

$$(F2) \quad X(x, y) = (1, 0), \quad Y(x, y) = (0, xe^{\phi(x,y)}),$$

$$(F3) \quad X(x, y) = (1, 0), \quad Y(x, y) = (0, (y - x^2\psi(x))e^{\xi(x,y)}),$$

where  $\phi, \xi$  and  $\psi$  are smooth real-valued functions such that  $\phi(0, y) = 0$  and  $\psi(0) > 0$ .

Let  $\mathcal{S}$  be a 2-ARS satisfying (H0). A point  $q \in M$  is said to be an *ordinary point* if  $\Delta(q) = T_q M$ , hence, if  $\mathcal{S}$  is locally described by (F1). We call  $q$  a *Grushin point* if  $\Delta(q)$  is one-dimensional and  $\Delta_2(q) = T_q M$ , i.e. if the local description (F2) applies. Finally, if  $\Delta(q) = \Delta_2(q)$  has dimension one and  $\Delta_3(q) = T_q M$  then we say that  $q$  is a *tangency point* and  $\mathcal{S}$  can be described near  $q$  by the normal form (F3). We define

$$\mathcal{T} = \{q \in \mathcal{Z} \mid q \text{ tangency point of } \mathcal{S}\}.$$

### 2.2. A Gauss–Bonnet formula for 2-ARSSs

Let  $M$  be an orientable two-dimensional manifold and  $\mathcal{S} = (E, f, \langle \cdot, \cdot \rangle)$  an orientable 2-ARS on  $M$ . Notice that  $\langle \cdot, \cdot \rangle$  defines a Riemannian structure on  $M \setminus \mathcal{Z}$ . Denote by  $K$  the Gaussian curvature of such a structure and by  $\omega$  a volume form for the Euclidean structure on  $E$ . Let  $dA_{\mathcal{S}}$  be the two-form on  $M \setminus \mathcal{Z}$  given by the pushforward of  $\omega$  along  $f$ . Once an orientation on  $M$  is fixed,  $M \setminus \mathcal{Z}$  is split into two open sets  $M^+$  and  $M^-$  such that the orientation on  $M$  coincides with the one defined by  $dA_{\mathcal{S}}$  on  $M^+$  and with its opposite on  $M^-$ .

For every  $\varepsilon > 0$  let  $M_{\varepsilon} = \{q \in M \mid d(q, \mathcal{Z}) > \varepsilon\}$ , where  $d(\cdot, \cdot)$  is the almost-Riemannian distance (see Eq. (4)). We say that  $K$  is  $\mathcal{S}$ -integrable if

$$\lim_{\varepsilon \rightarrow 0} \int_{M_{\varepsilon}} K dA_{\mathcal{S}}$$

exists and is finite. In this case we denote such limit by  $\int_M K dA_{\mathcal{S}}$ .

When  $\mathcal{S}$  has no tangency points  $K$  happens to be  $\mathcal{S}$ -integrable and  $\int_M K dA_{\mathcal{S}}$  is determined by the topology of  $M^+$  and  $M^-$ . This result, recalled in Theorem 4, can be seen as a generalization of Gauss–Bonnet formula to ARSSs.

**Theorem 4.** (See [1].) *Let  $M$  be a compact oriented two-dimensional manifold, endowed with an oriented 2-ARS  $\mathcal{S}$  for which condition (H0) holds true. Assume that  $\mathcal{S}$  has no tangency points. Then  $K$  is  $\mathcal{S}$ -integrable and  $\int_M K dA_{\mathcal{S}} = 2\pi(\chi(M^+) - \chi(M^-))$ .*

### 3. Number of revolutions of $\Delta$

Let  $M$  be a compact oriented two-dimensional manifold and  $W$  an oriented closed simple curve in  $M$ . Since  $M$  is oriented,  $TM|_W$  is isomorphic to the trivial bundle  $W \times \mathbb{R}^2$  and its projectivization is isomorphic to  $W \times S^1$ . Every subbundle  $\mathcal{Y}$  of  $TM|_W$  of rank one can be seen as a section of the projectivization of  $TM|_W$ , i.e. a smooth map  $\mathcal{Y} : W \rightarrow W \times S^1$  such that  $\pi_1 \circ \mathcal{Y} = \text{Id}_W$ , where  $\pi_1 : W \times S^1 \rightarrow W$  denotes the projection on the first component. We define  $\tau(\mathcal{Y}, W)$ , the *number of revolutions* of  $\mathcal{Y}$  along  $W$ , to be the degree of the map  $\pi_2 \circ \mathcal{Y} : W \rightarrow S^1$ , where




Fig. 2. Tangency points with opposite contributions.

$\pi_2 : W \times S^1 \rightarrow S^1$  is the projection on the second component. Notice that  $\tau(\mathcal{Y}, W)$  changes sign if we reverse the orientation of  $W$ .

**Remark 4.** Let  $M, W, \mathcal{Y}$  be as above and let us show how to compute  $\tau(\mathcal{Y}, W)$ . Let  $V \in \Gamma(TW)$  be a never-vanishing vector field along  $W$ . Then  $\text{span}(V)$  is a subbundle of  $TM|_W$  of rank one and there exists an isomorphism  $t : TM|_W \rightarrow W \times \mathbb{R}^2$  such that  $t \circ V(q) = (q, (1, 0))$ . The trivialization  $t$  induces an isomorphism, still denoted by  $t$ , between the projectivization of  $TM|_W$  and  $W \times S^1$  such that  $\pi_2 \circ t \circ V : W \rightarrow S^1$  is constant. To simplify notations, we omit  $t$  in the following. Let  $\pi_2 \circ V(q) \equiv \theta_0$ . Assume that  $\theta_0$  is a regular value of  $\pi_2 \circ \mathcal{Y}$  (otherwise there exists a smooth section  $\tilde{\mathcal{Y}}$  homotopic to  $\mathcal{Y}$  having  $\theta_0$  as a regular value). By definition,

$$\tau(\mathcal{Y}, W) = \sum_{q|\pi_2 \circ \mathcal{Y}(q) = \theta_0} \text{sign}(d_q(\pi_2 \circ \mathcal{Y})),$$

where  $d_q$  denotes the differential at  $q$  of a smooth map. Notice that a point  $q$  satisfies  $\pi_2 \circ \mathcal{Y}(q) = \theta_0$  if and only if  $\mathcal{Y}(q)$  is tangent to  $W$  at  $q$ .

Let  $M$  be a compact oriented two-dimensional manifold and  $\mathcal{S}$  be an oriented 2-ARS satisfying (H0). Then the singular locus  $\mathcal{Z}$  is the boundary of  $M^+$ . Fix on  $\mathcal{Z}$  the orientation induced by  $M^+$  and let  $C(\mathcal{Z})$  denote the set of connected component of  $\mathcal{Z}$ . Since  $\Delta$  is one-dimensional along  $\mathcal{Z}$ , we can define  $\tau(\mathcal{S}) = \sum_{W \in C(\mathcal{Z})} \tau(\Delta, W)$ .

**Remark 5.** Let  $\mathcal{S}$  be a 2-ARS satisfying hypothesis (H0). Let  $V, \theta_0, \pi_2$  be as in Remark 4 with  $\mathcal{Y} = \Delta$  and  $W \in C(\mathcal{Z})$ . Since  $\Delta_3(q) = T_q M$  for every point  $q \in M$ ,  $\theta_0$  is a regular value of  $\pi_2 \circ \Delta$ . Moreover, the set of points  $q \in W$  such that  $\pi_2 \circ \Delta(q) = \theta_0$  is the set of tangency points of  $\mathcal{S}$  belonging to  $W$ . Hence,

$$\tau(\Delta, W) = \sum_{q \in W \cap \mathcal{T}} \text{sign}(d_q(\pi_2 \circ \Delta)).$$

We define  $\tau_q = \text{sign}(d_q \pi_2 \circ \Delta)$  (see Fig. 2). Clearly,  $\tau(\mathcal{S}) = \sum_{q \in \mathcal{T}} \tau_q$ .

#### 4. Proof of Theorem 1

The idea of the proof is to find a section  $\sigma$  of  $SE$  with isolated singularities  $p_1, \dots, p_m$  such that  $\sum_{j=1}^m i(p_j, \sigma) = \chi(M^+) - \chi(M^-) + \tau(\mathcal{S})$ . In the sequel, we consider  $\mathcal{Z}$  to be oriented with the orientation induced by  $M^+$ .

We start by defining  $\sigma$  on a neighborhood of  $\mathcal{Z}$ . Let  $W$  be a connected component of  $\mathcal{Z}$ . Since  $M$  is oriented, there exists an open tubular neighborhood  $\mathbf{W}$  of  $W$  and a diffeomorphism  $\Psi : S^1 \times (-1, 1) \rightarrow \mathbf{W}$  that preserves the orientation and  $\Psi|_{S^1 \times \{0\}}$  is an orientation-preserving diffeomorphism between  $S^1$  and  $W$ . Remark that  $f : E|_{\mathbf{W}^+} \rightarrow T\mathbf{W}^+$  is an orientation-preserving isomorphism of vector bundles, while  $f : E|_{\mathbf{W}^-} \rightarrow T\mathbf{W}^-$  is an orientation-reversing isomorphism of vector bundles, where  $\mathbf{W}^\pm = \mathbf{W} \cap M^\pm$ . For every  $s \neq 0$ , lift the tangent vector to  $\theta \mapsto \Psi(\theta, s)$  to  $E$  using  $f^{-1}$ , rotate it by the angle  $\pi/2$  and normalize it:  $\sigma$  is defined as this unit vector (belonging to  $E_{\Psi(\theta, s)}$ ) if  $s > 0$ , its opposite if  $s < 0$ . In other words,  $\sigma : \mathbf{W} \setminus W \rightarrow SE$  is given by



$$\sigma(q) = \text{sign}(s) \frac{R_{\pi/2} f^{-1}(\frac{\partial \Psi}{\partial \theta}(\theta, s))}{\sqrt{\langle f^{-1}(\frac{\partial \Psi}{\partial \theta}(\theta, s)), f^{-1}(\frac{\partial \Psi}{\partial \theta}(\theta, s)) \rangle}}, \quad (\theta, s) = \Psi^{-1}(q), \tag{5}$$

where  $R_{\pi/2}$  denotes the rotation (with respect to the Euclidean structure) in  $E$  by angle  $\pi/2$  in the counterclockwise sense. The following lemma shows that  $\sigma$  can be extended to a continuous section from  $W \setminus T$  to  $SE$ .

**Lemma 1.**  $\sigma$  can be continuously extended to every point  $q \in W \setminus T$ .

**Proof.** Let  $q \in W \setminus T$ ,  $U$  be a neighborhood of  $q$  in  $M$  and  $(x, y)$  be a system of coordinates on  $U$  centered at  $q$  such the almost-Riemannian structure has the form (F2) (see Theorem 3). Assume, moreover, that  $U$  is a trivializing neighborhood of both  $E$  and  $TM$  and the pair of vector fields  $(X, Y)$  is the image under  $f$  of a positively-oriented local orthonormal frame of  $E$ . Then  $W \cap U = \{(x, y) \mid x = 0\}$ . Since  $\frac{\partial \Psi}{\partial \theta}(\theta, 0)$  is non-zero and tangent to  $W$ ,  $\frac{\partial \Psi}{\partial \theta}(\theta, 0)$  is tangent to the  $y$ -axis. Hence, thanks to the Preparation Theorem [14], there exist  $h_2 : \mathbb{R} \rightarrow \mathbb{R}$ ,  $h_1, h_3 : \mathbb{R}^2 \rightarrow \mathbb{R}$  smooth functions such that  $h_2(y) \neq 0$  for every  $y \in \mathbb{R}$  and for  $\Psi(\theta, s) \in U$

$$\frac{\partial \Psi}{\partial \theta}(\theta, s) = (xh_1(x, y), h_2(y) + xh_3(x, y)),$$

where  $(x, y)$  are the coordinates of the point  $\Psi(\theta, s)$ . Let us compute  $\sigma$  at a point  $p \in (W \cap U) \setminus W$ . Since

$$\frac{\partial \Psi}{\partial \theta}(\theta, s) = xh_1(x, y)X(x, y) + \frac{h_2(y) + xh_3(x, y)}{xe^{\phi(x, y)}}Y(x, y),$$

then

$$f^{-1}\left(\frac{\partial \Psi}{\partial \theta}(\theta, s)\right) = xh_1(x, y)\zeta(x, y) + \frac{h_2(y) + xh_3(x, y)}{xe^{\phi(x, y)}}\rho(x, y),$$

where  $(\zeta, \rho)$  is the unique local orthonormal basis of  $E|_U$  such that  $f \circ \zeta = X$  and  $f \circ \rho = Y$ . Notice that  $U \cap M^+ = \{(x, y) \mid x > 0\}$  and  $U \cap M^- = \{(x, y) \mid x < 0\}$ . Using formula (5), for  $(x, y) = \Psi(\theta, s) \in U \setminus W$  one easily gets

$$\sigma(x, y) = \frac{\text{sign}(x)}{l(x, y)} \left( -\frac{h_2(y) + xh_3(x, y)}{xe^{\phi(x, y)}}\zeta + xh_1(x, y)\rho \right),$$

where

$$l(x, y) = \sqrt{x^2 h_1(x, y)^2 + \frac{(h_2(y) + xh_3(x, y))^2}{x^2 e^{2\phi(x, y)}}}.$$

Since

$$\lim_{x \rightarrow 0} \frac{\text{sign}(x)(h_2(y) + xh_3(x, y))}{l(x, y)xe^{\phi(x, y)}} = \frac{h_2(y)}{|h_2(y)|} \quad \text{and} \quad \lim_{x \rightarrow 0} \frac{\text{sign}(x)xh_1(x, y)}{l(x, y)} = 0,$$

$\sigma$  can be continuously extended to the set  $\{x = 0\} = W \cap U$ .  $\square$

The next step of the proof is to show that for every  $q \in W \cap T$ ,  $i(q, \sigma) = \tau_q$ .

**Lemma 2.** Let  $\sigma : W \setminus T \rightarrow SE$  be the continuous section obtained in Lemma 1. Then, for every  $q \in W \cap T$  the index of  $\sigma$  at  $q$  is equal to  $\tau_q$  and, consequently,

$$\sum_{q \in T \cap W} i(q, \sigma) = \tau(\Delta, W). \tag{6}$$

**Proof.** Let  $q \in W \cap T$ ,  $U$  be a neighborhood of  $q$  in  $M$  and  $(x, y)$  be a system of coordinates on  $U$  centered at  $q$  such the almost-Riemannian structure has the form (F3) (see Theorem 3), i.e. a local orthonormal frame  $(X, Y)$  is given by

$$X = (1, 0), \quad Y = (0, (y - x^2\psi(x))e^{\xi(x, y)}).$$

Define  $\alpha = 1$ , respectively  $-1$ , if  $(X, Y)$  is the image under  $f$  of a positively-oriented, respectively negatively-oriented, local orthonormal frame of  $E$ . One can check that  $\tau_q = -\alpha$ . Let us make the following change of coordinates

$$\tilde{x} = x, \quad \tilde{y} = \alpha(y - x^2\psi(x)).$$

In these new coordinates,  $X$  and  $Y$  become

$$X = (1, -\alpha(2\tilde{x}\psi(\tilde{x}) + \tilde{x}^2\psi'(\tilde{x}))), \quad Y = (0, \tilde{y}e^{\xi(\tilde{x}, \alpha\tilde{y} + \tilde{x}^2\psi(\tilde{x}))})$$

and  $W \cap U$  is the  $\tilde{x}$ -axis. In the following, to simplify notations, we omit the tildes and denote the function  $\xi(\tilde{x}, \alpha\tilde{y} + \tilde{x}^2\psi(\tilde{x}))$  by  $\xi(x, y)$ . Since  $\frac{\partial\psi}{\partial\theta}(\theta, 0)$  is tangent to  $W$ , by the Preparation Theorem [14] there exist  $h_1 : \mathbb{R} \rightarrow \mathbb{R}, h_2, h_3 : \mathbb{R}^2 \rightarrow \mathbb{R}$  smooth functions such that  $h_1(x) \neq 0$  for every  $x \in \mathbb{R}$  and for  $\Psi(\theta, s) \in U$

$$\frac{\partial\psi}{\partial\theta}(\theta, s) = (h_1(x) + yh_2(x, y), yh_3(x, y)),$$

where  $(x, y)$  are the coordinates of the point  $\Psi(\theta, s)$ . This implies that

$$\begin{aligned} \frac{\partial\psi}{\partial\theta}(\theta, s) &= (h_1(x) + yh_2(x, y))X(x, y) \\ &\quad + \frac{yh_3(x, y) + \alpha(h_1(x) + yh_2(x, y))(2x\psi(x) + x^2\psi'(x))}{ye^{\xi(x, y)}}Y(x, y). \end{aligned}$$

Let  $(\zeta, \rho)$  be the local orthonormal frame of  $E$  such that  $X = f \circ \zeta$  and  $Y = f \circ \rho$ . From Eq. (5), it follows that

$$\begin{aligned} \sigma(x, y) &= -\alpha \frac{\text{sign}(y)}{l(x, y)} \frac{yh_3(x, y) + \alpha(h_1(x) + yh_2(x, y))(2x\psi(x) + x^2\psi'(x))}{ye^{\xi(x, y)}}\zeta \\ &\quad + \alpha \frac{\text{sign}(y)}{l(x, y)}(h_1(x) + yh_2(x, y))\rho, \end{aligned}$$

where

$$l(x, y) = \sqrt{(h_1(x) + yh_2(x, y))^2 + \left(\frac{yh_3(x, y) + \alpha(h_1(x) + yh_2(x, y))(2x\psi(x) + x^2\psi'(x))}{ye^{\xi(x, y)}}\right)^2}.$$

Notice that for  $x = 0, y \neq 0$  we have

$$\sigma(0, y) = \frac{\alpha \text{sign}(y)}{\sqrt{(h_1(0) + yh_2(0, y))^2 + h_3(0, y)^2 e^{-2\xi(0, y)}}} (-e^{-\xi(0, y)}h_3(0, y)\zeta + (h_1(0) + yh_2(0, y))\rho),$$

whence the limit of  $\sigma$  as  $(x, y)$  tends to  $(0, 0)$  does not exist. Let us compute the index of  $\sigma$  at  $q = (0, 0)$ . Using Taylor expansions of the components of  $\sigma$  in the basis  $(\zeta, \rho)$  we find

$$\sigma(x, y) = \alpha \frac{\text{sign}(y)}{l(x, y)} \left( -\left(\frac{yh_3(0, 0) + 2\alpha x h_1(0)\psi(0)}{ye^{\xi(0, 0)}} + O(\sqrt{x^2 + y^2})\right)\zeta + (h_1(0) + O(\sqrt{x^2 + y^2}))\rho \right). \tag{7}$$

Take a circle  $t \mapsto (r \cos t, r \sin t)$  of radius  $r$  centered at  $(0, 0)$  and assume  $r$  so small that  $(0, 0)$  is the unique singularity of  $\sigma$  on the closed disk of radius  $r$ . By definition,  $i((0, 0), \sigma)$  is half the degree of the map from the circle  $S^1$  to  $\mathbb{R}/\pi\mathbb{Z}$  that associates to each point the angle between  $\text{span}(\sigma)$  and the  $\zeta$ . Using (7), this angle is

$$a(x, y) = -\arctan \left( \frac{yh_1(0)e^{\xi(0, 0)}}{yh_3(0, 0) + 2\alpha x\psi(0)h_1(0)} + O(\sqrt{x^2 + y^2}) \right).$$

Computing  $a$  along the curve  $x(t) = r \cos t, y(t) = r \sin t$ , we find

$$a(r \cos t, r \sin t) = -\arctan \left( \frac{\sin(t)h_1(0)e^{\xi(0, 0)}}{\sin(t)h_3(0, 0) + 2\alpha \cos(t)\psi(0)h_1(0)} + O(r) \right).$$

Hence, by letting  $r$  go to zero, we are left to compute the degree of the map  $\tilde{a} : [0, 2\pi) \rightarrow [0, \pi)$  where

$$\tilde{a}(t) = -\arctan \left( \frac{\sin(t)h_1(0)e^{\xi(0, 0)}}{\sin(t)h_3(0, 0) + 2\alpha \cos(t)\psi(0)h_1(0)} \right).$$

Since zero is a regular value of  $\tilde{a}$ , the degree of  $\tilde{a}$  is

$$\sum_{t \in [0, 2\pi] \setminus \{\tilde{a}(t)=0\}} \text{sign}(\tilde{a}'(t)) = \text{sign}(\tilde{a}'(0)) + \text{sign}(\tilde{a}'(\pi)) = -2\alpha,$$

where the last equality follows from  $\tilde{a}'(0) = \tilde{a}'(\pi) = -\alpha e^{\xi(0,0)} / (2\psi(0))$ . Hence,  $i(q, \sigma) = -\alpha$ . Since  $\tau_q = -\alpha$ , the lemma is proved (see Remark 5).  $\square$

Let  $C(\mathcal{Z})$  denote the set of connected components of  $\mathcal{Z}$ . Let  $\tilde{\mathcal{Z}} = \coprod_{W \in C(\mathcal{Z})} S^1$  and consider an orientation-preserving diffeomorphism  $\Psi : \tilde{\mathcal{Z}} \times (-1, 1) \rightarrow \coprod_{W \in C(\mathcal{Z})} \mathbf{W}$  such that  $\Psi|_{\tilde{\mathcal{Z}} \times \{0\}}$  is an orientation-preserving diffeomorphism onto  $\mathcal{Z}$ . Applying Lemma 1 to every  $W \in C(\mathcal{Z})$  and reducing, if necessary, the cylinders  $\mathbf{W}$ , we can assume that the set of singularities of  $\sigma$  on  $\mathbf{U} = \coprod_{W \in C(\mathcal{Z})} \mathbf{W}$  is  $\mathcal{T}$ . Then  $\sigma : \mathbf{U} \setminus \mathcal{T} \rightarrow SE$  is continuous. Moreover, by Eq. (6),

$$\sum_{q \in \mathcal{T}} i(q, \sigma) = \tau(S).$$

Extend  $\sigma$  to  $M \setminus \mathbf{U}$ . By a transversality argument, we can assume that the extended section has only isolated singularities  $\{p_1, \dots, p_k\} \in M \setminus \mathcal{Z}$ . Since

$$e(E) = \sum_{j=1}^k i(p_j, \sigma) + \sum_{q \in \mathcal{T}} i(q, \sigma) = \sum_{j=1}^k i(p_j, \sigma) + \tau(S),$$

we are left to prove that

$$\sum_{j=1}^k i(p_j, \sigma) = \chi(M^+) - \chi(M^-). \tag{8}$$

To this aim, consider the vector field  $F = f \circ \sigma$ .  $F$  satisfies  $G(F, F) \equiv 1$ , where  $G(\cdot, \cdot)$  is defined as in Remark 3 and the set of singularities of  $F|_{M \setminus \mathcal{Z}}$  is exactly  $\{p_1, \dots, p_k\}$ . Let us compute the index of  $F$  at a singularity  $p \in \{p_1, \dots, p_k\}$ . Since  $f : E|_{M^+} \rightarrow TM^+$  preserves the orientation and  $f : E|_{M^-} \rightarrow TM^-$  reverses the orientation, it follows that  $i(p, F) = \pm i(p, \sigma)$ , if  $p \in M^\pm$ . Therefore,

$$\sum_{j=1}^k i(p_j, \sigma) = \sum_{j|p_j \in M^+} i(p_j, F) - \sum_{j|p_j \in M^-} i(p_j, F). \tag{9}$$

The theorem is proved if we show that

$$\sum_{j|p_j \in M^+} i(p_j, F) = \chi(M^+), \quad \sum_{j|p_j \in M^-} i(p_j, F) = \chi(M^-). \tag{10}$$

To deduce Eq. (10), define  $N^+ = M^+ \setminus \Psi(\tilde{\mathcal{Z}} \times (0, 1/2))$ . Notice that, by construction,  $\sigma|_{\Psi(\tilde{\mathcal{Z}} \times \{1/2\})}$  is non-singular, hence the same is true for  $F|_{\Psi(\tilde{\mathcal{Z}} \times \{1/2\})}$ . Moreover, the almost-Riemannian angle between  $T_q(\partial N^+)$  and  $\text{span}(F(q))$  is constantly equal to  $\pi/2$ . Hence  $F|_{\partial N^+}$  points towards  $N^+$  and applying the Hopf's Index Formula to every connected component of  $N^+$  we conclude that

$$\sum_{j|p_j \in M^+} i(p_j, F) = \sum_{j|p_j \in N^+} i(p_j, F) = \chi(N^+) = \chi(M^+).$$

Similarly, we find

$$\sum_{j|p_j \in M^-} i(p_j, F) = \chi(M^-).$$




Fig. 3. Graph of  $K$  for  $\Delta = \text{span}((1, 0), (0, y - x^2))$ .

## 5. $\mathcal{S}$ -integrability in presence of tangency points

### 5.1. Numerical simulations

In this section we provide some numerical simulations hinting that, when  $T \neq \emptyset$ ,

$$\int_{M_\varepsilon} K dA_s$$

does not converge, in general, as  $\varepsilon$  tends to zero.

From the proof of Theorem 4 we know that far from tangency points the integral of the geodesic curvature along  $\partial M_\varepsilon^+$  and  $\partial M_\varepsilon^-$  offset each other for  $\varepsilon$  going to zero. Hence, to understand whether the presence of a tangency point may lead to non- $\mathcal{S}$ -integrability of  $K$  it is sufficient to compute the geodesic curvature of  $\partial M_\varepsilon^+$  and  $\partial M_\varepsilon^-$  in a neighborhood of such a point. More precisely consider the almost-Riemannian structure  $(E, f, \langle \cdot, \cdot \rangle)$  on  $M = \mathbb{R}^2$  for which  $E = \mathbb{R}^2 \times \mathbb{R}^2$ ,  $f((x, y), (a, b)) = ((x, y), (a, b(y - x^2)))$  and  $\langle \cdot, \cdot \rangle$  is the canonical scalar product. For this system one has

$$K = \frac{-2(3x^2 + y)}{(x^2 - y)^2}.$$

The graph of  $K$  is illustrated in Fig. 3. Notice that  $\limsup_{q \rightarrow (0,0)} K(q) = +\infty$  and  $\liminf_{q \rightarrow (0,0)} K(q) = -\infty$ . This situation is different from the Grushin case where  $K(q)$  diverges to  $-\infty$  as  $q$  approaches  $\mathcal{Z}$ .

For every  $\varepsilon > 0$ , the sets  $\partial M_\varepsilon^+$  and  $\partial M_\varepsilon^-$  are smooth manifolds except at their intersections with the vertical axis  $x = 0$ , which is the cut locus for the problem of minimizing the distance from  $\mathcal{Z} = \{(x, x^2) \mid x \in \mathbb{R}\}$ . Fix  $0 < a < 1$  and consider the two geodesics starting from the point  $(a, a^2)$  and minimizing (locally) the distance from  $\mathcal{Z}$ . Let  $P^+$  and  $P^-$  be the two points along these geodesics at distance  $\varepsilon$  from  $\mathcal{Z}$ . Denote by  $\gamma^+$  and  $\gamma^-$  the portions of  $\partial M_\varepsilon^+$  and  $\partial M_\varepsilon^-$  connecting the vertical axis to the points  $P^+$  and  $P^-$ , oriented as in Fig. 4. It is easy to approximate numerically  $\gamma^+$  and  $\gamma^-$  by broken lines, but the evaluation of the integral of their geodesic curvatures is very unstable since its computation involves the second derivative of the curve parameterized by arclength. To avoid this problem, we rather apply the Riemannian Gauss–Bonnet formula on the regions  $\Omega^+$  and  $\Omega^-$  introduced in Fig. 4. This works better since the integral of the Gaussian curvature on  $\Omega^+$  and  $\Omega^-$  is numerically stable, and the integral of the geodesic curvature on horizontal and vertical segments can be computed analytically (in particular it is always zero on horizontal segments). Fig. 5 shows the value of




Fig. 4. Regions  $\Omega^\pm$  where to apply Riemannian Gauss–Bonnet formula.




Fig. 5. Divergence of the  $\mathcal{S}$ -integral of  $K$ .

$$\varepsilon \left( \int_{\gamma^+} K_g ds - \int_{\gamma^-} K_g ds \right)$$

for  $a = 0.1$  and  $\varepsilon$  varying in the interval  $[0.01, 0.04]$ . The graph seems to converge as  $\varepsilon$  tends to zero to a non-zero constant, strongly hinting at the divergence of  $\int_{M_\varepsilon} K dA_s$ .

### 5.2. More general notion of $\mathcal{S}$ -integrability

The simulations of the previous section lead us to introduce the following alternative notion of integrability.

**Definition 3** (3-scale  $\mathcal{S}$ -integrability). Let  $q \in \mathcal{T}$  and  $U^q$  be a neighborhood of  $q$  such that an orthonormal frame for  $G$  on  $U^q$  is given by the normal form (F3). For  $\delta_1, \delta_2 > 0$  sufficiently small the rectangle  $[-\delta_1, \delta_1] \times [-\delta_2, \delta_2]$  is a subset of  $U^q$  denoted by  $B_{\delta_1, \delta_2}^q$ . For every  $\varepsilon > 0$ , define

$$M_{\varepsilon, \delta_1, \delta_2} = M_\varepsilon \setminus \bigcup_{q \in \mathcal{T}} B_{\delta_1, \delta_2}^q. \tag{11}$$

We say that  $K$  is 3-scale  $\mathcal{S}$ -integrable if

$$\lim_{\delta_1 \rightarrow 0} \lim_{\delta_2 \rightarrow 0} \lim_{\varepsilon \rightarrow 0} \int_{M_{\varepsilon, \delta_1, \delta_2}} K dA_s \tag{12}$$

exists, is finite and does not depend on the choice of the normal form. In this case we denote such limit by  $\oint_M K dA_s$ .

**Remark 6.** Notice that if  $\mathcal{T} = \emptyset$ , then the concepts of  $\mathcal{S}$ -integrability and 3-scale  $\mathcal{S}$ -integrability coincide.

The order in which the limits are taken in (12) is important. Indeed, if the order is permuted, then the result given in Theorem 2 does not hold anymore.

Recall that the normal form (F3) is not totally intrinsic, since the functions  $\psi$  and  $\phi$  depend on the choice of a parameterized smooth curve passing through the tangency point and transversal to the distribution at the point.

### 5.3. Proof of Theorem 2

Let us recall the following Gauss–Bonnet-like formula for domains whose boundary is  $\mathcal{C}^2$  in a neighborhood of  $\mathcal{Z}$ .

**Theorem 5.** (See [9, Theorem 5.2].) Let  $U$  be an open bounded connected subset of  $M$  such that

- (i)  $\bar{U}$  contains only ordinary and Grushin points,
- (ii)  $\partial U$  is piecewise  $\mathcal{C}^2$ ,
- (iii)  $\partial U$  is  $\mathcal{C}^2$  in a neighborhood of  $\mathcal{Z}$ ,
- (iv)  $\partial U$  is the union of the supports of a finite set of curves  $\gamma^1, \dots, \gamma^m$  that are admissible for  $\Delta$  and of finite length.

Define  $U_\varepsilon^\pm = M_\varepsilon^\pm \cap U$ . Then the following limits exist and are finite

$$\int_U K dA_s := \lim_{\varepsilon \rightarrow 0} \int_{U_\varepsilon^+ \cup U_\varepsilon^-} K dA_s, \tag{13}$$

$$\int_{\partial U} k_g d\sigma_s := \lim_{\varepsilon \rightarrow 0} \left( \int_{\partial U \cap \partial U_\varepsilon^+} k_g d\sigma - \int_{\partial U \cap \partial U_\varepsilon^-} k_g d\sigma \right), \tag{14}$$

where we interpret each integral  $\int_{\partial U \cap \partial U_\varepsilon^\pm} k_g d\sigma$  as the sum of the integrals along the  $\mathcal{C}^2$  portions of  $\partial U \cap \partial U_\varepsilon^\pm$ , plus the sum of the angles at the points of  $\partial U \cap \partial U_\varepsilon^\pm$  where  $\partial U$  is not  $\mathcal{C}^1$ . Moreover, we have

$$\int_U K dA_s + \int_{\partial U} k_g d\sigma_s = 2\pi (\chi(U^+) - \chi(U^-)). \tag{15}$$

Fix  $\delta_1$  and  $\delta_2$  in such a way that the rectangles  $B_{\delta_1, \delta_2}^q$  are pairwise disjoint and  $\mathcal{Z} \cap \partial B_{\delta_1, \delta_2}^q \subset [-\delta_1, \delta_1] \times \{\delta_2\}$ , for every  $q \in \mathcal{T}$ . By construction,  $\partial B_{\delta_1, \delta_2}^q$  is the union of the supports of four admissible curves for every  $q \in \mathcal{T}$ . Hence we can take  $M_\varepsilon \setminus \bigcup_{q \in \mathcal{T}} B_{\delta_1, \delta_2}^q$  as  $U$  in Theorem 5. As a consequence, we have

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \int_{M_{\varepsilon, \delta_1, \delta_2}} K dA_s + \sum_{q \in \mathcal{T}} \int_{\partial B_{\delta_1, \delta_2}^q} k_g d\sigma_s &= 2\pi \left( \chi \left( M^+ \setminus \bigcup_{q \in \mathcal{T}} B_{\delta_1, \delta_2}^q \right) - \chi \left( M^- \setminus \bigcup_{q \in \mathcal{T}} B_{\delta_1, \delta_2}^q \right) \right) \\ &= 2\pi (\chi(M^+) - \chi(M^-)) = 2\pi (e(E) - \tau(S)), \end{aligned}$$

where the last equality follows from Theorem 1. We are left to prove that, for a fixed  $q \in \mathcal{T}$ ,

$$\lim_{\delta_1 \rightarrow 0} \lim_{\delta_2 \rightarrow 0} \int_{\partial B_{\delta_1, \delta_2}^q} k_g d\sigma_s = -2\pi \tau_q$$

(see Remark 5). In order to prove it, let us work with the normal form (F3) and assume that  $M^+ \cap U^q$  is the set  $\{y - x^2 \psi(x) < 0\} \cap U^q$ , the proof for the opposite situation being analogous. On one hand, one can check that  $\tau_q = 1$ . On the other hand, the geodesic curvature along  $[-\delta_1, \delta_1] \times \{\delta_2\}$  and along  $[-\delta_1, \delta_1] \times \{-\delta_2\}$  is zero, the two segments being the support of geodesics. Hence

$$\int_{\partial B_{\delta_1, \delta_2}^q} k_g d\sigma_s = \int_{\{\delta_1\} \times [-\delta_2, \delta_2]} k_g d\sigma_s + \int_{\{-\delta_1\} \times [-\delta_2, \delta_2]} k_g d\sigma_s + \sum_{j=1}^4 \alpha_j,$$

where the last term is the sum of the values of the angles of the box and is equal to  $-2\pi$ . Indeed, because of the diagonal form of the metric with respect to the chosen coordinates, each angle has value  $-\frac{\pi}{2}$ . The first two terms are well defined and tend to zero when  $\delta_2$  tends to zero. Hence

$$\lim_{\delta_1 \rightarrow 0} \lim_{\delta_2 \rightarrow 0} \int_{\partial B_{\delta_1, \delta_2}^q} k_g d\sigma_s = -2\pi = -2\pi \tau_q.$$

## References

- [1] A. Agrachev, U. Boscain, M. Sigalotti, A Gauss–Bonnet-like formula on two-dimensional almost-Riemannian manifolds, *Discrete Contin. Dyn. Syst.* 20 (4) (2008) 801–822.
- [2] A.A. Agrachëv, A “Gauss–Bonnet formula” for contact sub-Riemannian manifolds, *Dokl. Akad. Nauk* 381 (5) (2001) 583–585.
- [3] A.A. Agrachev, Y.L. Sachkov, *Control Theory from the Geometric Viewpoint*, Encyclopaedia Math. Sci., vol. 87, Springer-Verlag, Berlin, 2004, *Control Theory and Optimization, II*.
- [4] A. Bellaïche, The tangent space in sub-Riemannian geometry, in: *Sub-Riemannian Geometry*, in: *Progr. Math.*, vol. 144, Birkhäuser, Basel, 1996, pp. 1–78.
- [5] B. Bonnard, J.-B. Caillaud, R. Sinclair, M. Tanaka, Conjugate and cut loci of a two-sphere of revolution with application to optimal control, *Ann. Inst. H. Poincaré Anal. Non Linéaire* 26 (4) (2009) 1081–1098.
- [6] U. Boscain, T. Chambrion, G. Charlot, Nonisotropic 3-level quantum systems: Complete solutions for minimum time and minimum energy, *Discrete Contin. Dyn. Syst. Ser. B* 5 (4) (2005) 957–990.
- [7] U. Boscain, G. Charlot, J.-P. Gauthier, S. Guérin, H.-R. Jauslin, Optimal control in laser-induced population transfer for two- and three-level quantum systems, *J. Math. Phys.* 43 (5) (2002) 2107–2132.
- [8] U. Boscain, B. Piccoli, A short introduction to optimal control, in: T. Sari (Ed.), *Contrôle Non Linéaire et Applications*, Hermann, Paris, 2005, pp. 19–66.
- [9] U. Boscain, M. Sigalotti, High-order angles in almost-Riemannian geometry, in: *Actes de Séminaire de Théorie Spectrale et Géométrie*, vol. 24, Année 2005–2006, in: *Sémin. Théor. Spectr. Géom.*, vol. 25, Univ. Grenoble I, 2008, pp. 41–54.
- [10] B. Franchi, E. Lanconelli, Une métrique associée à une classe d’opérateurs elliptiques dégénérés, in: *Conference on Linear Partial and Pseudodifferential Operators*, Torino, 1982, *Rend. Semin. Mat. Univ. Politec. Torino (Special Issue)* (1984) 105–114, 1983.
- [11] V.V. Grušin, A certain class of hypoelliptic operators, *Mat. Sb. (N.S.)* 83 (125) (1970) 456–473.
- [12] M.W. Hirsch, *Differential Topology*, *Grad. Texts in Math.*, vol. 33, Springer-Verlag, New York, 1994.
- [13] V. Jurdjevic, *Geometric Control Theory*, *Cambridge Stud. Adv. Math.*, vol. 52, Cambridge University Press, Cambridge, 1997.
- [14] B. Malgrange, *Ideals of Differentiable Functions*, *Tata Inst. Fund. Res. Stud. Math.*, vol. 3, Tata Institute of Fundamental Research, Bombay, 1967.
- [15] R. Montgomery, *A tour of subriemannian geometries, their geodesics and applications*, *Math. Surveys Monogr.*, vol. 91, American Mathematical Society, Providence, RI, 2002.
- [16] F. Pelletier, Quelques propriétés géométriques des variétés pseudo-riemanniennes singulières, *Ann. Fac. Sci. Toulouse Math.* (6) 4 (1) (1995) 87–199.
- [17] F. Pelletier, L. Valère Bouche, The problem of geodesics, intrinsic derivation and the use of control theory in singular sub-Riemannian geometry, in: *Actes de la Table Ronde de Géométrie Différentielle*, Luminy, 1992, in: *Sémin. Congr.*, vol. 1, Soc. Math. France, Paris, 1996, pp. 453–512.