

Asymptotic behaviour of solutions to the Korteweg-deVries-Burgers system

by

M. E. SCHONBEK * and S. V. RAJOPADHYE

Department of Mathematics,
University of California at Santa Cruz

ABSTRACT. – We consider the large time behaviour of solutions of the Korteweg-deVries-Burgers system of equations to obtain lower and upper bounds for the rates of decay of the solution. These decay rates extend the work of Amick *et al.* [1], where the scalar case was considered and that of Zhang Linghai [8]. An important tool in the analysis is the so called Fourier Splitting method developed by M. E. Schonbek for obtaining algebraic upper bounds for the solution to the system of parabolic conservation laws. This tool was later used to establish algebraic upper and lower bounds for the Navier Stokes and Magneto Hydrodynamic equations. The lower bounds show that in the far field the behaviour of the solutions to the KdVB system and those of the heat system are very different. This behaviour is believed to be due to the nonlinearity and not to the dispersive nature of the equation, since such behaviour is also present in non-dispersive systems like the Navier-Stokes and the Magneto-Hydrodynamic equations.

Key words: Korteweg-deVries-Burgers equation, decay of solutions.

RÉSUMÉ. – Dans ce travail nous obtenons des bornes supérieures et inférieures pour le taux de décroissance des solutions du système de Korteweg-deVries-Burger. Ce travail prolonge celui de Amick *et al.* [1] dans lequel le cas scalaire est considéré, ainsi que le travail de Zhang Linghai. La technique qui est fondamentale pour les résultats que nous obtenons est le « Fourier splitting ». Cette méthode a été développée par

AMS (MOS) Classification: 35, 76 B

* The work of this author was partially supported by NSF Grant No. DMS 9307497.

Annales de l'Institut Henri Poincaré - Analyse non linéaire - 0294-1449

Vol. 12/95/04/§ 4.00/

M. E. Schonbek pour obtenir des bornes supérieures algébriques pour un système de lois paraboliques de conservation. Elle a été utilisée ensuite pour établir les bornes algébriques supérieures et inférieures pour les solutions des équations de Navier-Stokes et de Magnéto-Hydrodynamique. La borne inférieure montre que pour t tendant vers l'infini, le comportement des solutions pour les systèmes de Korteweg-deVries-Burger et le comportement des solutions de l'équation de la chaleur sont différents : ce comportement proviendrait des termes non linéaires et non de la nature dispersive du système car il est aussi présent dans des systèmes non dispersifs comme les équations de Navier-Stokes et les équations de Magnéto-Hydrodynamique.

1. INTRODUCTION

In this work we study the asymptotic behaviour of solutions to the Korteweg-deVries-Burgers system (henceforth referred to as the KdVB system), in n -space dimensions. This equation can be expressed in the form

$$\begin{aligned}
 (1) \quad U_t + \sum_{i=1}^n \frac{\partial}{\partial x_i} \nabla \Phi(U) + \sum_{i=1}^n \frac{\partial}{\partial x_i} \sum_{i=1}^n \frac{\partial}{\partial x_i^k} U \\
 = \alpha \Delta U + \beta \sum_{i=1}^n \frac{\partial}{\partial x_i} U, \\
 U(x, 0) = U_0,
 \end{aligned}$$

where $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$, $U(x, t) = (U_1(x, t), \dots, U_n(x, t))$ is the n -dimensional vector valued function, $\Phi(U)$ is a scalar function of the vector variable U which satisfies certain growth conditions which will be specified below. The gradient operator with respect to U is denoted by the symbol "grad", $\delta_k = \sum_{i=1}^n \frac{\partial^k}{\partial x_i^k}$, $\delta = \delta_1$ and p, n are integers greater than or equal to 1 and $\alpha > 0$, β are constants and $2p > n > 1$. Note that we can without loss of generality, choose β equal to zero. This is possible by a change of coordinates, from a stationary to a moving frame of reference, which enables us to absorb the term $\beta \delta U$ into the term U_t . Moreover, we can for simplicity assume $\alpha = 1$. Thus we consider the system 1 which

can be rewritten as,

$$(2) \quad \begin{aligned} U_t + \delta \operatorname{grad} \Phi(U) + \delta \delta_{2p} U &= \delta_2 U, \\ U(x, 0) &= U_0, \end{aligned}$$

In one dimension, the equation reduces to the generalized KdVB equation. When the nonlinearity is given by UU_x , Amick *et al.* [1] have obtained sharp rates of decay for the solutions to the KdVB equation. For the scalar case, Zhang Linghai has also obtained decay rates in $L_2 \cap L_\infty$ for a class of equations more general than the KdVB equation; e.g. the Benjamin-Ono-Burgers equation.

In higher dimensions, existence results have been obtained by Zhang [8] following the lines of the proof of [7] for the generalized K-dV equation. Moreover, he obtains rates of decay for the L_2 - and L_∞ -norms of the solution as $t \rightarrow \infty$.

In this work, we first obtain decay rates on the H^m -norm and by an application of standard Sobolev inequalities obtain the L_∞ -decay rate. In addition, lower bounds on the energy decay rates of the solutions are also obtained. It is shown that for a certain class of the initial data, the solution $U(x, t)$ to the KdVB system admits an algebraic lower bound on the energy decay. Two distinct cases have to be considered. First, when the average of the initial data is non-zero and second, when the average equals zero. In the first case, it can be shown that

$$|U(\cdot, t)|_{L_2}^2 \geq C(t+1)^{-\alpha},$$

where $\alpha = \frac{n}{2}$. In the second case, if the average is zero, that is the Fourier transform at the origin is zero, two cases are considered. If the zero is of order one, and the data U_0 lies in $L_1 \cap H^{p+1}$ as well as in suitably weighted L_s spaces, with $s = 1, 2$; then the lower bound is of the form

$$|U(\cdot, t)|_{L_2}^2 \geq C(t+1)^{-\alpha_1},$$

where $\alpha_1 = \frac{n}{2} + 1$ and C depends on the initial data and certain initial parameters. If the zero is of order greater than one, the data lies outside a set of equidistributed energy and the nonlinear function Φ lies in a large class of polynomials, then the lower bound is again of order α_1 . It is this second case which is the more subtle one and which shows the differences between the behaviour of solutions to the KdVB and heat systems in the far field. This explains our interest in studying the lower bound of rates of decay.

This difference in the behaviour of the KdVB and the heat system shows that the nonlinear term produces some mixing of the Fourier modes creating

long waves even when the initial data is highly oscillatory. More precisely if the initial data for the heat system is highly oscillatory the solution will have an exponential rate of decay. Moreover, depending on the data chosen for the heat equation the decay in the energy norm can vary from order $(t + 1)^{-\frac{n}{4}}$ (when no oscillations are present at the origin in Fourier space), to all possible algebraic orders up to exponential decay, depending on how oscillatory the data is. We restrict attention to the case where the nonlinearity is polynomial. By obtaining an algebraic lower bound for the solutions of the KdVB system we show that long waves are produced which slow down the decay. We believe that this phenomenon is due to the nonlinear term and not due to the dispersive term, since this behaviour is also present in non-dispersive systems like the Navier-Stokes and the Magneto-Hydrodynamic equations [4], [5].

The method used here is based on the Fourier Splitting technique, developed by Schonbek [2], to obtain upper bounds for solutions to the Navier-Stokes equation and parabolic conservation laws as well as for obtaining lower bounds on the solution to the Navier-Stokes and Magneto-Hydrodynamics equations.

The paper is organized as follows. In section 2, we briefly review the notational conventions. Section 3 deals with obtaining the H_m decay rates of the solution of the KdVB system. By a simple corollary using Sobolev inequalities the L_∞ decay rate follows. This rate coincides with the one obtained by Zhang [8]. For the sake of completeness, we present some results for the heat equation in section 4. In section 5, upper bounds for the difference of the solution to the heat system and the KdVB system are obtained, when both solutions correspond to the same initial data. Finally the lower bounds for the solution to the KdVB system are derived in section 6.

2. NOTATION

The notation that we use is mostly standard. For the sake of completeness, it is briefly reviewed here. For $1 \leq p \leq \infty$ we denote by $L_p = L_p(\mathbb{R}^n)$ the Banach space of measurable real-valued functions defined on \mathbb{R}^n which are p th power Lebesgue integrable. (essentially bounded in the case $p = \infty$). The usual norm on the space is denoted by $|\cdot|_p$. For non-negative integers s , $H^s = H^s(\mathbb{R}^n)$ is the Sobolev space of functions in L_2 whose generalized derivatives up to order s also belong to L_2 . The space is equipped with

the norm,

$$\|f\|_s^2 = \sum_{j=0}^s |f^{(j)}(\cdot)|_2, \quad \text{where } f^{(j)} = \sum_{|\alpha|=j} D^\alpha f.$$

Of course $H^0 = L_2$ and the L_2 -norm $|\cdot|_2 = \|\cdot\|_0$ will be denoted by the symbol $|\cdot|_2$. We also define a norm on H^s , equivalent to the usual one by,

$$\|U(\cdot, t)\|_s = \sum_{r=0}^s k_{r,s} \sum_{|\beta|=r} \int_{\mathbb{R}^n} |D^\beta U(x, t)|^2 dx,$$

where the $k_{r,s}$'s are positive constants that depend only on r , s and β .

The space $C^r(\mathbb{R}^n, \mathbb{R})$ is the space of all continuous functions from $\mathbb{R}^n \rightarrow \mathbb{R}$ which are r times differentiable with continuous derivatives. In addition we also define some weighted spaces as follows.

$$W_1 = \left\{ U : \int_{\mathbb{R}^n} |x|^2 |U| dx < \infty \right\}$$

and

$$W_2 = \left\{ U : \int_{\mathbb{R}^n} |x| |U|^2 dx < \infty \right\}.$$

The spaces are equipped with the norms $|U|_{W_1} = \int_{\mathbb{R}^n} |x|^2 |U| dx$; $|U|_{W_2} = \left(\int_{\mathbb{R}^n} |x| |U|^2 dx \right)^{\frac{1}{2}}$ respectively.

The Fourier transform of a function $f(x)$ is denoted by $\mathcal{F}(f(\xi))$ and is given by

$$\mathcal{F}(f(\xi)) = \int_{\mathbb{R}^n} e^{-2\pi i \xi \cdot x} f(x) dx.$$

The notation D^β denotes the derivative of order β where β is a multi-index *i.e.*, if

$$\beta = (\beta_1, \beta_2, \dots, \beta_n)$$

then

$$D^\beta = \partial_{x_1}^{\beta_1} \partial_{x_2}^{\beta_2} \dots \partial_{x_n}^{\beta_n}.$$

3. BOUNDS ON THE H^m AND L_∞ -NORM OF THE SOLUTION

Equation (1) has been studied by Zhang [8] to obtain results on the existence of weak solutions under the hypothesis that the nonlinearity (defined by Φ) satisfies the following growth conditions.

$$(3) \quad \left\{ \begin{array}{l} \Phi(U) \leq C|U|^{\frac{4p}{n}+2-q} + C|U|^2, \\ 0 < q < 1, \\ \left| \frac{\partial^2 \Phi(U)}{\partial U_i \partial U_j} \right| \leq C|U|^{\frac{4p}{n}} + C, \\ \text{for all } U \in \mathbb{R}^n, \quad i, j = 1, 2, \dots, n, \\ \left| \frac{\partial^2 \Phi(U)}{\partial U_i \partial U_j} \right| \leq C|U|^{\frac{4p}{n}+1}, \\ \text{for } |U| \text{ small, } \quad i, j = 1, 2, \dots, n. \end{array} \right.$$

In addition, Zhang uses estimates on the L_2 - and the H^s -norms of the solution to show that if $U_0 \in L_1 \cap H^{p+1}$ the weak solution is a strong solution. In the present work, we further restrict Φ to satisfy the condition

$$(4) \quad \Phi(U) = \sum_{i,k} m_{ik} U_i^{k_i},$$

where m_i are real scalars and $3 < k_i \leq \frac{4p}{n} + 2 - q$, with $p > 1$. The case when $k_i = 0, 1, 2$ corresponds to the linear equation and will not be considered for the lower bounds. Hence it also follows that,

$$(5) \quad \left\{ \begin{array}{l} |\Phi(U)| \leq C|U|^{\frac{4p}{n}+2-q}, \quad 0 < q < 1, \\ \left| \frac{\partial^2 \Phi(U)}{\partial U_i \partial U_j} \right| \leq C|U|^{\frac{4p}{n}}, \\ \left| \frac{\partial^2 \Phi(U)}{\partial U_i} \right| \leq C|U|^{\frac{4p}{n}+1}. \end{array} \right.$$

for all $U \in \mathbb{R}^n$ and $i, j = 1, 2, \dots, n$. Zhang has shown (cf. Theorem 6 in [8]) that if $U_0 \in L_1 \cap H^{p+1}$; $\Phi(U) \in C^3(\mathbb{R}^n)$ and satisfies (5), the following decay properties hold for the solution to the KdVB system:

$$(6) \quad \left\{ \begin{array}{l} \|U(t)\|_2 \leq C(t+1)^{-\frac{n}{4}}, \\ \|U(t)\|_\infty \leq C(t+1)^{-\frac{n}{4}}. \end{array} \right.$$

Here we first obtain the decay rate for the H^s -norm of the solution. Then, making use of standard Sobolev inequalities, the decay rate obtained in [8] for the L_∞ -norm follows.

THEOREM 3.1. – *If $\Phi(U) \in C^{m+1}(\mathbb{R}^n, \mathbb{R})$ and satisfies equation (5) with $U_0 \in L_1 \cap H^{p+1}(\mathbb{R}^n)$, then the solution U to the KdVB system with initial data U_0 is such that,*

$$(7) \quad \sum_{|\alpha| \leq m} \int_{\mathbb{R}^n} |D^\alpha U|^2 dx \leq C(t+1)^{-\frac{n}{2}}.$$

Moreover, if $m > \left[\frac{n}{2} \right]$ then,

$$\sum_{|\alpha| \leq m - \left[\frac{n}{2} \right]} |D^\alpha U|_\infty \leq C(t+1)^{-\frac{n}{4}},$$

where the constant C depends only on Φ , n , m and U_0 .

Proof. – To obtain decay rates for the H^s -norm of the solution of the KdVB system, we use induction. We begin by giving an equivalent definition for the H^s -norm. Define,

$$\| \| U(t) \| \|_s^2 := \sum_{r=0}^s k_{r,s} \sum_{|\beta|=r} \int_{\mathbb{R}^n} |D^\beta U(x,t)|^2 dx,$$

where the $k_{r,s}$'s are positive constants depending on r , s , β and $\|U\|_\infty$ which will be determined below. We will show by induction that

$$(8) \quad \frac{d}{dt} \| \| U \| \|_s^2 \leq -C \sum_{r=1}^{s+1} k_{r,s} \sum_{|\beta|=r} \int |D^\beta U|^2 dx$$

and

$$(9) \quad \| \| U \| \|_s^2 \leq C(t+1)^{-\frac{n}{2}}.$$

For $|s| = 0$ inequality (8) follows by multiplying the equation by U and integrating in space and inequality (9) reduces with $k_{0,0} = 1$ to

$$(10) \quad \int_{\mathbb{R}^n} |U(t)|^2 dx \leq C(t+1)^{-\frac{n}{2}},$$

a result obtained in [8]. For the sake of completeness, we give a sketch of the proof here. Multiply equation (2) by U and integrate over space to get,

$$(11) \quad \frac{1}{2} \frac{d}{dt} \int |U(x, t)|^2 dx + \int U \delta \Phi(U) dx \\ + \int U \delta \delta_{2p} U dx = \int U \delta_2 U dx.$$

On simplification, we have,

$$\frac{d}{dt} \int |U(x, t)|^2 dx = -2 \int |\nabla U|^2 dx.$$

By Plancherel's theorem the equality can be rewritten as

$$(12) \quad \frac{d}{dt} \int |\hat{U}|^2 d\xi = -2 \int |\xi|^2 |\hat{U}|^2 d\xi.$$

Using the Fourier Splitting technique introduced by Schonbek [2], the Fourier space is divided into two time dependent disjoint sets $A(t)$ and $A(t)^c$, where

$$(13) \quad A(t) := \left\{ \xi : |\xi| \leq \left[\frac{n}{2(t+1)} \right]^{\frac{1}{2}} \right\}.$$

Inequality (12) can then be rewritten as,

$$(14) \quad \frac{d}{dt} \int_{\mathbb{R}^n} |U|^2 d\xi \\ \leq -2 \int_{A(t)} |\xi|^2 |\hat{U}|^2 d\xi - 2 \int_{A(t)^c} |\xi|^2 |\hat{U}|^2 d\xi \\ \leq -\frac{n}{t+1} \int_{A^c(t)} |\hat{U}|^2 d\xi \\ \leq -\frac{n}{t+1} \int_{\mathbb{R}^n} |\hat{U}|^2 d\xi + \frac{n}{t+1} \int_{A(t)} |\hat{U}|^2 d\xi.$$

Hence it follows that,

$$(15) \quad \frac{d}{dt} \left[(t+1)^n \int_{\mathbb{R}^n} |\hat{U}|^2 d\xi \right] \leq n(t+1)^{n-1} \int_{A(t)} |\hat{U}|^2 d\xi.$$

Moreover, it has been shown in [8] that if $U_0 \in L_1 \cap H^{p+1}$, then

$$(16) \quad |\hat{U}(\cdot, t)|_\infty \leq \|U_0\|_1 + \|U_0\|_p^2.$$

Equation (15) together with this time independent bound for the Fourier transform yields,

$$(17) \quad \left\{ \begin{aligned} & \frac{d}{dt} \left[(t+1)^n \int_{\mathbb{R}^n} |\hat{U}|^2 d\xi \right] \\ & \leq n (t+1)^{n-1} (|U_0|_1^2 + \|U_0\|_p^2)^2 \int_{A(t)} d\xi \\ & \leq (|U_0|_1 + \|U_0\|_p^2)^2 n (t+1)^{n-1} \omega_n \left(\frac{n}{2(t+1)} \right)^{\frac{n}{2}}, \end{aligned} \right.$$

where ω_n denotes the volume of the n -dimensional unit sphere. Integrating (17) over the interval $[0, t]$ inequality (10) follows. This proves the estimate in the case $|s| = 0$.

Define,

$$\| \| U(t) \| \| _s^2 := \sum_{r=0}^s k_{r,s} \sum_{|\beta|=r} \int_{\mathbb{R}^n} |D^\beta U(x,t)|^2 dx.$$

By the induction hypothesis, it follows that,

$$(18) \quad \frac{d}{dt} \| \| U \| \|_{s-1}^2 \leq -C \sum_{r=0}^s k_{r,s-1} \sum_{|\beta|=r} \int |D^\beta U|^2 dx$$

and

$$\| \| U \| \|_{s-1}^2 \leq C (t+1)^{-\frac{n}{2}}.$$

Hence, for an appropriate choice of constants $k_{r,s}$, it is necessary to show that

$$\| \| U(t) \| \| _s^2 \leq C (t+1)^{-\frac{n}{2}}$$

which would prove the desired result. Multiply equation (2) by $D^{2\beta} U$ and integrate over space. This yields,

$$(19) \quad \begin{aligned} & (-1)^\beta \frac{1}{2} \frac{d}{dt} \int_{\mathbb{R}^n} |D^\beta U|^2 dx + \int_{\mathbb{R}^n} D^{2\beta} U \delta \text{grad } \Phi(U) dx \\ & + \int_{\mathbb{R}^n} D^{2\beta} U \delta \delta_{2p} U dx = (-1)^{\beta+1} \sum_{j=1}^n \int_{\mathbb{R}^n} |D^{\beta+e_j} U|^2 dx \end{aligned}$$

Note that the third term on the left-hand side integrates to zero. Using the definition of Φ given by (4) and hence determining the growth condition on $\text{grad } \Phi$, the second term on the right-hand side of (19) can be bounded above by

$$C \int_{\mathbb{R}^n} \delta \left(\sum U_i^{k_i-1} \right) D^{2\beta} U \, dx$$

where $k_i \leq \frac{4p}{n} + 2 - q$. Thus it follows that $2 < k = k_i - 1 \leq \frac{4p}{n} + 1$.

Combining this bound with equation (19) yields,

$$\begin{aligned} (20) \quad & \frac{d}{dt} \int_{\mathbb{R}^n} |D^\beta U|^2 \, dx \\ & \leq -2 \sum_{j=1}^n \int_{\mathbb{R}^n} |D^{\beta+\varepsilon_j} U|^2 \, dx \\ & \quad + C \sum_{j=1}^n \int_{\mathbb{R}^n} D^{\beta+\varepsilon_j} U D^\beta U^k \, dx \\ & \leq -2 \sum_{j=1}^n \int_{\mathbb{R}^n} |D^{\beta+\varepsilon_j} U|^2 \, dx \\ & \quad + C \left(\sum_{j=1}^n \frac{1}{C} \int_{\mathbb{R}^n} |D^{\beta+\varepsilon_j} U|^2 \, dx \right. \\ & \quad \left. + \sum_{j=1}^n \frac{C}{4} \int_{\mathbb{R}^n} |D^\beta U^k|^2 \, dx \right) \\ & \leq - \sum_{j=1}^n \int_{\mathbb{R}^n} |D^{\beta+\varepsilon_j} U|^2 \, dx + \frac{C^2}{4} \int_{\mathbb{R}^n} |D^\beta U^k|^2 \, dx \\ & \leq - \sum_{j=1}^n \int_{\mathbb{R}^n} |D^{\beta+\varepsilon_j} U|^2 \, dx + \frac{C^2}{4} C_0 \int_{\mathbb{R}^n} |D^\beta U|^2 \, dx \\ & \quad + \sum_{|\alpha| \leq \beta} C_{\alpha, \beta} \int_{\mathbb{R}^n} |D^\alpha U|^2 \, dx. \end{aligned}$$

In the second step we have made use of Schwartz's inequality, and in the last step we use the fact that if $s > k$ we have

$$D^s U^k = \sum_{j=1}^s C_{k, s} U^{k-s-j} \left(\sum_{\alpha r=j} (D^\alpha U)^r \right), \quad r \text{ integer}$$

and if $s \leq k$

$$D^s U^k = \sum_{\alpha_1 + \dots + \alpha_k = s} D^{\alpha_1} U \dots D^{\alpha_k} U,$$

and that $j \leq s$ and $\alpha, r \leq s$. In addition, by the induction hypothesis, since derivatives up to order $s - 1$ are bounded in L_2 , the last inequality in (20) follows by using a simple interpolation inequality and C_0 denotes a constant depending on Φ and the L_∞ -norms of U and its derivatives up to order $\beta - \left[\frac{n}{2} \right]$. Hence inequality (20) yields,

$$(21) \quad \begin{aligned} \frac{d}{dt} \int_{\mathbb{R}^n} |D^\beta U|^2 dx &\leq - \sum_{j=1}^n \int_{\mathbb{R}^n} |D^{\beta+\epsilon_j} U|^2 dx \\ &+ C_1 \int_{\mathbb{R}^n} |D^\beta U|^2 dx \\ &+ \sum_{|\alpha| \leq \beta} C_{\alpha, \beta} \int_{\mathbb{R}^n} |D^\alpha U|^2 dx. \end{aligned}$$

Summing over all β such that $|\beta| = s$ leads to

$$(22) \quad \begin{aligned} \frac{d}{dt} \left[\int_{\mathbb{R}^n} \sum_{|\beta|=s} |D^\beta U|^2 dx \right] \\ \leq C_s \left(C_1 \int_{\mathbb{R}^n} \sum_{|\beta|=s} |D^\beta U|^2 dx \right) \\ - \sum_{|\beta|=s+1} \left(\int_{\mathbb{R}^n} |D^\beta U|^2 dx \right). \end{aligned}$$

By the inductive hypothesis,

$$(23) \quad \frac{d}{dt} [||| U(t) |||_{s-1}^2] \leq -C \sum_{r=1}^s k_{r,s} \sum_{|\beta|=r} \int_{\mathbb{R}^n} |D^\beta U|^2 dx.$$

Let $m_{s-1} = C k_{r, s-1}$ $r \leq s-1$. Choose $k_{s+1} = m_{s-1} (2C_s C_1)^{-1}$. Adding together inequalities (23) and (22) times k_{s+1} we get,

$$\begin{aligned} \frac{d}{dt} ||| U |||_s^2 &\leq -C \sum_{r=0}^s k_r \sum_{|\beta|=s} \int_{\mathbf{R}^n} |D^\beta U|^2 dx \\ &\quad + C \max_{r \leq s} k_{r, s-1} \int_{\mathbf{R}^n} \sum_{|\beta|=s} |D^\beta U|^2 dx \\ &\quad - R_{s+1} \sum_{|\beta|=s+1} \int_{\mathbf{R}^n} |D^\beta U|^2 dx \\ &\leq -K_s \left(\sum_{r=1}^{s+1} k_r \sum_{|\beta|=r} \int_{\mathbf{R}^n} |D^\beta U|^2 dx \right), \end{aligned}$$

where $K_s = \min \{1, K_{s-1}\}$, and R_{s+1} is proportional to m_{s-1} . Let $k_{r, s} = k_{r, s-1}/2$, $k_{s, s} = k_{s+1}$ and $k_{s+1, s} = k_{s, s}$. This establishes the inequality

$$(24) \quad \frac{d}{dt} ||| U(t) |||_s^2 \leq -C \sum_{r=1}^{s+1} k_{r, s} \sum_{|\beta|=r} \int_{\mathbf{R}^n} |D^\beta U|^2 dx.$$

If $|\beta| = 0$ (24) reduces to obtaining the L_2 -rate of decay for U , which has been established in (10). Suppose that the decay rate has been obtained for β with $0 \leq |\beta| < s$. Let $|\beta| = s$. Taking the Fourier transform of inequality (24) we obtain,

$$\begin{aligned} (25) \quad \frac{d}{dt} ||| \hat{U} |||_s^2 &= \frac{d}{dt} \sum_{r=0}^s k_{r, s} \sum_{|\beta|=r} \int_{\mathbf{R}^n} |\widehat{D^\beta U}|^2 d\xi \\ &\leq -C \sum_{r=1}^{s+1} k_{r, s} \sum_{|\beta|=r} \int_{\mathbf{R}^n} |\widehat{D^\beta U}|^2 d\xi \\ &= -C \sum_{r=1}^s k_{r, s} \sum_{|\beta|=r} \int_{\mathbf{R}^n} |\xi|^2 |\widehat{D^\beta U}|^2 d\xi. \end{aligned}$$

Repeating the same argument in (25) as that used in obtaining (17), we have,

$$\begin{aligned} & \frac{d}{dt} [(t + 1)^n ||| U(t) |||_s] \\ &= \frac{d}{dt} \left[(t + 1)^n \sum_{r=0}^s k_{r,s} \sum_{|\beta|=r} \int_{\mathbf{R}^n} |D^{\widehat{\beta}} U|^2 d\xi \right] \\ &\leq K_s (t + 1)^{n-1} \sum_{r=0}^s k_{r,s} \sum_{|\beta|=r} \int_{A(t)} |D^{\widehat{\beta}} U|^2 d\xi, \end{aligned}$$

where $A(t)$ is as defined earlier in (13). Hence, if $K_s = \max k_{r,s}$

$$\begin{aligned} (26) \quad \frac{d}{dt} [(t + 1)^n ||| \hat{U} |||_s^2] &\leq C (t + 1)^{n-1} K_s \sum_{|\beta|=r} \int_{A(t)} |D^{\widehat{\beta}} U|^2 d\xi \\ &\leq C (t + 1)^{n-1} K_s \sum_{|\beta|<s} \int_{\mathbf{R}^n} |D^{\widehat{\beta}} U|^2 d\xi \\ &\quad + C (t + 1)^{n-1} \int_{A(t)} |\xi|^2 |D^{s-1} U|^2 \\ &\leq C (t + 1)^{n-1} K_s \sum_{|\beta|<s} \int_{\mathbf{R}^n} |D^{\widehat{\beta}} U|^2 d\xi \\ &\quad + C (t + 1)^{n-2} \int_{\mathbf{R}^n} |D^{s-1} U|^2 d\xi. \end{aligned}$$

By the inductive hypothesis for $|\beta| < s$,

$$\int_{\mathbf{R}^n} |D^{\widehat{\beta}} U|^2 d\xi \leq C (t + 1)^{-\frac{n}{2}}.$$

Using this in (26) we obtain,

$$\frac{d}{dt} [(t + 1)^n ||| \hat{U} |||_s^2] \leq C (t + 1)^{\frac{n}{2}-1}.$$

Integrating this last expression over the interval $[0, t]$ yields

$$(27) \quad (t + 1)^n ||| U(t) |||_s^2 \leq C (t + 1)^{\frac{n}{2}} + ||| U(0) |||_s^2,$$

which proves the claim. The bounds for the L_∞ -norm of U follow from inequality (27) and Sobolev's inequality. Hence we obtain,

$$\sum_{|\beta|=j} |D^\beta U|_\infty \leq C \|U\|_m, \quad j < m - \left[\frac{n}{2}\right].$$

which in turn yields,

$$\sum_{|\beta|=j} |D^\beta U|_\infty \leq C (t+1)^{-\frac{n}{4}}, \quad \text{with } j < m - \left[\frac{n}{2}\right].$$

4. PRELIMINARY ESTIMATES

In this section we begin with some preliminary estimates for the heat system. If $V(x, t)$ is the solution to the homogeneous heat system, *i.e.*, V satisfies

$$(28) \quad \begin{cases} V_t(x, t) = \Delta V(x, t), \\ V(x, 0) = U_0(x). \end{cases}$$

Then we have the following

THEOREM 4.1. – Let $U_0 \in L_1 \cap L_2(\mathbb{R}^n) \cap R_\alpha^{\delta_1}$ for some $\alpha, \delta_1 > 0$, where

$$R_\alpha^{\delta_1} = \{U : |\widehat{U}(\xi)| \geq \alpha, \text{ for } |\xi| \leq \delta_1\}.$$

Then

$$\int_{\mathbb{R}^n} |V(x, t)|^2 dx \geq C (t+1)^{-\frac{n}{2}} \quad \text{and} \quad |\nabla V|_\infty \leq C t^{-\left(\frac{n}{4}+1\right)}.$$

Proof. – See Schonbek [3].

THEOREM 4.2. – *Let V be a solution to the heat system with data U_0 . Suppose that*

$$\|V(t)\|_2^2 \leq C(t+1)^{-\rho}.$$

Then,

$$(29) \quad \|D^s V\|_\infty < C(t+1)^{-\frac{n}{4} - \frac{\rho}{2} - \frac{s}{2}} \quad s \geq 0.$$

Proof. – Let $\|V(t)\|_2^2 \leq C(t+1)^{-\rho}$ then it follows that

$$\begin{aligned} \|D^s V\|_\infty &\leq \int |\hat{D}^s V| d\xi \\ &\leq \int |\xi|^{2s} \hat{V}(t/2) e^{-|\xi|^2 t/2} d\xi \\ &\leq \left(\int |\xi|^{2s} e^{-|\xi|^2 t} d\xi \right)^{\frac{1}{2}} \left(\int_{\mathbb{R}^n} |\hat{V}(t/2)|^2 d\xi \right)^{\frac{1}{2}} \\ &\leq C_1 ((t+1)^{-\frac{n}{2} - s})^{\frac{1}{2}} (t+1)^{-\frac{\rho}{2}} \\ &\leq C_1 (t+1)^{-\frac{n}{4} - \frac{s}{2} - \frac{\rho}{2}}. \end{aligned}$$

PROPOSITION 4.3. – *Let $V_0 \in H^s \cap L_1$. This implies that if V is a solution to the heat system with data V_0 then*

$$(30) \quad \|D^s V\|_2^2 \leq C(t+1)^{-\frac{n}{2} - s}.$$

Proof. – The proof is standard and we include it for completeness. The s -derivatives of the solution to the heat equation can be explicitly given by

$$D^s V(x, t) = (4\pi t)^{-\frac{n}{2}} \int_{\mathbb{R}^n} D^s \exp\left(-\frac{|x-y|^2}{4\pi t}\right) |V_0(y)|^2 dy.$$

Hence

$$\begin{aligned}
 \int_{\mathbf{R}^n} |D^s V(x, t)|^2 dx &\leq (4\pi t)^{-\frac{n}{2}} \int_{\mathbf{R}^n} \int_{\mathbf{R}^n} (4\pi t)^{-\frac{n}{2}} D^s \\
 &\quad \times \exp\left(-\frac{|x-y|^2}{4\pi t}\right) |V_0(y)|^2 dy dx \\
 &\leq (4\pi t)^{-\left(\frac{n}{2}+s\right)} \int_{\mathbf{R}^n} (4\pi t)^{-\frac{n}{2}} \\
 &\quad \times \exp\left(-\frac{|x-y|^2}{2\pi t}\right) dx \int_{\mathbf{R}^n} |V_0(y)|^2 dy \\
 &\leq C(t+1)^{-\frac{n}{2}-s},
 \end{aligned}$$

which completes the proof of the proposition.

COROLLARY 4.4. - If $|V|_2^2 \leq C(t+1)^{-\rho}$, then

$$(31) \quad \int_{\mathbf{R}^n} |\widehat{D^s V}|^2 d\xi \leq C(t+1)^{-\frac{n}{2}-\rho-s}.$$

5. UPPER BOUNDS FOR THE DIFFERENCE

In this section we discuss bounds for the difference of the solution to the KdVB system and that of the heat system corresponding to the same initial data. There are several ways to approach this problem. Our approach, will be the Fourier splitting method ([3], [4]).

We first consider the case when the average of U is non-zero. That is, we have $\int U_0 dx \neq 0$. Let R_1 be a set defined by

$$R_1 = \left\{ U_0 : \int_{\mathbf{R}^n} U_0 dx \neq 0 \right\}.$$

This implies that $U \in R_\chi^{\delta_1}$ where $R_\chi^{\delta_1}$ is defined by

$$R_\chi^{\delta_1} = \{U : |\widehat{U}(\xi)| \geq \chi, \text{ for } |\xi| \leq \delta_1\}.$$

We now obtain upper bounds for the solution to the KdVB system. But first, we need the following result.

THEOREM 5.1. – *Let the initial data U_0 be such that $U_0 \in L_1 \cap R_1 \cap H^{p+1}$. Then there exists an upper bound for the difference between the solutions to the KdVB system and the heat system, both corresponding to the same initial data U_0 . Thus if $p > 1$ and if $W = U - V$ where U is the solution to the KdVB system and V that to the heat system, then W satisfies,*

$$\|W(t)\|_2^2 \leq C_0 (t+1)^{-(\frac{n}{2}+1)}.$$

Proof. – We consider the difference $W = U - V$, where U is the solution to the KdVB system, and V that of the heat system, both corresponding to the same initial data U_0 . Then W satisfies the equation

$$(32) \quad W_t - \Delta W = -\delta \delta_{2p} U - \delta \operatorname{grad} \Phi(U).$$

Multiply equation (32) by W and integrate over space. Noting that $\int U \delta \delta_{2p} U = 0$ and $\int U \delta \operatorname{grad} \Phi(U) = 0$, we obtain

$$(33) \quad \begin{aligned} \frac{d}{dt} \int_{\mathbf{R}^n} |W|^2 dx &\leq -2 \int_{\mathbf{R}^n} |\delta W|^2 dx \\ &\quad + 2 |\delta V|_\infty \int_{\mathbf{R}^n} |\operatorname{grad} \Phi(U)| dx \\ &\quad + 2 \left(\int_{\mathbf{R}^n} |\delta_{2p+1} V|^2 dx \right)^{\frac{1}{2}} \left(\int_{\mathbf{R}^n} |U|^2 dx \right)^{\frac{1}{2}}. \end{aligned}$$

Using Plancherel's theorem the above inequality can be written in the form,

$$(34) \quad \begin{aligned} \frac{d}{dt} \int_{\mathbf{R}^n} |\hat{W}|^2 dx &\leq -2 \int_{\mathbf{R}^n} |\xi|^2 |\hat{W}|^2 d\xi \\ &\quad + 2 |\delta V|_\infty \int_{\mathbf{R}^n} |\operatorname{grad} \Phi(U)| dx \\ &\quad + 2 \left(\int_{\mathbf{R}^n} |\delta_{2p+1} V|^2 dx \right)^{\frac{1}{2}} \left(\int_{\mathbf{R}^n} |U|^2 dx \right)^{\frac{1}{2}}. \end{aligned}$$

Using the Fourier Splitting method, with

$$S(t) = \left\{ \xi : |\xi| \leq \left(\frac{\gamma}{2(t+1)} \right)^{\frac{1}{2}} \right\},$$

with γ sufficiently large, equation (34) can be rewritten as,

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}^n} |\hat{W}|^2 d\xi &= -2 \int_{S(t)} |\xi|^2 |\hat{W}|^2 d\xi - 2 \int_{S^c(t)} |\xi|^2 |\hat{W}|^2 d\xi \\ &\quad + 2 |\delta V|_\infty \int_{\mathbb{R}^n} |\text{grad } \Phi(U)|^2 dx \\ &\quad + 2 \left(\int_{\mathbb{R}^n} |D^{2p+1} V|^2 dx \right)^{\frac{1}{2}} \left(\int_{\mathbb{R}^n} |U|^2 dx \right)^{\frac{1}{2}}. \end{aligned}$$

Hence, this can be simplified to yield,

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}^n} |\hat{W}|^2 d\xi &\leq -\frac{\gamma}{(t+1)} \int_{S^c(t)} |\hat{W}|^2 d\xi - 2 \int_{S(t)} |\xi|^2 |\hat{W}|^2 d\xi \\ &\quad + 2 |DV|_\infty \int_{\mathbb{R}^n} |\text{grad } \Phi(U)| dx \\ &\quad + 2 \left(\int_{\mathbb{R}^n} |D^{2p+1} V|^2 dx \right)^{\frac{1}{2}} \left(\int_{\mathbb{R}^n} |U|^2 dx \right)^{\frac{1}{2}}. \end{aligned}$$

Further simplification yields,

$$\begin{aligned} &\frac{d}{dt} \left[(t+1)^\gamma \int_{\mathbb{R}^n} |W|^2 dx \right] \\ &\leq n(t+1)^{\gamma-1} \int_{S(t)} |\hat{W}|^2 d\xi \\ &\quad + C(t+1)^\gamma |DV|_\infty \int_{\mathbb{R}^n} |\text{grad } \Phi(U)| dx \\ &\quad + C(t+1)^\gamma \left(\int_{\mathbb{R}^n} |D^{2p+1} V|^2 dx \right)^{\frac{1}{2}} \left(\int_{\mathbb{R}^n} |U|^2 dx \right)^{\frac{1}{2}} \end{aligned}$$

Using the bounds obtained in (30) and (31) for the solution of the heat equation and the growth condition for Φ expressed in (5), we obtain,

$$\begin{aligned}
 (35) \quad & \frac{d}{dt} \left[(t+1)^\gamma \int_{\mathbb{R}^n} |W|^2 dx \right] \\
 & \leq \gamma (t+1)^{\gamma-1} \int_{S(t)} |\hat{W}|^2 d\xi \\
 & \quad + C (t+1)^\gamma (t+1)^{-\left(\frac{n}{4} + \frac{1}{2}\right)} C_0 \cdot (|U|_2^2 |U|_\infty^{\frac{4p}{n}-1}) \\
 & \quad + C (t+1)^\gamma (t+1)^{-\left(\frac{n}{4} + p + \frac{1}{2}\right)} |U|_2 \\
 & \leq \gamma (t+1)^{\gamma-1} \int_{S(t)} |\hat{W}|^2 d\xi + C_1 (t+1)^{\gamma_1} + C_2 (t+1)^{\gamma_2},
 \end{aligned}$$

supposing first that $p > 1$ then $\gamma_1 \geq \gamma_2 = \gamma - \left(\frac{n}{2} + p + \frac{1}{2}\right)$. We now need to estimate $|\hat{W}|$ on the set $S(t)$. Taking the Fourier transform of (32) we have,

$$(36) \quad \begin{cases} \hat{W}_t + |\xi|^2 \hat{W} = -\delta \widehat{\delta_{2p} U} - \delta \widehat{\text{grad} \Phi(U)} = -\hat{H} \\ \hat{W}(\xi, 0) = 0 \end{cases}$$

Hence, it follows that

$$\hat{W}(\xi, t) = - \int_0^t e^{-|\xi|^2(t-s)} \hat{H}(\xi, s) ds.$$

If ν denotes the minimum power of U_i in the definition of Φ then,

$$(37) \quad |\hat{H}| \leq C |\xi| (|U|_2^2 (|U|_\infty^{\nu-2} + |U|_\infty^{\frac{4p}{n}-1}) + |\xi|^{2p} |\hat{U}|_\infty).$$

Since $|\hat{U}|$ is bounded by a constant depending only on the initial data and since $\xi \in S(t)$, $|\xi| \leq \left(\frac{n}{(t+1)}\right)^{\frac{1}{2}}$. Moreover, with $p \geq 1$ and $n \geq 2$ it follows that

$$\begin{aligned}
 |\widehat{W}(\xi, t)|_\infty & \leq \frac{C}{(t+1)^{\frac{1}{2}}} \int_0^t [(s+1)^{-(p+\frac{n}{4})} + (s+1)^{-\frac{\nu n}{4}} + C(s+1)^{-p}] ds \\
 & \leq \frac{C}{(t+1)^{\frac{1}{2}}}.
 \end{aligned}$$

Thus,

$$(38) \quad |W(\xi, t)|_\infty \leq \frac{C}{(t+1)^{\frac{1}{2}}}.$$

Substituting (38) into (35) and simplifying the result we obtain

$$\frac{d}{dt} \left[(t+1)^\gamma \int_{\mathbf{R}^n} |W|^2 dx \right] \leq C(t+1)^{\gamma-1} (t+1)^{-1-\frac{n}{2}} + C_1(t+1)^{\gamma_2},$$

since $\gamma_1 > \gamma_2$. Integrating over the interval $[0, t]$ yields,

$$(t+1)^\gamma \int_{\mathbf{R}^n} |W(t)|^2 dx \leq C_1(t+1)^{\gamma-(\frac{n}{2}+1)}$$

or equivalently,

$$\int_{\mathbf{R}^n} |W(t)|^2 dx \leq (t+1)^{-(\frac{n}{2}+1)}.$$

This proves the theorem.

6. LOWER BOUNDS

In obtaining the lower bounds for the solution to the KdVB system, two cases have to be considered. In the first case the mean of the initial data is different from zero, that is long waves are present. In this case the lower bound for the decay rate for the solution to the heat system is $(t+1)^{-\frac{n}{2}}$. Hence from the previous section we note that as a consequence of the upper bounds for the difference of solutions to the heat equation and the KdVB system, which is of order $(t+1)^{-(\frac{n}{2}+1)}$ when $p > 1$, the lower bounds for the solution to the KdVB system follow easily. That is since

$$|U(t)|_2^2 \geq |V|_2^2 - |W|_2^2,$$

it follows that,

$$|U(t)|_2^2 \geq C(t+1)^{-\frac{n}{2}}.$$

The second case to be considered is when the mean of the initial data equals zero. In Fourier space this is equivalent to the statement that the Fourier transform of the initial data has a zero at the origin. The bounds that are obtained in this case for the solution to the heat system depend on the order of the zero. We will show that outside of a class of data to be defined below the solution to the KdVB system will have a uniform rate of decay of order $\frac{n}{2} + 1$, showing that in the far-field the behaviour of the solutions to the Heat system and that to the KdVB system are very different, *i.e.*, long waves can be created for such systems, a phenomenon which is absent in the Heat system. Before we begin estimates for the lower bound, some preliminary results are needed. We begin by stating the following result, proved by Schonbek [4].

THEOREM 6.1. – *Let $V_0 \in L_2(\mathbb{R}^n)$. Let V be a solution to the heat system with initial data V_0 . Suppose that there exists functions l and h such that the Fourier transform of V_0 admits, for $|\xi| \leq \delta_1$ with $\delta_1 > 0$ the representation*

$$(39) \quad \hat{V}_0(\xi) = \xi \cdot l(\xi) + h(\xi), \quad l = (l_1, \dots, l_n),$$

where l and h satisfy the following conditions:

- (i) $|h(\xi)| \leq M_0 |\xi|^2$, for some $M_0 > 0$.
- (ii) l is homogeneous of degree zero.
- (iii) $\gamma = \int_{|\omega|=1} |\omega \cdot l(\omega)|^2 d\omega > 0$.

Let $M_1 = \sup_{|y|=1} |l(y)|$, $M_2 = \sup_{\frac{\delta_1}{2} \leq |y| \leq 1} |\nabla l(y)|$, $K = \max(M_0, M_1, M_2)$. Then there exist constants C_0 and C_1 such that

$$C_0 (t+1)^{-\left(\frac{n}{2}+1\right)} \leq |V(\cdot, t)|_2^2 \leq C_1 (t+1)^{-\left(\frac{n}{2}+1\right)}$$

where C_0 and C_1 both depend on n , M_0 , M_1 , δ_1 and $|V_0|_2$ and C_0 also depends on K and α_1 . Note that condition (iii) is not necessary for the upper bound of $|V(\cdot, t)|_2$.

Proof. – See Schonbek [4].

Let

$$\begin{aligned}\mathcal{M}_{\mathcal{R}} &= \{U : \operatorname{Re} \widehat{\nabla U}(0) = 0\} \\ \mathcal{M}_{\mathcal{I}} &= \{U : \operatorname{Im} \widehat{\nabla U}(0) = 0\} \\ P_k &= \int_0^\infty \widehat{\Phi}_{U_k}(U(0, s)) ds = \int_0^\infty a_k^0 ds \\ \bar{P}_k &= (P_k, \dots, P_k) \quad \text{and} \\ \hat{\Phi}(U) &= \left[\int \Phi e^{-2\pi i x \xi} dx \right]\end{aligned}$$

If the Fourier transform of the initial data has a zero of order one and $U_0 \in W_1$ then, this implies that either $U_0 \notin \mathcal{M}_{\mathcal{R}}$ and $U_0 \notin \mathcal{M}_{\mathcal{I}}$.

THEOREM 6.2. – *Let $U_0 \in W_1 \cap L_1 \cap H^{p+1}$. Suppose that $\int U_0 dx = 0$ and \hat{U}_0 has a zero of order one. Let $p > 1$ and let U be a solution to the KdVB system with data U_0 . Suppose that*

- (i) $U_0 \notin \mathcal{M}_{\mathcal{R}}$ or
- (ii) For some k ,

$$\operatorname{Im} \widehat{\nabla U}_k(0) - P_k \neq 0.$$

Then there exist constants C_1 and C_2 such that

$$C_1(1+t)^{-\left(\frac{n}{2}+1\right)} \leq |V(\cdot, t)|_2^2 \leq C_2(1+t)^{-\left(\frac{n}{2}+1\right)}$$

for all $t \geq 0$ where V is the solution to the heat equation with data U_0 and there exist constants K_1 and K_2 such that

$$K_1(1+t)^{-\left(\frac{n}{2}+1\right)} \leq |U(\cdot, t)|_2^2 \leq K_2(1+t)^{-\left(\frac{n}{2}+1\right)}$$

where C_1, C_2, K_1, K_2 depend only on the norms of the data.

Proof. – We first need to explain our hypotheses. They ensure that long waves persists, *i.e.*, we associate long waves with the terms of order one in the Fourier expansion. Note that

$$U(\widehat{\xi}, t) = \hat{U}_0 + \xi \widehat{\nabla U}_0 + i \xi P + \mathcal{O}(|\xi|^2).$$

If we choose data which satisfies *A* or *B* below, then the first order terms in the Taylor expansion of the Fourier transform persists, *i.e.*, long waves

will persist. Since the hypotheses are such that $\hat{U}_0(\xi)$ has a zero of order one, we need that

$$(40) \quad P_k - \text{Im } \widehat{\nabla} U_0 \neq 0. \quad \text{or } \text{Re } \widehat{\nabla} U_0 \neq 0.$$

Note that the following conditions on the data and on the function Φ ensure that (40) holds.

A) $U_0 \notin \mathcal{M}_R^c$.

B) $U_0 \in \mathcal{M}_I$ and $\Phi(U) = \sum_i \sum_k m_{k_i} U_i^{k_i}$ with $m_{k_i} = 0$ for k odd.

We begin by considering the KsVB system. The nonlinear term $\delta \text{grad } \Phi(U)$ is such that

$$\text{grad } \widehat{\Phi}(U) = (\widehat{\Phi}_{U_1}, \widehat{\Phi}_{U_2}, \dots, \widehat{\Phi}_{U_n}).$$

We introduce the notation

$$a_i = \widehat{\Phi}_{U_i}$$

Hence we have

$$\text{grad } \widehat{\Phi}(U) = (a_1, \dots, a_n).$$

Taking the Fourier transform of the KdVB system we arrive at,

$$\begin{aligned} \hat{U}_t + |\xi|^2 \hat{U} &= -\hat{H} = -\delta \text{grad } \widehat{\Phi}(U) - \delta \widehat{\delta_{2p}} U \\ \hat{U}(\xi, 0) &= \hat{g}(\xi). \end{aligned}$$

Treating this as an ordinary differential equation for U and writing down the formal solution we obtain for each component, the equation

$$\hat{U}_k(\xi, t) = g_k(\xi) e^{-|\xi|^2 t} - \int_0^t \hat{H}_k(\xi, s) e^{-|\xi|^2(t-s)} ds$$

where,

$$\hat{H}_k(\xi, s) = -i \sum_{j=1}^n \xi_j (\text{grad } \widehat{\Phi}(U))_k - i \sum_{j=1}^n \xi_j |\xi|^{2p} \hat{U}_k.$$

This enables us to rewrite it in the form,

$$\hat{H}_k(\xi, s) = -i \sum_s \xi_s \cdot [a_k + \sum_j \xi_j^{2p} \hat{U}_k],$$

which implies that

$$\hat{H} = (\hat{H}_1, \dots, \hat{H}_n) = i \xi \cdot (\tilde{a}_k - \sum_j \xi_j^{2p} \tilde{U}_k)$$

where

$$\tilde{a}_k = (a_k, \dots, a_k)$$

and

$$\tilde{U}_k = (\hat{U}_k, \dots, \hat{U}_k)$$

By the hypothesis on the form of $\delta \text{grad } \Phi$ it follows that,

$$\hat{H}_k(\xi, s) = -i \sum_{j=1}^n \xi \cdot [a_k + |\xi|^{2p} \hat{U}_k].$$

Let $a_i^0 = a_i(0, t)$. Then a_i can be represented as

$$a_i = a_i^0 + \xi \cdot \nabla_\xi a_i(\bar{\xi}), \quad 0 \leq \bar{\xi} \leq \xi.$$

In the appendix (Theorem A1) we show that for $|\xi| \leq \delta_1$

$$(41) \quad |\nabla_\xi a_i(\xi, t)| \leq C(t)$$

where $C(t)$ denotes a constant independent of ξ but which depends on $|g|_2$, $|g|_{W_2}$, δ_1 and t . Therefore it is possible to write \hat{H}_k as

$$\hat{H}_k(\xi, s) = -i \sum_{j=1}^n \xi_j \cdot [a_k^0 + \xi \cdot \nabla_\xi a_k(\bar{\xi}) + |\xi|^{2p} \hat{U}_k].$$

This yields

$$\int_0^t \hat{H}_k(\xi, s) e^{-|\xi|^2(t-s)} ds = i \sum_j \xi_j \cdot \int_0^t a_k^0 ds + H.O.T.$$

where

$$H.O.T. = \int_0^t a_k^0 (e^{-|\xi|^2(t-s)} - 1) ds + i \xi \cdot \int_0^t [\nabla_\xi a_k + |\xi|^{2p} \hat{U}_k] ds.$$

Hence it follows that

$$\hat{U}_k(\xi, t) = -i\xi \cdot L_k(\xi, t_0) + (H.O.T.)_k$$

with

$$L_k(\xi, t_0) = \left(\int_0^{t_0} a_k^0 ds, \dots, \int_0^{t_0} a_k^0 ds \right),$$

and

$$(H.O.T.)_k = \hat{g}_k(\xi) e^{-|\xi|^2 t} + a_k^0 (e^{-|\xi|^2 (t-s)} - 1) + i\xi \cdot [\nabla_\xi a_k + |\xi|^{2p} \hat{U}_k],$$

where

$$\int_0^t a_k^0 = \int_0^t \Phi_{U_k} ds = \int_0^t U_k^{jk-1} ds.$$

To apply Theorem 6.1 we need a lower bound for α_k , for at least one k , where

$$\alpha_k(t) = \alpha_k = \int_{|\omega|=1} |\omega \cdot L_k(\omega(t))|^2 d\sigma,$$

and

$$L_k(\omega, t) = \left[\int_0^t a_k^0, \dots, \int_0^t a_k^0 \right].$$

Conditions (i), (ii) ensure that there exists a sequence $t_n \rightarrow \infty$ such that (making use of the notation introduced above), for some ξ_0

$$(42) \quad U_k(\xi_0, t_n) = \xi_0 \cdot L_k(t_n) + H.O.T.$$

where $L_k(t_n) = \text{Re } \widehat{\nabla U}(0) + i(P_k - \text{Im } \widehat{\nabla U}(0))$ and for some ξ_0

$$(43) \quad \xi_0 \cdot L_k(t_n) \geq \alpha > 0,$$

with α given by $\alpha = \max \{ |\text{Re } \widehat{\nabla U}(0)|, |P_k - \text{Im } \widehat{\nabla U}(0)| \}$, that is α is independent of t_n . Moreover, ξ_0 can be chosen to be of the form

$$\xi_0 = (0, \dots, 1, \dots, 0) = e_j$$

for $j = k$ so that (40) holds.

A) and B) above show that the class of data producing solutions which satisfy (42) and (43) is large. Moreover, note that there exists $N(\xi_0)$ independent of t_n such that $\xi \cdot L_k(t_n) \geq \frac{\alpha}{2}$ for $\xi \in N(\xi_0) \cap S^{n-1}$. This can be shown as follows. First we show that for $k - 1 > 2$, $|L_k(t)| \leq C$, which follows since,

$$\int_0^t \int |a_k^0| \leq C \int_0^t \int |U|^{k_i-1} \leq C \int_0^t \frac{1}{(s+1)^{\frac{n}{2}+1}} \leq C.$$

Let

$$\xi = (\varepsilon, \dots, \varepsilon, 1 - \sqrt{n-1}\varepsilon, \dots, \varepsilon).$$

and it follows that $|\xi| = 1$ and $|\xi - \xi_0| \leq \tilde{\varepsilon} = 2\sqrt{n-1}\varepsilon$. This implies that

$$(44) \quad \xi \cdot L_k(t_n) > \frac{\alpha}{2}.$$

Note that if $\text{Re } \widehat{\nabla U}_0 \neq 0$ then it follows that

$$|\xi \cdot L_k|^2 = |\text{Re } \widehat{\nabla U}_0|^2 + \left| \text{Im } \widehat{\nabla U}_0 - \int_0^t \bar{P}_k \right|^2 \geq |\text{Re } \widehat{\nabla U}_0|^2 \geq \alpha/2$$

Also note that,

$$\alpha(t_n) = \alpha_1 = \int_{|\omega|=1} |\xi \cdot L_k(t_n)|^2 \geq \frac{\alpha^2}{4} \omega_n = \rho$$

where ω_n denotes the volume of the unit sphere. By the above remarks

$$\widehat{U}(\xi, T) = \xi \cdot L(\xi, T) + H(\xi, T),$$

for $T = t_n$ and $|\xi L(\xi, T)| \geq C_0$ for all n . Recall that for $t \geq \delta_0$, (see Theorem 6.1) $\delta_0 = (\delta_0(\alpha_1(t_n)))$ where δ is defined by the requirement that $4\delta M_0 M_1 \leq \alpha_1$, since we can clearly choose $\delta_1 \leq \delta_0(t_n)$ for all n by letting $4\delta_1 M_0 M_1 = \rho$. (see Schonbek [4].) Then $V(x, t)$ the solution to the heat system with data U_0 satisfies

$$|V(\cdot, t)|_2^2 \geq \chi_\rho (t+1)^{-\left(\frac{n}{2}+1\right)}$$

for $t \geq t_k$, with $\chi_\rho = \frac{\alpha \omega_n e^{-1}}{2(n+2)}$ (cf. Schonbek [4]), which, by the computations above is independent of t_k .

Let v be the solution to the heat system with initial data $v(x, 0) = U(x, T_1)$ such that $K(1 + T_1)^{-\theta} \leq \frac{\chi_\rho}{8} \omega_0$, where $K = K_1 K_2^{\frac{4p}{n}-1} K_3$ and χ_ρ is as above. The constants K_1 , K_2 and K_3 are defined as below.

$$\begin{aligned} |\text{grad } \Phi| &\leq K_1 |U|_2^2 (|U|_\infty^{\frac{4p+1}{n}} + |U|_\infty^\nu) \\ |U|_2^2 &\leq K_3 (1+t)^{-\frac{n}{2}} \\ |U|_\infty &\leq K_2 (1+t)^{-\frac{n}{4}} \end{aligned}$$

and ω_0 denotes the surface area of S^{n-1} , and

$$\theta = \min \left\{ \frac{n}{4} + \frac{n}{2} - 1, p + \frac{n}{4} - 1 \right\}.$$

Thus by Theorem 6.1 it follows that for $t \geq \delta_1 = \delta_1(\rho)$

$$(45) \quad \chi_\rho (t+1)^{-\left(\frac{n}{2}+1\right)} \leq |v(\cdot, t)|_2^2 \leq K_0 (t+1)^{-\left(\frac{n}{2}+1\right)}$$

where K_0 depends on the L_2 norm of U_0 and χ_ρ depends on β and C_0 . δ_0 is independent of T_1 since δ depends on ρ by the uniformity condition (44).

Let

$$u(x, t) = U(x, t + T_1).$$

The difference $w = v - u$ is now studied. The decay rates for the KdVB system will imply that $|w(\cdot, t)| \leq C(t+1)^{-\left(\frac{n}{2}+1\right)}$ with C sufficiently small. W satisfies an inhomogeneous heat system. The Fourier splitting method will yield

$$\begin{aligned} (46) \quad \frac{d}{dt} \left[(t+1)^{8n} \int_{\mathbb{R}^n} |w|^2 dx \right] &\leq (t+1)^{8n-1} \int_{S(t)} |\hat{w}|^2 d\xi + K_* |D^{2p+1} v|_2 |u|_2 \\ &\quad + K_2 |\nabla v|_\infty \int_{\mathbb{R}^n} |u|^{\frac{4p}{n}+1} dx. \end{aligned}$$

The second and third terms on the right-hand side of the above inequality will be of higher order and will as such be easily bounded as follows:

$$(47) \quad \begin{aligned} K_2 |\nabla v|_\infty \int_{\mathbb{R}^n} |u|^{\frac{4p}{n}+1} dx &\leq M_1 (t+1)^{-\gamma_1} \\ K_* |D^{2p+1} v|_2 |u|_2 &\leq M_2 (t+1)^{-\gamma_2} \end{aligned}$$

where $\gamma_1 = \frac{n}{2} + p + \frac{1}{2} = \gamma_2$. In obtaining these bounds, we make use of the bounds on the derivatives of the solution to the heat equation and the L_∞ and L_2 bounds for the solution to the KdVB system. To bound the first term on the right-hand side of (46) the Fourier transform of the inhomogeneous heat equation yields

$$\hat{w} + |\xi|^2 \hat{w} = -\delta \widehat{\text{grad}} \Phi(u) - \delta \widehat{\delta_{2p}} u = -\hat{H}.$$

As before, in (37)

$$|\hat{H}(\xi, t)| \leq K_1 |\xi| (|u(\cdot, t)|_\infty^{\frac{4p}{n}-1} + |u(\cdot, t)|_\infty^{\nu-2}) |u(\cdot, t)|_2^2 + K_* |\xi|^{2p+1} |\hat{u}(\cdot, t)|_\infty.$$

Let $\nu = \min$ power of U_i in $\Phi(U)$. Therefore,

$$\begin{aligned} |\hat{w}(\xi, t)| &\leq \int_0^t |\hat{H}(\xi, s)| ds \\ &\leq K_1 |\xi| \int_0^t (|u(\cdot, s)|_\infty^{\nu-2} + |u(\cdot, s)|_\infty^{\frac{4p}{n}-1}) |u(\cdot, s)|_2^2 ds \\ &\quad + K_* |\xi|^{2p+1} \int_0^t |\hat{u}(\cdot, s)|_\infty ds \\ &= K_1 |\xi| \int_{T_1}^{t+T_1} (|U(\cdot, s)|_\infty^{\nu-2} + |U(\cdot, s)|_\infty^{\frac{4p}{n}-1}) |U(\cdot, s)|_2^2 ds \\ &\quad + K_2 |\xi|^{2p+1} \int_{T_1}^{t+T_1} |\hat{U}(\cdot, s)|_\infty ds \\ &\leq K_1 |\xi| \int_{T_1}^{t+T_1} \left(\frac{K_2^{\nu-2}}{(s+1)^{\frac{n}{4}\nu}} + \frac{K_1^{\frac{4p}{n}-1}}{(1+s)^{p+\frac{n}{4}}} \right) ds \\ &\quad + CK_* |\xi|^{2p+1} \int_{T_1}^{t+T_1} C^* ds \\ &\leq K_1 |\xi| \left(\frac{1}{(1+T_1)} \right)^\theta \omega_0 + CK_* |\xi|^{2p+1} t \end{aligned}$$

where $\theta = \min \left\{ \frac{n}{4} \nu - 1, p + \frac{n}{4} - 1 \right\}$. Since $n \geq 2$ and $p \geq 1$ we have $\theta > 0$. Here we have made use of the fact that the L_∞ -norm of the solution to the KdVB system decays at the rate of $(1 + t)^{-\frac{n}{4}}$.

Thus we have,

$$|\hat{w}(\xi, t) \leq C_1 |\xi| + C_2 |\xi|^{2p+1} t$$

where $C_1 = C_1(T_1) = \frac{CK}{(1 + T_1)^\theta}$ and $C_2 = CK_*$. Note that T_1 can be chosen as large as needed since χ_ρ is independent of T_1 . Using the definition of the set $S(t)$, this implies that

$$\begin{aligned} \int_{S(t)} |\hat{w}|^2 d\xi &\leq C_1 \frac{\omega_0}{n} \int_{S(t)} |\xi|^2 d\xi + C_2 t^2 \int_{S(t)} |\xi|^{4p+2} d\xi \\ &\leq C_1 (t + 1)^{-\left(\frac{n}{2} + 1\right)} + C_2 (t + 1)^{-\left(\frac{n}{2} + 2p - 1\right)} \end{aligned}$$

Hence the first term on the right-hand side of (46) can be bounded as follows.

$$\begin{aligned} (t + 1)^{8n-1} \int_{S(t)} |\hat{w}|^2 d\xi &\leq C_1 (t + 1)^{8n - \frac{n}{2} - 2} + C_2 (t + 1)^{8n - \frac{n}{2} - 2p} \\ &\leq \frac{\chi_\rho}{8} (t + 1)^{\frac{15n}{2} - 2} + C_2 (t + 1)^{\frac{15n}{2} - 2p}. \end{aligned}$$

The last inequality follows from the choice of T_1 . Combining this last estimate with (46) and (47) yields

$$\begin{aligned} \frac{d}{dt} \left[(t + 1)^{8n} \int_{\mathbb{R}^n} |w|^2 dx \right] &\leq \frac{\chi_\rho}{8} (t + 1)^{\frac{15n}{2} - 2} \\ &\quad + C_2 (t + 1)^{\frac{15n}{2} - 2} + M_1 (t + 1)^{(8n-1) - \gamma_1} \\ &\quad + M_2 (t + 1)^{(8n-1) - \gamma_2}. \end{aligned}$$

Integrating over the interval $[\delta_1(\rho), t]$ gives

$$\begin{aligned} \int_{\mathbb{R}^n} |w|^2 dx &\leq \frac{\chi_\rho}{8} (t + 1)^{-\left(\frac{n}{2} + 1\right)} \\ &\quad + C_2 (t + 1)^{-\left(\frac{n}{2} + 2p - 1\right)} + M_1 (t + 1)^{-\left(\frac{n}{2} + p + \frac{1}{2}\right)} \\ &\quad + M_2 (t + 1)^{-\left(\frac{n}{2} + p + \frac{1}{2}\right)} + (t + 1)^{-8n} \int_{\mathbb{R}^n} |w(x, \delta_1)|^2 dx \end{aligned}$$

Note that

$$\int_{\mathbb{R}^n} |w(x, \delta_1)|^2 dx \leq \int_{\mathbb{R}^n} |u(x, 0)|^2 dx + \int_{\mathbb{R}^n} |v(x, 0)|^2 dx \leq 2 \int_{\mathbb{R}^n} |U_0|^2 dx.$$

Hence, it follows that

$$\int_{\mathbb{R}^n} |w|^2 \leq \frac{\chi_\rho}{8} (t+1)^{-\mu} + H.O.T.$$

for t large enough, where μ is defined by

$$\mu = \frac{n}{2} + 1.$$

The rest of the terms are of higher order since $2p > n$, $\frac{n}{2} + 1 < \frac{n}{2} + p + \frac{1}{2}$ and $\frac{n}{2} + 1 < \frac{n}{2} + 2p - 1$. That is for $t \geq T_1$

$$\begin{aligned} |u(\cdot, t)|_2 &\geq |v(\cdot, t)|_2 - |w(\cdot, t)|_2 \\ &\geq \chi_\rho (t+1)^{-(\frac{n}{2}+1)} - \frac{\chi_\rho}{8} (t+1)^{-(\frac{n}{2}+1)} + H.O.T \end{aligned}$$

and T_2 is such that

$$H.O.T. < \frac{\chi_\rho}{8} (t+1)^{-\frac{n}{2}-1}.$$

Hence, for $t \geq T_3 = T_1 + T_2$

$$|u(\cdot, t)|_2 \geq \frac{\chi_\rho}{4} (t+1)^{-\frac{1}{2}(\frac{n}{2}+1)}$$

For $t < T_3$ the decay of energy of U yields

$$|u(\cdot, t)|_2^2 \geq |u(\cdot, T_3)|_2^2 \geq \chi_\rho/4 \left[\frac{1+t}{1+T_3} \right]^{\frac{n}{2}+1} (1+t)^{-(\frac{n}{2}+1)}$$

and the result follows. This completes the proof.

THEOREM 6.3. — *Let $U_0 \in L_1 \cap H^{p+1} \cap \mathcal{M}_I \cap \mathcal{M}_R$. Let U be a solution of the KdVB system with data U_0 where U_0 is such that $\int U_0 = 0$ and \hat{U}_0 has a zero of order > 1 . If $|P_k| \neq 0$ for some k then, there exists constants K_1, K_2 such that*

$$K_1 (t+1)^{-(\frac{n}{2}+1)} \leq |U(\cdot, t)|_2^2 \leq K_2 (t+1)^{-(\frac{n}{2}+1)}$$

where K_1, K_2 depend only on the norms of the data.

Proof. – The proof is similar to the case when the data is of order one. Note that if $U_0 \in \mathcal{M}_{\mathcal{I}} \cap \mathcal{M}_{\mathcal{R}}$ we only need

$$\begin{aligned} \Phi(U) &= \sum \alpha_{k_i} U_i^{k_i} \quad k_i \text{ odd or} \\ \Phi(U) &= \sum \alpha_{k_i} U_i^{k_i} \quad k_i \text{ even,} \end{aligned}$$

Then, $P_k \neq 0$. As in the proof of Theorem 6.6, there exists a sequence of t_n such that

$$\hat{U}(\xi, t) = \xi \cdot L(t_n) + H.O.T.$$

This implies the uniform condition for the lower bounds,

$$\xi \cdot L_k(t_n) \geq \alpha/2 \quad \text{for } \xi \in N(\xi_0) \cap S^{k-1}$$

for some ξ_0 , and the proof now is a repetition of the proof of Theorem 6.2.

7. APPENDIX

THEOREM A1. – Let $U_0 \in H^{p+1} \cap W_2$. Then if U is the solution to the KdVB system with data U_0 then,

$$\nabla_{\xi} a_i(\bar{\xi}) \leq C(t)$$

where $C(t)$ depends only on the norm of U_0 in the spaces H^{p+1} and W_2 .

Proof. – By definition,

$$a_i = \hat{\Phi}_{U_i}$$

and

$$\hat{\Phi}_{U_i} = \left(\sum_i \sum_k m_{k_i} U_i^{k_i} \right)_{U_i} = \sum m_{k_i} U_i^{k_i-1}.$$

We carry out the analysis of one term of the sum. Note that $k > 2$ since otherwise the equation is linear. Thus, for $k > 2$

$$|\nabla_{\xi_k} a_i| \leq C \int |U_i^{k_j-1}| |x_k| \leq C \int U_i^2 |x_k|.$$

When $k_j > 2$ it suffices to bound $\int U_i^2 |x| dx$. We have,

$$\begin{aligned}
 (48) \quad \frac{d}{dt} \int U_i^2 |x_k| &= \int U_i |x_k| U_{it} \\
 &= \int |x_k| U_i (\delta \operatorname{grad} \Phi(U))_i \\
 &\quad + \int |x_k| U_i \delta_{2p+1} U_i - \int |x_k| U_i (\Delta U)_i \\
 &= I + II + III.
 \end{aligned}$$

Note that

$$\begin{aligned}
 I &= \int |x_k| U_i (\delta \operatorname{grad} \Phi)_i = \int \operatorname{sgn}(x_k) U_i U_i^{k_i-1} \\
 &\quad + \sum_j \int |x_k| U_i^{k_i-1} U_i x_j \\
 &\leq C_0 \int U_i^2 + \frac{1}{k} \int \operatorname{sgn}(x_k) U_i^{k_j} \\
 &\leq C_0 + \frac{C_0}{k} \int |U_i|^2
 \end{aligned}$$

and

$$\begin{aligned}
 II &= \int |x_k| U_i \delta_{2p+1} U_i = - \int \operatorname{sgn}(x_k) U_i \delta_{2p} U_i \\
 &\quad - \int |x_k| \delta U_i \delta_{2p} U_i \\
 &= - \int \operatorname{sgn}(x_k) U_i \delta_{2p} U_i + \int \operatorname{sgn}(x_k) \delta_{2p-1} U_i \\
 &\quad + \int |x_k| \delta_2 U_i \delta_{2p-1} U_i
 \end{aligned}$$

Hence it follows that

$$\begin{aligned}
 (49) \quad II &= \int \operatorname{sgn}(x_k) \sum_{k < 2p+1} \alpha_k \delta_{2p+1-k} U_i \delta_k U_i \\
 &\quad + (-1)^{2p+1} \int |x_k| \delta_{2p+1} U_i U_i
 \end{aligned}$$

and we can conclude that

$$(50) \quad 2II = \int \operatorname{sgn}(x_k) \sum_{k < 2p+1} \alpha_k \delta_{2p+1-k} U_i \delta_k U_i \\ \leq \alpha_k \left(\int |D^s U_i|^2 \right)^{\frac{1}{2}} \left(\int |D^j U_i|^2 \right)^{\frac{1}{2}} \leq C$$

where $s, j \leq 2p + 1$. Finally,

$$III = \int |x_k| U_i \Delta U_i = - \sum_j \int |x_k| U_{ix_j}^2 + \int \operatorname{sgn}(x_k) U_i \nabla U \\ \leq \left(\int U_i^2 \right)^{\frac{1}{2}} \left(\int |\nabla U_i|^2 \right)^{\frac{1}{2}} \\ \leq C$$

Combining these estimates in (48) we obtain

$$\frac{d}{dt} \int U_i^2 |x_k| \leq C.$$

Integrating this with respect to time the desired result follows.

REFERENCES

- [1] C. J. AMICK, J. L. BONA and M. E. SCHONBEK, Decay of solutions of some nonlinear wave-equations, *J. Differential Equations*, Vol. **81**, 1989, pp. 1-49.
- [2] M. E. SCHONBEK, Decay of parabolic conservation laws, *Comm. in Partial Differential Equations*, Vol. **7** (1), 1980, pp. 449-473.
- [3] M. E. SCHONBEK, Large time behaviour of the solution to the Navier-Stokes equations, *Comm. in Partial Differential Equations*, Vol. **11** (7), 1986, pp. 733-763.
- [4] M. E. SCHONBEK, Lower bounds on the rates of decay of the solutions to the Navier-Stokes equations, *J. of the American Math. Soc.*, Vol. **4**, # 3, July 1991.
- [5] M. E. SCHONBEK, T. SCHONBEK and E. SÜLLI, Large time behaviour of solutions to the Magneto-Hydrodynamics equation, Preprint.
- [6] M. WIEGNER, Decay results for weak solutions of the Navier-Stokes equation on \mathbb{R}^n , *J. London, Math. Soc.* Vol. **35** (2), 1987, pp. 303-313.
- [7] Z. YULIN and GUO BOLING, The periodic boundary-value problems and the initial value problems of the generalised Korteweg-deVries equation, *Acta Mathematica Sinica*, Vol. **27**, 1984, pp. 154-176.
- [8] Zhang LINGHAI, Decay of solutions of higher order multi-dimensional non-linear KdV-Burgers system, To appear in *the Proc. of the Royal Soc. of Edinburgh*.

(Manuscript received January 31, 1994;
Revised version received September 7, 1994.)