

THE CRITICAL VALUES OF CERTAIN DIRICHLET SERIES

GORO SHIMURA

Received: May 27, 2008

Communicated by Don Blasius

ABSTRACT. We investigate the values of several types of Dirichlet series $D(s)$ for certain integer values of s , and give explicit formulas for the value $D(s)$ in many cases. The easiest types of D are Dirichlet L -functions and their variations; a somewhat more complex case involves elliptic functions. There is one new type that includes $\sum_{n=1}^{\infty} (n^2+1)^{-s}$ for which such values have not been studied previously.

2000 Mathematics Subject Classification: 11B68, 11M06, 30B50, 33E05.

INTRODUCTION

By a Dirichlet character modulo a positive integer d we mean as usual a \mathbf{C} -valued function χ on \mathbf{Z} such that $\chi(x) = 0$ if x is not prime to d , and χ induces a character on $(\mathbf{Z}/d\mathbf{Z})^\times$. In this paper we always assume that χ is primitive and nontrivial, and so $d > 1$. For such a χ we put

$$(0.1) \quad L(s, \chi) = \sum_{n=1}^{\infty} \chi(n)n^{-s}.$$

It is well known that if k is a positive integer such that $\chi(-1) = (-1)^k$, then $L(k, \chi)$ is π^k times an algebraic number, or equivalently, $L(1-k, \chi)$ is an algebraic number. In fact, there is a well-known formula, first proved by Hecke in [3]:

$$(0.2) \quad kd^{1-k}L(1-k, \chi) = -\sum_{a=1}^{d-1} \chi(a)B_k(a/d),$$

where $B_k(t)$ is the Bernoulli polynomial of degree k . Actually Hecke gave the result in terms of $L(k, \chi)$, but here we state it in the above form. Hecke's proof is based on a classical formula

$$(0.3) \quad B_k(t) = -k!(2\pi i)^{-k} \sum_{0 \neq h \in \mathbf{Z}} h^{-k} \mathbf{e}(ht) \quad (0 < k \in \mathbf{Z}, 0 < t < 1).$$

There is also a well-known proof of (0.2), which is essentially the functional equation of $L(s, \chi)$ combined with a proof of (0.3). We will not discuss it in the present paper, as it is not particularly inspiring.

In [9] we gave many formulas for $L(1-k, \chi)$ different from (0.2). The primary purpose of the present paper is to give elementary proofs for some of them, as well as (0.2), and discuss similar values of a few more types of Dirichlet series. The point of our new proofs can be condensed to the following statement: *We find infinite sum expressions for $L(s, \chi)$, which are valid for all $s \in \mathbf{C}$ and so can be evaluated at $s = 1 - k$, whereas the old proof of Hecke and our proofs in [9] employ calculations at $s = k$ and involve the Gauss sum of $\bar{\chi}$.*

To make our exposition smooth we put

$$(0.4) \quad \mathbf{e}(z) = \exp(2\pi iz) \quad (z \in \mathbf{C}),$$

$$(0.5) \quad H = \{z \in \mathbf{C} \mid \operatorname{Im}(z) > 0\}.$$

The three additional types of Dirichlet series we consider naturally involve a complex variable s , and are defined as follows:

$$(0.6) \quad D^\nu(s; a, p) = \sum_{-a \neq n \in \mathbf{Z}} (n+a)^\nu |n+a|^{-\nu-s} \mathbf{e}(p(n+a)),$$

where $a \in \mathbf{R}$, $p \in \mathbf{R}$, and $\nu = 0$ or 1 ,

$$(0.7) \quad \mathcal{L}_k(s, z) = \sum_{m \in \mathbf{Z}} \mathbf{e}(mr) (z+m)^{-k} |z+m|^{-2s} \quad (k \in \mathbf{Z}, r \in \mathbf{Q}, z \in H),$$

$$(0.8) \quad \varphi_\nu(u, s; L) = \sum_{\alpha \in L} (u+\alpha)^{-\nu} |u+\alpha|^{\nu-2s},$$

where L is a lattice in \mathbf{C} , $0 \leq \nu \in \mathbf{Z}$, and $u \in \mathbf{C}$, $u \notin L$. We should also note

$$(0.9) \quad \mathfrak{E}(z, s) = \operatorname{Im}(z)^s \sum_{(m, n)} (mz+n)^{-k} |mz+n|^{-2s} \quad (0 \leq k \in \mathbf{Z}, z \in H),$$

where (m, n) runs over the nonzero elements of \mathbf{Z}^2 . The value $\mathfrak{E}(z, \mu)$ for an integer μ such that $1-k \leq \mu \leq 0$ was already discussed in [9], and so it is not the main object of study in this paper, but we mention it because (0.8) is a natural analogue of (0.9). We will determine in Section 3 the value $\varphi_\nu(u, \kappa/2; L)$ for an integer κ such that $2-\nu \leq \kappa \leq \nu$, which may be called a *nearly holomorphic elliptic function*. Now (0.6) is closely connected with $L(s, \chi)$. In [9, Theorem 4.2] we showed that $D^\nu(k; a, p)$ for $0 < k \in \mathbf{Z}$ is elementary factors times the value of a generalized Euler polynomial $E_{c, k-1}(t)$ at $t = p$. In Section 2 we will reformulate this in terms of $D^\nu(1-k; a, p)$. Finally, the nature of the series of (0.7) is quite different from the other types. We will show in Section 4 that $i^k \mathcal{L}_k(\beta, z)$ is a \mathbf{Q} -rational expression in π , $\mathbf{e}(z/N)$, and $\operatorname{Im}(z)$, if $\beta \in \mathbf{Z}$ and $-k < \beta \leq 0$, where N is the smallest positive integer such that $Nr \in \mathbf{Z}$. Similar results will also be given under other conditions on β . In the final section we will make some comments in the case where the base field is an algebraic number field.

1. $L(1 - k, \chi)$

1.1. We start with an elementary proof of (0.2). Strange as it may sound, the main idea is the binomial theorem. We first note

$$(1.1) \quad B_n(t) = \sum_{\nu=0}^n \binom{n}{\nu} B_\nu t^{n-\nu} \quad (0 \leq n \in \mathbf{Z}),$$

$$(1.2a) \quad B_0 = 1, \quad \zeta(0) = -1/2 = B_1,$$

$$(1.2b) \quad n\zeta(1-n) = -B_n \quad (1 < n \in \mathbf{Z}),$$

where B_n is the n th Bernoulli number. Formulas (1.1) and (1.2a) are well-known; (1.2b) is usually given only for even n , but actually true also for odd n , since $\zeta(-2m) = 0 = B_{2m+1}$ for $0 < m \in \mathbf{Z}$.

To prove (0.2), we first make a trivial calculation:

$$\begin{aligned} L(s, \chi) - \sum_{a=1}^{d-1} \chi(a) a^{-s} &= \sum_{m=1}^{\infty} \sum_{a=1}^{d-1} \chi(dm+a) (dm+a)^{-s} \\ &= \sum_{m=1}^{\infty} \sum_{a=1}^{d-1} \chi(a) (dm)^{-s} \left(1 + \frac{a}{dm}\right)^{-s}. \end{aligned}$$

Now we apply the binomial theorem to $(1+X)^{-s}$. Thus the last double sum equals

$$(1.3) \quad \begin{aligned} &\sum_{m=1}^{\infty} \sum_{a=1}^{d-1} \chi(a) (dm)^{-s} \sum_{r=0}^{\infty} \binom{-s}{r} \left(\frac{a}{dm}\right)^r \\ &= \sum_{r=0}^{\infty} \binom{-s}{r} \sum_{m=1}^{\infty} m^{-s-r} d^{-s} \sum_{a=1}^{d-1} \chi(a) (a/d)^r, \end{aligned}$$

where

$$\binom{\tau}{r} = \frac{\tau(\tau-1)\dots(\tau-r+1)}{r!},$$

which is of course understood to be 1 if $r=0$. So far our calculation is formal, but can be justified at least for $\operatorname{Re}(s) > 1$. Indeed, put $\operatorname{Re}(s) = \sigma$ and $|s| = \alpha$. Then

$$(1.4) \quad \left| \binom{-s}{r} \right| \leq \frac{\alpha(\alpha+1)\dots(\alpha+r-1)}{r!} = (-1)^r \binom{-\alpha}{r}.$$

Therefore the triple sum obtained from (1.3) by taking the absolute value of each term is majorized by

$$\begin{aligned} &\sum_{a=1}^{d-1} \sum_{r=0}^{\infty} \sum_{m=1}^{\infty} m^{-\sigma-r} d^{-\sigma} \binom{-\alpha}{r} \left(\frac{a}{d}\right)^r \\ &\leq \zeta(\sigma) d^{-\sigma} \sum_{a=1}^{d-1} \sum_{r=0}^{\infty} \binom{-\alpha}{r} \left(\frac{a}{d}\right)^r = \zeta(\sigma) d^{-\sigma} \sum_{a=1}^{d-1} \left(1 - \frac{a}{d}\right)^{-\alpha} \end{aligned}$$

if $\sigma > 1$. Thus, for $\operatorname{Re}(s) > 1$, (1.3) can be justified, and so

$$(1.5) \quad L(s, \chi) - \sum_{a=1}^{d-1} \chi(a)a^{-s} = \sum_{r=0}^{\infty} \binom{-s}{r} \zeta(s+r)d^{-s} \sum_{a=1}^{d-1} \chi(a)(a/d)^r.$$

We can show that the last sum $\sum_{r=0}^{\infty}$ defines a meromorphic function in s on the whole \mathbf{C} . For that purpose given $s \in \mathbf{C}$, take a positive integer μ so that $\operatorname{Re}(s) > -\mu$ and decompose the sum into $\sum_{r=0}^{\mu+1}$ and $\sum_{r=\mu+2}^{\infty}$. There is no problem about the first sum, as it is finite. As for the latter, we have $|\zeta(s+r)| \leq \zeta(2)$ for $r \geq \mu+2$. Putting $\varepsilon = (d-1)/d$, we have $|\sum_{a=1}^{d-1} \chi(a)(a/d)^r| \leq (d-1)\varepsilon^r$. Therefore for $\operatorname{Re}(s) > -\mu$ the infinite sum $\sum_{r=\mu+2}^{\infty}$ can be majorized by

$$d^\mu(d-1)\zeta(2) \sum_{r=0}^{\infty} \binom{-\alpha}{r} (-\varepsilon)^r = d^\mu(d-1)\zeta(2)(1-\varepsilon)^{-\alpha}.$$

This proves the desired meromorphy of the right-hand side of (1.5).

Now, for $0 < k \in \mathbf{Z}$, without assuming that $\chi(-1) = (-1)^k$, we evaluate (1.5) at $s = 1 - k$. We easily see that $\binom{-s}{r} = 0$ and $\zeta(s+r)$ is finite for $s = 1 - k$ if $r > k$. We have to be careful about the term for $r = k$, as $\zeta(s+k)$ has a pole at $s = 1 - k$. Since

$$(1.6) \quad \lim_{s \rightarrow 1-k} \binom{-s}{k} \zeta(s+k) = \lim_{s \rightarrow 1-k} \binom{-s}{k} \frac{1}{s-1+k} = \frac{-1}{k},$$

the term for $r = k$ at $s = 1 - k$ produces $-(d^{k-1}/k) \sum_{a=1}^{d-1} \chi(a)(a/d)^k$. Thus the evaluation of (1.5) at $s = 1 - k$ gives

$$(1.7) \quad kd^{1-k}L(1-k, \chi) = k \sum_{a=1}^{d-1} \chi(a)(a/d)^{k-1} - \sum_{a=1}^{d-1} \chi(a)(a/d)^k + \sum_{r=0}^{k-1} k \binom{k-1}{r} \zeta(1+r-k) \sum_{a=1}^{d-1} \chi(a)(a/d)^r.$$

By (1.2b) we have, for $0 \leq r < k-1$,

$$k \binom{k-1}{r} \zeta(1+r-k) = \frac{-k}{k-r} \binom{k-1}{r} B_{k-r} = -\binom{k}{r} B_{k-r}.$$

The term for $r = k-1$ produces $k\zeta(0) \sum_{a=1}^{d-1} \chi(a)(a/d)^{k-1}$, which combined with the first term on the right-hand side of (1.7) gives $-kB_1 \sum_{a=1}^{d-1} \chi(a)(a/d)^{k-1}$. Thus we obtain

$$kd^{1-k}L(1-k, \chi) = -\sum_{r=0}^k \binom{k}{r} B_{k-r} \sum_{a=1}^{d-1} \chi(a)(a/d)^r,$$

which together with (1.1) proves (0.2). Notice that we did not assume that $\chi(-1) = (-1)^k$, and so we proved (0.2) for every positive integer k . If $\chi(-1) = (-1)^{k-1}$, we have $L(1-k, \chi) = 0$, which means that the right-hand side of (0.2) is 0 if $\chi(-1) = (-1)^{k-1}$. This can be proved more directly; see [9, (4.28)].

In the above calculation the term for $r = 0$ actually vanishes, as $\sum_{a=1}^{d-1} \chi(a) = 0$. However, we included the term for the following reason. In later subsections we will consider similar infinite sums with r ranging from 0 to ∞ , of which the terms for $r = 0$ are not necessarily zero.

1.2. By the same technique as in §1.1 (that is, employing the binomial theorem) we will express $L(1 - k, \chi)$ explicitly in terms of a polynomial Φ_{k-1} of degree $k - 1$. Writing n for $k - 1$, the polynomial is defined by

$$(1.8) \quad \Phi_n(t) = t^n - \sum_{\nu=1}^{\lfloor (n+1)/2 \rfloor} \binom{n}{2\nu-1} (2^{2\nu}-1) \frac{B_{2\nu}}{\nu} t^{n+1-2\nu} \quad (0 \leq n \in \mathbf{Z}),$$

where B_ν denotes the Bernoulli number as before. We understand that $\Phi_0(t) = 1$. We will eventually show that Φ_n is the classical Euler polynomial of degree n , but we prove Theorem 1.4 below with this definition of Φ_n , with no knowledge of the Euler polynomial. We first prove:

LEMMA 1.3. *Let χ be a primitive Dirichlet character of conductor $4d_0$ with $0 < d_0 \in \mathbf{Z}$. Then $\chi(a - 2d_0) = -\chi(a)$ for every $a \in \mathbf{Z}$.*

Proof. We may assume that a is prime to $2d_0$, as the desired equality is trivial otherwise. Then we can find an integer b such that $ab - 1 \in 4d_0\mathbf{Z}$, and we have $\chi(a - 2d_0) = \chi(a)\chi(1 - 2d_0b)$. Since $(1 - 2d_0b)^2 - 1 \in 4d_0\mathbf{Z}$, we have $\chi(1 - 2d_0b) = \pm 1$. Suppose $\chi(1 - 2d_0b) = 1$; let $x = 1 - 2d_0y$ with $y \in \mathbf{Z}$. Then $x^b - (1 - 2d_0b)^y \in 4d_0\mathbf{Z}$, and so $\chi(x)^b = 1$. Thus $\chi(x) = 1$, as b is odd. This shows that the conductor of χ is a divisor of $2d_0$, a contradiction. Therefore $\chi(1 - 2d_0b) = -1$, which proves the desired fact.

THEOREM 1.4. *Let χ be a nontrivial primitive Dirichlet character modulo d , and let k be a positive integer such that $\chi(-1) = (-1)^k$.*

(i) *If $d = 2q + 1$ with $0 < q \in \mathbf{Z}$, then*

$$(1.9) \quad L(1 - k, \chi) = \frac{d^{k-1}}{2^k \chi(2) - 1} \sum_{b=1}^q (-1)^b \chi(b) \Phi_{k-1}(b/d).$$

(ii) *If $d = 4d_0$ with $1 < d_0 \in \mathbf{Z}$, then*

$$(1.10) \quad L(1 - k, \chi) = (2d_0)^{k-1} \sum_{a=1}^{d_0-1} \chi(a) \Phi_{k-1}(2a/d).$$

Before proving these, we note that these formulas are *better than* (0.2) in the sense that $\Phi_{k-1}(t)$ is of degree $k - 1$, whereas $B_k(t)$ is of degree k .

Proof. We first put

$$Z(s) = \sum_{n=1}^{\infty} (-1)^n n^{-s}, \quad \Lambda(s) = \sum_{n=1}^{\infty} (-1)^n \chi(n) n^{-s}.$$

We easily see that

$$\Lambda(s) + L(s, \chi) = 2 \sum_{n=1}^{\infty} \chi(2n)(2n)^{-s} = \chi(2)2^{1-s}L(s, \chi),$$

and a similar equality holds for $Z(s)$. Thus

$$Z(s) = \zeta(s)(2^{1-s} - 1), \quad \Lambda(s) = L(s, \chi)\{\chi(2)2^{1-s} - 1\}.$$

We prove (i) by computing $\Lambda(1-k)$ for a given k in the same elementary way as we did in §1.1. With q as in (i) we observe that every positive integer m not divisible by d can be written uniquely $m = nd + a$ with $0 \leq n \in \mathbf{Z}$ or $m = nd - a$ with $0 < n \in \mathbf{Z}$, where in either case a is in the range $0 < a \leq q$. Therefore

$$\begin{aligned} \Lambda(s) &= \sum_{a=1}^q (-1)^a \chi(a) a^{-s} \\ &+ \sum_{a=1}^q \sum_{n=1}^{\infty} \left\{ (-1)^{nd+a} \chi(nd+a)(nd+a)^{-s} + (-1)^{nd-a} \chi(nd-a)(nd-a)^{-s} \right\}. \end{aligned}$$

The last double sum can be written

$$\sum_{a=1}^q \sum_{n=1}^{\infty} (-1)^{n+a} d^{-s} n^{-s} \left\{ \chi(a) \left(1 + \frac{a}{nd}\right)^{-s} + \chi(-a) \left(1 - \frac{a}{nd}\right)^{-s} \right\}.$$

Applying the binomial theorem to $(1 \pm X)^{-s}$, we obtain

$$\begin{aligned} \Lambda(s) &- \sum_{a=1}^q (-1)^a \chi(a) a^{-s} \\ &= \sum_{n=1}^{\infty} \sum_{r=0}^{\infty} (-1)^n (dn)^{-r-s} \binom{-s}{r} \{1 + (-1)^{r+k}\} \sum_{a=1}^q (-1)^a \chi(a) a^r \\ &= \sum_{r=0}^{\infty} d^{-s} \binom{-s}{r} Z(s+r) \{1 + (-1)^{r+k}\} \sum_{a=1}^q (-1)^a \chi(a) (a/d)^r. \end{aligned}$$

By the same technique as in §1.1, we can justify this for $\operatorname{Re}(s) > 1$. We can even show that the last sum $\sum_{r=0}^{\infty}$ is absolutely convergent for every $s \in \mathbf{C}$ as follows. We first note that Z is an entire function. Take a positive integer μ and s so that $\operatorname{Re}(s) > -\mu$. Then for $r \geq \mu+2$ we have $|Z(s+r)| \leq \zeta(2)$. Put $|s| = \alpha$. We have also $|\sum_{a=1}^q (-1)^a \chi(a) (a/d)^r| \leq 2^{-r} q$. Therefore for $\operatorname{Re}(s) > -\mu$ the infinite sum $\sum_{r=\mu+2}^{\infty}$ can be majorized by

$$2d^\mu \zeta(2) q \sum_{r=0}^{\infty} \binom{-\alpha}{r} (-2)^{-r} = 2d^\mu \zeta(2) q (1 - 2^{-1})^{-\alpha}.$$

This proves the desired convergence of $\sum_{r=0}^{\infty}$. Substituting $1-k$ for s in the above equality, we obtain $\Lambda(1-k)$ as an infinite sum, which is actually a finite sum, because $\binom{k-1}{r} = 0$ if $r \geq k$. (This time, the term $r = k$ causes no problem.) Also, we need only those r such that $k-r \in 2\mathbf{Z}$. Putting $k-r = 2\nu$, we find that

$$d^{1-k}L(1-k, \chi)\{\chi(2)^{2^k}-1\} = d^{1-k}\Lambda(1-k) \\ = \sum_{a=1}^q (-1)^a \chi(a)(a/d)^{k-1} + 2 \sum_{\nu=1}^{\lfloor k/2 \rfloor} \binom{k-1}{2\nu-1} Z(1-2\nu) \sum_{a=1}^q (-1)^a \chi(a)(a/d)^{k-2\nu}.$$

From (1.2b) we obtain

$$(1.11) \quad 2\nu Z(1-2\nu) = 2\nu(2^{2\nu}-1)\zeta(1-2\nu) = (1-2^{2\nu})B_{2\nu}.$$

Using this expression for $Z(1-2\nu)$, we obtain a formula for $L(1-k, \chi)$. Then comparison of it with our definition of Φ_n proves (1.9).

Next, let $d = 4d_0$ with $1 < d_0 \in \mathbf{Z}$ as in (ii). Observe that the set of all positive integers greater than d_0 and not divisible by d_0 is the disjoint union of the sets

$$\{4\nu d_0 \pm a \mid 0 < a < d_0, 0 < \nu \in \mathbf{Z}\} \sqcup \{(4\nu+2)d_0 \pm a \mid 0 < a < d_0, 0 \leq \nu \in \mathbf{Z}\}.$$

Clearly $\chi(4\nu d_0 \pm a) = \chi(\pm a)$; also $\chi((4\nu+2)d_0 \pm a) = -\chi(\pm a)$ by Lemma 1.3. Therefore we have

$$L(s, \chi) = \sum_{a=1}^{d_0-1} \chi(a)a^{-s} \\ + \sum_{\nu=1}^{\infty} \sum_{a=1}^{d_0-1} \{\chi(a)(4\nu d_0 + a)^{-s} + \chi(-a)(4\nu d_0 - a)^{-s}\} \\ - \sum_{\nu=0}^{\infty} \sum_{a=1}^{d_0-1} \{\chi(a)((4\nu+2)d_0 + a)^{-s} + \chi(-a)((4\nu+2)d_0 - a)^{-s}\}.$$

Employing the binomial theorem in the same manner as before, we have

$$L(s, \chi) - \sum_{a=1}^{d_0-1} \chi(a)a^{-s} \\ = \sum_{\nu=1}^{\infty} \sum_{r=0}^{\infty} \binom{-s}{r} (4\nu d_0)^{-s-r} \{1 + (-1)^{k+r}\} \sum_{a=1}^{d_0-1} \chi(a)a^r \\ - \sum_{\nu=0}^{\infty} \sum_{r=0}^{\infty} \binom{-s}{r} ((4\nu+2)d_0)^{-s-r} \{1 + (-1)^{k+r}\} \sum_{a=1}^{d_0-1} \chi(a)a^r.$$

Notice that $\sum_{\nu=1}^{\infty} (4\nu)^{-s} - \sum_{\nu=0}^{\infty} (4\nu+2)^{-s} = 2^{-s}Z(s)$. Therefore

$$L(s, \chi) = \sum_{a=1}^{d_0-1} \chi(a)a^{-s} + \sum_{r=0}^{\infty} \binom{-s}{r} (2d_0)^{-s-r} Z(s+r) \{1 + (-1)^{k+r}\} \sum_{a=1}^{d_0-1} \chi(a)a^r.$$

The validity of this formula for all $s \in \mathbf{C}$ can be proved in the same way as in the previous case. The last infinite sum $\sum_{r=0}^{\infty}$ evaluated at $s = 1 - k$ becomes a finite sum $\sum_{r=0}^{k-1}$, which is actually extended only over those r such that $k - r = 2\nu$ with $\nu \in \mathbf{Z}$. Therefore, using (1.11), we obtain (1.10).

1.5. Let us now show that Φ_n coincides with the classical Euler polynomial. In [9, (4.2)] we defined polynomials $E_{c,n}(t)$ for $c = -\mathbf{e}(\alpha)$ with $\alpha \in \mathbf{R}$, $\notin \mathbf{Z}$, by

$$(1.12) \quad \frac{(1+c)e^{tz}}{e^z+c} = \sum_{n=0}^{\infty} \frac{E_{c,n}(t)}{n!} z^n.$$

If $c = 1$, the polynomial $E_{1,n}(t)$ is the classical Euler polynomial of degree n . Our task is to prove

$$(1.13) \quad E_{1,n} = \Phi_n.$$

We first note here some basic formulas:

$$(1.14) \quad E_{c,n}(t) = (1+c^{-1})n!(2\pi i)^{-n-1} \sum_{h \in \mathbf{Z}} (h+\alpha)^{-n-1} \mathbf{e}((h+\alpha)t)$$

$$(c = -\mathbf{e}(\alpha), \alpha \in \mathbf{R}, \notin \mathbf{Z}; 0 < t < 1 \text{ if } n = 0; 0 \leq t \leq 1 \text{ if } 0 < n \in \mathbf{Z}),$$

$$(1.15) \quad E_{c,n}(t+r) = \sum_{k=0}^n \binom{n}{k} E_{c,k}(r) t^{n-k} \quad (0 \leq n \in \mathbf{Z}),$$

$$(1.16) \quad E_{1,0}(0) = 1, \quad E_{1,n}(0) = 2(1-2^{n+1})(n+1)^{-1} B_{n+1} \quad (0 < n \in \mathbf{Z}).$$

Formula (1.14) was given in [9, (4.5)]; the sum $\sum_{h \in \mathbf{Z}}$ means $\lim_{m \rightarrow \infty} \sum_{|h| \leq m}$ if $n = 0$. Replacing t in (1.12) by $t+r$ and making an obvious calculation, we obtain (1.15). We have $E_{c,0}(t) = 1$ as noted in [9, (4.3h)]. Clearly $E_{1,n}(0) = 0$ if n is even. Assuming n to be odd, take $t = 0$ and $\alpha = 1/2$ in (1.14), and recall that $2 \cdot m!(2\pi i)^{-m} \zeta(m) = -B_m$ if $0 < m \in 2\mathbf{Z}$. Then we obtain $E_{1,n}(0)$ as stated in (1.16). Taking $r = 0$ in (1.15) and using (1.16), we obtain (1.13). The value $E_{c,n}(0)$ for an arbitrary c is given in [9, (4.6)].

1.6. In [9, Theorem 4.14] we proved, for χ , d , and k as in Theorem 1.4,

$$(1.17) \quad L(1-k, \chi) = \frac{d^{k-1}}{2^k - \overline{\chi}(2)} \sum_{a=1}^q \chi(a) E_{1,k-1}(2a/d),$$

where $q = [(d-1)/2]$, and derived (i) and (ii) above, with $E_{1,k-1}$ in place of Φ_{k-1} , from (1.17). In fact, (i) and (ii) combined are equivalent to (1.17). Though this is essentially explained in [9, p. 36], here let us show that (1.17) for even d follows from (ii). With $d = 4d_0$ as before, we have $[(d-1)/2] = 2d_0 - 1$ and

$$\sum_{a=1}^{2d_0-1} \chi(a) \Phi_{k-1}(2a/d)$$

$$= \sum_{a=1}^{d_0-1} \left\{ \chi(a) \Phi_{k-1}(2a/d) + \chi(2d_0 - a) \Phi_{k-1}(2(2d_0 - a)/d) \right\}.$$

We have $E_{1,n}(1-t) = (-1)^n E_{1,n}(t)$ as noted in [9, (4.3f)]. This combined with (1.13) shows that $\Phi_{k-1}(1-t) = (-1)^{k-1} \Phi_{k-1}(t)$. By Lemma 1.3, we have $\chi(2d_0 - a) = -\chi(-a) = (-1)^{k+1} \chi(a)$, and so the last sum equals

$$2 \sum_{a=1}^{d_0-1} \chi(a) \Phi_{k-1}(2a/d).$$

Therefore (1.17) follows from (1.10) if $d = 4d_0$. Similarly we can derive(1.17) for odd d from (1.9), which, in substance, is shown in the last paragraph of [9, p. 36].

1.7. Our technique is applicable even to $\zeta(1-k)$. Instead of $\zeta(s)$ we consider $W(s) = \sum_{m=0}^{\infty} (2m+1)^{-s}$. We have clearly

$$\begin{aligned} W(s) &= 1 + \sum_{m=1}^{\infty} (2m+1)^{-s} = 1 + \sum_{m=1}^{\infty} (2m)^{-s} \left(1 + \frac{1}{2m}\right)^{-s} \\ &= 1 + \sum_{m=1}^{\infty} (2m)^{-s} \sum_{r=0}^{\infty} \binom{-s}{r} (2m)^{-r} = 1 + \sum_{r=0}^{\infty} \zeta(s+r) \binom{-s}{r} 2^{-s-r}. \end{aligned}$$

We evaluate this at $s = 1 - k$ with $0 < k \in \mathbf{Z}$. Our calculation is similar to that of §1.1; we use (1.6) for determining the term for $r = k$, which produces $-(2k)^{-1}$. Thus

$$(1 - 2^{k-1})\zeta(1 - k) = W(1 - k) = 1 - \frac{1}{2k} + \sum_{r=0}^{k-1} \binom{k-1}{r} 2^{k-1-r} \zeta(1 - k + r).$$

Taking $k = 1$, we find a well-known fact $\zeta(0) = -1/2$. Also, $\zeta(1 - k)$ appears on both sides. Therefore, putting $k - r = t$ and rearranging our sum, we obtain

$$(1 - 2^k)\zeta(1 - k) = \frac{k-1}{2k} + \sum_{t=2}^{k-1} \binom{k-1}{t-1} 2^{t-1} \zeta(1 - t).$$

This holds for every even or odd integer $k > 1$. Recall that $\zeta(-m) = 0$ for $0 < m \in 2\mathbf{Z}$. Thus, taking $k = 2n$ with $0 < n \in \mathbf{Z}$, we obtain a formula for $\zeta(1 - 2n)$ as a linear combination of $\zeta(1 - 2\nu)$ for $1 \leq \nu < n$ (which is 0 if $n = 1$) plus a constant as follows:

$$(1.18) \quad (1 - 2^{2n})\zeta(1 - 2n) = \frac{2n-1}{4n} + \sum_{\nu=1}^{n-1} \binom{2n-1}{2\nu-1} 2^{2\nu-1} \zeta(1 - 2\nu).$$

Similarly, taking $k = 2n + 1$ and putting $t = 2\nu$, we obtain

$$(1.19) \quad \sum_{\nu=1}^n \binom{2n}{2\nu-1} 2^{2\nu-1} \zeta(1 - 2\nu) = \frac{-n}{2n+1}.$$

Either of these equalities (1.18) and (1.19) expresses $\zeta(1 - 2n)$ as a \mathbf{Q} -linear combination of $\zeta(1 - 2\nu)$ for $1 \leq \nu < n$ plus a constant. The two expressions are different, as can easily be seen.

In [9, (11.8)] we gave a similar recurrence formula which can be written

$$(1.20) \quad 4(1 - 2^{n+1})\zeta(-n) = 1 + 2 \sum_{k=2}^n \binom{n}{k-1} (2^k - 1) \zeta(1 - k) \quad (0 < n \in \mathbf{Z}).$$

Taking n to be even or odd, we again obtain two different recurrence formulas for $\zeta(1 - 2n)$. It should be noted that the technique of using the binomial theorem is already in §68 of Landau [5], in which $(s - 1)\zeta(s)$ is discussed, while we employ $W(s)$.

2. EXTENDING THE PARAMETERS c AND n IN $E_{c,n}$

2.1. The function $E_{c,n}(t)$ is a polynomial in t of degree n , and involves $c = -\mathbf{e}(\alpha)$ with $\alpha \in \mathbf{R}$. We now extend this in two ways: first, we take $\alpha \in \mathbf{C}$, $\notin \mathbf{Z}$; second, we consider $(h + \alpha)^{-s}$ instead of $(h + \alpha)^{-n-1}$. The first case is simpler. Since $E_{c,n}(t)$ is a polynomial in t and $(1 + c)^{-1}$ as noted in [9, p. 26], we can define a function $\mathcal{E}_n(\alpha, t)$ by

$$(2.1) \quad \mathcal{E}_n(\alpha, t) = E_{c,n}(t), \quad c = -\mathbf{e}(\alpha), \quad \alpha \in \mathbf{C}, \notin \mathbf{Z}, \quad 0 \leq n \in \mathbf{Z}.$$

This is a polynomial in t , whose coefficients are holomorphic functions in $\alpha \in \mathbf{C}$, $\notin \mathbf{Z}$. Now equality (1.14) can be extended to

$$(2.2) \quad \mathcal{E}_n(\alpha, t) = (1 - \mathbf{e}(-\alpha))n!(2\pi i)^{-n-1} \sum_{h \in \mathbf{Z}} (h + \alpha)^{-n-1} \mathbf{e}((h + \alpha)t)$$

for all $\alpha \in \mathbf{C}$, $\notin \mathbf{Z}$, where $0 < t < 1$ if $n = 0$, and $0 \leq t \leq 1$ if $n > 0$. Indeed, if $n > 0$, the right-hand side is absolutely convergent, and defines a holomorphic function. Since (2.2) holds for $\alpha \in \mathbf{R}$, $\notin \mathbf{Z}$, we obtain (2.2) as expected. If $n = 0$, we have to consider $\lim_{m \rightarrow \infty} \sum_{|h| \leq m} (h + \alpha)^{-1} \mathbf{e}((h + \alpha)t)$. Clearly

$$\sum_{h=-m}^m \frac{\mathbf{e}(ht)}{\alpha + h} = \frac{1}{\alpha} + \sum_{h=1}^m \frac{2\alpha \cdot \cos(2\pi ht)}{\alpha^2 - h^2} + 2i \sum_{h=1}^m \frac{h \cdot \sin(2\pi ht)}{h^2 - \alpha^2}.$$

The last sum on the right-hand side equals

$$\sum_{h=1}^m \frac{\sin(2\pi ht)}{h} + \sum_{h=1}^m \frac{\sin(2\pi ht)\alpha^2}{h(h^2 - \alpha^2)}.$$

It is well-known that the first sum tends to a finite value as $m \rightarrow \infty$. Obviously the last sum converges to a holomorphic function in $\alpha \in \mathbf{C}$, $\notin \mathbf{Z}$ as $m \rightarrow \infty$. Thus we can justify (2.2) for $n = 0$.

Formula (2.2) for $n = 0$ (with $-\alpha$ in place of α) can be written

$$(2.3) \quad \frac{\mathbf{e}(t\alpha)}{1 - \mathbf{e}(\alpha)} = \frac{1}{2\pi i} \sum_{h \in \mathbf{Z}} \frac{\mathbf{e}(th)}{h - \alpha} \quad (\alpha \in \mathbf{C}, \notin \mathbf{Z}, \quad 0 < t < 1).$$

This was first given by Kronecker [4].

2.2. We next ask if the power $(h + a)^{-n-1}$ in (1.14) can be replaced by $(h + a)^{-s}$ with a complex parameter s . Since $h + a$ can be negative, $(h + a)^{-s}$ is not suitable. Thus, for $s \in \mathbf{C}$, $a \in \mathbf{R}$, $p \in \mathbf{R}$, and $\nu = 0$ or 1 we put

$$(2.4) \quad D^\nu(s; a, p) = \sum_{-a \neq n \in \mathbf{Z}} (n + a)^\nu |n + a|^{-\nu-s} \mathbf{e}(p(n + a)),$$

$$(2.5) \quad T^\nu(s; a, p) = \Gamma((s + \nu)/2)\pi^{-(s+\nu)/2}D^\nu(s; a, p).$$

Clearly the infinite series of (2.4) is absolutely convergent for $\text{Re}(s) > 1$, and defines a holomorphic function of s there. Notice that if $k - \nu \in 2\mathbf{Z}$, then $D^\nu(k; 0, t) = \sum_{0 \neq n \in \mathbf{Z}} n^{-k} \mathbf{e}(nt)$, which is the infinite sum of (0.3). Thus the Bernoulli polynomials are included in our discussion.

THEOREM 2.3. *The function $T^\nu(s; a, p)$ can be continued as a meromorphic function of s to the whole \mathbf{C} . It is entire if $\nu = 1$. If $\nu = 0$, then $T^0(s; a, p)$ is*

$$\frac{-2\delta(a)}{s} + \frac{2\mathbf{e}(ap)\delta(p)}{s-1}$$

plus an entire function, where $\delta(x) = 1$ if $x \in \mathbf{Z}$ and $\delta(x) = 0$ if $x \notin \mathbf{Z}$. Moreover,

$$(2.6) \quad T^\nu(1-s; a, p) = i^{-\nu} \mathbf{e}(ap)T^\nu(s; -p, a).$$

Proof. Put $\varphi(x) = x^\nu \mathbf{e}(-x^2 z^{-1}/2 + px)$ for $x \in \mathbf{R}$ and $z \in H$. Denote by $\widehat{\varphi}$ the Fourier transform of φ . Then from [9, (2.25)] we easily obtain $\widehat{\varphi}(x) = i^{-\nu} (-iz)^\kappa (x-p)^\nu \mathbf{e}((x-p)^2 z/2)$, where $\kappa = \nu + 1/2$. Put also

$$f(z) = \sum_{n \in \mathbf{Z}} (n+a)^\nu \mathbf{e}((n+a)^2 z/2 + p(n+a)),$$

and $f^\#(z) = (-iz)^{-\kappa} f(-z^{-1})$. Then $f(-z^{-1}) = \sum_{n \in \mathbf{Z}} \varphi(n+a)$, which equals $\sum_{m \in \mathbf{Z}} \mathbf{e}(ma) \widehat{\varphi}(m)$ by virtue of the Poisson summation formula. In this way we obtain

$$f^\#(z) = i^{-\nu} \sum_{m \in \mathbf{Z}} \mathbf{e}(ma)(m-p)^\nu \mathbf{e}((m-p)^2 z/2).$$

Now $T^\nu(2s - \nu; a, p)$ is the Mellin transform of $f(iy)$, and so we obtain our theorem by the general principle of Hecke, which is given as Theorem 3.2 in [9].

THEOREM 2.4. *For $\nu = 0$ or 1 , $0 \leq a \leq 1$, and a positive integer k such that $k - \nu \in 2\mathbf{Z}$ we have*

$$(2.7) \quad D^0(0; a, p) = -\delta(a),$$

$$(2.8) \quad D^\nu(\nu - 2m; a, p) = 0 \quad \text{if } 0 < m \in \mathbf{Z},$$

$$(2.9) \quad D^\nu(1 - k; a, p) = 2(2\pi i)^{-k} (k-1)! \mathbf{e}(ap) D^\nu(k; -p, a),$$

$$(2.10) \quad D^\nu(1 - k; a, p) = -2\mathbf{e}(ap) B_k(a)/k \quad \text{if } p \in \mathbf{Z},$$

$$(2.11) \quad D^\nu(1 - k; a, p) = \frac{2\mathbf{e}(ap)}{1 - \mathbf{e}(p)} E_{c,k-1}(a) \quad \text{if } p \notin \mathbf{Z},$$

where $c = -\mathbf{e}(-p)$, and we have to assume that $0 < a < 1$ in (2.10) and (2.11) if $k = 1$.

Proof. By Theorem 2.3, $[sT^0(s; a, p)]_{s=0} = -2\delta(a)$, from which we obtain (2.7). Next, let $0 < m \in \mathbf{Z}$. Since $\Gamma((s + \nu)/2)D^\nu(s; a, p)$ is finite and

$\Gamma((s + \nu)/2)$ has a pole at $s = \nu - 2m$, we obtain (2.8). We easily see that $\Gamma(1/2 - m) = \pi^{1/2}(-2)^m \prod_{t=1}^m (2t - 1)^{-1}$. Therefore from (2.6) we obtain (2.9). If $p \in \mathbf{Z}$, then $D^\nu(k; -p, a) = \sum_{0 \neq n \in \mathbf{Z}} n^{-k} \mathbf{e}(an)$. The well-known classical formula, stated in [9, (4.9)] (and also as (0.3)), shows that the last sum equals $-(2\pi i)^k B_k(a)/k!$ for $0 \leq a \leq 1$ if $k > 1$, and for $0 < a < 1$ if $k = 1$. If $p \notin \mathbf{Z}$, then $D^\nu(k; -p, a) = \sum_{n \in \mathbf{Z}} (n - p)^{-k} \mathbf{e}(a(n - p))$. By (1.14), this equals $(1 + c^{-1})^{-1} (2\pi i)^k E_{c,k-1}(a)/(k - 1)!$, where $c = -\mathbf{e}(-p)$, under the same condition on a . Combining these with (2.9), we obtain (2.10) and (2.11).

We note here a special case of (2.10):

$$(2.12) \quad D^\nu(1 - k; 0, 0) = \begin{cases} -2B_k/k & \text{if } k > 1, \\ 0 & \text{if } k = 1. \end{cases}$$

It should be noted that $D^1(s; 0, 0) = 0$.

3. NEARLY HOLOMORPHIC ELLIPTIC FUNCTIONS

3.1. Let L be a lattice in \mathbf{C} . As an analogue of (2.4) we put

$$(3.1) \quad \varphi_\nu(u, s; L) = \sum_{\alpha \in L} (u + \alpha)^{-\nu} |u + \alpha|^{\nu - 2s}$$

for $0 \leq \nu \in \mathbf{Z}$, $u \in \mathbf{C}$, $\notin L$, and $s \in \mathbf{C}$. Clearly

$$(3.2a) \quad \varphi_\nu(\lambda u, s; \lambda L) = \lambda^{-\nu} |\lambda|^{\nu - 2s} \varphi_\nu(u, s; L) \quad \text{for every } \lambda \in \mathbf{C}^\times,$$

$$(3.2b) \quad \varphi_\nu(u + \alpha, s; L) = \varphi_\nu(u, s; L) \quad \text{for every } \alpha \in L.$$

If L is a \mathbf{Z} -lattice in an imaginary quadratic field K and $u \in K$, (3.1) is the same as the series of [9, (7.1)]. The analytic properties of the series that we proved there can easily be extended to the case of (3.1). First of all, the right-hand side of (3.1) is absolutely convergent for $\text{Re}(s) > 1$, and defines a holomorphic function of s there.

THEOREM 3.2. *Put $\Phi(u, s) = \pi^{-s} \Gamma(s + \nu/2) \varphi_\nu(u, s; L)$. Then $\Phi(u, s)$ can be continued to the whole s -plane as a meromorphic function in s , which is entire if $\nu > 0$. If $\nu = 0$, then $\Phi(u, s)$ is an entire function of s plus $v(L)^{-1}/(s - 1)$, where $v(L) = \text{vol}(\mathbf{C}/L)$. Moreover, $\Phi(u, s)$ is a C^∞ function in u , except when $\nu = 0$ and $s = 1$, and each derivative $(\partial/\partial u)^a (\partial/\partial \bar{u})^b \Phi(u, s)$ is meromorphic in s on the whole \mathbf{C} .*

Proof. This can be proved by the same argument as in [9, §7.2], except for the differentiability with respect to u and the last statement about the derivatives, which can be shown as follows. As shown in the proof of [9, Theorem 3.2], the product $\pi^{-s} \Gamma(s) \varphi_\nu(u, s - \nu/2; L)$ minus the pole part can be written

$$\int_p^\infty F(u, y)y^{s-1}dy + \int_p^\infty G(u, y)y^{\nu-s}dy,$$

where

$$F(u, y) = \sum_{\alpha \in L} (\bar{u} + \alpha)^\nu \exp(-\pi|u + \alpha|^2 y),$$

$$G(u, y) = A \sum_{\beta \in B} \exp(\pi i(\beta \bar{u} + \bar{\beta} u)) \sum_{\xi - \beta \in M} \xi^\nu \exp(-\pi|\xi|^2 y)$$

with a constant A , a finite subset B of \mathbf{C} , a positive constant p , and lattices L and M in \mathbf{C} . Therefore the differentiability and the last statement follow from the standard fact on differentiation under the integral sign.

3.3. Before stating the next theorem, we note a few elementary facts. Take L in the form $L = \mathbf{Z}\omega_1 + \mathbf{Z}\omega_2$ with complex numbers ω_1 and ω_2 such that $\omega_1/\omega_2 \in H$. We put then $v(\omega_1, \omega_2) = v(L)$. It can easily be seen that

$$(3.3) \quad v(\omega_1, \omega_2) = |\omega_2|^2 \text{Im}(\omega_1/\omega_2) = (2i)^{-1}(\omega_1 \bar{\omega}_2 - \bar{\omega}_1 \omega_2),$$

and in particular, $v(z, 1) = \text{Im}(z)$. We also recall the function ζ of Weierstrass defined by

$$(3.4) \quad \zeta(u) = \zeta(u; \omega_1, \omega_2) = \frac{1}{u} + \sum_{0 \neq \alpha \in L} \left\{ \frac{1}{u - \alpha} + \frac{1}{\alpha} + \frac{u}{\alpha^2} \right\},$$

where $L = \mathbf{Z}\omega_1 + \mathbf{Z}\omega_2$. It is well known that

$$(3.5) \quad \zeta(-u) = -\zeta(u), \quad (\partial/\partial u)\zeta(u; \omega_1, \omega_2) = -\wp(u; \omega_1, \omega_2)$$

with the Weierstrass function \wp . We put as usual

$$(3.6a) \quad \eta_\mu(\omega_1, \omega_2) = 2\zeta(\omega_\mu/2) \quad (\mu = 1, 2).$$

Then

$$(3.6b) \quad \zeta(u + \omega_\mu) = \zeta(u) + \eta_\mu(\omega_1, \omega_2).$$

We also need the classical nonholomorphic Eisenstein series E_2 of weight 2, which can be given by

$$(3.7) \quad E_2(z) = \frac{1}{8\pi y} - \frac{1}{24} + \sum_{n=1}^\infty \left(\sum_{0 < d|n} d \right) \mathbf{e}(nz).$$

We are interested in the value of $\varphi_\nu(u, s; L)$ at $s = \nu/2$, which is meaningful for every $\nu \in \mathbf{Z}, > 0$, by Theorem 3.2. The results can be given as follows.

THEOREM 3.4. *For $L = \mathbf{Z}\omega_1 + \mathbf{Z}\omega_2$ with $\omega_1/\omega_2 \in H$ we have*

$$(3.8) \quad \varphi_\nu(u, \nu/2; L) = \frac{(-1)^\nu}{(\nu - 1)!} \frac{\partial^{\nu-2}}{\partial u^{\nu-2}} \wp(u; \omega_1, \omega_2) \quad (2 < \nu \in \mathbf{Z}),$$

$$(3.9) \quad \varphi_2(u, 1; L) = \wp(u; \omega_1, \omega_2) - 8\pi^2 \omega_2^{-2} E_2(\omega_1/\omega_2),$$

$$(3.10) \quad \varphi_1(u, 1/2; L) = \zeta(u) + 8\pi^2 \omega_2^{-2} E_2(\omega_1/\omega_2)u - \pi v(L)^{-1} \bar{u},$$

$$(3.11) \quad \eta_\mu(\omega_1, \omega_2) = \pi \bar{\omega}_\mu v(L)^{-1} - 8\pi^2 \omega_2^{-2} E_2(\omega_1/\omega_2) \omega_\mu \quad (\mu = 1, 2).$$

Proof. If $\nu > 2$, then clearly $\varphi_\nu(u, \nu/2; L) = \sum_{\alpha \in L} (u + \alpha)^{-\nu}$, from which we obtain (3.8). The cases $\nu = 2$ and $\nu = 1$ are more interesting. We first note that

$$(3.12a) \quad (\partial/\partial u)\varphi_\nu(u, s; L) = (-s - \nu/2)\varphi_{\nu+1}(u, s + 1/2; L),$$

$$(3.12b) \quad (\partial/\partial \bar{u})\varphi_\nu(u, s; L) = (-s + \nu/2)\varphi_{\nu-1}(u, s + 1/2; L),$$

at least for sufficiently large $\operatorname{Re}(s)$. Since both sides of (3.12a, b) are meromorphic in s on the whole \mathbf{C} , we obtain (3.12a, b) for every s . The first formula with $\nu = 2$ produces

$$(\partial/\partial u)\varphi_2(u, 1; L) = -2\varphi_3(u, 3/2; L) = (\partial/\partial u)\wp(u; \omega_1, \omega_2),$$

from which we obtain $\varphi_2(u, 1; L) = \wp(u; \omega_1, \omega_2) + c(\bar{u})$ with an anti-holomorphic function $c(\bar{u})$. Since (3.12b) shows that $\varphi_2(u, 1; L)$ is holomorphic in u , we see that $c(\bar{u})$ does not involve u or \bar{u} , that is, it is a constant depending only on L . Suppose $L = \mathbf{Z}z + \mathbf{Z}$ with $z \in H$. For $0 < N \in \mathbf{Z}$ and $(p, q) \in \mathbf{Z}^2, \notin N\mathbf{Z}^2$ define a standard Eisenstein series $\mathfrak{E}_\nu^N(z, s; p, q)$ of level N by [9, (9.1)]. Then we easily see that

$$\varphi_\nu((pz + q)/N, s; L) = N^{2s} y^{\nu/2 - s} \mathfrak{E}_\nu^N(z, s - \nu/2; p, q),$$

$$\varphi_\nu((pz + q)/N, \nu/2; L) = N^\nu \mathfrak{E}_\nu^N(z, 0; p, q).$$

Define F_ν and \mathcal{F}_2 as in [9, (10.10b, c, d)]. Taking $\nu = 2$, we obtain

$$\varphi_2((pz + q)/N, 1; L) = N^2 \mathfrak{E}_2^N(z, 0; p, q) = (2\pi i)^2 \mathcal{F}_2(z; p/N, q/N).$$

By [9, (10.13)], $\mathcal{F}_2(z; a, b) = (2\pi i)^{-2} \wp(az + b; z, 1) + 2E_2(z)$ with E_2 of (3.7). Therefore we can conclude that

$$(3.13) \quad \varphi_2(u, 1; \mathbf{Z}z + \mathbf{Z}) = \wp(u; z, 1) - 8\pi^2 E_2(z).$$

More generally, using (3.2a) we obtain (3.9).

We next consider the case $\nu = 1$. Since $(\partial/\partial u)\zeta(u; \omega_1, \omega_2) = -\wp(u; \omega_1, \omega_2)$, from (3.9) and (3.12a) we obtain

$$\begin{aligned} (\partial/\partial u)\varphi_1(u, 1/2; L) &= -\varphi_2(u, 1; L) \\ &= (\partial/\partial u)\zeta(u; \omega_1, \omega_2) + 8\pi^2 \omega_2^{-2} E_2(\omega_1/\omega_2). \end{aligned}$$

We have also

$$(\partial/\partial \bar{u})\varphi_1(u, 1/2; L) = \lim_{\sigma \rightarrow 1} (1 - \sigma)\varphi_0(u, \sigma; L) = -\pi/v(L),$$

since the residue of $\pi^{-s} \Gamma(s)\varphi_0(u, s; L)$ at $s = 1$ is $v(L)^{-1}$ as shown in Theorem 3.2. Therefore $\varphi_1(u, 1/2; L) = -\pi \bar{u}/v(L) + g(u)$ with a function g holomorphic in u . Clearly $\partial g/\partial u = (\partial/\partial u)\varphi_1(u, 1/2; L)$, and so we can conclude that

$$(3.14) \quad \varphi_1(u, 1/2; L) = \zeta(u) + 8\pi^2\omega_2^{-2}E_2(\omega_1/\omega_2)u - \pi v(L)^{-1}\bar{u} + \xi(L)$$

with a constant $\xi(L)$ independent of u . From (3.2a) we obtain $\varphi_1(-u, s; L) = -\varphi_1(u, s; L)$. Also $\zeta(-u) = -\zeta(u)$. Thus $\xi(L) = 0$, and consequently we obtain (3.10). Since $\varphi_1(u, 1/2; L)$ is invariant under $u \mapsto u + \omega_\mu$, we obtain (3.11) from (3.10) and (3.6b).

3.5. In [9] we discussed the value of an Eisenstein series $E(z, s)$ of weight k at $s = -m$ for an integer m such that $0 \leq m \leq k - 1$, and observed that it is nearly holomorphic in the sense that it is a polynomial in y^{-1} with holomorphic functions as coefficients; for a precise statement, see [9, Theorem 9.6]. As an analogue we investigate $\varphi_\nu(u, \kappa/2; L)$ for an integer κ such that $2 - \nu \leq \kappa \leq \nu$ and $\kappa - \nu \in 2\mathbf{Z}$. From (3.12b) we obtain, for $0 \leq a \in \mathbf{Z}$,

$$(3.15) \quad (\partial/\partial\bar{u})^a \varphi_\nu(u, (\nu/2) - a; L) = a! \cdot \varphi_{\nu-a}(u, (\nu - a)/2; L).$$

THEOREM 3.6. *Let κ be an integer such that $2 - \nu \leq \kappa \leq \nu$ and $\kappa - \nu \in 2\mathbf{Z}$. Then $\varphi_\nu(u, \kappa/2; L)$ is a polynomial in \bar{u} of degree d with holomorphic functions in u as coefficients, where $d = (\nu - \kappa)/2$ if $\nu + \kappa \geq 4$ and $d = (\nu - \kappa + 2)/2$ if $\nu + \kappa = 2$. The leading term is $\bar{u}^d \varphi_{(\nu+\kappa)/2}(u, (\nu + \kappa)/4; L)$ or $-\pi d^{-1}v(L)^{-1}\bar{u}^d$ according as $\nu + \kappa \geq 4$ or $\nu + \kappa = 2$.*

Proof. Given κ as in the theorem, put $a = (\nu - \kappa)/2$. Then $(\nu/2) - a = \kappa/2$ and $\nu - a = (\nu + \kappa)/2 \geq 1$. If $\nu - a \geq 2$, then by Theorem 3.4, $\varphi_{\nu-a}(u, (\nu - a)/2; L)$ is holomorphic in u , and so (3.15) shows that $\varphi_\nu(u, \kappa/2; L)$ is a polynomial in \bar{u} of degree a with holomorphic functions in u as coefficients. If $\nu - a = 1$, the function $\varphi_1(u, 1/2; L)$ is linear in \bar{u} as given in (3.10). Therefore we obtain our theorem.

Thus, we may call $\varphi_\nu(u, \kappa/2; L)$ a *nearly holomorphic elliptic function*. In the higher-dimensional case it is natural to consider theta functions instead of periodic functions. For details of the basic ideas and results on this the reader is referred to [6] and [7].

4. THE SERIES WITH A PARAMETER IN H

4.1. To state the following lemma, we first define a confluent hypergeometric function $\tau(y; \alpha, \beta)$ for $y > 0$ and $(\alpha, \beta) \in \mathbf{C}^2$ by

$$(4.1) \quad \tau(y; \alpha, \beta) = \int_0^\infty e^{-yt}(1+t)^{\alpha-1}t^{\beta-1}dt.$$

This is convergent for $\text{Re}(\beta) > 0$. It can be shown that $\Gamma(\beta)^{-1}\tau(y; \alpha, \beta)$ can be continued to a holomorphic function in (α, β) on the whole \mathbf{C}^2 ; see [9, Section A3], for example. Also, for $v \in \mathbf{C}^\times$ and $\alpha \in \mathbf{C}$ we define v^α by

$$(4.2) \quad v^\alpha = \exp(\alpha \log(v)), \quad -\pi < \text{Im}[\log(v)] \leq \pi.$$

LEMMA 4.2. For $\alpha, \beta \in \mathbf{C}$ such that $\operatorname{Re}(\alpha + \beta) > 1$, $0 \leq r < 1$, and $z = x + iy \in H$ we have

$$\begin{aligned} & i^{\alpha-\beta}(2\pi)^{-\alpha-\beta}\Gamma(\alpha)\Gamma(\beta) \sum_{m \in \mathbf{Z}} \mathbf{e}(mr)(z+m)^{-\alpha}(\bar{z}+m)^{-\beta} \\ &= \sum_{n=1}^{\infty} \mathbf{e}((n-r)z)(n-r)^{\alpha+\beta-1}\tau(4\pi(n-r)y; \alpha, \beta) \\ &+ \sum_{n=1}^{\infty} \mathbf{e}(-(n+r)\bar{z})(n+r)^{\alpha+\beta-1}\tau(4\pi(n+r)y; \beta, \alpha) \\ &+ \begin{cases} (4\pi y)^{1-\alpha-\beta}\Gamma(\alpha+\beta-1) & \text{if } r=0, \\ \mathbf{e}(-r\bar{z})r^{\alpha+\beta-1}\tau(4\pi ry; \beta, \alpha) & \text{if } r \neq 0. \end{cases} \end{aligned}$$

Proof. If $r = 0$, this is Lemma A3.4 of [9]. The case with nontrivial r can be proved in the same way as follows. Define two functions $f(x)$ and $f_1(x)$ of $x \in \mathbf{R}$ by $f(x) = (x+iy)^{-\alpha}(x-iy)^{-\beta}$ with a fixed $y > 0$ and $f_1(x) = \mathbf{e}(rx)f(x)$. Then $\hat{f}_1(x) = \hat{f}(x-r)$, and so the Poisson summation formula (see [9, (2.9)]) shows that

$$\mathbf{e}(-rx) \sum_{m \in \mathbf{Z}} f_1(x+m) = \mathbf{e}(-rx) \sum_{n \in \mathbf{Z}} \hat{f}_1(n)\mathbf{e}(nx) = \sum_{n \in \mathbf{Z}} \mathbf{e}((n-r)x)\hat{f}(n-r).$$

In [9, p. 133] we determined \hat{f} explicitly in terms of τ as follows:

$$i^{\alpha-\beta}(2\pi)^{-\alpha-\beta}\Gamma(\alpha)\Gamma(\beta)\hat{f}(t) = \begin{cases} \mathbf{e}(ity)t^{\alpha+\beta-1}\tau(4\pi ty; \alpha, \beta) & (t > 0), \\ \mathbf{e}(-ity)|t|^{\alpha+\beta-1}\tau(4\pi|t|y; \beta, \alpha) & (t < 0), \\ (4\pi y)^{1-\alpha-\beta}\Gamma(\alpha+\beta-1) & (t = 0). \end{cases}$$

Therefore we obtain our lemma.

4.3. We now need an elementary result:

$$(4.3) \quad \sum_{n=1}^{\infty} n^{k-1}x^n = \frac{xP_k(x)}{(1-x)^k} \quad (1 \leq k \in \mathbf{Z}).$$

Here x is an indeterminate and P_k is a polynomial. We have $P_1 = P_2 = 1$ and $P_{k+1} = (kx - x + 1)P_k - (x^2 - x)P'_k$ for $k \geq 2$. Thus P_k is of degree $k-2$ for $k \geq 2$. These are easy; see [9, p. 17]. We also need two formulas and an estimate given as (A3.11), (A3.14), and Lemma A3.2 in [9]:

$$(4.4) \quad \tau(y; n, \beta) = \sum_{\mu=0}^{n-1} \binom{n-1}{\mu} \Gamma(\beta+\mu)y^{-\mu-\beta} \quad (0 < n \in \mathbf{Z}),$$

$$(4.5) \quad [\tau(y; \alpha, \beta)/\Gamma(\beta)]_{\beta=0} = 1,$$

(4.6) $\Gamma(\beta)^{-1}y^\beta\tau(y; \alpha, \beta)$ is bounded when (α, β) belongs to a compact subset of \mathbf{C}^2 and $y > c$ with a positive constant c .

Our principal aim of this section is to study the nature of the series

$$(4.7) \quad \mathcal{L}_k(s, z) = \sum_{m \in \mathbf{Z}} \mathbf{e}(mr)(z+m)^{-k}|z+m|^{-2s},$$

for certain integer values of s . Here $k \in \mathbf{Z}$, $r \in \mathbf{R}$, $s \in \mathbf{C}$, and $z \in H$. The sum depends only on r modulo \mathbf{Z} , and so we may assume that $0 \leq r < 1$. Clearly this series is absolutely convergent for $\text{Re}(2s+k) > 1$, and defines a holomorphic function of s there.

THEOREM 4.4. *The function $\mathcal{L}_k(s, z)$ can be continued as a meromorphic function of s to the whole \mathbf{C} , which is entire if $r \notin \mathbf{Z}$. If $r \in \mathbf{Z}$, the locations of the poles of $\mathcal{L}_k(s, z)$ are the same as those of $\Gamma(2s+k-1)/\{\Gamma(s+k)\Gamma(s)\}$.*

Proof. Our function is the infinite series of Lemma 4.2 defined with $\alpha = s+k$ and $\beta = s$. Therefore our assertion can easily be verified by means of the formula of Lemma 4.2 and the estimate given by (4.6).

THEOREM 4.5. *Assuming that $r \in \mathbf{Q}$, let N be the smallest positive integer such that $Nr \in \mathbf{Z}$ and let $\beta \in \mathbf{Z}$. Then the following assertions hold:*

- (i) *If $\beta > 0$ or $\beta+k > 0$, then $\mathcal{L}_k(s, z)$ is finite at $s = \beta$ and $i^k \mathcal{L}_k(\beta, z)$ is a rational function in π , $\mathbf{e}(z/N)$, $\mathbf{e}(-\bar{z}/N)$, and $\text{Im}(z)$ with coefficients in \mathbf{Q} .*
- (ii) *If $-k < \beta \leq 0$, then $i^k \{1 - \mathbf{e}(z)\}^{k+\beta} \mathcal{L}_k(\beta, z)$ is a polynomial in π , $\mathbf{e}(z/N)$, $\text{Im}(z)$, and $\text{Im}(z)^{-1}$ with coefficients in \mathbf{Q} .*
- (iii) *If $0 < \beta \leq -k$, then $i^k \{1 - \mathbf{e}(-\bar{z})\}^\beta \mathcal{L}_k(\beta, z)$ is a polynomial in π , $\mathbf{e}(-\bar{z}/N)$, $\text{Im}(z)$, and $\text{Im}(z)^{-1}$ with coefficients in \mathbf{Q} .*

Proof. As we already said, we may assume that $0 \leq r < 1$. Put $\alpha = \beta + k$. We first have to study the nature of $\Gamma(2s+k-1)/\{\Gamma(s+k)\Gamma(s)\}$ at $s = \beta$. This is clearly finite at $s = \beta$ if $\alpha + \beta > 1$. Suppose $\alpha + \beta \leq 1$; then $\alpha \leq 0$ if $\beta > 0$, and $\beta \leq 0$ if $\alpha > 0$. In all cases the value is finite, and in fact is a rational number. We now evaluate the formula of Lemma 4.2 divided by $\Gamma(\alpha)\Gamma(\beta)$. If $\alpha > 0$, we have, by (4.4),

$$\tau(4\pi(n-r)y; \alpha, \beta)/\Gamma(\beta) = \sum_{\mu=0}^{\alpha-1} \binom{\alpha-1}{\mu} (n-r)^{-\mu-\beta} (4\pi y)^{-\mu-\beta} \prod_{\kappa=0}^{\mu-1} (\beta + \kappa).$$

Thus an infinite sum of the form $\sum_{n=1}^{\infty} \mathbf{e}((n-r)z)(n-r)^{\alpha-\mu-1}$ appears. Applying the binomial theorem to the power of $n-r$, we see that the sum is a \mathbf{Q} -linear combination of $\mathbf{e}(-rz) \sum_{n=1}^{\infty} \mathbf{e}(nz)n^\nu$ for $0 \leq \nu \leq \alpha - \mu - 1$. We can handle $\tau(4\pi(n+r)y; \beta, \alpha)/\Gamma(\alpha)$ in a similar way if $\beta > 0$. Put $\mathbf{q} = \mathbf{e}(z)$ and $\mathbf{q}_r = \mathbf{e}(rz)$. Then, assuming that $0 < r < 1$, $\alpha > 0$, and $\beta > 0$, we have

$$\begin{aligned} i^k \mathcal{L}_k(\beta, z) &= \mathbf{q}_r^{-1} \sum_{\mu=0}^{\alpha-1} \sum_{\nu=0}^{\alpha-\mu-1} a_{\mu\nu} \pi^{\alpha-\mu} y^{-\mu-\beta} \sum_{n=1}^{\infty} n^\nu \mathbf{q}^n \\ &\quad + \bar{\mathbf{q}}_r \sum_{\mu=0}^{\beta-1} \sum_{\nu=0}^{\beta-\mu-1} b_{\mu\nu} \pi^{\beta-\mu} y^{-\mu-\alpha} \sum_{n=0}^{\infty} n^\nu \bar{\mathbf{q}}^n, \end{aligned}$$

where $a_{\mu\nu}$ and $b_{\mu\nu}$ are rational numbers depending on β , k , and r . Applying (4.3) to $\sum_{n=1}^{\infty} n^{\nu} X^n$ with $X = \mathbf{q}$ and $X = \bar{\mathbf{q}}$, we obtain (i). Suppose $\beta \leq 0$ and $\beta + k > 0$; then the sum involving $\tau(4\pi(n+r)y; \beta, \alpha) / \{\Gamma(\alpha)\Gamma(\beta)\}$ vanishes and we obtain (ii). The case in which $\beta > 0$ and $\beta + k \leq 0$ is similar and produces (iii). If $r = 0$, the constant term of $i^k \mathcal{L}_k(\beta, z)$ is $2\pi(2y)^{1-\alpha-\beta} \Gamma(\alpha + \beta - 1) / [\Gamma(\alpha)\Gamma(\beta)]$, which causes no problem. This completes the proof.

One special case is worthy of attention. Taking $\beta = 0$ and $1 < k = \alpha \in \mathbf{Z}$, and using (4.5), we obtain, for $0 \leq r < 1$,

$$(4.8) \quad \sum_{m \in \mathbf{Z}} \frac{\mathbf{e}(r(z+m))}{(z+m)^k} = \frac{(2\pi i)^k}{(k-1)!} \sum_{\nu=1}^k \binom{k-1}{\nu-1} r^{k-\nu} \frac{\mathbf{q} P_{\nu}(\mathbf{q})}{(\mathbf{q}-1)^{\nu}},$$

where $\mathbf{q} = \mathbf{e}(z)$. We assume $0 < r < 1$ and $\sum_{m \in \mathbf{Z}} = \lim_{h \rightarrow \infty} \sum_{|m| \leq h}$ when $k = 1$. In (4.7) we take z in H , but in (4.8) we can take $z \in \mathbf{C}$, $\notin \mathbf{Z}$, since both sides of (4.8) are meaningful for such z . If $k = 1$, the result is the same as (2.3).

We can mention another special case. Namely, take $z = ia$ with a positive rational number a . Then we see that the values

$$(4.9) \quad \sum_{m \in \mathbf{Z}} (a^2 + m^2)^{-\beta}$$

for $0 < \beta \in \mathbf{Z}$ belong to the field generated by π and $e^{-2\pi a}$ over \mathbf{Q} , and therefore any three such values satisfy a nontrivial algebraic equation over \mathbf{Q} .

5. THE RATIONALITY OVER A TOTALLY REAL BASE FIELD

5.1. Throughout this section we fix a totally real algebraic number field F . The algebraicity of $\pi^{-k} L(k, \chi)$ can be generalized to the case of L -functions over F , but there is no known formulas similar to (0.2), (1.9), (1.10), except that Siegel proved some such formulas in [10] and [11] when $[F : \mathbf{Q}] = 2$. The paper [1] of Hecke may be mentioned in this connection. In this section we merely consider a generalization of (2.4) and prove an algebraicity result on its critical values, without producing explicit expressions.

We denote by \mathfrak{g} , D_F , and \mathfrak{a} the maximal order of F , the discriminant of F , and the set of archimedean primes of F . We also put $\text{Tr}(x) = \text{Tr}_{F/\mathbf{Q}}(x)$ for $x \in F$ and $[F : \mathbf{Q}] = g$. For $\alpha \in F$ and a fractional ideal \mathfrak{a} in F we put $\alpha + \mathfrak{a} = \{\alpha + x \mid x \in \mathfrak{a}\}$ and $\tilde{\mathfrak{a}} = \{\xi \in F \mid \text{Tr}(\xi\mathfrak{a}) \subset \mathbf{Z}\}$.

Given α and \mathfrak{a} as above, $\xi \in F$, $0 < \mu \in \mathbf{Z}$, and a (sufficiently small) subgroup U of \mathfrak{g}^{\times} of finite index, we put

$$(5.1) \quad D_{\mu}(s; \xi, \alpha, \mathfrak{a}) = r_U \sum_{0 \neq h \in U \setminus (\alpha + \mathfrak{a})} \mathbf{e}_{\mathfrak{a}}(h\xi) h^{-\mu\mathfrak{a}} |h|^{(\mu-s)\mathfrak{a}},$$

$$(5.1a) \quad r_U = [\mathfrak{g}^{\times} : U]^{-1},$$

where $\mathbf{e}_a(\xi) = \mathbf{e}(\sum_{v \in \mathbf{a}}(\xi_v))$ for $\xi \in F$ and $x^{t\mathbf{a}} = \prod_{v \in \mathbf{a}} x_v^t$ for $x \in \mathbf{C}^{\mathbf{a}}$ and $U \backslash X$ means a complete set of representatives for X modulo multiplication by the elements of U . We have to take U so small that the sum of (5.1) is meaningful. For instance, it is sufficient to take

$$U \subset \{u \in \mathfrak{g}^\times \mid u^{\mathbf{a}} = 1, u\xi - \xi \in \alpha^{-1}\tilde{\mathfrak{g}} \cap \tilde{\mathfrak{a}}\}.$$

The factor r_U makes the quantity of (5.1) independent of the choice of U . Clearly the sum is convergent for $\text{Re}(s) > 1$. Now $D_\mu(s; \xi, \alpha, \mathbf{a})$ is a special case of the series of [8, (18.1)], and so from Lemma 18.2 of [8] we see that it can be continued as a holomorphic function in s to the whole \mathbf{C} .

THEOREM 5.2. *For $0 < \mu \in \mathbf{Z}$ we have*

$$(5.2) \quad (2\pi i)^{-\mu g} D_F^{1/2} D_\mu(\mu; \xi, 0, \mathbf{a}) \in \mathbf{Q},$$

$$(5.3) \quad D_\mu(1 - \mu; 0, \alpha, \mathbf{a}) \in \mathbf{Q}.$$

Proof. The last formula is a restatement of Proposition 18.10(2) of [8]. To prove (5.2), let $\mathfrak{b} = \{x \in \mathfrak{a} \mid \mathbf{e}_a(x\xi) = 1\}$ and let R be a complete set of representatives for $\mathfrak{a}/\mathfrak{b}$. Then

$$D_\mu(s; \xi, 0, \mathbf{a}) = \sum_{\beta \in R} \mathbf{e}_a(\beta\xi) D_\mu(s; 0, \beta, \mathfrak{b}).$$

Put $Q_\mu(\beta, \mathfrak{b}) = (2\pi i)^{-\mu g} D_F^{1/2} D_\mu(\mu; 0, \beta, \mathfrak{b})$. Then the quantity of (5.2) equals $\sum_{\beta \in R} \mathbf{e}_a(\beta\xi) Q_\mu(\beta, \mathfrak{b})$. Let $t \in \prod_p \mathbf{Z}_p^\times$ and let σ be the image of t under the canonical homomorphism of \mathbf{Q}_A^\times onto $\text{Gal}(\mathbf{Q}_{\text{ab}}/\mathbf{Q})$. Our task is to show that the last sum is invariant under σ . By [8, Proposition 18.10(1)] we have $Q_\mu(\beta, \mathfrak{b})^\sigma = Q_\mu(\beta_1, \mathfrak{b})$ with $\beta_1 \in F$ such that $(t\beta_1 - \beta)_v \in \mathfrak{b}_v$ for every nonarchimedean prime v of F . For $\beta \in R$ there is a unique $\beta_1 \in R$ with that property. Now $\mathbf{e}(c)^\sigma = \mathbf{e}(t^{-1}c)$ for every $c \in \mathbf{Q}/\mathbf{Z} = \prod_p (\mathbf{Q}_p/\mathbf{Z}_p)$; see [8, (8.2)]. Since $\mathbf{e}_a(\beta\xi) = \mathbf{e}(\text{Tr}(\beta\xi))$, we easily see that $\mathbf{e}_a(\beta\alpha)^\sigma = \mathbf{e}_a(\beta_1\alpha)$, which gives the desired fact.

REFERENCES

[1] E. Hecke, Bestimmung der Klassenzahl einer neuen Reihe von algebraischen Zahlkörpern, Nachr. der K. Gesellschaft der Wissenschaften zu Göttingen math.-phys. Klasse 1921, 1–23 (= Werke, 290–312).
 [2] E. Hecke, Zur Theorie der elliptischen Modulfunktionen, Math. Ann. 97 (1926), 210–242 (= Werke, 428–460).
 [3] E. Hecke, Analytische Arithmetik der positiven Quadratischen Formen, 1940, *Mathematische Werke*, 789–918.
 [4] Kronecker, Bemerkungen über die Darstellung von Reihen durch Integrale, Journal für r. angew. Math. 105 (1889), 157–159, 345–354 (=Werke V, 329–342).

- [5] E. Landau, Handbuch der Lehre von der Verteilung der Primzahlen, 1909.
- [6] G. Shimura, On the derivatives of theta functions and modular forms, *Duke Math. J.* 44 (1977), 365–387.
- [7] G. Shimura, Nearly holomorphic functions on hermitian symmetric spaces, *Math. Ann.* 278 (1987), 1–28.
- [8] G. Shimura, Arithmeticity in the theory of automorphic forms, *Mathematical Surveys and Monographs*, vol. 82, American Mathematical Society, 2000.
- [9] G. Shimura, *Elementary Dirichlet Series and Modular Forms*, Springer Monog. in Math., Springer, 2007.
- [10] C. L. Siegel, Bernoullische Polynome und quadratische Zahlkörper, *Nachr. der Akademie der Wissenschaften in Göttingen math.-phys. Klasse* 1968, Nr. 2, 7–38 (=Abhandlungen IV, 9–40).
- [11] C. L. Siegel, Zur Summation von L -Reihen, *Nachr. der Akademie der Wissenschaften in Göttingen math.-phys. Klasse* 1975, Nr. 18, 269–292 (=Abhandlungen IV, 305–328).

Goro Shimura
Department of Mathematics
Princeton University
Princeton
NJ 08544–1000
USA
goro@Math.Princeton.EDU