

FUNCTION FIELDS OF ONE VARIABLE OVER PAC FIELDS

MOSHE JARDEN AND FLORIAN POP

Received: July 8, 2009

Communicated by Peter Schneider

ABSTRACT. We give evidence for a conjecture of Serre and a conjecture of Bogomolov.

2000 Mathematics Subject Classification: 12E30, 20G15

Keywords and Phrases: Function fields of one variable, PAC fields, Conjecture II of Serre,

Conjecture II of Serre considers a field F of characteristic p with $\text{cd}(\text{Gal}(F)) \leq 2$ such that either $p = 0$ or $p > 0$ and $[F : F^p] \leq p$ and predicts that $H^1(\text{Gal}(F), G) = 1$ (i.e. each principal homogeneous G -space has an F -rational point) for each simply connected semi-simple linear algebraic group G [Ser97, p. 139].

As Serre notes, the hypothesis of the conjecture holds in the case where F is a field of transcendence degree 1 over a perfect field K with $\text{cd}(\text{Gal}(K)) \leq 1$. Indeed, in this case $\text{cd}(\text{Gal}(F)) \leq 2$ [Ser97, p. 83, Prop. 11] and $[F : F^p] \leq p$ if $p > 0$ (by the theory of p -bases [FrJ08, Lemma 2.7.2]). We prove the conjecture for F in the special case, where K is PAC of characteristic 0 that contains all roots of unity.

One of the main ingredients of the proof is the projectivity of $\text{Gal}(K(x)_{\text{ab}})$ (where x is transcendental over K and $K(x)_{\text{ab}}$ is the maximal Abelian extension of $K(x)$). We also use the same ingredient to establish an analog to the wellknown open problem of Shafarevich that $\text{Gal}(\mathbb{Q}_{\text{ab}})$ is free. Under the assumption that K is PAC and contains all roots of unity we prove that $\text{Gal}(K(x)_{\text{ab}})$ is not only projective but even free. This proves a stronger version of a conjecture of Bogomolov for a function field of one variable F over a PAC field that contains all roots of unity [Pos05, Conjecture 1.1].

1. THE PROJECTIVITY OF $\text{Gal}(K(x)_{\text{ab}})$

We denote the separable (resp. algebraic) closure of a field K by K_s (resp. \tilde{K}) and its absolute Galois group by $\text{Gal}(K)$. The field K is said to be PAC if every absolutely irreducible variety defined over K has a K -rational point. The proof of the projectivity result applies a local-global principle for Brauer groups to reduce the statement to Henselian fields.

For a prime number p and an Abelian group A , we say that A is p' -DIVISIBLE, if for each $a \in A$ and every positive integer n with $p \nmid n$ there exists $b \in A$ such that $a = nb$. Note that if $p = 0$, then “ p' -divisible” is the same as “divisible”.

LEMMA 1.1: *Let p be 0 or a prime number, B a torsion free Abelian group, and A is a p' -divisible subgroup of B of finite index. Then B is also p' -divisible.*

Proof: First suppose $p = 0$ and let $m = (B : A)$. Then, for each $b \in B$ and a positive integer n there exists $a \in A$ such that $mb = mna$. Since B is torsion free, $m = na$. Thus, B is divisible.

Now suppose p is a prime number, let $mp^k = (B : A)$, with $p \nmid m$ and $k \geq 0$, and consider $b \in B$. Then $mp^k b \in A$. Hence, for each positive integer n with $p \nmid n$ there exists $a \in A$ with $mp^k b = mna$. Thus, $p^k b = na$. Since $p \nmid n$, there exist $x, y \in \mathbb{Z}$ such that $xp^k + yn = 1$. It follows from $xp^k b = xna$ that $b = n(xa + yb)$, as claimed. \square

COROLLARY 1.2: *Let L/K be an algebraic field extension, v a valuation of L , and $p = 0$ or p is a prime number. Suppose that $v(K^\times)$ is p' -divisible. Then $v(L^\times)$ is p' -divisible.*

Proof: Let $x \in L^\times$ and n a positive integer with $p \nmid n$. Then $v(K(x)^\times)$ is a torsion free Abelian group and $v(K^\times)$ is a subgroup of index at most $[K(x) : K]$. Since $v(K^\times)$ is p' -divisible, Lemma 1.1 gives $y \in K(x)^\times$ such that $v(x) = nv(y)$. It follows that $v(L^\times)$ is p' -divisible. \square

Given a Henselian valued field (M, v) we use v also for its unique extension to M_s . We use a bar to denote the residue with respect to v of objects associated with M , let O_M be the valuation ring of M , and let $\Gamma_M = v(M^\times)$ be the value group of M .

We write $\text{cd}_l(K)$ and $\text{cd}(K)$ for the l th cohomological dimension and the cohomological dimension of $\text{Gal}(K)$ and note that $\text{cd}(K) \leq 1$ if and only if $\text{Gal}(K)$ is projective [Ser97, p. 58, Cor. 2].

LEMMA 1.3: *Let (M, v) be a Henselian valued field. Suppose $p = \text{char}(M) = \text{char}(\bar{M})$, $\text{Gal}(\bar{M})$ is projective, and Γ_M is p' -divisible. Then $\text{Gal}(M)$ is projective.*

Proof: We denote the INERTIA FIELD of M by M_u . It is determined by its absolute Galois group: $\text{Gal}(M_u) = \{\sigma \in \text{Gal}(M) \mid v(\sigma x - x) > 0 \text{ for all } x \in M_s \text{ with } v(x) \geq 0\}$. The map $\sigma \mapsto \bar{\sigma}$ of $\text{Gal}(M)$ into $\text{Gal}(\bar{M})$ such that $\bar{\sigma}x = \overline{\sigma x}$ for each $x \in O_M$ is a well defined epimorphism [Efr06, Thm. 16.1.1] whose kernel is $\text{Gal}(M_u)$. It therefore defines an isomorphism

$$(1) \quad \text{Gal}(M_u/M) \cong \text{Gal}(\bar{M}).$$

CLAIM A: \bar{M}_u is separably closed. Let $g \in \bar{M}[X]$ be a monic irreducible separable polynomial of degree $n \geq 1$. Then there exists a monic polynomial $f \in O_{M_u}[X]$ of degree n such that $\bar{f} = g$. We observe that f is also irreducible and separable. Moreover, if $f(X) = \prod_{i=1}^n (X - x_i)$ with $x_1, \dots, x_n \in M_s$, then $g(X) = \prod_{i=1}^n (X - \bar{x}_i)$. Given $1 \leq i, j \leq n$ there exists $\sigma \in \text{Gal}(M_u)$ such that $\sigma x_i = x_j$. By definition, $\bar{x}_j = \overline{\sigma x_i} = \bar{\sigma x_i} = \bar{x}_i$. Since g is separable, $i = j$, so $n = 1$. We conclude that \bar{M}_u is separably closed.

CLAIM B: Each l -Sylow group of $\text{Gal}(M_u)$ with $l \neq p$ is trivial. Indeed, let L be the fixed field of an l -Sylow group of $\text{Gal}(M_u)$ in M_s . If $l = 2$, then $\zeta_l = -1 \in L$. If $l \neq 2$, then $[L(\zeta_l) : L] | l - 1$ and $[L(\zeta_l) : L]$ is a power of l , so $\zeta_l \in L$.

Assume that $\text{Gal}(L) \neq 1$. By the theory of finite l -groups, L has a cyclic extension L' of degree l . By the preceding paragraph and Kummer theory, there exists $a \in L^\times$ such that $L' = L(\sqrt[l]{a})$. By Corollary 1.2, there exists $b \in L^\times$ such that $lv(b) = v(a)$. Then $c = \frac{a}{b^l}$ satisfies $v(c) = 0$. By Claim A, \bar{L} is separably closed. Therefore, \bar{c} has an l th root in \bar{L} . By Hensel's lemma, c has an l th root in L . It follows that a has an l -root in L . This contradiction implies that $L = M_s$, as claimed.

Having proved Claim B, we consider again a prime number $l \neq p$ and let G_l be an l -Sylow subgroup of $\text{Gal}(M)$. By the Claim, $G_l \cap \text{Gal}(M_u) = 1$, hence the map $\text{res} : \text{Gal}(M) \rightarrow \text{Gal}(M_u/M)$ maps G_l isomorphically onto an l -Sylow subgroup of $\text{Gal}(M_u/M)$. By (1), G_l is isomorphic to an l -Sylow subgroup of $\text{Gal}(\bar{M})$. Since the latter group is projective, so is G_l , i.e. $\text{cd}_l(G) \leq 1$ [Ser97, p. 58, Cor. 2].

Finally, if $p \neq 0$, then $\text{cd}_p(M) \leq 1$ [Ser97, p. 75, Prop. 3], because then $\text{char}(M) = p$. It follows that $\text{cd}(M) \leq 1$ [Ser97, p. 58, Cor. 2]. □

LEMMA 1.4: Let F be an extension of a PAC field K of transcendence degree 1 and characteristic p . Suppose $v(F^\times)$ is p' -divisible for each valuation v of F/K . Then $\text{Gal}(F)$ is projective.

Proof: Let K_{ins} be the maximal purely inseparable algebraic extension of K and set $F' = FK_{\text{ins}}$. Then K_{ins} is PAC [FrJ08, Cor. 11.2.5], $\text{trans.deg}(F'/K_{\text{ins}}) = 1$, and $v((F')^\times)$ is p' -divisible for every valuation v of F' (by Corollary 1.2). Moreover, $\text{Gal}(F') = \text{Gal}(F)$. Thus, we may replace K by K_{ins} and F by F' , if necessary, to assume that K is perfect.

Let $V(F/K)$ be a system of representatives of the equivalence classes of valuations of F that are trivial on K . For each $v \in V(F/K)$ we choose a Henselian closure F_v of F at v . By [Efr01, Thm. 3.4], there is an injection of Brauer groups,

$$(2) \quad \text{Br}(F) \rightarrow \prod_{v \in V(F/K)} \text{Br}(F_v).$$

For each $v \in V(F/K)$ we have, $v(F_v^\times) = v(F^\times)$ is p' -divisible. Also, the residue field \bar{F}_v is an algebraic extension of K . Since K is PAC, a theorem of Ax says

that $\text{Gal}(K)$ is projective [FrJ08, Thm. 11.6.2], hence $\text{Gal}(\bar{F}_v)$ is projective [FrJ08, Prop. 22.4.7]. Finally, $\text{char}(F_v) = \text{char}(\bar{F}_v)$. Therefore, by Lemma 1.3, $\text{Gal}(F_v)$ is projective, hence $\text{Br}(F_v) = 0$ [Ser97, p. 78, Prop. 5]. It follows from the injectivity of (2) that $\text{Br}(F) = 0$.

If F_1 is a finite separable extension of F , $v_1 \in V(F_1/K)$, and $v = v_1|_F$, then $v(F^\times)$ is p' -divisible. Hence, by Corollary 1.2, $v_1((F_1)^\times)$ is p' -divisible. It follows from the preceding paragraph that $\text{Br}(F_1) = 0$. Consequently, by [Ser97, p. 78, Prop. 5], $\text{cd}(\text{Gal}(F)) \leq 1$. \square

LEMMA 1.5: *Let p be either 0 or a prime number and let Γ be an additive subgroup of \mathbb{Q} . Suppose $\frac{1}{n} \in \Gamma$ for each positive integer n with $p \nmid n$. Then Γ is p' -divisible.*

Proof: We consider $\gamma \in \Gamma$. If $p = 0$, we write $\gamma = \frac{a}{b}$, with $a \in \mathbb{Z}$ and $b \in \mathbb{N}$. Given $n \in \mathbb{N}$, we have $\frac{\gamma}{n} = a \cdot \frac{1}{nb} \in \Gamma$.

If $p > 0$, we write $\gamma = \frac{a}{bp^k}$, where $a \in \mathbb{Z}$, $b \in \mathbb{N}$, $k \in \mathbb{Z}$, and $p \nmid a, b$. Let $n \in \mathbb{N}$ with $p \nmid n$. If $k \leq 0$, then $\frac{\gamma}{n} = ap^{-k} \cdot \frac{1}{nb} \in \Gamma$. If $k > 0$, we may choose $x, y \in \mathbb{Z}$ such that $xp^k + ynb = 1$. Then $\frac{\gamma}{n} = \frac{a}{nbp^k} = \frac{axp^k + aynb}{nbp^k} = ax \cdot \frac{1}{nb} + by \cdot \frac{a}{bp^k} \in \Gamma$, as claimed. \square

PROPOSITION 1.6: *Let K be a PAC field that contains all roots of unity and let E be an extension of K of transcendence degree 1. Then $\text{Gal}(E_{\text{ab}})$ is projective.*

Proof: First we consider the case where $E = K(x)$, where x is transcendental over K , and set $F = E_{\text{ab}}$. In the notation of Lemma 1.4 we consider a valuation $v \in V(F/K)$ normalized in such a way that $v(E^\times) = \mathbb{Z}$. Then $v(F^\times) \leq \mathbb{Q}$. On the other hand, let $p = \text{char}(K)$ and consider a positive integer n with $p \nmid n$. Let $e \in E$ with $v(e) = 1$. Then $e^{1/n} \in F$ (because K contains a root of 1 of order n). Therefore, $\frac{1}{n} = v(e^{1/n}) \in v(F^\times)$. By Lemma 1.5, $v(F^\times)$ is p' -divisible. We conclude from Lemma 1.4 that $\text{Gal}(F)$ is projective.

In the general case we choose $x \in E$ transcendental over K . By the preceding paragraph, $\text{Gal}(K(x)_{\text{ab}})$ is projective. Since taking purely inseparable extensions of a field does not change its absolute Galois group, $\text{Gal}(K(x)_{\text{ab,ins}})$ is projective. Now note that $\text{Gal}(E_{\text{ab,ins}})$ as a subgroup of $\text{Gal}(K(x)_{\text{ab,ins}})$ is also projective. Hence, $\text{Gal}(E_{\text{ab}})$ is projective. \square

Remark 1.7: Proposition 1.6 is false if K does not contain all roots of unity. Indeed, the authors will elsewhere provide an example of a prime number l and a PAC field K of characteristic 0 that contains all roots of unity of order n with $l \nmid n$ but not ζ_l such that $\text{Gal}(K(x)_{\text{ab}})$ is not projective. \square

2. SERRE AND SHAFAREVICH

We refer to a simply connected semi-simple linear algebraic group G as a SIMPLY CONNECTED GROUP. In this case $H^1(\text{Gal}(K), G)$ will be also denoted by $H^1(K, G)$. Since each element of $H^1(K, G)$ is represented by a principal homogeneous space V of G and V is an absolutely irreducible variety defined over

K , V has a K -rational point if K is PAC. Hence, V is equivalent to G [LaT58, Prop. 4]. Thus, $H^1(K, G) = 1$.

The proof of Serre’s Conjecture II in our case is based on the following consequence of a theorem of Colliot-Thélène, Gille, and Parimala:

PROPOSITION 2.1: *Let F be a field and G a simply connected group defined over F . Suppose F is a C_2 -field of characteristic 0, $\text{cd}(F) \leq 2$, and $\text{cd}(F_{\text{ab}}) \leq 1$. Then $H^1(F, G) = 1$.*

Proof: Let F' be a finite extension of F . Since F is C_2 , [CGP04, Thm. 1.1(vi)] implies that if the exponent e of a central simple algebra A over F' is a power of 2 or a power of 3, then e is equal to the index of A . Since $\text{cd}(F) \leq 2$ and $\text{cd}(F_{\text{ab}}) \leq 1$, [CGP04, Thm. 1.2(v)] implies that $H^1(F, G) = 1$. □

Remark 2.2: By Merkuriev-Suslin, the assumption that F is a C_2 -field implies that $\text{cd}(F) \leq 2$ [Ser97, end of page 88]. However, we will be able to prove both properties of F directly in the application we have in mind. □

The following result establishes the first condition on F .

LEMMA 2.3: *Let F be an extension of transcendence degree 1 over a perfect PAC field K . Suppose either $\text{char}(K) > 0$ and K contains all roots of unity or $\text{char}(K) = 0$. Then $\text{cd}(F) \leq 2$ and F is a C_2 -field.*

Proof: By Ax, $\text{cd}(K) \leq 1$ [FrJ08, Thm. 11.6.2]. Hence, by [Ser97, p. 83, Prop. 11], $\text{cd}(F) \leq 2$.

A conjecture of Ax from 1968 says that every perfect PAC field K is C_1 [FrJ08, Problem 21.2.5]. The conjecture holds if K contains an algebraically closed field [FrJ08, Lemma 21.3.6(a)]. In particular, if $p = \text{char}(K) > 0$ and K contains all roots of unity, then $\overline{\mathbb{F}}_p \subseteq K$, so K is C_1 . If $\text{char}(K) = 0$, K is C_1 , by [Kol07, Thm. 1]. It follows that in each case, F is C_2 [FrJ08, Prop. 21.2.12]. □

THEOREM 2.4: *Let F be an extension of transcendence degree 1 of a PAC field K of characteristic 0. Suppose K contains all roots of unity. Then F satisfies Serre’s conjecture II. That is, $H^1(F, G) = 1$ for each simply connected group G defined over F .*

Proof: By Lemma 2.3, $\text{cd}(F) \leq 2$ and F is a C_2 -field. By Proposition 1.6, $\text{cd}(F_{\text{ab}}) \leq 1$. It follows from Proposition 2.1 that $H^1(F, G) = 1$ for each simply connected group G . □

Remark 2.5: All of the ingredients of the proof of Theorem 2.4 except possibly Proposition 2.1 work also when $\text{char}(K) > 0$. □

The proof of the freeness of $\text{Gal}(K(x)_{\text{ab}})$ applies the notion of ”quasi-freeness” due to Harbater and Stevenson. To this end recall that a FINITE SPLIT EMBEDDING PROBLEM \mathcal{E} for a profinite group G is a pair $(\varphi: G \rightarrow A, \alpha: B \rightarrow A)$, where A, B are finite groups, φ, α are epimorphisms, and α has a group theoretic section. A SOLUTION of \mathcal{E} is an epimorphism $\gamma: G \rightarrow B$ such that

$\alpha \circ \gamma = \varphi$. We say that G is QUASI-FREE if its rank m is infinite and every finite split embedding problem for G has m distinct solutions.

THEOREM 2.6: *Let F be a function field of one variable over a PAC field K of cardinality m containing all roots of unity and let x be a variable. Then $\text{Gal}(F_{\text{ab}})$ is isomorphic to the free profinite group of rank m .*

Proof: Since K is PAC, K is AMPLE, that is every absolutely irreducible curve defined over K with a K -rational simple point has infinitely many K -rational points. By [HaS05, Cor. 4.4], $\text{Gal}(F)$ is quasi-free of rank $m = \text{card}(K)$. Hence, by [Har09, Thm. 2.4], $\text{Gal}(F_{\text{ab}})$ is also quasi-free of rank m . Since by Proposition 1.6, $\text{Gal}(F_{\text{ab}})$ is projective, it follows from a result of Chatzidakis and Melnikov [FrJ08, Lemma 25.1.8] that $\text{Gal}(F_{\text{ab}})$ is free of rank m . \square

Acknowledgment: The authors thank Jean-Louis Colliot-Thélène for stimulating discussions, in particular for pointing out Proposition 2.1 to them. \square

REFERENCES

- [CGP04] J.-L. Colliot-Thélène, P. Gille, and R. Parimala, *Arithmetic of linear algebraic groups over two-dimensional geometric fields*, Duke Mathematical Journal 121 (2004), 285–341.
- [Efr01] I. Efrat, *A Hasse principle for function fields over PAC fields*, Israel Journal of Mathematics 122 (2001), 43–60.
- [Efr06] I. Efrat, *Valuations, Orderings, and Milnor K -Theory*, Mathematical surveys and monographs 124, American Mathematical Society, Providence, 2006.
- [FrJ08] M. D. Fried and M. Jarden, *Field Arithmetic, Third Edition, revised by Moshe Jarden*, Ergebnisse der Mathematik (3) 11, Springer, Heidelberg, 2008.
- [Har09] D. Harbater, *On function fields with free absolute Galois groups*, Journal für die reine und angewandte Mathematik 632 (2009), 85–103.
- [HaS05] D. Harbater and K. Stevenson, *Local Galois theory in dimension two*, Advances in Mathematics 198 (2005), 623–653.
- [Kol07] J. Kollár, *A conjecture of Ax and degenerations of Fano varieties*, Israel Journal of Mathematics 162 (2007), 235–251.
- [LaT58] S. Lang and J. Tate, *Principal homogeneous spaces over abelian varieties*, American Journal of Mathematics 80 (1958), 659–684.
- [Pos05] L. Positselski, *Koszul property and Bogomolov’s conjecture*, International Mathematics Research Notices 31 (2005), 1901–1936.
- [Ser97] J.-P. Serre, *Galois Cohomology*, Springer, Berlin, 1997.

Moshe Jarden
School of Mathematics
Tel Aviv University
Ramat Aviv, Tel Aviv 69978, Israel
jarden@post.tau.ac.il

Florian Pop
Department of Mathematics
University of Pennsylvania
Philadelphia, Pennsylvania 19104,
USA
pop@math.upenn.edu

