

THE FIRST L^2 -BETTI NUMBER
AND APPROXIMATION IN ARBITRARY CHARACTERISTIC¹

MIKHAIL ERSHOV, WOLFGANG LÜCK

Received: September 23, 2012

Communicated by Stefan Schwede

ABSTRACT. Let G be a finitely generated group and $G = G_0 \supseteq G_1 \supseteq G_2 \supseteq \cdots$ a descending chain of finite index normal subgroups of G . Given a field K , we consider the sequence $\frac{b_1(G_i; K)}{[G:G_i]}$ of normalized first Betti numbers of G_i with coefficients in K , which we call a K -approximation for $b_1^{(2)}(G)$, the first L^2 -Betti number of G . In this paper we address the questions of when \mathbb{Q} -approximation and \mathbb{F}_p -approximation have a limit, when these limits coincide, when they are independent of the sequence (G_i) and how they are related to $b_1^{(2)}(G)$. In particular, we prove the inequality $\lim_{i \rightarrow \infty} \frac{b_1(G_i; \mathbb{F}_p)}{[G:G_i]} \geq b_1^{(2)}(G)$ under the assumptions that $\bigcap G_i = \{1\}$ and each G/G_i is a finite p -group.

2010 Mathematics Subject Classification: Primary: 20F65; Secondary: 46Lxx

Keywords and Phrases: First L^2 -Betti number, approximation in prime characteristic

1. INTRODUCTION

1.1. \mathbb{Q} -APPROXIMATION FOR THE FIRST L^2 -BETTI NUMBER. Let G be a finitely generated group. Given a field K , we let $b_1(G; K) = \dim_K(H_1(G; K))$ be the first Betti number of G with coefficients in K and $b_1(G) = b_1(G; \mathbb{Q})$ where \mathbb{Q} denotes the field of rational numbers. Denote by $b_1^{(2)}(G)$ the *first L^2 -Betti number of G* . Assuming that G is finitely presented and residually finite, by Lück Approximation Theorem (see [13]), $b_1^{(2)}(G)$ can be approximated by normalized rational first Betti numbers of finite index subgroups of G :

¹The first author is supported in part by the NSF grant DMS-0901703 and the Sloan Research Fellowship grant BR 2011-105. This paper is financially supported by the Leibniz-Preis of the second author. We are grateful to Andrei Jaikin-Zapirain for this observation.

THEOREM 1.1 (Lück approximation theorem). *Let G be a finitely presented residually finite group and $G = G_0 \supseteq G_1 \supseteq \dots$ a descending chain of finite index normal subgroups of G , with $\bigcap_{i \in \mathbb{N}} G_i = \{1\}$. Then*

$$(1.2) \quad b_1^{(2)}(G) = \lim_{i \rightarrow \infty} \frac{b_1(G_i)}{[G : G_i]}.$$

In the sequel we will occasionally refer to a descending chain (G_i) of finite index normal subgroups of G as a *finite index normal chain* in G and to the associated sequence $\left(\frac{b_1(G_i)}{[G : G_i]}\right)_i$ as \mathbb{Q} -*approximation*.

If we drop the assumption that G is finitely presented, but still require that $\bigcap_{i \in \mathbb{N}} G_i = \{1\}$, one still has inequality $b_1^{(2)}(G) \geq \limsup_{i \rightarrow \infty} \frac{b_1(G_i)}{[G : G_i]}$ by [16, Theorem 1.1], but equality need not hold [16, Theorem 1.2]. The latter is proved in [16] by constructing an example where $b_1^{(2)}(G) > 0$, but $\limsup_{i \rightarrow \infty} \frac{b_1(G_i)}{[G : G_i]} = 0$ for any chain (G_i) as above. In Section 5 we will describe a variation of this construction showing that the \mathbb{Q} -approximation $\left(\frac{b_1(G_i)}{[G : G_i]}\right)_i$ may not even have a limit:

THEOREM 1.3. *There exists a finitely generated residually finite group G and a descending chain $(G_i)_{i \in \mathbb{N}}$ of finite index normal subgroups of G , with $\bigcap_{i \in \mathbb{N}} G_i = \{1\}$, such that $\lim_{i \rightarrow \infty} \frac{b_1(G_i)}{[G : G_i]}$ does not exist.*

Another sequence we shall be interested in is \mathbb{F}_p -*approximation*, that is, $\left(\frac{b_1(G_i; \mathbb{F}_p)}{[G : G_i]}\right)_i$, where \mathbb{F}_p is the finite field of prime order p . This sequence is particularly important under the additional assumption that (G_i) is a *p-chain*, that is, each G_i has p -power index (equivalently, G/G_i is a finite p -group). In this case, $\left(\frac{b_1(G_i; \mathbb{F}_p)}{[G : G_i]}\right)_i$ is monotone decreasing and therefore has a limit, often called *p-gradient* or *mod p homology gradient* (see, e.g., [11]).

Since obviously $b_1(H) \leq b_1(H; \mathbb{F}_p)$ for any group H , one always has inequality

$$(1.4) \quad \limsup_{i \rightarrow \infty} \frac{b_1(G_i)}{[G : G_i]} \leq \limsup_{i \rightarrow \infty} \frac{b_1(G_i; \mathbb{F}_p)}{[G : G_i]},$$

and it is natural to ask for sufficient conditions under which equality holds. Of particular interest is the case when G is finitely presented and $\bigcap_{i \in \mathbb{N}} G_i = \{1\}$ when \mathbb{Q} -approximation does have a limit by Theorem 1.1.

QUESTION 1.5 (\mathbb{Q} -approximation and \mathbb{F}_p -approximation). For which finitely presented groups G and finite index normal chains (G_i) with $\bigcap_{i \in \mathbb{N}} G_i = \{1\}$ do we have equality

$$\lim_{i \rightarrow \infty} \frac{b_1(G_i)}{[G : G_i]} = \lim_{i \rightarrow \infty} \frac{b_1(G_i; \mathbb{F}_p)}{[G : G_i]}?$$

If G is not finitely presented, the above equality need not hold even if we require that (G_i) is a p -chain. Indeed, as proved in [18] and independently in [20], there exists a p -torsion residually- p group G with $\lim_{i \rightarrow \infty} \frac{b_1(G_i; \mathbb{F}_p)}{[G : G_i]} > 0$

for any p -chain (G_i) in G (and since G is residually- p , we can choose a p -chain with $\bigcap G_i = \{1\}$). Since $b_1(H) = 0$ for any torsion group H , we have $\lim_{i \rightarrow \infty} \frac{b_1(G_i)}{[G:G_i]} = 0$ for such group G .

In Section 4 we give an example showing that the answer to Question 1.5 would also become negative if we drop the assumption $\bigcap_{i \in \mathbb{N}} G_i = \{1\}$, even if G is finitely presented and (G_i) is a p -chain which has infinitely many distinct terms.

1.2. COMPARING \mathbb{F}_p -APPROXIMATION AND FIRST L^2 -BETTI NUMBER. Since both \mathbb{F}_p -approximation and the first L^2 -Betti number provide upper bounds for \mathbb{Q} -approximation, it is natural to ask how the former two quantities are related to each other. We address this question in the case of p -chains.

THEOREM 1.6. *Let p be a prime number. Let G be a finitely generated group and $G = G_0 \supseteq G_1 \supseteq G_2 \supseteq \dots$ a descending chain of normal subgroups of G of p -power index. Then*

- (1) *The sequence $\left(\frac{b_1(G_i; \mathbb{F}_p)}{[G:G_i]}\right)_i$ is monotone decreasing and therefore converges;*
- (2) *Assume that $\bigcap_{i \in \mathbb{N}} G_i = \{1\}$. Then*

$$b_1^{(2)}(G) \leq \lim_{i \rightarrow \infty} \frac{b_1(G_i; \mathbb{F}_p)}{[G : G_i]}.$$

We note that for finitely presented groups Theorem 1.6(2) is a straightforward consequence of Theorem 1.1.

We provide two different proofs of Theorem 1.6. First, Theorem 1.6 is a special case of Theorem 2.2, which will be proved in Section 2. An alternative proof of Theorem 1.6 given in Section 3 will be based on Theorem 3.1. The latter may be of independent interest and has another important corollary, which can be considered as an extension of Theorem 1.1 to groups which are finitely presented, but not necessarily residually finite. Here is a slightly simplified version of Theorem 3.1.

THEOREM 1.7. *Let G be a finitely presented group, and let K be the kernel of the canonical map from G to its profinite completion or pro- p completion for some prime p . Let (G_i) be a descending chain of finite index normal subgroups of G such that $\bigcap_{i \in \mathbb{N}} G_i = K$ (note that such a chain always exists). Then*

$$b_1^{(2)}(G/K) = \lim_{i \rightarrow \infty} \frac{b_1(G_i)}{[G : G_i]}.$$

1.3. CONNECTION WITH RANK GRADIENT. Let G be a finitely generated group. In the sequel we denote by $d(G)$ the minimal number of generators, sometimes also called *the rank of G* . Let $(G_i)_{i \in \mathbb{N}}$ be a descending chain of finite index normal subgroups of G . The *rank gradient of G* (with respect to (G_i)), denoted by $\text{RG}(G; (G_i))$, is defined by

$$(1.8) \quad \text{RG}(G; (G_i)) = \lim_{i \rightarrow \infty} \frac{d(G_i) - 1}{[G : G_i]}.$$

The above limit always exists since for any finite index subgroup H of G one has $\frac{d(H)-1}{[G:H]} \leq d(G) - 1$ by the Schreier index formula.

Rank gradient was originally introduced by Lackenby [10] as a tool for studying 3-manifold groups, but is also interesting from a purely group-theoretic point of view (see, e.g., [1, 2, 18, 20]).

Provided that G is infinite and $\bigcap_{i \in \mathbb{N}} G_i = \{1\}$, the following inequalities are known to hold:

$$(1.9) \quad \text{RG}(G; (G_i)) \geq \text{cost}(G) - 1 \geq b_1^{(2)}(G).$$

The first inequality was proved by Abért and Nikolov [2, Theorem 1], and the second one is due to Gaboriau [8, Corollaire 3.16, 3.23] (see [7, 8, 9] for the definition and some key results about cost).

It is not known if either inequality in (1.9) can be strict. In particular, the following question is open.

QUESTION 1.10. Let G be an infinite finitely generated residually finite group and (G_i) a descending chain of finite index normal subgroups of G with $\bigcap_{i \in \mathbb{N}} G_i = \{1\}$. Is it always true that

$$\text{RG}(G; (G_i)) = b_1^{(2)}(G)?$$

Theorem 1.6 provides a potentially new approach for answering Question 1.10 in the negative, as explained below.

In view of the obvious inequality $d(H) \geq b_1(H; K)$ for any group H and any field K , one always has $\text{RG}(G; (G_i)) \geq \limsup_{i \rightarrow \infty} \frac{b_1(G_i; K)}{[G:G_i]}$.

QUESTION 1.11. For which infinite finitely generated groups G , finite index normal chains $(G_i)_{i \in \mathbb{N}}$ with $\bigcap_{i \in \mathbb{N}} G_i = \{1\}$ and fields K , do we have

$$(1.12) \quad \text{RG}(G; (G_i)) = \limsup_{i \rightarrow \infty} \frac{b_1(G_i; K)}{[G:G_i]}?$$

REMARK 1.13. Since for a group H , the first Betti number $b_1(H; K)$ depends only on the characteristic of K , one can assume that $K = \mathbb{Q}$ or $K = \mathbb{F}_p$ for some p . The same remark applies to Question 1.14 below.

Note that if $K = \mathbb{Q}$, equality (1.12) does not hold in general – if it did, Theorem 1.3 would have implied the existence of a group G and a finite index normal chain (G_i) in G for which the sequence $\left(\frac{d(G_i)-1}{[G:G_i]}\right)_i$ has no limit, which is impossible since this sequence is monotone decreasing. If one can find a group G for which (1.12) fails with $K = \mathbb{F}_p$ and (G_i) a p -chain, then in view of Theorem 1.6 such group G would answer Question 1.10 in the negative.

The answer to Question 1.11 would become negative if we drop the assumption $\bigcap G_i = \{1\}$ even if G is finitely presented and (G_i) is a p -chain (with infinitely many distinct terms), as we will see in Section 4.

1.4. INDEPENDENCE OF THE CHAIN. So far we discussed the dependence of the quantity $\limsup_{i \rightarrow \infty} \frac{b_1(G_i; K)}{[G:G_i]}$ on the field K , but perhaps an even more important question is when it is independent of the chain. Again it is reasonable to require that $\bigcap_{i \in \mathbb{N}} G_i = \{1\}$ since without this restriction the answer would be negative already for very nice groups like $F \times \mathbb{Z}$, where F is a non-abelian free group. Note that independence of $\limsup_{i \rightarrow \infty} \frac{b_1(G_i; K)}{[G:G_i]}$ of the chain (G_i) as above automatically implies that $\lim_{i \rightarrow \infty} \frac{b_1(G_i; K)}{[G:G_i]}$ must exist.

QUESTION 1.14. For which finitely generated residually finite groups G and fields K does the limit $\lim_{i \rightarrow \infty} \frac{b_1(G_i; K)}{[G:G_i]}$ exist for all finite index normal chains $(G_i)_{i \in \mathbb{N}}$ with $\bigcap_{i \in \mathbb{N}} G_i = \{1\}$ and is independent of the choice of the chain (G_i) ?

The answer to Question 1.14 is known to be positive if $K = \mathbb{Q}$ and either G is finitely presented (by Theorem 1.1) or G is a limit of left orderable amenable groups in the space of marked group presentations, in which case equality (1.2) holds by [19, Corollary 1.5]. Question 1.14 remains open if G is finitely presented and $K = \mathbb{F}_p$. If G is arbitrary, the answer may be negative for any K – this follows directly from Theorem 1.3 if $K = \mathbb{Q}$ and from its stronger version Theorem 5.1 if $K = \mathbb{F}_p$. In the latter case, however, it is natural to impose the additional assumption that (G_i) is a p -chain, which does not hold in our examples.

Essentially the only case when answer to Question 1.14 is known to be positive for all fields is when G contains a normal infinite amenable subgroup (e.g., if G itself is infinite amenable). In this case, $\text{RG}(G; (G_i)) = 0$ for all finite index normal chains (G_i) with trivial intersection, as proved by Lackenby [10, Theorem 1.2] when G is finitely presented and by Abért and Nikolov [2, Theorem 3] in general. This, of course, implies that in such groups $\lim_{i \rightarrow \infty} \frac{b_1(G_i; K)}{[G:G_i]} = 0$ for any such chain (G_i) and hence the answer to Questions 1.11 and 1.14 is positive.

Finally, we comment on the status of a more general version of Question 1.14:

QUESTION 1.15. For which residually finite groups G , fields K , finite index normal chains (G_i) with $\bigcap_{i \in \mathbb{N}} G_i = \{1\}$, free G -CW-complexes X of finite type and natural numbers n , does the limit $\lim_{i \rightarrow \infty} \frac{b_n(G_i \setminus X; K)}{[G:G_i]}$ exist and is independent of the chain?

Again, if K has characteristic zero, the answer is always yes and the limit can be identified with the n -th L^2 -Betti number $b_n^{(2)}(X; \mathcal{N}(G))$ (see [13] or [14, Theorem 13.3 (2) on page 454], which is a generalization of Theorem 1.1). If K has positive characteristic, the answer is yes if G is virtually torsion-free elementary amenable, in which case the limit can be identified with the Ore dimension of $H_n(X; K)$ (see [12, Theorem 5.3]); the answer is also yes for any finitely generated amenable group G – this follows from [1, Theorem 17] or [12, Theorem 2.1] – and the limit can be described using Elek dimension function (see [5]). There are examples for $G = \mathbb{Z}$ of finite G -CW-complexes X where

the limits $\lim_{i \rightarrow \infty} \frac{b_n(G_i \backslash X; K)}{[G : G_i]}$ are different for $K = \mathbb{Q}$ and $K = \mathbb{F}_p$ (but X is not EG), see [12, Example 6.2].

1.5. ACKNOWLEDGMENTS. The authors want to thank the American Institute of Mathematics for its hospitality during their stay at the Workshop “ L^2 -invariants and their relatives for finitely generated groups” organized by Miklós Abért, Mark Sapir, and Dimitri Shlyakhtenko in September 2011, where some of the ideas of this paper were developed. The authors are very grateful to Denis Osin for proposing several improvements in Section 4 and other useful discussions. The first author is very grateful to Andrei Jaikin-Zapirain for many helpful discussions related to the subject of this paper, sending his unpublished work “On p -gradient of finitely presented groups” and suggesting a stronger version of Theorem 3.1(2).

2. THE FIRST L^2 -BETTI NUMBER AND APPROXIMATION IN PRIME CHARACTERISTIC

If G is a group and X a G -CW-complex, we denote by

$$(2.1) \quad b_n^{(2)}(X; \mathcal{N}(G)) = \dim_{\mathcal{N}(G)}(H_n(\mathcal{N}(G) \otimes_{\mathbb{Z}G} C_*(X)))$$

its n -th L^2 -Betti number. Here $C_*(X)$ is the cellular $\mathbb{Z}G$ -chain complex of X , $\mathcal{N}(G)$ is the group von Neumann algebra and $\dim_{\mathcal{N}(G)}$ is the dimension function for (algebraic) $\mathcal{N}(G)$ -modules in the sense of [14, Theorem 6.7 on page 239]. Notice that $b_1^{(2)}(G) = b_1^{(2)}(EG; \mathcal{N}(G))$.

The goal of this section is to prove the following theorem which generalizes Theorem 1.6:

THEOREM 2.2 (The first L^2 -Betti number and \mathbb{F}_p -approximation). *Let p be a prime number. Let G be a finitely generated group and (G_i) a descending chain of normal subgroups of p -power index in G . Let $K = \bigcap_{i \in \mathbb{N}} G_i$. Then the sequence $\left(\frac{b_1(G_i; \mathbb{F}_p)}{[G : G_i]} \right)_i$ is monotone decreasing, the limit $\lim_{i \rightarrow \infty} \frac{b_1(G_i; \mathbb{F}_p)}{[G : G_i]}$ exists and satisfies*

$$b_1^{(2)}(K \backslash EG; \mathcal{N}(G/K)) \leq \lim_{i \rightarrow \infty} \frac{b_1(G_i; \mathbb{F}_p)}{[G : G_i]}.$$

For its proof we will need the following lemma, which is proved in [3, Lemma 4.1], although it was probably well known before.

LEMMA 2.3. *Let p be a prime and m, n positive integers. Let H be a finite p -group. Consider an $\mathbb{F}_p H$ -map $\alpha: \mathbb{F}_p H^m \rightarrow \mathbb{F}_p H^n$. Define the \mathbb{F}_p -map*

$$\bar{\alpha} = \text{id}_{\mathbb{F}_p} \otimes_{\mathbb{F}_p H} \alpha: \mathbb{F}_p^m = \mathbb{F}_p \otimes_{\mathbb{F}_p H} \mathbb{F}_p H^m \rightarrow \mathbb{F}_p^n = \mathbb{F}_p \otimes_{\mathbb{F}_p H} \mathbb{F}_p H^n,$$

where we consider \mathbb{F}_p as $\mathbb{F}_p H$ -module by the trivial H -action. Then

$$\dim_{\mathbb{F}_p}(\text{im}(\alpha)) \geq |H| \cdot \dim_{\mathbb{F}_p}(\text{im}(\bar{\alpha})).$$

Notice that the assertion of Lemma 2.3 is not true if we do not require that H is a p -group or if we replace \mathbb{F}_p by a field of characteristic not equal to p .

Proof of Theorem 2.2. Since G is finitely generated, there is a CW -model for BG with one 0-cell and a finite number, let us say s , of 1-cells. Let $EG \rightarrow BG$ be the universal covering. Put $X = K \setminus EG$ and $Q = G/K$. Then X is a free Q - CW -complex with finite 1-skeleton. Its cellular $\mathbb{Z}Q$ -chain complex $C_*(X)$ looks like

$$\cdots \rightarrow C_2(X) = \bigoplus_{j=1}^r \mathbb{Z}Q \xrightarrow{c_2} C_1(X) = \bigoplus_{j=1}^s \mathbb{Z}Q \xrightarrow{c_1} C_0(X) = \mathbb{Z}Q$$

where r is a finite number or infinity.

For $m = 0, 1, 2, \dots$ we define a $\mathbb{Z}Q$ -submodule of $C_2(X)$ by $C_2(X)|_m = \bigoplus_{j=1}^{\max\{m,r\}} \mathbb{Z}Q$. Denote by $c_{2|m}: C_2(X)|_m \rightarrow C_1(X)$ the restriction of c_2 to $C_2(X)|_m$.

Consider a $\mathbb{Z}Q$ -map $f: M \rightarrow N$. Denote by $f^{(2)}: M^{(2)} \rightarrow N^{(2)}$ the $\mathcal{N}(Q)$ -homomorphism $\text{id}_{\mathcal{N}(G)} \otimes_{\mathbb{Z}Q} f: \mathcal{N}(Q) \otimes_{\mathbb{Z}Q} M \rightarrow \mathcal{N}(Q) \otimes_{\mathbb{Z}Q} N$. Put $Q_i = G_i/K$. Let $f[i]: M[i] \rightarrow N[i]$ be the \mathbb{Q} -homomorphism $\text{id}_{\mathbb{Q}} \otimes f: \mathbb{Q} \otimes_{\mathbb{Z}[Q_i]} M \rightarrow \mathbb{Q} \otimes_{\mathbb{Z}[Q_i]} N$. Denote by $f[i, p]: M[i, p] \rightarrow N[i, p]$ the \mathbb{F}_p -homomorphism $\text{id}_{\mathbb{F}_p} \otimes_{\mathbb{Z}[Q_i]} f: \mathbb{F}_p \otimes_{\mathbb{Z}[Q_i]} M \rightarrow \mathbb{F}_p \otimes_{\mathbb{Z}[Q_i]} N$. If $M = \bigoplus_{j=1}^t \mathbb{Z}Q$, then $M^{(2)} = \bigoplus_{j=1}^t \mathcal{N}(Q)$, $M[i] = \bigoplus_{j=1}^t \mathbb{Z}[Q/Q_i]$ and $M[i, p] = \bigoplus_{j=1}^t \mathbb{F}_p[Q/Q_i]$.

Note that

$$b_1(Q_i \setminus X; \mathbb{F}_p) = b_1(G_i \setminus EG; \mathbb{F}_p) = b_1(BG_i; \mathbb{F}_p) = b_1(G_i; \mathbb{F}_p).$$

Since all dimension functions are additive (see [14, Theorem 6.7 on page 239]), we conclude

$$(2.4) \quad b_1^{(2)}(X; \mathcal{N}(Q)) = s - 1 - \dim_{\mathcal{N}(Q)}(\text{im}(c_2^{(2)}));$$

$$(2.5) \quad \frac{b_1(G_i; \mathbb{F}_p)}{[Q : Q_i]} = s - 1 - \frac{\dim_{\mathbb{F}_p}(\text{im}(c_{2|i,p}))}{[Q : Q_i]};$$

$$(2.6) \quad \dim_{\mathcal{N}(Q)}(\text{im}(c_{2|m}^{(2)})) = m - \dim_{\mathcal{N}(Q)}(\ker(c_{2|m}^{(2)}));$$

$$(2.7) \quad \frac{\dim_{\mathbb{Q}}(\text{im}(c_{2|m}[i]))}{[Q : Q_i]} = m - \frac{\dim_{\mathbb{Q}}(\ker(c_{2|m}[i]))}{[Q : Q_i]};$$

$$(2.8) \quad \frac{\dim_{\mathbb{F}_p}(\text{im}(c_{2|m}[i, p]))}{[Q : Q_i]} = m - \frac{\dim_{\mathbb{F}_p}(\ker(c_{2|m}[i, p]))}{[Q : Q_i]}.$$

There is an isomorphism of \mathbb{F}_p -chain complexes $\mathbb{F}_p \otimes_{\mathbb{F}_p[Q_{i+1} \setminus Q_i]} C_*(X)[(i + 1), p] \xrightarrow{\cong} C_*(X)[i, p]$, where the $Q_{i+1} \setminus Q_i$ -operation on $C_*(X)[i + 1]$ comes from the identification $C_*(X)[i + 1] = \mathbb{F}_p \otimes_{\mathbb{F}_p[Q_{i+1}]} C_*(X) = \mathbb{F}_p[Q_{i+1} \setminus Q] \otimes_{\mathbb{F}_p Q} C_*(X)$. This is compatible with the passage from $C_2(X)$ to $C_2(X)|_m$. Hence $c_{2|m}[i, p]$ can be identified with $\text{id}_{\mathbb{F}_p} \otimes_{\mathbb{F}_p[Q_{i+1} \setminus Q_i]} c_{2|m}[(i + 1), p]$. Since $Q_{i+1} \setminus Q_i$ is a finite p -group, Lemma 2.3 implies

$$\dim_{\mathbb{F}_p}(\text{im}(c_{2|m}[(i + 1), p])) \geq [Q_i : Q_{i+1}] \cdot \dim_{\mathbb{F}_p}(\text{im}(c_{2|m}[i, p])).$$

We conclude

$$(2.9) \quad \frac{\dim_{\mathbb{F}_p}(\text{im}(c_{2|m}[(i + 1), p]))}{[Q : Q_{i+1}]} \geq \frac{\dim_{\mathbb{F}_p}(\text{im}(c_{2|m}[i, p]))}{[Q : Q_i]}.$$

Since $\text{im}(c_2^{(2)}) = \bigcup_m \text{im}(c_2|_m^{(2)})$ and $\text{im}(c_2[i, p]) = \bigcup_m \text{im}(c_2|_m[i, p])$ and the dimension functions are compatible with directed unions (see [14, Theorem 6.7 on page 239]), we get

$$(2.10) \quad \dim_{\mathcal{N}(Q)}(\text{im}(c_2^{(2)})) = \lim_{m \rightarrow \infty} \dim_{\mathcal{N}(Q)}(\text{im}(c_2|_m^{(2)}));$$

$$(2.11) \quad \dim_{\mathbb{F}_p}(\text{im}(c_2[i, p])) = \lim_{m \rightarrow \infty} \dim_{\mathbb{F}_p}(\text{im}(c_2|_m[i, p])).$$

We conclude from [14, Theorem 13.3 (2) on page 454 and Lemma 13.4 on page 455]

$$\lim_{i \rightarrow \infty} \frac{\dim_{\mathbb{Q}}(\ker(c_2|_m[i]))}{[Q : Q_i]} = \dim_{\mathcal{N}(Q)}(\ker(c_2|_m^{(2)})).$$

This implies together with (2.6) and (2.7)

$$(2.12) \quad \lim_{i \rightarrow \infty} \frac{\dim_{\mathbb{Q}}(\text{im}(c_2|_m[i]))}{[Q : Q_i]} = \dim_{\mathcal{N}(Q)}(\text{im}(c_2|_m^{(2)})).$$

Finally, it is easy to see that

$$(2.13) \quad \dim_{\mathbb{Q}}(\text{im}(c_2|_m[i])) \geq \dim_{\mathbb{F}_p}(\text{im}(c_2|_m[i, p])).$$

Putting everything together, we can now prove both assertions of Theorem 2.2.

First, for a fixed m , the sequence $\left(\frac{\dim_{\mathbb{F}_p}(\text{im}(c_2|_m[i, p]))}{[Q : Q_i]} \right)_i$ is monotone increasing by (2.9), whence the sequence $\left(\frac{\dim_{\mathbb{F}_p}(\text{im}(c_2[i, p]))}{[Q : Q_i]} \right)_i$ is also monotone increasing by (2.11) and therefore the sequence $\left(\frac{b_1(G_i; \mathbb{F}_p)}{[Q : Q_i]} \right)_i$ is monotone decreasing by (2.5). This proves the first assertion of Theorem 2.2 since clearly $[Q : Q_i] = [G : G_i]$.

Inequality (2.9) also implies that $\lim_{i \rightarrow \infty} \frac{\dim_{\mathbb{F}_p}(\text{im}(c_2|_m[i, p]))}{[Q : Q_i]} \geq \frac{\dim_{\mathbb{F}_p}(\text{im}(c_2|_m[j, p]))}{[Q : Q_j]}$ for any fixed j and m , and so

$$(2.14) \quad \lim_{m \rightarrow \infty} \lim_{i \rightarrow \infty} \frac{\dim_{\mathbb{F}_p}(\text{im}(c_2|_m[i, p]))}{[Q : Q_i]} \geq \sup_{i \geq 0} \left\{ \lim_{m \rightarrow \infty} \frac{\dim_{\mathbb{F}_p}(\text{im}(c_2|_m[i, p]))}{[Q : Q_i]} \right\}.$$

Therefore,

$$\begin{aligned}
 b_1^{(2)}(X; \mathcal{N}(Q)) &\stackrel{(2.4)}{=} s - 1 - \dim_{\mathcal{N}(Q)}(\text{im}(c_2^{(2)})) \\
 &\stackrel{(2.10)}{=} s - 1 - \lim_{m \rightarrow \infty} \dim_{\mathcal{N}(Q)}(\text{im}(c_2|_m^{(2)})) \\
 &\stackrel{(2.12)}{=} s - 1 - \lim_{m \rightarrow \infty} \lim_{i \rightarrow \infty} \frac{\dim_{\mathbb{Q}}(\text{im}(c_2|_m[i]))}{[Q : Q_i]} \\
 &\stackrel{(2.13)}{\leq} s - 1 - \lim_{m \rightarrow \infty} \lim_{i \rightarrow \infty} \frac{\dim_{\mathbb{F}_p}(\text{im}(c_2|_m[i, p]))}{[Q : Q_i]} \\
 &\stackrel{(2.14)}{\leq} s - 1 - \sup_{i \geq 0} \left\{ \lim_{m \rightarrow \infty} \frac{\dim_{\mathbb{F}_p}(\text{im}(c_2|_m[i, p]))}{[Q : Q_i]} \right\} \\
 &\stackrel{(2.11)}{=} s - 1 - \sup_{i \geq 0} \left\{ \frac{\dim_{\mathbb{F}_p}(\text{im}(c_2[i, p]))}{[Q : Q_i]} \right\} \\
 &= \inf_{i \geq 0} \left\{ s - 1 - \frac{\dim_{\mathbb{F}_p}(\text{im}(c_2[i, p]))}{[Q : Q_i]} \right\} \\
 &\stackrel{(2.5)}{=} \inf_{i \geq 0} \left\{ \frac{b_1(G_i; \mathbb{F}_p)}{[Q : Q_i]} \right\}.
 \end{aligned}$$

This finishes the proof of Theorem 2.2. □

3. ALTERNATIVE PROOF OF THEOREM 1.6

In this section we give an alternative proof of Theorem 1.6. Namely, Theorem 1.6 is an easy consequence of the following result, which may be useful in its own right.

THEOREM 3.1. *Let G be a finitely presented group, let (G_i) be a descending chain of finite index normal subgroups of G , and let $K = \bigcap_{i=1}^{\infty} G_i$.*

(1) *The following inequalities hold:*

$$\lim_{i \rightarrow \infty} \frac{b_1(G_i/K)}{[G : G_i]} \leq b_1^{(2)}(G/K) \leq b_1^{(2)}(K \backslash EG; \mathcal{N}(G/K)) = \lim_{n \rightarrow \infty} \frac{b_1(G_i)}{[G : G_i]}.$$

(2) *Let \mathcal{C} be any class of finite groups which is closed under subgroups, extensions (and isomorphisms) and contains at least one non-trivial group (for instance, \mathcal{C} could be the class of all finite groups or all finite p -groups for a fixed prime p). Assume that K is the kernel of the canonical map from G to its pro- \mathcal{C} completion. Then*

$$b_1^{(2)}(G/K) = \lim_{i \rightarrow \infty} \frac{b_1(G_i)}{[G : G_i]}.$$

If in addition all groups G/G_i are in \mathcal{C} , then

$$(3.2) \quad \lim_{i \rightarrow \infty} \frac{b_1(G_i/K)}{[G : G_i]} = b_1^{(2)}(G/K) = b_1^{(2)}(K \backslash EG; \mathcal{N}(G/K)) = \lim_{i \rightarrow \infty} \frac{b_1(G_i)}{[G : G_i]}.$$

Proof. (1) Since G is finitely presented, there is a G -CW-model for the classifying space BG whose 2-skeleton is finite. Let $EG \rightarrow BG$ be the universal covering. Then EG is a free G -CW-complex with finite 2-skeleton. Put

$$\begin{aligned} Q &= G/K; \\ Q_i &= G_i/K. \end{aligned}$$

Then $Q = Q_0 \supseteq Q_1 \supseteq \dots$ is a descending chain of finite index normal subgroups of Q with $\bigcap_{i=0}^{\infty} Q_i = \{1\}$ and we have for $i = 0, 1, 2, \dots$

$$(3.3) \quad [G : G_i] = [Q : Q_i].$$

The quotient $X = K \backslash EG$ is a free Q -CW-complex whose 2-skeleton is finite. Let X_2 be the 2-skeleton of X . Since the first L^2 -Betti number and the first Betti number depend only on the 2-skeleton, from [13, Theorem 0.1] applied to the G -covering $X_2 \rightarrow X_2/G$ (we do not need X_2 to be simply connected) or directly from [14, Theorem 13.3 on page 454], we obtain

$$(3.4) \quad b_1^{(2)}(X; \mathcal{N}(Q)) = \lim_{i \rightarrow \infty} \frac{b_1(Q_i \backslash X)}{[Q : Q_i]}.$$

Let $f: X \rightarrow EQ$ be the classifying map. Since EQ is simply connected, this map is 1-connected. This implies by [14, Theorem 6.54 (1a) on page 265]

$$(3.5) \quad b_1^{(2)}(X; \mathcal{N}(Q)) \geq b_1^{(2)}(EQ; \mathcal{N}(Q)).$$

The group Q is finitely generated (but not necessarily finitely presented), so by [16, Theorem 1.1] we have

$$(3.6) \quad \lim_{i \rightarrow \infty} \frac{b_1(Q_i)}{[Q : Q_i]} \leq b_1^{(2)}(Q).$$

Notice that $b_1^{(2)}(Q) = b_1^{(2)}(EQ; \mathcal{N}(Q))$ by definition and we obviously have $Q_i \backslash X = G_i \backslash EG = BG_i$ and hence $b_1(Q_i \backslash X) = b_1(G_i)$. Combining (3.3), (3.4), (3.5), and (3.6), we get

$$\lim_{i \rightarrow \infty} \frac{b_1(Q_i)}{[Q : Q_i]} \leq b_1^{(2)}(Q) \leq b_1^{(2)}(X; \mathcal{N}(Q)) = \lim_{i \rightarrow \infty} \frac{b_1(Q_i \backslash X)}{[Q : Q_i]} = \lim_{i \rightarrow \infty} \frac{b_1(G_i)}{[G : G_i]}.$$

This finishes the proof of assertion (1).

(2) First observe that since $b_1^{(2)}(K \backslash EG; \mathcal{N}(G/K)) = \lim_{i \rightarrow \infty} \frac{b_1(G_i)}{[G : G_i]}$ by (1), the limit $\lim_{i \rightarrow \infty} \frac{b_1(G_i)}{[G : G_i]}$ is the same for all finite index normal chains (G_i) with $\bigcap_{i \in \mathbb{N}} G_i = K$. By definition of K , there exists at least one such chain with $G/G_i \in \mathcal{C}$ for all i (e.g., we can let (G_i) be a base of neighborhoods of 1 for the pro- \mathcal{C} topology on G), so it suffices to prove (3.2). Thus, from now on we will assume that $G/G_i \in \mathcal{C}$ for $i \in \mathbb{N}$.

For a finitely generated group H we denote by H' the kernel of the composite of canonical projections $H \rightarrow H_1(H) \rightarrow H_1(H)/\text{tors}(H_1(H))$, so that H/H' is a free abelian group of rank $b_1(H)$.

As in the proof of (1), we put $Q_i = G_i/K$ for $i \in \mathbb{N}$. It is sufficient to prove that $K \subseteq G'_i$ for $i \in \mathbb{N}$. Indeed, this would imply that $Q_i/Q'_i \cong G_i/G'_i$, whence $b_1(Q_i) = b_1(G_i)$ and therefore $\lim_{i \rightarrow \infty} \frac{b_1(Q_i)}{[G:G_i]} = \lim_{i \rightarrow \infty} \frac{b_1(G_i)}{[G:G_i]}$, which proves (2) in view of (1).

Fix $i \in \mathbb{N}$ and let $H = G_i$. Since \mathcal{C} contains at least one non-trivial finite group and is closed under subgroups, it contains a finite cyclic group, say of order k . Since \mathcal{C} is closed under extensions, it contains $(\mathbb{Z}/k^m\mathbb{Z})^b$ for all $m, b \in \mathbb{N}$. Setting $b = b_1(H)$, we get that $H/H'H^{k^m} \in \mathcal{C}$ for all $m \in \mathbb{N}$, and since \mathcal{C} is closed under extensions, we obtain $G/H'H^{k^m} \in \mathcal{C}$. By definition, K is the intersection of all normal subgroups L of G with $G/L \in \mathcal{C}$. Therefore, $K \subseteq \bigcap_{m \in \mathbb{N}} H'H^{k^m} = H'$. \square

Second proof of Theorem 1.6.

(1) This is a direct consequence of the following well-known fact: if H is a normal subgroup of p -power index in G , then $b_1(H; \mathbb{F}_p) - 1 \leq [G : H](b_1(G; \mathbb{F}_p) - 1)$ (see, e.g., [11, Proposition 3.7]).

(2) Choose an epimorphism $\pi: F \rightarrow G$, where F is a finitely generated free group. Fix $n \in \mathbb{N}$, let $F_n = \pi^{-1}(G_n)$ and $H = [F_n, F_n]F_n^p$. Then H is a finite index subgroup of F , so we can choose a presentation (X, R) of G associated with π such that $R = R_1 \sqcup R_2$, where R_1 is finite and $R_2 \subseteq H$.

Consider the finitely presented group $\tilde{G} = \langle X \mid R_1 \rangle$. We have natural epimorphisms $\phi: \tilde{G} \rightarrow G$ and $\psi: F \rightarrow \tilde{G}$, with $\phi\psi = \pi$. If we let $\tilde{G}_i = \phi^{-1}(G_i)$ and $\tilde{K} = \bigcap_{i=1}^\infty \tilde{G}_i$, then $\tilde{G}/\tilde{K} \cong G$. Thus, applying Theorem 3.1 (1) to the group \tilde{G} and its subgroups (\tilde{G}_i) , we get $b_1^{(2)}(G) \leq \lim_{i \rightarrow \infty} \frac{b_1(\tilde{G}_i)}{[\tilde{G}:\tilde{G}_i]}$. Clearly,

$$\lim_{i \rightarrow \infty} \frac{b_1(\tilde{G}_i)}{[\tilde{G}:\tilde{G}_i]} \leq \lim_{i \rightarrow \infty} \frac{b_1(\tilde{G}_i; \mathbb{F}_p)}{[\tilde{G}:\tilde{G}_i]},$$

$$\lim_{i \rightarrow \infty} \frac{b_1(\tilde{G}_i; \mathbb{F}_p)}{[\tilde{G}:\tilde{G}_i]} \leq \frac{b_1(\tilde{G}_n; \mathbb{F}_p)}{[\tilde{G}:\tilde{G}_n]} = \frac{b_1(\tilde{G}_n; \mathbb{F}_p)}{[G:G_n]}.$$

Since $G \cong \tilde{G}/\langle\langle \psi(R_2) \rangle\rangle$ and by construction $\psi(R_2) \subseteq \psi(H) = [\tilde{G}_n, \tilde{G}_n]\tilde{G}_n^p$, we have $\ker \phi \subseteq [\tilde{G}_n, \tilde{G}_n]\tilde{G}_n^p$, and therefore $b_1(\tilde{G}_n; \mathbb{F}_p) = b_1(\phi(\tilde{G}_n); \mathbb{F}_p) = b_1(G_n; \mathbb{F}_p)$.

Combining these inequalities, we get $b_1^{(2)}(G) \leq \frac{b_1(G_n; \mathbb{F}_p)}{[G:G_n]}$. Since n is arbitrary, the proof is complete. \square

4. A COUNTEREXAMPLE WITH NON-TRIVIAL INTERSECTION

In this section we show that the answer to Questions 1.5 and 1.11 could be negative for a finitely presented group G and a strictly descending chain $(G_i)_{i \in \mathbb{N}}$

of normal subgroups of p -power index if the intersection $\bigcap_{i \in \mathbb{N}} G_i$ is non-trivial (see inequalities (4.2) below).

We start with a finitely generated group H (which will be specified later) and let $G = H * \mathbb{Z}$. Choose a strictly increasing sequence of positive integers n_1, n_2, \dots with $n_i \mid n_{i+1}$ for each i , and let $G_i \subseteq G$ be the preimage of $n_i \cdot \mathbb{Z}$ under the natural projection $\text{pr}: G = \mathbb{Z} * H \rightarrow \mathbb{Z}$. Then $(G_i)_{i \in \mathbb{N}}$ is a descending chain of normal subgroups of G with $\bigcap_{i \geq 1} G_i = \ker(\text{pr})$. Let $BG_i \rightarrow BG$ be the covering of BG associated to $G_i \subseteq G$. Then BG_i is homeomorphic to $S^1 \vee \left(\bigvee_{j=1}^{n_i} BH \right)$. We have

$$G_i \cong \pi_1(BG_i) \cong \pi_1 \left(S^1 \vee \left(\bigvee_{j=1}^{n_i} BH \right) \right) \cong \mathbb{Z} * (*_{j=1}^{n_i} H).$$

Since for any groups A and B we have $A * B / [A * B, A * B] \cong A / [A, A] \oplus B / [B, B]$ and $d(A * B) = d(A) + d(B)$ by Grushko-Neumann theorem (see [4, Corollary 2 in Section 8.5 on page 227], we conclude

$$\begin{aligned} H_1(G_i; K) &= K \oplus \bigoplus_{j=1}^{n_i} H_1(H; K); \\ H_1(G_i) &= \mathbb{Z} \oplus \bigoplus_{j=1}^{n_i} H_1(H); \\ d(G_i) &= 1 + n_i \cdot d(H); \\ \lim_{i \rightarrow \infty} \frac{b_1(G_i; K)}{n_i} &= b_1(H; K); \\ \lim_{i \rightarrow \infty} \frac{d(H_1(G_i))}{n_i} &= d(H_1(H)); \\ \text{RG}(G; (G_i)_{i \geq 1}) &= d(H). \end{aligned}$$

Now let $p \neq q$ be distinct primes and $H = \mathbb{Z}/p\mathbb{Z} * \mathbb{Z}/q\mathbb{Z} * \mathbb{Z}/q\mathbb{Z}$. Clearly we have

$$(4.1) \quad b_1(H) = 0, \quad b_1(H; \mathbb{F}_p) = 1, \quad d(H_1(H)) = 2, \quad d(H) = 3.$$

Hence we obtain

$$(4.2) \quad \lim_{i \rightarrow \infty} \frac{b_1(G_i)}{[G : G_i]} < \lim_{i \rightarrow \infty} \frac{b_1(G_i; \mathbb{F}_p)}{[G : G_i]} < \lim_{i \rightarrow \infty} \frac{d(H_1(G_i))}{[G : G_i]} < \text{RG}(G; (G_i)_{i \geq 1}).$$

Using a different H we can produce an example of this type where G has a very strong finiteness property, namely, G has finite 2-dimensional BG . The construction below is due to Denis Osin and is simpler and more explicit than the original version of our example.

Again, let $p \neq q$ be two primes. Consider the group

$$H = \langle x, y, z \mid x^p = u, y^q = v, z^q = w \rangle,$$

where u, v, w are words from the commutator subgroup of the free group F with basis x, y, z such that the presentation of H satisfies the $C'(1/6)$ small

cancellation condition. Such words are easy to find explicitly. Note that $G = H * \mathbb{Z}$ is a torsion-free $C'(1/6)$ group, hence it has a finite 2-dimensional BG . Since $u, v, w \in [F, F]$, we have $b_1(H) = 0$, $b_1(H; \mathbb{F}_p) = 1$, $d(H_1(H)) = 2$. Further it follows from [6, Corollary 2] that the exponential growth rate of H can be made arbitrarily close to $2 \cdot 3 - 1 = 5$, the exponential growth rate of the free group of rank 3, by taking sufficiently long words u, v, w . As the exponential growth rate of an m -generated group is bounded from above by $2m - 1$, we obtain $d(H) = 3$ whenever u, v, w are sufficiently long. (For details about the exponential growth rate we refer to [6].)

By using a more elaborated construction from [21], one can make such a group G the fundamental group of a compact 2-dimensional $CAT(-1)$ CW -complex. Other examples of this type can be found in [3] and [15].

5. \mathbb{Q} -APPROXIMATION WITHOUT LIMIT

In this section we prove the following theorem, which trivially implies Theorem 1.3.

THEOREM 5.1. *Let $d \geq 2$ be a positive integer, let p be a prime and let ε be a real number satisfying $0 < \varepsilon < 1$. Then there exist a group G with d generators and a descending chain $G = G_0 \supseteq G_1 \supseteq G_2 \dots$ of normal subgroups of G of p -power index with $\bigcap_{i=1}^\infty G_i = \{1\}$ with the following properties:*

- (i) $\liminf_{i \rightarrow \infty} \frac{b_1(G_{2i})}{[G:G_{2i}]} \geq d - 1 - \varepsilon$;
- (ii) $\lim_{i \rightarrow \infty} \frac{b_1(G_{2i-1})}{[G:G_{2i-1}]} = 0$.

Moreover, if q is a prime different from p , we can replace (ii) by a stronger condition (ii)':

(ii)' $\lim_{i \rightarrow \infty} \frac{b_1(G_{2i-1}; \mathbb{F}_q)}{[G:G_{2i-1}]} = 0$.

Note that the last assertion of Theorem 5.1 shows that the answer to Question 1.14 can be negative when $\text{char}(K) = q > 0$ if we do not require that (G_i) is a q -chain.

5.1. PRELIMINARIES. Throughout this section p will be a fixed prime number. Given a finitely generated group G , we will denote by $G_{\hat{p}}$ the pro- p completion of G and by $G_{(p)}$ the image of G in $G_{\hat{p}}$ (which is isomorphic to the quotient of G by the intersection of normal subgroups of p -power index). Given a set X , by $F(X)$ we denote the free group on X .

Let F be a free group and $w \in F$ a non-identity element. Given $n \in \mathbb{N}$, denote by $\sqrt[n]{w}$ the unique element of F whose n^{th} power is equal to w (if such element exists). Define $e_p(w, F)$ to be the largest natural number e with the property that $\sqrt[e]{w}$ exists in F .

LEMMA 5.2. *Let (X, R) be a presentation of a group G with X finite, $F = F(X)$ and $\pi: F \rightarrow G$ the natural projection. Let H be a normal subgroup of p -power index in G , and let $F_H = \pi^{-1}(H)$. Then $H = F_H / \langle\langle R_H \rangle\rangle$ where R_H contains $\frac{[G:H]}{p^{e_p(r, F)} - e_p(r, F_H)}$ F -conjugates of r for each $r \in R$ and no other elements.*

Proof. Very similar results are proved in both [18] and [20], but for completeness we give a proof. For each $r \in R$, write $r = w(r)^{p^{e_p(r,F)}}$, and choose a right transversal $T = T(r)$ for $\langle w(r) \rangle F_H$ in F . Then, since $w(r)$ commutes with r , by [17, Lemma 2.3] we have $\langle r \rangle^F = \langle \{t^{-1}rt : t \in T\} \rangle^{F_H}$. Hence $\langle \{t^{-1}rt : r \in R, t \in T(R)\} \rangle^{F_H} = \langle R \rangle^F = \ker \pi = \ker(F_H \rightarrow H)$, and so it suffices to prove that $|T(r)| = \frac{[G:H]}{p^{e_p(r,F) - e_p(r,F_H)}}$.

We have

$$|T(r)| = [F : \langle w(r) \rangle F_H] = \frac{[F : F_H]}{[\langle w(r) \rangle F_H : F_H]} = \frac{[G : H]}{[\langle w(r) \rangle : \langle w(r) \rangle \cap F_H]}$$

Finally note that $[\langle w(r) \rangle : \langle w(r) \rangle \cap F_H]$ is equal to p^k for some k (as it divides $[F : F_H] = p^n$), so $\langle w(r) \rangle \cap F_H = \langle w(r) \rangle^{p^k}$. But then from definition of $e_p(r, F_H)$ we easily conclude that $(\langle w(r) \rangle^{p^k})^{p^{e_p(r, F_H)}} = r = w(r)^{p^{e_p(r, F)}}$. Hence $k = e_p(r, F) - e_p(r, F_H)$ and $|T(r)| = \frac{[G:H]}{p^{e_p(r,F) - e_p(r,F_H)}}$, as desired. \square

The following definition was introduced by Schlage-Puchta in [20].

DEFINITION 5.3. Given a group presentation by generators and relators (X, R) , where X is finite, its p -deficiency $\text{def}_p(X, R) \in \mathbb{R} \cup \{-\infty\}$ is defined by

$$\text{def}_p(X, R) = |X| - 1 - \sum_{r \in R} \frac{1}{p^{e_p(r, F(X))}}.$$

The p -deficiency of a finitely generated group G is the supremum of the set $\{\text{def}_p(X, R)\}$ where (X, R) ranges over all presentations of G .

The main motivation for introducing p -deficiency in [20] was to construct a finitely generated p -torsion group with positive rank gradient. Indeed, it is clear that there exist p -torsion groups with positive p -deficiency, and in [20] it is proved that a group with positive p -deficiency has positive rank gradient (in fact, positive p -gradient). This is one of the results indicating that groups of positive p -deficiency behave similarly to groups of deficiency greater than 1 (all of which trivially have positive p -deficiency for any p).

Lemma 5.5 below shows that a finitely presented group G of positive p -deficiency actually contains a normal subgroup of p -power index with deficiency greater than 1, provided that the presentation of G yielding positive p -deficiency is finite and satisfies certain technical condition.

DEFINITION 5.4. A presentation (X, R) of a group G will be called p -regular if for any $r \in R$ such that $\sqrt[p]{r}$ exists in $F(X)$, the image of $\sqrt[p]{r}$ in $G_{(p)}$ is non-trivial. This is equivalent to saying that if we write each $r \in R$ as $r = v^{p^e}$, where v is not a p^{th} power in $F(X)$, then the image of v in $G_{(p)}$ has order p^e .

LEMMA 5.5. Let (X, R) be a finite p -regular presentation of a group G . Then there exists a normal subgroup of p -power index H of G with $\frac{\text{def}(H)-1}{[G:H]} \geq \text{def}_p(X, R)$.

Proof. Let $F = F(X)$. Let r_1, \dots, r_m be the elements of R and let $s_i = \sqrt[p]{r_i}$, whenever it is defined in $F(X)$.

Let $\pi: F \rightarrow G_{(p)}$ be the natural projection. Since the presentation (X, R) is p -regular, $\pi(s_i)$ is non-trivial whenever s_i is defined, and since the group $G_{(p)}$ is residually- p , there exists a normal subgroup H' of $G_{(p)}$ of p -power index which contains none of the elements $\pi(s_i)$.

Let $F_H = \pi^{-1}(H')$. By construction, $s_i \notin F_H$, but $r_i \in F_H$, and therefore $e_p(r_i, F_H) = 0$ for each i . Let H be the image of F_H in G . Then by Lemma 5.2, H has a presentation with $d(F_H)$ generators and $\sum_{i=1}^m \frac{[G:H]}{p^{e_p(r_i, F)}}$ relators. Since $d(F_H) - 1 = (|X| - 1)[F : F_H] = (|X| - 1)[G : H]$ by the Schreier formula, we get

$$\text{def}(H) - 1 \geq [G : H] \cdot \left(|X| - 1 - \sum_{i=1}^m p^{-e_p(r_i, F)} \right) = [G : H] \cdot \text{def}_p(X, R).$$

□

LEMMA 5.6. *Let (X, R) be a finite p -regular presentation, and let $G = \langle X | R \rangle$. Let $f \in F(X)$ be such that the image of f in the pro- p completion of G has infinite order. Then there exists $N \in \mathbb{N}$ such that for all $n \geq N$ the presentation $(X, R \cup \{f^{p^n}\})$ is p -regular.*

Proof. Let r_1, \dots, r_m be the elements of R . By assumption there is a normal subgroup of p -power index H of G such that $\sqrt[p]{r_i}$ does not vanish in G/H (whenever $\sqrt[p]{r_i}$ exists in $F(X)$). Let $\pi: F(X) \rightarrow G$ be the natural projection, and choose $N \in \mathbb{N}$ satisfying $\pi(f^{p^N}) \in H$.

Let $n \geq N$, let $g = \pi(f)$, and let $G' = G / \langle\langle g^{p^n} \rangle\rangle = \langle X | R \cup \{f^{p^n}\} \rangle$. We claim that the presentation $(X, R \cup \{f^{p^n}\})$ is p -regular. We need to check that

- (i) each $\sqrt[p]{r_i}$ does not vanish in G'_p
- (ii) $f^{p^{n-1}}$ does not vanish in G'_p

The kernel of the natural map $G \rightarrow G'_p$ is contained in H since $g^{p^n} \in H$ and G/H is a finite p -group. Since $\pi(\sqrt[p]{r_i}) \notin H$, this implies (i). Further, an element $x \neq 1$ of a pro- p group cannot lie in the closed normal subgroup generated by x^p . Hence if \hat{g} is the image of g (also the image of f) in G'_p , then $\hat{g}^{p^{n-1}}$ does not lie in the closed normal subgroup of G'_p generated by \hat{g}^{p^n} , call this subgroup C . Finally, by definition of G' , there is a canonical isomorphism from G'_p/C to G'_p , which maps the image of f in G'_p/C to the image of f in G'_p . Thus, we verified (ii). □

COROLLARY 5.7. *Let (X, R) be a finite p -regular presentation, and let $G = \langle X | R \rangle$. Let $H \subseteq K$ be normal subgroups of $F(X)$ of p -power index, and let $\delta > 0$ be a real number. Then there exists a finite set $R' \subset [K, K]$ with $\sum_{r \in R'} p^{-e_p(r, F(X))} < \delta$ such that*

- (1) the presentation $(X, R \cup R')$ is p -regular;

(2) if $G' = \langle X \mid R \cup R' \rangle$ and H' is the image of H in G' , then $b_1(H') \leq d(K)$.

Moreover, if q is a prime different from p , we can require that $b_1(H'; \mathbb{F}_q) \leq d(K)$.

Proof. If $b_1(H; \mathbb{F}_q) \leq d(K)$, we can choose $R' = \emptyset$. Hence we can assume without loss of generality that $b_1(H; \mathbb{F}_q) > d(K)$. Clearly, it suffices to prove a weaker statement, where inequality $b_1(H'; \mathbb{F}_q) \leq d(K)$ is replaced by $b_1(H'; \mathbb{F}_q) < b_1(H; \mathbb{F}_q)$. The assertion of Corollary 5.7 then follows by repeated applications with δ replaced by $\delta/(b_1(H; \mathbb{F}_q) - d(K))$.

Let Y be any free generating set for H . Obviously $K/[K, K]$ is a free abelian group of rank $d(K)$. Any (finite) matrix over the integers can be transformed by elementary row and column operations to a diagonal matrix. Hence by applying elementary transformations to Y , we can arrange that Y is a disjoint union $Y_1 \sqcup Y_2$ where $|Y_1| \leq d(K)$ and $Y_2 \subseteq [K, K]$.

Let $L = \langle Y_2 \rangle$, the subgroup generated by Y_2 . Since $b_1(H; \mathbb{F}_q) > d(K)$, there exists $f \in Y_2$ whose image in $H/[H, H]H^q \cong H_1(H, \mathbb{F}_q)$ is non-trivial. Now apply Lemma 5.6 to this f , choose n such that $\frac{1}{p^n} < \delta$ and let $R' = \{f^{p^n}\}$. The choice of f ensures that $b_1(H'; \mathbb{F}_q) < b_1(H; \mathbb{F}_q)$, so R' has the required properties. \square

5.2. PROOF OF THEOREM 5.1. To simplify the notations, we will give a proof of the main part of Theorem 5.1. The last part of Theorem 5.1 is proved in the same way by using the last assertion of Corollary 5.7.

We start by giving an outline of the construction. Let $F = F(X)$ be a free group of rank $d = |X|$. Below we shall define a descending chain $F = F_0 \supseteq F_1 \supseteq \dots$ of normal subgroups of F of p -power index and a sequence of finite subsets R_1, R_2, \dots of F . Let $R = \bigcup_{i=1}^{\infty} R_i$. For each $n \in \mathbb{Z}_{\geq 0}$ we let $G(n) = F/\langle\langle \bigcup_{i=1}^n R_i \rangle\rangle$, $G(\infty) = \varinjlim G(i) = F/\langle\langle R \rangle\rangle$ and let G be the image of $G(\infty)$ in its pro- p completion. Denote by $G(n)_i$, $G(\infty)_i$ and G_i the canonical image of F_i in $G(n)$, $G(\infty)$ and G , respectively. We will show that the group G and its subgroups (G_i) satisfy the conclusion of Theorem 5.1.

Fix a sequence of positive real numbers (δ_n) which converges to zero and a descending chain (Φ_n) of normal subgroups of p -power index in F which form a base of neighborhoods of 1 for the pro- p topology. The subgroups F_n and relator sets R_n will be constructed inductively so that the following properties hold:

(i) For $n \geq 0$ we have

$$\frac{b_1(G(n)_{2n})}{[G(n) : G(n)_{2n}]} > d - 1 - \varepsilon;$$

(ii) For $n \geq 1$ we have

$$\frac{b_1(G(n)_{2n-1})}{[G(n) : G(n)_{2n-1}]} < \delta_n;$$

(iii) R_n is contained in $[F_{2n-2}, F_{2n-2}]$ for $n \geq 1$;

- (iv) $F_{2n} \subseteq \Phi_n$ for $n \geq 1$;
- (v) $\text{def}_p(X, \cup_{i=1}^n R_i) > d - 1 - \varepsilon$ for $n \geq 1$;
- (vi) The presentation $(X, \cup_{i=1}^n R_i)$ is p -regular for $n \geq 1$.

We first explain why properties (i)-(vi) will imply that the group G and its subgroups (G_n) have the desired properties. Each G_n is normal of p -power index in G since F_n is normal of p -power index in F . Condition (iv) implies that (G_n) is a base of neighborhoods of 1 for the pro- p topology on G , and since G is residually- p by construction, we have $\bigcap_{n=1}^\infty G_n = \{1\}$.

Condition (iii) implies that $[G(n) : G(n)_i] = [G(\infty) : G(\infty)_i]$ and $b_1(G(n)_i) = b_1(G(\infty)_i)$ for $i \leq 2n$. Since $G(\infty)_i$ is normal of p -power index in $G(\infty)$, the group $G(\infty)/[G(\infty)_i, G(\infty)_i]$ is residually- p , so both the index and the first Betti number of $G(\infty)_i$ do not change under passage to the image in the pro- p completion of $G(\infty)$: $[G : G_i] = [G(\infty) : G(\infty)_i]$ and $b_1(G_i) = b_1(G(\infty)_i)$. In view of these equalities, conditions (i) and (ii) yield the corresponding conditions in Theorem 5.1.

We now describe the construction of the sets R_n and subgroups F_n . The base case $n = 0$ is obvious: we set $F_0 = F$ and $G(0) = F$, and the only condition we require for $n = 0$ (condition (i)) clearly holds.

Suppose now that $N \in \mathbb{N}$ and we constructed subsets $(R_i)_{i=1}^N$ and subgroups $(F_i)_{i=1}^{2N}$ such that (i)-(vi) hold for all $n \leq N$.

Let $F_{2N+1} = [F_{2N}, F_{2N}]F_{2N}^{p^e}$ where e is specified below. Then F_{2N+1} is a normal subgroup of p -power index in F and $F_{2N} \supseteq F_{2N+1} \supset [F_{2N}, F_{2N}]$. Since $b_1(G(N)_{2N}) > 0$ by (i) for $n = N$ and hence

$$\begin{aligned} p^e &\leq |H_1(G(N)_{2N})/p^e \cdot H_1(G(N)_{2N})| \\ &= |G(N)_{2N}/[G(N)_{2N}, G(N)_{2N}]G(N)_{2N}^{p^e}| \\ &= |G(N)_{2N}/G(N)_{2N+1}| \\ &= [G(N)_{2N} : G(N)_{2N+1}] \\ &\leq [G(N) : G(N)_{2N+1}], \end{aligned}$$

so we can arrange

$$\frac{d(F_{2N})}{[G(N) : G(N)_{2N+1}]} < \delta_{N+1}$$

by choosing e large enough.

Now applying Corollary 5.7 with $H = F_{2N+1}$, $K = F_{2N}$ and $\delta = \text{def}_p(X, \cup_{i=1}^N R_i) - (d - 1 - \varepsilon)$, we get that there is a finite subset $R_{N+1} \subseteq [F_{2N}, F_{2N}]$ such that the presentation $(X, \cup_{i=1}^{N+1} R_i)$ is p -regular and $\text{def}_p(X, \cup_{i=1}^{N+1} R_i) > d - 1 - \varepsilon$. Hence conditions (iii),(v),(vi) hold for $n = N + 1$. The subgroup H' in the notations of Corollary 5.7 is equal to $G(N + 1)_{2N+1}$, so $b_1(G(N + 1)_{2N+1}) \leq d(F_{2N})$. Since condition (iii) implies $[G(N + 1) : G(N + 1)_{2N+1}] = [G(N) : G(N)_{2N+1}]$, we conclude

$$\frac{b_1(G(N + 1)_{2N+1})}{[G(N + 1) : G(N + 1)_{2N+1}]} \leq \frac{d(F_{2N})}{[G(N) : G(N)_{2N+1}]} < \delta_{N+1}.$$

Thus we have shown that conditions (ii),(iii),(v),(vi) hold for $n = N + 1$.

It remains to construct F_{2N+2} and to verify (i) and (iv) for $n = N + 1$. We apply Lemma 5.5 to $G(N + 1) = \langle X \mid \cup_{i=1}^{N+1} R_i \rangle$ and obtain using (v) a normal subgroup H of $G(N + 1)$ of p -power index satisfying

$$\frac{\text{def}(H) - 1}{[G(N + 1) : H]} > d - 1 - \varepsilon.$$

Let $F_{2N+2} \subseteq F_{2N+1} \cap \Phi_{N+1}$ be the intersection of the preimage of H under the projection $p_{N+1}: F_{N+1} \rightarrow G(N + 1)$ with $F_{2N+1} \cap \Phi_{N+1}$. Obviously (iv) for holds $n = N + 1$. Then $G(N + 1)_{2N+2}$ is a subgroup of H of finite index. The quantity $\text{def}(\cdot) - 1$ is supermultiplicative, i.e., if L is a finite index subgroup of H , then $\text{def}(L) - 1 \geq [H : L] \cdot (\text{def}(H) - 1)$, see for instance [18, Lemma 2.2]. Hence we conclude

$$\frac{\text{def}(G(N + 1)_{2N+2}) - 1}{[G(N + 1) : G(N + 1)_{2N+2}]} \geq \frac{\text{def}(H) - 1}{[G(N + 1) : H]} > d - 1 - \varepsilon.$$

Since $b_1(G(N + 1)_{2N+2}) \geq \text{def}(G(N + 1)_{2N+2})$, condition (i) holds for $n = N + 1$. This finishes the proof of Theorem 5.1.

REFERENCES

- [1] M. Abért, A. Jaikin-Zapirain, and N. Nikolov. The rank gradient from a combinatorial viewpoint. *Groups Geom. Dyn.*, 5(2):213–230, 2011.
- [2] M. Abért and N. Nikolov. Rank gradient, cost of groups and the rank versus Heegaard genus problem. *J. Eur. Math. Soc. (JEMS)*, 14(5):1657–1677, 2012.
- [3] N. Bergeron, P. Linnell, W. Lück, and R. Sauer. On the growth of Betti numbers in p -adic analytic towers. Preprint, arXiv:1204.3298v1 [math.GT], to appear in *Groups, Geometry, and Dynamics*, 2012.
- [4] D. E. Cohen. *Combinatorial group theory: a topological approach*. Cambridge University Press, Cambridge, 1989.
- [5] G. Elek. The rank of finitely generated modules over group algebras. *Proc. Amer. Math. Soc.*, 131(11):3477–3485 (electronic), 2003.
- [6] A. Erschler. Growth rates of small cancellation groups. In *Random walks and geometry*, pages 421–430. Walter de Gruyter GmbH & Co. KG, Berlin, 2004.
- [7] D. Gaboriau. Coût des relations d'équivalence et des groupes. *Invent. Math.*, 139(1):41–98, 2000.
- [8] D. Gaboriau. Invariants l^2 de relations d'équivalence et de groupes. *Publ. Math. Inst. Hautes Études Sci.*, 95:93–150, 2002.
- [9] D. Gaboriau. On orbit equivalence of measure preserving actions. In *Rigidity in dynamics and geometry (Cambridge, 2000)*, pages 167–186. Springer, Berlin, 2002.
- [10] M. Lackenby. Expanders, rank and graphs of groups. *Israel J. Math.*, 146:357–370, 2005.
- [11] M. Lackenby. Covering spaces of 3-orbifolds. *Duke Math. J.*, 136(1):181–203, 2007.

- [12] P. Linnell, W. Lück, and R. Sauer. The limit of \mathbb{F}_p -Betti numbers of a tower of finite covers with amenable fundamental groups. *Proc. Amer. Math. Soc.*, 139(2):421–434, 2011.
- [13] W. Lück. Approximating L^2 -invariants by their finite-dimensional analogues. *Geom. Funct. Anal.*, 4(4):455–481, 1994.
- [14] W. Lück. *L²-Invariants: Theory and Applications to Geometry and K-Theory*, volume 44 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 2002.
- [15] W. Lück. Approximating L^2 -invariants and homology growth. *Geom. Funct. Anal.*, 23(2):622–663, 2013.
- [16] W. Lück and D. Osin. Approximating the first L^2 -Betti number of residually finite groups. *J. Topol. Anal.*, 3(2):153–160, 2011.
- [17] A. Y. Olshanskii and D. V. Osin. Large groups and their periodic quotients. *Proc. Amer. Math. Soc.*, 136(3):753–759, 2008.
- [18] D. Osin. Rank gradient and torsion groups. *Bull. Lond. Math. Soc.*, 43(1):10–16, 2011.
- [19] D. Osin and A. Thom. Normal generation and ℓ^2 -Betti numbers of groups. *Math. Ann.*, 355(4):1331–1347, 2013.
- [20] J.-C. Schlage-Puchta. A p -group with positive rank gradient. *J. Group Theory*, 15(2):261–270, 2012.
- [21] D. T. Wise. Incoherent negatively curved groups. *Proc. Amer. Math. Soc.*, 126(4):957–964, 1998.

Mikhail Ershov
Department of Mathematics
University of Virginia
PO Box 400137
Charlottesville
VA 22904-4137
U.S.A.
ershov@virginia.edu

Wolfgang Lück
Mathematisches Institut der
Universität
Bonn
Endenicher Allee 60
53115 Bonn
Germany
wolfgang.lueck@him.uni-
bonn.de

