

PARTIAL CLASSIFICATION OF THE BAUMSLAG-SOLITAR  
GROUP VON NEUMANN ALGEBRAS

NIELS MEESSCHAERT<sup>1</sup> AND STEFAAN VAES<sup>2</sup>

Received: March 28, 2013

Revised: March 12, 2014

Communicated by Joachim Cuntz

ABSTRACT. We prove that the rational number  $|n/m|$  is an invariant of the group von Neumann algebra of the Baumslag-Solitar group  $BS(n, m)$ . More precisely, if  $L(BS(n, m))$  is isomorphic with  $L(BS(n', m'))$ , then  $|n'/m'| = |n/m|^{\pm 1}$ . We obtain this result by associating to abelian, but not maximal abelian, subalgebras of a  $\text{II}_1$  factor, an equivalence relation that can be of type III. In particular, we associate to  $L(BS(n, m))$  a canonical equivalence relation of type  $\text{III}_{|n/m|}$ .

2010 Mathematics Subject Classification: Primary 46L36; Secondary 20E06, 22D25.

1. INTRODUCTION AND STATEMENT OF THE MAIN RESULT

Some of the deepest open problems in functional analysis center around the classification of group von Neumann algebras  $L(G)$  associated with certain natural families of countable groups  $G$ . In the case of the free groups, this becomes the famous free group factor problem asking whether  $L(\mathbb{F}_n) \cong L(\mathbb{F}_m)$  when  $n, m \geq 2$  and  $n \neq m$ . For property (T) groups with infinite conjugacy classes (icc), this leads to Connes's rigidity conjecture ([Co80]) asserting that an isomorphism  $L(G) \cong L(\Lambda)$  between the property (T) factors entails an isomorphism  $G \cong \Lambda$  of the groups.

As a consequence of Connes's uniqueness theorem of injective  $\text{II}_1$  factors ([Co75]), the group von Neumann algebra  $L(G)$  of an amenable icc group  $G$  is

---

<sup>1</sup>Supported by ERC Starting Grant VNALG-200749 and Research Programme G.0639.11 of the Research Foundation – Flanders (FWO).

<sup>2</sup>Supported by ERC Starting Grant VNALG-200749, Research Programme G.0639.11 of the Research Foundation – Flanders (FWO) and KU Leuven BOF research grant OT/13/079.

isomorphic with the unique hyperfinite  $\text{II}_1$  factor  $R$ . In the nonamenable case, many nonisomorphic groups  $G$  are known to have nonisomorphic group von Neumann algebras  $L(G)$ . Nevertheless, concerning the classification of group von Neumann algebras of natural families of groups, e.g. lattices in simple Lie groups, little is known. A notable exception however is [CH88] where it is shown that for  $n \neq m$ , lattices in  $\text{Sp}(n, 1)$ , respectively  $\text{Sp}(m, 1)$ , have nonisomorphic group von Neumann algebras.

Since 2001, Popa has been developing a new arsenal of techniques to study  $\text{II}_1$  factors, called deformation/rigidity theory. This theory has provided several classes  $\mathcal{G}$  of groups such that an isomorphism  $L(G) \cong L(\Lambda)$  with both  $G, \Lambda \in \mathcal{G}$  entails the isomorphism  $G \cong \Lambda$ . By [Po04], this holds in particular when  $\mathcal{G}$  is the class of wreath product groups of the form  $(\mathbb{Z}/2\mathbb{Z}) \wr \Gamma$  with  $\Gamma$  an icc property (T) group.

In [IPV10], the first  $W^*$ -superrigidity theorems for group von Neumann algebras were discovered, yielding icc groups  $G$  such that an isomorphism  $L(G) \cong L(\Lambda)$  with  $\Lambda$  an *arbitrary* countable group, implies that  $G \cong \Lambda$ . The groups  $G$  discovered in [IPV10] are generalized wreath products of a special form. In [BV12], it was then shown that one can actually take  $G = (\mathbb{Z}/2\mathbb{Z})^{(\Gamma)} \rtimes (\Gamma \times \Gamma)$  with  $\Gamma$  ranging over a large family of nonamenable groups including the free groups  $\mathbb{F}_n$ ,  $n \geq 2$ .

In this article, we apply Popa's deformation/rigidity theory to partially classify the group von Neumann algebras of the Baumslag-Solitar groups  $\text{BS}(n, m)$ . Recall that for all  $n, m \in \mathbb{Z} - \{0\}$ , this group is defined as the group generated by  $a$  and  $b$  subject to the relation  $ba^nb^{-1} = a^m$ . So,

$$\text{BS}(n, m) := \langle a, b \mid ba^nb^{-1} = a^m \rangle .$$

The Baumslag-Solitar groups were introduced in [BS62] as the first examples of finitely presented non-Hopfian groups. Ever since, they have been used as examples and counterexamples for numerous group theoretic phenomena. Therefore, it is a natural problem to classify the group von Neumann algebras  $L(\text{BS}(n, m))$ .

Whenever  $|n| = 1$  or  $|m| = 1$ , the group  $\text{BS}(n, m)$  is solvable, hence amenable. So we always assume that  $|n| \geq 2$  and  $|m| \geq 2$ . In that case,  $\text{BS}(n, m)$  contains a copy of the free group  $\mathbb{F}_2$  and hence, is nonamenable. In [Mo91], the Baumslag-Solitar groups were classified up to isomorphism:  $\text{BS}(n, m) \cong \text{BS}(n', m')$  if and only if  $\{n, m\} = \{\varepsilon n', \varepsilon m'\}$  for some  $\varepsilon \in \{-1, 1\}$ . So, up to isomorphism, we only consider  $2 \leq n \leq |m|$ . Finally by [St05, Exemple 2.4], the group  $\text{BS}(n, m)$  is icc if and only if  $|n| \neq |m|$ . Therefore, we always assume that  $2 \leq n < |m|$ .

Using Popa's deformation/rigidity theory and in particular his spectral gap rigidity ([Po06]) and the work on amalgamated free products ([IPP05]), several structural properties of the  $\text{II}_1$  factors  $M = L(\text{BS}(n, m))$  were proven. In particular, it was shown in [Fi10] that  $M$  is not solid, that  $M$  is prime and that  $M$  has no Cartan subalgebra. More generally, it is proven in [Fi10] that any amenable regular von Neumann subalgebra of  $M$  must have a nonamenable relative commutant.

Our main result is the following partial classification theorem for the Baumslag-Solitar group von Neumann algebras  $L(\text{BS}(n, m))$ . Whenever  $M$  is a  $\text{II}_1$  factor and  $t > 0$ , we denote by  $M^t$  the *amplification* of  $M$ . Up to unitary conjugacy,  $M^t$  is defined as  $p(M_n(\mathbb{C}) \otimes M)p$  where  $p$  is a projection satisfying  $(\text{Tr} \otimes \tau)(p) = t$ . The  $\text{II}_1$  factors  $M$  and  $N$  are called *stably isomorphic* if there exists a  $t > 0$  such that  $M \cong N^t$ .

**THEOREM A.** *Let  $n, m, n', m' \in \mathbb{Z}$  such that  $2 \leq n < |m|$  and  $2 \leq n' < |m'|$ . If  $L(\text{BS}(n, m))$  is stably isomorphic with  $L(\text{BS}(n', m'))$ , then  $\frac{n}{|m|} = \frac{n'}{|m'|}$ .*

Note that Theorem A formally resembles, but is independent of, the results in [Ki11] on orbit equivalence relations of essentially free ergodic probability measure preserving actions of Baumslag-Solitar groups, especially [Ki11, Proposition B.2 and Theorem 1.2]. It would be very interesting to find a framework that unifies both types of results.

We prove our Theorem A by associating a canonical equivalence relation to  $L(\text{BS}(n, m))$  and proving that it is of type  $\text{III}_{n/|m|}$ . More precisely, assume that  $(M, \tau)$  is a von Neumann algebra with separable predual, equipped with a faithful normal tracial state. Whenever  $A \subset M$  is an abelian von Neumann subalgebra, the normalizer

$$\mathcal{N}_M(A) := \{u \in \mathcal{U}(M) \mid uAu^* = A\}$$

induces a group of trace preserving automorphisms of  $A$ . Writing  $A = L^\infty(X, \mu)$  with  $\mu$  being induced by  $\tau|_A$ , the corresponding orbit equivalence relation is a countable probability measure preserving (pmp) equivalence relation on  $(X, \mu)$ .

More generally, we can consider the set of partial isometries

$$\{u \in M \mid u^*u \text{ and } uu^* \text{ are projections in } A' \cap M \text{ and } uAu^* = Auu^*\}. \quad (1)$$

Every such partial isometry induces a partial automorphism of  $A$  and hence a partial automorphism of  $(X, \mu)$ . We denote by  $\mathcal{R}(A \subset M)$  the equivalence relation generated by all these partial automorphisms. When  $A \subset M$  is maximal abelian, i.e.  $A' \cap M = A$ , then  $\mathcal{R}(A \subset M)$  coincides with the orbit equivalence relation induced by the normalizer  $\mathcal{N}_M(A)$ . In particular, in that case the equivalence relation  $\mathcal{R}(A \subset M)$  preserves the probability measure  $\mu$ .

If however  $A \subset M$  is not maximal abelian, the partial automorphisms of  $A$  induced by the partial isometries in the set (1) need not be trace preserving. So in general,  $\mathcal{R}(A \subset M)$  can be an equivalence relation of type III.

Our main technical result is Theorem 3.3 below, roughly saying the following. If  $A, B \subset M$  are abelian subalgebras such that  $\mathcal{Z}(A' \cap M) = A$  and  $\mathcal{Z}(B' \cap M) = B$ , and if there exist intertwining bimodules  $A \prec B$  and  $B \prec A$  (in the sense of Popa, see [Po03] and Theorem 2.3 below), then the equivalence relations  $\mathcal{R}(A \subset M)$  and  $\mathcal{R}(B \subset M)$  must be stably isomorphic. In particular, their types must be the same.

In Section 4, we apply this to  $M = L(\text{BS}(n, m))$  and  $A$  equal to the abelian von Neumann subalgebra generated by the unitary  $u_a$ . We prove that  $\mathcal{R}(A \subset M)$  is the unique hyperfinite ergodic equivalence relation of type  $\text{III}_{n/|m|}$ .

The proof of Theorem A can then be outlined as follows. First we note that the von Neumann algebra  $A' \cap M$  is nonamenable. Conversely if  $Q \subset M$  is a nonamenable subalgebra, it was proven in [CH08], using spectral gap rigidity ([Po06]) and the structure theory of amalgamated free product factors ([IPP05]), that  $Q' \cap M \prec A$ . So, up to intertwining-by-bimodules, the position of  $A$  inside  $M$  is “canonical”. Therefore a stable isomorphism between  $L(\text{BS}(n, m))$  and  $L(\text{BS}(n', m'))$  will preserve, up to intertwining-by-bimodules, these canonical abelian subalgebras. Hence their associated equivalence relations are stably isomorphic and, in particular, have the same type. This gives us the equality  $n/|m| = n'/|m'|$ .

## 2. PRELIMINARIES

We denote by  $(M, \tau)$  a von Neumann algebra equipped with a faithful normal tracial state  $\tau$ . We always assume that  $M$  has a *separable predual*. If  $B$  is a von Neumann subalgebra of  $(M, \tau)$ , we denote by  $E_B$  the unique trace preserving *conditional expectation* of  $M$  onto  $B$ .

Whenever  $x \in M$  is a normal element, we denote by  $\text{supp}(x)$  its support, i.e. the smallest projection  $p \in M$  that satisfies  $xp = x$  (or equivalently,  $px = x$ ).

Let  $\mathcal{R}$  be a countable nonsingular (i.e. measure class preserving) equivalence relation on a standard probability space  $(X, \mu)$ . We denote by  $[[\mathcal{R}]]$  the *full pseudogroup* of  $\mathcal{R}$ , i.e. the pseudogroup of all partial nonsingular automorphisms  $\varphi$  of  $X$  such that the graph of  $\varphi$  is contained in  $\mathcal{R}$ . We denote the domain of  $\varphi$  by  $\text{dom}(\varphi)$  and its range by  $\text{ran}(\varphi)$ . We denote by  $[x]$  the equivalence class of  $x \in X$ .

Assume that also  $\mathcal{R}'$  is a countable nonsingular equivalence relation on the standard probability space  $(X', \mu')$ . The equivalence relations  $\mathcal{R}$  and  $\mathcal{R}'$  are called

- *isomorphic*, if there exists a nonsingular isomorphism  $\Delta : X \rightarrow X'$  such that  $\Delta([x]) = [\Delta(x)]$  for almost every  $x \in X$  ;
- *stably isomorphic*, if there exist Borel subsets  $Z \subset X$  and  $Z' \subset X'$  that meet almost every orbit and a nonsingular isomorphism  $\Delta : Z \rightarrow Z'$  such that  $\Delta([x] \cap Z) = [\Delta(x)] \cap Z'$  for almost every  $x \in Z$ .

### 2.1. HNN EXTENSIONS AND BAUMSLAG-SOLITAR GROUPS

Let  $G$  be a group,  $H < G$  a subgroup and  $\theta : H \rightarrow G$  an injective group homomorphism. The *HNN extension*  $\text{HNN}(G, H, \theta)$  is defined as the group generated by  $G$  and an additional element  $t$  subject to the relation  $\theta(h) = tht^{-1}$  for all  $h \in H$ . So,

$$\text{HNN}(G, H, \theta) = \langle G, t \mid \theta(h) = tht^{-1} \text{ for all } h \in H \rangle .$$

Elements of  $\text{HNN}(G, H, \theta)$  can be canonically written as “reduced words” using as letters the elements of  $G$  and the letters  $t^{\pm 1}$ . More precisely, we have the following lemma.

LEMMA 2.1 (Britton’s lemma, [Br63]). *Consider the expression  $g = g_0 t^{n_1} g_1 t^{n_2} \dots t^{n_k} g_k$  with  $k \geq 0$ ,  $g_0, g_k \in G$ ,  $g_1, \dots, g_{k-1} \in G - \{e\}$  and  $n_1, \dots, n_k \in \mathbb{Z} - \{0\}$ . We call this expression reduced if the following two conditions hold:*

- for every  $i \in \{1, \dots, k - 1\}$  with  $n_i > 0$  and  $n_{i+1} < 0$ , we have  $g_i \notin H$ ,
- for every  $i \in \{1, \dots, k - 1\}$  with  $n_i < 0$  and  $n_{i+1} > 0$ , we have  $g_i \notin \theta(H)$ .

If the above expression for  $g$  is reduced, then  $g \neq e$  in the group  $\text{HNN}(G, H, \theta)$ , unless  $k = 0$  and  $g_0 = e$ . In particular, the natural homomorphism of  $G$  to  $\text{HNN}(G, H, \theta)$  is injective.

Recall from the introduction that the Baumslag-Solitar group  $\text{BS}(n, m)$  is defined for all  $n, m \in \mathbb{Z} - \{0\}$  as

$$\text{BS}(n, m) := \langle a, b \mid ba^n b^{-1} = a^m \rangle .$$

It is one of the easiest examples of an HNN extension. We also recall from the introduction that the  $\text{BS}(n, m)$  with  $2 \leq n < |m|$  form a complete list of all nonamenable icc Baumslag-Solitar groups up to isomorphism. Since we only want to consider the case where  $L(\text{BS}(n, m))$  is a nonamenable  $\text{II}_1$  factor, we always assume that  $2 \leq n < |m|$ .

2.2. HILBERT BIMODULES AND INTERTWINING-BY-BIMODULES

If  $M$  and  $N$  are tracial von Neumann algebras, then a *left  $M$ -module* is a Hilbert space  $\mathcal{H}$  endowed with a normal  $*$ -homomorphism  $\pi : M \rightarrow \text{B}(\mathcal{H})$ . A *right  $N$ -module* is a left  $N^{\text{op}}$ -module. An  *$M$ - $N$ -bimodule* is a Hilbert space  $\mathcal{H}$  endowed with commuting normal  $*$ -homomorphisms  $\pi : M \rightarrow \text{B}(\mathcal{H})$  and  $\varphi : N^{\text{op}} \rightarrow \text{B}(\mathcal{H})$ . For  $x \in M, y \in N$  and  $\xi \in \mathcal{H}$ , we write  $x\xi y$  instead of  $\pi(x)\varphi(y^{\text{op}})(\xi)$ . We denote an  $M$ - $N$ -bimodule  $\mathcal{H}$  by  ${}_M\mathcal{H}_N$ . We call an  $M$ - $N$ -bimodule *bifinite* if it is finitely generated both as a left Hilbert  $M$ -module and a right Hilbert  $N$ -module.

Let  $A$  and  $B$  be abelian von Neumann algebras. We denote by  $\text{PIso}(A, B)$  the set of all partial isomorphisms from  $A$  to  $B$ , i.e. isomorphisms  $\alpha : Aq \rightarrow Bp$ , where  $q \in A$  and  $p \in B$  are projections. We write  $\text{PAut}(A)$  instead of  $\text{PIso}(A, A)$ . Note that to every  $\alpha \in \text{PIso}(A, B)$  we can associate an  $A$ - $B$ -bimodule  ${}_A\mathcal{H}(\alpha)_B$  given by  $\mathcal{H}(\alpha) = L^2(Bp)$  and  $a\xi b = \alpha(aq)\xi bp$ . The composition of two partial isomorphisms is defined as follows: if  $\alpha \in \text{PIso}(B, C)$  and  $\beta \in \text{PIso}(A, B)$  are given by  $\alpha : Bp \rightarrow Cr$  and  $\beta : Aq \rightarrow Bp'$  for projections  $q \in A, p, p' \in B$  and  $r \in C$ , then the composition  $\alpha \circ \beta \in \text{PIso}(A, C)$  is defined by  $x \mapsto \alpha(\beta(x))$  for all  $x \in Aq\beta^{-1}(pp')$ .

The following is a well known result.

LEMMA 2.2. *Let  $A$  and  $B$  be abelian von Neumann algebras. Then every bifinite  $A$ - $B$ -bimodule  ${}_A\mathcal{H}_B$  is isomorphic to a direct sum of bimodules of the form  ${}_A\mathcal{H}(\alpha)_B$  with  $\alpha \in \text{PIso}(A, B)$ .*

We finally recall Popa's *intertwining-by-bimodules* theorem.

THEOREM 2.3 ([Po03, Theorem 2.1 and Corollary 2.3]). *Let  $(M, \tau)$  be a tracial von Neumann algebra and let  $A, B \subset M$  be possibly nonunital von Neumann subalgebras. Denote their respective units by  $1_A$  and  $1_B$ . The following three conditions are equivalent:*

1.  $1_A L^2(M) 1_B$  admits a nonzero  $A$ - $B$ -subbimodule that is finitely generated as a right  $B$ -module.
2. There exist nonzero projections  $p \in A$ ,  $q \in B$ , a normal unital  $*$ -homomorphism  $\psi : pAp \rightarrow qBq$  and a nonzero partial isometry  $v \in pMq$  such that  $av = v\psi(a)$  for all  $a \in pAp$ .
3. There is no sequence of unitaries  $u_n \in \mathcal{U}(A)$  satisfying  $\|E_B(xu_n y^*)\|_2 \rightarrow 0$  for all  $x, y \in 1_B M 1_A$ .

If one of these equivalent conditions holds, we write  $A \prec_M B$ .

### 2.3. QUASI-REGULARITY

Let  $(M, \tau)$  be a tracial von Neumann algebra and  $N \subset M$  a von Neumann subalgebra. We denote by  $\text{QN}_M(N)$  the *quasi-normalizer* of  $N$  inside  $M$ , i.e. the unital  $*$ -algebra defined by

$$\left\{ a \in M \mid \exists b_1, \dots, b_k \in M, \exists d_1, \dots, d_r \in M \right. \\ \left. \text{such that } Na \subset \sum_{i=1}^k b_i N \text{ and } aN \subset \sum_{j=1}^r N d_j \right\}.$$

We call  $N \subset M$  *quasi-regular* if  $\text{QN}_M(N)'' = M$ .

If  $A, B \subset M$  are abelian von Neumann subalgebras, we define  $\text{Q}_M(A, B)$  as

$$\text{Q}_M(A, B) := \{v \in M \mid vv^* \in A' \cap M, v^*v \in B' \cap M \text{ and } Av = vB\}.$$

Whenever  $v \in \text{Q}_M(A, B)$ , we define  $q_v = \text{supp}(E_A(vv^*))$  and  $p_v = \text{supp}(E_B(v^*v))$ , and we denote by  $\alpha_v : Aq_v \rightarrow Bp_v$  the unique  $*$ -isomorphism satisfying  $av = v\alpha_v(a)$  for all  $a \in Aq_v$ .

Note that the set  $\text{Q}_M(A, B)$  can be  $\{0\}$ . In Lemma 2.4, we will see that  $\text{Q}_M(A, B) \neq \{0\}$  if and only if there exists a bifinite  $A$ - $B$ -subbimodule  ${}_A\mathcal{H}_B$  of  ${}_A L^2(M)_B$ .

We denote  $\text{Q}_M(A, A)$  by  $\text{Q}_M(A)$ .

LEMMA 2.4. *Let  $(M, \tau)$  be a tracial von Neumann algebra and  $A, B \subset M$  abelian von Neumann subalgebras. Then the following statements hold.*

1. If  $\alpha \in \text{PIso}(A, B)$  and if  $\theta : {}_A\mathcal{H}(\alpha)_B \rightarrow {}_A L^2(M)_B$  is an  $A$ - $B$ -bimodular isometry, then there exists a partial isometry  $v \in \mathcal{Q}_M(A, B)$  such that  $\alpha = \alpha_v$  and such that

$$\theta(\mathcal{H}(\alpha)) \subset \overline{v(B' \cap M)}^{\|\cdot\|_2} \subset \overline{\text{span}}^{\|\cdot\|_2} \mathcal{Q}_M(A, B).$$

2. Every bifinite  $A$ - $B$ -subbimodule  ${}_A\mathcal{H}_B$  of  ${}_A L^2(M)_B$  is contained in  $\overline{\text{span}}^{\|\cdot\|_2} \mathcal{Q}_M(A, B)$ .
3.  $\mathcal{Q}_M(A)'' = \mathcal{QN}_M(A)''$ .
4. We have  $\mathcal{Q}_M(A, B) \neq \{0\}$  if and only if  ${}_A L^2(M)_B$  admits a nonzero bifinite  $A$ - $B$ -subbimodule.

*Proof.* 1. Let  $\alpha : Aq \rightarrow Bp$  be an element of  $\text{PIso}(A, B)$ . Define  $\xi := \theta(p) \in L^2(M)$  and let  $\xi = v|\xi|$  be its polar decomposition. For all  $a \in A$ , we have  $a\xi = \xi\alpha(a)$  and hence,  $av = v\alpha(a)$ . Furthermore  $p = \text{supp}(E_B(v^*v))$  and  $q = \alpha^{-1}(p) = \text{supp}(E_A(vv^*))$ . So we find that  $v \in \mathcal{Q}_M(A, B)$  and  $\alpha = \alpha_v$ . Because  $|\xi| \in L^2(B' \cap M)$ , we have that  $\xi = v|\xi|$  is an element of  $\overline{v(B' \cap M)}^{\|\cdot\|_2}$ . Since  $p$  generates  ${}_A\mathcal{H}(\alpha)_B$  as a right Hilbert  $B$ -module, we have proven the first inclusion  $\theta(\mathcal{H}(\alpha)) \subset \overline{v(B' \cap M)}^{\|\cdot\|_2}$ . Since  $v \in \mathcal{Q}_M(A, B)$ , also  $v(B' \cap M) \subset \mathcal{Q}_M(A, B)$  and the second inclusion in statement 1 is proven as well.

2. Let  ${}_A\mathcal{H}_B$  be a bifinite  $A$ - $B$ -subbimodule of  ${}_A L^2(M)_B$ . By Lemma 2.2,  ${}_A\mathcal{H}_B$  is isomorphic to a direct sum of bimodules of the form  ${}_A\mathcal{H}(\alpha_i)_B$  with  $\alpha_i \in \text{PIso}(A, B)$ . Using statement 1 of the lemma, we find that  $\mathcal{H}$  is generated by subspaces of  $\overline{\text{span}}^{\|\cdot\|_2} \mathcal{Q}_M(A, B)$ . This proves statement 2.

3. By definition, we have  $\mathcal{Q}_M(A)'' \subset \mathcal{QN}_M(A)''$ . On the other hand,  ${}_A L^2(\mathcal{QN}_M(A)'')_A$  is a direct sum of bifinite  $A$ - $A$ -subbimodules of  ${}_A L^2(M)_A$ . So by statement 2, we have that  $L^2(\mathcal{QN}_M(A)'') \subset \overline{\text{span}}^{\|\cdot\|_2}(\mathcal{Q}_M(A))$ . Therefore we conclude that  $\mathcal{QN}_M(A)'' = \mathcal{Q}_M(A)''$ .

Finally, 4 is an immediate consequence of 2. □

We end this subsection with the following lemma, clarifying why later, we will consider abelian subalgebras  $A \subset M$  satisfying  $\mathcal{Z}(A' \cap M) = A$ . Note that since  $A$  is abelian, the condition  $\mathcal{Z}(A' \cap M) = A$  is equivalent with the “bicommutant” property  $(A' \cap M)' \cap M = A$ . Also note that the composition of two partial isomorphisms was defined before Lemma 2.2.

LEMMA 2.5. *Let  $(M, \tau)$  be a tracial von Neumann algebra and  $A, B, C \subset M$  abelian von Neumann subalgebras. If  $v \in \mathcal{Q}_M(A, B)$ ,  $w \in \mathcal{Q}_M(B, C)$  and if  $\mathcal{Z}(B' \cap M) = B$ , then there exists an element  $u \in \mathcal{Q}_M(A, C)$  such that  $\alpha_w \circ \alpha_v = \alpha_u$ .*

*Proof.* Choose  $v \in \mathcal{Q}_M(A, B)$  and  $w \in \mathcal{Q}_M(B, C)$ . Note that  $vbw \in \mathcal{Q}_M(A, C)$  for every  $b \in B' \cap M$  and  $\alpha_{vbw} = \alpha_w \circ \alpha_v|_{Aq_{vbw}}$ . We claim that  $\bigvee_{b \in B' \cap M} q_{vbw} = \alpha_v^{-1}(q_w p_v)$ .

Denote  $\alpha_v^{-1}(q_w p_v) - \bigvee_{b \in B' \cap M} q_{vbw}$  by  $r$ . We need to prove that  $r$  is zero. Since  $(B' \cap M)' \cap M = B$ , we have that for every  $x \in M$ , the projection  $\text{supp}(E_B(xx^*))$  equals the projection of  $L^2(M)$  onto the closed linear span of  $(B' \cap M)xM \subset L^2(M)$ . Since  $w^*bv^*r = 0$  for every  $b \in B' \cap M$ , it follows that  $w^*q_{v^*r} = 0$ . Therefore  $q_w$  is orthogonal to  $q_{v^*r}$ . Because  $q_{v^*r} = \alpha_v(r)$  and  $\alpha_v(r) \leq q_w$ , it follows that  $\alpha_v(r) = 0$ . Hence  $r = 0$  and our claim that  $\bigvee_{b \in B' \cap M} q_{vbw} = \alpha_v^{-1}(q_w p_v)$  is proven.

By cutting down with appropriate projections, we find  $b_n \in B' \cap M$  such that the projections  $q_{vb_n w}$  are orthogonal and sum up to  $\alpha_v^{-1}(q_w p_v)$ . In particular, the left supports, resp. right supports, of the elements  $vb_n w$  are orthogonal. So we can define  $u = \sum_n vb_n w$ . It follows that  $u \in \mathcal{Q}_M(A, C)$  and  $\alpha_w \circ \alpha_v = \alpha_u$ .  $\square$

#### 2.4. THE TYPE OF AN ERGODIC NONSINGULAR COUNTABLE EQUIVALENCE RELATION

Let  $\mathcal{R}$  be a nonsingular ergodic countable Borel equivalence relation on a standard probability space  $(X, \mu)$ . Using the map  $\pi : \mathcal{R} \rightarrow X : \pi(x, y) = x$ , we define the measure  $\mu^{(1)}$  on  $\mathcal{R}$  given by

$$\mu^{(1)}(U) = \int_X \#(U \cap \pi^{-1}(x)) d\mu(x) \quad \text{for all Borel sets } U \subset \mathcal{R}.$$

We define  $\mathcal{R}^{(2)} := \{(x, y, z) \in X^3 \mid (x, y), (y, z) \in \mathcal{R}\}$ . Similarly, using the map  $\rho : \mathcal{R}^{(2)} \rightarrow X : \rho(x, y, z) = x$ , we define the measure  $\mu^{(2)}$  on  $\mathcal{R}^{(2)}$  given by

$$\mu^{(2)}(V) = \int_X \#(V \cap \rho^{-1}(x)) d\mu(x) \quad \text{for all Borel sets } V \subset \mathcal{R}^{(2)}.$$

The *Radon-Nikodym 1-cocycle* of  $\mathcal{R}$  is the  $\mu^{(1)}$ -a.e. uniquely defined Borel map  $\omega : \mathcal{R} \rightarrow \mathbb{R}$  such that

$$\omega(\varphi(x), x) = \log\left(\frac{d\mu \circ \varphi}{d\mu}(x)\right) \quad \text{for all } \varphi \in [[\mathcal{R}]] \text{ and almost every } x \in \text{dom } \varphi.$$

Note that  $\omega$  satisfies the 1-cocycle relation  $\omega(x, z) = \omega(x, y) + \omega(y, z)$  for  $\mu^{(2)}$ -a.e.  $(x, y, z) \in \mathcal{R}^{(2)}$ . One then defines the *Maharam extension*  $\tilde{\mathcal{R}}$  of  $\mathcal{R}$  as the equivalence relation on  $(X \times \mathbb{R}, \mu \times \exp(-t)dt)$  defined by

$$(x, t) \sim (y, s) \quad \text{if and only if } (x, y) \in \mathcal{R} \text{ and } t - s = \omega(x, y).$$

Note that  $\mu \times \exp(-t)dt$  is an infinite invariant measure for  $\tilde{\mathcal{R}}$ . Denote the von Neumann algebra of all  $\tilde{\mathcal{R}}$ -invariant functions in  $L^\infty(X \times \mathbb{R})$  by  $L^\infty(X \times \mathbb{R})^{\tilde{\mathcal{R}}}$ . Since  $\mathcal{R}$  was assumed to be ergodic, one can easily check that the action of  $\mathbb{R}$  on  $L^\infty(X \times \mathbb{R})^{\tilde{\mathcal{R}}}$  given by translation of the second variable, is also ergodic. Depending on how this action of  $\mathbb{R}$  looks like, we define as follows the type of  $\mathcal{R}$ .

- I or II, if the action is conjugate with  $\mathbb{R} \curvearrowright \mathbb{R}$ ;

- III $_{\lambda}$  ( $0 < \lambda < 1$ ), if the action is conjugate with  $\mathbb{R} \curvearrowright \mathbb{R}/\mathbb{Z} \log(\lambda)$  ;
- III $_1$ , if the action is on one point ;
- III $_0$ , if the action is properly ergodic, i.e. is ergodic and has orbits of measure zero.

REMARK 2.6. Denote by  $L(\mathcal{R})$  the von Neumann algebra associated with  $\mathcal{R}$ . Denote by  $\varphi$  the normal semifinite faithful state on  $L(\mathcal{R})$  that is induced by  $\mu$ . Finally denote by  $(\sigma_t^\varphi)_{t \in \mathbb{R}}$  its modular automorphism group. There is a canonical identification  $L(\tilde{\mathcal{R}}) \cong L(\mathcal{R}) \rtimes_{\sigma^\varphi} \mathbb{R}$ . Under this identification, the dual action of  $\mathbb{R}$  on  $L(\mathcal{R}) \rtimes_{\sigma^\varphi} \mathbb{R}$  corresponds to the action of  $\mathbb{R}$  on  $L(\tilde{\mathcal{R}})$  that we defined above. Also, the center of  $L(\mathcal{R}) \rtimes_{\sigma^\varphi} \mathbb{R}$  corresponds to  $L^\infty(X \times \mathbb{R})^{\tilde{\mathcal{R}}}$ . Altogether it follows that the type of the equivalence relation  $\mathcal{R}$  coincides with the type of the factor  $L(\mathcal{R})$ .

LEMMA 2.7. *Let  $\mathcal{R}$  be a nonsingular ergodic countable Borel equivalence relation on the standard probability space  $(X, \mu)$ . Denote by  $\omega$  its Radon-Nikodym 1-cocycle. If the essential image  $\text{Im}(\omega)$  of  $\omega$  equals  $\log(\lambda)\mathbb{Z}$  for some  $0 < \lambda < 1$  and if the kernel  $\text{Ker}(\omega)$  of  $\omega$  is an ergodic equivalence relation, then  $\mathcal{R}$  is of type III $_{\lambda}$ .*

*Proof.* Since  $\text{Ker}(\omega)$  is an ergodic equivalence relation on  $(X, \mu)$ , we have

$$L^\infty(X \times \mathbb{R})^{\tilde{\mathcal{R}}} \subset L^\infty(X \times \mathbb{R})^{\text{Ker}(\omega)} = 1 \otimes L^\infty(\mathbb{R}) .$$

For a given  $F \in L^\infty(\mathbb{R})$ , we have that  $1 \otimes F$  is  $\tilde{\mathcal{R}}$ -invariant if and only if  $F$  is invariant under translation by the essential image of  $\omega$ . So,

$$L^\infty(X \times \mathbb{R})^{\tilde{\mathcal{R}}} = 1 \otimes L^\infty(\mathbb{R}/\log(\lambda)\mathbb{Z}) .$$

□

### 3. EQUIVALENCE RELATIONS ASSOCIATED TO SUBALGEBRAS THAT ARE ABELIAN, BUT NOT MAXIMAL ABELIAN

Throughout this section, we fix a tracial von Neumann algebra  $(M, \tau)$  with separable predual. We also fix an abelian von Neumann subalgebra  $A \subset M$  satisfying  $\mathcal{Z}(A' \cap M) = A$ . Choose a standard probability space  $(X, \mu)$  such that  $A = L^\infty(X, \mu)$ . For every nonsingular partial automorphism  $\varphi$  of  $(X, \mu)$ , we denote by  $\alpha_\varphi$  the corresponding partial automorphism of  $A$ .

We first prove that  $\text{Q}_M(A)$  induces a nonsingular countable Borel equivalence relation  $\mathcal{R}(A \subset M)$  on  $(X, \mu)$ . For this, we introduce the notation

$$\mathcal{G}(A \subset M) := \{\alpha_v \mid v \in \text{Q}_M(A)\} . \tag{2}$$

PROPOSITION 3.1. *There exists a nonsingular countable Borel equivalence relation  $\mathcal{R}$  on  $(X, \mu)$  with the following property: a nonsingular partial automorphism  $\varphi$  of  $X$  satisfies  $\alpha_\varphi \in \mathcal{G}(A \subset M)$  if and only if  $(x, \varphi(x)) \in \mathcal{R}$  for a.e.  $x \in \text{dom}(\varphi)$ .*

Moreover,  $\mathcal{R}$  is essentially unique: if a nonsingular countable Borel equivalence relation  $\mathcal{R}'$  on  $(X, \mu)$  satisfies the same property, then there exists a Borel subset  $X_0 \subset X$  with  $\mu(X - X_0) = 0$  and  $\mathcal{R}'|_{X_0} = \mathcal{R}|_{X_0}$ .

We denote  $\mathcal{R}(A \subset M) := \mathcal{R}$ . The equivalence relation  $\mathcal{R}(A \subset M)$  is ergodic if and only if  $\text{QN}_M(A)''$  is a factor.

Before proving Proposition 3.1, we introduce some terminology and a lemma. To every  $\alpha \in \text{PAut}(A)$  are associated the support projections  $q_\alpha, p_\alpha \in A$  such that  $\alpha : Aq_\alpha \rightarrow Ap_\alpha$  is a  $*$ -isomorphism. Assume that  $\alpha \in \text{PAut}(A)$  and  $\mathcal{F} \subset \text{PAut}(A)$ . We say that  $\alpha$  is a *gluing* of elements in  $\mathcal{F}$ , if there exists a sequence of elements  $\alpha_n \in \mathcal{F}$  and projections  $q_n \in A$  such that  $q_\alpha = \sum_n q_n$  and such that  $q_n \leq q_{\alpha_n}$  and  $\alpha|_{Aq_n} = \alpha_n|_{Aq_n}$  for all  $n$ .

LEMMA 3.2. *Let  $\mathcal{J} \subset \text{Q}_M(A)$  and  $v \in \text{Q}_M(A)$  such that  $v \in \overline{\text{span}}^{\|\cdot\|_2} \mathcal{J}$ . Then  $\alpha_v$  is a gluing of elements in  $\{\alpha_w \mid w \in \mathcal{J}\}$ .*

*Proof.* By a standard maximality argument, it suffices to prove that for every nonzero projection  $q \in Aq_v$ , there exists a nonzero subprojection  $q_0 \in Aq$  and a  $w \in \mathcal{J}$  such that  $q_0 \leq q_w$  and  $\alpha_v|_{Aq_0} = \alpha_w|_{Aq_0}$ .

So fix a nonzero projection  $q \in Aq_v$ . It follows that  $qE_A(vv^*) \neq 0$ . Since  $v \in \overline{\text{span}}^{\|\cdot\|_2} \mathcal{J}$ , we can pick a  $w \in \mathcal{J}$  such that  $qE_A(vv^*) \neq 0$ . Define  $q_1 := \text{supp}(E_A(vv^*))$  and note that  $q_1 \in q_v Aq_w = Aq_v q_w$ . Also note that  $qq_1 \neq 0$ . For all  $a \in A$ , we have

$$\alpha_v^{-1}(ap_v)vv^* = vaw^* = vw^*\alpha_w^{-1}(ap_w).$$

Applying the conditional expectation onto  $A$  and using that  $A$  is abelian, we find that

$$\alpha_v^{-1}(ap_v)q_1 = \alpha_w^{-1}(ap_w)q_1 \quad \text{for all } a \in A.$$

This means that  $\alpha_v|_{Aq_1} = \alpha_w|_{Aq_1}$ . We put  $q_0 := qq_1$ . We already showed that  $q_0 \neq 0$ . Since  $q_0 \leq q_1$ , we have that  $\alpha_v|_{Aq_0} = \alpha_w|_{Aq_0}$ .  $\square$

*Proof of Proposition 3.1.* We say that a subpseudogroup  $\mathcal{G} \subset \text{PAut}(A)$  is of *countable type* if there exists a countable subset  $\mathcal{J} \subset \mathcal{G}$  such that every  $\alpha \in \mathcal{G}$  is a gluing of elements in  $\mathcal{J}$ . To prove the first part of the proposition, we must show that  $\mathcal{G}(A \subset M)$  is a subpseudogroup of countable type of  $\text{PAut}(A)$ . From Lemma 2.5, it follows that  $\mathcal{G}(A \subset M)$  is indeed a subpseudogroup. Since  $M$  has a separable predual, we can choose a countable  $\|\cdot\|_2$ -dense subset  $\mathcal{J} \subset \text{Q}_M(A)$ . By Lemma 3.2, every  $\alpha \in \mathcal{G}(A \subset M)$  is a gluing of elements in  $\{\alpha_w \mid w \in \mathcal{J}\}$ . Hence  $\mathcal{G}(A \subset M)$  is of countable type. So the first part of the proposition is proven and we can essentially uniquely define the nonsingular countable Borel equivalence relation  $\mathcal{R}$  on  $(X, \mu)$ .

Since  $A' \cap M \subset \text{QN}_M(A)''$  and since we assumed that  $(A' \cap M)' \cap M = A$ , the center of  $\text{QN}_M(A)''$  is a subalgebra of  $A$ . By Lemma 2.4.3, we have  $\text{QN}_M(A)'' = \text{Q}_M(A)''$ . Therefore,

$$\mathcal{Z}(\text{QN}_M(A)'' ) = \{a \in A \mid av = va \text{ for all } v \in \text{Q}_M(A)\}.$$

The right hand side equals  $A^{\mathcal{R}}$ , the subalgebra of  $\mathcal{R}$ -invariant functions in  $A$ . So  $\mathcal{R}$  is ergodic if and only if  $\text{QN}_M(A)''$  is a factor.  $\square$

For our application, the following theorem is crucial. It says that  $\mathcal{R}(A \subset M)$  remains the same, up to stable isomorphism, if we replace  $A$  by an abelian subalgebra  $B$  that has a mutual intertwining bimodule into  $A$ .

**THEOREM 3.3.** *Let  $M$  be a  $II_1$  factor with separable predual. Let  $A, B \subset M$  be abelian, quasi-regular von Neumann subalgebras satisfying  $\mathcal{Z}(A' \cap M) = A$  and  $\mathcal{Z}(B' \cap M) = B$ . If  $A \prec_M B$  and  $B \prec_M A$ , then the equivalence relations  $\mathcal{R}(A \subset M)$  and  $\mathcal{R}(B \subset M)$  are stably isomorphic.*

*Proof.* Since  $A, B$  are quasi-regular and since  $A \prec_M B$  as well as  $B \prec_M A$ , there exists a nonzero bifinite  $A$ - $B$ -subbimodule of  $L^2(M)$ . So by Lemma 2.4.4, there exists a nonzero element  $v \in \text{Q}_M(A, B)$  with corresponding  $\alpha_v \in \text{PAut}(A, B)$ . Using the notation in (2) and using Lemma 2.5, we find that

$$\begin{aligned} \alpha_v \circ \beta \circ \alpha_v^{-1} &\in \mathcal{G}(B \subset M) \quad \text{for all } \beta \in \mathcal{G}(A \subset M) \quad \text{and} \\ \alpha_v^{-1} \circ \gamma \circ \alpha_v &\in \mathcal{G}(A \subset M) \quad \text{for all } \gamma \in \mathcal{G}(B \subset M). \end{aligned}$$

So  $\alpha_v$  implements a stable isomorphism between  $\mathcal{R}(A \subset M)$  and  $\mathcal{R}(B \subset M)$ .  $\square$

The following lemma will allow us to easily compute  $\mathcal{R}(A \subset M)$  in concrete examples.

**LEMMA 3.4.** *Let  $(M, \tau)$  be a tracial von Neumann algebra and  $A \subset M$  an abelian von Neumann subalgebra satisfying  $\mathcal{Z}(A' \cap M) = A$ . Let  $\mathcal{F} \subset M$  be a subset such that*

- $M = (\mathcal{F} \cup \mathcal{F}^* \cup (A' \cap M))''$ ,
- as an  $A$ - $A$ -bimodule,  $\overline{\text{span}}^{\|\cdot\|_2} A\mathcal{F}A$  is isomorphic to a direct sum of bimodules of the form  ${}_A\mathcal{H}(\alpha_n)_A$  with  $\alpha_n \in \text{PAut}(A)$ .

Choose nonsingular partial automorphisms  $\varphi_n$  of  $(X, \mu)$  such that  $\alpha_n = \alpha_{\varphi_n}$  for all  $n$ . Up to measure zero,  $\mathcal{R}(A \subset M)$  is generated by the graphs of the partial automorphisms  $\varphi_n$ .

*Proof.* We again use the notation (2). By Lemma 2.4.1, we find  $v_n \in \text{Q}_M(A)$  such that  $\alpha_n = \alpha_{v_n}$  and

$$\overline{\text{span}}^{\|\cdot\|_2} A\mathcal{F}A \subset \overline{\text{span}}^{\|\cdot\|_2} \{v_n(A' \cap M) \mid n \in \mathbb{N}\}. \tag{3}$$

In particular, we have  $\alpha_n \in \mathcal{G}(A \subset M)$ . Choose nonsingular partial automorphisms  $\varphi_n$  of  $(X, \mu)$  such that  $\alpha_n = \alpha_{\varphi_n}$  for all  $n$ .

Denote by  $\mathcal{R}$  the smallest (up to measure zero) equivalence relation on  $(X, \mu)$  that contains the graphs of all the partial automorphisms  $\varphi_n$ . By the previous

paragraph, we know that  $\mathcal{R}$  is a subequivalence relation of  $\mathcal{R}(A \subset M)$ . Denote by  $\mathcal{J}$  the set of all products of elements in

$$\{v_n \mid n \in \mathbb{N}\} \cup \{v_n^* \mid n \in \mathbb{N}\} \cup (A' \cap M).$$

By construction, the graph of every  $\alpha_w$ ,  $w \in \mathcal{J}$ , belongs to  $\mathcal{R}$ . Combining our assumption that  $M = (\mathcal{F} \cup \mathcal{F}^* \cup (A' \cap M))''$  with (3), it follows that  $\text{span } \mathcal{J}$  is  $\|\cdot\|_2$ -dense in  $L^2(M)$ . By Lemma 3.2, every  $\alpha \in \mathcal{G}(A \subset M)$  is a gluing of elements in  $\{\alpha_w \mid w \in \mathcal{J}\}$ . So the graph of every  $\alpha \in \mathcal{G}(A \subset M)$  belongs to  $\mathcal{R}$  a.e. Hence  $\mathcal{R}$  equals  $\mathcal{R}(A \subset M)$  almost everywhere.  $\square$

We finally note in the following proposition that every nonsingular countable Borel equivalence relation  $\mathcal{R}$  arises as  $\mathcal{R}(A \subset M)$ .

**PROPOSITION 3.5.** *Let  $\mathcal{R}$  be a nonsingular countable Borel equivalence relation. Then there exists a quasi-regular inclusion of an abelian von Neumann algebra  $A$  in a tracial von Neumann algebra  $(M, \tau)$  satisfying  $\mathcal{Z}(A' \cap M) = A$  and such that  $\mathcal{R} \cong \mathcal{R}(A \subset M)$ .*

*Proof.* Let  $\mathcal{R}$  be a nonsingular countable Borel equivalence relation on a standard probability space  $(X, \mu)$ . Denote by  $(P, \text{Tr})$  the unique hyperfinite  $\text{II}_\infty$  factor and choose a trace-scaling action  $(\alpha_t)_{t \in \mathbb{R}}$  of  $\mathbb{R}$  on  $P$ . This means that  $\text{Tr} \circ \alpha_t = e^{-t} \text{Tr}$ . The corresponding action of  $\mathbb{R}$  on  $L^2(P)$  will also be denoted by  $(\alpha_t)$ . We denote by  $\omega : \mathcal{R} \rightarrow \mathbb{R}$  the Radon-Nikodym 1-cocycle of  $\mathcal{R}$  (see Section 2.4).

In the same way as with the Maharam extension of a nonsingular group action, the equivalence relation  $\mathcal{R}$  admits a natural trace preserving action on  $L^\infty(X) \overline{\otimes} P$ . We denote by  $(\mathcal{M}, \text{Tr})$  the crossed product. For completeness, we recall the construction of  $(\mathcal{M}, \text{Tr})$ . To every  $\varphi \in [[\mathcal{R}]]$ , we associate the operator  $W_\varphi$  on  $L^2(\mathcal{R}, L^2(P))$  given by

$$(W_\varphi \xi)(x, y) = \begin{cases} \alpha_{\omega(x, \varphi^{-1}(x))}(\xi(\varphi^{-1}(x), y)) & \text{if } x \in \text{dom}(\varphi^{-1}), \\ 0 & \text{otherwise,} \end{cases}$$

for every  $\xi \in L^2(\mathcal{R}, L^2(P))$ . One checks that  $W_\varphi W_\psi = W_{\varphi \circ \psi}$  and  $W_\varphi^* = W_{\varphi^{-1}}$ . We represent  $L^\infty(X) \overline{\otimes} P = L^\infty(X, P)$  on  $L^2(\mathcal{R}, L^2(P))$  by

$$(F\xi)(x, y) = F(x)\xi(x, y) \text{ for all } \xi \in L^2(\mathcal{R}, L^2(P)) \text{ and } F \in L^\infty(X, P).$$

Note that the partial isometries  $W_\varphi$ ,  $\varphi \in [[\mathcal{R}]]$ , normalize  $L^\infty(X, P)$  and that

$$(W_\varphi^* F W_\varphi)(x) = \begin{cases} \alpha_{\omega(x, \varphi(x))}(F(\varphi(x))) & \text{if } x \in \text{dom } \varphi \\ 0 & \text{otherwise.} \end{cases}$$

Define  $\mathcal{M}$  as the von Neumann algebra generated by  $L^\infty(X, P)$  and the partial isometries  $W_\varphi$ ,  $\varphi \in [[\mathcal{R}]]$ . Denoting by  $\Delta \subset \mathcal{R}$  the diagonal subset, the orthogonal projection onto  $L^2(\Delta, L^2(P))$  implements a normal faithful conditional expectation  $E : \mathcal{M} \rightarrow L^\infty(X) \overline{\otimes} P$  satisfying

$$E(W_\varphi) = \chi_{\{x \mid \varphi(x)=x\}} \otimes 1 \text{ for all } \varphi \in [[\mathcal{R}]].$$

The formula  $\text{Tr} := (\mu \otimes \text{Tr}) \circ E$  defines a normal semifinite faithful trace on  $\mathcal{M}$ . Fix a nonzero projection  $q \in P$  with  $\text{Tr}(q) = 1$ . Define the projection  $p \in L^\infty(X) \overline{\otimes} P$  given by  $p = 1 \otimes q$ . Write  $A := L^\infty(X)p$  and  $M := p\mathcal{M}p$ . Then  $A$  is a quasi-regular abelian von Neumann subalgebra of  $M$  and the restriction of  $\text{Tr}$  to  $M$  gives a normal faithful tracial state  $\tau$  on  $M$ . The relative commutant  $L^\infty(X)' \cap \mathcal{M}$  equals  $L^\infty(X) \overline{\otimes} P$ . Since  $P$  is a factor, it follows that  $\mathcal{Z}(A' \cap M) = A$ .

We finally prove that  $\mathcal{R} \cong \mathcal{R}(A \subset M)$ . Write  $\mathcal{R} = \bigcup_k \text{graph}(\varphi_k)$ , with  $\varphi_k \in [\mathcal{R}]$ . Then  $\varphi_k$  induces an automorphism of  $L^\infty(X)$  and hence of  $A = L^\infty(X)p$  that we denote by  $\beta_k \in \text{Aut}(A)$ . Since  $P$  is a  $\text{II}_\infty$  factor and  $q \in P$  is a finite projection, we can choose partial isometries  $w_n \in P$  such that  $\sum_n w_n^* w_n = 1$  and  $w_n w_n^* = q$  for all  $n$ . Define the elements

$$v_{n,k} := (1 \otimes w_n) W_{\varphi_k} p .$$

All  $v_{n,k}$  belong to  $\mathcal{Q}_M(A)$  and  $\alpha_{v_{n,k}}$  equals the restriction of  $\beta_k$  to  $Ap_{n,k}$  for projections  $p_{n,k} \in A$ . Since the sum of all  $w_n^* w_n$  equals 1, we also have that  $\bigvee_n p_{n,k} = p$ . Therefore the graphs of the partial automorphisms  $\alpha_{v_{n,k}}$  generate an equivalence relation that is isomorphic with  $\mathcal{R}$ . To conclude the proof, we put  $\mathcal{F} := \{v_{n,k} \mid n, k \in \mathbb{N}\}$  and observe that  $M = (\mathcal{F} \cup \mathcal{F}^* \cup (A' \cap M))''$ . By Lemma 3.4, the equivalence relation  $\mathcal{R}(A \subset M)$  is generated by the graphs of the partial automorphisms  $\alpha_{v_{n,k}}$ .  $\square$

4. PROOF OF THEOREM A

Throughout this section, we assume that  $n$  and  $m$  are integers satisfying  $2 \leq n < |m|$ . As explained in the introduction, the corresponding groups  $\text{BS}(n, m)$  form a complete list of the nonamenable icc Baumslag-Solitar groups up to isomorphism.

Throughout this section, we write  $M = L(\text{BS}(n, m))$  and  $A = \{u_a, u_a^*\}''$ . We start with the following observation.

PROPOSITION 4.1. *We have that  $A \subset M$  is a quasi-regular abelian von Neumann subalgebra satisfying  $\mathcal{Z}(A' \cap M) = A$ . Moreover,  $A' \cap M$  has no amenable direct summand.*

*Proof.* It is clear that  $A \subset M$  is a quasi-regular abelian von Neumann subalgebra, because the element  $a \in \text{BS}(n, m)$  generates an almost normal abelian subgroup of  $\text{BS}(n, m)$  : for every  $g \in \text{BS}(n, m)$ , the group  $ga^{\mathbb{Z}}g^{-1} \cap a^{\mathbb{Z}}$  has finite index in  $a^{\mathbb{Z}}$ .

To prove that  $\mathcal{Z}(A' \cap M) = A$ , we define the finite index subalgebra  $A_0 := \{u_a^n, u_a^{-n}\}''$  of  $A$ . We will first prove that  $\mathcal{Z}(A'_0 \cap M) = A_0$ . Afterwards we will show that this implies that  $\mathcal{Z}(A' \cap M) = A$ .

Define  $G := \langle a^{\mathbb{Z}}, b^{-1}a^{\mathbb{Z}}b \rangle \subset \text{BS}(n, m)$ . Then  $L(G)$  is a subalgebra of  $A'_0 \cap M$ . So  $\mathcal{Z}(A'_0 \cap M) \subset L(G)' \cap M$ . Using Lemma 2.1, one can easily see that  $\{g\gamma g^{-1} \mid g \in G\}$  is an infinite set for every  $\gamma \in \text{BS}(n, m) - a^{\mathbb{Z}}$ . Therefore  $L(G)' \cap M \subset A_0$ . This shows that  $\mathcal{Z}(A'_0 \cap M) \subset A_0$ . Since the converse inclusion is obvious, we find that  $A_0 = \mathcal{Z}(A'_0 \cap M)$ .

Since  $A_0 \subset A$  has finite index, there exist orthogonal projections  $p_j \in A$  such that  $Ap_j = A_0p_j$  and  $\sum_j p_j = \mathbf{1}$ . But then

$$\begin{aligned} \mathcal{Z}(A' \cap M)p_j &= \mathcal{Z}((A' \cap M)p_j) = \mathcal{Z}((Ap_j)' \cap p_jMp_j) \\ &= \mathcal{Z}((A_0p_j)' \cap p_jMp_j) = \mathcal{Z}(p_j(A'_0 \cap M)p_j) \\ &= \mathcal{Z}(A'_0 \cap M)p_j = A_0p_j \subset A \end{aligned}$$

Therefore  $\mathcal{Z}(A' \cap M) \subset A$ . The converse inclusion being obvious, we have proven that  $A = \mathcal{Z}(A' \cap M)$ .

Using Lemma 2.1, it follows that  $G$  is an amalgamated free product of two copies of  $\mathbb{Z}$  over a copy of  $\mathbb{Z}$  embedded as  $n\mathbb{Z}$  and  $m\mathbb{Z}$  respectively. In particular,  $G$  is nonamenable and  $L(G)$  has no amenable direct summand. Since  $L(G) \subset A'_0 \cap M$ , it follows that  $A'_0 \cap M$  has no amenable direct summand either. As above, we have that

$$(A' \cap M)p_j = p_j(A'_0 \cap M)p_j$$

for all  $j$ . Hence  $A' \cap M$  has no amenable direct summand.  $\square$

We now identify the associated countable equivalence relation  $\mathcal{R}(A \subset M)$ .

**PROPOSITION 4.2.** *The equivalence relation  $\mathcal{R}(A \subset M)$  is isomorphic with the unique hyperfinite ergodic countable equivalence relation of type III $_{n/|m|}$ .*

*Proof.* Let  $k$  be the greatest common divisor of  $n$  and  $|m|$ . Write  $n = n_0k$  and  $m = m_0k$ . By our assumptions on  $n$  and  $m$ , we have that  $1 \leq n_0 < |m_0|$ . Define the countable Borel equivalence relation  $\mathcal{R}_{n,m}$  on the circle  $\mathbb{T}$  given by

$$\mathcal{R}_{n,m} := \left\{ (y, z) \in \mathbb{T} \times \mathbb{T} \mid \exists a, b \in \mathbb{N} \text{ such that } a + b > 0 \text{ and } y^{(n_0^a m_0^b k)} = z^{(m_0^a n_0^b k)} \right\}.$$

Equip  $\mathbb{T}$  with its Lebesgue measure  $\lambda$  and note that  $\mathcal{R}_{n,m}$  is a nonsingular countable Borel equivalence relation on  $(\mathbb{T}, \lambda)$ .

Define  $\mathcal{R}_0 := \{(y, z) \in \mathbb{T} \times \mathbb{T} \mid y^m = z^n\}$ . Note that  $\mathcal{R}_0 \subset \mathcal{R}_{n,m}$  and that  $\mathcal{R}_{n,m}$  is the smallest equivalence relation containing  $\mathcal{R}_0$ . Define  $\pi : \mathcal{R}_0 \rightarrow \mathbb{T} : \pi(y, z) = y^m$ . Note that  $\pi$  is  $n|m|$ -to-1. Define the probability measure  $\mu$  on  $\mathcal{R}_0$  given by

$$\mu(U) = \frac{1}{n|m|} \int_{\mathbb{T}} \#(U \cap \pi^{-1}(\{x\})) d\lambda(x).$$

For all  $k, l \in \mathbb{Z}$ , we define the function  $P_{k,l} : \mathcal{R}_0 \rightarrow \mathbb{T} : P_{k,l}(y, z) = y^k z^l$ . A direct computation yields a unique unitary

$$T : L^2(\mathcal{R}_0, \mu) \rightarrow \overline{\text{span}}^{\|\cdot\|_2} Au_b A : P_{k,l} \mapsto u_a^k u_b u_a^l.$$

We turn  $L^2(\mathcal{R}_0, \mu)$  into an  $L^\infty(\mathbb{T})$ - $L^\infty(\mathbb{T})$ -bimodule by the formula

$$(F \cdot \xi \cdot F')(y, z) = F(y) \xi(y, z) F'(z).$$

Under the natural identification of  $L^\infty(\mathbb{T})$  and  $A$ , the unitary  $T$  is  $A$ - $A$ -bimodular.

By construction  ${}_A L^2(\mathcal{R}_0, \mu)_A$  is isomorphic with a direct sum of bimodules of the form  ${}_A \mathcal{H}(\alpha_j)_A$  where the union of the graphs of the partial automorphisms  $\alpha_j$  equals  $\mathcal{R}_0$  and hence generates the equivalence relation  $\mathcal{R}_{n,m}$ . Applying Lemma 3.4 to  $\mathcal{F} = \{u_b\}$ , we conclude that  $\mathcal{R}(A \subset M) \cong \mathcal{R}_{n,m}$  up to measure zero.

As in Section 2.4, denote by  $\omega : \mathcal{R}_{n,m} \rightarrow \mathbb{R}$  the Radon-Nikodym 1-cocycle. Denote by  $\Lambda \subset \mathbb{T}$  the subgroup given by

$$\Lambda := \left\{ \exp\left(\frac{2\pi i s}{(n_0 m_0)^b}\right) \mid s \in \mathbb{Z}, b \in \mathbb{N} \right\}.$$

For every  $z \in \Lambda$ , we denote by  $\alpha_z : \mathbb{T} \rightarrow \mathbb{T}$  the rotation  $\alpha_z(y) = zy$ . We have  $\text{graph } \alpha_z \subset \mathcal{R}_{n,m}$  for all  $z \in \Lambda$ . Since all  $\alpha_z$  are measure preserving, we actually have  $\text{graph } \alpha_z \subset \text{Ker}(\omega)$ . Since  $\Lambda \subset \mathbb{T}$  is a dense subgroup, it follows that  $\text{Ker}(\omega)$  is an ergodic equivalence relation. In particular,  $\mathcal{R}_{n,m}$  is ergodic. A direct computation shows that  $\omega(y, z) = \log(n/|m|)$  for all  $(y, z) \in \mathcal{R}_0$ . Since  $\mathcal{R}_0$  generates the equivalence relation  $\mathcal{R}_{n,m}$ , it follows that the essential image of  $\omega$  equals  $\log(n/|m|)\mathbb{Z}$ . Using Lemma 2.7, we conclude that  $\mathcal{R}_{n,m}$  is of type III $_{n/|m|}$ . By construction,  $\mathcal{R}_{n,m}$  is amenable and hence, hyperfinite.  $\square$

We are now ready to prove our main theorem.

*Proof of Theorem A.* Fix for  $i = 1, 2$ , integers  $n_i, m_i \in \mathbb{Z}$  with  $2 \leq n_i < |m_i|$ . Put  $M_i = L(\text{BS}(n_i, m_i))$  and denote by  $A_i \subset M_i$  the abelian von Neumann subalgebra generated by  $u_a$ , where  $a \in \text{BS}(n_i, m_i)$  is the first canonical generator. Assume that  $M_1$  and  $M_2$  are stably isomorphic. We must prove that

$$\frac{n_1}{|m_1|} = \frac{n_2}{|m_2|}. \tag{4}$$

Interchanging if necessary the roles of  $M_1$  and  $M_2$ , we can take a nonzero projection  $p_1 \in A_1$  and a  $*$ -isomorphism  $\alpha : p_1 M_1 p_1 \rightarrow M_2$ .

We claim that inside  $M_2$ , we have  $\alpha(A_1 p_1) \prec A_2$ . From Proposition 4.1, we know that

$$P := \alpha(A_1 p_1)' \cap M_2 = \alpha((A_1' \cap M_1) p_1)$$

has no amenable direct summand. By Proposition 3.1 in [Ue07], the HNN extension  $M_2$  can be viewed as the corner of an amalgamated free product of tracial von Neumann algebras. Since  $P$  has no amenable direct summand, it then follows from [CH08, Theorem 4.2] that  $P' \cap M_2 \prec A_2$ . So our claim that  $\alpha(A_1 p_1) \prec A_2$  follows.

By symmetry, we also have the intertwining  $\alpha^{-1}(A_2) \prec A_1 p_1$  inside  $p_1 M_1 p_1$ . Applying  $\alpha$ , we find that  $A_2 \prec \alpha(A_1 p_1)$  inside  $M_2$ .

Having proven that inside  $M_2$  we have the intertwining relations  $\alpha(A_1 p_1) \prec A_2$  and  $A_2 \prec \alpha(A_1 p_1)$ , it follows from Theorem 3.3 that the equivalence relations  $\mathcal{R}(A_1 p_1 \subset p_1 M_1 p_1)$  and  $\mathcal{R}(A_2 \subset M_2)$  are stably isomorphic. By construction,

$\mathcal{R}(A_1 p_1 \subset p_1 M_1 p_1)$  is the restriction of  $\mathcal{R}(A_1 \subset M_1)$  to the support of  $p_1$ . So we conclude that the equivalence relations  $\mathcal{R}(A_1 \subset M_1)$  and  $\mathcal{R}(A_2 \subset M_2)$  are stably isomorphic. In particular, these ergodic nonsingular equivalence relations must have the same type. Using Proposition 4.2, we find (4).  $\square$

## REFERENCES

- [BS62] G. Baumslag and D. Solitar, Some two-generator one-relator non-Hopfian groups. *Bull. Amer. Math. Soc.* 68 (1962) 199-201.
- [BV12] M. Berbec and S. Vaes,  $W^*$ -superrigidity for group von Neumann algebras of left-right wreath products. To appear in *Proc. Lond. Math. Soc.* arXiv:1210.0336
- [Br63] J.L. Britton, The word problem. *Ann. of Math.* 77 (1963), 16-32.
- [CH88] M. Cowling and U. Haagerup, Completely bounded multipliers of the Fourier algebra of a simple Lie group of real rank one. *Invent. Math.* 96 (1989), 507-549.
- [CH08] I. Chifan, C. Houdayer, Bass-Serre rigidity results in von Neumann algebras. *Duke Math. J.* 153 (2010), 23-54.
- [Co75] A. Connes, Classification of injective factors. *Ann. of Math.* 104 (1976), 73-115.
- [Co80] A. Connes, Classification des facteurs. In *Operator algebras and applications, Part 2 (Kingston, 1980)*, Proc. Sympos. Pure Math. 38, Amer. Math. Soc., Providence, 1982, pp. 43-109.
- [Fi10] P. Fima, A note on the von Neumann algebra of a Baumslag-Solitar group. *C. R. Acad. Sci. Paris, Ser. I* 349 (2011), 25-27.
- [IPP05] A. Ioana, J. Peterson, S. Popa, Amalgamated free products of  $w$ -rigid factors and calculation of their symmetry groups. *Acta Math.* 200 (2008), 85-153.
- [IPV10] A. Ioana, S. Popa and S. Vaes, A class of superrigid group von Neumann algebras. *Ann. Math.* 178 (2013), 231-286.
- [Ki11] Y. Kida, Invariants for orbit equivalence relations of Baumslag-Solitar groups. To appear in *Tohoku Math. J.* arXiv:1111.3701
- [Mo91] D.I. Moldavanskii, Isomorphism of the Baumslag-Solitar groups. *Ukrainian Math. J.* 43 (1991), 1569-1571.
- [Po03] S. Popa, Strong rigidity of  $II_1$  factors arising from malleable actions of  $w$ -rigid groups, I. *Invent. Math.* 165 (2006), 369-408.
- [Po04] S. Popa, Strong rigidity of  $II_1$  factors arising from malleable actions of  $w$ -rigid groups, II. *Invent. Math.* 165 (2006), 409-452.

- [Po06] S. Popa, On the superrigidity of malleable actions with spectral gap. *J. Amer. Math. Soc.* 21 (2008), 981-1000.
- [St05] Y. Stalder, Moyennabilité intérieure et extensions HNN. *Ann. Inst. Fourier* 56 (2006), 309-323.
- [Ue07] Y. Ueda, Remarks on HNN extensions in operator algebras. *Illinois J. Math.* 52 (2008), 705-725.

Niels Meesschaert  
KU Leuven  
Department of Mathematics  
Leuven (Belgium)  
niels.meesschaert@wis.kuleuven.be

Stefaan Vaes  
KU Leuven  
Department of Mathematics  
Leuven (Belgium)  
stefaan.vaes@wis.kuleuven.be

