

## A REMARK ON THE GRADIENT MAP

LEONARDO BILIOTTI, ALESSANDRO GHIGI, AND PETER HEINZNER<sup>1</sup>

Received: February 7, 2014

Communicated by Thomas Peternell

ABSTRACT. For a Hamiltonian action of a compact group  $U$  of isometries on a compact Kähler manifold  $Z$  and a compatible subgroup  $G$  of  $U^{\mathbb{C}}$ , we prove that for any closed  $G$ -invariant subset  $Y \subset Z$  the image of the gradient map  $\mu_{\mathfrak{p}}(Y)$  is independent of the choice of the invariant Kähler form  $\omega$  in its cohomology class  $[\omega]$ .

2010 Mathematics Subject Classification: 53D20

## 1. INTRODUCTION

Let  $(Z, \omega)$  be a compact Kähler manifold and let  $U$  be a compact connected semisimple Lie group such that  $U^{\mathbb{C}}$  acts holomorphically on  $Z$ ,  $U$  preserves  $\omega$  and there is a momentum map  $\mu : Z \rightarrow \mathfrak{u}^*$ . Let  $G \subset U^{\mathbb{C}}$  be a *compatible* subgroup. By this we mean a subgroup which is compatible with the Cartan involution  $\Theta$  of  $U^{\mathbb{C}}$  which defines  $U$ , i.e. if  $\mathfrak{p} = \mathfrak{g} \cap i\mathfrak{u}$  and  $K = U \cap G$ , then  $G = K \cdot \exp \mathfrak{p}$ . Let  $\mu_{\mathfrak{p}} : Z \rightarrow \mathfrak{p}$  be the associated *gradient map* (see [4, 5] or section 2).

In this note we prove the following.

**THEOREM 1.** *Let  $Y \subset Z$  be a closed  $G$ -stable subset. Then up to translation the set  $\mu_{\mathfrak{p}}(Y)$  is independent of the choice of the invariant Kähler form  $\omega$  in the cohomology class  $[\omega]$ .*

---

<sup>1</sup>The first two authors were partially supported by a grant of Max-Planck Institute für Mathematik, Bonn and by FIRB 2012 MIUR “Geometria differenziale e teoria geometrica delle funzioni”. author was partially supported also by PRIN 2009 MIUR “Moduli, strutture geometriche e loro applicazioni”. The third author was partially supported by DFG-priority program SPP 1388 (Darstellungstheorie).

Since  $Z$  is compact and  $G$  is compatible there is a stratification of  $Z$  analogous to the Kirwan stratification, see [4]. This gives a stratification of any closed  $G$ -invariant subset  $Y$  of  $Z$ , by intersecting the strata in  $Z$  with  $Y$ . It follows from Theorem 1 that when the momentum map is properly normalized (see Lemma 2) this stratification does not depend on the choice of  $\omega$  in its cohomology class. When  $Z$  is a projective manifold and  $\omega$  is the pull-back of a Fubini-Study form via an equivariant embedding of  $Z$  in  $\mathbb{P}^N$ , Kirwan [6, §12] proved that the stratification in terms of a properly normalized  $\mu$  can be defined purely in terms of algebraic geometry. In the present note we give a proof of this fact for a general compact Kähler manifold  $Z$  in the more general setting of *gradient* maps for actions of compatible subgroups on closed  $G$ -invariant subsets of  $Z$ . Another consequence of the above is the following. Assume that  $Z$  is a projective manifold and that  $[\omega]$  is an integral class. Let  $Y \subset Z$  be a closed  $G$ -invariant real semi-algebraic subset whose real algebraic Zariski closure is irreducible. Let  $\mathfrak{a} \subset \mathfrak{p}$  be a maximal subalgebra and let  $\mathfrak{a}_+$  be a closed Weyl chamber in  $\mathfrak{a}$ . Then  $A(Y)_+ := \mu_{\mathfrak{p}}(Y) \cap \mathfrak{a}_+$  is convex (see [2], which deals with the case when  $\omega$  is the restriction of a Fubini-Study metric).

ACKNOWLEDGEMENTS. The first two authors are grateful to the Fakultät für Mathematik of Ruhr-Universität Bochum for the wonderful hospitality during several visits. They also wish to thank the Max-Planck Institut für Mathematik, Bonn for excellent conditions provided during their visit at this institution, where part of this paper was written.

## 2. BACKGROUND

Let  $(Z, \omega)$  be a compact Kähler manifold and let  $U$  be a compact Lie group. Assume that  $U$  acts on  $Z$  by holomorphic Kähler isometries. Since  $Z$  is compact the  $U$ -action extends to a holomorphic action of the complexified group  $U^{\mathbb{C}}$ . Assume also that there is a momentum map  $\mu : Z \rightarrow \mathfrak{u}^* \cong \mathfrak{u}$ , where  $\mathfrak{u}^*$  is identified with  $\mathfrak{u}$  using a fixed  $U$ -invariant scalar product on  $\mathfrak{u}$  that we denote by  $\langle \cdot, \cdot \rangle$ . We also denote by  $\langle \cdot, \cdot \rangle$  the scalar product on  $i\mathfrak{u}$  such that multiplication by  $i$  is an isometry of  $\mathfrak{u}$  onto  $i\mathfrak{u}$ . If  $\xi \in \mathfrak{u}$  we denote by  $\xi_Z$  the fundamental vector field on  $Z$  and we let  $\mu^\xi \in C^\infty(Z)$  be the function  $\mu^\xi(z) := \langle \mu(z), \xi \rangle$ . That  $\mu$  is the momentum map means that it is  $U$ -equivariant and that  $d\mu^\xi = i_{\xi_Z}\omega$ . For a closed subgroup  $G \subset U^{\mathbb{C}}$  let  $K := G \cap U$  and  $\mathfrak{p} := \mathfrak{g} \cap i\mathfrak{u}$ . The group  $G$  is called *compatible* if  $G = K \cdot \exp \mathfrak{p}$  [4, 5]. In the following we fix a compatible subgroup  $G \subset U^{\mathbb{C}}$ . If  $z \in Z$ , let  $\mu_{\mathfrak{p}}(z) \in \mathfrak{p}$  denote  $-i$  times the component of  $\mu(z)$  in the direction of  $i\mathfrak{p}$ . In other words we require that  $\langle \mu_{\mathfrak{p}}(z), \beta \rangle = -\langle \mu(z), i\beta \rangle$  for any  $\beta \in \mathfrak{p}$ . The map

$$\mu_{\mathfrak{p}} : Z \rightarrow \mathfrak{p}$$

is called the *gradient map* (see [3]) or *restricted momentum map*. Let  $\mu_{\mathfrak{p}}^\beta \in C^\infty(Z)$  be the function  $\mu_{\mathfrak{p}}^\beta(z) = \langle \mu_{\mathfrak{p}}(z), \beta \rangle = \mu^{-i\beta}(z)$ . Let  $(\cdot, \cdot)$  be the Kähler metric associated to  $\omega$ , i.e.  $(v, w) = \omega(v, Jw)$ . Then  $\beta_Z$  is the gradient of  $\mu_{\mathfrak{p}}^\beta$  with respect to  $(\cdot, \cdot)$ .

EXAMPLE 1. (1) For any compact subgroup  $K \subset U$ , both  $K$  and its complexification  $G = K^{\mathbb{C}}$  are compatible. In particular  $G = U^{\mathbb{C}}$  is a compatible subgroup. (2) If  $G$  is a real form of  $U^{\mathbb{C}}$ , then  $G$  is compatible. (3) For any  $\xi \in \mathfrak{iu}$ , the subgroup  $G = \exp(\mathbb{R}\xi)$  is compatible.

Next we recall the Stratification Theorem for actions of compatible subgroups. Given a maximal subalgebra  $\mathfrak{a} \subset \mathfrak{p}$  and a Weyl chamber  $\mathfrak{a}^+ \subset \mathfrak{a}$  define

$$\begin{aligned} \eta_{\mathfrak{p}} : X &\rightarrow \mathbb{R} & \eta_{\mathfrak{p}}(x) &:= \frac{1}{2} \|\mu_{\mathfrak{p}}(x)\|^2 \\ C_{\mathfrak{p}} &:= \text{Crit}(\eta_{\mathfrak{p}}) & \mathcal{B}_{\mathfrak{p}} &:= \mu_{\mathfrak{p}}(C_{\mathfrak{p}}) & \mathcal{B}_{\mathfrak{p}}^+ &:= \mathcal{B}_{\mathfrak{p}} \cap \mathfrak{a}^+ \\ X(\mu) &:= \{x \in X : \overline{G \cdot x} \cap \mu_{\mathfrak{p}}^{-1}(0) \neq \emptyset\} \end{aligned}$$

where  $X$  is a compact  $G$ -invariant subset of  $Z$ . Points lying in  $X(\mu)$  are called *semistable*. Using semistability and the function  $\eta_{\mathfrak{p}}$  one can define a stratification of  $X$  in the following way, see [6] and [4]. For  $\beta \in \mathcal{B}_{\mathfrak{p}}^+$  set

$$\begin{aligned} X_{\|\beta\|^2} &:= \{x \in X : \overline{\exp(\mathbb{R}\beta) \cdot x} \cap (\mu^{\beta})^{-1}(\|\beta\|^2) \neq \emptyset\} \\ X^{\beta} &:= \{x \in X : \beta_X(x) = 0\} \\ X_{\|\beta\|^2}^{\beta} &:= X^{\beta} \cap X_{\|\beta\|^2} \\ X_{\|\beta\|^2}^{\beta+} &:= \{x \in X_{\|\beta\|^2} : \lim_{t \rightarrow -\infty} \exp(t\beta) \cdot x \text{ exists and it lies in } X_{\|\beta\|^2}^{\beta}\} \\ G^{\beta+} &:= \{g \in G : \text{the limit } \lim_{t \rightarrow -\infty} \exp(t\beta)g \exp(-t\beta) \text{ exists in } G\}. \end{aligned}$$

Set also

$$G^{\beta} := \{g \in G : \text{Ad } g(\beta) = \beta\} \quad \mathfrak{p}^{\beta} := \{\xi \in \mathfrak{p} : [\xi, \beta] = 0\}.$$

The group  $G^{\beta} = K^{\beta} \cdot \exp(\mathfrak{p}^{\beta})$  is a compatible subgroup of  $U^{\mathbb{C}}$  and the set  $X_{\|\beta\|^2}^{\beta+}$  is  $G^{\beta+}$ -invariant. Denote by  $\mu_{\mathfrak{p}^{\beta}}$  the composition of  $\mu_{\mathfrak{p}}$  with the orthogonal projection  $\mathfrak{p} \rightarrow \mathfrak{p}^{\beta}$ . Then  $\mu_{\mathfrak{p}^{\beta}}$  is a gradient map for the  $G^{\beta}$ -action on  $X_{\|\beta\|^2}^{\beta+}$ . We set  $\widehat{\mu}_{\mathfrak{p}^{\beta}} := \mu_{\mathfrak{p}^{\beta}} - \beta$ . Since  $\beta$  lies in the center of  $\mathfrak{g}^{\beta}$  and since  $G^{\beta}$  is a compatible subgroup of  $(U^{\beta})^{\mathbb{C}} = (U^{\mathbb{C}})^{\beta}$ , it is a gradient map too. We let  $S^{\beta+}$  denote the set of  $G^{\beta}$ -semistable points in  $X_{\|\beta\|^2}^{\beta+}$  with respect to  $\widehat{\mu}_{\mathfrak{p}^{\beta}}$ , i.e.

$$S^{\beta+} := \{x \in X_{\|\beta\|^2}^{\beta+} : \overline{G^{\beta} \cdot x} \cap \mu_{\mathfrak{p}^{\beta}}^{-1}(\beta) \neq \emptyset\}.$$

The set  $S^{\beta+}$  coincides with the set of semistable points of the group  $G^{\beta}$  in  $X_{\|\beta\|^2}^{\beta+}$  after shifting. By definition the  $\beta$ -stratum is given by  $S_{\beta} := G \cdot S^{\beta+}$ .

STRATIFICATION THEOREM. (See [4, Thm. 7.3]) *Assume that  $X$  is a compact  $G$ -invariant subset of  $Z$ . Then  $\mathcal{B}_{\mathfrak{p}}^+$  is finite and*

$$X = \bigsqcup_{\beta \in \mathcal{B}_{\mathfrak{p}}^+} S_{\beta}.$$

Moreover

$$\overline{S_\beta} \subset S_\beta \cup \bigcup_{\|\gamma\| > \|\beta\|} S_\gamma.$$

### 3. PROOF OF THEOREM 1

For a  $U$ -invariant function  $f$  on  $Z$  we set

$$\tilde{\omega} := \omega + dd^c f$$

where  $d^c f := -2J^*df$ . Since  $Z$  is compact and  $U$  acts by holomorphic transformations, any  $U$ -invariant Kähler form  $\tilde{\omega}$  in the Kähler class  $[\omega]$  can be written in this way. Since pluriharmonic functions on  $Z$  are constant, the function  $f$  is unique up to a constant.

LEMMA 2. *If  $\mu : Z \rightarrow \mathfrak{u}$  is a momentum map for the  $U$ -action on  $Z$  with respect to  $\omega$ , then the map  $\tilde{\mu} : Z \rightarrow \mathfrak{u}$  defined by*

$$\tilde{\mu}^\xi := \mu^\xi - d^c f(\xi_Z) \tag{3}$$

*is a momentum map for the  $U$ -action on  $Z$  with respect to  $\tilde{\omega}$ .*

*Proof.* That  $\tilde{\mu}$  is a momentum map follows from Cartan formula using that  $L_{\xi_Z} d^c f = d^c L_{\xi_Z} f = 0$ . This in turn follows from the assumption that the action of  $U$  is holomorphic and  $f$  is  $U$ -invariant.  $\square$

A more precise version of Theorem 1 is the following.

THEOREM 4. *For any closed  $G$ -stable subset  $Y \subset Z$  we have  $\mu_{\mathfrak{p}}(Y) = \tilde{\mu}_{\mathfrak{p}}(Y)$ .*

*Proof.* Let  $\mathfrak{a} \subset \mathfrak{p}$  be a maximal subalgebra and set  $A := \exp \mathfrak{a}$ . The group  $A$  is a compatible subgroup. Let  $\mu_{\mathfrak{a}} : Z \rightarrow \mathfrak{a}$  be the restricted gradient map. Any connected subgroup  $B \subset A$  is compatible. Given such a  $B$ , set  $Z^{(B)} := \{z \in Z : A_z = B\}$ . A connected component  $S$  of  $Z^{(B)}$  will be called an  $A$ -stratum of type  $\mathfrak{b}$ . For a given  $S$  let  $C$  denote the connected component of  $Z^B$  containing  $S$ . Then  $C$  is a complex submanifold of  $Z$  and the Slice Theorem (see Theorem 14.10 and 14.21 in [3] or Theorem 2.2 in [2]) applied to the  $A$ -action on  $C$  shows that  $S$  is open and dense in  $C$ .

Let  $A^c$  be the Zariski closure of  $A$  in  $U^c$ . The group  $A^c$  is a compatible subgroup of  $U^c$ ,  $A^c \cap U = T$  is a torus and  $A^c = T \exp(it)$ , where  $\mathfrak{t}$  denotes the Lie algebra of  $T$ . Moreover  $\overline{S}$  is  $A^c$ -stable [2, Lemma 3.3 (1)]. Denote by  $\mu_{\mathfrak{t}} : Z \rightarrow \mathfrak{t}$  the momentum map obtained by projecting  $\mu : Z \rightarrow \mathfrak{u}$  to  $\mathfrak{t}$ , and denote by  $\Pi : it \rightarrow \mathfrak{a}$  the orthogonal projection. Then  $\mu_{\mathfrak{a}} = \Pi \circ i\mu_{\mathfrak{t}}$  and  $\mu_{\mathfrak{a}}(\overline{S}) = \Pi(i\mu_{\mathfrak{t}}(\overline{S}))$ . By the convexity theorem of Atiyah-Guillemin-Sternberg  $\mu_{\mathfrak{t}}(\overline{S})$  is a convex polytope and its vertices are images of points fixed by  $A^c$ . It follows that  $\mu_{\mathfrak{a}}(\overline{S})$  is a convex polytope as well. Since  $\Pi$  is linear, any vertex of  $\mu_{\mathfrak{a}}(\overline{S})$  is the projection of at least one vertex of  $i\mu_{\mathfrak{t}}(\overline{S})$ . Therefore  $\mu_{\mathfrak{a}}(\overline{S})$  is the convex hull of  $\mu_{\mathfrak{a}}(\overline{S}^A)$ . Now we use Lemma 2: if  $x \in \overline{S}^A$ , then  $\xi_Z(x) = 0$ , so  $\tilde{\mu}^\xi(x) = \mu^\xi(x)$ , for any  $\xi \in \mathfrak{a}$ . Therefore  $\tilde{\mu}_{\mathfrak{a}}(x) = \mu_{\mathfrak{a}}(x)$  for every  $A$ -fixed point

$x$ . It follows that both  $\mu_{\mathfrak{a}}(\overline{S})$  and the affine subspace spanned by  $\mu_{\mathfrak{a}}(S)$  do not depend on the choice of the Kähler form  $\omega$ .

Let  $\Sigma$  be the collection of affine hyperplanes of  $\mathfrak{a}$  that are affine hulls of  $\mu_{\mathfrak{a}}(\overline{S})$  for some  $A$ -stratum  $S$ . Set  $P := \mu_{\mathfrak{a}}(Z)$  and

$$P_0 := P - \bigcup_{H \in \Sigma} P \cap H.$$

(This construction is similar to the one in [2, §§4-5].) The set  $P_0$  is an open subset of  $\mathfrak{a}$ . Let  $C(P_0)$  denote the set of its connected components. This is a finite set. For  $\gamma \in C(P_0)$  let  $P(\gamma)$  be the closure of the connected component  $\gamma$ . Then  $P(\gamma)$  is a convex polytope. Since both  $P$  and the hyperplanes  $H$  are independent of  $\omega$ , also the polytopes  $P(\gamma)$  do not depend on  $\omega$ . By [2, Corollary 5.8]

$$\mu_{\mathfrak{p}}(Y) \cap \mathfrak{a} = \bigcup_{\gamma \in F(\omega)} P(\gamma),$$

where  $F(\omega) \subset \Gamma$  is some subset of  $C(P_0)$ . One can join  $\omega$  to  $\tilde{\omega}$  continuously, e.g. by  $\omega_t := \omega + tdd^c f$ . Then  $\tilde{\mu}_t := \mu - td^c f(\cdot_Z)$  also depends continuously on  $t$ . So  $P(\gamma) \subset \mu_{\mathfrak{p}}(Y) \cap \mathfrak{a}$  if and only if  $P(\gamma) \subset \mu_{t,\mathfrak{p}}(Y) \cap \mathfrak{a}$ . Therefore  $F(\omega_t)$  is independent of  $t$ . Thus  $\mu_{\mathfrak{p}}(Y) \cap \mathfrak{a} = \tilde{\mu}_{\mathfrak{p}}(Y) \cap \mathfrak{a}$ . Since  $\mu_{\mathfrak{p}}(Y) = K(\mu_{\mathfrak{p}}(Y) \cap \mathfrak{a})$  this implies  $\mu_{\mathfrak{p}}(Y) = \tilde{\mu}_{\mathfrak{p}}(Y)$ .  $\square$

**COROLLARY 5.** *Assume that  $Z$  is connected and let  $\omega$  and  $\tilde{\omega}$  be two cohomologous Kähler forms with momentum maps  $\mu$  and  $\tilde{\mu}$  respectively as in Lemma 2. Then  $\tilde{\mu}$  is the unique momentum map such that  $\mu(Z) = \tilde{\mu}(Z)$ .*

*Proof.* Since two momentum maps with respect to  $\tilde{\omega}$  differ by addition of an element of the center of  $\mathfrak{u}$ , it is clear that there is at most one such map with the image equal to  $\mu(Z)$ . To complete the proof it is therefore enough to check that  $\tilde{\mu}(Z) = \mu(Z)$ . This is a special case of the previous theorem.  $\square$

**THEOREM 6.** *Let  $\omega$  and  $\tilde{\omega}$  be two cohomologous Kähler forms on  $Z$ , with momentum maps  $\mu$  and  $\tilde{\mu}$  respectively as in Lemma 2. Then the set  $\mathcal{B}_{\mathfrak{p}}^+$  is the same for both momentum maps and the two stratifications of  $X$  coincide.*

*Proof.* By [4, Corollary 7.6]

$$\mathcal{B}_{\mathfrak{p}} = \{ \beta \in \mathfrak{p} : \text{there exists } x \in X : \frac{\|\beta\|^2}{2} = \inf_{G \cdot x} \eta_{\mathfrak{p}} \text{ and } \beta \in \mu_{\mathfrak{p}}(\overline{G \cdot x}) \}. \quad (7)$$

Moreover for  $\beta \in \mathcal{B}_{\mathfrak{p}}$

$$S_{\beta} = \{ x \in X : \frac{\|\beta\|^2}{2} = \inf_{G \cdot x} \eta_{\mathfrak{p}} \text{ and } \beta \in \mu_{\mathfrak{p}}(\overline{G \cdot x}) \}. \quad (8)$$

For any point  $x \in X$ , the set  $\overline{G \cdot x}$  is closed and  $G$ -invariant. Hence by Theorem 4  $\mu_{\mathfrak{p}}(\overline{G \cdot x}) = \tilde{\mu}_{\mathfrak{p}}(\overline{G \cdot x})$ . From this it follows that  $\inf_{G \cdot x} \eta_{\mathfrak{p}} = \inf_{G \cdot x} \tilde{\eta}_{\mathfrak{p}}$ , where  $\tilde{\eta}_{\mathfrak{p}} := \|\tilde{\mu}_{\mathfrak{p}}\|^2/2$ . The result follows from (7) and (8).  $\square$

From the above we obtain the following generalization.

COROLLARY 9. *If  $Z$  is a complex projective manifold,  $U$  is a compact connected semisimple Lie group acting on  $Z$ ,  $\omega$  is a  $U$ -invariant Hodge metric and  $Y \subset Z$  is a closed  $G$ -invariant real semi-algebraic subset whose real algebraic Zariski closure is irreducible, then  $A(Y)_+$  is convex. Moreover if  $G$  is semisimple, then  $X(\mu)$  is dense (if it is nonempty).*

*Proof.* By assumption there is a very ample line bundle  $L \rightarrow Z$  such that  $[\omega] = 2\pi c_1(L)/m$  for an integer  $m > 0$ . Let  $\omega_{FS}$  be a  $U$ -invariant Fubini-Study metric on  $\mathbb{P}(H^0(Z, L)^*)$ . Let  $\mu_{FS}$  be the momentum map with respect to  $\omega_{FS}|_Z$ . In [2] the convexity theorem has been proved for  $\mu_{FS}$ . A rescaling in the symplectic form yields a corresponding rescaling in the momentum map. Therefore the convexity theorem also holds for the momentum map  $\tilde{\mu}$  relative to the symplectic form  $\tilde{\omega} := \omega_{FS}/m$ . So it holds also for  $\mu$ , since  $\mu_{\mathfrak{p}}(Y) = \tilde{\mu}_{\mathfrak{p}}(Y)$  by Theorem 4. The proof of the last statement is similar: see [2] and Corollary 5.  $\square$

COROLLARY 10. *Under the same assumptions, any local minimum of  $\|\mu_{\mathfrak{p}}\|^2$  is a global minimum.*

*Proof.* This follows since  $\|\mu_{\mathfrak{p}}\|^2$  is  $K$ -invariant and  $\mu(Z)_+$  is a convex subset of  $\mathfrak{a}_+$ .  $\square$

COROLLARY 11. *If  $\omega$  and  $\omega'$  are cohomologous Kähler forms on  $Z$  with momentum maps  $\mu$  and  $\tilde{\mu}$  as in Lemma 2, then  $X(\mu) = X(\tilde{\mu})$ .*

*Proof.* It is enough to observe that  $X(\mu) = S_0$ .  $\square$

#### REFERENCES

- [1] P. Heinzner and A. Huckleberry. Kählerian potentials and convexity properties of the moment map. *Invent. Math.*, 126(1):65–84, 1996.
- [2] P. Heinzner and P. Schützdeller. Convexity properties of gradient maps. *Adv. Math.*, 225(3):1119–1133, 2010.
- [3] P. Heinzner and G. W. Schwarz. Cartan decomposition of the moment map. *Math. Ann.*, 337(1):197–232, 2007.
- [4] P. Heinzner, G. W. Schwarz, and H. Stötzel. Stratifications with respect to actions of real reductive groups. *Compos. Math.*, 144(1):163–185, 2008.
- [5] P. Heinzner and H. Stötzel. Critical points of the square of the momentum map. In *Global aspects of complex geometry*, pages 211–226. Springer, Berlin, 2006.
- [6] F. C. Kirwan. *Cohomology of quotients in symplectic and algebraic geometry*, volume 31 of *Mathematical Notes*. Princeton University Press, Princeton, NJ, 1984.

Leonardo Biliotti  
Università di Parma  
leonardo.biliotti@unipr.it

Alessandro Ghigi  
Università di Milano  
Bicocca  
alessandro.ghigi@unimib.it

Peter Heinzner  
Ruhr Universität Bochum  
peter.heinzner@rub.de

