

FREE CURVES ON VARIETIES

FRANK GOUNELAS

Received: June 26, 2014

Revised: January 30, 2016

Communicated by Thomas Peternell

ABSTRACT. We study various generalisations of rationally connected varieties, allowing the connecting curves to be of higher genus. The main focus will be on free curves $f : C \rightarrow X$ with large unobstructed deformation space as originally defined by Kollár, but we also give definitions and basic properties of varieties X covered by a family of curves of a fixed genus g so that through any two general points of X there passes the image of a curve in the family. We prove that the existence of a free curve of genus $g \geq 1$ implies the variety is rationally connected in characteristic zero and initiate a study of the problem in positive characteristic.

2010 Mathematics Subject Classification: 14M20, 14M22, 14H10.

1. INTRODUCTION

Let k be an algebraically closed field. A smooth projective rationally connected variety, originally defined in [Cam92] and [KMM92], is a variety such that through every two general points there passes the image of a rational curve. In characteristic zero this is equivalent to the notion of a separably rationally connected variety, given by the existence of a rational curve $f : \mathbb{P}^1 \rightarrow X$ such that $f^* \mathcal{T}_X$ is ample. In characteristic p , however, one has to distinguish between these two notions. Deformations of a morphism $f : \mathbb{P}^1 \rightarrow X$ are controlled by the sheaf $f^* \mathcal{T}_X$, hence studying positivity conditions of this bundle is intimately tied to deformation theory and the existence of many rational curves on X . Rationally connected varieties have especially nice properties and an introduction to the theory is contained in [Kol96] and [Deb01]. Note in particular the important theorem of Graber-Harris-Starr [GHS03] (and de Jong-Starr [dJS03] in positive characteristic) which we will make repeated use of throughout this paper, which says that a separably rationally connected fibration over a curve admits a section. An equivalent statement in characteristic zero is that the maximal rationally connected (MRC) quotient $R(X)$ is not uniruled (see [Kol96, IV.5.6.3]), although this can fail in positive characteristic.

In this paper we study various ways in which a variety can be connected by higher genus curves. After an introductory section with auxiliary results on vector bundles on curves and Frobenius, we consider first varieties which admit a morphism from a family of curves of fixed arithmetic genus g whose product with itself dominates the product of the variety with itself and call these varieties “genus g connected”, generalising the notion of there being a rational curve through two general points. We also consider C -connected varieties, where there exists a family $C \times U \rightarrow X$ of a single smooth genus g curve C such that $C \times C \times U \rightarrow X \times X$ is dominant. Mori’s Bend and Break result allows us to produce rational curves going through a fixed point given a higher genus curve which has large enough deformation space. For example, in Proposition 3.6 as an easy corollary, we show that over any characteristic, if for any two general points of a smooth projective variety X with $\dim X \geq 3$ there passes the image of a morphism from a fixed curve C of genus g , then X is uniruled. This fails for surfaces, where an example is provided.

A stronger condition than the aforementioned is the existence of a morphism from a curve which deforms a lot without obstructions, as discussed for separably rationally connected varieties above. Namely, for $f : C \rightarrow X$ a morphism to a variety X where C is of any genus g , Kollár [Kol96] defines f to be free if $f^* \mathcal{T}_X$ is globally generated as a vector bundle on C and also $H^1(C, f^* \mathcal{T}_X) = 0$. In the case of genus $g = 0$ one must distinguish between free and very free curves. Geometrically, the former implies that $f : \mathbb{P}^1 \rightarrow X$ deforms so that its image covers all points in X (hence X is uniruled) whereas the latter that it can do so even fixing a point $x \in X$ (X rationally connected). If $g \geq 1$, however, after defining an r -free curve to be one which deforms keeping any r points fixed, we show that the notions of the existence of a free (0-free) and very free (1-free) curve coincide and in fact are equivalent with the existence of a curve $f : C \rightarrow X$ such that $f^* \mathcal{T}_X$ is ample.

THEOREM. (see 5.5) *Let X be a smooth projective variety and C a smooth projective curve of genus $g \geq 1$ over an algebraically closed field k . Then for any $r \geq 0$, there exists an $f : C \rightarrow X$ which is r -free if and only if there exists a morphism $f' : C \rightarrow X$ such that $f'^* \mathcal{T}_X$ is ample.*

Work of Bogomolov-McQuillan (see [BM01], [KSCT07]) on foliations which restrict to an ample bundle on a smooth curve sitting inside a complex variety X shows that the leaves of such a foliation are not only algebraic but in fact have rationally connected closures. From the above, one deduces this result in the case of the foliation $\mathcal{F} = \mathcal{T}_X$, complementing the currently known connections between existence of curves with large deformation space and rationally connected varieties (cf. the uniruledness criterion of Miyaoka [Miy87]). Our proof emphasises the use of free curves and C -connected varieties, in particular with a view towards similar results in positive characteristic.

THEOREM. (see 5.2) *Let X be a smooth projective variety over an algebraically closed field of characteristic zero and let $f : C \rightarrow X$ be a smooth projective curve of genus $g \geq 1$ such that $f^* \mathcal{T}_X$ is globally generated and $H^1(C, f^* \mathcal{T}_X) = 0$. Then X is rationally connected.*

In the sixth section we study the particular case of elliptically connected varieties (i.e. genus one connected varieties) where, even allowing a covering family of genus 1 curves to vary in moduli, one can prove the following theorem.

THEOREM. (Theorem 6.2) *Let X be a smooth projective variety over an algebraically closed field of characteristic zero. Then the following two statements are equivalent*

- (1) *There exists $\mathcal{C} \rightarrow U$ a flat projective family of irreducible genus 1 curves with a map $\mathcal{C} \rightarrow X$ such that $\mathcal{C} \times_U \mathcal{C} \rightarrow X \times X$ is dominant.*
- (2) *X is either rationally connected or a rationally connected fibration over a curve of genus one.*

In positive characteristic, at this point we have not been able to prove that the existence of a higher genus free curve implies the existence of a very free rational curve (which would mean that X is separably rationally connected). We work however in this direction, establishing this result in dimensions two (with a short discussion about dimension three) and furthermore by studying algebraic implications of the existence of a free higher genus curve, such as the vanishing of pluricanonical forms and triviality of the Albanese variety. In the final section we give an example of a threefold in characteristic p whose MRC quotient is rationally connected and which has infinite fundamental group.

The study of rational curves on varieties is an important and active area of research, and shedding light on the existence of rational curves coming from the deformation theory of higher genus curves is a theme explored in a variety of sources, for example the minimal model program or [BDPP13]. Aside from the unresolved difficulties arising in positive characteristic, the author expects uniruledness and rational connected results of the type presented in this article to be of use in moduli theory.

ACKNOWLEDGEMENTS. The contents of this paper are from the author's thesis under the supervision of Victor Flynn, whom I would like to thank for his continuous encouragement. I am indebted to Damiano Testa for the many hours spent helping with the material of this paper and to Johan de Jong not only for the hospitality at Columbia University but also for helping improve the contents of this paper. I would also like to thank Jason Starr and Yongqi Liang for comments, János Kollár for pointing out a similar construction to that in the last section and Mike Roth for showing me how abelian surfaces are C -connected. The anonymous referee's numerous suggestions and corrections also significantly improved this paper. This research was completed under the support of EPSRC grant number EP/F060661/1 at the University of Oxford.

2. AMPLE VECTOR BUNDLES AND FROBENIUS

We begin with some results concerning positivity of vector bundles on curves. Recall that a locally free sheaf \mathcal{E} on a scheme X is called ample if $\mathcal{O}_{\mathbb{P}(\mathcal{E})}(1)$ has this property. Equivalent definitions involving global generation of $\mathcal{F} \otimes S^n(\mathcal{E})$ for \mathcal{F} a coherent sheaf and n large enough, and also cohomological vanishing criteria can be found in [Har66]. Ampleness on curves can be checked using various criteria such as the following.

LEMMA 2.1. *Let C be a smooth projective curve of genus $g \geq 2$ over an algebraically closed field of characteristic zero and \mathcal{E} a locally free sheaf on C such that $H^1(C, \mathcal{E}) = 0$. It follows that \mathcal{E} is ample.*

PROOF. From [Har71, Theorem 2.4], it suffices to show that every non-trivial quotient locally free sheaf of \mathcal{E} has positive degree. Let $\mathcal{E} \rightarrow \mathcal{E}' \rightarrow 0$ be a quotient. From the long exact sequence in cohomology we see that $H^1(C, \mathcal{E}')$ is also 0. From the Riemann-Roch formula $\deg \mathcal{E}' = h^0(C, \mathcal{E}') + (\text{rk } \mathcal{E}')(g - 1)$ and since $g \geq 2$ we deduce that $\deg \mathcal{E}' > 0$. \square

Note that Hartshorne's ampleness criterion only works in characteristic zero. More generally, over any characteristic if we further assume that our locally free sheaf is globally generated then the same result holds so long as the genus is at least one.

PROPOSITION 2.2. *Let C be a smooth projective curve of genus $g \geq 1$ over an algebraically closed field k and \mathcal{E} a globally generated locally free sheaf on C such that $H^1(C, \mathcal{E}) = 0$. Then \mathcal{E} is ample.*

PROOF. Since \mathcal{E} is globally generated, there exists a positive integer n such that $\mathcal{O}_C^{\oplus n} \rightarrow \mathcal{E} \rightarrow 0$ is exact. This gives (see [Har77, ex. II.3.12]) a closed immersion of the respective projective bundles $\mathbb{P}(\mathcal{E}) \hookrightarrow \mathbb{P}^{n-1}$. By projecting onto the first factor we have the following diagram

$$\begin{array}{ccccc} \mathbb{P}(\mathcal{E}) & \xrightarrow{i} & \mathbb{P}^{n-1} \times C & \xrightarrow{\text{pr}_1} & \mathbb{P}^{n-1} \\ & \searrow \pi & \downarrow \text{pr}_2 & & \\ & & C & & \end{array}$$

and from [Har77, II.5.12] we have $\text{pr}_1^* \mathcal{O}_{\mathbb{P}^{n-1}}(1) = \mathcal{O}_{\mathbb{P}^{n-1} \times C}(1)$. Also, since i is a closed immersion it follows that $i^* \mathcal{O}_{\mathbb{P}^{n-1} \times C}(1) = \mathcal{O}_{\mathbb{P}^{n-1} \times C}(1)|_{\mathbb{P}(\mathcal{E})} = \mathcal{O}_{\mathbb{P}(\mathcal{E})}(1)$ which concludes that $i^* \text{pr}_1^* \mathcal{O}_{\mathbb{P}^{n-1}}(1) = \mathcal{O}_{\mathbb{P}(\mathcal{E})}(1)$. To show that \mathcal{E} is an ample locally free sheaf on C it is enough to show that this invertible sheaf is ample. Since we know that $\mathcal{O}_{\mathbb{P}^{n-1}}(1)$ is ample though, it is sufficient to show that $i \circ \text{pr}_1$ is a finite morphism. Since it is projective, we need only show that it is quasi-finite. Hence assuming that the fibre of $i \circ \text{pr}_1$ over a general point $p \in \mathbb{P}^{n-1}$ is not finite, it must be the whole of C . We now embed this fibre $j : C \rightarrow \mathbb{P}(\mathcal{E})$ as a section to π and pull back the surjection $\pi^* \mathcal{E} \rightarrow \mathcal{O}_{\mathbb{P}(\mathcal{E})}(1)$ via j , obtaining $j^* \mathcal{O}_{\mathbb{P}(\mathcal{E})}(1)$ as a quotient of $j^* \pi^* \mathcal{E} = \mathcal{E}$ (see [Har77, II.7.12]).

However $\text{pr}_1 \circ i \circ j : C \rightarrow \mathbb{P}^{n-1}$ is a constant map so $j^* \mathcal{O}_{\mathbb{P}(\mathcal{E})}(1) = \mathcal{O}_C$. Taking cohomology of the corresponding short exact sequence given by this quotient, we obtain a contradiction since $H^1(C, \mathcal{E}) = 0$ whereas $H^1(C, \mathcal{O}_C)$ is not trivial for $g \geq 1$. \square

In Proposition 2.4 below we will prove that given an ample bundle on a curve in positive characteristic, then after pulling back by Frobenius, we can make this bundle be globally generated and have vanishing first cohomology.

LEMMA 2.3. *Let C be a smooth projective curve over an algebraically closed field k , $d \geq 0$ an integer and \mathcal{E} a locally free sheaf on C . If $H^1(C, \mathcal{E}(-D)) = 0$ for all effective divisors D of fixed degree d then for $d' < d$ it follows that $H^1(C, \mathcal{E}(-D')) = 0$ and $\mathcal{E}(-D')$ is globally generated for all effective divisors D' of degree d' .*

PROOF. The first result follows from the short exact sequence

$$0 \rightarrow \mathcal{E}(-D' - R) \rightarrow \mathcal{E}(-D') \rightarrow \mathcal{E}(-D')|_R \rightarrow 0$$

where R is an effective divisor of degree $d - d'$. For the second, let $p \in C$. From the first part we have $H^1(C, \mathcal{E}(-D' - p)) = 0$ since $D' + p$ is an effective divisor of degree $d' + 1 \leq d$ so the following sequence is exact

$$0 \rightarrow H^0(C, \mathcal{E}(-D' - p)) \rightarrow H^0(C, \mathcal{E}(-D')) \rightarrow \mathcal{E}(-D') \otimes k(p) \rightarrow 0.$$

Hence $\mathcal{E}(-D')$ is globally generated at p and the result follows. \square

A partial converse to Proposition 2.2 in characteristic p is given in [KSCT07, Proposition 9], using \mathbb{Q} -twisted vector bundles as in [Laz04, II.6.4]. We prove the following different version of this result.

PROPOSITION 2.4. *Let C be a smooth projective curve of genus g over an algebraically closed field k of characteristic p and let \mathcal{E} be an ample locally free sheaf on C . Let $B \subset C$ be a closed subscheme of length b and ideal sheaf \mathcal{I}_B . Then there exists a positive integer n such that $H^1(C^{(n)}, F_n^* \mathcal{E} \otimes \mathcal{I}_B) = 0$ and $F_n^* \mathcal{E} \otimes \mathcal{I}_B$ is globally generated on $C^{(n)}$ where $F_n : C^{(n)} \rightarrow C$ the n -fold composition of the k -linear Frobenius morphism.*

PROOF. We proceed by induction. First, assume we can write \mathcal{E} as an extension

$$0 \rightarrow \mathcal{M} \rightarrow \mathcal{E} \rightarrow \mathcal{Q} \rightarrow 0$$

where \mathcal{M} is an ample line bundle. If \mathcal{Q} is not torsion free, consider the saturation of \mathcal{M} in \mathcal{E} instead and take \mathcal{Q} as that quotient. Since \mathcal{E} is ample, so is its quotient \mathcal{Q} . Note also that the rank of \mathcal{Q} is one less than that of \mathcal{E} and that if we can prove the result for \mathcal{Q} then we will have it for \mathcal{E} too by considering cohomology of the appropriate exact sequences. We thus reduce to the case of $\mathcal{E} = \mathcal{L}$ an invertible sheaf of positive degree (since it is ample). An invertible sheaf \mathcal{L} pulls back under the n -fold composition of the linear Frobenius morphism to an invertible sheaf $F_n^* \mathcal{L}$ of degree $p^n \text{deg } \mathcal{L}$. To show that $H^1(C^{(n)}, F_n^* \mathcal{L} \otimes \mathcal{I}_B) = 0$, it is equivalent by Serre duality to

show that $\mathrm{Hom}_{C^{(n)}}(F_n^* \mathcal{L}, \mathcal{O}_{C^{(n)}}(B) \otimes \omega_{C^{(n)}}) = 0$. Since the invertible sheaf $\mathcal{O}_{C^{(n)}}(B) \otimes \omega_{C^{(n)}}$ has degree $b + 2g - 2$ and by picking n large enough, we can ensure $p^n \deg \mathcal{L} > b + 2g - 2$ from which we obtain $H^1(C^{(n)}, F_n^* \mathcal{L} \otimes \mathcal{I}_B) = 0$ and hence $H^1(C^{(n)}, F_n^* \mathcal{E} \otimes \mathcal{I}_B) = 0$ for a locally free sheaf of any rank.

To show that $F_n^* \mathcal{E} \otimes \mathcal{I}_B$ is globally generated, pick a point $q \in C$. Then $\mathcal{I}_B \otimes \mathcal{I}_q$ has length $b + 1$ and from the discussion above $H^1(C^{(n)}, F_n^* \mathcal{E} \otimes \mathcal{I}_B \otimes \mathcal{I}_q)$ vanishes when $p^n \deg L > b + 1 + 2g - 2$ so we can just pick n large enough to fit this condition. Now, by taking the long exact sequence in cohomology of

$$0 \rightarrow F_n^* \mathcal{E} \otimes \mathcal{I}_B \otimes \mathcal{I}_q \rightarrow F_n^* \mathcal{E} \otimes \mathcal{I}_B \rightarrow (F_n^* \mathcal{E} \otimes \mathcal{I}_B) \otimes k(q) \rightarrow 0$$

we conclude that $F_n^* \mathcal{E} \otimes \mathcal{I}_B$ is globally generated.

That \mathcal{E} can not be written as an extension of \mathcal{M} an ample line bundle and a quotient locally free sheaf \mathcal{Q} is equivalent to $H^0(C, \mathcal{E} \otimes \mathcal{M}^{-1}) = 0$. However there exists a positive integer m and an ample line bundle $\mathcal{M}_{C^{(m)}}$ on $C^{(m)}$ for which $H^0(C^{(m)}, (F_m^* \mathcal{E}) \otimes \mathcal{M}_{C^{(m)}}^{-1}) \neq 0$ and we proceed as before with the sheaf $(F_m^* \mathcal{E})$. \square

3. DEFINITION OF CURVE CONNECTEDNESS: COVERING FAMILIES

We now define various ways in which a variety can be covered by curves, generalising the notion of a rationally connected varieties (see [Kol96, IV]).

DEFINITION 3.1. We say that a variety X over a field k is *connected by genus $g \geq 0$ curves* (resp. *chain connected by genus g curves*) if there exists a proper flat morphism $\mathcal{C} \rightarrow Y$, for a variety Y , whose geometric fibres are irreducible genus g curves (resp. connected genus g curves) such that there is a morphism $u : \mathcal{C} \rightarrow X$ making the induced morphism $u^{(2)} : \mathcal{C} \times_Y \mathcal{C} \rightarrow X \times_k X$ dominant.

We say X is separably (chain) connected by genus g curves if $u^{(2)}$ is smooth at the generic point. Note that the notion of separability is redundant in characteristic zero due to generic smoothness. A genus zero connected variety is rationally connected. A variety which is connected by genus one curves will be called (with a slight abuse of notation) elliptically connected. The relevant moduli spaces which we will be considering are the following. Let $\pi : \mathcal{C} \rightarrow S$ be a flat projective curve over an irreducible scheme S and let $B \subset \mathcal{C}$ be a closed subscheme that is flat and finite over S . Let $p : X \rightarrow S$ be a smooth quasi-projective scheme and $g : B \rightarrow X$ an S -morphism. The space (see [Kol96, II.1.5] and [Mor79]) $\mathrm{Hom}_S(\mathcal{C}, X, g)$ parametrises S -morphisms from \mathcal{C} to X keeping the points given by g fixed. Restricting to the case where S is the spectrum of an algebraically closed field k we fix some notation of the following evaluation morphisms to be used in later sections

$$\begin{aligned} F : \mathcal{C} \times \mathrm{Hom}(\mathcal{C}, X, g) &\rightarrow X \\ \phi(p, f) : H^0(\mathcal{C}, f^* \mathcal{T}_X \otimes \mathcal{I}_B) &\rightarrow f^* \mathcal{T}_X \otimes k(p) \end{aligned}$$

and similarly the double evaluation morphisms $F^{(2)}$ and $\phi^{(2)}(p, q, f)$ as in [Kol96, II.3.3]. Secondly we consider the relative moduli space of genus g degree d stable curves with base point $t : P \rightarrow X$, denoted by $\overline{\mathcal{M}}_g(X/S, d, t)$ as in [AK03] (originally [FP97]). By Bertini, we can always find a genus g such that a projective X is genus g connected, the minimal such g however is an interesting invariant of the variety. Finding higher genus covering families is an easy operation.

LEMMA 3.2. *Let X be a genus g (chain) connected smooth projective variety over an algebraically closed field k . Then if $g' \geq 2g - 1$, X is also genus g' (chain) connected.*

PROOF. Let $\mathcal{C}/Y \rightarrow X$ be a family making X a genus g (chain) connected variety. From [AK03, Theorem 50] we have a projective algebraic space $Y' = \overline{\mathcal{M}}'_g(\mathcal{C}/Y, d)$ of finite type over Y parametrising stable families of degree d curves of genus g' over $\mathcal{C} \rightarrow Y$. The condition $g' \geq 2g - 1$ coming from the Riemann-Hurwitz formula ensures that this moduli space is non-empty. From [ACG11, 12.9.2] there exists a normal scheme Z finite and surjective over Y' and a flat and proper family $\mathcal{X} \rightarrow Z$ of stable genus g curves of degree d . Restricting to a suitable open subset $W \subset Z$ parametrising irreducible curves we compose the family $\mathcal{X}|_W \rightarrow W$ with the evaluation morphism to X and the result follows. \square

An example of an elliptically connected variety over a non-algebraically closed field is given after the proof of Theorem 6.2. A much stronger condition is the existence of a family of curves which is constant in moduli.

DEFINITION 3.3. We say that a variety X over a field k is C -connected for a curve C if there exists a variety Y and a map $u : C \times Y \rightarrow X$ such that the induced map $u^{(2)} : C \times C \times Y \rightarrow X \times X$ is dominant. If $u^{(2)}$ is also smooth at the generic point, then we say that X is *separably C -connected*.

Projective space is C -connected for every smooth projective curve C whereas an example of a C -connected variety which is not rationally connected is $C \times \mathbb{P}^n$ where $g(C) \geq 1$. To see this let $(c_1, x_1), (c_2, x_2)$ be any two points in $C \times \mathbb{P}^n$ and let $f : C \rightarrow \mathbb{P}^n$ a morphism which sends $c_i \mapsto x_i$. Considering the graph of f in $C \times \mathbb{P}^n$ we have found a curve isomorphic to C which goes through our two points. Using parts (3) and (4) from Lemma 3.4 below, the result follows. More generally, examples can also be constructed from Proposition 3.5 below. The following are mostly straight forward generalisations of various results in [Kol96, IV.3].

LEMMA 3.4. *The following statements hold for a variety X over a field k and C a smooth projective curve.*

- (1) *If X is genus g connected and $X \dashrightarrow Y$ a dominant rational map to a proper variety Y , then Y is also genus g connected. The same holds if X is C -connected.*

- (2) A variety X is C -connected if and only if there is a variety W , closed in $\mathrm{Hom}(C, X)$ such that $u^{(2)} : C \times C \times W \rightarrow X \times X$ is dominant.
- (3) If X is defined over a field k and K/k is an extension of fields, then $X_K := X \times_k K$ is C -connected if and only if X_k is.
- (4) A variety X over an uncountable algebraically closed field is C -connected if and only if for all very general $x_1, x_2 \in X$ there exists a morphism $C \rightarrow X$ which passes through x_1, x_2 .
- (5) A variety X over an uncountable algebraically closed field is genus g connected if and only if for all very general $x_1, x_2 \in X$ there exists a smooth irreducible genus g curve containing them.
- (6) Being rationally or elliptically connected is closed under connected finite étale covers of varieties.

PROOF. To prove (1), let $u : \mathcal{C}/M \rightarrow X$ be a family making X genus g connected and denote by $u' : \mathcal{C}/M \dashrightarrow Y$ the composition. Restricting u' to the generic fibre $\mathcal{C}_{k(M)}$ we have a rational map $\phi : \mathcal{C}_{k(M)} \dashrightarrow Y$. Since Y is proper, by the valuative criterion of properness we can extend ϕ to a morphism $\phi : \mathcal{C}_{k(M)} \rightarrow Y$. By spreading out to an open subset $M' \subseteq M$ (see [DG67, IV₃ 8.10.5] for properness and 11.2.6 for flatness of the family) we obtain a family $\mathcal{C}|_{M'} \rightarrow M'$ which makes Y also genus g connected.

Since being C -connected or connected by genus g curves is a birational property, we may assume by compactifying that X is projective. For (2), consider $\mathrm{Hom}(C, X) = \cup R_i$ the decomposition into irreducible components. One direction of the statement is obvious, whereas for the other let $C \times W \rightarrow W$ be a family which makes X a C -connected variety. If $u_i : C \times R_i \rightarrow X$ is the evaluation morphism, then for some i there is a morphism $h : W \rightarrow R_i$ such that $h(w) = [C_w \rightarrow X]$ for general $w \in W$. This implies that $u_i^{(2)} : C \times C \times R_i \rightarrow X \times X$ is also dominant. For one direction of (3), pullback by $\mathrm{Spec} K \rightarrow \mathrm{Spec} k$. For the other, if X_K is C_K -connected then from (2) there is a positive integer d such that the evaluation morphism $\mathrm{ev}_K^d : C_K \times C_K \times \mathrm{Hom}_d(C_K, X_K) \rightarrow X_K \times X_K$ is dominant. Because of the universal property of the Hom-scheme, we have that $\mathrm{Hom}(C, X) \times_k K = \mathrm{Hom}(C_K, X_K)$ and $(\mathrm{ev}^d)_K = \mathrm{ev}_K^d$ so ev^d is also dominant.

If through every two very general points there passes the image of C under some morphism, then the map $u^{(2)} : C \times C \times \mathrm{Hom}(C, X) \rightarrow X \times X$ is dominant. Since $\mathrm{Hom}(C, X)$ has at most countably many irreducible components the restriction of $u^{(2)}$ to at least one of the components R_i must be dominant, which proves (4). Similarly for (5) working instead with the Kontsevich moduli of curves $\mathcal{M}_{g,1}(X) \rightarrow \mathcal{M}_{g,0}(X)$ the result follows. For (6), the proof for rationally connected varieties is contained in [Deb01, 4.4.(5)]. Let $\mathcal{C} \rightarrow U$ be a family which makes X elliptically connected and let $X' \rightarrow X$ be a connected finite étale cover. Consider the pullback diagram and $\mathcal{C}' \rightarrow U' \rightarrow U$ the Stein

factorisation

$$\begin{array}{ccccc}
 & & \mathcal{C}' = \mathcal{C} \times_X X' & \longrightarrow & X' \\
 & \swarrow & \downarrow & & \downarrow \\
 & & \mathcal{C} & \longrightarrow & X \\
 & \swarrow & \downarrow & & \\
 U' & & U & &
 \end{array}$$

After possibly restricting U' to the open subset of curves in \mathcal{C}' which are irreducible, the family $\mathcal{C}' \rightarrow U'$ makes X' elliptically connected. \square

PROPOSITION 3.5. *Let X be a smooth projective variety over an algebraically closed field k and $f : X \rightarrow C$ a flat morphism to a smooth projective curve whose geometric generic fibre is separably rationally connected. Then X is C -connected.*

PROOF. From [dJS03], there is a section $\sigma : C \rightarrow X$ to f . Now from [KMM92, Theorem 2.13] we can find a section to f passing through any two points in different smooth fibres over C , hence we can find a copy of C passing through two general points. The result now follows from Lemma 3.4 parts (4) and (5) above after possibly passing to an uncountable extension K/k . \square

We now come to the main theme of this paper, which is that varieties covered by higher genus curves in a strong sense are also covered by rational curves. This is illustrated in the following proposition, and continues in the next sections.

PROPOSITION 3.6. *Let X be a C -connected variety of dimension at least 3 over an algebraically closed field k . Then X is uniruled.*

PROOF. We may assume X is projective. Let $u : C \times Y \rightarrow X$ be a family such that $u^{(2)} : C \times C \times Y \rightarrow X \times X$ is dominant. We have $\dim Y + 2 \geq 2 \dim X$ and so if $\dim X \geq 3$ we obtain $\dim Y \geq 4$. Now, pick general points $x \in X, c \in C$ and denote by $Z \subset Y$ the locus of curves $u_z : C_z \rightarrow X$ such that $x = u_z(c)$ for all $z \in Z$. We have that $\dim Z \geq \dim Y - (\dim X - 1) - 1$ and so for $\dim X \geq 3, \dim Z \geq 1$. Since any two general points in X can be connected by the image of a C_y , it follows that Z does not get contracted to a point when mapped to $\text{Hom}(C, X; c \mapsto x)$. From Bend and Break (see [Deb01, Prop. 3.1]) we obtain a rational curve through x and hence through every general point. After possibly an extension to an uncountable algebraically closed field this implies that X is uniruled (see [Deb01, Remark 4.2(5)]). \square

If C has genus one, the above result is also proved in Section 6, even allowing the curve C to vary in moduli and with the dimension of X assumed greater or equal to two. On the other hand, a C -connected surface does not have to be uniruled when C has genus at least two. Consider $C \subset A$ a curve in an abelian

surface such that C contains the identity 0 of A and the genus of C is at least two. Consider the map $\phi : C \times C \rightarrow X$ sending (p, q) to $p - q$. If the image is one dimensional, it has to be isomorphic to C since it has to be irreducible and contains the image of $C \times \{0\}$. On the other hand, the image will be closed under the group operation, hence would have to be abelian itself, which is a contradiction. Hence ϕ is surjective, and we obtain that for any $x \in A$, there is a $(p, q) \mapsto x$, hence a morphism $C \cong C \times \{q\} \rightarrow X$ passing through x and 0 (for (q, q)). Take any two points $x, y \in A$, and consider the image of a morphism from C through 0 and the point $x - y$ that we just constructed. Translate this curve by y and obtain an image of C through x, y .

Denoting by $X \dashrightarrow R(X)$ the maximal rationally chain connected (MRC) fibration, we let $R^0(X) = X$, $R^i(X) = R(R^{i-1}X)$ and obtain a tower of MRC fibrations

$$X \dashrightarrow R^1(X) \dashrightarrow \cdots \dashrightarrow R^n(X).$$

This tower eventually stabilises, and if $R^i(X)$ is uniruled then $\dim R^{i+1}(X) < \dim R^i(X)$. In characteristic zero, we in fact have $R(X) = \cdots = R^n(X)$ (see discussion below). In positive characteristic it can be that the tower has length greater than one - see the example given in the last section of this paper.

PROPOSITION 3.7. *Let X be a normal and proper C -connected variety over an algebraically closed field where C is a smooth projective curve. Then the tower $X \dashrightarrow R^1(X) \dashrightarrow \cdots \dashrightarrow R^n(X)$ of MRC quotients terminates in either a point, a curve or a surface.*

PROOF. Let $C \times Y \rightarrow X$ be a family which makes X a C -connected variety. From Lemma 3.4 part (1) it follows that $R^i(X)$ are also C -connected. From Proposition 3.6 we obtain that $R^i(X)$ is uniruled if $\dim R^i(X) \geq 3$. This implies that $R^{i+1}(X)$ must have dimension strictly less than $R^i(X)$ and so the result follows. \square

Note that if k is algebraically closed of characteristic zero then we know from [GHS03] that the MRC quotient $R(X)$ is not uniruled, so if X is C -connected of dimension at least three, $R(X)$ must be a surface, curve or point, in which case X is respectively a rationally connected fibration over a surface or curve, or a point (and so X is rationally connected). From Proposition 3.5 the converse holds too for a fibration over a curve.

REMARK 3.8. As observed in [Occ06, Remark 4], if the MRC quotient of a smooth complex projective variety X is a curve, then the MRC fibration extends to the whole variety and coincides with the Albanese map.

4. DEFINITION OF CURVE CONNECTEDNESS: FREE MORPHISMS

In this section we define ways in which a morphism from a curve C to a variety X can deform enough to give a large family of morphisms from C so as to cover X . A notion studied extensively by Hartshorne [Har70] is that of a (local

complete intersection) subvariety Y in a smooth projective variety X such that the normal bundle $\mathcal{N}_{Y/X}$ is ample. Hartshorne proved in [Har70, III.4] that for some $g \geq 0$ there exists a curve $C \subset X$ of genus g such that $\mathcal{N}_{C/X}$ is ample. Alternatively, Ottem [Ott12] defines an ample closed subscheme $Y \subset X$ of codimension r to be one where the exceptional divisor $\mathcal{O}(E)$ of the blowup $\text{Bl}_Y X$ of X along Y is an $(r - 1)$ -ample line bundle in the sense that for every coherent sheaf \mathcal{F} there is an integer $m_0 > 0$ such that $H^i(X, \mathcal{F} \otimes \mathcal{O}(E)^m) = 0$ for all $m > m_0$ and $i > r - 1$. One can then prove that if Y is a local complete intersection subscheme of X which is ample, then the normal bundle $\mathcal{N}_{Y/X}$ is an ample bundle. We impose the following stronger positivity condition.

DEFINITION 4.1. ([Kol96, II.3.1]) Let C be a smooth proper curve and X a smooth variety over a field k . Let $f : C \rightarrow X$ a morphism and $B \subset C$ a closed subscheme with ideal sheaf \mathcal{I}_B and $g = f|_B$. The morphism f is called *free over g* if it is non-constant and one of the following two equivalent conditions is satisfied:

- (1) for every $p \in C$ we have $H^1(C, f^* \mathcal{I}_X \otimes \mathcal{I}_B(-p)) = 0$ or,
- (2) $H^1(C, f^* \mathcal{I}_X \otimes \mathcal{I}_B) = 0$ and $f^* \mathcal{I}_X \otimes \mathcal{I}_B$ is generated by global sections.

Note that there is also a relative version of the above definition discussed in [KSCT07].

DEFINITION 4.2. We say that a curve $f : C \rightarrow X$ is *r-free* if for all effective divisors D of degree $r \geq 0$, $H^1(C, f^* \mathcal{I}_X \otimes \mathcal{O}_C(-D)) = 0$ and $f^* \mathcal{I}_X \otimes \mathcal{O}_C(-D)$ is generated by global sections. A 0-free curve is called *free* whereas a 1-free curve is called *very free*.

The condition of *r-freeness* makes formal the notion that the curve C deforms in X while keeping any general r points fixed. The following follows immediately from Lemma 2.3.

LEMMA 4.3. *If $f : C \rightarrow X$ is an r -free curve then f is r' -free for all $r' \leq r$.*

In the case of $C = \mathbb{P}^1$, $f^* \mathcal{I}_X = \oplus_{i=1}^n \mathcal{O}_{\mathbb{P}^1}(a_i)$ with $a_1 \leq \dots \leq a_n$ so it follows that $f : \mathbb{P}^1 \rightarrow X$ is r -free if and only if $a_1 \geq r$.

REMARK 4.4. We should remark at this point that there do not exist complete intersection curves of large enough degree which are free on a general smooth hypersurface. For example, let X be a degree d smooth hypersurface in \mathbb{P}^n . Assume $d \leq n$ since otherwise X will be of general type or Calabi-Yau and will not have any free curves. Let Y_i be $n - 2$ suitably general hypersurfaces in \mathbb{P}^n all of degree e and let $C = X \cap_{i=1}^{n-2} Y_i$ be the resulting curve. The degree of C is de^{n-2} and the normal bundle is

$$\mathcal{N}_{C/X} = \oplus_{i=1}^{n-2} \mathcal{O}_{\mathbb{P}^n}(Y_i)|_C = \oplus_{i=1}^{n-2} \mathcal{O}_{\mathbb{P}^n}(e)|_C.$$

By adjunction, we compute

$$\text{deg } \mathcal{I}_C = -\text{deg } \omega_C = -d(-n - 1 + d + \sum_{i=1}^{n-2} e).$$

Even setting $e = 1$ to make $\deg \mathcal{T}_C$ as large as possible, and taking into account that $\deg \mathcal{N}_{C/X} = e(n - 2)$, we see that $\deg \mathcal{T}_X|_C = \deg \mathcal{T}_C + \deg \mathcal{N}_{C/X}$ is not going to be positive for large values of d and n . Positivity of the degree of $\mathcal{T}_X|_C$ would be necessary for any ampleness conditions. See [Gou14] for a discussion on separable rational connectedness of Fano complete intersections.

A result of Kollár ([Kol96, II.1.8]) implies that if the dimension of X is at least 3, a general deformation of a 2-free morphism is an embedding into X . We will see (Theorem 5.5) that if the genus of C is at least one, this holds for any free morphism too. From [Kol96, II.3.2], if a family of curves mapping to a variety has a member which is free over g , then the locus of all such curves in this family is open.

LEMMA 4.5. *Let X be a smooth variety over an algebraically closed field k , $D \subset X$ a divisor and $f : C \rightarrow X$ a free morphism. If $p \in C$ then there exists a deformation $f' : C \rightarrow X$ with $f'(p) \notin D$.*

PROOF. By semicontinuity let $U \subset \text{Hom}(C, X)$ be a connected open neighbourhood of $[f]$ such that $H^1(C, f_t^* \mathcal{T}_X) = 0$ for all $[f_t] \in U$. From [Mor79] it follows that the dimension of U is $h^0(C, f^* \mathcal{T}_X)$. Denote by \mathcal{I}_p the ideal sheaf on C of the closed subscheme with unique point p . Since f is free, we have $H^1(C, f_t^* \mathcal{T}_X \otimes \mathcal{I}_p) = 0$ for all $[f_t] \in U$ and so by fixing a point $x \in X$ such that $p \mapsto x$, we have

$$\begin{aligned} \dim(\text{Hom}(C, X; p \mapsto x) \cap U) &= h^0(C, f^* \mathcal{T}_X \otimes \mathcal{I}_p) \\ &= h^0(C, f^* \mathcal{T}_X) - \dim X \\ &= \dim U - \dim X. \end{aligned}$$

Next, denote by

$$V = \{[f_t] \in U \mid f_t(p) \in D\} = \bigcup_{x \in D} \{[f_t] \in U \mid f_t(p) = x\}$$

the subspace of all morphisms in U which send p to a point in the divisor D . It follows that

$$\begin{aligned} \text{codim}(V, U) &\geq \dim U - \dim V \\ &= h^0(C, f^* \mathcal{T}_X) - (h^0(C, f^* \mathcal{T}_X) - \dim X + \dim X - 1) = 1 \end{aligned}$$

and hence there exists an $[f'] \in U \setminus V$ such that $f'(p) \notin D$. □

PROPOSITION 4.6. *Let X be a smooth variety over an algebraically closed field k and $f : C \rightarrow X$ a smooth projective curve which is free over $B \subset C$ a closed subscheme with ideal sheaf \mathcal{I}_B . Let $g : X \dashrightarrow Y$ be a generically smooth dominant rational map to a smooth proper variety Y . Then it follows that $f' := g \circ f : C \dashrightarrow Y$ can be deformed to a morphism free over B .*

PROOF. Deform $f : C \rightarrow X$ so that it misses the codimension 2 exceptional locus of g (from [Kol96, II.3.7]) so we can assume that the composition $g \circ f : C \dashrightarrow Y$ is in fact a non-constant morphism. Starting with the standard exact

sequence of tangent bundles on X and applying f^* and tensoring with \mathcal{I}_B we obtain

$$(4.1) \quad 0 \rightarrow f^* \mathcal{T}_{X/Y} \otimes \mathcal{I}_B \rightarrow f^* \mathcal{T}_X \otimes \mathcal{I}_B \rightarrow (g \circ f)^* \mathcal{T}_Y \otimes \mathcal{I}_B.$$

From [Liu02, Ex. 6.2.10] this is exact on the right and we conclude. \square

In the case of higher genus curves there exist genus g connected varieties which do not have a free or very free curve for all $g \geq 1$, for example consider $E \times \mathbb{P}^1$ where E is an elliptic curve. As pointed out after Definition 3.3, $E \times \mathbb{P}^1$ is E -connected yet it is not possible that there exists a morphism $f : C \rightarrow E \times \mathbb{P}^1$ from a curve C such that $f^* \mathcal{T}_{E \times \mathbb{P}^1}$ is ample since this bundle is isomorphic to $\mathcal{O}_C \oplus \mathcal{O}_C(2)$ which has a non-ample quotient \mathcal{O}_C . One can however prove the following proposition.

PROPOSITION 4.7. *Let X be a smooth variety over an algebraically closed field and $f : C \rightarrow X$ a very free morphism for some smooth projective curve C . Then X is separably C -connected.*

PROOF. Let $[f] \in Y \subset \text{Hom}(C, X)$ be an open and smooth neighbourhood with cycle map $u : C \times Y \rightarrow X$. We first show that the evaluation map

$$\phi^{(2)}(p, q, f) : H^0(C, f^* \mathcal{T}_X) \rightarrow f^* \mathcal{T}_X \otimes k(p) \oplus f^* \mathcal{T}_X \otimes k(q)$$

is surjective for $p \neq q$ general points in C . Consider the following exact sequences of sheaves

$$\begin{aligned} 0 \rightarrow f^* \mathcal{T}_X(-p-q) \rightarrow f^* \mathcal{T}_X \rightarrow (f^* \mathcal{T}_X \otimes k(p)) \oplus (f^* \mathcal{T}_X \otimes k(q)) \rightarrow 0 \\ 0 \rightarrow f^* \mathcal{T}_X(-p-q) \rightarrow f^* \mathcal{T}_X(-p) \rightarrow f^* \mathcal{T}_X(-p) \otimes k(q) \rightarrow 0 \end{aligned}$$

and note that by taking the long exact sequence in cohomology of the first, to show that $\phi^{(2)}(p, q, f)$ is surjective, we need to show that $H^1(C, f^* \mathcal{T}_X(-p-q)) = 0$. Since f is very free we have from the second sequence that $H^0(C, f^* \mathcal{T}_X(-p)) \rightarrow f^* \mathcal{T}_X(-p) \otimes k(q)$ is surjective and also that $H^1(C, f^* \mathcal{T}_X(-p)) = 0$ from which it follows that $H^1(C, f^* \mathcal{T}_X(-p-q)) = 0$. Since $\phi^{(2)}(p, q, f)$ is surjective, it follows from [Kol96, II.3.5] that $u^{(2)} : C \times C \times Y \rightarrow X \times X$ is smooth at $(p, q, [f])$. We conclude that X is separably C -connected and thus also separably connected by genus g curves. \square

REMARK 4.8. It follows that in the setting above that a very free curve (or in fact even a C such that X is C -connected) has the property that it intersects non-trivially all but a finite number of divisors. This follows from the fact that we can cover an open subset by images of C , whose complement will be a proper closed subset of X and so contains a finite number of divisors.

5. PROVING UNIRULEDNESS AND RATIONAL CONNECTEDNESS

In this section we prove that the existence of a free curve of genus $g \geq 1$ is equivalent to the existence of an r -free curve of genus g for all $r \geq 1$, and that in

characteristic zero this is also equivalent to the existence of a very free rational curve. This is in stark contrast to rational curves, where uniruled varieties (possessing free rational curves) are not always rationally connected (possessing very free rational curves). We begin by noting that there is another type of positive curve one can consider for a smooth projective variety X , namely $f : C \rightarrow X$ such that $f^* \mathcal{T}_X$ is ample. Note that such a curve automatically has $\mathcal{N}_{C/X}$ ample. Such curves have traditionally been studied in terms of foliations (cf. Theorem 5.3). We will also prove that the existence of a curve such that $f^* \mathcal{T}_X$ is ample is in fact equivalent to the existence of a free curve of the same genus.

PROPOSITION 5.1. *Let X be a smooth projective variety over an algebraically closed field k and $f : C \rightarrow X$ a morphism from a smooth projective curve of genus g such that $f^* \mathcal{T}_X$ ample. Then X is uniruled.*

PROOF. The proof follows the usual Mori argument so we present only a sketch (cf. Theorem 5.3). Note that if X is a curve, then since a bundle is ample if and only if its pullback under a finite morphism is ample, we obtain that $X = \mathbb{P}^1$. In characteristic zero, after spreading out over a finitely generated extension $\text{Spec } S$ of $\text{Spec } \mathbb{Z}$, one can reduce to any closed prime and consider the equivalent set-up in positive characteristic. After pulling back by Frobenius, Lemma 2.4 implies that there is a morphism $f_p^{(n)} : C_p \rightarrow X_p$ such that $(f_p^{(n)})^* \mathcal{T}_{X_p}$ is very free (or r -free even), where $f_p : C_p \rightarrow X_p$ the reduction of $f : C \rightarrow X$. Bend and Break now produces a rational curve passing through a general point, of bounded degree independent of p (see [Deb01, Prop. 3.5]). These are points in fibres over $\text{Spec } S$ of a finite type relative moduli $\text{Hom}_S^d(\mathbb{P}_S^1, \mathcal{X}/S, s)$, for $s : \text{Spec } S \rightarrow \mathcal{X}$ a section specifying the general point the rational curve goes through. Hence by Chevalley's Theorem the generic fibre over $\text{Spec } S$ is also non-empty, and there is a rational curve through a general point of X . \square

THEOREM 5.2. *Let X be a smooth projective variety over an algebraically closed field k and $f : C \rightarrow X$ a morphism from a smooth projective curve of genus g such that $f^* \mathcal{T}_X$ is ample.*

- (1) *If the characteristic p of k is zero, then X is rationally connected.*
- (2) *If $p > 0$ then the tower of MRC fibrations terminates with a point.*

PROOF. From 5.1, we conclude that X is uniruled, regardless of the characteristic. Denote by $\pi : X \rightarrow R(X)$ the MRC fibration ($R(X)$ is defined up to birational transformation so we may assume π is a morphism). In characteristic zero, the composition $g : C \rightarrow X \rightarrow R(X)$ again has $g^* \mathcal{T}_{R(X)}$ ample, since from the proof of 4.6 the quotient of an ample bundle is ample. So by the Graber-Harris-Starr Theorem, since $R(X)$ is uniruled by Proposition 5.1, it must be a point. In positive characteristic, it may not be the case that the composition $g : X \rightarrow R(X)$ is generically smooth, in which case $g^* \mathcal{T}_{R(X)}$ might not be ample. From Lemma 2.4 however there is a morphism $h : C' \rightarrow X$ such that $h^* \mathcal{T}_X$ is very free (here C' is a Frobenius pullback of C so has the same

genus). From 4.7, X is separably C' -connected, and so by 3.7 we do obtain that the tower of MRC quotients $X \rightarrow R(X) \rightarrow \cdots R^n(X)$ ends in a point, curve or surface. If $\pi : X \rightarrow T$ where $T := R^n(X)$ is a smooth projective curve, then by Lemma 4.5, for a point $p \in C'$, we can deform h so that the image of p misses the inverse image under π of $\pi(h(p))$. Hence $\text{Hom}(C', T)$ is at least one dimensional and from de Franchis' Theorem [ACG11, 8.27] it follows that T has genus zero or one. One excludes the case where C', T both of genus one, by using the fact that there are only countably many isogenies between two elliptic curves. Also, T cannot be rational since we have assumed the tower is maximal. If now $R^n(X) = S$ is a smooth projective surface, we may assume by pulling back by Frobenius from 2.4 and deforming, that there is an at least one dimensional family of morphisms sending a fixed point on C to a fixed point on S . Hence by Bend and Break [Deb01, Prop. 3.1] the surface would have to be uniruled and we are reduced to the case of a point again. \square

Assuming ampleness and regularity of a foliation on a smooth curve in characteristic zero, results of this type have been demonstrated in the work of various people, starting with Miyaoka's uniruledness criterion [Miy87, Theorem 8.5]. A short summary of recent results follows.

THEOREM 5.3. ([BM01, Theorem 0.1], [KSCT07, Theorem 1]) *Let X be a normal complex projective variety and $C \subset X$ a complete curve in the smooth locus of X . Assume that $\mathcal{F} \subset \mathcal{T}_X$ is a foliation regular along C and such that $\mathcal{F}|_C$ is ample. If $x \in C$ is any point, the leaf through x is algebraic and if $x \in C$ is general then the closure of the leaf is also rationally connected.*

Using [BDPP13, Corollary 0.3], Peternell proved a weaker version of Mumford's conjecture on numerical characterisation of rationally connected varieties from which one can deduce the following theorem.

THEOREM 5.4. ([Pet06, 5.4, 5.5]) *Let X/\mathbb{C} be a projective manifold and $C \subset X$ a possibly singular curve. If $\mathcal{T}_X|_C$ is ample then X is rationally connected. If $\mathcal{T}_X|_C$ is nef and $-K_X.C > 0$ then X is uniruled.*

The precise relation between r -free morphisms and morphisms $f : C \rightarrow X$ such that $f^*\mathcal{T}_X$ is ample is given in the following.

THEOREM 5.5. *Let X be a smooth projective variety over an algebraically closed field k and $r \geq 0$ any integer. Then there exists a morphism $f : C \rightarrow X$ from a smooth projective genus $g \geq 1$ curve C such that $f^*\mathcal{T}_X$ is ample if and only if there is an r -free morphism $h : C' \rightarrow X$ from a genus g smooth projective curve C' .*

PROOF. Assuming the existence of h , we obtain from Lemma 2.3 that h is also free, and so by Proposition 2.2, $h^*\mathcal{T}_X$ is ample. If $f^*\mathcal{T}_X$ is ample, one needs to separate between characteristic $p > 0$ or equal to zero. In the former case, as in the proof of 5.1 we get $h : C' \rightarrow X$ (here again C' is a Frobenius

pullback of C so of genus g) which is r -free. When the characteristic is zero, X will be rationally connected from 5.2. The idea now is to attach many very free rational curves to C , apply standard smoothing of combs techniques and prove that the resulting general smooth deformations of the comb will be r -free genus g curves (cf [Kol96, II.7.10]). This proceeds as follows. Assemble a comb $D = C \cup \cup_{i=1}^m C_i$ with m rational teeth that are $(r+1)$ -free like in [Kol96, II.7]. For m large enough, D is smoothable to a flat proper family $Y \rightarrow T$ where the general fibre is isomorphic to C , the central fibre is a subcomb of D with a large number of teeth depending on $C \subset X$ and m , and there is a morphism $F : Y \rightarrow X$ which extends $D \rightarrow X$. To show that the general nearby fibre $f_t : Y_t \rightarrow X$ is r -free, it suffices to show that $H^1(Y_t, f_t^* \mathcal{F}_X(-\sum_{i=0}^r p_i))$ for p_0, p_1, \dots, p_r any points on $Y_t \subset X$ (see Definition 4.1). Pick sections $s_0, s_1, \dots, s_r : T \rightarrow Y$ with $s_i(t) = p_i$. Let $E = F^* \mathcal{F}_X(-\sum_{i=1}^r s_i(T))$. By Riemann-Roch, for m large enough, we have that $H^1(C, M \otimes E|_C) = 0$ for all line bundles M of degree larger than m , and also that $E|_{C_i}$ is ample since C_i is $(r+1)$ -free. Now apply [Kol96, II.7.10.1] for m large enough. \square

Using any of Theorems 5.3, 5.4 or 5.2, a smooth projective variety X over an algebraically closed field of characteristic zero with a free genus g curve $f : C \rightarrow X$ such that $g \geq 1$ is automatically rationally connected.

REMARK 5.6. At this point we cannot prove that in positive characteristic, assuming that we have a free curve $f : C \rightarrow X$ of genus $g \geq 1$ implies that X is separably rationally connected or even rationally chain connected. It is tempting to hope that both statements are true though. Jason Starr informs us that his maximal free rational quotient (MFRC) [Sta06] gives a generically (on the source) smooth morphism $X \rightarrow R_f(X)$ over any algebraically closed field k , so if X contained a free rational curve $f : \mathbb{P}^1 \rightarrow X$, then $\dim R_f(X) < \dim X$. Hence, if $f : C \rightarrow X$ a free curve of genus $g \geq 1$ implied that we have a free rational curve $\mathbb{P}^1 \rightarrow X$ (we do not know how to show this), taking successive MFRC quotients and using Proposition 4.6 would reduce the tower of MFRC quotients to a point. This does not mean that X will necessarily be rationally connected, but since there is a free rational curve on X , it will at least be separably uniruled. Even though Bend and Break arguments give us the existence of many rational curves, the author does not know any general techniques to construct free rational curves in positive characteristic. See the last two sections for results in this direction.

6. ELLIPTICALLY CONNECTED VARIETIES

In this section we will study more carefully the case of genus one. Denoting RC and EC to mean rationally and elliptically connected (genus one connected) respectively, we have the following inclusions of sets of varieties

$$\{\text{rational}\} \subsetneq \{\text{unirational}\} \subsetneq \{\text{RC}\} \subsetneq \{\text{EC}\} \subsetneq \{\text{uniruled}\}.$$

It is an open problem whether there exists a non-unirational rationally connected variety but it is widely expected these do exist. The following result

is in the spirit of 3.6. The following proof was suggested by the anonymous referee.

PROPOSITION 6.1. *Let X be an elliptically connected smooth projective variety of $\dim X \geq 2$ over an algebraically closed field k . Then X is uniruled.*

PROOF. Like in 3.6, for $\mathcal{C} \rightarrow U$ a family of genus one curves mapping to X such that $\mathcal{C} \times_U \mathcal{C} \rightarrow X \times X$ is dominant, there is an at least one dimensional locus $Z \subset U$ parametrising curves which pass through a (general) point $x \in X$. In fact, after fixing a general hyperplane H , we obtain a morphism $Z \rightarrow \mathcal{M}_{1,2}(X)$ where for $z \in Z$, the two marked points are the point $p_z \in \mathcal{C}_z$ sent to x , and a point $q_z \in \mathcal{C}_z$ which is sent to H . Denote also by $\mathcal{C} \rightarrow Z$ the restriction of the family from U . Consider now a compactification and the induced rational map to X

$$\begin{array}{c} \bar{\mathcal{C}} \dashrightarrow X \\ \downarrow \pi \\ \bar{Z} \end{array}$$

and let $\mu : \bar{Z} \rightarrow \bar{\mathcal{M}}_{1,2}(X)$ be the moduli map. Since $\mathcal{M}_{1,2}$ contains no proper subvarieties which do not get contracted when mapped to \mathcal{M}_1 , either the image of μ meets the boundary, which implies that there is a rational curve through x , or μ is a contraction to a point. In the latter case, we thus have that the family π is isotrivial, so after passing to a finite flat cover \bar{Z}' of \bar{Z} we obtain $C \times \bar{Z}' \rightarrow \bar{Z}'$, with $f' : C \times \bar{Z}' \dashrightarrow X$ the induced morphism. From the construction, we also obtain a point $p \in C$ (mapped to each p_z under the map $C \times \bar{Z}' \rightarrow \bar{\mathcal{C}}$) such that f' contracts $\{p\} \times \bar{Z}'$ to x . If f' were defined everywhere, Mumford's Rigidity Theorem would imply that all fibres $\{s\} \times \bar{Z}'$ are contracted, which contradicts the fact that images of our initial family dominate H . Hence f' is not defined everywhere and like in Bend and Break, we obtain a rational curve through x . \square

THEOREM 6.2. *Let X be a smooth projective variety over an algebraically closed field k of characteristic zero. Then X is elliptically connected if and only if it is rationally connected or a rationally connected fibration over an elliptic curve.*

PROOF. Consider the MRC fibration $\pi : X \dashrightarrow R(X)$ where $R(X)$ is elliptically connected as π is dominant. Since $R(X)$ is elliptically connected and not uniruled, it follows from Proposition 6.1 that it must be either of dimension 0 and thus X is rationally connected, or of dimension 1 and so an elliptic curve E by Riemann-Hurwitz. By Remark 3.8, the MRC fibration coincides with the map to the Albanese and so fibres of $X \rightarrow E'$ are rationally connected. Conversely, we have seen that a rationally connected variety is elliptically connected in Lemma 3.2. If on the other hand X is a rationally connected fibration over

an elliptic curve E then from Proposition 3.5 we know that it is E -connected. \square

If k is of positive characteristic, using the same methods as in Lemma 3.7 we deduce that for an elliptically connected variety, the tower of MRC fibrations terminates with a point or a curve.

REMARK 6.3. Note that Bjorn Poonen [Poo10] has constructed non-trivial examples over an arbitrary field, of elliptically connected threefolds which are not rationally connected. These are Châtelet surface fibrations over an elliptic curve.

7. TOWARDS A POSITIVE CHARACTERISTIC ANALOGUE

From Remark 5.6 and the work preceding it, we would like to demonstrate that the existence of a free higher genus curve implies the existence of a free rational curve in positive characteristic, something which holds in characteristic zero from Theorem 5.2. In this section we make the first steps in this direction. If $f : C \rightarrow X$ is a very free morphism from a smooth projective curve of genus $g \geq 2$ to a smooth projective variety X , then $K_X.C = -\deg f^*T_X < 0$ from the ampleness of f^*T_X . In fact, a Riemann-Roch calculation gives a better bound of $K_X.C \leq -n(g-1)$ where $n = \dim X$.

PROPOSITION 7.1. *Let X be a smooth projective surface over an algebraically closed field k with $f : C \rightarrow X$ a free morphism from a smooth projective curve C of genus $g > 0$ or a very free morphism of genus zero. It follows that X is separably rationally connected.*

PROOF. If C is of genus zero then X is separably rationally connected by definition. From the discussion above we have that K_X is not nef. Also, any surface Y which is birational to X admits a morphism $C \rightarrow Y$ from 4.6, which is again free, so K_Y is also not nef. From the classification of surfaces this means that X is either rational or ruled. If ruled, X would admit a birational morphism to $\mathbb{P}^1 \times C$. The free morphism $f : C \rightarrow X$ would give a free morphism $C \rightarrow \mathbb{P}^1 \times C$ which would mean C is \mathbb{P}^1 and X was rational. \square

REMARK 7.2. Some remarks about the case of dimension three, where the minimal model program is incomplete in positive characteristic. From the main theorem in [Kol91], assuming X is smooth and that it admits a free morphism from a curve, we can contract extremal rays in the cone of curves in arbitrary characteristic, to obtain a Fano fibration over a curve, surface or point. In the case where there exists a conic fibration $X \rightarrow S$ where S is a smooth surface, Kollár proves that if the characteristic of k is not 2 then the general fibre is smooth. From Proposition 4.6 it follows that the composition morphism $C \rightarrow S$ is free and so from the above proposition for the case of surfaces, S is a rational surface. Hence X is a conic bundle over a rational surface hence separably rationally connected. If $X \rightarrow Y$ a Fano fibration over a curve, to the author's knowledge, it is not known whether the fibres of the del Pezzo

surface fibration over Y obtained in this way must be smooth. Assuming for the time being that they were, they would be separably rationally connected and from the deformation theory argument in Theorem 5.2 and de Franchis' Theorem [ACG11, 8.27], Y would be \mathbb{P}^1 . From the de Jong-Starr Theorem we would obtain sections $\mathbb{P}^1 \rightarrow X$ from which we could assemble combs with very free teeth to be smoothed to very free rational curves in X , showing that X is separably rationally connected. Finally, even though it is open whether Fano threefolds are separably rationally connected (this result is not true in higher dimensions however), Shepherd-Barron [SB97] proved that Fano threefolds of Picard rank one are liftable to characteristic zero, hence admitting a very free morphism implies they are separably rationally connected.

The following result is well known in the case of \mathbb{P}^1 (see [Deb01, 4.18]) and easily extends to higher genus.

PROPOSITION 7.3. *Let $f : C \rightarrow X$ be a very free morphism from a smooth projective curve C to a smooth projective variety X over an algebraically closed field k . Then for all positive integers m, ℓ*

$$H^0(X, (\Omega_X^\ell)^{\otimes m}) = 0.$$

PROOF. Since $f : C \rightarrow X$ is very free, from Proposition 4.7 there is a variety U such that $C \times U \rightarrow X$ makes X separably C -connected. Being very free is an open property ([Kol96, II.3.2]) so we can assume that the general morphism $f_u : C_u \rightarrow X$ for $u \in U$ is very free and also an immersion from [Kol96, II.1.8], and so $f_u^* \mathcal{T}_X$ is ample from Proposition 2.2 (and by definition of a very free curve in the genus zero case). We conclude that for a general point $x \in X$ there is a morphism $f_u : C_u \rightarrow X$ such that $f_u^* \mathcal{T}_X$ is ample and whose image passes through x . Hence since $f_u^* \Omega_X^1$ is negative, any section of $(\Omega_X^\ell)^{\otimes m}$ must vanish on the image $f(C_u)$ hence on a dense open subset of X , and so on X . \square

COROLLARY 7.4. *Let $f : C \rightarrow X$ as above. Then the Albanese variety $\text{Alb } X$ is trivial.*

PROOF. Note that we have that $\dim \text{Alb } X \leq \dim H^1(X, \mathcal{O}_X) = h^{0,1}$. In characteristic zero Hodge duality gives that $h^{1,0} = h^{0,1}$ but more generally over any algebraically closed field we have (see [Igu55]) that $\dim \text{Alb } X \leq h^{1,0} = h^0(X, \Omega_X^1)$. The result follows from Proposition 7.3. \square

The above also follows from the result in [Gou14], which says that in the above situation $H^1(X, \mathcal{O}_X) = 0$. See *ibid.* for a discussion around the vanishing of $H^i(X, \mathcal{O}_X)$ for separably rationally connected varieties in positive characteristic. Note also that if X is C -connected, since any map $C \rightarrow \text{Alb } X$ must factor through the Jacobian, and there are only countably many homomorphisms between abelian varieties, one concludes that the image of X in $\text{Alb } X$ is either a point or a curve.

8. AN EXAMPLE IN POSITIVE CHARACTERISTIC

Let X be the Fermat quintic surface $x_0^5 + x_1^5 + x_2^5 + x_3^5 = 0$ in \mathbb{P}^3 over an algebraically closed field of characteristic p . In [Shi74] it is proven that if $p \neq 5$ and p is not congruent to 1 modulo 5, then X is a unirational general type surface and if we quotient by the action of the group G of 5-th roots of unity $x_i \mapsto \zeta^i x_i$, then we obtain a Godeaux surface which is again unirational but has algebraic fundamental group $\pi_1^{\text{et}}(X/G, \bar{y}) \cong \mathbb{Z}/5\mathbb{Z}$. Note that in characteristic zero, the notions of rationally chain connected, rationally connected, freely rationally connected (see [She10]) and separably rationally connected all coincide and it is known that each variety in this class is simply connected. In positive characteristic however these notions are in decreasing generality and can differ. A rationally chain connected variety always has finite fundamental group (see [CL03]) whereas a freely rationally connected variety is simply connected (see [She10]). Note that Shioda's example above gives a unirational and hence rationally connected variety over a characteristic p algebraically closed field which is not simply connected.

We show there is a smooth projective variety in characteristic p which has infinite étale fundamental group but after a finite number of MRC quotients we terminate with a point. Let C be a smooth 5 to 1 cover of \mathbb{P}^1 , with defining affine equation of the form $y^5 = f(x)$ where f is a general polynomial of high degree. We have an action of $G = \mathbb{Z}/5\mathbb{Z}$ on C which we can extend to the product $X \times C$ of the above Fermat quintic X with C . Projecting from the quotient onto the second factor we have a morphism $(X \times C)/G \rightarrow \mathbb{P}^1$ where we have identified C/G with \mathbb{P}^1 . The general fibre of this morphism is isomorphic to X . We have a short exact sequence

$$1 \rightarrow \pi_1^{\text{et}}(X, \bar{x}) \times \pi_1^{\text{et}}(C, \bar{c}) \rightarrow \pi_1^{\text{et}}((X \times C)/G, \bar{z}) \rightarrow G \rightarrow 1.$$

Hence we have constructed an example of a smooth projective variety over an algebraically closed field of characteristic p whose fundamental group is infinite yet whose tower of MRC quotients terminates with a point.

REFERENCES

- [ACG11] Enrico Arbarello, Maurizio Cornalba, and Phillip A. Griffiths. *Geometry of algebraic curves. Volume II*, volume 268 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer, Heidelberg, 2011. With a contribution by Joseph Daniel Harris.
- [AK03] Carolina Araujo and János Kollár. Rational curves on varieties. In *Higher dimensional varieties and rational points (Budapest, 2001)*, volume 12 of *Bolyai Soc. Math. Stud.*, pages 13–68. Springer, Berlin, 2003.
- [BDPP13] Sébastien Boucksom, Jean-Pierre Demailly, Mihai Păun, and Thomas Peternell. The pseudo-effective cone of a compact Kähler

- manifold and varieties of negative Kodaira dimension. *J. Algebraic Geom.*, 22(2):201–248, 2013.
- [BM01] Fedor Bogomolov and Michael McQuillan. Rational curves on foliated varieties. *preprint*, 2001.
- [Cam92] F. Campana. Connexité rationnelle des variétés de Fano. *Ann. Sci. École Norm. Sup. (4)*, 25(5):539–545, 1992.
- [CL03] Antoine Chambert-Loir. A propos du groupe fondamental des variétés rationnellement connexes. 2003.
- [Deb01] Olivier Debarre. *Higher-dimensional algebraic geometry*. Universitext. Springer-Verlag, New York, 2001.
- [DG67] Jean Dieudonné and Alexander Grothendieck. Éléments de géométrie algébrique. *Inst. Hautes Études Sci. Publ. Math.*, 4, 8, 11, 17, 20, 24, 28, 32, 1961–1967.
- [dJS03] A. J. de Jong and J. Starr. Every rationally connected variety over the function field of a curve has a rational point. *Amer. J. Math.*, 125(3):567–580, 2003.
- [FP97] W. Fulton and R. Pandharipande. Notes on stable maps and quantum cohomology. In *Algebraic geometry—Santa Cruz 1995*, volume 62 of *Proc. Sympos. Pure Math.*, pages 45–96. Amer. Math. Soc., Providence, RI, 1997.
- [GHS03] Tom Graber, Joe Harris, and Jason Starr. Families of rationally connected varieties. *J. Amer. Math. Soc.*, 16(1):57–67 (electronic), 2003.
- [Gou14] Frank Gounelas. The first cohomology of separably rationally connected varieties. *C. R. Math. Acad. Sci. Paris*, 352(11):871–873, 2014.
- [Har66] Robin Hartshorne. Ample vector bundles. *Inst. Hautes Études Sci. Publ. Math.*, (29):63–94, 1966.
- [Har70] Robin Hartshorne. *Ample subvarieties of algebraic varieties*. Notes written in collaboration with C. Musili. Lecture Notes in Mathematics, Vol. 156. Springer-Verlag, Berlin, 1970.
- [Har71] Robin Hartshorne. Ample vector bundles on curves. *Nagoya Math. J.*, 43:73–89, 1971.
- [Har77] Robin Hartshorne. *Algebraic geometry*. Springer-Verlag, New York, 1977. Graduate Texts in Mathematics, No. 52.
- [Igu55] Jun-ichi Igusa. A fundamental inequality in the theory of Picard varieties. *Proc. Nat. Acad. Sci. U.S.A.*, 41:317–320, 1955.
- [KMM92] János Kollár, Yoichi Miyaoka, and Shigefumi Mori. Rationally connected varieties. *J. Algebraic Geom.*, 1(3):429–448, 1992.
- [Kol91] János Kollár. Extremal rays on smooth threefolds. *Ann. Sci. École Norm. Sup. (4)*, 24(3):339–361, 1991.
- [Kol96] János Kollár. *Rational curves on algebraic varieties*, volume 32 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 1996.

- [KSCT07] Stefan Kebekus, Luis Solá Conde, and Matei Toma. Rationally connected foliations after Bogomolov and McQuillan. *J. Algebraic Geom.*, 16(1):65–81, 2007.
- [Laz04] Robert Lazarsfeld. *Positivity in algebraic geometry. II*, volume 49 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 2004. Positivity for vector bundles, and multiplier ideals.
- [Liu02] Qing Liu. *Algebraic geometry and arithmetic curves*, volume 6 of *Oxford Graduate Texts in Mathematics*. Oxford University Press, Oxford, 2002. Translated from the French by Reinie Ern e, Oxford Science Publications.
- [Miy87] Yoichi Miyaoka. Deformations of a morphism along a foliation and applications. In *Algebraic geometry, Bowdoin, 1985 (Brunswick, Maine, 1985)*, volume 46 of *Proc. Sympos. Pure Math.*, pages 245–268. Amer. Math. Soc., Providence, RI, 1987.
- [Mor79] Shigefumi Mori. Projective manifolds with ample tangent bundles. *Ann. of Math. (2)*, 110(3):593–606, 1979.
- [Occ06] Gianluca Occhetta. Extending rationally connected fibrations. *Forum Math.*, 18(5):853–867, 2006.
- [Ott12] John Christian Ottem. Ample subvarieties and q -ample divisors. *Adv. Math.*, 229(5):2868–2887, 2012.
- [Pet06] Thomas Peternell. Kodaira dimension of subvarieties. II. *Internat. J. Math.*, 17(5):619–631, 2006.
- [Poo10] Bjorn Poonen. Insufficiency of the Brauer-Manin obstruction applied to  tale covers. *Ann. of Math. (2)*, 171(3):2157–2169, 2010.
- [SB97] N. I. Shepherd-Barron. Fano threefolds in positive characteristic. *Compositio Math.*, 105(3):237–265, 1997.
- [She10] Mingmin Shen. Foliations and rational connectedness in positive characteristic. *J. Algebraic Geom.*, 19(3):531–553, 2010.
- [Shi74] Tetsuji Shioda. An example of unirational surfaces in characteristic p . *Math. Ann.*, 211:233–236, 1974.
- [Sta06] Jason Starr. The maximal free rational quotient. *preprint*, 2006.

Frank Gounelas
Institut f ur Mathematik
Humboldt Universit at zu Berlin
Unter den Linden 6
10099 Berlin
gounelas@mathematik.hu-berlin.de