

CATEGORIFIED TRACE FOR MODULE TENSOR CATEGORIES OVER BRAIDED TENSOR CATEGORIES

ANDRÉ HENRIQUES, DAVID PENNEYS, AND JAMES TENER

Received: September 11, 2015

Revised: April 22, 2016

Communicated by Eckhard Meinrenken

ABSTRACT. Given a braided pivotal category \mathcal{C} and a pivotal module tensor category \mathcal{M} , we define a functor $\mathrm{Tr}_{\mathcal{C}} : \mathcal{M} \rightarrow \mathcal{C}$, called the associated categorified trace. By a result of Bezrukavnikov, Finkelberg and Ostrik, the functor $\mathrm{Tr}_{\mathcal{C}}$ comes equipped with natural isomorphisms $\tau_{x,y} : \mathrm{Tr}_{\mathcal{C}}(x \otimes y) \rightarrow \mathrm{Tr}_{\mathcal{C}}(y \otimes x)$, which we call the traciators. This situation lends itself to a diagrammatic calculus of ‘strings on cylinders’, where the traciator corresponds to wrapping a string around the back of a cylinder. We show that $\mathrm{Tr}_{\mathcal{C}}$ in fact has a much richer graphical calculus in which the tubes are allowed to branch and braid. Given algebra objects A and B , we prove that $\mathrm{Tr}_{\mathcal{C}}(A)$ and $\mathrm{Tr}_{\mathcal{C}}(A \otimes B)$ are again algebra objects. Moreover, provided certain mild assumptions are satisfied, $\mathrm{Tr}_{\mathcal{C}}(A)$ and $\mathrm{Tr}_{\mathcal{C}}(A \otimes B)$ are semisimple whenever A and B are semisimple.

2010 Mathematics Subject Classification: 18D10

CONTENTS

1	INTRODUCTION	1090
2	BACKGROUND	1095
2.1	Tensor categories	1096
2.2	The Drinfel’d center	1100
2.3	Synoptic chart of tensor categories	1101
2.4	Algebra objects	1102
3	CATEGORIFIED TRACES FOR MODULE TENSOR CATEGORIES	1104

3.1	Categorified traces	1104
3.2	Module tensor categories	1107
3.3	The categorified trace associated to a module tensor category	1108
4	PROPERTIES OF THE TRACE FUNCTOR	1112
4.1	The attaching map	1112
4.2	The multiplication map	1114
4.3	The unit of the adjunction	1116
4.4	Construction of the traciator	1118
4.5	Interaction between traciator and braiding	1122
5	TRACES OF ALGEBRAS AND SEMISIMPLICITY	1127
5.1	The algebra $\mathrm{Tr}_{\mathcal{C}}(A)$	1127
5.2	The algebra $\mathrm{Tr}_{\mathcal{C}}(A \otimes B)$	1132
A	APPENDIX	1138
A.1	Commutative algebras in braided tensor categories	1138
A.2	Braided pivotal categories	1141

1 INTRODUCTION

A *tensor category* is a linear category \mathcal{M} , equipped with a functor $\otimes : \mathcal{M} \times \mathcal{M} \rightarrow \mathcal{M}$ along with extra data encoding the ideas of associativity and unitality. If x is an object of \mathcal{M} , then its dual x^* is characterized, assuming it exists, by adjunctions $\mathrm{Hom}(y, x \otimes z) \cong \mathrm{Hom}(x^* \otimes y, z)$ and $\mathrm{Hom}(y \otimes x, z) \cong \mathrm{Hom}(y, z \otimes x^*)$. The category \mathcal{M} is *pivotal* if it comes equipped with certain isomorphisms $\varphi_x : x \rightarrow x^{**}$ from every object to its double dual. It is interesting to note that, in a pivotal category, the functor $x \mapsto \mathrm{Hom}(1, x)$ satisfies the following cyclic invariance property:

$$\begin{aligned} \mathrm{Hom}(1, x \otimes y) &\cong \mathrm{Hom}(x^* \otimes 1, y) \cong \mathrm{Hom}(1 \otimes x^*, y) \\ &\cong \mathrm{Hom}(1, y \otimes x^{**}) \cong \mathrm{Hom}(1, y \otimes x). \end{aligned}$$

We think of $\mathrm{Hom}(1, -)$ as a vector space valued trace $\mathrm{Tr} : \mathcal{M} \rightarrow \mathrm{Vec}$. This is our prototypic example of a *categorified trace*.


Given a tensor category \mathcal{C} , and two objects x, y of some module category \mathcal{M} , the internal hom $\underline{\mathrm{Hom}}(x, y)$ is the object of \mathcal{C} that represents the exact functor $c \mapsto \mathcal{M}(c \cdot x, y)$ [Ost03, Def. 3.4]. If in addition to being a module category \mathcal{M} is also a tensor category in its own right, then we may consider the functor

$$\mathrm{Tr}_{\mathcal{C}} := \underline{\mathrm{Hom}}(1_{\mathcal{M}}, -) : \mathcal{M} \rightarrow \mathcal{C},$$

and ask whether it has a similar cyclic invariance property.

The appropriate compatibility between the \mathcal{C} -module structure and the tensor structure of \mathcal{M} can only be formulated when \mathcal{C} is braided, and the resulting notion is what we call a *module tensor category* (Definition 3.5). We write

$\Phi : \mathcal{C} \rightarrow \mathcal{M}$ for the functor that sends $c \in \mathcal{C}$ to $c \cdot 1_{\mathcal{M}} \in \mathcal{M}$. By definition, equipping \mathcal{M} with the structure of a module tensor category over \mathcal{C} is the same thing as equipping the functor Φ with a factorization $\mathcal{C} \xrightarrow{\Phi^z} \mathcal{Z}(\mathcal{M}) \rightarrow \mathcal{M}$, where Φ^z is a braided functor to the Drinfel'd center of \mathcal{M} [Bez04, Def.1] [DMNO13, Def.2.4]. The trace functor can be alternatively described as the right adjoint of Φ :




A *categorified trace* (Definition 3.1), also known as a commutator functor [BFO09, §6], [Ost14, Def. 2.1], is a functor $\text{Tr} : \mathcal{M} \rightarrow \mathcal{C}$ equipped with natural isomorphisms


$$\tau_{x,y} : \text{Tr}(x \otimes y) \rightarrow \text{Tr}(y \otimes x),$$

which we call the *traciators*, subject to the axiom $\tau_{x,y \otimes z} = \tau_{z \otimes x,y} \circ \tau_{x \otimes y,z}$. At the level of Grothendieck groups, the functor Tr induces a map $K_0(\text{Tr}) : K_0(\mathcal{M}) \rightarrow K_0(\mathcal{C})$. The traciator isomorphism descends to an equality $K_0(\text{Tr})([x][y]) = K_0(\text{Tr})([y][x])$ in $K_0(\mathcal{C})$, so $K_0(\text{Tr})$ satisfies the defining property of a trace.

We denote a categorified trace graphically by lifting objects and morphisms up from the plane (corresponding to \mathcal{M}) onto a cylinder in 3-space (corresponding to \mathcal{C})



Following [PS13]¹, the traciator is represented graphically by a strand wrapping around the cylinder:



When \mathcal{C} and \mathcal{M} are pivotal and Φ^z is a pivotal functor, then $\text{Tr}_{\mathcal{C}}$ is a categorified trace. In fact, this only depends on Φ factoring through the Drinfel'd center, and not on it being a tensor functor, as proven in [BFO09, Prop. 5] and [Ost14, Prop. 2.5].

¹See Remark 3.2 for a subtle difference between our notion of categorified trace and the one used in [PS13].

The fact that Φ is a tensor functor contributes to the structure of $\mathrm{Tr}_{\mathcal{C}}$ in a different way. Adjoints of tensor functors are lax monoidal [Kel74], and so we have unit and multiplication maps $i : 1_{\mathcal{C}} \rightarrow \mathrm{Tr}_{\mathcal{C}}(1_{\mathcal{M}})$ and $\mu_{x,y} : \mathrm{Tr}_{\mathcal{C}}(x) \otimes \mathrm{Tr}_{\mathcal{C}}(y) \rightarrow \mathrm{Tr}_{\mathcal{C}}(x \otimes y)$ which we represent graphically as follows:

$$i = \text{cup} \qquad \mu_{x,y} = \text{Y-junction with strands } x \text{ and } y$$

The novelty of this paper is the rich interplay between the above two structures. Using everything we have, we can assign a morphism to any picture of strands on tubes. These, in turn, can be used to formulate a number of non-trivial identities, such as

$$\begin{array}{ccc}
 \text{Y-junction with crossing} = \text{Y-junction with twist} & \longleftrightarrow & \begin{array}{ccc}
 \mathrm{Tr}_{\mathcal{C}}(x) \otimes \mathrm{Tr}_{\mathcal{C}}(y) & \xrightarrow{\mu} & \mathrm{Tr}_{\mathcal{C}}(x \otimes y) \\
 \beta \circ (\mathrm{id} \otimes \theta) \downarrow & & \downarrow \tau \\
 \mathrm{Tr}_{\mathcal{C}}(y) \otimes \mathrm{Tr}_{\mathcal{C}}(x) & \xrightarrow{\mu} & \mathrm{Tr}_{\mathcal{C}}(y \otimes x)
 \end{array}
 \end{array}$$

and

$$\text{Complex Y-junction with multiple strands} = \text{Another configuration of the same strands} \longleftrightarrow$$

$$\begin{array}{ccc}
 \mathrm{Tr}_{\mathcal{C}}(w) \otimes \mathrm{Tr}_{\mathcal{C}}(x \otimes y) \otimes \mathrm{Tr}_{\mathcal{C}}(z) & \xrightarrow{\mathrm{id} \otimes \tau \otimes \mathrm{id}} & \mathrm{Tr}_{\mathcal{C}}(w) \otimes \mathrm{Tr}_{\mathcal{C}}(y \otimes x) \otimes \mathrm{Tr}_{\mathcal{C}}(z) \\
 \downarrow \mu \otimes \mathrm{id} & & \downarrow \mathrm{id} \otimes \mu \\
 \mathrm{Tr}_{\mathcal{C}}(w \otimes x \otimes y) \otimes \mathrm{Tr}_{\mathcal{C}}(z) & & \mathrm{Tr}_{\mathcal{C}}(w) \otimes \mathrm{Tr}_{\mathcal{C}}(y \otimes x \otimes z) \\
 \downarrow \tau \otimes \mathrm{id} & & \downarrow \mathrm{id} \otimes \tau^{-1} \\
 \mathrm{Tr}_{\mathcal{C}}(y \otimes w \otimes x) \otimes \mathrm{Tr}_{\mathcal{C}}(z) & & \mathrm{Tr}_{\mathcal{C}}(w) \otimes \mathrm{Tr}_{\mathcal{C}}(x \otimes z \otimes y) \\
 \downarrow \mu & & \downarrow \mu \\
 \mathrm{Tr}_{\mathcal{C}}(y \otimes w \otimes x \otimes z) & \xrightarrow{\tau^{-1}} & \mathrm{Tr}_{\mathcal{C}}(w \otimes x \otimes z \otimes y)
 \end{array}$$

which combine the traciator and multiplication maps with the braiding and the twist of \mathcal{C} (see Lemmas 4.26 and 4.19). We summarize the situation in Figure 1.




Figure 1: Our setup (in the middle), in comparison to previously studied situations.

In our sequel paper [HPT16], we will prove that the relations established in this paper imply that the morphism assigned to a diagram is invariant under *all* isotopies. This verification will be performed in the language of planar algebras internal to braided tensor categories. In our forthcoming paper, we will later

classify planar algebras internal to \mathcal{C} in terms of module tensor categories for \mathcal{C} .

It is well known that lax tensor functors send algebras to algebras. As a result, if A is an algebra object in \mathcal{M} , then $\mathrm{Tr}_{\mathcal{C}}(A)$ is an algebra object in \mathcal{C} . In our situation, more is true. If A and B are two algebra objects, then $\mathrm{Tr}_{\mathcal{C}}(A \otimes B)$ is again an algebra object (actually, this only requires \mathcal{C} and Φ^z to be monoidal, as opposed to braided). The structure maps are best illustrated by putting the strand corresponding to A on the front of the cylinders, and the one corresponding to B on the back of the cylinders:

$$m_{\mathrm{Tr}_{\mathcal{C}}(A \otimes B)} = \text{diagram} \qquad i_{\mathrm{Tr}_{\mathcal{C}}(A \otimes B)} = \text{diagram}$$

Let us now assume that \mathcal{C} and \mathcal{M} are fusion categories over a field of characteristic zero. One of our main results is that if A and B are semisimple algebras then so is $\mathrm{Tr}_{\mathcal{C}}(A \otimes B)$ (Theorem 5.6 and Corollary 5.9):

THEOREM A. *Let \mathcal{C} and \mathcal{M} be tensor categories subject to the above assumptions, and let $\Phi^z : \mathcal{C} \rightarrow \mathcal{Z}(\mathcal{M})$ be a tensor functor. If $A, B \in \mathcal{M}$ are semisimple algebras, then $\mathrm{Tr}_{\mathcal{C}}(A \otimes B)$ is also semisimple.*

Alternatively, we can trade the extra assumptions on \mathcal{C} and \mathcal{M} for the assumption that A and B are separable algebras (Theorem 5.6). We illustrate our construction by computing the algebra $\mathrm{Tr}_{\mathcal{C}}(A \otimes B)$ for certain algebra objects related to the Coxeter–Dynkin diagrams E_7 and D_{10} (Example 5.10). Note that given three algebras $A, B, C \in \mathcal{M}$, we see no reason why $\mathrm{Tr}_{\mathcal{C}}(A \otimes B \otimes C)$ should carry an algebra structure. Unfortunately, we cannot supply a counterexample at this time.

In the special case in which we have only one algebra, our result holds in the greater generality of an arbitrary tensor functor $\Phi : \mathcal{C} \rightarrow \mathcal{M}$ between rigid semisimple tensor categories (Theorem 5.1 and Corollary 5.3):

THEOREM B. *Let \mathcal{C} and \mathcal{M} be tensor categories subject to the above assumptions, and let $\Phi : \mathcal{C} \rightarrow \mathcal{M}$ be a tensor functor with right adjoint $\mathrm{Tr}_{\mathcal{C}}$. If $A \in \mathcal{M}$ is a semisimple algebra, then $\mathrm{Tr}_{\mathcal{C}}(A)$ is as well. In particular, if A is semisimple connected, and thus Frobenius, then so is $\mathrm{Tr}_{\mathcal{C}}(A)$.*

We finish our introduction by listing three applications of categorified traces. The first occurrence of categorified traces in the literature goes back to [MV88], where a certain functor was constructed between categories of B -equivariant sheaves on G/B and G -equivariant sheaves on G for the adjoint action (G is a reductive group and B a Borel subgroup). It was recognized in [BFO09, §6] that this functor satisfies the defining properties of a categorified trace, and this was later used in [Ost14] to analyze the structure of tensor categories attached to cells in Weyl groups.

The first author recently defined higher categorical analogs of von Neumann algebras called a bicommutant categories [Hen15], and examples from fusion categories were constructed in [HP15]. These bicommutant categories admit categorified traces with values in the category of Hilbert spaces, a fact that was crucial to the study of their structural properties.² In the special case of the category of bimodules over a von Neumann factor, the categorified trace is the ‘cyclic fusion’, first introduced in the context of conformal nets [BDH14, App. A]. For more general bicommutant categories, we expect categorified traces with values in the Drinfel’d center of the bicommutant category. In their recent work [JL16], Jaffe and Liu introduced a novel notion of ‘planar para algebra’. Their main examples are the parafermion algebras. In our forthcoming article [HPT16], we will show that planar para algebras can be constructed using categorified traces, and that the vector spaces that constitute the parafermion algebras are given by $\mathrm{Tr}_{\mathcal{C}}(m^{\otimes n})$, where m is the generating object of a Tambara-Yamagami category [TY98] (a \mathcal{C} -module tensor category for \mathcal{C} a $\mathrm{Vec}(\mathbb{Z}/N\mathbb{Z})$ ribbon category). Our categorified trace will provide a direct connection between Izumi’s calculation of the center of the Tambara-Yamagami categories [Izu01], and Jaffe-Liu’s work on the parafermion planar para algebras.

1.1 ACKNOWLEDGEMENTS

The authors would like to thank Bojko Bakalov, Bruce Bartlett, Alexandru Chirvasitu, Vaughan Jones, Scott Morrison, Richard Ng, Victor Ostrik, Mathew Pugh, Ingo Runkel, Chris Schommer-Pries, Noah Snyder, and Kevin Walker for helpful discussions.

The authors would like to thank Mathematisches Forschungsinstitut Oberwolfach for their hospitality during the workshop on “Subfactors and conformal field theory.” André Henriques gratefully acknowledges the Leverhulme trust for financing his visiting position in Oxford, and the Hausdorff Institute for its hospitality while visiting James Tener. James Tener would like to thank the Max Planck Institute for Mathematics for its hospitality and support during the course of this work. David Penneys and James Tener were partially supported by NSF DMS grant 0856316. David Penneys was partially supported by an AMS-Simons travel grant and NSF DMS grant 1500387.

2 BACKGROUND

The natural setting to study the categorified trace is when \mathcal{C} is a braided pivotal category, \mathcal{M} is a pivotal category, and $\Phi^z : \mathcal{C} \rightarrow \mathcal{Z}(\mathcal{M})$ is a braided pivotal functor. This is perhaps an unfamiliar setting, since most readers are more likely to work with either braided tensor categories or ribbon categories,

²These categorified traces are only partially defined, see [BDH14, Warn. A.6] for more details.

and braided pivotal categories are a strange intermediate (see Figure 2 in Section 2.3).

For these reasons, it is important to carefully understand what we can and cannot do in a braided pivotal category. Moreover, in the literature there are sometimes redundancies within definitions, as well as incorrect statements, so we begin with a short, comprehensive background on various flavours of tensor categories. Various technical lemmas about braided pivotal categories are deferred to Appendix A.2. Finally, we provide a short section on algebras in tensor categories.

2.1 TENSOR CATEGORIES

We will usually assume that all our categories are linear over some field k (the morphism spaces are finite dimensional vector spaces), and that all functors are linear. However, the large majority of our results do not require this linearity assumption. The only place where this is really required is for statements involving semisimplicity of certain algebra objects (Theorems 5.1 and 5.6).

DEFINITION 2.1. A tensor category³ consists of the data $(\mathcal{C}, 1, \otimes, \alpha, \lambda, \rho)$ where \mathcal{C} is a category, $1 \in \mathcal{C}$ is the distinguished unit object, $\otimes : \mathcal{C} \boxtimes \mathcal{C} \rightarrow \mathcal{C}$ is a bifunctor, the associator $\alpha : (a \otimes b) \otimes c \rightarrow a \otimes (b \otimes c)$ is a natural isomorphism, and the left and right unitors $\lambda : 1 \otimes a \rightarrow a$ and $\rho : a \otimes 1 \rightarrow a$ are natural isomorphisms. This data must satisfy the well known pentagon and triangle axioms.

A tensor category \mathcal{C} is called:⁴

- rigid (or has duals) if for every $a \in \mathcal{C}$ there is an object $a^* \in \mathcal{C}$, and there are maps $\text{coev}_a : 1 \rightarrow a \otimes a^*$ and $\text{ev}_a : a^* \otimes a \rightarrow 1$ satisfying the zig-zag axioms

$$\begin{aligned} (\text{id}_a \otimes \text{ev}_a) \circ (\text{coev}_a \otimes \text{id}_a) &= \text{id}_a \\ (\text{ev}_a \otimes \text{id}_{a^*}) \circ (\text{id}_{a^*} \otimes \text{coev}_a) &= \text{id}_{a^*} . \end{aligned}$$

Moreover, for every $a \in \mathcal{C}$, there should exist an object ${}^*a \in \mathcal{C}$ such that $({}^*a)^* \cong a$.

Note that being rigid is not data; it is just a property of the category.⁵ Given $f : a \rightarrow b$ in \mathcal{C} , we write $f^* : b^* \rightarrow a^*$ for $(\text{ev}_b \otimes \text{id}_{a^*}) \circ (\text{id}_{b^*} \otimes f \otimes \text{id}_{a^*}) \circ (\text{id}_{b^*} \otimes \text{coev}_a)$.

³The adjectives ‘tensor’ and ‘monoidal’ are essentially synonymous. The first one is usually only used when the category is linear; the second one is used regardless of whether the category is linear or not.

⁴Alternative terminologies include [Sel11]: rigid = autonomous, pivotal = sovereign, ribbon = tortile.

⁵In any category, an object a is said to *have a dual* if there exist solutions of the zig-zag equations. The dual object a^* is then unique up to canonical isomorphism, and may thus be referred to as *the* dual of a .


- **fusion** if \mathcal{C} is rigid, semisimple, $\mathcal{C}(1, 1) \cong k$, and there are only finitely many isomorphism classes of simple objects.
- **pivotal** if \mathcal{C} is rigid, and there is a monoidal natural isomorphism φ from the identity functor to the double dual functor.⁶ The left and right pivotal traces of a morphism $f : a \rightarrow a$ are then defined by

$$\begin{aligned} \text{tr}_L(f) &= \text{ev}_a \circ (\text{id}_{a^*} \otimes f) \circ (\text{id}_{a^*} \otimes \varphi_a^{-1}) \circ \text{coev}_{a^*} \\ \text{tr}_R(f) &= \text{ev}_{a^*} \circ (\varphi_a \otimes \text{id}_{a^*}) \circ (f \otimes \text{id}_{a^*}) \circ \text{coev}_a. \end{aligned}$$

- **spherical** if \mathcal{C} is pivotal and for every $c \in \mathcal{C}$ and $f \in \mathcal{C}(c, c)$, we have $\text{tr}_L(f) = \text{tr}_R(f)$.
- **braided** if there is a family of natural isomorphisms $\beta_{a,b} : a \otimes b \rightarrow b \otimes a$ satisfying the two well known hexagon axioms. We also write $\beta_{a,b}^+$ for $\beta_{a,b}$ and $\beta_{a,b}^-$ for $\beta_{b,a}^{-1}$.
- **balanced** if \mathcal{C} is braided and there are twist isomorphisms $\theta_a : a \rightarrow a$ for $a \in \mathcal{C}$, natural in a , satisfying $\theta_{a \otimes b} = \beta_{b,a} \circ \beta_{a,b} \circ (\theta_a \otimes \theta_b)$ for all $a, b \in \mathcal{C}$.
- **ribbon**⁷ if \mathcal{C} is balanced and rigid, and the twist maps satisfy $\theta_{a^*} = \theta_a^*$ for all $a \in \mathcal{C}$.

There exist graphical calculi for morphisms in various kinds of monoidal or tensor categories. We may draw different types of diagrams and perform different types of isotopies based on the properties and structures of our category. We give a helpful guide below (see [Sel11] for a comprehensive survey). If \mathcal{C} is...


- **monoidal** (Joyal–Street [JS91a]), we may draw our string diagrams so that strands only go up and down. We read the diagram from bottom to top.




⁶The original definition of Freyd and Yetter [FY92] contains the redundant axiom $\varphi_{a^*} = (\varphi_a^{-1})^*$, later reproduced by other authors. See [Sel11, Lem. 4.11] for a proof of that property.

⁷We warn the reader that [BK01, Def. 2.2.1] defines the notion of a braided pivotal category (equivalently, a balanced rigid category – see Appendix A.2), rather than that of a ribbon category. In a braided pivotal category, it is not necessarily the case that $\theta_{a^*} = \theta_a^*$. The same problem appears in [KO02, Eq. (1.6)].

- rigid, then we can draw any planar diagram. Strings may bend up and down, however coupons are not allowed to rotate.




- pivotal (Freyd–Yetter [FY92]), we may rotate coupons by 2π without changing the morphism. In defining the right hand side below, we must use the pivotal structure:



- spherical, then the value of a closed diagram is invariant under spherical isotopy. In particular:


$$\text{tr}_R(f) = \left(\begin{array}{c} \circlearrowleft \\ f \end{array} \right) = \left(\begin{array}{c} f \\ \circlearrowright \end{array} \right) = \text{tr}_L(f).$$

- braided (Joyal–Street [JS91a]), then our diagrams are now in three dimensions, and we draw them projected to the plane. If \mathcal{C} is not rigid, then coupons again may only travel up and down, and are not allowed to rotate.




- balanced (Shum [Shu94]), then strands are replaced by ribbons, which can twirl on themselves, but cannot bend up and down. Coupons may

rotate around the z -axis.




- **braided pivotal** (Freyd–Yetter [FY92]), then the second and third Reidemeister moves are allowed, but not the first. Strands behave like ribbons pressed flat against the plane.



There are two ways of making such a category into a balanced one, by letting θ be either $|$ or $|$ (see Lemma A.2 in the appendix).

- **ribbon** (Shum [Shu94]), then the strands are replaced by ribbons, and all three dimensional isotopies are allowed.



Given a functor between categories with extra structure, one can ask for the functor to be compatible with that structure. We list the relevant conditions for the notions mentioned above. Let \mathcal{C} and \mathcal{D} be tensor categories, and let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a functor.

- We say that F is a **tensor functor** if it comes equipped with an invertible⁸ natural transformation $\nu_{x,y} : F(x) \otimes F(y) \rightarrow F(x \otimes y)$ and an isomorphism

⁸ If ν and i are not assumed invertible, then this is called a *lax tensor functor*. We warn the reader that some people use ‘strong tensor functor’ for tensor functors, and ‘tensor functor’ for lax tensor functors.

PROPOSITION 2.3. *Let \mathcal{C} be a rigid tensor category. Then a pivotal structure on \mathcal{C} induces a pivotal structure on $\mathcal{Z}(\mathcal{C})$.*

Proof (communicated by Noah Snyder). The dual of an object $(a, e_a) \in \mathcal{Z}(\mathcal{C})$ is given by (a^*, e_{a^*}) , with $e_{a^*,b} := e_{a^*,b}^*$. The somewhat non-trivial claim is that the map $\varphi_a : a \rightarrow a^{**}$ provided by the pivotal structure of \mathcal{C} defines a morphism $(a, e_a) \rightarrow (a, e_a)^{**}$ in $\mathcal{Z}(\mathcal{C})$. We need to show that $(\text{id}_b \otimes \varphi_a) \circ e_{a,b} = e_{a^{**},b} \circ (\varphi_a \otimes \text{id}_b)$ for every $b \in \mathcal{C}$. This is done as follows:

$$\begin{aligned} e_{a^{**},b} \circ (\varphi_a \otimes \text{id}_b) &= e_{a^{**},b}^* \circ \varphi_{a \otimes b} \circ (\text{id}_a \otimes \varphi_{b^{**}}^{-1}) \\ &= \varphi_{b \otimes a} \circ e_{a,b} \circ (\text{id}_a \otimes \varphi_{b^{**}}^{-1}) \\ &= \varphi_{b \otimes a} \circ (\varphi_{b^{**}}^{-1} \otimes \text{id}_a) \circ e_{a,b} = (\text{id}_b \otimes \varphi_a) \circ e_{a,b}. \quad \square \end{aligned}$$

2.3 SYNOPSIS CHART OF TENSOR CATEGORIES

We present here a chart (Figure 2) that summarizes the various notions of tensor category. The two ways of going between balanced rigid categories and braided pivotal categories will be discussed in Appendix A.2.

In Figure 2, we use the following notations:

- An arrow $\textcircled{A} \longrightarrow \textcircled{B}$ indicates that notion B can be obtained from notion A by forgetting part of the data.
- An arrow $\textcircled{A} \longleftarrow \textcircled{B}$ indicates that notion A can be obtained from notion B by imposing extra axioms.
- A dashed arrow $\textcircled{A} \overset{\mathcal{Z}}{\dashrightarrow} \textcircled{B}$ indicates that the Drinfel'd center construction goes from notion A to notion B.
- A double arrow $\textcircled{A} \overset{\cong}{\longleftrightarrow} \textcircled{B}$ indicates an equivalence between notions A and B.

As mentioned above, there are *two* ways of going between balanced rigid categories and braided pivotal categories. To avoid any confusion, we will only use *one* of those two equivalences in the remainder of this paper. Specifically, given a braided pivotal category $(\mathcal{C}, \beta, \varphi)$, we will always equip it with the twists given by

$$\theta_a := (\text{id}_a \otimes \text{ev}_{a^*}) \circ (\beta_{a^{**},a} \otimes \text{id}_{a^*}) \circ (\text{id}_{a^{**}} \otimes \text{coev}_a) \circ \varphi_a = \begin{array}{c} a \\ | \\ \textcircled{\varphi_a} \\ | \\ a \end{array} \quad (3)$$

See Appendix A.2 for more details.




Figure 2: Synoptic chart of tensor categories

2.4 ALGEBRA OBJECTS

We now discuss algebra objects in tensor categories. General references include [KO02, §1], [DMNO13, §2.4], [Ost03, §3], and [Müg03a].

DEFINITION 2.4. An algebra object in a tensor category \mathcal{C} is a triple $(A, m : A \otimes A \rightarrow A, i : 1 \rightarrow A)$ which satisfies the following axioms:

$$\begin{aligned}
 m(m \otimes \text{id}_A) &= m(\text{id}_A \otimes m) && \text{(Associativity)} \\
 m(i \otimes \text{id}_A) &= \text{id}_A = m(\text{id}_A \otimes i) && \text{(Unitality)}
 \end{aligned}$$

Dually, a coalgebra is a triple $(A, \Delta : A \rightarrow A \otimes A, \epsilon : A \rightarrow 1)$ satisfying analogous coassociativity and counitality axioms.

We represent the multiplication m and unit i by a trivalent, respectively univalent, vertex

$$m_A = \begin{array}{c} |^A \\ \bullet \\ / \quad \backslash \\ A \quad A \end{array} \qquad i_A = \begin{array}{c} |^A \\ \bullet \end{array} .$$

An algebra object is called *connected* if $\mathcal{C}(1, A)$ is one dimensional. If the ambient category \mathcal{C} is semisimple, then an algebra object is:

- *semisimple* if the category $\text{Mod}_{\mathcal{C}}(A)$ of A -modules in \mathcal{C} is semisimple,
- *separable* if $m_A : A \otimes A \rightarrow A$ splits as a morphism of A - A -bimodules.

At first glance, the notions of semisimple and separable algebras do not seem very related. However, a special case of the following proposition (namely setting $B = 1$) shows that separable always implies semisimple:

PROPOSITION 2.5. *Let A and B be separable algebras in a semisimple tensor category \mathcal{C} . Then the category $\text{Bimod}_{\mathcal{C}}(A, B)$ of A - B -bimodules is semisimple.*

Proof (communicated by Victor Ostrik). Let $s_A : A \rightarrow A \otimes A$ and $s_B : B \rightarrow B \otimes B$ be bimodule maps that split m_A and m_B . Given $M \in \text{Bimod}_{\mathcal{C}}(A, B)$, the action map $A \otimes M \otimes B \rightarrow M$ admits a splitting

$$M \cong A \otimes_A M \otimes_B B \xrightarrow{s_A \otimes \text{id} \otimes s_B} (A \otimes A) \otimes_A M \otimes_B (B \otimes B) \cong A \otimes M \otimes B,$$

which is furthermore a bimodule map. M is therefore isomorphic to a direct summand of $A \otimes M \otimes B$. As the latter is projective, so is M . Every object of $\text{Bimod}_{\mathcal{C}}(A, B)$ is projective, every exact sequence splits, and the category is semisimple. \square

We also get the following well-known equivalent definition of separability:

PROPOSITION 2.6 ([DMNO13, Prop. 2.7]). *An algebra object A in a semisimple tensor category \mathcal{C} is separable if and only if $\text{Bimod}_{\mathcal{C}}(A, A)$ is semisimple.*

Proof. The multiplication map $m_A : A \otimes A \rightarrow A$ admits a right inverse (given by $\text{id}_A \otimes i_A$) and is thus always an epimorphism. If $\text{Bimod}_{\mathcal{C}}(A, A)$ is semisimple, m_A therefore splits. The other direction follows from the previous proposition by setting $B = A$. \square

REMARK 2.7. In positive characteristic there exist semisimple algebras which are not separable, e.g., the group algebra $\mathbb{F}_p[G]$ viewed as a G -graded vector space, for some p -group G .

However, if \mathcal{C} is a fusion category over a field of characteristic zero, then algebra objects in \mathcal{C} are semisimple if and only if they are separable. To see that, consider a semisimple algebra object A , which we assume without loss of generality to be simple (not a direct sum of smaller algebras). Then $\mathcal{M} := \text{Mod}_{\mathcal{C}}(A)$ is an indecomposable module category, and the category $\text{Func}_{\mathcal{C}}(\mathcal{M}, \mathcal{M}) = \text{Bimod}_{\mathcal{C}}(A, A)$ of \mathcal{C} -linear endofunctors of \mathcal{M} is semisimple by [ENO05, Lem. 2.15].

We now briefly introduce the notion of a Frobenius algebra object:

DEFINITION 2.8. A Frobenius algebra in a tensor category \mathcal{C} is a quintuplet $(A, m, \Delta, i, \epsilon)$ such that (A, m, i) is an algebra, (A, Δ, ϵ) is a coalgebra, and the following compatibility condition is satisfied:

$$(m \otimes \text{id}_A) \circ (\text{id}_A \otimes \Delta) = \Delta \circ m = (\text{id}_A \otimes m) \circ (\Delta \otimes \text{id}_A) \quad (\text{Frobenius})$$

It is interesting to note that a Frobenius algebra is entirely determined by its underlying algebra (A, m, i) and counit ϵ . Moreover, for a given algebra $A = (A, m, i)$, a morphism $\epsilon : A \rightarrow 1$ is the counit of a Frobenius structure if and only if the pairing $\epsilon \circ m : A \otimes A \rightarrow 1$ is *non-degenerate* (a pairing $p : A \otimes B \rightarrow 1$ is non-degenerate if A and B are dualizable and $(p \otimes \text{id}_{B^*}) \circ (\text{id}_A \otimes \text{coev}_B) : A \rightarrow B^*$ is an isomorphism):

PROPOSITION 2.9 ([FRS02, Lem. 3.7], [FS08, Prop. 8]). *Let \mathcal{C} be a tensor category, and let (A, m, i) be an algebra object in \mathcal{C} . Let $\epsilon : A \rightarrow 1$ be such that the pairing $\epsilon \circ m$ is non-degenerate. Then, there exists a unique comultiplication $\Delta : A \rightarrow A \otimes A$ making the quintuplet $(A, m, \Delta, i, \epsilon)$ into a Frobenius algebra.*

3 CATEGORIFIED TRACES FOR MODULE TENSOR CATEGORIES

In this section, we explain the general notion of categorified trace, along with some basic properties thereof. We then explain how to construct a categorified trace $\text{Tr}_{\mathcal{C}} : \mathcal{M} \rightarrow \mathcal{C}$ from a module tensor category \mathcal{M} over a braided tensor category \mathcal{C} .

3.1 CATEGORIFIED TRACES

Recall that a trace on an algebra is a linear map to the ground field satisfying $\text{tr}(xy) = \text{tr}(yx)$. The categorification of the notion of algebra is that of tensor category, where notably the associativity constraint gets replaced by the additional data of associators. The notion of a *categorified trace* (also known as a *commutator functor* [BFO09, §6], [Ost14, Def. 2.1]) is defined in the same spirit:

DEFINITION 3.1. Let \mathcal{M} be a tensor category, and let \mathcal{C} be a category. A *categorified trace* is a functor $\text{Tr} : \mathcal{M} \rightarrow \mathcal{C}$ along with natural isomorphisms

$$\tau_{x,y} : \text{Tr}(x \otimes y) \rightarrow \text{Tr}(y \otimes x) \quad x, y \in \mathcal{M}$$

which we call the *traciators*, subject to the following axiom:

$$\tau_{x,y \otimes z} = \tau_{z \otimes x,y} \circ \tau_{x \otimes y,z}. \tag{4}$$

Here, we have suppressed the associators for readability. The axiom including the necessary associators reads as follows: $\text{Tr}(\alpha_{y,z,x}) \circ \tau_{x,y \otimes z} \circ \text{Tr}(\alpha_{x,y,z}) = \tau_{z \otimes x,y} \circ \text{Tr}(\alpha_{z,x,y})^{-1} \circ \tau_{x \otimes y,z}$.

We will sometimes write $\tau_{x,y}^+$ for $\tau_{x,y}$, and $\tau_{x,y}^-$ for $\tau_{y,x}^{-1}$. The latter is called the inverse traciator. We point out that τ^- does not in general satisfy equation (4), but instead a mirrored version of that axiom.

REMARK 3.2. A slightly less general notion of categorified trace was studied in [PS13] under the name “shadow”. It differs from Definition 3.1 in that the authors also included the axioms⁹


$$\tau_{x,1} = \text{id} \quad \text{and} \quad \tau_{1,x} = \text{id}.$$

The first axiom above is redundant: see Lemma 3.3 below. The second axiom is not satisfied by the trace functors we will consider (see Remark 3.15), and we prefer not to include it in the general definition of a categorified trace.


LEMMA 3.3 ([Ost14, Rem. 2.2.]). *Let (Tr, τ) be a categorified trace. Then $\tau_{x,1} = \text{id}_{\text{Tr}(x)}$.*

Proof. Omitting unitors and associators, we have $\tau_{x,1} = \tau_{x,1 \otimes 1} = \tau_{1 \otimes x, 1} \circ \tau_{x \otimes 1, 1} = \tau_{x,1} \circ \tau_{x,1}$, where the second equality holds by Equation (4). As $\tau_{x,1}$ is invertible, it follows that $\tau_{x,1} = \text{id}_{\text{Tr}(x)}$. \square

In Section 2.1 we have reviewed the classical string diagram notation for objects and morphisms in tensor categories. As explained in [PS13], a good graphical notation for categorified traces is to take the strands that represent the objects of \mathcal{M} , and place them on the surface of a cylinder. For example, if we denote $x \in \mathcal{M}$ as a strand on a horizontal plane, then we represent $\text{Tr}(x) \in \mathcal{C}$ as the same strand on the surface of a cylinder:




Morphisms $f \in \mathcal{M}(x, y)$ are denoted by coupons in the horizontal plane, and we represent $\text{Tr}(f) : \text{Tr}(x) \rightarrow \text{Tr}(y)$ as the same coupon, but now on a cylinder:




⁹We have again suppressed the coherences for readability. The actual axioms are $\text{Tr}(\lambda_x) \circ \tau_{x,1} = \text{Tr}(\rho_x)$ and $\text{Tr}(\rho_x) \circ \tau_{1,x} = \text{Tr}(\lambda_x)$.

The traciators $\tau^\pm : \text{Tr}_{\mathcal{C}}(x \otimes y) \rightarrow \text{Tr}_{\mathcal{C}}(y \otimes x)$ are depicted



The naturality of the traciator and the traciator axiom (4) are given in diagrams by




To illustrate the graphical notation, we prove a general basic property of categorified traces.


LEMMA 3.4. *Let $\text{Tr} : \mathcal{M} \rightarrow \mathcal{C}$ be a categorified trace. Then for any dualizable object $x \in \mathcal{M}$, the following equality holds:*

$$\tau_{y,x^*}^+ = \text{Tr}(\text{id}_{x^* \otimes y} \otimes \text{ev}_x) \circ \tau_{x,x^* \otimes y \otimes x^*}^- \circ \text{Tr}(\text{coev}_x \otimes \text{id}_{y \otimes x^*}).$$

In diagrams:



Proof. We prove this identity after composing with the inverse traciator $\tau_{x^*,y}^-$:



The first equality holds by naturality of τ^- . The second one is the analog of (4) for inverse traciators. Finally, the third equality follows from naturality, along with Lemma 3.3. \square

The main focus of this paper is the construction of a particular class of examples of categorical traces, which have the further property that the cylinders can branch and braid (see Figure 1). To obtain such a categorified trace $\text{Tr} : \mathcal{M} \rightarrow \mathcal{C}$, we must assume that \mathcal{M} is equipped with the structure of a pivotal module tensor category over \mathcal{C} , a notion we describe below.

3.2 MODULE TENSOR CATEGORIES

If \mathcal{M} and \mathcal{C} are tensor categories, a functor $\Phi : \mathcal{C} \rightarrow \mathcal{M}$ equips \mathcal{M} with the structure of a left module category via $c \cdot m := \Phi(c) \otimes m$. If \mathcal{C} is braided, then Φ actually equips \mathcal{M} with *two* distinct left \mathcal{C} -module structures (one coming from the left action above, and one coming from the right action, twisted by the braiding). These two module structures are equivalent precisely when Φ is given the structure of a *braided central functor*:

DEFINITION 3.5. Let \mathcal{C} be a category and \mathcal{M} a tensor category. A *central functor*¹⁰ from \mathcal{C} to \mathcal{M} is a functor $\Phi : \mathcal{C} \rightarrow \mathcal{M}$ equipped with a factorisation

$$\Phi : \mathcal{C} \xrightarrow{\Phi^z} \mathcal{Z}(\mathcal{M}) \xrightarrow{F} \mathcal{M}$$

where F is the forgetful functor [BFO09][Ost14, §2.1]. When \mathcal{C} is monoidal, a central functor $\Phi : \mathcal{C} \rightarrow \mathcal{M}$ is called monoidal if Φ^z is monoidal. When \mathcal{C} is braided, a central functor is called braided if the functor Φ^z is braided.

Equivalently, a functor Φ is a central functor if it is equipped with the additional data of half-braidings $e_{\Phi(c),x} : \Phi(c) \otimes x \rightarrow x \otimes \Phi(c)$, natural in c and x , satisfying

$$e_{\Phi(c),x \otimes y} = (\text{id}_x \otimes e_{\Phi(c),y}) \circ (e_{\Phi(c),x} \otimes \text{id}_y) \quad (\text{Central functor}).$$

To go from a central functor to a monoidal central functor, one imposes the additional axiom

$$e_{\Phi(c \otimes d),x} = (e_{\Phi(c),x} \otimes \text{id}_{\Phi(d)}) \circ (\text{id}_{\Phi(c)} \otimes e_{\Phi(d),x}) \quad (\text{Monoidal central functor}).$$

Finally, a braided central functor satisfies the further axiom

$$e_{\Phi(c),\Phi(d)} = \Phi(\beta_{c,d}) \quad (\text{Braided central functor}).$$

Note that in formulating the above axioms we have omitted the associators and the isomorphisms $\Phi(x) \otimes \Phi(y) \rightarrow \Phi(x \otimes y)$, for readability.

Let \mathcal{C} be a braided tensor category. A braided central functor $\Phi : \mathcal{C} \rightarrow \mathcal{M}$ makes the tensor category \mathcal{M} into a \mathcal{C} -module category in way that is compatible with the tensor structure of \mathcal{M} and the braiding of \mathcal{C} . We call such an \mathcal{M} a *module tensor category* over \mathcal{C} :

¹⁰ The reader is cautioned that our usage of the term ‘central functor’ agrees with [BFO09, Ost14], but differs from [Bez04, DMNO13] as these require that a central functor be braided.

DEFINITION 3.6. If \mathcal{M} is a tensor category and \mathcal{C} is a braided category, then \mathcal{M} is a *module tensor category* over \mathcal{C} if it is equipped with a braided central functor $\Phi : \mathcal{C} \rightarrow \mathcal{M}$.

If \mathcal{M} , \mathcal{C} and Φ^z are pivotal, then we say that \mathcal{M} is a *pivotal module tensor category*.

As module tensor categories might be an unfamiliar concept, we provide a couple of alternative viewpoints:

A tensor category can be thought of as a 2-category with a single object. Similarly, a pair consisting of a tensor category \mathcal{C} and module category \mathcal{M} can be encoded by a 2-category with exactly two objects 1 and $*$, and only non-trivial homs given by $\mathcal{C} := \text{Hom}(1, 1)$ and $\mathcal{M} := \text{Hom}(1, *)$.

On the other hand, a braided tensor category can be thought of as a 3-category with one object and one 1-morphism [BD95, ‘Periodic Table’, Fig. 21]. The pair consisting of a braided category \mathcal{C} and module tensor category \mathcal{M} can be encoded by a 3-category with exactly two objects 1 and $*$, and with only non-trivial homs given by the 2-categories $\text{Hom}(1, 1)$ and $\text{Hom}(1, *)$, which are furthermore required to have only one object. We recover the braided category and its module tensor category as $\mathcal{C} = \text{Hom}_{\text{Hom}(1,1)}(1_1, 1_1)$ and $\mathcal{M} = \text{Hom}_{\text{Hom}(1,*)}(\cdot, \cdot)$, respectively, where $\cdot : 1 \rightarrow *$ is the unique morphism.

Alternatively, for those who like to think of a braided tensor category as a category which is an algebra over the little discs operad, then a pair consisting of a braided tensor category \mathcal{C} and a module tensor category \mathcal{M} is the same thing as an algebra over Voronov’s Swiss-cheese operad [Vor99].

REMARK 3.7. By [Ost03, Thm. 3.1], if \mathcal{C} is a fusion category and \mathcal{M} is an indecomposable module category, then \mathcal{M} is equivalent to $\text{Mod}_{\mathcal{C}}(a)$ for a semi-simple algebra $a \in \mathcal{C}$. This theorem applies when \mathcal{C} is braided pivotal and \mathcal{M} is a module tensor category which is indecomposable as a \mathcal{C} -module category. In this case, $\Phi : \mathcal{C} \rightarrow \mathcal{M}$ is a dominant tensor functor, so by [BN11, Prop. 6.1] there is a canonical commutative algebra $a^z \in \mathcal{Z}(\mathcal{C})$ such that Φ is tensor equivalent to the free module functor $\mathcal{C} \rightarrow \text{Mod}_{\mathcal{C}}(a^z)$.

3.3 THE CATEGORIFIED TRACE ASSOCIATED TO A MODULE TENSOR CATEGORY

We now introduce the main construction of this paper, the categorified trace associated to a module tensor category.

DEFINITION 3.8. Let \mathcal{C} be a braided pivotal category, and let \mathcal{M} be a pivotal module tensor category. The *associated categorified trace* $\text{Tr}_{\mathcal{C}} : \mathcal{M} \rightarrow \mathcal{C}$ is the right adjoint of the action functor Φ .

The existence of a right adjoint is a mild assumption, which can be easily checked in many circumstances (see, e.g., [DSPS14, Cor. 1.9]):

LEMMA 3.9. *If \mathcal{C} and \mathcal{M} are semisimple (linear over some field k) and \mathcal{C} has finitely many types of simple objects, then any linear functor $F : \mathcal{C} \rightarrow \mathcal{M}$ admits a right adjoint.*

Proof. Let $\{c_i\}$ be a basis of \mathcal{C} (representatives of the isomorphism classes of simple objects), and let $\{m_j\}$ be a basis of \mathcal{M} . Let $A = (A_{ij})$ be the matrix of finite dimensional vector spaces given by $A_{ij} := \mathcal{M}(m_j, F(c_i))$. Note that A has only finitely many non-zero entries.

We have canonical isomorphisms $F(c_i) \cong \bigoplus_j A_{ij} \otimes m_j$, and the functor F can be recovered by the formula

$$F\left(\bigoplus_i V_i \otimes c_i\right) \cong \bigoplus_{i,j} V_i \otimes A_{ij} \otimes m_j.$$

Let A^* be the ‘conjugate transpose’ matrix of vector spaces, given by $A_{ji}^* = \text{Hom}_k(A_{ij}, k)$. It is then an easy exercise to check that the functor $G\left(\bigoplus_j W_j \otimes m_j\right) := \bigoplus_{i,j} W_j \otimes A_{ji}^* \otimes c_i$ is a right adjoint (also a left adjoint) of F . \square

Alternatively one can define $\text{Tr}_{\mathcal{C}} := \underline{\text{Hom}}(1_{\mathcal{M}}, -)$, where $\underline{\text{Hom}}$ is the internal hom for module categories [Ost03]. By definition, $\underline{\text{Hom}}(1_{\mathcal{M}}, x)$ represents the functor $c \mapsto \text{Hom}(c \cdot 1_{\mathcal{M}}, x)$ and so, recalling that $\Phi(c) = c \cdot 1_{\mathcal{M}}$, we indeed have the adjunction $\text{Hom}(\Phi(c), x) \cong \text{Hom}(c, \underline{\text{Hom}}(1_{\mathcal{M}}, x))$.

In Section 4.4 we will construct traciators for $\text{Tr}_{\mathcal{C}}$ and prove that they satisfy the axioms of a categorified trace (Lemmas 4.16 and 4.17), a result which first appeared in [BFO09, Prop.5] and [Ost14, Prop.2.5].

We illustrate our construction by a couple of examples.

EXAMPLE 3.10. Any pivotal category \mathcal{M} is naturally a pivotal module tensor category over Vec . For every $x \in \mathcal{M}$ we have $\mathcal{C}(1, \text{Tr}_{\mathcal{C}}(x)) \cong \mathcal{M}(\Phi(1_{\mathcal{C}}), x) = \mathcal{M}(1_{\mathcal{M}}, x)$, the invariant vectors. This means that we can naturally identify $\text{Tr}_{\mathcal{C}}(x)$ with the vector space $\mathcal{M}(1_{\mathcal{M}}, x)$.

EXAMPLE 3.11. Using the conventions of [Shi11, §3], we let \mathcal{M} be the Tambara-Yamagami fusion category [TY98] associated to the abelian group $A = \mathbb{Z}/N\mathbb{Z}$, the bicharacter $\langle a, b \rangle := q^{ab}$, $q = \exp(2\pi i/N)$, and the sign $\tau = +N^{-1/2}$. The category \mathcal{M} has N invertible objects $1, \dots, N$, and one other simple object m satisfying the fusion rule $m \otimes m \cong \bigoplus_{a \in A} a$. By [Izu01], again in the conventions of [Shi11, §3], there are $2N$ invertible elements in $\mathcal{Z}(\mathcal{M})$, parametrized by (a, ε) where $a \in \mathbb{Z}/N\mathbb{Z}$ and ε is a square root of $\langle a, a \rangle$, which gives the half-braiding of (a, ε) with m .

Using the same bicharacter as above, we get a ribbon structure on $\mathcal{C} := \text{Vec}(\mathbb{Z}/N\mathbb{Z})$ with braiding $\beta_{a,b} = \langle a, b \rangle \text{id}_{a+b}$, twist $\theta_a = \langle a, a \rangle \text{id}_a$, and pivotal structure $\varphi_a = \text{id}_a$. The functor $\Phi^z : \mathcal{C} \rightarrow \mathcal{Z}(\mathcal{M})$ given by $a \mapsto (a, q^{a^2(N+1)/2})$ is braided pivotal, and equips the Tambara-Yamagami category with the structure of a module tensor category over \mathcal{C} . The composite $\Phi = F \circ \Phi^z$ is the forgetful functor onto the copy of $\text{Vec}(\mathbb{Z}/N\mathbb{Z})$ in \mathcal{M} , and its right adjoint $\text{Tr}_{\mathcal{C}} : \mathcal{M} \rightarrow \text{Vec}(\mathbb{Z}/N\mathbb{Z})$ is the projection from \mathcal{M} onto its subcategory $\text{Vec}(\mathbb{Z}/N\mathbb{Z})$.

EXAMPLE 3.12. Let \mathcal{C} be a braided pivotal tensor category, let $a \in \mathcal{C}$ be a commutative algebra object, and let $\mathcal{M} = \text{Mod}_{\mathcal{C}}(a)$ be the category of a -modules in \mathcal{C} . Then \mathcal{M} is both a tensor category [Par95], and a \mathcal{C} -module category [KO02, §1]. Moreover, it is a \mathcal{C} -module tensor category, as noted in [DMNO13, §3.4]. The functor $\Phi : \mathcal{C} \rightarrow \mathcal{M}$ is the *free module* functor $\Phi(x) := a \otimes x$, and the half-braidings are given by

$$e_{\Phi(x),y} : \Phi(x) \otimes_a y \cong x \otimes y \xrightarrow{\beta_{x,y}} y \otimes x \cong y \otimes_a \Phi(x).$$

The associated categorified trace $\text{Tr}_{\mathcal{C}} : \text{Mod}_{\mathcal{C}}(a) \rightarrow \mathcal{C}$ is given by the forgetful functor, so that $\text{Tr}_{\mathcal{C}}(x) = x$ for all $x \in \mathcal{M}$. Indeed, using the adjunction, we have

$$\mathcal{C}(c, \text{Tr}_{\mathcal{C}}(x)) \cong \mathcal{M}(\Phi(c), x) = \mathcal{M}(a \otimes c, x) := \mathcal{C}_a(a \otimes c, x) \cong \mathcal{C}(c, x),$$

where $\mathcal{C}_a(-, -)$ denotes the a -module maps.

When a is furthermore separable and with trivial twist, then $\text{Mod}_{\mathcal{C}}(a)$ is a pivotal module tensor category. For more details on this example, and in particular for the proof that Φ^z is a pivotal functor, we refer the reader to Appendix A.1.

SUB-EXAMPLE 3.13. Take \mathcal{C} to be $SU(2)_k$ (the semi-simplification of the category of tilting modules for Lusztig’s integral form of $U_q(\mathfrak{sl}(2))$, specialized at $q = \exp(\frac{2\pi i}{2(k+2)})$ ¹¹ – see [Saw06] for details), which is an A_{k+1} modular tensor category. The commutative algebra objects $a \in \mathcal{C}$ are completely classified [Ocn02, KO02], and yield A_n, D_{2n}, E_6 , and E_8 fusion categories, whose simple objects are in bijective correspondence with the nodes of the respective Dynkin diagrams. Combining Example 3.12 with the computation at the end of [KO02, §6], we get the following explicit description of the trace functor:

Example: D_4 . This is a module tensor category over $SU(2)_4$ (with simple objects $\mathbf{1}, \dots, \mathbf{5}$). Its unit object is $\mathbf{1}_{\mathcal{M}} = \bullet \circ \circ$, and the adjacency matrix of the Dynkin diagram encodes the operation of tensoring by $\Phi(\mathbf{2}) = \circ \circ \bullet \circ$. The trace functor is given (at the level of isomorphism classes of simple objects) by:

$x \in \mathcal{M}$					
$\text{Tr}_{\mathcal{C}}(x) \in SU(2)_4$	$\mathbf{1} \oplus \mathbf{5}$	$\mathbf{2} \oplus \mathbf{4}$	$\mathbf{3}$	$\mathbf{3}$	

Example: E_6 . This is a module tensor category over $SU(2)_{10}$ (with simple objects $\mathbf{1}, \dots, \mathbf{11}$). The unit object is $\bullet \circ \circ \circ$, and the adjacency matrix of the Dynkin diagram encodes the operation of tensoring by $\Phi(\mathbf{2}) = \circ \bullet \circ \circ \circ$. The trace functor is:

x											
$\text{Tr}_{\mathcal{C}}(x)$	$\mathbf{1} \oplus \mathbf{7}$	$\mathbf{2} \oplus \mathbf{6} \oplus \mathbf{8}$	$\mathbf{3} \oplus \mathbf{5} \oplus \mathbf{7} \oplus \mathbf{9}$	$\mathbf{4} \oplus \mathbf{8}$	$\mathbf{4} \oplus \mathbf{6} \oplus \mathbf{10}$	$\mathbf{5} \oplus \mathbf{11}$					

¹¹We use the convention according to which $[2]_q = q + q^{-1}$.

4 PROPERTIES OF THE TRACE FUNCTOR

In this section we establish all the basic properties of the categorified trace $\mathrm{Tr}_{\mathcal{C}}$ associated to a pivotal module tensor category (Definition 3.8). In Sections 4.1–4.3, we discuss the properties of $\mathrm{Tr}_{\mathcal{C}}$ which follow from the fact that it is the right adjoint of a tensor functor, notably the existence of the multiplication map $\mu_{x,y} : \mathrm{Tr}_{\mathcal{C}}(x) \otimes \mathrm{Tr}_{\mathcal{C}}(y) \rightarrow \mathrm{Tr}_{\mathcal{C}}(x \otimes y)$. The construction of the traciator $\tau_{x,y} : \mathrm{Tr}_{\mathcal{C}}(x \otimes y) \rightarrow \mathrm{Tr}_{\mathcal{C}}(y \otimes x)$ is done in Section 4.4. This only depends on the fact that \mathcal{M} is pivotal and that Φ factors through $\mathcal{Z}(\mathcal{M})$. Finally, in Section 4.5 we assume our full set of hypothesis: that \mathcal{M} is a pivotal module tensor category over \mathcal{C} . In this context we establish various compatibility relations which combine the traciator and multiplication maps with the braiding and the twist of \mathcal{C} .

In terms of Figure 1 from the introduction, Sections 4.1–4.3 deal with the right column, Section 4.4 deals with the left column, and Section 4.5 deals with the middle column.

4.1 THE ATTACHING MAP

Adjoint of tensor functors have been studied systematically [Kel74] and arise in many contexts (e.g. [BN11], [DMNO13]). We will reprove several known properties as a way of introducing our graphical notation for $\mathrm{Tr}_{\mathcal{C}}$. For Sections 4.1–4.3, we fix the following notation:

NOTATION 4.1.

- \mathcal{C}, \mathcal{M} are tensor categories,
- $\Phi : \mathcal{C} \rightarrow \mathcal{M}$ is a tensor functor, and
- $\mathrm{Tr}_{\mathcal{C}} : \mathcal{M} \rightarrow \mathcal{C}$ is the right adjoint of Φ .


The reader is cautioned that under these limited assumptions (compare Notation 4.12) the functor $\mathrm{Tr}_{\mathcal{C}}$ may fail to be a categorified trace. Nevertheless, we shall use the notation $\mathrm{Tr}_{\mathcal{C}}$ for notational consistency.

Recall from Section 3.1 that we depict an object $x \in \mathcal{M}$ by a strand in the plane, and $\mathrm{Tr}_{\mathcal{C}}(x)$ by a cylinder with an x -strand on its surface. Given an object $c \in \mathcal{C}$, the adjunction $\mathcal{C}(c, \mathrm{Tr}_{\mathcal{C}}(x)) \cong \mathcal{M}(\Phi(c), x)$ is represented diagrammatically by:


$$\begin{array}{ccc}
 \begin{array}{c} \text{Cylinder with } x \text{-strand} \\ \text{on box } g \\ \text{with } c \text{-strand} \end{array} \in \mathcal{C}(c, \mathrm{Tr}_{\mathcal{C}}(x)) & \longleftrightarrow & \begin{array}{c} \text{Green strand } c \\ \text{over red strand } x \\ \text{on box } g \end{array} \in \mathcal{M}(\Phi(c), x)
 \end{array}
 \tag{6}$$

Visually, we can think of the above adjunction as the operation of opening the tube (like an umbrella) and then flattening it to a plane.

DEFINITION 4.2 (the attaching map ε). Given $x \in \mathcal{M}$, we define $\varepsilon_x : \Phi(\mathrm{Tr}_{\mathcal{C}}(x)) \rightarrow x$ to be the image of $\mathrm{id}_{\mathrm{Tr}_{\mathcal{C}}(x)}$ under the correspondence (6). Equivalently, $\varepsilon : \Phi \circ \mathrm{Tr}_{\mathcal{C}} \rightarrow \mathrm{id}_{\mathcal{M}}$ is the counit of the adjunction $\Phi \dashv \mathrm{Tr}_{\mathcal{C}}$. We represent the morphism ε_x as follows:



A possibly better graphical notation for ε_x is that of a tube coming out of the plane:



but in order to show that this graphical depiction of ε_x is valid, we would first need to prove properties such as (8), or (13). The next two lemmas are general results that hold for any adjunction.


LEMMA 4.3. Let $a \in \mathcal{C}$ and $x \in \mathcal{M}$. Then, under the adjunction $\mathcal{C}(a, \mathrm{Tr}_{\mathcal{C}}(x)) \cong \mathcal{M}(\Phi(a), x)$, a morphism $f \in \mathcal{C}(a, \mathrm{Tr}_{\mathcal{C}}(x))$ corresponds to $\varepsilon_x \circ \Phi(f) \in \mathcal{M}(\Phi(a), x)$.

Proof. Since the adjunction is natural, the following diagram commutes:

$$\begin{array}{ccccccc}
 \varepsilon_x & \in & \mathcal{M}(\Phi(\mathrm{Tr}_{\mathcal{C}}(x)), x) & \cong & \mathcal{C}(\mathrm{Tr}_{\mathcal{C}}(x), \mathrm{Tr}_{\mathcal{C}}(x)) & \ni & \mathrm{id}_{\mathrm{Tr}_{\mathcal{C}}(x)}, \\
 \downarrow & & \Phi(f)^* \downarrow & & f^* \downarrow & & \downarrow \\
 \varepsilon_x \circ \Phi(f) & \in & \mathcal{M}(\Phi(a), x) & \cong & \mathcal{C}(a, \mathrm{Tr}_{\mathcal{C}}(x)) & \ni & f
 \end{array}$$

where $f^*(g) = g \circ f$. □


LEMMA 4.4. Suppose $f \in \mathcal{M}(x, y)$. Then $\varepsilon_y \circ \Phi(\mathrm{Tr}_{\mathcal{C}}(f)) = f \circ \varepsilon_x$, i.e.,



Proof. The counit of an adjunction is always a natural transformation [ML98, §IV.1]. □

4.2 THE MULTIPLICATION MAP


DEFINITION 4.5. For $x, y \in \mathcal{M}$, there is a canonical map $\mu_{x,y} : \text{Tr}_{\mathcal{C}}(x) \otimes \text{Tr}_{\mathcal{C}}(y) \rightarrow \text{Tr}_{\mathcal{C}}(x \otimes y)$ given as the mate of



under the isomorphisms


$$\begin{aligned} \mathcal{M}(\Phi(\text{Tr}_{\mathcal{C}}(x)) \otimes \Phi(\text{Tr}_{\mathcal{C}}(y)), x \otimes y) &\cong \mathcal{M}(\Phi(\text{Tr}_{\mathcal{C}}(x) \otimes \text{Tr}_{\mathcal{C}}(y)), x \otimes y) \\ &\cong \mathcal{C}(\text{Tr}_{\mathcal{C}}(x) \otimes \text{Tr}_{\mathcal{C}}(y), \text{Tr}_{\mathcal{C}}(x \otimes y)), \end{aligned}$$

where we used that Φ is a tensor functor. Diagrammatically, we denote μ by a pair of pants:



Our next task will be to show that μ is associative. We begin by showing that it is compatible with the attaching map ε :


LEMMA 4.6. For any $x, y \in \mathcal{M}$, we have $\varepsilon_{x \otimes y} \circ \Phi(\mu_{x,y}) = \varepsilon_x \otimes \varepsilon_y$.



Proof. By definition, $\varepsilon_x \otimes \varepsilon_y$ is the mate of $\mu_{x,y}$ under the adjunction. Now apply Lemma 4.3 to see that $\mu_{x,y}$ is also the mate of $\varepsilon_{x \otimes y} \circ \Phi(\mu_{x,y})$. \square


As a corollary, we see $\mu_{x,y}$ that is natural in x and y :

COROLLARY 4.7. For all $x, y, z \in \mathcal{M}$ and all $f : y \rightarrow z$ in \mathcal{M} , we have $\mu_{x,z} \circ (\text{id}_{\text{Tr}_{\mathcal{C}}(x)} \otimes \text{Tr}_{\mathcal{C}}(f)) = \text{Tr}_{\mathcal{C}}(\text{id}_x \otimes f) \circ \mu_{x,y}$. In diagrams,



Similarly, for $g : w \rightarrow x$, we have $\mu_{x,y} \circ (\text{Tr}_C(g) \otimes \text{id}_{\text{Tr}_C(y)}) = \text{Tr}_C(g \otimes \text{id}_y) \circ \mu_{w,y}$.

Proof. We only show the first statement; the other is similar. By taking mates, and using Lemma 4.3, Equation (9) becomes $\varepsilon \circ \Phi(\mu_{x,z} \circ (\text{id}_{\text{Tr}_C(x)} \otimes \text{Tr}_C(f))) = \varepsilon \circ \Phi(\text{Tr}_C(\text{id}_x \otimes f) \circ \mu_{x,y})$




which follows readily from Lemmas 4.6 and 4.4, along with the fact that Φ is a tensor functor. \square


LEMMA 4.8. *The multiplication map μ is associative, i.e., the following diagram commutes:*

$$\begin{array}{ccc} \text{Tr}_C(x) \otimes \text{Tr}_C(y) \otimes \text{Tr}_C(z) & \xrightarrow{\text{id}_{\text{Tr}_C(x)} \otimes \mu} & \text{Tr}_C(x) \otimes \text{Tr}_C(y \otimes z) \\ \mu \otimes \text{id}_{\text{Tr}_C(z)} \downarrow & & \downarrow \mu \\ \text{Tr}_C(x \otimes y) \otimes \text{Tr}_C(z) & \xrightarrow{\mu} & \text{Tr}_C(x \otimes y \otimes z) \end{array}$$

In diagrams,



Proof. We claim that, under the adjunction, each map individually is the mate of



We only prove this for the right hand side of Equation (10) (the left hand side is similar). The mate of the right hand side is given by

$$\begin{aligned} \varepsilon_{x \otimes y \otimes z} \circ \Phi(\mu_{x,y \otimes z} \circ (\text{id}_{\text{Tr}_C(x)} \otimes \mu_{y,z})) &= \varepsilon_{x \otimes y \otimes z} \circ \Phi(\mu_{x,y \otimes z}) \circ \Phi(\text{id}_{\text{Tr}_C(x)} \otimes \mu_{y,z}) \\ &= (\varepsilon_x \otimes \varepsilon_{y \otimes z}) \circ (\Phi(\text{id}_{\text{Tr}_C(x)}) \otimes \Phi(\mu_{y,z})) = \varepsilon_x \otimes (\varepsilon_{y \otimes z} \circ \Phi(\mu_{y,z})) \\ &= \varepsilon_x \otimes (\varepsilon_y \otimes \varepsilon_z), \end{aligned}$$

where the first equality is the functoriality of Φ , the second equality is Lemma 4.6 and the fact that Φ is a tensor functor, and the fourth equality is again Lemma 4.6. \square

4.3 THE UNIT OF THE ADJUNCTION


In Section 4.1 we studied the attaching map ε , which is the counit of the adjunction $\Phi \dashv \text{Tr}_{\mathcal{C}}$. We now study the unit of the adjunction, which we denote η .

Thus, for any object $c \in \mathcal{C}$, we have a morphism $\eta_c : c \rightarrow \text{Tr}_{\mathcal{C}}(\Phi(c))$. We will also write $i : 1_{\mathcal{C}} \rightarrow \text{Tr}_{\mathcal{C}}(1_{\mathcal{M}})$ for the unit η evaluated on the unit object $1_{\mathcal{C}} \in \mathcal{C}$. We represent η and i by the following diagrams:




Since we have an adjunction $\Phi \dashv \text{Tr}_{\mathcal{C}}$, the counit ε and the unit η interact in the following way:

- (1) For $x \in \mathcal{M}$, the mate of ε_x is $\text{id}_{\text{Tr}_{\mathcal{C}}(x)}$. Hence $\text{Tr}_{\mathcal{C}}(\varepsilon_x) \circ \eta_{\text{Tr}_{\mathcal{C}}(x)} = \text{id}_{\text{Tr}_{\mathcal{C}}(x)}$. We represent this diagrammatically by the following relation.



- (2) For $c \in \mathcal{C}$, the mate of η_c is $\text{id}_{\Phi(c)}$. So by Lemma 4.3, $\varepsilon_{\Phi(c)} \circ \Phi(\eta_c) = \text{id}_{\Phi(c)}$. We represent this diagrammatically by the following relation.




The following lemmas demonstrate how η and i interact with the graphical calculus introduced so far.

LEMMA 4.9. *The following diagram commutes:*

$$\begin{array}{ccccc}
 1_{\mathcal{C}} \otimes \text{Tr}_{\mathcal{C}}(x) & \xrightarrow{\quad} & \text{Tr}_{\mathcal{C}}(x) & \xleftarrow{\quad} & \text{Tr}_{\mathcal{C}}(x) \otimes 1_{\mathcal{C}} \\
 \downarrow i \otimes \text{id}_{\text{Tr}_{\mathcal{C}}(x)} & & \downarrow \text{id}_{\text{Tr}_{\mathcal{C}}(x)} & & \downarrow \text{id}_{\text{Tr}_{\mathcal{C}}(x)} \otimes i \\
 \text{Tr}_{\mathcal{C}}(1) \otimes \text{Tr}_{\mathcal{C}}(x) & \xrightarrow{\mu} & \text{Tr}_{\mathcal{C}}(1 \otimes x) & \xrightarrow{\quad} & \text{Tr}_{\mathcal{C}}(x \otimes 1) \xleftarrow{\mu} \text{Tr}_{\mathcal{C}}(x) \otimes \text{Tr}_{\mathcal{C}}(1)
 \end{array}$$

In pictures, this is




Proof. We check that the left square commutes upon taking mates (the right one is similar). Using Lemma 4.3, we compute:

$$\begin{aligned} \varepsilon_x \circ \Phi(\mu_{1,x} \circ (i \otimes \text{id}_{\text{Tr}_C(x)})) &= \varepsilon_x \circ \Phi(\mu_{1,x}) \circ \Phi(i \otimes \text{id}_{\text{Tr}_C(x)}) \\ &= (\varepsilon_1 \otimes \varepsilon_x) \circ (\Phi(i) \otimes \Phi(\text{id}_{\text{Tr}_C(x)})) = (\varepsilon_1 \circ \Phi(i)) \otimes \varepsilon_x = \varepsilon_x. \end{aligned}$$


We have used Lemma 4.6 for the second equality, and Equation (12) for the last one. \square

Note that a lot of what we have done so far is to reprove in our special case the well known fact that the adjoint of a tensor functor is a lax tensor functor [Kel74]. The data is provided by the natural transformations μ and η , and the axioms are verified in Lemmas 4.8 and 4.9.

LEMMA 4.10. For $c, d \in \mathcal{C}$, we have $\eta_{c \otimes d} = \mu_{\Phi(c), \Phi(d)} \circ (\eta_c \otimes \eta_d)$. In diagrams:



Proof. By Equation (12), we know that $\Phi(\eta_{c \otimes d}) \circ \varepsilon_{\Phi(c \otimes d)} = \text{id}_{\Phi(c \otimes d)}$, so it suffices to show that the mate of the right hand side is equal to $\text{id}_{\Phi(c \otimes d)}$. The result now follows from Lemma 4.6 together with two applications of the relation (12).



LEMMA 4.11. *The following two maps $\mathrm{Tr}_{\mathcal{C}}(x) \otimes c \rightarrow \mathrm{Tr}_{\mathcal{C}}(x \otimes \Phi(c))$ are equal:*

$$\text{Diagram 1} = \text{Diagram 2} \tag{13}$$

Moreover, when \mathcal{C} is rigid, the above map (13) is invertible.

Proof. To show that the two sides of (13) are equal, apply Lemma 4.10 to the left hand side and use Equation (11).

If we assume that c is dualizable, then we can write down an inverse to the above map:

$$\text{Diagram} : \mathrm{Tr}_{\mathcal{C}}(x \otimes \Phi(c)) \rightarrow \mathrm{Tr}_{\mathcal{C}}(x) \otimes c$$

In formulas, this is:

$$[(\mathrm{Tr}_{\mathcal{C}}(\mathrm{ev}_{\Phi(*c)}) \circ \mu_{x \otimes \Phi(c), \Phi(*c)} \circ (\mathrm{id}_{\mathrm{Tr}_{\mathcal{C}}(x \otimes \Phi(c))} \otimes \eta_{*c})) \otimes \mathrm{id}_c] \circ (\mathrm{id}_{\mathrm{Tr}_{\mathcal{C}}(x \otimes \Phi(c))} \otimes \mathrm{coev}_{*c}),$$

where we have used the canonical identification $\Phi(*c) \cong * \Phi(c)$ provided by the fact that Φ is a tensor functor. The two maps compose to the identity by a straightforward application of Lemma 4.8 (the associativity of μ), Lemma 4.10, and the naturality of η . \square

4.4 CONSTRUCTION OF THE TRACIATOR

So far we have constructed the multiplication $\mu_{x,y} : \mathrm{Tr}_{\mathcal{C}}(x) \otimes \mathrm{Tr}_{\mathcal{C}}(y) \rightarrow \mathrm{Tr}_{\mathcal{C}}(x \otimes y)$ and the corresponding unit map under the assumption that \mathcal{C} and \mathcal{M} are tensor categories and that $\Phi : \mathcal{C} \rightarrow \mathcal{M}$ is a tensor functor. However, the construction of the traciator, along with the proof that it equips $\mathrm{Tr}_{\mathcal{C}}$ with the structure of a categorified trace, only depends on $\Phi : \mathcal{C} \rightarrow \mathcal{M}$ factoring through $\mathcal{Z}(\mathcal{M})$, and not on it being a tensor functor. The material of this section can be found in [BFO09, Prop. 5] and [Ost14, Prop. 2.5].

We fix the following notation for Section 4.4.

NOTATION 4.12.

- \mathcal{C} is a category


- \mathcal{M} is a pivotal category,
- $\Phi^{\mathcal{Z}} : \mathcal{C} \rightarrow \mathcal{Z}(\mathcal{M})$ is a functor,
- the trace functor $\text{Tr}_{\mathcal{C}} : \mathcal{M} \rightarrow \mathcal{C}$ is the right adjoint of $\Phi := F \circ \Phi^{\mathcal{Z}}$.

DEFINITION 4.13. For $x, y \in \mathcal{M}$, we define the traciator $\tau_{x,y} : \text{Tr}_{\mathcal{C}}(x \otimes y) \rightarrow \text{Tr}_{\mathcal{C}}(y \otimes x)$ in the following way. Let $c = \text{Tr}_{\mathcal{C}}(x \otimes y)$, and let us write $\tilde{\text{ev}}_y : y \otimes y^* \rightarrow 1$ for the composite $\text{ev}_{y^*} \circ (\varphi_y \otimes \text{id}_{y^*})$, with $\varphi_y : y \rightarrow y^{**}$ the pivotal structure. Then $\tau_{x,y}$ is the mate of

$$(\text{id}_{y \otimes x} \otimes \tilde{\text{ev}}_y) \circ (\text{id}_y \otimes \varepsilon_{x \otimes y} \otimes \text{id}_{y^*}) \circ (e_{\Phi(c),y} \otimes \text{id}_{y^*}) \circ (\text{id}_{\Phi(c)} \otimes \text{coev}_y)$$

under the adjunction $\mathcal{M}(\Phi(\text{Tr}_{\mathcal{C}}(x \otimes y)), y \otimes x) \cong \mathcal{C}(\text{Tr}_{\mathcal{C}}(x \otimes y), \text{Tr}_{\mathcal{C}}(y \otimes x))$.

As we will see shortly, the traciator is always invertible. We will sometimes write $\tau_{x,y}^+$ for $\tau_{x,y}$, and $\tau_{x,y}^-$ for $\tau_{y,x}^{-1}$. In terms of 3-dimensional diagrams, the traciators $\tau^{\pm} : \text{Tr}_{\mathcal{C}}(x \otimes y) \rightarrow \text{Tr}_{\mathcal{C}}(y \otimes x)$ are depicted




LEMMA 4.14. The traciator $\tau_{y,x}$ is invertible, and its inverse $\tau_{x,y}^-$ is the mate of

$$\in \mathcal{M}(\Phi(\text{Tr}_{\mathcal{C}}(x \otimes y)), y \otimes x) \tag{14}$$

Proof. Let us define $\tilde{\tau}_{x,y}^-$ to be the mate of (14) and let us agree, for the purpose of this proof, to reserve the graphical notation for $\tilde{\tau}_{x,y}^-$. We need to show that $\tilde{\tau}_{x,y}^- = \tau_{x,y}^-$. Equivalently, we need to show that the following two equations hold:



$$\tau_{y,x} \circ \tilde{\tau}_{x,y}^- = \text{id}_{\text{Tr}_{\mathcal{C}}(x \otimes y)} = \tilde{\tau}_{y,x}^- \circ \tau_{x,y} : \quad \text{Cylinder 1} = \text{Cylinder 2} = \text{Cylinder 3}$$

We only treat the first one (the other is entirely similar): upon taking mates, we get



where we have used Lemma 4.3 for the first two equalities. The last equal sign follows from the zig-zag equations satisfied by the (co)evaluation morphisms ev , $\bar{\text{ev}}$, coev , and $\bar{\text{coev}}$. \square


By Lemmas 4.14 and 4.3, the traciators satisfy



REMARK 4.15. In the event that \mathcal{M} is rigid but not pivotal, we only get isomorphisms $\tau^+ : \text{Tr}_{\mathcal{C}}(x \otimes y) \rightarrow \text{Tr}_{\mathcal{C}}(**y \otimes x)$ and $\tau^- : \text{Tr}_{\mathcal{C}}(x \otimes y) \rightarrow \text{Tr}_{\mathcal{C}}(y \otimes x^{**})$. All of our theorems can be generalised to this more general situation, provided double duals are inserted in the appropriate places.

We now prove the naturality of the traciator, and that it satisfies the axiom of a categorified trace.

LEMMA 4.16. *The two maps $\text{Tr}_{\mathcal{C}}(x \otimes y \otimes z) \rightarrow \text{Tr}_{\mathcal{C}}(y \otimes z \otimes x)$ given by $\tau_{x,y \otimes z}$ and by $\tau_{z \otimes x,y} \circ \tau_{x \otimes y,z}$ are equal. In diagrams:*




Similarly, we have $\tau_{x \otimes y, z}^- = \tau_{y, z \otimes x}^- \circ \tau_{x, y \otimes z}^-$.

Proof. The first statement is equivalent, upon taking mates, to the equation

$$\varepsilon_{y \otimes z \otimes x} \circ \Phi(\tau_{z \otimes x, y} \circ \tau_{x \otimes y, z}) = \varepsilon_{y \otimes z \otimes x} \circ \Phi(\tau_{x, y \otimes z}).$$

The argument is similar to the one in the previous lemma:



Here, we have used Equation (15.a) for the first two equalities, and we have used Equation (15.a) along with the identities

$$\begin{aligned} \tilde{e}v_{y \otimes z} &= \tilde{e}v_y \circ (\text{id}_y \otimes \tilde{e}v_z \otimes \text{id}_{y^*}) \\ \text{coev}_{y \otimes z} &= (\text{id}_y \otimes \text{coev}_z \otimes \text{id}_{y^*}) \circ \text{coev}_y \end{aligned}$$


for the last equality. The second statement follows from the first one by taking inverses. \square

Naturality of the traciator is given by the following lemma.


LEMMA 4.17. For $f \in \mathcal{M}(x, w)$ and $g \in \mathcal{M}(y, z)$, we have $\tau_{w, z} \circ \text{Tr}_C(f \otimes g) = \text{Tr}_C(g \otimes f) \circ \tau_{x, y}$, and similarly for τ^- . In diagrams:

(16)

Proof. By Equation (15.a) and Lemma 4.4, the two sides of (16) are mates of



and



respectively. The latter are equal by pivotality in \mathcal{M} . □

4.5 INTERACTION BETWEEN TRACIATOR AND BRAIDING

In Sections 4.1–4.3, we saw that when $\text{Tr}_{\mathcal{C}}$ is the adjoint of a tensor functor, it can be described by a graphical calculus of strings on tubes. The tubes are allowed to split, but the strings must remain on the fronts of the tubes. In Section 4.4, we also saw that when Φ is a central functor (not assumed monoidal) $\text{Tr}_{\mathcal{C}}$ admits a graphical calculus of strings winding around a single tube.

In this section, we start by assuming that Φ is a monoidal central functor. The tubes, with the strings on their surface are now allowed to branch, but they must remain in a single plane. Finally, we go on to assume our strongest hypothesis, namely that Φ is a braided central functor. The tubes can now braid freely in three dimensions (see Figure 1 in the introduction).

We begin with the following assumptions:

NOTATION 4.18.

- \mathcal{C} and \mathcal{M} are pivotal categories
- $\Phi^{\mathcal{Z}} : \mathcal{C} \rightarrow \mathcal{Z}(\mathcal{M})$ is a pivotal tensor functor,
- the trace functor $\text{Tr}_{\mathcal{C}} : \mathcal{M} \rightarrow \mathcal{C}$ is the right adjoint of $\Phi := F \circ \Phi^{\mathcal{Z}}$.

Under these assumptions, we can establish a generalization of the associativity of μ :


LEMMA 4.19. *Given $w, x, y, z \in \mathcal{M}$, the following two morphisms in*

$\mathcal{C}(\text{Tr}_{\mathcal{C}}(w) \otimes \text{Tr}_{\mathcal{C}}(x \otimes y) \otimes \text{Tr}_{\mathcal{C}}(z), \text{Tr}_{\mathcal{C}}(w \otimes x \otimes z \otimes y))$ are equal:

(17)


REMARK 4.20. In the above equation, it is best to imagine the blue strands as running along the backs of the tubes: pre- and postcomposing by the “half-traciators” and , Equation (17) becomes

Proof. We start from the left hand side, whose mate is given by



By successive application of Equations (15.b), (8), (15.a), and then once again

(8), we can rewrite this as:




(18)


Starting instead from the right hand side of (17), if we take its mate as above, and apply (8), then (15.b), then (8), and finally (15.a), we obtain the same picture (18). \square

We now prove a compatibility between the traciator and the unit η .


LEMMA 4.21. For $x \in \mathcal{M}$ and $c \in \mathcal{C}$, the following diagram commutes:




In diagrams,



Proof. We only prove that the left hand diagram commutes (the other one is similar). The mate of the leftmost map can be simplified in the following way:



where we have used Equation (15.a), then Lemma 4.4 (naturality of ε), and finally Equation (12). It agrees with the mate of the middle map:




We now assume that \mathcal{C} and $\Phi^{\mathcal{Z}}$ are braided, so that $\text{Tr}_{\mathcal{C}}$ is the categorified trace associated to a pivotal module tensor category. In this context, we establish certain relations between $\text{Tr}_{\mathcal{C}}$ and the braiding and twist of \mathcal{C} . We begin by examining the traciator and the twist.


LEMMA 4.22. *Let $c \in \mathcal{C}$ and $x \in \mathcal{M}$ be objects. Then for every $f \in \mathcal{M}(\Phi(c), x)$, we have*

$$f \circ \Phi(\theta_c) = (\text{id}_x \otimes \tilde{e}v_x) \circ (\text{id}_x \otimes f \otimes \text{id}_{x^*}) \circ (e_{\Phi(c),x} \otimes \text{id}_{x^*}) \circ (\text{id}_{\Phi(c)} \otimes \text{coev}_x),$$

where $\tilde{e}v_x : x \otimes x^* \rightarrow 1$ is as in Definition 4.13. In diagrams:




Proof. Starting from the right hand side, we bring f under the crossing using pivotality:



To conclude the argument, note that $(\text{id}_{\Phi(c)} \otimes \tilde{e}v_{\Phi(c)}) \circ (e_{\Phi(c),\Phi(c)} \otimes \text{id}_{\Phi(c)^*}) \circ (\text{id}_{\Phi(c)} \otimes \text{coev}_{\Phi(c)})$ is the image of $(\text{id}_c \otimes \tilde{e}v_c) \circ (\beta_{c,c} \otimes \text{id}_{c^*}) \circ (\text{id}_c \otimes \text{coev}_c) = \theta_c$ under $\Phi = F \circ \Phi^{\mathcal{Z}}$, because $\Phi^{\mathcal{Z}} : \mathcal{C} \rightarrow \mathcal{Z}(\mathcal{M})$ is a balanced functor (by Lemma A.6). □

PROPOSITION 4.23. *The map $\tau_{1_{\mathcal{M}},x} : \text{Tr}_{\mathcal{C}}(1_{\mathcal{M}} \otimes x) \rightarrow \text{Tr}_{\mathcal{C}}(x \otimes 1_{\mathcal{M}})$ is equal to the twist $\theta_{\text{Tr}_{\mathcal{C}}(x)}$. In diagrams:*



Similarly, we have $\theta_{\text{Tr}_C(x)}^{-1} = \tau_{x,1\mathcal{M}}^-$.

Proof. Taking $c = \text{Tr}_C(x)$ and $f = \varepsilon_x$ in the previous lemma, and using Equation (15.a), we get $\varepsilon_x \circ \Phi(\theta_{\text{Tr}_C(x)}) = \varepsilon_x \circ \Phi(\tau_{1\mathcal{M},x})$, which is the mate of our equation. \square

COROLLARY 4.24. *For $x, y \in \mathcal{M}$, the map $\theta_{\text{Tr}_C(x \otimes y)}$ is equal to $\tau_{y,x} \circ \tau_{x,y} : \text{Tr}_C(x \otimes y) \rightarrow \text{Tr}_C(x \otimes y)$.*


Proof. By Lemma 4.16, $\tau_{y,x} \circ \tau_{x,y}$ is equal to $\tau_{x \otimes y, 1\mathcal{M}} : 1\mathcal{M} \otimes x \otimes y \rightarrow x \otimes y \otimes 1\mathcal{M}$ (modulo unitors and associators which we suppress). The latter is equal to $\theta_{\text{Tr}_C(x \otimes y)}$ by the previous proposition. \square

We now establish the relationship between the traciator, the braiding and the twist.

LEMMA 4.25. *Given morphisms $f : \Phi(c) \rightarrow x$ and $g : \Phi(d) \rightarrow y$ in \mathcal{M} , the following equation holds:*

$$(\text{id}_{x \otimes y} \otimes \bar{e}v_x) \circ (\text{id}_x \otimes g \otimes f \otimes \text{id}_{x^*}) \circ e_{\Phi(d \otimes c), x} \circ (\text{id}_{\Phi(d \otimes c)} \otimes \text{coev}_x) = (f \otimes g) \circ \Phi(\beta_{d,c} \circ (\text{id}_d \otimes \theta_c)).$$

In diagrams:




Proof. This follows from Lemma 4.22 applied to f , together with passing the morphism f under the d strand. \square

LEMMA 4.26. *Given $x, y \in \mathcal{M}$, we have*

$$\tau_{x,y} \circ \mu_{x,y} = \mu_{y,x} \circ \beta_{\text{Tr}_C(x), \text{Tr}_C(y)} \circ (\text{id}_{\text{Tr}_C(x)} \otimes \theta_{\text{Tr}_C(y)}).$$

In diagrams:



Similarly, we have $\tau_{x,y}^- \circ \mu_{x,y} = \mu_{y,x} \circ \beta_{\text{Tr}_C(x), \text{Tr}_C(y)}^- \circ (\theta_{\text{Tr}_C(x)}^{-1} \otimes \text{id}_{\text{Tr}_C(y)})$.

Proof. We only prove the first claim. By taking mates, it is equivalent to the equation

where we have used (15.a) and Lemma 4.6 for the left hand side, and Lemma 4.6 for the right hand side. Equation (19) now follows from Proposition 4.23 and Lemma 4.25. \square

5 TRACES OF ALGEBRAS AND SEMISIMPLICITY

We have seen in Section 4 that our categorified trace is a lax monoidal functor. As a result, it takes algebra objects to algebra objects. In fact, given two algebras A and B , we will see that $\text{Tr}_C(A \otimes B)$ is also naturally equipped with an algebra structure. What is more, under mild additional assumptions, we will see that $\text{Tr}_C(A \otimes B)$ is semisimple when both A and B are. We begin by investigating the semisimplicity of $\text{Tr}_C(A)$ under weaker assumptions than those necessary to study $\text{Tr}_C(A \otimes B)$.

5.1 THE ALGEBRA $\text{Tr}_C(A)$

If $A = (A, m_A, i_A)$ is an algebra object in \mathcal{M} , then $\text{Tr}_C(A)$ is an algebra object in \mathcal{C} , where the multiplication and unit maps are given by:

$$m_{\text{Tr}_C(A)} := \text{Tr}_C(m_A) \circ \mu_{A,A} = \text{diagram of a cup with two strands} \quad i_{\text{Tr}_C(A)} := \text{Tr}_C(i_A) \circ i = \text{diagram of a cylinder with a red strand}$$

Recall that an algebra object A in a semisimple tensor category \mathcal{M} is called semisimple if its category of A -modules $\text{Mod}_{\mathcal{M}}(A)$ is semisimple. The following is the main result of this subsection:

THEOREM 5.1. *Let \mathcal{C} and \mathcal{M} be rigid semisimple tensor categories. Let $\Phi : \mathcal{C} \rightarrow \mathcal{M}$ be a tensor functor, with right adjoint Tr_C . If A is a semisimple algebra object in \mathcal{M} , then $\text{Tr}_C(A)$ is a semisimple algebra object in \mathcal{C} .*

Proof. Let us write B for $\text{Tr}_C(A)$. We prove the theorem by finding an equivalent, manifestly semisimple, description of the category $\text{Mod}_{\mathcal{C}}(B)$ of B -modules. Let \mathcal{N}_0 be the essential image of the functor $\mathcal{C} \rightarrow \text{Mod}_{\mathcal{M}}(A) : c \mapsto A \otimes \Phi(c)$, and let \mathcal{N} be the idempotent completion of \mathcal{N}_0 . The category \mathcal{N} can equivalently be

described as the full subcategory of $\text{Mod}_{\mathcal{M}}(A)$ generated by the simple objects that occur as summands of $A \otimes \Phi(c)$ for $c \in \mathcal{C}$. The trace functor induces a functor

$$\text{Tr}_{\mathcal{C}}^{\text{Mod}(A)} : \text{Mod}_{\mathcal{M}}(A) \rightarrow \text{Mod}_{\mathcal{C}}(B),$$

which in turn restricts to a functor

$$\mathcal{N} \hookrightarrow \text{Mod}_{\mathcal{M}}(A) \xrightarrow{\text{Tr}_{\mathcal{C}}^{\text{Mod}(A)}} \text{Mod}_{\mathcal{C}}(B).$$

We will prove that the latter is an equivalence of categories $\mathcal{N} \cong \text{Mod}_{\mathcal{C}}(B)$. Since \mathcal{N} is semisimple, this will immediately imply the theorem.

- *The functor $\text{Tr}_{\mathcal{C}}^{\text{Mod}(A)}|_{\mathcal{N}}$ is fully faithful.* It suffices to show that the restriction of $\text{Tr}_{\mathcal{C}}^{\text{Mod}(A)}$ to \mathcal{N}_0 is fully faithful. By definition, any object of \mathcal{N}_0 is of the form $A \otimes \Phi(c)$ for some $c \in \mathcal{C}$. We must prove that the map

$$\text{Tr}_{\mathcal{C}}^{\text{Mod}(A)} : \mathcal{N}_0(A \otimes \Phi(c), A \otimes \Phi(d)) \rightarrow \mathcal{C}_B(\text{Tr}_{\mathcal{C}}(A \otimes \Phi(c)), \text{Tr}_{\mathcal{C}}(A \otimes \Phi(d)))$$

given by

is an isomorphism.

We claim that its inverse is provided by

$$g \mapsto (m_A \otimes \text{id}_{\Phi(d)}) \circ (\text{id}_A \otimes [\varepsilon_{A \otimes \Phi(d)} \circ \Phi(g \circ \text{Tr}_{\mathcal{C}}(i_A \otimes \text{id}_{\Phi(c)}) \circ \eta_c)])$$

We show that (20) and (21) compose both ways to the identity. One direction

goes as follows:

$$\begin{aligned}
 ((21) \circ (20))(f) &= \text{Diagram 1} \\
 &= \text{Diagram 2} \\
 &= \text{Diagram 3} \\
 &= \text{Diagram 4} = f.
 \end{aligned}$$

The second equality holds by the naturality of ε , the third equality is Equation (12), and the fourth one follows from f being an A -module map.

For the other direction, it is convenient to precompose both sides of the equation by the isomorphism

$$\text{Diagram} : B \otimes c \rightarrow \text{Tr}_c(A \otimes \Phi(c)) \tag{22}$$


provided by Lemma 4.11:

$$\begin{aligned}
 ((20) \circ (21))(g) \circ (22) &= \text{[Diagram 1]} = \text{[Diagram 2]} \\
 &= \text{[Diagram 3]} = \text{[Diagram 4]} \\
 &= \text{[Diagram 5]} = \text{[Diagram 6]} = g \circ (22).
 \end{aligned}$$

Here, the second and third equations follows by the naturality of μ and η . The fourth equality is the content of Equation (11), and the fifth one holds because g is a B -module map.

- The functor $\text{Tr}_{\mathcal{C}}^{\text{Mod}(A)}|_{\mathcal{N}}$ is essentially surjective. Every B -module $c \in \mathcal{C}$ fits

in a coequalizer diagram $B \otimes B \otimes c \rightrightarrows B \otimes c \rightarrow c$:



Using the isomorphism (22), we see that $B \otimes B \otimes c$ and $B \otimes c$ are in the essential image of \mathcal{N} . We are now finished by Lemma 5.2 below. \square

LEMMA 5.2. *Suppose $\mathcal{N}_1, \mathcal{N}_2$ are linear categories with \mathcal{N}_1 semisimple. Let $T : \mathcal{N}_1 \rightarrow \mathcal{N}_2$ be a fully faithful linear functor. Assume that every $b \in \mathcal{N}_2$ sits in a coequalizer diagram $T(a_1) \rightrightarrows T(a_2) \rightarrow b$. Then T is essentially surjective (and thus an equivalence of categories).*

Proof. We may replace the coequalizer diagram by an exact sequence

$$T(a_1) \xrightarrow{f} T(a_2) \rightarrow b \rightarrow 0.$$

Let $g : a_1 \rightarrow a_2$ be the map such that $T(g) = f$. Using semisimplicity, the map g is isomorphic to one of the form

$$\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} : a_0 \oplus a'_1 \rightarrow a_0 \oplus a'_2.$$

The same holds after applying the functor T , since linear functors preserve direct sums. But the cokernel of this map is clearly $T(a'_2)$, which is in the image of T . \square

COROLLARY 5.3. *Assume the hypotheses of Theorem 5.1. If $A \in \mathcal{M}$ is a connected (i.e., $\dim \mathcal{C}(1, A) = 1$) semisimple Frobenius algebra, then $\text{Tr}_{\mathcal{C}}(A)$ is as well.*

Proof. Using that $\Phi(1_{\mathcal{C}}) = 1_{\mathcal{M}}$, it is easy to check that $\text{Tr}_{\mathcal{C}}(A)$ is connected. It is semisimple by Theorem 5.1. Define the counit by

$$\epsilon_{\text{Tr}_{\mathcal{C}}(A)} := \epsilon \circ \text{Tr}_{\mathcal{C}}(\epsilon_A) : \text{Tr}_{\mathcal{C}}(A) \rightarrow 1_{\mathcal{C}},$$

where $\epsilon : \text{Tr}_{\mathcal{C}}(1_{\mathcal{M}}) \rightarrow 1_{\mathcal{C}}$ is the left inverse of i . The pairing $p_{\text{Tr}_{\mathcal{C}}(A)} := \epsilon_{\text{Tr}_{\mathcal{C}}(A)} \circ m_{\text{Tr}_{\mathcal{C}}(A)}$ is nondegenerate by [Ost03, Prop. 3.1.ii], and so $\text{Tr}_{\mathcal{C}}(A)$ has a natural Frobenius algebra structure by Proposition 2.9. \square

REMARK 5.4. More generally, we would like to construct a comultiplication

$$\Delta_{x,y} = \text{[Diagram: A pair of pants with two top openings and one bottom opening. The left opening is red, the right is blue, and the bottom is dashed. The bottom opening is labeled 'x y' below it.]} : \text{Tr}_{\mathcal{C}}(x \otimes y) \rightarrow \text{Tr}_{\mathcal{C}}(x) \otimes \text{Tr}_{\mathcal{C}}(y)$$

which is compatible with the multiplication and all the other structures on $\text{Tr}_{\mathcal{C}}$. The graphical calculus suggests that there should be one, and Corollary 5.3 is a partial step in that direction. The main missing ingredient is the non-degeneracy of the pairing

$$\text{[Diagram: A pair of pants with one top opening and two bottom openings. The top opening is dashed. The left bottom opening is red, the right is blue, and both are dashed. The bottom openings are labeled 'x*' and 'x' below them.]} = \epsilon \circ \text{Tr}_{\mathcal{C}}(\text{ev}_x) \circ \mu_{x^*,x}$$

(compare with Proposition 2.9).


5.2 THE ALGEBRA $\text{Tr}_{\mathcal{C}}(A \otimes B)$

We now assume that \mathcal{C} and \mathcal{M} are pivotal tensor categories and that $\Phi^{\mathcal{Z}} : \mathcal{C} \rightarrow \mathcal{Z}(\mathcal{M})$ is a tensor functor. As always, $\text{Tr}_{\mathcal{C}}$ is the right adjoint of Φ . In the previous section, we saw that, under weaker hypotheses, if $A \in \mathcal{M}$ is an algebra then so is $\text{Tr}_{\mathcal{C}}(A)$. With our current assumptions, the same holds true for $\text{Tr}_{\mathcal{C}}(A \otimes B)$:

PROPOSITION 5.5. *Let \mathcal{C} and \mathcal{M} be pivotal tensor categories, let $\Phi^{\mathcal{Z}} : \mathcal{C} \rightarrow \mathcal{Z}(\mathcal{M})$ be a pivotal functor, and let $\text{Tr}_{\mathcal{C}}$ be the right adjoint of $\Phi = F \circ \Phi^{\mathcal{Z}}$. If A and B are algebra objects in \mathcal{M} , then $\text{Tr}_{\mathcal{C}}(A \otimes B)$ is an algebra object in \mathcal{C} .*

Proof. The structure morphisms of $\text{Tr}_{\mathcal{C}}(A \otimes B)$ are given by

$$m_{\text{Tr}_{\mathcal{C}}(A \otimes B)} := \text{Tr}_{\mathcal{C}}(m_A \otimes m_B) \circ \tau_{B,A \otimes A \otimes B}^- \circ \mu_{B \otimes A, A \otimes B} \circ (\tau_{A,B}^+ \otimes \text{id}_{\text{Tr}_{\mathcal{C}}(A \otimes B)})$$



and

$$i_{\mathrm{Tr}_C(A \otimes B)} := \mathrm{Tr}_C(i_A \otimes i_B) \circ i = \text{[Diagram: A cylinder with two vertical strands, one red and one blue, entering from the top and exiting from the bottom.]}.$$

We leave it as an exercise to check that these structure maps satisfy the necessary associativity and unitality axioms (Lemma 4.19 gets used for associativity). \square

Similarly to Remark 4.20, the pictures for $m_{\mathrm{Tr}_C(A \otimes B)}$ and $i_{\mathrm{Tr}_C(A \otimes B)}$ become more intuitive if we allow to draw strands on the back of the tubes. The structure morphisms become:

$$m_{\mathrm{Tr}_C(A \otimes B)} = \text{[Diagram: A pair of pants with two bottom openings and one top opening. Red and blue strands enter from the bottom openings and exit from the top opening. Dotted lines on the back represent strands on the back of the tubes.]} \quad \text{and} \quad i_{\mathrm{Tr}_C(A \otimes B)} = \text{[Diagram: A cylinder with a red strand on the front and a blue strand on the back, both entering from the top and exiting from the bottom.]} \quad (23)$$

We now prove our main theorem, about the semisimplicity of $\mathrm{Tr}_C(A \otimes B)$. Recall that for an algebra object in a semisimple category, separability is a somewhat stronger condition than semisimplicity — see Section 2.4 for a discussion.

THEOREM 5.6. *Let \mathcal{C} and \mathcal{M} be as in Proposition 5.5, and let us assume that they are both semisimple. Let $A, B \in \mathcal{M}$ be algebra objects such that the category $\mathrm{Bimod}_{\mathcal{M}}(A, B)$ of A - B -bimodules is semisimple. Then $\mathrm{Tr}_C(A \otimes B)$ is semisimple.*


In particular, if A and B are separable algebras (see Proposition 2.5), then $\mathrm{Tr}_C(A \otimes B)$ is semisimple.

Proof. The proof follows the same outline as that of Theorem 5.1. Let \mathcal{N}_0 be the essential image of the functor $\mathcal{C} \rightarrow \mathrm{Bimod}_{\mathcal{M}}(A, B)$ given by $c \mapsto A \otimes \Phi(c) \otimes B$, and let \mathcal{N} be the idempotent completion of \mathcal{N}_0 . Since $\mathrm{Bimod}_{\mathcal{M}}(A, B)$ is semisimple, \mathcal{N} is the full subcategory of $\mathrm{Bimod}_{\mathcal{M}}(A, B)$ generated by the simple objects that occur as direct summands of $A \otimes \Phi(c) \otimes B$ for $c \in \mathcal{C}$. The categorified trace induces a functor

$$\mathrm{Tr}_C^{\mathrm{Bimod}(A, B)} : \mathrm{Bimod}_{\mathcal{M}}(A, B) \rightarrow \mathrm{Mod}_{\mathcal{C}}(\mathrm{Tr}_C(A \otimes B)),$$

where the $\mathrm{Tr}_C(A \otimes B)$ -module structure on $\mathrm{Tr}_C(z)$ for $z \in \mathrm{Bimod}_{\mathcal{M}}(A, B)$ is

given by



We will show that the composite

$$\mathcal{N} \hookrightarrow \text{Bimod}_{\mathcal{M}}(A, B) \xrightarrow{\text{Tr}_C^{\text{Bimod}(A, B)}} \text{Mod}_C(\text{Tr}_C(A \otimes B))$$

is an equivalence of categories. As \mathcal{N} is semisimple, this will complete the proof.

- *The functor $\text{Tr}_C^{\text{Bimod}(A, B)}|_{\mathcal{N}}$ is fully faithful.* It is enough to show that the restriction to \mathcal{N}_0 is fully faithful, so we must show that the map

$$\mathcal{N}_0(A \otimes \Phi(c) \otimes B, A \otimes \Phi(d) \otimes B) \rightarrow \mathcal{C}_{\text{Tr}_C(A \otimes B)}(\text{Tr}_C(A \otimes \Phi(c) \otimes B), \text{Tr}_C(A \otimes \Phi(d) \otimes B))$$


$$\text{3D diagram of } f \text{ with strands } A, B, c, d \mapsto \text{Tr}_C \left(\text{3D diagram of } f \text{ with strands } A, B, c, d \right) = \text{Cylinder diagram of } f \tag{24}$$

is an isomorphism. The inverse of (24) is given by

$$\text{Cylinder diagram with } g \text{ and } \epsilon \mapsto \text{3D diagram with strands } A, B, c, d \tag{25}$$

One can check the equation (25)◦(24)=id directly as in the proof of Theorem 5.1. On the other hand, the equation (24)◦(25)=id is best checked after

precomposition by the isomorphism



$$: \mathrm{Tr}_{\mathcal{C}}(A \otimes B) \otimes c \rightarrow \mathrm{Tr}_{\mathcal{C}}(A \otimes \Phi(c) \otimes B). \quad (26)$$

Overall, the argument is very similar to the one used in the proof of Theorem 5.1.

- The functor $\mathrm{Tr}_{\mathcal{C}}^{\mathrm{Bimod}(A,B)}|_{\mathcal{N}}$ is essentially surjective. Once again, the argument is completely parallel to the one in Theorem 5.1. Every $\mathrm{Tr}_{\mathcal{C}}(A \otimes B)$ -module c fits in a coequalizer diagram

$$\mathrm{Tr}_{\mathcal{C}}(A \otimes B) \otimes \mathrm{Tr}_{\mathcal{C}}(A \otimes B) \otimes c \rightrightarrows \mathrm{Tr}_{\mathcal{C}}(A \otimes B) \otimes c \rightarrow c.$$

Using the isomorphism (26), we see that $\mathrm{Tr}_{\mathcal{C}}(A \otimes B) \otimes \mathrm{Tr}_{\mathcal{C}}(A \otimes B) \otimes c$ and $\mathrm{Tr}_{\mathcal{C}}(A \otimes B) \otimes c$ are in the essential image of \mathcal{N} . We are finished by Lemma 5.2. \square

REMARK 5.7. At this time, we do not know whether the algebra $\mathrm{Tr}(A \otimes B)$ is separable under the hypotheses of Theorem 5.6. The issues are similar to the ones encountered in Remark 5.4.

REMARK 5.8. The proofs of 5.5 and 5.6 go through if we only assume \mathcal{C} and \mathcal{M} to be rigid, as opposed to pivotal. When we encounter a traciator (as for example in the definition of $m_{\mathrm{Tr}_{\mathcal{C}}(A \otimes B)}$, or in that of the $\mathrm{Tr}_{\mathcal{C}}(A \otimes B)$ -algebra structure on $\mathrm{Tr}_{\mathcal{C}}(z)$ for $z \in \mathrm{Bimod}_{\mathcal{M}}(A, B)$, or in the isomorphism (26)) we need to insert a double dual at the appropriate place, as explained in Remark 4.15. It so happens that every such double duals gets undone by an inverse traciator later on.

Combining Theorem 5.6 with Remarks 2.7 and 5.8, we get the following

COROLLARY 5.9. *Let \mathcal{C} and \mathcal{M} be fusion categories over a field of characteristic zero, and let $\Phi^z : \mathcal{C} \rightarrow \mathcal{Z}(\mathcal{M})$ be a tensor functor. If $A, B \in \mathcal{M}$ are semisimple algebras, then so is $\mathrm{Tr}_{\mathcal{C}}(A \otimes B)$.*

EXAMPLE 5.10. Let $\mathcal{C} := \mathrm{SU}(2)_{16}$, with simple objects $\mathbf{1}, \dots, \mathbf{17}$. As explained in [Ost03] this tensor category has three indecomposable module categories, denoted A_{17} , D_{10} , and E_7 . The first one, A_{17} , is just \mathcal{C} acting on itself. A_{17} and D_{10} are module tensor categories, whereas E_7 is just a module category.

The simple objects of A_{17} , D_{10} , and E_7 correspond to the vertices of the Dynkin diagrams

$$A_{17} : \circ-\circ-\circ-\circ-\circ-\circ-\circ-\circ-\circ-\circ-\circ-\circ-\circ-\circ-\circ-\circ-\circ$$

$$D_{10} : \circ-\circ-\circ-\circ-\circ-\circ-\circ-\circ-\circ \quad E_7 : \circ-\circ-\circ-\circ-\circ-\circ-\circ$$

where the edges encode the action of $\mathbf{2} \in \mathcal{C}$. Let $\underline{\text{Hom}}_{\mathcal{C}}$ denote the \mathcal{C} -valued internal hom, explained at the beginning of the introduction. The simple algebra objects in \mathcal{C} are all of the form $\underline{\text{End}}_{\mathcal{C}}(m) := \underline{\text{Hom}}_{\mathcal{C}}(m, m)$, for some (not necessarily simple) object m in one of the above module categories.

Let us write $\underline{1}, \dots, \underline{9}, \underline{9}'$ for the simple objects of D_{10} . Triality is an action¹² of the symmetric group S_3 on the subcategory of D_{10} spanned by $\underline{1}, \underline{3}, \underline{5}, \underline{7}, \underline{9}, \underline{9}'$ [MPS11, Thm. 4.3]. The objects $\underline{1}, \underline{5}, \underline{7}$ are fixed, whereas the objects $\underline{3}, \underline{9}, \underline{9}'$ are permuted:



The algebra $\underline{\text{End}}_{D_{10}}(\underline{2}) = \underline{1} \oplus \underline{3}$ yields, under triality, the interesting algebras $A := \underline{1} \oplus \underline{9}$ and $B := \underline{1} \oplus \underline{9}'$. Let $\text{Tr}_{\mathcal{C}} : D_{10} \rightarrow \mathcal{C}$ be our categorified trace functor. At the level of underlying objects, one easily computes

$$\begin{aligned} \text{Tr}_{\mathcal{C}}(A) &\cong \mathbf{1} \oplus \mathbf{9} \oplus \mathbf{17} \\ \text{Tr}_{\mathcal{C}}(A \otimes A) &\cong \mathbf{1} \oplus \mathbf{1} \oplus \mathbf{5} \oplus \mathbf{9} \oplus \mathbf{9} \oplus \mathbf{9} \oplus \mathbf{13} \oplus \mathbf{17} \oplus \mathbf{17} \\ \text{Tr}_{\mathcal{C}}(A \otimes B) &\cong \mathbf{1} \oplus \mathbf{3} \oplus \mathbf{7} \oplus \mathbf{9} \oplus \mathbf{9} \oplus \mathbf{11} \oplus \mathbf{15} \oplus \mathbf{17} \end{aligned}$$

By compiling a list of semisimple algebra objects in \mathcal{C} , one notes that the above objects have a unique such structure. They are given by:

$$\begin{aligned} \text{Tr}_{\mathcal{C}}(A) &\cong \underline{\text{End}}_{\mathcal{C}}(\bullet-\circ-\circ-\circ-\circ) \\ \text{Tr}_{\mathcal{C}}(A \otimes A) &\cong \underline{\text{End}}_{\mathcal{C}}(\underline{1} \oplus \underline{9}) \\ \text{Tr}_{\mathcal{C}}(A \otimes B) &\cong \underline{\text{End}}_{\mathcal{C}}(\circ-\bullet-\circ-\circ-\circ). \end{aligned}$$

The algebras $\text{Tr}_{\mathcal{C}}(A)$ and $\text{Tr}_{\mathcal{C}}(A \otimes B)$ lie in the Morita equivalence class E_7 , whereas the algebra $\text{Tr}_{\mathcal{C}}(A \otimes A)$ lies in the Morita equivalence class D_{10} . Our computation should be contrasted with the one performed in [FRS02, (5.51)], where it was found that $[E_7] \times [E_7] = [D_{10}] + [E_7]$.

Given an algebra A and an object z in \mathcal{M} , then $z \otimes A \otimes z^*$ has a canonical algebra structure given by ‘protecting’ the multiplication and unit maps of A

¹²Here, an ‘action’ is a homomorphism to the group of isomorphism classes of tensor auto-equivalences.

by z -strands as in [MPS15, Proof of Prop. 5.4, p. 20]:

$$m_{z \otimes A \otimes z^*} = \text{[Diagram: A red strand splits into two green strands, with a green arc below them]} \quad \text{and} \quad i_{z \otimes A \otimes z^*} = \text{[Diagram: A red strand splits into two green strands, with a green arc above them]}$$

A straightforward calculation shows the following:

PROPOSITION 5.11. *Given two algebras A, B in \mathcal{M} and an object $z \in \mathcal{M}$, the two algebras $\text{Tr}_{\mathcal{C}}((z \otimes A \otimes z^*) \otimes B)$ and $\text{Tr}_{\mathcal{C}}(A \otimes (z^* \otimes B \otimes z))$ agree up to conjugation by the traciator.*

The structure morphisms can be drawn as in (23), with the A -strand on the front, the B -strand on the back, and the ‘protecting’ z -strands on the sides:

$$m_{\text{Tr}_{\mathcal{C}}(z \otimes A \otimes z^* \otimes B)} = \text{[Diagram: A 3D representation of the multiplication morphism with red, green, and blue strands]} \quad \text{and} \quad i_{\text{Tr}_{\mathcal{C}}(z \otimes A \otimes z^* \otimes B)} = \text{[Diagram: A 3D representation of the comultiplication morphism with red, green, and blue strands]}$$

The reader may wonder whether the two algebras $\text{Tr}_{\mathcal{C}}(A \otimes B)$ and $\text{Tr}_{\mathcal{C}}(B \otimes A)$ are related. We show that they are, under the assumption that \mathcal{C} is braided pivotal and $\Phi^z : \mathcal{C} \rightarrow \mathcal{Z}(\mathcal{M})$ is a braided pivotal functor. Let us define the *opposite algebra* of $A = (A, m, i)$ to be the algebra $A^{\text{op}} := (A, m \circ \beta, i)$.

PROPOSITION 5.12. *The traciator $\tau_{B,A} : \text{Tr}_{\mathcal{C}}(B \otimes A) \rightarrow \text{Tr}_{\mathcal{C}}(A \otimes B)$ induces an algebra isomorphism $\text{Tr}_{\mathcal{C}}(B \otimes A) \cong \text{Tr}_{\mathcal{C}}(A \otimes B)^{\text{op}}$.*

Proof. We need to show that $m_{\text{Tr}_{\mathcal{C}}(B \otimes A)} = \tau^- \circ m_{\text{Tr}_{\mathcal{C}}(A \otimes B)} \circ \beta \circ (\tau^+ \otimes \tau^+)$. We rewrite $m_{\text{Tr}_{\mathcal{C}}(A \otimes B)}$ as follows

$$m_{\text{Tr}_{\mathcal{C}}(A \otimes B)} = \text{[Diagram 1: A 3D representation of the multiplication morphism with red and blue strands]} = \text{[Diagram 2: A 3D representation of the multiplication morphism with red and blue strands]} = \text{[Diagram 3: A 3D representation of the multiplication morphism with red and blue strands, showing a crossing]} = \text{[Diagram 4: A 3D representation of the multiplication morphism with red and blue strands, showing a crossing]} = \text{[Diagram 5: A 3D representation of the multiplication morphism with red and blue strands, showing a crossing]}$$

and compute:

$$\tau_{A,B}^- \circ m_{\text{Tr}_{\mathcal{C}}(A \otimes B)} \circ \beta \circ (\tau_{B,A}^+ \otimes \tau_{B,A}^+) = \text{[Diagram 1]} = \text{[Diagram 2]} = m_{\text{Tr}_{\mathcal{C}}(B \otimes A)}.$$

The remaining statement $i_{\text{Tr}_{\mathcal{C}}(B \otimes A)} = \tau^- \circ i_{\text{Tr}_{\mathcal{C}}(A \otimes B)}$ is straightforward, and left to the reader. □

REMARK 5.13. Unless \mathcal{C} is symmetric, there are actually two opposite algebras: $(A, m \circ \beta^+, i)$ and $(A, m \circ \beta^-, i)$, the (+)-opposite and the (−)-opposite algebras. However, if \mathcal{C} is balanced then the twist map $\theta_A : A \rightarrow A$ provides an isomorphism between (−)-opposite and the (+)-opposite algebras, so that there is, in fact, only one opposite algebra.

A APPENDIX

A.1 COMMUTATIVE ALGEBRAS IN BRAIDED TENSOR CATEGORIES

In this appendix, we expand on and fill in the details of Example 3.12. To begin with, we introduce a convenient graphical notation for tensor products over algebra objects. Let a be an algebra object in a tensor category \mathcal{C} , and let $x, y \in \mathcal{C}$ be right and left a -modules. Denoting a by an orange strand and x, y by green and blue strands, the tensor product over a is the coequalizer $x \otimes_a y$ of the two morphisms

$$\left| \begin{array}{c} \text{[Green strand]} \\ \text{[Orange strand]} \end{array} \right|, \left| \begin{array}{c} \text{[Blue strand]} \\ \text{[Orange strand]} \end{array} \right| : x \otimes a \otimes y \rightarrow x \otimes y.$$

(we assume that \mathcal{C} has all the necessary colimits). It is convenient to denote $x \otimes_a y$ by an orange ribbon bordered by green and blue edges:

$$\begin{array}{c} \text{[Green and Blue ribbon]} \\ x \otimes_a y \end{array}$$


We draw the natural projection $\pi_{x,y} = \left| \begin{array}{c} \text{orange strand} \\ \text{green and blue strands} \end{array} \right| : x \otimes y \rightarrow x \otimes_a y$. It makes the equation

graphically motivated as the orange strand (which we might as well have represented by an orange ribbon) is allowed to slide along the orange edge of the surface. As in usual algebra, there are canonical isomorphisms $x \otimes_a a \cong x$ and $a \otimes_a y \cong y$.

When \mathcal{C} is a braided tensor category and $a \in \mathcal{C}$ is a commutative algebra object

then it makes sense to tensor two left a -modules, and the result is again an a -module. Specifically, if x, y are left a -modules, then $x \otimes_a y$ is the coequalizer of the two morphisms

Instead of using a ribbon as above, it is preferable in this situation to represent $x \otimes_a y$ by a thick filled orange ‘rope’ with green and blue strands on its surface:



Extrapolating the notation, we can also denote a single a -module x by , where the presence of the orange rope is just an indication that x is an a -module.

Let us now assume that a is commutative and separable, also known as étale [DMNO13, §3] (such an algebra is automatically Frobenius – combine [DMNO13, Rem. 3.2.ii], [Ost03, Prop. 3.1.ii], and Proposition 2.9 for a proof). The dual of an a -module x is then naturally also an a -module. The evaluation $\text{ev}_x : x^* \otimes x \rightarrow 1$ induces a corresponding evaluation morphism $\bar{\text{ev}}_x : x^* \otimes_a x \rightarrow a$ in the category of a -modules, and the same holds for coevaluations [DMNO13, §3.3]. We write

$$\mathcal{M} := \text{Mod}_{\mathcal{C}}(a)$$

for the category of a -modules in \mathcal{C} . This is a rigid tensor category under the operation $x, y \mapsto x \otimes_a y$.

Let $\Phi : \mathcal{C} \rightarrow \mathcal{M}$ be the free module functor, given by $\Phi(x) := a \otimes x$ on objects and $\Phi(f : x \rightarrow y) = \text{id}_a \otimes f : a \otimes x \rightarrow a \otimes y$ on morphisms. We then have a functor $\Phi^z : \mathcal{C} \rightarrow \mathcal{Z}(\mathcal{M})$ given by $\Phi^z(x) := (\Phi(x), e_{\Phi(x)})$, where the half-braiding $e_{\Phi(x)}$ is given by

$$e_{\Phi(x),y} : \Phi(x) \otimes_a y \cong x \otimes y \xrightarrow{\beta_{x,y}} y \otimes x \cong y \otimes_a \Phi(x).$$

In diagrams, this is denoted

$$e_{\Phi(x),y} = \text{[Diagram of a rope with a blue strand on the surface and a green strand inside]} \tag{27}$$

The blue strand represents $y \in \mathcal{M}$, and lies on the surface of the rope. The green strand represents $x \in \mathcal{C}$, and doesn't touch the surface. The above map is visibly invertible, and natural in y . The hexagon axiom (2) for the half-braiding is the equation

$$\text{[Diagrammatic equation (28) showing two rope configurations with three strands (blue, green, red) and an equals sign]} \tag{28}$$

It holds as both sides of (28) fit into the same commutative diagram

$$\begin{array}{ccc} y \otimes z \otimes x & \xrightarrow{\pi_{yz} \otimes 1} & y \otimes_a z \otimes x \cong y \otimes_a z \otimes_a \Phi(x) \\ \uparrow (1 \otimes \beta_{x,z})(\beta_{x,y} \otimes 1) = \beta_{x,y \otimes z} & & \uparrow (28) \\ x \otimes y \otimes z & \xrightarrow{1 \otimes \pi_{yz}} & x \otimes y \otimes_a z \cong \Phi(x) \otimes_a y \otimes_a z \end{array}$$

with surjective horizontal maps. The isomorphism $(a \otimes x) \otimes_a (a \otimes y) \cong a \otimes x \otimes y$ endows Φ^z with the structure of a tensor functor. Indeed, the half braidings $e_{\Phi(x) \otimes_a \Phi(y)} = (e_{\Phi(x)} \otimes_a 1) \circ (1 \otimes_a e_{\Phi(y)})$ and $e_{\Phi(x \otimes y)}$ are naturally isomorphic, as required for Φ^z to be a tensor functor:


$$\text{[Diagrammatic equation showing two rope configurations with three strands (blue, green, red) and an equals sign]} \tag{28}$$

Finally, the functor $\Phi^{\mathcal{Z}}$ is braided because the morphism  is both the image of $\beta_{x,y}$ under $\Phi^{\mathcal{Z}}$, and a special case of (27).

Let us now furthermore assume that \mathcal{C} is pivotal, and that $\theta_a = 1$, where the twist maps are given by (3). Then for any a -module x , the pivotal map $\varphi_x : x \rightarrow x^{**}$ is a map of a -modules [KO02, Thm. 1.17]¹³. This equips the category $\text{Mod}_{\mathcal{C}}(a)$ of a -modules with the structure of a pivotal tensor category. We wish to check that $\Phi^{\mathcal{Z}} : \mathcal{C} \rightarrow \mathcal{Z}(\mathcal{M})$ is a pivotal functor, i.e., that the equation $\delta_x^* \circ \varphi_{\Phi^{\mathcal{Z}}(x)} = \delta_{x^*} \circ \Phi^{\mathcal{Z}}(\varphi_x)$ holds. Let $\omega : a \rightarrow a^*$ be the isomorphism induced by the Frobenius pairing $\epsilon \circ \mu : a \otimes a \rightarrow 1$. The canonical isomorphism $\delta_x : \Phi^{\mathcal{Z}}(x^*) \rightarrow \Phi^{\mathcal{Z}}(x)^*$ is then given by

$$\Phi(x^*) = a \otimes x^* \xrightarrow{\beta_{a,x^*}^-} x^* \otimes a \xrightarrow{1 \otimes \omega} x^* \otimes a^* \cong (a \otimes x)^* = \Phi(x)^*.$$

We need to check that



holds. By the monoidal property of φ , this is equivalent to the equation $\varphi_a = (\omega^*)^{-1} \circ \omega$, which is itself a consequence of $\theta_a = 1$ [KO02, Lem. 1.13]. This finishes the proof that $\Phi^{\mathcal{Z}}$ is a pivotal functor.

A.2 BRAIDED PIVOTAL CATEGORIES

In this second appendix, we provide an overview of some basic properties of braided pivotal categories, since these have not received as much attention as many related notions. Some of these results can be found in the literature [Sel11, §4.6 and 4.7] [Yet92, Prop. 2.11] (see also [BK01, §2.2] and [DGNO10, §2.8.2]). We provide them here along with short proofs for the convenience of the reader.

LEMMA A.1. *Let \mathcal{C} be balanced rigid category. Then there are two ways of identifying each object with its double dual, each one making \mathcal{C} into a pivotal category.*

Proof. For $a \in \mathcal{C}$, define $\varphi_a^{(1)} : a \rightarrow a^{**}$ by

$$\varphi_a^{(1)} = (\text{ev}_a \otimes \text{id}_{a^{**}}) \circ (\text{id}_{a^*} \otimes \beta_{a^{**},a}^-) \circ (\text{coev}_{a^*} \otimes \text{id}_a) \circ \theta_a = \begin{array}{c} a^{**} \\ | \\ \text{---} \circ \text{---} \\ | \\ \theta_a \\ | \\ a \end{array}. \quad (29)$$

¹³To match our conventions, one should replace all over-crossings by under-crossings in [KO02].

This is clearly invertible with inverse $\theta_a^{-1} \circ (\text{id}_a \otimes \text{ev}_{a^*}) \circ (\beta_{a^{**},a} \otimes \text{id}_{a^*}) \circ (\text{id}_{a^{**}} \otimes \text{coev}_a)$. The naturality of $\varphi_a^{(1)}$ is left as an exercise, and it is straightforward to show $\varphi_{a \otimes b}^{(1)} = \varphi_a^{(1)} \otimes \varphi_b^{(1)}$ using that $\theta_{a \otimes b} = \beta_{b,a} \circ \beta_{a,b} \circ (\theta_a \otimes \theta_b)$. The other pivotal structure is given by

$$\varphi_a^{(2)} = (\text{ev}_a \otimes \text{id}_{a^{**}}) \circ (\text{id}_{a^*} \otimes \beta_{a^{**},a}^+) \circ (\text{coev}_{a^*} \otimes \text{id}_a) \circ \theta_a^{-1} = \begin{array}{c} a^{**} \\ | \\ \circlearrowleft \\ | \\ \boxed{\theta_a^{-1}} \\ | \\ a \end{array}. \quad \square$$

The two pivotal structures are related by

$$\varphi_a^{(2)} = \begin{array}{c} a^{**} \\ | \\ \circlearrowleft \\ | \\ \boxed{(\varphi_a^{(1)})^{-1}} \\ | \\ a^{**} \\ | \\ \circlearrowleft \\ | \\ a \end{array} = \begin{array}{c} a^{**} \\ | \\ \circlearrowleft \\ | \\ \boxed{(\varphi_a^{(1)})^{-1}} \\ | \\ a^{**} \\ | \\ \circlearrowleft \\ | \\ a \end{array} \quad \text{and} \quad \varphi_a^{(1)} = \begin{array}{c} a^{**} \\ | \\ \circlearrowleft \\ | \\ \boxed{(\varphi_a^{(2)})^{-1}} \\ | \\ a^{**} \\ | \\ \circlearrowleft \\ | \\ a \end{array} = \begin{array}{c} a^{**} \\ | \\ \circlearrowleft \\ | \\ \boxed{(\varphi_a^{(2)})^{-1}} \\ | \\ a^{**} \\ | \\ \circlearrowleft \\ | \\ a \end{array}. \quad (30)$$

LEMMA A.2. *A braided pivotal category has two sets of twists, each one making it a balanced category.*

Proof. For $a \in \mathcal{C}$, let $\varphi_a : a \rightarrow a^{**}$ be the natural isomorphism, and define

$$\theta_a^{(1)} = (\text{id}_a \otimes \text{ev}_{a^*}) \circ (\beta_{a^{**},a} \otimes \text{id}_{a^*}) \circ (\text{id}_{a^{**}} \otimes \text{coev}_a) \circ \varphi_a = \begin{array}{c} a \\ | \\ \circlearrowright \\ | \\ \boxed{\varphi_a} \\ | \\ a^{**} \\ | \\ \circlearrowleft \\ | \\ a \end{array}. \quad (31)$$

The other alternative is

$$\theta_a^{(2)} = \varphi_a^{-1} \circ (\text{ev}_a \otimes \text{id}_{a^{**}}) \circ (\text{id}_{a^*} \otimes \beta_{a^{**},a}^+) \circ (\text{coev}_{a^*} \otimes \text{id}_a) = \begin{array}{c} a \\ | \\ \boxed{\varphi_a^{-1}} \\ | \\ a^{**} \\ | \\ \circlearrowleft \\ | \\ a \end{array}. \quad (32)$$

The naturality of $\theta^{(1)}$ and $\theta^{(2)}$ follows from that of φ . Using the monoidal property of φ , respectively φ^{-1} , one verifies that the balance axiom holds between these twist maps and the braiding. \square

The two twists are related by

$$\theta_a^{(2)} = \left[\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \theta_{a^*}^{(1)} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \right] = \left[\begin{array}{c} \text{---} \\ \text{---} \\ \theta_{a^*}^{(1)} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \right] \quad \text{and} \quad \theta_a^{(1)} = \left[\begin{array}{c} \text{---} \\ \text{---} \\ \theta_{a^*}^{(2)} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \right] = \left[\begin{array}{c} \text{---} \\ \text{---} \\ \theta_{a^*}^{(2)} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \right]. \quad (33)$$

Lemmas A.1 and A.2 now combine to give the following corollary:

COROLLARY A.3. *Let \mathcal{C} be a braided rigid category. Then there are two ways¹⁴ of establishing a one-to-one correspondence between pivotal structures on \mathcal{C} , and twists making \mathcal{C} into a balanced category (see Figure 2 in Section 2.3).*

PROPOSITION A.4. *Let $(\mathcal{C}, \beta, \varphi^{(1)})$ be a braided pivotal category. Using $\varphi^{(1)}$, let us define $\varphi^{(2)}, \theta^{(1)}, \theta^{(2)}$ by means of Equations (30), (31), (32), respectively. Then the following three properties are equivalent*

- (1) $\theta^{(1)} = \theta^{(2)}$
- (2) $(\mathcal{C}, \beta, \theta^{(i)})$ is ribbon for either $i = 1, 2$
- (3) $\varphi^{(1)} = \varphi^{(2)}$

and imply this fourth one:

- (4) $(\mathcal{C}, \beta, \varphi^{(i)})$ is spherical for either $i = 1, 2$.

If moreover \mathcal{C} is semisimple, then the fourth property is equivalent to the first three.

Proof.

(1) \Leftrightarrow (2): By definition, $(\mathcal{C}, \beta, \theta^{(1)})$ is ribbon if and only if $(\theta_a^{(1)})^* = \theta_{a^*}^{(1)}$ holds for every $a \in \mathcal{C}$. By Equation (33), we have $(\theta_a^{(1)})^* = \theta_{a^*}^{(2)}$. Therefore $(\mathcal{C}, \beta, \theta^{(1)})$ is ribbon iff $\theta_{a^*}^{(1)} = \theta_{a^*}^{(2)}$ holds $\forall a$ iff $\theta_a^{(1)} = \theta_a^{(2)}$ holds $\forall a$. The same argument works with $(\mathcal{C}, \beta, \theta^{(2)})$ in place of $(\mathcal{C}, \beta, \theta^{(1)})$.

(2) \Leftrightarrow (3): Letting $\theta := \theta^{(1)}$, then for every $a \in \mathcal{C}$, we have

$$\left[\begin{array}{c} \text{---} \\ \theta_{a^*}^* \\ \text{---} \\ a^* \end{array} \right] = \left[\begin{array}{c} \text{---} \\ \theta_a \\ \text{---} \\ a \end{array} \right] = \left[\begin{array}{c} \text{---} \\ \theta_a \\ \text{---} \\ a \end{array} \right] = \left[\begin{array}{c} \text{---} \\ \varphi_a^{(1)} \\ \text{---} \\ a \end{array} \right]$$

¹⁴In [Sel11, Rem. 4.22], it is incorrectly stated that there are \mathbb{Z} many ways of establishing such a correspondence. In fact, the construction only produces $\mathbb{Z}/2\mathbb{Z}$ many distinct correspondences.

and

The diagram shows four stages of a string diagram transformation.
 1. A box labeled θ_{a^*} with a vertical line passing through it, labeled a^* on the left and a on the right.
 2. An equality sign followed by a box labeled θ_{a^*} with a loop on the left side.
 3. An equality sign followed by a box labeled $(\varphi_{a^*}^{(2)})^{-1}$ with a loop on the left side.
 4. An equality sign followed by a box labeled $\varphi_{a^*}^{(2)}$ with a loop on the right side.

It follows that $(\theta_a^{(1)})^* = \theta_{a^*}^{(1)}$ holds if and only if $\varphi_a^{(1)} = \varphi_{a^*}^{(2)}$ holds. Using Equations (33), the former is also easily seen to be equivalent to $(\theta_a^{(2)})^* = \theta_{a^*}^{(2)}$. (3) \Rightarrow (4): Assuming $\varphi := \varphi^{(1)} = \varphi^{(2)}$, we show that $(\mathcal{C}, \beta, \varphi)$ is spherical:

The diagram shows a sequence of five stages of a string diagram transformation.
 1. A box labeled φ above a box labeled f , with a vertical line passing through them.
 2. An equality sign followed by a box labeled f to the left of a box labeled φ^{-1} , with a loop on the left side.
 3. An equality sign followed by a box labeled f to the left of a box labeled φ , with a loop on the right side.
 4. An equality sign followed by a box labeled φ to the left of a box labeled f , with a loop on the left side.
 5. An equality sign followed by a box labeled f above a box labeled φ^{-1} , with a vertical line passing through them.
 The final expression is $= \text{tr}_L(f)$.

We have used Equation (30) for the third equality above.

Let now \mathcal{C} be semisimple.

(4) \Rightarrow (2): We again write $\varphi := \varphi^{(1)}$, and assume that $(\mathcal{C}, \beta, \varphi)$ is spherical. By naturality of the twist, it is enough to verify $\theta_a^* = \theta_{a^*}$ on simple objects. Let ϑ_a^* and ϑ_{a^*} be the scalars defined by $\theta_a^* = \vartheta_a^* \text{id}_{a^*}$ and $\theta_{a^*} = \vartheta_{a^*} \text{id}_{a^*}$. By Lemma A.5, the quantum dimension $\dim(a) := \text{tr}_L(1_a)$ of a simple object is always non-zero. So we have $\theta_a^* = \theta_{a^*}$ if and only if $\text{tr}_L(\theta_a^*) = \dim(a^*)\vartheta_a^*$ equals $\text{tr}_L(\theta_{a^*}) = \dim(a^*)\vartheta_{a^*}$:

The diagram shows a sequence of three stages of a string diagram transformation.
 1. A box labeled θ above a box labeled φ^{-1} , with a vertical line passing through them.
 2. An equality sign followed by a box labeled φ above a box labeled θ , with a loop on the left side.
 3. An equality sign followed by a box labeled φ above a box labeled θ , with a vertical line passing through them.
 The final expression is $= \text{tr}_R(\theta_{a^*}) = \text{tr}_L(\theta_{a^*})$. \square

LEMMA A.5. *Let \mathcal{C} be a rigid semisimple tensor category (linear over some field k). Then for every simple object a and non-zero morphisms $c : 1 \rightarrow a \otimes a^*$ and $e : a \otimes a^* \rightarrow 1$, we have $e \circ c \neq 0$.*

Proof. By the adjunction $\mathcal{C}(1, a \otimes a^*) \cong \mathcal{C}(a, a) \cong k$, any morphism $1 \rightarrow a \otimes a^*$ is a multiple of c . Decompose $a \otimes a^*$ as $u \oplus x$ with u the image of c , and decompose 1 as $1' \oplus 1''$ with $1'$ the coimage of c . We have $\mathcal{C}(1', x) = \mathcal{C}(1'', x) = \mathcal{C}(1'', u) = 0$, $\mathcal{C}(1', u) = k$, and c restricts to an isomorphism $1' \rightarrow u$.

By semisimplicity, we also have $\mathcal{C}(x, 1') = \mathcal{C}(x, 1'') = \mathcal{C}(u, 1'') = 0$, $\mathcal{C}(u, 1') = k$, and e restricts to an isomorphism $u \rightarrow 1'$. The composite $e \circ c$ is now visibly non-zero. \square

In Lemma A.2, we have discussed the two ways of equipping a braided pivotal category with a balanced structure. We provide the corresponding result at the level of functors:

LEMMA A.6. *Let \mathcal{C}, \mathcal{D} be braided pivotal categories, equipped with the balanced structure (31). Then a braided pivotal functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is automatically balanced. The same result holds using the other balanced structure (32).*


Proof. Recall from Section 2.1 that the isomorphism $\delta_a : F(a^*) \rightarrow F(a)^*$ is defined by requiring that

$$F(\text{ev}_a) = i \circ \text{ev}_{F(a)} \circ (\delta_a \otimes \text{id}_{F(a)}) \circ \nu_{a^*,a}^{-1}.$$

It follows that $\text{ev}_{F(a)} = i^{-1} \circ F(\text{ev}_a) \circ \nu_{a^*,a} \circ (\delta_a^{-1} \otimes \text{id}_{F(a)})$. The map $\text{coev}_{F(a)}$ is characterised by the zig-zag equation $(\text{id} \otimes \text{ev}_{F(a)}) \circ (\text{coev}_{F(a)} \otimes \text{id}) = \text{id}_{F(a)}$. As the morphism $(\text{id}_{F(a)} \otimes \delta_a) \circ \nu_{a,a^*}^{-1} \circ F(\text{coev}_a) \circ i$ satisfies that equation, it must be equal to $\text{coev}_{F(a)}$. So we get that

$$F(\text{coev}_a) = \nu_{a,a^*} \circ (\text{id}_{F(a)} \otimes \delta_a^{-1}) \circ \text{coev}_{F(a)} \circ i^{-1}.$$

We can now prove that $F : (\mathcal{C}, \theta^{(1)}) \rightarrow (\mathcal{D}, \theta^{(1)})$ is a balanced functor:



The proof that $F : (\mathcal{C}, \theta^{(2)}) \rightarrow (\mathcal{D}, \theta^{(2)})$ is a balanced functor is similar. □

REFERENCES

- [BD95] John C. Baez and James Dolan, *Higher-dimensional algebra and topological quantum field theory*, J. Math. Phys. 36 (1995), no. 11, 6073–6105, [MR1355899](#) [arXiv:q-alg/9503002](#) [DOI:10.1063/1.531236](#).
- [BDH14] Arthur Bartels, Chris Douglas, and André Henriques, *Conformal nets II: conformal blocks*, 2014, [arXiv:1409.8672](#).
- [Bez04] Roman Bezrukavnikov, *On tensor categories attached to cells in affine Weyl groups*, Representation theory of algebraic groups and quantum groups, Adv. Stud. Pure Math., vol. 40, Math. Soc. Japan, Tokyo, 2004, [MR2074589](#), pp. 69–90.
- [BFO09] Roman Bezrukavnikov, Michael Finkelberg, and Victor Ostrik, *On tensor categories attached to cells in affine Weyl groups. III*, Israel J. Math. 170 (2009), 207–234, [MR2506324](#) [DOI:10.1007/s11856-009-0026-9](#) [arXiv:math/0605628](#).
- [Big10] Stephen Bigelow, *Skein theory for the ADE planar algebras*, J. Pure Appl. Algebra 214 (2010), no. 5, 658–666, [arXiv:math.QA/0903.0144](#) [MR2577673](#) [DOI:10.1016/j.jpaa.2009.07.010](#).
- [BK01] Bojko Bakalov and Alexander Kirillov, Jr., *Lectures on tensor categories and modular functors*, University Lecture Series, vol. 21, American Mathematical Society, Providence, RI, 2001, [MR1797619](#).
- [BN11] Alain Bruguières and Sonia Natale, *Exact sequences of tensor categories*, Int. Math. Res. Not. IMRN (2011), no. 24, 5644–5705, [MR2863377](#) [DOI:10.1093/imrn/rnq294](#) [arXiv:1006.0569](#).
- [DGNO10] Vladimir Drinfeld, Shlomo Gelaki, Dmitri Nikshych, and Victor Ostrik, *On braided fusion categories. I*, Selecta Math. (N.S.) 16 (2010), no. 1, 1–119, [MR2609644](#) [DOI:10.1007/s00029-010-0017-z](#) [arXiv:0906.0620](#).
- [DMNO13] Alexei Davydov, Michael Müger, Dmitri Nikshych, and Victor Ostrik, *The Witt group of non-degenerate braided fusion categories*, J. Reine Angew. Math. 677 (2013), 135–177, [MR3039775](#) [arXiv:1009.2117](#).
- [DSPA14] Chris Douglas, Chris Schommer-Pries, and Noah Snyder, *The balanced tensor product of module categories*, 2014, [arXiv:1406.4204](#).
- [ENO05] Pavel Etingof, Dmitri Nikshych, and Viktor Ostrik, *On fusion categories*, Ann. of Math. (2) 162 (2005), no. 2, 581–642, [MR2183279](#) [DOI:10.4007/annals.2005.162.581](#) [arXiv:math.QA/0203060](#).

- [EO04] Pavel Etingof and Viktor Ostrik, *Module categories over representations of $SL_q(2)$ and graphs*, Math. Res. Lett. 11 (2004), no. 1, 103–114, [MR2046203](#) [arXiv:math.QA/0302130](#).
- [FKST95] Jürg Fröhlich, Thomas Kerler, Urban M. Studer, and Emmanuel Thiran, *Structuring the set of incompressible quantum Hall fluids*, Nuclear Phys. B 453 (1995), no. 3, 670–704, [MR1358783](#) [DOI:10.1016/0550-3213\(95\)00426-S](#) [arXiv:cond-mat/9505156](#).
- [FRS02] Jürgen Fuchs, Ingo Runkel, and Christoph Schweigert, *TFT construction of RCFT correlators. I. Partition functions*, Nuclear Phys. B 646 (2002), no. 3, 353–497, [MR1940282](#) [DOI:10.1016/S0550-3213\(02\)00744-7](#) [arXiv:hep-th/0204148](#).
- [FS08] Jürgen Fuchs and Carl Stigner, *On Frobenius algebras in rigid monoidal categories*, Arab. J. Sci. Eng. Sect. C Theme Issues 33 (2008), no. 2, 175–191, [MR2500035](#) [arXiv:0901.4886](#).
- [FY92] Peter Freyd and David N. Yetter, *Coherence theorems via knot theory*, J. Pure Appl. Algebra 78 (1992), no. 1, 49–76, [MR1154897](#) [DOI:10.1016/0022-4049\(92\)90018-B](#).
- [Hen15] André Henriques, *What Chern-Simons theory assigns to a point*, 2015, [arXiv:1503.06254](#).
- [HP15] André Henriques and David Penneys, *Bicommutant categories from fusion categories*, 2015.
- [HPT16] André Henriques, David Penneys, and James E. Tener, *Planar algebras in braided tensor categories*, 2016, In preparation.
- [Izu01] Masaki Izumi, *The structure of sectors associated with Longo-Rehren inclusions. II. Examples*, Rev. Math. Phys. 13 (2001), no. 5, 603–674, [MR1832764](#) [DOI:10.1142/S0129055X01000818](#).
- [JL16] Arthur Jaffe and Zhengwei Liu, *Planar para algebras, reflection positivity*, 2016, [arXiv:1602.02662](#).
- [JS91a] André Joyal and Ross Street, *The geometry of tensor calculus. I*, Adv. Math. 88 (1991), no. 1, 55–112, [MR1113284](#).
- [JS91b] ———, *Tortile Yang-Baxter operators in tensor categories*, J. Pure Appl. Algebra 71 (1991), no. 1, 43–51, [MR1107651](#) [DOI:10.1016/0022-4049\(91\)90039-5](#).
- [Kac90] Victor G. Kac, *Infinite-dimensional Lie algebras*, third ed., Cambridge University Press, Cambridge, 1990, [MR1104219](#) [DOI:10.1017/CB09780511626234](#).
- [Kel74] G. M. Kelly, *Doctrinal adjunction*, Category Seminar (Proc. Sem., Sydney, 1972/1973), Springer, Berlin, 1974, [MR0360749](#), pp. 257–280. Lecture Notes in Math., Vol. 420.

- [KO02] Alexander Kirillov, Jr. and Viktor Ostrik, *On a q -analogue of the McKay correspondence and the ADE classification of \mathfrak{sl}_2 conformal field theories*, Adv. Math. 171 (2002), no. 2, 183–227, [MR1936496](#) [arXiv:math.QA/0101219](#) DOI:10.1006/aima.2002.2072.
- [ML98] Saunders Mac Lane, *Categories for the working mathematician*, second ed., Graduate Texts in Mathematics, vol. 5, Springer-Verlag, New York, 1998, [MR1712872](#).
- [MPS10] Scott Morrison, Emily Peters, and Noah Snyder, *Skein theory for the D_{2n} planar algebras*, J. Pure Appl. Algebra 214 (2010), no. 2, 117–139, [arXiv:0808.0764](#) [MR2559686](#) DOI:10.1016/j.jpaa.2009.04.010.
- [MPS11] ———, *Knot polynomial identities and quantum group coincidences*, Quantum Topol. 2 (2011), no. 2, 101–156, [MR2783128](#) DOI:10.4171/QT/16 [arXiv:1003.0022](#).
- [MPS15] Scott Morrison, Emily Peters, and Noah Snyder, *Categories generated by a trivalent vertex*, 2015, [arXiv:1501.06869](#).
- [Müg03a] Michael Müger, *From subfactors to categories and topology. I. Frobenius algebras in and Morita equivalence of tensor categories*, J. Pure Appl. Algebra 180 (2003), no. 1-2, 81–157, [MR1966524](#) DOI:10.1016/S0022-4049(02)00247-5 [arXiv:math.CT/0111204](#).
- [Müg03b] ———, *From subfactors to categories and topology. II. The quantum double of tensor categories and subfactors*, J. Pure Appl. Algebra 180 (2003), no. 1-2, 159–219, [MR1966525](#) DOI:10.1016/S0022-4049(02)00248-7 [arXiv:math.CT/0111205](#).
- [MV88] I. Mirković and K. Vilonen, *Characteristic varieties of character sheaves*, Invent. Math. 93 (1988), no. 2, 405–418, [MR948107](#) DOI:10.1007/BF01394339.
- [Ocn02] Adrian Ocneanu, *The classification of subgroups of quantum $SU(N)$* , Quantum symmetries in theoretical physics and mathematics (Bariloche, 2000), Contemp. Math., vol. 294, Amer. Math. Soc., Providence, RI, 2002, [MR1907188](#), pp. 133–159.
- [Ost03] Victor Ostrik, *Module categories, weak Hopf algebras and modular invariants*, Transform. Groups 8 (2003), no. 2, 177–206, [MR1976459](#) [arXiv:math/0111139](#).
- [Ost14] ———, *Tensor categories attached to exceptional cells in Weyl groups*, Int. Math. Res. Not. IMRN (2014), no. 16, 4521–4533, [MR3250042](#) [arXiv:1108.3060](#).
- [Par95] Bodo Pareigis, *On braiding and dyslexia*, J. Algebra 171 (1995), no. 2, 413–425, [MR1315904](#) DOI:10.1006/jabr.1995.1019.
- [PS13] Kate Ponto and Michael Shulman, *Shadows and traces in bicategories*, J. Homotopy Relat. Struct. 8 (2013), no. 2, 151–200, [MR3095324](#) DOI:10.1007/s40062-012-0017-0 [arXiv:0910.1306](#).

- [Saw06] Stephen F. Sawin, *Quantum groups at roots of unity and modularity*, J. Knot Theory Ramifications 15 (2006), no. 10, 1245–1277, [MR2286123](#) [arXiv:math.QA/0308281](#).
- [Sel11] P. Selinger, *A survey of graphical languages for monoidal categories*, New structures for physics, Lecture Notes in Phys., vol. 813, Springer, Heidelberg, 2011, [MR2767048](#) [DOI:10.1007/978-3-642-12821-9_4](#), pp. 289–355.
- [Shi11] Kenichi Shimizu, *Frobenius-Schur indicators in Tambara-Yamagami categories*, J. Algebra 332 (2011), 543–564, [MR2774703](#) [arXiv:1005.4500](#) [DOI:10.1016/j.jalgebra.2011.02.002](#).
- [Shu94] Mei Chee Shum, *Tortile tensor categories*, J. Pure Appl. Algebra 93 (1994), no. 1, 57–110, [MR1268782](#).
- [TY98] Daisuke Tambara and Shigeru Yamagami, *Tensor categories with fusion rules of self-duality for finite abelian groups*, J. Algebra 209 (1998), no. 2, 692–707, [MR1659954](#).
- [Vor99] Alexander A. Voronov, *The Swiss-cheese operad*, Homotopy invariant algebraic structures (Baltimore, MD, 1998), Contemp. Math., vol. 239, Amer. Math. Soc., Providence, RI, 1999, [MR1718089](#) [arXiv:math.QA/9807037](#), pp. 365–373.
- [Yet92] David N. Yetter, *Framed tangles and a theorem of Deligne on braided deformations of Tannakian categories*, Deformation theory and quantum groups with applications to mathematical physics (Amherst, MA, 1990), Contemp. Math., vol. 134, Amer. Math. Soc., Providence, RI, 1992, [MR1187296](#) [DOI:10.1090/conm/134/1187296](#), pp. 325–349.

André Henriques
 Mathematical Institute
 University of Oxford
 Andrew Wiles Building
 Radcliffe Observatory Quarter
 Woodstock Road
 Oxford
 OX2 6GG
 United Kingdom
 a.g.henriques@uu.nl

David Penneys
 Department of Mathematics
 Ohio State University
 231 W. 18th Ave.
 Columbus, OH 43210
 USA
 penneys.2@osu.edu

James Tener
 Department of Mathematics
 University of California, Santa
 Barbara
 Santa Barbara, CA 93106
 USA
 jtener@math.ucsb.edu

