

CONCORDANCE INVARIANCE OF  
LEVINE-TRISTRAM SIGNATURES OF LINKS

MATTHIAS NAGEL, MARK POWELL

Received: September 7, 2016

Revised: December 14, 2016

Communicated by Ursula Hamenstädt

ABSTRACT. We determine for which complex numbers on the unit circle the Levine-Tristram signature and the nullity give rise to link concordance invariants.

2010 Mathematics Subject Classification: primary: 57M25, secondary: 57M27, 57N70

Keywords and Phrases: link concordance, Levine-Tristram signatures, nullity

1. INTRODUCTION

Let  $L \subset S^3$  be an  $m$ -component oriented link in the 3-sphere. Each connected, oriented Seifert surface  $F$  for  $L$  has a bilinear Seifert form defined by

$$V: H_1(F; \mathbb{Z}) \times H_1(F; \mathbb{Z}) \rightarrow \mathbb{Z}$$

$$(p[x], q[y]) \mapsto pq \operatorname{lk}(x^-, y),$$

where  $p, q \in \mathbb{Z}$ ,  $x, y$  are simple closed curves on  $F$  with associated homology classes  $[x], [y]$ , and  $x^-$  is a push-off of  $x$  in the negative normal direction of  $F$ . Given a unit modulus complex number  $z \in S^1 \setminus \{1\}$ , choose a basis for  $H_1(F; \mathbb{Z})$  and define the hermitian matrix

$$B(z) := (1 - z)V + (1 - \bar{z})V^T.$$

The *Levine-Tristram signature*  $\sigma_L(z)$  of  $L$  at  $z$  is defined to be the signature of  $B(z)$ , namely the number of positive eigenvalues minus the number of negative eigenvalues. The *nullity*  $\eta_L(z)$  of  $L$  at  $z$  is the dimension of the null space of  $B(z)$ . Both quantities can be shown to be invariants of the  $S$ -equivalence class of the Seifert matrix, and are therefore link invariants [Lev69, Tri69].

We say that two oriented  $m$ -component links  $L$  and  $J$  are *concordant* if there is a flat embedding into  $S^3 \times I$  of a disjoint union of  $m$  annuli  $A \subset S^3 \times I$ , such that the oriented boundary of  $A$  satisfies

$$\partial A = -L \sqcup J \subset -S^3 \sqcup S^3 = \partial(S^3 \times I).$$

An  $m$ -component link  $L$  is *slice* if it is concordant to the  $m$ -component unlink. The purpose of this paper is to answer the following question: *for which values of  $z$  are  $\sigma_L(z)$  and  $\eta_L(z)$  link concordance invariants?* We work in the topological category, in order to obtain the strongest possible results. In order to state our main theorem, we need one more definition.

DEFINITION 1.1. A complex number  $z \in S^1 \setminus \{1\}$  is a *Knotennullstelle* if there exists a Laurent polynomial  $p(t) \in \mathbb{Z}[t, t^{-1}]$  with  $p(1) = \pm 1$  and  $p(z) = 0$ .

Note that a complex number  $z \in S^1 \setminus \{1\}$  is a Knotennullstelle if and only if there exists a knot  $K$  whose Alexander polynomial  $\Delta_K$  has the property that  $\Delta_K(z) = 0$ . This follows from the fact that all Laurent polynomials  $q \in \mathbb{Z}[t, t^{-1}]$  with  $q(1) = \pm 1$  and  $q(t) = q(t^{-1})$  can be realised as Alexander polynomials of knots [BZ03, Theorem 8.13]. Here is our main theorem.

THEOREM 1.2. *The link invariants  $\sigma_L(z)$  and  $\eta_L(z)$  are concordance invariants if and only if  $z \in S^1 \setminus \{1\}$  does not arise as a Knotennullstelle.*

*Discussion of previously known results.* The first point to note is that, due to J. C. Cha and C. Livingston [CL04], when  $z$  is a Knotennullstelle neither  $\sigma_L(z)$  nor  $\eta_L(z)$  are link concordance invariants.

THEOREM 1.3 (Cha, Livingston). *For any Knotennullstelle  $z \in S^1 \setminus \{1\}$ , there exists a slice knot  $K$  with  $\sigma_K(z) \neq 0$  and  $\eta_K(z) \neq 0$ .*

Given a polynomial  $p(t)$  with  $p(1) = \pm 1$  and  $p(z) = 0$ , Cha and Livingston construct a matrix  $V$  with  $V - V^T$  nonsingular, with  $\det(tV - V^T)$  equal to  $p(t)p(t^{-1})$ , such that the upper left half-size block contains only zeroes, and such that  $\sigma(B(z)) \neq 0$ . Such a matrix can easily be realised as the Seifert matrix of a slice knot.

Some positive results on concordance invariance are also known. For  $z$  a prime power root of unity,  $\sigma_L(z)$  and  $\eta_L(z)$  are concordance invariants; see [Mur65], [Tri69] and [Kau78]. D. Cimasoni and V. Florens [CF08] dealt with multivariable signature and nullity concordance invariants, but again only at prime power roots of unity.

For the signature and nullity at algebraic numbers away from prime power roots of unity, we could not find any statements or results in the literature pertaining to our question. Levine [Lev07] studied the question in terms of  $\rho$ -invariants, but only discussed concordance invariance away from the roots of the Alexander polynomial.

By changing the rules slightly, one can obtain a concordance invariant for all  $z$ . The usual method is to define a function that is the average of the two

one-sided limits of the Levine-Tristram signature function. Let  $z = e^{i\theta} \in S^1$ , and consider:

$$\bar{\sigma}_L(z) := \frac{1}{2} \left( \lim_{\omega \rightarrow \theta_+} \sigma(B(e^{i\omega})) + \lim_{\omega \rightarrow \theta_-} \sigma(B(e^{i\omega})) \right).$$

Since prime power roots of unity are dense in  $S^1$ , this averaged signature function yields a concordance invariant at every  $z \in S^1$ . The earliest explicit observation of this that we could find was by Gordon in the survey article [Gor78]. One can also consider the averaged nullity function, to which similar remarks apply:

$$\bar{\eta}_L(z) := \frac{1}{2} \left( \lim_{\omega \rightarrow \theta_+} \eta(B(e^{i\omega})) + \lim_{\omega \rightarrow \theta_-} \eta(B(e^{i\omega})) \right).$$

In particular this is also a link concordance invariant.

Note that the function  $\sigma_L: S^1 \setminus \{1\} \rightarrow \mathbb{Z}$  is continuous away from roots of the Alexander polynomial  $\det(tV - V^T)$  of  $L$ . More generally one can consider the torsion Alexander polynomial  $\Delta_L^{\text{Tor}}$  of  $L$ , which by definition is the greatest common divisor of the  $(n - r) \times (n - r)$  minors of  $tV - V^T$ , where  $n$  is the size of  $V$  and  $r$  is the minimal nonnegative integer for which the set of minors contains a nonzero polynomial. The function  $\sigma_L$  is continuous away from the roots of the torsion Alexander polynomial  $\Delta_L^{\text{Tor}}$ , by [GL15, Theorem 2.1] (their  $A_L$  is our  $\Delta_L^{\text{Tor}}$ ).

Thus if  $z$  is not a root of the torsion Alexander polynomial of *any* link, the signature cannot jump at that value, and the signature function  $\sigma_L(z)$  equals the averaged signature function  $\bar{\sigma}_L(z)$  there. Since the averaged function is known to be a concordance invariant, the non-averaged function is also an invariant when  $z$  is not the root of any link's Alexander polynomial. The excitement happens when  $z$  is the root of the Alexander polynomial of some link, but is not the root of an Alexander polynomial of any knot. The averaged and non-averaged signature functions can differ at such  $z$ , but nevertheless *both* are concordance invariants. In Section 2 we will give an example which illustrates this difference, and gives an instance where the non-averaged function is more powerful. Similar examples were given in [GL15], but only with jumps occurring at prime power roots of unity.

Finally we remark that our proof of Theorem 1.2 covers the previously known cases of prime power roots of unity and transcendental numbers, as well as the new cases.

*Organisation of the paper.* The rest of the paper is organised as follows. In Section 2, we give an example of two links that are not concordant, where we use the signature and nullity functions at a root of their Alexander polynomials, which is not a prime power root of unity, to detect this fact. Section 4 proves that the nullity is a concordance invariant, and the corresponding fact for signatures is proven in Sections 5 and 6.

*Acknowledgements.* We thank Stefan Friedl, Pat Gilmer, Chuck Livingston and Andrew Ranicki for helpful discussions. In particular [GL15] inspired the question that led to this paper. We also thank the referee for helpful feedback. M.

Powell is supported by an NSERC Discovery grant. M. Nagel is supported by a CIRGET Postdoctoral Fellowship.

## 2. AN APPLICATION

In the introduction, for a link  $L$  we defined the signature function  $\sigma_L(z)$  and the nullity function  $\eta_L(z)$ , for each  $z \in S^1 \setminus \{1\}$ . From the characterisation in Theorem 1.2, one easily finds new values  $z$  for which it was not previously known that  $\sigma(z)$  and  $\eta(z)$  are concordance invariants. In Proposition 2.3, by exhibiting the obligatory explicit example, we show that these values give obstructions to concordance that are independent from previously known obstructions coming from the signature and nullity functions. We finish the section by constructing, in Proposition 2.5, a family of such examples for any algebraic number on  $S^1$ . Before the construction, we collect some facts on the set of roots of Alexander polynomials of links. We say that a complex number  $z \in S^1 \setminus \{1\}$  is a *Linknullstelle* if  $z$  is a root of a non-vanishing single variable Alexander polynomial of some link. We have the following inclusions:

$$\begin{aligned} \{\text{Knotennullstellen}\} &\subset \{\text{Linknullstellen}\} \subset S^1 \setminus \{1\} \\ &\cup \\ &\left\{ \begin{array}{l} \text{prime power} \\ \text{roots of 1} \end{array} \right\} \end{aligned}$$

We will see that these inclusions are strict. The two subsets of the set of Linknullstellen are disjoint, since no prime power root of unity can be a root of a polynomial that augments to  $\pm 1$ , because the corresponding cyclotomic polynomial augments to the prime. Moreover, the union of the Knotennullstellen and the prime power roots of unity is not exhaustive.

LEMMA 2.1.

- (1) *The set of Linknullstellen coincides with the set of algebraic numbers in  $S^1 \setminus \{1\}$ .*
- (2) *The number  $z_0 = \frac{3+4i}{5} \in S^1$  is an algebraic number, which is neither a Knotennullstelle nor a root of unity.*

*Proof.* Let  $z \in S^1 \setminus \{1\}$  be an algebraic number, so that  $p(z) = 0$  for some  $p \in \mathbb{Z}[t]$ . Let

$$q(t) := (t-1)^3 p(t) p(t^{-1}) \in \mathbb{Z}[t, t^{-1}].$$

We claim that there is a link  $L$  with single variable Alexander polynomial  $\Delta_L(t) = q(t)$ . Choose a 2-variable polynomial  $P(x, y) \in \mathbb{Z}[x^{\pm 1}, y^{\pm 1}]$  with  $P(t, t) = p(t)$ . Let

$$Q(x, y) := (x-1)(y-1)P(x, y)P(x^{-1}, y^{-1}).$$

A corollary [Hil12, Corollary 8.4.1] to Bailey's theorem [Bai77] states that any polynomial  $Q(x, y)$  in  $\mathbb{Z}[x^{\pm 1}, y^{\pm 1}]$ , with  $Q = \overline{Q}$  up to multiplication by  $\pm x^k y^\ell$ , and such that  $(x-1)(y-1)$  divides  $Q$ , is the Alexander polynomial of some 2-component link of linking number zero. Thus there exists a 2-component link  $L$  with 2-variable Alexander polynomial  $Q(x, y)$ .

The single variable Alexander polynomial  $\Delta_L(t)$  is obtained from the 2-variable Alexander polynomial of a 2-component link  $Q(x, y)$  as  $(t - 1)Q(t, t)$  [BZ03, Remark 9.18]. But

$$(t - 1)Q(t, t) = (t - 1)^3 P(t, t)P(t^{-1}, t^{-1}) = (t - 1)^3 p(t)p(t^{-1}) = q(t).$$

This completes the proof of the claim and therefore of (1): the set of Linknullstellen is the set of algebraic numbers lying on  $S^1 \setminus \{1\}$ .

For (2), first observe that the complex number  $z_0 := \frac{3+4i}{5}$  has unit modulus and that  $z_0$  is a zero of the polynomial

$$p(t) := 5t^2 - 6t + 5,$$

and therefore is an algebraic number. Note that no cyclotomic polynomial divides the polynomial  $p(t)$ . This can be checked for the first six by hand, and the rest have degree larger than 2. From Abel's irreducibility theorem, we learn that  $z_0$  is not a zero of a cyclotomic polynomial and thus is not a root of unity. Since  $p(1) = 4$  and  $p(t)$  is irreducible over  $\mathbb{Z}[t]$ ,  $z_0$  is not the root of any polynomial that augments to  $\pm 1$ . As a result,  $z_0$  is not a Knotennullstelle.  $\square$

Next we describe links  $L$  and  $L'$  whose signature and nullity functions are equal everywhere on  $S^1 \setminus \{1\}$  apart from at  $z_0$ , which will be a root of the Alexander polynomials of  $L$  and  $L'$ . We find these links by realising suitable Seifert forms.

EXAMPLE 2.2. Consider the following Seifert matrix:

$$V := \begin{pmatrix} 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 5 & -4 & 4 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 5 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -4 & 1 & -1 & 0 & 0 & 0 & 0 \\ 0 & 4 & 0 & 1 & 0 & 0 & 0 & 1 \end{pmatrix}.$$

This matrix represents the Seifert form of the 3-component link  $L$  given by the boundary of the Seifert surface shown in Figure 1. As usual, a box with  $n \in \mathbb{Z}$  inside denotes  $n$  full right-handed twists between two bands, made without introducing any twists into the individual bands. To see what we mean, observe that there are three instances in the figure of one full left-handed twist, otherwise known as  $-1$  full right-handed twists. The left-most twist is between the bands labelled  $e_1$  and  $e_5$ . To obtain the Seifert matrix, note that the beginning of each of the eight bands is labelled  $e_i$ , for  $i = 1, \dots, 8$ . Orient the bands clockwise and compute using  $V_{ij} = \text{lk}(e_i^-, e_j)$ , where the picture is understood to show the positive side of the Seifert surface.

Produce a link  $L'$  from  $L$  by removing the single twist in the right-most band, labelled  $e_8$  in Figure 1. This gives rise to a Seifert matrix  $V'$  for  $L'$  which is the same as  $V$ , except that the bottom right entry is a 0 instead of a 1.




FIGURE 1. Realisation of the Seifert form  $V$ .

Consider the sesquilinear form  $B$  over  $\mathbb{Q}[t^{\pm 1}]$  determined by the matrix

$$(1 - t)V + (1 - t^{-1})V^T.$$

The form  $B$  splits into a direct sum of sesquilinear forms. For a Laurent polynomial  $p(t) \in \mathbb{Q}[t^{\pm 1}]$ , abbreviate the form given by the  $2 \times 2$  matrix

$$\begin{pmatrix} 0 & p(t) \\ p(t^{-1}) & 0 \end{pmatrix}.$$

by  $[p(t)]$ . A calculation shows that  $B$  is congruent to the form

$$[t - 1] \oplus [t - 1] \oplus [t - 1] \oplus \begin{pmatrix} 0 & q(t) \\ q(t^{-1}) & -t^{-1} + 2 - t \end{pmatrix},$$

where the polynomial  $q(t)$  is

$$q(t) = t^{-1} \cdot (t - 1)^3 \cdot (5t^2 - 6t + 5).$$

On the other hand the corresponding sesquilinear form  $B'$  over  $\mathbb{Q}[t^{\pm 1}]$  for  $L'$  is equivalent to

$$[t - 1] \oplus [t - 1] \oplus [t - 1] \oplus [q(t)].$$

**PROPOSITION 2.3.** *Let  $z_0$  denote the algebraic number  $\frac{3+4i}{5}$ . The links  $L$  and  $L'$  constructed in Example 2.2 have the following properties.*

- (1) *If  $z$  is a root of unity, then  $\sigma_L(z) = \sigma_{L'}(z)$  and  $\eta_L(z) = \eta_{L'}(z)$ .*

(2) The averaged signature and nullity functions agree, i.e.

$$\bar{\sigma}_L(z) = \bar{\sigma}_{L'}(z) \text{ and } \bar{\eta}_L(z) = \bar{\eta}_{L'}(z)$$

for all  $z \in S^1 \setminus \{1\}$ .

(3) The signatures and nullities of  $L$  and  $L'$  at  $z_0$  differ:

$$\sigma_L(z_0) \neq \sigma_{L'}(z_0) \text{ and } \eta_L(z_0) \neq \eta_{L'}(z_0),$$

and so  $L$  is not concordant to  $L'$ .

*Proof.* Note that for any  $z \in \mathbb{C} \setminus \{0, 1\}$  with  $q(z) \neq 0$ , the form  $B(z)$  over  $\mathbb{C}$  is nonsingular and metabolic. The same holds for  $B'(z)$ . This implies that the signatures  $\text{sign } B(z)$  and  $\text{sign } B'(z)$  vanish. The nullities  $\eta_L(z), \eta_{L'}(z)$  are also both zero. Since the roots of  $q(z)$  are exactly  $z_0$  and  $\bar{z}_0$ , which are not roots of unity by Lemma 2.1, we obtain the first statement of the proposition. We also see that the averaged signature function on  $S^1 \setminus \{1\}$  and the averaged nullity function are identically zero, so we obtain the second statement.

From Lemma 2.1, we know that  $z_0 := \frac{3+4i}{5}$  is not a Knotennullstelle, and

$$\text{sign } B(z_0) = \text{sign} \begin{pmatrix} 0 & 0 \\ 0 & \frac{4}{5} \end{pmatrix} = 1.$$

Thus  $\sigma_L(z_0) = 1 = \eta_L(z_0)$ . On the other hand, for  $L'$  the matrix  $B'(z_0)$  is a  $2 \times 2$  zero matrix, so we have that  $\sigma_{L'}(z_0) = 0$  and  $\eta_{L'}(z_0) = 2$ . Both signatures and the nullities at  $z_0$  differ, so  $L$  and  $L'$  are not concordant by Theorem 1.2.  $\square$

*Remark 2.4.* One can also see that  $L$  and  $L'$  are not concordant using linking numbers.

A more systematic study of the construction of the example above leads to the following proposition.

PROPOSITION 2.5. *Let  $q(t) \in \mathbb{Z}[t]$  be a polynomial. Then there exists a natural number  $k > 0$  and a link  $L$  with Alexander polynomial  $\Delta_L(t) \doteq q(t^{-1})q(t)(t - 1)^k$  up to units in  $\mathbb{Z}[t, t^{-1}]$  such that*

- (1) the form  $B(z)$  of  $L$  is metabolic and nonsingular for all  $z \in S^1 \setminus \{1\}$  which are not roots of  $q(t)$ , so  $\sigma_L(z) = 0$ .
- (2) if  $z_0 \neq 1$  is a root of  $q(t)$  of unit modulus, then  $\sigma_L(z_0) \neq 0$ .

The proof of this proposition is based on ideas from [CL04].

*Proof.* Consider the size  $n + 1$  square matrix  $P$  with entries in  $\mathbb{Z}[y]$  given by

$$P(y) := \begin{pmatrix} 1 & y & 0 & & & ya_1 \\ 0 & 1 & y & & & ya_2 \\ \vdots & & \ddots & \ddots & & \vdots \\ & & & 1 & y & ya_{n-1} \\ 0 & & & 0 & 1 & ya_n \\ y & 0 & \dots & 0 & 0 & 0 \end{pmatrix},$$

with  $a_i$  integers. Over  $\mathbb{Z}[y^{\pm 1}]$ , the matrix  $P$  can be transformed via invertible row operations and column operations to the matrix

$$A(y) = \begin{pmatrix} 1 & 0 & 0 & & p(y) \\ 0 & 1 & 0 & & 0 \\ \vdots & & \ddots & \ddots & \vdots \\ & & & 1 & 0 & 0 \\ 0 & & & 0 & 1 & 0 \\ y & 0 & \dots & 0 & 0 & 0 \end{pmatrix}$$

with  $p(y) = b_1(y)$  where  $b_k(y) \in \mathbb{Z}[y]$  is defined by the recursion  $b_{k-1}(y) := y \cdot (a_k - b_k(y))$  and  $b_n(y) := y \cdot a_n$ . Notice that, up to units, we can arrange  $p(y)$  to be any polynomial in  $\mathbb{Z}[y^{\pm 1}]$  by choosing  $n$  sufficiently large and then suitable entries  $a_k \in \mathbb{Z}$ . That is, multiply by  $y^\ell$  so that the lowest order term is the linear term, and take  $(-1)^i a_i$  to be the coefficient of  $y^{i-1}$  in  $p(y)$ , for  $i = 2, \dots, n+1$ .

Pick the entries  $a_k$  so that if we evaluate  $p(y)$  at  $(t-1)$  we get the equality  $p(t-1) = q(t)(t-1)^k$  for a suitable integer  $k$ . Now consider the block matrix

$$V := \begin{pmatrix} 0 & V^u \\ V^b & Q(1) \end{pmatrix}$$

with

$$V^u = \begin{pmatrix} 0 & 1 & 0 & & a_1 \\ 0 & 0 & 1 & & a_2 \\ \vdots & & \ddots & \ddots & \vdots \\ & & & 0 & 1 & a_{n-1} \\ 0 & & & 0 & 0 & a_n \\ 1 & 0 & \dots & 0 & 0 & 0 \end{pmatrix} \quad V^b = \begin{pmatrix} -1 & 0 & 0 & & 1 \\ 1 & -1 & 0 & & 0 \\ 0 & 1 & \ddots & \ddots & \vdots \\ \vdots & & \ddots & -1 & 0 & 0 \\ 0 & & & 1 & -1 & 0 \\ a_1 & a_2 & \dots & a_{n-1} & a_n & 0 \end{pmatrix}$$

and

$$Q(y) = \begin{pmatrix} 0 & \dots & 0 & 0 \\ \vdots & \ddots & \vdots & \vdots \\ 0 & \dots & 0 & 0 \\ 0 & \dots & 0 & y \end{pmatrix}.$$

The matrix  $V$  is the Seifert matrix of a link as  $V - V^T$  is the intersection form of a genus  $n$  oriented surface with three boundary components. Let  $L$  be such a link, necessarily a 3-component link. We remark in passing that the matrix  $V$  from Example 2.2 is not a special case of the matrix  $V$  defined in the current proof, although it is close to being so.

Recall that  $B(z) = (1-z)V + (1-\bar{z})V^T = (\bar{z}-1) \cdot (zV - V^T)$ . The matrix  $V$  was constructed in such a way that

$$B(z) = \begin{pmatrix} 0 & (\bar{z}-1) \cdot P(z-1) \\ (z-1) \cdot P^T(\bar{z}-1) & Q(-\bar{z}-z+2) \end{pmatrix}.$$

Using the transformations associated to the above row and column operations, we see that  $B(z)$  is congruent to

$$B(z) \sim \begin{pmatrix} 0 & (\bar{z} - 1) \cdot A(z - 1) \\ (z - 1) \cdot A^T(\bar{z} - 1) & Q(-\bar{z} - z + 2) \end{pmatrix}.$$

Note that the matrix  $Q$  is unchanged by this congruency, because in the corresponding sequence of row and column operations, it never happens that the last row or column is added to another row or column.

We complete the proof of the proposition by showing that indeed the link  $L$  has the required properties. If  $z \in S^1 \setminus \{1\}$  is not a zero of  $q(t)$ , then also  $p(z) \neq 0$ . Consequently, the form  $B(z)$  is nonsingular and metabolic. On the other hand, if  $z \in S^1 \setminus \{1\}$  is a root of  $q(t)$ , then also  $p(z) = 0$ . In this case the Levine-Tristram form  $B(z)$  is a sum

$$B(z) = M \oplus \begin{pmatrix} 0 & 0 \\ 0 & -\bar{z} - z + 2 \end{pmatrix}$$

with  $M$  nonsingular and metabolic. Thus  $\sigma_L(z) = 1$ . □

*Remark 2.6.* Replace  $Q(1)$  with  $Q(0)$  in the construction of the matrix  $V$  in the proof of Proposition 2.5, to obtain a matrix  $V'$ . Using the same construction as in Example 2.2, the matrices  $V$  and  $V'$  give rise to links  $L$  and  $L'$  respectively, such that

$$\eta_L(z) = \eta_{L'}(z) \text{ and } \sigma_L(z) = \sigma_{L'}(z)$$

for every  $z \in S^1$  that is not a root of  $q(t)$ . Analogously to Example 2.2,  $L$  and  $L'$  are not concordant, but again this can also be seen using linking numbers. This leads to the following question. Does there exist a pair of links  $L$  and  $L'$ , with the same pairwise linking numbers, whose signature and nullity functions can only tell the concordance classes of the links apart at an isolated algebraic numbers  $z, \bar{z} \in S^1$  that are roots of the Alexander polynomial  $\Delta_L = \Delta_{L'}$ .

### 3. TWISTED HOMOLOGY AND INTEGRAL HOMOLOGY ISOMORPHISMS

Now we begin working towards the proof of Theorem 1.2. Fix  $z \in S^1 \setminus \{1\}$  to be a unit complex number that is not the root of any polynomial  $p(t) \in \mathbb{Z}[t]$  with  $p(1) = \pm 1$  i.e.  $z$  is not a Knotennullstelle. We denote the classifying space for the integers  $\mathbb{Z}$  by  $B\mathbb{Z}$ , which has the homotopy type of the circle  $S^1$ . Given a CW complex  $X$ , a map  $X \rightarrow B\mathbb{Z}$  induces a homomorphism  $\pi_1(X) \rightarrow \mathbb{Z}$ . This determines a representation

$$\alpha: \mathbb{Z}[\pi_1(X)] \rightarrow \mathbb{Z}[\mathbb{Z}] \xrightarrow{\text{ev}_z} \mathbb{C}$$

of the group ring of the fundamental group of  $X$ , with respect to which we can consider the twisted homology

$$H_i(X; \mathbb{C}^\alpha) := H_i \left( \mathbb{C} \otimes_{\mathbb{Z}[\pi_1(X)]} C_*(\tilde{X}) \right).$$

Let  $\Sigma \subset \mathbb{Z}[\mathbb{Z}]$  be the multiplicative subset of polynomials that map to  $\pm 1$  under the augmentation  $\varepsilon: \mathbb{Z}[\mathbb{Z}] \rightarrow \mathbb{Z}$ , that is  $\Sigma = \{p(t) \in \mathbb{Z}[\mathbb{Z}] \mid p(1) = \pm 1\}$ .

By inverting this subset we obtain the localisation  $\Sigma^{-1}\mathbb{Z}[\mathbb{Z}]$  of the Laurent polynomial ring. This has the following properties.

- (i) The canonical map  $\mathbb{Z}[\mathbb{Z}] \rightarrow \Sigma^{-1}\mathbb{Z}[\mathbb{Z}]$  is an inclusion, since  $\mathbb{Z}[\mathbb{Z}]$  is an integral domain.
- (ii) For any  $\mathbb{Z}[\mathbb{Z}]$ -module morphism  $f: M \rightarrow N$  of finitely generated free  $\mathbb{Z}[\mathbb{Z}]$ -modules such that the augmentation

$$\varepsilon(f) = \text{Id} \otimes f: \mathbb{Z} \otimes_{\mathbb{Z}[\mathbb{Z}]} M \rightarrow \mathbb{Z} \otimes_{\mathbb{Z}[\mathbb{Z}]} N$$

is an isomorphism, we have that

$$\text{Id} \otimes f: \Sigma^{-1}\mathbb{Z}[\mathbb{Z}] \otimes_{\mathbb{Z}[\mathbb{Z}]} M \rightarrow \Sigma^{-1}\mathbb{Z}[\mathbb{Z}] \otimes_{\mathbb{Z}[\mathbb{Z}]} N$$

is also an isomorphism.

The second property can be reduced to the following. Assume  $A$  is a matrix over  $\mathbb{Z}[\mathbb{Z}]$  such that  $\varepsilon(A)$  is invertible. Consequently, we have  $\det(\varepsilon(A)) = \pm 1$  and as  $\varepsilon(\det(A)) = \det(\varepsilon(A))$ , we deduce that  $\det(A) \in \Sigma$ . Therefore, the determinant  $\det(A)$  is invertible in the localisation  $\Sigma^{-1}\mathbb{Z}[\mathbb{Z}]$  and so is the matrix  $A$  over  $\Sigma^{-1}\mathbb{Z}[\mathbb{Z}]$ .

As the unit modulus complex number  $z$  that we have fixed is not a Knoten-nullstelle, the representation  $\alpha$  defined above factors through the localisation, i.e. evaluation at  $z$  determines a ring homomorphism  $\Sigma^{-1}\mathbb{Z}[\mathbb{Z}] \xrightarrow{\Sigma^{-1}\text{ev}_z} \mathbb{C}$  such that the ring homomorphisms  $\mathbb{Z}[\mathbb{Z}] \xrightarrow{\text{ev}_z} \mathbb{C}$  and

$$\mathbb{Z}[\mathbb{Z}] \rightarrow \Sigma^{-1}\mathbb{Z}[\mathbb{Z}] \xrightarrow{\Sigma^{-1}\text{ev}_z} \mathbb{C}$$

coincide.

LEMMA 3.1. *Let  $f: X \rightarrow Y$  be a map of finite CW complexes over  $S^1$ , that is there are maps  $g: X \rightarrow S^1$  and  $h: Y \rightarrow S^1$  such that  $h \circ f = g$ , and suppose that*

$$f_*: H_i(X; \mathbb{Z}) \xrightarrow{\cong} H_i(Y; \mathbb{Z})$$

*is an isomorphism for all  $i$ . Then*

$$f_*: H_i(X; \mathbb{C}^\alpha) \xrightarrow{\cong} H_i(Y; \mathbb{C}^\alpha)$$

*is also an isomorphism for all  $i$ .*

The lemma follows [COT03, Proposition 2.10]. The difference is that we use the well-known refinement that one does not need to invert all nonzero elements. We give the proof for the convenience of the reader. This is adapted from the proof given in [FP12].

*Proof.* The algebraic mapping cone  $D_* := \mathcal{C}(f_*: C_*(X; \mathbb{Z}) \rightarrow C_*(Y; \mathbb{Z}))$  has vanishing homology, and comprises finitely generated free  $\mathbb{Z}$ -modules. Therefore it is chain contractible. We claim that the chain contraction can be lifted to a chain contraction for  $\mathcal{C}(f_*: C_*(X; \Sigma^{-1}\mathbb{Z}[\mathbb{Z}]) \rightarrow C_*(Y; \Sigma^{-1}\mathbb{Z}[\mathbb{Z}]))$ , the mapping cone over the localisation  $\Sigma^{-1}\mathbb{Z}[\mathbb{Z}]$ .

To see this, let  $s: D_* \rightarrow D_{*+1}$  be a chain contraction, that is we have that  $\partial s_i + s_{i-1} \partial = \text{Id}_{D_i}$  for each  $i$ . Define  $\tilde{D}_* := \mathcal{C}(f_*: C_*(X; \mathbb{Z}[\mathbb{Z}]) \rightarrow C_*(Y; \mathbb{Z}[\mathbb{Z}]))$

and consider  $\varepsilon: \tilde{D}_* \rightarrow D_* = \mathbb{Z} \otimes_{\mathbb{Z}[\mathbb{Z}]} \tilde{D}_*$ , induced by the augmentation map. Denote  $E_* := \mathcal{C}(f_*: C_*(X; \Sigma^{-1}\mathbb{Z}[\mathbb{Z}]) \rightarrow C_*(Y; \Sigma^{-1}\mathbb{Z}[\mathbb{Z}]))$  and note that there is an inclusion  $\tilde{D}_i \rightarrow E_i = \Sigma^{-1}\mathbb{Z}[\mathbb{Z}] \otimes_{\mathbb{Z}[\mathbb{Z}]} \tilde{D}_i$ , induced by the localisation. Lift  $s$  to a map  $\tilde{s}: \tilde{D}_* \rightarrow \tilde{D}_{*+1}$ , as in the next diagram

$$\begin{array}{ccc} \tilde{D}_* & \overset{\tilde{s}}{\dashrightarrow} & \tilde{D}_{*+1} \\ \downarrow \varepsilon & & \downarrow \varepsilon \\ D_* & \xrightarrow{s} & D_{*+1}. \end{array}$$

The lifts exist since all modules are free and  $\varepsilon$  is surjective. But then we have that

$$f := d\tilde{s} + \tilde{s}d: \tilde{D}_* \rightarrow \tilde{D}_*$$

is a morphism of free  $\mathbb{Z}[\mathbb{Z}]$ -modules whose augmentation  $\varepsilon(f)$  is an isomorphism. Thus by property (ii) of  $\Sigma^{-1}\mathbb{Z}[\mathbb{Z}]$ ,  $f$  is also an isomorphism over  $\Sigma^{-1}\mathbb{Z}[\mathbb{Z}]$ , and so  $\tilde{s}$  determines a chain contraction for  $E_*$ . We therefore have that  $E_* = C_*(Y, X; \Sigma^{-1}\mathbb{Z}[\mathbb{Z}]) \simeq 0$  as claimed.

Next, tensor  $E_*$  with  $\mathbb{C}$  over the representation  $\alpha$ , to get that

$$\mathbb{C}^\alpha \otimes_{\Sigma^{-1}\mathbb{Z}[\mathbb{Z}]} C_*(Y, X; \Sigma^{-1}\mathbb{Z}[\mathbb{Z}]) = C_*(Y, X; \mathbb{C}^\alpha) \simeq 0.$$

Thus  $H_i(Y, X; \mathbb{C}^\alpha) = 0$  for all  $i$  and so  $f_*: H_i(X; \mathbb{C}^\alpha) \xrightarrow{\cong} H_i(Y; \mathbb{C}^\alpha)$  is an isomorphism for all  $i$  as desired.  $\square$

#### 4. CONCORDANCE INVARIANCE OF THE NULLITY

In this section we show concordance invariance of the nullity function away from the set of Knotennullstellen.

**DEFINITION 4.1** (Homology cobordism). A cobordism  $(W^{n+1}; M^n, N^n)$  between  $n$ -manifolds  $M$  and  $N$  is said to be a  $\mathbb{Z}$ -homology cobordism if the inclusion induced maps  $H_i(M; \mathbb{Z}) \rightarrow H_i(W; \mathbb{Z})$  and  $H_i(N; \mathbb{Z}) \rightarrow H_i(W; \mathbb{Z})$  are isomorphisms for all  $i \in \mathbb{Z}$ .

**THEOREM 4.2.** *Suppose that oriented  $m$ -component links  $L$  and  $J$  are concordant and that  $z \in S^1 \setminus \{1\}$  is not a Knotennullstelle. Then  $\eta_L(z) = \eta_J(z)$ .*

*Proof.* As in the statement suppose that  $z \in S^1 \setminus \{1\}$  is not a Knotennullstelle. Denote the exterior of the link  $L$  by  $X_L := S^3 \setminus \nu L$ . As above, let  $V$  be a matrix representing the Seifert form of  $L$  with respect to a Seifert surface  $F$  and a basis for  $H_1(F; \mathbb{Z})$ .

We assert that the matrix  $zV - V^T$  presents the homology  $H_1(X_L; \mathbb{C}^\alpha)$ . This can be seen as follows. Consider the infinite cyclic cover  $\overline{X}_L$  corresponding to the kernel of the homomorphism  $\pi_1(X_L) \rightarrow \mathbb{Z}$ , defined as the composition of the abelianisation  $\pi_1(X_L) \rightarrow H_1(X_L; \mathbb{Z}) \cong \mathbb{Z}^m$ , followed by the map  $(x_1, \dots, x_m) \mapsto \sum_{i=1}^m x_i$  i.e. each oriented meridian is sent to  $1 \in \mathbb{Z}$ . A decomposition of  $\overline{X}_L$  and the associated Mayer-Vietoris sequence [Lic97, Theorem

6.5] give rise the following presentation

$$\mathbb{C}[t^{\pm 1}] \otimes_{\mathbb{C}} H_1(F; \mathbb{C}) \xrightarrow{tV - V^T} \mathbb{C}[t^{\pm 1}] \otimes_{\mathbb{C}} H_1(F; \mathbb{C})^{\vee} \rightarrow H_1(\overline{X}_L; \mathbb{C}) \rightarrow 0,$$

where  $H_1(F; \mathbb{C})^{\vee}$  is the dual module  $\text{Hom}_{\mathbb{C}}(H_1(F; \mathbb{C}), \mathbb{C})$ . Apply the right-exact functor  $\mathbb{C}^{\alpha} \otimes_{\mathbb{C}[t^{\pm 1}]}$  to this sequence, to obtain the sequence

$$\mathbb{C}^{\alpha} \otimes_{\mathbb{C}} H_1(F; \mathbb{C}) \xrightarrow{zV - V^T} \mathbb{C}^{\alpha} \otimes_{\mathbb{C}} H_1(F; \mathbb{C})^{\vee} \rightarrow \mathbb{C}^{\alpha} \otimes_{\mathbb{C}[t^{\pm 1}]} H_1(\overline{X}_L; \mathbb{C}) \rightarrow 0.$$

As  $H_0(\overline{X}_L; \mathbb{C}) \cong \mathbb{C}$ , we have that  $\text{Tor}_1^{\mathbb{C}[t^{\pm 1}]}(H_0(\overline{X}_L; \mathbb{C}), \mathbb{C}^{\alpha}) = 0$  by the projective resolution

$$0 \rightarrow \mathbb{C}[t^{\pm 1}] \xrightarrow{(1-t)} \mathbb{C}[t^{\pm 1}] \rightarrow \mathbb{C} \rightarrow 0$$

and  $z \neq 1$ . Since  $\mathbb{C}[t^{\pm 1}]$  is a principal ideal domain, we can apply the universal coefficient theorem for homology to deduce that  $\mathbb{C}^{\alpha} \otimes_{\mathbb{C}[t^{\pm 1}]} H_1(\overline{X}_L; \mathbb{C}) = H_1(X_L; \mathbb{C}^{\alpha})$ . This completes the proof of the assertion that  $zV - V^T$  presents the homology  $H_1(X_L; \mathbb{C}^{\alpha})$ .

Next observe that  $(\bar{z} - 1)(zV - V^T) = (1 - z)V + (1 - \bar{z})V^T$  presents the same module as  $zV - V^T$ , since  $\bar{z} - 1$  is nonzero. The dimension of  $H_1(X_L; \mathbb{C}^{\alpha})$  therefore coincides with the nullity  $\eta_L(z)$ , which is by definition the nullity of the matrix  $(1 - z)V + (1 - \bar{z})V^T$ .

Now, let  $A \subset S^3 \times I$  be a union of annuli giving a concordance between  $L$  and  $J$ , and let  $W := S^3 \times I \setminus \nu A$ . Then  $W$  is a  $\mathbb{Z}$ -homology bordism between  $X_L$  and  $X_J$ ; this is a straightforward computation with Mayer-Vietoris sequences or with Alexander duality; see for example [FP14, Lemma 2.4]. Thus by two applications of Lemma 3.1, with  $Y = W$  and  $X = X_L$  and  $X = X_J$  respectively, we see that  $H_1(X_L; \mathbb{C}^{\alpha}) \cong H_1(W; \mathbb{C}^{\alpha}) \cong H_1(X_J; \mathbb{C}^{\alpha})$ , and so the nullities of  $L$  and  $J$  agree. We need that  $z$  is not a Knotennullstelle in order to apply Lemma 3.1. □

### 5. IDENTIFICATION OF THE SIGNATURE WITH THE SIGNATURE OF A 4-MANIFOLD

In the proof of Theorem 4.2, a key step was to reexpress the nullity  $\eta(z)$  of the form  $B(z)$  as a topological invariant of a 3-manifold, and then to use the bordism constructed from a concordance to relate the invariants. An analogous approach is used here to obtain the corresponding statement for the signature. Everything in this section is independent of whether  $z$  is a Knotennullstelle.

Recall that we fixed an oriented  $m$ -component link  $L \subset S^3$ , and that we picked a connected Seifert surface  $F$  for  $L$ . Denote the link complement by  $X_L := S^3 \setminus \nu L$ . First note that the fundamental class  $[F] \in H_2(F, \partial F; \mathbb{Z})$  of the Seifert surface  $F$  is independent of the choice of  $F$ . This follows from the fact that its Poincaré dual is characterised as the unique cohomology class  $\xi \in H^1(X_L; \mathbb{Z})$  mapping each meridian  $\mu$  to  $\xi(\mu) = 1$ .

The boundary of  $F \subset S^3 \setminus \nu L$  is a collection of embedded curves in the boundary tori that we refer to as the *attaching curves*. The attaching curves together with the meridians determine a framing of each boundary torus of  $X_L$ . Also,

this framing depends solely on  $[F]$ , since the connecting homomorphism of the pair  $(X_L, \partial X_L)$  maps  $\partial[F] = [\partial F]$ .

With respect to this framing, we can consider the Dehn filling of slope zero, resulting in the closed 3-manifold  $M_L$ . By definition, to obtain  $M_L$  attach a disc to each of the attaching curves, and then afterwards fill each of the resulting boundary spheres with a 3-ball.

DEFINITION 5.1. The framing of the boundary tori of  $X_L$  constructed above is called the *Seifert framing*. The *Seifert surgery* on  $L$  is the 3-manifold  $M_L$  constructed above.

Remark 5.2. For links there is no reason for this framing to agree with the zero-framing of each individual component.

Collapsing the complement of a tubular neighbourhood of the Seifert surface  $F$  gives rise to map  $S^3 \setminus \nu L \rightarrow S^1 = B\mathbb{Z}$ , which extends to a map from the Seifert surgery  $\phi: M_L \rightarrow B\mathbb{Z}$ . To see this in more detail, parametrise a regular neighbourhood of  $F$  as  $F \times [-1, 1]$ , with  $F$  as  $F \times \{0\}$ . The intersection of this parametrised neighbourhood with each component of  $\partial F$  determines a parametrised subset  $S^1 \times [-1, 1] \subset S^1 \times S^1 \subseteq \partial F$ . Extend this to a subset  $D^2 \times [-1, 1] \subset D^2 \times S^1$  for each of the Dehn filling solid tori  $D^2 \times S^1$  in  $M_L$ . Now define

$$\begin{aligned} \phi: M_L &\rightarrow S^1 = B\mathbb{Z} \\ x &\mapsto \begin{cases} e^{\pi i t} & x = (f, t) \in (F \cup \bigsqcup^m D^2) \times [-1, 1] \\ -1 & \text{otherwise.} \end{cases} \end{aligned}$$

The map  $\phi$  classifies the image of the fundamental class of the capped-off Seifert surface in  $M_L$ , in the sense that  $[\phi]$  maps to  $[F \cup \bigsqcup^m D^2]$  under  $[M_L, S^1] \xrightarrow{\cong} H^1(M_L; \mathbb{Z}) \xrightarrow{\cong} H_2(M_L; \mathbb{Z})$ . Recall that the homology class  $[F \cup \bigsqcup^m D^2] \in H_2(M_L; \mathbb{Z})$  only depends on the isotopy class of  $L$  and so also the homotopy class of  $\phi$  does not depend on the Seifert surface  $F$ . The manifold  $M_L$  together with the map  $\phi$  defines an element  $[(M_L, \phi)] \in \Omega_3(B\mathbb{Z})$ , where  $\Omega_k(X)$  denotes the bordism group of oriented, topological  $k$ -dimensional manifolds with a map to  $X$ . Recall that cobordism is a generalised homology theory fulfilling the suspension axiom, see e.g. [tD08, Chapter 21] and [May99, Section 14.4]. As a consequence, we obtain

$$\tilde{\Omega}_3(B\mathbb{Z}) = \tilde{\Omega}_3(S^1) = \tilde{\Omega}_3(\Sigma S^0) \cong \tilde{\Omega}_2(S^0) = \Omega_2(\text{pt}) = 0.$$

Thus  $\Omega_3(B\mathbb{Z}) \cong \Omega_3(\text{pt}) = 0$  [Roh53].

The group  $\Omega_3(B\mathbb{Z}) \cong \Omega_3 \oplus \Omega_2 = 0 \oplus 0 = 0$  is trivial, and we can make use of this fact to define a signature defect invariant, as follows.

For any oriented 3-manifold  $M$  with a map  $\phi: M \rightarrow B\mathbb{Z}$ , we will define an integer for each complex number  $z \in S^1$ . Since  $\Omega_3(B\mathbb{Z}) = 0$ , there exists a 4-manifold  $W$  with boundary  $M$  and a map  $\Phi: W \rightarrow B\mathbb{Z}$  extending the map  $M \rightarrow B\mathbb{Z}$  on the boundary. Similarly to before, an element  $z \in S^1$  determines a representation

$$\alpha: \mathbb{Z}[\pi_1(W)] \xrightarrow{\Phi} \mathbb{Z}[\mathbb{Z}] \xrightarrow{t \mapsto z} \mathbb{C}.$$

Consider the twisted homology  $H_i(W; \mathbb{C}^\alpha)$ , and consider the intersection form  $\lambda_\alpha(W)$  on the quotient  $H_2(W; \mathbb{C}^\alpha)/\text{im } H_2(M; \mathbb{C}^\alpha)$ . Define the promised integer

$$\sigma(M, \phi, z) := \sigma(\lambda_\alpha(W)) - \sigma(W),$$

where  $\sigma(W)$  is the ordinary signature of the intersection form on  $W$ .

The proof of the following proposition is known for the coefficient system  $\mathbb{Q}(t)$ , e.g. [Pow16]. For the convenience of the reader, we sketch the key steps for an adaptation to  $\mathbb{C}^\alpha$ .

PROPOSITION 5.3.

- (i) *The intersection form  $\lambda_\alpha(W)$  is nonsingular.*
- (ii) *The signature defect  $\sigma(M, \phi, z)$  is independent of the choice of 4-manifold  $W$ .*

*Proof.* The long exact sequence of the pair  $(W, \partial W) = (W, M)$  gives rise to the following commutative diagram

$$\begin{array}{ccccccc} \dots & \longrightarrow & H_2(\partial W; \mathbb{C}^\alpha) & \longrightarrow & H_2(W; \mathbb{C}^\alpha) & \longrightarrow & H_2(W, \partial W; \mathbb{C}^\alpha) \longrightarrow \dots \\ & & & & \searrow & & \downarrow \text{PD}_W^{-1} \\ & & & & & & H^2(W; \mathbb{C}^\alpha) \\ & & & & & & \downarrow \kappa \\ & & & & & & (H_2(W; \mathbb{C}^\alpha))^\vee, \end{array}$$

where for a  $\mathbb{C}$ -module  $P$  we denote its dual module by  $P^\vee := \text{Hom}_{\mathbb{C}}(P, \mathbb{C})$ . Since Poincaré-Lefschetz duality  $\text{PD}_W$  and the Kronecker pairing  $\kappa$  are isomorphisms, we obtain an injective map  $H_2(W; \mathbb{C}^\alpha)/\text{im } H_2(M; \mathbb{C}^\alpha) \rightarrow H_2(W; \mathbb{C}^\alpha)^\vee$ . This map descends to

$$\lambda_\alpha: H_2(W; \mathbb{C}^\alpha)/\text{im } H_2(M; \mathbb{C}^\alpha) \rightarrow (H_2(W; \mathbb{C}^\alpha)/\text{im } H_2(\partial W; \mathbb{C}^\alpha))^\vee,$$

so that the diagram below commutes:

$$\begin{array}{ccc} H_2(W; \mathbb{C}^\alpha)/\text{im } H_2(M; \mathbb{C}^\alpha) & \xrightarrow{\quad} & H_2(W; \mathbb{C}^\alpha)^\vee \\ & \searrow \lambda_\alpha & \uparrow \\ & & (H_2(W; \mathbb{C}^\alpha)/\text{im } H_2(\partial W; \mathbb{C}^\alpha))^\vee. \end{array}$$

Consequently, the form  $\lambda_\alpha$  is nondegenerate, and so it is nonsingular since it is a form over the field  $\mathbb{C}$ .

We proceed with the second statement of the proposition, namely independence of  $\sigma(M, \phi, z)$  on the choice of  $W$ . Suppose that we are given two 4-manifolds  $W^+, W^-$ , both with boundary  $\partial W^\pm = M$ , and a map  $\Phi^\pm: W^\pm \rightarrow B\mathbb{Z}$  extending  $\phi: M \rightarrow B\mathbb{Z}$ . Temporarily, define the signature defects arising from the two choices to be

$$\sigma(W^\pm, \Phi^\pm, z) := \sigma(\lambda_\alpha(W^\pm)) - \sigma(W^\pm).$$

We will show that  $\sigma(W^+, \Phi^+, z) = \sigma(W^-, \Phi^-, z)$ , and thus that  $\sigma(M, \phi, z)$  is a well-defined integer, so our original notation was justified.

Glue  $W^+$  and  $\overline{W^-}$  together along  $M$ , to obtain a closed manifold  $U$ , together with a map  $\Phi: U \rightarrow B\mathbb{Z}$ . By Novikov additivity, we learn that

$$\sigma_z(U, \Phi) := \sigma(\lambda_\alpha(U)) - \sigma(U) = \sigma(W^+, \Phi^+, z) - \sigma(W^-, \Phi^-, z).$$

This defect  $\sigma_z(U, \Phi)$  can be promoted to a bordism invariant  $\sigma_z: \Omega_4(B\mathbb{Z}) \rightarrow \mathbb{Z}$ , see e.g. [Pow16, Proof of Lemma 3.2] and replace  $\mathbb{Q}(t)$  coefficients with  $\mathbb{C}^\alpha$  coefficients.

*Claim.* The map  $\sigma_z: \Omega_4(B\mathbb{Z}) \rightarrow \mathbb{Z}$  is the zero map.

Let  $U$  be a closed 4-manifold together with a map  $\Phi: U \rightarrow S^1$ , representing an element of  $\Omega_4(B\mathbb{Z})$ . By the axioms of generalised homology theories, we have

$$\tilde{\Omega}_4(S^1) = \tilde{\Omega}_4(\Sigma S^0) \cong \tilde{\Omega}_3(S^0) = \Omega_3(\text{pt}) = 0.$$

Thus an inclusion  $\text{pt} \rightarrow S^1$  induced an isomorphism  $\Omega_4(\text{pt}) \xrightarrow{\cong} \Omega_4(S^1)$ . So  $(U, \Phi)$  is bordant over  $S^1$  to a 4-manifold  $U'$  with a null-homotopic map  $\Phi'$  to  $S^1$ . In this case the local coefficient system  $\mathbb{C}^\alpha$  is just the trivial representation  $\mathbb{C}$ . Consequently, we have  $\lambda_\alpha(U') = \lambda(U')$ , so  $\sigma_z(U', \Phi') = 0$ . By bordism invariance,  $\sigma_z(U, \Phi) = 0$ , which completes the proof of the claim.

Now the independence of  $\sigma(M, \Phi, z)$  on the choice of  $W$  follows from

$$0 = \sigma_z(U, \Phi) = \sigma(W^+, \Phi^+, z) - \sigma(W^-, \Phi^-, z).$$

□

Now that we have constructed an invariant, we need to relate it to the Levine-Tristram signatures. Recall that  $L$  is an oriented link, that  $M_L$  is the Seifert surgery, and that we constructed a canonical map  $\phi: M_L \rightarrow S^1$ , well-defined up to homotopy.

LEMMA 5.4. *Suppose that  $z \in S^1 \setminus \{1\}$  and let  $\phi: M_L \rightarrow S^1$  be the map defined at the beginning of this section. Then we have*

$$\sigma(M_L, \phi, z) = \sigma_L(z).$$

*Proof.* Construct a 4-manifold with boundary  $M_L$  as follows. Let  $F$  be a connected Seifert surface for  $L$ . Push the Seifert surface into  $D^4$  and consider its complement  $V_F := D^4 \setminus \nu F$ . Note that if we cap  $F$  off with  $m$  2-discs, we obtain a closed surface. Let  $H$  be a 3-dimensional handlebody whose boundary is this surface. Note that  $\partial V_F = X_L \cup F \times S^1$ . Then define

$$W_F := V_F \cup_{F \times S^1} H \times S^1.$$

Note that  $\partial W_F = M_L$ . By [Ko89, pp. 538-9] and [COT04, Lemma 5.4], we have that  $\sigma(W_F) = 0$  and  $\lambda_z(W_F) = (1 - z)V + (1 - \bar{z})V^T$ . Therefore

$$\sigma(\lambda_z(W_F)) - \sigma(W_F) = \sigma((1 - z)V + (1 - \bar{z})V^T) = \sigma(B(z)).$$

□

## 6. CONCORDANCE INVARIANCE OF THE SIGNATURE

We start with a straightforward lemma, then we prove the final part of the main theorem. Recall that the complement  $X_L$  and the Seifert surgery  $M_L$  are both equipped with a homotopy class of a map to  $S^1$ , or equivalently with a cohomology class. For the link complement  $X_L$ , this class  $\xi_L \in H^1(X_L; \mathbb{Z})$  is characterised by the property that it sends each oriented meridian to 1.

LEMMA 6.1. *Let  $L$  and  $J$  be concordant links. Their Seifert surgeries  $M_L$  and  $M_J$  are homology bordant over  $S^1$ .*

*Proof.* Denote the maps to  $S^1$  by  $\phi_L: M_L \rightarrow S^1$  and  $\phi_J: M_J \rightarrow S^1$ , and denote the corresponding cohomology classes by  $\xi_L \in H^1(M_L; \mathbb{Z})$  and  $\xi_J \in H^1(M_J; \mathbb{Z})$ . Define  $X_L := S^3 \setminus \nu L$  and  $X_J := S^3 \setminus \nu J$ . Let  $A \subset S^3 \times I$  be an embedding of a disjoint union of annuli giving a concordance between  $L$  and  $J$ .

Fix a tubular neighbourhood  $\nu A = A \times D^2$  of the annulus  $A$  with a trivialisation. Denote  $W_A := S^3 \times I \setminus \nu A$ , whose boundary consists of the union of  $X_L$ ,  $X_J$ , and a piece identified with the total space of the unit sphere bundle  $A \times S^1$  of  $\nu A$ . As usual, we refer to a representative  $\{\text{pt}\} \times S^1$  for the  $S^1$  factor in  $A \times S^1$  as a *meridian* of  $A$ . Note that the inclusions  $X_L \subset W_A$  and  $X_J \subset W_A$  map the meridians in the link complements to the meridians in  $W_A$ .

*Claim.* There exists a cohomology class  $\xi_A \in H^1(W_A; \mathbb{Z})$  mapping each meridian  $\mu_A$  of  $A$  to 1.

This can be seen by the Mayer-Vietoris sequence

$$H^1(\nu A; \mathbb{Z}) \oplus H^1(W_A; \mathbb{Z}) \rightarrow H^1(\partial \nu A; \mathbb{Z}) \rightarrow H^2(S^3 \times I; \mathbb{Z}) = 0,$$

in which the map  $H^1(\nu A; \mathbb{Z}) \cong \mathbb{Z}^m \rightarrow H^1(\partial \nu A; \mathbb{Z}) \cong (\mathbb{Z} \oplus \mathbb{Z})^m$  is given by  $1 \mapsto (1, 0)$  on each of the  $m$  summands. That is, the homology classes of the meridians of  $\partial \nu A \cong A \times S^1$  do not lie in the image of this surjective map, so they must lie in the image of  $H^1(W_A; \mathbb{Z})$ . This completes the proof of the claim.

It follows that  $\xi_A$  is pulled back to the unique classes  $\xi_L$  and  $\xi_J$  that map the meridians in the link complements to 1. Using the natural isomorphism between the functors  $[-, S^1]$  and  $H^1(-; \mathbb{Z})$ , find a map  $\phi_W: W_A \rightarrow S^1$  that restricts to the prescribed map  $\phi_L \sqcup \phi_J: X_L \sqcup X_J \rightarrow S^1$  on the boundary.

Up to isotopy, there is a unique product structure on an annulus  $A = S^1 \times I$ . Having fixed such a structure, we consider the manifold

$$Y := W_A \cup_{A \times S^1} \bigsqcup^m (D^2 \times S^1 \times I).$$

The gluing is done in such a way as to restrict on  $\bigsqcup^m S^1 \times S^1 \times \{i\}$ , for  $i = 0, 1$ , to the gluing of the Seifert surgery on  $X_L$  and  $X_J$ . By construction, this gives a bordism between  $M_L$  and  $M_J$ .

Note that the map  $\phi_W$  and the projection  $A \times S^1 \rightarrow S^1$  glue together to give a map  $\phi_Y: Y \rightarrow S^1$ . Equipped with this map,  $(Y, \phi_Y)$  is an  $S^1$ -bordism between  $(M_L, \phi_L)$  and  $(M_J, \phi_J)$ .

Finally, we assert that  $Y$  is a homology bordism. To see this, first observe, as in the proof of Theorem 4.2, that  $W_A$  is a homology bordism from  $X_L$  to  $X_J$ . Flagrantly,  $A \times S^1$  is a homology bordism from  $S^1 \times S^1$  to itself, and  $\bigsqcup^m (D^2 \times S^1 \times I)$  is a homology bordism from  $\bigsqcup^m D^2 \times S^1$  to itself. Gluing two homology bordisms together along a homology bordism, with the same maps on homology induced by the gluings for  $M_L$ ,  $M_J$  and  $Y$ , it follows easily from the Mayer-Vietoris sequence and the five lemma that  $Y$  is a homology bordism.  $\square$

**THEOREM 6.2.** *Suppose that oriented  $m$ -component links  $L$  and  $J$  are concordant and that  $z \in S^1 \setminus \{1\}$  is not a Knotennullstelle. Then  $\sigma_L(z) = \sigma_J(z)$ .*

*Proof.* As in the statement of the theorem, suppose that  $z \in S^1 \setminus \{1\}$  is not a Knotennullstelle. Let  $W_{LJ}$  be a homology bordism between the Seifert surgeries  $M_L$  and  $M_J$ , whose existence is guaranteed by Lemma 6.1. Let  $W_J$  be a 4-manifold that gives a nullbordism of  $M_J$  over  $B\mathbb{Z}$ , and define  $W_L := W_{LJ} \cup_{M_J} W_J$ .

The signature of the intersection form on  $H_2(W_L; \mathbb{C}^\alpha)/H_2(M_L; \mathbb{C}^\alpha)$ , together with the ordinary signature over  $\mathbb{Z}$ , determines the signature  $\sigma_L(z)$  by Section 5. Similarly, the signature of the intersection form on the quotient  $H_2(W_J; \mathbb{C}^\alpha)/H_2(M_J; \mathbb{C}^\alpha)$  and the ordinary signature of  $W_J$  determine the signature  $\sigma_J(z)$ . By Lemma 3.1, we have homology isomorphisms

$$H_2(M_L; \mathbb{C}^\alpha) \xrightarrow{\cong} H_2(W_{LJ}; \mathbb{C}^\alpha) \text{ and } H_2(M_J; \mathbb{C}^\alpha) \xrightarrow{\cong} H_2(W_{LJ}; \mathbb{C}^\alpha).$$

It follows that every class in  $H_2(W_L; \mathbb{C}^\alpha)$  has a representative in  $W_J$ , that

$$H_2(W_L; \mathbb{C}^\alpha)/H_2(M_L; \mathbb{C}^\alpha) \cong H_2(W_J; \mathbb{C}^\alpha)/H_2(M_J; \mathbb{C}^\alpha),$$

and that this isomorphism induces an isometry of the intersection forms. Thus the twisted signatures of both intersection forms are equal. We needed that  $z$  is not a Knotennullstelle in order to apply Lemma 3.1 in the preceding argument. The same argument over  $\mathbb{Z}$  implies that the ordinary signatures also coincide, that is  $\sigma(W_L) = \sigma(W_J)$ . Therefore  $\sigma(M_L, \phi_L, z) = \sigma(M_J, \phi_J, z)$ , and so  $\sigma_L(z) = \sigma_J(z)$  by Lemma 5.4. Thus the Levine-Tristram signature at  $z$  is a concordance invariant, as desired.  $\square$

#### REFERENCES

- [Bai77] James L. Bailey. *Alexander invariants of links*. Thesis – The University of British Columbia. ProQuest LLC, Ann Arbor, MI, 1977.
- [BZ03] Gerhard Burde and Heiner Zieschang. *Knots*, volume 5 of *de Gruyter Studies in Mathematics*. Walter de Gruyter & Co., Berlin, second edition, 2003.
- [CF08] David Cimasoni and Vincent Florens. Generalized Seifert surfaces and signatures of colored links. *Trans. Amer. Math. Soc.*, 360(3):1223–1264 (electronic), 2008.

- [CL04] Jae Choon Cha and Charles Livingston. Knot signature functions are independent. *Proc. Amer. Math. Soc.*, 132(9):2809–2816 (electronic), 2004.
- [COT03] Tim D. Cochran, Kent E. Orr, and Peter Teichner. Knot concordance, Whitney towers and  $L^2$ -signatures. *Ann. of Math. (2)*, 157(2):433–519, 2003.
- [COT04] Tim D. Cochran, Kent E. Orr, and Peter Teichner. Structure in the classical knot concordance group. *Comment. Math. Helv.*, 79(1):105–123, 2004.
- [FP12] Stefan Friedl and Mark Powell. An injectivity theorem for Casson-Gordon type representations relating to the concordance of knots and links. *Bull. Korean Math. Soc.*, 49(2):395–409, 2012.
- [FP14] Stefan Friedl and Mark Powell. Cobordisms to weakly splittable links. *Proc. Amer. Math. Soc.*, 142:703–712, 2014.
- [GL15] Patrick M. Gilmer and Charles Livingston. Signature jumps and Alexander polynomials for links. *To appear in Proc. Amer. Math. Soc.*, 2015. Preprint available at arXiv:1508.04394.
- [Gor78] Cameron Gordon. Some aspects of classical knot theory. In *Knot theory (Proc. Sem., Plans-sur-Bex, 1977)*, pages 1–60. Springer, Berlin, 1978.
- [Hil12] Jonathan Hillman. *Algebraic invariants of links*, volume 52 of *Series on Knots and Everything*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, second edition, 2012.
- [Kau78] Louis H. Kauffman. Signature of branched fibrations. In Jean-Claude Hausmann, editor, *Knot Theory*, volume 685 of *Lecture Notes in Mathematics*, pages 203–217. Springer Berlin Heidelberg, 1978.
- [Ko89] Ki Hyoung Ko. A Seifert-matrix interpretation of Cappell and Shaneson’s approach to link cobordisms. *Math. Proc. Cambridge Philos. Soc.*, 106(3):531–545, 1989.
- [Lev69] Jerome P. Levine. Invariants of knot cobordism. *Invent. Math.* 8 (1969), 98–110; addendum, *ibid.*, 8:355, 1969.
- [Lev07] Jerome P. Levine. Concordance of boundary links. *J. Knot Theory Ramifications*, 16(9):1111–1120, 2007.
- [Lic97] W. B. Raymond Lickorish. *An introduction to knot theory*, volume 175 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1997.
- [May99] J. Peter May. *A concise course in algebraic topology*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 1999.
- [Mur65] Kunio Murasugi. On a certain numerical invariant of link types. *Trans. Amer. Math. Soc.*, 117:387–422, 1965.
- [Pow16] Mark Powell. The four genus of a link, Levine-Tristram signatures and satellites. *To appear in Journal of Knot Theory and its Ramifications, Special issue in honour of Tim Cochran*, 2016. Preprint available at arXiv:1605.06833.

- [Roh53] Vladimir A. Rohlin. Intrinsic homologies. *Doklady Akad. Nauk SSSR (N.S.)*, 89:789–792, 1953.
- [tD08] Tammo tom Dieck. *Algebraic topology*. EMS Textbooks in Mathematics. European Mathematical Society (EMS), Zürich, 2008.
- [Tri69] Andrew G. Tristram. Some cobordism invariants for links. *Proc. Cambridge Philos. Soc.*, 66:251–264, 1969.

Matthias Nagel  
Département de Mathématiques  
Université du Québec à Montréal  
Canada  
nagel@cirget.ca

Mark Powell  
Département de Mathématiques  
Université du Québec à Montréal  
Canada  
mark@cirget.ca

