

PSEUDO-DIFFERENTIAL OPERATORS, WIGNER TRANSFORM
AND WEYL SYSTEMS ON TYPE I LOCALLY COMPACT GROUPSMARIUS MĂNTOIU ¹ AND MICHAEL RUZHANSKY ²

Received: February 20, 2016

Revised: April 27, 2017

Communicated by Stefan Teufel

ABSTRACT. Let G be a unimodular type I second countable locally compact group and let \widehat{G} be its unitary dual. We introduce and study a global pseudo-differential calculus for operator-valued symbols defined on $G \times \widehat{G}$, and its relations to suitably defined Wigner transforms and Weyl systems. We also unveil its connections with crossed products C^* -algebras associated to certain C^* -dynamical systems, and apply it to the spectral analysis of covariant families of operators. Applications are given to nilpotent Lie groups, in which case we relate quantizations with operator-valued and scalar-valued symbols.

2010 Mathematics Subject Classification: Primary 46L65, 47G30; Secondary 22D10, 22D25.

Keywords and Phrases: locally compact group, nilpotent Lie group, non-commutative Plancherel theorem, pseudo-differential operator, C^* -algebra, dynamical system.

1 INTRODUCTION

Let G be a locally compact group with unitary dual \widehat{G} , composed of classes of unitary equivalence of strongly continuous irreducible representations. To have a manageable

¹MM was supported by Núcleo Milenio de Física Matemática RC120002 and the Fondecyt Project 1160359.

²MR was supported by the EPSRC Grant EP/K039407/1 and by the Leverhulme Research Grant RPG-2014-02.

The authors were also partly supported by EPSRC Mathematics Platform grant EP/I019111/1.

Fourier transformation, it will be assumed second countable, unimodular and postliminal (type I). The formula

$$[\text{Op}(a)u](x) = \int_G \left(\int_{\widehat{G}} \text{Tr}_\xi [\xi(y^{-1}x)a(x, \xi)] d\widehat{m}(\xi) \right) u(y) dm(y) \quad (1)$$

is our starting point for a global pseudo-differential calculus on G ; it involves operator-valued symbols defined on $G \times \widehat{G}$. In (1) dm is the Haar measure of the group G , $d\widehat{m}$ is the Plancherel measure on the space \widehat{G} and for the pair (x, ξ) formed of an element x of the group and a unitary irreducible representation $\xi : G \rightarrow \mathbb{B}(\mathcal{H}_\xi)$, the symbol $a(x, \xi)$ is essentially assumed to be a trace-class operator in the representation Hilbert space \mathcal{H}_ξ . In further extensions of the theory it is important to also include densely defined symbols to cover, for example, differential operators on Lie groups (in which case one can make sense of (1) for such $a(x, \xi)$ by letting it act on the dense in \mathcal{H}_ξ subspace of smooth vectors of the representation ξ , see [19]).

Particular cases of (1) have been previously initiated in [39, 41] and then intensively developed further in [8, 9, 11, 12, 17, 42, 43] for compact Lie groups, and in [18, 19, 20] for large classes of nilpotent Lie groups (graded Lie groups), as far-reaching versions of the usual Kohn-Nirenberg quantization on $G = \mathbb{R}^n$, cf. [22]. The idea to use the irreducible representation theory of a type I group in defining pseudo-differential operators seems to originate in [44, Sect. I.2], but it has not been developed before in such a generality. All the articles cited above already contain historical discussions and references to the literature treating pseudo-differential operators (quantization) in group-like situations, so we are not going to try to put this subject in a larger perspective.

Let us just say that an approach involving pseudo-differential operators with representation-theoretical operator-valued symbols has the important privilege of being global. On most of the smooth manifolds there is no notion of full scalar-valued symbol for a pseudo-differential operator defined using local coordinates. This is unfortunately true even in the rather simple case of a compact Lie group, for which the local theory, only leading to a principal symbol, has been shown to be equivalent to the global operator-valued one (cf. [39, 42]). On the other hand, in the present article we are not going to rely on compactness, on the nice properties implied by nilpotency, not even on the smooth structure of a Lie group. In the category of type I second countable locally compact groups one has a good integration theory on G and a manageable integration theory on \widehat{G} , allowing a general form of the Plancherel theorem, and this is enough to develop the basic features of a quantization. Unimodularity has been assumed, for simplicity, but by using tools from [10] it might be possible to develop the theory without it.

More structured cases (still more general than those studied before) will hopefully be analysed in the close future, having the present paper as a framework and a starting point. In particular, classes of symbols of Hörmander type would need more than a smooth structure on G . The smooth theory, still to be developed, seems technically difficult if the class of Lie groups is kept very general. Of course, only in this setting

one could hope to cover differential operators and certain types of connected applications. On the other hand, the setting of our article allows studying multiplication and invariant operators as very particular cases, cf. Subsection 7.3.

The formula (1) is a generalisation of the Kohn-Nirenberg quantization rule for the particular case $G = \mathbb{R}^n$. But for \mathbb{R}^n there are also the so-called τ -quantizations

$$[\text{Op}^\tau(a)u](x) = \int_{\mathbb{R}^n} \left(\int_{\mathbb{R}^n} a((1 - \tau)x + \tau y, \eta) e^{i(x-y)\eta} d\eta \right) u(y) dy,$$

related to ordering issues, with $\tau \in [0, 1]$, and the Kohn-Nirenberg quantization is its special case for $\tau = 0$. It is possible to provide extensions of the pseudo-differential calculus on type I groups corresponding to any measurable function $\tau : G \rightarrow G$. The general formula turns out to be

$$[\text{Op}^\tau(a)u](x) = \int_G \left(\int_{\widehat{G}} \text{Tr}_\xi \left[\xi(y^{-1}x) a(x\tau(y^{-1}x)^{-1}, \xi) \right] d\widehat{m}(\xi) \right) u(y) dm(y), \quad (2)$$

from which (1) can be recovered putting $\tau(x) = e$ (the identity) for every $x \in G$. This formula and its integral version will be summarised in (37). The case $\tau(x) = x$ is also related to a standard choice

$$[\text{Op}^{\text{id}_G}(a)u](x) = \int_G \left(\int_{\widehat{G}} \text{Tr}_\xi \left[\xi(y^{-1}x) a(y, \xi) \right] d\widehat{m}(\xi) \right) u(y) dm(y), \quad (3)$$

familiar at least in the case $G = \mathbb{R}^n$ (derivatives to the left, positions to the right). In the presence of τ some formulae are rather involved, but the reader can take the basic case $\tau(\cdot) = e$ as the main example. Anyhow, for the function spaces we consider in this paper, the formalisms corresponding to different mappings τ are actually isomorphic. Having in mind the Weyl quantization for $G = \mathbb{R}^n$ we deal in Section 4 with the problem of a symmetric quantization, for which one has $\text{Op}^\tau(a)^* = \text{Op}^\tau(a^*)$, where a^* is an operator version of complex conjugation. We also note that if the symbol $a(x, \xi) = a(\xi)$ is independent of x , the operator $\text{Op}^\tau(a)$ is left-invariant and independent of τ , and can be rewritten in the form of the Fourier multiplier

$$\mathcal{F} [\text{Op}^\tau(a)u](\xi) = a(\xi) [\mathcal{F}u](\xi), \quad \xi \in \widehat{G}, \quad (4)$$

at least for sufficiently well-behaved functions u , i.e. as an operator of “multiplication” of the operator-valued Fourier coefficients from the left.

One of our purposes is to sketch two justifications of formula (2), which both hold without a Lie structure on G (we refer to [3, 30] to similar strategies in quite different situations). They also enrich the formalism and have certain applications, some of them included here, others subject of subsequent developments. Let us say some words about the two approaches.

1. A locally compact group G being given, we have a canonical action by (left) translations on various C^* -algebras of functions on G . There are crossed product constructions associated to such situations, presented in Section 7.1: One gets $*$ -algebras of

scalar-valued functions on $G \times G$ involving a product which is a convolution in one variable and a pointwise multiplication in the other variable, suitably twisted by the action by translations. A C^* -norm with an operator flavour is also available, with respect to which one takes a completion. Since we have to accommodate the parameter τ , we were forced to outline an extended version of crossed products.

Among the representations of these C^* -algebras there is a distinguished one presented and used in Subsection 7.2, the Schrödinger representation, in the Hilbert space $L^2(G)$. If G is type I, second countable and unimodular, there is a nicely-behaved Fourier transform sending functions on G into operator-valued sections defined over \widehat{G} . This can be augmented to a partial Fourier transform sending functions on $G \times G$ into sections over $G \times \widehat{G}$. Starting from the crossed products, this partial Fourier transform serves to define, by transport of structure, $*$ -algebras of symbols with a multiplication generalising the Weyl-Moyal calculus as well as Hilbert space representations of the form (2). They are shown to be generated by products of suitable multiplication and convolution operators.

The C^* -background can be used, in a slightly more general context, to generate covariant families of pseudo-differential operators, cf. Subsection 7.4. It also leads to results about the spectrum of certain bounded or unbounded pseudo-differential operators, as it is presented in Subsection 7.4 and will be continued in a subsequent paper.

2. A second approach relies on Weyl systems. If $G = \mathbb{R}^n$ one can write

$$\text{Op}(a) = \int_{\mathbb{R}^{2n}} \widehat{a}(\xi, x) W(\xi, x) dx d\xi,$$

where the Weyl system (phase-space shifts)

$$\{W(\xi, x) := V(\xi)U(x) \mid (x, \xi) \in \mathbb{R}^{2n}\}$$

is a family of unitary operators in $L^2(\mathbb{R}^n)$ obtained by putting together translations and modulations. This is inspired by the Fourier inversion formula, but notice that W is only a projective representation; this is a precise way to codify the canonical commutation relations between positions Q (generating V) and momenta P (generating U) and Op can be seen as a non-commutative functional calculus $a \mapsto a(Q, P) \equiv \text{Op}(a)$. Besides its phase-space quantum mechanical interest, this point of view also opens the way to some new topics or tools such as the Bargmann transform, coherent states, the anti-Wick quantization, coorbit spaces, etc.

In Section 3 we show that such a “Weyl system approach” and its consequences are also available in the context of second countable, unimodular type I groups; in particular it leads to (2). The Weyl system in this general case, adapted in Definitions 3.1 and 3.3 to the existence of the quantization parameter τ , has nice technical properties (including a fibered form of square integrability) that are proven in Subsection 3.1. This has useful consequences at the level of the quantization process, as shown in Subsection 3.2. In particular, it is shown that Op^τ is a unitary map from a suitable class of square integrable sections over $G \times \widehat{G}$ to the Hilbert space of all Hilbert-Schmidt operators on $L^2(G)$. The intrinsic $*$ -algebraic structure on the level of symbols is briefly

treated in Subsection 3.3. In Subsection 3.4 we rely on complex interpolation and non-commutative L^p -spaces to put into evidence certain families of Schatten-class operators.

Without assuming that G is a Lie group we do not have the usual space of smooth compactly supported functions readily available as the standard space of test functions. So, in Section 5, we will be using its generalisation to the setting of locally compact groups by Bruhat [4], and these Bruhat spaces $\mathcal{D}(G)$ and $\mathcal{D}'(G)$ will replace the usual spaces of test functions and distributions in our setting. An important fact is that they are nuclear. Taking suitable tensor products one also gets a space $\mathcal{D}(G \times \widehat{G})$ of regularising symbols and (by duality) a space $\mathcal{D}'(G \times \widehat{G})$ of “distributions”, allowing to define unbounded pseudo-differential operators.

In Subsection 5.3 we show that pseudo-differential operators with regularising operator-valued symbols can be used to describe compactness of families of vectors or operators in $L^2(G)$.

Besides the usual ordering issue (derivatives to the left or to the right), already appearing for \mathbb{R}^n and connected to the Heisenberg commutation relations and the symplectic structure of phase space, for general groups there is a second ordering problem coming from the intrinsic non-commutativity of G . The Weyl system used in Section 3 relies on translations to the right, aiming at a good correspondence with the previously studied compact and nilpotent cases. Another Weyl system, involving left translations, is introduced in Section 6 and used in defining a left quantization. It turns out that this one is directly linked to crossed product C^* -algebras.

We dedicated the last section to a brief overview of quantization on (connected, simply connected) nilpotent Lie groups. Certain subclasses have been thoroughly examined in references cited above, so we are going to concentrate on some new features. Besides the extra generality of the present setting (non-graded nilpotent groups, τ -quantizations, C^* -algebras), we are also interested in the presence of a second formalism, involving scalar-valued symbols. We show that it is equivalent to the one involving operator-valued symbols, emerging as a particular case of the previous sections. This is a rather direct consequence of the excellent behaviour of the exponential function in the nilpotent case. On one hand, the analysis in this paper here outlines a τ -extension of the scalar-valued calculus on nilpotent Lie groups initiated by Melin [33], see also [24, 25] for further developments on homogeneous and general nilpotent Lie groups. On the other hand, it relates this to the operator-valued calculus developed in [18, 19].

After some basic constructions involving various types of Fourier transformations are outlined, the detailed development of the pseudo-differential operators with scalar-valued symbol follows along the lines already indicated. So, to save space and avoid repetitions, we will be rather formal and sketchy and leave many details to the reader. Actually the Lie structure of a nilpotent group permits a deeper investigation that was treated in [19] and should be still subject of future research.

Thus, to summarise, the main results of this paper are as follows:

- We develop a rigorous framework for the analysis of pseudo-differential operators on locally compact groups of type I, which we assume also unimodular for technical simplicity.
- We introduce notions of Wigner and Fourier-Wigner transforms, and of Weyl systems, adapted to this general setting. These notions are used to define and analyse τ -quantizations (or quantization by Weyl systems) of operators modelled on families of quantizations on \mathbb{R}^n that include the Kohn-Nirenberg and Weyl quantizations.
- We develop the C^* -algebraic formalism to put τ -quantizations in a more general perspective, also allowing analysis of operators with ‘coefficients’ taking values in different C^* -algebras. The link with a left form of τ -quantization is given via a special covariant representation, the Schrödinger representation. This is further applied to investigate spectral properties of covariant families of operators.
- Although the initial analysis is set for operators bounded on $L^2(G)$, this can be extended further to include densely defined operators and, more generally, operators from $\mathcal{D}(G)$ to $\mathcal{D}'(G)$. Since G does not have to be a Lie group (i.e. there may be no compatible smooth differential structure on G) we show how this can be done using the so-called Bruhat space $\mathcal{D}(G)$, an analogue of the space of smooth compactly supported functions in the setting of general locally compact groups.
- The results are applied to a deeper analysis of τ -quantizations on nilpotent Lie groups. On one hand, this extends the setting of graded Lie groups developed in depth in [18, 19] to more general nilpotent Lie groups, also introducing a possibility for Weyl-type quantizations there. On the other hand, it extends the invariant Melin calculus [33, 25] on homogeneous groups to general non-invariant operators with the corresponding τ -versions of scalar-symbols on the dual of the Lie algebra;
- We give a criterion for the existence of Weyl-type quantizations in our framework, namely, to quantizations in which real-valued symbols correspond to self-adjoint operators. We show the existence of such quantizations in several settings, most interestingly in the setting of general groups of exponential type.

In this paper we are mostly interested in symbolic understanding of pseudo-differential operators. Approaches through kernels exist as well, see e.g. Meladze and Shubin [32] and further works by these authors on operators on unimodular Lie groups, or Christ, Geller, Głowacki and Polin [6] on homogeneous groups – but see also an alternative (and earlier) symbolic approach to that on the Heisenberg group by Taylor [44].

As we have explained, there exist several approaches to global quantizations of operators on groups, such as those worked out in detail in [39] and [18] in the settings of compact and nilpotent groups, respectively, as well as the approach by Melin [33, 25]

for nilpotent groups. As both points of view are effective in different applications, one motivation for this paper is to describe a link between them explaining how one could go from one description to the other. As a byproduct of such a link we managed to extend the Melin's formalism to non-invariant operators. Observing the similarities between the compact and nilpotent cases in [39] and [18], respectively, and based on Taylor's observation [44], another motivation for this paper is to put both approaches in a single framework, that of locally compact groups of type I. While losing the results depending on the differential structure of the group as a manifold, this framework is still effective in handling a scope of spectral questions. Some applications to this end are given in Section 4. Moreover, it greatly extends the variety of settings where such pseudo-differential analysis becomes available. Furthermore, as the Weyl quantization is particularly effective for certain problems, an additional motivation for our analysis comes from the desire to understand the nature of the Weyl quantization in the settings when it is not even clear how to define the midpoint $\frac{x+y}{2}$ for two points x, y in the group. This problem becomes apparent already on the torus when such a 'midpoint' mapping is not continuous. Thus, in Section 4 we show that such Weyl-type quantizations are still available in a large class of groups, including the class of the exponential groups. The C^* -algebraic approach of Section 7 has been used in [27] to prove Fredholm properties and to evaluate essential spectra of global pseudo-differential operators. Other applications of the obtained constructions will appear elsewhere. In particular, see [31] for applications in the case of nilpotent Lie groups having flat coadjoint orbits, where further connections with Pedersen's quantization [37] are established.

2 FRAMEWORK

In this section we set up a general framework of this paper, also recalling very briefly necessary elements of the theory of type I groups and their Fourier analysis.

2.1 GENERAL

For a given (complex, separable) Hilbert space \mathcal{H} , the scalar product $\langle \cdot, \cdot \rangle_{\mathcal{H}}$ will be linear in the first variable and anti-linear in the second. One denotes by $\mathbb{B}(\mathcal{H})$ the C^* -algebra of all linear bounded operators in \mathcal{H} and by $\mathbb{K}(\mathcal{H})$ the closed bi-sided $*$ -ideal of all the compact operators. The Hilbert-Schmidt operators form a two-sided $*$ -ideal $\mathbb{B}^2(\mathcal{H})$ (dense in $\mathbb{K}(\mathcal{H})$) which is also a Hilbert space with the scalar product $\langle A, B \rangle_{\mathbb{B}^2(\mathcal{H})} := \text{Tr}(AB^*)$. This Hilbert space is unitarily equivalent to the Hilbert tensor product $\mathcal{H} \otimes \overline{\mathcal{H}}$, where $\overline{\mathcal{H}}$ is the Hilbert space opposite to \mathcal{H} . The unitary operators form a group $\mathbb{U}(\mathcal{H})$. The commutant of a subset \mathcal{N} of $\mathbb{B}(\mathcal{H})$ is denoted by \mathcal{N}' .

Let G be a locally compact group with unit e and fixed left Haar measure m . Our group will soon be supposed unimodular, so m will also be a right Haar measure. By $\mathcal{C}_c(G)$ we denote the space of all complex continuous compactly supported functions on G . For $p \in [1, \infty]$, the Lebesgue spaces $L^p(G) \equiv L^p(G; m)$ will always refer to

the Haar measure. We denote by $C^*(G)$ the full (universal) C^* -algebra of G and by $C_{\text{red}}^*(G) \subset \mathbb{B}[L^2(G)]$ the reduced C^* -algebra of G . Recall that any representation π of G generates canonically a non-degenerate representation Π of the C^* -algebra $C^*(G)$. The notation $A(G)$ is reserved for Eymard's Fourier algebra of the group G .

The canonical objects in representation theory [14, 23] will be denoted by $\text{Rep}(G)$, $\text{Irrep}(G)$ and \widehat{G} . An element of $\text{Rep}(G)$ is a Hilbert space representation $\pi : G \rightarrow \mathbb{U}(\mathcal{H}_\pi) \subset \mathbb{B}(\mathcal{H}_\pi)$, always supposed to be strongly continuous. If it is irreducible, it belongs to $\text{Irrep}(G)$ by definition. Unitary equivalence of representations will be denoted by \cong . We set $\widehat{G} := \text{Irrep}(G)/\cong$ and call it *the unitary dual of G* . If G is Abelian, the unitary dual \widehat{G} is the Pontryagin dual group; if not, \widehat{G} has no group structure. A *primary (factor) representation* π satisfies, by definition, $\pi(G)' \cap \pi(G)'' = \mathbb{C} \text{id}_{\mathcal{H}_\pi}$.

DEFINITION 2.1. *The locally compact group G will be called admissible if it is second countable, type I and unimodular.*

Admissibility will be a standing assumption and it is needed for most of the main constructions and results. There are hopes to extend at least parts of this paper to non-unimodular groups, by using techniques of [10].

REMARK 2.2. We assume that the reader is familiar with the concept of *type I group*. Let us only say that for such a group every primary representation is a direct sum of copies of some irreducible representation; for the full theory we refer to [14, 23, 21]. In [23, Th. 7.6] (see also [14]), many equivalent characterisations are given for a *second countable* locally compact group to be type I. In particular, in such a case, the notion is equivalent to postliminality (GCR). Thus G is type I if and only if for every irreducible representation π one has $\mathbb{K}(\mathcal{H}_\pi) \subset \Pi[C^*(G)]$.

The single way we are going to use the fact that G is type I is through one main consequence of this property, to be outlined below: the existence of a measure on the unitary dual \widehat{G} for which a Plancherel Theorem holds.

EXAMPLE 2.3. Compact and Abelian groups are type I. So are the Euclidean and the Poincaré groups. Among the connected groups, real algebraic, exponential (in particular nilpotent) and semi-simple Lie groups are type I. Not all the solvable groups are type I; see [23, Th. 7.10] for a criterion. A discrete group is type I [45] if and only if it is the finite extension of an Abelian normal subgroup. So the non-trivial free groups or the discrete Heisenberg group are not type I.

REMARK 2.4. We recall that, being second countable, G will be separable, σ -compact and completely metrizable; in particular, as a Borel space it will be standard. The Haar measure m is σ -finite and $L^p(G)$ is a separable Banach space if $p \in [1, \infty)$. In addition, all the cyclic representations have separable Hilbert spaces; this applies, in particular, to irreducible representations.

A second countable discrete group is at most countable.

We mention briefly some harmonic analysis concepts; full treatment is given in [14, 23]. The precise definitions and properties will either be outlined further on, when needed, or they will not be explicitly necessary.

Both $\text{Irrep}(G)$ and the unitary dual $\widehat{G} := \text{Irrep}(G)/\cong$ are endowed with (standard) Borel structures [14, 18.5]. The structure on \widehat{G} is the quotient of that on $\text{Irrep}(G)$ and is called the Mackey Borel structure. There is a measure on \widehat{G} , called the Plancherel measure associated to m and denoted by \widehat{m} [14, 18.8]. Its basic properties, connected to the Fourier transform, will be briefly discussed below.

The unitary dual \widehat{G} is also a separable locally quasi-compact Baire topological space having a dense open locally compact subset [14, 18.1]. Very often this topological space is not Hausdorff (this is the difference between "locally quasi-compact" and "locally compact").

REMARK 2.5. We are going to use a systematic abuse of notation that we now explain. There is a \widehat{m} -measurable field $\{\mathcal{H}_\xi \mid \xi \in \widehat{G}\}$ of Hilbert spaces and a measurable section $\widehat{G} \ni \xi \mapsto \pi_\xi \in \text{Irrep}(G)$ such that each $\pi_\xi : G \rightarrow \mathbb{B}(\mathcal{H}_\xi)$ is a irreducible representation belonging to the class ξ . In various formulae, instead of π_ξ we will write ξ , making a convenient identification between irreducible representations and classes of irreducible representations. The measurable field of irreducible representations $\{(\pi_\xi, \mathcal{H}_\xi) \mid \xi \in \widehat{G}\}$ is fixed and other choices would lead to equivalent constructions and statements.

One introduces the direct integral Hilbert space

$$\mathcal{B}^2(\widehat{G}) := \int_{\widehat{G}}^{\oplus} \mathbb{B}^2(\mathcal{H}_\xi) d\widehat{m}(\xi) \cong \int_{\widehat{G}}^{\oplus} \mathcal{H}_\xi \otimes \overline{\mathcal{H}_\xi} d\widehat{m}(\xi), \tag{5}$$

with the obvious scalar product

$$\langle \phi_1, \phi_2 \rangle_{\mathcal{B}^2(\widehat{G})} := \int_{\widehat{G}} \langle \phi_1(\xi), \phi_2(\xi) \rangle_{\mathbb{B}^2(\mathcal{H}_\xi)} d\widehat{m}(\xi) = \int_{\widehat{G}} \text{Tr}_\xi[\phi_1(\xi)\phi_2(\xi)^*] d\widehat{m}(\xi), \tag{6}$$

where Tr_ξ refers to the trace in $\mathbb{B}(\mathcal{H}_\xi)$. More generally, for $p \in [1, \infty)$ one defines $\mathcal{B}^p(\widehat{G})$ as the family of measurable fields $\phi \equiv (\phi(\xi))_{\xi \in \widehat{G}}$ for which $\phi(\xi)$ belongs to the Schatten-von Neumann class $\mathbb{B}^p(\mathcal{H}_\xi)$ for almost every ξ and

$$\|\phi\|_{\mathcal{B}^p(\widehat{G})} := \left(\int_{\widehat{G}} \|\phi(\xi)\|_{\mathbb{B}^p(\mathcal{H}_\xi)}^p d\widehat{m}(\xi) \right)^{1/p} < \infty. \tag{7}$$

They are Banach spaces. We also recall that the von Neumann algebra of decomposable operators $\mathcal{B}(\widehat{G}) := \int_{\widehat{G}}^{\oplus} \mathbb{B}(\mathcal{H}_\xi) d\widehat{m}(\xi)$ acts to the left and to the right in the Hilbert space $\mathcal{B}^2(\widehat{G})$ in an obvious way.

On $\Gamma := G \times \widehat{G}$, which might *not* be a locally compact space or a group, we consider the product measure $m \otimes \widehat{m}$. It is independent of our choice for m (if m is replaced by λm for some strictly positive number λ , the corresponding Plancherel measure will be $\lambda^{-1}\widehat{m}$). Very often we are going to need $\widehat{\Gamma} := \widehat{G} \times G$ (this notation should *not* suggest a duality) with the measure $\widehat{m} \otimes m$. We could identify it with Γ (by means of the map $(\xi, x) \mapsto (x, \xi)$) but in most cases it is better not to do this identification.

Associated to these two measure spaces, we also need the Hilbert spaces

$$\mathcal{B}^2(\Gamma) \equiv \mathcal{B}^2(\mathbb{G} \times \widehat{\mathbb{G}}) := L^2(\mathbb{G}) \otimes \mathcal{B}^2(\widehat{\mathbb{G}}) \quad (8)$$

and

$$\mathcal{B}^2(\widehat{\Gamma}) \equiv \mathcal{B}^2(\widehat{\mathbb{G}} \times \mathbb{G}) := \mathcal{B}^2(\widehat{\mathbb{G}}) \otimes L^2(\mathbb{G}), \quad (9)$$

also having direct integral decompositions.

2.2 THE FOURIER TRANSFORM

The Fourier transform [14, 18.2] of $u \in L^1(\mathbb{G})$ is given in weak sense by

$$(\mathcal{F}u)(\xi) \equiv \widehat{u}(\xi) := \int_{\mathbb{G}} u(x)\xi(x)^* dm(x) \in \mathbb{B}(\mathcal{H}_\xi). \quad (10)$$

Here and subsequently the interpretation of $\xi \in \widehat{\mathbb{G}}$ as a true irreducible representation is along the lines of Remark 2.5. Actually, by the compressed form (10) we mean that for $\varphi_\xi, \psi_\xi \in \mathcal{H}_\xi$ one has

$$\langle (\mathcal{F}u)(\xi)\varphi_\xi, \psi_\xi \rangle_{\mathcal{H}_\xi} := \int_{\mathbb{G}} u(x)\langle \varphi_\xi, \pi_\xi(x)\psi_\xi \rangle_{\mathcal{H}_\xi} dm(x).$$

Some useful facts [14, 18.2 and 3.3]:

- The Fourier transform $\mathcal{F} : L^1(\mathbb{G}) \rightarrow \mathcal{B}^2(\widehat{\mathbb{G}})$ is linear, contractive and injective.
- For every $\epsilon > 0$ there exists a quasi-compact subset $K_\epsilon \subset \widehat{\mathbb{G}}$ such that $\|(\mathcal{F}u)(\xi)\|_{\mathbb{B}(\mathcal{H}_\xi)} \leq \epsilon$ if $\xi \notin K_\epsilon$.
- The map $\widehat{\mathbb{G}} \ni \xi \mapsto \|(\mathcal{F}u)(\xi)\|_{\mathbb{B}(\mathcal{H}_\xi)} \in \mathbb{R}$ is lower semi-continuous. It is even continuous, whenever $\widehat{\mathbb{G}}$ is Hausdorff.

Recall [14, 22, 21] that the Fourier transform \mathcal{F} extends (starting from $L^1(\mathbb{G}) \cap L^2(\mathbb{G})$) to a unitary isomorphism $\mathcal{F} : L^2(\mathbb{G}) \rightarrow \mathcal{B}^2(\widehat{\mathbb{G}})$. This is the generalisation of Plancherel's Theorem to (maybe non-commutative) admissible groups and it will play a central role in our work.

REMARK 2.6. It is also known [26, 21] that \mathcal{F} restricts to a bijection

$$\mathcal{F}_{(0)} : L^2(\mathbb{G}) \cap A(\mathbb{G}) \rightarrow \mathcal{B}^2(\widehat{\mathbb{G}}) \cap \mathcal{B}^1(\widehat{\mathbb{G}}) \quad (11)$$

with inverse given by (the traces refer to \mathcal{H}_ξ)

$$\left(\mathcal{F}_{(0)}^{-1}\phi\right)(x) = \int_{\widehat{\mathbb{G}}} \text{Tr}_\xi[\xi(x)\phi(\xi)] d\widehat{m}(\xi). \quad (12)$$

Rephrasing this, the restriction of the inverse \mathcal{F}^{-1} to the subspace $\mathcal{B}^2(\widehat{G}) \cap \mathcal{B}^1(\widehat{G})$ has the explicit form (12), and this will be a useful fact. Note the consequence, valid for $u \in L^2(G) \cap A(G)$ and for m -almost every $x \in G$:

$$u(x) = \int_{\widehat{G}} \text{Tr}_{\xi}[(\mathcal{F}u)(\xi)\xi(x)]d\widehat{m}(\xi) = \int_{\widehat{G}} \text{Tr}_{\xi}[\xi(x)\widehat{u}(\xi)]d\widehat{m}(\xi). \tag{13}$$

In particular, this holds for $u \in C_c(G)$. The extension $\mathcal{F}_{(1)}$ of $\mathcal{F}_{(0)}$ to $A(G)$ makes sense as an isometry $\mathcal{F}_{(1)} : A(G) \rightarrow \mathcal{B}^1(\widehat{G})$.

Combining the quantization formula (1) with the Fourier transform (10), we can write (1) also as

$$[\text{Op}(a)u](x) = \int_{\widehat{G}} \text{Tr}_{\xi}[\xi(x)a(x, \xi)\widehat{u}(\xi)]d\widehat{m}(\xi), \tag{14}$$

which can be viewed as an extension of the Fourier inversion formula (13).

REMARK 2.7. By a formula analogous to (10), the Fourier transform is even defined (and injective) on bounded complex Radon measures μ on G . One gets easily

$$\sup_{\xi \in \widehat{G}} \|\mathcal{F}\mu\|_{\mathbb{B}(\mathcal{H}_{\xi})} \leq \|\mu\|_{M^1(G)} := |\mu|(G).$$

REMARK 2.8. There are many (full or partial) Fourier transformations that can play important roles, as

$$\mathcal{F} \otimes \text{id} : L^2(G \times G) \rightarrow \mathcal{B}^2(\widehat{\Gamma}), \quad \text{id} \otimes \mathcal{F} : L^2(G \times G) \rightarrow \mathcal{B}^2(\Gamma). \tag{15}$$

$$\mathcal{F} \otimes \mathcal{F}^{-1} : \mathcal{B}^2(\Gamma) \rightarrow \mathcal{B}^2(\widehat{\Gamma}), \quad \mathcal{F}^{-1} \otimes \mathcal{F} : \mathcal{B}^2(\widehat{\Gamma}) \rightarrow \mathcal{B}^2(\Gamma). \tag{16}$$

They might admit various extensions or restrictions.

3 QUANTIZATION BY A WEYL SYSTEM

In this section we introduce a notion of a Weyl system in our setting and outline its relation to Wigner and Fourier-Wigner transforms. This is then used to define pseudo-differential operators through τ -quantization for an arbitrary measurable function $\tau : G \rightarrow G$. The introduced formalism is then applied to study (involutive) algebra properties of symbols and operators as well as Schatten class properties in the setting of non-commutative L^p -spaces. One of the goals here is to give rigorous understanding to the τ -quantization formula (2).

3.1 WEYL SYSTEMS AND THEIR ASSOCIATED TRANSFORMATIONS

Let us fix a measurable function $\tau : G \rightarrow G$. We will often use the notation $\tau x \equiv \tau(x)$ to avoid writing too many brackets.

DEFINITION 3.1. For $x \in G$ and $\pi \in \text{Rep}(G)$ one defines a unitary operator $W^{\tau}(\pi, x)$ in the Hilbert space $L^2(G; \mathcal{H}_{\pi}) \equiv L^2(G) \otimes \mathcal{H}_{\pi}$ by

$$[W^{\tau}(\pi, x)\Theta](y) := \pi[y(\tau x)^{-1}]^* [\Theta(yx^{-1})] = \pi[\tau(x)]\pi(y)^*[\Theta(yx^{-1})]. \tag{17}$$

If $\pi \cong \rho$, i.e. if $\rho(x)U = U\pi(x)$ for some unitary operator $U : \mathcal{H}_\pi \rightarrow \mathcal{H}_\rho$ and for every $x \in G$, then it follows easily that

$$W^\tau(\rho, x) = (\text{id} \otimes U)W^\tau(\pi, x)(\text{id} \otimes U)^{-1}.$$

We record for further use the formula

$$\begin{aligned} W^{\tau'}(\pi, x) &= [\text{id} \otimes \pi(\tau'x)] [\text{id} \otimes \pi(\tau x)^*] W^\tau(\pi, x) \\ &= [\text{id} \otimes \pi((\tau'x)(\tau x)^{-1})] W^\tau(\pi, x) \end{aligned} \tag{18}$$

making the connection between operators defined by different parametres τ, τ' as well as the explicit form of the adjoint

$$[W^\tau(\pi, x)^* \Theta](y) = \pi[yx(\tau x)^{-1}] [\Theta(yx)].$$

One also notes that $W^\tau(\mathbf{1}, x) = R(x^{-1})$, where R is the right regular representation of G and $\mathbf{1}$ is the 1-dimensional trivial representation. In this case $\mathcal{H}_\mathbf{1} = \mathbb{C}$, so $L^2(G; \mathcal{H}_\mathbf{1})$ reduces to $L^2(G)$.

REMARK 3.2. One can not compose the operators $W^\tau(\pi, x)$ and $W^\tau(\rho, y)$ in general, since they act in different Hilbert spaces. Note, however, that the family $\text{Rep}(G)/\cong$ of all the unitary equivalence classes of representations form an Abelian monoid with the tensor composition

$$(\pi \otimes \rho)(x) := \pi(x) \otimes \rho(x), \quad x \in G,$$

and the unit $\mathbf{1}$ (after a suitable reinterpretation in terms of equivalence classes). The subset $\widehat{G} = \text{Irrep}(G)/\cong$ is not a submonoid in general, but the generated submonoid, involving finite tensor products of irreducible representations, could be interesting. It is instructive to compute the operator in $L^2(G; \mathcal{H}_\pi \otimes \mathcal{H}_\rho)$

$$[W(\pi, x) \otimes \text{id}_\rho] [W(\rho, y) \otimes \text{id}_\pi] = [\text{id}_{L^2(G)} \otimes \rho(x) \otimes \text{id}_\pi] W(\pi \otimes \rho, yx); \tag{19}$$

to get this result one has to identify $\mathcal{H}_\pi \otimes \mathcal{H}_\rho$ with $\mathcal{H}_\rho \otimes \mathcal{H}_\pi$. If G is Abelian, the unitary dual \widehat{G} is the Pontryagin dual group, the irreducible representations are 1-dimensional and for $\xi \equiv \pi \in \widehat{G}$ and $\eta \equiv \rho \in \widehat{G}$ the identity (19) reads

$$W(\xi, x)W(\eta, y) = \eta(x)W(\xi\eta, xy).$$

Thus $W : \widehat{G} \times G \rightarrow \mathbb{B}[L^2(G)]$ is a unitary projective representation with 2-cocycle (multiplier)

$$\varpi : (\widehat{G} \times G) \times (\widehat{G} \times G) \rightarrow \mathbb{T}, \quad \varpi((\xi, x), (\eta, y)) := \eta(x).$$

Similar computations can be done for W^τ with general τ .

From now on we mostly concentrate on the family of operators $W^\tau(\xi, x)$ where $x \in G$ and ξ is an irreducible representation. Extrapolating from the case $G = \mathbb{R}^n$, we call this family a *Weyl system*.

Below, for an operator T in $L^2(G; \mathcal{H}_\xi) \cong L^2(G) \otimes \mathcal{H}_\xi$ and a pair of vectors $u, v \in L^2(G)$, the action of $\langle Tu, v \rangle_{L^2(G)} \in \mathbb{B}(\mathcal{H}_\xi)$ on $\varphi_\xi \in \mathcal{H}_\xi$ is given by

$$\langle Tu, v \rangle_{L^2(G)} \varphi_\xi := \int_G [T(u \otimes \varphi_\xi)](y) \overline{v(y)} \, dm(y) \in \mathcal{H}_\xi. \tag{20}$$

DEFINITION 3.3. For $(x, \xi) \in G \times \widehat{G}$ and $u, v \in L^2(G)$ one sets

$$\mathcal{W}_{u,v}^\tau(\xi, x) := \langle W^\tau(\xi, x) \overline{u}, \overline{v} \rangle_{L^2(G)} \in \mathbb{B}(\mathcal{H}_\xi). \tag{21}$$

This definition is suggested by the standard notion of *representation coefficient function* from the theory of unitary group representations. However, in general, $\widehat{G} \times G$ is not a group, $\mathcal{W}_{u,v}^\tau$ is not scalar-valued, and $W^\tau(\xi, x)W^\tau(\eta, y)$ makes no sense.

REMARK 3.4. Note the identity

$$\langle \mathcal{W}_{u,v}^\tau(\xi, x) \varphi_\xi, \psi_\xi \rangle_{\mathcal{H}_\xi} = \langle W^\tau(\xi, x) (\overline{u} \otimes \varphi_\xi), \overline{v} \otimes \psi_\xi \rangle_{L^2(G; \mathcal{H}_\xi)}, \tag{22}$$

valid for $u, v \in L^2(G)$, $\varphi_\xi, \psi_\xi \in \mathcal{H}_\xi$, $(\xi, x) \in \widehat{\Gamma}$. It follows immediately from (21) and (20). In fact (22) can serve as a definition of $\mathcal{W}_{u,v}^\tau(\xi, x)$.

PROPOSITION 3.5. The mapping $(u, v) \mapsto \mathcal{W}_{u,v}^\tau$ defines a unitary map (denoted by the same symbol) $\mathcal{W}^\tau : \overline{L^2(G)} \otimes L^2(G) \rightarrow \mathcal{B}^2(\widehat{\Gamma})$, called the Fourier-Wigner τ -transformation.

Proof. Let us define the change of variables

$$cv^\tau : G \times G \rightarrow G \times G, \quad cv^\tau(x, y) := (x\tau(y^{-1}x)^{-1}, y^{-1}x) \tag{23}$$

with inverse

$$(cv^\tau)^{-1}(x, y) = (x\tau(y), x\tau(y)y^{-1}). \tag{24}$$

Using the definition and the interpretation (20), one has for $\varphi_\xi \in \mathcal{H}_\xi$

$$\begin{aligned} \mathcal{W}_{u,v}^\tau(\xi, x) \varphi_\xi &= \int_G [W^\tau(\xi, x) (\overline{u} \otimes \varphi_\xi)](z) v(z) \, dm(z) \\ &= \int_G v(z) \overline{u(zx^{-1})} \xi(z\tau(x)^{-1})^* \varphi_\xi \, dm(z) \\ &= \int_G v(y\tau(x)) \overline{u(y\tau(x)x^{-1})} \xi(y)^* \varphi_\xi \, dm(y) \\ &= \int_G (v \otimes \overline{u}) [(cv^\tau)^{-1}(y, x)] \xi(y)^* \varphi_\xi \, dm(y). \end{aligned}$$

By using the properties of the Haar measure and the unimodularity of G , it is easy to see that the composition with the map cv^τ , denoted by CV^τ , is a unitary operator in

$L^2(\mathbb{G} \times \mathbb{G}) \cong L^2(\mathbb{G}) \otimes L^2(\mathbb{G})$. On the other hand, the conjugation $\overline{L^2(\mathbb{G})} \ni w \mapsto \bar{w} \in L^2(\mathbb{G})$ is also unitary. Making use of the unitary partial Fourier transformation

$$(\mathcal{F} \otimes \text{id}) : L^2(\mathbb{G}) \otimes L^2(\mathbb{G}) \rightarrow \mathcal{B}^2(\widehat{\mathbb{G}}) \otimes L^2(\mathbb{G}),$$

one gets

$$\mathcal{W}_{u,v}^\tau = (\mathcal{F} \otimes \text{id})(\text{CV}^\tau)^{-1}(v \otimes \bar{u}) \tag{25}$$

and the statement follows. □

The unitarity of the Fourier-Wigner transformation implies the next irreducibility result:

COROLLARY 3.6. *Let \mathcal{K} be a closed subspace of $L^2(\mathbb{G})$ such that $W^\tau(\xi, x)(\mathcal{K} \otimes \mathcal{H}_\xi) \subset \mathcal{K} \otimes \mathcal{H}_\xi$ for every $(\xi, x) \in \widehat{\Gamma}$. Then $\mathcal{K} = \{0\}$ or $\mathcal{K} = L^2(\mathbb{G})$.*

Proof. Suppose that $\mathcal{K} \neq L^2(\mathbb{G})$ and let $\bar{v} \in \mathcal{K}^\perp \setminus \{0\}$.

Let us examine the identity (22), where $\bar{u} \in \mathcal{K}$, $(\xi, x) \in \widehat{\Gamma}$ and $\varphi_\xi, \psi_\xi \in \mathcal{H}_\xi$. Since $W^\tau(\xi, x)(\bar{u} \otimes \varphi_\xi) \in \mathcal{K} \otimes \mathcal{H}_\xi$, the right hand side is zero. So the left hand side is also zero for φ_ξ, ψ_ξ arbitrary, so $\mathcal{W}_{u,v}^\tau(\xi, x) = 0$. Then, by unitarity

$$\|u\|_{L^2(\mathbb{G})}^2 \|v\|_{L^2(\mathbb{G})}^2 = \|\mathcal{W}_{u,v}^\tau\|_{\mathcal{B}^2(\widehat{\Gamma})}^2 = \int_{\mathbb{G}} \int_{\widehat{\mathbb{G}}} \|\mathcal{W}_{u,v}^\tau(\xi, x)\|_{\mathbb{B}^2(\mathcal{H}_\xi)}^2 d\mathfrak{m}(x) d\widehat{\mathfrak{m}}(\xi) = 0$$

and since $v \neq 0$ one must have $u = 0$. □

Depending on the point of view, one uses one of the notations $\mathcal{W}_{u,v}^\tau$ or $W^\tau(u \otimes v)$. We also introduce

$$\begin{aligned} \mathcal{V}_{u,v}^\tau &\equiv \mathcal{V}^\tau(u \otimes v) := (\mathcal{F}^{-1} \otimes \mathcal{F})\mathcal{W}_{v,u}^\tau = \\ &(\text{id} \otimes \mathcal{F})(\text{CV}^\tau)^{-1}(v \otimes \bar{u}) \in L^2(\mathbb{G}) \otimes \mathcal{B}^2(\widehat{\mathbb{G}}), \end{aligned} \tag{26}$$

which reads explicitly

$$\mathcal{V}_{u,v}^\tau(x, \xi) = \int_{\mathbb{G}} \overline{u(x\tau(y)y^{-1})} v(x\tau(y)) \xi(y)^* d\mathfrak{m}(y).$$

One can name the unitary mapping $\mathcal{V}^\tau : \overline{L^2(\mathbb{G})} \otimes L^2(\mathbb{G}) \rightarrow \mathcal{B}^2(\Gamma)$ the Wigner τ -transformation. We record for further use the orthogonality relations, valid for $u, u', v, v' \in L^2(\mathbb{G})$:

$$\langle \mathcal{W}_{u,v}^\tau, \mathcal{W}_{u',v'}^\tau \rangle_{\mathcal{B}^2(\widehat{\Gamma})} = \langle u', u \rangle_{L^2(\mathbb{G})} \langle v, v' \rangle_{L^2(\mathbb{G})} = \langle \mathcal{V}_{u,v}^\tau, \mathcal{V}_{u',v'}^\tau \rangle_{\mathcal{B}^2(\Gamma)}. \tag{27}$$

3.2 PSEUDO-DIFFERENTIAL OPERATORS

Let, as before, $\tau : G \rightarrow G$ be a measurable map. The next definition should be seen as a rigorous way to give sense to the τ -quantization $\text{Op}^\tau(a)$ introduced in (2).

We note that in general, due to various non-commutativities (of the group, of the symbols), there are essentially two ways of introducing the quantization of this type - these will be given and discussed in the sequel in Section 6, see especially formulae (71) and (72). In the context of compact Lie groups these issues have been extensively discussed in [39], see e.g. Remark 10.4.13 there, and most of that discussion extends to our present setting. One advantage of the order of operators in the definition (2) is that the invariant operators can be viewed as Fourier multipliers with multiplication by the symbol from the left (4), which is perhaps a more familiar way of viewing such operators in non-commutative harmonic analysis. However, it will turn out that the other ordering has certain advantages from the point of view of C^* -algebra theories. We postpone these topics to subsequent sections.

DEFINITION 3.7. For $a \in \mathcal{B}^2(\Gamma)$ (with Fourier transform $\widehat{a} := (\mathcal{F} \otimes \mathcal{F}^{-1})a \in \mathcal{B}^2(\widehat{\Gamma})$) we define $\text{Op}^\tau(a)$ to be the unique bounded linear operator in $L^2(G)$ associated by the relation

$$\text{op}_a^\tau(u, v) = \langle \text{Op}^\tau(a)u, v \rangle_{L^2(G)} \tag{28}$$

to the bounded sesquilinear form $\text{op}_a^\tau : L^2(G) \times L^2(G) \rightarrow \mathbb{C}$

$$\text{op}_a^\tau(u, v) := \langle \widehat{a}, \mathcal{W}_{u,v}^\tau \rangle_{\mathcal{B}^2(\widehat{\Gamma})} = \int_G \int_{\widehat{G}} \text{Tr}_\xi [\widehat{a}(\xi, x) \mathcal{W}_{u,v}^\tau(\xi, x)^*] dm(x) d\widehat{m}(\xi) \tag{29}$$

or, equivalently,

$$\text{op}_a^\tau(u, v) := \langle a, \mathcal{V}_{u,v}^\tau \rangle_{\mathcal{B}^2(\Gamma)} = \int_G \int_G \text{Tr}_\xi [a(x, \xi) \mathcal{V}_{u,v}^\tau(x, \xi)^*] dm(x) d\widehat{m}(\xi). \tag{30}$$

One says that $\text{Op}^\tau(a)$ is the τ -pseudo-differential operator corresponding to the operator-valued symbol a while the map $a \rightarrow \text{Op}^\tau(a)$ will be called the τ -pseudo-differential calculus or τ -quantization.

To justify Definition 3.7, one must show that op_a^τ is indeed a well-defined bounded sesquilinear form. Clearly $\text{op}_a^\tau(u, v)$ is linear in u and antilinear in v . Using the Cauchy-Schwartz inequality in the Hilbert space $\mathcal{B}^2(\widehat{\Gamma})$, the Plancherel formula and Proposition 3.5, one gets

$$|\text{op}_a^\tau(u, v)| \leq \|\widehat{a}\|_{\mathcal{B}^2(\widehat{\Gamma})} \|\mathcal{W}_{u,v}^\tau\|_{\mathcal{B}^2(\widehat{\Gamma})} = \|a\|_{\mathcal{B}^2(\Gamma)} \|u\|_{L^2(G)} \|v\|_{L^2(G)}.$$

This implies in particular the estimation $\|\text{Op}^\tau(a)\|_{\mathbb{B}[L^2(G)]} \leq \|a\|_{\mathcal{B}^2(\Gamma)}$. This will be improved in the next result, in which we identify the rank-one, the trace-class and the Hilbert-Schmidt operators in $L^2(G)$ as τ -pseudo-differential operators.

THEOREM 3.8. 1. Let us define by

$$\Lambda_{u,v}(w) := \langle w, u \rangle_{L^2(G)} v, \quad \forall w \in L^2(G)$$

the rank-one operator associated to the pair of vectors (u, v) . Then one has

$$\Lambda_{u,v} = \text{Op}^\tau(\mathcal{V}_{u,v}^\tau), \quad \forall u, v \in L^2(\mathbb{G}). \quad (31)$$

2. Let T be a trace-class operator in $L^2(\mathbb{G})$. Then there exist orthonormal sequences $(u_n)_{n \in \mathbb{N}}$, $(v_n)_{n \in \mathbb{N}}$ and a sequence $(\lambda_n)_{n \in \mathbb{N}} \subset \mathbb{C}$ with $\sum_{n \in \mathbb{N}} |\lambda_n| < \infty$ such that

$$T = \sum_{n \in \mathbb{N}} \lambda_n \text{Op}^\tau(\mathcal{V}_{u_n, v_n}^\tau). \quad (32)$$

3. The mapping Op^τ sends unitarily $\mathcal{B}^2(\Gamma)$ onto the Hilbert space composed of all Hilbert-Schmidt operators in $L^2(\mathbb{G})$.

Proof. 1. By the definition (30) and the orthogonality relations (27), one has for $u', v' \in L^2(\mathbb{G})$

$$\begin{aligned} \langle \text{Op}^\tau(\mathcal{V}_{u,v}^\tau)u', v' \rangle_{L^2(\mathbb{G})} &= \langle \mathcal{V}_{u,v}^\tau, \mathcal{V}_{u',v'}^\tau \rangle_{\mathcal{B}^2(\Gamma)} \\ &= \langle u', u \rangle_{L^2(\mathbb{G})} \langle v, v' \rangle_{L^2(\mathbb{G})} \\ &= \langle \Lambda_{u,v}u', v' \rangle_{L^2(\mathbb{G})}. \end{aligned}$$

2. Follows from 1 and from the fact [46, pag. 494] that every trace-class operator T can be written as $T = \sum_{n \in \mathbb{N}} \lambda_n \Lambda_{u_n, v_n}$ with u_n, v_n, λ_n as in the statement.

3. One recalls that Λ defines (by extension) a unitary map $\overline{L^2(\mathbb{G})} \otimes L^2(\mathbb{G}) \rightarrow \mathbb{B}^2[L^2(\mathbb{G})]$ and that \mathcal{V}^τ is also unitary and note that

$$\text{Op}^\tau = \Lambda \circ (\mathcal{V}^\tau)^{-1} = \Lambda \circ (\mathcal{W}^\tau)^{-1} \circ (\mathcal{F} \otimes \mathcal{F}^{-1}). \quad (33)$$

Another proof consists in examining the integral kernel of $\text{Op}^\tau(a)$ given in Proposition 3.9. \square

The unitarity of the map Op^τ can be written in the form

$$\text{Tr}[\text{Op}^\tau(a)\text{Op}^\tau(b)^*] = \int_{\mathbb{G}} \int_{\widehat{\mathbb{G}}} \text{Tr}_\xi[a(x, \xi)b(x, \xi)^*] dm(x) d\widehat{m}(\xi),$$

where Tr refers to the trace in $\mathbb{B}[L^2(\mathbb{G})]$.

PROPOSITION 3.9. *If $a \in \mathcal{B}^2(\Gamma)$, then $\text{Op}^\tau(a)$ is an integral operator with kernel $\text{Ker}_a^\tau \in L^2(\mathbb{G} \times \mathbb{G})$ given by*

$$\text{Ker}_a^\tau := \text{CV}^\tau(\text{id} \otimes \mathcal{F}^{-1})a. \quad (34)$$

Proof. Using the definitions, Plancherel’s Theorem and the unitarity of CV^τ , one gets

$$\begin{aligned} \langle \text{Op}^\tau(a)u, v \rangle_{L^2(\mathbb{G})} &:= \langle a, \mathcal{V}_{u,v}^\tau \rangle_{\mathcal{B}^2(\Gamma)} \\ &= \langle a, (\text{id} \otimes \mathcal{F})(CV^\tau)^{-1}(v \otimes \bar{u}) \rangle_{L^2(\mathbb{G}) \otimes \mathcal{B}^2(\widehat{\mathbb{G}})} \\ &= \langle (\text{id} \otimes \mathcal{F}^{-1})a, (CV^\tau)^{-1}(v \otimes \bar{u}) \rangle_{L^2(\mathbb{G}) \otimes L^2(\mathbb{G})} \\ &= \langle CV^\tau(\text{id} \otimes \mathcal{F}^{-1})a, (v \otimes \bar{u}) \rangle_{L^2(\mathbb{G}) \otimes L^2(\mathbb{G})} \\ &= \int_{\mathbb{G}} \int_{\mathbb{G}} [CV^\tau(\text{id} \otimes \mathcal{F}^{-1})a](x, y)(\bar{v} \otimes u)(x, y) dm(y) dm(x) \\ &= \int_{\mathbb{G}} \left(\int_{\mathbb{G}} [CV^\tau(\text{id} \otimes \mathcal{F}^{-1})a](x, y) u(y) dm(y) \right) \overline{v(x)} dm(x), \end{aligned}$$

completing the proof. □

REMARK 3.10. We rephrase Proposition 3.9 as

$$\text{Op}^\tau = \text{Int} \circ \text{Ker}^\tau = \text{Int} \circ CV^\tau \circ (\text{id} \otimes \mathcal{F}^{-1}), \tag{35}$$

where $\text{Int} : L^2(\mathbb{G} \times \mathbb{G}) \rightarrow \mathbb{B}^2[L^2(\mathbb{G})]$ is given by

$$[\text{Int}(M)u](x) := \int_{\mathbb{G}} M(x, y)u(y) dm(y).$$

Now we see that Op^τ actually coincides with the one defined in (2), at least in a certain sense. Formally, using (34), one gets

$$\text{Ker}_a^\tau(x, y) = \int_{\widehat{\mathbb{G}}} \text{Tr}_\xi \left[a(x\tau(y^{-1}x)^{-1}, \xi) \xi(y^{-1}x) \right] d\widehat{m}(\xi) \tag{36}$$

and this should be compared to (2). The formula (36) is rigorously correct if, for instance, the symbol a belongs to $(\text{id} \otimes \mathcal{F})\mathcal{C}_c(\mathbb{G} \times \mathbb{G})$, since the explicit form (12) of the inverse Fourier transform holds on $\mathcal{FC}_c(\mathbb{G}) \subset \mathcal{F}[A(\mathbb{G}) \cap L^2(\mathbb{G})] = \mathcal{B}^1(\widehat{\mathbb{G}}) \cap \mathcal{B}^2(\widehat{\mathbb{G}})$. Thus we reobtain the formula (2) as

$$\begin{aligned} [\text{Op}^\tau(a)u](x) &= \int_{\mathbb{G}} \text{Ker}_a^\tau(x, y)u(y) dm(y) \\ &= \int_{\mathbb{G}} \left(\int_{\widehat{\mathbb{G}}} \text{Tr}_\xi \left[\xi(y^{-1}x) a(x\tau(y^{-1}x)^{-1}, \xi) \right] d\widehat{m}(\xi) \right) u(y) dm(y). \end{aligned} \tag{37}$$

REMARK 3.11. If $\tau, \tau' : \mathbb{G} \rightarrow \mathbb{G}$ are measurable maps, the associated pseudo-differential calculi are related by $\text{Op}^{\tau'}(a) = \text{Op}^\tau(a_{\tau\tau'})$ where, based on (35), one gets

$$(\text{id} \otimes \mathcal{F}^{-1})a_{\tau\tau'} = [(\text{id} \otimes \mathcal{F}^{-1})a] \circ c\nu^{\tau'} \circ (c\nu^\tau)^{-1}. \tag{38}$$

One computes easily

$$cv^{\tau'\tau}(x, y) := [cv^{\tau'} \circ (cv^\tau)^{-1}](x, y) = (x\tau(y)\tau'(y)^{-1}, y). \quad (39)$$

However, it seems difficult to turn this into a nice explicit formula for $a_{\tau\tau'}$, but this is already the case in the Euclidean space too. The crossed product realisation is nicer from this point of view (when “turned to the right”). Using (47) one can write

$$\text{Sch}^{\tau'}(\Phi) = \text{Sch}^\tau(\Phi_{\tau\tau'}), \quad (40)$$

with $\Phi_{\tau\tau'} = \Phi \circ cv^{\tau'\tau}$. See also Remark 7.4.

3.3 INVOLUTIVE ALGEBRAS OF SYMBOLS

Since our pseudo-differential calculus is one-to-one, we can define an involutive algebra structure on operator-valued symbols, emulating the algebra of operators. One defines a composition law $\#_\tau$ and an involution $\#_\tau$ on $\mathcal{B}^2(\widehat{\Gamma})$ by

$$\text{Op}^\tau(a\#_\tau b) := \text{Op}^\tau(a)\text{Op}^\tau(b),$$

$$\text{Op}^\tau(a^{\#_\tau}) := \text{Op}^\tau(a)^*.$$

The composition can be written in terms of integral kernels as

$$\text{Ker}_{a\#_\tau b}^\tau = \text{Ker}_a^\tau \bullet \text{Ker}_b^\tau,$$

where, by (35),

$$\text{Ker}^\tau := CV^\tau \circ (\text{id} \otimes \mathcal{F}^{-1})$$

and \bullet is the usual composition of kernels

$$(M \bullet N)(x, y) := \int_{\mathbf{G}} M(x, z)N(z, y)dm(z),$$

corresponding to $\text{Int}(M \bullet N) = \text{Int}(M)\text{Int}(N)$. It follows that for $a, b \in \mathcal{B}^2(\widehat{\Gamma})$

$$\begin{aligned} a\#_\tau b &= (\text{Ker}^\tau)^{-1}(\text{Ker}_a^\tau \bullet \text{Ker}_b^\tau) \\ &= (\text{id} \otimes \mathcal{F}) \circ (CV^\tau)^{-1} \left\{ [CV^\tau \circ (\text{id} \otimes \mathcal{F}^{-1})]a \bullet [CV^\tau \circ (\text{id} \otimes \mathcal{F}^{-1})]b \right\}. \end{aligned} \quad (41)$$

Similarly, in terms of the natural kernel involution $M^\bullet(x, y) := \overline{M(y, x)}$ (corresponding to $\text{Int}(M)^* = \text{Int}(M^\bullet)$), one gets

$$a^{\#_\tau} = (\text{Ker}^\tau)^{-1}[(\text{Ker}_a^\tau)^\bullet] = (\text{id} \otimes \mathcal{F}) \circ (CV^\tau)^{-1} \left\{ \left([CV^\tau \circ (\text{id} \otimes \mathcal{F}^{-1})]a \right)^\bullet \right\}. \quad (42)$$

REMARK 3.12. As a conclusion, $(\mathcal{B}^2(\Gamma), \#_\tau, \#_\tau)$ is a $*$ -algebra. This is part of a more detailed result, stating that $(\mathcal{B}^2(\Gamma), \langle \cdot, \cdot \rangle_{\mathcal{B}^2(\Gamma)}, \#_\tau, \#_\tau)$ is an H^* -algebra, i.e. a complete Hilbert algebra [14, App. A]. Among others, this contains the following compatibility relations between the scalar product and the algebraic laws

$$\begin{aligned} \langle a\#_\tau b, c \rangle_{\mathcal{B}^2(\Gamma)} &= \langle a, b\#_\tau c \rangle_{\mathcal{B}^2(\Gamma)}, \\ \langle a, b \rangle_{\mathcal{B}^2(\Gamma)} &= \langle b\#_\tau, a\#_\tau \rangle_{\mathcal{B}^2(\Gamma)}, \end{aligned}$$

valid for every $a, b, c \in \mathcal{B}^2(\Gamma)$. The simplest way to prove all these is to recall that $\mathbb{B}^2[L^2(\mathbb{G})]$ is an H^* -algebra with the operator multiplication, with the adjoint and with the complete scalar product $\langle S, T \rangle_{\mathbb{B}^2} := \text{Tr}[ST^*]$ and to invoke the algebraic and unitary isomorphism $\mathcal{B}^2(\Gamma) \stackrel{\text{Op}^\tau}{\cong} \mathbb{B}^2[L^2(\mathbb{G})]$.

Formulae (41) and (42) take a more explicit integral form on symbols particular enough to allow applying formula (12) for the inverse Fourier transform. Since, anyhow, we will not need such formulas, we do not pursue this here. Let us give, however, the simple algebraic rules satisfied by the Wigner τ -transforms defined in (26):

COROLLARY 3.13. *For every $u, v, u_1, u_2, v_1, v_2 \in L^2(\mathbb{G})$ one has*

$$\mathcal{V}_{u_1, v_1}^\tau \#_\tau \mathcal{V}_{u_2, v_2}^\tau = \langle v_2, u_1 \rangle \mathcal{V}_{u_2, v_1}^\tau \tag{43}$$

and

$$(\mathcal{V}_{u, v}^\tau)^{\#_\tau} = \mathcal{V}_{v, u}^\tau. \tag{44}$$


Proof. The first identity is a consequence of the first point of Theorem 3.8:

$$\begin{aligned} \text{Op}^\tau(\mathcal{V}_{u_1, v_1}^\tau \#_\tau \mathcal{V}_{u_2, v_2}^\tau) &= \text{Op}^\tau(\mathcal{V}_{u_1, v_1}^\tau) \text{Op}^\tau(\mathcal{V}_{u_2, v_2}^\tau) \\ &= \Lambda_{u_1, v_1} \Lambda_{u_2, v_2} \\ &= \langle v_2, u_1 \rangle \Lambda_{u_2, v_1} \\ &= \langle v_2, u_1 \rangle \text{Op}^\tau(\mathcal{V}_{u_2, v_1}^\tau), \end{aligned}$$

which implies (43) because Op^τ is linear and injective.

The relation (44) follows similarly, taking into account the identity $\Lambda_{u, v}^* = \Lambda_{v, u}$. \square

REMARK 3.14. It seems convenient to summarise the situation in the following commutative diagram of unitary mappings (which are even isomorphisms of H^* -algebras):



For completeness and for further use we also included two new maps. The first one is given by the formula $\text{Po}^\tau := \text{Op}^\tau \circ (\mathcal{F}^{-1} \otimes \mathcal{F})$ and it is the integrated form of the family of operators $\{W^\tau(x, \xi) \mid (x, \xi) \in \mathbb{G} \times \widehat{\mathbb{G}}\}$, defined formally by

$$\text{Po}^\tau(\mathfrak{a}) := \int_{\mathbb{G}} \int_{\widehat{\mathbb{G}}} \text{Tr}_\xi [\mathfrak{a}(\xi, x) W^\tau(\xi, x)^*] dm(x) d\widehat{m}(\xi). \tag{45}$$

Here we can think that $\mathfrak{a} = (\mathcal{F} \otimes \mathcal{F}^{-1})a$. It is treated rigorously in the same way as Op^τ ; the correct weak definition is to set for $u, v \in L^2(\mathbb{G})$

$$\langle \text{Po}^\tau(\mathfrak{a})u, v \rangle_{L^2(\mathbb{G})} = \langle \text{Op}^\tau[(\mathcal{F}^{-1} \otimes \mathcal{F})(\mathfrak{a})]u, v \rangle_{L^2(\mathbb{G})} = \langle \mathfrak{a}, \mathcal{W}_{u,v}^\tau \rangle_{\mathcal{B}^2(\widehat{\Gamma})}. \tag{46}$$

The second one is the Schrödinger representation $\text{Sch}^\tau := \text{Int} \circ \text{CV}^\tau$ defined for $\Phi \in L^2(\mathbb{G} \times \mathbb{G})$ by

$$[\text{Sch}^\tau(\Phi)v](x) := \int_{\mathbb{G}} \Phi(x\tau(y^{-1}x)^{-1}, y^{-1}x)v(y) dm(y). \tag{47}$$

It satisfies $\text{Op}^\tau = \text{Sch}^\tau \circ (\text{id} \otimes \mathcal{F}^{-1})$ and we put it into evidence because it is connected to the C^* -algebraic formalism described in Subsection 7.2.

3.4 NON-COMMUTATIVE L^p -SPACES AND SCHATTEN CLASSES

DEFINITION 3.15. For $p \in [1, \infty)$ we introduce the Banach space $\mathcal{B}^{p,p}(\widehat{\Gamma}) := L^p[\mathbb{G}; \mathcal{B}^p(\widehat{\mathbb{G}})]$ with the norm

$$\begin{aligned} \|\mathfrak{a}\|_{\mathcal{B}^{p,p}(\widehat{\Gamma})} &:= \left(\int_{\mathbb{G}} \|\mathfrak{a}(x)\|_{\mathcal{B}^p(\widehat{\mathbb{G}})}^p dm(x) \right)^{1/p} \\ &= \left(\int_{\mathbb{G}} \left[\int_{\widehat{\mathbb{G}}} \|\mathfrak{a}(\xi, x)\|_{\mathbb{B}^p(\mathcal{H}_\xi)}^p d\widehat{m}(\xi) \right] dm(x) \right)^{1/p}, \end{aligned}$$

where the convenient notation $\mathfrak{a}(\xi, x) := [\mathfrak{a}(x)](\xi)$ has been used.

Note that $\mathcal{B}^{1,1}(\widehat{\Gamma}) \cong \mathcal{B}^1(\widehat{\mathbb{G}}) \overline{\otimes} L^1(\mathbb{G})$ (projective completed tensor product), while $\mathcal{B}^{2,2}(\widehat{\Gamma}) \cong \mathcal{B}^2(\widehat{\Gamma}) = \mathcal{B}^2(\widehat{\mathbb{G}}) \otimes L^2(\mathbb{G})$ (Hilbert tensor product). The double index indicates that the spaces $\mathcal{B}^{p,q}(\widehat{\Gamma}) := L^p[\mathbb{G}; \mathcal{B}^q(\widehat{\mathbb{G}})]$ could also be taken into account for $p \neq q$.

To put the definition in a general context, we recall some basic facts about non-commutative L^p -spaces [38, 48]. A non-commutative measure space is a pair $(\mathcal{M}, \mathcal{T})$ formed of a von Neumann algebra \mathcal{M} with positive cone \mathcal{M}_+ , acting in a Hilbert space \mathcal{K} , endowed with a normal semifinite faithful trace $\mathcal{T} : \mathcal{M}_+ \rightarrow [0, \infty]$. One defines

$$\mathcal{S}_+ := \{m \in \mathcal{M}_+ \mid \mathcal{T}[s(m)] < \infty\},$$

where $s(m)$ is the support of m , i.e. the smallest orthogonal projection $e \in \mathcal{M}$ such that $eme = m$. Then \mathcal{S} , defined to be the linear span of \mathcal{S}_+ , is a w^* -dense $*$ -subalgebra of \mathcal{M} . For every $p \in [1, \infty)$, the map $\|\cdot\|_{(p)} : \mathcal{S} \rightarrow [0, \infty)$ given by

$$\|m\|_{(p)} := [\mathcal{T}(|m|^p)]^{1/p} = [\mathcal{T}((m^*m)^{p/2})]^{1/p}$$

is a well-defined norm. The completion of $(\mathcal{S}, \|\cdot\|_{(p)})$ is denoted by $\mathcal{L}^p(\mathcal{M}, \mathcal{T})$ and is called *the non-commutative L^p -space associated to the non-commutative measure space $(\mathcal{M}, \mathcal{T})$* . The scale is completed by setting $\mathcal{L}^\infty(\mathcal{M}, \mathcal{T}) := \mathcal{M}$. It can be shown that $\mathcal{L}^1(\mathcal{M}, \mathcal{T})$ can be viewed as the predual of \mathcal{M} and the elements of $\mathcal{L}^p(\mathcal{M}, \mathcal{T})$ can be interpreted as closed, maybe unbounded, operators in \mathcal{K} [38].

We are going to need two important properties of these non-commutative L^p -spaces.

- *Duality*: if $p \neq \infty$ and $1/p + 1/p' = 1$, then $[\mathcal{L}^p(\mathcal{M}, \mathcal{T})]^* \cong \mathcal{L}^{p'}(\mathcal{M}, \mathcal{T})$ isometrically; the duality is defined by $\langle m, n \rangle_{(p), (p')} := \mathcal{T}(mn^*)$ (consequence of a non-commutative Hölder inequality).
- *Interpolation*: the complex interpolation of these spaces follows the rule

$$[\mathcal{L}^{p_0}(\mathcal{M}, \mathcal{T}), \mathcal{L}^{p_1}(\mathcal{M}, \mathcal{T})]_\theta = \mathcal{L}^p(\mathcal{M}; \mathcal{T}), \quad \theta \in (0, 1), \quad \frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}.$$

In our case the non-commutative measure space can be defined as follows: The von Neumann algebra is

$$\mathcal{B}^{\infty, \infty}(\widehat{\Gamma}) = \mathcal{B}(\widehat{\mathbb{G}}) \widetilde{\otimes} L^\infty(\mathbb{G}) = \int_{\widehat{\mathbb{G}}}^{\oplus} \mathbb{B}(\mathcal{H}_\xi) d\widehat{m}(\xi) \widetilde{\otimes} L^\infty(\mathbb{G})$$

(weak*-completion of the algebraic tensor product). Denoting as before by Tr_ξ the standard trace in $\mathbb{B}(\mathcal{H}_\xi)$, then on $\mathcal{B}(\widehat{\mathbb{G}})$ one has [15, Sect II.5.1] the direct integral trace $\text{Tr} := \int_{\widehat{\mathbb{G}}}^{\oplus} \text{Tr}_\xi d\widehat{m}(\xi)$ and on $\mathcal{B}(\widehat{\mathbb{G}}) \widetilde{\otimes} L^\infty(\mathbb{G})$ the tensor product [48, 1.7.5] $\mathcal{T} := \text{Tr} \otimes \int_{\mathbb{G}}$ of Tr with the trace given by Haar integration in the commutative von Neumann algebra $L^\infty(\mathbb{G})$. Thus one gets the non-commutative measure space $(\mathcal{B}^{\infty, \infty}(\widehat{\Gamma}), \mathcal{T})$. It is not difficult to show that the associated non-commutative L^p -spaces are the Banach spaces $\mathcal{B}^{p, p}(\widehat{\Gamma})$ introduced in Definition 3.15 ([48, 1.7.5] is useful again). In particular, we have the following rule of complex interpolation:

$$[\mathcal{B}^{p_0, p_0}(\widehat{\Gamma}), \mathcal{B}^{p_1, p_1}(\widehat{\Gamma})]_\theta = \mathcal{B}^{p, p}(\widehat{\Gamma}), \quad \theta \in (0, 1), \quad \frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}.$$

On the other hand, the Schatten-von Neumann ideals $\mathbb{B}^p[L^2(\mathbb{G})]$ are the non-commutative L^p -spaces associated to the non-commutative measure space $(\mathbb{B}[L^2(\mathbb{G})], \text{Tr})$. So they interpolate in the same way.

PROPOSITION 3.16. *For every $p \in [2, \infty]$ one has a linear contraction*

$$\mathcal{W}^\tau : \overline{L^2(\mathbb{G})} \otimes L^2(\mathbb{G}) \rightarrow \mathcal{B}^{p, p}(\widehat{\Gamma}). \tag{48}$$

Proof. We have seen in Proposition 3.5 that \mathcal{W}^τ is unitary if $p = 2$. If we also check the case $p = \infty$, then (48) follows by complex interpolation. But the uniform estimate

$$\|\mathcal{W}_{u, v}^\tau(\xi, x)\|_{\mathbb{B}(\mathcal{H}_\xi)} \leq \|u\|_{L^2(\mathbb{G})} \|v\|_{L^2(\mathbb{G})}$$

is an immediate consequence of (22) and of the unitarity of $\mathcal{W}^\tau(\xi, x)$ in $L^2(\mathbb{G}; \mathcal{H}_\xi)$. □

For the next two results we switch our interest from Op^τ to Po^τ , given by (46), since for such general groups G there is no inversion formula for the Fourier transform at the level of the non-commutative L^p -spaces (the Hausdorff-Young inequality cannot be used for our purposes).

THEOREM 3.17. *If $\mathfrak{a} \in \mathcal{B}^{1,1}(\widehat{\Gamma}) = L^1[G; \mathcal{B}^1(\widehat{G})]$ then $\text{Po}^\tau(\mathfrak{a})$ is bounded in $L^2(G)$ and*

$$\|\text{Po}^\tau(\mathfrak{a})\|_{\mathbb{B}[L^2(G)]} \leq \|\mathfrak{a}\|_{\mathcal{B}^{1,1}(\widehat{\Gamma})}.$$

Proof. One modifies (46) to a similar definition by duality

$$\langle \text{Po}^\tau(\mathfrak{a})u, v \rangle_{L^2(G)} = \langle \mathfrak{a}, \mathcal{W}_{u,v}^\tau \rangle_{(1),(\infty)} := \mathcal{T}(\mathfrak{a} [\mathcal{W}_{u,v}^\tau]^*),$$

based on the case $p = \infty$ of Proposition 3.16 and on the duality of the non-commutative Lebesgue spaces. \square

By using complex interpolation between the end points $p_0 = 2$ and $p_1 = \infty$, one gets

COROLLARY 3.18. *If $p \in [1, 2]$, $\frac{1}{p} + \frac{1}{p'} = 1$ and $\mathfrak{a} \in L^p[G; \mathcal{B}^p(\widehat{G})]$, then $\text{Po}^\tau(\mathfrak{a})$ belongs to $\mathbb{B}^{p'}[L^2(G)]$ and*

$$\|\text{Po}^\tau(\mathfrak{a})\|_{\mathbb{B}^{p'}[L^2(G)]} \leq \|\mathfrak{a}\|_{\mathcal{B}^{p,p}(\widehat{\Gamma})}.$$

More refined results follow from real interpolation; the interested reader could write them down easily.

4 SYMMETRIC QUANTIZATIONS

Having in mind the well-known [22] Weyl quantization, we inquire about the existence of a parameter τ allowing a symmetric quantization; if it exists, for emphasis, we denote it by σ . By definition, this means that $a^{\#\sigma} = a^*$ for every $a \in \mathcal{B}^2(\Gamma)$, where of course $a^*(x, \xi) := a(x, \xi)^*$ (adjoint in $\mathbb{B}(\mathcal{H}_\xi)$) for every $(x, \xi) \in \Gamma$. At the level of pseudo-differential operators the consequence would be the simple relation $\text{Op}^\sigma(a)^* = \text{Op}^\sigma(a^*)$, so “real-valued symbols” are sent into self-adjoint operators.

4.1 AN EXPLICIT FORM FOR THE ADJOINT

In order to study symmetry it is convenient to give a more explicit form of the involution (42); we need to allow different values of the parameter τ . For any measurable map $\tau : G \rightarrow G$, let us define

$$\tilde{\tau} : G \rightarrow G, \quad \tilde{\tau}(x) := \tau(x^{-1})x. \quad (49)$$

It is worth mentioning that if $\tau(\cdot) = e$ then $\tilde{\tau} = \text{id}_G$ and if $\tau = \text{id}_G$ then $\tilde{\tau}(\cdot) = e$. In addition $\tilde{\tilde{\tau}} = \tau$ holds.

If $G = \mathbb{R}^n$ and $\tau := t \text{id}_{\mathbb{R}^n}$ with $t \in [0, 1]$, one has $\tilde{\tau} = (1 - t)\text{id}_{\mathbb{R}^n}$ and the next proposition is well-known for pseudo-differential operators on \mathbb{R}^n .

PROPOSITION 4.1. *One has $\text{Op}^\tau(a)^* = \text{Op}^{\tau'}(a^*)$ for every $a \in \mathcal{B}(\Gamma)$ if and only if $\tau' = \tilde{\tau}$.*

Proof. Hoping that $\text{Op}^\tau(a)^* = \text{Op}^{\tau'}(a^*)$ for some $\tau' : G \rightarrow G$, by (35), one has to examine the equality

$$\left([\text{CV}^\tau \circ (\text{id} \otimes \mathcal{F}^{-1})] a \right)^\bullet = [\text{CV}^{\tau'} \circ (\text{id} \otimes \mathcal{F}^{-1})] a^*.$$

This and the next identities should hold almost everywhere with respect to the product measure $m \otimes m$. Using the easy relation

$$[(\text{id} \otimes \mathcal{F}^{-1})a^*](y, z) = \overline{[(\text{id} \otimes \mathcal{F}^{-1})a](y, z^{-1})},$$

one gets immediately

$$\left([\text{CV}^{\tau'} \circ (\text{id} \otimes \mathcal{F}^{-1})] a^* \right)(y, z) = \overline{[(\text{id} \otimes \mathcal{F}^{-1})a](y\tau'(z^{-1}y)^{-1}, y^{-1}z)}. \tag{50}$$

On the other hand

$$\begin{aligned} \left([\text{CV}^\tau \circ (\text{id} \otimes \mathcal{F}^{-1})] a \right)^\bullet(y, z) &= \overline{\left([\text{CV}^\tau \circ (\text{id} \otimes \mathcal{F}^{-1})] a \right)(z, y)} \\ &= \overline{[(\text{id} \otimes \mathcal{F}^{-1})a](z\tau(y^{-1}z)^{-1}, y^{-1}z)} \end{aligned} \tag{51}$$

and the two expressions (50) and (51) always coincide $m \otimes m$ -almost everywhere if and only if

$$y\tau'(z^{-1}y)^{-1} = z\tau(y^{-1}z)^{-1}, \quad m \otimes m - \text{a.e. } (y, z) \in G \times G.$$

This condition can be transformed into

$$\tau'(z^{-1}y) = \tau(y^{-1}z)z^{-1}y, \quad m \otimes m - \text{a.e. } (y, z) \in G \times G,$$

which must be shown to be equivalent to $\tau' = \tilde{\tau}$ holding m -almost everywhere.

This follows if we prove that $A \subset G$ is m -negligible if and only if $M(A) := \{(y, z) \in G \times G \mid z^{-1}y \in A\}$ is $m \otimes m$ -negligible. Since m is σ -finite, there is a Borel partition $G = \sqcup_{n \in \mathbb{N}} B_n$ with $m(B_n) < \infty$ for every $n \in \mathbb{N}$. Thus $M(A) = \sqcup_{n \in \mathbb{N}} M_n(A)$, with

$$M_n(A) := \{(y, z) \in M(A) \mid z \in B_n\} = \{(y, z) \in G \times B_n \mid y \in zA\}.$$

Using the invariance of the Haar measure, one checks that $(m \otimes m)[M_n(A)] = m(A)m(B_n)$ and the conclusion follows easily. \square

4.2 SYMMETRY FUNCTIONS

The measurable function $\sigma : G \rightarrow G$ is called a *symmetry function* if one has $\text{Op}^\sigma(a)^* = \text{Op}^\sigma(a^*)$ for every $a \in \mathcal{B}^2(\Gamma)$. When G is admissible and a symmetry function exists we say that *the group G admits a symmetric quantization*. As a consequence of Proposition 4.1 one gets

COROLLARY 4.2. *The map $\sigma : \mathbb{G} \rightarrow \mathbb{G}$ is a symmetry function if and only if for almost every $x \in \mathbb{G}$*

$$\sigma(x) = \sigma(x^{-1})x. \quad (52)$$

In particular, if σ is a symmetry function and $a(\cdot, \cdot) \in \mathcal{B}^2(\Gamma)$ is self-adjoint pointwise (or $m \otimes \widehat{m}$ -almost everywhere) then $\text{Op}^\sigma(a)$ is a self-adjoint operator in $L^2(\mathbb{G})$.

The problem of existence of σ satisfying (52) seems rather obscure in general, so we only treat some particular cases.

PROPOSITION 4.3. *1. The product $\mathbb{G} := \prod_{k=1}^m \mathbb{G}_k$ of a family of groups admitting a symmetric quantization also admits a symmetric quantization.*

2. The admissible central extension of a group admitting a symmetric quantization by another group with this property is a group admitting a symmetric quantization.

3. Any exponential Lie group (in particular any connected simply connected nilpotent group) admits a symmetric quantization.

Proof. 1. Finite products of admissible groups are admissible. If σ_k is a symmetry function for \mathbb{G}_k , then $\sigma[(x_k)_k] := (\sigma_k(x_k))_k$ defines a symmetry function for \mathbb{G} .

2. The structure of central group extensions can be described in terms of 2-cocycles up to canonical isomorphisms. Let \mathbb{N} be an Abelian locally compact group, \mathbb{H} a locally compact group and $\varpi : \mathbb{H} \times \mathbb{H} \rightarrow \mathbb{N}$ a 2-cocycle. On $\mathbb{G} := \mathbb{H} \times \mathbb{N}$ one has the composition law and the inversion

$$(h_1, n_1)(h_2, n_2) := (h_1 h_2, n_1 n_2 \varpi(h_1, h_2)), \quad (h, n)^{-1} := (h^{-1}, \varpi(h^{-1}, h)^{-1} n^{-1}). \quad (53)$$

The properties of ϖ are normalisation $\varpi(h, e_{\mathbb{H}}) = e_{\mathbb{N}} = \varpi(e_{\mathbb{H}}, h)$, $\forall h \in \mathbb{H}$, and the 2-cocycle identity

$$\varpi(h_1, h_2)\varpi(h_1 h_2, h_3) = \varpi(h_2, h_3)\varpi(h_1, h_2 h_3), \quad \forall h_1, h_2, h_3 \in \mathbb{H}. \quad (54)$$

We are given symmetry functions $\sigma_{\mathbb{H}} : \mathbb{H} \rightarrow \mathbb{H}$ and $\sigma_{\mathbb{N}} : \mathbb{N} \rightarrow \mathbb{N}$ thus satisfying

$$\sigma_{\mathbb{H}}(h) = \sigma_{\mathbb{H}}(h^{-1})h, \quad \sigma_{\mathbb{N}}(n) = \sigma_{\mathbb{N}}(n^{-1})n = n\sigma_{\mathbb{N}}(n^{-1}).$$

We define the measurable map $\sigma : \mathbb{G} \rightarrow \mathbb{G}$ by

$$\begin{aligned} \sigma(h, n) &:= \left(\sigma_{\mathbb{H}}(h), \sigma_{\mathbb{N}}[\varpi(\sigma_{\mathbb{H}}(h^{-1}), h)n] \right) \\ &= \left(\sigma_{\mathbb{H}}(h^{-1})h, \varpi(\sigma_{\mathbb{H}}(h^{-1}), h)n\sigma_{\mathbb{N}}[\varpi(\sigma_{\mathbb{H}}(h^{-1}), h)^{-1}n^{-1}] \right); \end{aligned} \quad (55)$$

the second line has been deduced from the first by using the properties of $\sigma_{\mathbb{H}}$ and $\sigma_{\mathbb{N}}$ and the fact that \mathbb{N} is commutative. We compute using (53) and (55)

$$\begin{aligned} \sigma((h, n)^{-1})(h, n) &= \sigma(h^{-1}, \varpi(h^{-1}, h)^{-1}n^{-1})(h, n) \\ &= \left(\sigma_{\mathbb{H}}(h^{-1}), \sigma_{\mathbb{N}}[\varpi(\sigma_{\mathbb{H}}(h), h^{-1})\varpi(h^{-1}, h)^{-1}n^{-1}] \right)(h, n) \\ &= \left(\sigma_{\mathbb{H}}(h^{-1})h, n\sigma_{\mathbb{N}}[\varpi(\sigma_{\mathbb{H}}(h), h^{-1})\varpi(h^{-1}, h)^{-1}n^{-1}]\varpi(\sigma_{\mathbb{H}}(h^{-1}), h) \right). \end{aligned} \quad (56)$$

The first components in (55) and (56) are equal. The second ones would coincide if one shows

$$\varpi(\sigma_H(h^{-1}), h)^{-1} = \varpi(\sigma_H(h), h^{-1})\varpi(h^{-1}, h)^{-1},$$

which is equivalent to

$$\varpi(\sigma_H(h), h^{-1})\varpi(\sigma_H(h^{-1}), h) = \varpi(h^{-1}, h). \tag{57}$$

Taking in (54) $h_3 = h_2^{-1} = h$ and using the normalization of ϖ one gets

$$\varpi(h_1, h^{-1})\varpi(h_1 h^{-1}, h) = \varpi(h^{-1}, h), \quad \forall h_1, h \in H.$$

Choosing $h_1 := \sigma_H(h)$ we get (57), because σ_H is a symmetry function.

3. Assume that G is a Lie group with Lie algebra \mathfrak{g} . It is known that the exponential map $\exp : \mathfrak{g} \rightarrow G$ restricts to a diffeomorphism $\exp : \mathfrak{u} \rightarrow U$, where \mathfrak{u} is a neighborhood of $0 \in \mathfrak{g}$ (a ball centered in the origin, for example) and U is a neighborhood of $e \in G$. The inverse diffeomorphism is denoted by $\log : U \rightarrow \mathfrak{u}$. One defines “the midpoint mapping”

$$\sigma : U \rightarrow G, \quad \sigma(x) := \int_0^1 \exp[s \log x] ds. \tag{58}$$

We claim that

$$x \sigma(x^{-1}) = \sigma(x) = \sigma(x^{-1}) x, \quad \forall x \in G. \tag{59}$$

We prove the second equality; the first one is similar:

$$\begin{aligned} \sigma(x^{-1}) x &= \int_0^1 \exp[s \log(x^{-1})] ds x \\ &= \int_0^1 \exp[-s \log(x)] ds \exp[\log(x)] \\ &= \int_0^1 \exp[(1-s) \log(x)] ds \\ &= \int_0^1 \exp[s \log(x)] ds = \sigma(x). \end{aligned}$$

An exponential group G is, by definition, a Lie group for which one can take $\mathfrak{u} = \mathfrak{g}$ and $U = G$ (i.e. the exponential map is a global diffeomorphism), case in which the symmetry function σ is globally defined. In addition G is admissible. Connected simply connected nilpotent group are exponential, by [7, Th. 1.2.1]. □

EXAMPLE 4.4. For $G = \mathbb{R}^n$ one just sets $\sigma(x) := x/2$ (getting finally the Weyl quantization).

If σ is required to satisfy $\sigma(\cdot^{-1}) = \sigma(\cdot)^{-1}$ (e.g. being an endomorphism), (52) reduces to $x = \sigma(x)^2$ for almost every x , which is equivalent to “any” element in G having a square root; such a group does admit a symmetric quantization. This fails for many groups, as $G := \mathbb{Z}^n$ for instance.

5 EXTENSION TO DISTRIBUTIONS

Having started with the formalism involving symbols in $\mathcal{B}^2(\Gamma)$ and operators bounded on $L^2(G)$, it is useful to be able to extend it to e.g. unbounded symbols and to operators which are only densely defined on $L^2(G)$. If the group G is a Lie group (has a smooth manifold structure), we can, for example, think of operators acting from the space of test functions to the space of distributions, or of operators having e.g. polynomial growth of symbols. Without assuming that G is a Lie group we do not have the usual space of smooth compactly supported functions readily available as the standard space of test functions. So we will be using its generalisation to locally compact groups by Bruhat [4], and these Bruhat spaces $\mathcal{D}(G)$ and $\mathcal{D}'(G)$ will replace the usual spaces of test functions and distributions, respectively, in our setting.

5.1 SMOOTH FUNCTIONS AND DISTRIBUTIONS – BRUHAT SPACES

The Bruhat spaces $\mathcal{D}(G)$ and $\mathcal{D}'(G)$ have been introduced in [4], to which we refer for further details. Most of their properties hold for every locally compact group, but in some cases second countability is also used. Postliminarity, unimodularity or amenability are not needed.

A *good subgroup* of G is a compact normal subgroup H of G such that G/H is (isomorphic to) a Lie group. The family of all good subgroups of G will be denoted by $\text{good}(G)$; it is stable under intersections. We are going to assume first that

$$\bigcap_{H \in \text{good}(G)} H = \{e\}. \quad (60)$$

Denoting the connected component of the identity by G_0 , this happens, for instance, if G/G_0 is compact, in particular if G is connected. If (60) holds, every neighborhood of e contains an element of $\text{good}(G)$. Then G can be seen as the projective limit of the projective family of Lie groups

$$\{G/H \rightarrow G/K \mid H, K \in \text{good}(G), K \subset H\}.$$

For every good group H one sets $\mathcal{D}(G/H) := \mathcal{C}_c^\infty(G/H)$ with the usual inductive limit topology. Functions on quotients are identified with invariant functions on the group by the map $j_H(v) := v \circ q_H$, where $q_H : G \rightarrow G/H$ is the canonical surjection. Thus on

$$\mathcal{D}_H(G) := j_H[\mathcal{D}(G/H)] \subset \mathcal{C}_c(G) \subset \mathcal{C}_0(G)$$

one can transport the topology of $\mathcal{D}(G/H)$.

DEFINITION 5.1. *The Bruhat space $\mathcal{D}(G)$ of the locally compact group G is the topological inductive limit of the family of subspaces $\{\mathcal{D}_H(G) \mid H \in \text{good}(G)\}$ of $\mathcal{C}_c(G)$.*

The strong dual of $\mathcal{D}(G)$ is denoted by $\mathcal{D}'(G)$; it contains $\mathcal{D}(G)$ densely. Its elements are called distributions.

The space $\mathcal{D}(\mathbb{G})$ is barrelled and bornological. It is continuously and densely contained in $\mathcal{C}_c(\mathbb{G})$ and complete. If \mathbb{G} is already a Lie group, then $\{e\} \in \text{good}(\mathbb{G})$ and clearly $\mathcal{D}(\mathbb{G}) = \mathcal{C}_c^\infty(\mathbb{G})$.

The spaces $\mathcal{D}(\mathbb{G})$ and $\mathcal{D}'(\mathbb{G})$ are complete and Montel (thus reflexive) as well as nuclear.

We presented above the case in which our locally compact group satisfies $\bigcap_{H \in \text{good}(\mathbb{G})} H = \{e\}$. Following [4, Sect. 2], we briefly indicate what to do without this assumption.

There exists an open subgroup G_1 of \mathbb{G} such that $\bigcap_{H_1 \in \text{good}(G_1)} H_1 = \{e\}$. Thus the space $\mathcal{D}(G_1) \subset \mathcal{C}_c(G_1)$ is available. The group \mathbb{G} is partitioned in classes modulo G_1 to the left: $\mathbb{G} = \bigcup xG_1$ or to the right: $\mathbb{G} = \bigcup G_1x$. By left translations one generates the subspaces $\mathcal{D}(xG_1)$ of $\mathcal{C}_c(\mathbb{G})$ (with the transported topology); the elements are particular types of continuous functions on \mathbb{G} compactly supported in xG_1 . Then define $\mathcal{D}^L(\mathbb{G}) \subset \mathcal{C}_c(\mathbb{G})$ to be the topological direct sum $\mathcal{D}^L(\mathbb{G}) := \bigoplus \mathcal{D}(xG_1)$. It comes out that the similarly constructed $\mathcal{D}^R(\mathbb{G}) := \bigoplus \mathcal{D}(G_1x)$ is the same subspace of $\mathcal{C}_c(\mathbb{G})$ with the same topology. In addition, it does not depend on the choice of the open subgroup G_1 so it deserves the notation $\mathcal{D}(\mathbb{G})$.

Then the construction of the space $\mathcal{D}'(\mathbb{G})$ follows similarly and all the nice properties mentioned above still hold (cf. [4]). The main reason is the fact that topological direct sums are rather easy to control. Subsequently we will have recourse to these Bruhat spaces in the general case.

We are going to use the symbol $\overline{\otimes}$ for the projective tensor product of locally convex spaces; note, however, that this will only be applied to spaces known to be nuclear. By the Kernel Theorem for Bruhat spaces [4, Sect. 5] one has

$$\mathcal{D}(\mathbb{G} \times \mathbb{G}) \cong \mathcal{D}(\mathbb{G}) \overline{\otimes} \mathcal{D}(\mathbb{G}) \subset L^2(\mathbb{G} \times \mathbb{G})$$

continuously and densely. Soon we are going to need the next result:

LEMMA 5.2. *The mapping*

$$CV : \mathcal{D}(\mathbb{G} \times \mathbb{G}) \rightarrow \mathcal{D}(\mathbb{G} \times \mathbb{G}), \quad [CV(\Psi)](x, y) := \Psi[cv(x, y)] = \Psi(x, y^{-1}x) \tag{61}$$

is a well-defined topological isomorphism. Its inverse CV^{-1} is the operation of composing with

$$cv^{-1} : \mathbb{G} \times \mathbb{G} \rightarrow \mathbb{G} \times \mathbb{G}, \quad cv^{-1}(x, y) := (x, xy^{-1}).$$

By transposing the inverse one gets a topological isomorphism

$$CV := [CV^{-1}]^{\text{tr}} : \mathcal{D}'(\mathbb{G} \times \mathbb{G}) \rightarrow \mathcal{D}'(\mathbb{G} \times \mathbb{G}),$$

which is an extension of the one given in (61) (this explains the notational abuse).

Proof. The proof is quite straightforward, but rather long if all the details are included, so it is essentially left to the reader. Besides using the definitions and the standard tools of duality, one should also note the following:

- If H is a good subgroup of G , then $H \times H$ is a good subgroup of $G \times G$ and $(G \times G)/(H \times H)$ is canonically isomorphic to $(G/H) \times (G/H)$.
- For $H \in \text{good}(G)$ there is a *Lie group isomorphism*

$$cv_H : (G/H) \times (G/H) \rightarrow (G/H) \times (G/H),$$

$$cv_H(xH, yH) := ((xH), (yH)^{-1}xH) = (xH, y^{-1}xH)$$

and related to the initial change of variables cv through

$$cv_H \circ (q_H \times q_H) = (q_H \times q_H) \circ cv.$$

This and the fact that cv_H is a proper map easily allow us to conclude that $CV : \mathcal{D}_{H \times H}(G \times G) \rightarrow \mathcal{D}_{H \times H}(G \times G)$ is a well-defined isomorphism for every good subgroup H .

- Let G_1 be a subgroup of G ; then cv carries $G_1 \times G_1$ into itself isomorphically.
- Let G_1 be an open subgroup of G such that $\bigcap_{H_1 \in \text{good}(G_1)} H_1 = \{e\}$. Then the family $\{H_1 \times H_1 \mid H_1 \in \text{good}(G_1)\}$ is directed under inclusion and

$$\bigcap_{H_1 \in \text{good}(G_1)} H_1 \times H_1 = \{(e, e)\}.$$

□

REMARK 5.3. Of course, the case $\tau(x) = x$ can be treated the same way. If one tries to do the same for the change of variables cv^τ , in general one encounters rather complicated conditions relating the map τ to the family $\text{good}(G)$. However, if G is a Lie group, $\text{good}(G)$ has a smallest element $\{e\}$ and thus $\mathcal{D}(G)$ coincides with $\mathcal{C}_c^\infty(G)$. Then it is easy to see that the statements of the lemma hold if cv^τ is proper and $\tau : G \rightarrow G$ is a \mathcal{C}^∞ -function.

5.2 RESTRICTIONS AND EXTENSIONS OF THE PSEUDO-DIFFERENTIAL CALCULUS

Let us define $\mathcal{D}(\widehat{G}) := \mathcal{F}[\mathcal{D}(G)]$ with the locally convex topological structure transported from the Bruhat space $\mathcal{D}(G)$. One has $\mathcal{D}(G) \subset \mathcal{C}_c(G) \subset L^2(G) \cap A(G)$ (continuously and densely), so $\mathcal{D}(\widehat{G})$ is a dense subspace of $\mathcal{B}^2(\widehat{G}) \cap \mathcal{B}^1(\widehat{G})$ (with the intersection topology) and of $\mathcal{B}^2(\widehat{G})$. Thus the explicit form of the inverse (12) holds on $\mathcal{D}(\widehat{G})$. One also has

$$u(e) = \int_{\widehat{G}} \text{Tr}_\xi[(\mathcal{F}u)(\xi)] d\widehat{m}(\xi), \quad \forall u \in \mathcal{D}(G).$$

We are going to use the dense subspace

$$\mathcal{D}(\Gamma) \equiv \mathcal{D}(G \times \widehat{G}) := \mathcal{D}(G) \overline{\otimes} \mathcal{D}(\widehat{G}) \subset \mathcal{B}^2(\Gamma),$$

possessing its own locally convex topology, obtained by transport of structure and the completed projective tensor product construction. Taking also into account the strong dual, one gets a Gelfand triple $\mathcal{D}(\Gamma) \hookrightarrow \mathcal{B}^2(\Gamma) \hookrightarrow \mathcal{D}'(\Gamma)$.

PROPOSITION 5.4. *The calculus $\text{Op} : L^2(\mathbb{G}) \otimes \mathcal{B}^2(\widehat{\mathbb{G}}) \rightarrow \mathbb{B}^2[L^2(\mathbb{G})]$*

- *restricts to a topological isomorphism $\text{Op} : \mathcal{D}(\mathbb{G}) \overline{\otimes} \mathcal{D}(\widehat{\mathbb{G}}) \rightarrow \mathbb{B}[\mathcal{D}'(\mathbb{G}); \mathcal{D}(\mathbb{G})]$,*
- *extends to a topological isomorphism $\text{Op} : \mathcal{D}'(\mathbb{G}) \overline{\otimes} \mathcal{D}'(\widehat{\mathbb{G}}) \rightarrow \mathbb{B}[\mathcal{D}(\mathbb{G}); \mathcal{D}'(\mathbb{G})]$.*

Proof. The proof can essentially be read in the diagrams

$$\begin{array}{ccc}
 \mathcal{D}(\mathbb{G}) \overline{\otimes} \mathcal{D}(\widehat{\mathbb{G}}) & \xrightarrow{\text{id} \overline{\otimes} \mathcal{F}^{-1}} & \mathcal{D}(\mathbb{G}) \overline{\otimes} \mathcal{D}(\mathbb{G}) \cong \mathcal{D}(\mathbb{G} \times \mathbb{G}) \\
 \downarrow \text{Op} & & \downarrow \text{CV} \\
 \mathbb{B}[\mathcal{D}'(\mathbb{G}); \mathcal{D}(\mathbb{G})] & \xleftarrow{\text{Int}} & \mathcal{D}(\mathbb{G} \times \mathbb{G})
 \end{array}$$

and

$$\begin{array}{ccc}
 \mathcal{D}'(\mathbb{G}) \overline{\otimes} \mathcal{D}'(\widehat{\mathbb{G}}) & \xrightarrow{\text{id} \overline{\otimes} \mathcal{F}^{-1}} & \mathcal{D}'(\mathbb{G}) \overline{\otimes} \mathcal{D}'(\mathbb{G}) \cong \mathcal{D}'(\mathbb{G} \times \mathbb{G}) \\
 \downarrow \text{Op} & & \downarrow \text{CV} \\
 \mathbb{B}[\mathcal{D}(\mathbb{G}); \mathcal{D}'(\mathbb{G})] & \xleftarrow{\text{Int}} & \mathcal{D}'(\mathbb{G} \times \mathbb{G})
 \end{array}$$

The vertical arrows to the right are justified by Lemma 5.2. We leave the details to the reader. □

Techniques from [29] could be applied to define and study large Moyal algebras of vector-valued symbols corresponding to the spaces $\mathbb{B}[\mathcal{D}(\mathbb{G})]$ and $\mathbb{B}[\mathcal{D}'(\mathbb{G})]$ of operators.

5.3 COMPACTNESS CRITERIA

The next result shows that compactness of sets, operators and families of operators in the Hilbert space $L^2(\mathbb{G})$ can be characterised by localisation with pseudo-differential operators with symbols in $\mathcal{D}(\Gamma)$. We adapt the methods of proof from [28], whose framework cannot be applied directly.

THEOREM 5.5. 1. A bounded subset Δ of $L^2(\mathbb{G})$ is relatively compact if and only if for every $\epsilon > 0$ there exists $a \in \mathcal{D}(\Gamma)$ such that

$$\sup_{u \in \Delta} \|\text{Op}^\tau(a)u - u\|_{L^2(\mathbb{G})} \leq \epsilon. \quad (62)$$

2. Let \mathcal{X} be a Banach space. An element $T \in \mathbb{B}[\mathcal{X}, L^2(\mathbb{G})]$ is a compact operator if and only if for every $\epsilon > 0$ there exists $a \in \mathcal{D}(\Gamma)$ such that

$$\|\text{Op}^\tau(a)T - T\|_{\mathbb{B}[\mathcal{X}, L^2(\mathbb{G})]} \leq \epsilon.$$

3. Let $\mathcal{L} \subset \mathbb{B}[L^2(\mathbb{G})]$ be a family of bounded operators. Then $\mathcal{L} \subset \mathbb{K}[L^2(\mathbb{G})]$ and its closure in $\mathbb{K}[L^2(\mathbb{G})]$ is compact if and only if for every $\epsilon > 0$ there exists $a \in \mathcal{D}(\Gamma)$ such that

$$\sup_{T \in \mathcal{L}} (\|\text{Op}^\tau(a)T - T\|_{\mathbb{B}[L^2(\mathbb{G})]} + \|\text{Op}^\tau(a)T^* - T^*\|_{\mathbb{B}[L^2(\mathbb{G})]}) \leq \epsilon.$$

Proof. 1. If Δ is relatively compact, it is totally bounded. Thus, for every $\epsilon > 0$, there is a finite set F such that for each $u \in \Delta$ there exists $u' \in F$ with $\|u - u'\|_{L^2(\mathbb{G})} \leq \epsilon/4$. This finite subset generates a finite-dimensional subspace $\mathcal{H}_F \subset L^2(\mathbb{G})$ with finite-rank corresponding projection P_F . Then for every $u \in \Delta$, recalling our choice for u' and the fact that $P_F u' = u'$, one gets

$$\begin{aligned} \|P_F u - u\|_{L^2(\mathbb{G})} &\leq \|P_F u - P_F u'\|_{L^2(\mathbb{G})} + \|P_F u' - u'\|_{L^2(\mathbb{G})} + \|u' - u\|_{L^2(\mathbb{G})} \\ &\leq 2 \|u - u'\|_{L^2(\mathbb{G})} \leq \epsilon/2. \end{aligned}$$

Let now $M := \sup_{u \in \Delta} \|u\|_{L^2(\mathbb{G})}$; if we find $a \in \mathcal{D}(\Gamma)$ such that

$$\|\text{Op}^\tau(a) - P_F\|_{\mathbb{B}[L^2(\mathbb{G})]} \leq \epsilon/2M \quad (63)$$

one writes for every $u \in \Delta$

$$\|\text{Op}^\tau(a)u - u\|_{L^2(\mathbb{G})} \leq \|\text{Op}^\tau(a)u - P_F u\|_{L^2(\mathbb{G})} + \|P_F u - u\|_{L^2(\mathbb{G})} \leq \epsilon/2 + \epsilon/2 = \epsilon$$

and the formula (62) is proved.

Since $\mathcal{D}(\mathbb{G} \times \mathbb{G})$ is dense in $L^2(\mathbb{G} \times \mathbb{G})$, the subspace $\mathcal{D}(\Gamma)$ is dense in $\mathcal{B}^2(\Gamma)$. Consequently, $\text{Op}^\tau : \mathcal{B}^2(\Gamma) \rightarrow \mathbb{B}[L^2(\mathbb{G})]$ being unitary, one even gets an improved version of (63) with the operator norm replaced by the Hilbert-Schmidt norm. This finishes the ‘‘only if’’ implication.

We now prove the converse assertion. Fix $\epsilon > 0$ and choose $a \in \mathcal{D}(\Gamma)$ such that

$$\sup_{u \in \Delta} \|\text{Op}^\tau(a)u - u\|_{L^2(\mathbb{G})} \leq \epsilon/2.$$

Since $\mathcal{D}(\Gamma) \subset \mathcal{B}^2(\Gamma)$, the operator $\text{Op}^\tau(a)$ is Hilbert-Schmidt, in particular compact. The set Δ is assumed bounded, thus the range $\text{Op}^\tau(a)\Delta$ is totally bounded.

Consequently, there is a finite set E such that for each $u \in \Delta$ there exists $u'' \in E$ satisfying $\|\text{Op}^\tau(a)u - u''\|_{L^2(\mathbb{G})} \leq \epsilon/2$. Therefore

$$\|u - u''\|_{L^2(\mathbb{G})} \leq \|u - \text{Op}^\tau(a)u\|_{L^2(\mathbb{G})} + \|\text{Op}^\tau(a)u - u''\|_{L^2(\mathbb{G})} \leq \epsilon/2 + \epsilon/2 = \epsilon,$$

so the set Δ is totally bounded, thus relatively compact.

2. The operator T is compact if and only if $\Delta := T(\{w \in \mathcal{X} \mid \|w\|_{\mathcal{X}} \leq 1\})$ is relatively compact in the Hilbert space $L^2(\mathbb{G})$. By 1, this happens exactly when for every $\epsilon > 0$ there is a symbol a in $\mathcal{D}(\Gamma)$ such that

$$\|\text{Op}^\tau(a)T - T\|_{\mathbb{B}[\mathcal{X}, L^2(\mathbb{G})]} = \sup_{\|w\|_{\mathcal{X}} \leq 1} \|\text{Op}^\tau(a) - 1\|(Tw)\|_{L^2(\mathbb{G})} \leq \epsilon.$$

3. The set \mathcal{L} is called *collectively compact* if $\bigcup_{T \in \mathcal{L}} T(\{u \in L^2(\mathbb{G}) \mid \|u\|_{L^2(\mathbb{G})} \leq 1\})$ is relatively compact in $L^2(\mathbb{G})$. It is a rather deep fact [1, 34] that \mathcal{L} is a relatively compact subset of $\mathbb{K}[L^2(\mathbb{G})]$ with respect to the operator norm if and only if both \mathcal{L} and $\mathcal{L}^* := \{T^* \mid T \in \mathcal{L}\}$ are collectively compact. This and the point 2 lead to the desired conclusion. \square

REMARK 5.6. In Subsection 7.3 we are going to introduce multiplication operators $\text{Mult}(f)$ and left convolution operators $\text{Conv}_L(g)$. Completing Theorem 5.5, one can easily prove that a bounded subset Δ of $L^2(\mathbb{G})$ is relatively compact if and only if for every $\epsilon > 0$ there exist $f, g \in \mathcal{C}_c(\mathbb{G})$ such that

$$\sup_{u \in \Delta} (\|\text{Mult}(f)u - u\|_{L^2(\mathbb{G})} + \|\text{Conv}_L(g)u - u\|_{L^2(\mathbb{G})}) \leq \epsilon.$$

Such type of results, in a much more general setting, have been proved in [16]. They are not depending on the existence of a pseudo-differential calculus. On the other hand, essentially by the same proof, we could assign f, g to the Bruhat space $\mathcal{D}(\mathbb{G})$, which is not covered by [16].

6 RIGHT AND LEFT QUANTIZATIONS

Our construction of the pseudo-differential calculus Op^τ started from a concrete expression (17) for the Weyl system $W^\tau \equiv W_R^\tau$; we set for $x, y \in \mathbb{G}$, $\xi \in \widehat{\mathbb{G}}$ and $\Theta \in L^2(\mathbb{G}, \mathcal{H}_\xi)$

$$[W_R^\tau(\xi, x)\Theta](y) := \xi[y(\tau x)^{-1}]^* [\Theta(yx^{-1})]. \tag{64}$$

The extra index R hints to the fact that right translations are used in (64). Building on (64) we constructed a “right” pseudo-differential calculus $\text{Op}^\tau \equiv \text{Op}_R^\tau$ given on suitable symbols a by

$$[\text{Op}_R^\tau(a)u](x) = \int_{\mathbb{G}} \left(\int_{\widehat{\mathbb{G}}} \text{Tr}_\xi \left[\xi(y^{-1}x)a(x\tau(y^{-1}x)^{-1}, \xi) \right] d\widehat{m}(\xi) \right) u(y) dm(y). \tag{65}$$

We recall that one gets integral operators, i.e. one can write

$$\text{Op}_R^\tau = \text{Int} \circ \text{Ker}_R^\tau = \text{Int} \circ \text{CV}_R^\tau \circ (\text{id} \otimes \mathcal{F}^{-1}), \quad (66)$$

in terms of a partial Fourier transformation and the change of variables

$$\text{cv}^\tau \equiv \text{cv}_R^\tau : \mathbb{G} \times \mathbb{G} \rightarrow \mathbb{G} \times \mathbb{G}, \quad \text{cv}_R^\tau(x, y) := (x\tau(y^{-1}x)^{-1}, y^{-1}x). \quad (67)$$

Besides (64) there is (at least) another version of a Weyl system, involving translations to the left, given by

$$[W_L^\tau(\xi, x)\Theta](y) := \xi[(\tau x)^{-1}y]^* [\Theta(x^{-1}y)]. \quad (68)$$

Using it, by arguments similar to those of Subsections 3.1 and 3.2, one gets a *left pseudo-differential calculus*

$$[\text{Op}_L^\tau(a)u](x) = \int_{\mathbb{G}} \left(\int_{\widehat{\mathbb{G}}} \text{Tr}_\xi [\xi(xy^{-1})a(\tau(xy^{-1})^{-1}x, \xi)] d\widehat{m}(\xi) \right) u(y) dm(y), \quad (69)$$

which can also be written as

$$\text{Op}_L^\tau = \text{Int} \circ \text{Ker}_L^\tau = \text{Int} \circ \text{CV}_L^\tau \circ (\text{id} \otimes \mathcal{F}^{-1}), \quad (70)$$

in terms of a different change of variables

$$\text{cv}_L^\tau : \mathbb{G} \times \mathbb{G} \rightarrow \mathbb{G} \times \mathbb{G}, \quad \text{cv}_L^\tau(x, y) := (\tau(xy^{-1})^{-1}x, xy^{-1}).$$

Once again we get a unitary map $\text{Op}_L^\tau : \mathcal{B}^2(\Gamma) \rightarrow \mathbb{B}^2[L^2(\mathbb{G})]$ and all the results obtained above have, mutatis mutandis, analogous versions in the left calculus. In particular, Op_L^τ also have extensions to distribution spaces connected to the Bruhat space, as in Section 5.

REMARK 6.1. The parameter τ is connected to ordering issues even in the standard case $\mathbb{G} = \mathbb{R}^n$. In general, another ordering problem comes from the non-commutativity of the group \mathbb{G} and the non-commutativity of $\mathbb{B}(\mathcal{H}_\xi)$ for each irreducible representation $\xi : \mathbb{G} \rightarrow \mathbb{B}(\mathcal{H}_\xi)$. It is to this problem that we refer now. It is worth writing again the two quantizations for the simple case $\tau(x) = e$:

$$[\text{Op}_R(a)u](x) = \int_{\widehat{\mathbb{G}}} \text{Tr}_\xi [\xi(x)a(x, \xi)\widehat{u}(\xi)] d\widehat{m}(\xi), \quad (71)$$

following easily from (65), and

$$[\text{Op}_L(a)u](x) = \int_{\widehat{\mathbb{G}}} \text{Tr}_\xi [\xi(x)\widehat{u}(\xi)a(x, \xi)] d\widehat{m}(\xi), \quad (72)$$

following from (69). The two expressions coincide if \mathbb{G} is Abelian, since then each \mathcal{H}_ξ will be 1-dimensional. We will say more on this in Subsections 7.2 and 7.3.

REMARK 6.2. We note that the choices of left or right quantizations as in (71) and (72) may lead to parallel equivalent (as in the case of compact Lie groups) or non-equivalent (as in the case of graded Lie groups) theories. In the main body of the article we adopted the conventions leading to Op_R , mainly to recover the compact [39] and the nilpotent [18] case (both already exposed in book form) as particular cases. But the left quantization is connected to the formalism of crossed product C^* -algebras, as will be seen subsequently, and this can be very useful for certain applications. We also note that there may be no canonical way of calling the quantization “left” or “right”. Thus, the terminology was opposite in [39, Remark 10.4.13], although it was natural in that context. In the present paper, the adopted terminology is related to the group actions in (64) and (68), respectively. It also seems natural from the formula $\text{Op}_L^\tau = \text{Sch}_L^\tau \circ (\text{id} \otimes \mathcal{F})^{-1}$ obtained later in (83), where $\text{Sch}_L^\tau = r \rtimes^\tau L$ appears in (81) as the integrated form of the Schrödinger representation, with the left-regular group action L given by $[L(y)v](x) = v(y^{-1}x)$, and multiplication $r(f)v = fv$. In any case, we refer to Section 7.2 for further interpretation of the quantization formulae in terms of the appearing Schrödinger representations.

REMARK 6.3. The measurable map $\sigma_L : G \rightarrow G$ is called a *symmetry function with respect to the left quantization* if one has $\text{Op}_L^{\sigma_L}(a)^* = \text{Op}_L^{\sigma_L}(a^*)$ for every $a \in \mathcal{B}^2(\Gamma)$. As in Proposition 4.1, one shows that $\text{Op}_L^\tau(a)^* = \text{Op}_L^{\widehat{\tau}}(a^*)$, for $\widehat{\tau}(x) := x\tau(x^{-1})$, so σ_L must satisfy this time $\sigma_L(x) = x\sigma_L(x^{-1})$ (almost everywhere). An analog of Proposition 4.3 also holds. For central extensions, instead of (55), one must set

$$\sigma_L(h, n) := \left(\sigma_H(h), \sigma_N[\varpi(h, \sigma_H(h^{-1}))n] \right).$$

Note that, in the Lie exponential case, the function (58) is a symmetry function simultaneously to the left and to the right, cf. (59).

Let a be an element of $\mathcal{B}^2(\Gamma)$ and $\tau, \tau' : G \rightarrow G$ two measurable functions. One has

$$\text{Op}_L^{\tau'}(a) = \text{Int}[\text{Ker}_{L,a}^{\tau'}] \quad \text{and} \quad \text{Op}_R^\tau(a) = \text{Int}[\text{Ker}_{R,a}^\tau].$$

It is easy to deduce from (66) and (70) the connection between the left and the right kernel:

$$\text{Ker}_{L,a}^{\tau'} = \text{CV}_L^{\tau'} (\text{CV}_R^\tau)^{-1} [\text{Ker}_{R,a}^\tau],$$

meaning that one has $\text{Ker}_{L,a}^{\tau'} = \text{Ker}_{R,a}^\tau \circ \text{cv}_{R,L}^{\tau,\tau'}$, where $\text{cv}_{R,L}^{\tau,\tau'} := (\text{cv}_R^\tau)^{-1} \circ \text{cv}_L^{\tau'}$ is explicitly

$$\text{cv}_{R,L}^{\tau,\tau'}(x, y) = (\tau'(xy^{-1})^{-1}x\tau(xy^{-1}), \tau'(xy^{-1})^{-1}x\tau(xy^{-1})yx^{-1}).$$

This relation looks frightening, but particular cases are nicer. Setting $\tau(x) = e = \tau'(x)$ for instance, one gets $\text{cv}_{R,L}^{e,e}(x, y) = (x, xyx^{-1})$, while $\tau = \text{id} = \tau'$ leads to $\text{cv}_{R,L}^{\text{id},\text{id}}(x, y) = (yxy^{-1}, y)$.

Investigating when $\text{cv}_{R,L}^{\tau,\tau'} = \text{id}_{G \times G}$ holds (leading to $\text{Op}_L^{\tau'}(a) = \text{Op}_R^\tau(a)$ for every a), one could be disappointed. It comes out quickly that $xyx^{-1} = y$ for all x, y is a necessary condition, so the group G must be Abelian! Then $\tau = \tau'$ is the remaining condition.

7 THE C^* -ALGEBRAIC FORMALISM

In this section we describe a general formalism in terms of C^* -algebras that becomes useful as a background setting for pseudo-differential operators, in particular allowing working with operators with coefficients taking values in various Abelian C^* -algebras. Especially, an interpretation in terms of crossed product C^* -algebras become handy making use of C^* -dynamical systems and their covariant representations. We introduce an analogue of the Schrödinger representation and its appearance in τ -quantizations. Consequently, we investigate the role of multiplication and convolution operators in describing general families of pseudo-differential operators. The formalism is then used to investigate covariant families of pseudo-differential operators and establish several results concerning their spectra.

7.1 CROSSED PRODUCT C^* -ALGEBRAS

We change now the point of view and place the pseudo-differential calculus in the setting of C^* -algebras generated by actions of our group G on suitable function algebras. For a full general treatment of C^* -dynamical systems and their crossed products we refer to [36, 47].

DEFINITION 7.1. A C^* -dynamical system is a triple (\mathcal{A}, θ, G) where

- G is a locally compact group with Haar measure m ,
- \mathcal{A} is a C^* -algebra,
- $\theta : G \rightarrow \text{Aut}(\mathcal{A})$ is a strongly continuous action by automorphisms.

The third condition means that each $\theta_x : \mathcal{A} \rightarrow \mathcal{A}$ is a C^* -algebra isomorphism, the map $G \ni x \mapsto \theta_x(f) \in \mathcal{A}$ is continuous for every $f \in \mathcal{A}$ and one has $\theta_x \circ \theta_y = \theta_{xy}$ for all $x, y \in G$.

DEFINITION 7.2. 1. To a C^* -dynamical system (\mathcal{A}, θ, G) we associate the Banach $*$ -algebra structure on $L^1(G; \mathcal{A})$ (the space of all Bochner integrable functions $G \rightarrow \mathcal{A}$) given by

$$\begin{aligned} \|\Phi\|_{(1)} &:= \int_G \|\Phi(x)\|_{\mathcal{A}} \, dm(x), \\ (\Phi \diamond \Psi)(x) &:= \int_G \Phi(y) \theta_y[\Psi(y^{-1}x)] \, dm(y), \\ \Phi^\diamond(x) &:= \theta_x[\Phi(x^{-1})^*]. \end{aligned}$$

2. Then the crossed product C^* -algebra $\mathcal{A} \rtimes_\theta G := \text{Env}[L^1(G; \mathcal{A})]$ is the enveloping C^* -algebra of this Banach $*$ -algebra, i.e its completion in the universal norm

$$\|\Phi\|_{\text{univ}} := \sup_{\Pi} \|\Pi(\Phi)\|_{\mathbb{B}(\mathcal{H})},$$

where the supremum is taken over all the $*$ -representations $\Pi : L^1(G, \mathcal{A}) \rightarrow \mathbb{B}(\mathcal{H})$.

The Banach space $L^1(\mathbb{G}; \mathcal{A})$ can be identified with the projective tensor product $\mathcal{A} \overline{\otimes} L^1(\mathbb{G})$, and $\mathcal{C}_c(\mathbb{G}; \mathcal{A})$, the space of all \mathcal{A} -valued continuous compactly supported function on \mathbb{G} , is a dense $*$ -subalgebra of $L^1(\mathbb{G}, \mathcal{A})$ and of $\mathcal{A} \rtimes_{\theta} \mathbb{G}$ (cf. [47]).

DEFINITION 7.3. Let $(\mathcal{A}, \theta, \mathbb{G})$ be a C^* -dynamical system. A covariant representation is a triple (r, T, \mathcal{H}) where

- \mathcal{H} is a Hilbert space,
- $T : \mathbb{G} \rightarrow \mathbb{U}(\mathcal{H})$ is a (strongly continuous) unitary representation,
- $r : \mathcal{A} \rightarrow \mathbb{B}(\mathcal{H})$ is a $*$ -representation,
- $T(x)r(f)T(x)^* = r[\theta_x(f)]$, for every $f \in \mathcal{A}$ and $x \in \mathbb{G}$.

It is known that there is a one-to-one correspondence between

- covariant representations of the C^* -dynamical system $(\mathcal{A}, \theta, \mathbb{G})$,
- non-degenerate $*$ -representations of the crossed product $\mathcal{A} \rtimes_{\theta} \mathbb{G}$.

We only need the direct correspondence: *The integrated form* of the covariant representation (r, T, \mathcal{H}) is uniquely defined by $r \rtimes T : L^1(\mathbb{G}; \mathcal{A}) \rightarrow \mathbb{B}(\mathcal{H})$, with

$$(r \rtimes T)(\Phi) := \int_{\mathbb{G}} r[\Phi(x)]T(x)dm(x);$$

then the unique continuous extension $r \rtimes T : \mathcal{A} \rtimes_{\theta} \mathbb{G} \rightarrow \mathbb{B}(\mathcal{H})$ is justified by the universal property of the envelopping C^* -algebra.

This is the formalism one usually encounters in the references treating crossed products [36, 47]; in terms of pseudo-differential operators this would only cover the case $\tau(\cdot) = e$, i.e. the Kohn-Nirenberg type quantization. To treat the general case of a measurable map $\tau : \mathbb{G} \rightarrow \mathbb{G}$, one needs the modifications

$$(\Phi \diamond^{\tau} \Psi)(x) := \int_{\mathbb{G}} \theta_{\tau(x)^{-1}\tau(y)}[\Phi(y)] \theta_{\tau(x)^{-1}y\tau(y^{-1}x)}[\Psi(y^{-1}x)] dm(y), \tag{73}$$

$$\Phi \diamond^{\tau}(x) := \theta_{\tau(x)^{-1}x\tau(x^{-1})}[\Phi(x^{-1})]^*, \tag{74}$$

in the $*$ -algebra structure of $L^1(\mathbb{G}; \mathcal{A})$ and the next modification of the integrated form of a covariant representation (r, T, \mathcal{H}) as

$$(r \rtimes^{\tau} T)(\Phi) := \int_{\mathbb{G}} r[\theta_{\tau(x)}(\Phi(x))] T(x) dm(x). \tag{75}$$

By taking enveloping C^* -algebras, one gets a family $\{\mathcal{A} \rtimes_{\theta}^{\tau} \mathbb{G}\}_{\tau}$ of C^* -algebras indexed by all the measurable mappings $\tau : \mathbb{G} \rightarrow \mathbb{G}$.

REMARK 7.4. In fact all these C^* -algebras are isomorphic: $\mathcal{A} \rtimes_{\theta}^{\tau'} \mathbb{G} \xrightarrow{\nu_{\tau\tau'}} \mathcal{A} \rtimes_{\theta}^{\tau} \mathbb{G}$ is an isomorphism, uniquely determined by its action on $L^1(\mathbb{G}; \mathcal{A})$ defined as

$$(\nu_{\tau\tau'}\Phi)(x) := \theta_{\tau(x)^{-1}\tau'(x)}[\Phi(x)].$$

The family of isomorphisms satisfy

$$\nu_{\tau_1\tau_2} \circ \nu_{\tau_2\tau_3} = \nu_{\tau_1\tau_3}, \quad \nu_{\tau\tau'}^{-1} = \nu_{\tau'\tau},$$

and the relation $r \rtimes^{\tau'} T = (r \rtimes^{\tau} T) \circ \nu_{\tau\tau'}$ is easy to check. An important ingredient is the fact that $\nu_{\tau\tau'}$ leaves the space $L^1(\mathbb{G}; \mathcal{A})$ invariant (actually it is an isometry).

REMARK 7.5. For further use, let us also examine $*$ -morphisms in the setting of crossed products (cf [47]). Assume that $(\mathcal{A}, \theta, \mathbb{G})$ and $(\mathcal{A}', \theta', \mathbb{G})$ are C^* -dynamical systems and $\gamma : \mathcal{A} \rightarrow \mathcal{A}'$ is an equivariant $*$ -morphism, i.e. a $*$ -morphism satisfying

$$\gamma \circ \theta_x = \theta'_x \circ \gamma, \quad \forall x \in \mathbb{G}. \tag{76}$$

One defines

$$\gamma^{\times} : L^1(\mathbb{G}; \mathcal{A}) \rightarrow L^1(\mathbb{G}; \mathcal{A}'), \quad [\gamma^{\times}(\Phi)](x) := \gamma[\Phi(x)]. \tag{77}$$

It is easy to check that γ^{\times} is a $*$ -morphism of the two Banach $*$ -algebra structures and thus it extends to a $*$ -morphism $\gamma^{\times} : \mathcal{A} \rtimes_{\theta}^{\tau} \mathbb{G} \rightarrow \mathcal{A}' \rtimes_{\theta'}^{\tau} \mathbb{G}$. If γ is injective, γ^{\times} is also injective.

7.2 THE SCHRÖDINGER REPRESENTATION AND τ -QUANTIZATIONS

It will be convenient to assume that \mathcal{A} is a C^* -subalgebra of $\mathcal{LUC}_b(\mathbb{G})$ (bounded, left uniformly continuous functions on \mathbb{G}) invariant under left translations and that $[\theta_y(f)](x) := f(y^{-1}x)$. The maximal choice $\mathcal{A} = \mathcal{LUC}_b(\mathbb{G})$ is very convenient, but studying “pseudo-differential operators with coefficients of a certain type, modelled by \mathcal{A} ”, can sometimes be useful. Applications and extensions will appear elsewhere.

For \mathcal{A} -valued functions Φ defined on \mathbb{G} and for elements x, q of the group, we are going to use notations as $[\Phi(x)](q) = \Phi(q, x)$, interpreting Φ as a function of two variables. The strange order of these variables is convenient to make the connection with previous sections. We can also understand the action as given by $\theta_y(\Phi(x))(q) = \Phi(y^{-1}q, x)$.

Thus on the dense subset $L^1(\mathbb{G}; \mathcal{A}) \subset \mathcal{A} \rtimes_{\theta}^{\tau} \mathbb{G}$ the composition law (73) becomes more explicit

$$(\Phi \diamond^{\tau} \Psi)(q, x) := \int_{\mathbb{G}} \Phi(\tau(y)^{-1}\tau(x)q, y) \Psi(\tau(y^{-1}x)^{-1}y^{-1}\tau(x)q, y^{-1}x) dm(y), \tag{78}$$

while the involution (74) becomes

$$\Phi \diamond^{\tau}(q, x) := \overline{\Phi(\tau(x^{-1})^{-1}x^{-1}\tau(x)q, x^{-1})}. \tag{79}$$

REMARK 7.6. If G admits a symmetric quantization to the left and $\tau \equiv \sigma_L$ is a symmetry function to the left, as in Remark 6.3, the involution boils down to $\Phi^{\diamond \sigma_L}(q, x) := \overline{\Phi(q, x^{-1})}$.

In the situation described above, we always have a natural covariant representation (r, L, \mathcal{H}) , called *the Schrödinger representation*, given in $\mathcal{H} := L^2(G)$ by

$$[L(y)v](x) := v(y^{-1}x), \quad r(f)v := fv; \tag{80}$$

thus $L(y)$ is the unitary left-translation by y^{-1} in $L^2(G)$ and $r(f)$ is just the operator of multiplication by the bounded function f . The corresponding (modified) integrated form

$$\text{Sch}_L^\tau := r \rtimes^\tau L, \tag{81}$$

computed as in (75), is given for $\Phi \in L^1(G; \mathcal{A})$ and $v \in L^2(G)$ by the formula

$$\begin{aligned} [\text{Sch}_L^\tau(\Phi)v](x) &= \int_G \Phi(\tau(z)^{-1}x, z)v(z^{-1}x) \, dm(z) \\ &= \int_G \Phi(\tau(xy^{-1})^{-1}x, xy^{-1})v(y) \, dm(y). \end{aligned} \tag{82}$$

The good surprise is that if we compose Sch_L^τ with the inverse of the partial Fourier transform one finds again, at least formally, the left pseudo-differential representation (69) and (70):

$$\text{Op}_L^\tau = \text{Sch}_L^\tau \circ (\text{id} \otimes \mathcal{F})^{-1} = \text{Int} \circ \text{CV}_L^\tau \circ (\text{id} \otimes \mathcal{F}^{-1}). \tag{83}$$

It is worth comparing this expression of Sch_L^τ in (82) with (47).

To extend the meaning of (83) beyond the L^2 -theory and to take full advantage of the C^* -algebraic formalism, one needs to be more careful. Recall that the Fourier transform defines an injective linear contraction $\mathcal{F} : L^1(G) \rightarrow \mathcal{B}(\widehat{G})$. We already mentioned that $L^1(G; \mathcal{A})$ can be identified with the completed projective tensor product $\mathcal{A} \overline{\otimes} L^1(G)$. Then, by [46, Ex.43.2], one gets a linear continuous injection

$$\text{id}_{\mathcal{A}} \overline{\otimes} \mathcal{F} : \mathcal{A} \overline{\otimes} L^1(G) \rightarrow \mathcal{A} \overline{\otimes} \mathcal{B}(\widehat{G})$$

and we endow the image $(\text{id}_{\mathcal{A}} \overline{\otimes} \mathcal{F})[\mathcal{A} \overline{\otimes} L^1(G)]$ with the Banach $*$ -algebra structure transported from $L^1(G; \mathcal{A}) \cong \mathcal{A} \overline{\otimes} L^1(G)$ through $\text{id}_{\mathcal{A}} \overline{\otimes} \mathcal{F}$.

Let us denote by $\mathcal{C}_{\mathcal{A}}^\tau$ the envelopping C^* -algebra $\mathcal{A} \rtimes_\theta^\tau G$ of the Banach $*$ -algebra $L^1(G; \mathcal{A})$ (with the τ -structure indicated above). Similarly, we denote by $\mathfrak{B}_{\mathcal{A}}^\tau$ the envelopping C^* -algebra of the Banach $*$ -algebra $(\text{id}_{\mathcal{A}} \overline{\otimes} \mathcal{F})[\mathcal{A} \overline{\otimes} L^1(G)]$. By the universal property of the envelopping functor, $\text{id}_{\mathcal{A}} \overline{\otimes} \mathcal{F}$ extends to an isomorphism $\mathfrak{F}_{\mathcal{A}} : \mathcal{C}_{\mathcal{A}}^\tau \rightarrow \mathfrak{B}_{\mathcal{A}}^\tau$.

Now $\text{Op}_L^\tau := \text{Sch}_L^\tau \circ \mathfrak{F}_{\mathcal{A}}^{-1}$ defines a $*$ -representation of the C^* -algebra $\mathfrak{B}_{\mathcal{A}}^\tau$ in the Hilbert space $L^2(G)$, which is compatible with (83) when both expressions make sense. It seems pointless to use different notations for the two basically identical quantizations, one defined in the C^* -algebraical setting, starting from the Schrödinger

representation, and the other introduced in Section 6, built on the family (68) of unitary operators. The difference is merely a question of domains, but each of them can be still extended or restricted (at least for particular classes of groups G), being constructed on the versatile operations Int , CV^τ , \mathcal{F}^{-1} .

REMARK 7.7. Starting from (78) and (79), one also considers the transported composition

$$a\#^\tau b := \mathfrak{F}_A [(\mathfrak{F}_A^{-1}a) \diamond^\tau (\mathfrak{F}_A^{-1}b)]$$

defined to satisfy $\text{Op}_L^\tau(a\#^\tau b) = \text{Op}_L^\tau(a) \text{Op}_L^\tau(b)$ as well as the involution

$$a\#^\tau := \mathfrak{F}_A [(\mathfrak{F}_A^{-1}a) \diamond^\tau]$$

verifying $\text{Op}_L^\tau(a\#^\tau) = \text{Op}_L^\tau(a)^*$.

REMARK 7.8. As a consequence of Remark 7.4 (see also Remark 38) and of the properties of envelopping C^* -algebras, there are isomorphisms $\mu_{\tau, \tau'} : \mathfrak{B}_A^\tau \rightarrow \mathfrak{B}_A^{\tau'}$ leaving $(\text{id}_A \otimes \mathcal{F}) [L^1(G; \mathcal{A})]$ invariant and satisfying

$$\text{Op}_L^\tau = \text{Op}_L^{\tau'} \circ \mu_{\tau, \tau'}, \quad \tau, \tau' : G \rightarrow G.$$

Therefore, the C^* -subalgebra

$$\mathfrak{D}_A := \text{Op}_L^\tau(\mathfrak{B}_A^\tau) = \text{Sch}_L^\tau(\mathfrak{C}_A^\tau) \subset \mathbb{B}[L^2(G)] \tag{84}$$

is τ -independent. It could be called *the C^* -algebra of left global pseudo-differential operators with coefficients in \mathcal{A} on the admissible group G* .

PROPOSITION 7.9. *The C^* -algebra \mathfrak{D}_A is isomorphic to the reduced crossed product $(\mathcal{A} \rtimes_\theta^\tau G)_{\text{red}}$. If G is amenable, the representation $\text{Op}_L^\tau : \mathfrak{B}_A^\tau \rightarrow \mathfrak{D}_A \subset \mathbb{B}[L^2(G)]$ is faithful.*

Proof. Of course, it is enough to work with one of the mappings $\tau : G \rightarrow G$, for instance for $\tau(x) = e$. Since \mathfrak{F}_A is an isomorphism, it suffices to study the $*$ -representation $\text{Sch}_L := r \rtimes L : \mathfrak{C}_A := \mathcal{A} \rtimes_\theta G \rightarrow \mathbb{B}[L^2(G)]$ and its range. For this we are going to recall the left regular $*$ -representation Left of the crossed product in the Hilbert space $\mathcal{H} := L^2(G \times G) \cong L^2(G; L^2(G))$ and show that Sch_L and Left are "unitarily equivalent up to multiplicity". The range of Left in $\mathbb{B}[L^2(G \times G)]$ is, by definition, *the reduced crossed product $(\mathcal{A} \rtimes_\theta^\tau G)_{\text{red}}$* . Since Left is injective if (and only if) G is amenable [47], this would finish our proof.

The $*$ -representation $\text{Left} = r' \rtimes L'$ is the integrated form of the covariant representation (r', L', \mathcal{H}) given by

$$[r'(f)\nu](q, x) := f(xq)\nu(q, x), \quad x, q \in G, f \in \mathcal{A}, \nu \in L^2(G \times G),$$

$$[L'(y)\nu](q, x) := \nu(q, y^{-1}x), \quad x, y, q \in G, \nu \in L^2(G \times G).$$

Then the unitary operator $W : L^2(G \times G) \rightarrow L^2(G \times G)$ defined by

$$(W\nu)(q, x) := \nu(q, xq)$$

satisfies for all $f \in \mathcal{A}$ and $y \in G$

$$W^*r'(f)W = \text{id}_{L^2(G)} \otimes r(f), \quad W^*L'(y)W = \text{id}_{L^2(G)} \otimes L(y),$$

which readily implies the unitary equivalence at the level of *-representations

$$W^*\text{Left}(\Phi)W = \text{id}_{L^2(G)} \otimes \text{Sch}_L^\tau(\Phi), \quad \forall \Phi \in \mathcal{C}_\mathcal{A},$$

and we are done. □

REMARK 7.10. Explicit descriptions of the C^* -algebras $\mathfrak{B}_\mathcal{A}^\tau$ are difficult to achieve. Even for $G = \mathbb{R}^n$ some of the elements of $\mathfrak{B}_\mathcal{A}^\tau$ are not ordinary functions on $(\mathbb{R}^n)^* \times \mathbb{R}^n$.

REMARK 7.11. Let us denote by $\mathcal{C}_0(G)$ the C^* -algebra of all the continuous complex-valued functions on G which converge to 0 at infinity (they are arbitrarily small outside sufficiently large compact subsets). It is well-known [47] that the Schrödinger *-representation sends $\mathcal{C}_0(G) \rtimes_\theta G$ onto $\mathbb{K}[L^2(G)] \subset \mathbb{B}[L^2(G)]$; it is an isomorphism between $\mathcal{C}_0(G) \rtimes_\theta G$ and $\mathbb{K}[L^2(G)]$ if and only if G is amenable. This provides classes of compact global pseudo-differential operators:

$$\mathbb{K}[L^2(G)] = \text{Op}_L^\tau(\mathfrak{B}_{\mathcal{C}_0(G)}^\tau) = \text{Sch}_L^\tau[\mathcal{C}_0(G) \rtimes_\theta G], \tag{85}$$

giving a characterisation of compact operators.

REMARK 7.12. One sees that, in the process of construction of the crossed product C^* -algebra, taking the completion in the envelopping norm supplies a lot of interesting new elements. One has

$$L^2(G \times G) \cong \mathcal{B}^2(\widehat{G} \times G) \cong \mathbb{B}^2[L^2(G)] \subset \mathbb{K}[L^2(G)] \cong \mathcal{C}_0(G) \rtimes_\theta G = \overline{L^1(G; \mathcal{C}_0(G))}$$

(the last expression involves the closure in the enveloping norm), while $L^2(G \times G)$ and $L^1(G; \mathcal{C}_0(G))$ are incomparable as soon as G is an infinite group. There are many Hilbert-Schmidt operators whose symbols are not partial Fourier transforms of elements from the class $L^1(G; \mathcal{C}_0(G))$.

REMARK 7.13. Recall that $\mathbb{K}[L^2(G)] = \text{Op}_L^\tau(\mathfrak{B}_{\mathcal{C}_0(G)}^\tau)$ is an irreducible family of operators in $L^2(G)$. So if \mathfrak{B} is a space of symbols containing $(\text{id} \otimes \mathcal{F})[\mathcal{C}_c(G; \mathcal{C}_0(G))]$ or $\mathcal{B}^2(\widehat{\Gamma})$, and if $\text{Op}_L^\tau(b)$ makes sense for every $b \in \mathfrak{B}$, then $\text{Op}_L^\tau(\mathfrak{B})$ is irreducible. This happens, for instance, if $\mathfrak{B} = \mathfrak{B}_\mathcal{A}^\tau$ and $\mathcal{C}_0(G) \subset \mathcal{A}$. In many other situations $\text{Op}_L^\tau(\mathfrak{B})$ could be reducible. Let us set $[R(z)u](x) := u(xz)$; a simple computation shows that $R(z)\text{Op}_L^\tau(b)R(z^{-1}) = \text{Op}_L^\tau(b_z)$, where $b_z(x, \xi) := b(xz, \xi)$. Thus, if b does not depend on the first variable, $\text{Op}_L^\tau(b)$ commutes with the right translations and irreducibility is lost for $\mathcal{A} := \mathbb{C}$. The same happens if \mathcal{A} is defined through a periodicity (invariance) condition with respect to some nontrivial closed subgroup of G .

7.3 MULTIPLICATION AND CONVOLUTION OPERATORS

We reconsider the Schrödinger covariant representation $(r, L, L^2(\mathbb{G}))$ of the C^* -dynamical system $(\mathcal{A}, \theta, \mathbb{G})$ where, as before, $\mathcal{A} \subset \mathcal{LUC}_b(\mathbb{G})$ is invariant under left translations.

Let us define $\text{Conv}_L : L^1(\mathbb{G}) \rightarrow \mathbb{B}[L^2(\mathbb{G})]$ by the formula (interpreted in weak sense)

$$\text{Conv}_L(f) := \int_{\mathbb{G}} f(y)L(y)dm(y) = \int_{\mathbb{G}} f(y)W_L^T(\mathbf{1}, y)(y)dm(y). \quad (86)$$

Clearly $\text{Conv}_L(f)$ is the operator of left-convolution with f : one has $\text{Conv}_L(f)v = f * v$ for every $v \in L^2(\mathbb{G})$. It is easy to check the right-invariance:

$$\text{Conv}_L(f)R(x) = R(x)\text{Conv}_L(f), \quad \forall f \in L^1(\mathbb{G}), x \in \mathbb{G}. \quad (87)$$

The map Conv_L extends to a C^* -epimorphism from the group C^* -algebra $C^*(\mathbb{G}) = \mathbb{C} \rtimes_{\theta} \mathbb{G}$ to the reduced group C^* -algebra $C_{\text{red}}^*(\mathbb{G}) = (\mathbb{C} \rtimes_{\theta} \mathbb{G})_{\text{red}} \subset \mathbb{B}[L^2(\mathbb{G})]$, which is an isomorphism if and only if \mathbb{G} is amenable [47].

Of course, the family of right-convolution operators $\{u \mapsto \text{Conv}_R(f)u := u * f \mid f \in L^1(\mathbb{G})\}$ is also available and it has similar properties. The analog of (86) is in this case

$$\text{Conv}_R(f) := \int_{\mathbb{G}} \check{f}(y)R(y)dm(y), \quad \text{with } \check{f}(y) := f(y^{-1}).$$

Note the commutativity property

$$\text{Conv}_L(f)\text{Conv}_R(f') = \text{Conv}_R(f')\text{Conv}_L(f)$$

as well as the identities

$$\text{Conv}_L(f)\text{Conv}_L(f') = \text{Conv}_L(f * f'), \quad \text{Conv}_R(f)\text{Conv}_R(f') = \text{Conv}_R(f' * f).$$

REMARK 7.14. More generally, one can also define $\text{Conv}_L(\mu)$ and $\text{Conv}_R(\mu)$ for any bounded complex Radon measure $\mu \in M^1(\mathbb{G})$. Let us denote by $\mathfrak{L}(\mathbb{G}) := [L(\mathbb{G})]''$ the left von Neumann algebra of \mathbb{G} and by $\mathfrak{R}(\mathbb{G}) := [R(\mathbb{G})]''$ the right von Neumann algebra of \mathbb{G} . One has $\text{Conv}_L[M^1(\mathbb{G})] \subset \mathfrak{L}(\mathbb{G})$ and $\text{Conv}_R[M^1(\mathbb{G})] \subset \mathfrak{R}(\mathbb{G})$.

It is easy to check that

$$\mathcal{F} \circ \text{Conv}_L(f) \circ \mathcal{F}^{-1} = \text{Dec}_R(\mathcal{F}f) \quad \text{and} \quad \mathcal{F} \circ \text{Conv}_R(f) \circ \mathcal{F}^{-1} = \text{Dec}_L(\mathcal{F}f),$$

where

$$\text{Dec}_R(\mathcal{F}f), \text{Dec}_L(\mathcal{F}f) \in \mathcal{B}(\widehat{\mathbb{G}}) := \int_{\widehat{\mathbb{G}}}^{\oplus} \mathbb{B}(\mathcal{H}_{\xi}) d\widehat{m}(\xi) \subset \mathbb{B}[\mathcal{B}^2(\widehat{\mathbb{G}})]$$

are decomposable (multiplication) operators defined for every $\varphi \in \mathcal{B}^2(\widehat{\mathbb{G}})$ by

$$[\text{Dec}_R(\mathcal{F}f)\varphi](\xi) := \varphi(\xi)(\mathcal{F}f)(\xi), \quad [\text{Dec}_L(\mathcal{F}f)\varphi](\xi) := (\mathcal{F}f)(\xi)\varphi(\xi).$$

We want to compute $\text{Op}_L^\tau(g \otimes \beta) = \text{Sch}_L^\tau[g \otimes (\mathcal{F}^{-1}\beta)]$, where g is some bounded uniformly continuous function on G and the inverse Fourier transform of β belongs to $L^1(G)$. Of course we set $(g \otimes \beta)(x, \xi) := g(x)\beta(\xi) \in \mathbb{B}(\mathcal{H}_\xi)$. One gets the formula

$$([\text{Op}_L^\tau(g \otimes \beta)] u)(x) = \int_G g[\tau(xy^{-1})^{-1}x](\mathcal{F}^{-1}\beta)(xy^{-1})u(y) dm(y),$$

which is not very inspiring for general τ . But using the notation $\text{Mult}(g) := r(g)$ (a multiplication operator in $L^2(G)$ given in (80)), one gets the particular cases, for $\text{Op}_L \equiv \text{Op}_L^\varepsilon$:

$$\begin{aligned} ([\text{Op}_L(g \otimes \beta)] u)(x) &= g(x) \int_G (\mathcal{F}^{-1}\beta)(xy^{-1})u(y) dm(y) \\ &= g(x) \int_G (\mathcal{F}^{-1}\beta)(z)u(z^{-1}x) dm(z), \end{aligned}$$

which can be rewritten

$$\text{Op}_L(g \otimes \beta) = \text{Mult}(g) \text{Conv}_L(\mathcal{F}^{-1}\beta), \tag{88}$$

and

$$\begin{aligned} ([\text{Op}_L^{\text{id}}(g \otimes \beta)] u)(x) &= \int_G (\mathcal{F}^{-1}\beta)(xy^{-1})g(y)u(y) dm(y) \\ &= \int_G (\mathcal{F}^{-1}\beta)(z)(gu)(z^{-1}x) dm(y), \end{aligned}$$

i.e.

$$\text{Op}_L^{\text{id}}(g \otimes \beta) = \text{Conv}_L(\mathcal{F}^{-1}\beta) \text{Mult}(g). \tag{89}$$

Thus in the quantization $\text{Op}_L \equiv \text{Op}_L^\varepsilon$ the operators of multiplication stay at the left and those of left-convolution to the right and vice versa for the quantization Op_L^{id} .

REMARK 7.15. In both (88) and (89) left convolution operators appear. But using the right quantization Op_R^τ one gets

$$([\text{Op}_R^\tau(g \otimes \beta)] u)(x) = \int_G g[x\tau(y^{-1}x)^{-1}](\mathcal{F}^{-1}\beta)(y^{-1}x)u(y) dm(y),$$

with particular cases

$$\begin{aligned} \text{Op}_R(g \otimes \beta) &= \text{Mult}(g) \text{Conv}_R(\mathcal{F}^{-1}\beta), \\ \text{Op}_R^{\text{id}}(g \otimes \beta) &= \text{Conv}_R(\mathcal{F}^{-1}\beta) \text{Mult}(g), \end{aligned}$$

and this should be compared with (88) and (89).

REMARK 7.16. As mentioned in Remark 7.8, the represented C^* -algebra

$$\mathfrak{D}_A := \text{Sch}_L^\tau(\mathcal{A} \rtimes_\theta^\tau G) = \text{Op}_L^\tau[\mathfrak{F}_A(\mathcal{A} \rtimes_\theta^\tau G)] \subset \mathbb{B}[L^2(G)]$$

is independent of τ . Actually it coincides with the closed vector space spanned by products of the form $\text{Mult}(g) \text{Conv}_L(f)$ (respectively $\text{Conv}_L(f) \text{Mult}(g)$) with $g \in \mathcal{A}$ and, say, $f \in (L^1 \cap L^2)(G)$ (or even $f \in \mathcal{D}(G)$). So this closed vector space is automatically a C^* -algebra, although this is not clear at a first sight. The remote reason is the last axiom of Definition 7.3.

REMARK 7.17. In [11, 13], in the case of a compact Lie group G , precise characterisations of the convolution operators belonging to the Schatten-von Neumann classes $\mathbb{B}^p[L^2(G)]$ are given. The main result [11, Th. 3.7] holds, with the same proof, for arbitrary compact groups.

When G is not compact, the single compact convolution operator is $0 = \text{Op}_L^\tau(0) = \text{Conv}_L(0) = \text{Conv}_R(0)$. A way to see this is to recall Remark 7.11 and to note that the constant function $g = 1$ belongs to $\mathcal{C}_0(G)$ if and only if G is compact. Another, more direct, argument is as follows: If G is not compact then $R(x)$ converges weakly to 0 when $x \rightarrow \infty$. Multiplication to the left by a compact operator would improve this to strong convergence. But for $u \in L^2(G)$ and a compact $\text{Conv}_L(f)$ one has

$$\|\text{Conv}_L(f)u\|_{L^2(G)} = \|R(x)\text{Conv}_L(f)u\|_{L^2(G)} = \|\text{Conv}_L(f)R(x)u\|_{L^2(G)} \xrightarrow{x \rightarrow \infty} 0$$

and this implies $\text{Conv}_L(f) = 0$. Replacing $R(\cdot)$ by $L(\cdot)$, a similar argument shows that the single compact right convolution operator is the null operator.

7.4 COVARIANT FAMILIES OF PSEUDO-DIFFERENTIAL OPERATORS

An important ingredient in constructing the Schrödinger representation has been the fact that the C^* -algebra \mathcal{A} was an algebra of (bounded, uniformly continuous) functions on G . If \mathcal{A} is just an Abelian C^* -algebra endowed with the action ρ of our group G , by Gelfand theory, it is connected to a topological dynamical system (Ω, ϱ, G) . The locally compact space Ω is the Gelfand spectrum of \mathcal{A} and we have the G -equivariant isomorphism $\mathcal{A} \cong \mathcal{C}_0(\Omega)$ if the action ρ_x of $x \in G$ on $\mathcal{C}_0(\Omega)$ is given just by composition with $\varrho_{x^{-1}}$. In this section we are going to prove that to such a data one associates a covariant family of pseudo-differential calculi with operator-valued symbols. For convenient bundle sections h defined on $\Omega \times \widehat{G}$ one gets families $\{\text{Op}_{(\omega)}^\tau(h) \mid \omega \in \Omega\}$ of “usual” left pseudo-differential operators (the index L will be omitted). By covariance, modulo unitary equivalence, they are actually indexed by the orbits of the topological dynamical system, while their spectra are indexed by the quasi-orbits in Ω .

As before, the locally compact group G is supposed second countable, unimodular and type I, while $\tau : G \rightarrow G$ is measurable.

Since the Schrödinger covariant representation (80) no longer makes sense as it stands, we are going to construct for each point $\omega \in \Omega$ a covariant representation $(r_{(\omega)}, L, L^2(G))$ and then let the formalism act. One sets explicitly

$$[r_{(\omega)}(f)u](x) := f[\varrho_x(\omega)]u(x), \quad f \in \mathcal{C}_0(\Omega), u \in L^2(G), x \in G, \quad (90)$$

$$[L(y)u](x) := u(y^{-1}x), \quad u \in L^2(G), x, y \in G. \quad (91)$$

Proceeding as in Subsection 7.2, one constructs the integrated form $\text{Sch}^\tau_{(\omega)} := r_{(\omega)} \rtimes L$ associated to the covariant representation $(r_{(\omega)}, L, L^2(\mathbb{G}))$ and then sets

$$\text{Op}^\tau_{(\omega)} := \text{Sch}^\tau_{(\omega)} \circ \mathfrak{F}^{-1}_{\mathcal{C}_0(\Omega)}. \tag{92}$$

As in Subsection 7.2, the isomorphism $\mathfrak{F}_\Omega \equiv \mathfrak{F}_{\mathcal{C}_0(\Omega)}$ is the extension of the Banach $*$ -algebra monomorphism

$$\text{id}_{\mathcal{C}_0(\Omega)} \overline{\otimes} \mathcal{F} : L^1(\mathbb{G}; \mathcal{C}_0(\Omega)) \cong \mathcal{C}_0(\Omega) \overline{\otimes} L^1(\mathbb{G}) \rightarrow \mathcal{C}_0(\Omega) \overline{\otimes} \mathcal{B}(\mathbb{G})$$

to the enveloping C^* -algebra $\mathcal{C}_0(\Omega) \rtimes_\rho^\tau \mathbb{G}$; the fact that $\mathcal{A} = \mathcal{C}_0(\Omega)$ is more general as before is not important. Setting $\mathfrak{B}^\tau_\Omega \equiv \mathfrak{B}^\tau_{\mathcal{C}_0(\Omega)}$ for the enveloping C^* -algebra of $(\text{id}_{\mathcal{C}_0(\Omega)} \overline{\otimes} \mathcal{F}) [L^1(\mathbb{G}; \mathcal{C}_0(\Omega))]$ (with the transported structure), we have the isomorphism

$$\mathfrak{F}_\Omega : \mathcal{C}_0(\Omega) \rtimes_\rho^\tau \mathbb{G} \rightarrow \mathfrak{B}^\tau_\Omega.$$

One gets for every section

$$\{h(\omega, \xi) \in \mathbb{B}(\mathcal{H}_\xi) \mid \xi \in \widehat{\mathbb{G}}, \omega \in \Omega\}$$

from $(\text{id}_{\mathcal{C}_0(\Omega)} \overline{\otimes} \mathcal{F}) [L^1(\mathbb{G}; \mathcal{C}_0(\Omega))]$ a family of operators

$$\left\{ \text{Op}^\tau_{(\omega)}(h) = (r_{(\omega)} \rtimes L)(\mathfrak{F}_{\mathcal{C}_0(\Omega)} h) \in \mathbb{B}[L^2(\mathbb{G})] \mid \omega \in \Omega \right\}$$

given explicitly (but somewhat formally) by

$$\left[\text{Op}^\tau_{(\omega)}(h)u \right](x) = \int_{\mathbb{G}} \left(\int_{\widehat{\mathbb{G}}} \text{Tr}_\xi [\xi(xy^{-1})h(\varrho_{\tau(xy^{-1})^{-1}x}(\omega), \xi)] d\widehat{m}(\xi) \right) u(y) dm(y). \tag{93}$$

More generally, the family $\{ \text{Op}^\tau_{(\omega)}(h) \mid \omega \in \Omega \}$ makes sense for $h \in \mathfrak{B}^\tau_\Omega$, but it is no longer clear when the symbol h can still be interpreted as a function on $\Omega \times \widehat{\mathbb{G}}$.

PROPOSITION 7.18. *Let $h \in \mathfrak{B}^\tau_\Omega$. If ω, ω' belong to the same ϱ -orbit, then $\text{Op}^\tau_{(\omega)}(h)$ and $\text{Op}^\tau_{(\omega')}(h)$ are unitarily equivalent.*

Proof. The points ω, ω' are on the same orbit if and only if there exists $z \in \mathbb{G}$ such that $\omega' = \varrho_z(\omega)$. In terms of the unitary right translation $[R(z)u](\cdot) := u(\cdot z)$, the operatorial covariance relation

$$R(z)\text{Op}^\tau_{(\omega)}(h)R(z)^* = \text{Op}^\tau_{(\varrho_z(\omega))}(h) \tag{94}$$

follows by an easy but formal calculation relying on (93). This can be upgraded to a rigorous justification by a density argument, but it is better to argue as follows: Formula (94) for arbitrary $h \in \mathfrak{B}^\tau_\Omega$ is equivalent to

$$R(z)\text{Sch}^\tau_{(\omega)}(\Phi)R(z)^* = \text{Sch}^\tau_{(\varrho_z(\omega))}(\Phi), \quad \forall \Phi \in \mathcal{C}_0(\Omega) \rtimes_\rho^\tau \mathbb{G}.$$

Since $\text{Sch}_{(\omega')}^\tau$ is the integrated form of the covariant representation $(r_{(\omega')}, L, L^2(\mathbb{G}))$ indicated in (90) and (91), it is enough to prove

$$R(z)L(x)R(z)^* = L(x), \quad \forall x, z \in \mathbb{G}$$

and

$$R(z)r_{(\omega)}(f)R(z)^* = r_{(\varrho_z(\omega))}(f), \quad \forall z \in \mathbb{G}, f \in \mathcal{C}_0(\Omega).$$

The first one is trivial. The second one follows from

$$\begin{aligned} [R(z)r_{(\omega)}(f)R(z)^*u](x) &= [r_{(\omega)}(f)R(z^{-1})u](xz) \\ &= f[\varrho_{xz}(\omega)][R(z^{-1})u](xz) \\ &= f[\varrho_x(\varrho_z(\omega))]u(x) \\ &= [r_{(\varrho_z(\omega))}(f)u](x), \end{aligned}$$

completing the proof. \square

REMARK 7.19. In fact one has

$$\text{Op}_{(\omega)}^\tau(h) = \text{Op}_L^\tau(h_{(\omega)}), \quad \text{with } h_{(\omega)}(x, \xi) := h(\varrho_x(\omega), \xi). \quad (95)$$

This relation supplies another interpretation of the family $\{\text{Op}_{(\omega)}^\tau(h) \mid \omega \in \Omega\}$. We can see it as being obtained by applying the left quantization procedure Op_L^τ of the preceding sections to a family $\{h_{(\omega)} \mid \omega \in \Omega\}$ of symbols (classical observables) defined in $\mathbb{G} \times \widehat{\mathbb{G}}$, associated through the action ϱ to a single function h on $\Omega \times \widehat{\mathbb{G}}$. Note that this family satisfies *the covariance condition*

$$h_{\varrho_z(\omega)}(x, \xi) = h_{(\omega)}(xz, \xi), \quad x, z \in \mathbb{G}, \xi \in \widehat{\mathbb{G}}, \omega \in \Omega. \quad (96)$$

Using the reinterpretation (95), the unitary equivalence (94) can be reformulated only in terms of the quantization Op_L^τ as

$$R(z)\text{Op}_L^\tau(h_{(\omega)})R(z)^* = \text{Op}_L^\tau(h_{(\varrho_z(\omega))}),$$

which is easily proved directly using relation (96) if h is not too general.

We recall that a *quasi-orbit* for the action ϱ is the closure of an orbit. If $\overline{O_\omega} := \varrho_{\mathbb{G}}(\omega)$ is the orbit of the point $\omega \in \Omega$, we denote by $\mathbb{Q}_\omega := \overline{O_\omega} = \varrho_{\mathbb{G}}(\omega)$ the *quasi-orbit generated by ω* . As a preparation for Theorem 7.20, we decompose the correspondance $\Phi \mapsto \text{Sch}_\omega^\tau(\Phi)$ into several parts. The starting point is the chain

$$\mathcal{C}_0(\Omega) \xrightarrow{\gamma_\omega} \mathcal{C}_0(\mathbb{Q}_\omega) \xrightarrow{\beta_\omega} \mathcal{LUC}_u(\mathbb{G}),$$

involving the restriction $*$ -morphism

$$\gamma_\omega : \mathcal{C}_0(\Omega) \rightarrow \mathcal{C}_0(\mathbb{Q}_\omega), \quad \gamma_\omega(f) := f|_{\mathbb{Q}_\omega}$$

and the composition $*$ -morphism

$$\beta_\omega : \mathcal{C}_0(Q_\omega) \rightarrow \mathcal{LUC}_u(G), \quad [\beta_\omega(g)](x) := g[\varrho_x(\omega)].$$

Note that β_ω is injective, since $\varrho_G(\omega)$ is dense in Q_ω . Both these $*$ -morphisms are equivariant in the sense of Remark 7.5 if on $\mathcal{C}_0(\Omega)$ one has the action ρ , on $\mathcal{C}_0(Q_\omega)$ its obvious restriction and on $\mathcal{LUC}_u(G)$ the action θ of G by left translations, as in Subsection 7.2. Correspondingly, one gets the chain

$$\mathcal{C}_0(\Omega) \rtimes \xrightarrow{\gamma_\omega^\times} \mathcal{C}_0(Q_\omega) \rtimes \xrightarrow{\beta_\omega^\times} \mathcal{LUC}_u(G) \rtimes \xrightarrow{\text{Sch}_L^\tau} \mathbb{B}[L^2(G)].$$

We indicated crossed products of the form $\mathcal{B} \rtimes^\tau G$ by $\mathcal{B} \rtimes^\times$ (leaving the actions unnoticed) and the $*$ -morphism δ^\times acting between crossed products is deduced canonically from an equivariant $*$ -morphism δ by the procedure described in Remark 7.5. The arrow Sch_L^τ is just the left Schrödinger representation of Subsection 7.2. It is easy to check that

$$\text{Sch}_L^\tau \circ \beta_\omega^\times \circ \gamma_\omega^\times = \text{Sch}_{(\omega)}^\tau, \tag{97}$$

which also leads to recapturing (95) after a partial Fourier transformation.

Note that some points $\omega \in Q_{\omega'}$ could generate strictly smaller quasi-orbits $Q_\omega \subset Q_{\omega'}$. On the other hand a quasi-orbit can be generated by points belonging to different orbits, so Proposition 7.18 is not enough to prove the following result.

THEOREM 7.20. *Suppose that the group G is admissible and amenable and that $h \in \mathfrak{B}_\Omega^\tau$.*

1. *If ω, ω' generate the same ϱ -quasi-orbit, then $\text{Op}_{(\omega)}^\tau(h)$ and $\text{Op}_{(\omega')}^\tau(h)$ have the same spectrum.*
2. *If (Ω, ϱ, G) is a minimal dynamical system then all the operators $\text{Op}_{(\omega)}^\tau(h)$ have the same spectrum.*
3. *Assume that Ω is compact and metrizable and endowed with a Borel probability measure μ which is ϱ -invariant and ergodic. Then the topological support $\text{supp}(\mu)$ is a ϱ -quasi-orbit and one has $\mu[\{\omega \in \Omega \mid \overline{O_\omega} = \text{supp}(\mu)\}] = 1$. The operators $\text{Op}_{(\omega)}^\tau(h)$ corresponding to points generating this quasi-orbit have all the same spectrum; in particular $\text{sp}[\text{Op}_{(\omega)}^\tau(h)]$ is constant μ -a.e.*

Proof. 1. Let us denote by $Q_\omega := \overline{\varrho_G(\omega)}$ the quasi-orbit generated by ω and similarly for ω' . We show that if $Q_\omega \subset Q_{\omega'}$ then $\text{sp}[\text{Op}_{(\omega)}^\tau(h)] \subset \text{sp}[\text{Op}_{(\omega')}^\tau(h)]$ and this clearly implies the statement by changing the role of ω and ω' . Actually, by (92), under the stated inclusion of quasi-orbits, one needs to show that $\text{sp}[\text{Sch}_{(\omega)}^\tau(\Phi)] \subset \text{sp}[\text{Sch}_{(\omega')}^\tau(\Phi)]$ for every element Φ of the crossed product $\mathcal{C}_0(\Omega) \rtimes_\rho^\tau G$.

The basic idea, trivial consequence of the definitions, is the following: If $\Upsilon : \mathcal{E}' \rightarrow \mathcal{E}$ is a $*$ -morphism between two C^* -algebras and g' is an element of \mathcal{E}' , then $\text{sp}[\Upsilon(g') \mid \mathcal{E}] \subset \text{sp}[g' \mid \mathcal{E}']$, and we have equality of spectra if Υ is injective. The notation indicates the C^* -algebra in which each spectrum is computed.

In our case, by (97), one can write

$$\text{Sch}_{(\omega)}^{\tau}(\Phi) = [\text{Sch}_L^{\tau} \circ \beta_{\omega}^{\times}] [\gamma_{\omega}^{\times}(\Phi)] \quad \text{and} \quad \text{Sch}_{(\omega')}^{\tau}(\Phi) = [\text{Sch}_L^{\tau} \circ \beta_{\omega'}^{\times}] [\gamma_{\omega'}^{\times}(\Phi)].$$

Since G is amenable Sch_L^{τ} is injective and, as remarked before, β_{ω}^{\times} and $\beta_{\omega'}^{\times}$ are always injective. Thus we are left with proving that

$$\text{sp}[\gamma_{\omega}^{\times}(\Phi) | \mathcal{C}_0(\mathbb{Q}_{\omega})^{\times}] \subset \text{sp}[\gamma_{\omega'}^{\times}(\Phi) | \mathcal{C}_0(\mathbb{Q}_{\omega})^{\times}], \quad (98)$$

assuming the inclusion $\mathbb{Q}_{\omega} \subset \mathbb{Q}_{\omega'}$ of quasi-orbits. We use now

$$\Upsilon \equiv \gamma_{\omega',\omega}^{\times} : \mathcal{C}_0(\mathbb{Q}_{\omega'}) \rtimes_{\rho}^{\tau} G \rightarrow \mathcal{C}_0(\mathbb{Q}_{\omega}) \rtimes_{\rho}^{\tau} G,$$

which is obtained by applying the functorial construction of Remark 7.5 to the covariant restriction *-morphism

$$\gamma_{\omega',\omega} : \mathcal{C}_0(\mathbb{Q}_{\omega'}) \rightarrow \mathcal{C}_0(\mathbb{Q}_{\omega}), \quad \gamma_{\omega',\omega}(f) := f|_{\mathbb{Q}_{\omega}}.$$

Note that $\gamma_{\omega} = \gamma_{\omega',\omega} \circ \gamma_{\omega'}$ (successive restrictions), which functorially implies $\gamma_{\omega}^{\times} = \gamma_{\omega',\omega}^{\times} \circ \gamma_{\omega'}^{\times}$. Then $\gamma_{\omega}^{\times}(\Phi) = \gamma_{\omega',\omega}^{\times} [\gamma_{\omega'}^{\times}(\Phi)]$ and (98) and thus the result follows.

2. In a minimal dynamical system, by definition, all the orbits are dense. Thus any point generates the same single quasi-orbit $\mathbb{Q} = \Omega$ and one applies 1.

3. The statement concerning the properties of $\text{supp}(\mu)$ is contained in [2, Lemma 3.1]. Then the spectral information follows applying 1. once again. \square

The final point of Theorem 7.20 treats “a random Hamiltonian of pseudo-differential type”. Almost everywhere constancy of the spectrum in an ergodic random setting is a familiar property proved in many other situations [5, 35]. But note that a precise statement about the family of points giving the almost sure spectrum is available above.

8 THE CASE OF NILPOTENT LIE GROUPS

We now give the application of the introduced construction to the case of nilpotent Lie groups. Two previous main approaches seem to exist here. The first one uses the fact that, since the exponential mapping is a global diffeomorphism, one can introduce classes of symbols and the symbolic calculus on the group from the one on its Lie algebra. This allows for operators on a nilpotent Lie group G to have scalar-valued symbols which can be interpreted as functions on the dual \mathfrak{g}' of its Lie algebra. Such approach becomes effective mostly for invariant operators on general nilpotent Lie groups [33, 24, 25], see also [44] for the case of the Heisenberg group. The second approach applies also well to noninvariant operators on G and leads to operator-valued symbols, as developed in [18, 19]. This is also a special case (with $\tau(\cdot) = e$) of τ -quantizations developed in this paper.

We now extend both approaches to τ -quantizations with the link between them provided in Remark 8.5.

8.1 SOME MORE FOURIER TRANSFORMATIONS

Let us suppose that G is a nilpotent Lie group with unit e and Haar measure m ; it will also be assumed connected and simply connected. Such a group is unimodular, second countable and type I, so it fits in our setting and all the previous constructions and statements hold.

Let \mathfrak{g} be the Lie algebra of G and \mathfrak{g}' its dual. If $Y \in \mathfrak{g}$ and $X' \in \mathfrak{g}'$ we set $\langle Y|X' \rangle := X'(Y)$. We shall develop further the theory in this nilpotent setting, but only to the extent the next two basic properties are used:

1. the exponential map $\exp : \mathfrak{g} \rightarrow G$ is a diffeomorphism, with inverse $\log : G \rightarrow \mathfrak{g}$, [7, Th. 1.2.1];
2. under \exp the Haar measure on G corresponds to the Haar measure dX on \mathfrak{g} (normalised accordingly), cf [7, Th. 1.2.10].

It follows from the properties above that $L^p(G)$ is isomorphic to $L^p(\mathfrak{g})$. Actually, for each $p \in [1, \infty]$, one has a surjective linear isometry

$$L^p(G) \xrightarrow{\text{Exp}} L^p(\mathfrak{g}), \quad \text{Exp}(u) := u \circ \exp$$

with inverse

$$L^p(\mathfrak{g}) \xrightarrow{\text{Log}} L^p(G), \quad \text{Log}(u) := u \circ \log .$$

There is a unitary Fourier transformation $\mathcal{F} : L^2(\mathfrak{g}) \rightarrow L^2(\mathfrak{g}')$ associated to the duality $\langle \cdot | \cdot \rangle : \mathfrak{g} \times \mathfrak{g}' \rightarrow \mathbb{R}$. It is defined by

$$(\mathcal{F}u)(X') := \int_{\mathfrak{g}} e^{-i\langle X|X' \rangle} u(X) dX ,$$

with inverse given (for a suitable normalization of dX') by

$$(\mathcal{F}^{-1}u')(X) := \int_{\mathfrak{g}'} e^{i\langle X|X' \rangle} u'(X') dX' .$$

Now composing with the mappings Exp and Log one gets unitary Fourier transformations

$$\mathbf{F} := \mathcal{F} \circ \text{Exp} : L^2(G) \rightarrow L^2(\mathfrak{g}'), \quad \mathbf{F}^{-1} := \text{Log} \circ \mathcal{F}^{-1} : L^2(\mathfrak{g}') \rightarrow L^2(G) ,$$

the second one being the inverse of the first. They are defined essentially by

$$(\mathbf{F}u)(X') = \int_{\mathfrak{g}} e^{-i\langle X|X' \rangle} u(\exp X) dX = \int_G e^{-i\langle \log x|X' \rangle} u(x) dm(x) ,$$

$$(\mathbf{F}^{-1}u')(x) = \int_{\mathfrak{g}'} e^{i\langle \log x|X' \rangle} u'(X') dX' .$$

Recalling Plancherel’s Theorem for unimodular second countable type I groups, one gets finally a commuting diagram of unitary transformations

$$\begin{array}{ccc}
 L^2(\mathbb{G}) & \xrightarrow{\text{Exp}} & L^2(\mathfrak{g}) \\
 \mathcal{F} \downarrow & \searrow \mathbf{F} & \downarrow \mathcal{F} \\
 \mathcal{B}^2(\widehat{\mathbb{G}}) & \xrightarrow{\mathfrak{J}} & L^2(\mathfrak{g}')
 \end{array}$$

The lower horizontal arrow is defined as $\mathfrak{J} := \mathbf{F} \circ \mathcal{F}^{-1} = \mathcal{F} \circ \text{Exp} \circ \mathcal{F}^{-1}$ and is given explicitly on $\mathcal{B}^1(\widehat{\mathbb{G}}) \cap \mathcal{B}^2(\widehat{\mathbb{G}})$ by

$$(\mathfrak{J}\phi)(X') = \int_{\mathbb{G}} \int_{\widehat{\mathbb{G}}} e^{-i(\log x|X')} \text{Tr}_{\xi}[\phi(\xi)\xi(x)] dm(x)d\widehat{m}(\xi).$$

REMARK 8.1. If $\mathbb{G} = \mathbb{R}^n$ it is possible, by suitable interpretations, to identify $\mathbb{G} \sim \widehat{\mathbb{G}}$ with \mathfrak{g} and with \mathfrak{g}' (as vector spaces) and then the three Fourier transformations \mathcal{F} , \mathcal{F} and \mathbf{F} will coincide and \mathfrak{J} will become the identity.

8.2 A QUANTIZATION BY SCALAR SYMBOLS ON NILPOTENT LIE GROUPS

To get pseudo-differential operators one could start, as in Subsection 7.2, with a C^* -dynamical system $(\mathcal{A}, \theta, \mathbb{G})$ where \mathcal{A} is a C^* -algebra of bounded left-uniformly continuous functions on \mathbb{G} which is invariant under the action θ by left translations. We compose the left Schrödinger representation (82) with the inverse of the partial Fourier transform

$$\text{id} \otimes \mathbf{F} : (L^1 \cap L^2)(\mathbb{G}; \mathcal{A}) \rightarrow \mathcal{A} \otimes L^2(\mathfrak{g}'),$$

finding the pseudo-differential representation

$$\text{Op}_L^{\tau} := \text{Sch}_L^{\tau} \circ (\text{id} \otimes \mathbf{F}^{-1}) = \text{Int} \circ \text{CV}_L^{\tau} \circ (\text{id} \otimes \mathbf{F}^{-1}) \tag{99}$$

which can afterwards be extended to the relevant enveloping C^* -algebra. One gets

$$[\text{Op}_L^{\tau}(\mathfrak{s})u](x) = \int_{\mathbb{G}} \int_{\mathfrak{g}'} e^{i(\log(xy^{-1})|X')} \mathfrak{s}(\tau(xy^{-1})^{-1}x, X') u(y) dm(y)dX', \tag{100}$$

so $\text{Op}_L^{\tau}(\mathfrak{s})$ is an integral operator with kernel $\mathbf{Ker}_{(\mathfrak{s})}^{\tau} : \mathbb{G} \times \mathbb{G} \rightarrow \mathbb{C}$ given by

$$\mathbf{Ker}_{(\mathfrak{s})}^{\tau}(x, y) = \int_{\mathfrak{g}'} e^{i(\log(xy^{-1})|X')} \mathfrak{s}(\tau(xy^{-1})^{-1}x, X') dX'.$$

Examining this kernel, or using directly (99), one sees that (100) also defines a unitary mapping

$$\text{Op}_L^{\tau} : L^2(\mathfrak{g}' \times \mathbb{G}) \rightarrow \mathbb{B}^2[L^2(\mathbb{G})].$$

Actually there is a Weyl system on which the construction of pseudo-differential operators with symbols $\mathfrak{s} : \mathfrak{g}' \times \mathbb{G} \rightarrow \mathbb{C}$ can be based:

DEFINITION 8.2. For $(x, X') \in \mathbf{G} \times \mathfrak{g}'$ one defines a unitary operator $\mathbf{W}_L^\tau(x, X')$ in $L^2(\mathbf{G})$ by

$$\begin{aligned} [\mathbf{W}_L^\tau(x, X')u](z) &:= e^{i\langle \log[\tau(x)^{-1}z] | X' \rangle} u(x^{-1}z) \\ &= e^{i\langle \log[\tau(x)^{-1}z] | X' \rangle} [L(x)u](z). \end{aligned}$$

By direct computations, one shows the following

LEMMA 8.3. Let us denote by Q the operator of multiplication by the variable in $L^2(\mathbf{G})$. For any pairs $(x, X'), (y, Y') \in \mathbf{G} \times \mathfrak{g}'$ one has

$$\mathbf{W}_L^\tau(x, X') \mathbf{W}_L^\tau(y, Y') = \Upsilon^\tau[(x, X'), (y, Y'); Q] \mathbf{W}_L^\tau(xy, X' + Y'),$$

where $\Upsilon^\tau[(x, X'), (y, Y'); Q]$ is the operator of multiplication by the function

$$\begin{aligned} z \mapsto \Upsilon^\tau[(x, X'), (y, Y'); z] &= \exp \left\{ i \left[\langle \log[\tau(x)^{-1}z] - \log[\tau(xy)^{-1}z] | X' \rangle - \right. \right. \\ &\quad \left. \left. - \langle \log[\tau(xy)^{-1}z] - \log[\tau(y^{-1})x^{-1}z] | Y' \rangle \right] \right\}. \end{aligned}$$

REMARK 8.4. The family $\mathcal{C}(\mathbf{G}; \mathbb{T})$ of all continuous functions on \mathbf{G} with values in the torus is a Polish group and the mapping $\Upsilon : (\mathbf{G} \times \mathfrak{g}') \times (\mathbf{G} \times \mathfrak{g}') \rightarrow \mathcal{C}(\mathbf{G}; \mathbb{T})$ can be seen as a 2-cocycle. We are not going to pursue here the cohomological meaning and usefulness of these facts.

In terms of the Weyl system $\{\mathbf{W}_L^\tau(x, X') \mid (x, X') \in \mathbf{G} \times \mathfrak{g}'\}$ one can write

$$\mathbf{Op}_L^\tau(\mathbf{s}) := \int_{\mathbf{G}} \int_{\mathfrak{g}'} \tilde{\mathbf{s}}(X', x) \mathbf{W}_L^\tau(x, X') dm(x) dX'; \tag{101}$$

we used the notation $\tilde{\mathbf{s}} := (\mathbf{F} \otimes \mathbf{F}^{-1})\mathbf{s}$. The technical details are similar but simpler than those in Subsection 3.2 and are left to the reader.

REMARK 8.5. One also considers the composition \sharp_τ defined to satisfy the equality $\mathbf{Op}_L^\tau(\mathbf{r} \sharp_\tau \mathbf{s}) = \mathbf{Op}_L^\tau(\mathbf{r}) \mathbf{Op}_L^\tau(\mathbf{s})$, as well as the involution \sharp_τ verifying $\mathbf{Op}_L^\tau(\mathbf{s} \sharp_\tau) = \mathbf{Op}_L^\tau(\mathbf{s})^*$. Then $(L^2(\mathfrak{g}' \times \mathbf{G}), \sharp_\tau, \sharp_\tau)$ will be a $*$ -algebra. It is isomorphic to the $*$ -algebra $(\mathcal{B}^2(\widehat{\mathbf{G}} \times \mathbf{G}), \sharp_\tau, \sharp_\tau)$ defined in Subsection 3.3. Actually one has the following commutative diagram of isomorphisms:

$$\begin{array}{ccc} L^2(\mathbf{G}) \otimes L^2(\mathbf{G}) & \xrightarrow{\text{id} \otimes \mathcal{F}} & L^2(\mathbf{G}) \otimes \mathcal{B}^2(\widehat{\mathbf{G}}) \\ \text{id} \otimes \mathbf{F} \downarrow & \searrow \text{Sch}_L^\tau & \downarrow \mathbf{Op}_L^\tau \\ L^2(\mathbf{G}) \otimes L^2(\mathfrak{g}') & \xrightarrow{\mathbf{Op}_L^\tau} & \mathbb{B}^2[L^2(\mathbf{G})] \end{array}$$

One justifies this diagram by comparing (99) with (83). The conclusion of this diagram is that for simply connected nilpotent Lie groups the “operator-valued pseudo-differential calculus” \mathbf{Op}_L^τ with symbols defined on $\mathbf{G} \times \widehat{\mathbf{G}}$ can be obtained from the

“scalar-valued pseudo-differential calculus” \mathbf{Op}_L^τ (which provides a quantization on the cotangent bundle $\mathbf{G} \times \mathfrak{g}' \cong T'(\mathbf{G})$) just by composing at the level of symbols with the isomorphism $(\text{id} \otimes \mathbf{F}) \circ (\text{id} \otimes \mathcal{F})^{-1} = \text{id} \otimes (\mathbf{F} \circ \mathcal{F}^{-1})$.

REMARK 8.6. In Prop. 4.3 we have shown that a connected simply connected nilpotent Lie group \mathbf{G} admits a symmetric quantization, corresponding to the map $\tau = \sigma$ given by (58) globally defined. With this choice one also has $\mathbf{s}^{\sharp\sigma}(x, X') = \mathbf{s}(x, X')$ for every $(x, X') \in \mathbf{G} \times \mathfrak{g}'$.

REMARK 8.7. A right quantization \mathbf{Op}_R^τ with scalar symbols is also possible; for completeness, we list the main quantization formula

$$\mathbf{Op}_R^\tau := \text{Sch}_R^\tau \circ (\text{id} \otimes \mathbf{F}^{-1}) \equiv \text{Int} \circ \text{CV}_R^\tau \circ (\text{id} \otimes \mathbf{F}^{-1}), \tag{102}$$

where CV_R^τ is the change of variables given by the composition with the mapping

$$\text{cv}^\tau \equiv \text{cv}_R^\tau : \mathbf{G} \times \mathbf{G} \rightarrow \mathbf{G} \times \mathbf{G}, \quad \text{cv}_R^\tau(x, y) := (x\tau(y^{-1}x)^{-1}, y^{-1}x), \tag{103}$$


see also (23). Here, $\text{Sch}_R^\tau := \text{Int} \circ \text{CV}_R^\tau$ also allows for an integrated interpretation similar to (81). More explicitly, \mathbf{Op}_R^τ can be written as

$$[\mathbf{Op}_R^\tau(\mathbf{s})u](x) = \int_{\mathbf{G}} \int_{\mathfrak{g}'} e^{i(\log(y^{-1}x)|X')} \mathbf{s}(x\tau(y^{-1}x)^{-1}, X') u(y) \, dm(y) dX', \tag{104}$$

so $\mathbf{Op}_R^\tau(\mathbf{s})$ is an integral operator with kernel $\mathbf{Ker}_{(\mathbf{s}),R}^\tau : \mathbf{G} \times \mathbf{G} \rightarrow \mathbb{C}$ given by

$$\mathbf{Ker}_{(\mathbf{s}),R}^\tau(x, y) = \int_{\mathfrak{g}'} e^{i(\log(y^{-1}x)|X')} \mathbf{s}(x\tau(y^{-1}x)^{-1}, X') \, dX'.$$

Consequently, we have the commutative diagram of isomorphisms of H^* -algebras



where all the listed mappings in this diagram are unitary, and where $\text{Op}^\tau = \text{Op}_R^\tau$ is the τ -quantization formula (2) that we have started with, and so

$$\text{Op}^\tau = \text{Op}_R^\tau = \mathbf{Op}_R^\tau \circ [\text{id} \otimes (\mathbf{F} \circ \mathcal{F}^{-1})].$$

REFERENCES

[1] Ph. M. Anselone: *Compactness Properties of Sets of Operators and Their Adjoints*, Math. Z. 113, 233-236, (1970).

- [2] J. Bellissard, D. Herrmann and M. Zarrouati: *Hull of Aperiodic Solids and Gap Labelling Theorems*, in Directions in Mathematical Quasicrystals, CRM Monograph Series, 13, 207–259, 2000.
- [3] I. Beltiță, D. Beltiță and M. Măntoiu: *Quantization and Dequantization via Square-Integrable Families of Operators*, Rocky Mount. Math., 46 (6), 1795–1851, (2016).
- [4] F. Bruhat: *Distributions sur un groupe localement compact et applications à l'étude des représentations des groupes p -adiques*, Bull. Soc. Math. France 89, 43–75, (1961).
- [5] R. Carmona and J. Lacroix: *Spectral Theory of Random Schrödinger Operators*, Birkhäuser Boston Inc., Boston, MA, 1990.
- [6] M. Christ, D. Geller, P. Głowacki and L. Polin: *Pseudodifferential Operators on Groups with Dilations*, Duke Math. J., 68, no. 1, 31–65, (1992).
- [7] L. J. Corwin and F. P. Greenleaf: *Representations of Nilpotent Lie Groups and their Applications*, Cambridge University Press, 1990.
- [8] A. Dasgupta and M. Ruzhansky: *Gevrey Functions and Ultradistributions on Compact Lie Groups and Homogeneous Spaces*, Bull. Sci. Math., 138, 756–782, (2014).
- [9] A. Dasgupta and M. Ruzhansky: *Gohberg Lemma, Compactness, and Essential Spectrum of Operators on Compact Lie Groups*, J. Anal. Math., 128, 179–190, (2016).
- [10] M. Duflo and C.C. Moore: *On the Regular Representation of a Nonunimodular Locally Compact Group*, J. Funct. Anal. 21, 209–243 (1976).
- [11] J. Delgado and M. Ruzhansky: *Schatten Classes and Traces on Compact Lie Groups*, Preprint ArXiv and Math. Res. Lett., to appear.
- [12] J. Delgado and M. Ruzhansky: *Kernel and symbol criteria for Schatten classes and r -nuclearity on compact manifolds*, C. R. Acad. Sci. Paris, Ser. I 352, 779–784 (2014).
- [13] J. Delgado and M. Ruzhansky: *L^p -nuclearity, Traces, and Grothendieck-Lidskii Formula on Compact Lie Groups*, J. Math. Pures Appl., 102, no. 1, 153–172, (2014).
- [14] J. Dixmier: *Les C^* -algèbres et leurs représentations*, Cahiers scientifiques, XXIX, Gauthier-Villars Éditeurs, Paris, 1969.
- [15] J. Dixmier: *Von Neumann Algebras*, North Holland Math. Library, 1981.
- [16] H. G. Feichtinger: *Compactness in Translational Invariant Banach Spaces of Distributions and Compact Multipliers*, J. Math. Anal. Appl. 102, no. 2, 289–327, (1984).

- [17] V. Fischer: *Intrinsic Pseudo-differential Calculi on any Compact Lie Group*, J. Funct. Anal. 268, 3404–3477, (2015).
- [18] V. Fischer and M. Ruzhansky: *A Pseudo-differential Calculus on Graded Nilpotent Groups*, in *Fourier Analysis*, pp. 107–132, Trends in Mathematics, Birkhäuser, 2014.
- [19] V. Fischer and M. Ruzhansky: *Quantization on Nilpotent Lie Groups*, Progress in Mathematics, Vol. 314, Birkhäuser, 2016. (open access book)
- [20] V. Fischer and M. Ruzhansky: *A Pseudo-differential Calculus on the Heisenberg Group*, C. R. Acad. Sci. Paris, Ser I, 352, 197–204, (2014).
- [21] H. Führ: *Abstract Harmonic Analysis of Continuous Wavelet Transforms*, Lecture Notes in Mathematics 1863, Springer-Verlag Berlin Heidelberg, 2005.
- [22] G. B. Folland: *Harmonic Analysis in Phase Space*, Annals of Mathematics Studies, 122. Princeton University Press, Princeton, NJ, 1989.
- [23] G. B. Folland: *A Course in Abstract Harmonic Analysis*, CRC Press, Boca Raton Ann Arbor London Tokio, 1995.
- [24] P. Głowacki, *Invertibility of convolution operators on homogeneous groups*, Rev. Mat. Iberoam. 28, no. 1, 141–156, (2012).
- [25] P. Głowacki, *The Melin calculus for general homogeneous groups*, Ark. Mat., 45, no. 1, 31–48, (2007).
- [26] R.L. Lipsman: *Non-Abelian Fourier Analysis*, Bull. Sci. Math. 98, 209–233, (1974).
- [27] M. Măntoiu: *Essential Spectrum and Fredholm Properties for Operators on Locally Compact Groups*, J. Operator Theory, 77 (2), 481–501, (2017).
- [28] M. Măntoiu and D. Parra: *Compactness Criteria in Banach Spaces in the Setting of Continuous Frames*, Banach J. of Math. Anal. 8 (2), 30–48, (2014).
- [29] M. Măntoiu and R. Purice: *On Frechet-Hilbert Algebras*, Ark. Math, 103, no. 2, 157–166, (2014).
- [30] M. Măntoiu, R. Purice and S. Richard: *Twisted Crossed Products and Magnetic Pseudodifferential Operators*, Proc. of the Conference of Sinaia, Theta Foundation (2005).
- [31] M. Măntoiu and M. Ruzhansky: *Quantizations on Nilpotent Lie Groups and Algebras Having Flat Coadjoint Orbits*, arXiv:1611.07581
- [32] G.A. Meladze and M.A. Shubin: *Algebras of Pseudodifferential Operators on Unimodular Lie Groups*, (Russian) Dokl. Akad. Nauk SSSR, 279, no. 3, 542–545, (1984).

- [33] A. Melin: *Parametrix Constructions for Right Invariant Differential Operators on Nilpotent Groups*, Ann. Global Anal. Geom., 1, no. 1, 79–130, (1983).
- [34] W. Palmer: *Totally Bounded Sets of Precompact Linear Operators*, Proc. AMS, 20, 101-106, (1969).
- [35] L. A. Pastur and A. Figotin: *Spectra of Random and Almost Periodic Operators*, Springer Verlag, Berlin, 1992.
- [36] G. Pedersen: *C^* -Algebras and their Automorphisms Groups*, LMS Monographs, 14, Academic Press, San Diego, 1979.
- [37] N. V. Pedersen: *Matrix Coefficients and a Weyl Correspondence for Nilpotent Lie Groups*. Invent. Math. 118, 1–36, (1994).
- [38] G. Pisier and Q. Xu: *Non-commutative L^p -spaces*, unpublished manuscript, (1999).
- [39] M. Ruzhansky and V. Turunen: *Pseudo-differential Operators and Symmetries*, Pseudo-Differential Operators: Theory and Applications 2, Birkhäuser Verlag, 2010.
- [40] M. Ruzhansky and V. Turunen: *Quantization of Pseudo-differential Operators on the Torus*, J. Fourier Anal. Appl. 16, 943–982, (2010).
- [41] M. Ruzhansky and V. Turunen: *Global Quantization of Pseudo-differential Operators on Compact Lie Groups, $SU(2)$ and 3-Sphere*, Int. Math. Res. Not., no. 11, 2439–2496 (2013).
- [42] M. Ruzhansky, V. Turunen and J. Wirth: *Hörmander-Class of Pseudo-Differential Operators on Compact Lie Groups and Global Hypoellipticity*, J. Fourier Anal. Appl., 20, 476–499, (2014).
- [43] M. Ruzhansky and J. Wirth: *Global Functional Calculus for Operators on Compact Lie Groups*, J. Funct. Anal. 267, 144–172, (2014).
- [44] M. Taylor: *Noncommutative Microlocal Analysis*, Mem. Amer. Math. Soc., 313, 1984.
- [45] E. Thoma: *Eine Charakterisierung diskreter Gruppen vom Typ I*, Invent. Math. 6, 190–196, (1968).
- [46] F. Trèves, *Topological Vector Spaces, Distributions and Kernels*, Academic Press, 1967.
- [47] D. Williams: *Crossed Products of C^* -Algebras*, Mathematical Surveys and Monographs, 134, American Mathematical Society, 2007.
- [48] Q. Xu: *Operator Spaces and Noncommutative L^p . The Part on Noncommutative L^p -Spaces*, Summer School on Banach Spaces and Operator Spaces, Nankai University, China, (2007).

Marius Măntoiu
Dept. de Matemáticas,
Facultad de Ciencias,
Universidad de Chile
Casilla 653,
Las Palmeras 3425,
Santiago, Chile
mantoiu@uchile.cl

Michael Ruzhansky
Department of Mathematics,
Imperial College London
180 Queen's Gate, London
SW7 2AZ,
United Kingdom
m.ruzhansky@imperial.ac.uk