

FROBENIUS GAUGES AND A NEW THEORY OF
 p -TORSION SHEAVES IN CHARACTERISTIC p

JEAN-MARC FONTAINE[†] AND UWE JANNSEN

Received: July 1, 2019

Revised: February 26, 2021

Communicated by Thomas Geisser

ABSTRACT. We develop a new cohomology theory in characteristic $p > 0$, the so called F-gauge cohomology, a cohomology with values in the category of so-called F-gauges, which refines the crystalline cohomology. We mainly discuss the theory for smooth projective varieties over perfect fields of characteristic p . We also compare our theory with other existing structures like the F-zips of Moonen and Wedhorn and the displays of Langer and Zink.

2020 Mathematics Subject Classification: 14F20, 14F30

Keywords and Phrases: Gauges, p -torsion sheaves, gauge cohomology

1 INTRODUCTION

Let k be a field, k_s a separable closure of k and $G_k = \text{Gal}(k_s/k)$. Let X be a proper and smooth variety over k . This defines a morphism of topoi $\pi : X_{\text{ét}} \rightarrow (\text{Spec } k)_{\text{ét}}$. Let l be a prime number and F be any finite l -torsion abelian sheaf over $(\text{Spec } k)_{\text{ét}}$, such as $\mathbb{Z}/l^n\mathbb{Z}$ or μ_{l^n} (with $n \in \mathbb{N}$), then, for any $j \in \mathbb{N}$, $\mathcal{F} = R^j\pi_*\pi^*F$ is again a finite abelian sheaf over $(\text{Spec } k)_{\text{ét}}$. To know \mathcal{F} amounts to the same as knowing its *Galois module*, that is the finite abelian group $\mathcal{F}(k_s)$ (the fiber of \mathcal{F} at the geometric point k_s) with its natural linear and discrete action of G_k .

If l is different from the characteristic of k , this gives rise to a nice theory, the *étale l -adic cohomology* which has many useful applications.

[†]The author passed away on 29. Januar 2019.

The editors would like to thank E. Lau for helping with the final version of the manuscript.

Assume now k is a perfect field of characteristic $p > 0$ and $l = p$. Then the étale topology is not fine enough for many applications. For instance, the restriction of μ_{p^n} to the small étale site of k is just the trivial sheaf. If we now view π as a morphism for the fppf topology

$$\pi : X_{flat} \rightarrow (\text{Spec } k)_{flat}$$

then $\mathcal{F} = R^1\pi_*\pi^*\mu_{p^n}$ is a finite commutative group scheme over k . To know \mathcal{F} amounts to the same as knowing its *Dieudonné module*, an elementary object whose definition involves some linear algebra.

Now $R^1\pi_*\pi^*\mu_{p^n}$ is a *good* object to consider. We could also consider $R^1\pi_*\pi^*\mathbb{Z}/p^n\mathbb{Z}$ but it would be too small for some applications. Too small as well would be the sheaves $R^j\pi_*\pi^*\mu_{p^n}$ for $j > 1$. Roughly speaking we want to introduce some sheaves of p -torsion, the S_n^r (for $n, r \in \mathbb{N}$ (we shall have $S_n^0 = \mathbb{Z}/p^n\mathbb{Z}$ and $S_n^1 = \mu_{p^n}$) and to consider the $R^j\pi_*\pi^*S_n^r$. They will belong to a nice class of p -torsion sheaves which are classified by their *Dieudonné modules* (elementary objects generalizing the classical Dieudonné modules). Technically,

- i) we will have to use a topology which is weaker than the flat topology but also stronger than the étale topology,
- ii) the generalized Dieudonné modules give more complicated objects, so-called gauges, or rather φ -(or Frobenius-)gauges.

The aim of this paper is to give the definition of these φ -gauges, relate them to previous constructions like Dieudonné modules, F -zips, or displays, and to define a cohomology theory with values in φ -gauges, which refines crystalline cohomology. For examples and applications we will rather concentrate on the case of varieties over fields.

In following papers we will construct and discuss the functors between φ -gauges and certain p -torsion sheaves for the syntomic cohomology, will develop a relative theory, and will discuss relations with p -adic Hodge theory over discrete valuation rings.

Remark: Let K be a field of characteristic 0, complete with respect to a discrete valuation, with perfect residue field k of characteristic p . Let \overline{K} be an algebraic closure of K and $G_K = \text{Gal}(\overline{K}/K)$. Let V be a crystalline representation of G_K with non negative Hodge-Tate weights and T a G_K -stable \mathbb{Z}_p -lattice of V . Then V/T may be viewed in a natural way as one of the fibers of a sheaf Γ for the syntomic-étale topology over $\text{Spec } \mathcal{O}_K$ (for instance, if the Hodge-Tate weights are 0 and 1 this is a Barsotti-Tate group (or p -divisible group) over \mathcal{O}_K . This Γ has a special fiber Γ_k and, roughly speaking the kernel of the multiplication by p^n on Γ_k is one of these nice p -torsion sheaves that we are constructing. Somehow, the theory we develop here is the *special fiber of classical p -adic Hodge theory*.

One application we expect from this theory is to deformation of Galois representations coming from algebraic geometry. As we just said, these may often

be extended in a natural way to a p -adic sheaf for the syntomic-étale topology. The knowledge of the p -adic sheaf is equivalent to the knowledge of the Galois representation. But when we take sub-quotients killed by a power of p , this is no more true in general and it may be wise to look at the deformations of the sheaves rather than just at the deformations of the Galois representations. In the case of Barsotti-Tate groups, this idea has already been used by Kisin [Ki]. A long time ago, just after our joint work on p -adic Hodge theory, William Messing and one of us (JMF) started to think about this kind of things. What follows is just a continuation of this old work which has never been completed and we want to thank Bill heartily for the old and new discussions we had with him on that.

This paper is organized as follows: In section 1, we introduce the notion of gauges, φ -modules, and φ -gauges, which are the basic objects from linear algebra which give rise to the notion of generalized Dieudonné modules. In section 2, we study how these structures arise from (virtual) W -crystals, where $W = W(k)$ is the ring of Witt vectors for a perfect field of characteristic p , and we discuss properties of the new category in this case. In sections 3 and 4 we consider arbitrary schemes in positive characteristic and show that our theory contains and extends the theories of F -zips and displays. In sections 5 and 6 we discuss the different Grothendieck topologies that we are going to use and their basic properties. In section 7 we recall the definitions and properties of the syntomic sheaves of rings \mathcal{O}_n^{cris} and we explain how one can use these rings to get a gauge of rings \mathcal{G} , also called the universal gauge, which is central for our theory. In section 8 we define the gauge cohomology (cohomology theory with values in the category of gauges) which is a refinement of crystalline cohomology, and we give some first properties.

2 GRADED OBJECTS, GAUGES, φ -MODULES, φ -GAUGES, AND φ -RINGS

2.1 Graded objects and p -gauges

By a graded object in an abelian category \mathcal{A} we mean a \mathbb{Z} -graded object, which is just a collection $A = (A^n)$ of objects indexed by \mathbb{Z} . If direct sums exist in \mathcal{A} , we may also think of the direct sum $\bigoplus_n A^n$. A morphism $f : A \rightarrow B$ of graded objects of degree d is a collection of morphisms $f_n : A^n \rightarrow B^{n+d}$. A morphism of graded objects is a morphism of degree 0. Graded objects in \mathcal{A} form again an abelian category.

Fix a prime p . A p -gauge in \mathcal{A} is a graded object M in \mathcal{A} together with a morphism f of degree 1 and a morphism v of degree -1

$$\dots \rightrightarrows M^{r-1} \underset{v}{\overset{f}{\rightleftarrows}} M^r \underset{v}{\overset{f}{\rightleftarrows}} M^{r+1} \rightrightarrows \dots$$

with $fv = p = vf$. Morphisms of p -gauges are morphisms α of graded object which are compatible with f and v (i.e., $\alpha f = f\alpha$ and $\alpha v = v\alpha$).

If, for example, \mathcal{A} is the category of modules over a ring, then a p -gauge is simply a module over the commutative graded ring $D(R) = R[f, v]/(fv - p)$, where $R[f, v]$ is the free graded ring generated by f in degree 1 and v in degree -1 .

For $-\infty \leq a \leq b \leq \infty$ we say that the p -gauge (M, f, v) is concentrated in the interval $[a, b]$, if v is an isomorphism to the left of M^a and f is an isomorphism to the right of M^b . If a and b are finite, then such a p -gauge is just determined by the finite diagram

$$M^a \rightleftarrows \dots \rightleftarrows M^{r-1} \begin{array}{c} \xrightarrow{f} \\ \xleftarrow{v} \end{array} M^r \begin{array}{c} \xrightarrow{f} \\ \xleftarrow{v} \end{array} M^{r+1} \rightleftarrows \dots \rightleftarrows M^b,$$

because everything is determined outside the interval $[a, b]$.

Call a p -gauge over a ring R (i.e., in the category of R -modules) of finite type, if it is finitely generated as a module over $D(R) = R[f, v]/(fv - p)$.

LEMMA 2.1.1. *Let R be a noetherian ring, and let M be a p -gauge of R -modules. Then the following conditions are equivalent.*

(a) M is of finite type.

(b) Each M^r is finitely generated as an R -module, and M is concentrated in a finite interval (i.e., $f : M^r \rightarrow M^{r+1}$ is an isomorphism for $r \gg 0$ and $v : M^r \rightarrow M^{r-1}$ is an isomorphism for $r \ll 0$).

Proof. (Compare the positively graded case in [GW], Lemma 13.10.) Assume (a) and let m_1, \dots, m_r be generators of M , without restriction each m_i homogenous of degree $d_i \in \mathbb{Z}$, say. Let d_{min} be the minimum of the d_i , and let d_{max} be their maximum. Then every element of M^n is a R -linear combination of the elements in the set

$$S_n = \{ f^a m_i \mid a \geq 0, n = a + d_i \} \cup \{ v^b m_i \mid b \geq 0, n = d_i - b \}.$$

(Note that $fv = p = vf$.) Since

$$d_{min} - n \leq b = d_i - n \leq d_{max} - n \quad \text{and} \quad n - d_{max} \leq a - d_i \leq n - d_{min},$$

these are finitely many elements, which shows the first claim in (a).

For the second claim we first note that $f : M^n \rightarrow M^{n+1}$ is surjective for $n \geq d_{max}$. In fact, for the elements $v^b m_i$ with $n + 1 = d_i - b$ and $b \geq 0$ in the generating set S_{n+1} above we would have $d_i - b = n + 1 \geq d_{max} + 1$, i.e., $d_i \geq d_{max} + 1$, a contradiction. Hence these elements do not appear. Moreover, for the elements $f^a m_i$ with $a + d_i = n + 1 \geq d_{max} + 1$ we must have $a \geq d_{max} - d_i + 1 \geq 1$.

Next, for $d = d_{max}$ the sequence of surjections $M^d \xrightarrow{f} M^{d+1} \xrightarrow{f} M^{d+2} \xrightarrow{f} \dots$ becomes stationary, because R is noetherian and all M^i are finitely generated. Thus $f : M^n \rightarrow M^{n+1}$ is bijective for $n \gg 0$. In a similar (dual) way one proves that $v : M^{n+1} \rightarrow M^n$ is an isomorphism for $n \ll 0$.

The converse implication from (b) to (a) is easier: If M is concentrated in the finite interval $[a, b]$, and F is a finite generating set for the R -module $M^a \oplus M^{a+1} \oplus \dots \oplus M^{b-1} \oplus M^b$, then F is a generating set for the graded $D(R)$ -module M . \square

In the following we will always fix a prime p and will omit it in the notation. Of course we could make a more general definition and replace the multiplication by p by any natural transformation $t : id_{\mathcal{A}} \rightarrow id_{\mathcal{A}}$.

2.2 φ -modules and φ -gauges

For any gauge (M, f, v) in \mathcal{A} we define

$$M^{+\infty} = \varinjlim_{r \rightarrow +\infty} M^r \quad \text{and} \quad M^{-\infty} = \varprojlim_{r \rightarrow -\infty} M^r,$$

where the transition morphisms are given by the morphisms f and the morphisms v , respectively, and where these objects either exist as direct limits in \mathcal{A} or as objects in the *Ind*-category of \mathcal{A} .

Now let $\sigma : \mathcal{A} \rightarrow \mathcal{A}$ be some endomorphism of \mathcal{A} . A φ -module (with respect to σ) is a gauge (M, f, v) together with a morphisms

$$\varphi : \sigma(M^{+\infty}) \longrightarrow M^{-\infty}.$$

Morphisms of φ -modules are morphisms α of gauges which are compatible with φ (i.e., for the morphisms $\alpha^{+\infty}$ and $\alpha^{-\infty}$ induced by α on the limit terms one has $\varphi\sigma(\alpha^{+\infty}) = \alpha^{-\infty}\varphi$). A φ -gauge is a φ -module for which φ is an isomorphism. If the formation of inductive limits is exact, then it is easy to see that the φ -modules form an abelian category if σ is right-exact, and that the category of φ -gauges is abelian if σ is exact.

2.3 Tensor products of graded modules and gauges

To fix ideas let \mathcal{T} be a topos. Assume that \mathcal{T} is the topos of sheaves over \mathcal{C} for the topology E .

If \mathcal{F} is a sheaf of sets (or groups, or ...), a *global section of \mathcal{F}* is a collection $(s_U)_{U \in \text{Ob } \mathcal{C}}$ such that, for any morphism $f : V \rightarrow U$ of \mathcal{C} , we have $f^{-1}(s_U) = s_V$. The global sections of \mathcal{F} form a set (resp. a group,...) $\Gamma(\mathcal{F})$. If \mathcal{C} has a final object S , we have $\Gamma(\mathcal{F}) = \mathcal{F}(S)$.

Below we will consider groups, rings, modules in \mathcal{T} , but for simplicity, we will omit \mathcal{T} and do as if we were just using plain groups, rings, modules (which would correspond to the trivial topos), by thinking of local sections,... at least when the extension of the things considered to the general setting is completely straightforward. In the whole paper, a graded ring is a commutative graded ring with grading indexed by \mathbb{Z} .

Let $R = \bigoplus_{r \in \mathbb{Z}} R^r$ be such a graded ring. A *graded R -module* is an R -module M together with a decomposition $M = \bigoplus_{r \in \mathbb{Z}} M^r$ of M into a direct sum of abelian groups such that, if $r, s \in \mathbb{Z}$, $\lambda \in R^r$ and $x \in M^s$, then $\lambda x \in M^{r+s}$. The graded R -modules form a $\Gamma(R^0)$ -linear abelian category.

Let M and N be two graded R -modules. For $r \in \mathbb{Z}$, set $(M \otimes_{R^0} N)^r = \bigoplus_{i+j=r} M^i \otimes_{R^0} N^j$. Let L^r be the sub-group of $(M \otimes_{R^0} N)^r$ (locally) generated by the $\lambda x \otimes y - x \otimes \lambda y$, for $\lambda \in R^i$, $x \in M^j$, $y \in N^k$ and $i + j + k = r$. We have $M \otimes_{R^0} N = \bigoplus_{r \in \mathbb{Z}} (M \otimes_{R^0} N)^r$ and

$$M \otimes_R N = \bigoplus_{r \in \mathbb{Z}} (M \otimes_R N)^r \text{ with } (M \otimes_R N)^r = (M \otimes_{R^0} N)^r / L^r$$

This endows $M \otimes_R N$ with a structure of graded R -module. In this way, graded- R -modules form a tensor category.

If M is a graded R -module, for any $i \in \mathbb{Z}$, the *i -th Tate twist of M* is the graded R -module whose underlying R -Module is M and with $M(i)^r = M^{r+i}$. The functor $M \mapsto M(1)$, from the category of graded R -Modules to itself, is an equivalence of categories, with $M \mapsto M(-1)$ as a quasi-inverse. We have $M(0) = M$, $M(i+j) = M(i)(j)$ ($\forall i, j \in \mathbb{Z}$) and $M(i) = R(i) \otimes_R M$.

A *free graded R -module of rank 1* is a graded R -module isomorphic to an $R(i)$, for some $i \in \mathbb{Z}$. A graded R -module M is called *free* if it can be written as a direct sum of free R -modules of rank 1. For any graded R -module M there is a canonical bijection

$$\text{Hom}(R(i), M) \xrightarrow{\sim} \Gamma(M^{-i}),$$

sending a morphism $R(i) \rightarrow M$ to the image of $1 \in \Gamma(R(i)^{-i}) = R$. Thus an R -Module can be written as a quotient of a free graded R -module if and only if it is generated by global sections (this is always the case, if the topos is trivial!).

If R is a graded ring, and $I \subset R$ is a graded ideal, i.e., generated by homogeneous elements, then R/I is naturally a graded ring. We can apply this to the category of gauges: if R_0 is a ring (in \mathcal{T}), then the category of R_0 -gauges in \mathcal{T} is equivalent to the category of graded $D(R_0)$ -modules, where $D(R_0) = R_0[f, v]/(fv - p)$ is the graded ring (in \mathcal{T}) defined similarly as in section 1.1. Therefore there is a natural tensor product $M \otimes N$ of R_0 -gauges M and N in \mathcal{T} , defined as the tensor product $M \otimes_{D(R_0)} N$.

One easily sees that, for R_0 -gauges M and N , one has canonical isomorphisms

$$(M \otimes_{D(R_0)} N)^{+\infty} \cong M^{+\infty} \otimes_{R_0} N^{+\infty} \text{ and } (M \otimes_{D(R_0)} N)^{-\infty} \cong M^{-\infty} \otimes_{R_0} N^{-\infty}.$$

As a consequence, if σ is an endomorphism of the category of R_0 -modules, and is a tensor morphism, there is a canonical tensor product on the category of φ -modules of R_0 -modules with respect to σ , by endowing $M \times N$ with the following φ :

$$\sigma(M^{+\infty} \otimes_{R_0} M^{+\infty}) \cong \sigma(M^{+\infty}) \otimes_{R_0} \sigma(M^{+\infty}) \xrightarrow{\varphi_M \otimes \varphi_N} M^{-\infty} \otimes_{R_0} N^{-\infty}.$$

Obviously, this respects the subcategory of φ -gauges.

2.4 φ -rings, and φ -modules and φ -gauges over them

We need a certain generalization of the considerations in the previous section. As there, we consider objects (rings, modules, etc.) in some topos \mathcal{T} , and suppress the mentioning of \mathcal{T} . Consider a triple (R, f, v) where $R = \bigoplus_{n \in \mathbb{Z}} R^n$ is a \mathbb{Z} -graded commutative ring (in \mathcal{T}) with $f \in \Gamma(R^1)$ and $v \in \Gamma(R^{-1})$. Set

$$R^{+\infty} = R/(f - 1) \quad \text{and} \quad R^{-\infty} = R/(v - 1)$$

Observe that we may identify these two rings, as R^0 -modules, to the direct limits

$$R^{+\infty} = \varinjlim_{r \rightarrow +\infty} R^r \quad \text{and} \quad R^{-\infty} = \varinjlim_{r \rightarrow -\infty} R^r$$

(the transition maps being given by multiplication by f (resp v)).

We define a φ -ring as a quadruple (R, f, v, φ) with (R, f, v) as above and

$$\varphi : R^{+\infty} \longrightarrow R^{-\infty}$$

a morphism of rings. If this is an isomorphism, we call (R, f, v, φ) a perfect φ -ring.

Let $R = (R, f, v, \varphi)$ a φ -ring. If M is a graded R -module, we may consider the $R^{+\infty}$ -module $M^{+\infty} = R^{+\infty} \otimes_R M = M/(f - 1)M$ and the $R^{-\infty}$ -Module $M^{-\infty} = R^{-\infty} \otimes_R M = M/(v - 1)M$. Observe that, as R^0 -modules, we also have the identifications

$$M^{+\infty} = \varinjlim_{r \rightarrow +\infty} M^r \quad \text{and} \quad M^{-\infty} = \varinjlim_{r \rightarrow -\infty} M^r$$

A φ - R -module is a pair (M, φ) where M is a graded R -module and

$$\varphi : M^{+\infty} \longrightarrow M^{-\infty}$$

is a morphism of groups such that $\varphi(\lambda x) = \varphi(\lambda)\varphi(x)$ for $\lambda \in R^{+\infty}$ and $x \in M^{+\infty}$.

A φ - R -gauge is a φ - R -module (M, φ_M) such that the canonical morphism of $R^{-\infty}$ -modules

$$\varphi'_M : R^{-\infty} \otimes_{\varphi_R R^{+\infty}} M^{+\infty} \longrightarrow M^{-\infty}$$

induced by φ_M is an isomorphism. If (R, φ_R) is a perfect φ -ring, this holds if and only if φ_M is an isomorphism.

With obvious definitions of morphisms, the graded R -modules and the φ - R -modules are abelian categories with enough injectives. As a full sub-category of the category of φ - R -modules, the category of φ - R -gauges is stable under direct sums and direct factors. If R is a perfect φ -ring, it is also stable under kernels and cokernels and therefore also abelian (here we use the exactness of the formation of $M^{+\infty}$ and $M^{-\infty}$).

If M and N are two φ - R -modules, we have

$$(M \otimes_R N)^{+\infty} = M^{+\infty} \otimes_{R^{+\infty}} N^{+\infty} \quad \text{and} \quad (M \otimes_R N)^{-\infty} = M^{-\infty} \otimes_{R^{-\infty}} N^{-\infty}.$$

Therefore, the morphism $\varphi \otimes \varphi$ endows $M \otimes_R N$ with a structure of φ - R -Module (which is a φ - R -gauge if M and N are φ - R -gauges). With this tensor product, φ - R -modules and φ - R -gauges become tensor categories.

For any graded R -module M and any $i \in \mathbb{Z}$, we have $M(i)^{+\infty} = M^{+\infty}$ and $M(i)^{-\infty} = M^{-\infty}$. This allows us to extend the definition of Tate twists to φ - R -modules and φ - R -gauges in an obvious way.

3 GAUGES OVER A PERFECT FIELD

3.1 Preliminaries

Assume R is a graded ring with $f \in R^1$ and $v \in R^{-1}$ such that $R = R^0[f, v]$. If fv is not a zero divisor in R^0 , the natural maps $R^0 \rightarrow R^{+\infty}$ and $R^0 \rightarrow R^{-\infty}$ are isomorphisms and to give a map $\varphi : R^{+\infty} \rightarrow R^{-\infty}$ such that (R, f, v, φ) is a perfect φ -ring is the same as giving an automorphism φ of the ring R^0 .

If moreover fv is invertible in R^0 , the correspondence $M \mapsto M^0$ induces an equivalence between the category of graded R -modules and the category of R^0 -modules. Let $R^0[\varphi]$ the (non-commutative if $\varphi \neq id_{R^0}$) ring generated by R^0 and an element φ , with the relation that $\varphi\lambda = \varphi(\lambda).\varphi$, for $\lambda \in R^0$. Similarly, let $R^0[\varphi, \varphi^{-1}]$ the ring generated over $R^0[\varphi]$ by an element φ^{-1} with the relations $\varphi.\varphi^{-1} = \varphi^{-1}\varphi = 1$ and $\lambda\varphi^{-1} = \varphi^{-1}\varphi(\lambda)$ for $\lambda \in R^0$. The previous equivalence of categories induces an equivalence between

- φ - R -modules and left $R^0[\varphi]$ -modules,
- φ - R -gauges and left $R^0[\varphi, \varphi^{-1}]$ -modules.

In these equivalences, the tensor product becomes the tensor product over R^0 .

The situation is slightly more complicated when fv is not invertible. This is the situation for our generalized Dieudonné modules : We chose a perfect field k of characteristic p and we let $W = W(k)$ be the ring of Witt vectors over k and, for any $n \in \mathbb{N}$, we let $W_n = W_n(k) = W/p^n$ be the ring of n -th truncated Witt vectors. To cover both cases, we write W_n for $n \in \mathbb{N} \cup \{\infty\}$, where $W_\infty := W$. Then W_n -gauges, i.e., gauges of W_n -modules, are simply graded modules over the ring $D_n = D_n(k) = W_n[f, v]/(fv - p)$.

We turn D_n into a perfect φ -ring by taking for φ the absolute Frobenius $\sigma : D_n^{+\infty} = W_n \rightarrow W_n = D_n^{-\infty}$, which is an isomorphism by perfectness of k .

Hence a φ - W_n -module is a W_n -gauge (M, f, v) together with a group homomorphism $\varphi : M^{+\infty} \rightarrow M^{-\infty}$ which is semi-linear with respect to the absolute Frobenius σ on W_n .

We say that a W_n -gauge or φ - W_n -gauge M is of finite type, if the associated D_n -module is finitely generated. As we have seen in section 2.1, M is then

concentrated in a finite interval $[a, b]$ and is just given by the finite diagram of finitely generated W_n -modules

$$M^a \rightleftarrows \dots \rightleftarrows M^{r-1} \begin{matrix} \xrightarrow{f} \\ \xleftarrow{v} \end{matrix} M^r \begin{matrix} \xrightarrow{f} \\ \xleftarrow{v} \end{matrix} M^{r+1} \rightleftarrows \dots \rightleftarrows M^b,$$

such that $fv = vf = p$ (For $r > b$, we use the multiplication by f^{b-r} to identify M^b to M^r . Similarly, for $r < a$, we use multiplication by v^{a-r} to identify M^a to M^r). The structure of a φ -module is obtained by adding a σ -semi-linear map $\varphi : M^b \rightarrow M^a$ (because, we have canonical identifications $M^{+\infty} = M^b$ and $M^{-\infty} = M^a$). The φ -module M will be a φ -gauge if and only if $\varphi : M^b \rightarrow M^a$ is bijective.

Let $\mathcal{G}_{ft}^{[a,b]}(W_n)$ be the category of finite-type W_n -gauges which are concentrated in $[a, b]$. If M is an object of $\mathcal{G}_{ft}^{[a,b]}(W_n)$ and N an object of $\mathcal{G}_{ft}^{[a',b']}(W_n)$, then $M \otimes N$ is an object of $\mathcal{G}_{ft}^{[a+a',b+b']}(W_n)$. If M is an object of $\mathcal{G}_{ft}^{[a,b]}(W_n)$ and $i \in \mathbb{Z}$, then $M(i)$ is an object of $\mathcal{G}_{ft}^{[a-i,b-i]}(W_n)$.

3.2 The standard construction: p -divisibility of Frobenius

The idea of gauges is related to the following construction, going back to ideas of Mazur and Kato. Let B be the fraction field of W .

Let D be an isocrystal over k , i.e., a finite dimensional B -vector space with a σ -semi-linear isomorphism $\phi : D \rightarrow D$, and let M be a lattice in D , i.e., a finitely generated W -submodule with $M \otimes_W B \cong D$. We call such object a virtual crystal over k (and a crystal if $\phi(M) \subset M$). For $r \in \mathbb{Z}$ define

$$M^r = \{ m \in M \mid \phi(m) \in p^r M \},$$

let $f : M^r \rightarrow M^{r+1}$ be the multiplication by p , and let $v : M^{r+1} \rightarrow M^r$ be the inclusion. Then (M^r, f, v) is a W_n -gauge. Moreover, by finite generation of M one has integers $a \leq b$ with $p^b M \subseteq \phi(M) \subseteq p^a M$. The last inclusion implies that $M \subset M^a$ and hence the inclusions $M^r \subset M \subset M^a \subset M$ are isomorphisms for $r \leq a$. The first inclusion implies that $M^r = p^{r-b} M^b$ for $r \geq b$. In fact, if $x \in M^r$, i.e., $\phi(x) = p^r y = p^{r-b} p^b y$ with $y \in M$, then $p^b y = \phi(z)$ with $z \in M$. This implies $\phi(x) = \phi(p^{r-b} z)$ and hence $x = p^{r-b} z$, where $z \in M^b$. We conclude that the gauge is concentrated in the interval $[a, b]$.

Moreover, we get a canonical structure of a φ - W -gauge. In fact, we have natural σ -semi-linear homomorphisms

$$\varphi_r : M^r \rightarrow M = M^{-\infty},$$

by sending $x \in M^r$ to $p^{-r} \phi(x)$. These are compatible ($\varphi_{r+1}(fx) = p^{-r-1} \varphi(p^r x) = p^{-r} \varphi(x) = \varphi_r(x)$) and thus define a σ -semi-linear morphism

$$\varphi : M^{+\infty} = \varinjlim_{r \rightarrow +\infty} M^r \longrightarrow M^{-\infty},$$

which is easily seen to be an isomorphism. Moreover, one sees

THEOREM 3.2.1. *The above construction gives a fully faithful embedding of categories*

$$\begin{aligned} & (\text{virtual crystals } (D, \phi, M) \text{ over } k) \longrightarrow \\ & (\text{finite type } \varphi\text{-}W\text{-gauges with free components}). \end{aligned}$$

Now we want to characterize the essential image of this functor. If (M, f, v) is a gauge, then we let

$$f_r : M^r \longrightarrow M^{+\infty} \quad \text{and} \quad v_r : M^r \longrightarrow M^{-\infty}$$

be the canonical morphisms into the respective inductive limits. We introduce the following definitions, which will also be of use later.

DEFINITION 3.2.2. *Let \mathcal{A} be an \mathbb{F}_p -linear abelian category. A gauge (M, f, v) in \mathcal{A} is called*

(a) *strict, if the morphism*

$$(f_r, v_r) : M^r \longrightarrow M^{+\infty} \oplus M^{-\infty}$$

is a monomorphism for all $r \in \mathbb{Z}$,

(b) *quasi-rigid, if the sequence*

$$M^r \xrightarrow{f} M^{r+1} \xrightarrow{v} M^r \xrightarrow{f} M^{r+1}$$

is exact for all $r \in \mathbb{Z}$,

(c) *rigid, if M is strict and quasi-rigid.*

We note that, in general, the notion of (quasi-)rigidity makes sense only if the objects are annihilated by p , since $vf = p = vf$.

LEMMA 3.2.3. *Let M be a quasi-rigid gauge, and assume that one M^s has finite length. Then all M^r have finite length and have the same length.*

Proof. For the morphisms $f^{(r)} : M^r \rightarrow M^{r+1}$ and $v^{(r+1)} : M^{r+1} \rightarrow M^r$ we have exact sequences

$$0 \rightarrow \text{im}(v^{(r+1)}) \hookrightarrow M^r \xrightarrow{f} \text{im}(f^{(r)}) \rightarrow 0;$$

$$0 \rightarrow \text{im}(f^{(r)}) \hookrightarrow M^{r+1} \xrightarrow{v} \text{im}(v^{(r+1)}) \rightarrow 0.$$

This implies that M^r and M^{r+1} have the same length, hence the claim. \square

Now we consider gauges over a field k of characteristic $p > 0$. We have the following Nakayama-type lemma:

LEMMA 3.2.4. *Let M be a gauge of finite type over k . If $M/(f, v)M = 0$, then $M = 0$. As a consequence, if m_1, \dots, m_r are homogeneous elements in M whose residue classes generate $M/(f, v)$ as a k -vector space, then these elements generate M (as a $D(k) = k[f, v]/(fv)$ -module).*

Proof. Assume that $M/(f, v)M = 0$ and let $m \in M^s$ for some s . Then there exist elements $m_{s-1}^1 \in M^{s-1}$ and $m_{s+1}^2 \in M^{s+1}$ with $m = fm_{s-1}^1 + vm_{s+1}^2$. By induction, and noting that $fv = vf = p = 0$, for each $n > 0$ we get elements $m_{s-n}^1 \in M^{s-n}$ and $m_{s+n}^2 \in M^{s+n}$ with

$$m = f^n m_{s-n}^1 + v^n m_{s+n}^2.$$

We see that $m = 0$, since $f = 0$ on M^{s-n} for $n \gg 0$ and $v = 0$ on M^{s+n} for $n \gg 0$, because v is an isomorphism on M^r for $r \ll 0$ and f is an isomorphism on M^r for $r \gg 0$, and $fv = vf = 0$. The second claim follows in a standard way, by looking at M/N , where N is the sub- $D(k)$ -module generated by m_1, \dots, m_r . \square

We derive from this the following criterion.

LEMMA 3.2.5. *Let M be a k -gauge of finite type. Then the following are equivalent.*

- (a) M is free.
- (b) M is rigid.
- (c) The maps $M^r/v \xrightarrow{f} M^{r+1}/v$ and $M^r/f \xleftarrow{v} M^{r+1}/f$ are injective for all r .

Here we have used the short notation M^r/v for $(M/(v))^r$ or, explicitly, M^r/vM^{r+1} ; similarly for M^r/f .

Proof. Obviously, (b) holds for the free gauge $k = k(0)$ (see the definition in section 2.3), i.e., the module $M = D = k[f, v]/(fv)$, where M/v is a free $k[f]$ -module and M/f is a free $k[v]$ -module. In fact, this gauge corresponds to the diagram

$$\dots \begin{array}{cccc} 0 & 0 & id & id \\ \xrightarrow{\quad} & k & \xrightarrow{\quad} & k & \xrightarrow{\quad} & k & \xrightarrow{\quad} & \dots \\ id & id & 0 & 0 \end{array},$$

where the middle k is placed in degree 0. One immediately sees strictness ($\ker(v) \cap \ker(f) = 0$) and quasi-rigidity ($\ker(f) = \text{im}(v)$ and $\ker(v) = \text{im}(f)$) at all places. Thus (b) holds for gauges $k(i)$ by degree shifting, and for arbitrary free gauges by taking sums.

On the other hand, (b) implies (c). In fact, for the injectivity of $M^r/v \xrightarrow{f} M^{r+1}/v$ assume that $f(x) = v(y) =: a$ for $x \in M^r$ and $y \in M^{r+2}$. By rigidity we have $0 = \ker(v) \cap \ker(f) = \text{im}(f) \cap \text{im}(v)$, so that $a = 0$. Since $\ker(f) = \text{im}(v)$, $f(x) = 0$ implies $x \in \text{im}(v)$ as claimed. The injectivity of $M^{r+1}/f \xrightarrow{v} M^r/f$ follows dually: If $v(y) = f(x)$, then $y \in \ker(v) = \text{im}(f)$. We also note that (c) immediately implies that M is quasi-rigid: We have a factorization $M^r/v \xrightarrow{f} M^{r+1} \twoheadrightarrow M^{r+1}/v$, so that (c) implies $\ker(f) = \text{im}(v)$. Similarly, (c) implies $\ker(v) = \text{im}(f)$.

Finally we show that (c) implies (a). We may assume that M is concentrated in the finite interval $[a, b]$, and then (c) gives a sequence of injections of finite-dimensional k -vector spaces

$$\dots \rightarrow 0 \rightarrow M^a/v \hookrightarrow M^{a+1}/v \hookrightarrow \dots \hookrightarrow M^b/v = M^b \xrightarrow[\sim]{f} M^{b+1} \xrightarrow[\sim]{f} \dots$$

Note that $M^{r-1} \xleftarrow{v} M^r$ is an isomorphism for $r \leq a$, and that $M^r \xleftarrow{v} M^{r+1}$ is zero for $r \geq b$, because the map f in the other direction is bijective, and $fv = p = 0$. Now take a k basis $m_1^a, \dots, m_{d_a}^a$ of M^a/v , a k -basis $m_1^{a+1}, \dots, m_{d_{a+1}}^{a+1}$ of $(M^{a+1}/v)/f(M^a/v) = M^{a+1}/(v, f)$ etc. up to a k -basis m_1^b, \dots of $M^b/(f, b)$, and lift these elements to elements $\hat{m}_1^a, \dots, \hat{m}_{d_a}^a, \hat{m}_1^{a+1}, \dots$ in M^a, M^{a+1} etc. Then the universal property of the free gauges $k(i)$ (see section 2.3) gives a morphism of k -gauges

$$g : F = \bigoplus_{i=a}^b k(-i)^{d_i} \longrightarrow M$$

mapping the canonical elements of $k(-i)^i$ to the elements $\hat{m}_1^i, \dots, \hat{m}_{d_i}^i$. Since this map is surjective modulo (f, v) , it is surjective by lemma 3.2.4. Moreover, by construction the map $F^b \rightarrow M^b/v = M^b$ is bijective: both spaces have dimension $d = d_a + \dots + d_b$. But, as remarked above, (c) implies that M is quasi-rigid, so that each M^r has dimension d , and the same is true for each F^r . Therefore the surjective map g is an isomorphism, and we have shown (a). \square

We draw the following consequences for $W(k)$ -gauges for a perfect field k .

COROLLARY 3.2.6. *Let k be a perfect field of characteristic $p > 0$, and let M be gauge of finite type over $W = W(k)$. If $M/(p, f, v)M = 0$, then $M = 0$. Consequently, M is generated by homogeneous elements m_1, \dots, m_r if and only if their residue classes generate $M/(p, f, v)M$.*

Proof. This follows from lemma 3.2.4 by the usual Nakayama lemma for the local ring W , because the components M^s of M are finitely generated W -modules. \square

Next we characterize free W -gauges of finite type. Obviously, their components are free W -modules. But this condition does not suffice.

THEOREM 3.2.7. *Let M be a W -gauge of finite type with free components. Then the following conditions are equivalent.*

- (a) M is a free W -gauge.
- (b) $N = M/pM$ is a free k -gauge.
- (c) The map $M^r/v \xrightarrow{f} M^{r+1}/v$ is injective for all r .
- (d) The map $M^{r+1}/f \xrightarrow{v} M^r/f$ is injective for all r .

Proof. (a) trivially implies (b), but (b) also implies (a): Assume $N = M/pM$ is free, say isomorphic to $\oplus_i k(i)^{d_i}$. By the universal property of free gauges (= free modules over $D(k)$ and $D(W)$, respectively) we can lift the isomorphism modulo p to a morphism $F = \oplus_i W(i)^{d_i} \rightarrow M$, which is surjective by corollary 3.2.6. Since this map is an isomorphism modulo p , and all components are free W -modules, it is an isomorphism.

Next we remark that the maps in (c) and (d) can be identified with the maps

$$N^r/v \xrightarrow{f} N^{r+1}/v \quad \text{and} \quad N^{r+1}/f \xrightarrow{v} N^r/f,$$

respectively, because pM is contained in both fM and vM , by the equality $fv = vf = p$.

Therefore (b) is equivalent to the conjunction of (c) and (d), by Lemma 3.2.5. But (c) and (d) are in fact equivalent in our situation: Assume (c). To show the injectivity in (d) let $y \in M^{r+1}$ with $v(y) = f(x)$, where $x \in M^{r-1}$. Then (c) implies $x = v(z)$ with $z \in M^r$. We get $v(y) = f(v(z)) = v(f(z))$, and hence that $y = f(z)$, because v is injective ($fv = p$, and M is torsion-free as W -module). A similar reasoning shows that also (d) implies (c), again since M is a torsion-free W -module. Therefore properties (a) to (d) are equivalent. \square

COROLLARY 3.2.8. *A W -gauge of finite type M is free if and only if it comes from a virtual crystal over k , i.e., is the underlying gauge of a φ - W -gauge in the essential image of the functor in theorem 3.2.1.*

Proof. Assume that M comes from the virtual k -crystal (D, ϕ, L) . Then M has free components, and we show condition (c) in theorem 3.2.7. We have

$$M^r = \{ x \in L \mid \phi(x) \in p^r L \}, \quad f(x) = px, \quad \text{and} \quad v(x) = x,$$

by definition. Now let $x \in M^r$ with $f(x) = v(y)$ for $y \in M^{r+2}$. Then, by definition, we have $x, y \in L$ satisfying $\phi(x) = p^r z$ with $z \in L$ and $\phi(y) = p^{r+2}t$ with $t \in L$, and $px = y$. Then $p^{r+1}z = p\phi(x) = \phi(px) = \phi(y) = p^{r+2}t$ and hence $z = pt$, since L is torsion-free. This implies $\phi(x) = p^{r+1}t$, i.e., $x \in vM^{r+2}$. Conversely we show that any free W -gauge arises from a virtual crystal. By considering sums in both categories, we may consider the case $M = W(i)$ for some $i \in \mathbb{Z}$. But this gauge arises from the virtual crystal (B, ϕ, W) where $\phi(b) = p^{-i}\sigma(b)$. \square

We can now strengthen this result and characterize the image of the functor in theorem 3.2.1.

THEOREM 3.2.9. *A φ - W -gauge (M, f, v, φ) of finite type comes from a virtual crystal over k , i.e., lies in the essential image of the functor in theorem 3.2.1, if and only if the underlying gauge is a free W -gauge.*

Proof. One direction follows from corollary 3.2.8. For the other direction assume that (M, f, v) is free. Assume that M is concentrated in the finite interval

$[a, b]$. The functor in theorem 3.2.1 is compatible with twists: If ϕ is multiplied by p^i , then the associated gauge M is replaced by $M(-i)$. Therefore we may assume that $a = 0$. Then the φ - W -gauge corresponds to the finite diagram

$$M^0 \begin{array}{c} \xrightarrow{f} \\ \xleftarrow{v} \end{array} \dots \begin{array}{c} \xrightarrow{f} \\ \xleftarrow{v} \end{array} M^{r-1} \begin{array}{c} \xrightarrow{f} \\ \xleftarrow{v} \end{array} M^r \begin{array}{c} \xrightarrow{f} \\ \xleftarrow{v} \end{array} M^{r+1} \begin{array}{c} \xrightarrow{f} \\ \xleftarrow{v} \end{array} \dots \begin{array}{c} \xrightarrow{f} \\ \xleftarrow{v} \end{array} M^b,$$

together with a σ -semi-linear isomorphism

$$\varphi : M^{+\infty} = M^b \xrightarrow{\sim} M^0 = M^{-\infty}.$$

All maps f and v are injective, because $fv = vf = p$. Let $L = M^0$, and define the σ -linear endomorphism

$$\phi = \varphi \circ f^b : L = M^0 \xrightarrow{f^b} M^b \xrightarrow{\varphi} M^0 = L.$$

Then (L, φ) is a crystal over k , and we claim that the associated φ - W -gauge is canonically isomorphic to (M, f, v, φ) . In fact, first we claim that, for $0 \leq r \leq b$, the injective map $v^r : M^r \rightarrow L$ has the image $L^r = \{x \in L \mid \phi(x) \in p^r L\}$. First of all, we have $\phi(v^r x) = \varphi(f^b v^r x) = \varphi(p^r f^{b-r} x) \in p^r L$. Conversely, by the assumption and criterion (c) in theorem 3.2.7, all maps

$$M^0/vM^1 \xrightarrow{f} M^1/vM^2 \xrightarrow{f} M^2/vM^3 \xrightarrow{f} \dots$$

are injective. If now $x \in L = M^0$ with $\phi(x) = \varphi(f^b x) = p^r y$ with $y \in L$, then $f^b x = p^r \varphi^{-1} y = f^r v^r \varphi^{-1} y$ and hence $f^{b-r} x = v^r \varphi^{-1} y$ by injectivity of f . By the sequence of injective maps above this implies inductively $x = vx_1$ with $x_1 \in M^1$, hence $f^{b-r} vx_1 = v^r \varphi^{-1} y$, hence $f^{b-r} x_1 = v^{r-1} \varphi^{-1} y$ by injectivity of v on M , hence $x_1 = vx_2$ with $x_2 \in M^2$ etc. and inductively $x = v^r x_r$ with $x_r \in M^r$. Thus we have $L^r = v^r M^r$ as claimed.

Identifying L^r with M^r via v^r the maps v become inclusions, and the maps f become multiplication by p , because $fv = p$. Finally one sees that the map $\varphi : M^b \rightarrow M^0$ identifies with $p^{-b} \phi : L^b \rightarrow L^0$, sending y to $p^{-b} \phi(y)$ as in the construction of the φ -gauge associated to (L, ϕ) . \square

COROLLARY 3.2.10. *The functor in theorem 3.2.1 induces an equivalence of categories*

$$\begin{aligned} & (\text{virtual crystals } (D, \phi, M) \text{ over } k) \longrightarrow \\ & \hspace{10em} (\text{finite type free } \varphi\text{-}W\text{-gauges } (M, f, v, \varphi)), \end{aligned}$$

where we call a φ - W -gauge free if the underlying gauge is free.

REMARK 3.2.11 Let M be a W -gauge of finite type with free components. If M is concentrated in a point, i.e., in an interval $[a, a]$, then it is obviously free, viz., isomorphic to $W(-a)^d$ for some d . If M is concentrated in an interval

of length 1, it is free as well. For this we use criterion (c) from theorem 3.2.7. If M is represented by

$$M^a \begin{matrix} \xrightarrow{f} \\ \xleftarrow{v} \end{matrix} M^{a+1},$$

we only have to show the injectivity of $f : M^a/vM^{a+1} \rightarrow M^{a+1}/vM^{a+2}$; the injectivity at the other places is clear. But the image of $v : M^{a+2} \rightarrow M^{a+1}$ is pM^{a+1} , and if $x \in M^a$ and $fx = vy = pz = fvz$ with $z \in M^{a+1}$, then $x = vz$ by injectivity of f .

But already if M is concentrated in an interval of length 2, M is not in general free. A counterexample is the W -gauge

$$N \begin{matrix} \xrightarrow{p} \\ \supset \end{matrix} pN \begin{matrix} \xrightarrow{p} \\ = \end{matrix} pN,$$

for any free W -module $N \neq 0$, since here $N/v = N/p \rightarrow pN/v = 0$ is not injective.

3.3 Gauges and Dieudonné modules.

A Dieudonné-module of finite type over k is a W -module of finite type M endowed with two additive endomorphisms F and V such that

$$FV = VF = p \text{ and } F(\lambda x) = \sigma(\lambda)Fx, \quad V(\sigma(\lambda)x) = \lambda Vx \quad (\forall \lambda \in W, \forall x \in M)$$

With an obvious definition of morphisms, Dieudonné modules of finite type over k form a \mathbb{Z}_p -linear abelian category $Dieud(k)$.

Let M be an object of the category $\varphi\mathcal{G}_{ft}^{[-1,0]}(W)$ of finite-type φ - W -gauges which are concentrated in $[-1,0]$. To give M is the same as giving $(M^{-1}, M^0, f, v, \varphi)$ where M^{-1} and M^0 are W -modules of finite type, $f : M^{-1} \rightarrow M^0$ and $v : M^0 \rightarrow M^{-1}$ are W -linear maps such that $fv = pid_{M^0}$ and $vf = pid_{M^{-1}}$ and $\varphi : M^0 \rightarrow M^{-1}$ is a bijective φ -linear map. We define $F, V : M^{-1} \rightarrow M^{-1}$ by $F = \varphi f$ and $V = v\varphi^{-1}$. This turns M^{-1} into a Dieudonné module of finite type over k . In this way, we get a functor

$$\varphi\mathcal{G}_{ft}^{[-1,0]}(W) \longrightarrow Dieud(k),$$

which is an equivalence of categories.

Dieudonné-modules arise from p -divisible groups over k or from the first crystalline cohomology of smooth projective varieties X over k . By the theory of the de Rham-Witt complex, the i -th crystalline cohomology of X gets the structure of what could be called a ‘Dieudonné-module of weight i ’: a finitely generated W -module M together with a σ -linear endomorphism F and a σ^{-1} -linear endomorphism V such that $FV = VF = p^i$. Such a structure can also

be obtained by a φ -gauge with free components which is concentrated in an interval of length i ,

$$M^a \rightleftarrows \dots \rightleftarrows M^{r-1} \begin{array}{c} \xleftarrow{f} \\ \xrightarrow{v} \end{array} M^r \begin{array}{c} \xleftarrow{f} \\ \xrightarrow{v} \end{array} M^{r+1} \rightleftarrows \dots \rightleftarrows M^{a+i} \xrightarrow[\sim]{\varphi} M^a,$$

by letting $M = M^a$, $F = \varphi f^i$, and $V = v^i \varphi^{-1}$. This gives a functor

$$\mathcal{G}_{ft}^{[a, a+i]}(W) \longrightarrow \text{Dieud}^i(k),$$

where $\text{Dieud}^i(k)$ is the category of Dieudonné-modules of weight i , whose morphisms are linear maps compatible with F and V . However, for $i > 1$ this functor is no longer an equivalence of categories, because it forgets all information concerning the modules $M^{a+1}, \dots, M^{a+i-1}$.

One aim of this paper is to establish a canonical cohomology theory giving φ - W -gauges of finite type $H_g^i(X/W)^\bullet$ for each i , concentrated in the interval $[0, i]$, whose associated Dieudonné-module of weight i is the i -th crystalline cohomology $H_{cris}^i(X/W)$. This new cohomology theory thus refines the crystalline cohomology.

3.4 Effective, coeffective modules and truncations

We say that a gauge M over $W_n = W_n(k)$ (for $1 \leq n \leq \infty$) is *effective* (resp. *coeffective*) if $v : M^r \rightarrow M^{r-1}$ is an isomorphism for $r \leq 0$ (resp. $f : M^r \rightarrow M^{r+1}$ is an isomorphism for $r \geq 0$).

For any object M of $\mathcal{G}_{ft}(W_n)$, $M(i)$ is effective for $i \ll 0$ and coeffective for $i \gg 0$.

To any W_n -gauge M , we may associate the co-effective W_n -gauge $M_{\leq 0}$ defined as follows: we have $M_{\leq 0}^r = M^r$ for $r < 0$ and $M_{\leq 0}^r = M^0$ for $r \in \mathbb{N}$, with $fx = x$ if $x \in M_{\leq 0}^r$ and $r \geq 0$ and $vx = px$ if $x \in M_{\leq 0}^r$ and $r > 0$.

If M is of finite type, so is $M_{\leq 0}$ and we may view $M \mapsto M_{\leq 0}$ as a functor from $\mathcal{G}_{ft}(W_n)$ to the full sub-category $\mathcal{G}_{ft}^{\leq 0}(W_n)$ of coeffective gauges of finite type, which is a right adjoint of the inclusion functor. We observe that the obvious map $M_{\leq 0} \rightarrow M$ is not in general injective.

For any W_n -gauge M , the natural maps $M^0 \rightarrow (M_{\leq 0})^{+\infty}$ and $(M_{\leq 0})^{-\infty} \rightarrow M^{-\infty}$ are isomorphisms. Therefore, we may also view $M \mapsto M_{\leq 0}$ as a functor from the category $\varphi\text{-}\mathcal{M}_{ft}(W_n)$ of φ - W_n -modules of finite type to the full sub-category $\varphi\text{-}\mathcal{M}_{ft}^{\leq 0}(W_n)$ of coeffective φ - W_n -modules, by defining $\varphi : M_{\leq 0}^{+\infty} \rightarrow M_{\leq 0}^{-\infty}$ as the compositum φ^0 of the natural map $M^0 \rightarrow M^{+\infty}$ with the original φ .

Again this functor is a right adjoint of the inclusion functor. We observe that, when M is a φ -gauge, $M_{\leq 0}$ is not always a φ -gauge.

4 ZARISKI-GAUGES AND F -ZIPS OVER SCHEMES OF CHARACTERISTIC p

Let S be a scheme of characteristic p .

4.1 Zariski-gauges and Zariski- φ -gauges

The φ -ring associated to S in the Zariski topology is defined as the commutative ring (in this topology) $D(S) = \mathcal{O}_S[f, v]/(fv)$ together with the ring morphism $\varphi : D(S)^{+\infty} = \mathcal{O}_S \rightarrow \mathcal{O}_S = D(S)^{-\infty}$ which is given by the absolute Frobenius $F = F_S$ on \mathcal{O}_S (which is the identity on S and the map $x \mapsto x^p$ on the sections. Note that this is only an isomorphism if S is a perfect scheme, so $D(S)$ is not in general a perfect φ -ring.

A Zariski gauge M over S is a $D(S)$ -module, which is obviously just a gauge in the category of \mathcal{O} -modules. It is called coherent if it is locally of finite presentation over $D(S)$. Hence, if S is noetherian, it is coherent if and only if M is concentrated in a finite interval and each component M^r is a coherent \mathcal{O}_S -module.

In accordance with the definitions in section 2.4, a Zariski φ -module (M, φ) is a Zariski gauge M together with an \mathcal{O}_S -linear morphism

$$\varphi : (M^{+\infty})^{(p)} = \mathcal{O}_{S \text{ Fr} \otimes_{\mathcal{O}_S}} M^{+\infty} \rightarrow M^{-\infty} .$$

Here $\mathcal{N}^{(p)} = \mathcal{O}_{S \text{ Fr} \otimes_{\mathcal{O}_S}} \mathcal{N}$ is the usual Frobenius twist (twist with the absolute Frobenius Fr) of an \mathcal{O}_S -module \mathcal{N} . Since this operation is a right exact functor, the φ - \mathcal{O}_S -modules form an abelian category, see section 2.4. A φ -module is a φ -gauge if φ is an isomorphism. They form a subcategory which is closed under direct sums and direct factors.

4.2 The relationship with F -zips

An F -zip over S is defined as a locally free coherent \mathcal{O}_S -module \mathcal{M} together with

- (a) A descending filtration $C = (C^i)_{i \in \mathbb{Z}}$ on $\mathcal{M}^{(p)}$ by locally direct summands,
- (b) An increasing filtration $D = (D_i)_{i \in \mathbb{Z}}$ on \mathcal{M} by locally direct summands,
- (c) a family $(\varphi_i)_{i \in \mathbb{Z}}$ of \mathcal{O}_S -linear isomorphisms $\varphi_i : C^i/C^{i+1} \xrightarrow{\sim} D_i/D_{i-1}$.

A morphism of F -zips $(\mathcal{M}, C, D) \rightarrow (\mathcal{M}', C', D)$ is an \mathcal{O}_S -linear morphism $\mathcal{M} \rightarrow \mathcal{M}'$ respecting the filtrations. (This is the modified definition in [Wed], improving the original definition of Moonen and Wedhorn [MW]. If S is perfect, then both definitions are equivalent.)

Then one has the following result.

THEOREM 4.2.1. (see [Schn]) *There is a canonical full embedding of categories*

$$(F\text{-zips over } S) \rightarrow (\varphi\text{-}\mathcal{O}_S\text{-gauges}).$$

The essential image consist of the φ - \mathcal{O}_S -gauges which are rigid, coherent, and have locally free components.

5 ZARISKI-GAUGES AND DISPLAYS OVER SCHEMES OF CHARACTERISTIC p

Langer and Zink defined the notion of a display over a ring R of positive characteristic. Let $W(R)$ be the ring of Witt vectors for R , and let $I(R) = VW(R)$, the image of the Verschiebung V on $W(R)$. Then a *predisplay* consists of the following data:

- 1) A chain of morphisms of $W(R)$ -modules

$$\dots \longrightarrow P_{i+1} \xrightarrow{\iota_i} P_i \longrightarrow \dots \longrightarrow P_1 \xrightarrow{\iota_0} P_0,$$

- 2) For each $i \geq 0$ a $W(R)$ -linear map

$$\alpha_i : I_R \otimes_{W(R)} P_i \longrightarrow P_{i+1},$$

- 3) For each $i \geq 0$ a Frobenius-linear map

$$F_i : P_i \longrightarrow P_0.$$

These are required to satisfy the following conditions: The composition $\iota_i \circ \alpha_i$ is the multiplication $I_R \otimes P_i \longrightarrow P_i$, and one has

$$F_{i+1}(\alpha_i(V(\eta) \otimes x)) = \eta F_i x, \quad \text{for } \eta \in I_R, x \in P_i.$$

Predisplays form an abelian category, in an obvious way.

Finally, a predisplay is called a *display* of degree d if there are finitely generated projective $W(R)$ -modules L_0, \dots, L_d such that

$$P_i = (I \otimes L_0) \oplus \dots \oplus (I \otimes L_{i-1}) \oplus L_i \oplus \dots \oplus L_d,$$

and such that the structural maps ι_i, α_i and F_i come from Frobenius-linear maps

$$\Phi_i : L_i \longrightarrow L_0 \oplus \dots \oplus L_d$$

with the property that $\oplus_i \Phi_i$ is a Frobenius-linear automorphism of $L_0 \oplus \dots \oplus L_d$. (See [LZ] Definition 2.5 for the precise prescription how to get a predisplay out of these data). These data are not part of the datum of a display, only the existence matters. Thus the displays form a full subcategory of the category of predisplays.

Then one has the following result.

THEOREM 5.0.1. (see [Wid]) *Let R be a perfect ring. There is a fully faithful embedding of categories*

$$(\text{predisplays over } R) \longrightarrow (\varphi\text{-}W(R)\text{-modules}),$$

which maps the category of displays to the category of φ - $W(R)$ -modules for which φ is an epimorphism.

6 TOPOLOGIES

Recall that a morphism of schemes is said to be *syntomic* if it is flat and locally a complete intersection (which implies that it is locally of finite type).

We say that a morphism $\text{Spec } B \rightarrow \text{Spec } A$ of affine schemes of characteristic p is an *extraction of p -th root* if one may write $B = A[t]/(t^p - a)$, for some $a \in A$.

We say that a morphism $X \rightarrow Y$ of \mathbb{F}_p -schemes is a *p -root-morphism* (resp. is *quiet*¹) if locally for the Zariski topology (resp. for the étale topology), it may be written as a successive extractions of p -th roots.

In characteristic p , we have the following inclusions :

$$\begin{array}{ccccc} \text{open immersions} & \subset & \text{étale morphisms} & \subset & \text{flat morphisms} \\ \cap & & \cap & & \cup \\ p\text{-morphisms} & \subset & \text{quiet morphisms} & \subset & \text{syntomic morphisms} \end{array}$$

A ring A of characteristic p is *perfect* if the Frobenius $\varphi : A \rightarrow A, a \mapsto a^p$ is bijective. A scheme of characteristic p is perfect if \mathcal{O}_X is a sheaf of perfect rings.

LEMMA 6.0.1. *Let A be a noetherian ring of characteristic p . Then the following holds.*

- (1) *If $\varphi : A \rightarrow A$ is surjective, then A is perfect.*
- (2) *The subring $A_{per} = \bigcap_n \varphi^n(A)$ is a perfect ring.*

Proof. Consider the ideals $A(p^n) = \{ a \in A \mid a^{p^n} = 0 \}$. Since A is noetherian, the ascending sequence $A(p) \subseteq A(p^2) \subseteq A(p^3) \dots$ becomes stationary. Assume that the union of all these ideals is equal to $A(p^N)$ for some $N \geq 1$, say.

(1): Assume that φ is surjective and that $a \in A$ with $a^p = 0$. Then there exists an element $b \in A$ with $b^{p^N} = a$. Hence $b^{p^{N+1}} = a^p = 0$ and thus $0 = b^{p^N} = a$.

(2): Let $x \in A_{per}$. Then there exist elements $x_N, x_{N+1}, x_{N+2}, \dots$ in A such that

$$x = x_N^{p^{N+1}} = x_{N+1}^{p^{N+2}} = x_{N+2}^{p^{N+3}} = \dots$$

This implies $(x_N - x_{N+1}^p)^{p^{N+1}} = 0$ and hence $(x_N - x_{N+1}^p)^{p^N} = 0$, i.e., $x_N^{p^N} = (x_{N+1}^p)^{p^N}$. Setting $y_i = x_i^{p^N}$ for $i \geq N$, we similarly get $y_i = y_{i+1}^p$ for all $i \geq N$, so that $y_N \in A_{per}$ and $x = y_N$. Hence φ is surjective on A_{per} . If now $a \in A_{per}$ with $a^p = 0$, we have an element $b \in A_{per}$ with $a = b^{p^N}$, and we conclude as before, arguing inside A , that $a = 0$. □

As a consequence, if X is a locally noetherian scheme of characteristic p , there is a unique morphism $X \rightarrow X_{per}$ with a perfect scheme X_{per} such that any morphism $X \rightarrow Y$ with Y perfect factors uniquely through X_{per} . We say that a scheme X of characteristic p is *absolutely syntomic* if it is locally noetherian and if the morphism $X \rightarrow X_{per}$ is syntomic.

¹quiet abbreviates quasi-étale

(We do not know whether or not X locally noetherian implies X_{per} locally noetherian, but we do not care).

A field k of characteristic p is absolutely syntomic if and only if k is perfect. If Y is absolutely syntomic and if $X \rightarrow Y$ is syntomic, X is absolutely syntomic. We denote by \mathcal{C} the full subcategory of the category of schemes of characteristic p whose objects are absolutely syntomic schemes.

In this paper, an *admissible class of morphisms of \mathcal{C}* is a class of syntomic morphisms of \mathcal{C} containing all the open immersions and stable under composition and base change. A *p -admissible class of morphisms of \mathcal{C}* is an admissible class containing the extractions of p -th roots.

For any admissible class E , call \mathcal{C}_E the site whose underlying category is \mathcal{C} , with surjective families of E -morphisms (that is of morphisms of \mathcal{C} belonging to E) as coverings.

We set $E = \text{Zar}$ (resp. p , ét, quiet, synt) for the class of open immersions (resp. p -morphisms, étale morphisms, quiet morphisms, syntomic morphisms).

Let X be any object of \mathcal{C} . For any admissible class E of morphisms of \mathcal{C} , we call $\mathcal{C}_{X,E}$ the site whose underlying category is the full subcategory of X -schemes $Y \rightarrow X$ such that Y is an object of \mathcal{C} , with surjective families of E -morphisms as covering. The corresponding sheaves shall be called *sheaves (over \mathcal{C}) for the E -topology*.

If moreover X is noetherian, we call X_{quiet} the site whose underlying category is the full subcategory of X -schemes $Y \rightarrow X$ such that the structural morphism is quiet of finite type, with finite surjective families of quiet morphisms as coverings.

REMARKS 6.0.2: Let k be a field of characteristic p such that the extension $k/\varphi(k)$ is finite.

(1) If k is not perfect, $k_{quiet} = k_{quiet}$ is the *smallest Grothendieck topology able to deal with all finite extensions of k* . More precisely:

- i) any finite extension of k is a quiet k -algebra,
- ii) if $X \rightarrow k$ is a quiet morphism of finite type, there is a surjective quiet morphism of finite type $U \rightarrow X$, such that $U = \text{Spec}(k_1 \otimes_k \otimes_k k_2 \otimes_k \dots \otimes_k k_d)$ with k_1, k_2, \dots, k_d finite fields extensions of k .

(2) If k is perfect, the functor which associates to any finite commutative group scheme over k the sheaf it defines on k_{quiet} induces an equivalence of categories between the category of finite and flat commutative group schemes over k and the category of abelian groups over k_{quiet} which are representable. This is due to the fact that any finite commutative group scheme over k is quiet. Observe that we have a similar statement for fields of characteristic 0 and the étale topology.

7 THE RINGS \mathcal{O}_n^{cris} AND THE φ -RINGS \mathcal{G}_n

We continue to call a ring object in a topos \mathcal{T} simply a ring in \mathcal{T} , or a ring over \mathcal{C} with respect to the topology E if the topos given by the category \mathcal{C} and the topology E .

7.1 The p -adic ring $\mathcal{O}^{cris} = (\mathcal{O}_n)_n$

A p -adic ring R in a fixed topos \mathcal{T} consists of giving, for each $n \in \mathbb{N}$ a commutative ring R_n in \mathcal{T} , together with an isomorphism $R_{n+1}/p^n \simeq R_n$. A p -adic ring R in \mathcal{T} is flat if R_n is flat over \mathbb{Z}/p^n , for all $n \in \mathbb{N}$, i.e., if the sequence

$$R_n \xrightarrow{p} R_n \xrightarrow{p^{n-1}} R_n$$

is exact. In this case, for $m, n \in \mathbb{N}$, we have exact sequences

$$0 \rightarrow R_n \rightarrow R_{n+m} \xrightarrow{p^n} R_{n+m} \rightarrow R_n \rightarrow 0,$$

and, in particular, exact sequences

$$0 \rightarrow R_n \rightarrow R_{n+m} \rightarrow R_m \rightarrow 0.$$

Let $\mathcal{T} = (\mathcal{C}, E)$ be a ringed topos with \mathcal{C} as in section 6, a topology E , and the structural sheaf of rings \mathcal{O} , defined by $\mathcal{O}(X) = \mathcal{O}_X(X)$ for a scheme X in \mathcal{C} . For $n \in \mathbb{N}$, a \mathbb{Z}/p^n -divided power thickening of \mathcal{O} (for the E -topology) is a triple $(\mathcal{G}, \theta, \gamma)$ where \mathcal{G} is a \mathbb{Z}/p^n -ring on \mathcal{C}_E , $\theta : \mathcal{G} \rightarrow \mathcal{O}$ is an epimorphism of rings and γ is a divided power structure on the kernel of θ such that, for any object U of \mathcal{C} , any $x \in \mathcal{G}(U)$ and any $m \in \mathbb{N}$, we have $\gamma_m(px) = (p^m/m!)x^m$. The \mathbb{Z}/p^n -divided power thickenings of \mathcal{O} for the E -topology form, in an obvious way, a category. For $E = p$, this category has an initial object that we call \mathcal{O}_n^{cris} . This can be shown

- either by working on the crystalline site and showing that

$$X \mapsto \mathcal{O}_n^{cris}(X) := H^0((X \rightarrow \text{Spec } W_n(\mathcal{O}_{X_{per}}))_{crys}, \text{structural sheaf})$$

is a solution of the universal problem,

- or by constructing \mathcal{O}_n^{cris} directly as the syntomic sheaf associated to the presheaf

$$X \mapsto W_n^{DP}(X),$$

the divided power envelope of the ring of Witt vectors of X (see [FM]).

Moreover, for any admissible class E , \mathcal{O}_n^{cris} is also a sheaf for the E -topology. Therefore, if E is p -admissible, \mathcal{O}_n^{cris} is also an initial object of the category of $\mathbb{Z}/p^n\mathbb{Z}$ -divided power thickenings of \mathcal{O} for the E -topology.

Under the same assumption on E , the natural morphism $\mathcal{O}_{n+1}^{cris}/p^n \rightarrow \mathcal{O}_n^{cris}$ is an isomorphism and the p -adic Ring $\mathcal{O}^{cris} = (\mathcal{O}_n^{cris})_{n \in \mathbb{N}}$ is flat [FM], i.e., we have natural exact sequences for all n and m

$$0 \rightarrow \mathcal{O}_n^{cris} \rightarrow \mathcal{O}_{n+m}^{cris} \rightarrow \mathcal{O}_m^{cris} \rightarrow 0.$$

The Frobenius $\varphi : a \mapsto a^p$ is an endomorphism of the structural Ring \mathcal{O} . By functoriality, it induces an endomorphism of \mathcal{O}_n^{cris} , that we still denote by φ . This is also an endomorphism of \mathcal{O}^{cris} , i.e. the projection $\mathcal{O}_{n+1}^{cris} \rightarrow \mathcal{O}_n^{cris}$ commutes with φ .

If A is a perfect (noetherian) ring of characteristic p , we have $\mathcal{O}_n^{cris}(A) = W_n(A)$, with the usual Frobenius.

7.2 The φ -Ring \mathcal{G}

For each $n \in \mathbb{N}$, we want to define a φ -ring \mathcal{G}_n , such that the \mathcal{G}_n form a p -adic φ -ring $\mathcal{G} = (\mathcal{G}_n)_{n \in \mathbb{N}}$.

Morally, we get it by the standard construction introduced in section 3.2, from the p -adic ring $\mathcal{O}^{cris} = (\mathcal{O}_n^{cris})$ and the Frobenius $\varphi : \mathcal{O}^{cris} \rightarrow \mathcal{O}^{cris}$ on it, by defining

$$\mathcal{G}^r = \ker(\mathcal{O}^{cris} \xrightarrow{\varphi} \mathcal{O}^{cris} \longrightarrow \mathcal{O}^{cris}/p^r) = \{x \in \mathcal{O}^{cris} \mid \varphi(x) \in p^r \mathcal{O}^{cris}\}.$$

This makes perfect sense in the setting of pro-objects, and then gives rise to the objects $\mathcal{G}_n^r = \mathcal{G}^r/p^n$ which are essentially constant pro-objects.

For a more elementary and direct approach we proceed as follows. For all n we set $\mathcal{G}_n^0 = \mathcal{O}_n^{cris}$, and the sub-ring $\oplus_{r \leq 0} \mathcal{G}_n^r$ is the ring of polynomials in an indeterminate, called v , with coefficients in \mathcal{O}_n^{cris} and v is in degree -1 . In other words, for any object U of \mathcal{C} and any integer $r \leq 0$, $\mathcal{G}_n^r(U)$ is the free $\mathcal{O}_n^{cris}(U)$ -module of rank one generated by v^{-r} .

If $r \geq 0$ and $m \in \mathbb{N}$ with $m \geq r$, set

$$\hat{\mathcal{G}}_m^r = \ker(\mathcal{O}_m^{cris} \xrightarrow{\varphi} \mathcal{O}_m^{cris} \xrightarrow{proj} \mathcal{O}_m^{cris}/p^r = \mathcal{O}_r^{cris}),$$

so this is the sub-sheaf whose sections x are such that $\varphi(x)$ is locally (for the p -topology) divisible by p^r . For any $m \geq n + r$ we define

$$\mathcal{G}_n^r = \hat{\mathcal{G}}_m^r/p^n.$$

It is easily seen that this definition is independent of the choice of $m \geq n + r$, and that this definition agrees with the previous definition $\mathcal{G}_n^0 = \mathcal{O}_n^{cris}$ for $r = 0$. If U is an object of \mathcal{C} , if $m, r \in \mathbb{N}$ with $m \geq r$ and if $x \in \hat{\mathcal{G}}_m^r(U)$, $y \in \hat{\mathcal{G}}_m^s(U)$, then $xy \in \hat{\mathcal{G}}_m^{r+s}(U)$. By going to the quotient this gives a map $\mathcal{G}_n^r \times \mathcal{G}_n^s \rightarrow \mathcal{G}_n^{r+s}$ which define the multiplication on the sub-ring $\oplus_{n \in \mathbb{N}} \mathcal{G}_n^r$.

To complete the definition of the multiplication on \mathcal{G}_n , it is enough to define the multiplication by $v : \mathcal{G}_n^r \rightarrow \mathcal{G}_n^{r-1}$ for $r \geq 1$. It is induced by the inclusion $\hat{\mathcal{G}}_m^r \subset \hat{\mathcal{G}}_m^{r-1}$, via passing to the quotients modulo p^n . In this way \mathcal{G}_n becomes a graded ring.

To get the structure of φ -ring, we have to introduce f, v and φ . We have already defined the global section $v \in \mathcal{G}_n^{-1}(\mathbb{F}_p)$. For all $m \in \mathbb{N}$, we have $\mathcal{O}_m^{cris}(\mathbb{F}_p) = \mathbb{Z}/p^m\mathbb{Z}$. For $n \in \mathbb{N}$ and $m \geq n + 1$, the image of p belongs to $\mathcal{G}_{m,1}(\mathbb{F}_p)$ and we call f its image in $\mathcal{G}_n^1(\mathbb{F}_p)$. Observe that $fv = p$.

The natural morphism $\mathcal{O}_n^{cris} = \mathcal{G}_n^0 \rightarrow \mathcal{G}_n^{-\infty}$ is an isomorphism of rings and we use it to identify $\mathcal{G}_n^{-\infty}$ with \mathcal{O}_n^{cris} .

Finally we define φ . Let $n, r \in \mathbb{N}$, let U be an object of \mathcal{C} and $x \in \mathcal{G}_n^r(U)$. If we choose an integer $m \geq n+r$, we may find a covering V of U , a lifting y of x in $\hat{\mathcal{G}}_m^r(V)$, a covering W of V and $z \in \mathcal{O}_m^{cris}(W)$ such that $\varphi(y) = p^r z$. The image $\varphi_r(x)$ of z in $\mathcal{O}_n^{cris}(W)$ is independent of the choices made and belongs to $\mathcal{O}_n^{cris}(U)$. In this way we have $\varphi_r : \mathcal{G}_n^r \rightarrow \mathcal{O}_n^{cris}$ such that “ $\varphi_r(x) = \varphi(x)/p^r$ ”. Obviously, we have $\varphi_{r+1}(fx) = \varphi_r(x)$, therefore the morphisms φ_r define a map

$$\varphi : \varinjlim_{r \in \mathbb{N}, f} \mathcal{G}_n^r = \mathcal{G}_n^\infty \rightarrow \mathcal{O}_n^{cris} = \mathcal{G}_n^{-\infty}$$

One sees that φ is a morphism of rings; this corresponds to the equality

$$“\varphi(x)/p^r \cdot \varphi(y)/p^s = \varphi(xy)/p^{r+s}”$$

for sections $x \in \hat{\mathcal{G}}_m^r$ and $y \in \hat{\mathcal{G}}_m^s$ and the ring endomorphism φ of \mathcal{O}_m^{cris} .

THEOREM 7.2.1. *For each $n \geq 1$, the map*

$$\varphi : \mathcal{G}_n^{+\infty} \rightarrow \mathcal{G}_n^{-\infty}$$

is an isomorphism of rings.

In other words, \mathcal{G}_n is a perfect φ -ring. The proof of this theorem is given in sections 7.3–7.5 below.

Here we note the following properties.

LEMMA 7.2.2. *The following holds for the gauges \mathcal{G}_n .*

- (i) *One has natural isomorphisms $\mathcal{G}_{n+1}/p^n \rightarrow \mathcal{G}_n$ for all n , i.e., $(\mathcal{G}_n)_n$ is a p -adic ring.*
- (ii) *The p -adic ring $\mathcal{G} = (\mathcal{G}_n)_{n \in \mathbb{N}}$ is flat.*
- (ii) *The gauge \mathcal{G}_1 is rigid.*

Proof. Morally, all statements are proved in the same way as for the standard construction in section 3, and could be proved by noting that \mathcal{O}^{cris} is a flat (= “torsion-free”) p -adic ring. But for concreteness we give a proof by the above explicit construction. We argue with local sections, i.e., all statements hold after possible passing to some cover. Moreover, we constantly use that the flatness of \mathcal{O}^{cris} implies the exactness of the sequence

$$\mathcal{O}_m^{cris} \xrightarrow{p^{m-i}} \mathcal{O}_m^{cris} \xrightarrow{p^i} \mathcal{O}_m^{cris}$$

for all $i \leq m$. Recall that $\mathcal{G}_n^r = \hat{\mathcal{G}}_m^r/p^n$ for some fixed $m \gg 0$ ($m \geq n+r$ suffices), where $\hat{\mathcal{G}}_m^r = \{x \in \mathcal{O}_m^{cris} \mid \varphi(x) \in p^r \mathcal{O}_m^{cris}\}$. Unless specified explicitly, we calculate inside \mathcal{O}_m^{cris} .

- (i): This is trivial: $(\hat{\mathcal{G}}_m^r/p^{n+1})/p^n = \hat{\mathcal{G}}_m^r/p^n$.

(ii): For each $i \leq n$ and $m \geq n + r$ we show the exactness of

$$\hat{\mathcal{G}}_m^r/p^n \xrightarrow{p^{n-i}} \hat{\mathcal{G}}_m^r/p^n \xrightarrow{p^i} \hat{\mathcal{G}}_m^r/p^n$$

as follows: If $x \in \hat{\mathcal{G}}_m^r$ and $p^i x = p^n y$ with $y \in \hat{\mathcal{G}}_m^r$, then we have $p^i(x - p^{n-i}y) = 0$ and hence $x - p^{n-i}y = p^{m-i}z$ with $z \in \mathcal{O}_m^{cris}$ by flatness of \mathcal{O}^{cris} . Thus $x = p^{n-i}(y + p^{m-n}z)$ with $y, p^{m-n}z \in \hat{\mathcal{G}}_m^r$ (note that $m - n \geq r$).

(iii): Recall that v and f are induced by the inclusion $\hat{\mathcal{G}}_m^r \hookrightarrow \hat{\mathcal{G}}_m^{r-1}$ and the p -multiplication $\hat{\mathcal{G}}_m^r \xrightarrow{p} \hat{\mathcal{G}}_m^r$, respectively. First we show quasi-rigidity. If $x \in \hat{\mathcal{G}}_m^r$ and $x = py$ for $y \in \hat{\mathcal{G}}_m^{r-1}$, then $[x] = f[y]$ for the class $[x]$ of x in \mathcal{G}_1^r and the class $[y]$ of y in \mathcal{G}_1^{r-1} . On the other hand, if $x \in \hat{\mathcal{G}}_m^r$ with $px = pz$ for $z \in \hat{\mathcal{G}}_m^{r+1}$, then $p(x - z) = 0$ and hence $x - z = p^{m-1}t$ with $t \in \mathcal{O}_m^{cris}$. Thus $x = z + p^{m-1}t \in \hat{\mathcal{G}}_m^{r+1}$ for $m \geq r + 2$, i.e., $[x] = v[x]$. Now we show strictness. Let $x \in \hat{\mathcal{G}}_m^r$ with $f[x] = 0 = v[x]$. This means that $x = py$ with $y \in \hat{\mathcal{G}}_m^{r-1}$ and $px = pz$ with $z \in \hat{\mathcal{G}}_m^{r+1}$. This implies $pz = px = p^2y$, hence $p^2\varphi(y) = p\varphi(z) = p^{r+2}t$ for some $t \in \mathcal{O}_m^{cris}$. This implies $p^2(\varphi(y) - p^r t) = 0$ and thus $\varphi(y) - p^r t = p^{m-2}u$ for some $u \in \mathcal{O}_m^{cris}$ by flatness of \mathcal{O}^{cris} . We conclude $y \in \hat{\mathcal{G}}_m^r$ for $m \geq r + 2$ and therefore $[x] = [py] = 0$. \square

7.3 The structure of a (generalized) F -zip on \mathcal{O}_1^{cris}

Let $\mathcal{T} = (\mathcal{C}, E, \mathcal{O})$ be a ringed topos with the category \mathcal{C} , an admissible topology E , and the structure ring \mathcal{O} as in section 7.1.

An effective generalized F -zip (over \mathcal{O}) is an \mathcal{O} -module \mathcal{M} together with

- a decreasing filtration $(F^r \mathcal{M})_{r \in \mathbb{N}}$ by sub- \mathcal{O} -modules indexed by \mathbb{N} such that $F^0 \mathcal{M} = \mathcal{M}$ and $\bigcap_{r \in \mathbb{N}} F^r \mathcal{M} = 0$,
- an increasing filtration $(F_r \mathcal{M})_{r \geq -1}$ by sub- \mathcal{O} -modules indexed by the natural integers $r \geq -1$ such that $F_{-1} \mathcal{M} = 0$ and $\bigcup_{r \in \mathbb{N}} F_r \mathcal{M} = \mathcal{M}$,
- for all $r \in \mathbb{N}$, an isomorphism

$$\varphi_r : F^r \mathcal{M} / F^{r+1} \mathcal{M} \xrightarrow{\sim} F_r \mathcal{M} / F_{r-1} \mathcal{M}$$

of abelian sheaves which is semi-linear with respect to the absolute Frobenius.

REMARK 7.3.1 Assume that E is p -admissible. Then the absolute Frobenius F_r is an epimorphism on \mathcal{O} , and it follows that the associated morphism

$$(F^r \mathcal{M} / F^{r+1} \mathcal{M})^{(p)} = \mathcal{O}_{F_r \otimes \mathcal{O}} (F^r \mathcal{M} / F^{r+1} \mathcal{M}) \longrightarrow F_r \mathcal{M} / F_{r-1} \mathcal{M}$$

is an isomorphism. This ties this definition with the definition in section 4.2.

We shall now define such a structure on \mathcal{O}_1^{cris} (with \mathcal{O} the structural sheaf). We define $F_r = F_r \mathcal{O}_1^{cris} = im(\varphi_r)$ for $\varphi_r : \mathcal{G}_1^r \rightarrow \mathcal{O}_1^{cris}$ and $F^r = F^r \mathcal{O}_1^{cris} = im(v^r)$ for $v^r : \mathcal{G}_1^r \rightarrow \mathcal{O}_1^{cris}$.

PROPOSITION 7.3.2. *There are canonical exact sequences for all $r \geq 1$*

$$\begin{aligned} 0 \longrightarrow F_r \longrightarrow \mathcal{G}_1^{r+1} \xrightarrow{v} \mathcal{G}_1^r \xrightarrow{\varphi_r} F_r \longrightarrow 0 \\ 0 \longrightarrow F^r \longrightarrow \mathcal{G}_1^{r-1} \xrightarrow{f} \mathcal{G}_1^r \xrightarrow{v^r} F^r \longrightarrow 0. \end{aligned}$$

Proof. For the first sequence we claim that there is an exact sequence

$$0 \rightarrow \hat{\mathcal{G}}_{r+2}^{r+1} \hookrightarrow \hat{\mathcal{G}}_{r+2}^r \xrightarrow{\varphi_r} \mathcal{O}_1^{cris},$$

where the morphism on the right is the composition $\hat{\mathcal{G}}_{r+2}^r \rightarrow \mathcal{G}_1^r \xrightarrow{\varphi_r} \mathcal{O}_1^{cris}$. In fact, if $x \in \hat{\mathcal{G}}_{r+2}^r$, so that $\varphi(x) = p^r y$ with $y \in \mathcal{O}_{r+2}^{cris}$, then $\varphi_r([x]) = [y] \in \mathcal{O}_{r+2}^{cris}/p = \mathcal{O}_1^{cris}$ for the corresponding classes modulo p . If $\varphi_r([x]) = 0$, then $y = py'$ with $y' \in \mathcal{O}_{r+2}^{cris}$. This implies $\varphi(x) = p^{r+1}y'$ and hence $x \in \hat{\mathcal{G}}_{r+2}^{r+1}$. Applying now the snake lemma to the multiplication by p on the exact sequence

$$0 \rightarrow \hat{\mathcal{G}}_{r+2}^{r+1} \hookrightarrow \hat{\mathcal{G}}_{r+2}^r \xrightarrow{\varphi_r} F_r \rightarrow 0,$$

we get an exact sequence

$$\hat{\mathcal{G}}_{r+2}^r[p] \rightarrow F_r \rightarrow \mathcal{G}_1^{r+1} \hookrightarrow \mathcal{G}_1^r \xrightarrow{\varphi_r} F_r \rightarrow 0,$$

where $A[p]$ means the p -torsion subsheaf of a sheaf A . But the map on the left is zero: If $x \in \hat{\mathcal{G}}_{r+2}^r$ with $px = 0$, then $x = p^{r+1}u$ for some $u \in \mathcal{O}_{r+2}^{cris}$ by flatness of \mathcal{O}^{cris} . Then $\varphi(x) = p^{r+1}\varphi(u)$, and by definition, $\varphi_r([x]) = [p\varphi(u)] = 0 \in \mathcal{O}_1^{cris}$.

For the second sequence we claim that we have an exact sequence

$$p\hat{\mathcal{G}}_{r+1}^{r-1} \hookrightarrow \hat{\mathcal{G}}_{r+1}^r \xrightarrow{v^r} \mathcal{O}_1^{cris},$$

where the morphism on the right is the composition $\hat{\mathcal{G}}_{r+1}^r \rightarrow \mathcal{G}_1^r \xrightarrow{v^r} \mathcal{O}_1^{cris}$. In fact, if $x \in \hat{\mathcal{G}}_{r+1}^r$, so that $\varphi(x) = p^r y$ with $y \in \mathcal{O}_{r+1}^{cris}$, and $x = pz$ with $z \in \mathcal{O}_{r+1}^{cris}$, then $p^r y = \varphi(x) = p\varphi(z)$, hence $p(\varphi(z) - p^{r-1}y) = 0$, so that $\varphi(z) - p^{r-1}y = p^r t$ with $t \in \mathcal{O}_{r+1}^{cris}$. This implies $z \in \hat{\mathcal{G}}_{r+1}^{r-1}$, hence the claim, because $x = pz$. Applying the snake lemma to the multiplication by p to the exact sequence

$$0 \rightarrow p\hat{\mathcal{G}}_{r+1}^{r-1} \rightarrow \hat{\mathcal{G}}_{r+1}^r \xrightarrow{v^r} F^r \rightarrow 0,$$

we get an exact sequence

$$\hat{\mathcal{G}}_{r+1}^r[p] \rightarrow F^r \rightarrow (p\hat{\mathcal{G}}_{r+1}^{r-1})/p \rightarrow \mathcal{G}_1^r \xrightarrow{v^r} F^r \rightarrow 0.$$

Now we claim that the first map is the zero map. In fact, if $x \in \hat{\mathcal{G}}_{r+1}^r$ with $px = 0$, then $x = p^r y$ with $y \in \mathcal{O}_{r+1}^{cris}$ by flatness of \mathcal{O}^{cris} . Hence x is mapped to zero in \mathcal{O}_1^{cris} .

Moreover we claim that the exact sequence

$$0 \rightarrow \hat{\mathcal{G}}_{r+1}^{r-1}[p] \rightarrow \hat{\mathcal{G}}_{r+1}^{r-1} \xrightarrow{P} p\hat{\mathcal{G}}_{r+1}^{r-1} \rightarrow 0$$

taken modulo p induces an isomorphism

$$\mathcal{G}_1^{r-1} = \hat{\mathcal{G}}_{r+1}^{r-1}/p \xrightarrow{\sim} (p\hat{\mathcal{G}}_{r+1}^{r-1})/p.$$

In fact, for this it suffices to show that the induced morphism $(\hat{\mathcal{G}}_{r+1}^{r-1}[p])/p \rightarrow (\hat{\mathcal{G}}_{r+1}^{r-1})/p$ is zero. But if $x \in \hat{\mathcal{G}}_{r+1}^{r-1}[p]$, then $px = 0$, so that $x = p^r t$ with $t \in \mathcal{O}_{r+1}^{cris}$ by flatness of \mathcal{O}^{cris} . Then $x = py$ with $y = p^{r-1}t \in \mathcal{G}_{r+1}^{r-1}$ as claimed.

Both claims together imply the second exact sequence in the proposition. \square

PROPOSITION 7.3.3. (*Cartier isomorphism*) *The subsheaves $F_r = F_r \mathcal{O}_1^{cris}$ form an increasing filtration of \mathcal{O}_1^{cris} (i.e., $F_r \subseteq F_{r+1}$), and the subsheaves $F^r = F^r \mathcal{O}_1^{cris}$ form a decreasing filtration of \mathcal{O}_1^{cris} (i.e., $F^{r+1} \subseteq F^r$). For each $r \geq 0$, the morphism $\varphi_r : \mathcal{G}_1^r \rightarrow \text{im}(\varphi_r) = F_r$ induces an isomorphism*

$$\bar{\varphi}_r : F^r/F^{r+1} \xrightarrow{\sim} F_r/F_{r-1}.$$

Proof. The first two claims are clear. For the third look at the diagram with exact rows

$$\begin{array}{ccccccc} \mathcal{G}_1^{r+1} & \xrightarrow{v} & \mathcal{G}_1^r & \xrightarrow{\varphi_r} & F_r & \longrightarrow & 0 \\ & & \parallel & & & & \\ \mathcal{G}_1^{r-1} & \xrightarrow{f} & \mathcal{G}_1^r & \xrightarrow{v^r} & F^r & \longrightarrow & 0 \end{array}.$$

Since $\varphi_r f = \varphi_{r-1}$ the diagram induces an epimorphism

$$\varphi'_r : F^r \longrightarrow F_r/F_{r-1}.$$

Explicitly: If $x \in F^r$, and $x = v^r y$ for $y \in \mathcal{G}_1^r$, let $\varphi'_r(x)$ be the class of $\varphi_r(y)$, which is well-defined modulo F_{r-1} , because the kernel of v^r is the image of f . The kernel of this morphism is F^{r+1} : If $\varphi'_r(x) = \varphi_r(y) = \varphi_{r-1}(z) = \varphi_r(fz)$, then $\varphi_r(y - fz) = 0$, so that $y - fz = vt$ with $t \in \mathcal{G}_1^{r+1}$ by the exactness of the upper row. But then $x = v^r y = v^r fz + v^{r+1}t = v^{r+1}t \in F^{r+1}$, because $vf = 0$. \square

From Proposition 7.3.2 we already obtain one half of Theorem 7.2.1.

COROLLARY 7.3.4. *The morphism $\varphi : \mathcal{G}_1^{+\infty} \rightarrow \mathcal{G}_1^{-\infty}$ is injective.*

Proof. In view of Lemma 7.2.2 we may assume that $n = 1$. Let $x \in \mathcal{G}_1^{+\infty}$ with $\varphi(x) = 0$. Suppose x is the image of an element x_r under $f_r : \mathcal{G}_1^r \rightarrow \mathcal{G}_1^{+\infty}$. Then $\varphi_r(x_r) = 0$. By the first sequence in 7.3.2, we have $x_r = vx_{r+1}$. But then $fx_r = fv x_{r+1} = px_{r+1} = 0$, which implies $x = f_r x_r = 0$. \square

To obtain the fact that $\cup_r F_r = \mathcal{O}_1^{cris}$ (which gives the surjectivity of φ above and hence the second half of Theorem 7.2.1), and that $\cap_r F^r = 0$, we need some explicit calculations.

7.4 Some calculations for the universal F -zip

We first observe that we have a morphism of rings

$$f : \mathcal{O} \rightarrow \mathcal{O}_1^{cris} .$$

In fact, if X is an object of \mathcal{C} and if $a \in \mathcal{O}(X)$, one may find a quiet covering $Y \rightarrow X$ such that a is the image of some $b \in \mathcal{O}_1^{cris}(Y)$ and $f(a) = b^p$ is independent of the choice of b and belongs to $\mathcal{O}_1^{cris}(X)$.

The kernel $\tilde{F}^1 \mathcal{O}_1^{cris}$ (often called $J_1^{[1]}$) of the canonical epimorphism $\mathcal{O}_1^{cris} \rightarrow \mathcal{O}$ is a divided power Ideal and, for all $r \in \mathbb{N}$, we set $\tilde{F}^r \mathcal{O}_1^{cris} = J_1^{[r]}$, the r -th divided power of $\tilde{F}^1 \mathcal{O}_1^{cris} = J_1^{[1]}$. We set $I = J_1^{[1]}/J_1^{[2]}$ and call it *the cotangent space*. We denote $G^r \mathcal{O}_1^{cris}$ the abelian sheaf $\tilde{F}^r \mathcal{O}_1^{cris} / \tilde{F}^{r+1} \mathcal{O}_1^{cris} = J_1^{[r]}/J_1^{[r+1]}$ (hence $G^0 \mathcal{O}_1^{cris} = \mathcal{O}$ and $G^1 \mathcal{O}_1^{cris} = I$). On each $G^r \mathcal{O}_1^{cris}$, we have two different structures of \mathcal{O} -Modules :

- the *naive structure* which comes from the fact that $J_1^{[r+1]} \subset J_1^{[r]}$ are sub- \mathcal{O} -modules of \mathcal{O}_1^{cris} ,
- the *nice structure* which comes from the fact that $J_1^{[r+1]} \subset J_1^{[r]}$ are sub- \mathcal{O}_1^{cris} -modules of \mathcal{O}_1^{cris} , but that $J_1^{[1]} \cdot J_1^{[r]} \subset J_1^{[r+1]}$ and $\mathcal{O}_1^{cris}/J_1^{[1]} = \mathcal{O}$.

If λ is a local section of \mathcal{O} and a a local section of $F^r \mathcal{O}_1^{cris}$, $\lambda \cdot_{naive} a = \lambda^p \cdot_{nice} a$. In what follows, we will always consider $J_1^{[r]}/J_1^{[r+1]}$ as a \mathcal{O} -module via the nice structure (but we have to keep in mind that, in the structure of F -zip, it is the naive structure which matters).

If A is a ring, $A \langle \theta_1, \theta_2, \dots, \theta_d \rangle$ is the divided power algebra in the indeterminates $\theta_1, \theta_2, \dots, \theta_d$ with coefficients in A . If t belongs to some divided power ideal in some $\mathbb{Z}_{(p)}$ -algebra, and if $m = qp + r$, with $q, r \in \mathbb{N}$ and $r < p$, we have $\gamma_m(t) = c_m t^r \gamma_q(\gamma_p(t))$ with $c_m = \frac{q!(p!)^q}{m!}$ a unit in $\mathbb{Z}_{(p)}$. The following result is the key for many explicit computations with \mathcal{O}_1^{cris} :

PROPOSITION 7.4.1. *Let k be a perfect ring of characteristic $p > 0$, let \mathcal{A} be a smooth k -algebra, $t_1, t_2, \dots, t_d \in \mathcal{A}$ a regular sequence, $A = \mathcal{A}/(t_1^p, t_2^p, \dots, t_d^p)$ and \bar{t}_i the image of t_i in A . The unique homomorphism of divided power A -algebras*

$$A \langle \theta_1, \theta_2, \dots, \theta_d \rangle \longrightarrow \mathcal{O}_1^{cris}(A)$$

sending θ_i to $\gamma_p(f(\bar{t}_i))$ is an isomorphism. Moreover

- i) For any $m = qp + r$ with $q, r \in \mathbb{N}$, $r < p$, set $\gamma_m(t_i) = c_m \bar{t}_i^r \gamma_q(\theta_i)$. For any $r \in \mathbb{N}$, $J_1^{[r]}(A)$ is the image of the sub- A -module generated by the $\gamma_{m_1}(t_1) \gamma_{m_2}(t_2) \dots \gamma_{m_d}(t_d)$ for $\sum m_i \geq r$.
- ii) Let \bar{I} be the ideal of A generated by the \bar{t}_i and let $\bar{A} = A/\bar{I}$. The A -module quotient $J_1^{[r]}(A)/J_1^{[r+1]}(A)$ is annihilated by \bar{I} and is a free \bar{A} -module with the images of the elements

$$\gamma_{m_1}(t_1) \gamma_{m_2}(t_2) \dots \gamma_{m_d}(t_d) \quad \text{for} \quad \sum m_i = r$$

as a basis.

iii) The natural map $J_1^{[r]}(A)/J_1^{[r+1]}(A) \rightarrow (J_1^{[r]}/J_1^{[r+1]})(A)$ is injective and, for the nice structure, $(J_1^{[r]}/J_1^{[r+1]})(A)$ is a free A -module with the images of the elements

$$\gamma_{m_1}(t_1)\gamma_{m_2}(t_2)\dots\gamma_{m_d}(t_d) \quad \text{for } \Sigma m_i = r$$

as a basis.

iv) Let ρ be an endomorphism of the k -algebra A such that one can find $\lambda_1, \lambda_2, \dots, \lambda_d \in A$ with $\rho(\bar{t}_i) = \lambda_i t_i$. Then the endomorphism induced by ρ on $A \langle \theta_1, \theta_2, \dots, \theta_d \rangle$ (by functoriality and the isomorphism $A \langle \theta_1, \theta_2, \dots, \theta_d \rangle \simeq \mathcal{O}_1^{cris}(A)$) is compatible with the divided power structure and sends θ_i to $\lambda_i^p \theta_i$.

We start with a lemma:

LEMMA 7.4.2. Let \mathcal{A}' be a k -algebra, $t_1, t_2, \dots, t_d \in \mathcal{A}'$ a regular sequence and $A' = \mathcal{A}'/(t_1^p, t_2^p, \dots, t_d^p)$. Let \mathcal{D}' the divided power envelope of \mathcal{A}' with respect to the ideal generated by the t_i 's and $\xi: \mathcal{A}' \rightarrow \mathcal{D}'$ the structural map. Then:

i) each t_i^p belongs to the kernel of ξ (and therefore \mathcal{D}' may be viewed as an A' -algebra),

ii) the unique homomorphism of divided power A' -algebras

$$\eta: A' \langle \theta_1, \theta_2, \dots, \theta_d \rangle \rightarrow \mathcal{D}'$$

sending θ_i to $\gamma_p(t_i)$ is an isomorphism.

Proof of the lemma: We have $t_i^p = p! \gamma_p(t_i) = 0$, which proves (i).

As an A' -algebra, \mathcal{D}' is generated by the $\Pi \gamma_{m_i}(t_i)$. But, if $m_i = pq_i + r_i$ with $q_i, r_i \in \mathbb{N}$ and $r_i < p$, we have

$$\Pi \gamma_{m_i}(t_i) = \Pi c_{m_i} \Pi t_i^{r_i} \Pi \gamma_{m_i}(\gamma_p(t_i))$$

hence \mathcal{D}' is also generated by the $\Pi \gamma_{m_i}(\gamma_p(t_i))$ and η is surjective.

For $m \in \mathbb{N}$, let $\mathcal{J}^{[m]}$ the m -th divided power of the structural divided power ideal of \mathcal{D}' , let \mathcal{I} the ideal of \mathcal{A}' generated by the t_i 's and $\overline{\mathcal{A}'} = \mathcal{A}'/\mathcal{I}$. As the t_i 's form a regular sequence, $\mathcal{I}/\mathcal{I}^2$ is a free $\overline{\mathcal{A}'}$ -module with the image of the t_i 's as a basis and the canonical map $\Gamma_{\overline{\mathcal{A}'}}^m(\mathcal{I}/\mathcal{I}^2) \rightarrow \mathcal{J}^{[m]}/\mathcal{J}^{[m+1]}$ is an isomorphism, where $\Gamma_{\overline{\mathcal{A}'}}^r(M)$ denotes the r -th divided power of an $\overline{\mathcal{A}'}$ -module M [Be] I, 3.4.4. If $\mathcal{J}_1^{[m]}$ denote the inverse image under η of $\mathcal{J}^{[m]}$, we see that, for all m , the induced map $\mathcal{J}_1^{[m]}/\mathcal{J}_1^{[m+1]} \rightarrow \mathcal{J}^{[m]}/\mathcal{J}^{[m+1]}$ is an isomorphism. As $\bigcap_{m \in \mathbb{N}} \mathcal{J}_1^{[m]} = 0$, the map η is injective. \square

Proof of proposition 7.4.1 Let $\mathcal{A}' = \mathcal{A}$, but viewed as an \mathcal{A} -algebra via the absolute Frobenius, that we use to identify \mathcal{A} to a sub-ring of \mathcal{A}' . Therefore, any element of \mathcal{A} has a unique p -th root in \mathcal{A}' and, for an $b \in \mathcal{A}'$, we have $b^p \in \mathcal{A}$. We denote by t'_i the element t_i viewed as an element of \mathcal{A}' , hence $(t'_i)^p = t_i$.

As \mathcal{A} is smooth, \mathcal{A}' is a syntomic \mathcal{A} -algebra and t_1, t_2, \dots, t_d is still a regular sequence in \mathcal{A}' . The map $\varphi : \mathcal{A}' \rightarrow \mathcal{A}$ sending b to b^p is an isomorphism and the compositum $\overline{\varphi}$ with the projection onto A is surjective, with kernel the ideal generated by the t_i 's. As \mathcal{A}' is smooth, if \mathcal{D}' is as in the lemma, we have an exact sequence

$$0 \rightarrow \mathcal{O}_1^{cris}(A) \rightarrow \mathcal{D}' \rightarrow \mathcal{D}' \otimes_{\mathcal{A}'} \Omega_{\mathcal{A}'/k}^1$$

which, granted to the previous lemma, can be rewritten as

$$0 \rightarrow \mathcal{O}_1^{cris}(A) \rightarrow A' \langle \theta_1, \theta_2, \dots, \theta_d \rangle \rightarrow A' \langle \theta_1, \theta_2, \dots, \theta_d \rangle \otimes_{\mathcal{A}'} \Omega_{\mathcal{A}'/k}^1$$

with θ_i mapping to $\gamma_p((t'_i)^p)$. Therefore, we have $d\theta_i = \gamma_{p-1}((t'_i)^p) \cdot p(t'_i)^{p-1} = 0$. For $m = (m_1, m_2, \dots, m_d) \in \mathbb{N}^d$ set $\gamma_m(\theta) = \prod \gamma_{m_i}(\theta_i)$. We have also $d\gamma_m(\theta) = 0$, for all m .

But $A' \langle \theta_1, \theta_2, \dots, \theta_d \rangle$ is a free A' -module with basis the $\gamma_m(\theta)$'s. If $\sum a_m \gamma_m(\theta) \in A' \langle \theta_1, \theta_2, \dots, \theta_d \rangle$, we have $d(\sum a_m \gamma_m(\theta)) = \sum da_m \gamma_m(\theta)$. It is easy to check that we have $\Omega_{\mathcal{A}'/k}^1 = A' \otimes_{\mathcal{A}'} \Omega_{\mathcal{A}'/k}^1$ and that the sequence

$$0 \rightarrow A \rightarrow A' \rightarrow \Omega_{\mathcal{A}'/k}^1$$

is exact. This proves that the map

$$A \langle \theta_1, \theta_2, \dots, \theta_d \rangle \rightarrow \mathcal{O}_1^{cris}(A)$$

is an isomorphism.

The proof of (i) and (ii) and (iv) are straightforward. Let us prove (iii). We have to understand the sheaf F associated to the presheaf $P : X \mapsto J_1^{[r]}(X)/J_1^{[r+1]}(X)$. It is enough to consider the restriction of the functor P to the objects X of \mathcal{C} of the form $X = Spec A$, with A as in the proposition. Let M be the set of the $\underline{m} = \{m_1, m_2, \dots, m_d\} \in \mathbb{N}^d$ such that $\sum m_i = r$. For any such \underline{m} , let $\gamma_{\underline{m}}$ the image of $\prod \gamma_{m_i}(t_i)$ in $P(A)$ and $\tilde{\gamma}_{\underline{m}}$ its image in $F(A)$. Checking carefully, we see that the map $P(A) \rightarrow F(A)$ is A -linear if $F(A)$ is equipped with the naive structure. We may express this fact as follows: Let A_1 the image of $\varphi : A \rightarrow A$. The absolute Frobenius induces an isomorphism $f : \overline{A} \rightarrow A_1$ and we may use it to view $P(A)$ as a free A_1 -module. Now the map $P(A) \rightarrow F(A)$ is A_1 -linear when we endow $F(A)$ with the A_1 -structure coming from the inclusion $A_1 \subset A$ and the nice structure of A -module on $F(A)$. Therefore, if we set $P'(A) = A \otimes_{A_1} P(A)$, we have a natural A -linear map $P'(A) \rightarrow F(A)$ and what we want to prove is that this map is an isomorphism. Because there are enough algebras of this kind, it makes sense to speak of the sheaf associated to the presheaf $A \rightarrow P'(A)$ and all what we have to prove is that this presheaf is a sheaf. We are easily reduced to checking that, if B is any faithfully flat A -algebra of the type $A[x_1, x_2, \dots, x_r]/(u_1, u_2, \dots, u_n)$ with $u_1, u_2, \dots, u_n \in A[x_1, x_2, \dots, x_r]$ a sequence which is transverse regular with respect to A , then the sequence

$$0 \rightarrow P'(A) \rightarrow P'(B) \rightrightarrows P'(B \otimes_A B)$$

is exact. Replacing B with a covering if necessary, we may assume that $u_i = v_i^p$ and the v_i^p 's are transverse regular as well. Set $\mathcal{B} = \mathcal{A}[x_1, x_2, \dots, x_r]$ and $\mathcal{C} = \mathcal{B} \otimes_A \mathcal{B} = \mathcal{A}[x_1, x_2, \dots, x_r, y_1, y_2, \dots, y_r]$ if we set $x_i = x_i \otimes 1$ and $y_i = 1 \otimes x_i$. If for $1 \leq i \leq n$, we choose a lifting t_{d+i} of v_i in \mathcal{B} and if, for $1 \leq i \leq n$, we set $t_{d+i} = t_{d+i} \otimes 1$ and $t_{d+n+i} = 1 \otimes t_{d+i}$. The sequence $(t_i)_{1 \leq i \leq d}$ (resp. $(t_i)_{1 \leq i \leq d+n}$, $(t_i)_{1 \leq i \leq d+2n}$) is regular in \mathcal{A} (resp. \mathcal{B} , resp. \mathcal{C}) and

$$A = \mathcal{A}/((t_i^p)_{1 \leq i \leq d}), \quad B = \mathcal{B}/((t_i^p)_{1 \leq i \leq d+n}), \quad B \otimes_A B = \mathcal{C}/((t_i^p)_{1 \leq i \leq d+2n})$$

Let M (resp. N , resp. L) the set of the $\underline{m} = (m_i)_{1 \leq i \leq d}$ (resp. $(m_i)_{1 \leq i \leq d+n}$, resp. $(m_i)_{1 \leq i \leq d+2n}$) such that $\sum m_i = r$ and, for $\underline{m} \in M$ (resp. N, L), let $\gamma_{\underline{m}}(t)$ the image of $\Pi \gamma_{m_i}(t_i)$ in $P'(A)$ (resp. $P'(B), P'(B \otimes_A B)$). These $\gamma_{\underline{m}}(t)$ form a basis respectively of the free A -module $P'(A)$, the free B -module $P'(B)$, the free $B \otimes_A B$ -module $P'(B \otimes_A B)$ and we are reduced to prove the exactness of the sequence

$$0 \rightarrow \bigoplus_{\underline{m} \in M} A \gamma_{\underline{m}}(t) \rightarrow \bigoplus_{\underline{m} \in N} B \gamma_{\underline{m}}(t) \xrightarrow{\rightarrow} \bigoplus_{\underline{m} \in L} B \otimes_A B \gamma_{\underline{m}}(t)$$

This is easily reduced to the exactness of

$$0 \rightarrow A \rightarrow B \rightarrow B \otimes_A B$$

which comes from the fact that B is faithfully flat over A . \square

Call J the kernel of the absolute Frobenius φ on \mathcal{O} (hence, with the usual notations $J = \alpha_p$).

PROPOSITION 7.4.3. *We have a commutative diagram of abelian sheaves*

$$\begin{array}{ccccccccc} 0 & \rightarrow & J & \rightarrow & \mathcal{O} & \xrightarrow{\varphi} & \mathcal{O} & \rightarrow & 0 \\ & & \downarrow & & \downarrow f & & \parallel & & \\ 0 & \rightarrow & J_1^{[1]} & \rightarrow & \mathcal{O}_1^{cris} & \rightarrow & \mathcal{O} & \rightarrow & 0 \end{array}$$

whose lines are exact and whose vertical arrows are monomorphisms.

Proof. The commutativity of the diagram is clear. We already know that the second line is exact. As our topology allows us to extract p -th roots, the Frobenius is an epimorphism and the first line is also exact. We are left to check that f is a monomorphism. As we can always cover any object X of \mathcal{C} by affine k -schemes $Y = \text{Spec } A$, with A as in the previous proposition, we are reduced to check the injectivity of $A \rightarrow \mathcal{O}_1^{cris}(A)$ which is a consequence of that proposition. \square

If A is a commutative ring and M is an A -module, we may consider the symmetric algebra $Sym_A M = \bigoplus_{r \in \mathbb{N}} Sym_A^r M$ and the divided power algebra $\Gamma_A M = \bigoplus_{r \in \mathbb{N}} \Gamma_A^r M$. They are graded algebras, with grading indexed by \mathbb{N} . For $r \in \mathbb{N}$, $\Gamma_A^r M$ is the sub- A -module of $\Gamma_A M$ generated by the $\gamma_{r_1}(x_1) \gamma_{r_2}(x_2) \dots \gamma_{r_d}(x_d)$ with $r_1, r_2, \dots, r_d \in \mathbb{N}$ satisfying $\sum r_i = r$ and $x_1, x_2, \dots, x_d \in M$. This extends to rings and modules in a topos.

PROPOSITION 7.4.4. *We have $\cap_{r \in \mathbb{N}} J_1^{[r]} = 0$. Moreover, for any $r \in \mathbb{N}$, the obvious map*

$$\Gamma_{\mathcal{O}}^r I \rightarrow J_1^{[r]} / J_1^{[r+1]}$$

is an isomorphism of \mathcal{O} -modules.

Proof. From proposition 7.4.1, we see that for any k -algebra A satisfying the condition of 7.4.1, we have $\cap_{r \in \mathbb{N}} J_1^{[r]}(A) = 0$ and the map $\Gamma_{\mathcal{O}}^r I(A) \rightarrow (J_1^{[r]} / J_1^{[r+1]})(A)$ is an isomorphism. The proposition therefore follows from the fact that any object X of \mathcal{C} has a covering by affine k -schemes $Y = \text{Spec } A$, with A of this kind. \square

In view of proposition 7.4.3, we may use f to regard \mathcal{O} as a sub-ring of \mathcal{O}_1^{cris} and J as a sub \mathcal{O} -Module of \mathcal{O}_1^{cris} . For any $r \in \mathbb{N}$, we call $\tilde{F}_r \mathcal{O}_1^{cris}$ the sub- \mathcal{O} -module of \mathcal{O}_1^{cris} generated locally by the elements of the form $\gamma_{pr_1}(x_1)\gamma_{pr_2}(x_2)\dots\gamma_{pr_d}(x_d)$ with $r_1, r_2, \dots, r_d \in \mathbb{N}$ satisfying $\sum r_i \leq r$ and $x_1, x_2, \dots, x_d \in J$. We have $\tilde{F}_0 \mathcal{O}_1^{cris} = \mathcal{O}$ and we set $\tilde{F}_{-1} \mathcal{O}_1^{cris} = 0$.

The next theorem gives the second construction for the structure of F -zip on \mathcal{O}_1^{cris} :

THEOREM 7.4.5. *i) We have $\cup_{r \in \mathbb{N}} \tilde{F}_r \mathcal{O}_1^{cris} = \mathcal{O}_1^{cris}$.*

ii) (Cartier isomorphism, second version) Let $r \in \mathbb{N}$. There is a unique morphism of \mathcal{O} -modules

$$f_r : J_1^{[r]} / J_1^{[r+1]} \rightarrow \tilde{F}_r \mathcal{O}_1^{cris} / \tilde{F}_{r-1} \mathcal{O}_1^{cris}$$

such that, if $r_1, r_2, \dots, r_d \in \mathbb{N}$ satisfies $\sum r_i = r$ and if $x_1, x_2, \dots, x_d \in J_1$, then f_r sends the image of $\gamma_{r_1}(x_1)\gamma_{r_2}(x_2)\dots\gamma_{r_d}(x_d)$ to the image of $(-1)^r \gamma_{pr_1}(x_1)\gamma_{pr_2}(x_2)\dots\gamma_{pr_d}(x_d)$.

Moreover f_r is an isomorphism.

Proof. Again, it is enough to prove the corresponding assertion for the sections of these sheaves with values in A for any k -algebra A of the kind considered in proposition 7.4.1.

We use the notation of that proposition. Then $J(A)$ is the A -module generated by the \bar{t}_i . For $m_1, m_2, \dots, m_d \in \mathbb{N}$, we have

$$\gamma_{pm_1}(\bar{t}_1)\gamma_{pm_2}(\bar{t}_2)\dots\gamma_{pm_d}(\bar{t}_d) = c_{m_1, m_2, \dots, m_d} \gamma_{m_1}(\theta_1)\gamma_{m_2}(\theta_2)\dots\gamma_{m_d}(\theta_d)$$

with $c_{m_1, m_2, \dots, m_d} = \prod \frac{(pm_i)!}{m_i!(p!)^{m_i}}$, a unit of $\mathbb{Z}_{(p)}$. Hence, $\tilde{F}_r \mathcal{O}_1^{cris}(A)$ is the free A -module with basis the $\gamma_{m_1}(\theta_1)\gamma_{m_2}(\theta_2)\dots\gamma_{m_d}(\theta_d)$ with $\sum m_i \leq r$ and the first assertion is clear. We see also that the A -module $\tilde{F}_r \mathcal{O}_1^{cris}(A) / \tilde{F}_{r-1} \mathcal{O}_1^{cris}(A)$ is free with the images of the $\gamma_{m_1}(\theta_1)\gamma_{m_2}(\theta_2)\dots\gamma_{m_d}(\theta_d)$ (or, equivalently of the $\gamma_{pm_1}(\bar{t}_1)\gamma_{pm_2}(\bar{t}_2)\dots\gamma_{pm_d}(\bar{t}_d)$) for $\sum m_i = r$ as a basis.

Let $x, y \in J(A)$. For all $m \in \mathbb{N}$, we have $\gamma_{pm}(x+y) = \sum_{i+j=m} \gamma_{pi}(x)\gamma_{pj}(y) + e(x, y)$ with $e(x, y) \in \tilde{F}_{m-1} \mathcal{O}_1^{cris}(A)$. From that and from the fact that if $a \in$

$\tilde{F}_m \mathcal{O}_1^{cris}(A)$ and $b \in \tilde{F}_n \mathcal{O}_1^{cris}(A)$, then $xy \in \tilde{F}_{n+m} \mathcal{O}_1^{cris}(A)$, we deduce the fact that f_r is well defined. The A -module $(J_1^{[r]}/J_1^{[r+1]})(A)$ is free with the images of the $\gamma_{m_1}(\bar{t}_1)\gamma_{m_2}(\bar{t}_2)\dots\gamma_{m_d}(\bar{t}_d)$ for $\sum m_i = r$ as a basis. As the A -linear map f_r sends the basis onto a basis of the free A -module $\tilde{F}_r \mathcal{O}_1^{cris}(A)/\tilde{F}_{r-1} \mathcal{O}_1^{cris}(A)$, f_r is bijective. \square

REMARKS 7.4.6 - i) The reason for the sign $(-1)^r$ is that we want f_r to be “morally” the Frobenius divided out by p^r . In characteristic 0, we have $\gamma_m(x^p) = p^m u_m \gamma_{pm}(x)$, with $u_m \in \mathbb{Z}_{(p)}$, congruent to $(-1)^m \pmod p$.
 ii) We may say that \mathcal{O}_1^{cris} is a *ring object in the category of F -zips*, i.e this is a ring, which is a \mathcal{O} -Algebra and for $r, s \in \mathbb{N}$, we have $\tilde{F}^r \mathcal{O}_1^{cris} \times \tilde{F}^s \mathcal{O}_1^{cris} \subset \tilde{F}^{r+s} \mathcal{O}_1^{cris}$ and $\tilde{F}_r \mathcal{O}_1^{cris} \times \tilde{F}_s \mathcal{O}_1^{cris} \subset \tilde{F}_{r+s} \mathcal{O}_1^{cris}$.

7.5 *End of the proof of theorem 7.2.1 and of the construction of the structure of an F -zip on \mathcal{O}_1^{cris}*

By the flatness of the p -adic ring \mathcal{G} it suffices to show theorem 7.2.1 for $n = 1$. First it follows inductively from theorem 7.4.5 (ii) and remark 7.4 (i) that for all $r \in \mathbb{N}$ we have

$$\tilde{F}_r = F_r \quad \text{and} \quad \tilde{F}^r = F^r$$

(for the rings considered in the previous section): The start of the induction is the fact that we have $F^0 = \mathcal{O}_1^{cris} = \tilde{F}^0$ by definition, and that the Frobenius $\varphi = \varphi_0 : \mathcal{O}_1^{cris} \rightarrow \mathcal{O}_1^{cris}$ has image $F_0 \mathcal{O}_1^{cris} = \mathcal{O} \xrightarrow{f} \mathcal{O}_1^{cris}$ and kernel $J_1^{[1]}$. We conclude that $F_0 = \mathcal{O} = \tilde{F}_0$ and $F^1 = \ker(\varphi) = \tilde{F}^1$. Then, by remark 7.4 (i) the map f_r in theorem 7.4.5 (ii) can be identified with the map induced by φ_r , which gives the induction steps. In fact, if $F_s = \tilde{F}_s$ for $s \leq r - 1$ and $F^s = \tilde{F}^s$ for $s \leq r$, then the two exact sequences

$$0 \rightarrow F^{r+1} \rightarrow F^r \xrightarrow{\varphi_r} F_r/F_{r-1} \rightarrow 0$$

$$0 \rightarrow \tilde{F}^{r+1} \rightarrow \tilde{F}^r \xrightarrow{\varphi_r} \tilde{F}_r/\tilde{F}_{r-1} \rightarrow 0$$

imply that $F_s = \tilde{F}_s$ for $s \leq r$ and $F^s = \tilde{F}^s$ for $s \leq r + 1$. This now shows that $\cap_r F^r = 0$ (by proposition 7.4.4) and that $\cup_r F_r = \mathcal{O}_1^{cris}$ (by theorem 7.4.5 (i)), showing the surjectivity in theorem 7.2.1, and the remaining property of F -zips.

8 φ -GAUGE-COHOMOLOGY - A REFINEMENT OF CRYSTALLINE COHOMOLOGY

Let k be a perfect field of characteristic p , let $W_n = W_n(k)$ for $n \in \mathbb{N}$, and let X be a syntomic variety over k .

8.1 *The definition of gauge-cohomology*

We define the i -th φ -gauge cohomology of level n of X by

$$H_g^i(X, W_n) = H_{syn}^i(X, \mathcal{G}_n)$$

This is a φ - W_n -gauge by transport of structure: We let $H_g^i(X, W_n)^r := H_{syn}^i(X, \mathcal{G}_n^r)$, and let the structural maps

$$H_g^i(X, W_n)^r \xrightleftharpoons[v]{f} H_g^i(X, W_n)^{r+1}$$

be induced by the morphisms

$$\mathcal{G}_n^r \xrightleftharpoons[v]{f} \mathcal{G}_n^{r+1}.$$

Moreover, the σ -linear map of W_n -modules

$$\begin{aligned} \varphi : H_g^i(X, W_n)^{+\infty} = \lim_{r \rightarrow +\infty, f} H_g^i(X, W_n)^r \longrightarrow \\ \lim_{r \rightarrow -\infty, v} H_g^i(X, W_n) = H_g^i(X, W_n)^{-\infty} \end{aligned}$$

is induced by the isomorphism $\varphi : \mathcal{G}_n^{+\infty} \longrightarrow \mathcal{G}_n^{-\infty}$, i.e., the isomorphisms

$$\begin{aligned} \lim_{r \rightarrow +\infty, f} H_{syn}^i(X, \mathcal{G}_n^r) \cong H_{syn}^i(X, \mathcal{G}_n^{+\infty}) \xrightarrow{\varphi} \\ \sim H_{syn}^i(X, \mathcal{G}_n^{-\infty}) \cong \lim_{r \rightarrow -\infty, v} H_{syn}^i(X, \mathcal{G}_n^r) \end{aligned}$$

where the outer isomorphisms come from the commuting of cohomology with limits.

We note that

$$H_g^i(X, W_n)^{-\infty} = H_{syn}^i(X, \mathcal{G}^{-\infty}) = H_{syn}^i(X, \mathcal{O}^{cris}) \cong H_{cris}^i(X/W_n)$$

by the comparison theorem of Fontaine and Messing [FM]. In this way, we achieved the refinement of crystalline cohomology announced in section 3.3.

We note also that

$$H_g^i(X, W_n)^{r-1} \xleftarrow[\sim]{v} H_g^i(X, W_n)^r$$

is an isomorphism for $r \leq 0$, because this holds for $\mathcal{G}_n^{r-1} \xleftarrow[v]{v} \mathcal{G}_n^r$. Hence the gauge $H_g^i(X, W_n)$ is effective (concentrated in degrees ≥ 0), and we have $H_g^i(X, W_n)^{-\infty} = H_g^i(X, W_n)^0 = H_{cris}^i(X/W_n)$.

8.2 The de Rham gauge of a variety X over k

We denote by $\mathcal{C}_b^{(p)}(\mathcal{O}_X)$ the following category:

- An object is a bounded complex

$$(C') \quad \dots \rightarrow C^{n-1} \rightarrow C^n \xrightarrow{d^n} C^{n+1} \rightarrow \dots$$

of \mathcal{O}_X -modules of finite type, whose differentials (which are additive) are \mathcal{O}_X^p -linear (i.e., satisfy $d^n(a^p x) = a^p d^n x$ for any $n \in \mathbb{Z}$, any local section a of \mathcal{O}_X and x of C^n for any $n \in \mathbb{Z}$).

- A morphism $\alpha : C' \rightarrow D'$ is a collection of \mathcal{O}_X -linear maps $\alpha^n : C^n \rightarrow D^n$, for $n \in \mathbb{Z}$, such that the diagram

$$\begin{array}{ccccccc} \dots & \rightarrow & C^{n-1} & \rightarrow & C^n & \rightarrow & C^{n+1} & \rightarrow & \dots \\ & & \downarrow & & \downarrow & & \downarrow & & \\ \dots & \rightarrow & D^{n-1} & \rightarrow & D^n & \rightarrow & D^{n+1} & \rightarrow & \dots \end{array}$$

is commutative.

This is an abelian (\mathcal{O}_X^p -linear) category.

Because k is perfect, we have $\Omega_{X/k}^1 = \Omega_X^1 := \Omega_{X/\mathbb{Z}}^1$. The de Rham complex

$$(\Omega'_X) \quad \mathcal{O}_X \rightarrow \Omega_X^1 \rightarrow \dots \rightarrow \Omega_X^{n-1} \rightarrow \Omega_X^n \rightarrow \Omega_X^{n+1} \rightarrow \dots$$

is an object of the above category (We adopt the following convention: if a complex C' starts with a given term on the left, this term is in degree 0 and $C^n = 0$ for $n < 0$).

The restriction of scalars via the absolute Frobenius $\sigma : \mathcal{O}_X \rightarrow \mathcal{O}_X$ defines an additive, exact and faithful functor from $\mathcal{C}_b^{(p)}(\mathcal{O}_X)$ to itself: If C' is an object of $\mathcal{C}_b^{(p)}(\mathcal{O}_X)$, we let $(\sigma C)^n = {}_\sigma(C^n)$ and take the same differentials. In other words, the underlying complex of sheaves of abelian groups is C' , but we change the structure of \mathcal{O}_X -module on each C^n : the multiplication of a local section x of σC^n by a local section a of \mathcal{O}_X is the multiplication of x by a^p in C^n .

Let us recall that, for any integer $n \in \mathbb{N}$, the *Cartier isomorphism* is an isomorphism of sheaves of abelian groups

$$c : \mathcal{H}^n(\Omega_X^\cdot) \rightarrow \Omega_X^n$$

satisfying $c(a^p \alpha) = ac(\alpha)$ for a (resp. α) any local section of \mathcal{O}_X (resp. $\mathcal{H}^n(\Omega_X^\cdot)$). It is characterized by the fact that, if a_0, a_1, \dots, a_n are local sections of \mathcal{O}_X , then $c^{-1}(a_0 da_1 \wedge \dots \wedge da_n)$ is the image of the closed form $a_0^p a_1^{p-1} a_2^{p-1} \dots a_n^{p-1} da_1 \wedge da_2 \wedge \dots \wedge da_n$.

We can also regard c as an \mathcal{O}_X -linear morphism

$$\sigma \mathcal{H}^n(\Omega_X^\cdot) \rightarrow \Omega_X^n.$$

If $Z\Omega_X^n$ denotes the kernel of $d : \Omega_X^n \rightarrow \Omega_X^{n+1}$, the map c induces an exact sequence of sheaves of \mathcal{O}_X -modules

$${}_{\sigma}\Omega_X^{n-1} \xrightarrow{d} {}_{\sigma}Z\Omega_X^n \xrightarrow{c} \Omega_X^n \rightarrow 0.$$

We observe that the fact that $d(a^p)\omega = a^pd\omega$ (for a local section of \mathcal{O}_X and ω local section of Ω_X^n) implies that ${}_{\sigma}Z\Omega_X^n$ is in fact a sub \mathcal{O}_X -module of ${}_{\sigma}\Omega_X^n$.

We are now ready to construct the de Rham φ -gauge $G_1(X) = (G_1^\cdot(X), f, v, \varphi)$ of X , which is a φ -gauge in $\mathcal{C}_b^{(p)}(\mathcal{O}_X)$:

– For $r < 0$, $G_1^r(X)$ is the de Rham complex Ω_X^\cdot :

$$\mathcal{O}_X \xrightarrow{d} \Omega_X^1 \xrightarrow{d} \dots \rightarrow \Omega_X^{n-1} \xrightarrow{d} \Omega_X^n \xrightarrow{d} \Omega_X^{n+1} \xrightarrow{d} \dots$$

– for $r \geq 0$, $G_1^r(X)$ is the complex

$${}_{\sigma}\mathcal{O}_X \xrightarrow{d} {}_{\sigma}\Omega_X^1 \xrightarrow{d} \dots \xrightarrow{d} {}_{\sigma}\Omega_X^{r-1} \xrightarrow{d} {}_{\sigma}Z\Omega_X^r \xrightarrow{dc} \Omega_X^{r+1} \xrightarrow{d} \dots \xrightarrow{d} \Omega_X^n \xrightarrow{d} \dots$$

– the map $f : G_1^r(X) \rightarrow G_1^{r+1}(X)$ is 0 for $r < 0$ and is, for $r \geq 0$,

$$\begin{array}{ccccccccccc} {}_{\sigma}\mathcal{O}_X & \xrightarrow{d} & \dots & {}_{\sigma}\Omega_X^{r-1} & \xrightarrow{d} & {}_{\sigma}Z\Omega_X^r & \xrightarrow{dc} & \Omega_X^{r+1} & \xrightarrow{d} & \Omega_X^{r+2} & \xrightarrow{d} & \Omega_X^{r+3} & \dots \\ \downarrow \parallel & & & \downarrow \parallel & & \downarrow \text{incl} & & \downarrow 0 & & \downarrow 0 & & \downarrow 0 & \\ {}_{\sigma}\mathcal{O}_X & \xrightarrow{d} & \dots & {}_{\sigma}\Omega_X^{r-1} & \xrightarrow{d} & {}_{\sigma}\Omega_X^r & \xrightarrow{d} & {}_{\sigma}Z\Omega_X^{r+1} & \xrightarrow{dc} & \Omega_X^{r+2} & \xrightarrow{d} & \Omega_X^{r+3} & \dots \end{array}$$

– the map $v : G_1^{r+1}(X) \rightarrow G_1^r(X)$ is the identity for $r < -1$ and is, for $r \geq -1$,

$$\begin{array}{ccccccccccc} {}_{\sigma}\mathcal{O}_X & \xrightarrow{d} & \dots & {}_{\sigma}\Omega_X^{r-1} & \xrightarrow{d} & {}_{\sigma}Z\Omega_X^r & \xrightarrow{dc} & \Omega_X^{r+1} & \xrightarrow{d} & \Omega_X^{r+2} & \xrightarrow{d} & \Omega_X^{r+3} & \dots \\ \uparrow 0 & & & \uparrow 0 & & \uparrow 0 & & \uparrow c & & \uparrow \parallel & & \uparrow \parallel & \\ {}_{\sigma}\mathcal{O}_X & \xrightarrow{d} & \dots & {}_{\sigma}\Omega_X^{r-1} & \xrightarrow{d} & {}_{\sigma}\Omega_X^r & \xrightarrow{d} & {}_{\sigma}Z\Omega_X^{r+1} & \xrightarrow{dc} & \Omega_X^{r+2} & \xrightarrow{d} & \Omega_X^{r+3} & \dots \end{array}$$

Clearly $vf = fv = 0$, hence we have a gauge. We see also that for all $r \in \mathbb{Z}$, the map $(f, v) : G_1^r(X) \rightarrow G_1^{r+1}(X) \oplus G_1^{r-1}(X)$ is injective. Hence we have a strict gauge.

Because $c : Z_{\sigma}\mathcal{O}_X \rightarrow \mathcal{O}_X$ is an isomorphism, v_{-1} is an isomorphism, hence we have an effective gauge.

–If d is the dimension of X , we have

$$G_1^r(X) = {}_{\sigma}\Omega_X^\cdot \quad \text{for } r \geq d$$

(for $r = d$, because ${}_{\sigma}Z\Omega_X^d = {}_{\sigma}\Omega_X^d$) and $f : G_1^r(X) \rightarrow G_1^{r+1}(X)$ is an isomorphism if $r \geq d$.

Hence, we have a gauge concentrated in the interval $[0, d]$, with $G_1^{-\infty}(X) = \Omega_X^\cdot$ and $G_1^{+\infty}(X) = {}_{\sigma}\Omega_X^\cdot$. We define the isomorphism

$$\varphi : G_1^{+\infty}(X) \rightarrow {}_{\sigma}G_1^{-\infty}(X)$$

as the identity on ${}_{\sigma}\Omega_X^\cdot$, which finishes the definition of the de Rham φ -gauge $G_1(X)$.

THEOREM 8.2.1. *Let $\alpha : X_{\text{syn}} \rightarrow X_{\text{Zar}}$ be the morphism of sites coming from the fact that the Zariski topology is coarser than the syntomic cohomology. There is a canonical isomorphism*

$$R\alpha_*\mathcal{G}_1 \xrightarrow{\sim} G_1 .$$

Proof. This follows via the same methods proving the Fontaine-Messing isomorphism

$$H_{\text{syn}}^i(X, \mathcal{O}_1^{\text{cris}}) \cong H_{\text{cris}}^i(X/k) = H_{dR}^i(X/k) ,$$

using canonical isomorphisms following from results of Berthelot [Be]

$$R\alpha_*I_1^{[r]} \cong \Omega_X^{\geq r} ,$$

where the complex on the right is obtained by naive truncation, i.e., is the upper part, starting with Ω_X^r , of the de Rham complex. \square

COROLLARY 8.2.2. *Let X be a proper variety over k . Then the following holds.*

(a) *The gauge cohomology $H_g^i(X, W_n)$ is of finite type and is concentrated in the interval $[0, i]$, and it vanishes for $i \geq 2d$.*

(b) *One has $H_g^i(X, W_n)^0 = H_{\text{cris}}^i(X/W_n)$.*

(c) *If X is smooth, proper, and irreducible of pure dimension d , then the Poincaré duality for crystalline cohomology extends to a perfect duality of φ - W_n -gauges*

$$H_g^i(X, W_n) \times H_g^{2d-i}(X, W_n) \rightarrow H_g^{2d}(X, W_n) \xrightarrow{\sim} W_n(-d) .$$

Here the W_n -gauge $W_n(-d)$ extends to a φ - W_n -gauge by defining

$$\varphi : W_n(-d)^{+\infty} = W_n \rightarrow W_n = W_n(-d)^{-\infty}$$

as the Frobenius σ on W_n , which is σ -linear.

Proof. All questions are easily reduced to the case $n = 1$. But by theorem 8.2.1, we have $H_g^i(X, W_1) = H_{\text{Zar}}^i(X, G_1^r)$, and thus the claim follows from classical Serre duality, and the explicit shape of the de Rham gauge $G_1(X)$ defined above. \square

REFERENCES

- [Be] P. Berthelot, *Cohomologie cristalline des schémas de caractéristique $p > 0$* , Lecture Notes in Mathematics, 407. Springer-Verlag, Berlin-New York, 604 pp, 1974.
- [FM] J.-M. Fontaine, W. Messing, *p -adic periods and p -adic étale cohomology*, in: Current trends in arithmetical algebraic geometry (Arcata, Calif., 1985), 179–207, Contemp. Math., 67, Amer. Math. Soc., Providence, RI, 1987.

- [GW] U. Görtz, T. Wedhorn, *Algebraic Geometry I*, Vieweg+Teubner, Wiesbaden, 2010.
- [Ha] R. Hartshorne, *Algebraic Geometry*, Graduate Texts in Math., 52, Springer, New York, 1977.
- [Ki] M. Kisin, *Modularity for some geometric Galois representations. With an appendix by Ofer Gabber*, London Math. Soc. Lecture Note Ser., 320, L-functions and Galois representations, 438–470, Cambridge Univ. Press, Cambridge, 2007.
- [LZ] A. Langer, T. Zink, *De Rham-Witt cohomology and displays*, Doc. Math. 12 (2007), 147–191.
- [MW] B. Moonen, T. Wedhorn, *Discrete invariants of varieties in positive characteristic*, Int. Math. Res. Not. 2004, no. 72, 3855–3903.
- [Schn] F. Schnellinger, *Rigid gauges and F -zips, and the fundamental sheaf of gauges G_n* , Ph.D. thesis, University of Regensburg, 2009, <http://epub.uni-regensburg.de/12290/>.
- [Wed] T. Wedhorn, *De Rham cohomology of varieties over fields of positive characteristic*, in: Higher-dimensional geometry over finite fields, 269–314, NATO Sci. Peace Secur. Ser. D Inf. Commun. Secur., 16, IOS, Amsterdam, 2008.
- [Wid] M. Wid, *Displays and gauges*, Ph.D. thesis, University of Regensburg, 2012, <http://epub.uni-regensburg.de/25489/>.

Jean-Marc Fontaine
Université Paris-Sud
France

Uwe Jannsen
Universität Regensburg
Germany

