

ALGEBRAIC SLICE SPECTRAL SEQUENCES

DOMINIC LEON CULVER, HANA JIA KONG, J.D. QUIGLEY

Received: December 15, 2020

Revised: July 28, 2021

Communicated by Mike Hill

ABSTRACT. For certain motivic spectra, we construct a square of spectral sequences relating the effective slice spectral sequence and the motivic Adams spectral sequence. We show the square can be constructed for connective algebraic K-theory, motivic Morava K-theory, and truncated motivic Brown–Peterson spectra. In these cases, we show that the \mathbb{R} -motivic effective slice spectral sequence is completely determined by the ρ -Bockstein spectral sequence. Using results of Heard, we also obtain applications to the Hill–Hopkins–Ravenel slice spectral sequences for connective Real K-theory, Real Morava K-theory, and truncated Real Brown–Peterson spectra.

2020 Mathematics Subject Classification: 14F42, 55P42, 55P91, 55T05, 55T15

Keywords and Phrases: Motivic Adams spectral sequence, slice spectral sequence, algebraic slice spectral sequence

1 INTRODUCTION

1.1 MOTIVATION AND MAIN THEOREMS

Two of the chief computational tools in motivic stable homotopy theory are the effective slice spectral sequence [27, 38, 41] and the motivic Adams spectral sequence [11, 31]. The purpose of this paper is to systematically relate the effective slice spectral sequence and the motivic Adams spectral sequence.

The effective slice spectral sequence has many applications, such as a new proof of Milnor’s Conjecture on quadratic forms [35] and the calculation of the first stable homotopy group of motivic spheres [36] which identifies certain motivic stable stems with variants of K-theory. In general, the effective slice spectral sequence is an excellent tool for working over general base schemes

because slices are often expressible in terms of arithmetic invariants like motivic cohomology.

The motivic Adams spectral sequence is a powerful resource for making large-scale computations over specific base fields, such as the complex numbers [22], the real numbers [6], and finite fields [43]. Computations over $\text{Spec}(\mathbb{C})$ have brought about remarkable progress on the classical stable homotopy groups of spheres, and applications of \mathbb{R} -motivic computations to C_2 -equivariant homotopy have been the topic of much recent work, such as [4, 5, 23].

Our first main theorem fits the effective slice spectral sequence and motivic Adams spectral sequence into a square of spectral sequences.

Theorem A (Comparison square, Theorem 3.4). Let $E \in \text{SH}(k)$ be a motivic spectrum which is algebraically n -sliceable over k (Definition 3.1). There is a square of spectral sequences of the form

$$\begin{array}{ccc}
 & \bigoplus_{q \geq 0} \text{Ext}_A^{***}(H_{**} s_q^n E) & \\
 \begin{array}{c} \swarrow \\ \text{\scriptsize } n\text{-aESSS} \end{array} & & \begin{array}{c} \searrow \\ \text{\scriptsize } \bigoplus_{q \geq 0} m\text{ASS} \end{array} \\
 \text{Ext}_A^{***}(H_{**} E) & & \bigoplus_{q \geq 0} \pi_{**} s_q^n E \quad (1) \\
 \begin{array}{c} \searrow \\ \text{\scriptsize } m\text{ASS} \end{array} & & \begin{array}{c} \swarrow \\ \text{\scriptsize } n\text{-ESSS} \end{array} \\
 & \pi_{**} E &
 \end{array}$$

The spectral sequences in the square are discussed in Sections 2 and 3.

Remark. In this paper, we refer to a “square of spectral sequences” whenever one can run the spectral sequences on one half of the square or the other half of the square to attempt to compute the same groups. In particular, a square of spectral sequences is not the same as a commuting square in the category of spectral sequences of abelian groups, and we make no claims about convergence to the same filtration.

The condition that $E \in \text{SH}(k)$ is algebraically n -sliceable is defined in terms of the homology of its slices and slice covers. Our second main theorem says that several familiar motivic spectra are algebraically n -sliceable.

Theorem B (Theorems 3.7, 3.9, and 3.16, and Remark 3.17).

Let k be a perfect field. Theorem A¹ applies to the following motivic spectra:

1. The effective cover of algebraic K-theory, kgl , if the characteristic of k is not two.
2. For all $n \geq 1$, the connective motivic Morava K-theory, $k(n)$, if the exponential characteristic of k is prime to the characteristic of the coefficients of H_{**} .
3. For all $n \geq 1$, the n -th truncated motivic Brown–Peterson spectrum, $BPGL\langle n \rangle$, if k has characteristic zero.

¹Or its very effective variant, see Remark 3.5.

4. The very effective cover of hermitian K-theory, kq , if the characteristic of k is not two.

The ρ -Bockstein spectral sequence [15] is another important device in computational motivic stable homotopy theory. Theorems A and B can be used to prove a precise relation between differentials in the ρ -Bockstein spectral sequence and the \mathbb{R} -motivic effective slice spectral sequence:

Theorem C (Theorems 4.10 and 4.12). Let $k = \mathbb{R}$. There are 1-to-1 correspondences between the following differentials:

1. ρ -Bockstein spectral sequence d_3 -differentials and effective slice spectral sequence d_1 -differentials for kgl .
2. ρ -Bockstein spectral sequence $d_{2^{n+1}-1}$ -differentials and type n effective slice spectral sequence d_1 -differentials for $k(n)$.
3. For each $1 \leq i \leq n$, ρ -Bockstein spectral sequence $d_{2^{i+1}-1}$ -differentials and effective slice spectral sequence d_{2^i-1} -differentials for $BPGL\langle n \rangle$.

Moreover, the (type n) effective slice spectral sequence differentials are completely determined by knowledge of the corresponding ρ -Bockstein spectral sequence differentials.

Formal similarities between the ρ -Bockstein and effective slice spectral sequence differentials have been observed in previous work, such as [14, Sec. 6]. Theorem C and its proof may be viewed as a rigorous explanation for these similarities.

Finally, our computations have consequences in the C_2 -equivariant stable homotopy theory. The Hill–Hopkins–Ravenel slice spectral sequence [16] has enjoyed many applications since its use in the solution to the Kervaire invariant one problem, and thus we are motivated to calculate differentials for common C_2 -spectra. In previous work, Heard [14] compared the \mathbb{R} -motivic effective slice filtration and the Hill–Hopkins–Ravenel slice filtration [16] and deduced certain differentials in the effective slice spectral sequence using previously computed differentials in the HHR slice spectral sequence. Using Heard’s comparison results and Theorem C, we reverse this logic to produce differentials in the HHR slice spectral sequences for the genuine C_2 -spectra $k\mathbb{R}$, $k\mathbb{R}(n)$, and $BPR\langle n \rangle$ from known slice spectral sequence differentials. Our results recover work of Dugger [10] and Hill–Hopkins–Ravenel [16].

Theorem D (Corollaries 5.7, 5.8, and 5.9). The following statements hold:

1. The nontrivial differentials in the HHR slice spectral sequence for $\pi_{**}^{C_2}k\mathbb{R}$ are determined via Leibniz rule by

$$d_3(\tau^2) = \rho^3 \bar{v}_1 \quad \text{and} \quad d_3\left(\frac{\gamma}{\rho^3 \tau^2}\right) = \frac{\gamma}{\tau^4} \bar{v}_1.$$

2. Let $n \geq 1$. There is a nontrivial differential in the HHR slice spectral sequence for $\pi_{**}^{C_2} k\mathbb{R}\langle n \rangle$ of the form $d_{2^{n+1}-1}(\tau^{2^{n+1}}) = \rho^{2^{n+1}-1}\bar{v}_n$.
3. Let $n \geq 1$. There are nontrivial differentials in the HHR slice spectral sequence for $\pi_{**}^{C_2} BP\mathbb{R}\langle n \rangle$ of the form $d_{2^{i+1}-1}(\tau^{2^{i+1}}) = \rho^{2^{i+1}-1}\bar{v}_i$, $1 \leq i \leq n$.

The relevant elements are recalled in Remark 5.1.

1.2 ORGANIZATION

In Section 2, we recall the functors and spectral sequences discussed in this paper, such as (equivariant) Betti realization, variants of the effective and very effective slice spectral sequences, and the motivic Adams spectral sequence.

In Section 3, we define when a motivic spectrum is algebraically n -sliceable and prove Theorem A. We then analyze several examples and non-examples of algebraically n -sliceable spectra, and in particular we prove Theorem B.

In Section 4, we recall the ρ -Bockstein spectral sequence and differentials in the motivic Adams spectral sequence; we then prove Theorem C.

In Section 5, we recall Heard's comparison of the motivic and C_2 -equivariant slice filtrations and apply it to prove Theorem D.

1.3 CONVENTIONS

1. After Section 2, everything is implicitly (p, η) -complete.²
2. k is a perfect field of exponential characteristic prime to p .
3. SH denotes the classical stable homotopy category, $\mathrm{SH}(k)$ denotes the motivic stable homotopy category over $\mathrm{Spec}(k)$, and SH^{C_2} denotes the C_2 -equivariant stable homotopy category.
4. H denotes the classical, equivariant, or motivic mod p Eilenberg-MacLane spectrum.
5. \mathbb{M}_p^k denotes the mod p motivic cohomology of $\mathrm{Spec}(k)$.
6. A (resp. A^\vee) denotes the motivic Steenrod algebra (resp. its \mathbb{M}_p^k -linear dual).
7. If M is an A -module, M^\vee denotes its \mathbb{M}_p^k -linear dual.
8. The homotopy groups of $E \in \mathrm{SH}(k)$ are denoted $\pi_{m,u}^k(E)$, or if there is no confusion about the base field, $\pi_{m,u}(E)$. Our bigrading convention is that $\pi_{m,u}^k(E) \cong \mathrm{SH}(k)[\Sigma^\infty(S_\bullet^1)^{\wedge m-u} \wedge \mathbb{G}_m^{\wedge u}, E]$.

²This agrees with p -completion in many cases we consider. See e.g. [20, Thm. 1].

$\pi_{**}^{C_2} H\mathbb{F}_2$			
element	alt. name	bidegree	$RO(C_2)$ -degree
ρ	a_σ	$(-1, -1)$	$-\sigma$
τ	u_σ	$(0, -1)$	$1 - \sigma$
$\frac{\gamma}{\tau}$	θ	$(0, 2)$	$2\sigma - 2$
$\pi_{**}^{C_2} H\mathbb{Z}_2$			
element	alt. name	bidegree	$RO(C_2)$ -degree
ρ	a_σ	$(-1, -1)$	$-\sigma$
τ^2	$u_{2\sigma}$	$(0, -2)$	$2 - 2\sigma$
$\frac{\gamma}{\tau}$	$2u_{2\sigma}^{-1}$	$(0, 2)$	$2\sigma - 2$
$\frac{\gamma}{\tau^2}$	θ	$(0, 3)$	$3\sigma - 3$

Table 1: Names, alternate names, and degrees of elements in C_2 -equivariant homotopy groups.

9. The homotopy groups of $E \in SH^{C_2}$ are denoted $\pi_{m,u}^{C_2}(E)$. Our bigrading convention is that $\pi_{m,u}^{C_2}(E) \cong SH^{C_2}[\Sigma^\infty S^{m-u} \wedge S^{u\sigma}, E]$ where σ is the one-dimensional real sign representation of C_2 .
10. We will often write $\text{Ext}_A^{***}(E)$ in place of $\text{Ext}_A^{***}(H_{**}E)$.
11. We use the following abbreviations for spectral sequences:
 - (a) mASS: motivic Adams spectral sequence.
 - (b) ESSS and VSSS: effective and very effective slice spectral sequence.
 - (c) aESSS and aVSSS: algebraic effective and very effective slice spectral sequence.
 - (d) n -(-)SSS: type n (-)SSS, with (-) either E, V, aE, or aV.
 - (e) H_* - n -(-)SSS: homological type n (-)SSS, with (-) as above.
 - (f) ρ -BSS: ρ -Bockstein spectral sequence.
12. Notation for C_2 -equivariant homotopy elements is contained in Table 1.

ACKNOWLEDGEMENTS

The authors thank the organizers of the BIRS workshop “Equivariant stable homotopy theory and p -adic Hodge theory”, where some early stages of this work were completed. The authors thank Mark Behrens, Bob Bruner, Drew Heard, Dan Isaksen, XiaoLin Danny Shi, Inna Zakharevich, and an anonymous referee for helpful comments and questions. Some of the work on this paper was done while the first author was at the Max Planck Institute for Mathematics. He would like to thank the Institute for their support.

2 MOTIVIC SLICE AND ADAMS SPECTRAL SEQUENCES

We start by discussing the functors and spectral sequences which we will study in the rest of the paper.

2.1 (EQUIVARIANT) BETTI REALIZATION FUNCTORS

Since we will mention them in our discussion of the various spectral sequences below, we start by recalling the Betti realization and equivariant Betti realization functors which relate the motivic, classical, and C_2 -equivariant stable homotopy categories.

Suppose the base field k has a complex embedding. The Betti realization functor [32]

$$\mathrm{Re} : \mathrm{SH}(k) \rightarrow \mathrm{SH} \quad (2)$$

is defined by sending $X \in \mathrm{SH}(k)$ to $X(\mathbb{C}) \in \mathrm{SH}$. Similarly, if k has a real embedding, the equivariant Betti realization functor [32]

$$\mathrm{Re}_{C_2} : \mathrm{SH}(k) \rightarrow \mathrm{SH}^{C_2} \quad (3)$$

is defined by sending $X \in \mathrm{SH}(k)$ to $X(\mathbb{C}) \in \mathrm{SH}^{C_2}$, where C_2 acts on $X(\mathbb{C})$ via complex conjugation.

2.2 THE TYPE n EFFECTIVE SLICE SPECTRAL SEQUENCE

We now introduce one of our main objects of study, the type n effective slice spectral sequence, which generalizes the effective slice spectral sequence. Throughout this section, we fix a prime p .

Let $\mathrm{SH}^{\mathrm{eff}}(k)$ be the localizing subcategory of $\mathrm{SH}(k)$ generated by $\{\Sigma_+^\infty X : X \in \mathrm{Sm}_k\}$. These subcategories form the *effective slice filtration*

$$\dots \subset \Sigma^{2q+2, q+1} \mathrm{SH}^{\mathrm{eff}}(k) \subset \Sigma^{2q, q} \mathrm{SH}^{\mathrm{eff}}(k) \subset \Sigma^{2q-2, q-1} \mathrm{SH}^{\mathrm{eff}}(k) \subset \dots \quad (4)$$

The inclusion $i_q : \Sigma^{2q, q} \mathrm{SH}^{\mathrm{eff}}(k) \subset \mathrm{SH}(k)$ is left adjoint to the functor $r_q : \mathrm{SH}(k) \rightarrow \Sigma^{2q, q} \mathrm{SH}^{\mathrm{eff}}(k)$.

DEFINITION 2.1. The composite $f_q := i_q \circ r_q$ is the q -th *effective slice cover* and $s_q := \mathrm{cofib}(f_{q+1} \rightarrow f_q)$ is the q -th *effective slice*.

For each $n \geq 1$, we define the q -th *type n effective slice cover* $f_q^n := f_{q(p^n-1)}$ and the q -th *type n effective slice* $s_q^n := \mathrm{cofib}(f_{q+1}^n \rightarrow f_q^n)$. Observe that $f_q^1 = f_q$ if $p = 2$.

For any motivic spectrum $E \in \mathrm{SH}(k)$ and any integer $n \geq 1$, we obtain a

\mathbb{Z} -indexed tower of fibrations

$$\begin{array}{ccc}
 \vdots & & \\
 \downarrow & & \\
 f_{q+1}^n E & \longrightarrow & s_{q+1}^n E \\
 \downarrow & & \\
 f_q^n E & \longrightarrow & s_q^n E \\
 \downarrow & & \\
 f_{q-1}^n E & \longrightarrow & s_{q-1}^n E \\
 \downarrow & & \\
 \vdots & &
 \end{array} \tag{5}$$

called the *type n effective slice tower*. Applying motivic homotopy groups gives rise to the *type n effective slice spectral sequence*

$${}^{n,\text{eff}}E_{m,q,u}^1(E) = \pi_{m,u} s_q^n E \Rightarrow \pi_{m,u} \text{sc}(E) \tag{6}$$

with differentials $d^r(E) : {}^{n,\text{eff}}E_{m,q,u}^r(E) \rightarrow {}^{n,\text{eff}}E_{m-1,q+r,u}^r(E)$. Let $f_n^q E$ denote the cofiber of $f_{q+1}^n E \rightarrow E$. The term $\text{sc}(E)$ in the abutment is the *slice completion* of E defined as the homotopy inverse limit $\text{sc}(E) := \lim_q f_q E$. For each n , we have $\text{sc}(E) \simeq \lim_q f_q^n E$ since the type n effective slice filtration is a sped up version of the ordinary effective slice filtration.

For each $E \in \text{SH}(k)$ we study below, there is an equivalence $\text{sc}(E) \simeq E$ by [17, Thm. 8.12]. More generally, [36, Thm. 3.50] implies that in very general situations, $\text{sc}(E)$ may be identified with the η -completion of E .

2.3 THE TYPE n VERY EFFECTIVE SLICE SPECTRAL SEQUENCE

We now discuss a variant of the type n effective slice spectral sequence. Throughout this section, fix an integer $n \geq 1$ and prime p .

Let $\text{SH}^{\text{veff}}(k)$ be the smallest full subcategory of $\text{SH}(k)$ that contains all suspension spectra of smooth k -schemes of finite type and which is closed under all homotopy colimits and extensions. Repeating the construction from Section 2.2 with $\text{SH}^{\text{veff}}(k)$ in place of $\text{SH}^{\text{eff}}(k)$ yields the q -th type n very effective slice cover \tilde{f}_q^n and the q -th type n very effective slice \tilde{s}_q^n . These fit into a tower analogous to (5) which gives rise to the *type n very effective slice spectral sequence*

$${}^{n,\text{veff}}E_{m,q,u}^1(E) = \pi_{m,u} \tilde{s}_q^n E \Rightarrow \pi_{m,u} \text{vsc}(E) \cong \pi_{m,u} E \tag{7}$$

with differentials $d^r(E) : {}^{n,\text{veff}}E_{p,q,u}^r(E) \rightarrow {}^{n,\text{veff}}E_{p-1,q+r,u}^r(E)$. If we let $\tilde{f}_n^q E$ denote the cofiber of the map $f_{q+1}^n E \rightarrow E$, then the *very effective slice com-*

pletion $\text{vsc}(E) := \lim_q \tilde{f}_1^q E \simeq \lim_q \tilde{f}_n^q E$ is equivalent to E because $\tilde{s}_q^n E$ is $q(p^n - 1)$ -connected in Morel’s homotopy t-structure on $\text{SH}(k)$ for all $q \in \mathbb{Z}$.³

Remark 2.2. Suppose the ground field k admits a complex embedding. Then the very effective slice spectral sequence over $\text{Spec}(k)$ is compatible with the double-speed Postnikov filtration under Betti realization [13], so there is a map of spectral sequences from the very effective slice spectral sequence for $X \in \text{SH}(k)$ to the double-speed Postnikov spectral sequence of $\text{Re}(X)$.

Remark 2.3. Suppose the ground field k admits a real embedding. Then the very effective slice spectral sequence over $\text{Spec}(k)$ for a localized quotient of MGL is compatible with the Hill–Hopkins–Ravenel slice spectral sequence [16] of its equivariant Betti realization by [14, Thms. 5.15-5.16].

Remark 2.4. The effective and very effective slice filtrations coincide for localized quotients of MGL [14, Prop. 4.3] and Landweber exact motivic spectra [14, Prop. 4.11].

2.4 THE MOTIVIC ADAMS SPECTRAL SEQUENCE

Let $E \in \text{SH}(k)$ be a unital motivic spectrum and let $\overline{E} := \text{cofib}(S^{0,0} \rightarrow E)$. The canonical E -based Adams resolution of the sphere spectrum \mathbb{S} is given by

$$\begin{array}{ccccccc}
 \mathbb{S} & \longleftarrow & \Sigma^{-1,0}\overline{E} & \longleftarrow & \Sigma^{-2,0}\overline{E}^{\wedge 2} & \longleftarrow & \dots \\
 \downarrow & & \downarrow & & \downarrow & & \\
 E & & \Sigma^{-1,0}E \wedge \overline{E} & & \Sigma^{-2,0}E \wedge \overline{E}^{\wedge 2} & &
 \end{array} \tag{8}$$

The canonical E -based Adams resolution over a motivic spectrum $X \in \text{SH}(k)$ is obtained by smashing every term in (8) with X .

When $E = H$ is mod p motivic cohomology, we have $H_{**}(S^{0,0}) = \mathbb{M}_p^k$ and $\pi_{**}(H \wedge H) = A^\vee$ is the dual motivic Steenrod algebra [18, 42]. Applying motivic homotopy groups to (8) gives the motivic Adams spectral sequence

$$\text{mASS} E_2^{s,t,u} = \text{Ext}_{A^\vee}^{s,t,u}(\mathbb{M}_p^k, H_{**}(X)) \Rightarrow \pi_{t-s,u}(X_H^\wedge) \tag{9}$$

with $d_r : \text{mASS} E_r^{s,t,u} \rightarrow \text{mASS} E_r^{s+r,t+r-1,u}$. Here, X_H^\wedge is the H -nilpotent completion of E . By [20, Thm. 1][26, Cor. 6.1][29, Thm. 1.0.1], there is an equivalence $X_H^\wedge \simeq X_{(p,\eta)}^\wedge$ for X sufficiently nice⁴. Moreover, if k is a field of characteristic zero, and if $p > 2$ and $\mathcal{D}_p(k) < \infty$ and $-1 \in k$ is a sum of squares, or if $p = 2$ and k has finite virtual cohomological dimension, then $X_{(p,\eta)}^\wedge \simeq X_p^\wedge$ [20].

Remark 2.5. The motivic homology \mathbb{M}_p^k is reviewed in Section 4.1.

³See [38, Sec. 5] for a full discussion of convergence of the very effective slice spectral sequence.

⁴In this paper, “sufficiently nice” includes all motivic spectra X which are connective and cellular of finite type.

Remark 2.6. As in Remark 2.2, if the ground field k admits a complex embedding, the motivic Adams and Adams–Novikov filtrations of $X \in \text{SH}(k)$ are compatible with the Adams and Adams–Novikov filtrations of $\text{Re}(X) \in \text{SH}$. Thus Betti realization induces a map from the motivic Adams(–Novikov) spectral sequence to the classical Adams(–Novikov) spectral sequence. Similarly (c.f. Remark 2.3), if the ground field k admits a real embedding, then the motivic Adams and Adams–Novikov filtrations of $X \in \text{SH}(k)$ are compatible with the C_2 -equivariant Adams and Adams–Novikov filtrations of $\text{Re}_{C_2}(X) \in \text{SH}^{C_2}$. Thus equivariant Betti realization induces a map between the corresponding spectral sequences.

3 COMPARISON SQUARE AND EXAMPLES

In this section, we introduce a condition on a motivic spectrum which allows for the construction of a comparison square relating the effective slice spectral sequence and the motivic Adams spectral sequence. We prove that many common motivic spectra satisfy this condition, and we find an example which does not. From now on, we will implicitly p -complete everywhere.

3.1 ALGEBRAIC SLICE SPECTRAL SEQUENCES

The type n effective slice tower (5) is built out of cofiber sequences

$$f_{q+1}^n E \rightarrow f_q^n E \rightarrow s_q^n E.$$

Applying mod p motivic homology induces a long exact sequence

$$\cdots \rightarrow H_{i,j} f_{q+1}^n E \rightarrow H_{i,j} f_q^n E \rightarrow H_{i,j} s_q^n E \rightarrow H_{i-1,j} f_{q+1}^n E \rightarrow \cdots \quad (10)$$

for each $j \in \mathbb{Z}$.

DEFINITION 3.1. We say that $E \in \text{SH}(k)$ is *algebraically n -sliceable over k* if the long exact sequences (10) split into short exact sequences of A^\vee -comodules

$$0 \rightarrow H_{**} f_q^n E \xrightarrow{p} H_{**} s_q^n E \xrightarrow{l} H_{*-1,*} (f_{q+1}^n E) \rightarrow 0 \quad (11)$$

for each $q \in \mathbb{Z}$.

Similarly, we say that E is *algebraically sliceable over k* if the long exact sequences associated to the effective slice tower split into short exact sequences of A^\vee -comodules

$$0 \rightarrow H_{**} f_q E \xrightarrow{p} H_{**} s_q E \xrightarrow{l} H_{*-1,*} (f_{q+1} E) \rightarrow 0 \quad (12)$$

for each $q \in \mathbb{Z}$.

When the field k is clear, we will omit the phrase “over k ” for brevity.

Remark 3.2. If $p = 2$, being algebraically sliceable is the same as being algebraically 1-sliceable.

Applying the functor $\text{Ext}_{A^\vee}^{***}(\mathbb{M}_p^k, -)$ to (12) gives rise to long exact sequences which assemble into the unrolled exact couple

$$\begin{array}{ccccc}
 \dots \text{Ext}^{s,t,u}(f_0^n E) & \xleftarrow{\delta} & \text{Ext}^{s-1,t-1,u}(f_1^n E) & \xleftarrow{\delta} & \text{Ext}^{s-2,t-2,u}(f_2^n E) \dots \\
 \downarrow p_* & \nearrow \iota_* & \downarrow p_* & \nearrow \iota_* & \downarrow p_* \\
 \dots \text{Ext}^{s,t,u}(s_0^n E) & & \text{Ext}^{s-1,t-1,u}(s_1^n E) & & \text{Ext}^{s-2,t-2,u}(s_2^n E) \dots
 \end{array} \tag{13}$$

in which the dotted arrows have tridegree $(0, -1, 0)$. This gives rise to the algebraic type n effective slice spectral sequence

$${}^{n,\text{aESSS}} E_1^{q,s,t,u}(E) = \text{Ext}_{A^\vee}^{s,t,u}(\mathbb{M}_p^k, H_{**} s_q^n E) \tag{14}$$

with differentials

$$d_r : E_r^{q,s,t,u} \rightarrow E_r^{q+r,s-r+1,t-r,u}.$$

PROPOSITION 3.3. *The spectral sequence (15) converges conditionally to the Ext groups of $H_{**}(E)$:*

$${}^{n,\text{aESSS}} E_1^{q,s,t,u}(E) \implies \text{Ext}^{s+q,t+q,u}(H_{**} E). \tag{15}$$

Moreover, if E is effective, i.e. $E = f_0 E$, and the spectral sequence collapses at a finite stage, then the spectral sequence converges strongly.

Proof. We will verify that the spectral sequence (15) converges conditionally to the colimit $\text{Ext}^{***}(H_{**}(E))$ in the sense of Boardman [7, Def. 5.10]: for each fixed tridegree (s, t, u) , we will show that

$$\lim_k \text{Ext}^{s-k,t-k,u}(H_{**} f_k^n E) = 0, \tag{16}$$

$$\lim_k^1 \text{Ext}^{s-k,t-k,u}(H_{**} f_k^n E) = 0. \tag{17}$$

The limits above are taken along the maps δ in the unrolled exact couple. Fix a tridegree (s, t, u) . Observe that

$$\text{Ext}^{s-k,t-k,u}(H_{**} f_k^n E) = 0$$

whenever $k > s$, so (16) is clear. For (17), note that

$$\text{Ext}^{s-k,t-k,u}(H_{**} s_k^n E) = 0$$

for all $k > s$, so

$$p_* : \text{Ext}^{s-k,t-k,u}(H_{**} f_k^n E) \rightarrow \text{Ext}^{s-k,t-k,u}(H_{**} s_k^n E)$$

is zero for all $k > s$ and thus

$$\delta : \text{Ext}^{s-(k+1),t-(k+1),u}(H_{**} f_{k+1}^n E) \rightarrow \text{Ext}^{s-k,t-k,u}(H_{**} f_k^n E)$$

is surjective for all $k > s$. Therefore the inverse system $(\text{Ext}^{s-k,t-k,u}(H_{**}f_k^n E), \delta)$ is Mittag-Leffler, cf. [7, Pg. 9], and the equality (17) holds.

We have now shown that (15) converges conditionally to the colimit. We now justify strong convergence under the hypotheses that E is effective and the spectral sequence collapses at a finite stage. Since E is effective, (15) is a half-plane spectral sequence with entering differentials [7, Pg. 20].⁵ By [7, Remark, Pg. 20], we have $RE_\infty = 0$ whenever the spectral sequence collapses at a finite stage. The spectral sequence therefore converges strongly under our hypotheses by [7, Thm. 7.1]. \square

The same discussion carries over to the type n very effective slice filtration. In particular, we can define when $E \in \text{SH}(k)$ is *very algebraically n -sliceable over k* , and for such E we may define the *algebraic type n very effective slice spectral sequence*

$${}^{n,a}\text{VSSS} E_1^{q,s,t,u}(E) = \text{Ext}_{A\vee}^{s,t,u}(\mathbb{M}_p^k, H_{**}\tilde{s}_q^n E) \Rightarrow \text{Ext}_{A\vee}^{s,t,u}(\mathbb{M}_p^k, H_{**}\text{sc}(E)). \quad (18)$$

The spectral sequence converges conditionally. If E is very effective and the spectral sequence collapses at a finite stage, the spectral sequence converges strongly.

3.2 COMPARISON SQUARE

Our main tool for relating the type n effective slice spectral sequence and the motivic Adams spectral sequence is the following theorem, which assembles the spectral sequences (6), (9), and (15) discussed above into a square.

THEOREM 3.4 (Comparison square). *Let $E \in \text{SH}(k)$ be a k -motivic spectrum. If E is algebraically n -sliceable, then there is a square of spectral sequences of the form*

$$\begin{array}{ccc}
 & \bigoplus_{q \geq 0} \text{Ext}_A^{s,f,u}(H_{**}s_q^n E) & \\
 \begin{array}{c} \swarrow \\ n\text{-aESSS} \end{array} & & \begin{array}{c} \searrow \\ \bigoplus_{q \geq 0} m\text{ASS} \end{array} \\
 \text{Ext}_A^{s+q,f+q,u}(H_{**}E) & & \bigoplus_{q \geq 0} \pi_{s,u} s_q^n E \\
 \begin{array}{c} \searrow \\ m\text{ASS} \end{array} & & \begin{array}{c} \swarrow \\ n\text{-ESSS} \end{array} \\
 & \pi_{s+q,u} E &
 \end{array} \quad (19)$$

Remark 3.5. The obvious analog holds if we work with the type n very effective slice filtration instead of the type n effective slice filtration. Similarly, we may work with the ordinary (very) effective slice filtration instead of the type n (very) effective slice filtration.

⁵More precisely, the restriction to each fixed weight u is a half-plane spectral sequence with entering differentials.

3.3 EXAMPLES

We now present some examples of motivic spectra which are algebraically sliceable.

3.3.1 ZERO SLICES

We begin with a simple example.

EXAMPLE 3.6. If $E \in \text{SH}(k)$ is effective and the canonical map $f_0^n E \rightarrow s_0^n E$ is an equivalence, then the type n effective slice tower of E is just an equivalence at the bottom level and zero elsewhere. The only nontrivial long exact sequence in homology then splits into a short exact sequence of the form

$$0 \rightarrow H_{**} f_0^n E \xrightarrow{\cong} H_{**} s_0^n E \rightarrow 0 \rightarrow 0,$$

so E is algebraically n -sliceable. This applies, for example, if $E = H\mathbb{F}_p$ or $E = H\mathbb{Z}_p$, where n can be any positive integer, and k can be any base field.

3.3.2 CONNECTIVE ALGEBRAIC K-THEORY

Most spectra of interest are not concentrated in a single type n slice. However, the following example shows that some more complicated spectra are algebraically sliceable.

THEOREM 3.7. *Let $p = 2$ and let k be any base field of characteristic not two. The effective cover of algebraic K-theory, kgl , is algebraically and very algebraically sliceable over k .*

Proof. The effective and very effective slice filtrations of kgl coincide by [14, Prop. 4.3] since it is a localized quotient of MGL , so it suffices to prove that kgl is algebraically sliceable. Recall that for any $E \in \text{SH}(k)$, we have by [3, Lem. 8] that $\Sigma^{2,1} f_n E \simeq f_{n+1}(\Sigma^{2,1} E)$. Bott periodicity for algebraic K-theory [40, Sec. 6.2] implies that $\Sigma^{2,1} KGL \simeq KGL$, so $\Sigma^{2,1} f_n KGL \simeq f_{n+1} KGL$ for all $n \in \mathbb{Z}$. Therefore

$$f_n kgl \simeq f_n f_0 KGL \simeq f_n KGL \simeq \Sigma^{2n,n} f_0 KGL \simeq \Sigma^{2n,n} kgl \tag{20}$$

for all $n \geq 0$ since $kgl = f_0 KGL$. We also recall that $s_q kgl \simeq \Sigma^{2q,q} H\mathbb{Z}_p$ if $q \geq 0$ and $s_q kgl = 0$ if $q < 0$.

The cofiber sequences of motivic spectra

$$f_q kgl \rightarrow s_q kgl \rightarrow \Sigma^{1,0} f_{q+1} kgl$$

thus can be rewritten as

$$\Sigma^{2q,q} kgl \rightarrow \Sigma^{2q,q} H\mathbb{Z}_p \rightarrow \Sigma^{2q+3,q+1} kgl.$$

Applying homology gives a short exact sequence

$$0 \rightarrow \Sigma^{2q,q}(A//E(1))^\vee \xrightarrow{i} \Sigma^{2q,q}(A//E(0))^\vee \xrightarrow{j} \Sigma^{2q+3,q+1}(A//E(1))^\vee \rightarrow 0$$

where i is the evident inclusion of A^\vee -comodules and j is the evident projection of A^\vee -comodules. This follows from the identification of $f_1kgl \rightarrow f_0kgl$ with $v_1 : \Sigma^{2,1}kgl \rightarrow kgl$, along with the fact that $H_{**}(v_1) = 0$. It follows that kgl is algebraically sliceable. \square

3.3.3 CONNECTIVE MOTIVIC MORAVA K-THEORY

We now include an example to demonstrate the necessity of the type n effective slice filtration. Throughout this section, let p be a prime and k a perfect field of exponential characteristic prime to p .

PROPOSITION 3.8. *Let $k(n)$ denote the n -th connective motivic Morava K-theory [28, Def. 4.3].⁶ If $n > 1$, then $k(n)$ is not algebraically or very algebraically sliceable.*

Proof. The effective and very effective slice filtrations of $k(n)$ coincide by [14, Prop. 4.3] since it is a localized quotient of MGL , so it suffices to prove that $k(n)$ is not algebraically sliceable. By [28, Cor. 4.7], we have

$$s_qk(n) \simeq \begin{cases} \Sigma^{2q,q}H\mathbb{F}_p & \text{if } q \equiv 0 \pmod{p^n - 1}, \\ * & \text{if else.} \end{cases} \quad (21)$$

In particular, we have $s_1k(n) = 0$ and $f_1k(n) \simeq f_2k(n)$ whenever $n > 1$. Applying homology of cofiber sequence

$$f_1k(n) \rightarrow s_1k(n) \rightarrow \Sigma^{1,0}f_2k(n)$$

therefore yields

$$A^\vee \rightarrow 0 \rightarrow \Sigma^{1,0}A^\vee$$

which cannot be a short exact sequence since A^\vee is bounded below in each fixed motivic weight. Thus $k(n)$ is not algebraically sliceable. \square

Although $k(n)$ is not algebraically sliceable if $n > 1$, the arguments showing kgl is algebraically sliceable extend to show that $k(1)$ is algebraically sliceable: the key point is that $s_qk(1) \neq *$ for all $q \geq 0$, and thus we avoid the issues in the previous proof. More generally, the type n effective slice filtration allows us to avoid these trivial slices:

THEOREM 3.9. *Let n be a positive integer. The n -th connective motivic Morava K-theory, $k(n)$, is algebraically and very algebraically n -sliceable.*

⁶The definition of $k(n)$ given by Levine–Tripathi in [28] as a localized quotient of algebraic cobordism works for all base fields with exponential characteristic prime to p . This definition generalizes the earlier definition of the auxiliary spectrum $k'(n)$ given by Borghesi over fields with a complex embedding [8, Sec. 4.2] and certain perfect fields [9, Sec. 4].

Proof. It suffices to prove $k(n)$ is algebraically n -sliceable. The slices of $k(n)$ are given by Equation (21), so we have

$$s_q^n k(n) \simeq \Sigma^{q \cdot (2p^n - 2), q \cdot (p^n - 1)} H\mathbb{F}_p$$

for all $q \geq 0$. We also observe that by periodicity of $k(n)$, we have

$$f_{q+1}^n k(n) \simeq \Sigma^{2p^n - 2, (p^n - 1)} f_q^n k(n) \simeq \dots \simeq \Sigma^{(q+1)(2p^n - 2), (q+1)(p^n - 1)} k(n).$$

By [17, Lem. 6.10], the cohomology of $k(n)$ is $A//E(Q_n)$, where $E(Q_n)$ is the sub-Hopf algebroid of motivic Steenrod algebra generated by the motivic Milnor primitive Q_n . We are now in a situation mirroring that of Theorem 3.7. Just like in that proof, the long exact sequences in homology associated to the cofiber sequences

$$f_{q+1}^n k(n) \rightarrow f_q^n k(n) \rightarrow s_q^n k(n)$$

split into short exact sequences of A^\vee -comodules of the form

$$\begin{aligned} 0 \rightarrow \Sigma^{q(2p^n - 2), q(p^n - 1)} (A//E(Q_n))^\vee &\xrightarrow{i} \Sigma^{q(2p^n - 2), q(p^n - 1)} A^\vee \\ &\xrightarrow{j} \Sigma^{(q+1)(2p^n - 2) + 1, (q+1)(p^n - 1)} (A//E(Q_n))^\vee \rightarrow 0 \end{aligned}$$

since $H_{**}(v_n) = 0$. Therefore $k(n)$ is algebraically n -sliceable. □

3.3.4 TRUNCATED MOTIVIC BROWN–PETERSON SPECTRA

Like $k(n)$, the truncated motivic Brown–Peterson spectrum $BPGL\langle m \rangle$ is not algebraically 1-sliceable over certain base fields when $m > 1$, but it is algebraically m -sliceable over certain base fields. The proof is more subtle in this case, though.

LEMMA 3.10. *Consider a tower of fibrations*

$$\begin{array}{ccc} & \vdots & \\ & \downarrow & \\ & f_2 E & \longrightarrow s_2 E \\ & \downarrow & \\ & f_1 E & \longrightarrow s_1 E \\ & \downarrow & \\ E & \xleftarrow{\cong} f_0 E & \longrightarrow s_0 E \end{array}$$

where $f_q E \rightarrow f_{q-1} E \rightarrow s_{q-1} E$ is a fibration for each $q \geq 1$. If $H_*(f_q E \rightarrow f_{q-1} E)$ is zero for each $q \geq 1$, then the associated spectral sequence

$$E_1^{**} = H_*(s_* E) \implies H_* E$$

collapses at the E_2 -page.

Proof. Take any class $x \in H_*s_qE$ with $d_1(x) = 0$. We show that it is a permanent cycle, i.e. $d_i(x) = 0$ for any $i \geq 1$. Consider the sequence of maps

$$H_*f_qE \hookrightarrow H_*s_qE \xrightarrow{\delta_q} H_{*-1}f_{q+1}E \hookrightarrow H_{*-1}s_{q+1}E,$$

where the first and last arrows are inclusion by assumption. Therefore $d_1(x) = 0$ implies that $\delta_q(x) = 0$. Hence we have that the class x lifts to H_*f_qE . When $q = 0$, the class x lifts to H_*f_0E and is therefore a permanent cycle. When $q \geq 1$, since the connecting homomorphism $H_{*+1}s_{q-1}E \rightarrow H_*f_qE$ is surjective, x is the target of a d_1 differential. In particular, it is a permanent cycle. \square

Remark 3.11. From the proof of Lemma 3.10, we can see that when $H_*(f_qE \rightarrow f_{q-1}E)$ is zero for each $q \geq 1$, the elements in H_*E are all detected by the classes in filtration zero. For higher filtrations s_qE , the classes either support a differential (i.e. have nonzero image under the connecting homomorphism), or are hit by a d_1 differential (i.e. lift to f_qE).

With some extra assumptions, the converse of Lemma 3.10 is true.

LEMMA 3.12. *We use the same notation as in Lemma 3.10. The map $H_*(f_qE \rightarrow f_{q-1}E)$ is zero for each $q \geq 1$ if the tower satisfies the following conditions:*

1. *the map $H_*(f_1E \rightarrow f_0E)$ is zero;*
2. *the associated spectral sequence collapses at E_2 -page;*
3. $\varprojlim_n f_nE = *$.

Proof. We prove the statement by induction. Assume the statement is true for $q \leq n + 1$. We prove that $H_*(f_{n+2}E \rightarrow f_{n+1}E)$ is zero, or equivalently, $H_*(j_{n+1} : f_{n+1}E \rightarrow s_{n+1}E)$ is injective.

We show this by contradiction. Suppose there is a nonzero class $0 \neq x$ in $H_*f_{n+1}E$ such that $j_{n+1}(x) = 0$. The connecting homomorphism $\delta_n : H_{*+1}s_nE \rightarrow H_*f_{n+1}E$ is surjective by inductive hypotheses. We can choose a preimage $y \in \delta_n^{-1}(x)$. It follows that $d_1(y) = j_{n+1}\delta_n(y) = 0$. By the collapsing condition, the class y is a permanent cycle. Therefore, the class x lifts to $\varprojlim_n H_*(f_nE)$. Since the spectral sequence collapses at a finite page, this inverse limit $\varprojlim_n H_*(f_nE) \cong H_*(\varprojlim_n f_nE) \cong 0$. This contradicts the assumption that x is nonzero. \square

We now discuss the sliceability of the m -th motivic Brown–Peterson spectrum $BPGL\langle m \rangle$. For the sake of simplicity, we restrict to the case when $p = 2$. The odd primary cases work similarly.

THEOREM 3.13. *Over any base field k of characteristic zero, we have that*

$$\begin{aligned} H_{**}BPGL\langle m \rangle &\cong A//E(m)_{**} \\ &\cong \mathbb{M}_2^k[\overline{\tau}_{m+1}, \dots, \overline{\xi}_1, \dots] / (\overline{\tau}_i^2 - \tau\overline{\xi}_{i+1} - \rho\overline{\tau}_{i+1}, i \geq m + 1). \end{aligned}$$

In particular,

$$H_{**}HZ_p \cong A//E(0)_{**} \cong \mathbb{M}_2^k[\overline{\tau}_1, \dots, \overline{\xi}_1, \dots] / (\overline{\tau}_i^2 - \tau\overline{\xi}_{i+1} - \rho\overline{\tau}_{i+1}, i \geq 1).$$

Proof. The first isomorphism is [33, Thm. 3.9]. The second isomorphism can be obtained by direct computation. For example, see [15, 18, 25, 34] for the formulas of the k -motivic A_{**} , $E(m)$, and conjugation. \square

Take the mod 2 homology of the type 1 effective slice tower of $BPGL\langle m \rangle$. By [28, Cor. 4.7], the q -th slice is a wedge sum of suspensions of $H\mathbb{Z}_p$:

$$s_q^1(BPGL\langle m \rangle) \simeq \Sigma^{2q,q}H\mathbb{Z}_2 \otimes \pi_{2q}BP\langle m \rangle.$$

We obtain a homological effective slice spectral sequence:

$$E_1^{***} = H_{**}(s_*^1BPGL\langle m \rangle) \cong H_{**}H\mathbb{Z}_2[v_1, \dots, v_m] \implies H_{**}BPGL\langle m \rangle. \tag{22}$$

For a fixed bidegree (s, t) , we have that $H_{s,t}(s_q^1BPGL\langle n \rangle) \simeq 0$ when $s - t < q$. There are only finitely many slices that can contribute to a degree. On the other hand, recall that we have $sc(BPGL\langle n \rangle) \simeq BPGL\langle m \rangle$ by [17, Thm. 8.12]. Therefore the spectral sequence strongly converges to $H_{**}BPGL\langle m \rangle$.

We will refer to this spectral sequence as the H_* -ESSS below. We have the following result about this spectral sequence:

THEOREM 3.14. *Let k be either \mathbb{Q} , \mathbb{R} , or \mathbb{C} . The differentials in the spectral sequence (22) are determined by $d_{2i-1}(\overline{\tau}_i) = v_i$, $1 \leq i \leq m$, and \mathbb{M}_2^k -linearity.*

Proof. We first treat the case $k = \mathbb{R}$. We prove the result by induction. When $m = 1$, the spectral sequence (22) takes the form

$$H_{**}H\mathbb{Z}_2[v_1] \Rightarrow H_{**}kgl,$$

where by Theorem 3.13, the abutment is

$$\mathbb{M}_2^k[\overline{\tau}_2, \dots, \overline{\xi}_1, \dots] / (\overline{\tau}_i^2 - \tau\overline{\xi}_{i+1} - \rho\overline{\tau}_{i+1}, i \geq 2),$$

and the E_1 -page is

$$\mathbb{M}_2^k[v_1][\overline{\tau}_1, \dots, \overline{\xi}_1, \dots] / (\overline{\tau}_i^2 - \tau\overline{\xi}_{i+1} - \rho\overline{\tau}_{i+1}, i \geq 1).$$

The degree (s, q, u) of the elements and the differentials are as follows:

$$|\overline{\tau}_i| = (2^{i+1} - 1, 0, 2^i - 1), \quad |\overline{\xi}_i| = (2^{i+1} - 2, 0, 2^i - 1),$$

$$|v_i| = (2^{i+1} - 2, 2^i - 1, 2^i - 1), \quad i \geq 1;$$

$$|\rho| = (-1, 0, -1), \quad |\tau| = (0, 0, -1); \quad |d_r| = (-1, r, 0), \quad r \geq 1$$

By comparing degrees, the only possible target of $\bar{\tau}_1$ is v_1 . That determines all the differentials.

Now suppose that the statement holds for $m - 1$. Consider the quotient map

$$BPGL\langle m \rangle \rightarrow BPGL\langle m - 1 \rangle$$

and the induced map between spectral sequences. We have $d_{2^i-1}(\bar{\tau}_i) = v_i$ for $1 \leq i \leq m - 1$ by naturality, so the $E_{2^{m-1}}$ -page is

$$\mathbb{M}_2^k[v_m][\bar{\tau}_m, \dots, \bar{\xi}_1, \dots]/(\bar{\tau}_i^2 - \tau\bar{\xi}_{i+1} - \rho\bar{\tau}_{i+1}; \quad i \geq m).$$

By comparing degrees, the generator $\bar{\tau}_m$ supports a d_{2^m-1} differential and hits v_m . This completes the proof for $k = \mathbb{R}$.

We now consider the other base fields. The collections of bidegrees where $\mathbb{M}_2^{\mathbb{C}}$ or $\mathbb{M}_2^{\mathbb{Q}}$ are nonzero are contained in the collection of bidegrees where $\mathbb{M}_2^{\mathbb{R}}$ is nonzero, c.f. [34, Sec. 5]. Since we only used degree arguments to deduce the differentials above, we obtain the desired result by the same proof. \square

THEOREM 3.15. *Let $m \geq 2$ and let $k \subseteq \mathbb{C}$ be a field which admits a complex embedding. The m -th motivic Brown–Peterson spectrum $BPGL\langle m \rangle \in \text{SH}(k)$ is not algebraically 1-sliceable.*

Proof. The $k = \mathbb{C}$ case follows from Theorem 3.14 and the contrapositive of Lemma 3.10.

More generally, let $k \hookrightarrow \mathbb{C}$ be a field which admits a complex embedding. Base change induces a map between the H_* -ESSS in $\text{SH}(k)$ and the H_* -ESSS in $\text{SH}(\mathbb{C})$. The target does not collapse at E_2 , so it follows from inspection of the map that the source cannot collapse at E_2 . Indeed, base change sends the generators $\bar{\tau}_i$ and v_i to the generators with the same name. Therefore we may apply Lemma 3.10 to conclude that $BPGL\langle m \rangle \in \text{SH}(k)$ is not algebraically sliceable. \square

If we work with the type m effective slice tower, then in the H_* - m -ESSS⁷ for $BPGL\langle m \rangle$, the q -th filtration quotient in the E_1 term, $H_{**}s_q^m BPGL\langle m \rangle$, is an extension of the type 1 filtration quotients $H_{**}s_i^1 BPGL\langle m \rangle$ with $q(2^m - 1) \leq i \leq (q + 1)(2^m - 1) - 1$. In other words, the homology of the type m slices can be computed using the truncated H_* -ESSS. As a result, in the H_* - m -ESSS of $BPGL\langle m \rangle$, the longest differential has length 1.

Although $BPGL\langle m \rangle$, $m \geq 2$, is not algebraically 1-sliceable, it is algebraically m -sliceable. We have the following result.

THEOREM 3.16. *Let k be a field of characteristic zero. The m -th motivic Brown–Peterson spectrum $BPGL\langle m \rangle \in \text{SH}(k)$ is algebraically m -sliceable.*

⁷The spectral sequence obtained by applying homology to the type m effective slice tower.

Proof. We first show the case $k = \mathbb{Q}$. We show the three assumptions in Lemma 3.12 are satisfied.

1. By Remark 3.11, we have that the map in the type m tower

$$BPGL\langle m \rangle \rightarrow s_0^m(BPGL\langle m \rangle)$$

induces an inclusion on homology. Therefore $H_{**}(f_1 BPGL\langle m \rangle \rightarrow f_0 BPGL\langle m \rangle)$ is zero.

2. By the discussion above, the longest differential is of length 1. As a result, the spectral sequence collapses at the E_2 page.
3. Since $sc(BPGL\langle m \rangle) \simeq BPGL\langle m \rangle$, the inverse limit of the effective slice tower of $BPGL\langle m \rangle$ is contractible. Equivalently, the inverse limit of the type m effective slice tower is contractible.

The result for $k = \mathbb{Q}$ follows by applying Lemma 3.12.

If k is any field of characteristic zero, then there exists a field homomorphism $i : \mathbb{Q} \rightarrow k$. We have

$$i^*(f_{q+1}^m BPGL\langle m \rangle \rightarrow f_q^m BPGL\langle m \rangle) \simeq f_{q+1}^m BPGL\langle m \rangle \rightarrow f_q^m BPGL\langle m \rangle$$

by naturality of the slice filtration, where the left-hand side is i^* applied to part of the type m slice tower over \mathbb{Q} and the right-hand side is part of the type m slice tower over k . The left-hand side is zero, so the right-hand side is zero and thus $BPGL\langle m \rangle$ is algebraically m -sliceable over k . \square

3.3.5 VERY EFFECTIVE COVER OF HERMITIAN K-THEORY

We conclude by mentioning an example where the effective and very effective slice filtrations differ.

Remark 3.17. Let $p = 2$ and let k be a perfect field of characteristic not two. A quadruple speed very effective slice filtration can be defined by setting $\tilde{f}_q = \tilde{f}_{4q}$ and $\tilde{s}_q = \text{cofib}(\tilde{f}_{q+1} \rightarrow \tilde{f}_q)$. By [3], the very effective cover of hermitian K-theory kq [1] satisfies $\tilde{f}_q kq \simeq \Sigma^{8q, 4q} kq$. Observe that $\tilde{f}_{q+1} kq \rightarrow \tilde{f}_q kq$ may be identified with the Bott map $\beta : \Sigma^{8q+8, 4q+4} kq \rightarrow \Sigma^{8q, 4q} kq$ which induces the zero map in homology. It follows (c.f. the proof for kgl) that kq is “algebraically sliceable” with respect to the quadruple speed very effective slice filtration. Unfortunately, we cannot compute $H_{**}(\tilde{s}_q kq)$ for any $q \geq 0$, so have been unsuccessful in attempts to apply Theorem 3.4 to understanding the VSSS or mASS for kq .

It is also worth noting that the very effective slice filtration of kq does *not* coincide with the effective slice filtration. It seems unlikely that kq is algebraically sliceable with respect to a quadruple speed effective slice filtration.

3.4 NON-EXAMPLE

We mention a non-example to indicate the limitations of our techniques.

PROPOSITION 3.18. *The algebraic cobordism spectrum MGL is not algebraically sliceable.*

Proof. The slices of MGL are [17, Theorem 8.5][37, Thm. 4.7]

$$s_q MGL \simeq \Sigma^{2q,q} H\pi_{2q} MU.$$

In particular, we have that $s_0 MGL \simeq H\mathbb{Z}_p$, and that the map

$$MGL \rightarrow s_0 MGL \simeq H\mathbb{Z}_p$$

coincides with the Thom class.

Furthermore, by [17, Theorem 6.5], we have an isomorphism of left A_{**} -comodules

$$H_{**} MGL \simeq P_{**} \otimes \mathbb{F}_p[x_i \mid i \neq p^r - 1],$$

where

$$P_{**} = \mathbb{M}_p^k[\xi_1, \xi_2, \dots] \subseteq A_{**}$$

is the even subalgebra of the motivic dual Steenrod algebra. The map

$$H_{**} MGL \rightarrow H_{**} H\mathbb{Z}_p \cong A/E(0)_{**}$$

is the inclusion of P_{**} and 0 on the other factor. Thus the fiber sequence

$$f_1 MGL \rightarrow MGL \rightarrow s_0 MGL$$

induces a long exact sequence in mod p motivic homology which is not short exact. This shows that MGL is not algebraically sliceable. \square

3.5 THE SPHERE SPECTRUM

We conclude our discussion of examples and non-examples by discussing the sphere spectrum $S^{0,0} \in \text{SH}(k)$.

First, we observe that as in the case of the very effective cover of hermitian K-theory kq (see Remark 3.17), the effective and very effective slice filtrations of the sphere spectrum are different. Unlike kq , however, we do not know if the sphere spectrum is algebraically sliceable with respect to any variant of either slice filtration.

In any case, it is interesting to suppose that $S^{0,0}$ is algebraically sliceable and to speculate on the behavior of its algebraic slice spectral sequence. Such a spectral sequence would have the form

$$E_1^{q,s,t,u} = \bigoplus_{i \geq 0} \text{Ext}_{A^\vee}^{s,t,u}(H_{**} s_q S^{0,0}) \Rightarrow \text{Ext}_{A^\vee}^{s+q,t+q,u}(\mathbb{M}_p^k). \tag{23}$$

By [36, Thm. 2.12], the effective slices of the sphere spectrum are intimately connected to the E_2 -page of the classical Adams–Novikov spectral sequence. That is, there is an equivalence of motivic spectra

$$s_q((S^{0,0})_p^\wedge) \simeq \bigvee_{i \geq 0} \Sigma^{2q-i,q} H(\text{Ext}_{BP_*BP}^{i,2q}(BP_*, BP_*)), \tag{24}$$

where BP is the classical p -primary Brown–Peterson spectrum. Therefore the algebraic slice spectral sequence for the sphere spectrum, if it existed, would begin with the Ext-groups of the entries in the E_2 -term of the classical Adams–Novikov spectral sequence and end with the E_2 -term of the k -motivic Adams spectral sequence.

4 COMPARISON OF DIFFERENTIALS

We now apply Theorem 3.4 to compare differentials in the effective slice spectral sequence, the motivic Adams spectral sequence, and the ρ -Bockstein spectral sequence.

4.1 THE ρ -BOCKSTEIN SPECTRAL SEQUENCE

When $k = \mathbb{C}$, the groups $\text{Ext}_{A^\vee}^{***}(\mathbb{M}_p^{\mathbb{C}}, H_{**}(X))$ are roughly as complicated as their classical counterparts $\text{Ext}_{A^\vee}^{***}(\mathbb{F}_p, H_*(\text{Re}(X)))$. When k is an arbitrary base field, the groups $\text{Ext}_{A^\vee}^{***}(\mathbb{M}_p^k, H_{**}(X))$ contain more complicated arithmetic data coming from \mathbb{M}_p^k :

THEOREM 4.1 (Voevodsky). [21, Thm. 2.7] *Suppose that p and $\text{char}(k)$ are coprime, and that k contains a primitive p -th root of unity. Then there is an isomorphism*

$$\mathbb{M}_p^k \cong K_*^M(k)/p[\tau]$$

where $K_n^M(k)$ has degree $(-n, -n)$ and $|\tau|$ has degree $(0, -1)$.⁸

EXAMPLE 4.2. [21, Exs. 2.1, 2.2, 2.6][39, Prop. 2.4.2] We record \mathbb{M}_p^k for several cases of k considered later:

1. $\mathbb{M}_p^{\mathbb{C}} \cong \mathbb{F}_p[\tau]$ with $|\tau| = (0, -1)$.
- 2.

$$\mathbb{M}_p^{\mathbb{R}} \cong \begin{cases} \mathbb{F}_2[\tau, \rho] & \text{if } p = 2, \\ \mathbb{F}_p[\theta] & \text{if } p \neq 2, \end{cases}$$

where $|\rho| = (-1, -1)$, $|\tau| = (0, -1)$, and $|\theta| = (0, -2)$.

⁸Recall that we have defined \mathbb{M}_p^k as the mod p motivic *homology* of a point, so the bidegree of elements in $K_n^M(k)$, as well as τ , are negative instead of positive.

3. $\mathbb{M}_2^{\mathbb{F}_q} \cong \mathbb{F}_2[\tau, u]/u^2$ with $|u| = (-1, -1)$ and $|\tau| = (0, -1)$. We note that the A^\vee -coaction on u depends on the equivalence class of q modulo 4, and if $q \equiv 3 \pmod 4$, then $u = \rho$.

Recall that $\text{Ext}_{A^\vee}^{***}(\mathbb{M}_p^k, H_{**}(X))$ may be calculated using the cobar complex $C_\bullet(\mathbb{M}_p^k, A^\vee, H_{**}(X))$. When $k = \mathbb{R}$ and $p = 2$, filtering the cobar complex by powers of ρ gives rise to the ρ -Bockstein spectral sequence [15]

$$E_1^{****} = \text{Ext}_{A^\vee}^{***}(\mathbb{M}_2^{\mathbb{C}}, H_{**}(X))[\rho] \Rightarrow \text{Ext}_{A^\vee}^{***}(\mathbb{M}_2^{\mathbb{R}}, H_{**}(X)) \tag{25}$$

with differentials $d_r : E_r^{q,s,t,u} \rightarrow E_r^{q+r,s+1,t,u}$ calculated using the coaction of A^\vee on $\mathbb{M}_2^{\mathbb{R}}$. The ρ -Bockstein spectral sequence can be built over more general base fields (e.g. $k = \mathbb{F}_q$ [43]) using analogous constructions.

Hill applied the ρ -Bockstein spectral sequence to compute $\text{Ext}_{E(n)}^{***}(\mathbb{M}_2^{\mathbb{R}}, \mathbb{M}_2^{\mathbb{R}})$ for all $n \geq 0$ [15, Thm. 3.1] and $\text{Ext}_{A(1)}^{***}(\mathbb{M}_2^{\mathbb{R}}, \mathbb{M}_2^{\mathbb{R}})$ [15, Fig. 6]. In both cases, the differentials were computed using an explicit analysis of the cobar complex and technical Massey product arguments. The groups $\text{Ext}_{A(1)}^{***}(\mathbb{M}_2^{\mathbb{R}}, \mathbb{M}_2^{\mathbb{R}})$ were recomputed by Guillou–Hill–Isaksen–Ravenel [12, Sec. 6] using a result comparing the ρ -inverted \mathbb{R} -motivic and classical Ext groups [12, Prop. 4.1] to force all of the necessary differentials.

As mentioned in the proof of Theorem 3.9, $H^{**}(k(n)) \cong A//E(Q_n)$. Its mASS E_2 -page is the the trigraded group $\text{Ext}_{E(Q_n)}^{***}(\mathbb{M}_2, \mathbb{M}_2)$. Restricting to \mathbb{C} and \mathbb{R} , we have the following result on the ρ -Bockstein spectral sequence associated to $k(n)$; compare with [14, Thm. 6.3] for the case $n = 1$.

PROPOSITION 4.3. *For all $n \geq 0$, the E_1 -page of the ρ -Bockstein spectral sequence converging to $\text{Ext}_{E(Q_n)}^{***}(\mathbb{M}_2^{\mathbb{R}}, \mathbb{M}_2^{\mathbb{R}})$ is given by*

$$E_1 = \text{Ext}_{E(Q_n)}^{***}(\mathbb{M}_2^{\mathbb{C}}, \mathbb{M}_2^{\mathbb{C}})[\rho] \cong \mathbb{F}_2[\rho, \tau, v_n].$$

The nontrivial differentials are generated under ρ - and v_n -linearity by the differentials

$$d_{2^{n+1}-1}(\tau^{2^n}) = \rho^{2^{n+1}-1}v_n.$$

Proof. The result about the E_1 page follows from direct computation. The result about the differentials can be obtained by a similar computation as the ρ -Bockstein spectral sequence computation for $E(n)$ ([15, Thm. 3.2]). Here we analyze it by comparison. The ρ -Bockstein spectral sequence for $E(n)$ has a differential hitting $\rho^{2^{n+1}-1}v_n$ and its v_n - and ρ -power multiples. Therefore, in our case, the elements with the same names must be killed. By degree reasons, the only possibility is $d_{2^{n+1}-1}(\tau^{2^n}) = \rho^{2^{n+1}-1}v_n$ and differentials generated by it under ρ and v_n -linearity. \square

Remark 4.4. Although the ρ -Bockstein spectral sequence for $\text{Ext}_A^{***}(\mathbb{M}_2^{\mathbb{R}}, \mathbb{M}_2^{\mathbb{R}})$ is multiplicative [15, Prop. 2.3], the ρ -Bockstein spectral sequence for $\text{Ext}_{E(Q_n)}^{***}(\mathbb{M}_2^{\mathbb{R}}, \mathbb{M}_2^{\mathbb{R}})$ is not. Thus no issues arise from the fact that τ does not

support a differential while τ^{2^n} does. This is analogous to the topological situation, where the ESSS for the sphere spectrum is multiplicative [36, Prop. 2.24], but the ESSS for $k(n)$ cannot be multiplicative: the differentials in the $n = 1$ case of [14, Thm. 6.3] are incompatible with a Leibniz rule.

4.2 REVIEW OF ADAMS DIFFERENTIALS

In this section we record some facts about differentials in the motivic Adams spectral sequence. For the remainder of this section, we focus on the case $p = 2$ since $\rho = 0$ when p is odd.

LEMMA 4.5. *If k is an algebraically closed field or $k = \mathbb{R}$, then the motivic Adams spectral sequences converging to $\pi_{**}(k(n))$ and $\pi_{**}(BPGL\langle n \rangle)$, $n \geq 0$, collapse at E_2 .*

Proof. Recall $H^{**}k(n) \cong A//E(Q_n)$ and $H^{**}BPGL\langle n \rangle \cong A//E(n)$. If k is algebraically closed, then

$${}^{\text{mASS}}E_2(k(n)) = \text{Ext}_{E(Q_n)}^{***}(\mathbb{M}_2^k, \mathbb{M}_2^k) \cong \text{Ext}_{E(Q_n)}^{***}(\mathbb{M}_2^{\mathbb{C}}, \mathbb{M}_2^{\mathbb{C}}) \cong \mathbb{F}_2[\tau, v_n],$$

$$\begin{aligned} {}^{\text{mASS}}E_2(BPGL\langle n \rangle) &= \text{Ext}_{E(n)}^{***}(\mathbb{M}_2^k, \mathbb{M}_2^k) \\ &\cong \text{Ext}_{E(n)}^{***}(\mathbb{M}_2^{\mathbb{C}}, \mathbb{M}_2^{\mathbb{C}}) \cong \mathbb{F}_2[\tau, v_1, \dots, v_n]. \end{aligned}$$

Both spectral sequences collapse for tridegree reasons. When $k = \mathbb{R}$, this follows from [15, Thm. 5.3] for $BPGL\langle n \rangle$. Similarly, the spectral sequence collapses at E_2 for $k(n)$ for tridegree reasons. □

On the other hand, there are possible differentials when $k = \mathbb{F}_q$.

LEMMA 4.6. *When $q \equiv 1 \pmod{4}$, the E_2 -page of the \mathbb{F}_q -motivic Adams spectral sequence for kgl is given by*

$$E_2 = \text{Ext}_{E(1)}^{***}(\mathbb{M}_2^{\mathbb{F}_q}, \mathbb{M}_2^{\mathbb{F}_q}) \cong \mathbb{F}_2[\tau, u, v_0, v_1]/u^2.$$

When $q \equiv 3 \pmod{4}$, the E_2 -page is given by

$$E_2 = \text{Ext}_{E(1)}^{***}(\mathbb{M}_2^{\mathbb{F}_q}, \mathbb{M}_2^{\mathbb{F}_q}) \cong \mathbb{F}_2[\tau^2, \rho, [\rho\tau], v_0, v_1]/(\rho^2, \rho[\rho\tau], [\rho\tau]^2, \rho v_0).$$

Proof. When $q \equiv 1 \pmod{4}$, the class $u \in \mathbb{M}_2^{\mathbb{F}_q}$ satisfies $\eta_R(u) = \eta_L(u)$. Therefore it is a permanent cycle in the cobar complex and we see that $\text{Ext}^{\mathbb{F}_q} \cong \text{Ext}^{\mathbb{C}}[u]/u^2$.

When $q \equiv 3 \pmod{4}$, the result follows from the ρ -Bockstein spectral sequence⁹. In particular, the ρ -Bockstein d_1 -differentials are generated under the Leibniz rule by $d_1(\tau) = \rho v_0$, and the spectral sequence collapses at E_2 . □

⁹In this case, just a long exact sequence.

Kylling showed that the \mathbb{F}_q -motivic Adams spectral sequences for $H\mathbb{Z}_2$, kgl , and kq do not collapse [24, Sec. 4.2].

THEOREM 4.7. *Let ν_2 denote 2-adic valuation. When $q \equiv 1 \pmod{4}$, the nontrivial differentials in the motivic Adams spectral sequence converging to $\pi_{**}^{\mathbb{F}_q}(BPGL\langle m \rangle)$ are generated under the Leibniz rule by*

$$d_{\nu_2(q-1)+s}(\tau^{2^s}) = u\tau^{2^s-1}h_0^{\nu_2(q-1)+s}, \quad s \geq 0.$$

When $q \equiv 3 \pmod{4}$, they are generated by

$$d_{\nu_2(q^2-1)+s-1}(\tau^{2^s}) = \rho\tau^{2^s-1}h_0^{\nu_2(q^2-1)+s}, \quad s \geq 1.$$

Proof. For all $n \geq 1$, there is a map

$$BPGL\langle m \rangle \rightarrow BPGL\langle 0 \rangle \simeq H\mathbb{Z}_2.$$

The claimed differentials then follow by naturality of the motivic Adams spectral sequence from the differentials for $H\mathbb{Z}_2$ which were calculated in [24, Lems. 4.2.1, 4.2.2]. \square

4.3 RELATING ρ -BOCKSTEIN AND SLICE DIFFERENTIALS OVER $\text{Spec}(\mathbb{R})$

Throughout this section, we fix $k = \mathbb{R}$. The square (19) allows us to describe a precise relationship between differentials in the ρ -BSS and n -ESSS for kgl and $k(n)$. We do so by first relating the ρ -BSS differentials with n -aESSS differentials, and then we will relate n -aESSS differentials to n -ESSS differentials.

PROPOSITION 4.8. *There are 1-to-1 correspondences between the following differentials:*

1. ρ -BSS d_3 -differentials and aESSS d_1 -differentials for kgl .
2. ρ -BSS d_{2n+1-1} -differentials and n -aESSS d_1 -differentials for $k(n)$.

Moreover, the aESSS and n -aESSS differentials are forced by knowledge of the corresponding ρ -BSS differentials.

Proof. Throughout this proof, we will write $\text{Ext}_A^{***}(E)$ in place of $\text{Ext}_A^{***}(H_{**}E)$ to avoid clutter.

Let E denote kgl or $k(n)$, $n \geq 1$, let $i^* : \text{SH}(\mathbb{R}) \rightarrow \text{SH}(\mathbb{C})$ denote base change along the inclusion $\mathbb{R} \rightarrow \mathbb{C}$, and let \bar{s}_q denote s_q if $E = kgl$ or s_q^n if $E = k(n)$. We will prove the proposition by considering the square of spectral sequences

$$\begin{array}{ccc} \bigoplus_{q \geq 0} \text{Ext}_{A^{\mathbb{C}}}^{***}(\bar{s}_q i^* E)[\rho] & \xrightarrow{\rho\text{-BSS}} & \bigoplus_{q \geq 0} \text{Ext}_{A^{\mathbb{R}}}^{***}(\bar{s}_q E) \\ \Downarrow & & \Downarrow \\ \text{Ext}_{A^{\mathbb{C}}}^{***}(i^* E)[\rho] & \xrightarrow{\rho\text{-BSS}} & \text{Ext}_{A^{\mathbb{R}}}^{***}(E). \end{array} \tag{26}$$

where the rows are ρ -Bockstein spectral sequences, the left column is the direct sum over powers of ρ of the aESSS or n -aESSS, and the right column is the aESSS or n -aESSS.¹⁰

Let $E = kgl$. We can identify each corner of (26). We have

$$\begin{aligned} \bigoplus_{q \geq 0} \text{Ext}_{A^c}^{***}(i^* s_q kgl)[\rho] &\cong \bigoplus_{q \geq 0} \text{Ext}_{A^c}^{***}(\Sigma^{2q,q} i^* H\mathbb{Z}_2)[\rho] \cong \bigoplus_{q \geq 0} \Sigma^{2q,q} \mathbb{F}_2[\tau, \rho, v_0], \\ \bigoplus_{q \geq 0} \text{Ext}_{A^{\mathbb{R}}}^{***}(s_q kgl) &\cong \bigoplus_{q \geq 0} \text{Ext}_{A^{\mathbb{R}}}^{***}(\Sigma^{2q,q} H\mathbb{Z}_2) \cong \bigoplus_{q \geq 0} \Sigma^{2q,q} \mathbb{F}_2[\tau^2, \rho, v_0]/(\rho v_0), \\ \text{Ext}_{A^c}^{***}(i^* kgl)[\rho] &\cong \mathbb{F}_2[\tau, \rho, v_0, v_1], \\ \text{Ext}_{A^{\mathbb{R}}}^{***}(kgl) &\cong \mathbb{F}_2[\tau^4, \rho, v_0, v_1]/(\rho v_0, \rho^3 v_1) \end{aligned}$$

where $|\tau| = (0, 0, -1)$, $|\rho| = (0, -1, -1)$, $|v_0| = (1, 0, 0)$, and $|v_1| = (1, 3, 1)$. The left column collapses for quad-degree reasons: the E_1 -page of the aESSS for $i^* kgl$ is concentrated in quad-degrees with $t-s \equiv 0 \pmod 2$, but d_r decreases $t-s$ by 1 for all $r \geq 1$. The differentials in the top row are generated by

$$d_1(\tau) = \rho v_0$$

and the differentials in the bottom row are generated by

$$d_1(\tau) = \rho v_0, \quad d_3(\tau^2) = \rho^3 v_1.$$

Since the upper right and lower left composites must arrive at the same answer, we must have

$$d_1(\tau^2) = \rho^3 v_1$$

in the right column, where by abuse of notation v_1 denotes the generator of $\text{Ext}_{A^{\mathbb{R}}}^{***}(s_1 kgl)$.

Now fix $n \geq 1$ and let $E = k(n)$. To declutter the notation, let $r = 2^n - 1$. For the corners of (26), we have

$$\begin{aligned} \bigoplus_{q \geq 0} \text{Ext}_{A^c}^{***}(i^* s_q^n k(n))[\rho] &\cong \bigoplus_{q \geq 0} \text{Ext}_{A^c}^{***}(\Sigma^{2qr,qr} i^* H\mathbb{F}_2) \cong \bigoplus_{q \geq 0} \Sigma^{2qr,qr} \mathbb{F}_2[\tau, \rho], \\ \bigoplus_{q \geq 0} \text{Ext}_{A^{\mathbb{R}}}^{***}(s_q^n k(n)) &\cong \bigoplus_{q \geq 0} \text{Ext}_{A^{\mathbb{R}}}^{***}(\Sigma^{2qr,qr} H\mathbb{F}_2) \cong \bigoplus_{q \geq 0} \Sigma^{2qr,qr} \mathbb{F}_2[\tau, \rho], \\ \text{Ext}_{A^c}^{***}(k(n))[\rho] &\cong \mathbb{F}_2[\tau, \rho, v_n], \\ \text{Ext}_{A^{\mathbb{R}}}^{***}(k(n)) &\cong \mathbb{F}_2[\tau, \rho, t_{n+1}, v_n]/(\tau^{2^n}, \rho^{2^{n+1}-1} v_n) \end{aligned}$$

where $|\tau| = (0, 0, -1)$, $|\rho| = (0, -1, -1)$, $|v_n| = (1, 2^{n+1} - 1, 2^n - 1)$ and $|t_n| = (0, 0, -2^{n+1})$. Here, t_n is represented by $\tau^{2^{n+1}}$ on the E_1 -page of the ρ -BSS, c.f. [14, Thm. 6.3].

¹⁰The top left corner is the E_1 -page of the ρ -BSS because $\bar{s}_q i^* E \simeq i^* \bar{s}_q E$, which follows from the observation that $\bar{s}_q E$ is Eilenberg–MacLane for all $q \in \mathbb{Z}$ and base change preserves Eilenberg–MacLane spectra.

The left column and top row both collapse for quad-degree reasons, and the differentials in the bottom row are determined by

$$d_{2^{n+1}-1}(\tau^{2^n}) = \rho^{2^{n+1}-1}v_n.$$

We conclude by commutativity of (26) that we must have

$$d_1(\tau^{2^n}) = \rho^{2^{n+1}-1}v_n$$

in the n -aESSS, where by abuse of notation v_n denotes the generator of $\text{Ext}_{A^{\mathbb{R}}}^{***}(s_1^n k(n))$. \square

LEMMA 4.9. *There are 1-to-1 correspondences between the following differentials:*

1. *aESSS d_1 -differentials and ESSS d_1 -differentials for kgl .*
2. *n -aESSS d_1 -differentials and n -ESSS d_1 -differentials for $k(n)$.*

Moreover, the ESSS and n -ESSS differentials are forced by knowledge of the corresponding aESSS and n -aESSS differentials.

Proof. We have shown kgl is algebraically sliceable and $k(n)$ is algebraically n -sliceable, so in the notation of the previous proof, we have squares of spectral sequences

$$\begin{array}{ccc}
 & \bigoplus_{q \geq 0} \text{Ext}_A^{***}(H_{**}\bar{s}_q E) & \\
 \begin{array}{c} \text{aESSS} \\ \swarrow \\ \text{Ext}_A^{***}(H_{**}E) \end{array} & & \begin{array}{c} \bigoplus_{q \geq 0} \text{mASS} \\ \searrow \\ \bigoplus_{q \geq 0} \pi_{**}\bar{s}_q E \end{array} \\
 \begin{array}{c} \text{mASS} \\ \searrow \\ \pi_{**}E \end{array} & & \begin{array}{c} \text{ESSS} \\ \swarrow \\ \pi_{**}E \end{array}
 \end{array} \tag{27}$$

for $E = kgl$ and $E = k(n)$.

The mASS's for kgl , $k(n)$, $s_q kgl \simeq \Sigma^{2q,q} H\mathbb{Z}_2$, and $s_q^n k(n) \simeq \Sigma^{2q^n, q^n} H\mathbb{F}_2$ all collapse. The only nontrivial differentials in the aESSS for kgl and the n -aESSS for $k(n)$ have length one, so for quad-degree reasons, we must have identical differentials in the ESSS for kgl and n -ESSS for $k(n)$. \square

Putting together Proposition 4.8 and Lemma 4.9, we have proven the following theorem.

THEOREM 4.10. *There are 1-to-1 correspondences between the following differentials:*

1. *ρ -BSS d_3 -differentials and ESSS d_1 -differentials for kgl .*
2. *ρ -BSS $d_{2^{n+1}-1}$ -differentials and n -ESSS d_1 -differentials for $k(n)$.*

Moreover, the ESSS and n -ESSS differentials are completely determined by knowledge of the corresponding ρ -BSS differentials.

Remark 4.11. As the n -ESSS is obtained by increasing the speed of the effective slice filtration, a d_1 -differential in the n -ESSS corresponds to a d_{2^n-1} -differential in the ESSS. Therefore part (2) of Theorem 4.10 may be restated as a 1-to-1 correspondence between ρ -BSS $d_{2^{n+1}-1}$ -differentials and d_{2^n-1} -differentials in the ESSS.

Theorem 4.10 can be used to prove a correspondence between more complicated differentials. The following theorem is closely related to Yagita's analysis of the effective slice spectral sequence for $BPGL$ in [44, Sec. 4].

THEOREM 4.12. *Let $m \geq 1$. For each $1 \leq i \leq m$, there is a 1-to-1 correspondence between the ρ -BSS $d_{2^{i+1}-1}$ -differentials and ESSS d_{2^i-1} -differentials for $BPGL\langle m \rangle$.*

Proof. We proceed by induction on m . At $p = 2$, there is an equivalence $kgl \simeq BPGL\langle 1 \rangle$, so the base case $m = 1$ is handled by Theorem 4.10.

Suppose now that the result holds for all $k < m$ and we wish to show the theorem holds for $BPGL\langle m \rangle$. We begin by using the induction hypothesis to show that the theorem holds for $i < m$ (but not necessarily $i = m$).

Observe that the $d_{2^{i+1}-1}$ -differentials in the ρ -BSS's for $BPGL\langle m-1 \rangle$ and $BPGL\langle m \rangle$ are determined by the same differential

$$d_{2^{i+1}-1}(\tau^{2^i}) = \rho^{2^{i+1}-1}v_i \quad (28)$$

for $1 \leq i \leq m-1$ by [15, Thm. 3.2]. This implies that the $d_{2^{i+1}-1}$ -differentials for $BPGL\langle m \rangle$ are precisely the $d_{2^{i+1}-1}$ -differentials for $BPGL\langle m-1 \rangle$ extended v_m -linearly.

On the other hand, naturality of the ESSS allows us to completely determine the d_{2^i-1} -differentials in the ESSS for $BPGL\langle m \rangle$. Indeed, the quotient map $BPGL\langle m \rangle \rightarrow BPGL\langle m-1 \rangle$ implies that the d_{2^i-1} -differentials between slices indexed on monomials not divisible by v_m coincide for $BPGL\langle m-1 \rangle$ and $BPGL\langle m \rangle$, and the quotient map $BPGL\langle m \rangle \rightarrow k(n)$ implies that v_m is a d_{2^i-1} -cycle for $i < m$ by Theorem 4.10. Applying the Leibniz rule for the ESSS [36, Prop. 2.24] proves that the d_{2^i-1} -differentials in the ESSS for $BPGL\langle m \rangle$ are precisely the d_{2^i-1} -differentials in the ESS for $BPGL\langle m-1 \rangle$ extended v_m -linearly. Therefore we have proven a 1-to-1 correspondence between ρ -BSS and ESSS differentials for $i < m$.

We now consider the case $i = m$. Theorem 3.16 shows that $BPGL\langle m \rangle$ algebraically m -sliceable, so we may consider the comparison square (19) relating its m -aESSS and m -ESSS. The m ASS's for $BPGL\langle m \rangle$ and its type m slices (extensions of $(2j, j)$ -suspensions of $H\mathbb{Z}_2$) all collapse, so we obtain a 1-to-1 correspondence between d_1 -differentials in the m -aESSS and m -ESSS.

We are therefore reduced to relating the $d_{2^{m+1}-1}$ -differentials in the ρ -BSS and the d_1 -differentials in the m -aESSS, but this follows from essentially the same

arguments as Proposition 4.8. Indeed, consider the $BPGL\langle m \rangle$ -analog of (26):

$$\begin{array}{ccc}
 \bigoplus_{q \geq 0} \text{Ext}_{A^c}^{***}(s_q^m i^* BPGL\langle m \rangle)[\rho] & \xrightarrow{\rho\text{-BSS}} & \bigoplus_{q \geq 0} \text{Ext}_{A^{\mathbb{R}}}^{***}(s_q^m BPGL\langle m \rangle) \\
 \Downarrow & & \Downarrow \\
 \text{Ext}_{A^c}^{***}(i^* BPGL\langle m \rangle)[\rho] & \xrightarrow{\rho\text{-BSS}} & \text{Ext}_{A^{\mathbb{R}}}^{***}(BPGL\langle m \rangle).
 \end{array} \tag{29}$$

The left column collapses for degree reasons and the differentials in the bottom row were determined in [15, Thm. 3.2]. The differentials in the top row coincide with the differentials in the bottom row, with two types of exceptions:

1. All of the $d_{2^{m+1}-1}$ -differentials, i.e. the longest nontrivial differentials, in the bottom row cannot occur in the top row since their target in the bottom is divisible by v_m (which is zero in the top). We refer to these as *long differentials*.
2. The d_r -differentials for $r < 2^{m+1} - 1$ in the bottom row of the form $d_r(x) = y$ where x lies in stem s with $k(2^{m+1} - 2) \leq s < (k + 1)(2^{m+1} - 2)$ and y lies in stem s' with $s' \geq (k + 1)(2^{m+1} - 2)$ cannot support differentials in the top row for degree reasons. We refer to these as *fringe differentials* since they involve elements from adjacent type m slices.

A discussion fringe differentials for the case $m = 2$ appears in Example 4.13. The long and fringe differentials in the bottom row which do not occur in the top row both are accounted for by d_1 -differentials in the m -aESSS. The d_1 -differentials in the m -aESSS forced by the fringe differentials in the ρ -BSS give rise to d_r -differentials in the ESSS with $r < 2^m - 1$, which are completely understood by the induction hypothesis. On the other hand, the d_1 -differentials in the m -aESSS forced by the long differentials in the ρ -BSS correspond to d_{2^m-1} -differentials in the ESSS; this is precisely the correspondence we wished to show. \square

EXAMPLE 4.13. We analyze the fringe d_r -differentials in the ρ -BSS for $BPGL\langle 2 \rangle$. In this case, we have the cofiber sequence

$$f_3 BPGL\langle 2 \rangle \rightarrow BPGL\langle 2 \rangle \rightarrow s_0^2 BPGL\langle 2 \rangle.$$

Observe that

$$s_0^2 BPGL\langle 2 \rangle \simeq s_0^2 BPGL\langle 1 \rangle \simeq BPGL\langle 1 \rangle / v_1^3.$$

Furthermore, the associated graded motivic spectrum $s_*^2 BPGL\langle 2 \rangle$ is given by

$$s_*^2 BPGL\langle 2 \rangle \simeq s_0^2 BPGL\langle 2 \rangle [v_1^3, v_2].$$

In particular, to understand the top ρ -BSS of (29) it is enough to understand the ρ -BSS for $s_0^2 BPGL\langle 2 \rangle$. This can be done using the naturality of the ρ -BSS

which produces the diagram

$$\begin{array}{ccc}
 \text{Ext}_{A^c}^{***}(BPGL\langle 2 \rangle)[\rho] & \Longrightarrow & \text{Ext}_{A^{\mathbb{R}}}(BPGL\langle 2 \rangle) \\
 \downarrow & & \downarrow \\
 \text{Ext}_{A^c}^{***}(s_0^2 BPGL\langle 2 \rangle)[\rho] & \Longrightarrow & \text{Ext}_{A^{\mathbb{R}}}^{***}(s_0^2 BPGL\langle 2 \rangle).
 \end{array} \tag{30}$$

On E_1 -pages, the left-hand map is the projection map

$$\mathbb{F}_2[\tau, \rho, v_0, v_1, v_2] \rightarrow \mathbb{F}_2[\tau, \rho, v_0, v_1, v_2]/(v_1^3, v_2).$$

This allows us to import differentials from the ρ -BSS for $BPGL\langle 2 \rangle$. In the top and bottom rows, the d_1 -differentials are determined by the differential $d_1(\tau) = \rho v_0$, and the morphism of spectral sequences is the obvious map on E_2 -pages. There are no d_2 -differentials so $E_2 = E_3$. We encounter a fringe differential on the E_3 -page. In the top, there is a d_3 -differential

$$d_3(\tau^2) = \rho^3 v_1$$

which implies a differential of the same form in the bottom. However, in the top there is a differential $d_3(\tau^2 v_1^2) = \rho^3 v_1^3$ which cannot occur in the bottom since $v_1^3 = 0$. This is precisely the phenomena described by *fringe differentials* in the previous proof.

After running d_3 -differentials, both rows collapse. Thus, in $\text{Ext}_{A^{\mathbb{R}}}(s_0^2 BPGL\langle 2 \rangle)$, there is an extra class which we denote by $[\tau^2 v_1^2]$ to remind us that it is indecomposable in $\text{Ext}_{A^{\mathbb{R}}}$. Since there are ρ -BSS differentials $d_3(\tau^2 v_1^2) = \rho^3 v_1^3$ in the ρ -BSS for $BPGL\langle 2 \rangle$, it follows that we must have differentials of the form

$$d_1(\rho^\ell [\tau^2 v_1^2](v_1^3)^j v_2^k) = \rho^{\ell+3} (v_1^3)^{j+1} v_2^k$$

in the m -aESS, and hence in the m -ESSS. These correspond to the d_1 -differentials in the ESSS which are already understood by reduction to $BPGL\langle 1 \rangle \simeq kgl$.

Remark 4.14. Naturality of the ESSS can be used to prove an analogous correspondence between ρ -BSS and ESSS differentials for $BPGL$.

Remark 4.15. Although we have only considered the case $k = \mathbb{R}$ here, it would be interesting to consider more general base fields of characteristic zero. For example, the mASS for $BPGL\langle n \rangle$ is completely understood when $k = \mathbb{Q}$ by work of Ormsby and Østvær [34, Thm. 5.8]. If one extends Theorem 3.16 to $k = \mathbb{Q}$, then it may be possible to understand the ESSS for $BPGL\langle n \rangle$ over $\text{Spec}(\mathbb{Q})$ using Theorem 3.4.

4.4 COMPARISON OVER $\text{Spec}(\mathbb{F}_q)$

We now turn to the case when $k = \mathbb{F}_q$.

Unlike the \mathbb{R} -motivic case, the comparison is trivial for $k(n)$, $n \geq 1$. Since $\mathbb{M}_2^{\mathbb{F}_p}$ is zero in bidegrees (s, u) with $s < -1$, the ρ -BSS, mASS, and n -ESSS all collapse for degree reasons.

However, the analysis is still interesting for kgl . Recall from Example 4.2(3) that the A^\vee -coaction depends on the equivalence class of q modulo 4. We discuss the case when $q \equiv 1 \pmod{4}$, and the other case when $q \equiv 3 \pmod{4}$ is similar.

4.4.1 $q \equiv 1 \pmod{4}$

By Lemma 4.6, we observe that in the motivic Adams spectral sequence of kgl and $H\mathbb{Z}_2$, the E_2 -pages have the relation that

$$\mathrm{Ext}_A^{***}(kgl) \cong \mathrm{Ext}_A^{***}(H\mathbb{Z}_2)[v_1].$$

Since $\mathrm{Ext}_A^{***}(H\mathbb{Z}_2)[v_1]$ is exactly the E_1 -page of the algebraic effective slice spectral sequence for kgl , we conclude that the algebraic effective slice spectral sequence collapses.

By Theorem 4.7, the nontrivial differentials in both motivic Adams spectral sequences for kgl and $H\mathbb{Z}_2$ are determined under Leibniz rule by

$$d_{\nu_2(q-1)+s}(\tau^{2^s}) = u\tau^{2^s-1}h_0^{\nu_2(q-1)+s}, s \geq 0.$$

Therefore, using the motivic comparison square (19), we conclude that the effective slice spectral sequence for kgl collapses.

4.4.2 $q \equiv 3 \pmod{4}$

When $q \equiv 3 \pmod{4}$, we also get that the algebraic effective slice spectral sequence and the effective slice spectral sequence for kgl both collapse.

Remark 4.16. The work of Ormsby [33] shows that the situation over local fields is very similar. The above analysis can be carried over local fields to show that the effective slice spectral sequence for kgl , as well as $BPGL\langle m \rangle$ for all $m \geq 0$, are trivial.

5 APPLICATIONS TO C_2 -EQUIVARIANT STABLE HOMOTOPY THEORY

We now make some calculations in C_2 -equivariant stable homotopy theory. Let $k \subseteq \mathbb{R}$ be a field with a real embedding.

Remark 5.1. Throughout this section, we will freely use the names of elements in $\pi_{**}^{C_2}(H\mathbb{F}_2)$ and $\pi_{**}^{C_2}(H\mathbb{Z}_2)$ used in [12]. We briefly describe the relevant elements here.

The C_2 -equivariant elements $\tau \in \pi_{0,-1}^{C_2}(H\mathbb{F}_2)$ and $\rho \in \pi_{-1,-1}^{C_2}(H\mathbb{F}_2)$ are the images under equivariant Betti realization of the \mathbb{R} -motivic elements with the

same names.¹¹ The subalgebra $\mathbb{F}_2[\tau, \rho] \subseteq \pi_{**}^{C_2}(H\mathbb{F}_2)$ is sometimes called the *positive cone*.

There is another element, $\frac{\gamma}{\tau} \in \pi_{0,2}^{C_2}(H\mathbb{F}_2)$, which is infinitely ρ - and τ -divisible but is not in the image of equivariant Betti realization. The infinitely divisible part of $\pi_{**}^{C_2}(H\mathbb{F}_2)$ is sometimes called the *negative cone*.

There are two sets of naming conventions for C_2 -equivariant homotopy elements. We refer the reader to tables in Section 1.3 for the correspondence.

We will use the following result of Heard to obtain C_2 -equivariant slice differentials from known motivic differentials. We note that Heard actually proves a more general result, but we only use the following specialization in our work.

THEOREM 5.2 (Heard). [14, Thms. 5.15-5.16] *Let $E \in \text{SH}(k)$ be a motivic spectrum which is a localized quotient of MGL. Then $\text{Re}_{C_2} : (\text{SH}(k), \Sigma_T^q \text{SH}(k)^{eff}) \rightarrow (\text{SH}^{C_2}, \Sigma^{2q}(\text{SH}^{C_2})^{HHR})$ is compatible with the slice filtration at E .*¹²

Heard applied this theorem to deduce effective slice spectral sequence differentials from the existence of HHR slice differentials. We will apply Heard’s theorem in the opposite direction: starting with the \mathbb{R} -motivic effective slice spectral sequence differentials computed above, we will deduce the existence of differentials in the HHR slice spectral sequence.

COROLLARY 5.3. *Let $E \in \text{SH}(k)$ be a motivic spectrum which is a localized quotient of MGL. Every d_{2r-1-1} -differential in the ESSS for E uniquely determines a d_{2r-1} -differentials in the HHR slice spectral sequence for $\text{Re}_{C_2}(E)$.*

Remark 5.4. The difference in the lengths of the differentials in Corollary 5.3 stems from a difference in the indexing conventions between the motivic slice spectral sequence and the equivariant slice spectral sequence. In particular, motivic d_1 -differentials correspond to equivariant d_3 -differentials. See the proof of [14, Proposition 6.1] for more details.

We will apply Corollary 5.3 to the following C_2 -spectra:

DEFINITION 5.5. We define the following C_2 -spectra using equivariant Betti realization.¹³

1. Let

$$k\mathbb{R} := \text{Re}_{C_2}(kgl)$$

denote the C_2 -equivariant slice cover of Atiyah’s K -theory with reality.

2. Let

$$k\mathbb{R}(n) := \text{Re}_{C_2}(k(n))$$

denote the n -th connective Real Morava K -theory.

¹¹There are also geometric models for τ and ρ , cf. [30, Section 4].

¹²We refer the reader to [14, 16] for details on the C_2 -equivariant slice spectral sequences discussed in this section.

¹³Each spectrum can also be defined directly in C_2 -spectra. See [2, 10] for $k\mathbb{R}$ and [16, 19] for $k\mathbb{R}(n)$ and $BP\mathbb{R}(n)$.

3. Let

$$B\mathbb{P}\mathbb{R}\langle n \rangle := \text{Re}_{C_2}(BPGL\langle n \rangle)$$

denote the n -th truncated Real Brown–Peterson spectrum.

By [14, Ex. 3.5], the motivic spectra kgl , $k(n)$, and $BPGL\langle n \rangle$ are localized quotients of MGL . Combining Theorem 5.2 and Corollary 5.3, we obtain:

COROLLARY 5.6. *Let E be kgl , $k(n)$, or $BPGL\langle n \rangle$. Every d_{2r-1-1} -differential in the ESSS for E uniquely determines a d_{2r-1} -differential in the HHR slice spectral sequence for $\text{Re}_{C_2}(E)$.*

We identified the differentials in the ESSS for kgl , $k(n)$, and $BPGL\langle n \rangle$ with ρ -Bockstein spectral sequence differentials in Theorems 4.10 and 4.12. Coupled with Corollary 5.6, this allows us to produce differentials in the HHR slice spectral sequences for $k\mathbb{R}$, $k\mathbb{R}(n)$, and $B\mathbb{P}\mathbb{R}\langle n \rangle$. We spell out these consequences in the next few corollaries.

COROLLARY 5.7. *The nontrivial differentials in the HHR slice spectral sequence for $\pi_{**}^{C_2}k\mathbb{R}$ are determined via the Leibniz rule by*

$$d_3(\tau^2) = \rho^3 \bar{v}_1 \quad \text{and} \quad d_3\left(\frac{\gamma}{\rho^3 \tau^2}\right) = \frac{\gamma}{\tau^4} \bar{v}_1.$$

Proof. The differential $d_1(\tau^2) = \rho^3 v_1$ in the ESSS for kgl arising from Theorem 4.10 gives rise to the differential $d_3(\tau^2) = \rho^3 \bar{v}_1$ in the HHR slice spectral sequence for $k\mathbb{R}$ using Corollary 5.6.

We now prove the second differential. By the Leibniz rule,

$$0 = d\left(\frac{\gamma}{\rho^3 \tau^2} \cdot \tau^2\right) = d\left(\frac{\gamma}{\rho^3 \tau^2}\right) \cdot \tau^2 + \frac{\gamma}{\rho^3 \tau^2} \cdot d(\tau^2)$$

Therefore we have that

$$d\left(\frac{\gamma}{\rho^3 \tau^2}\right) \cdot \tau^2 = \frac{\gamma}{\rho^3 \tau^2} \cdot d(\tau^2).$$

Since the element τ^2 support a d_3 -differential, we deduce that the element $\frac{\gamma}{\rho^3 \tau^2}$ lives to the E_3 -page and supports a differential with target $\frac{\gamma}{\tau^4} \bar{v}_1$. This differential generates a family of d_3 -differentials by multiplicity.

By inspection, there is no room for further differentials. This completes the proof. \square

COROLLARY 5.8. *There is a nontrivial differential in the HHR slice spectral sequence for $\pi_{**}^{C_2}k\mathbb{R}(n)$ of the form*

$$d_{2^{n+1}-1}(\tau^{2^n}) = \rho^{2^{n+1}-1} \bar{v}_n.$$

Proof. By Theorem 4.10 and Remark 4.11, there is a 1-to-1 correspondence between $d_{2^{n+1}-1}$ -differentials in the ρ -BSS and d_{2^n-1} -differentials in the ESSS for $k\mathbb{R}\langle n \rangle$. By Proposition 4.3, the nontrivial differentials in the ρ -BSS are generated under ρ - and v_n -linearity by the differentials

$$d_{2^{n+1}-1}(\tau^{2^n}) = \rho^{2^{n+1}-1}v_n.$$

By Corollary 5.6, these \mathbb{R} -motivic differentials uniquely determine the stated $d_{2^{n+1}-1}$ -differentials in the HHR slice spectral sequence for $k\mathbb{R}\langle n \rangle$. \square

Similar arguments also apply to truncated Real Brown–Peterson spectra, where one uses Theorem 4.12 and [15, Thm. 3.2] to identify the differentials in the ESSS for $BPGL\langle n \rangle$.

COROLLARY 5.9. *There are nontrivial differentials in the HHR slice spectral sequence for $\pi_{**}^{C_2}BP\mathbb{R}\langle n \rangle$ of the form*

$$d_{2^{i+1}-1}(\tau^{2^i}) = \rho^{2^{i+1}-1}\bar{v}_i$$

for $1 \leq i \leq n$.

Remark 5.10. Using naturality and the quotient maps $BP\mathbb{R} \rightarrow BP\mathbb{R}\langle n \rangle$, the corollary recovers the p -local version of the $G = C_2$ -case of the Hill–Hopkins–Ravenel Slice Differentials Theorem [16, Thm. 9.9].¹⁴

REFERENCES

- [1] A. Ananyevskiy, O. Röndigs, and P. A. Østvær. On very effective Hermitian K -theory. *Math. Z.* 294(3-4):1021–1034, 2020.
- [2] M. F. Atiyah. K -theory and reality. *Quart. J. Math. Oxford Ser. (2)* 17(1):367–386, 1966.
- [3] T. Bachmann. The generalized slices of Hermitian K -theory. *J. Topol.* 10(4):1124–1144, 2017.
- [4] M. Behrens and J. Shah. C_2 -equivariant stable homotopy from real motivic stable homotopy. *Ann. K-Theory* 5(3):411–464, 2020.
- [5] E. Belmont, B. Guillou, and D. Isaksen. C_2 -equivariant and \mathbb{R} -motivic stable stems, II. *Proc. Amer. Math. Soc.* 149(1):53–61, 2021.
- [6] E. Belmont and D. C. Isaksen. \mathbb{R} -motivic stable stems. *arXiv preprint arXiv:2001.03606*, 2020.

¹⁴They note this originally appeared in the work of Hu–Kriz [19] and in unpublished work of Araki.

- [7] J. M. Boardman. Conditionally convergent spectral sequences. In *Homotopy invariant algebraic structures (Baltimore, MD, 1998)*, volume 239 of *Contemp. Math.*, pages 49–84. Amer. Math. Soc., Providence, RI, 1999.
- [8] S. Borghesi. Algebraic Morava K -theories. *Invent. Math.* 151(2):381–413, 2003.
- [9] S. Borghesi. Algebraic Morava K -theory spectra over perfect fields. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* 8(2):369–390, 2009.
- [10] D. Dugger. An Atiyah-Hirzebruch spectral sequence for KR -theory. *K-Theory* 35(3-4):213–256, 2005.
- [11] D. Dugger and D. C. Isaksen. Motivic cell structures. *Algebr. Geom. Topol.* 5(2):615–652, 2005.
- [12] B. J. Guillou, M. A. Hill, D. C. Isaksen, and D. C. Ravenel. The cohomology of C_2 -equivariant $A(1)$ and the homotopy of ko_{C_2} . *Tunis. J. Math.* 2(3):567–632, 2019.
- [13] J. J. Gutiérrez, O. Röndigs, M. Spitzweck, and P. A. Østvær. Motivic slices and coloured operads. *J. Topol.* 5(3):727–755, 2012.
- [14] D. Heard. On equivariant and motivic slices. *Algebr. Geom. Topol.* 19(7):3641–3681, 2019.
- [15] M. A. Hill. Ext and the motivic Steenrod algebra over \mathbb{R} . *J. Pure Appl. Algebra* 215(5):715–727, 2011.
- [16] M. A. Hill, M. J. Hopkins, and D. C. Ravenel. On the nonexistence of elements of Kervaire invariant one. *Ann. of Math. (2)* 184(1):1–262, 2016.
- [17] M. Hoyois. From algebraic cobordism to motivic cohomology. *J. Reine Angew. Math.* 702:173–226, 2015.
- [18] M. Hoyois, S. Kelly, and P. A. Østvær. The motivic Steenrod algebra in positive characteristic. *J. Eur. Math. Soc. (JEMS)* 19(12):3813–3849, 2017.
- [19] P. Hu and I. Kriz. Real-oriented homotopy theory and an analogue of the Adams–Novikov spectral sequence. *Topology* 40(2):317–399, 2001.
- [20] P. Hu, I. Kriz, and K. Ormsby. Convergence of the motivic Adams spectral sequence. *J. K-Theory* 7(3):573–596, 2011.
- [21] D. C. Isaksen and P. A. Østvær. Motivic stable homotopy groups. In *Handbook of homotopy theory*, CRC Press/Chapman Hall Handb. Math. Ser., pages 757–791. CRC Press, Boca Raton, FL [2020].
- [22] D. C. Isaksen, G. Wang, and Z. Xu. More stable stems. *arXiv preprint arXiv:2001.04511*, 2020.

- [23] H. J. Kong. The C_2 -effective spectral sequence for C_2 -equivariant connective real K-theory. *arXiv preprint arXiv:2004.00806*, 2020.
- [24] J. I. Kylling. Hermitian K-theory of finite fields via the motivic Adams spectral sequence. Master's thesis, University of Oslo, Norway, 2015.
- [25] J. I. Kylling. Recursive formulas for the motivic Milnor basis. *New York J. Math.* 23(250399):49–58, 2017.
- [26] J. I. Kylling and G. M. Wilson. Strong convergence in the motivic Adams spectral sequence. *arXiv preprint arXiv:1901.03399*, 2019.
- [27] M. Levine. The homotopy coniveau tower. *J. Topol.* 1(1):217–267, 2008.
- [28] M. Levine and G. S. Tripathi. Quotients of MGL, their slices and their geometric parts. *Doc. Math., Extra Vol. Alexander S. Merkurjev's sixtieth birthday*, 407–442, 2015.
- [29] L. Mantovani. Localizations and completions in motivic homotopy theory. *arXiv preprint arXiv:1810.04134*, 2018.
- [30] C. May. A structure theorem for $RO(C_2)$ -graded Bredon cohomology. *Algebr. Geom. Topol.* 20(4):1691–1728, 2020.
- [31] F. Morel. Suite spectrale d'Adams et invariants cohomologiques des formes quadratiques. *C. R. Acad. Sci. Paris Sér. I Math.* 328(11):963–968, 1999.
- [32] F. Morel and V. Voevodsky. A^1 -homotopy theory of schemes. *Inst. Hautes Études Sci. Publ. Math.* 90:45–143, 1999.
- [33] K. M. Ormsby. Motivic invariants of p -adic fields. *J. K-Theory* 7(3):597–618, 2011.
- [34] K. M. Ormsby and P. A. Østvær. Motivic Brown-Peterson invariants of the rationals. *Geom. Topol.* 17(3):1671–1706, 2013.
- [35] O. Röndigs and P. A. Østvær. Slices of Hermitian K-theory and Milnor's conjecture on quadratic forms. *Geom. Topol.* 20(2):1157–1212, 2016.
- [36] O. Röndigs, M. Spitzweck, and P. A. Østvær. The first stable homotopy groups of motivic spheres. *Ann. of Math. (2)* 189(1):1–74, 2019.
- [37] M. Spitzweck. Relations between slices and quotients of the algebraic cobordism spectrum. *Homology Homotopy Appl.* 12(2):335–351, 2010.
- [38] M. Spitzweck and P. A. Østvær. Motivic twisted K-theory. *Algebr. Geom. Topol.* 12(1):565–599, 2012.
- [39] S.-T. Stahn. The motivic Adams-Novikov spectral sequence at odd primes over \mathbb{C} and \mathbb{R} . *arXiv preprint arXiv:1606.06085*, 2016.

- [40] V. Voevodsky. A1-homotopy theory. In *Proceedings of the International Congress of Mathematicians*, Doc. Math., Extra Vol. ICM Berlin 1998, vol. I, pages 579–604. Berlin, 1998.
- [41] V. Voevodsky. Open problems in the motivic stable homotopy theory. I. *Motives, polylogarithms and Hodge theory, Part I (Irvine, CA, 1998)*, Int. Press Lect. Ser., 3, I, pages 3–34, 1998.
- [42] V. Voevodsky. Reduced power operations in motivic cohomology. *Publ. Math. Inst. Hautes Études Sci.* 98(1):1–57, 2003.
- [43] G. M. Wilson and P. A. Østvær. Two-complete stable motivic stems over finite fields. *Algebr. Geom. Topol.* 17(2):1059–1104, 2017.
- [44] N. Yagita. Applications of Atiyah–Hirzebruch spectral sequences for motivic cobordism. *Proc. London Math. Soc. (3)* 90(3):783–816, 2005.

Dominic Leon Culver
Max Planck Institute for
Mathematics
Vivatsgasse 7
53111 Bonn
Germany
dominic.culver@gmail.com

Hana Jia Kong
Department of Mathematics
5734 S University Ave
Chicago IL, 60637
U.S.A
hana.jia.kong@gmail.com

J. D. Quigley
Department of Mathematics
Cornell University
Ithaca, NY
U.S.A.
jdq27@cornell.edu

