

## Splittings of triangle Artin groups

Kasia Jankiewicz

**Abstract.** We show that a triangle Artin group  $\text{Art}_{MNP}$ , where  $M \leq N \leq P$ , splits as an amalgamated product or an HNN extension of finite rank free groups, provided that either  $M > 2$  or  $N > 3$ . We also prove that all even 3-generator Artin groups are residually finite.

A triangle Artin group is given by the presentation

$$\text{Art}_{MNP} = \langle a, b, c \mid (a, b)_M = (b, a)_M, (b, c)_N = (c, b)_N, (c, a)_P = (a, c)_P \rangle,$$

where  $(a, b)_M$  denote the alternating word  $aba \dots$  of length  $M$ . Squier showed that the Euclidean triangle Artin group, i.e.,  $\text{Art}_{236}$ ,  $\text{Art}_{244}$  and  $\text{Art}_{333}$ , split as amalgamated product or an HNN extension of finite rank free groups along finite index subgroups [25]. We generalize that result to other triangle Artin groups.

**Theorem A.** *Suppose that  $M \leq N \leq P$ , where either  $M > 2$  or  $N > 3$ . Then the Artin group  $\text{Art}_{MNP}$  splits as an amalgamated product or an HNN extension of finite rank free groups.*

The assumptions of the above theorem are satisfied for all triples of numbers except for  $(2, 2, P)$  and  $(2, 3, P)$ . An Artin group is *spherical*, if the associated Coxeter group is finite. A 3-generator Artin group  $\text{Art}_{MNP}$  is spherical exactly when  $\frac{1}{M} + \frac{1}{N} + \frac{1}{P} > 1$ , i.e.,  $(M, N, P) = (2, 2, P)$  or  $(2, 3, 3)$ ,  $(2, 3, 4)$ ,  $(2, 3, 5)$ . None of 3-generator spherical Artin groups splits as a graph of finite rank free groups (see Proposition 2.10). The remaining cases are  $(2, 3, P)$ , where  $P \geq 6$ . The above theorem holds for triple  $(2, 3, 6)$  by [25]. It remains unknown for  $(2, 3, P)$  with  $P \geq 7$ . The cases where  $M > 2$  were considered in [18, Theorem B], and it was proven that they all split as amalgamated products of finite rank free groups.

Graphs of free groups form an important family of examples in geometric group theory. Graphs of free groups with cyclic edge groups that contain no Baumslag–Solitar subgroups are virtually special [15], and contain quasiconvex surface subgroups [27]. Graphs of free groups with arbitrary edge groups can exhibit various behaviors. For example, an amalgamated product  $A *_C B$  of finite rank free groups, where  $C$  is malnormal

in  $A$  and  $B$ , is hyperbolic [1], and virtually special [16]. On the other hand, there are examples of amalgamated products of finite rank free groups that are not residually finite [2,28], and even simple [8]. The last two arise as lattices in the automorphism group of a product of two trees.

By further analysis of the splitting, we are also able to show that some of the considered Artin groups are residually finite.

**Theorem B.** *The Artin group  $\text{Art}_{2MN}$ , where  $M, N \geq 4$  and at least one of  $M, N$  is even, is residually finite.*

An Artin group  $\text{Art}_{MNP}$  is even if all  $M, N, P$  are even. The above theorem combined with our result in [18] (and the fact that  $\text{Art}_{22P} = \mathbb{Z} \times \text{Art}_P$  is linear) gives us the following.

**Corollary C.** *All even Artin groups on three generators are residually finite.*

All linear groups are residually finite [21], so residual finiteness can be viewed as testing for linearity. Spherical Artin groups are known to be linear ([3, 19] for braid groups, and [9, 11] for other spherical Artin groups). The right-angled Artin groups are also well known to be linear, but not much more is known about linearity of Artin groups. In last years, a successful approach in proving that groups are linear is by showing that they are virtually special. Artin groups whose defining graphs are forests are the fundamental groups of graph manifolds with boundary [7, 14], and so they are virtually special [20, 23]. Many Artin groups in certain classes (including 2-dimensional, or 3-generator) are not cocompactly cubulated even virtually, unless they are sufficiently similar to RAAGs [12, 17]. In particular, the only (virtually) cocompactly cubulated 3-generator Artin groups are  $\text{Art}_{22M} = \mathbb{Z} \times \text{Art}_M$ ,  $\text{Art}_{MN\infty}$ , where  $M$  and  $N$  are both even,  $\text{Art}_{M\infty\infty} = \mathbb{Z} * \text{Art}_M$ , and  $\text{Art}_{\infty\infty\infty} = F_3$ . Some triangle-free Artin groups act properly but not cocompactly on locally finite, finite-dimensional CAT(0) cube complexes [13].

In [18], we showed that  $\text{Art}_{MNP}$  are residually finite when  $M, N, P \geq 3$ , except for the cases where  $(M, N, P) = (3, 3, 2p + 1)$  with  $p \geq 2$ . Few more families of Artin groups are known to be residually finite, e.g., even FC type Artin groups [5], and certain triangle-free Artin groups [4].

## Organization

In Section 1, we provide some background. In Section 2, we prove Theorem A as Proposition 2.5 and Corollary 2.9. We also show that the only irreducible spherical Artin groups splitting as graph of finite rank free groups are dihedral. In Section 3, we recall a criterion for residual finiteness of amalgamated products and HNN extensions of free groups from [18] and prove Theorem B.

## 1. Background

### 1.1. Graphs

Let  $X$  be a finite graph with directed edges. We denote the vertex set of  $X$  by  $V(X)$  and the edge set of  $X$  by  $E(X)$ . The vertices incident to an edge  $e$  are denoted by  $e^+$  and  $e^-$ . A map of graphs  $f: X_1 \rightarrow X_2$  sends vertices to vertices, and edges to concatenations of edges. A map  $f$  is a *combinatorial map* if single edges are mapped to single edges. A combinatorial map  $f$  is a *combinatorial immersion* if given two edges  $e_1, e_2$  such that  $e_1^- = e_2^-$ , we have  $f(e_1) = f(e_2)$  (as oriented edges) if and only if  $e_1 = e_2$ . Consider two edges  $e_1, e_2$  with  $e_1^- = e_2^-$ . A *fold* is the natural combinatorial map  $X \rightarrow \bar{X}$ , where

$$V(\bar{X}) = V(X)/e_1^+ \sim e_2^+ \quad \text{and} \quad E(\bar{X})/e_1 \sim e_2.$$

Stallings showed that every combinatorial map  $X \rightarrow X'$  factors as  $X \rightarrow \bar{X} \rightarrow X'$ , where  $X \rightarrow \bar{X}$  is a composition of finitely many folds, and  $\bar{X} \rightarrow X'$  is a combinatorial immersion [26]. We refer to  $X \rightarrow \bar{X}$  as a *folding map*.

### 1.2. Maps between free groups

Let  $H, G$  be finite rank free groups. Let  $Y$  be a bouquet of  $n = \text{rk } G$  circles. We can identify  $\pi_1 Y \simeq F_n$  with  $G$  by orienting and labeling edges of  $Y$  with the generators of  $G$ . Every homomorphism  $\varphi: H \rightarrow G$  can be represented by a combinatorial immersion of graphs. Indeed, start with a map of graphs  $X \rightarrow Y$ , where  $X$  is a bouquet of  $m = \text{rk } H$  circles. We think of each circle in  $X$  as subdivided with edges oriented and labeled by the generators of  $G$ , so that each circle is labeled by a word from a generating set of  $H$ . By Stallings, the map  $X \rightarrow Y$  factors as  $X \rightarrow \bar{X} \rightarrow Y$ , where  $X \rightarrow \bar{X}$  is a folding map, and  $\bar{X} \rightarrow Y$  is a combinatorial immersion. Indeed,  $\bar{X}$  is obtained by identifying two edges with the same orientation and label that share an endpoint.

Note that the rank of  $\varphi(H)$  is equal to  $\text{rk } \pi_1 \bar{X} = 1 - \chi(\bar{X})$ , where  $\chi$  denotes the Euler characteristic. In particular, a homomorphism  $\varphi$  is injective if and only if the folding map  $X \rightarrow \bar{X}$  is a homotopy equivalence. In that case,  $\bar{X}$  is a precover of  $Y$  which can be completed to a cover of  $Y$  corresponding to the subgroup  $H$  of  $G$  via the Galois correspondence. In particular, every subgroup of  $G$  is uniquely represented by a combinatorial immersion  $(X, x) \rightarrow (Y, y)$ , where  $y$  is the unique vertex of  $Y$ , and  $X$  is a folded graph with basepoint  $x$ . We refer to [26] for more details.

### 1.3. Intersections of subgroups of a free group

Let  $Y$  be a graph, and let  $\rho_i: (X_i, x_i) \rightarrow (Y, y)$  be a combinatorial immersion for  $i = 1, 2$ . The *fiber product of  $X_1$  and  $X_2$  over  $Y$* , denoted by  $X_1 \otimes_Y X_2$  is a graph with the vertex set

$$V(X_1 \otimes_Y X_2) = \{(v_1, v_2) \in V(X_1) \times V(X_2): \rho_1(v_1) = \rho_2(v_2)\},$$

and the edge set

$$E(X_1 \otimes_Y X_2) = \{(e_1, e_2) \in E(X_1) \times E(X_2) : \rho_1(e_1) = \rho_2(e_2)\}.$$

The graph  $X_1 \otimes_Y X_2$  often has several connected components. There is a natural combinatorial immersion  $X_1 \otimes_Y X_2 \rightarrow Y$ , and it induces an embedding  $\pi_1(X_1 \otimes_Y X_2, (x_1, x_2)) \rightarrow \pi_1(Y, y)$ . We have the following.

**Theorem 1.1** ([26, Theorem 5.5]). *Let  $H_1, H_2$  be two subgroups of  $G = \pi_1 Y$ , and for  $i = 1, 2$ , let  $(X_i, x_i) \rightarrow (Y, y)$  be a combinatorial immersion of graphs inducing the inclusion  $H_i \hookrightarrow G$ . The intersection  $H_1 \cap H_2$  is represented by a combinatorial immersion  $(X_1 \otimes_Y X_2, (x_1, x_2)) \rightarrow (Y, y)$ .*

In particular, when  $Y$  is a bouquet of circles with  $\pi_1 Y = G$ , and  $(X, x) \rightarrow (Y, y)$  is a combinatorial immersion inducing  $H = \pi_1 X \hookrightarrow G$ , then for every pair of (not necessarily distinct) vertices  $x_1, x_2 \in X$ , the group  $\pi_1(X \otimes_Y X, (x_1, x_2))$  is an intersection  $H^{g_1} \cap H^{g_2}$  for some  $g_1, g_2 \in G$ . In fact, every non-trivial intersection  $H \cap H^g$  is equal to  $\pi_1(X \otimes_Y X, (x_1, x_2))$ , where  $x_1 = x$ , and  $x_2$  is some (possibly the same) vertex in  $X$ . The connected component of  $X \otimes_Y X$  containing  $(x, x)$  is a copy of  $X$ , which we refer to as a *diagonal component*. The group  $\pi_1(X \otimes_Y X, (x, x))$  is the intersection  $H \cap H^g = H$ , i.e., where  $g \in H$ . A connected component of  $X \otimes_Y X$  that has no edges is called *trivial*.

#### 1.4. Graph of groups and spaces

We recall the definitions of a graph of groups and a graph of spaces, following [24].

A *graph of spaces* consists of

- a graph  $\Gamma$ , called the *underlying graph*,
- a collection of CW-complexes  $X_v$  for each  $v \in V(\Gamma)$ , called *vertex spaces*,
- a collection of CW-complexes  $X_e$  for each  $e \in E(\Gamma)$ , called *edge spaces*,
- a collection of continuous  $\pi_1$ -injective maps  $f_{(e, \pm)}: X_e \rightarrow X_{e^\pm}$  for each  $e \in E(\Gamma)$ .

The *total space*  $X(\Gamma)$  is defined as

$$X(\Gamma) = \bigsqcup_{v \in V(\Gamma)} X_v \sqcup \bigsqcup_{e \in E(\Gamma)} X_e \times [-1, 1] / \sim,$$

where  $(x, \pm 1) \sim f_{(e, \pm)}(x)$  for  $x \in X_e$ .

Similarly, a *graph of groups* consists of

- the *underlying graph*  $\Gamma$ ,
- a collection of *vertex groups*  $G_v$  for each  $v \in V(\Gamma)$ ,
- a collection of *edge groups*  $G_e$  for each  $e \in E(\Gamma)$ ,
- a collection of injective homomorphisms  $\varphi_{(e, \pm)}: G_e \rightarrow G_{e^\pm}$  for each  $e \in E(\Gamma)$ .

The *fundamental group of a graph of groups* is defined as the fundamental group of the graph of spaces  $X(\Gamma)$ , where  $X_v = K(G_v, 1)$  for each  $v \in V(\Gamma)$ ,  $X_e = K(G_e, 1)$  for each edge  $e \in E(\Gamma)$ , and  $f_{(e,\pm)}$  induces homomorphism  $\varphi_{(e,\pm)}$  on the fundamental groups. Note that the fundamental group  $\pi_1 \Gamma$  is a subgroup of  $G$ .

### 1.5. HNN extensions and doubles

We will denote the *HNN extension of  $A$  relative to  $\beta: B \rightarrow A$* , where  $B \subseteq A$ , by  $A *_B \beta$ , that is,

$$A *_B \beta = \langle A, t \mid t^{-1}xt = \beta(x) \text{ for all } x \in B \rangle.$$

The generator  $t$  is called the *stable letter*. Note that  $A *_B \beta$  can be viewed as a graph of group  $G(\Gamma)$ , where  $\Gamma$  is a single vertex  $v$  with a single loop  $e$ ,  $G_v = A$ ,  $G_e = B$ ,  $\varphi_{(e,-)}$  is the inclusion of  $B$  in  $A$ , and  $\varphi_{(e,+)} = \beta$ .

A *double of  $A$  along  $C$  twisted by an automorphism  $\beta: C \rightarrow C$* , denoted by  $D(A, C, \beta)$ , is an amalgamated product  $A *_C A$ , where  $C$  is mapped to the first factor via the standard inclusion, and to the second via the standard inclusion precomposed with  $\beta$ . As usual,  $D(A, C, \beta)$  depends only on the outer automorphism class of  $\beta$ , and not a particular representative. A double  $D(A, C, \beta)$  can be viewed as a graph of groups  $G(\Gamma)$ , where  $\Gamma$  is a single edge  $e$  with distinct endpoints,  $G_{e^\pm} = A$ ,  $G_e = C$ , and  $\varphi_{(e,-)}$  is the inclusion of  $C$  in  $A$ , and  $\varphi_{(e,+)}$  is the inclusion precomposed with  $\beta$ . Note that an amalgamated product  $A *_C B$  with  $[B : C] = 2$  has an index 2 subgroup  $D(A, C, \beta)$ . The homomorphism  $\beta: C \rightarrow C$  is conjugation by some (any) representative  $g \in B$  of the non-trivial coset of  $B/C$ .

In both situations where there is unique edge in the underlying graph  $\Gamma$ , we will skip the label  $e$  and denote  $\varphi_{(e,\pm)}$  simply by  $\varphi_\pm$ . Similarly, in a graph of spaces with a unique edge  $e$ , we will write  $f_\pm$  instead of  $f_{(e,\pm)}$ .

### 1.6. Triangle groups

A *triangle (Coxeter) group* is given by the presentation

$$W_{MNP} = \langle a, b, c \mid a^2, b^2, c^2, (ab)^M, (bc)^N, (ca)^P \rangle.$$

The group  $W_{MNP}$  acts as a reflection group on

- the sphere if  $\frac{1}{M} + \frac{1}{N} + \frac{1}{P} > 1$ ,
- the Euclidean plane if  $\frac{1}{M} + \frac{1}{N} + \frac{1}{P} = 1$ ,
- the hyperbolic plane if  $\frac{1}{M} + \frac{1}{N} + \frac{1}{P} < 1$ .

Hyperbolic triangle groups are commensurable with the fundamental groups of negatively curved surfaces, and therefore they are locally quasiconvex, virtually special, and Gromov-hyperbolic. A *von Dyck triangle group* is an index 2 subgroup of  $W_{MNP}$  with the presentation

$$\langle x, y \mid x^M, y^N, (x^{-1}y)^P \rangle$$

obtained by setting  $x = ba$  and  $y = bc$ .

## 2. Splittings

The goal of this section is to prove Theorem A. In [18], we proved the following.

**Theorem 2.1** ([18, Theorem B]). *The Artin group  $\text{Art}_{MNP}$  with  $M, N, P \geq 3$  splits as an amalgamated product or an HNN extension of finite rank free groups.*

The remaining cases in Theorem A are  $\text{Art}_{2MN}$ , where  $M, N \geq 4$ . The case where  $M, N$  are both even is Proposition 2.5, and the other case is Corollary 2.9. We start with considering a non-standard presentation of  $\text{Art}_{2MN}$ . In the next two subsections, we construct the splittings. Then we show that the only spherical Artin groups that split as graphs of free groups are dihedral Artin groups. Spherical Artin groups on three generators  $\text{Art}_{MNP}$  correspond to one of the following triples:  $(2, 2, P)$  for any  $P \geq 2$ ,  $(2, 3, 3)$ ,  $(2, 3, 4)$  or  $(2, 3, 5)$ . For completeness, in the last subsection we include the proof that the 3-generator Artin groups with at least one  $\infty$  label admit splittings as HNN extensions or amalgamated products of finite rank free groups.

The Artin group  $\text{Art}_{236}$  splits as  $F_3 *_{F_7} F_4$  by Squier [25]. The only remaining 3-generator Artin groups are  $\text{Art}_{23M}$ , where  $M \geq 7$ , and the following remains unanswered.

**Question 2.2.** *Does the Artin group  $\text{Art}_{23M}$ , where  $M \geq 7$ , split as a graph of finite rank free groups?*

We conjecture that the answer is positive. More generally, we ask the following.

**Question 2.3.** *Do all 2-dimensional Artin groups split as a graph of finite rank free groups?*

### 2.1. Presentations of $\text{Art}_{2MN}$

Here is the standard presentation of Artin group  $\text{Art}_{2MN}$ :


$$\langle a, b, c \mid (a, b)_M = (b, a)_M, (b, c)_N = (c, b)_N, ac = ca \rangle.$$

Let  $x = ab$  and  $y = cb$ , and consider a new presentation of  $\text{Art}_{2MN}$  with generators  $b, x, y$ . The relation  $(a, b)_M = (b, a)_M$  is replaced by  $bx^mb^{-1} = x^m$  when  $M = 2m$ , and by  $bx^mb = x^{m+1}$  when  $M = 2m + 1$ . We denote this relation by  $r_M(b, x)$ . Note that  $yx^{-1} = ca^{-1}$ , so relation  $ac = ca$  can be replaced by  $yx^{-1} = bx^{-1}yb^{-1}$ . See Figure 1(a).

This gives us the following presentation:

$$\text{Art}_{2MN} = \langle b, x, y \mid r_M(b, x), r_N(b, y), bx^{-1}yb^{-1} = yx^{-1} \rangle. \quad (*)$$

Let  $X_{2MN}$  be the presentation complex associated to presentation (\*). Let  $X_A$  be the bouquet of two loops labeled by  $x$  and  $y$ . The complex  $X_{2MN}$  can be viewed as a union of the graph  $X_A$  and for each relation in (\*), a cylinder (for relations  $bx^{-1}yb^{-1} = yx^{-1}$  and each  $r_M(b, x)$  with  $M$  even) or a Möbius strip (for each relation  $r_M(b, x)$  with  $M$  odd) with boundary cycles are glued to  $X_A$ . We can metrize them so that the height of each cylinder/Möbius strip equals 2.



**Figure 1.** (a) The relation  $R_M(b, x)$ , where  $M$  is even (left) and odd (middle), and the relation  $b x^{-1} y b^{-1} = y x^{-1}$  (right) with the projection  $p$  of the cells onto the interval  $[0, 1]$ . The horizontal graphs  $X_A, X_B, X_C$  are the preimages  $p^{-1}(1), p^{-1}(0), p^{-1}(\frac{1}{2})$ , respectively. (b) Graphs  $X_A, X_B$  and two versions of  $X_C$  depending on whether  $M, N$  are both odd (left green), or one of  $M, N$  is even (right green).

We now define a map  $p: X_{2MN} \rightarrow [0, 1]$  by describing the restriction of  $p$  to each cylinder/Möbius strip of  $X_{2MN}$ . Each point of the cylinder/Möbius strip is mapped to its distance from the center circle of that cylinder/Möbius strip. In particular, the center circle of each cylinder or Möbius strip is mapped to 0, and the boundary circles of the cylinder or Möbius strip are mapped to 1. See Figure 1 (a).

We can identify  $X_A$  with the preimage  $p^{-1}(1)$ . We define a graph  $X_B$  as the union of all the center circles, i.e., the preimage  $p^{-1}(0)$ . We also define a subgraph  $X_C$  of  $X_{2MN}$  as the preimage  $p^{-1}(\frac{1}{2})$ . The graphs  $X_A, X_B$  and  $X_C$  are illustrated in Figure 1(b). The graph  $X_C$  has two vertices, which are its intersections with the edge  $b$ . We denote them by  $b_-, b_+$ , so that  $b_-$ , the midpoint of the edge  $b$ ,  $b_+$  are ordered consistently with the orientation of the edge  $b$ . When  $M, N$  are both even, then the graph  $X_C$  is not connected. Indeed, each of its connected components is a copy of  $X_B$ . We denote the connected component containing the vertex  $b_-$  by  $X_B^-$ , and the component containing the vertex  $b_+$  by  $X_B^+$ . Otherwise, if at least one  $M, N$  is odd, then  $X_C$  is a connected double cover of  $X_B$ . In the next two sections, we describe the graph of spaces decomposition of  $X_{2MN}$  associated to the map  $p$ , and the induced graph of groups decomposition of  $\text{Art}_{2MN}$ . We consider separately the case where  $M, N$  are both even, and the case where at least one of them is odd.

## 2.2. Both even

In the case where both  $M, N$  are even and  $\geq 4$ , presentation  $(*)$  of  $\text{Art}_{2MN}$  is the standard presentation of an HNN-extension.

**Proposition 2.4.** *Let  $M = 2m, N = 2n$  and both  $\geq 4$ . Then  $\text{Art}_{2MN}$  splits as an HNN-extension  $A *_B \beta$ , where  $A = \langle x, y \rangle \simeq F_2$  and  $B = \langle x^m, y^n, x^{-1}y \rangle \simeq F_3$ , and  $\beta: B \rightarrow A$  is an injective homomorphism given by  $\beta(x^m) = x^m, \beta(y^n) = y^n$  and  $\beta(x^{-1}y) = yx^{-1}$ .*

Proposition 2.4 follows from the following.

**Proposition 2.5.** *Let  $M = 2m, N = 2n$  and both  $\geq 4$ . Then  $X_{2MN}$  is a graph of spaces  $X(\Gamma)$ , where  $\Gamma$  is a single vertex with a single loop. The vertex space is the graph  $X_A$ , the edge space is the graph  $X_B$ , and the two maps  $X_B \rightarrow X_A$  are given in the Figure 2.*

*Proof.* Indeed, the map  $p$  factors as

$$X_{2MN} \xrightarrow{\tilde{p}} S^1 = [-1, 1]/(-1 \sim 1) \rightarrow [0, 1],$$

where the second map is the absolute value, and where  $\tilde{p}$  sends the loop  $b$  isometrically onto  $S^1$  and is extended linearly. By construction, the preimage  $\tilde{p}^{-1}(t)$  is homeomorphic to  $X_B$  when  $t \in (-1, 1)$ , and to  $X_A$  when  $t = 1$ . In particular,  $X_{2MN}$  can be expressed as a graph of spaces, induced by  $\tilde{p}$ , where the cellular structure of  $S^1$  consists of a single vertex  $v = 1$  and a single edge  $e$ . Indeed,

$$X_{2MN} = X_A \cup X_B \times [-1, 1]/(x, -1) \sim f_-(x), (x, 1) \sim f_+(x),$$


where  $f_-, f_+ : X_B \rightarrow X_A$  are the two maps obtained by “pushing” the graph  $X_B$  in Figure 1 (a) “upwards” and “downwards”, respectively. See Figure 2 for  $f_-, f_+$  expressed as a composition of Stallings fold and a combinatorial immersion. ■

**Remark 2.6.** The subgroups  $B$  and  $\beta(B)$  are conjugate. See Figure 2. Indeed, the graphs  $\bar{X}_B^-$  and  $\bar{X}_B^+$  are identical (but have different basepoints).

**Example 2.7** (Group  $\text{Art}_{244}$ ). In the case where  $M = N = 4$ , Proposition 2.5 provides the splitting of  $\text{Art}_{244} = A *_{B, \beta}$ , where

$$A = \langle x, y \rangle \quad \text{and} \quad B = \langle x^2, y^2, x^{-1}y \rangle,$$

and  $\beta : B \rightarrow B$  is given by  $\beta(x^2) = x^2, \beta(y^2) = y^2, \beta(x^{-1}y) = yx^{-1}$ . In particular,  $B$  has index 2 in  $A$ . This splitting was first proven by Squier [25].



**Figure 2.** The map  $f_- : X_B \rightarrow X_B^- \rightarrow \bar{X}_B^- \rightarrow X_A$  (top), and the map  $f_+ : X_B \rightarrow X_B^+ \rightarrow \bar{X}_B^+ \rightarrow X_A$  (bottom).

### 2.3. At least one odd

We now assume that at least one  $M, N$  is odd. We have the following description of the complex  $X_{2MN}$ .

**Proposition 2.8.** *The complex  $X_{2MN}$  is a graph of spaces whose underlying graph is an interval, the vertex spaces are graphs  $X_A$  and  $X_B$ , and the edge space is  $X_C$ . The attaching map  $X_C \rightarrow X_B$  is a double cover, and the attaching map  $X_C \rightarrow X_A$  factors as  $X_C \rightarrow \bar{X}_C \rightarrow X_A$ , illustrated in Figure 3, where the first map is a homotopy equivalence and the second map is a combinatorial immersion.*


*Proof.* Indeed,  $X_{2MN}$  can be obtained as a union of  $X_A, X_B$  and  $X_C \times [0, 1]$ , where  $X_C \times \{1\}$  is glued to  $X_A$ , and  $X_C \times \{0\}$  is glued to  $X_B$ . Note that the preimage  $p^{-1}([0, \frac{1}{2}])$  is a union of cylinders and Möbius strips, and its boundary is the graph  $X_C$ . The projection onto the center circle of the boundary of each cylinder or Möbius strip is a (connected or not) double cover of the center circle. It follows that  $X_C \rightarrow X_B$  is a double cover. The map  $X_C \rightarrow X_A$  is induced by “pushing”  $X_C$  “downwards” and “upwards” onto  $X_A$ , and it can be described by the labeling of the right graphs in Figure 3. The factorization  $X_C \rightarrow \bar{X}_C$  is obtained by performing Stallings folds. Note that the middle graphs in Figure 3 are fully folded, provided that  $m - 1, n - 1 > 0$ , which is equivalent to the condition that  $M, N \geq 4$ . It follows that the map  $\bar{X}_C \rightarrow X_A$  is a combinatorial immersion. Since the ranks of  $\pi_1 X_C$  and  $\pi_1 \bar{X}_C$  are both equal to 5, the folding map is a homotopy equivalence. ■

**Corollary 2.9.** *Suppose at most one of  $M, N$  is even and  $M, N \geq 4$ . Then  $\text{Art}_{2MN}$  splits as a free product with amalgamation  $A *_C B$ , where  $A = F_2$  and  $B = F_3$ , and  $C = F_5$ .*

*Proof.* This directly follows from Proposition 2.8. We get that

$$\text{Art}_{2MN} = \pi_1 X_{2MN} = A *_C B,$$

where  $A = \pi_1 X_A, B = \pi_1 X_B$  and  $C = \pi_1 X_C$ . From Figure 1, we see that  $\text{rk } A = 2, \text{rk } B = 3$  and  $\text{rk } C = 5$ . ■



**Figure 3.** The maps  $X_C \rightarrow \bar{X}_C \rightarrow X_A$  when (a):  $M = 2m + 1, N = 2n + 1$ ; and (b):  $M = 2m + 1, N = 2n$ .

## 2.4. Splittings of spherical Artin groups

All spherical Artin groups have non-trivial center and their cohomological dimension is equal to the number of standard generators [6, 10]. We now give a characterization of graph of finite rank free groups with non-trivial center. This allows us to deduce that the only spherical Artin groups that split as graphs of finite rank free groups are the dihedral Artin groups (i.e., on two generators).

**Proposition 2.10.** *The only irreducible spherical Artin groups that split as non-trivial graphs of free groups are the dihedral Artin groups and  $\mathbb{Z}$ .*

*Proof.* Clearly,  $\mathbb{Z}$  is an HNN-extension of a trivial group. Let  $\text{Art}_M$  be a dihedral Artin group with the presentation  $\langle a, b \mid (a, b)_M = (b, a)_M \rangle$ . Let  $M = 2m$ , and let  $x = ab$ . Then  $\text{Art}_M \simeq \langle a, x \mid ax^m a^{-1} = x^m \rangle$ . In particular,  $\text{Art}_M = \langle x \rangle *_{\langle x^m \rangle} \mathbb{Z} *_{\mathbb{Z}}$ . Now if  $M = 2m + 1$ , let  $x = ab$  and  $y = (a, b)_M$ . Then  $\text{Art}_M \simeq \langle x, y \mid x^M = y^2 \rangle$ . In particular,  $\text{Art}_M \simeq \langle x \rangle *_{\langle x^M \rangle} \langle y \rangle = \mathbb{Z} *_{\mathbb{Z}} \mathbb{Z}$ . Conversely, non-trivial graphs of free groups have cohomological dimension at most 2 since the corresponding graphs of spaces are aspherical. The only irreducible spherical Artin groups of cohomological dimension at most 2 are dihedral Artin groups and  $\mathbb{Z}$ . ■

## 2.5. Splittings of 3-generator Artin groups with $\infty$ labels

To complete the picture, we prove that the remaining 3-generator Artin groups, i.e., those with at least one  $\infty$  label, also split as graphs of finite rank free groups. The Artin group  $\text{Art}_{\infty\infty\infty}$  is the free group on three generators. The group  $\text{Art}_{M\infty\infty} = \text{Art}_M * \mathbb{Z}$  can be described as

$$\langle x, c \rangle *_{x^M=y^2} \langle y \rangle = F_2 *_{\mathbb{Z}} \mathbb{Z},$$

where  $x = ab$  and  $y = (a, b)_M$ . Finally, for the Artin group  $\text{Art}_{MN\infty}$ , we can use presentation (\*) skipping the relation  $bx^{-1}yb^{-1} = yx^{-1}$ , i.e.,

$$\text{Art}_{MN\infty} = \langle x, y, b \mid r_M(b, x), r_N(b, y) \rangle.$$

We get that  $\text{Art}_{MN\infty}$  splits as

- $A *_{B, \beta}$ , where  $A = \langle x, y \rangle \simeq F_2$ ,  $B = \langle x^m, y^n \rangle \simeq F_2$ ,  $\beta = \text{id}_C$ , when  $M = 2m$  and  $N = 2n$ ,
- $A *_{\beta} B$ , where  $A = \langle x, y \rangle \simeq F_2$ ,  $B \simeq F_2$  and  $C \simeq F_3$ , when at least one of  $M, N$  is odd. The splitting is obtained in the same way as in the case of  $\text{Art}_{2MN}$ .

## 3. Residual finiteness

In the section, we prove Theorem B. We do so separately in the case where  $M, N$  are both even (Theorem 3.6), and where exactly one of  $M, N$  is odd (Theorem 3.7). We start with recalling a criterion for residual finiteness of amalgamated products and HNN extensions

of finite rank free groups, proven in [18], which relies on Wise’s result on residual finiteness of *algebraically clean* graphs of free groups [29]. In Section 3.2, we compute the intersections of conjugates of the amalgamating subgroup in the factor groups. Finally, we give proofs of the main theorems.

We start with a motivating example.

**Example 3.1** (Group  $\text{Art}_{244}$ ). By Example 2.7, the group  $\text{Art}_{244}$  fits in the short exact sequence of groups

$$1 \rightarrow C \rightarrow \text{Art}_{244} \rightarrow \mathbb{Z}/2\mathbb{Z} * \mathbb{Z} \rightarrow 1.$$

In particular,  $\text{Art}_{244}$  is a (finite rank free group)-by-(virtually free group), and so it is virtually (finite rank free group)-by-(free group). The residual finiteness of  $\text{Art}_{244}$  follows from the fact that every split extension of a finitely generated residually finite group by residually finite group is residually finite [22].

### 3.1. Criteria for residual finiteness

Recall that a subgroup  $C$  is *malnormal* in a group  $A$ , if  $C \cap g^{-1}Cg = \{1\}$  for every  $g \in A - C$ . Similarly,  $C$  is *almost malnormal* in  $A$ , if  $|C \cap g^{-1}Cg| < \infty$  for every  $g \in A - C$ . More generally, a collection of subgroup  $\{C_i\}_{i \in I}$  is an *almost malnormal* collection, if  $|C_i \cap gC_jg^{-1}| < \infty$  whenever  $g \notin C_i$  or  $i \neq j$ .

Assume that the inclusion of free groups  $C \rightarrow A$  is induced by a map  $f: X_C \rightarrow X_A$  of graphs, and the automorphism  $\beta: C \rightarrow C$  is induced by some graph automorphism  $X_C \rightarrow X_C$ . The following theorem was proven in [18].

**Theorem 3.2** ([18, Theorem 2.9]). *Let  $\hat{f}: \hat{X}_C \rightarrow \hat{X}_A$  be a map of 2-complexes that restricted to the 1-skeletons is equal to  $f$ , and let  $\pi: A \rightarrow \hat{A}$  be the natural quotient induced by the inclusion  $X_A \hookrightarrow \hat{X}_A$  of the 1-skeleton. Suppose that the following conditions hold:*

- (1)  $\hat{A}$  is a locally quasiconvex, virtually special hyperbolic group.
- (2)  $\pi(C) = \pi_1 \hat{X}_C$  and the lift of  $\hat{f}$  to the universal covers is an embedding.
- (3)  $\pi(C)$  is almost malnormal in  $\hat{A}$ .
- (4)  $\beta$  projects to an isomorphism  $\pi(C) \rightarrow \pi(C)$ .

Then  $D(A, C, \beta)$  is residually finite.

The theorem above is a combination of Theorem 2.9 and Lemma 2.6 from [18]. Condition (2) in the statement of [18, Theorem 2.9] is that  $\pi(C)$  is *malnormal* in  $\hat{A}$ . However, the proof is identical when we replace it with *almost malnormal*. Indeed, the Bestvina–Feighn combination theorem [1], as well as the Hsu–Wise combination theorem [16] only require almost malnormality.

We now state a version for HNN extension. Similarly as above, combining Theorem 2.12 and Lemma 2.6 from [18], we obtain the following.

**Theorem 3.3** ([18, Theorem 2.12]). *Let  $\hat{f}_-, \hat{f}_+: \hat{X}_B \rightarrow \hat{X}_A$  be two maps of 2-complexes that restricted to the 1-skeletons are equal to  $f_-, f_+$ , respectively, and let  $\pi: A \rightarrow \hat{A}$*

be the natural quotient induced by the inclusion  $X_A \hookrightarrow \widehat{X}_A$ . Suppose that the following conditions hold:

- (1)  $\widehat{A}$  is a locally quasiconvex, virtually special hyperbolic group.
- (2)  $\pi(B) = \widehat{f}_{-*}(\pi_1 \widehat{X}_B)$ , and  $\pi(\beta(B)) = \widehat{f}_{+*}(\pi_1 \widehat{X}_B)$  and the lifts of  $\widehat{f}_-$ ,  $\widehat{f}_+$  to the universal covers are both embeddings.
- (3) The collection  $\{\pi(B), \pi(\beta(B))\}$  is almost malnormal in  $\widehat{A}$ .
- (4)  $\beta: B \rightarrow \beta(B)$  projects to an isomorphism  $\pi(B) \rightarrow \pi(\beta(B))$ .

Then  $A *_B \beta$  is residually finite.

### 3.2. Intersection of the conjugates of the amalgamating subgroup

Let  $G = \text{Art}_{2MN}$ , where  $M, N \geq 4$  and at least one of them is odd. Let  $A, B, C$  be free groups of ranks 2, 3, 5, respectively, provided by Corollary 2.9. In this section, we describe intersections  $C \cap g^{-1}Cg$ , where  $g \in A$ .

**Proposition 3.4.** *Let  $M = 2m + 1$  be odd,  $N = 2n$  even, and let  $A, B, C$  be as in Corollary 2.9. Let  $g \in A - C$ . Then the intersection  $C \cap g^{-1}Cg$  is one of the sets:  $\langle x^{2m+1}, y^n, x^{-1}y \rangle$ ,  $\langle x^{2m+1}, y^n, yx^{-1} \rangle$ ,  $\langle x^{2m+1}, y^n \rangle$ ,  $\langle x^{2m+1} \rangle$ ,  $\langle y^n \rangle$ .*

*Proof.* Let  $X_A, \bar{X}_C$  be as in Figure 3 (b). By Theorem 1.1, the conjugates  $C \cap g^{-1}Cg$  can be represented by the connected components of the fiber product  $\bar{X}_C \otimes_{X_A} \bar{X}_C$ . Let  $x_1, x_2, x_3, x_4$  be the four vertices of valence 4 in  $\bar{X}_C$  such that  $x_1, x_2$  belong to the same  $y$ -cycle and  $x_3, x_4$  belong to the same  $y$ -cycle, and so that the ordering of the indices is consistent with the order of the vertices on the  $x$ -cycle. Then the connected component of  $\bar{X}_C \otimes_{X_A} \bar{X}_C$  containing one of  $(x_i, x_j)$  is

- the graph in Figure 4 (a), if  $|i - j| = 2$ , in which case  $C \cap g^{-1}Cg$  is  $\langle x^{2m+1}, y^n, x^{-1}y \rangle$  or  $\langle x^{2m+1}, y^n, yx^{-1} \rangle$ , or
- a bouquet of two circles, labeled by  $x^{2m+1}$  and  $y^n$  otherwise, in which case  $C \cap g^{-1}Cg$  is  $\langle x^{2m+1}, y^n \rangle$ .

Every other non-diagonal connected component of  $\bar{X}_C \otimes_{X_A} \bar{X}_C$  is either trivial or a single circle, which is labeled by either  $x^{2m+1}$  or  $y^n$ , in which case  $C \cap g^{-1}Cg$  is  $\langle x^{2m+1} \rangle$  or  $\langle y^n \rangle$ , respectively. ■



**Figure 4.** A connected component of  $\bar{X}_C \otimes_{X_A} \bar{X}_C$  when (a):  $M = 2m + 1, N = 2n$ , and (b):  $M = 2m + 1, N = 2n + 1$ .

We finish with the following observation regarding the case where  $M$  and  $N$  are both odd.

**Remark 3.5.** Let  $M = 2m + 1$ ,  $N = 2n + 1$ , and let  $A, B, C$  be as in Corollary 2.9. Let  $g \in A - C$ . Then the intersection  $C \cap g^{-1}Cg$  is one of the sets:  $\langle x^{2m+1}, y^n, x^{-1}y \rangle$ ,  $\langle x^{2m+1}, y^n \rangle$ ,  $\langle x^{2m+1} \rangle$ ,  $\langle y^n \rangle$ . Let  $X_A, \bar{X}_C$  be as in Figure 3. Let  $x_1, x_2, x_3, x_4$  be the four vertices of valence 4 in  $\bar{X}_C$  ordered consistently with the orientation of the  $x$ -cycle such that  $x_1$  and  $x_4$  are at distance  $m$  in the  $x$ -cycle. Then the connected component of  $\bar{X}_C \otimes_{X_A} \bar{X}_C$  containing vertices  $(x_1, x_3), (x_2, x_4), (x_4, x_1)$  (or vertices  $(x_3, x_1), (x_4, x_2), (x_1, x_4)$ ) looks like the graph in Figure 4 (b).

### 3.3. Proof of residual finiteness

We now use Theorem 3.3 to prove that  $\text{Art}_{2MN}$ , where  $M, N$  are even and at least 4, is residually finite.

**Theorem 3.6.** *Let  $M = 2m, N = 2n$  and both  $\geq 4$ . Then  $\text{Art}_{2MN}$  is residually finite.*

*Proof.* The case where  $M = N = 4$  is proven in Example 3.1, so we assume that at least one of  $M, N$ , say  $M$ , is at least 6. By Proposition 2.5,  $\text{Art}_{2MN}$  splits as an HNN-extension  $A *_B \beta$ , where  $A = \langle x, y \rangle$  and  $B = \langle x^m, y^n, x^{-1}y \rangle$ , and  $\beta: B \rightarrow A$  is given by  $x^m \mapsto x^m$ ,  $y^n \mapsto y^n$  and  $x^{-1}y \mapsto yx^{-1}$ . We deduce residual finiteness of  $\text{Art}_{2MN}$  from Theorem 3.3. We now check that all its assumptions are satisfied.

Let  $\hat{A} = \langle x, y \mid x^m, y^n, (x^{-1}y)^p \rangle$ , where  $p \geq 7$ , and let  $\hat{X}_A$  be the presentation complex of  $\hat{A}$ . Let  $\pi: A \rightarrow \hat{A}$  be the natural quotient. Since  $m \geq 3$ , the group  $\hat{A}$  is a hyperbolic von Dyck triangle group, and in particular, condition (1) of Theorem 3.3 is satisfied.


The image  $\pi(B)$  is a finite cyclic group  $\mathbb{Z}/p$  of order  $p$  generated by  $x^{-1}y$ , and the image  $\pi(\beta(B))$  is a copy of  $\mathbb{Z}/p$  generated by  $yx^{-1}$ . Since  $\pi(B), \pi(\beta(B))$  are finite groups, they form an almost malnormal collection in  $\hat{A}$ , so condition (3) in Theorem 3.3 is satisfied. Let  $\hat{X}_B$  be obtained from  $X_B$  by attaching a 2-cell to each of the left and the right loop of  $X_B$  (left and right in Figure 2) via a 1-to-1 map (corresponding to the relations  $x^m, y^n$ ), and one 2-cell to the middle loop via a  $p$ -to-1 map (corresponding to the relation  $(x^{-1}y)^p$ ). It is immediate that conditions (2) and (4) of Theorem 3.3 hold. See Figure 5. The proof is complete. ■

Similarly, we use Theorem 3.2 to prove that  $\text{Art}_{2MN}$ , where one of  $M, N$  is odd, is residually finite.


**Theorem 3.7.** *Let  $M = 2m + 1$  and  $N = 2n$  be both  $\geq 4$ . Then  $\text{Art}_{2MN}$  is residually finite.*

*Proof.* By Corollary 2.9,  $\text{Art}_{2MN}$  splits as  $A *_C B$ , and therefore  $\text{Art}_{2MN}$  has an index 2 subgroup  $D(A, C, \beta)$ . We use Theorem 3.2 to prove that  $D(A, C, \beta)$  is residually finite.

Let  $\hat{A} = \langle x, y \mid x^{2m+1}, y^n, (x^{-1}y)^p \rangle$ , where  $p \geq 6$ , is even, and let  $\hat{X}_A$  be its presentation 2-complex. Since  $n \geq 2$  and  $2m + 1 \geq 5$ , the group  $\hat{A}$  is a hyperbolic von Dyck



**Figure 5.** The maps  $\hat{f}_-: \hat{X}_B \rightarrow \hat{X}_B^- \rightarrow \hat{X}_A$  (top) and  $\hat{f}_+: \hat{X}_B \rightarrow \hat{X}_B^+ \rightarrow \hat{X}_A$  (bottom). White nodes are contained in the 2-cells whose boundary is mapped  $p$ -to-1.




**Figure 6.** The maps  $\hat{X}_C \rightarrow \hat{X}_C \rightarrow \hat{X}_A$ . White nodes are in the 2-cells whose boundary is mapped  $p$ -to-1.

triangle group, and in particular,  $\hat{A}$  satisfies condition (1) of Theorem 3.2. Let  $\pi: A \rightarrow \hat{A}$  be the natural quotient. Let  $\hat{X}_C$  be a 2-complex obtained from  $\bar{X}_C$  by attaching five 2-cells: one along the unique cycle labeled  $x^{2m+1}$ , one along each of the two cycles labeled by  $y^n$ , and one along each of two cycles  $xy^{-1}$  via a  $p$ -to-one map. See Figure 6. In Lemma 3.8 (below), we verify that condition (2) of Theorem 3.2 is satisfied.

By Proposition 3.4, the intersection of distinct conjugates of  $\pi(C)$  in  $\hat{A}$  is either  $\mathbb{Z}/p$  or trivial. In particular,  $\pi(C)$  is almost malnormal in  $\hat{A}$ , so condition (3) of Theorem 3.2 is satisfied. Finally, we note that the 2-cells of  $\hat{X}_C$  can be pulled back via  $X_C \rightarrow \bar{X}_C$ , and are preserved under the (unique) non-trivial deck transformation of  $X_C \rightarrow X_B$ . See Figure 6. Condition (4) of Theorem 3.2 follows. This proves that  $D(A, C, \beta)$ , and therefore  $\text{Art}_{2MN}$ , is residually finite. ■

**Lemma 3.8.** *The image  $\pi(C)$  is isomorphic to  $\mathbb{Z}/p * \mathbb{Z}/p$ . In particular,  $\pi(C) = \pi_1 \hat{X}_C$ . Moreover,  $\hat{f}$  lifts to an embedding in the universal covers.*

*Proof.* For simplicity, we set  $z = xy^{-1}$ . The image  $\pi(C)$  is generated by  $z$ , and  $z' = x^m z x^{-m}$ . The universal cover of the complex  $\hat{X}_A$  can be identified with the hyperbolic plane. Consider the tiling of  $\mathbb{H}^2$  by a triangle with angles  $\frac{\pi}{2m+1}, \frac{\pi}{n}, \frac{\pi}{p}$ . Each vertex of the tiling is a fixed point of a conjugate of one of  $x, y, z$ , and the action of  $\hat{A}$  preserves the type of a vertex (i.e., whether it is fixed by a conjugate of  $x, y$  or  $z$ ). We abuse the notation and identify each vertex  $v$  with the conjugate  $x^g, y^g$  or  $z^g$  which generates the stabilizer of  $v$  (where  $g$  is some element of  $\hat{A}$ ). The tiling is the dual of the universal cover of the complex  $\hat{X}_A$ , in the way that the vertices of types  $x, y, z$  correspond to the 2-cells with boundary words  $x^{2m+1}, y^n, (xy^{-1})^p$ , respectively.



**Figure 7.** The vertices  $z_1, z_5$  are  $\pi(C)$ -conjugates of  $z$ , and the vertex  $z_3$  is a  $\pi(C)$ -conjugate of  $z'$ . Thick edges are in the image of the tree  $T$ . Odd vertices  $q_1, q_3, q_5$  are either green or red, depending on parity of  $n$ . Even vertices  $q_0, q_2, q_4, q_6$  are either blue or red, depending on parity of  $m$ . Note that the orange segments are not necessarily edges in the tiling.

Consider the Bass–Serre tree  $T$  of the free product  $\mathbb{Z}/p * \mathbb{Z}/p$ , i.e., a regular tree of valence  $p$ , where each vertex is stabilized by a conjugate of one of two  $\mathbb{Z}/p$  factors. In order to prove that  $\pi(C)$  splits as a free product and that  $\hat{f}$  lifts to embedding of the universal covers, we show that there is  $\mathbb{Z}/p * \mathbb{Z}/p$ -equivariant embedding of  $T$  in  $\mathbb{H}^2$  where the action of  $\mathbb{Z}/p * \mathbb{Z}/p$  on  $\mathbb{H}^2$  is the action of the group  $\langle z, z' \rangle$ .

First consider the union of the orbits of  $z$  and  $z'$  under the action of  $\pi(C)$ . Note that it is a collection of vertices of type  $z$ . We join  $z^g$  and  $(z')^g$  by a path consisting of two edges of the tiling meeting at a vertex of type  $x$ . See Figure 7. Note that the image of the orbits of  $z$  and  $z'$  under the action of subgroup  $\langle z^{p/2}, (z')^{p/2} \rangle \simeq \mathbb{Z}/2 * \mathbb{Z}/2$  is contained in a geodesic line in  $\mathbb{H}^2$  (which is the line stabilized by  $\langle z^{p/2}, (z')^{p/2} \rangle$ ).

The map from  $T$  to  $\mathbb{H}^2$  is locally injective at every vertex. In order to see that  $T$  is embedded in  $\mathbb{H}^2$ , we verify that any bi-infinite path  $\gamma$  always turning rightmost in the image of  $T$  never crosses itself. Indeed, if  $T$  were not embedded, then a closed path in the image of  $T$  could be tracked by following a path turning rightmost in  $T$ . In Figure 7, a part of that path is presented as the path with vertices  $x_0, z_1, x_2, z_3, x_4, z_5, x_6$ . We will construct another path  $\gamma'$  (whose part is labeled by  $q_0, s_0, q_1, s_1, \dots, q_5, s_5, q_6$  in Figure 7) that stays at a finite distance from  $\gamma$ . We will think of  $\gamma'$  as an oriented path (where  $q_0$  arises before  $q_1$  etc.). By construction (describe below),  $\gamma'$  never intersects the image of  $T$ . Since the image of  $T$  contains a geodesic line and lies in the region on the left side of  $\gamma'$  with respect to its orientation, the region of  $\mathbb{H}^2$  on the left side of  $\gamma'$  is unbounded. It suffices to show that the region of  $\mathbb{H}^2$  on the right side of  $\gamma'$  is unbounded. We will do it by showing that the region on the right side of  $\gamma'$  is a union of halfspaces.

Let us explain how the vertices  $q_i, s_i$  are defined. The vertex  $s_i$  with odd  $i$  is the unique vertex other than  $z_i$  that forms a triangle with  $y_i$  and  $x_{i+1}$ . The vertex  $s_i$  with even  $i$  is the unique vertex other than  $z_{i+1}$  that forms a triangle with  $x_i$  and  $y_{i+1}$ . In particular, vertices  $s_i$  are always of type  $z$ . The vertices  $q_i$  with odd  $i$  are images of a  $z_i$  under the

$\pi$ -rotation at  $y_i$ , i.e., the vertex  $y_i$  is a midpoint of the segment  $[z_i, q_i]$ . The vertices  $q_i$  with even  $i$  are chosen so that the angles  $\angle s_{i-1}x_iq_i$  and  $\angle s_ix_iq_i$  are equal. The path  $\gamma'$  is obtained by joining each pair  $s_{i-1}, q_i$  and  $q_i, s_i$  by a geodesic segment (which are not necessarily edges of the tiling).

The vertices  $s_{i-1}, q_i, s_i$  are not necessarily distinct. If  $n = 2$ , then  $s_{i-1} = q_i = s_i$  for each odd  $i$ . Also, if  $2m + 1 = 5$ , then  $s_{i-1} = q_i = s_i$  for  $i = 4k + 2$ . In that extreme case,  $\gamma'$  is a geodesic line, as long as  $p \geq 6$ . In more general case, the “left-side” angle between the segments of  $\gamma'$  at each vertex  $q_i$  or  $s_i$  (i.e., the angle of the sector containing  $y_i$  or  $x_i$  depending on the parity of  $i$ ) is always at most  $\pi$ . Thus the “right-side” angle at each vertex  $q_i, s_i$  is always at least  $\pi$ . Consequently, the subspace of  $\mathbb{H}^2$  bounded by  $\gamma'$  which is on the right side of  $\gamma'$  is a union of halfspaces. This proves our claim. ■

Our approach in the last two theorems fails in the case where  $M = 2m + 1, N = 2n + 1$ . Indeed, the fiber product  $\bar{X}_C \otimes_{X_A} \bar{X}_C$  is too “large”. The fiber product was computed in Remark 3.5. After attaching 2-cells along  $x^{2m+1}, y^{2n+1}$  and  $(x^{-1}y)^p$ , the resulting 2-complex has fundamental group  $\mathbb{Z} * \mathbb{Z}/p$ .

### 3.4. Summary of residual finiteness of 3-generator Artin groups

To summarize, the only 3-generator Artin groups that are not known to be residually finite are  $\text{Art}_{33(2m+1)}$  for  $m \geq 2$ ,  $\text{Art}_{2(2m+1)(2n+1)}$  for  $m + n \geq 4$  and  $\text{Art}_{23(2m)}$  for  $m \geq 4$ . Indeed, if at least one label is  $\infty$ , then the defining graph is a tree, and hence virtually special [7, 14, 20, 23]. Artin groups  $\text{Art}_{22M}$  for any  $M \geq 2$ , and  $\text{Art}_{23M}$ , where  $M \in \{3, 4, 5\}$  are spherical, and so linear [9, 11]. The cases  $(3, 3, 3)$ ,  $(2, 4, 4)$  and  $(2, 3, 6)$  follow from [25]. The cases where  $M, N, P \geq 3$ , except the case of  $(3, 3, 2m + 1)$ , were covered by [18].

**Acknowledgments.** The author thanks the referee for the helpful suggestions.

**Funding.** This material is based upon work supported by the National Science Foundation under Grant No. DMS-2105548/2203307, and under Grant No. DMS-1928930 while the author participated in a program hosted by the Mathematical Sciences Research Institute in Berkeley, California, during the Fall 2020 semester.

## References

- [1] M. Bestvina and M. Feighn, [A combination theorem for negatively curved groups](#). *J. Differential Geom.* **35** (1992), no. 1, 85–102 Zbl 0724.57029 MR 1152226
- [2] M. Bhattacharjee, [Constructing finitely presented infinite nearly simple groups](#). *Comm. Algebra* **22** (1994), no. 11, 4561–4589 Zbl 0834.20020 MR 1284345
- [3] S. J. Bigelow, [Braid groups are linear](#). *J. Amer. Math. Soc.* **14** (2001), no. 2, 471–486 Zbl 0988.20021 MR 1815219

- [4] R. Blasco-García, A. Juhász, and L. Paris, [Note on the residual finiteness of Artin groups](#). *J. Group Theory* **21** (2018), no. 3, 531–537 Zbl [1425.20020](#) MR [3794930](#)
- [5] R. Blasco-García, C. Martínez-Pérez, and L. Paris, [Poly-freeness of even Artin groups of FC type](#). *Groups Geom. Dyn.* **13** (2019), no. 1, 309–325 Zbl [1498.20086](#) MR [3900773](#)
- [6] E. Brieskorn and K. Saito, [Artin-Gruppen und Coxeter-Gruppen](#). *Invent. Math.* **17** (1972), 245–271 Zbl [0243.20037](#) MR [323910](#)
- [7] A. M. Brunner, [Geometric quotients of link groups](#). *Topology Appl.* **48** (1992), no. 3, 245–262 Zbl [0774.57004](#) MR [1200426](#)
- [8] M. Burger and S. Mozes, [Finitely presented simple groups and products of trees](#). *C. R. Acad. Sci. Paris Sér. I Math.* **324** (1997), no. 7, 747–752 Zbl [0966.20013](#) MR [1446574](#)
- [9] A. M. Cohen and D. B. Wales, [Linearity of Artin groups of finite type](#). *Israel J. Math.* **131** (2002), 101–123 Zbl [1078.20038](#) MR [1942303](#)
- [10] P. Deligne, [Les immeubles des groupes de tresses généralisés](#). *Invent. Math.* **17** (1972), 273–302 Zbl [0238.20034](#) MR [422673](#)
- [11] F. Digne, [On the linearity of Artin braid groups](#). *J. Algebra* **268** (2003), no. 1, 39–57 Zbl [1066.20044](#) MR [2004479](#)
- [12] T. Haettel, [Virtually cocompactly cubulated Artin–Tits groups](#). *Int. Math. Res. Not. IMRN* **2021** (2021), no. 4, 2919–2961 Zbl [1499.20103](#) MR [4218342](#)
- [13] T. Haettel, [Cubulation of some triangle-free Artin groups](#). *Groups Geom. Dyn.* **16** (2022), no. 1, 287–304 Zbl [07531897](#) MR [4424971](#)
- [14] S. M. Hermiller and J. Meier, [Artin groups, rewriting systems and three-manifolds](#). *J. Pure Appl. Algebra* **136** (1999), no. 2, 141–156 Zbl [0936.20033](#) MR [1674774](#)
- [15] T. Hsu and D. T. Wise, [Cubulating graphs of free groups with cyclic edge groups](#). *Amer. J. Math.* **132** (2010), no. 5, 1153–1188 Zbl [1244.20040](#) MR [2732342](#)
- [16] T. Hsu and D. T. Wise, [Cubulating malnormal amalgams](#). *Invent. Math.* **199** (2015), no. 2, 293–331 Zbl [1320.20039](#) MR [3302116](#)
- [17] J. Huang, K. Jankiewicz, and P. Przytycki, [Cocompactly cubulated 2-dimensional Artin groups](#). *Comment. Math. Helv.* **91** (2016), no. 3, 519–542 Zbl [1401.20044](#) MR [3541719](#)
- [18] K. Jankiewicz, [Residual finiteness of certain 2-dimensional Artin groups](#). *Adv. Math.* **405** (2022), article no. 108487 Zbl [07557151](#) MR [4437605](#)
- [19] D. Krammer, [Braid groups are linear](#). *Ann. of Math. (2)* **155** (2002), no. 1, 131–156 Zbl [1020.20025](#) MR [1888796](#)
- [20] Y. Liu, [Virtual cubulation of nonpositively curved graph manifolds](#). *J. Topol.* **6** (2013), no. 4, 793–822 Zbl [1286.57002](#) MR [3145140](#)
- [21] A. Malcev, [On isomorphic matrix representations of infinite groups](#). *Rec. Math. (N.S.)* **8** (1940), 405–422 Zbl [66.0088.03](#)
- [22] A. Mal'cev, [On homomorphisms onto finite groups](#). In *Twelve papers in algebra*, pp. 67–79, Amer. Math. Soc. Transl. Ser. 2 119, American Mathematical Society, Providence, RI, 1983 Zbl [0511.20026](#) MR [693297](#)
- [23] P. Przytycki and D. T. Wise, [Graph manifolds with boundary are virtually special](#). *J. Topol.* **7** (2014), no. 2, 419–435 Zbl [1331.57023](#) MR [3217626](#)
- [24] P. Scott and T. Wall, [Topological methods in group theory](#). In *Homological group theory (Proc. Sympos., Durham, 1977)*, pp. 137–203, London Math. Soc. Lecture Note Ser. 36, Cambridge University Press, Cambridge, 1979 Zbl [0423.20023](#) MR [564422](#)
- [25] C. C. Squier, [On certain 3-generator Artin groups](#). *Trans. Amer. Math. Soc.* **302** (1987), no. 1, 117–124 Zbl [0637.20016](#) MR [887500](#)

- [26] J. R. Stallings, [Topology of finite graphs](#). *Invent. Math.* **71** (1983), no. 3, 551–565  
Zbl [0521.20013](#) MR [695906](#)
- [27] H. Wilton, [Essential surfaces in graph pairs](#). *J. Amer. Math. Soc.* **31** (2018), no. 4, 893–919  
Zbl [1452.20038](#) MR [3836561](#)
- [28] D. T. Wise, *No-positively curved squared complexes: Aperiodic tilings and non-residually finite groups*. Ph.D. thesis, 1996, Princeton University MR [2694733](#)
- [29] D. T. Wise, [The residual finiteness of negatively curved polygons of finite groups](#). *Invent. Math.* **149** (2002), no. 3, 579–617 Zbl [1040.20024](#) MR [1923477](#)

Received 28 August 2021.

**Kasia Jankiewicz**

Department of Mathematics, University of California – Santa Cruz, 1156 High Street, Santa Cruz,  
CA 95060, USA; [kasia@ucsc.edu](mailto:kasia@ucsc.edu)