

The weak hyperbolization conjecture for 3-dimensional CAT(0)-groups

Michael Kapovich and Bruce Kleiner

Abstract. We prove a weak hyperbolization conjecture for CAT(0) 3-dimensional Poincaré duality groups.

Mathematics Subject Classification (2000). Primary 20F65, 20F67; Secondary 57P10.

Keywords. Poincaré duality groups; CAT(0)-groups.

1. Introduction

For a variety of classes of groups, it is a well-known open problem whether the failure of Gromov hyperbolicity can be detected by the presence of special subgroups, e.g. rank 2 abelian groups or Baumslag–Solitar groups. This is of interest, for instance, for CAT(0)-groups (even for the fundamental groups of finite 2-dimensional locally CAT(0) square complexes), for 1-relator groups, and 3-dimensional Poincaré duality groups. We say that a class of groups *satisfies the weak hyperbolization conjecture* if every group in the class is either Gromov hyperbolic, or contains a copy of \mathbb{Z}^2 . We recall that the weak hyperbolization conjecture for 3-manifold groups was a part of the program for proving the Geometrization Conjecture for closed irreducible aspherical 3-manifolds, the other ingredient in the program being the Cannon conjecture. Although the work of Perelman has now resolved the full Geometrization Conjecture, the weak hyperbolization conjecture for PD(3)-groups is a potential step in an approach to the following open question of C. T. C. Wall:

Question 1 (Wall). Is every finitely presented PD(3)-group over \mathbb{Z} isomorphic to the fundamental group of a closed aspherical 3-manifold?

Our main result is that the weak hyperbolization conjecture holds for CAT(0) 3-dimensional Poincaré duality groups over hereditary rings:

Theorem 2. *Let G be a 3-dimensional Poincaré duality group over a commutative hereditary ring \mathcal{R} with a unit. Suppose in addition that G is a CAT(0)-group, i.e., a group which admits a cocompact isometric properly discontinuous action $G \curvearrowright X$ on a locally compact CAT(0) space X .*

Then G satisfies the weak hyperbolization conjecture.

We refer the reader to [9] for the definition of a hereditary ring; here we note only that every PID is hereditary.

We note that special cases of this theorem were proven earlier by various people: S. Buyalo [8] and V. Schroeder [18] independently have proven that this theorem holds provided that X is the universal cover \tilde{M} of a closed 3-manifold M , the CAT(0) structure on \tilde{M} is Riemannian and $G = \pi_1(M)$ acts on X by deck-transformations. L. Mosher [16] proved that Theorem 2 holds provided that $X = \tilde{M}$, $G = \pi_1(M)$, and the CAT(0) metric on X is obtained by lifting a piecewise-Euclidean (locally) CAT(0)-cubulation from M . M. Bridson and L. Mosher also have an unpublished proof of Theorem 2 under the assumption that $X = \tilde{M}$ has an arbitrary G -invariant CAT(0) structure. Unlike all these proofs, our proof takes place on the ideal boundary of X ; this allows us to treat 3-dimensional Poincaré duality groups and relax the assumptions on the CAT(0) space.

Outline of the proof of Theorem 2. Assume that G is not Gromov hyperbolic, i.e., that X contains a 2-flat. By the work of Bestvina [2], the ideal boundary of X is homeomorphic to S^2 . Our proof exploits the geometry of flats and parallel sets in X , and the pattern of their boundaries in the 2-sphere $\partial_\infty X$. The proof breaks into three cases.

Case 1. X contains a 3-flat, Section 5.1. This implies that X is at finite Hausdorff distance from the 3-flat, and we conclude that G is virtually \mathbb{Z}^3 .

Case 2. X contains no 3-flat but some parallel set $P \subset X$ has full ideal boundary, i.e. $\partial_\infty P = \partial_\infty X$, Section 5.3. We argue that P splits isometrically as $\mathbb{R} \times Y$, where $\partial_\infty Y$ is a circle, and G acts as a convergence group on $\partial_\infty Y$. We then deduce that a finite index subgroup of G is isomorphic to the fundamental group of a 3-dimensional Seifert manifold.

Case 3. X contains no 3-flat and no parallel set with full boundary, Section 5.4. This is the main case. We show that every parallel set P in X is isometric to a product $\mathbb{R} \times Y$, where Y is Gromov hyperbolic. The ideal boundary of P is a suspension of the boundary $\partial_\infty Y$; when P contains a 2-flat, we identify certain topological circles in $\partial_\infty P$ which we call peripheral, and show that peripheral circles cannot cross one another in the 2-sphere $\partial_\infty X$.

Next, we choose a flat $F \subset X$ whose boundary $\partial_\infty F \subset \partial_\infty X$ is a peripheral circle, and consider its orbit $\{g(F)\}_{g \in G}$. Because the circles $\{g(\partial_\infty F)\}_{g \in G}$ do not cross, we may use them to define a pretree \mathcal{T} on which G has a natural action. Using a Plante-type construction, we associate to \mathcal{T} an \mathbb{R} -tree T , which then inherits a

nontrivial small stable G -action. By applying Rips' theory [3], we conclude that G admits a small nontrivial action on a simplicial tree. Using the fact that G is a PD(3)-group, we deduce that the edge groups must be virtually \mathbb{Z}^2 .

Acknowledgements. The first author was supported in part by NSF Grants DMS-02-03045 and DMS-04-05180, the second author was supported in part by NSF Grants DMS-02-24104 and DMS-05-05610. The authors are grateful to the referee for useful suggestions.

2. Geometric preliminaries

In this section we briefly review several notions of metric geometry. We refer the reader to [1], [6] for the detailed discussion.

A *geodesic metric space* is a metric space (X, d) such that any two points $x, y \in X$ in X are connected by geodesic, i.e., if $D := d(x, y)$ then there exists an isometric embedding

$$\gamma: [0, D] \rightarrow X$$

so that $\gamma(0) = x, \gamma(D) = y$.

Let X be a metric space and $C \subset X$ be a subset. The r -neighborhood of C in X is defined as

$$N_r(C) := \{x \in X : d(x, C) < r\},$$

where $d(x, C) := \inf\{d(x, c) : c \in C\}$.

The *Hausdorff distance* between closed subsets of a metric space X is defined as

$$d_H(C_1, C_2) := \inf\{r : C_1 \subset N_r(C_2), C_2 \subset N_r(C_1)\}.$$

Note that this distance is allowed to take infinite values. If X has finite diameter, the Hausdorff distance defines the *Hausdorff topology* on the set $\mathcal{C}(X)$ of closed subsets of X . More generally, even for unbounded metric spaces X one defines the *Gromov–Hausdorff topology* on $\mathcal{C}(X)$ as follows. We say that a sequence $C_n \in \mathcal{C}(X)$ converges (in the Gromov–Hausdorff topology) to a closed set $C \in \mathcal{C}(X)$ if for each closed metric ball $B \subset X$ the intersections

$$C_n \cap B \in \mathcal{C}(B)$$

converge to $C \cap B$ in the Hausdorff topology on $\mathcal{C}(B)$. Equivalently, C_n 's converge to C if the corresponding distance functions $d(\cdot, C_n)$ converge to the distance function $d(\cdot, C)$ uniformly on bounded subsets in X .

Given a number $\kappa \in \mathbb{R}$ let M_κ denote the (unique up to isometry) complete simply-connected surface of the constant curvature κ . A geodesic metric space X is said to be a CAT(κ) space if X is complete as a metric space and geodesic triangles

in X are “thinner” than triangles in M_κ . More precisely, consider a geodesic triangle $T = [x, y, z] \subset X$ (with the vertices x, y, z), in case when $\kappa > 0$ (and M_κ is a sphere) we assume that the perimeter of this triangle is less than the circumference of the great circle in M_κ . Consider a triangle $T' = [x', y', z'] \subset M_\kappa$ whose side-lengths are equal to the corresponding side-lengths of the triangle T . Let p be a point in the geodesic side \overline{xy} of T and let $p' \in \overline{x'y'}$ be such that

$$d(x', p') = d(x, p).$$

Then we require

$$d(z, p) \leq d(z', p').$$

In this paper we will also need a generalization of the concept of a CAT(1) space to metric spaces X which are not geodesic. We assume that X is a disjoint union of geodesic metric spaces X_α , $\alpha \in J$, where each X_α is a geodesic CAT(1) metric space and if $\alpha \neq \beta$ the distance between any $x \in X_\alpha$, $y \in X_\beta$ equals π . Then X will be also referred to as a CAT(1) space. An example of such a space is a space with discrete metric where distance between any pair of distinct points equals π .

If X is a CAT(1) space, we call points $x, y \in X$ *antipodal* if $d(x, y) = \pi$.

Suppose that X is a CAT(0) space. Then the distance function on X is *convex*, i.e., its restriction to each geodesic in X is convex.

A space X is called CAT($-\infty$) if it is CAT(κ) for each $\kappa \in \mathbb{R}$. A *metric tree* is a CAT($-\infty$); in other words, it is a complete geodesic metric space where each geodesic triangle is isometric to a tripod.

A group G is called a CAT(0)-*group* if it admits an isometric properly discontinuous cocompact action on a locally compact CAT(0) space.

Suppose that X is a CAT(0) space and $F \subset X$ is a *k-flat*, i.e., an isometrically embedded copy of a Euclidean space \mathbb{R}^k . Then the *parallel set* P_F of F in X is the union of all k -flats $F' \subset X$ which are within finite distance from F . The parallel set P_F is closed, convex and is isometric to a product

$$F \times Y$$

where Y is a CAT(0) space, see for instance [6, Theorem II.2.14].

Remark 3. Theorem II.2.14 in [6] is stated in the case $k = 1$. The general case follows, for instance, by induction on the dimension of the flat.

We will say that a parallel set is *trivial* if $k = 1$ and Y is bounded.

Given a CAT(0) space one defines the *ideal boundary* of X as the collection of equivalence classes of geodesic rays in X , where rays are equivalent if they are within finite Hausdorff distance from each other. This boundary has two (typically distinct) topologies:

1. the *visual topology*, in which case the ideal boundary is denoted $\partial_\infty X$ and is called the *geometric boundary* of X ;
2. the *Tits topology*, which is defined via the *Tits angular metric*, in which case the ideal boundary is denoted $\partial_{\text{Tits}} X$.

The second boundary is called *Tits boundary* of X ; this boundary is always a CAT(1) space.

For instance, in the case when $X = \mathbb{H}^2$, $\partial_\infty X$ is homeomorphic to S^1 , while $\partial_{\text{Tits}} X$ has discrete metric: the distance between distinct points equals π . A CAT(0) space is called a *visibility space* if any pair of distinct points in $\partial_{\text{Tits}} X$ are antipodal.

A subset $C \subset Z := \partial_{\text{Tits}} X$ is called *convex* if for any two non-antipodal points $x, y \in Z$, the geodesic segment \overline{xy} connecting x to y , is entirely contained in C . Intersection of two convex subsets of Z is also convex. If $Y \subset X$ is a convex subset then $\partial_{\text{Tits}} Y \subset Z$ is convex as well.

Let $\delta \in [0, \infty)$ and consider a geodesic metric space X . A triangle $T \subset X$ is called δ -*thin* if there exists a point $p \in X$ which is within distance $\leq \delta$ from all three sides of T . A complete geodesic metric space X is called δ -*hyperbolic* if each geodesic triangle T in X is δ -thin. A space X is called *Gromov-hyperbolic* if it is δ -hyperbolic for some δ . A finitely generated group G is called *Gromov-hyperbolic* if its Cayley graph is Gromov-hyperbolic. One again defines the ideal boundary $\partial_\infty X$ by looking at the equivalence classes of geodesic rays in X .

Suppose that G is a group acting isometrically, properly discontinuously and cocompactly on a CAT(0) space X . Then the group G is Gromov-hyperbolic iff X is a visibility space.

Let X be a Gromov-hyperbolic geodesic metric space which admits a cocompact isometric group action. We assume that the ideal boundary of X consists of more than 2 points; it then follows that $\partial_\infty X$ has the cardinality of the continuum. The *displacement function* of an isometry $g: X \rightarrow X$ is

$$\text{dis}(g): x \rightarrow d(x, g(x)), \quad x \in X.$$

Lemma 4. *Under the above assumptions there exists a constant $D = D(X)$ such that for each $g \in \text{Isom}(X)$ which fixes $\partial_\infty X$ pointwise, the displacement of g is bounded from above by D .*

Proof. Let $G \curvearrowright X$ be a cocompact isometric group action; pick a metric ball $B = B(o, R) \subset X$ so that the G -orbit of B equals X . It then suffices to prove that there exists $D < \infty$ such that for each isometry g of X fixing $\partial_\infty X$ pointwise,

$$d(o, g(o)) \leq D.$$

Since the ideal boundary of X contains at least 4 points, there exists a pair of geodesics $\gamma_1, \gamma_2 \subset X$ which have disjoint ideal boundaries. Without loss of generality we may assume that both γ_1, γ_2 pass through the ball B .

Since X is δ -hyperbolic, there exists a number $r = r(\delta) < \infty$ such that if geodesics $\alpha, \beta \subset X$ are within finite Hausdorff distance, then

$$d_H(\alpha, \beta) \leq r,$$

see for instance [6]. For every isometry g as above, the geodesics

$$\gamma_i, g(\gamma_i)$$

are within finite Hausdorff distance from each other; therefore

$$d_H(\gamma_i, g(\gamma_i)) \leq r, \quad i = 1, 2.$$

Then

$$d(g(o), g(\gamma_i)) \leq R \implies d(g(o), \gamma_i) \leq R + r, \quad i = 1, 2.$$

However, since the geodesics γ_1, γ_2 have disjoint ideal boundaries, the diameter of

$$S := N_{R+r}(\gamma_1) \cap N_{R+r}(\gamma_2)$$

is finite. Therefore, if we take $D := \text{diam}(S)/2$, the distance between o and $g(o)$ is at most D . \square

Remark 5. An analogue of Lemma 4 holds for quasi-isometries of X with uniformly bounded quasi-isometry constants.

3. Pretrees

In what follows we will need definitions and basic facts about pretrees; the definitions which we give follow [5].

A *pretree* is a set T together with a ternary relation (the *betweenness relation*)

“ y is between x and z ”,

to be denoted $\beta(xyz)$, satisfying the following axioms:

Axiom 1. $\beta(xyz)$ implies that $x \neq y \neq z$.

Axiom 2. $\beta(xyz) \iff \beta(zyx)$.

Axiom 3. $\beta(xyz)$ and $\beta(yxz)$ cannot hold simultaneously.

Axiom 4. If $w \neq y$ then $\beta(xyz)$ implies that either $\beta(xyw)$ or $\beta(wyz)$.

Given a pretree T one can define *closed*, *open* and *half-open* intervals in T by

$$(x, z) := \{y \in T : \beta(xyz)\}, \quad [x, z] := (x, z) \cup \{x, z\}, \text{ etc.}$$

Given an increasing union of intervals

$$[x_1, y_1] \subset [x_2, y_2] \subset \cdots \subset [x_i, y_i] \subset \cdots$$

we will also refer to the union of these intervals as a (possibly infinite) interval in T .

We note that β defines an order on each interval in T .

Define a “triangle” in T with vertices a, b, c to be the union of the segments (called “sides” of the triangle) $[a, b], [b, c], [c, a]$.

Lemma 6. *Each triangle Δ in T is 0-thin, i.e., each side of Δ is contained in the union of the two other sides.*

Proof. Follows immediately from Axiom 4. □

Suppose that T is a pretree which is given a measure μ (without atoms) defined on closed intervals in T and the σ -algebra which these intervals generate. Define a function $d(x, y)$ on T by $d(x, y) := \mu([x, y])$.

Lemma 7. *d is a pseudo-metric on T .*

Proof. It is clear that d is symmetric and $d(x, x) = 0$ (since μ has no atoms). The triangle inequality follows because for each triangle with the vertices a, b, c we have (see Lemma 6)

$$[a, b] \subset [a, c] \cup [b, c]. \quad \square$$

We note that if for each interval $[a, b] \subset T$, with $a \neq b$, $\mu(a, b) > 0$ then d is a metric. Moreover, it follows that $(a, b) \neq \emptyset$ for each $a \neq b$. If the restriction of the metric d to each interval $[x, y]$ is complete then $[x, y]$ is order isomorphic to an interval in \mathbb{R} and moreover, $([x, y], d)$ is isometric to an interval in \mathbb{R} . We thus get:

Lemma 8. *Suppose that for each interval $[x, y] \subset T$, with $x \neq y$, $\mu[x, y] > 0$, and that the restriction of the metric d to each interval in T is complete. Then (T, d) is a metric tree.*

Proof. It is clear from the above discussion that T is a geodesic metric space. Since each triangle in T is 0-thin, it follows that each triangle in T is isometric to a tripod. Finally, let us check completeness of T : Suppose that $x_i, i \geq 0$, is a Cauchy sequence in T . Then there exists an increasing sequence of intervals $I_i \subset T$ such that

$$\lim_i \mu([x_0, x_i] \cap I_i) = \lim_i d(x_0, x_i).$$

Then completeness of d restricted to the union I of I_i 's implies that (x_i) converges to a point in the interval I . □

4. Ideal boundaries of CAT(0) Poincaré duality groups

Let $G \curvearrowright X$ be a discrete cocompact action of a PD(3)-group G on a CAT(0) space X . In this section we show that the ideal boundary of the CAT(0) space X is homeomorphic to S^2 .

We refer the reader to [4], [7] for the background on the cohomology of groups. Recall [4] that an n -dimensional Poincaré duality group over a ring \mathcal{R} (for short, PD(n)-group over \mathcal{R}), is an FP-group over \mathcal{R} such that $H^i(G, \mathcal{R}G)$ is isomorphic to \mathcal{R} as an \mathcal{R} -module when $i = n$ and is trivial otherwise.

Let $Z := \partial_\infty X$ be the ideal boundary of a locally compact CAT(0) space. M. Bestvina in [2] proved that the compactification

$$\bar{X} := X \cup Z$$

satisfies the axioms of the Z -set compactification. Instead of listing all the axioms of the Z -set compactification we mention only several properties:

1. If $G \curvearrowright X$ is an isometric group action then this action extends to a topological action of G on \bar{X} .
2. There exists a natural isomorphism

$$H_c^*(X) \rightarrow \tilde{H}_c^{*-1}(Z),$$

which is compatible with inclusions of closed convex subsets $X' \subset X$.

3. We state the third property as a lemma:

Lemma 9. *If G is a PD(3)-group acting isometrically, properly discontinuously and cocompactly on a CAT(0) space X , then the ideal boundary Z of X is homeomorphic to S^2 .*

Proof. Bestvina proves, [2, Theorem 2.8], that if G is a PD(3)-group over \mathcal{R} , then Z is homeomorphic to S^2 . We note that Bestvina proves the latter theorem under more restrictive assumptions than we are working with (although, his class of groups G includes 3-manifold groups):

1. Bestvina assumes that the commutative ring \mathcal{R} is a PID. However this assumption is used only to apply the Universal Coefficients Theorem, which works for hereditary rings as well, see [9].

2. Bestvina's definition of an n -dimensional Poincaré duality group is more restrictive than the usual one: Instead of the FP-property he assumes that a group G acts freely, properly discontinuously, cocompactly on a contractible cell complex Y . Note however that Bestvina in his proof uses only the fact that $G \curvearrowright Y^{(i)}$ is cocompact on each i -skeleton of Y . Then existence of such an action for the CAT(0)-groups follows from a general construction described in [14]. Namely, if a group G admits a properly discontinuous cocompact action on a contractible space X (e.g. the CAT(0) space in

our case) then it also admits a free, properly discontinuous action on a contractible cell complex Y (possibly of infinite dimension) such that $Y^{(i)}/G$ is compact for each i .

3. Bestvina assumes that the image of the orientation character χ of the Poincaré duality group G is finite (he then passes to a finite index subgroup in G which is the kernel of χ). However this assumption can be omitted from his theorem using *twisting* of the action $G \curvearrowright C_*(Y)$ by the character χ as it is done in [14, Section 5.1].

With the above modifications, Bestvina's arguments apply in our case and it follows that $\partial_\infty X$ is homeomorphic to the 2-sphere. \square

5. Proof of the main theorem

5.1. Case 1: X contains a 3-flat. The main goal of this section is to show that, in case X contains a 3-flat, the group G contains a finite index subgroup isomorphic to \mathbb{Z}^3 .

Lemma 10. *Suppose that S is a convex subset in X such that $\partial_\infty S = \partial_\infty X$. Then S is within finite Hausdorff distance from X .*

Proof. Pick a base-point $o \in X$. If S is not within finite Hausdorff distance from X then there exists a sequence of isometries $g_i \in G$ such that $d(o, g_i S)$ diverges to infinity. Consider the functions $f_i := d(x, g_i S) - d(o, g_i S)$. Then, according to Lemma 2.3 in [15], the sequence of functions f_i subconverges to a Busemann function b on X . Clearly, the sublevel sets $\{f_i \leq 0\}$ subconverge into the horoball $U := \{b \leq 0\}$ in X . Since $\partial_\infty \{f_i \leq 0\} = \partial_\infty g_i S = \partial_\infty X$, it follows that $\partial_\infty X = \partial_\infty U$.

Let F be a 2-flat in X . Then $\partial_\infty F \subset \partial_\infty U$ and the convexity of horoballs in X imply that for each $x \in F$,

$$t = f(x) \implies F \subset \{z : b(z) \leq t\}.$$

It follows that the restriction $b|_F$ is constant and thus F is contained in the horosphere $\{x : b(x) = t\}$ for some $t \in \mathbb{R}$. Then Lemma 2.2 in [15] implies that X contains a half-space $H := \mathbb{R}_+ \times F$. Then, by taking an appropriate limit of the half-spaces $h_j(H)$, $h_j \in G$, we see that X contains the 3-flat $F' := F \times \mathbb{R}$. By Lemma 9, $\partial_\infty F' = \partial_\infty X$. Suppose that F' is not within finite Hausdorff distance from X . Then, by repeating the same argument as above with S replaced with F' and then F replaced with F' , we see that X contains a 4-flat, which contradicts Lemma 9.

Therefore X is within finite Hausdorff distance from the 3-flat F' ; in particular, there are no horoballs in X which have the same ideal boundary as X . Contradiction. \square

Corollary 11. *If X contains a 3-flat then the group G is virtually abelian; in particular, it contains $\mathbb{Z} \times \mathbb{Z}$.*

Proof. If F is a 3-flat in X then, by Lemma 9, $\partial_\infty F = \partial_\infty X$ and, by Lemma 10, F is within finite Hausdorff distance from X . It follows that the group G is isomorphic to a lattice in $\text{Isom}(\mathbb{R}^3)$ and hence it is virtually abelian and contains \mathbb{Z}^3 as a subgroup of finite index. \square

Assumption. From now on we will assume that X contains no 3-flats.

5.2. Metric balls and parallel sets in X . In this section we establish certain geometric properties of X which follow from the above assumption.

Lemma 12. *There exists $r_0 \in \mathbb{R}$ such that the following holds. For each ball $B(x, r) \subset X$, isometric to a disk of the radius r in \mathbb{R}^3 , we have $r \leq r_0$.*

Proof. If the assertion is false then there exists a sequence of balls $B(x_i, r_i)$ with $\lim_i r_i = \infty$. Let $g_i \in G$ be such that $g_i(x_i)$ is a bounded sequence in X . Then the balls $g_i(B(x_i, r_i))$ subconverge to a 3-flat in X . Contradiction. \square

Corollary 13. *The set of 2-flats $F' \subset X$ which are parallel to a flat F is compact in the Gromov–Hausdorff topology.*

Proof. If not then X contains convex subsets isometric to $[0, r] \times \mathbb{R}^2$ for arbitrarily large r . This contradicts the previous lemma. \square

Lemma 14. *Suppose that $Y \times \mathbb{R}$ is a parallel set in X . Then Y is Gromov-hyperbolic.*

Proof. We repeat the arguments in [6, Theorem 9.33]. If Y is not Gromov-hyperbolic then there exists a pair of points $\xi, \eta \in \partial_\infty Y$ so that the Tits angle between ξ, η is positive but less than π . Pick a point $o \in Y$ and consider a sequence of points $y_i \in \overline{o\xi}$ which converge to ξ and the geodesic rays $\overline{y_i\eta}$. We identify the rays $\overline{y_i\xi}, \overline{y_i\eta}$ with geodesic rays in $Y \times \mathbb{R} \subset X$ (that share common point y_i). Then, by applying an appropriate sequence of elements $g_i \in G$ (for which $\{g_i(y_i)\}$ is bounded in X) to $Y \times \mathbb{R}$ and to the rays $\overline{y_i\xi}, \overline{y_i\eta}$ and passing to the limit of a subsequence, we get:

1. The sets $g_i(Y \times \mathbb{R})$ subconverge to a parallel set $Y' \times \mathbb{R}$.
2. Y' contains two geodesic rays $\overline{y\xi'}, \overline{y\eta'}$ (limits of the sequences of rays $g_i(\overline{y_i\xi}), g_i(\overline{y_i\eta})$) which bound a flat sector in Y' .

This contradicts Lemma 12. \square

5.3. Case 2: X contains a parallel set with the full boundary. In this section we prove the main theorem under the assumption that X contains a parallel set P whose ideal boundary is the entire $\partial_\infty X$.

Proposition 15. *Suppose that there is a convex product subset $P = \mathbb{R} \times Y$ such that $\partial_\infty S = \partial_\infty X$. Then G is commensurable to the fundamental group of a 3-dimensional Seifert manifold. In particular, G contains \mathbb{Z}^2 .*

Proof. We will assume that P is a maximal convex product subset in X . Since Y is Gromov-hyperbolic, it follows that the Tits boundary of S is the suspension of a discrete metric space which is the ideal boundary of Y . Therefore, since $\partial_\infty P = \partial_\infty X$, the group G preserves the ideal boundary of the geodesic $l = \mathbb{R} \times \{y\}$. Hence for each $g \in G$ the geodesic $g(l)$ is parallel to l , which (by the maximality assumption) implies that $g(P) = P$.

We have an induced isometric action $\rho: G \curvearrowright Y$. Since the suspension of $\partial_\infty Y$ is homeomorphic to the 2-sphere $\partial_\infty X$, the ideal boundary of Y is homeomorphic to S^1 . Thus the cocompact isometric action $\rho: G \curvearrowright Y$ extends to a uniform (topological) convergence action $G \curvearrowright \partial_\infty Y = S^1$. Therefore, according to [10], [12], [13], [19], the action $G \curvearrowright S^1$ is topologically conjugate to a Moebius action ρ' .

Let K denote the kernel of ρ' .

Lemma 16. *K contains an infinite cyclic subgroup of finite index.*

Proof. Let $D = D(Y)$ denote the constant given by Lemma 4. Pick a point $y \in Y$. Then for each $g \in K$,

$$d(y, g(y)) \leq D.$$

Therefore the K -orbit of y is contained in the metric ball $B(y, D)$. Thus for every $x \in X$, the K -orbit of x is contained in a D -neighborhood of the geodesic $l = \{y\} \times \mathbb{R}$ (passing through x). Therefore K is quasi-isometric to \mathbb{Z} and hence is virtually \mathbb{Z} .

Lemma 17. *The action $G \curvearrowright S^1$ is topologically conjugate to an action of a uniform lattice in $\text{Isom}(\mathbb{H}^2)$.*

Proof. The action $\rho'(G) \curvearrowright \mathbb{H}^2$ is cocompact, therefore we have the following possibilities:

- (a) $\rho'(G)$ is a cocompact discrete subgroup in $\text{Isom}(\mathbb{H}^2)$.
- (b) $\rho'(G)$ is a solvable subgroup in $\text{Isom}(\mathbb{H}^2)$, which fixes a point in S^1 . Then $\rho'(G)$ is not virtually abelian which contradicts the fact that G is a CAT(0)-group.
- (c) $\rho'(G)$ is dense in $\text{PSL}(2, \mathbb{R})$. Then, the group $\rho'(G)$ contains a nontrivial elliptic element \hat{g} and it also contains a sequence of elements \hat{h}_i which converge to $1 \in \text{PSL}(2, \mathbb{R})$. Let $g, h_i \in G$ be elements which map (via ρ') to \hat{g} and \hat{h}_i respectively. Clearly, $\rho(g) \in \text{Isom}(Y)$ is elliptic as well, let $y \in Y$ be its fixed point. By taking conjugates $g_i := h_i g h_i^{-1}$, we get an infinite collection of distinct elements $\{g_i : i \in \mathbb{N}\}$ of G such that for each $n \in \mathbb{Z}$, $g_i(y \times \mathbb{R})$ is contained in $N_R(y \times \mathbb{R})$ where $R \in \mathbb{R}_+$ is independent of i . We note that since all g_i are pairwise conjugate,

there exists $C < \infty$ such that $d(x, g_i(x)) < C$ for each $x \in y \times \mathbb{R}$ and $i \in \mathbb{N}$. This contradicts discreteness of the action of G on X .

The above two lemmas imply that the kernel of ρ is commensurable to \mathbb{Z} and the quotient $\rho(G)$ is commensurable to the fundamental group of a 2-dimensional hyperbolic surface. Thus, after passing to a finite index subgroup in G we obtain a short exact sequence

$$1 \rightarrow K \rightarrow G \rightarrow Q \rightarrow 1 \quad (18)$$

where Q is the fundamental group of a closed oriented surface.

Lemma 19. *Suppose that for a group H we have a short exact sequence*

$$1 \rightarrow \mathbb{Z}/n\mathbb{Z} \rightarrow H \rightarrow Q \rightarrow 1.$$

Then H contains a finite index surface subgroup.

Proof. Let t denote the generator of $\mathbb{Z}/n\mathbb{Z}$. Let $a_i, b_i, i = 1, \dots, n$, denote the lifts to H of the standard generators of Q . It suffices to consider the case when

$$[a_1, b_1] \dots [a_n, b_n] = t$$

and t belongs to the center of H . Consider the finite Heisenberg group

$$H_n := \langle a, b, t : [a, b] = t, a^n = b^n = t^n = 1, [a, t] = 1, [b, t] = 1 \rangle.$$

Define the homomorphism $\phi: H \rightarrow H_n$ by

$$\phi(a_1) = a, \quad \phi(b_1) = b, \quad \phi(a_i) = \phi(b_i) = 1 \quad \text{for all } i \geq 2.$$

Then the kernel H' of ϕ is a torsion-free subgroup of finite index in H . It follows that the map $H \rightarrow Q$ sends H' injectively to a finite index subgroup in Q . Therefore H' is a surface group.

We now return to the exact sequence (18). As in the above lemma we let $a_i, b_i, i = 1, \dots, n$, denote the lifts to G of the standard generators of Q . Let $H \subset G$ denote the subgroup generated by these elements. If

$$t := [a_1, b_1] \dots [a_n, b_n]$$

is an infinite order element of K then H is isomorphic to the fundamental group of a Seifert manifold (whose base is a surface with the fundamental group Q). It is clear that H has finite index in G .

If t has finite order then, according to Lemma 19, after passing to a finite index subgroup in Q we can assume that $t = 1$. Pick an infinite order element $k \in K$ which belongs to the center of G . Then the subgroups H and $\langle k \rangle$ generate the product

$$\mathbb{Z} \times Q \subset G.$$

Again, clearly, this subgroup has finite index in G . Thus, in the both cases, G is commensurable to the fundamental group of a 3-dimensional Seifert manifold. \square

Thus, the conclusion of Theorem 2 holds provided that X contains a parallel set with the full boundary.

Assumption. From now on we will assume that the ideal boundary of each parallel set of X is a proper subset of $\partial_\infty X$.

5.4. Case 3: The ideal boundary of every parallel set in X is a proper subset of $\partial_\infty X$. In this section we show that the *peripheral circles* of the ideal boundaries of nontrivial parallel sets in X can be used to construct a *small stable nontrivial isometric action* of G on an \mathbb{R} -tree. Then, by Rips theory, G admits a nontrivial splitting as an amalgam with virtually abelian edge groups. This, in turn, implies that the edge groups are virtually \mathbb{Z}^2 .

According to Eberlein's theorem (see [11] in the smooth case and [6, Theorem 9.33] in general), the CAT(0) space X is either a visibility space or it contains a 2-flat F . Since in the former case, G is Gromov-hyperbolic, we assume that X contains a 2-flat F . In particular, X contains *nontrivial parallel sets*.

Lemma 20. *Suppose that $P = Y \times \mathbb{R}$ is a nontrivial parallel set in X . Then $\partial_\infty P$ contains a topological circle S which is geodesic in the Tits metric so that S bounds a disk in $\partial_\infty X \setminus \partial_\infty P$.*

Proof. Let $\xi, \eta \in \partial_\infty P$ be the ideal points of a geodesic $y \times \mathbb{R} \subset Y \times \mathbb{R} = P$. Then the Tits boundary $\partial_{\text{Tits}} P$ is the metric join $S^0 \star_{\partial_{\text{Tits}}} Y$, which is the union of geodesic segments L_μ of length π connecting η and ξ and passing through $\mu \in \partial_{\text{Tits}} Y \subset \partial_{\text{Tits}} X$. Clearly, if $\mu \neq \mu'$ then $S := L_\mu \cup L_{\mu'}$ is a topological circle which is geodesic in the Tits metric.

Let D be a component of $\partial_\infty X \setminus \partial_\infty P$. Then there is a point $\zeta \in \partial D$ which belongs to $L_\mu \setminus \{\xi, \eta\}$ for some $\mu \in \partial_{\text{Tits}} Y$. Clearly, ∂D is not contained in L_μ , therefore there exists a point $\zeta' \in \partial D$ which belongs to $L_{\mu'} \setminus \{\xi, \eta\}$ for some $\mu' \in \partial_{\text{Tits}} Y \setminus \{\mu\}$. The reader will verify that the circle $S = L_\mu \cup L_{\mu'}$ bounds D . \square

We will refer to these circles S as in Lemma 20, as *peripheral circles* of $\partial_\infty P$. A flat in X whose boundary is a peripheral circle will be called a *peripheral flat*.

It follows from the properties of the Tits metric (discussed in Section 2) that if $F, F' \subset X$ are 2-flats then the intersection $\partial_{\text{Tits}} F \cap \partial_{\text{Tits}} F' \subset \partial_{\text{Tits}} X$ is convex and either consists of two antipodal points or is a circular arc in $\partial_{\text{Tits}} F$ of the length $\leq \pi$.

Definition 21. We say that totally-geodesic circles $S, S' \subset Z$ *cross* if S contains points from each component of $Z \setminus S'$ (in the visual topology). Note that *crossing* is a symmetric relation. We will say that the ideal boundaries of two parallel sets P, P' *cross* if at least one circle in $\partial_{\text{Tits}} P$ crosses a circle in $\partial_{\text{Tits}} P'$.

Observe that if S and S' cross, the intersection $S \cap S'$ consists of a pair of antipodal points.




Figure 1

Lemma 22. *Suppose that $P = l \times Y \subset X$ is a parallel set for which $\partial_\infty Y$ consists of at least 3 points (i.e., P is not within finite Hausdorff distance from a flat) and $F \subset X$ is a 2-flat which is not contained in P . Then $\partial_\infty P$ and $S = \partial_\infty F$ do not cross.*

Proof. Suppose to the contrary that $\partial_\infty P$ and $S = \partial_\infty F$ do cross. Recall that $\partial_\infty P$ is the metric join of $\{\eta, -\eta\} = \partial_\infty l$ and $\partial_\infty Y$. If S were to pass through η then, by convexity, S passes through $-\eta$ as well and hence F would be contained in the parallel set P . Therefore, S does not pass through $\partial_\infty l$ and the configuration $\{\partial_\infty P, S\}$ has to look like the one in Figure 1, where x, y, z denote the distances from η to the points of intersection between $\partial_\infty P$ and S . It follows that $x + y = \pi, y + z = \pi, x + z = \pi$ and thus

$$x = y = z = \pi/2.$$

This implies that the circle S is contained in $\partial_\infty Y$, thus Y cannot be Gromov-hyperbolic. This contradicts Lemma 14. \square

We observe that, since $G \curvearrowright X$ is properly discontinuous, the stabilizer of each flat $F \subset X$ in the group G is virtually abelian. We assume that this stabilizer is virtually cyclic (possibly finite) – otherwise G contains \mathbb{Z}^2 .

Suppose that we have three flats $F, F', F'' \subset X$ with pairwise distinct ideal boundaries. We will say that F' separates F from F'' if the following holds:

$$\partial_\infty F \subset \bar{D}, \partial_\infty F'' \subset \bar{D}'',$$

where $D \sqcup D'' = Z \setminus \partial_\infty F'$. We set the ternary relation β by: $\beta(FF'F'')$ if F' separates F from F'' .

We leave it to the reader to verify that with this ternary relation the set \mathcal{P} of all peripheral flats in X satisfies the axioms of a pretree.

Lemma 23. *If U_0 is a horoball in X then $W := \partial_\infty U_0$ does not separate $\partial_\infty X$.*

Proof. Let $\xi \in \partial_\infty X$ and consider the horoballs $U_t = \{b_\xi(x) \leq t\}$, $t \in \mathbb{R}$, where b_ξ is the appropriately normalized Busemann function at ξ . Clearly $\partial_\infty U_t = W$ for each t . Property (2) of the \mathcal{Z} -set compactification applied to the pairs (U_t, W) means that we have natural isomorphisms

$$H_c^i(U_t) \rightarrow \tilde{H}^{i-1}(W). \tag{24}$$

Suppose that $[\zeta] \in H_c^i(U_t)$. Then there exists $s < t$ such that U_s is disjoint from the support set of the cocycle ζ . Therefore $[\zeta]$ maps trivially to $H_c^i(U_s)$ and hence, by naturality of (24), it maps trivially to $\tilde{H}^{i-1}(W)$. We conclude that $\tilde{H}^*(W) = 0$. Therefore, by the Alexander duality on $\partial_\infty X$, the subset $W = \partial_\infty U_0$ cannot separate $\partial_\infty X$. \square

Proposition 25. *Let F, F'' be flats in X . Then the set $S(F, F'')$ of flats F' separating F from F'' is compact with respect to the Gromov–Hausdorff topology.*

Proof. If $\partial_\infty F = \partial_\infty F''$ then for each flat F' separating F and F'' we have: $\partial_\infty F' = \partial_\infty F$. Therefore, $S(F, F'')$ is compact by Corollary 13.

Whence we can assume that $\partial_\infty F' \neq \partial_\infty F''$. Suppose that F_i is a sequence of 2-flats in X which diverge to infinity, i.e.,

$$\lim_i d(o, F_i) = \infty$$

where $o \in X$ is a base-point. Then, as in the proof of Lemma 10, the limit of the distance functions to F_i (normalized at o) subconverge to a Busemann function b_ξ in X . Let U be the horoball $\{x : b_\xi(x) \leq 0\}$.

If, say, $\partial_\infty F \subset \partial_\infty U$ then the flat F is contained in the sublevel set of the Busemann function b_ξ and therefore X would contain a flat half-space \mathbb{R}_+^3 , which contradicts Lemma 12. Thus both complements

$$\partial_\infty F \setminus \partial_\infty U, \quad \partial_\infty F'' \setminus \partial_\infty U$$

are nonempty.

Lemma 26. 1. *In the Hausdorff topology on the set of closed subsets of $X \cup \partial_\infty X$, the sets $F_i \cup \partial_\infty F_i$ subconverge into $\partial_\infty U$.*

2. $\partial_\infty F \cap \partial_\infty F'' \subset \partial_\infty U$.

Proof. 1. Suppose that the assertion is false. Then there exists a sequence of points $x_i \in \partial_\infty F_i$ such that

$$\eta = \lim_i x_i \notin \partial_\infty U.$$

Clearly, $\eta \in \partial_\infty X$. Consider a parametrization $\rho(t)$, $t \in \mathbb{R}_+$ of the geodesic ray $\overline{o\eta}$. Then, since $\eta \notin \partial_\infty U$, there exists $T \geq 0$ such that

$$b_\xi(\rho(t)) \geq 1 \quad \text{for all } t \geq T. \quad (27)$$

The Busemann function b_ξ is the limit of the normalized distance functions

$$d_i(x) = d(x, F_i) - d(o, F_i).$$

Then $d_i(o) = 0$, $d_i(x_i) \leq 0$ for all i and hence, by convexity,

$$d_i(y_i) \leq 0 \quad \text{for all } y_i \in \overline{o x_i}.$$

This, together with the inequality (27), contradicts the assumption that the geodesics $\overline{o x_i}$ converge to the geodesic ray $\overline{o\eta}$.

2. Observe that $\partial_\infty F \cap \partial_\infty F'' \subset \partial_\infty F_i$ for each i . Thus (2) follows from (1).

We continue the proof of Proposition 25. Pick points

$$\eta \in \partial_\infty F \setminus \partial_\infty U, \quad \eta'' \in \partial_\infty F'' \setminus \partial_\infty U.$$

The previous lemma implies that

$$\eta, \eta'' \notin \partial_\infty F \cap \partial_\infty F''$$

and that (since $\partial_\infty U$ does not separate $\partial_\infty X$) for large i the points η, η'' belong to the same connected component of $\partial_\infty X \setminus \partial_\infty F_i$. This contradicts the assumption that F_i is between F, F'' for all i . \square

Now, let us pick a peripheral 2-flat $F_0 \in \mathcal{P}$, consider the set $\{gF_0, g \in G\}$ and its closure \mathcal{F} in the Gromov–Hausdorff topology. The elements of \mathcal{F} are peripheral 2-flats in X and the group G acts naturally on \mathcal{F} . We note that since no flat in \mathcal{F} has cocompact stabilizer, \mathcal{F} contains no isolated points. After passing to a smaller G -invariant subset in \mathcal{F} we may assume that the action $G \curvearrowright \mathcal{F}$ is minimal. The union

$$\tilde{\mathcal{L}} := \cup_{F \in \mathcal{F}} F$$

equipped with the Gromov–Hausdorff topology becomes a locally compact 2-dimensional lamination, the topological action $G \curvearrowright \tilde{\mathcal{L}}$ is properly discontinuous and cocompact. The lamination $\tilde{\mathcal{L}}$ has a continuous G -invariant leafwise flat metric. Therefore, since each leaf of $\tilde{\mathcal{L}}$ is amenable, Plante’s construction (see [17]) implies existence of a transversal G -invariant measure μ on $\tilde{\mathcal{L}}$; minimality of $G \curvearrowright \mathcal{F}$ implies that this measure has full support.

Lemma 28. *Suppose that $F \in \mathcal{F}$, $g_n \in G$ is a sequence such that $\lim_{n \rightarrow \infty} g_n F = F_\infty \in \mathcal{F}$. Then there exist $x_-, x_+ \in \mathcal{F}$ such that for all sufficiently large n , $g_n F \in [x_-, x_+]$ and $F_\infty \in [x_-, x_+]$.*

Proof. Since $\lim_{n \rightarrow \infty} g_n F = F_\infty$, the circles $\partial_{\text{Tits}}(g_n F)$ converge to the circle $\partial_{\text{Tits}} F_\infty$ in the Chabauty topology (we again are using here the visual topology on Z). The circles in the collection

$$\{\partial_{\text{Tits}}(g_n F), \partial_{\text{Tits}} F_\infty, n \in \mathbb{N}\}$$

are all peripheral and hence do not cross each other (by Lemma 22). This implies that for all large n, m either $\partial_{\text{Tits}}(g_n F)$ separates $\partial_{\text{Tits}}(g_m F)$ from $\partial_{\text{Tits}} F_\infty$ or $\partial_{\text{Tits}} F_\infty$ separates $\partial_{\text{Tits}}(g_n F)$ from $\partial_{\text{Tits}}(g_m F)$. \square

The above lemma implies that the natural projection $p: \tilde{\mathcal{L}} \rightarrow \mathcal{F}$ is continuous, where we give \mathcal{F} the order topology, whose basis consists of the open intervals (a, b) . It is also clear that p is a proper map in the sense that for each interval $[a, b]$ the inverse image $p^{-1}([a, b])$ consists of leaves of $\tilde{\mathcal{L}}$ which intersect a certain compact subset in X : If a sequence of flats F_j leaves every compact subset in X then this sequence subconverges to a point in $\partial_\infty X$, but a point cannot separate one circle in $\partial_{\text{Tits}} X$ from another.

The measure μ on the pretree \mathcal{F} has no atoms and (since the measure μ transversal to $\tilde{\mathcal{L}}$ has full support) for each pair of distinct points $x, x' \in \mathcal{F}$, $\mu([x, x']) = 0$ iff the corresponding flats F, F' in X are not separated by any flat in \mathcal{F} . We let T be the quotient of \mathcal{F} by the equivalence relation: Points $x, x' \in \mathcal{F}$ are equivalent iff $\mu([x, x']) = 0$. The G -action, the pretree structure, and the measure μ project to T (we retain the notation μ for the projection of the measure). As it was explained in Section 3, the measure μ yields a metric d on T . Local compactness of $\tilde{\mathcal{L}}$ implies that the restriction of d to each interval in T is a complete metric. It is clear that the group G acts isometrically on T .

Remark 29. The map $\mathcal{F} \rightarrow T$ has at most countable multiplicity. Moreover, all but countably many points in T have a unique preimage in \mathcal{F} .

Lemma 30. 1. T is an uncountable metric tree.

2. Stabilizers of nondegenerate arcs in T are virtually cyclic and the action $G \curvearrowright T$ is stable.

3. G does not have a global fixed point in T .

Proof. 1. Follows from Lemma 8.

2. By our hypothesis, for each point $F \in \mathcal{F}$ its G -stabilizer is virtually cyclic. Since \mathcal{F} is perfect, it is uncountable; hence, by Remark 29, uncountably many points in each nondegenerate arc $[x, y] \subset T$ have a virtually cyclic stabilizer. Thus the

action $G \curvearrowright T$ is *small*. Since G is a CAT(0)-group, each virtually cyclic subgroup of G is contained in a maximal virtually cyclic subgroup. Therefore, if $I_1 \supset I_2 \supset \dots$ is a descending chain of arcs in T , then the sequence of their stabilizers in the group G

$$G_{I_1} \subset G_{I_2} \subset \dots$$

is eventually constant. Thus the action $G \curvearrowright T$ is *stable*.

3. The action $G \curvearrowright \mathcal{F}$ is minimal, hence the action $G \curvearrowright T$ is minimal as well. Since T is not a point it follows that G cannot fix a point in T . \square

Since G acts properly discontinuously and cocompactly on the contractible space X , this group is finitely-presented. Therefore, by Lemma 30, we can apply [3] to conclude that the group G splits as an amalgam with a virtually solvable edge subgroup A . Since G is a CAT(0)-group, the subgroup A is virtually abelian and finitely generated; let $A' \subset A$ be a finite index free abelian subgroup. Since G splits over A , the pair (G, A) has at least two ends, and hence the same is true for the pair (G, A') . Since G is a 3-dimensional Poincaré duality group over \mathcal{R} this implies that A' has rank at least 2. This proves the main theorem. \square

References

- [1] W. Ballmann, *Lectures on spaces of nonpositive curvature*. DMV Sem. 25, Birkhäuser, Basel 1995. [Zbl 0834.53003](#) [MR 1377265](#)
- [2] M. Bestvina, Local homology properties of boundaries of groups. *Michigan Math. J.* **43** (1996), 123–139. [Zbl 0872.57005](#) [MR 1381603](#)
- [3] M. Bestvina and M. Feighn, Stable actions of groups on real trees. *Invent. Math.* **121** (1995), 287–321. [Zbl 0837.20047](#) [MR 1346208](#)
- [4] R. Bieri, *Homological dimension of discrete groups*. Queen Mary College Mathematics Notes, London 1976. [Zbl 0357.20027](#) [MR 0466344](#)
- [5] B. H. Bowditch and J. Crisp, Archimedean actions on median pretrees. *Math. Proc. Cambridge Philos. Soc.* **130** (2001), 383–400. [Zbl 1034.20022](#) [MR 1816800](#)
- [6] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*. Grundlehren Math. Wiss. 319, Springer-Verlag, Berlin 1999. [Zbl 0988.53001](#) [MR 1744486](#)
- [7] K. Brown, *Cohomology of groups*. Grad. Texts in Math. 87, Springer-Verlag, New York 1982. [Zbl 0584.20036](#) [MR 0672956](#)
- [8] S. Buyalo, Euclidean planes in three-dimensional manifolds of nonpositive curvature. *Mat. Zametki* **43** (1988), 103–114; *Math. Notes* (1988) **43**, 60–66. [Zbl 0644.53035](#) [MR 0932905](#)
- [9] H. Cartan and S. Eilenberg, *Homological algebra*. Princeton University Press, Princeton 1999. [Zbl 0933.18001](#) [MR 1731415](#)
- [10] A. Casson and D. Jungreis, Convergence groups and Seifert fibered 3-manifolds. *Invent. Math.* **118** (1994), 441–456. [Zbl 0840.57005](#) [MR 1296353](#)

- [11] P. Eberlein, Geodesic flow on certain manifolds without conjugate points. *Trans. Amer. Math. Soc.* **167** (1972), 151–170. [Zbl 0209.53304](#) [MR 0295387](#)
- [12] D. Gabai, Convergence groups are Fuchsian groups. *Ann. of Math. (2)* **136** (1992), 447–510. [Zbl 0785.57004](#) [MR 1189862](#)
- [13] A. Hinkkanen, Abelian and nondiscrete convergence groups on the circle. *Trans. Amer. Math. Soc.* **318** (1990), 87–121. [Zbl 0699.30017](#) [MR 1000145](#)
- [14] M. Kapovich and B. Kleiner, Geometry of quasi-planes. Preprint 2004.
- [15] M. Kapovich and B. Leeb, Quasi-isometries preserve the geometric decomposition of Haken manifolds. *Invent. Math.* **128** (1997), 393–416. [Zbl 0866.20033](#) [MR 1440310](#)
- [16] L. Mosher, Geometry of cubulated 3-manifolds. *Topology* **34** (1995), 789–814. [Zbl 0869.57015](#) [MR 1362788](#)
- [17] J. Plante, Foliations with measure preserving holonomy. *Ann. of Math. (2)* **102** (1975), 327–361. [Zbl 0314.57018](#) [MR 0391125](#)
- [18] V. Schroeder, Codimension one tori in manifolds of nonpositive curvature. *Geom. Dedicata* **33** (1990), 251–265. [Zbl 0698.53026](#) [MR 1050413](#)
- [19] P. Tukia, Homeomorphic conjugates of Fuchsian groups. *J. Reine Angew. Math.* **391** (1988), 1–54. [Zbl 0644.30027](#) [MR 0961162](#)

Received May 11, 2006; revised September 15, 2006

Department of Mathematics, University of California, Davis, CA 95616, U.S.A.

E-mail: kapovich@math.ucdavis.edu

Department of Mathematics, Yale University, New Haven, CT 06520-8283, U.S.A.

E-mail: bruce.kleiner@yale.edu