

Constructing pseudo-Anosovs from expanding interval maps

Ethan Farber

Abstract. We investigate a phenomenon observed by Thurston wherein one constructs a pseudo-Anosov homeomorphism on the limit set of a certain lift of a piecewise linear expanding interval map. We reconcile this construction with a special subclass of generalized pseudo-Anosovs, first defined by de Carvalho. From there we classify the circumstances under which this construction produces a pseudo-Anosov. As an application, we produce for each $g \geq 1$, a pseudo-Anosov ϕ_g on the closed surface of genus g that preserves an algebraically primitive translation structure and whose dilatation is a Salem number.

Contents

1. Introduction	1
2. Review of generalized pseudo-Anosovs	10
3. First constructions	19
4. Reconciling two constructions	27
5. The postcritical orbit of a zig-zag of pseudo-Anosov type	37
6. Classifying zig-zags of pseudo-Anosov type	42
7. The digit polynomial D_f	49
8. A family of pseudo-Anosovs with Salem dilatation	53
References	60

1. Introduction

In [14], Thurston investigates the properties of *uniform λ -expanders*, piecewise linear maps $f: I \rightarrow I$ of a compact interval with $|f'| = \lambda > 1$ wherever defined. Here λ is sometimes referred to as the *growth rate* of f . These expanders arise naturally as linear models for interval endomorphisms of positive entropy (see, e.g., [7]). In many ways, uniform λ -expanders may be seen as one-dimensional analogs of pseudo-Anosov homeomorphisms of a surface.

Recall that a *pseudo-Anosov* is a homeomorphism $\phi: S \rightarrow S$ of surface S such that there exist two singular transverse measured foliations $(\mathcal{F}^u, \mu^u)$, $(\mathcal{F}^s, \mu^s)$ and a constant $\lambda > 1$ such that $\phi_*(\mathcal{F}^u, \mu^u) = (\mathcal{F}^u, \lambda\mu^u)$ and $\phi_*(\mathcal{F}^s, \mu^s) = (\mathcal{F}^s, \lambda^{-1}\mu^s)$. The constant λ is called the *dilatation* of ϕ , and determines the topological entropy of ϕ , namely $h(\phi) = \log \lambda$. For a uniform λ -expander f , we also have $h(f) = \log \lambda$.

In [14], Thurston considers another parallel: it is shown by Fried [5] that the dilatation of a pseudo-Anosov is a *bi-Perron unit*, i.e., that λ is a real, positive algebraic unit whose Galois conjugates lie in the annulus $A_\lambda = \{z: \lambda^{-1} < |z| < \lambda\}$, with the exception of λ and possibly λ^{-1} . It is conjectured that this condition is also sufficient for λ to be the dilatation of some pseudo-Anosov on some surface. On the other hand, Thurston argues in [14] that λ is the growth rate of a postcritically finite (PCF, cf. Definition 3.3) uniform expander if and only if λ is a *weak Perron number*; that is to say, all Galois conjugates are contained in the disc $D_\lambda = \{z: |z| \leq \lambda\}$.

Given these and other dynamical similarities between uniform expanders and pseudo-Anosovs, it is reasonable to ask if one can lift a uniform expander to a pseudo-Anosov on some surface, or conversely if one can find a projection from a pseudo-Anosov to some uniform expander. Thurston gives a suggestive example near the end of [14]. Thurston's example concerns the tent map for $\lambda = (1 + \sqrt{5})/2$ defined by

$$f(x) = \begin{cases} \lambda x, & 0 \leq x < \lambda^{-1}, \\ 2 - \lambda x, & \lambda^{-1} \leq x \leq 1. \end{cases} \quad (1.1)$$

The sole Galois conjugate of λ is $\psi = (1 - \sqrt{5})/2 = -\lambda^{-1}$, and Thurston defines the map $f_G: I \times \mathbb{R} \rightarrow I \times \mathbb{R}$ by

$$f_G(x, y) = \begin{cases} (\lambda x, \psi y), & 0 \leq x < \lambda^{-1}, \\ (2 - \lambda x, 2 - \psi y), & \lambda^{-1} \leq x \leq 1. \end{cases}$$

Note that if $\pi: I \times \mathbb{R} \rightarrow I$ is the projection onto the first coordinate, then $\pi \circ f_G = f \circ \pi$. The *limit set* Λ_f of f_G is defined to be the smallest closed set containing all accumulation points of orbits under f_G . In this example, Λ_f is a finite union of rectangles in $I \times \mathbb{R}$, and while f_G is discontinuous across the line $x = \lambda^{-1}$, one may find gluings of the edges of Λ_f such that the induced map ϕ is a homeomorphism of a closed surface $\tilde{\Lambda}$. The definition of f_G implies that the foliations of Λ_f by horizontal and vertical lines are each preserved under the transformation, with the one-dimensional measure inherited from Lebesgue scaling in the horizontal direction by λ and in the vertical direction by $|\psi| = \lambda^{-1}$. This persists in the quotient $\tilde{\Lambda}$, which is homeomorphic to S^2 . In $\tilde{\Lambda}$ there are four cone points of angle π , each having a single preimage in Λ . These preimages have distinct x -coordinates and so project to four points of dynamical relevance for f : in particular, there is a correspondence between these points of I and the cone points of $\tilde{\Lambda}$, namely

- Three points lying over the unique postcritical orbit of f (shown in red in Figure 1).
- One point lying over the non-zero fixed point of f (shown in blue in the same figure).




Figure 1. Thurston's example. He gives gluings under which the limit set is a sphere with four cone points, corresponding to the postcritical orbit (red) and a fixed point (blue) of f . (a) A uniform λ -expander f for $\lambda = (1 + \sqrt{5})/2$. (b) The limit set of the Galois lift f_G .

Taken together, we see that the Galois lift f_G admits a quotient map ϕ which is an orientation-reversing pseudo-Anosov homeomorphism of the sphere with four marked points. The dilatation of ϕ is λ , and ϕ projects to the λ -expander f . It is not clear from the discussion in [14] whether this process is canonical, or whether it produces pseudo-Anosovs in greater generality. In this paper, we classify when this construction works for interval maps of a particular form, and use it to quickly generate pseudo-Anosovs on surfaces of any genus.

Remark 1.1. The fact that ϕ reverses orientation is because λ and ψ have opposite signs. Throughout this paper, we will restrict our attention to orientation-preserving pseudo-Anosovs, but the methods for understanding the orientation-reversing case are similar.

Remark 1.2. In general, we will see that for a pseudo-Anosov ϕ generated by thickening an interval map f , there is a one-pronged singularity of ϕ for each point in the forward orbit of a critical point for f . See Theorem 3.14 and the discussion in Section 2. These singularities lie on the vertical boundary of Λ , whereas on the horizontal boundary there exists a unique singularity corresponding to some periodic cycle of f . In Thurston's example this periodic cycle is a fixed point, marked in blue, but in general the cycle is longer, generating a singularity of higher rank.

The question of how to formalize this construction was also addressed by Baik, Rafiqi, and Wu in [2]. Their construction concerns a different class of uniform expanders, and together our work and theirs constitute special cases of a larger phenomenon wherein interval maps with certain combinatorics can be realized as train track maps.




Figure 2. The positive (a) and negative (b) zig-zag maps of slope λ equal to the Perron root of $x^2 - 4x - 1$.

Definition 1.3. We say that a PCF map $f: I \rightarrow I$ is *postcritically periodic* (or *PCP*) if each critical value $f(c)$ is periodic. In other words, f acts on the finite set

$$\text{PC}(f) = \bigcup_{c \text{ critical}} \{f^n(c): n \geq 1\}$$

by a permutation. It is possible for multiple critical points c_1, c_2 to have the same critical value.

As we will see (cf. Theorem 3.14), an interval map $f: I \rightarrow I$ being PCP is necessary for producing a pseudo-Anosov by thickening f .

Definition 1.4. We call a uniform λ -expander $f: I \rightarrow I$ a *zig-zag map* (or *zig-zag* or λ -*zig-zag*) if the only critical points of f are $c_i = i \cdot \lambda^{-1}$ for $i = 1, \dots, \lfloor \lambda \rfloor$. In other words, $f(0) \in \partial I$, and $c \in \text{int}(I)$ is a critical point of f if and only if $f(c) \in \partial I$. Note that for a fixed λ , there are two distinct λ -zig-zags: the *positive* zig-zag satisfies $f(0) = 0$, while the *negative* zig-zag satisfies $f(0) = 1$. See Figure 2.

Observe that a zig-zag map f is PCP if and only if $x = 1$ is periodic. Throughout this paper, we consider primarily the case of a PCP zig-zag map. For the sake of generality, however, we give the following two definitions for *any* PCF uniform λ -expander with coefficients in $\mathbb{Q}(\lambda)$, that is, after forming the Markov partition $\mathcal{P}_f = \{P_j\}$ for f by cutting I at the union of the critical points and their forward images (cf. Definition 3.3 and Remark 3.4), there exist polynomials $a_j(\lambda) \in \mathbb{Q}(\lambda)$ such that the linear map defining f on the subinterval P_j is given by $f_j(x) = a_j(\lambda) \pm \lambda x$. This assumption is mild, and allows us to make the following two definitions.

Definition 1.5. Let $f: I \rightarrow I$ be a uniform λ -expander with linear branches $f_j(x) = a_j(\lambda) \pm \lambda x$. Set $\tilde{f}_j(x) = a_j(\lambda^{-1}) \pm \lambda^{-1}x$. The *Galois lift* of f is defined to be the function $f_G: I \times \mathbb{R} \rightarrow I \times \mathbb{R}$ such that

$$f_G(x, y) = (f_j(x), \tilde{f}_j(y)) \quad \text{if } x \in P_j.$$

For concreteness, we define each P_j to be closed on the left, so that f_G is continuous from the right. The limit set of f_G will be denoted by Λ_f .

In this paper, we investigate when Λ_f satisfies the following property.

Definition 1.6. Let $f: I \rightarrow I$ be a PCF uniform λ -expander with canonical Markov partition $\mathcal{P}_f = \{P_j\}$, and let $\pi: \Lambda_f \rightarrow I$ be projection onto the first coordinate. We say that Λ_f is *rectangular* if it has connected interior and each $R_j = \pi^{-1}(P_j)$ is a Euclidean rectangle.

Remark 1.7. Note here that, despite the terminology “Galois lift”, we do not explicitly require λ and λ^{-1} to be Galois conjugate. There is, however, a natural integral polynomial $D_f(t)$ associated to a PCP λ -zig-zag f , called the *digit polynomial* (cf. Definition 1.8 below), and under the condition that $f_G: \Lambda_f \rightarrow \Lambda_f$ defines a pseudo-Anosov this polynomial has both λ and λ^{-1} as roots (cf. Theorem 1). Indeed, in all observed examples λ and λ^{-1} are Galois conjugate.

Definition 1.8. Let f be a PCP λ -zig-zag, and let $n \in \mathbb{N}$ be minimal such that $f^n(1) \in \partial I$. Let $\mathcal{P} = \{P_j\}$ be the Markov partition for f , and let $P_{j(k)}$ be the element of this partition containing $f^k(1)$ for each $k \geq 0$. Let $f_{j(k)}$ denote the linear branch of f defined on $P_{j(k)}$. For each $0 \leq k < n$, let $f_k: \mathbb{C}^2 \rightarrow \mathbb{C}$ be the map of the form $f_k(x, z) = c_k \pm zx$ such that the restriction $f_k|_\lambda: x \mapsto f_k(x, \lambda)$ coincides with the function $f_{j(k)}(x)$. Then the *digit polynomial* of f is the degree n polynomial $D_f: \mathbb{C} \rightarrow \mathbb{C}$ defined by

$$D_f(t) = \varepsilon[f_{n-1}|_{z=t} \circ \cdots \circ f_0|_{z=t}(1) - f^n(1)],$$

where $\varepsilon = \pm 1$ is a normalization factor to make D_f monic. By definition, $D_f(\lambda) = 0$.

Example 1.9. Let $f: I \rightarrow I$ be the tent map from Thurston’s example defined by (1.1). Then $f^3(1) = 1$, and we have $f(1) = \lambda^{-2}$ and $f^2(1) = \lambda^{-1}$. Explicitly,

$$f_0(x, t) = 2 - tx, \quad f_1(x, t) = tx, \quad \text{and} \quad f_2(x, t) = 2 - tx.$$

We compute

$$\begin{aligned} \varepsilon D_f(t) &= f_2(f_1(f_0(1, t), t), t) - f^3(1) \\ &= f_2(f_1(2 - t, t), t) - 1 \\ &= f_2(2t - t^2, t) - 1 \\ &= 2 - 2t^2 + t^3 - 1 \\ &= t^3 - 2t^2 + 1. \end{aligned}$$

Thus in this case $\varepsilon = 1$, and we find that

$$D_f(t) = t^3 - 2t^2 + 1 = (t - 1)(t^2 - t - 1)$$

is the digit polynomial of f . In particular, we verify that $\lambda = (1 + \sqrt{5})/2$ is a root of D_f . We remark that even though $D_f(t) \neq t^3 D_f(t^{-1})$, this does not provide a counterexample

to Theorem 3, since f is not technically of pseudo-Anosov type: it defines an orientation-reversing sphere homeomorphism, rather than an orientation-preserving one.

Example 1.10. Let $f: I \rightarrow I$ be the (unrestricted negative PCP) zig-zag map defined by

$$f(x) = \begin{cases} 1 - \lambda x, & 0 \leq x < \lambda^{-1}, \\ \lambda x - 1, & \lambda^{-1} \leq x < 2\lambda^{-1}, \\ 3 - \lambda x, & 2\lambda^{-1} \leq x < 3\lambda^{-1}, \\ \lambda x - 3, & 3\lambda^{-1} \leq x \leq 1, \end{cases}$$

where $\lambda > 1$ is the largest real root of $t^4 - 3t^3 - 3t^2 - 3t + 1$. In this case,

$$f^4(1) = 0, \quad f_0(x, t) = f_1(x, t) = tx - 3, \quad f_2(x, t) = 3 - tx, \quad \text{and} \quad f_3(x, t) = 1 - tx.$$

We compute as before

$$\begin{aligned} \varepsilon D_f(t) &= f_3(f_2(f_1(f_0(1, t), t), t), t) - f^4(1) \\ &= f_3(f_2(f_1(t - 3), t), t) \\ &= f_3(f_2(t^2 - 3t - 3, t), t) \\ &= f_3(3 + 3t + 3t^2 - t^3, 1) \\ &= 1 - 3t - 3t^2 - 3t^3 + t^4. \end{aligned}$$

Here again $\varepsilon = 1$, and we obtain the polynomial relation $D_f(\lambda) = 0$, where $D_f(t) = t^4 - 3t^3 - 3t^2 - 3t + 1$ coincides with the minimal polynomial of λ .

Remark 1.11. The term “digit” polynomial is chosen to reflect the relation between the coefficients of $D_f(t)$ and the digits of the f -expansion of $x = 1$ (cf. Section 7 and Theorem 3). Since $D_f(t) \in \mathbb{Z}[t]$, the minimal polynomial of λ divides the digit polynomial. The definition of $D_f(t)$ resembles that of the *Parry polynomial* of the λ -expander f (cf. [13]).

Our approach appeals to the machinery of so called *generalized pseudo-Anosovs*, defined in [3] by de Carvalho and further investigated in [4] by de Carvalho and Hall. In brief terms, a *generalized pseudo-Anosov* is a pseudo-Anosov except that we allow for infinitely many singularities of the measured foliations, as long as these only accumulate on finitely many points (cf. Definition 2.10).

The first goal of this paper is Theorem 1, which shows that Thurston’s construction for PCP zig-zags recovers the generalized pseudo-Anosov of de Carvalho and Hall in the case when the latter is a pseudo-Anosov.

For a given zig-zag f , there is only one thickening F_L of f that can produce a pseudo-Anosov, which we call the *exterior left-veering* thick interval map (cf. Definition 3.24 and Proposition 3.27). We say that f is of *pseudo-Anosov type* if F_L generates a pseudo-Anosov according to the construction of de Carvalho and Hall (cf. Definition 3.13). In this case, there is associated to F_L an *invariant train track* τ_L of a very explicit form (cf. Theorem 3.14).

Theorem 1. *Let $f: I \rightarrow I$ be a PCP λ -zig-zag map with $\lambda > 2$. Then f is of pseudo-Anosov type if and only if the following conditions are satisfied:*

- (1) *The digit polynomial D_f of f has λ^{-1} as a root.*
- (2) *The limit set Λ_f of f_G is rectangular (cf. Definition 1.6).*

In this case, the invariant generalized train track τ_L of F_L is finite, and recovers the action of f_G on Λ_f in the following way: Let S' be the closed topological disc obtained by performing the gluings indicated by the non-loop infinitesimal edges of τ_L . Let $\tilde{f}: S' \rightarrow S'$ be the map induced by F_L . Then there is an isometry $i: S' \rightarrow \Lambda_f$ such that the following diagram commutes:

$$\begin{array}{ccc} S' & \xrightarrow{\tilde{f}} & S' \\ \downarrow i & & \downarrow i \\ \Lambda_f & \xrightarrow{f_G} & \Lambda_f. \end{array}$$

Moreover, i sends the horizontal and vertical foliations of S' to those of Λ_f . Therefore, after identifying segments of boundary in each set so as to obtain pseudo-Anosovs

$$\phi_1: S \rightarrow S \quad \text{and} \quad \phi_2: \tilde{\Lambda}_f \rightarrow \tilde{\Lambda}_f,$$

these systems are conjugate via an isometry that sends the (un)stable foliation of ϕ_1 to the (un)stable foliation of ϕ_2 .

Remark 1.12. The content of Theorem 1 is to reveal that, for PCP zig-zags, Thurston's construction of a pseudo-Anosov from a zig-zag is essentially the construction of a generalized pseudo-Anosov following de Carvalho and Hall. Indeed, Thurston's construction is more direct: one need only examine Λ_f to determine if f is of pseudo-Anosov type, and if so, then the action of f_G on Λ_f recovers the pseudo-Anosov action given by applying the methods of de Carvalho and Hall. The proof of Theorem 1 is given in Section 4.

Remark 1.13. Theorem 3 below strengthens condition (1) of Theorem 1: rather than merely vanishing at λ^{-1} , D_f is in fact reciprocal, meaning that for any $\alpha \in \mathbb{C}^*$, $D_f(\alpha) = 0$ if and only if $D_f(\alpha^{-1}) = 0$.

The remainder of the paper classifies for each modality m the PCP zig-zags of pseudo-Anosov type, i.e., the maps whose Galois lift produces a pseudo-Anosov. These results, in particular Theorems 2 and 3, are extensions of the work of Hall, who in [6] classifies unimodal maps (i.e., tent maps) of pseudo-Anosov type, although not in these terms. In particular, Hall shows that there is an explicit bijection between the tent maps of pseudo-Anosov type and $\mathbb{Q} \cap (0, 1/2)$, defined dynamically by the action of the tent map on its postcritical set. Theorem 2 generalizes this to multimodal PCP zig-zags for each modality $m \geq 2$.

The key observation is that the outward winding of the exterior left-veering thickening F_L places strong restrictions on how f permutes the periodic orbit of the point $x = 1$ (cf. Section 5). We prove that these restrictions are also sufficient to determine a PCP zig-

zag of pseudo-Anosov type. The analysis is carried out in three separate cases, according to whether the modality m of f satisfies

- (a) $m \geq 4$ even,
- (b) $m \geq 3$ odd,
- (c) $m = 2$.

Definition 1.14. For $m \geq 2$ and $p \geq 3$, define $\text{PA}(m, p)$ to be the set of zig-zags f of pseudo-Anosov type such that

- (1) f has m critical points,
- (2) $\# \text{PC}(f) = p$.

We also define the set

$$\text{PA}(m) = \bigcup_{p \geq 4} \text{PA}(m, p).$$

As it turns out, the elements of $\text{PA}(m, 3)$ are relatively uninteresting: the growth rates of these maps are precisely the positive quadratic units, and all of the resulting pseudo-Anosovs are defined on the four-punctured sphere (cf. Corollary 5.6). In particular, we recover hyperbolic automorphisms of the torus for each trace $t \geq 3$. The case when $p \geq 4$ is more interesting.

Definition 1.15. Let f be a PCP interval map, and let $x_1 < x_2 < \dots < x_n = 1$ denote the elements of the forward orbit of $x = 1$. The *permutation type* of f is the permutation $\rho(f) \in S_n$ such that $f(x_i) = x_{\rho(f)(i)}$.

In Section 6, we show that for $m \geq 2$, if $f \in \text{PA}(m)$, then $\rho(f)$ has a particular form. Indeed, there exist integers k, n such that $2 \leq k \leq n - 1$ and $\gcd(n - 1, n - k) = 1$ so that we have

$$\rho(f) = \begin{cases} \rho_e(n, k) & \text{if } m \geq 4 \text{ even,} \\ \rho_o(n, k) & \text{if } m \geq 3 \text{ odd,} \\ \rho_2(n, k) & \text{if } m = 2. \end{cases}$$

The permutations ρ_e, ρ_o , and ρ_2 are defined in Section 6. We will write $\rho(f) = \rho_m(n, k)$ when m is not specified. A consequence of the following theorem is that for each $m \geq 2$, there is a bijection between the permutations $\rho_m(n, k)$ and the elements of $\text{PA}(m)$.

Theorem 2. Fix $m \geq 2$ and let $\Phi: \text{PA}(m) \rightarrow \mathbb{Q} \cap (0, 1)$ be the map defined by

$$\Phi(f) = \frac{n - k}{n - 1} \quad \text{if } \rho(f) = \rho_m(n, k).$$

Then Φ is a bijection. Moreover, for each $p \geq 4$, the image $\Phi(\text{PA}(m, p))$ consists of the set of reduced rationals in $(0, 1)$ of denominator $p - 2$.

Theorem 2 classifies PCP zig-zags of pseudo-Anosov type of modality $m \geq 2$. As we mentioned, the proof in the case of unimodal maps of pseudo-Anosov type is due to

Hall [6]; our results should be interpreted as an extension of his to higher modalities. Sections 5 and 6 are dedicated to proving Theorem 2, after which we pass to a discussion of their kneading theoretic implications in Section 7.

We also have the following conjecture.

Conjecture 1.16. *Let $f_1, f_2 \in \text{PA}(m)$ for $m \geq 2$. If λ_i is the growth rate of f_i and $\Phi(f_i) = q_i$ as in Theorem 2, then $\lambda_1 < \lambda_2 \Leftrightarrow q_1 < q_2$.*

We treat this conjecture in a forthcoming paper.

Remark 1.17. There is a certain symmetry in the choice of rational number representing a zig-zag of pseudo-Anosov type. Given such a zig-zag f , one may choose to define $\Phi(f)$ to be either $(n - k)/(n - 1)$ (as in Theorem 2) or else $(k - 1)/(n - 1)$. One may view this symmetry as a manifestation of the fact that $\rho(f) = \rho_m(n, k)$ is essentially a rotation by $n - k$ modulo $n - 1$, which may equivalently be seen as a rotation by $-(k - 1)$ modulo $n - 1$. Making a choice determines whether Conjecture 1.16 posits that the association $\lambda_i \mapsto q_i$ preserves or reverses linear order. Here we have defined $\Phi(f) = (n - k)/(n - 1)$ so that this map is order-preserving. This choice also makes the statement of Theorem 3 more natural.

Theorem 3. *Suppose $f \in \text{PA}(m)$ for $m \geq 2$ with $\Phi(f) = a/b \in \mathbb{Q} \cap (0, 1)$ in lowest terms. Define $L: [0, b] \rightarrow \mathbb{R}$ by $L(t) = (a/b) \cdot t$. Then*

$$D_f(t) = t^{b+1} + 1 - \sum_{i=1}^b c_i t^{b+1-i},$$

where the c_i satisfy

$$c_i = \begin{cases} m & \text{if } L(t) \in \mathbb{N} \text{ for some } t \in [i - 1, i], \\ m - 2 & \text{otherwise.} \end{cases}$$

In particular, $c_i = c_{b-i}$, so D_f is reciprocal, that is, $D_f(t) = t^{b+1} D_f(t^{-1})$.

Theorem 3 generalizes [6, Lemmas 2.5 and 2.6].

In Section 8, we consider an infinite family of zig-zags producing an algebraically primitive translation surface for each positive genus.

Theorem 4. *For each $g \geq 1$, define $f_g: I \rightarrow I$ to be the bimodal PCP zig-zag map of pseudo-Anosov type corresponding to $q_g = 1/(2g) \in \mathbb{Q} \cap (0, 1)$. Let λ_g be the growth rate of f_g . Then the following are true for each $g \geq 1$:*

- (1) λ_g is a Salem number of degree $2g$.
- (2) The pseudo-Anosov ϕ_g obtained from f_g is defined on a $(2g + 2)$ -punctured sphere $\Sigma_{0,2g+2}$.
- (3) The translation surface (X_g, ω_g) obtained as the hyperelliptic double cover of $\Sigma_{0,2g+2}$ is of genus g , and hence algebraically primitive.

To summarize: in Section 2, we review the construction of generalized pseudo-Anosovs, focusing particularly on the concept of a thick interval map. This is followed in Section 3 by a discussion on the different ways of thickening an interval map to a thick interval map. Here we prove that, when attempting to construct pseudo-Anosovs, the only thickening of a zig-zag we need consider is the exterior left-veering map F_L (cf. Proposition 3.27).

In Section 4, we prove Theorem 1, reconciling Thurston's construction with that of de Carvalho. Beginning in Section 5, we turn our attention to classifying zig-zag maps of pseudo-Anosov type, observing several necessary conditions on their postcritical orbit structure. Section 6 in effect demonstrates that these conditions are also sufficient, proving Theorem 2.

Section 7 investigates the digit polynomial $D_f(t)$, establishing Theorem 3.

Section 8 turns briefly to considerations of flat geometry, providing a terse introduction to part of the theory. As an application, we prove Theorem 4, demonstrating the existence of an infinite family of algebraically primitive translation surfaces arising from our construction, one for each genus $g \geq 1$.

2. Review of generalized pseudo-Anosovs

The purpose of this section is to review the theory of generalized pseudo-Anosovs, following the work of de Carvalho in [3] and de Carvalho and Hall in [4]. A reader who is already familiar with the theory may skip this section.

2.1. Thick intervals

In this subsection, we introduce thick interval maps and the procedure of *thickening* an interval map $f: I \rightarrow I$ to a thick interval map F .

A *thick interval* is a closed topological 2-disc $\mathbb{I} \subseteq S^2$ consisting of *decomposition elements* which come in two types: a *leaf*, homeomorphic to the interval $I = [0, 1]$, and a *junction*, homeomorphic to the closed 2-disc. The intersection of the boundary of a junction with $\mathbb{I}$ may consist of one or two connected components. We allow only finitely many junctions in a thick interval.

We denote by $\mathbb{V}$ the union of the junctions of $\mathbb{I}$, and we refer to the connected components of $\mathbb{I} \setminus \mathbb{V}$ as *strips*. Each strip is homeomorphic to $(0, 1) \times [0, 1]$, and is a union of leaves. We put coordinates $h_s: \bar{s} \rightarrow [0, 1] \times [0, 1]$ on the closure of each strip s such that the leaves of s are precisely the sets $h_s^{-1}(\{x\} \times [0, 1])$, $x \in (0, 1)$. Following [4], we denote by $\mathbb{E}$ the union of the closures of the strips. See Figure 3.

The notation $\mathbb{V}$, $\mathbb{E}$ is purposefully suggestive. As we shall see, a thick interval map is meant as a dynamical blow-up of an unrestricted interval map $f: I \rightarrow I$. The junctions correspond to elements of the weak postcritical set $\text{WPC}(f)$ (cf. Section 3), whereas the strips correspond to the subintervals between these points. It is for this reason that we sometimes refer to junctions as *fat vertices* and the strips as *thick edges*.




Figure 3. A typical thick interval consists of alternating thick edges and two-sided junctions, book-ended by a pair of one-sided junctions.




Figure 4. A thick interval map. The images of the junctions have been shaded darker for clarity. The first junction is mapped into itself, the second into the fifth, the third into the second, the fourth into the first, and the fifth into the third.

We refer to [4] for the following definition. We write $F: (X, A) \rightarrow (X, A)$ to represent a map $F: X \rightarrow X$ of topological spaces such that $F(A) \subseteq A$ for some $A \subseteq X$.

Definition 2.1 (See Figure 4 for an example). A *thick interval map* is an orientation-preserving homeomorphism $F: (S^2, \mathbb{I}) \rightarrow (S^2, \mathbb{I})$ such that

- (1) $F(\mathbb{I})$ is contained in the interior of $\mathbb{I}$,
- (2) if γ is a leaf of $\mathbb{I}$, then $F(\gamma)$ is contained in a decomposition element, and the diameter of $F^n(\gamma)$ with respect to the coordinates h_s tends to 0 as $n \rightarrow \infty$,
- (3) if J is a junction of $\mathbb{I}$, then $F(J)$ is contained in a junction,
- (4) F is linear with respect to the coordinates h_s , that is, in each connected component of $s_i \cap F^{-1}(s_j)$, where s_i and s_j are strips, F contracts vertical coordinates uniformly by a factor $\mu_j < 1$ and expands horizontal coordinates uniformly by a factor $\lambda_j > 1$,
- (5) if J and J' are junctions such that $F(J) \subseteq J'$, then $F(\partial J \setminus \partial \mathbb{I}) \subseteq (\partial J' \setminus \partial \mathbb{I})$,
- (6) if J is a junction with $F^n(J) \subseteq J$ for some $n \geq 1$, then J has an attracting periodic point of period n in its interior whose basin contains the interior of J .

Definition 2.2. We associate to a thick interval map F a *transition matrix* $M = (m_{i,j})$ such that if $s_1, \dots, s_n$ are the strips of $\mathbb{I}$, then $m_{i,j}$ is the number of times $F(s_j)$ crosses s_i . Note that since strips are separated by junctions and since junctions are mapped by F to other junctions, $m_{i,j}$ are integers; there are no partial crossings.

Definition 2.3. A non-negative matrix M is said to be *primitive* if there exists some $m \in \mathbb{N}$ such that M^m is positive, i.e., $(M^m)_{i,j}$ is positive for each i, j .

Definition 2.4. Let $F: (S^2, \mathbb{I}) \rightarrow (S^2, \mathbb{I})$ be a thick interval map. Collapse each decomposition element of $\mathbb{I}$ to a point, obtaining an interval $\tilde{I}$ and an induced map $\tilde{f}: \tilde{I} \rightarrow \tilde{I}$. If $x \in \tilde{I}$ corresponds to a leaf that F maps into a junction, then $\tilde{f}$ will be constant in a neighborhood of x . Further collapsing these intervals of constancy produces either a single point or a new interval I with an induced map $f: I \rightarrow I$. We say that f is a *thinning* of F . Similarly, we say that F is a *thickening* of f , and *thickening* f refers to the process of associating to f a thick interval map F that thins to f .

Proposition 2.5 ([3, Theorem 2]). *If F has primitive transition matrix M , then thinning F always produces a non-trivial interval I . Moreover, the induced map $f: I \rightarrow I$ also has transition matrix M .*

Remark 2.6. Observe that while there is a unique thinning of a thick interval map, there are multiple thickenings of an interval map with at least one critical point. In Section 3, we investigate the proper thickening to choose for a zig-zag map if one hopes to produce a pseudo-Anosov.

2.2. Train tracks and generalized pseudo-Anosovs

In this subsection, we describe how to associate to a given thick interval map F a branched 1-manifold τ invariant under F , up to isotopy. This τ is called a *generalized invariant train track* (cf. Definition 2.7), and provides the blueprints for constructing a *generalized pseudo-Anosov* $\phi: S^2 \rightarrow S^2$ (cf. Definition 2.10). In particular, τ dictates the structure of the singular invariant foliations of ϕ .

To a thick interval $\mathbb{I}$ we associate the data of a finite set A of points, called *punctures*. Each puncture is contained in a junction, and each junction contains at most one puncture. For a strip s , we define the arc γ_s to be the path $\gamma_s(t) = h_s^{-1}(t, 1/2)$. Let RE denote the set of such paths. The endpoints of each arc γ_s are on the boundary components of s and are called *switches*. We denote by L the set of switches.

We again take the following definition from [4].

Definition 2.7. Given a thick interval $\mathbb{I} \subseteq S^2$ with a set of punctures A , a *generalized train track* $\tau \subseteq \mathbb{I} \setminus A$ is a graph with vertex set L and countably many edges, each of which intersects $\partial\mathbb{V}$ only at L , such that

- (1) The edges of τ which intersect the interior of $\mathbb{E}$ are precisely the elements of RE .
- (2) No two edges e_1, e_2 contained in a given junction J are *parallel*, that is, e_1 and e_2 may only bound a disc if it contains a point of A or another edge of τ .

Two generalized train tracks τ and τ' are *equivalent*, denoted $\tau \sim \tau'$, if they are isotopic by an isotopy supported on $\mathbb{V} \setminus A$.

Condition (1) says that the edges of τ contained in the strips of $\mathbb{I}$ are simple to describe: they are elements γ_s of RE, which are called *real edges*. The more complicated edges are those contained in the junctions, which are called *infinitesimal edges*. The collection of infinitesimal edges will be denoted by IE.

The infinitesimal edges will provide extra information not already given by the (finite) incidence matrix M for some thick interval map F . Indeed, we associate to F a specific generalized train track as follows. Let τ_0 denote the (disconnected) generalized train track given by the real edges $\gamma_s \in \text{RE}$. We apply F to τ_0 and then perform the following series of pseudo-isotopies:

- (1) On each strip s , we define the pseudo-isotopy $\psi_s: \bar{s} \times [0, 1] \rightarrow \bar{s}$ by

$$\psi_s(x, y, t) = \left(x, (1-t)y + \frac{t}{2}\right).$$

- (2) Within each junction we define another pseudo-isotopy ψ_{e_1, e_2} for each pair of infinitesimal edges that are parallel, which homotopes e_1 and e_2 together.

The effect of the first set of pseudo-isotopies is to collapse all components of $F(\tau_0)$ contained within $\mathbb{E}$ to the real edges γ_s , while the second set homotopes parallel infinitesimal edges and is only supported on a disc containing the relevant junction. Composing these pseudo-isotopies produces a new generalized train track, denoted $\tau'_1 := F_*(\tau_0)$. One may check that τ'_1 is isotopic, relative to A , to a generalized train track τ_1 containing τ_0 . Continuing in this way, we obtain an increasing sequence $\tau_0 \subseteq \tau_1 \subseteq \dots$ of train tracks, and the union $\tau = \bigcup_{n \geq 0} \tau_n$ is F -invariant, i.e., $F_*(\tau)$ is isotopic to τ . See Figure 5.

Definition 2.8. Let $\psi: S^2 \times I \rightarrow S^2$ denote the composition of the pseudo-isotopies in steps (1) and (2) above. The generalized train track $\tau = \bigcup_{n \geq 0} \tau_n$ is called the *invariant generalized train track* for F . The *train track map* associated to F is the map $\phi: \tau \rightarrow \tau$ defined by $\phi(x) = \psi(F(x), 1)$.

Let $\mathbb{I}$ be a thick interval and $F: (S^2, \mathbb{I}, A) \rightarrow (S^2, \mathbb{I}, A)$ be a thick interval map with A , a finite invariant set of F . Let τ be the associated invariant generalized train track. Then τ has at most countably many edges. Label the finitely many real edges $e_1, \dots, e_n$ and then label the possibly infinitely many infinitesimal edges e_k for $k \geq n+1$. We form an *extended transition matrix* $N = (n_{i,j})$ by setting

$$n_{i,j} = \text{the number of times } \phi(e_j) \text{ crosses } e_i.$$

We may write this as the block matrix

$$N = \begin{pmatrix} M & 0 \\ B & \Pi \end{pmatrix},$$

where M is the incidence matrix of F , B records the transitions from real to infinitesimal edges, and Π records the transitions from infinitesimal edges to other infinitesimal edges.




Figure 5. The process for generating an invariant train track for a given thick interval map. One alternately applies the map to τ_n and the pseudo-isotopies to obtain τ_{n+1} . In this case, the process terminates with τ_4 , which is invariant. Since the invariant generalized train track is finite, the resulting surface homeomorphism will be a pseudo-Anosov (cf. Figure 7). In general, however, this process may continue indefinitely, producing a *generalized* pseudo-Anosov with infinitely many singularities. See Figures 8 and 9 for such an example.

We assume from now on that M is primitive. In this case, the Perron–Frobenius theorem states that the spectral radius $\rho(M) > 0$ of M is in fact a real eigenvalue of M , called the *dominant eigenvalue* of M . The Perron–Frobenius theorem further states that the dominant eigenvalue is simple, and that the associated one-dimensional eigenspace is spanned by a positive eigenvector $u = (u_i)$, while no other eigenspace contains a positive eigenvector. We normalize this eigenvector to have unit L^1 -norm, that is, $\sum_i u_i = 1$.

If $\lambda = \rho(M)$ is the Perron eigenvalue of M , let $x' = (x_1, \dots, x_n)$ denote the canonical positive left λ -eigenvector associated to M such that $\sum_i x_i = 1$, and let $y' = (y_1, \dots, y_n)$ denote the positive right λ -eigenvector of M such that $\sum_i x_i y_i = 1$. One shows that these can be extended to left and right λ -eigenvectors of N , i.e., that there exist possibly infinite vectors x and y such that $xN = \lambda x$ and $Ny = \lambda y$. Moreover, it is not hard to see that $x = (x_1, \dots, x_n, 0, 0, \dots)$.

For $i = 1, \dots, n$, construct rectangles R_i of dimensions $x_i \times y_i$. These are the building blocks of the surface on which the generalized pseudo-Anosov will act. The infinitesimal edges incident to one endpoint of a real edge e_i , along with their weights, encode how to identify segments of the corresponding vertical boundary of R_i . While this process is visually intuitive, a precise explanation is nonetheless elusive in the literature, so we describe it here for completeness.

In what follows, we fix a junction J between two adjacent real edges e_L and e_R . Denote by v_L the endpoint of e_L on ∂J , and similarly define v_R .

Definition 2.9. Let e be an infinitesimal edge contained in J , considered as a smooth parameterized arc $e: [0, 1] \rightarrow J$. Observe that

$$e(\{0, 1\}) \subseteq \{v_L, v_R\}.$$

We define an *end* of e to be an arc of the form

$$\alpha = e\left(\left[0, \frac{1}{2}\right]\right) \quad \text{or} \quad \alpha = e\left(\left[\frac{1}{2}, 1\right]\right).$$

If α is an end of e , then for $0 < \varepsilon < 1/2$, we define the ε -*subend* of α to be

$$\alpha_\varepsilon = \begin{cases} e([0, \varepsilon]), & \alpha = e([0, \frac{1}{2}]), \\ e([1 - \varepsilon, 1]), & \alpha = e([\frac{1}{2}, 1]). \end{cases}$$

If e, f are two infinitesimal edges of J , not necessarily distinct, with ends α, β incident to v_L , we set

$$\alpha \leq_L \beta \quad \text{if } \alpha_\delta \text{ is below } \beta_\delta \text{ for all } \delta > 0 \text{ sufficiently small.}$$

If instead α, β are incident to v_R , we set

$$\alpha \leq_R \beta \quad \text{if } \alpha_\delta \text{ is below } \beta_\delta \text{ for all } \delta > 0 \text{ sufficiently small.}$$

It is not difficult to see that $\leq_L$ is a total order on the set of ends incident to v_L . That is, for any two arcs α, β incident to v_L , we either have $\alpha \leq_L \beta$ or $\beta \leq_L \alpha$. Moreover, if

- $e_{j_1}, e_{j_2}, \dots$ are the infinitesimal edges with one end incident to e_L ,
- $e_{k_1}, e_{k_2}, \dots$ are the infinitesimal edges with two ends incident to e_L , and
- the rectangle R_{e_L} has height $y(L)$,

then the fact that y is a right- λ -eigenvector for N implies that

$$y(L) = \sum_i y_{j_i} + 2 \cdot \sum_l y_{k_l}. \quad (2.1)$$

Equation (2.1) is often referred to as the *switch condition*.

For an end α , denote by $e(\alpha)$ the infinitesimal edge of which α is an end, and for an infinitesimal edge e , let $y(e)$ denote the entry of the eigenvector y for N such that $y(e) = y_i$ if $e = e_i$. Then equation (2.1) and the fact that $\leq_L$ is a total order together imply that there is a unique way to partition the right vertical boundary of R_{e_L} into segments d_α of length $y(e(\alpha))$ such that for two ends α, β incident to e_L ,

$$d_\alpha \text{ is below } d_\beta \Leftrightarrow \alpha \leq_L \beta.$$

The same argument shows how to partition the left vertical boundary of R_{e_R} . It remains to describe how to identify these boundary segments. If e is an infinitesimal edge incident to both e_L and e_R , and α, β are the corresponding ends of e , then $y(e(\alpha)) = y(e(\beta)) = y(e)$ and we identify d_α on $\partial_V R_{e_L}$ with d_β on $\partial_V R_{e_R}$ by an orientation-preserving isometry. If instead both ends α, β of e are incident to e_L , then we identify the segments d_α, d_β on $\partial_V R_{e_L}$ by an orientation-reversing isometry, and similarly if both ends α, β are incident to e_R . See Figure 6 for an example with finitely many infinitesimal edges, and Figure 9 for an example with infinitely many.




Figure 6. The rectangle decomposition of a surface from the invariant train track. The black edges are real edges, while the blue edges are infinitesimal edges.

The train track map ϕ induces an endomorphism $\tilde{\Phi}: \mathcal{R} \rightarrow \mathcal{R}$ which stretches the foliation of $\mathcal{R}$ by horizontal lines by a factor of λ , and it scales the vertical foliation by a factor of λ^{-1} . This map $\tilde{\Phi}$ is a homeomorphism except on the boundary of $\mathcal{R}$. This boundary is a topological circle and contains a periodic orbit of $\tilde{\Phi}$. After identifying adjacent segments of this circle that eventually map to the same segment, we obtain a homeomorphism Φ in




Figure 7. The transformation $\tilde{\Phi}: \mathcal{R} \rightarrow \mathcal{R}$. Identifying points on the horizontal boundary that are eventually mapped to the same point produces a homeomorphism of S^2 . Since there are only finitely many singularities, this is a pseudo-Anosov.

the quotient. This new quotient surface is homeomorphic to S^2 , and the periodic orbit on $\partial\mathcal{R}$ becomes a single point, called the *point at infinity*. The induced map $\Phi: S^2 \rightarrow S^2$ inherits stable and unstable foliations, and is a *generalized pseudo-Anosov*.

Definition 2.10. A *generalized pseudo-Anosov* is a homeomorphism ϕ of a compact surface S such that the following hold:

- (1) There exist a real number $\lambda > 1$, called the *stretch factor* of ϕ , as well as two transverse singular measured foliations $(\mathcal{F}_u, \mu_u)$, $(\mathcal{F}_s, \mu_s)$ of S such that

$$\phi_*(\mathcal{F}_u, \mu_u) = (\mathcal{F}_u, \lambda\mu_u), \quad \phi_*(\mathcal{F}_s, \mu_s) = (\mathcal{F}_s, \lambda^{-1}\mu_s).$$

- (2) The singularities of $\mathcal{F}_u$ and $\mathcal{F}_s$, while potentially infinite in number, accumulate on only finitely many points of S .

Remark 2.11. Observe that in the case that the foliations have only finitely many singularities, ϕ is a pseudo-Anosov.

In what follows, we will investigate closely the construction of generalized pseudo-Anosovs from thick interval maps, focusing particularly on the circumstances under which this process produces a pseudo-Anosov. This occurs if and only if the generalized invariant train track is finite, i.e., has only finitely many edges. In this case, τ is simply a train track on a multiply-punctured sphere.

Example 2.12. It will be instructive to see an example where the invariant generalized train track has infinitely many edges, so that the resulting homeomorphism Φ is not a pseudo-Anosov. Let $\lambda = 1 + \sqrt{2}$ and $f: I \rightarrow I$ the uniform λ -expander defined by

$$f(x) = \begin{cases} \lambda x, & 0 \leq x < \lambda^{-1}, \\ 2 - \lambda x, & \lambda^{-1} \leq x < 2\lambda^{-1}, \\ \lambda x - 2, & 2\lambda^{-1} \leq x \leq 1. \end{cases}$$

We see that f has two critical points, $c_1 = \lambda^{-1}$ and $c_2 = 2\lambda^{-1}$. The first of these is 2-periodic, with $f(c_1) = 1$ and $f(1) = c_1$. On the other hand, $f(c_2) = 0$ is a fixed point. Thus the weak postcritical set of f is $\text{WPC}(f) = \{0, c_1, c_2, 1\}$. We thicken this set to junctions and the intermediate subintervals to thick edges.

Figure 8 shows an example of a thickening F of f . In this case, the generalized invariant train track for F has infinitely many edges, corresponding to infinitely many singularities for the resulting sphere homeomorphism. See Figure 9. It is not hard to show that any orientation-*preserving* thickening of f will fail to produce a pseudo-Anosov. Interestingly enough, there does exist an orientation-reversing map that accomplishes this, and indeed if we define the Galois lift f_G of f by replacing all instances of λ with its conjugate $-\lambda^{-1}$, then the limit set Λ_f is rectangular.




Figure 8. A uniform λ -expander for $\lambda = 1 + \sqrt{2}$ and a thickening of it. Note that in the first junction the tightening pseudo-isotopies will produce non-parallel loops of the generalized invariant train track.




Figure 9. The action of F on its generalized invariant train track, and the induced map on the corresponding surface.

3. First constructions

We turn our attention to the task of constructing a pseudo-Anosov on S^2 from a zig-zag map (cf. Definition 1.4). Our goal in this section is to show that for a zig-zag map, there is a unique pair of thickenings to consider when attempting to construct a pseudo-Anosov, that is, no other possible thickenings can give a pseudo-Anosov (cf. Theorem 3.27). Moreover, these thickenings produce conjugate generalized pseudo-Anosovs, so it suffices to only consider one of them.

Before we proceed to the proof of Theorem 3.27, we must consider how to thicken zig-zag maps, and whether they can be thickened in the first place. For this we will rely on Proposition 2.5.

3.1. Some ergodic theory for zig-zag maps

In this subsection, we show that every zig-zag map is weak-mixing. It follows that the transition matrix for a postcritically finite λ -zig-zag with $\lambda > 2$ is primitive.

Definition 3.1. A *critical point* of a map $f: I \rightarrow I$ is a point $x \in I$, where $f: I \rightarrow I$ is not a local homeomorphism. We will say that f is *multimodal* if it has finitely many critical points. If c is a critical point, then $f(c)$ is a *critical value*. Any point of the form $f^n(c)$ for $n \geq 1$ and c a critical point is a *postcritical point*.

Definition 3.2. The *critical set* $C(f)$ will mean the set of critical points of f , and the union of the postcritical points of f is the *postcritical set*, denoted $PC(f)$. Note in particular that a critical point need not be in the postcritical set, and so we define the *weak postcritical set* $WPC(f)$ to be the union of $C(f)$ and $PC(f)$.

If $\#WPC(f)$ is finite, we say f is *postcritically finite*, or *PCF*.

Throughout this paper, we will assume that all interval maps $f: I \rightarrow I$ are multimodal. In other words, $\#C(f) < \infty$ and f is monotone between each adjacent pair of critical points.

Definition 3.3. A *Markov partition* for a map $f: I \rightarrow I$ is a decomposition $I = I_1 \cup \dots \cup I_l$ into finitely many subintervals such that

- (1) $\text{int}(I_i) \cap \text{int}(I_j) = \emptyset$ if $i \neq j$,
- (2) for each i , $f(\overline{I_i})$ is a union of $\overline{I_{j_k}}$'s,
- (3) for each i , the restriction $f|_{\text{int}(I_i)}$ is injective.

In particular, if f has a Markov partition, then f is postcritically finite.

Remark 3.4. Since we will always assume that $\#C(f) < \infty$, it follows that f has a Markov partition if and only if f is PCF. Moreover, f has a unique minimal Markov partition in terms of inclusion: namely, the partition W obtained by cutting the interval at the points of $WPC(f)$. Unless otherwise specified, this is the Markov partition we will use.

Definition 3.5. Let $f: I \rightarrow I$ be a PCF multimodal map with Markov partition $\mathcal{P} = \{I_1, \dots, I_l\}$. The *transition matrix* of f is the $l \times l$ matrix $M = (m_{ij})$ such that

$$m_{ij} = \begin{cases} 1 & \text{if } f(\overline{I_j}) \supseteq \overline{I_i}, \\ 0 & \text{otherwise.} \end{cases}$$

Recall that a multimodal $f: I \rightarrow I$ is PCP if each critical value $f(c)$ is periodic. Observe that a zig-zag map is PCP if and only if the point $x = 1$ is periodic.

In [6], Hall classifies the λ -zig-zags of pseudo-Anosov type for $1 < \lambda \leq 2$. A unimodal zig-zag map is called a *tent map*. Much of our focus will be on PCP λ -zig-zags for $\lambda > 2$. It is therefore important for the generalized pseudo-Anosov construction that we ensure such maps have primitive transition matrix M . Theorem 3.6 essentially accomplishes this goal, and its proof uses a result by Wilkinson [15].

Theorem 3.6. *If $f: I \rightarrow I$ is a λ -zig-zag for some $\lambda > 2$, then f is weak-mixing with respect to Lebesgue measure.*

Proof. Let $\lfloor \lambda \rfloor = m \geq 2$ be the number of critical points of f and set $P_0 = [0, c_1)$, $P_m = [c_m, 1]$, and $P_i = [c_i, c_{i+1}]$ for $i = 1, \dots, m-1$. Following [15], set

$$\Delta(j_1, \dots, j_n) = P_{j_1} \cap f^{-1}P_{j_2} \cap \dots \cap f^{-(n-1)}P_{j_n}.$$

We say $\Delta(j_1, \dots, j_n)$ is full of rank n if $\mu(f^n(\Delta(j_1, \dots, j_n))) = 1$, where μ is Lebesgue measure; otherwise, $\Delta(j_1, \dots, j_n)$ is said to be non-full.

For the n -tuple $(j_1, \dots, j_n)$, let $\mathcal{I}(j_1, \dots, j_n)$ be the number of non-full intervals of positive measure of the form $\Delta(j_1, \dots, j_n, i)$ for $1 \leq i \leq m+1$. Define

$$\mathcal{I}_n = \sup \mathcal{I}(j_1, \dots, j_n),$$

where the supremum is taken over all n -tuples $(j_1, \dots, j_n)$ such that $\Delta(j_1, \dots, j_n)$ has positive Lebesgue measure. Wilkinson shows in [15] that f is weak-mixing as long as

$$\mathcal{I} = \sup_n \mathcal{I}_n < \lambda.$$

However, note that all P_i are full for $i = 0, \dots, m-1$. The full subsets of $\Delta(j_1, \dots, j_n)$ of rank $n+1$ and positive measure are of the form $\Delta(j_1, \dots, j_n, j)$ for $0 \leq j \leq J$, where J depends on the ordered n -tuple $(j_1, \dots, j_n)$. If $m \geq 2$, the only one of these that can be non-full is $\Delta(j_1, \dots, j_n, J)$, hence $\mathcal{I} \leq 1 < \lambda$. ■

Remark 3.7. If a given zig-zag map f is of pseudo-Anosov type (cf. Definition 3.13 below), then it follows that f is in fact *mixing*.

Definition 3.8. A subset $E \subseteq \mathbb{N}$ is said to have *density 0* if

$$\lim_{n \rightarrow \infty} \frac{\#(E \cap \{1, \dots, n\})}{n} = 0.$$

The union of two sets of density 0 also has density 0. It is well known that a measure-preserving dynamical system $(X, \mathcal{B}, \mu, T)$ is weak-mixing if and only if for every $A, B \in \mathcal{B}$, there exists a set $E = E(A, B) \subseteq \mathbb{N}$ of density 0 such that

$$\lim_{E \nrightarrow \infty} \mu(T^{-n} A \cap B) = \mu(A)\mu(B).$$

Recall that a non-negative matrix M is *primitive* if there is some $k \in \mathbb{N}$ such that every entry of M^k is positive.

Corollary 3.9. *If $f: I \rightarrow I$ is a PCP λ -zig-zag map for some $\lambda > 2$, then the transition matrix of f is primitive.*

Proof. Let I_j, I_k be two subintervals in the Markov partition for f obtained by cutting at the points of $\text{WPC}(f)$. Since f is weak-mixing and both I_j and I_k have positive measure, there exists a subset $E_{jk} \subseteq \mathbb{N}$ of density 0 such that $m(I_j \cap f^{-i}(I_k))$ is positive for large $i \notin E_{jk}$. The set $E = \bigcup_{j,k} E_{jk}$ also has density 0, so there exists some large i so that $m(I_j \cap f^{-i}(I_k)) > 0$ for all j, k . In particular, if M is the transition matrix associated to the Markov partition, then M^i is a positive matrix. ■

3.2. Passing between intervals and thick intervals

Definition 3.10. Recall that from a thick interval map $F: (S^2, \mathbb{I}) \rightarrow (S^2, \mathbb{I})$, we obtain an interval map $f: I \rightarrow I$ by first collapsing all decomposition elements of $\mathbb{I}$ to points, and then further collapsing any subintervals on which the induced map is constant. We call this composition of quotient maps the *thinning projection* and denote it by $\pi: \mathbb{I} \rightarrow I$.

Note that by definition, $\pi \circ F|_{\mathbb{I}} = f \circ \pi$.

Lemma 3.11. *Let $F: (S^2, \mathbb{I}) \rightarrow (S^2, \mathbb{I})$ be a thickening of $f: I \rightarrow I$. Then for each $x \in \text{PC}(f)$, there exists a junction $J_x \subseteq \mathbb{I}$ such that $\pi(J_x) = x$. In particular, since thick intervals have only finitely many junctions, f cannot be thickened unless it is PCF.*

Proof. Let $\pi: \mathbb{I} \rightarrow I$ denote the thinning projection. Let c be a critical point of f , and let $C = \pi^{-1}(c)$ be the set of decomposition elements projecting to c . Similarly, let $C' = \pi^{-1}(f(c))$.

We may assume without loss of generality that c is a local maximum of f . Therefore, there exist points x_1, x_2 satisfying $x_1 < c < x_2$ such that $f(x_1) = f(x_2) < f(c)$. Let

$$X_i = \pi^{-1}(x_i) \quad \text{for } i = 1, 2.$$

Then each $F(X_i)$ lies to the left of C' , whereas $F(C) \subseteq C'$. Since F is linear with respect to the coordinates h_s (cf. Definition 2.1), $F(\mathbb{I})$ must pass through a junction J after $F(X_1)$ and before $F(X_2)$. Picking x_i arbitrarily close to c shows that in fact $F(C) \subseteq J$. But now $J \cap C' \neq \emptyset$, and since C' is the collection of all decomposition elements that project to $f(c)$, it follows that $J \subseteq C'$.

The thickening F maps junctions into junctions, so $F^n(C')$ must be contained in a junction for each $n \geq 1$. But $\pi(F^n(C')) = f^n(f(c))$, so by the same argument as before, $\pi^{-1}(f^n(f(c)))$ contains a junction for each n . Repeating this procedure for all critical points shows that $\pi^{-1}(x)$ contains a junction for each $x \in \text{PC}(f)$. ■

Example 3.12. We revisit Example 2.12, demonstrated in Figure 8. Here $\text{PC}(f)$ consists of the first, second, and fourth heavily drawn points, and the thick interval map F has a junction for each of these. Note that F also has a junction for the third point $c_2 = 2\lambda^{-1}$, despite the fact that c_2 is not a postcritical point. This junction is not strictly necessary, since no junction maps into it, but it does allow us to identify the transition matrices of F and f : each is given by

$$M_W = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 1 & 0 \end{pmatrix}.$$

One checks that $\chi_W(t) = t(t^2 - 2t - 1)$ has dominant root $\lambda = 1 + \sqrt{2}$, the growth rate of f . Eliminating the extraneous junction over c_2 has the effect of combining the second and third subintervals, producing a new transition matrix

$$M_P = \begin{pmatrix} 1 & 2 \\ 1 & 1 \end{pmatrix}$$

with characteristic polynomial $\chi_P(t) = t^2 - 2t - 1$. See Remark 3.16 below.

Definition 3.13. We say that a map $f: I \rightarrow I$ is of *pseudo-Anosov type* if there is a thickening $F: \mathbb{I} \rightarrow \mathbb{I}$ of f whose invariant generalized train track is finite, i.e., is a train track in the classical sense of Thurston.

Theorem 3.14. *Let $f: I \rightarrow I$ be a PCF λ -expander of pseudo-Anosov type, and let F be a thickening of f that has finite invariant generalized train track τ . Then τ has exactly one loop in each fat vertex corresponding to an element of $\text{PC}(f)$. Moreover, these loops determine the one-pronged singularities of the pseudo-Anosov ϕ , and f is in fact PCP.*

Proof. Each loop of τ determines a one-pronged singularity of ϕ . Multiple loops in a single vertex V would imply that ϕ has singularities connected by a leaf of its stable (vertical) foliation. It is well known that such a thing, called a *saddle connection*, is impossible for the invariant foliation of a pseudo-Anosov. Thus each vertex V , corresponding to a point $v \in \text{WPC}(f)$, contains either 0 or 1 loops of τ . We claim that V contains a loop of τ if and only if v is in the subset $\text{PC}(f)$.

To see this, observe that V contains a loop if v is a critical value of f , since in this case $F(\tau)$ makes a turn through V , and after pseudo-isotopy this turn pinches to a loop. By the invariance of τ , it now follows that V contains a loop if v is in the forward orbit of a critical point, which is to say that $v \in \text{PC}(f)$. The only other possibility remaining for v is that it is an element of $\text{WPC}(f) \setminus \text{PC}(f)$, i.e., a critical point that is not in the forward orbit of any critical point. But then no junction of $\mathbb{I}$ maps into V , so no loop of τ

maps into V . Furthermore, $F(\tau)$ does not make a turn through V , since that would imply that v is a critical value, which is it not. These are the only ways a loop of τ will appear in V during the process of constructing τ , so in fact V does not contain a loop of τ if $v \notin \text{PC}(f)$.

To finish the proof, recall that a pseudo-Anosov permutes its 1-prong singularities. Hence f acts on the elements of $\text{PC}(f)$ by a permutation; in other words, f is PCP. ■

Definition 3.15. Let $f: I \rightarrow I$ be a PCP λ -expander of pseudo-Anosov type, and let τ be the invariant train track associated to a thickening F of f . We say an infinitesimal edge of τ is a *connecting edge* if it is not a loop, i.e., if it joins distinct adjacent real edges.

Remark 3.16. In a sense, Theorem 3.14 demonstrates that the important points of f to consider are the elements of $\text{PC}(f)$, rather than those of $\text{WPC}(f)$, since these are the points that correspond to the singularities of any generalized pseudo-Anosov obtained from a thickening of f . One could define a transition matrix according to the partition of I given by $\text{PC}(f)$, although this partition would not technically be Markov, since f might fail to be injective on each subinterval. Since injectivity will be helpful in our arguments, we will resort to using the weak postcritical Markov partition. In any case, there is a straightforward relationship between the two transition matrices and their characteristic polynomials. Specifically, the weakly postcritical transition matrix M_W is primitive if and only if the postcritical transition matrix M_P is as well, and the relationship between the characteristic polynomials is $\chi_W(t) = t^a \chi_P(t)$, where $a = \#(\text{WPC}(f) \setminus \text{PC}(f))$.

Example 3.17. Let $f: I \rightarrow I$ be the positive zig-zag map of growth rate $\lambda = (3 + \sqrt{5})/2$, the dominant root of $x^2 - 3x + 1$. The orbit of $x = 1$ is periodic of period 3 and includes the first critical point $c_1 = \lambda^{-1}$. The other postcritical orbit is the forward orbit of $x = c_2$, which maps to the fixed point at $x = 0$. See Figure 10. Thus

$$C(f) = \{c_1, c_2\}, \quad \text{PC}(f) = \{0, c_1, v, 2\}, \quad \text{WPC}(f) = \{0, c_1, v, c_2, 1\}.$$




Figure 10. The positive zig-zag map for $\lambda = (3 + \sqrt{5})/2$. Compare with Figures 5 and 11.

In particular, the map f is postcritically periodic, acting on $\text{PC}(f)$ by the permutation $(1)(2, 4, 3)$. The postcritical and weak postcritical transition matrices for f are

$$M_P = \begin{pmatrix} 1 & 0 & 2 \\ 1 & 1 & 1 \\ 1 & 1 & 0 \end{pmatrix}, \quad M_W = \begin{pmatrix} 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \end{pmatrix}$$

with characteristic polynomials $\chi_P(t) = (t + 1)(t^2 - 3t + 1)$ and $\chi_W(t) = t(t + 1)(t^2 - 3t + 1)$. The matrices M_P and M_W are each primitive, hence we have

$$M_P^2 = \begin{pmatrix} 3 & 2 & 2 \\ 3 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix}, \quad M_W^2 = \begin{pmatrix} 3 & 2 & 1 & 1 \\ 3 & 2 & 1 & 2 \\ 2 & 1 & 1 & 2 \\ 2 & 1 & 1 & 2 \end{pmatrix}.$$

Example 3.18. Here is an example of a tent map, i.e., a zig-zag of growth rate $1 < \lambda < 2$. It is not hard to show that positive tent maps are not ergodic: indeed, $x = 0$ is a repelling fixed point with only itself as a preimage. Nonetheless, restricting to the subinterval $[f(1), 1]$ produces an ergodic transformation with the same growth rate.

Let $g: I \rightarrow I$ be the positive zig-zag of growth rate $\lambda = (1 + \sqrt{5})/2$, the dominant root of $x^2 - x - 1$. The orbit of $x = 1$ is periodic of period 3 and includes the lone critical point $c = \lambda^{-1}$. See Figure 1 (a). Thus $C(f) = \{c\}$ and $\text{PC}(f) = \text{WPC}(f) = \{0, u, c, 1\}$. The postcritical and weak postcritical transition matrices for g coincide as

$$M = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix}$$

with characteristic polynomial $\chi_M(t) = (t - 1)(t^2 - t - 1)$. Note that this is equal to the digit polynomial $D_f(t)$ of the tent map (cf. Example 1.9). Since the only subinterval that maps to the first is itself, M cannot be primitive. However, the 2-by-2 minor describing the transitions between the second two subintervals is primitive:

$$M' = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}, \quad (M')^2 = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}.$$

3.3. The exterior left-veering thickening

In this subsection, we investigate the possible thickenings of a PCP zig-zag, and show that only two of these have a chance of producing a pseudo-Anosov. These are the *exterior left*- and *exterior right-veering* thickenings F_L and F_R (cf. Definition 3.24). These thickenings “swirl” out from the center, turning either always left or always right, respectively. Since these maps are topologically conjugate, we therefore restrict our analysis to F_L in future sections.

Let f be a zig-zag map of growth rate $\lambda > 2$ and critical points $c_i = i \cdot \lambda^{-1}$ for $i = 1, \dots, \lfloor \lambda \rfloor$. We assume that f is PCP, which is equivalent to the orbit of $x = 1$ being periodic. Let $0 \leq v_1 < v_2 < \dots < v_n = 1$ be the orbit of 1. These, along with $x = 0$ if it is not already among the v_i , are precisely the points to which the junctions of our thick interval $\mathbb{I}$ will project. To specify a particular thick interval map F projecting to f , however, it remains to determine how F folds $\mathbb{I}$, i.e., how the image $F(\mathbb{I})$ turns within $\mathbb{I}$.

If $F: \mathbb{I} \rightarrow \mathbb{I}$ is a thickening of $f: I \rightarrow I$ with thinning projection π , then for brevity we will denote by $\mathbf{0}$ the junction satisfying $\pi(\mathbf{0}) = 0$. Similarly, $\mathbf{1}$ will denote the junction satisfying $\pi(\mathbf{1}) = 1$.

Definition 3.19. Let $f: I \rightarrow I$ be a PCP λ -zig-zag map with $\lfloor \lambda \rfloor = m \geq 2$. Let $F: \mathbb{I} \rightarrow \mathbb{I}$ be a thickening of f . The image $F(\mathbb{I}) \subseteq \mathbb{I}$ is a collection of thick intervals stacked vertically stretching between $\mathbf{0}$ and $\mathbf{1}$ with at most a single exception. Orienting $\mathbb{I}$ from left to right, we number these thick subintervals $\mathbb{I}_j$ as we travel along $F(\mathbb{I})$, beginning at $\mathbf{0}$ if f is a positive zig-zag and at $\mathbf{1}$ if f is a negative zig-zag. We may also assign a *height* to each $\mathbb{I}_j$ by determining its vertical order among the other thick subintervals, counting from bottom to top. The height of $\mathbb{I}_j$ is defined to be $h(\mathbb{I}_j) = i$ if $\mathbb{I}_j$ is the i -th thick subinterval in this ordering. The *type* of the thick interval map F is then the element σ of the permutation group S_{m+1} such that

$$\sigma(j) = i \Leftrightarrow h(\mathbb{I}_j) = i.$$

For an example, see Figure 11.

Example 3.20. Let f be the positive zig-zag map for $\lambda = (3 + \sqrt{5})/2$ from Example 3.17. We have $\lfloor \lambda \rfloor = 2$, and f is postcritically periodic of length 3 with $\text{PC}(f)$ given by

$$0 < c_1 = v_1 < v_2 < c_2 < v_3 = 1.$$

More specifically, the orbit of $x = 1$ is $1 \mapsto v_2 \mapsto v_1 \mapsto 1$. The six possible thickenings of f are pictured in Figure 11. These are paired according to the topological conjugacy class of the resulting generalized pseudo-Anosov. Note that the conjugacy classes in this example are the elements of the orbit space of ι_3 acting on S_3 , where ι_3 is the order-reversing permutation $\tau_3 = (1, 3)(2)$. See Definition 3.22 below.

Example 3.21. Not every permutation $\sigma \in S_{m+1}$ gives a valid thick interval map. For example, let f be a PCP λ -zig-zag map such that $\lfloor \lambda \rfloor = 3$. Then regardless of the orbit of $x = 1$, there can be no thickening of f of type $\sigma = (1)(2, 3)(4)$ since the image of this thick interval map would necessarily intersect itself.

Definition 3.22. The *orientation-reversing permutation* $\iota_n \in S_n$ is the permutation such that $\iota_n(i) = n + 1 - i$ for all $1 \leq i \leq n$.

We will often suppress notation and write $\iota_n = \iota$ when n is understood from context. Note that $\iota^2 = \text{id}$, so if we let ι_n act on S_n on the left then the orbit space is parameterized




Figure 11. The six possible thickenings of the positive zig-zag for $\lambda = (3 + \sqrt{5})/2$. The permutation type of each is an element of S_3 , determined by the heights of the horizontal layers of $F(\mathbb{I})$. For example, the top left thick interval map has permutation type $\sigma_1 = (1, 2, 3)$, and the top right has permutation type $\sigma_2 = (1)(2, 3)$. Each column is an orbit of the action of τ_3 .

by pairs of elements of S_n . In general, ι_n is not central in S_n , so the orbit space is not a group. The following proposition says that the elements of the orbit space correspond to topological conjugacy classes of generalized pseudo-Anosovs.

Proposition 3.23. *Let f be a PCP zig-zag map and F_1, F_2 thickenings of f of types σ_1, σ_2 , respectively. If $\sigma_1 = \iota\sigma_2$, then F_1 is topologically conjugate to F_2 via the orientation-reversing homeomorphism $i: \mathbb{I} \rightarrow \mathbb{I}$ that reflects through the horizontal midline of $\mathbb{I}$. Consequently, if ϕ_i is the generalized pseudo-Anosov induced by F_i , then ϕ_1 is topologically conjugate to ϕ_2 .*

Proof. One immediately checks that if $\sigma_1 = \iota\sigma_2$, then $F_1 = i \circ F_2 \circ i^{-1}$. Following through the details of construction given in Section 2, we see that the invariant generalized train tracks τ_i of F_i satisfy $\tau_1 \sim i_*(\tau_2)$, and so the resulting generalized pseudo-Anosovs are conjugate by the homeomorphism induced by i . ■

Definition 3.24. The *positive exterior left-veering permutation* is the element σ_L^+ of S_n defined as follows: if n is even, then

$$\sigma_L^+(k) = \begin{cases} \frac{n+k}{2} & \text{if } k \text{ is even,} \\ \frac{n+1-k}{2} & \text{if } k \text{ is odd.} \end{cases}$$

If instead n is odd, then we define

$$\sigma_L^+(k) = \begin{cases} \frac{n+1+k}{2} & \text{if } k \text{ is even,} \\ \frac{n-k}{2} + 1 & \text{if } k \text{ is odd.} \end{cases}$$

The *negative exterior left-veering permutation* $\sigma_L^- \in S_n$ is defined by

$$\sigma_L^-(k) = n + 1 - \sigma_L^+(k).$$

The *positive (resp. negative) exterior right-veering permutation* is $\sigma_R^+ = \iota \circ \sigma_L^+$ (resp. $\sigma_R^- = \iota \circ \sigma_L^-$). If F is a thick interval map of type $\sigma_L^\pm$, we also say F is *exterior left-veering*, and write $F = F_L^\pm$. Similarly, we denote by $F_R^\pm$ the *exterior right-veering* thick interval of permutation type $\sigma_R^\pm$.

Example 3.25. For S_3 , σ_L^+ and σ_R^+ are shown in Figure 11 as the top and bottom of the leftmost column, respectively. For S_4 , we have $\sigma_L^+ = (1, 2, 3)(4)$ and $\sigma_R^+ = (1, 3, 4)(2)$.

Remark 3.26. Given a PCP zig-zag f , only one pair of thickenings is defined: either F_L^+ and F_R^+ if f is positive, or F_L^- and F_R^- if f is negative. In this case, we will drop the superscripts from the notation and simply refer to F_L and F_R .

Proposition 3.27. Let $f: I \rightarrow I$ be a PCP zig-zag map of pseudo-Anosov type with growth rate $\lambda > 2$, and let $F: \mathbb{I} \rightarrow \mathbb{I}$ be a thickening of f that induces a pseudo-Anosov. Then $F = F_L$ or $F = F_R$.

Proof. Since F induces a pseudo-Anosov, Theorem 3.14 implies that each of $\mathbf{0}$ and $\mathbf{1}$ contains a single loop of the invariant generalized train track τ . For each of these vertices, the turns of F that pass through them must be concentric, since otherwise a second loop would appear. Orienting $F(\mathbb{I})$ from $F(\mathbf{0})$ to $F(\mathbf{1})$, it follows inductively that each turn has to be in the same direction as the previous one. Thus if the first turn is to the left, all turns are to the left and $F = F_L$. Similarly, if the first turn is to the right then $F = F_R$. ■

It is always possible to construct a thick interval map F having permutation type σ_L or σ_R projecting to a given PCP zig-zag map. Indeed, such a thick interval map swirls out from the center, turning to the left if it has type σ_L and to the right if it has type σ_R . In particular, it follows from Propositions 3.23 and 3.27 that associated to a given PCP zig-zag map $f: I \rightarrow I$ is a canonical thick interval map to consider when investigating whether f is of pseudo-Anosov type, namely F_L .

4. Reconciling two constructions

In this section, we prove Theorem 1, establishing the connection between generalized pseudo-Anosovs and the Galois lift f_G of Thurston (cf. Definition 1.5). We recall the statement of the theorem.

Theorem 1. Let $f: I \rightarrow I$ be a PCP λ -zig-zag map with $\lambda > 2$. Then f is of pseudo-Anosov type if and only if the following conditions are satisfied:

- (1) The digit polynomial D_f of f has λ^{-1} as a root.
- (2) The limit set Λ_f of f_G is rectangular.

In this case, the invariant generalized train track τ_L of F_L is finite, and recovers the action of f_G on Λ_f in the following way: Let S' be the closed topological disc obtained by performing the gluings indicated by the non-loop infinitesimal edges of τ_L . Let $\tilde{f}: S' \rightarrow S'$ be the map induced by F_L . Then there is an isometry $i: S' \rightarrow \Lambda_f$ such that the following diagram commutes:

$$\begin{array}{ccc} S' & \xrightarrow{\tilde{f}} & S' \\ \downarrow i & & \downarrow i \\ \Lambda_f & \xrightarrow{f_G} & \Lambda_f. \end{array}$$

Moreover, i sends the horizontal and vertical foliations of S' to those of Λ_f . Therefore, after identifying segments of boundary in each set so as to obtain pseudo-Anosovs $\phi_1: S \rightarrow S$ and $\phi_2: \tilde{\Lambda}_f \rightarrow \tilde{\Lambda}_f$, these systems are conjugate via an isometry that sends the (un)stable foliation of ϕ_1 to the (un)stable foliation of ϕ_2 .

4.1. The reverse direction

Throughout this subsection, we assume that $f: I \rightarrow I$ is a PCP λ -zig-zag whose Galois lift f_G has rectangular limit set Λ_f and whose digit polynomial D_f satisfies $D_f(\lambda^{-1}) = 0$. We prove that the exterior left-veering thickening F_L of f has finite invariant generalized train track, and that the generalized pseudo-Anosov so obtained will recover the action of f_G on Λ_f .

Lemma 4.1. *Let μ be the finite measure on Λ_f inherited from Lebesgue measure on $\mathbb{R}^2$. Then $f_G: (\Lambda_f, \mu) \rightarrow (\Lambda_f, \mu)$ is measure-preserving.*

Proof. For each i , the affine map $\tilde{f}_i$ defining f_G on the rectangle R_i is measure-preserving, having Jacobian $\pm \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix}$. Indeed, these maps are invertible with measure-preserving inverse, so

$$\mu(\tilde{f}_i(R_i)) = \mu(R_i).$$

Therefore, we have

$$\mu(\Lambda_f) = \sum_{i=0}^k \mu(R_i) = \sum_{i=0}^k \mu(\tilde{f}_i(R_i)) \geq \mu(f_G(\Lambda_f)).$$

If the final inequality is strict, then $A = \Lambda_f \setminus f_G(\Lambda_f)$ has positive measure. But since Λ_f and $f_G(\Lambda_f)$ are each finite unions of rectangles, so is A , hence if $\mu(A) > 0$, then A has non-empty interior U . No point $x \in U \subseteq \Lambda_f$ can be a limit point of an orbit of f_G , contradicting the definition of Λ_f . Hence in fact $\mu(f_G(\Lambda_f)) = \mu(\Lambda_f)$, and in particular

$$\sum_{i=0}^k \mu(\tilde{f}_i(R_i)) = \mu(f_G(\Lambda_f)).$$

In other words, $\mu(\tilde{f}_i(R_i) \cap \tilde{f}_j(R_j)) = 0$ for all indices $i \neq j$, and so for any measurable $B \subseteq \Lambda_f$, we have

$$\mu(f_G^{-1}(B)) = \sum_{i=0}^k \mu(\tilde{f}_i^{-1}(B)) = \sum_{i=0}^k \mu(B \cap R_i) = \mu(B).$$

Thus f_G preserves μ . ■

Let M be the transition matrix for an ergodic PCF uniform λ -expander g . The map g has a unique invariant measure that is absolutely continuous, i.e., that is in the equivalence class of Lebesgue measure. Denote this measure by ν_0 . If M is primitive, then results of Parry in [10] on subshifts of finite type imply that the Perron eigenvectors of M determine ν_0 . Namely, let $u = (u_1, \dots, u_n)$ denote the left λ -eigenvector of M with $\sum_i u_i = 1$, and let $v = (v_1, \dots, v_n)$ denote the right λ -eigenvector of M such that $u \cdot v = 1$. Then the density of ν_0 is (up to a null set) a step function with height v_i on a subinterval of Lebesgue measure u_i . Note that in general, if we use the right λ -eigenvector $v' = cv$ for $c > 0$, then we obtain the invariant measure $\nu'_0 = c \cdot \nu_0$, which is a probability measure exactly when $c = 1$.

Lemma 4.2. *Let $\pi: \Lambda_f \rightarrow I$ denote projection onto the first coordinate, and we define $\nu = \pi_*\mu$. Then $\nu = \nu_0(I) \cdot \nu_0$.*

Proof. Since μ is f_G -invariant and $\pi \circ f_G = f \circ \pi$, ν is invariant with respect to f . Furthermore, because Λ_f is a connected finite union of rectangles of positive measure, we see that ν is equivalent to Lebesgue measure, and in fact has invariant density a step function whose values are given by the heights of the rectangles of Λ_f . The result now follows from uniqueness of ν_0 . Note in general that $\nu(I) = \mu(\Lambda_f) \neq 1$. ■

Lemma 4.3. *Let $\alpha = \alpha_0$ be a periodic point of f of period p . Set $\alpha_i = f^i(\alpha_0)$ for $0 \leq i \leq p-1$, and let $g_i: \mathbb{R} \rightarrow \mathbb{R}$ denote the affine map defining f at $x = \alpha_i$ so that $f(\alpha_i) = g_i(\alpha_i) = c_i \pm \lambda\alpha_i = \alpha_{i+1}$, where the indices are understood modulo p . Define $\tilde{g}_i: \mathbb{R} \rightarrow \mathbb{R}$ by $\tilde{g}_i(y) = \alpha_i \pm \lambda^{-1}y$.*

Then there is a unique periodic point of f_G projecting to α_0 , and it is given by (α_0, y_0) where y_0 is the unique solution to the equation $\tilde{g}_{p-1} \circ \dots \circ \tilde{g}_0(y) = y$. In the case $\alpha_0 = 1$, this periodic point is $(1, 1)$.

See Figure 12 for a few examples.

Proof. Observe first that $\tilde{g}_{p-1} \circ \dots \circ \tilde{g}_0: \mathbb{R} \rightarrow \mathbb{R}$ is a contraction by a factor of $\lambda^{-k} < 1$. This map has a unique fixed point y_0 . By definition,

$$f_G(\alpha_0, y_0) = (g_0(\alpha_0), \tilde{g}_0(y_0)).$$

Indeed, inductively defining $y_i = \tilde{g}_{i-1}(y_{i-1})$, we have $f_G^i(\alpha_0, y_0) = (\alpha_i, y_i)$ for $0 \leq i \leq p-1$. Since α_0 and y_0 are periodic of period p , it now follows that $f_G^p(\alpha_0, y_0) = (\alpha_0, y_0)$.




Figure 12. *Below:* The graph of the negative λ -zig-zag f , for λ the Perron root of $D_f(t) = t^4 - 3t^3 - t^2 - 3t + 1$. *Above:* The limit set Λ_f of the Galois lift of f . In each picture, heavily marked points of the same color belong to the same periodic orbit, and the points of one color in Λ_f are the unique periodic lifts of the points of I of the same color. This limit set was drawn by plotting the orbit under f_G of a single point with transcendental coordinates, to ensure that it is not eventually periodic. The same method was used to draw all limit sets in the paper.

Suppose that (α_0, z) is a periodic point of f_G of period p . Then

$$f_G^i(\alpha_0, z) = (\alpha_i, \tilde{g}_i \circ \cdots \circ \tilde{g}_0(z)) \quad \text{for } 0 \leq i \leq p-1.$$

In particular, z is a fixed point of $\tilde{g}_{k-1} \circ \cdots \circ \tilde{g}_0$, hence is equal to y_0 .

The fact that $(1, 1)$ is the periodic point of f_G projecting to $1 \in I$ follows immediately from the assumption that $D_f(\lambda^{-1}) = 0$. ■

Definition 4.4. Let $R = [a, b] \times [c, d]$. The *horizontal boundary* of R is $\partial_H R = [a, b] \times \{c, d\}$. The *vertical boundary* of R is $\partial_V R = \{a, b\} \times [c, d]$. Two rectangles

$$R_1 = [a_1, b_1] \times [c_1, d_1], \quad R_2 = [a_2, b_2] \times [c_2, d_2]$$

are *lower-aligned* (resp. *upper-aligned*) if $c_1 = c_2$ (resp. $d_1 = d_2$).

Definition 4.5. Let $f: I \rightarrow I$ be a PCP zig-zag whose Galois lift f_G has rectangular limit set Λ_f . Let $R_0, \dots, R_k$ be the rectangles defined by the canonical Markov partition of f which subdivide Λ_f . The *vertical boundary* of Λ_f is the set

$$\partial_V \Lambda_f = \left(\partial \Lambda_f \cap \bigcup_{i=0}^k \partial_V R_i \right) \setminus A,$$

where A is the set of isolated points of $\partial\Lambda_f \cap \bigsqcup_i \partial_V R_i$. The *horizontal boundary* of Λ_f is the set

$$\partial_H \Lambda_f = \bigsqcup_{i=0}^k \partial_H R_i.$$

A *vertical (horizontal) component* of $\partial\Lambda_f$ is a connected component of $\partial_V \Lambda_f$ ($\partial_H \Lambda_f$).

Lemma 4.6. *Vertical components of $\partial\Lambda_f$ project to postcritical points of f .*

Proof. If $x = a \in \text{WPC}(f) \setminus \text{PC}(f)$, then a is a critical point of f that is not in the forward image of any $b \in \text{WPC}(f)$. Therefore, any rectangle R projecting to an element of the weak postcritical Markov partition of f such that $f_G(R)$ intersects the line $x = a$ must in fact map across this line. In other words, if R_i and R_{i+1} are the rectangles bordering the line $x = a$, then for any rectangle R , the image $f_G(R)$ crosses R_i if and only if it also crosses R_{i+1} . Consequently, R_i and R_{i+1} are both upper- and lower-aligned, and hence there is no vertical component of $\partial\Lambda_f$ projecting to $a \in I$. ■

Lemma 4.7. *If $a \in \text{PC}(f)$, then the unique periodic point of f_G projecting to a is contained in $\partial_V \Lambda_f$.*

Proof. If $a = 1$, then the conclusion holds. Since the point $x = 1$ is periodic under f , its forward orbit contains a critical point $a \in \text{PC}(f)$ such that $f(a) = 1$. Since λ^{-1} is a root of $D_f(t)$, the periodic point of f_G projecting to $1 \in I$ is $(1, 1) \in I \times \mathbb{R}$.

Moreover, if f_i, f_{i+1} are the affine maps defining f on either side of $a \in I$, then in fact $f_i(x) = a_i + \lambda x$ and $f_{i+1}(x) = \lambda x + 2 - a_i$ for some integer a_i . Therefore, the affine maps defining f_G on either side of the line $x = a$ are

$$\tilde{f}_i(x, y) = (a_i + \lambda x, a_i - \lambda^{-1} y), \quad \tilde{f}_{i+1}(x, y) = (\lambda x + 2 - a_i, \lambda^{-1} y + 2 - a_i).$$

If $(a, y) \in \Lambda_f$, then the images $\tilde{f}_i(a, y) = (1, a_i - \lambda^{-1} y)$ and $\tilde{f}_{i+1}(1, \lambda^{-1} y + 2 - a_i)$ are on the line $x = 1$ and are symmetric about the point $(1, 1)$. In particular, the maps $\tilde{f}_i$ and $\tilde{f}_{i+1}$ agree precisely at the periodic point $\tilde{a}$ of f_G projecting to a . It follows that if $\tilde{a} \in \text{int}(R_i \cup R_{i+1})$ and $U \subseteq \text{int}(R_i \cup R_{i+1})$ is an open rectangle symmetric about $\tilde{a}$, then

$$f_G(U \cap \text{int}(R_i)) = f_G(U \cap \text{int}(R_{i+1})).$$

This contradicts the fact that f_G is measure-preserving, so in fact $\tilde{a} \notin \text{int}(R_i \cup R_{i+1})$. In other words, $\tilde{a} \in \partial_V \Lambda_f$.

We now proceed inductively, going through the periodic orbit of $(1, 1)$ in reverse order. Suppose that $\tilde{a} \in I \times \mathbb{R}$ is a point in the orbit of $(1, 1)$ that is contained in $\partial_V \Lambda_f$, and let $\tilde{b}$ be the periodic point such that $f_G(\tilde{b}) = \tilde{a}$. If $\pi(\tilde{b}) = b$ is a critical point of f , then we repeat the above argument. Otherwise, f_G is defined by a single affine map in a neighborhood of $\tilde{b}$. In particular, if R_j and R_{j+1} are the rectangles bordering the line $x = b$, if $\tilde{b} \in \text{int}(R_j \cup R_{j+1})$, then f_G maps an open rectangular neighborhood of $\tilde{b}$ to an




Figure 13. The weak postcritical set of f pulls back to a rectangular decomposition of Λ_f . Note that the vertical components of $\partial\Lambda_f$ project to the points of $\text{PC}(f)$, in red. Observe that each rectangle is either upper- or lower-aligned with its neighbors (cf. Lemma 4.8). Moreover, each vertical edge in this case contains at its metric center the periodic lift of the postcritical point to which it projects (cf. Figure 12 and Lemma 4.9).

open rectangular neighborhood of $f_G(\tilde{b}) = \tilde{a}$ in $\mathbb{R}^2$. This neighborhood cannot be a subset of Λ_f , since $\tilde{a} \in \partial_V \Lambda_f$ by assumption, but this contradicts the invariance of Λ_f .

Therefore, $\tilde{b} \notin \text{int}(R_j \cup R_{j+1})$, hence $\tilde{b} \in \partial_V \Lambda_f$. $\blacksquare$

Lemma 4.8. *There is only one vertical component of $\partial\Lambda_f$ projecting to a point $a \in \text{PC}(f)$. Therefore, all adjacent rectangles R_i, R_{i+1} of Λ_f are either upper- or lower-aligned.*

See Figure 13 for an example.

Proof of Lemma 4.8. If $a = 0$ or $a = 1$, then the claim follows immediately. Suppose otherwise. Since f is PCP, there is a unique $b \in \text{WPC}(f)$ such that $f(b) = a$. For each i , the connected components of $\partial_V R_i$ project to elements of $\text{WPC}(f)$, so if f_G maps such a component into the line $x = a$, then that component must project to b since $\pi \circ f_G = f \circ \pi$. In other words, the preimage of any vertical component of $\partial\Lambda_f$ is contained in the intersection of Λ_f with the line $x = b$.

Since $a \neq 0, 1$, b is not a critical point of f , and therefore if R_i, R_{i+1} are the rectangles of Λ_f intersecting the line $x = b$, then f_G acts on R_i and R_{i+1} by the same affine map $\tilde{f}$. The fact that $f_G(\partial_V R_j)$ does not intersect the line $x = b$ for any $j \neq i, i + 1$ now implies that the number of vertical components of $\partial\Lambda_f$ projecting to $a \in \text{PC}(f)$ is equal to the

number of vertical components of $\partial\Lambda_f$ projecting to $b \in \text{PC}(f)$, and f_G maps the latter homeomorphically onto the former.

We proceed inductively backwards through the periodic orbit of $a \in \text{PC}(f)$. Eventually, we will arrive at $c = f(1) \in \text{PC}(f)$. Since there is only a single vertical component of $\partial\Lambda_f$ projecting to $1 \in \text{PC}(f)$, the above argument shows that there is similarly a single vertical component projecting to c . Moreover, the same statement now follows for each point in the periodic orbit of a , including a itself.

That all adjacent rectangles are either upper- or lower-aligned now follows immediately: if two such rectangles are neither upper- nor lower-aligned, then there would exist at least two components of $\partial_V \Lambda_f$ along their intersection. ■

Lemma 4.9. *Let $a \in \text{PC}(f)$ and $\tilde{a} \in \Lambda_f$ be the unique periodic point projecting to it. Then $\tilde{a}$ is at the metric center of the vertical component of $\partial\Lambda_f$ containing it.*

Proof. We begin by proving the statement for $a = 1$. By Lemma 4.3, we know that $\tilde{a} = (1, 1)$ in this case. If $f(c) = 1$, then c is a critical point that either is or is not periodic.

If c is not periodic, then $c \notin \text{PC}(f)$ and the two rectangles of Λ_f intersecting the line $x = c$ are both upper- and lower-aligned by Lemma 4.6. Denote these two rectangles by R_i, R_{i+1} , and let $\tilde{f}_i, \tilde{f}_{i+1}$ denote the affine maps by which f_G acts on these rectangles, respectively. Then $\tilde{f}_i(R_i \cap R_{i+1})$ and $\tilde{f}_{i+1}(R_i \cap R_{i+1})$ are vertical line segments of equal length lying on the line $x = 1$. Furthermore, these line segments are contained in Λ_f and are symmetric about $\tilde{a}$.

If c' is another non-postcritical point satisfying $f(c') = 1$ and R_j, R_{j+1} are the rectangles intersecting the line $x = c'$, then the images $\tilde{f}_j(R_j \cap R_{j+1})$ and $\tilde{f}_{j+1}(R_j \cap R_{j+1})$ do not intersect $\tilde{f}_i(R_i \cap R_{i+1})$ and $\tilde{f}_{i+1}(R_i \cap R_{i+1})$ except perhaps at their endpoints: otherwise two rectangles, say R_i and R_j , would have

$$f_G(\text{int } R_i) \cap f_G(\text{int } R_j) \neq \emptyset,$$

contradicting the fact that f_G is measure-preserving.

If c is the unique periodic critical point of f , then there is a single vertical component of $\partial\Lambda_f$ projecting to c , and this component contains the periodic lift $\tilde{c}$ of c . Denote by R_l, R_{l+1} the rectangles of Λ_f intersecting at $x = c$. As before, $\tilde{f}_l(R_l \cap R_{l+1})$ and $\tilde{f}_{l+1}(R_l \cap R_{l+1})$ are vertical line segments symmetric about $\tilde{a}$. Furthermore, f_G maps the single vertical component V to a line segment containing $\tilde{a}$. The union $V \cup (R_l \cap R_{l+1})$ is a connected line segment, hence $\tilde{f}_l(V \cup (R_l \cap R_{l+1}))$ and $\tilde{f}_{l+1}(V \cup (R_l \cap R_{l+1}))$ are connected line segments such that

$$\tilde{f}_l(V \cup (R_l \cap R_{l+1})) \cap \tilde{f}_{l+1}(V \cup (R_l \cap R_{l+1})) = f_G(V).$$

It now follows that $f_G(V)$ is symmetric about $\tilde{a}$, and consequently, so is the entire component of $\partial_V \Lambda_f$ containing $\tilde{a}$.

We now prove the claim for all points in the forward orbit of $1 \in \text{PC}(f)$, proceeding inductively. As we argued in the proof of Lemma 4.8, f_G maps the component of $\partial_V \Lambda_f$

projecting to $a \in \text{PC}(f)$ homeomorphically onto the component projecting to $f(a)$. In particular, if the unique periodic lift $\tilde{a}$ of a lies at the metric center of the vertical component containing it, then so does $f_G(\tilde{a})$, since f_G contracts the vertical direction uniformly by λ^{-1} .

If f is a negative zig-zag, then it has a single postcritical orbit and so the proof is complete. If, however, f is a positive zig-zag, then we must still prove the claim for the fixed point $0 \in \text{PC}(f)$. The exact argument used for $a = 1$ applies here, after observing that the unique periodic lift of $a = 0$ is the point $\tilde{a} = (0, 0)$. ■

Lemma 4.10. *The action of f_G on Λ_f can be recovered via the de Carvalho–Hall construction in the sense of Theorem 1. In particular, f is of pseudo-Anosov type.*

Proof. We reverse engineer the invariant train track τ for the exterior left-veering thick interval map F projecting to f . Construct τ to be the train track with a real edge e_j for each rectangle R_j , a connecting infinitesimal edge between the real edges e_j, e_{j+1} for adjacent rectangles R_j, R_{j+1} , and an infinitesimal loop for each e_j corresponding to a p_i . The loop is on the left (resp. right) of e_j if p_i is on the left (resp. right) vertical boundary of R_j , and if the loop is on the same side of e_j as a connecting infinitesimal edge, then the loop is above (resp. below) the connecting edge if the vertical boundary component of L that contains p_i is above (resp. below) the adjacent rectangle. See Figure 14.

The map induced by f_G on τ leaves τ invariant under the pseudo-isotopies defined in Section 2, hence τ is the invariant train track for a thick interval map projecting to f . By Proposition 3.27, this thick interval map must be either F_L or F_R . One easily checks that the Galois conjugate coordinates imply that it must be F_L .

To complete the proof, observe that by Lemma 4.2, we may choose a right λ -eigenvector of the transition matrix M of f such that the rectangles obtained from τ have the exact dimensions of those in the rectangle decomposition of L . Therefore, we may define an isometry $i: S' \rightarrow \Lambda_f$ taking the horizontal (resp. vertical) foliation of S' to that of Λ_f . By construction, the map induced on Λ_f is f_G . ■

4.2. The forward direction

Lemma 4.10 completes the proof of the reverse direction of Theorem 1. Now it remains to prove the forward direction.

We consider the case when $f: I \rightarrow I$ a positive λ -zig-zag of pseudo-Anosov type, the case when f negative being essentially identical. Let F be the exterior left-veering thickening of f . Let S' be as in the statement of the theorem. The thickening F induces a map $\tilde{f}: S' \rightarrow S'$. Let $p \in S'$ be the unique fixed point of $\tilde{f}$ projecting to $x = 0 \in I$, and let $q \in S'$ be the unique periodic point projecting to $x = 1 \in I$.

We wish to define a map $i: S' \rightarrow I \times \mathbb{R}$ such that $i \circ \tilde{f} = f_G \circ i$ and such that i sends the horizontal and vertical foliations of S' to those of $I \times \mathbb{R}$ in an orientation-preserving way. Fix $s, t \in \mathbb{R}$ distinct and define $i(p) = (0, s)$ and $i(q) = (1, t)$. Observe that these two choices determine i , and moreover the number $d = t - s$ controls the area of $i(S')$:




Figure 14. Reverse-engineering the invariant train track τ for the exterior left-veering thickening of a zig-zag f with rectangular limit set Λ_f . On the top left is the rectangular decomposition of Λ_f projecting to $\text{WPC}(f)$, and on the top right is a train track with junctions corresponding to the vertical boundaries of the rectangles. From left to right, each junction corresponds to a vertical boundary of the same color. Moreover, a junction contains a loop if and only if it corresponds to a line projecting to an element of $\text{PC}(f)$. In this case, the loop is above (resp. below) the spine of τ if the rectangles adjoining the line are lower-aligned (resp. upper-aligned), and is on the left (resp. right) of the junction if the rectangle intersecting $\partial_V \Lambda_f$ is on the left (resp. right) of the line.

if A_d is the Lebesgue measure of $i(S')$ for any choice of i satisfying $t - s = d$, then $A_d = |d| \cdot A_1$.

By construction, there is a piecewise affine map $G: I \times \mathbb{R} \rightarrow I \times \mathbb{R}$ satisfying $i \circ \tilde{f} = G \circ i$. In particular, if $f_0, \dots, f_k$ denote the linear maps defining the original zig-zag f , then G is of the form

$$G(x, y) = (f_i(x), g_i(y)) \quad \text{if } f(x) = f_i(x),$$

where $g_i(y) = a_i(s, t) + (-1)^i \cdot \lambda^{-1}y$. This follows from the fact that $\tilde{f}$ is a piecewise affine orientation-preserving map projecting to f . The remainder of the proof is an analysis of the constants $a_i(s, t)$.

Note that $i(p) = (0, s)$ is a fixed point for G , and hence $y = s$ is fixed by $g_0(y) = a_0(s, t) + \lambda^{-1}y$. We thus have $a_0(s, t) = (1 - \lambda^{-1})s$. To compute the remaining a_i , we require the following lemma.

Lemma 4.11. *Let $c_i = i \cdot \lambda^{-1}$ be a critical point of f , and set g_{i-1}, g_i to be the defining y -coordinate maps of G on either side of the line $x = c_i$. Then $g_{i-1}(y_i) = g_i(y_i)$ for a unique number y_i , and moreover this number satisfies $G(c_i, y_i) = s$ or $G(c_i, y_i) = t$ depending on whether i is even or odd, respectively.*

Proof. Since g_{i-1} and g_i are affine with slopes of opposite signs they agree at a unique point y_i . To understand this number y_i , we consider what happens for $\tilde{f}: S' \rightarrow S'$.

If c_i is not a postcritical point of f , then the line $x = c_i$ partitions a rectangle of S' . Depending on the parity of i , the affine maps defining $\tilde{f}'$ on either side of the vertical line send the line to either $x = 0$ or $x = 1$, with opposite orientations. The fact that the unique periodic lifts of $x = 0$ and $x = 1$ are points of cone angle π lying in the center of the corresponding vertical sides of S' implies that these two affine maps would send the same point of $c_i \times \mathbb{R}$ to the singularity if their domains were extended to include this point.

If c_i is a postcritical point, then the unique periodic lift of c_i lies on the vertical boundary of exactly one of the corresponding rectangles of S' , and moreover it lies in the center of this boundary component. Since the image of this singularity is the sole singularity of $\tilde{f}$ on the corresponding vertical leaf of S' , the two affine maps defining $\tilde{f}$ on either side of $x = c_i$ must map each point of this line to points on $x = 0$ or $x = 1$ that are at equal distances from the singularity. Because all vertical distances are scaled by the same factor, it now follows that both maps agree on the singularity projecting to c_i .

In either case, the same is true for the maps g_{i-1} and g_i after mapping into $I \times \mathbb{R}$. ■

We return to computing the $a_i(s, t)$, thereby completing the proof of Theorem 1.

Proof of Theorem 1. By Lemma 4.11, $g_1(y) = 2t - g_0(y)$, and in particular

$$a_1(s, t) = 2t - a_0(s, t) = 2t - (1 - \lambda^{-1})s.$$

Similarly, we have $g_2(y) = 2s - g_1(y)$, hence $a_2(s, t) = 2s - a_1(s, t)$, and in general we have

$$a_i(s, t) = \begin{cases} (i + 1 - \lambda^{-1})s - it & \text{if } i \text{ is even,} \\ (\lambda^{-1} - i)s + (i + 1)t & \text{if } i \text{ is odd.} \end{cases}$$

Setting $s = 0$ and $t = 1$ (and hence $d = 1$), we specialize to the case

$$a_i(s, t) = \begin{cases} -i & \text{if } i \text{ is even,} \\ i + 1 & \text{if } i \text{ is odd.} \end{cases}$$

In other words, each affine piece of $G: I \times \mathbb{R} \rightarrow I \times \mathbb{R}$ is of the form

$$G_i(x, y) = \begin{cases} (\lambda x - i, \lambda^{-1}y - i) & \text{if } i \text{ is even,} \\ (i + 1 - \lambda x, i + 1 - \lambda^{-1}y) & \text{if } i \text{ is odd.} \end{cases}$$

Thus G is precisely the Galois lift f_G of Thurston. Condition (2) of the theorem is immediately verified, so it remains to argue that the digit polynomial D_f has λ^{-1} as a root. We can define a “vertical” digit polynomial using the orbit of $y = 1$ under the g_i . Observe that this new polynomial is precisely D_f , and is necessarily satisfied by λ^{-1} . This completes the proof of Theorem 1. ■

Remark 4.12. One quickly verifies that

$$-\frac{k}{1-\lambda^{-1}}a_0(s, t) + \sum_{i=1}^k a_i(s, t) = 0,$$

and hence the map $(s, t) \mapsto (a_0(s, t), \dots, a_k(s, t))$ is a linear map of $\mathbb{R}^2 \setminus \Delta$ into $\mathbb{R}^{k+1}$ whose image is a plane minus the line corresponding to $d = 0$. This missing line is the image of the diagonal $\Delta \subseteq \mathbb{R}^2$, and we can foliate the image plane with lines parallel to this one, each such line corresponding to a different value of $d = t - s$. The area of the limit set $i_d(S')$ of G_d scales linearly with $|d|$, i.e., if A_d is the area of $i_d(S')$, then $A_d = |d| \cdot A_1$. In this way, we can interpret the prohibited case $d = 0$ as a degenerate limit set of area zero. Moreover, the two half-planes defined by $d > 0$ and $d < 0$ correspond to the underlying train track map τ being left- or right-veering, respectively.

5. The postcritical orbit of a zig-zag of pseudo-Anosov type

In light of Theorem 1, it is natural to ask when an exterior left-veering thickening F_L has a finite generalized invariant train track. This section obtains necessary conditions on the structure of the thick interval map F_L associated to a PCP zig-zag of pseudo-Anosov type. In particular, Proposition 5.10 will be instrumental to our proof of Theorem 2 in Section 6.

Recall that if $F_L: (S^2, \mathbb{I}) \rightarrow (S^2, \mathbb{I})$ is a thickening of a PCF interval map f , then we denote by $\mathbf{0}$ the junction projecting to $0 \in I$, and similarly we denote by $\mathbf{1}$ the vertex projecting to $1 \in I$.

Definition 5.1. Let $f: I \rightarrow I$ be a λ -zig-zag map and $c_i = i \cdot \lambda^{-1}$ the critical points of f , for $i = 1, \dots, \lfloor \lambda \rfloor$. Let $F_L: (S^2, \mathbb{I}) \rightarrow (S^2, \mathbb{I})$ be the exterior left-veering thickening of f . For each i , we denote by C_i the junction projecting to c_i .

Definition 5.2. Let τ be the generalized invariant train track for the left-veering thickening F_L of the PCP zig-zag map f . By a *connecting infinitesimal edge* we will mean an infinitesimal edge of τ connecting two distinct real edges. The remaining infinitesimal edges of τ , namely those that connect a real edge to itself, are called *loops*.

We define the *spine* τ' of τ to be the union of all real edges and connecting infinitesimal edges of τ . We orient τ' from $\mathbf{0}$ to $\mathbf{1}$. A loop $\gamma \subseteq \tau$ contained in an intermediate vertex of $\mathbb{I}$ is called *interior* if γ is to the left of τ' and *exterior* otherwise. See Figure 15.




Figure 15. A (finite) generalized train track τ and an exterior left-veering thick interval map preserving it. Here the spine of τ is the union of the black edges. Of the intermediate loops, e_1 is interior while e_2 and e_3 are exterior. Observe that under the action of F_L , the image of an interior loop points in toward the horizontal midline of $\mathbb{I}$, hence will be enclosed by arcs after pseudo-isotopy unless it maps into $\mathbf{0}$ or $\mathbf{1}$.




Figure 16. An exterior loop in C_i produces two non-parallel loops in $\mathbf{1}$.

Remark 5.3. The interior loops γ of τ are precisely the loops whose image $F(\gamma)$ points toward the horizontal midline of $\mathbb{I}$. One can see this by noting that F is exterior left-veering and preserves orientation. It is for this reason that we refer to such loops as “interior”. In the arguments of this section, we will see that interior loops are rare, and the existence of more than one for τ is an obstruction to the finiteness of τ .

Recall (cf. Definition 1.4) that a zig-zag f is *positive* if $f(0) = 0$, and *negative* if $f(0) = 1$.

Proposition 5.4. *Let $f: I \rightarrow I$ be a PCP zig-zag map of growth rate $\lambda > 2$. If f is of pseudo-Anosov type, then $c_1 = \lambda^{-1}$ is in the forward orbit of $x = 1$. Moreover, the loop in the vertex C_1 projecting to c_1 must be interior.*

Proof. Suppose first that f is positive, so that $f(0) = 0$. Since f is PCP, some $c_i = i \cdot \lambda^{-1}$ is in the forward orbit of $x = 1$. By Theorem 3.14, the corresponding junction C_i contains an edge ε and a loop γ of the invariant train track τ . If γ is exterior, then after pseudo-isotopy, the images $F(\varepsilon)$ and $F(\gamma)$ will be non-parallel loops in $\mathbf{1}$, contradicting Theorem 3.14. See Figure 16. Thus γ must be interior.




Figure 17. The image of all edges in C_1 is further interior than that of any other C_i mapping into $\mathbf{1}$, so any loop in C_i generates a loop in $\mathbf{1}$ that persists after pseudo-isotopy.




Figure 18. The image of an interior loop points inward, and because all horizontal layers of $F(\tau)$ span the full length of $\mathbb{I}$ except for the last, the image of an interior loop will be trapped by other edges after pseudo-isotopy unless it maps into $\mathbf{0}$ or $\mathbf{1}$, where it can be absorbed into the parallel loop formed by the turn.

If $i \neq 1$, then F will map the connecting edge ε' of τ that is within C_1 into $\mathbf{1}$ further interior than the images of ε and γ , and thus after pseudo-isotopy, we will obtain two non-parallel loops enclosed by a third, again contradicting Lemma 3.14. See Figure 17. On the other hand, if $i = 1$, then the images of ε and γ will be the furthest interior in $\mathbf{1}$ and all other edges that map into $\mathbf{1}$ will pinch to parallel loops, which then combine under pseudo-isotopy.

Now suppose that f is negative, so that $f(0) = 1$. The same argument applies, except that the role of $\mathbf{1}$ is taken by $\mathbf{0}$, which maps into $\mathbf{1}$. ■

Proposition 5.5. *Let $f: I \rightarrow I$ be a PCP zig-zag map of growth rate $\lambda > 2$. If f is of pseudo-Anosov type, then the invariant train track $\tau = \tau_L$ has exactly one interior loop, namely the loop contained in the vertex C_1 .*

Proof. Proposition 5.4 tells us that C_1 contains an interior loop, so it remains to show that τ has no other interior loops. Furthermore, our argument in the proof of Proposition 5.4 demonstrates that no other critical vertex C_i can contain an interior loop, so if τ contains another interior loop γ , then $F(\gamma)$ cannot lie in $\mathbf{0}$ or in $\mathbf{1}$. In other words, $F(\gamma)$ lies in some intermediate vertex V . However, since γ is interior, the image $F(\gamma)$ faces toward the center of $\mathbb{I}$. See Figure 18. Thus after pseudo-isotopy, $F(\gamma)$ will be enclosed by multiple connecting edges, contradicting the structure of τ described in Theorem 3.14. ■

Corollary 5.6. *Let $f: I \rightarrow I$ be a PCP zig-zag map of growth rate $\lambda > 2$, and suppose f is of pseudo-Anosov type.*

- (1) *If f is positive and $\lfloor \lambda \rfloor = m$ is odd, then the orbit of $x = 1$ under f is $1 \mapsto \lambda^{-1} \mapsto 1$.*
- (2) *If f is negative and $\lfloor \lambda \rfloor = m$ is even, then the orbit of $x = 1$ under f is $1 \mapsto \lambda^{-1} \mapsto 0 \mapsto 1$.*

In either case, λ has minimal polynomial $p(x) = x^2 - (m + 1)x + 1$.

Proof. First suppose f is positive and m is odd. The image $F_L(\gamma)$ of the loop $\gamma \subseteq \tau$ contained in **1** is above the other strands of $F_L(\tau)$, and so after pseudo-isotopy it will be an interior loop γ' . This interior loop must necessarily be contained in C_1 and must map into **1** by Proposition 5.5. It follows that $x = 1$ has period 2. The linear branch of f that applies to $x = 1$ is $x \mapsto (m + 1) - \lambda x$, hence we have $\lambda^{-1} = m + 1 - \lambda$.

The case when f is negative and m is even is similar. The image $F_L(\gamma)$ is again an interior loop after pseudo-isotopy, and so must be contained in C_1 . Since the linear branch of f that applies to $x = 1$ is $x \mapsto (m + 1) - \lambda x$, it follows that $\lambda^{-1} = m + 1 - \lambda$. ■

Remark 5.7. The examples described in Corollary 5.6 are the simplest examples of uniform expanders of pseudo-Anosov type. The corresponding pseudo-Anosovs all live on the four-punctured sphere $\Sigma_{0,4}$ and lift via a branched double cover to hyperbolic automorphisms of the torus. To avoid these simple cases, we make the following definition (see Remark 5.9 below).

Definition 5.8. For $m \geq 2$ and $p \geq 3$, define $\text{PA}(m, p)$ to be the set of zig-zags f of pseudo-Anosov type such that

- (1) f has m critical points,
- (2) $\#\text{PC}(f) = p$.

We also define the set

$$\text{PA}(m) = \bigcup_{p \geq 4} \text{PA}(m, p).$$

Remark 5.9. An element $f \in \text{PA}(m, p)$ defines a pseudo-Anosov on the $(p + 1)$ -punctured sphere, where the point at infinity is fixed. Ripping open this point to a boundary component, we may also think about this pseudo-Anosov as a braid on the p -punctured disc, up to multiplication by a full twist Δ^2 around the boundary. The simple examples described in Corollary 5.6 are precisely the maps $f \in \text{PA}(m, 3)$, and the definition of $\text{PA}(m)$ allows us to avoid these examples in the future.

Recall that we use the notation $c_i = i \cdot \lambda^{-1}$. For a λ -zig-zag map f , we have the subintervals of definition

$$I_j = [c_j, c_{j+1}) \quad \text{for } 0 \leq j \leq m - 1, \quad \text{and} \quad I_m = [c_m, 1].$$




Figure 19. The positive zig-zag for λ the Perron root of $t^4 - 4t^3 - 2t^2 - 4t + 1$. The map f is of pseudo-Anosov type, as demonstrated by Figure 15, and the orbit of $x = 1$ is shown in red. In particular, $\text{PC}(f)$ has points of types E , C , R , and P_{m-2} .

Proposition 5.10. *Suppose $f \in \text{PA}(m)$ with slope λ . Then $\text{PC}(f)$ always contains the following three types of points:*

- (E) *The extremal points $x = 0$ and $x = 1$.*
- (C) *The critical point $c_1 = \lambda^{-1}$.*
- (R) *Exactly one point in the interior of I_{m-1} .*

If $\text{PC}(f)$ contains other points, then they are of the following two types:

- (P_{m-2}) *Points in the interior of I_{m-2} .*
- (P_m) *Points in the interior of I_m .*

Proof. The existence of type E points is obvious, and the existence of type C points follows from Proposition 5.4. The single point of type R is precisely the postcritical point that maps to c_1 . Since the loop of τ in C_1 must be interior, it follows that the image of the loop of τ that maps into C_1 must be interior, hence lies over a postcritical point at which f is orientation-reversing. Since there is only one interior loop of τ , this is the only postcritical point at which f reverses orientation.

To finish the proof, suppose f has another critical point $x = p$ other than type E , C , and R . Then the loop of τ over this critical point must have an image that is exterior so that $F(\tau)$ does not trap the loop after tightening. Since F is exterior veering, it follows that the image of this loop must come after the third-to-last turn of F , hence p is in the interior of $I_{m-2} \cup I_{m-1} \cup I_m$. Moreover, since p is not of type R , f preserves orientation at p , hence $p \notin I_{m-1}$. ■

Example 5.11. Consider the thick interval map in Figure 15. This is the exterior left-veering thickening of the positive λ zig-zag f , where λ is the Perron root of $D_f(t) = t^4 - 4t^3 - 2t^2 - 4t + 1$. The graph of f is shown in Figure 19.

Corollary 5.12. *Suppose $f \in \text{PA}(m)$ has slope λ . If $\text{PC}(f)$ contains no points of type P_{m-2} or P_m , then λ is a quadratic integer.*

Proof. If f is positive and has no postcritical points of type P_i , then the orbit of $x = 1$ is

$$1 \mapsto \lambda - m \mapsto m + m\lambda - \lambda^2 \mapsto 2 - m\lambda - m\lambda^2 + \lambda^3 = 1,$$

hence λ satisfies the relation

$$0 = \lambda^3 - m\lambda^2 - m\lambda + 1 = (\lambda + 1)(\lambda^2 - (m + 1)\lambda + 1).$$

Since $\lambda > 1$ and $x^2 - (m + 1)x + 1$ is irreducible over $\mathbb{Q}$, we see that λ is a quadratic integer. Similarly, if f is negative and has no postcritical points of type P_i , then from the orbit of $x = 1$, we obtain the relation

$$0 = \lambda(\lambda + 1)(\lambda^2 - (m + 1)\lambda + 1).$$

Hence λ is again a quadratic integer. ■

6. Classifying zig-zags of pseudo-Anosov type

In this section, we consider the set $\text{PA}(m)$ of zig-zags of pseudo-Anosov type with modality $m \geq 2$. The case $m = 1$ was considered by Hall in [6, Theorem 2.1], in which he gave an explicit bijection between $\text{PA}(1)$ and $\mathbb{Q} \cap (0, 1/2)$. We give an explicit bijection between $\text{PA}(m)$ and $\mathbb{Q} \cap (0, 1)$ for each $m \geq 2$.

The proof of Theorem 2 naturally breaks into three cases: (1) $m \geq 4$ even, (2) $m \geq 3$ odd, and (3) $m = 2$. In preparation for the proof, we investigate each of these cases separately. We treat case (1) first since it is essential to understanding cases (2) and (3).

Throughout this section, f is a PCP λ -zig-zag map with modality $m = \lfloor \lambda \rfloor$. Moreover, we always assume that $\text{PC}(f) \geq 4$, so that it is possible for f to belong to $\text{PA}(m)$ (cf. Definition 5.8).

6.1. The case $m \geq 4$ even

Definition 6.1. Given $n \geq 3$ and $2 \leq k \leq n - 1$, define $\rho_e(n, k) \in S_n$ to be the permutation such that

$$\rho_e(n, k)(i) = \begin{cases} n, & i = 1, \\ i + (n - k), & 2 \leq i \leq k - 1, \\ i - (k - 1), & k \leq i \leq n. \end{cases}$$

Example 6.2. Here are some examples of $\rho_e(n, k)$ for $n = 7$:

$$\rho_e(7, 2) = (7, 6, 5, 4, 3, 2, 1),$$

$$\rho_e(7, 3) = (7, 5, 3, 1)(6, 4, 2),$$

$$\rho_e(7, 6) = (7, 2, 3, 4, 5, 6, 1).$$

Note that $\rho_e(7, 2)$ and $\rho_e(7, 6)$ are full 7-cycles, whereas $\rho_e(7, 3)$ is not. As we will see, $\rho_e(n, k)$ is an n -cycle if and only if $n - k$ and $k - 1$ are coprime (cf. Proposition 6.7).




Figure 20. The layers of $F(\tau)$ for the exterior left-veering thickening of a bimodal zig-zag.

Definition 6.3. Let F be a thickening of a PCF map f , and let τ denote the invariant generalized train track of F . As usual, denote by $\mathbf{0}$ and $\mathbf{1}$ the junctions projecting to $0, 1 \in I$, respectively, and let τ' be the spine of τ . We define a *layer* of the image $F(\tau)$ to be a connected component of the complement of $\mathbf{0} \cup \mathbf{1}$ in $F(\tau')$. See Figure 20.

Remark 6.4. If f is an m -modal zig-zag and F is any thickening of f , then $F(\tau)$ has $m + 1$ layers. Indeed, if L is a layer of $F(\tau)$, then $F^{-1}(L)$ projects to one of the $m + 1$ different intervals of monotonicity I_j for f .

Definition 6.5. Let f be an m -modal zig-zag and F the exterior left-veering thickening of f . For each $0 \leq j \leq m$, the j -th layer of $F(\tau)$ is the layer L_j such that

$$\pi(F^{-1}(L_j)) = I_j.$$

Recall (cf. Definition 1.15) that the *permutation type* of a PCP interval map f is the permutation $\rho(f)$ such that $f(x_i) = x_{\rho(f)(i)}$.

Proposition 6.6. Let $f: I \rightarrow I$ be a PCP λ -zig-zag map with $\lfloor \lambda \rfloor = m \geq 4$ even. Then $f \in \text{PA}(m)$ if and only if the following hold:

- (1) Under the ordering $x_1 < x_2 < \dots < x_n = 1$ of the orbit of $x = 1$, we have $\rho(f) = \rho_e(n, k)$ for some k .
- (2) The $\{x_i\}$ satisfy the conclusion of Proposition 5.10, with x_1 being the sole point of type C and x_k being the sole point of type R .

Proof. We begin first by assuming that f is of pseudo-Anosov type. Let τ be the invariant train track for the exterior left-veering thick interval map F . By Proposition 5.4, the junction C_1 over $c_1 = \lambda^{-1}$ contains the sole interior loop γ of τ , and this loop must map into $\mathbf{1}$. Moreover, since $\lfloor \lambda \rfloor \geq 4$, the layers L_0, L_1 , and L_2 stretch fully between $\mathbf{0}$ and $\mathbf{1}$, with L_1 above L_3 and below L_2 . In particular, there can be no intermediate loops of τ before γ , since otherwise these would be trapped by L_2 or L_3 . Thus if ρ is the permutation describing the action of f on the orbit of $x = 1$, we must have $\rho(1) = n$.

Now set $\rho^{-1}(1) = k$. By Proposition 5.5, all loops in τ corresponding to $i \neq k$ must have exterior image, so their images must lie in L_{m-2} or L_m . In particular, if $k = 2$, then the remaining $n - k$ loops must be sent in order to the remaining $k - 2$ junctions containing a loop. In particular, we have

$$\rho = (1, n, n - 1, n - 2, \dots, 3, 2) = \rho_e(n, 2).$$

If $k > 2$, then the loops labeled 2 through $k - 1$ must map to the last $k - 3$ loops not in $\mathbf{1}$, in order. This is again because these loops must all have exterior images, and so in particular, cannot be covered from below by loops on L_m . Now the loops labeled $k + 1$ through n must map to the remaining loops labeled 2 through $n - k + 1$ in order, and thus we see that $\rho = \rho_e(n, k)$.

Suppose instead that f satisfies properties (1) and (2) above. Let F be the exterior left-veering thickening of f . We construct the invariant generalized train track τ of F as described in Section 2. We begin with τ_0 , the (disconnected) union of real edges, and after the tightening pseudo-isotopies, $\tau_1 = F_*(\tau_0)$ contains a loop γ projecting to $x_n = 1$. Consider the successive images $F_*^j(\gamma)$, which by conditions (1) and (2) are exterior until some minimal index $j = a$. Moreover, each of these initial forward images is guaranteed to be uncovered from below by layers of $F(\tau_{j-1})$, and so are the only infinitesimal edges within the corresponding junctions aside from a single connecting edge between the two adjacent real edges.

The image $F_*^a(\gamma)$ is an interior loop, and by assumption must project to $x_1 = \lambda^{-1}$, since x_k is the sole postcritical point of type R and $\rho_e(n, k)(k) = 1$. In particular, $F_*^{a+1}(\gamma)$ is a loop over x_n and since $x_1 = \lambda^{-1}$, this loop is interior to every turn of $F_*(\tau_a)$. Hence after tightening, we recover a single loop over x_n and the resulting generalized train track $\tau_{a+1} = \tau$ is invariant. It follows that τ is finite and that f is of pseudo-Anosov type. ■

In order to completely characterize positive PCP zig-zags with modality $m \geq 4$ even, it remains to determine when $\rho_e(n, k)$ is an n -cycle.

Proposition 6.7. *The permutation $\rho_e(n, k) \in S_n$ is an n -cycle if and only if*

$$(n - k, n - 1) = 1.$$

Proof. Set $\rho = \rho_e(n, k)$, and let ρ' be the permutation on $\{2, \dots, n\}$ defined by

$$\rho'(i) = \begin{cases} \rho(i), & 2 \leq i \neq k, \\ n, & i = k. \end{cases}$$

Thus ρ' is the element of S_{n-1} obtained by deleting the symbol 1 from the cycle decomposition of ρ . Observe that, since 1 is in the orbit of n , ρ is an n -cycle if and only if ρ' is an $(n - 1)$ -cycle. Shifting all labels down by 1, we have

$$\rho'(i) = \begin{cases} i + (n - k), & 1 \leq i \leq k - 1, \\ i - (k - 1), & k \leq i \leq n - 1. \end{cases}$$

Interpreting this modulo $n - 1$, we see that ρ' acts via addition by $n - k$: indeed, subtraction by $k - 1 \pmod{n - 1}$ is equivalent to addition by $n - 1 - (k - 1) \equiv n - k \pmod{n - 1}$. Thus this action is transitive if and only if $(n - k, n - 1) = 1$. ■

6.2. The case $m \geq 3$ odd

We next consider the case when f is a PCP zig-zag of modality $m \geq 3$ odd. In this case, we observe that the orbit of $x = 1$ ends with $c_1 \mapsto 0 \mapsto 1$.

Definition 6.8. Given $n \geq 3$ and $2 \leq k \leq n - 1$, define $\rho_o(n, k) \in S_{n+1}$ to be the permutation such that

$$\rho_o(n, k)(i) = \begin{cases} n, & i = 0, \\ 0, & i = 1, \\ i + (n - k), & 2 \leq i \leq k - 1, \\ i - (k - 1), & k \leq i \leq n. \end{cases}$$

Remark 6.9. Note that if we define $\rho' \in S_n$ to be the permutation obtained from $\rho = \rho_o(n, k)$ by deleting 0 from the orbit of n , then ρ' is an n -cycle if and only if ρ is an $(n + 1)$ -cycle. Since $\rho' = \rho_e(n, k)$, we see that this is again the case if and only if $(n - k, n - 1) = 1$. We have therefore proven the following statement.

Proposition 6.10. *The permutation $\rho_o(n, k) \in S_{n+1}$ is an $(n + 1)$ -cycle if and only if $(n - k, n - 1) = 1$.*

A nearly identical argument to that of Proposition 6.6 proves the following analog.

Proposition 6.11. *Let $f: I \rightarrow I$ be a PCP λ -zig-zag map with $\lfloor \lambda \rfloor = m \geq 3$ odd. Then $f \in \text{PA}(m)$ if and only if the following hold:*

- (1) *Under the ordering $0 = x_0 < x_1 < \dots < x_n = 1$ of the orbit of $x = 1$, we have $\rho(f) = \rho_o(n, k)$ for some k .*
- (2) *The $\{x_i\}$ satisfy the conclusion of Proposition 5.10, with x_1 being the sole point of type C and x_k being the sole point of type R.*

6.3. The case $m = 2$

It remains to consider the case of unrestricted PCP zig-zags of modality $m = 2$.

Definition 6.12. Let $n \geq 3$. For $2 \leq k \leq n - 1$, set

$$\kappa(n, k) = (1, 2, \dots, k - 1)(k) \cdots (n) \in S_n$$

and define

$$\rho_2(n, k) = [\kappa(n, k)]^{-1} \circ \rho_e(n, k) \circ \kappa(n, k)$$

Lemma 6.13. Fix $n \geq 3$ and $2 \leq k \leq n - 1$ and set $\rho_2 = \rho_2(n, k)$. Then the following conditions hold:

- (1) $\rho_2(k) = k - 1$.
- (2) $\rho_2(k - 1) = n$.
- (3) If $i \leq j < k$, then $\rho_2(i) \leq \rho_2(j)$.
- (4) If $k < i \leq j$, then $\rho_2(i) \leq \rho_2(j)$.
- (5) If $i < k < j$, then $\rho_2(j) < \rho_2(i)$.

Moreover, $\rho_2(n, k)$ is the only element of S_n to satisfy these conditions. Finally, $\rho_2(n, k)$ is a full n -cycle if and only if $(n - k, n - 1) = 1$.

Proof. Set $\rho = \rho_e(n, k)$ and $\kappa = \kappa(n, k)$ so that $\rho_2 = \kappa^{-1}\rho\kappa$. Using this relation, one readily verifies conditions (1) and (2). Now suppose $i \leq j < k$. Then we have $\rho_2(i) = \kappa^{-1}(i + 1 + (n - k))$ and $\rho_2(j) = \kappa^{-1}(j + 1 + (n - k))$. Since $i \leq j$ and κ preserves this ordering except at k , we see that $\rho_2(i) \leq \rho_2(j)$ unless $i + 1 + (n - k) = 1$, which is impossible. Similarly, if $k < i \leq j$, then $\rho_2(i) = \kappa^{-1}(i - (k - 1))$, and $\rho_2(j) = \kappa^{-1}(j - (k - 1))$. Again, we see that $\rho_2(i) \leq \rho_2(j)$ unless $i - (k - 1) = 1$, i.e., that $i = k$. This is also impossible.

To prove condition (5), it is enough by (4) to argue that if $i < k$, then $\rho_2(n) < \rho_2(i)$. In this direction, note that $\rho_2(i) = \kappa^{-1}(i + 1 + (n - k))$, whereas $\rho_2(n) = \kappa^{-1}\rho(n) = \kappa^{-1}(n - k + 1)$. Again, by the fact that κ preserves ordering except at k , it follows that $\rho_2(n) < \rho_2(i)$.

Suppose that $\omega \in S_n$ satisfies conditions (1)–(5) above. We wish to show that $\omega = \rho_2(n, k)$. Condition (1) implies that this second set has image including n , and condition (2) determines the image of the last remaining index. The monotonicity conditions (3) and (4) impose a strict ordering on the two subsets $\{1, \dots, k - 1\}$ and $\{k + 1, \dots, n\}$ of remaining indices, and condition (5) implies that the image of the second subset must be completely below that of the first. It follows that $\omega(i)$ is determined for each i , and hence $\omega = \rho_2(n, k)$.

Finally, since ρ_2 is conjugate to ρ , the two share the same cycle type, and hence by Proposition 6.7, ρ_2 is an n -cycle if and only if $(n - k, n - 1) = 1$. ■

Proposition 6.14. Let $f: I \rightarrow I$ be a PCP λ -zig-zag map with $[\lambda] = 2$. Then $f \in \text{PA}(2)$ if and only if the following hold:

- (1) Under the ordering $x_1 < x_2 < \dots < x_n = 1$ of the orbit of $x = 1$, we have $\rho(f) = \rho_2(n, k)$ for some k .
- (2) The $\{x_i\}$ satisfy the conclusion of Proposition 5.10, with x_{k-1} being the sole point of type C and x_k being the sole point of type R .

Proof. Suppose that f is of pseudo-Anosov type. By Proposition 5.10, f has exactly one postcritical point in the interior of I_1 , as well as another at $x = \lambda^{-1}$. Let $\rho \in S_n$ describe the action of f on the periodic orbit of $x = 1$. Let F be the exterior left-veering

thick interval map projecting to f and let τ be the invariant generalized train track for F . Since f is of pseudo-Anosov type, τ is finite and has structure described by Lemma 3.14 and Proposition 5.5. In particular, τ has a single interior loop, and hence there is a single loop γ of τ whose image is this interior loop. Let x_k be the postcritical point of f to which this loop projects. Since the single interior loop of τ necessarily maps to the loop projecting to $x_n = 1$, we have $\rho^2(k) = n$.

Since γ maps to the unique interior loop of τ , γ is the unique loop such that $F(\gamma) \in L_2$. Moreover, since $\rho^2(k) = n$, we see that $F(F_*(\gamma))$ is after L_1 and before L_2 . In particular, there are no loops of τ after $F_*(\gamma)$ and before γ . Therefore, $F_*(\gamma)$ projects to x_{k-1} , and so $\rho(k) = k - 1$. Since $\rho^2(k) = n$, it now follows that $\rho(k - 1) = n$. It is now readily checked that ρ satisfies the five conditions of Lemma 6.13, and hence $\rho = \rho_2(n, k)$.

Suppose on the other hand that conditions (1) and (2) above hold. Let F be the exterior left-veering thick interval map projecting to f , and let τ be the invariant generalized train track of F . Note that τ has a loop γ projecting to $x = 1$. Let F^a be the first iterate that sends γ to an interior loop. All forward images of γ before this must be exterior loops, by the definition of ρ_2 and the fact that F is exterior left-veering. Note that $a \geq 2$ and that $F^{a-1}(\gamma)$ is a loop projecting to x_k since its image is an interior loop.

We claim that $F^{a+1}(\gamma) = \gamma$. Indeed, $x = \lambda^{-1}$ must be the $(k - 1)$ -st postcritical point of f from left to right, since $\rho_2(n, k)(k - 1) = n$. Moreover, since f has a single postcritical point in I_1 , this point must be the k -th postcritical point, immediately proceeding $x = \lambda^{-1}$, and since $\rho_2(n, k)(k - 1) = n$, we see that this point maps to $x = \lambda^{-1}$. Thus the single interior loop of τ is mapped into the fat vertex over $x = 1$, and hence after isotopy becomes simply γ . In particular, τ is finite, and hence f is of pseudo-Anosov type. ■

Remark 6.15. Propositions 6.6, 6.11, and 6.14 imply that if $f \in \text{PA}(m)$, then

$$\rho(f) = \begin{cases} \rho_e(n, k) & \text{if } m \geq 4 \text{ even,} \\ \rho_o(n, k) & \text{if } m \geq 3 \text{ odd,} \\ \rho_2(n, k) & \text{if } m = 2. \end{cases}$$

Therefore, when m is unspecified, we will write $\rho(f) = \rho_m(n, k)$.

6.4. The proof of Theorem 2

We are now ready to prove Theorem 2, which we restate here for convenience.

Theorem 2. Fix $m \geq 2$ and let $\Phi: \text{PA}(m) \rightarrow \mathbb{Q} \cap (0, 1)$ be the map defined by

$$\Phi(f) = \frac{n - k}{n - 1} \quad \text{if } \rho(f) = \rho_m(n, k).$$

Then Φ is a bijection. Moreover, for each $p \geq 4$ the image $\Phi(\text{PA}(m, p))$ consists of the set of reduced rationals in $(0, 1)$ of denominator $p - 2$.

Proof. To fix ideas, let $m = 2$. Suppose f and g are two bimodal unrestricted PCP zig-zags of pseudo-Anosov type, and that $\rho(f) = \rho(g) = \rho_2(n, k)$ for some n, k satisfying $(n - k, n - 1) = 1$. We claim that $f = g$. Indeed, $\rho_2(n, k)$ and bimodality determine the transition matrix M of f , by Proposition 6.14. In particular, f and g are uniform expanders with the same topological entropy, and hence the same slope. It follows that $f = g$, and so the map is injective.

To prove surjectivity, it is enough to show that for every $\rho_2(n, k)$ with $(n - k, n - 1)$, there is a bimodal unrestricted PCP zig-zag f of pseudo-Anosov type that acts on the orbit of $x = 1$ as $\rho_2(n, k)$. We do this by constructing the exterior left-veering thick interval map and then projecting onto the horizontal coordinate to obtain a zig-zag with the necessary combinatorics. Indeed, fix some $\rho_2(n, k)$ satisfying $(n - k, n - 1) = 1$ and let τ be a train track consisting of real edges $e_1, \dots, e_n$ and infinitesimal edges $f_1, \dots, f_{n-1}$, with f_j joining e_j to e_{j+1} . Further, adorn τ with infinitesimal loops as follows:

- (1) Loops γ_0 and γ_n on the left of e_1 and the right of e_n , respectively.
- (2) An upward-pointing loop γ_{k-1} attached to the left of e_k .
- (3) Downward-pointing loops γ_j attached to the right of e_j for $j = 1, \dots, n - 1$, except for $j = k - 1$.

Define F to be the exterior left-veering train track map permuting the γ_k according to $\rho_2(n, k)$, while also fixing γ_0 . It is not hard to see by the construction of F and the structure of $\rho_2(n, k)$ that F preserves τ .

We claim that the transition matrix M of the e_j is irreducible, i.e., for every pair of edges e_{j_1}, e_{j_2} some iterate of F maps e_{j_1} across e_{j_2} . Since each γ_j for $j \geq 1$ maps to γ_n , every e_j eventually maps across e_n , and so it is enough to prove that each e_j is eventually covered by some forward image of e_n . Since γ_n is on the right of e_n and γ_n maps to γ_j for each $j \geq 1$, it follows that e_n must map across the edge to the left of γ_j for each positive $j \neq k$, that is, e_n eventually maps across e_j for each $j \neq k - 1$.

It remains to prove that e_n eventually maps across e_{k-1} . But some e_j does this, and since e_n already maps to every edge besides e_{k-1} , we will be done unless e_{k-1} is the only edge that maps across e_{k-1} . This is impossible: the image of e_{k-1} is entirely before the first turn of τ , since γ_{k-1} maps to γ_n , and so the image of some other e_j covers e_{k-1} between the first and second turns. Thus M is irreducible.

Let λ be the spectral radius of M . By the Perron–Frobenius theorem, λ is a simple eigenvalue for M , and M has positive left- and right-eigenvectors for λ . Let $u = (u_1, \dots, u_n)$ be the unique left eigenvector such that $\sum_i u_i = 1$. Then assigning each e_j the length u_j (and declaring each infinitesimal edge to be length 0), we obtain a uniform λ -expander f by projecting the action of F onto the horizontal coordinate. Since u is positive, none of the e_j are collapsed, and so f is a PCP zig-zag having the necessary combinatorics: $x = 1$ is periodic and f acts on this orbit by $\rho_2(n, k)$. Thus the map is surjective for $m = 2$.

Finally, consider the image $\Phi(\text{PA}(2, p))$. For a map $f \in \text{PA}(2, p)$, the fraction $\Phi(f)$ is reduced and has denominator $p - 2$ since $\rho(f) = \rho_2(p - 1, k)$ for some k such that

$k - 1$ is coprime to $p - 2$. The fact that $\Phi(\text{PA}(2, p))$ contains all such fractions follows because Φ is a bijection onto $\mathbb{Q} \cap (0, 1)$.

A similar argument works for $m \geq 3$ odd and $m \geq 4$ even. In this case, the placement of infinitesimal loops as in steps (1)–(3) above is chosen accordingly to ensure that τ is invariant under the exterior left-veering thickening. ■

7. The digit polynomial D_f

In this section, we prove Theorem 3, restated below for convenience. This theorem generalizes [6, Lemma 2.5], which treats the unimodal case. In [6], the result is phrased in terms of the kneading sequence of the unique critical point.

Theorem 3. *Suppose $f \in \text{PA}(m)$ for $m \geq 2$ with $\Phi(f) = a/b \in \mathbb{Q} \cap (0, 1)$ in lowest terms. Define $L: [0, b] \rightarrow \mathbb{R}$ by $L(t) = (a/b) \cdot t$. Then*

$$D_f(t) = t^{b+1} + 1 - \sum_{i=1}^b c_i t^{b+1-i},$$

where the c_i satisfy

$$c_i = \begin{cases} m & \text{if } L(t) \in \mathbb{N} \text{ for some } t \in [i-1, i], \\ m-2 & \text{otherwise.} \end{cases}$$

In particular, $c_i = c_{b-i}$, so D_f is reciprocal, that is,

$$D_f(t) = t^{b+1} D_f(t^{-1}).$$

Example 7.1. Let $f, g \in \text{PA}(2)$ be the zig-zags such that $\Phi(f) = 1/7$ and $\Phi(g) = 6/7$. According to Theorem 3 and Figure 21, we have

$$\begin{aligned} D_f(t) &= t^8 - 2t^7 - 0t^6 - 0t^5 - 0t^4 - 0t^3 - 0t^2 - 2t + 1, \\ D_g(t) &= t^8 - 2t^7 - 2t^6 - 2t^5 - 2t^4 - 2t^3 - 2t^2 - 2t + 1. \end{aligned}$$

Example 7.2. Let $p, q \in \text{PA}(7)$ be the zig-zags such that $\Phi(p) = 4/13$ and $\Phi(q) = 9/13$. According to Theorem 3 and Figure 22, we have

$$\begin{aligned} D_p(t) &= t^{14} - 7t^{13} - 5t^{12} - 5t^{11} - 7t^{10} - 5t^9 - 5t^8 - 7t^7 - 5t^6 - 5t^5 \\ &\quad - 7t^4 - 5t^3 - 5t^2 - 7t + 1, \\ D_q(t) &= t^{14} - 7t^{13} - 7t^{12} - 7t^{11} - 5t^{10} - 7t^9 - 7t^8 - 5t^7 - 7t^6 - 7t^5 \\ &\quad - 5t^4 - 7t^3 - 7t^2 - 7t + 1. \end{aligned}$$

We first prove Theorem 3 for the case that f has modality $m \geq 4$ even. This case being completed, we will then use it to deduce the cases for $m \geq 3$ odd and $m = 2$. We proceed via a sequence of lemmas.




Figure 21. Computing the digit polynomials of bimodal maps using Theorem 3. (a) Computing the coefficients of D_f for the unique $f \in \text{PA}(2)$ such that $\Phi(f) = 1/7$. (b) Computing the coefficients of D_g for the unique $g \in \text{PA}(2)$ such that $\Phi(g) = 6/7$.




Figure 22. Computing the digit polynomials of 7-modal maps using Theorem 3. (a) Computing the coefficients of D_p for the unique $p \in \text{PA}(7)$ such that $\Phi(p) = 4/13$. (b) Computing the coefficients of D_q for the unique $q \in \text{PA}(7)$ such that $\Phi(q) = 9/13$.

Lemma 7.3. Suppose $f \in \text{PA}(m)$ for $m \geq 4$ even. Let n be minimal such that $f^n(1) = 1$, and let $f_i(x)$ be the defining linear branch of f at $f^{i-1}(1)$, that is, $f^i(1) = f_i(f^{i-1}(1))$. Then there exist constants c_i for $i = 1, \dots, n-2$ such that

$$f_i(x) = \begin{cases} \lambda x - c_i, & i = 1, \dots, n-2, \\ m - \lambda x, & i = n-1, \\ 2 - \lambda x, & i = n. \end{cases}$$

Proof. The sole point of type R (cf. Proposition 5.10) in the orbit of $x = 1$ must map to the point of type C , by Proposition 5.5, and this latter point is precisely $x = \lambda^{-1}$, and hence maps to $x = 1$ by $f_n(x) = 2 - \lambda x$. At the point of type R , we have $f_{n-1}(x) = m - \lambda x$.

All other points in the orbit of $x = 1$ are of type P_1 or P_2 , and the corresponding f_i are $f_i(x) = \lambda x - (m-2)$ and $f_i(x) = \lambda x - m$, respectively. ■

Lemma 7.4. Suppose $f \in \text{PA}(m)$ for $m \geq 4$ even. Let $D_f(t)$ be the digit polynomial of f . Then

$$D_f(t) = t^n + 1 - \sum_{i=1}^{n-1} c_i t^{n-i},$$

where, as in Lemma 7.3, the c_i are defined by $f_i(x) = \lambda x - c_i$ for $1 \leq i \leq n-2$ and $f_{n-1}(x) = c_{n-1} - \lambda x$. In particular, $c_1 = c_{n-1} = m$.

Proof. For $i = 0, \dots, n$, define $g_i(t) \in \mathbb{Z}[t]$ by $g_i(\lambda) = f^i(\lambda)$. Thus, for example, $g_0(\lambda) = 1$ and $g_1(\lambda) = \lambda - m$. Here it is important that we are treating λ as a formal variable, rather than an algebraic integer satisfying a polynomial relation.

Observe that $g_{i+1}(\lambda) = f_{i+1}(g_i(\lambda))$ by definition. In particular, by Lemma 7.3 for $i = 0, \dots, n-3$ we have $g_{i+1}(\lambda) = \lambda g_i(\lambda) - c_i$, and so inductively we see that

$$g_i(\lambda) = \lambda^i - c_1 \lambda^{i-1} - \dots - c_{i-1} \lambda - c_i \quad \text{for } 0 \leq i \leq n-2.$$

We therefore have the equalities

$$g_{n-1}(\lambda) = c_{n-1} - \lambda^{n-1} + c_1 \lambda^{n-2} + \dots + c_{n-3} \lambda^2 + c_{n-2} \lambda$$

and

$$g_n(\lambda) = 2 - c_{n-1} \lambda - c_{n-2} \lambda^2 - \dots - c_1 \lambda^{n-1} + \lambda^n.$$

By the definition of $D_f(t)$ and the fact that $g_n(\lambda) = f^n(1) = 1$, we have

$$D_f(\lambda) = g_n(\lambda) - 1. \quad \blacksquare$$

Lemma 7.5. Suppose $f \in \text{PA}(m)$ for $m \geq 4$ even. Let n be minimal such that $f^n(1) = 1$ and suppose that f acts on the orbit of $x = 1$ by the permutation $\rho(f) = \rho_e(n, k)$ with $\gcd(n-k, k-1) = 1$. Let $D_f(t) = t^n + 1 - \sum_{i=1}^{n-1} c_i t^{n-i}$. Then for $i = 1, \dots, n-1$, we have

$$c_i = \begin{cases} m & \text{if } \rho(f)^{i-1}(n) \geq k, \\ m-2 & \text{if } 2 \leq \rho(f)^{i-1}(n) \leq k-1. \end{cases}$$

Proof. The permutation $\rho(f)$ is defined such that postcritical points corresponding to symbols i satisfying $k+1 \leq i \leq n$ are of type P_2 , whereas those corresponding to i satisfying $2 \leq i \leq k-1$ are of type P_1 . Finally, the point corresponding to the symbol k is of type R , specifically $f^{n-2}(1)$. The result follows from Lemmas 7.3 and 7.4. $\blacksquare$

Lemma 7.6. Theorem 3 holds for $f \in \text{PA}(m)$ with $m \geq 4$ even.

Proof. We have already shown that $c_0 = c_n = 1$ and that $c_i \in \{m-2, m\}$ for $1 \leq i \leq n-1$. By Lemma 7.5, it is enough to understand when $\rho^i(n) \geq k$. Let $\rho' \in S_{n-1}$ be the permutation obtained by deleting the symbol 1 from the cycle decomposition of ρ and decreasing all remaining labels by 1, as in the proof of Proposition 6.7. As observed

previously, ρ' acts on $\{1, \dots, n-1\}$ as addition by $n-k$ modulo $n-1$. Therefore, if we set $q = (n-k)/(n-1)$, the values $L_q(t) = qt$ for $t = 1, \dots, n-1$ satisfy

$$L_q(t) - \lfloor L_q(t) \rfloor = \frac{(\rho')^t(n-1)}{n-1}.$$

To see why this is true, observe first that the left-hand side is the fractional part of $L_q(t)$. This quantity is a rational number with denominator $n-1$, and the numerator increases by $n-k$ modulo $n-1$. Since $(\rho')^t(n-1)$ also changes in this way, it remains to note that

$$L_q(1) = \frac{n-k}{n-1} = \frac{(\rho')(n-1)}{n-1}.$$

Now we observe that

$$\begin{aligned} c_i = m &\Leftrightarrow \rho^{i-1}(n) \geq k \\ &\Leftrightarrow (\rho')^{i-1}(n-1) \geq k-1 \\ &\Leftrightarrow L_q(i-1) - \lfloor L_q(i-1) \rfloor \geq \frac{k-1}{n-1} \\ &\Leftrightarrow L_q(t) \in \mathbb{N} \text{ for some } t \in [i-1, i]. \end{aligned}$$

The last equivalence holds because $L_q(t) = qt$ is a line of slope $(n-k)/(n-1)$. The proof is complete. ■

It remains to prove Theorem 3 for the case when the modality of f is $m = 2$ or $m \geq 3$ odd. Recall that

$$\rho_2(n, k) = \kappa^{-1}(n, k) \circ \rho_e(n, k) \circ \kappa(n, k),$$

where $\kappa(n, k) = (1, 2, \dots, k-1) \in S_n$.

Lemma 7.7. *For any $l \geq 0$, $\rho_2^l(n) \geq k$ if and only if $\rho_e^l(n) \geq k$, and in this case, $\rho_2^l(n) = \rho_e^l(n)$.*

Proof. Suppose $\rho_e^l(n) \geq k$. Then $\rho_2^l(n) = \kappa^{-1} \rho_e^l \kappa(n) = \rho_e^l(n)$, since $\kappa(j) = j$ for all $j \geq k$. Similarly, if $\rho_2^l(n) \geq k$, then it follows that $\rho_e^l(n) \geq k$ as well. ■

We now complete the proof of Theorem 3.

Proof of Theorem 3. One quickly verifies that Lemmas 7.3 through 7.5 hold for f an m -modal zig-zag of pseudo-Anosov type for $m = 2$ and $m \geq 3$ odd, after replacing all instances of $\rho_e(n, k)$ with either $\rho_2(n, k)$ or $\rho_o(n, k)$. Lemmas 7.5 and 7.7 now imply the Theorem for $m = 2$.

For $m \geq 3$ odd, recall that if we delete the symbol 0 from the cycle type of $\rho_o(n, k)$, we obtain the permutation $\rho_e(n, k)$. Deleting this symbol corresponds to ignoring the linear map $f_{n+1}(x) = 1 - \lambda x$. This linear branch is not used to compute D_f , since the fact that $f \in \text{PA}(m)$ for $m \geq 3$ odd implies that $f^n(1) = 0$, terminating the process of constructing D_f before the $(n+1)$ -st step (cf. Definition 1.8).

The arguments of Lemmas 7.3 through 7.6 now prove the theorem in this case. ■

8. A family of pseudo-Anosovs with Salem dilatation

In this section, we provide an application of the theory developed over the course of this paper. Recall that a *Salem number*, introduced in [11], is a real algebraic integer $\lambda > 1$ such that all Galois conjugates of λ are contained within the closed unit disc, with at least one of these conjugate lying on the unit circle. It is not hard to show that λ^{-1} must be among the Galois conjugates of λ in this case, and that all other conjugates lie on the unit circle. In particular, a Salem number is a Perron number of even degree $d = 2g$. If $p(x) \in \mathbb{Z}[x]$ is the minimal polynomial of a Salem number, then $p(x)$ is *reciprocal*, i.e.,

$$p_*(x) := x^{\deg(p)} p(x^{-1}) = p(x).$$

It is well known that if $f(x) \in \mathbb{Z}[x]$ is a reciprocal polynomial of degree $d = 2g$, then $f(x) = x^g q(x + x^{-1})$ for some integral polynomial q . We call $q(x)$ the *companion polynomial* to $f(x)$. If $\deg(f) = 2g + 1$, then $f(x) = (x + 1)f_1(x)$ for $f_1(x)$ reciprocal of even degree, and therefore $f(x) = (x + 1)x^g q(x + x^{-1})$ for some $q(x) \in \mathbb{Z}[x]$. In this case, we again call q the companion polynomial of f .

Note that there is a bijection between roots of q and pairs of roots of f : if λ, λ^{-1} are roots of f , then $\lambda + \lambda^{-1}$ is a root of q . Moreover, if $|\lambda| = 1$, then $\lambda^{-1} = \bar{\lambda}$, and so the root $\lambda + \lambda^{-1}$ of q is a real number contained in the interval $[-2, 2]$. We refer to this interval, with or without its endpoints, as the *critical interval*.

In the case of a Salem number λ of degree $2g$, the companion polynomial is irreducible of degree g with dominant root $\lambda + \lambda^{-1} > 2$ and the remaining $g - 1$ roots in the critical interval. In particular, the companion polynomial has all real roots, so $\lambda + \lambda^{-1}$ is a totally real algebraic integer of degree g .

Recall that a *translation surface* is a pair (X, ω) of a Riemann surface X equipped with a non-zero abelian differential ω , i.e., a holomorphic one-form. If X is of genus g , then ω has $2g - 2$ zeros, counting multiplicity. Let Σ be the collection of these zeros. Then away from Σ , X has a Euclidean structure: in other words, $X \setminus \Sigma$ admits an atlas of charts to $\mathbb{C}$ whose transition functions are translations. In the neighborhood of a zero p of order k , X has the structure of $2(k + 1)$ metric half-discs glued together, so that the total angle about p is $2\pi(k + 1)$.

Fixing $g > 1$, let μ be a positive integer partition of $2g - 2$. We think of μ as describing the multiplicities of the zeros of an abelian differential ω on X . The *stratum* $\mathcal{H}(\mu)$ is the collection of genus g translations surfaces with zero orders specified by μ .

Given some $(X, \omega) \in \mathcal{H}(\mu)$ and $A \in \mathrm{SL}_2(\mathbb{R})$, define $A \cdot (X, \omega)$ to be the translation surface obtained by post-composing the charts of (X, ω) into $\mathbb{R}^2 \cong \mathbb{C}$ by A .

The *Veech group* of a translation surface (X, ω) , denoted $\mathrm{SL}(X, \omega)$, is the stabilizer of (X, ω) under the action by $\mathrm{SL}_2(\mathbb{R})$. The *trace field* of (X, ω) is the field K obtained by adjoining to $\mathbb{Q}$ the traces of all elements of $\mathrm{SL}(X, \omega)$. By a result of Möller in [9], the degree of the trace field satisfies

$$[K : \mathbb{Q}] \leq g(X).$$

If the degree of this extension is maximal, i.e., is equal to the genus of X , then we say that (X, ω) is *algebraically primitive*. One remarks that such surfaces cannot arise as covers of translation surfaces of lower genus: if $\pi: (X, \omega) \rightarrow (Y, \eta)$ is a translation cover, then the trace fields of X and Y coincide, and the result of Möller mentioned above now shows that if X is algebraically primitive, then $g(X) = g(Y)$.

A translation surface is called *Veech* if its Veech group is a lattice in $\mathrm{SL}_2(\mathbb{R})$, i.e., as large as possible. The $\mathrm{GL}_2(\mathbb{R})$ -orbit of a Veech surface is called a *Teichmüller curve*. By results of Möller [8] and Apisa [1], there are only finitely many algebraically primitive Teichmüller curves in any genus $g \geq 3$. Therefore, it is interesting to find algebraically primitive surfaces with non-trivial Veech group of arbitrarily high genus.

We now restate our last main result.

Theorem 4. *For each $g \geq 1$, define $f_g: I \rightarrow I$ to be the bimodal PCP zig-zag map of pseudo-Anosov type corresponding to $r_g = 1/(2g) \in \mathbb{Q} \cap (0, 1)$. Let λ_g be the growth rate of f_g . Then the following are true for each $g \geq 1$:*

- (1) λ_g is a Salem number of degree $2g$.
- (2) The pseudo-Anosov ϕ_g obtained from f_g is defined on a $(2g + 2)$ -punctured sphere $\Sigma_{0,2g+2}$.
- (3) The translation surface (X_g, ω_g) obtained as the hyperelliptic double cover of $\Sigma_{0,2g+2}$ is of genus g , and hence algebraically primitive.

Remark 8.1. One might reasonably object that there is no such thing as a Salem number of degree two. Indeed, Salem numbers are normally defined to have at least one Galois conjugate on the unit circle, in which case all Salem numbers must be of degree at least four: if λ is a Salem number and $\alpha \in S^1$ a Galois conjugate, then $\bar{\alpha} = \alpha^{-1}$ is also a conjugate of λ , and hence so is λ^{-1} . However, one may think of the quadratic units $\lambda > 1$ as degenerate Salem numbers. That is what we choose to do here.

8.1. λ_g is Salem of degree $2g$

As before, the proof of Theorem 4 will proceed in a sequence of lemmas. The bulk of our efforts will be focused on establishing statement (1) of Theorem 4.

Lemma 8.2. *Let $D_g(t)$ be the digit polynomial of f_g . Then for all $g \geq 1$, we have*

$$D_g(t) = t^{2g+1} - 2t^{2g} - 2t + 1 = (t + 1)d_g(t),$$

where

$$d_g(t) = t^{2g} + 1 + 3 \sum_{i=1}^{2g-1} (-1)^i t^i.$$

Thus, for example,

$$D_1(t) = (t + 1)(t^2 - 3t + 1) \quad \text{and} \quad D_2(t) = (t + 1)(t^4 - 3t^3 + 3t^2 - 3t + 1).$$

Proof of Lemma 8.2. The fact that $D_g(t) = t^{2g+1} - 2t^g - 2t + 1$ follows from Theorem 3. One readily checks that $(t + 1)d_g(t) = D_g(t)$. ■

We wish to show that $d_g(t)$ is the minimal polynomial of a Salem number. To do this, we must prove that $d_g(t)$ has $2g - 2$ roots on the unit circle, and also that the polynomial is irreducible. As we have seen, to accomplish the first task it will be enough to show that the companion polynomial $q_g(t)$, defined by $d_g(t) = t^g q_g(t + t^{-1})$, has $g - 1$ roots in the critical interval. This is the content of the next two lemmas.

Lemma 8.3. *The companion polynomials $q_g(w)$ satisfy the following properties:*

- (1) *For all $g \geq 4$, we have the recurrence relation*

$$q_{g+2}(w) = wq_{g+1}(w) - q_g(w). \quad (8.1)$$

- (2) *$q_g(2) = -1$ for all n .*

- (3) *Each q_g has a real root $\alpha_g > 2$.*

- (4) *For all $g \geq 2$, we have $(-1)^g q_g(-2) > 0$. In particular, $q_g(-2)$ and $q_{g+1}(-2)$ have opposite signs.*

The first few companion polynomials q_g are the following:

$$\begin{aligned} q_2(w) &= w^2 - 3w + 1, \\ q_3(w) &= w^3 - 3w^2 + 3, \\ q_4(w) &= w^4 - 3w^3 - w^2 + 6w - 1, \\ q_5(w) &= w^5 - 3w^4 - 2w^3 + 9w^2 - w - 3. \end{aligned}$$

Proof of Lemma 8.3. By definition, q_g satisfies $q_g(t + t^{-1}) = t^{-g} d_g(t)$, so relation (8.1) is equivalent to the recurrence

$$d_{g+2}(t) = t^2[d_{g+1}(t) - d_g(t)] + d_{g+1}(t).$$

This formula is a straightforward consequence of the pattern of the coefficients of d_g , proving statement (1). Statement (2) follows inductively after noting that it holds for q_2 and q_3 . Now note q_2 and q_3 are monic, so by equation (8.1) q_g is monic for all g . In particular, $\lim_{w \rightarrow \infty} q_g(w) = \infty$ for all g , so the intermediate value theorem and statement (2) together imply statement (3).

Finally, to prove statement (4), observe that

$$d_g(-1) = 1 + 3(2g - 1) + 1 = 6g - 1 > 0$$

for all $g \geq 2$. Since $d_g(-1) = (-1)^g q_g(-2)$, the result follows. ■

Lemma 8.4. *For each $g \geq 2$, the polynomial q_g has $g - 1$ roots in $(-2, 2)$. Moreover, if these roots are denoted $-2 < a_1 < \dots < a_{g-1} < 2$, then the g roots of q_{g+1} in $(-2, 2)$, denoted $b_1, \dots, b_g$, satisfy the ordering*

$$-2 < b_1 < a_1 < b_2 < a_2 < \dots < b_{g-1} < a_{g-1} < b_g < 2.$$




Figure 23. The interlacing property of the roots of the companion polynomials q_g .

Thus, the roots of q_{g+1} “interlace” the roots of q_g . See Figure 23.

Proof of Lemma 8.4. We proceed inductively. The quadratic polynomial q_2 has a single root $a_1 = (3 - \sqrt{5})/2$ in $(-2, 2)$. By Lemma 8.3 (3), this is in fact the only root of q_2 in the critical interval. Since $q_3(-1) = -1$, $q_3(0) = 3$, $q_3(1) = 1$, and $q_3(2) = -1$, we see that q_3 has two roots $b_1, b_2 \in (-2, 2)$ satisfying $-1 < b_2 < 0$ and $1 < b_2 < 2$. Since $0 < a_1 < 1$, the claim is satisfied in this case.

Suppose now that the claim holds for all $n \leq g + 1$. We may assume without loss of generality that $q_{g+1}(-2) < 0$: the other case is essentially identical. Lemma 8.3 (4) implies that both q_g and q_{g+2} are positive at $w = -2$. Let $b_1 < \dots < b_g$ be the roots of q_{g+1} in the critical interval. Then we have

$$q_{g+2}(b_1) = b_1 q_{g+1}(b_1) - q_g(b_1) = -q_g(b_1).$$

By assumption, the smallest root a_1 of q_g in $(-2, 2)$ is greater than b_1 . Since $q_g(-2) > 0$, it follows that $q_g(b_1) > 0$ and thus $q_{g+2}(b_1) < 0$. Therefore, q_{g+2} has a root $c_1 \in (-2, b_1)$. Next we observe that

$$q_{g+2}(b_2) = b_2 q_{g+1}(b_2) - q_g(b_2) = -q_g(b_2).$$

Since $a_1 < b_2 < a_2$, we see that $q_g(b_2) < 0$, so $q_{g+2}(b_2) > 0$, implying that q_{g+2} has a root $c_2 \in (b_1, b_2)$. Continuing in this fashion, we find roots $c_i \in (b_{i-1}, b_i)$ for $2 \leq i \leq g$. Finally, note that since $q_g(2) = -1$, we have $q_g(w) < 0$ for all $w \in (a_{g-1}, 2]$. Therefore, since $a_{g-1} < b_g < 2$, we have

$$q_{g+2}(b_g) = -q_g(b_g) > 0.$$

Since $q_{g+2}(2) = -1$, it follows that q_{g+2} has a root $c_{g+1} \in (b_g, 2)$. Lemma 8.3 (4) implies that q_{g+2} has another root $\alpha_{g+2} > 2$, so there cannot be any other roots of q_{g+2} . The proof is complete. $\blacksquare$

Lemma 8.4 implies that d_g has $g - 1$ pairs of roots on the unit circle in addition to a pair of positive real roots λ_g and λ_g^{-1} . This is already enough to conclude that λ_g is a Salem number. Indeed, λ_g must be Galois conjugate to λ_g^{-1} , since otherwise the conjugates of λ_g^{-1} would be contained in the closed unit disc, implying that they are all roots of unity, by Kronecker's theorem. This last is impossible, since $\lambda_g^{-1} < 1$. Hence λ_g and λ_g^{-1} are Galois conjugate and λ_g is a Salem number. It remains to determine whether d_g is irreducible.

We follow a similar proof given by Shin in [12]. Interestingly, this proof shows that the “dual” Perron roots ψ_g corresponding to the fraction

$$\tilde{r}_g = 1 - r_g = \frac{2g - 1}{2g}$$

are also Salem of degree $2g$.

Lemma 8.5. *Each $d_g(t)$ is irreducible over $\mathbb{Z}[t]$. Consequently, λ_g is a Salem number of degree $2g$ for all $g \geq 1$.*

Proof. Since λ_g is necessarily Galois conjugate to λ_g^{-1} , any factor of $d_g(t)$ other than the minimal polynomial of λ_g must have all roots on the unit circle, and therefore must be cyclotomic by Kronecker's theorem. Suppose therefore that $e^{2\pi i/m}$ is a root of $d_g(t)$. Then we have

$$D_g(e^{2\pi i/m}) = e^{(2g+1) \cdot 2\pi i/m} - 2e^{2g \cdot 2\pi i/m} - 2e^{2\pi i/m} + 1 = 0.$$

We take real and imaginary parts to obtain the system of equations

$$\begin{cases} \cos\left(\frac{(2g+1)2\pi}{m}\right) - 2\cos\left(\frac{2g \cdot 2\pi}{m}\right) - 2\cos\left(\frac{2\pi}{m}\right) + 1 = 0, \\ \sin\left(\frac{(2g+1)2\pi}{m}\right) - 2\sin\left(\frac{2g \cdot 2\pi}{m}\right) - 2\sin\left(\frac{2\pi}{m}\right) = 0. \end{cases} \quad (8.2)$$

For the first equation, we use the formula $\cos(2x) = 2\cos^2(x) - 1$ for the first cosine term and the formula $\cos(a) + \cos(b) = 2\cos((a+b)/2)\cos((a-b)/2)$ for the latter two terms to obtain

$$\cos\left(\frac{(2g+1)\pi}{m}\right) \left[\cos\left(\frac{(2g+1)\pi}{m}\right) - 2\cos\left(\frac{(2g-1)\pi}{m}\right) \right] = 0.$$

Similarly, for the second equation in (8.2), we use the formula $\sin(2x) = 2\sin(x)\cos(x)$ and the formula $\sin(a) + \sin(b) = 2\sin((a+b)/2)\cos((a-b)/2)$ to find

$$\sin\left(\frac{(2g+1)\pi}{m}\right) \left[\cos\left(\frac{(2g+1)\pi}{m}\right) - 2\cos\left(\frac{(2g-1)\pi}{m}\right) \right] = 0.$$

Since $\sin(x)$ and $\cos(x)$ do not have any common roots, it must be the case that

$$\cos\left(\frac{(2g+1)\pi}{m}\right) - 2\cos\left(\frac{(2g-1)\pi}{m}\right) = 0.$$

Setting $\varphi = (2g - 1)\pi/m$, we rewrite this as

$$\cos\left(\varphi + \frac{2\pi}{m}\right) - 2\cos(\varphi) = 0. \quad (8.3)$$

It follows that $-1/2 \leq \cos(\varphi) \leq 1/2$. In other words, as $\cos(\pi/3) = 1/2$ and $\cos(2\pi/3) = -1/2$, we must have

$$-\frac{2\pi}{3} \leq \varphi \leq -\frac{\pi}{3} \quad \text{or} \quad \frac{\pi}{3} \leq \varphi \leq \frac{2\pi}{3}.$$

Moreover, by equation (8.3), φ and $\varphi + 2\pi/m$ are angles on the same side of the y -axis, since their cosines have the same sign. We claim that φ must be in either the second or fourth quadrant. Suppose for contradiction that φ is in the first quadrant, so that $\pi/3 \leq \varphi \leq \pi/2$. We may assume $m \geq 3$, since $x = \pm 1$ are clearly not roots of d_g , and thus $2\pi/m < \pi$. In particular, $\varphi + 2\pi/m < \pi/2 + \pi = 3\pi/2$, hence cannot be in the fourth quadrant.

Since both φ and $\varphi + 2\pi/m$ are on the same side of the y -axis and we assumed φ is in the first quadrant, it follows that $\pi/3 \leq \varphi + 2\pi/m \leq \pi/2$ as well. Since $\cos(x)$ is decreasing on this interval, we have

$$\cos(\varphi) > \cos\left(\varphi + \frac{2\pi}{m}\right) > 0 \Rightarrow 2\cos(\varphi) > \cos\left(\varphi + \frac{2\pi}{m}\right),$$

contradicting equation (8.3). A similar argument shows that φ cannot be in the third quadrant. Thus we revise the restrictions on φ to be

$$-\frac{\pi}{2} < \varphi \leq -\frac{\pi}{3} \quad \text{or} \quad \frac{\pi}{2} < \varphi \leq \frac{2\pi}{3}.$$

Appealing to the formula $\cos(\theta) = \sin(\theta + \pi/2)$, we may equivalently consider the equation $\sin(\psi + 2\pi/m) - 2\sin(\psi) = 0$ with

$$0 < \psi \leq \frac{\pi}{6} \quad \text{or} \quad \pi < \psi \leq \frac{7\pi}{6}.$$

Suppose ψ is in the first quadrant and write

$$\psi = \varphi + \frac{\pi}{2} = \frac{(2g-1)\pi}{m} + \frac{\pi}{2} \equiv \frac{j\pi}{2m} \pmod{2\pi}$$

for some positive integer $j \leq 2m-1$ such that $0 < j\pi/(2m) \leq \pi/6$. Using the subadditivity of $\sin(x)$ on $[0, \pi]$ now gives

$$\sin\left(\psi + \frac{2\pi}{m}\right) - 2\sin(\psi) \leq \sin(\psi) + \sin\left(\frac{2\pi}{m}\right) - 2\sin(\psi) = \sin\left(\frac{2\pi}{m}\right) - \sin\left(\frac{j\pi}{2m}\right).$$

This expression cannot be zero unless $j = 4$ because of the restriction on j . In this case,

$$\begin{aligned} \sin\left(\psi + \frac{2\pi}{m}\right) - 2\sin(\psi) &= \sin\left(\frac{4\pi}{m}\right) - 2\sin\left(\frac{2\pi}{m}\right) \\ &= 2\sin\left(\frac{2\pi}{m}\right)\cos\left(\frac{2\pi}{m}\right) - 2\sin\left(\frac{2\pi}{m}\right) \\ &= 2\sin\left(\frac{2\pi}{m}\right)\left[\cos\left(\frac{2\pi}{m}\right) - 1\right]. \end{aligned}$$

This quantity can only be zero if $m = 1$, which we quickly rule out after observing that $d_g(1) = 6g - 1$. The same argument works if we assume ψ is in the third quadrant. Therefore, $d_g(t)$ has no cyclotomic factor, and hence is irreducible. ■

8.2. Completing the proof of Theorem 4

Proof of Theorem 4. Lemma 8.5 proves statement (1). Since the λ_g -uniform expander f_g corresponds to the fraction $q_g = 1/(2g)$, the point $x = 1$ is periodic of length $2g + 1$. Each of these points lifts to a one-pronged singularity of the pseudo-Anosov ϕ_g on a punctured sphere, as does the fixed postcritical point $x = 0$. There are no other one-pronged singularities, so taking the double cover of the surface by branching at each of these $2g + 2$ points produces a surface on which the lift of each point is a flat point, i.e., has cone angle 2π . The only other cone point downstairs is the fixed point at infinity, with cone angle $2g \cdot \pi$. This point lifts to two points of angle $g \cdot 2\pi$.

The derivative of the lifted pseudo-Anosov is

$$D\widetilde{\phi}_g = \begin{pmatrix} \lambda_g & 0 \\ 0 & \lambda_g^{-1} \end{pmatrix},$$

an element of the Veech group $\mathrm{SL}(X_g, \omega_g)$. The trace of this matrix is $\lambda_g + \lambda_g^{-1}$, contained in the trace field K_g by definition. But now

$$g \geq [K_g : \mathbb{Q}] \geq [\mathbb{Q}(\lambda_g + \lambda_g^{-1}) : \mathbb{Q}] = g,$$

so in fact we have $[K_g : \mathbb{Q}] = g$, implying that (X_g, ω_g) is algebraically primitive. ■

Figures 24, 25, and 26 display the limit sets for the Galois lifts of the zig-zags f_1 , f_2 , and f_3 , respectively.




Figure 24. The limit set of the Galois lift of f_1 . This glues to a sphere with 4 marked points, which then lifts to a torus. The lift of the pseudo-Anosov ϕ_1 is a linear Anosov diffeomorphism of the torus with stretch factor $\lambda_1 = (3 + \sqrt{5})/2$, hence is conjugate to Arnold's cat map.




Figure 25. The limit set of the Galois lift of f_2 . This glues to a sphere with 6 cone points of angle π , one at the center of each vertical edge. Taking the double cover of this sphere branched at the 6 cone points produces a genus 2 surface.




Figure 26. The limit set of the Galois lift of f_3 . This glues to a sphere with 8 marked points, which then lifts to a genus 3 surface.

Acknowledgments. I gratefully acknowledge André de Carvalho and Hyungryul Baik for their helpful and insightful comments. I am also particularly grateful to my advisor Kathryn Lindsey for introducing me to this subject and participating in many extremely helpful conversations.

References

- [1] P. Apisa, [GL₂ \$\mathbb{R}\$ orbit closures in hyperelliptic components of strata](#). *Duke Math. J.* **167** (2018), no. 4, 679–742 Zbl [1436.32053](#) MR [3769676](#)
- [2] H. Baik, A. Rafiqi, and C. Wu, [Constructing pseudo-Anosov maps with given dilatations](#). *Geom. Dedicata* **180** (2016), 39–48 Zbl [1364.37094](#) MR [3451455](#)

- [3] A. de Carvalho, [Extensions, quotients and generalized pseudo-Anosov maps](#). In *Graphs and patterns in mathematics and theoretical physics*, pp. 315–338, Proc. Sympos. Pure Math. 73, American Mathematical Society, Providence, RI, 2005 Zbl [1330.37045](#) MR [2131019](#)
- [4] A. de Carvalho and T. Hall, [Unimodal generalized pseudo-Anosov maps](#). *Geom. Topol.* **8** (2004), 1127–1188 Zbl [1057.37041](#) MR [2087080](#)
- [5] D. Fried, [Growth rate of surface homeomorphisms and flow equivalence](#). *Ergodic Theory Dynam. Systems* **5** (1985), no. 4, 539–563 Zbl [0603.58020](#) MR [829857](#)
- [6] T. Hall, [The creation of horseshoes](#). *Nonlinearity* **7** (1994), no. 3, 861–924 Zbl [0806.58015](#) MR [1275533](#)
- [7] J. Milnor and W. Thurston, [On iterated maps of the interval](#). In *Dynamical systems (College Park, MD, 1986–87)*, pp. 465–563, Lecture Notes in Math. 1342, Springer, Berlin, 1988 Zbl [0664.58015](#) MR [970571](#)
- [8] M. Möller, [Finiteness results for Teichmüller curves](#). *Ann. Inst. Fourier (Grenoble)* **58** (2008), no. 1, 63–83 Zbl [1140.14010](#) MR [2401216](#)
- [9] M. Möller, [Teichmüller curves, mainly from the viewpoint of algebraic geometry](#). In *Moduli spaces of Riemann surfaces*, pp. 267–318, IAS/Park City Math. Ser. 20, American Mathematical Society, Providence, RI, 2013 Zbl [1279.14031](#) MR [3114688](#)
- [10] W. Parry, [Intrinsic Markov chains](#). *Trans. Amer. Math. Soc.* **112** (1964), 55–66 Zbl [0127.35301](#) MR [161372](#)
- [11] R. Salem, *Algebraic numbers and Fourier analysis*. Wadsworth Math. Ser., Wadsworth, Belmont, CA, 1983 Zbl [0505.00033](#) MR [732447](#)
- [12] H. Shin, [Algebraic degrees of stretch factors in mapping class groups](#). *Algebr. Geom. Topol.* **16** (2016), no. 3, 1567–1584 Zbl [1356.57017](#) MR [3523049](#)
- [13] D. J. Thompson, [Generalized \$\beta\$ -transformations and the entropy of unimodal maps](#). *Comment. Math. Helv.* **92** (2017), no. 4, 777–800 Zbl [1380.37080](#) MR [3718487](#)
- [14] W. P. Thurston, Entropy in dimension one. In *Frontiers in complex dynamics*, pp. 339–384, Princeton Math. Ser. 51, Princeton University Press, Princeton, NJ, 2014 Zbl [1408.37031](#) MR [3289916](#)
- [15] K. M. Wilkinson, [Ergodic properties of a class of piecewise linear transformations](#). *Z. Wahrscheinlichkeitstheorie Verw. Geb.* **31** (1975), 303–328 Zbl [0299.28015](#) MR [374390](#)

Received 13 December 2021.

Ethan Farber

Department of Mathematics, Boston College, Maloney Hall, MA 02467-3806 Chestnut Hill, USA;
farbereth@gmail.com