

A recursive presentation for Mihailova's subgroup

Oleg Bogopolski and Enric Ventura

Abstract. An explicit recursive presentation for Mihailova's subgroup $M(H)$ of $F_n \times F_n$ corresponding to a finite, concise and Peiffer aspherical presentation $H = \langle x_1, \dots, x_n \mid R_1, \dots, R_m \rangle$ is given. This partially answers a question of R. I. Grigorchuk. As a corollary, we construct a finitely generated recursively presented orbit undecidable subgroup of $\text{Aut}(F_3)$.

Mathematics Subject Classification (2010). 20F05, 20F10.

Keywords. Mihailova's subgroup, recursive presentation, asphericity, orbit undecidability.

1. Introduction

For all the paper, let $n \geq 2$, let F_n be the free group with basis $\{x_1, \dots, x_n\}$, and let $H = \langle x_1, \dots, x_n \mid R_1, \dots, R_m \rangle$ be a finite presentation of a quotient H of F_n (although most of what follows will depend on the specific presentation, we shall make the usual abuse of notation which consists on denoting by H both the group and its given presentation).

K. A. Mihailova, in her influential paper [12], associated to the presentation H the *Mihailova subgroup* of $F_n \times F_n$, namely

$$M(H) = \{(w_1, w_2) \in F_n \times F_n \mid w_1 =_H w_2\} \leq F_n \times F_n,$$

i.e., the subgroup of pairs of words in F_n determining the same element in H . It is clear that (x_i, x_i) and $(1, R_j)$ belong to $M(H)$ for all $i = 1, \dots, n$ and $j = 1, \dots, m$, and it is not difficult to see that, in fact, these pairs generate $M(H)$. The important observation made in [12] says that the membership problem for $M(H)$ in $F_n \times F_n$ is solvable (i.e., there exists an algorithm to decide whether a given $(w_1, w_2) \in F_n \times F_n$ belongs to $M(H)$ or not) if and only if the word problem for H is solvable.

By a result of P. S. Novikov [14] and W. W. Boone [3] (see also [4]), there exist finitely presented groups with unsolvable word problem. Thus, there also exist finitely generated subgroups of $F_n \times F_n$ with unsolvable membership problem.

Clearly, $M(H)$ has solvable word problem for every H (because $F_n \times F_n$ also does). In particular, $M(H)$ is recursively presented. More interestingly, F. J. Grunewald proved, in [9, Theorem B], that if H is infinite then $M(H)$ cannot be finitely

presented. In [1], G. Baumslag and J. E. Roseblade completely described the structure of finitely presented subgroups of $F_n \times F_n$, a result that was later reproved by H. Short [15] and M. R. Bridson and D. T. Wise [5], and that implies Grunewald's result.

In this context, a natural problem is to look for recursive presentations for Mihailova's group $M(H)$, in terms of the original presentation H . This was recently posted as Problem 4.14 in [8] by R. I. Grigorchuk: "What kind of presentations can be obtained for Mihailova's subgroups of $F_n \times F_n$ determined by finite automata?"

The main result in the present paper (Theorem 1.1 below) gives a partial answer to this problem: under certain technical conditions on the initial H we give an explicit recursive presentation for $M(H)$ with finitely many generators and a one-parametric family of relations.

Theorem 1.1. *Let F_n be the free group on x_1, \dots, x_n , and let $H = \langle x_1, \dots, x_n \mid R_1, \dots, R_m \rangle$ be a finite, concise and Peiffer aspherical presentation. Then Mihailova's group $M(H) \leq F_n \times F_n$ admits the following presentation*

$$\langle d_1, \dots, d_n, t_1, \dots, t_m \mid [t_j, d^{-1}t_i^{-1}r_i d], [t_i, \text{root}(r_i)] \quad (1 \leq i, j \leq m, d \in D_n) \rangle,$$

where D_n is the free group with basis d_1, \dots, d_n , where r_i denotes the word in D_n obtained from R_i by replacing each x_k to d_k , and where $\text{root}(r_i)$ denotes the unique element $s_i \in D_n$ such that r_i is a positive power of s_i but s_i itself is not a proper power.

In this presentation the elements d_i and t_j correspond, respectively, to the elements (x_i, x_i) and $(1, R_j)$ of $M(H)$.

As a corollary we deduce the existence of a finitely generated, orbit undecidable subgroup of $\text{Aut}(F_3)$ (see [2] for details), which has the recursive presentation given in Theorem 1.1. Another application of Mihailova's construction can be found in [10], Proposition 5.4.

The structure of the paper is as follows. In Section 2 we recall some definitions and discuss some properties of concise and Peiffer aspherical presentations that will be used later. In Section 3 we prove Theorem 1.1. And in Section 4 we recall the relationship between Mihailova's subgroup and orbit undecidability, recently discovered in [2], and deduce the announced corollary (Theorem 4.2).

2. Asphericity

As stated, let $\langle x_1, \dots, x_n \mid R_1, \dots, R_m \rangle$ be a finite presentation. Formally, R_1, \dots, R_m is a list of words in the alphabet $\{x_1, \dots, x_n\}^{\pm 1}$ which may contain trivial and non-reduced words, possible repetitions, and even possible members conjugated to each other or to the inverse of each other.

A presentation $\langle x_1, \dots, x_n \mid R_1, \dots, R_m \rangle$ is called *concise* if every relation R_i is non-trivial and reduced, and every two relations $R_i, R_j, i \neq j$, are not conjugate to each other, or to the inverse of each other. Given an arbitrary finite presentation, $\langle x_1, \dots, x_n \mid R_1, \dots, R_m \rangle$, one can always reduce the relations and eliminate some of them, to obtain another presentation of the same group, which is concise. We call this a *concise refinement* of $\langle x_1, \dots, x_n \mid R_1, \dots, R_m \rangle$.

Now we recall the definition of Peiffer transformations. Consider some elements $U_1, \dots, U_l \in F_n$, some relators $R_{i_1}, \dots, R_{i_l} \in \{R_1, \dots, R_n\}$, and some numbers $\varepsilon_1, \dots, \varepsilon_l \in \{-1, 1\}$ such that the equation

$$(U_1 R_{i_1}^{\varepsilon_1} U_1^{-1}) \dots (U_l R_{i_l}^{\varepsilon_l} U_l^{-1}) = 1$$

holds in F_n . In this situation, the sequence of elements $(U_1 R_{i_1}^{\varepsilon_1} U_1^{-1}, \dots, U_l R_{i_l}^{\varepsilon_l} U_l^{-1})$ of F_n is called an *identity among relations* of length l . For $l = 0$ we have the *empty identity among relations*, $()$.

In such a sequence, let us replace two consecutive terms, say $U_p R_{i_p}^{\varepsilon_p} U_p^{-1}$ and $U_{p+1} R_{i_{p+1}}^{\varepsilon_{p+1}} U_{p+1}^{-1}$ for some $1 \leq p \leq l - 1$, by the new ones $U_{p+1} R_{i_{p+1}}^{\varepsilon_{p+1}} U_{p+1}^{-1}$ and $(U_{p+1} R_{i_{p+1}}^{-\varepsilon_{p+1}} U_{p+1}^{-1} U_p) R_{i_p}^{\varepsilon_p} (U_p^{-1} U_{p+1} R_{i_{p+1}}^{\varepsilon_{p+1}} U_{p+1}^{-1})$. Since the product of the two old terms do coincide with that of the two new ones, the new sequence is again an identity among relations. This transformation is called a *Peiffer transformation of the first kind* or, shortly, an *exchange*.

Suppose now that in the sequence $(U_1 R_{i_1}^{\varepsilon_1} U_1^{-1}, \dots, U_l R_{i_l}^{\varepsilon_l} U_l^{-1})$ there are two consecutive terms, say $U_p R_{i_p}^{\varepsilon_p} U_p^{-1}$ and $U_{p+1} R_{i_{p+1}}^{\varepsilon_{p+1}} U_{p+1}^{-1}$ for some $1 \leq p \leq l - 1$, whose product equals 1. Then we can obtain a new identity among relations by just deleting these two terms. This transformation and the inverse one are called *Peiffer transformations of the second kind* or shortly, *deletion* and *insertion*, respectively.

Definition 2.1. We say that a presentation is *Peiffer aspherical* if every identity among relations can be carried to the empty one by a sequence of Peiffer transformations.

In particular, a presentation admitting identities among relations of odd length is automatically not Peiffer aspherical.

A large class of Peiffer aspherical presentations can be obtained by using Theorems 3.1 and 4.2, and Lemma 5.1 from [6]. They state, respectively, that Peiffer asphericity is preserved under certain HNN extensions, under free products, and under Tietze transformations.

In the next section we shall argue using Peiffer asphericity. However, for completeness, we mention that in the literature there are (at least) three concepts of asphericity for presentations, which do not agree in general: *Peiffer asphericity* (called *combinatorial asphericity* in [6], see Proposition 1.5 there); *diagrammatical asphericity* defined in [6] like Peiffer asphericity but without allowing insertions (and also considered in Chapter III.10 of [11]); and *topological asphericity*.

Let $H = \langle x_1, \dots, x_n \mid R_1, \dots, R_m \rangle$ be a presentation and $\mathcal{K}(H)$ be the two-dimensional CW-complex with a single 0-cell, n 1-cells corresponding to the generators x_1, \dots, x_n , and m 2-cells each one being attached to the 1-skeleton along the path determined by the spelling of the corresponding relation. The presentation H is said to be *topologically aspherical* if $\pi_2(\mathcal{K}(H)) = 0$. As is indicated in Proposition 1.1 of [6], this is equivalent to the triviality of the second homology group of the universal cover of $\mathcal{K}(H)$.

The relations between these three concepts are as follows (for more details, see the introduction and Proposition 1.3 of [6]):

- (i) topological asphericity implies Peiffer asphericity,
- (ii) diagrammatical asphericity implies Peiffer asphericity,
- (iii) for presentations where every relation is reduced, topological asphericity is equivalent to Peiffer asphericity plus conciseness and “no relator being a proper power”.

3. Proof of Theorem 1.1

Back to Mihailova’s construction for $H = \langle x_1, \dots, x_n \mid R_1, \dots, R_m \rangle$, we recall that $M(H) \leq F_n \times F_n$ is generated by (x_i, x_i) and $(1, R_j)$, $i = 1, \dots, n$, $j = 1, \dots, m$. So, letting F_{n+m} be the free group with basis $\{d_1, \dots, d_n, t_1, \dots, t_m\}$, we have an epimorphism $\pi: F_{n+m} \rightarrow M(H)$ defined by $d_i \mapsto (x_i, x_i)$ and $t_j \mapsto (1, R_j)$, $i = 1, \dots, n$, $j = 1, \dots, m$. Now, for proving Theorem 1.1 we have to show that $\ker \pi$ is precisely the normal closure of the relations shown in the pretended presentation for $M(H)$. Note that the images of elements d_1, \dots, d_n generate the diagonal subgroup of $F_n \times F_n$, denoted $\text{Diag}(F_n \times F_n)$, which is isomorphic to F_n ; hence, π restricts to an isomorphism from $D_n = \langle d_1, \dots, d_n \rangle \leq F_{n+m}$ onto $\text{Diag}(F_n \times F_n) \leq M(H) \leq F_n \times F_n$.

We will keep the following notational convention in the proof: capital letters will always mean words in x_1, \dots, x_n , like R_j ; with this in mind, if u is a word on d_1, \dots, d_n , then its capitalization U will denote the word obtained from u by replacing each occurrence of d_i to x_i . Thus, U is just the projection of $\pi(u)$ to the first (or the second) coordinate.

Proof of Theorem 1.1. Recall that in the statement, r_j is the word in D_n obtained from R_j by replacing each x_i to d_i , $j = 1, \dots, m$.

Let \mathcal{N} be the normal closure (in the free group F_{n+m}) of the recursive family of commutators

$$\{[t_j, d^{-1}t_i^{-1}r_i d], [t_i, \text{root}(r_i)] \mid i, j = 1, \dots, m, d \in D_n\}.$$

Our goal is to show that $\mathcal{N} = \ker \pi$. The inclusion $\mathcal{N} \leq \ker \pi$ is straightforward from the following computations:

$$\begin{aligned} \pi([t_j, d^{-1}t_i^{-1}r_i d]) &= [(1, R_j), (u, u)^{-1}(R_i, 1)(u, u)] \\ &= [(1, R_j), (u^{-1}R_i u, 1)] = (1, 1), \\ \pi([t_i, \text{root}(r_i)]) &= [(1, R_i), (\text{root}(R_i), \text{root}(R_i))] = (1, 1). \end{aligned}$$

In order to prove $\ker \pi \leq \mathcal{N}$, we shall use the following strategy: to each word $w \in \ker \pi$ we will associate an identity among relations for the presentation $\langle x_1, \dots, x_n \mid R_1, \dots, R_m \rangle$ of H , in such a way that if $w \neq 1$ then the associated identity is non-empty; then we will show that, after applying an arbitrary Peiffer transformation, the resulting identity among relations is again the one associated to some other word $w' \in \ker \pi$ satisfying, additionally, that $w^{-1}w' \in \mathcal{N}$.

Having seen this, let $w \in \ker \pi$ and consider the associated identity among relations. Since, by hypothesis, the presentation $\langle x_1, \dots, x_n \mid R_1, \dots, R_m \rangle$ is Peiffer aspherical, there exists a sequence of Peiffer transformations reducing such identity to the empty one. Now, repeatedly using the result mentioned in the previous paragraph, we obtain a list of words (ending with the trivial one because the last identity is empty), $w, w', w'', \dots, 1$, and such that the difference between every two consecutive ones belongs to \mathcal{N} . This shows that $w \in \mathcal{N}$ concluding the proof.

So we are reduced to construct such an association. Let $w \in \ker \pi \leq F_{n+m}$ and write it in the form $w = u_1 t_{i_1}^{\varepsilon_1} u_2 \dots u_l t_{i_l}^{\varepsilon_l} u_{l+1}$, where $l \geq 0$ and u_1, \dots, u_{l+1} are words in d_1, \dots, d_n . Then, projecting $\pi(w)$ to each coordinate, we have

$$U_1 U_2 \dots U_{l+1} = 1 \quad \text{and} \quad U_1 R_{i_1}^{\varepsilon_1} U_2 \dots U_l R_{i_l}^{\varepsilon_l} U_{l+1} = 1. \tag{1}$$

Denote the accumulative products by $\mathbb{U}_i = U_1 U_2 \dots U_i, i = 1, \dots, l + 1$ (note that $\mathbb{U}_{l+1} = 1$). By (1), we have

$$\mathbb{U}_1 R_{i_1}^{\varepsilon_1} \mathbb{U}_1^{-1} \mathbb{U}_2 R_{i_2}^{\varepsilon_2} \mathbb{U}_2^{-1} \dots \mathbb{U}_l R_{i_l}^{\varepsilon_l} \mathbb{U}_l^{-1} = 1$$

in the free group F_n . In other words,

$$(\mathbb{U}_1 R_{i_1}^{\varepsilon_1} \mathbb{U}_1^{-1}, \mathbb{U}_2 R_{i_2}^{\varepsilon_2} \mathbb{U}_2^{-1}, \dots, \mathbb{U}_l R_{i_l}^{\varepsilon_l} \mathbb{U}_l^{-1}) \tag{2}$$

is an identity among relations for the presentation $\langle x_1, \dots, x_n \mid R_1, \dots, R_m \rangle$ of H . This is the *identity associated to* $w \in \ker \pi$. Note that if this identity is empty, that is $l = 0$, then $w = u_1 \in \langle d_1, \dots, d_n \rangle \cap \ker \pi$ and so $w = 1$.

Let us analyze the situation when we apply an arbitrary Peiffer transformation to this identity.

Case 1: Consider the exchange which, for some $1 \leq p \leq l - 1$, replaces the consecutive terms

$$\mathbb{U}_p R_{i_p}^{\varepsilon_p} \mathbb{U}_p^{-1} \quad \text{and} \quad \mathbb{U}_{p+1} R_{i_{p+1}}^{\varepsilon_{p+1}} \mathbb{U}_{p+1}^{-1},$$

in (2), by the terms

$$\mathbb{U}_{p+1} R_{i_{p+1}}^{\varepsilon_{p+1}} \mathbb{U}_{p+1}^{-1} \quad \text{and} \quad (\mathbb{U}_{p+1} R_{i_{p+1}}^{-\varepsilon_{p+1}} \mathbb{U}_{p+1}^{-1} \mathbb{U}_p) R_{i_p}^{\varepsilon_p} (\mathbb{U}_p^{-1} \mathbb{U}_{p+1} R_{i_{p+1}}^{\varepsilon_{p+1}} \mathbb{U}_{p+1}^{-1}), \quad (3)$$

respectively. We claim that the identity among relations obtained in this way is precisely the one corresponding to the word

$$w' = v_1 t_{i_1}^{\varepsilon_1} \cdots v_{p-1} t_{i_{p-1}}^{\varepsilon_{p-1}} v_p t_{i_{p+1}}^{\varepsilon_{p+1}} v_{p+1} t_{i_p}^{\varepsilon_p} v_{p+2} t_{i_{p+2}}^{\varepsilon_{p+2}} \cdots v_l t_{i_l}^{\varepsilon_l} v_{l+1},$$

where

$$\begin{aligned} v_1 &= u_1, & v_p &= u_p u_{p+1}, & v_{p+3} &= u_{p+3}, \\ \vdots & & v_{p+1} &= r_{i_{p+1}}^{-\varepsilon_{p+1}} u_{p+1}^{-1}, & \vdots & \\ v_{p-1} &= u_{p-1}, & v_{p+2} &= u_{p+1} r_{i_{p+1}}^{\varepsilon_{p+1}} u_{p+2}, & v_{l+1} &= u_{l+1}. \end{aligned}$$

And we also claim that $w^{-1}w' \in \mathcal{N}$. This second assertion is easy to verify since we can obtain back w from w' by permuting the two consecutive subwords $u_{p+1} t_{i_{p+1}}^{\varepsilon_{p+1}} r_{i_{p+1}}^{-\varepsilon_{p+1}} u_{p+1}^{-1}$ and $t_{i_p}^{\varepsilon_p}$. But the commutator of these two words is an element of \mathcal{N} : for $\varepsilon_{p+1} = -1$ this is immediate; and for $\varepsilon_{p+1} = 1$ it follows from the facts that, modulo \mathcal{N} , t_{i_p} (and so $t_{i_p}^{\varepsilon_p}$) commutes with $u_{p+1} (t_{i_{p+1}}^{-1} r_{i_{p+1}})^{\pm 1} u_{p+1}^{-1}$, but also $t_{i_{p+1}}$ commutes with $t_{i_{p+1}}^{-1} r_{i_{p+1}}$ (and so, $t_{i_{p+1}}^{-1}$ with $r_{i_{p+1}}$). Therefore, w' equals w modulo \mathcal{N} .

To see the first part of the claim, let us capitalize the v_i 's:

$$\begin{aligned} V_1 &= U_1, & V_p &= U_p U_{p+1}, & V_{p+3} &= U_{p+3}, \\ \vdots & & V_{p+1} &= R_{i_{p+1}}^{-\varepsilon_{p+1}} U_{p+1}^{-1}, & \vdots & \\ V_{p-1} &= U_{p-1}, & V_{p+2} &= U_{p+1} R_{i_{p+1}}^{\varepsilon_{p+1}} U_{p+2}, & V_{l+1} &= U_{l+1}. \end{aligned}$$

And let us compute the $\mathbb{V}_i = V_1 V_2 \dots V_i$'s:

$$\begin{aligned} \mathbb{V}_1 &= \mathbb{U}_1, & \mathbb{V}_p &= \mathbb{U}_{p+1}, & \mathbb{V}_{p+3} &= \mathbb{U}_{p+3}, \\ \vdots & & \mathbb{V}_{p+1} &= \mathbb{U}_{p+1} R_{i_{p+1}}^{-\varepsilon_{p+1}} \mathbb{U}_{p+1}^{-1} \mathbb{U}_p, & \vdots & \\ \mathbb{V}_{p-1} &= \mathbb{U}_{p-1}, & \mathbb{V}_{p+2} &= \mathbb{U}_{p+2}, & \mathbb{V}_{l+1} &= \mathbb{U}_{l+1}. \end{aligned}$$

Finally, the identity among relations associated to w' is

$$\begin{aligned}
 (\mathbb{V}_1 R_{i_1}^{\varepsilon_1} \mathbb{V}_1^{-1} &= \mathbb{U}_1 R_{i_1}^{\varepsilon_1} \mathbb{U}_1^{-1}, \\
 &\vdots \\
 \mathbb{V}_{p-1} R_{i_{p-1}}^{\varepsilon_{p-1}} \mathbb{V}_{p-1}^{-1} &= \mathbb{U}_{p-1} R_{i_{p-1}}^{\varepsilon_{p-1}} \mathbb{U}_{p-1}^{-1}, \\
 \mathbb{V}_p R_{i_{p+1}}^{\varepsilon_{p+1}} \mathbb{V}_p^{-1} &= \mathbb{U}_{p+1} R_{i_{p+1}}^{\varepsilon_{p+1}} \mathbb{U}_{p+1}^{-1}, \\
 \mathbb{V}_{p+1} R_{i_p}^{\varepsilon_p} \mathbb{V}_{p+1}^{-1} &= \mathbb{U}_{p+1} R_{i_{p+1}}^{-\varepsilon_{p+1}} \mathbb{U}_{p+1}^{-1} \mathbb{U}_p R_{i_p}^{\varepsilon_p} \mathbb{U}_p^{-1} \mathbb{U}_{p+1} R_{i_{p+1}}^{\varepsilon_{p+1}} \mathbb{U}_{p+1}^{-1}, \\
 \mathbb{V}_{p+2} R_{i_{p+2}}^{\varepsilon_{p+2}} \mathbb{V}_{p+2}^{-1} &= \mathbb{U}_{p+2} R_{i_{p+2}}^{\varepsilon_{p+2}} \mathbb{U}_{p+2}^{-1}, \\
 &\vdots \\
 \mathbb{V}_l R_{i_l}^{\varepsilon_l} \mathbb{V}_l^{-1} &= \mathbb{U}_l R_{i_l}^{\varepsilon_l} \mathbb{U}_l^{-1}),
 \end{aligned}$$

which does coincide with the identity among relations obtained from (2) after applying the Peiffer transformation (3).

Case 2: Consider the deletion which, for some $1 \leq p \leq l - 1$, deletes the consecutive terms

$$\mathbb{U}_p R_{i_p}^{\varepsilon_p} \mathbb{U}_p^{-1} \quad \text{and} \quad \mathbb{U}_{p+1} R_{i_{p+1}}^{\varepsilon_{p+1}} \mathbb{U}_{p+1}^{-1}, \tag{4}$$

in (2), assuming that its product equals 1. We claim that the identity among relations obtained in this way is precisely the one corresponding to the word

$$w' = v_1 t_{i_1}^{\varepsilon_1} \dots v_{p-1} t_{i_{p-1}}^{\varepsilon_{p-1}} v_p t_{i_{p+2}}^{\varepsilon_{p+2}} v_{p+1} \dots v_{l-2} t_{i_l}^{\varepsilon_l} v_{l-1},$$

where

$$\begin{aligned}
 v_1 &= u_1, & v_{p+1} &= u_{p+3}, \\
 \vdots & & v_p &= u_p u_{p+1} u_{p+2}, & \vdots \\
 v_{p-1} &= u_{p-1}, & v_{l-1} &= u_{l+1}.
 \end{aligned}$$

And we also claim that $w^{-1}w' \in \mathcal{N}$. This second assertion follows from the hypothesis that $(\mathbb{U}_p R_{i_p}^{\varepsilon_p} \mathbb{U}_p^{-1}) \cdot (\mathbb{U}_{p+1} R_{i_{p+1}}^{\varepsilon_{p+1}} \mathbb{U}_{p+1}^{-1}) = 1$. In fact, conciseness implies that $i_p = i_{p+1}$, $\varepsilon_p = -\varepsilon_{p+1}$ and so $\mathbb{U}_p^{-1} \mathbb{U}_{p+1} = U_{p+1}$ commutes with $R_{i_{p+1}}$; hence, u_{p+1} commutes with $r_{i_{p+1}}$ and so $u_{p+1} \in \langle \text{root}(r_{i_{p+1}}) \rangle$. Now w' can be obtained from w by replacing the subword $t_{i_p}^{\varepsilon_p} u_{p+1} t_{i_{p+1}}^{\varepsilon_{p+1}}$ to u_{p+1} . But $(t_{i_p}^{\varepsilon_p} u_{p+1} t_{i_{p+1}}^{\varepsilon_{p+1}})^{-1} u_{p+1} \in \mathcal{N}$ since $t_{i_{p+1}}$ commutes with $\text{root}(r_{i_{p+1}})$ modulo \mathcal{N} .

To see the first part of the claim, let us capitalize the v_i 's:

$$\begin{aligned}
 V_1 &= U_1, & V_{p+1} &= U_{p+3}, \\
 \vdots & & V_p &= U_p U_{p+1} U_{p+2}, & \vdots \\
 V_{p-1} &= U_{p-1}, & V_{l-1} &= U_{l+1}.
 \end{aligned}$$

And let us compute the $\mathbb{V}_i = V_1 V_2 \dots V_i$'s:

$$\begin{aligned} \mathbb{V}_1 &= \mathbb{U}_1, & \mathbb{V}_{p+1} &= \mathbb{U}_{p+3}, \\ \vdots & & \mathbb{V}_p &= \mathbb{U}_{p+2}, & \vdots \\ \mathbb{V}_{p-1} &= \mathbb{U}_{p-1}, & \mathbb{V}_{l-1} &= \mathbb{U}_{l+1}. \end{aligned}$$

Finally, the identity among relations associated to w' is

$$\begin{aligned} (\mathbb{V}_1 R_{i_1}^{\varepsilon_1} \mathbb{V}_1^{-1} &= \mathbb{U}_1 R_{i_1}^{\varepsilon_1} \mathbb{U}_1^{-1}, \\ &\vdots \\ \mathbb{V}_{p-1} R_{i_{p-1}}^{\varepsilon_{p-1}} \mathbb{V}_{p-1}^{-1} &= \mathbb{U}_{p-1} R_{i_{p-1}}^{\varepsilon_{p-1}} \mathbb{U}_{p-1}^{-1}, \\ \mathbb{V}_p R_{i_{p+2}}^{\varepsilon_{p+2}} \mathbb{V}_p^{-1} &= \mathbb{U}_{p+2} R_{i_{p+2}}^{\varepsilon_{p+2}} \mathbb{U}_{p+2}^{-1}, \\ \mathbb{V}_{p+1} R_{i_{p+3}}^{\varepsilon_{p+3}} \mathbb{V}_{p+1}^{-1} &= \mathbb{U}_{p+3} R_{i_{p+3}}^{\varepsilon_{p+3}} \mathbb{U}_{p+3}^{-1}, \\ &\vdots \\ \mathbb{V}_{l-2} R_{i_l}^{\varepsilon_l} \mathbb{V}_{l-2}^{-1} &= \mathbb{U}_l R_{i_l}^{\varepsilon_l} \mathbb{U}_l^{-1}), \end{aligned}$$

which coincides with the identity among relations obtained from (2) after applying the Peiffer transformation (4).

Case 3: Consider an insertion, and argue in a similar way as in Case 2.

This concludes the proof. □

4. A recursively presented orbit undecidable subgroup of $\text{Aut}(F_3)$

In [2], O. Bogopolski, A. Martino and E. Ventura studied the conjugacy problem for extensions of groups. In that context, the notion of orbit decidability is crucial and we recall it here.

Let F be a group, and $A \leq \text{Aut}(F)$. We say that A is *orbit decidable* if and only if there exists an algorithm such that, given $u, v \in F$, decides whether v is conjugate to $\alpha(u)$ for some $\alpha \in A$.

The main result in [2] states that, given a short exact sequence of groups

$$1 \rightarrow F \rightarrow G \rightarrow P \rightarrow 1$$

with some conditions on F and P , the group G has solvable conjugacy problem if and only if the action subgroup

$$A_G = \{\gamma_g : F \rightarrow F, x \mapsto g^{-1}xg \mid g \in G\} \leq \text{Aut}(F)$$

is orbit decidable (see [2], Theorem 3.1, for details).

In particular, this applies to the case where F and P are finitely generated free groups, giving a characterization of the solvability of the conjugacy problem within the family of [finitely generated free]-by-[finitely generated free] groups. This family of groups is interesting because C. F. Miller, back in the 1970s, already showed the existence of [finitely generated free]-by-[finitely generated free] groups with unsolvable conjugacy problem (see [13]). Via [2], Theorem 3.1, this can be restated by saying that $\text{Aut}(F_n)$ contains finitely generated orbit undecidable subgroups (for some n).

Question 6 in the last section of [2] asks whether finitely presented subgroups $A \leq \text{Aut}(F_n)$ are orbit decidable or not. The answer is known to be positive in rank 2 (every finitely generated subgroup of $\text{Aut}(F_2)$ is orbit decidable, see [2], Proposition 6.21), but open for bigger rank. The comment made in [2] after this question says that if H is a finitely generated group with unsolvable word problem, then Mihailova’s group $M(H)$ is isomorphic to an orbit undecidable subgroup of $\text{Aut}(F_3)$. And, as mentioned in the introduction, this subgroup is then finitely generated, and recursively presented, but it cannot be finitely presented.

In the rest of the paper, we will recall how $M(H)$ can be embedded into $\text{Aut}(F_3)$, in such a way that the image becomes an orbit undecidable subgroup of $\text{Aut}(F_3)$. Then we will choose an appropriate H and prove Theorem 4.2 by applying Theorem 1.1 to $A = M(H)$.

Of course, Theorem 4.2 does not answer the above mentioned Question 6, but shows its tightness in the sense that orbit undecidability is already showing up in the class of one-parametric recursively presented subgroups of $\text{Aut}(F_3)$.

First, let $F_3 = \langle q, a, b \mid \rangle$ be the free group on $\{q, a, b\}$, and let us embed $F_2 \times F_2$ into $\text{Aut}(F_3)$ in the following natural way. For every $u, v \in \langle a, b \rangle$, consider the automorphism

$${}_u\theta_v : F_3 \rightarrow F_3, \quad q \mapsto uqv, \quad a \mapsto a, \quad b \mapsto b.$$

Clearly, ${}_u\theta_1 \cdot {}_{u_2}\theta_1 = {}_{u_1u_2}\theta_1$ and ${}_1\theta_{v_1} \cdot {}_1\theta_{v_2} = {}_1\theta_{v_2v_1}$, which means that $\{{}_u\theta_1 \mid u \in \langle a, b \rangle\} \simeq F_2$ and $\{{}_1\theta_v \mid v \in \langle a, b \rangle\} \simeq F_2^{\text{op}} \simeq F_2$. It is also clear that ${}_u\theta_1 \cdot {}_1\theta_v = {}_u\theta_v = {}_1\theta_v \cdot {}_u\theta_1$. So we have an embedding $\theta : F_2 \times F_2 \simeq F_2^{\text{op}} \times F_2^{\text{op}} \hookrightarrow \text{Aut}(F_3)$ given by $(u, v) \mapsto {}_{u^{-1}}\theta_v$, whose image is

$$F_2 \times F_2 \simeq B = \langle {}_{a^{-1}}\theta_1, {}_{b^{-1}}\theta_1, {}_1\theta_a, {}_1\theta_b \rangle = \{{}_u\theta_v \mid u, v \in \langle a, b \rangle\} \leq \text{Aut}(F_3).$$

As shown in [2], Section 7.2, the element $qaqbq$ satisfies the technical condition required in [2], Proposition 7.3. Hence, we have

Lemma 4.1 (7.3 in [2]). *For the above defined subgroup $B \leq \text{Aut}(F_3)$ and for every subgroup $A \leq B$, undecidability of the membership problem for A in B implies orbit undecidability for A in $\text{Aut}(F_3)$.*

We are ready to deduce the main result of this section.

Theorem 4.2. *There exists a finitely generated (and not finitely presented) orbit undecidable subgroup $A \leq \text{Aut}(F_3)$ admitting a one-parametric recursive presentation as in Theorem 1.1.*

Proof. In [7], D. J. Collins and C. F. Miler III proved that there exists a finite, concise and Peiffer aspherical presentation $\langle x_1, \dots, x_n \mid R_1, \dots, R_m \rangle$ of a group H with unsolvable word problem. The corresponding Mihailova's group $M(H)$ is a subgroup of $F_n \times F_n$ and the membership problem for $M(H)$ in $F_n \times F_n$ is unsolvable.

Now, denoting $A = M(H)$ and using a finite index embedding of $F_n \times F_n$ in $B \cong F_2 \times F_2$, we have that $A \leq B$ and the membership problem for A in B is unsolvable. By Lemma 4.1, A is an orbit undecidable subgroup of $\text{Aut}(F_3)$.

Moreover, as it was discussed in the introduction, A is finitely generated, and is not finitely presented. But Theorem 1.1 provides an explicit one-parametric recursive presentation for A . This concludes the proof. \square

We end by reproducing [2], Question 6, again:

Question 4.3. Does there exist a finitely presented orbit undecidable subgroup of $\text{Aut}(F_n)$ for $n \geq 3$?

Acknowledgements. The first named author thanks the MPIM at Bonn for its support and excellent working conditions during the fall 2008, while this research was finished. The second author gratefully acknowledges partial support from the MEC (Spain) and the EFRD (EC) through project number MTM-2008-01550.

References

- [1] G. Baumslag and J. E. Roseblade, Subgroups of direct products of free groups. *J. London Math. Soc.* (2) **30** (1984), 44–52. [Zbl 0559.20018](#) [MR 760871](#)
- [2] O. Bogopolski, A. Martino, and E. Ventura, Orbit decidability and the conjugacy problem for some extensions of groups. *Trans. Amer. Math. Soc.* **362** (2010), 2003–2036. [Zbl 05692501](#) [MR 2574885](#)
- [3] W. W. Boone, The word problem. *Ann. of Math.* (2) **70** (1959), 207–265. [Zbl 0102.00902](#) [MR 0179237](#)
- [4] V. V. Borisov, Simple examples of groups with unsolvable word problem. *Mat. Zametki* **6** (1969), 521–532; English transl. *Math. Notes* **6** (1969), 768–775. [Zbl 0215.10803](#) [MR 0260851](#)
- [5] M. R. Bridson and D. T. Wise, $\mathcal{V}\mathcal{H}$ complexes, towers and subgroups of $F \times F$. *Math. Proc. Cambridge Philos. Soc.* **126** (1999), 481–497. [Zbl 0942.20009](#) [MR 1684244](#)
- [6] I. M. Chiswell, D. J. Collins, and J. Huebschmann, Aspherical group presentations. *Math. Z.* **178** (1981), 1–36. [Zbl 0443.20030](#) [MR 627092](#)

- [7] D. J. Collins and C. F. Miller, III, The word problem in groups of cohomological dimension 2. In *Groups St. Andrews 1997 in Bath*, I, London Math. Soc. Lecture Note Ser. 260, Cambridge University Press, Cambridge 1999, 211–218. [Zbl 1001.20030](#) [MR 1676618](#)
- [8] R. Grigorchuk, Solved and unsolved problems around one group. In *Infinite groups: geometric, combinatorial and dynamical aspects*, Progr. Math. 248, Birkhäuser, Basel 2005, 117–218. [Zbl 1165.20021](#) [MR 2195454](#)
- [9] F. J. Grunewald, On some groups which cannot be finitely presented. *J. London Math. Soc.* (2) **17** (1978), 427–436. [Zbl 0385.20020](#) [MR 500627](#)
- [10] F. Grunewald and V. Platonov, Rigidity results for groups with radical cohomology of finite groups and arithmeticity problems. *Duke Math. J.* **100** (1999), 321–358. [Zbl 1007.11029](#) [MR 1722957](#)
- [11] R. C. Lyndon and P. E. Schupp, *Combinatorial group theory*. Ergeb. Math. Grenzgeb. 89, Springer-Verlag, Berlin 1977. [Zbl 0368.20023](#) [MR 0577064](#)
- [12] K. A. Mihailova, The occurrence problem for direct products of groups. *Dokl. Akad. Nauk SSSR* **119** (1958), 1103–1105 (in Russian). [MR 0100018](#)
- [13] C. F. Miller III, *On group-theoretic decision problems and their classification*. Ann. of Math. Stud. 68, Princeton University Press, Princeton 1971. [Zbl 0277.20054](#) [MR 0310044](#)
- [14] P. S. Novikov, On the algorithmic insolvability of the word problem in group theory. *Trudy Mat. Inst. Steklov.* **44** 1955, 1–143; English transl. in Amer. Math. Soc. Transl. (2) 9, Amer. Math. Soc., Providence, R.I., 1958, 1–122. [Zbl 0093.01304](#) [MR 0075197](#)
- [15] H. Short, Finitely presented subgroups of a product of two free groups. *Q. J. Math.* **52** (2001), 127–131. [Zbl 0984.20021](#) [MR 1820907](#)

Received October 15, 2008; revised August 10, 2009

O. Bogopolski, Mathematisches Institut der Heinrich-Heine-Universität Düsseldorf, Universitätsstr. 1, 40225 Düsseldorf, Germany, and Institute of Mathematics of Siberian Branch of Russian Academy of Sciences, Koptjuga 4, Novosibirsk 630090, Russia
E-mail: Oleg_Bogopolski@yahoo.com

E. Ventura, Department of Applied Mathematics III, Escola Politècnica Superior d'Enginyeria de Manresa, Universitat Politècnica de Catalunya, Av. Bases de Manresa 61–73, 08242 Manresa, Barcelona, Catalonia, Spain
E-mail: enric.ventura@upc.edu