

On hyperbolicity of free splitting and free factor complexes

Ilya Kapovich and Kasra Rafi*

Abstract. We show how to derive hyperbolicity of the free factor complex of F_N from the Handel–Mosher proof of hyperbolicity of the free splitting complex of F_N , thus obtaining an alternative proof of a theorem of Bestvina–Feighn. We also show that under the natural map τ from the free splitting complex to free factor complex, a geodesic $[x, y]$ maps to a path that is uniformly Hausdorff-close to a geodesic $[\tau(x), \tau(y)]$.

Mathematics Subject Classification (2010). 20F65, 57M07.

Keywords. Free group, curve complex, outer automorphism group of the free group.

1. Introduction

The notion of a *curve complex*, introduced by Harvey [17] in late 1970s, plays a key role in the study of hyperbolic surfaces, mapping class group and the Teichmüller space.

If S is a compact connected oriented surface, the *curve complex* $\mathcal{C}(S)$ of S is a simplicial complex whose vertices are isotopy classes of essential non-peripheral simple closed curves. A collection $[\alpha_0], \dots, [\alpha_n]$ of $(n + 1)$ distinct vertices of $\mathcal{C}(S)$ spans an n -simplex in $\mathcal{C}(S)$ if there exist representatives $\alpha_0, \dots, \alpha_n$ of these isotopy classes such that for all $i \neq j$ the curves α_i and α_j are disjoint. (The definition of $\mathcal{C}(S)$ is a little different for several surfaces of small genus). The complex $\mathcal{C}(S)$ is finite-dimensional but not locally finite, and it comes equipped with a natural action of the mapping class group $\text{Mod}(S)$ by simplicial automorphisms. It turns out that the geometry of $\mathcal{C}(S)$ is closely related to the geometry of the Teichmüller space $\mathcal{T}(S)$ and also of the mapping class group itself. The curve complex is a basic tool in modern Teichmüller theory, and has also found numerous applications in the study of 3-manifolds and of Kleinian groups. A key general result of Masur and Minsky [23] says that the curve complex $\mathcal{C}(S)$, equipped with the simplicial metric, is a Gromov-hyperbolic space. Hyperbolicity of the curve complex was an important ingredient in the solution by Masur, Minsky, Brock and Canary of the

*The first author was partially supported by the NSF grant DMS-0904200. The second author was partially supported by the NSF grant DMS-1007811.

Ending Lamination Conjecture [24], [27], [10] (see [26] for detailed background discussion about this solution).

The outer automorphism group $\text{Out}(F_N)$ of a free group F_N is a cousin of the mapping class group. However the group $\text{Out}(F_N)$ is much less well understood and, in general, more difficult to study than the mapping class group. A free group analog of the Teichmüller space is the Culler–Vogtmann Outer space cv_N , introduced by Culler and Vogtmann in [11]. The points of cv_N are free minimal discrete isometric actions of F_N on \mathbb{R} -trees, considered up to F_N -equivariant isometry. The Outer space comes equipped with a natural action of $\text{Out}(F_N)$. It is known that cv_N is finite-dimensional and contractible; as a result, quite a bit is known about homotopy properties of $\text{Out}(F_N)$. However, the geometry of cv_N and of $\text{Out}(F_N)$ proved to be much more difficult to tackle, particularly because cv_N lacks the various useful analytic and geometric structures present in the Teichmüller space case. Another problem is that many geometric dualities from the world of Riemann surfaces and their homeomorphisms break down for automorphisms of free groups.

In the case of a compact connected oriented surface S , an essential non-peripheral simple closed curve α on S can be viewed in several other ways. Thus one can view $[\alpha]$ as a conjugacy class in the fundamental group $\pi_1(S)$. We may also think of α as corresponding to the (possibly disconnected) subsurface K_α of S obtained by cutting S open along α . Third, α determines a splitting of $\pi_1(S)$ as an amalgamated product or an HNN-extension (depending on whether the curve α is separating or non-separating) over the infinite cyclic subgroup $\langle \alpha \rangle$. We can interpret adjacency of vertices in $\mathcal{C}(S)$ using each of these points of views – or a combination of them, providing several essentially equivalent descriptions of the curve complex. Thus two distinct vertices $[\alpha], [\beta]$ of $\mathcal{C}(S)$ are adjacent if and only if α is conjugate in $\pi_1(S)$ to an element of a vertex group of the cyclic splitting of $\pi_1(S)$ corresponding to β . Equivalently, $[\alpha]$ and $[\beta]$ of $\mathcal{C}(S)$ are adjacent if and only if the cyclic splittings of $\pi_1(S)$ corresponding to $[\alpha]$ and $[\beta]$ admit a common refinement, that is, a splitting of $\pi_1(S)$ as the fundamental group of a graph of groups with two edges and cyclic edge groups, such that collapsing one of the edges produces a splitting corresponding to $[\alpha]$ and collapsing the other edge produces a splitting corresponding to $[\beta]$. Also, $[\alpha]$ and $[\beta]$ are adjacent in $\mathcal{C}(S)$ if and only if there are connected components K'_α of K_α and K'_β of K_β such that $K'_\alpha \subseteq K'_\beta$ or $K'_\beta \subseteq K'_\alpha$.

In the case of F_N these different points of view produce several possible analogs of the notion of a curve complex that are no longer essentially equivalent. The first of these is the *free splitting complex* FS_N . The vertices of FS_N are nontrivial splittings of the type $F_N = \pi_1(\mathbb{A})$ where \mathbb{A} is a graph of groups with a single edge (possibly a loop edge) and the trivial edge group; two such splittings are considered to be the same if their Bass–Serre covering trees are F_N -equivariantly isometric. Two distinct vertices \mathbb{A} and \mathbb{B} of FS_N are joined by an edge if these splittings admit a common refinement, that is, a splitting $F_N = \pi_1(\mathbb{D})$ where \mathbb{D} is a graph of groups with two edges and trivial edge groups, such that collapsing one edge gives the splitting \mathbb{A} and collapsing the other edge produces the splitting \mathbb{B} . Higher-dimensional simplices

are defined in a similar way, see Definition 3.2 below for a careful formulation. For example, if $F_N = A * B * C$, where A, B, C are nontrivial, then the splittings $F_N = (A * B) * C$ and $F_N = A * (B * C)$ are adjacent in FS_N . There is a natural action of $\text{Out}(F_N)$ on FS_N by simplicial automorphisms. The above definition of FS_N has a variation [29], called the *edge-splitting complex*, denoted ES_N , where in the definition of vertices only splittings \mathbb{A} with a single *non-loop* edge are allowed.

A rather different free group analog of the curve complex is the *free factor complex* FF_N , originally introduced by Hatcher and Vogtmann [18]. The vertices of FF_N are conjugacy classes $[A]$ of proper free factors A of F_N . Two distinct vertices $[A], [B]$ are joined by an edge in FF_N if there exist representatives A of $[A]$ and B of $[B]$ such that $A \leq B$ or $B \leq A$. Higher-dimensional simplices are defined similarly, see Definition 3.1 below. Note that this definition does not work well for $N = 2$ as it produces a graph consisting of isolated vertices corresponding to conjugacy classes of primitive elements in F_2 . However, there is a natural modification of the definition of FF_N for $N = 2$ (see [4]) such that FF_2 becomes the standard Farey graph (and in particular FF_2 is hyperbolic).

A closely related object to FF_N is the *simplicial intersection graph* I_N . The graph I_N is a bipartite graph with two types of vertices: single-edge free splittings $F_N = \pi_1(\mathbb{A})$ (that is, vertices of FS_N) and conjugacy classes of *simple* elements of F_N . Here an element $a \in F_N$ is *simple* if a belongs to some proper free factor of F_N . A free splitting \mathbb{A} and a conjugacy class $[a]$ of a simple element a are adjacent if a is conjugate to an element of a vertex group of \mathbb{A} . The graph is a subgraph of a more general “intersection graph” defined in [20].

Both FF_N and I_N admit natural $\text{Out}(F_N)$ -actions. It is also not hard to check that for $N \geq 3$ the graph I_N is quasi-isometric to the free factor complex FF_N . By contrast, the free factor complex FF_N and the free splitting complex FS_N are rather different objects geometrically. By construction, the vertex set $V(FS_N)$ is a 1-dense subset of $V(I_N)$. Also, the inclusion map $\iota: (V(FS_N), d_{FS_N}) \rightarrow (I_N, d_{I_N})$ is 2-Lipschitz. However the distance between two free splittings in I_N is generally much smaller than the distance between them in FS_N . Intuitively, it is “much easier” for \mathbb{A} and \mathbb{B} to share a common elliptic simple element (which would make $d_{I_N}(\mathbb{A}, \mathbb{B}) \leq 2$) then for these splittings to admit a common refinement.

Until recently, basically nothing was known about the geometry of the above complexes. Several years ago Kapovich–Lustig [20] and Behrstock–Bestvina–Clay [2] showed that for $N \geq 3$ the (quasi-isometric) complexes FF_N and I_N have infinite diameter. Since the inclusion map ι above is Lipschitz, this implies that FS_N has infinite diameter as well. A subsequent result of Bestvina–Feighn [3] implies that every fully irreducible element $\varphi \in \text{Out}(F_N)$ acts on FF_N with positive asymptotic translation length (hence the same is true for the action of φ on FS_N). It is easy to see from the definitions that if $\varphi \in \text{Out}(F_N)$ is not fully irreducible then some positive power of φ fixes a vertex of FF_N , so that φ acts on FF_N with bounded orbits.

Sabalka and Savchuk proved [29] in 2010 that the edge-splitting complex ES_N is not Gromov-hyperbolic, because it possesses some quasi-flats. Aramayona and

Souto [1] showed that every automorphism of FS_N is induced by some element of $\text{Out}(F_N)$.

In 2011, two significant further advances occurred. First, Bestvina and Feighn [4] proved that for $N \geq 2$ the free splitting complex is Gromov-hyperbolic (as noted above, for $N = 2$ this essentially follows from the definition of FF_2 , so the main case of the Bestvina–Feighn result is for $N \geq 3$). Then Handel and Mosher [16] proved that for all $N \geq 2$ the free splitting complex FS_N is also Gromov-hyperbolic. The two proofs are rather different in nature, although both are quite complicated. Recently Hilion and Horbez [19] produced another proof of hyperbolicity of FS_N , using “surgery paths” in the sphere complex model of FS_N . Bestvina and Reynolds [6] and Hamenstädt [15] gave a description of the hyperbolic boundary of FF_N . Also, Bestvina and Feighn [5] and then Sabalka and Savchuk [30] investigated analogs of subsurface projections in the FS_N and FF_N contexts.

In the present paper we show how to derive hyperbolicity of the free factor complex from the Handel–Mosher proof of hyperbolicity of the free splitting complex. This gives a new proof of the Bestvina–Feighn result [4] about hyperbolicity of FF_N .

There is a natural “almost canonical” Lipschitz projection from the free splitting complex to a free factor complex. Namely, for any free splitting $v = \mathbb{A} \in V(FS_N)$ choose a vertex u of \mathbb{A} and put $\tau(v) := [A_u]$, where A_u is the vertex group of u in \mathbb{A} . This defines a map (easily seen to be Lipschitz) $\tau: V(FS_N) \rightarrow V(FF_N)$. Extend this map to a graph-map $\tau: FS_N^{(1)} \rightarrow FF_N^{(1)}$ by sending every edge in FS_N to a geodesic in $FF_N^{(1)}$ joining the τ -images of the endpoints of that edge. Although the map τ is not quite canonically defined (since it involves choosing a vertex group in a free splitting \mathbb{A} when defining $\tau(\mathbb{A})$), it is easy to check that, for $N \geq 3$, if $\tau': V(FS_N) \rightarrow V(FF_N)$ is another map constructed by the above procedure, then $d(\tau(v), \tau'(v)) \leq 2$ for all $v \in V(FS_N)$.

We prove:

Theorem 1.1. *Let $N \geq 3$. Then the free factor complex FF_N is Gromov-hyperbolic. Moreover, there exists a constant $C > 0$ such that for any two vertices x, y of FS_N and any geodesic $[x, y]$ in $FS_N^{(1)}$ the path $\tau([x, y])$ is C -Hausdorff close to a geodesic $[\tau(x), \tau(y)]$ in $FF_N^{(1)}$.*

To prove Theorem 1.1, we first introduce a new object, called the *free bases graph*, and denoted FB_N , see Definition 4.2 below. The vertices of FB_N are free bases of F_N , up to some natural equivalence. Informally, adjacency in FB_N corresponds to two free bases sharing a common element. We then prove (Proposition 4.3) that the natural map from FB_N to FF_N is a quasi-isometry. Thus to show that FF_N is hyperbolic it suffices to establish hyperbolicity of FB_N . To do the latter we use a hyperbolicity criterion for graphs (Proposition 2.3 below) due to Bowditch [8]. Roughly, this criterion requires that there exists a family of paths $\mathcal{S} = \{g_{x,y}\}_{x,y}$ (where $x, y \in VX$) joining x to y and that there exists a “center”-like map $\Phi: V(X) \times V(X) \times V(X) \rightarrow V(X)$, such that

the pair (\mathcal{G}, Φ) satisfies some nice “thin triangle” type properties, see Definition 2.2 below. Using Bowditch’s criterion we obtain Corollary 2.4 saying that if X, Y are connected graphs, with X hyperbolic and if $f: X \rightarrow Y$ is a surjective Lipschitz graph-map with the property that if $d(f(x), f(y))$ is small then $f([x, y])$ has bounded diameter, then Y is also hyperbolic. Moreover, in this case $f([x, y])$ is uniformly Hausdorff-close to any geodesic $[f(x), f(y)]$ in Y . (See also Theorem 3.11 in [25] and Proposition 3.1 in [9].)

We then construct a surjective Lipschitz map $f: \text{FS}'_N \rightarrow \text{FB}_N$, where FS'_N is the barycentric subdivision of FS_N . The map f restricts to a natural bijection from a subset S of $V(\text{FS}'_N)$, corresponding to N -roses, to the set $V(\text{FB}_N)$ of vertices of FB_N . Thus we may, by abuse of notation, say that $S = V(\text{FB}_N)$ and that $f|_S = \text{Id}_S$. In [16] Handel and Mosher constructed nice paths $g_{x,y}$ given by “folding sequences” between arbitrary vertices x and y of FS'_N , and proved that these paths are quasigeodesics in FS'_N . To apply Corollary 2.4 to the map $f: \text{FS}'_N \rightarrow \text{FB}_N$ it turns out to be enough to show that $f(g_{x,y})$ has bounded diameter if $x, y \in S$ and $d(f(x), f(y)) \leq 1$ in FB_N . To do that we analyze the properties of the Handel–Mosher folding sequences in this specific situation. The construction of $g_{x,y}$ for arbitrary $x, y \in V(\text{FS}'_N)$ is fairly complicated. However, in our situation, we have $x, y \in S$, so that x, y correspond to free bases of F_N . In this case the construction of $g_{x,y}$ becomes much easier and boils down to using standard Stallings foldings (in the sense of [21], [31]) to get from x to y . Verifying that $f(g_{x,y})$ has bounded diameter in FB_N , assuming $d(f(x), f(y)) \leq 1$, becomes a much simpler task. Thus we are able to conclude that FB_N is Gromov-hyperbolic, and, moreover, that $f([x, y])$ is uniformly Hausdorff-close to any geodesic $[f(x), f(y)]$ in FB_N . Using the quasi-isometry between FB_N and FF_N provided by Proposition 4.3, we then obtain the conclusion of Theorem 1.1.

Moreover, as we note in Remark 6.2, our proof of Theorem 1.1 provides a fairly explicit description of quasigeodesics joining arbitrary vertices (i.e. free bases) in FB_N in terms of Stallings foldings.

Acknowledgement. This paper grew out of our discussions during the November 2011 AIM SQuaRE Workshop “Subgroups of $\text{Out}(F_N)$ and their action on outer space” at the American Institute of Mathematics in Palo Alto. We thank AIM and the other participants of the workshop, Thierry Coulbois, Matt Clay, Arnaud Hilion, Martin Lustig and Alexandra Pettet, for the stimulating research environment at the workshop. We are also grateful to Patrick Reynolds for the many helpful comments on the initial version of this paper and to Saul Schleimer for a helpful discussion regarding references for Proposition 2.6. We especially thank the referee for the numerous useful comments and suggestions, and for providing Figure 1 (see Section 6 below) illustrating the relationships and the maps between the main objects considered in this paper.

2. Hyperbolicity criteria for graphs

Convention 2.1. From now on, unless specified otherwise, every connected graph X will be considered as a geodesic metric space with the simplicial metric (where every edge has length 1). As in the introduction, we denote the vertex set of X by $V(X)$. Also, when talking about a connected simplicial complex Z as a metric space, we will in fact mean the 1-skeleton $Z^{(1)}$ of Z endowed with the simplicial metric.

Let X, Y be connected graphs. A *graph-map* from X to Y is a continuous function $f: X \rightarrow Y$ such that $f(V(X)) \subseteq V(Y)$ (so that f takes vertices to vertices), and such that for every edge e of X $f(e)$ is an edge-path in Y (where we allow for an edge-path to be degenerate and to consist of a single vertex). Note that if $f: X \rightarrow Y$ is a graph-map and X' is a subgraph of X then $f|_{X'}: X' \rightarrow Y$ is also a graph-map and $f(X')$ is a subgraph of Y .

We say that a graph-map $f: X \rightarrow Y$ is *L -Lipschitz* (where $L \geq 0$) if for every edge e of X the edge-path $f(e)$ has simplicial length $\leq L$.

We use a characterization of hyperbolicity for a geodesic metric space (X, d_X) that is due to Bowditch [8]. A similar hyperbolicity conditions have been originally stated by Masur and Minsky (see Theorem 2.3 in [23]). A related statement was also obtained by Hamenstädt [14].

Definition 2.2 (Thin triangles structure). Let X be a connected graph. Let $\mathcal{G} = \{g_{x,y} | x, y \in V(X)\}$ be a family of edge-paths in X such that for any vertices x, y of X $g_{x,y}$ is a path from x to y in X . Let $\Phi: V(X) \times V(X) \times V(X) \rightarrow V(X)$ be a function such that for any $a, b, c \in V(X)$,

$$\Phi(a, b, c) = \Phi(b, c, a) = \Phi(c, a, b).$$

Assume, for constants B_1 and B_2 that \mathcal{G} and Φ have the following properties:

- (1) For $x, y \in V(X)$, the Hausdorff distance between $g_{x,y}$ and $g_{y,x}$ is at most B_2 .
- (2) For $x, y \in V(X)$, $g_{x,y}: [0, l] \rightarrow X$, $s, t \in [0, l]$ and $a, b \in V(X)$, assume that

$$d_X(a, g(s)) \leq B_1 \quad \text{and} \quad d_X(b, g(t)) \leq B_1.$$

Then, the Hausdorff distance between $g_{a,b}$ and $g_{x,y}|_{[s,t]}$ is at most B_2 .

- (3) For any $a, b, c \in V(X)$, the vertex $\Phi(a, b, c)$ is contained in a B_2 -neighborhood of $g_{a,b}$.

Then, we say that the pair (\mathcal{G}, Φ) is a (B_1, B_2) -thin triangles structure on X .

The following result is a slightly restated special case of Proposition 3.1 in [8].

Proposition 2.3 (Bowditch). *Let X be a connected graph. For every $B_1 > 0$ and $B_2 > 0$, there is $\delta > 0$ and $H > 0$ so that if (\mathcal{G}, Φ) is a (B_1, B_2) -thin triangles structure on X then X is δ -hyperbolic. Moreover, every path $g_{x,y}$ in \mathcal{G} is H -Hausdorff-close to any geodesic segment $[x, y]$.*

Corollary 2.4. *Let X and Y be connected graphs and assume that X is δ_0 -Gromov-hyperbolic. Let $f : X \rightarrow Y$ be a L -Lipschitz graph-map and $f(V(X)) = V(Y)$. Suppose there are integers $M_1 > 0$ and $M_2 > 0$ so that, for $x, y \in V(X)$, if $d_Y(f(x), f(y)) \leq M_1$ then $\text{diam}_Y f([x, y]) \leq M_2$.*

Then, there exists $\delta_1 > 0$ such that Y is δ_1 -hyperbolic. Moreover, there exists $H > 0$ such that for any vertices x, y of X the path $f([x, y])$ is H -Hausdorff close to any geodesic $[f(x), f(y)]$ in Y .

Proof. For every pair of vertices $a, b \in X$, let $g_{a,b}$ be any geodesic segment $[a, b]$ and let \mathcal{G} be the set of all these paths. Also, for any vertices a, b, c of X let

$$\Phi(a, b, c) = \Phi(b, c, a) = \Phi(c, a, b)$$

be any vertex of X that is at most δ_0 away from each of $[a, b], [b, c], [a, c]$. The hyperbolicity of X implies that (\mathcal{G}, Φ) forms a (b_1, b_2) -thin triangles structure on X for some b_1 and b_2 depending on δ_0 . We now *push* this structure (\mathcal{G}, Φ) forward via the map f .

For any vertex y of Y choose a vertex v_y of X such that $f(v_y) = y$.

For any vertices $y, z \in Y$ choose a geodesic g_{v_y, v_z} from v_y to v_z in X (note that such a geodesic is generally not unique) and let $g'_{y,z} := f(g_{v_y, v_z})$. Then let $\mathcal{G}' = \{g'_{y,z} \mid y, z \in V(Y)\}$. Now, for any vertices $w, y, z \in Y$ put

$$\Phi'(w, y, z) := f(\Phi(v_w, v_y, v_z)).$$

We claim that, the pair (\mathcal{G}', Φ') is a (B_1, B_2) -thin triangles structure for Y for some B_1 and B_2 . The conditions (1) and (3) of Definition 2.2 are satisfied as long as $B_2 \geq L b_2$ since f is L -Lipschitz. Thus we only need to verify that condition (2) of Definition 2.2 holds for (\mathcal{G}', Φ') .

Let y, z be vertices of Y , v_y, v_z be the associated vertices in X , $g_{v_y, v_z} : [0, l] \rightarrow X$ be the path in \mathcal{G} connecting v_x to v_y and $g'_{y,z} : [0, l'] \rightarrow Y$ be the f -image of g_{v_y, v_z} . In the interest of brevity, we denote these paths simply by g and g' .

Let $B_1 = M_1$ and let, $a, b \in Y$ and $s', t' \in [0, l']$ be such that

$$d_Y(a, g'(s')) \leq B_1 \quad \text{and} \quad d_Y(b, g'(t')) \leq B_1.$$

We need to bound the Hausdorff distance between $g'_{a,b} = f(g_{v_a, v_b})$ and $g'|_{[s', t']}$.

Let $s, t \in [0, l]$ be such that $fg(s) = g'(s')$ and $fg(t) = g'(t')$. Let u be a vertex of $g'_{a,b}$. From hyperbolicity, we have v_u is contained in a $2\delta_0$ -neighborhood of the union

$$g|_{[s, t]} \cup [g(s), v_a] \cup [g(t), v_b].$$

Thus u is $(2L \delta_0)$ -close to the union

$$g'|_{[s', t']} \cup f([g(s), v_a]) \cup f([g(t), v_b]).$$

But the d_Y -diameters of $f([g(s), v_a])$ $f([g(t), v_b])$ are less than M_2 . Hence, u is in $(2L\delta_0 + M_2)$ -neighborhood of $g'|_{[s',t']}$. Similarly, $g'|_{[s',t']}$ is in the same size neighborhood of $g'_{a,b}$. The condition (2) of Definition 2.2 holds for $B_2 = (2L\delta_0 + M_2)$.

Therefore, by Proposition 2.3, the graph Y is δ_1 -hyperbolic, and, moreover, for any two vertices y, z of Y the path $g'_{y,z} = f(g_{v_y,v_z})$ is H' -Hausdorff close to $[y, z]$ for some constant $H' \geq 0$ independent of y, z . Since g_{v_y,v_z} was chosen to be a geodesic from v_y to v_z in X and any two such geodesics are δ_0 -Hausdorff close, by increasing the constant H' we also get that for any geodesic $[y, z]$ from y to z in Y and any geodesic $[v_y, v_z]$ from v_y to v_z in X the paths $[y, z]$ and $f([v_y, v_z])$ are H' -Hausdorff close in Y .

Now let y, z be any vertices of Y and let v'_y, v'_z be arbitrary vertices of X such that $f(v'_y) = y$ and $f(v'_z) = z$. A geodesic $[v'_y, v'_z]$ in X is contained in the $2\delta_0$ -neighborhood of $[v'_y, v_y] \cup [v_y, v_z] \cup [v_z, v'_z]$. Since $f(v_y) = f(v'_y) = y$ and $f(v_z) = f(v'_z) = z$, the assumptions on f imply that $f([v'_y, v_y])$ is contained in the M_2 -ball around y and $f([v_z, v'_z])$ is contained in the M_2 -ball around z in Y . Moreover, we have already shown that $f([v_y, v_z])$ is H' -Hausdorff close to $[y, z]$. Therefore $f([v'_y, v'_z])$ is contained in the H -neighborhood of $[y, z] = [f(v'_y), f(v'_z)]$ with $H = 2L\delta_0 + M_2 + H'$. A similar argument shows that $[y, z] = [f(v'_y), f(v'_z)]$ is contained in the H -neighborhood of $f([v'_y, v'_z])$. Thus $[y, z] = [f(v'_y), f(v'_z)]$ and $f([v'_y, v'_z])$ are H -Hausdorff close, as required. \square

It is also possible to derive Corollary 2.4 using Proposition 3.1 from a new paper of Bowditch (December 2012, [9]) instead of Bowditch’s hyperbolicity criterion from [8]. Proposition 3.1 of [9] gives a strengthened form of Theorem 3.11 of Masur–Schleimer [25]. Theorem 3.11 of [25] is sufficient to conclude that the graph Y in Corollary 2.4 is Gromov-hyperbolic; however, the existence of the constant H as in the conclusion of Corollary 2.4 requires Proposition 3.1 of [9].

Proposition 2.5. *For any positive integers δ_0, L, M and D there exist $\delta_1 > 0$ and $H > 0$ so that the following holds.*

Let X, Y be connected graphs, such that X is δ_0 -hyperbolic. Let $f : X \rightarrow Y$ be a L -Lipschitz graph map for some $L \geq 0$. Let $S \subseteq V(X)$ be such that the following holds.

- (1) *We have $f(S) = V(Y)$.*
- (2) *The set S is D -dense in X for some $D > 0$.*
- (3) *For $x, y \in S$, if $d(f(x), f(y)) \leq 1$ then for any geodesic $[x, y]$ in X we have*

$$\text{diam}_Y(f([x, y])) \leq M.$$

Then Y is δ_1 -hyperbolic and, for any $x, y \in V(X)$ and any geodesic $[x, y]$ in X , the path $f([x, y])$ is H -Hausdorff close to any geodesic $[f(x), f(y)]$ in Y .

Proof. First we show that, for every $m_1 > 0$, there is $m_2 > 0$ so that whenever $x, y \in S$ satisfy $d_Y(f(x), f(y)) \leq m_1$ then $\text{diam}_Y(f([x, y])) \leq m_2$. Indeed let $x, y \in S$ be as above and consider a geodesic path $[f(x), f(y)]$ in Y . Let

$$f(x) = z_0, z_1, \dots, z_t = f(y), \quad t < m_1,$$

be the sequence of consecutive vertices on $[f(x), f(y)]$. Let $x_0 = x, x_t = y$ and for $1 \leq i \leq t - 1$ let $x_i \in S$ be such that $f(x_i) = z_i$. Such x_i exist since by assumption $f(S) = V(Y)$. We have $\text{diam}_Y f([x_i, x_{i+1}]) \leq M$. By hyperbolicity, the geodesic $[x, y]$ is contained in the $(m_1\delta_0)$ -neighborhood of the union

$$\bigcup_{i=0}^{t-1} [x_i, x_{i+1}].$$

Since f is L -Lipschitz, $f([x, y])$ is contained in the $(Lm_1\delta_0)$ -neighborhood of

$$\bigcup_{i=0}^{t-1} f([x_i, x_{i+1}]).$$

But each $f([x_i, x_{i+1}])$ has diameter $\leq M$. Therefore, $f([x, y])$ has a diameter of at most $m_2 = (m_1M + 2Lm_1\delta_0)$.

Now let $M_1 \geq 0$ and $x, y \in V(X)$ be arbitrary vertices with $d_Y(f(x), f(y)) \leq M_1$. Since S is D -dense in X , there exist $x', y' \in S$ such that $d(x, x'), d(y, y') \leq D$. The fact that f is L -Lipschitz implies that $d(f(x'), f(y')) \leq M_1 + 2DL$. Therefore, by the above claim, it follows that the

$$\text{diam}_Y f([x', y']) \leq m_2(M_1 + 2DL).$$

Since X is δ_0 -hyperbolic and $d(x, x'), d(y, y') \leq D$, we have that $[x, y]$ and $[x', y']$ are $(2\delta + 2D)$ -Hausdorff close. Again, using that f is L -Lipschitz, we conclude that $f([x, y])$ has a diameter of at most

$$M_2 = m_2(M_1 + 2DL) + 4L(2\delta_0 + 2D).$$

The assumption of Corollary 2.4 are now satisfied for constants δ_0, L, M_1 and M_2 . Proposition 2.5 now follows from Corollary 2.4. □

Proposition 2.5 easily implies the well-known fact that “coning-off” or “electrifying” a family of uniformly quasiconvex subsets in a hyperbolic space produces a hyperbolic space. Various versions of this statement have multiple appearances in the literature; see, for example, Lemma 4.5 and Proposition 4.6 in [13], Proposition 7.12 in [7], Lemma 2.3 in [28], Theorem 3.4 in [12], etc. Maher and Schleimer [22] appear to be the first to explicitly note that after “electrifying” a family of uniformly quasiconvex subsets in a hyperbolic space, not only is the resulting space again hyperbolic, but the image of a geodesic is a reparameterized quasigeodesic.

We give a version of the “coning-off” statement here phrased in the context of graphs with simplicial metrics.

Proposition 2.6. *Let X be a connected graph with simplicial metric d_X such that (X, d_X) is δ_0 -hyperbolic. Let $C > 0$ and $(X_j)_{j \in J}$ be a family of subgraphs of X such that each X_j is a C -quasiconvex subset of X . Let Y be the graph obtained from X by adding to X the new edges $e_{x,y,j}$ with endpoints x, y whenever $j \in J$ and x, y are vertices of X_j (thus X is a subgraph of Y). Let d_Y be the simplicial metric on Y .*

Then Y is δ_1 -hyperbolic for some constant $\delta_1 > 0$ depending only on C and δ_0 . Moreover there exists $H = H(C, \delta_0) > 0$ such that whenever $x, y \in V(X)$, $[x, y]_X$ is a d_X -geodesic from x to y in X and $[x, y]_Y$ is a d_Y -geodesic from x to y in Y then $[x, y]_X$ and $[x, y]_Y$ are H -Hausdorff close in (Y, d_Y) .

Proof. Let $f : X \rightarrow Y$ be the inclusion map and put $S = V(X)$. We claim that the conditions of Proposition 2.5 are satisfied.

By construction $V(X) = V(Y)$, so $f(S) = V(Y)$. Also, the map f is obviously 1-Lipschitz. Suppose now that $x, y \in V(X)$ are such that $d_Y(x, y) \leq 1$. If $d_Y(x, y) < 1$ then $d_Y(x, y) = 0$, so that $x = y$. In this case it is obvious that condition (3) of Proposition 2.5 holds for $f([x, y]_X)$. Suppose now that $d_Y(x, y) = 1$. Thus there exists an edge e in Y with endpoints x, y . If e is an edge of X then $d_X(x, y) = 1$ and it is again obvious that condition (3) of Proposition 2.5 holds for $f([x, y]_X)$. Suppose now that $e = e_{x,y,j}$ for some $j \in J$. Thus $x, y \in V(X_j)$. Since X_j is C -quasiconvex in X , we see that for any point u on $[x, y]_X$ there exists a vertex z of X_j with $d_X(u, z) \leq C$. Hence $d_Y(u, z) \leq C$ as well. We have $d_Y(x, z) \leq 1$ and $d_Y(z, y) \leq 1$ since x, y, z are vertices of X_j . Hence $d_Y(f(u), x) = d_Y(u, x) \leq C + 1$. Thus $f([x, y]_X)$ is contained in the d_Y -ball of radius $C + 1$ centered at x in (Y, d_Y) , and again condition (3) of Proposition 2.5 holds.

Thus Proposition 2.5 applies and the conclusion of Proposition 2.6 follows. \square

Note that the assumptions of Proposition 2.6 do not require the family $(X_j)_{j \in J}$ to be “sufficiently separated”. Such a requirement is present in many versions of Proposition 2.6 available in the literature, although this assumption is not in fact necessary and, in particular, Proposition 7.12 in [7] does not impose the “sufficiently separated” requirement. We have derived Proposition 2.6 from Proposition 2.5, which in turn was a consequence of Corollary 2.4. A close comparison of these statements show that the converse implication does not work, and that Corollary 2.4 is a more general statement than Proposition 2.6.

3. Free factor complex and free splitting complex

Definition 3.1 (Free factor complex). Let F_N be a free group of finite rank $N \geq 3$.

The *free factor complex* FF_N of F_N is a simplicial complex defined as follows. The set of vertices $V(\text{FF}_N)$ of FF_N is defined as the set of all F_N -conjugacy classes

$[A]$ of proper free factors A of F_N . Two distinct vertices $[A]$ and $[B]$ of FF_N are joined by an edge whenever there exist proper free factors A, B of F_N representing $[A]$ and $[B]$ respectively, such that either $A \leq B$ or $B \leq A$.

More generally, for $k \geq 1$, a collection of $k + 1$ distinct vertices $[A_0], \dots, [A_k]$ of FF_N spans a k -simplex in FF_N if, up to a possible re-ordering of these vertices there exist representatives A_i of $[A_i]$ such that $A_0 \leq A_1 \leq \dots \leq A_k$.

There is a canonical action of $\text{Out}(F_N)$ on FF_N by simplicial automorphisms: If

$$\Delta = \{[A_0], \dots, [A_k]\}$$

is a k simplex and $\varphi \in \text{Out}(F_N)$, then

$$\varphi(\Delta) := \{[\varphi(A_0)], \dots, [\varphi(A_k)]\}.$$

It is not hard to check that for $N \geq 3$ the complex FF_N is connected, has dimension $N - 2$ and that $\text{FF}_N/\text{Out}(F_N)$ is compact.

Definition 3.2 (Free splitting complex). Let F_N be a free group of finite rank $N \geq 3$.

The *free splitting complex* FS_N is a simplicial complex defined as follows. The vertex set $V(\text{FS}_N)$ consists of equivalence classes of splittings $F_N = \pi_1(\mathbb{A})$, where \mathbb{A} is a graph of groups with a single topological edge e (possibly a loop edge) and the trivial edge group such that the action of F_N on the Bass–Serre tree $T_{\mathbb{A}}$ is minimal (i.e. such that if e is a non-loop edge then both vertex groups in \mathbb{A} are nontrivial). Two such splittings $F_N = \pi_1(\mathbb{A})$ and $F_N = \pi_1(\mathbb{B})$ are equivalent if there exists an F_N -equivariant isometry between $T_{\mathbb{A}}$ and $T_{\mathbb{B}}$. We denote the equivalence class of a splitting $F_N = \pi_1(\mathbb{A})$ by $[\mathbb{A}]$.

The edges in FS_N correspond to two splittings admitting a common refinement. Thus two distinct vertices $[\mathbb{A}]$ and $[\mathbb{B}]$ of FS_N are joined by an edge whenever there exists a splitting $F_N = \pi_1(\mathbb{D})$ such that the graph of groups \mathbb{D} has exactly two topological edges, both with trivial edge groups, and such that collapsing one of these edges produces a splitting of F_N representing $[\mathbb{A}]$ and collapsing the other edge produces a splitting representing $[\mathbb{B}]$.

More generally, for $k \geq 1$ a collection of $k + 1$ distinct vertices $[\mathbb{A}_0], \dots, [\mathbb{A}_k]$ of FS_N spans a k -simplex in FS_N whenever there exists a splitting $F_N = \pi_1(\mathbb{D})$ such that the graph of groups \mathbb{D} has the following properties:

- (a) The underlying graph of \mathbb{D} has exactly $k + 1$ topological edges, e_0, \dots, e_k .
- (b) The edge group of each e_i is trivial.
- (c) For each $i = 0, \dots, k$ collapsing all edges except for e_i in \mathbb{D} produces a splitting of F_N representing $[\mathbb{A}_i]$.

The complex FS_N comes equipped with a natural action of $\text{Out}(F_N)$ by simplicial automorphisms.

Again, it is not hard to check that for $N \geq 3$ the complex FS_N is finite-dimensional, connected and that the quotient FS_N/F_N is compact.

We denote the barycentric subdivision of FS_N by FS'_N .

Definition 3.3 (Marking). Let $N \geq 2$. Recall that a *marking* on F_N is an isomorphism $\alpha: F_N \rightarrow \pi_1(\Gamma, v)$ where Γ is a finite connected graph without any degree-one and degree-two vertices and v is a vertex of Γ . By abuse of notation, if α is specified, we will often refer to Γ as a marking.

Two markings $\alpha: F_N \rightarrow \pi_1(\Gamma, v)$ and $\alpha': F_N \rightarrow \pi_1(\Gamma', v')$ are said to be equivalent, if there exists an F_N -equivariant isometry $(\widetilde{\Gamma}, v) \rightarrow (\widetilde{\Gamma'}, v')$. The equivalence class of a marking $\alpha: F_N \rightarrow \pi_1(\Gamma, v)$ is denoted by $[\alpha]$ or, if α is already specified, just $[\Gamma]$.

Convention 3.4 (Barycenters). Note that for $N \geq 3$ any marking $\alpha: F_N \rightarrow \pi_1(\Gamma)$ corresponds to a simplex Δ_α in FS_N , as follows. We can view Γ as a graph of groups by assigning trivial groups to all the vertices and edges of Γ . Then the vertices of Δ_α correspond to the (topological) edges of Γ and come from choosing an edge e of Γ and collapsing all the other edges of Γ . It is easy to see that Δ_α depends only on the equivalence class $[\alpha]$ of the marking α .

We denote the vertex of FS'_N given by the barycenter of Δ_α by $z(\alpha)$ or, if it is more convenient, by $z(\Gamma)$. Note that if $[\alpha] = [\beta]$ then $z(\alpha) = z(\beta)$. We will sometimes refer to a marking Γ as a vertex of FS'_N ; when that happens, we always mean the vertex $z(\Gamma)$.

4. The free bases graph

If Γ is a graph (i.e. a one-dimensional CW-complex), then any topological edge (i.e. a closed 1-cell) of Γ is homeomorphic to either $[0, 1]$ or to S^1 and thus admits exactly two orientations. An *oriented edge* of Γ is a topological edge together with a choice of an orientation on this edge. If e is an oriented edge of Γ , we denote by e^{-1} the oriented edge obtained by changing the orientation on e to the opposite one. Note that $(e^{-1})^{-1} = e$ for any oriented edge e . For an oriented edge e we denote the initial vertex of e by $o(e)$ and the terminal vertex of e by $t(e)$. Then $o(e^{-1}) = t(e)$ and $t(e^{-1}) = o(e)$. We will denote by $E\Gamma$ the set of oriented edges of Γ and by $V\Gamma$ the set of vertices of Γ .

Let $N \geq 2$. We denote by W_N the graph with a single vertex v_0 and N distinct oriented loop-edges e_1, \dots, e_N .

Definition 4.1 (\mathcal{A} -rose). Let $\mathcal{A} = \{a_1, \dots, a_N\}$ be a free basis of F_N . Define the \mathcal{A} -rose $R_{\mathcal{A}}$ as the marking $\alpha_{\mathcal{A}}: F_N \rightarrow \pi_1(W_N, v_0)$ where $\alpha_{\mathcal{A}}$ sends a_i to the loop at v_0 in Γ corresponding to e_i , traversed in the direction given by the orientation of e_i .

Definition 4.2 (Free bases graph). Let $N \geq 3$. The *free bases graph* FB_N of F_N is a simple graph defined as follows. The vertex set $V(\text{FB}_N)$ consists of equivalence classes free bases \mathcal{A} of F_N . Two free bases \mathcal{A} and \mathcal{B} of F_N are considered equivalent if the Cayley graphs $T_{\mathcal{A}}$ and $T_{\mathcal{B}}$ of F_N with respect to \mathcal{A} and \mathcal{B} are F_N -equivariantly isometric. We denote the equivalence class of a free basis \mathcal{A} of F_N by $[\mathcal{A}]$.

Note that for free bases $\mathcal{A} = \{a_1, \dots, a_N\}$ and $\mathcal{B} = \{b_1, \dots, b_N\}$ of F_N we have $[\mathcal{A}] = [\mathcal{B}]$ if and only if there exist a permutation $\sigma \in S_N$, an element $g \in F_N$ and numbers $\varepsilon_i \in \{1, -1\}$ (where $i = 1, \dots, N$) such that

$$b_i = g^{-1} a_{\sigma(i)}^{\varepsilon_i} g$$

for $i = 1, \dots, N$. Thus $[\mathcal{A}] = [\mathcal{B}]$ if and only if the roses $R_{\mathcal{A}}$ and $R_{\mathcal{B}}$ are equivalent as markings. Note also that for any free basis \mathcal{A} of F_N and any $g \in F_N$ we have $[g^{-1} \mathcal{A} g] = [\mathcal{A}]$.

The edges in FB_N are defined as follows. Let $[\mathcal{A}]$ and $[\mathcal{B}]$ be two distinct vertices of FB_N . These vertices are adjacent in FB_N whenever there exists $a \in \mathcal{A}$ such that some element $b \in \mathcal{B}$ is conjugate to a or a^{-1} . Thus two distinct vertices v_1, v_2 of FB_N are adjacent if and only if there exist free bases \mathcal{A} and \mathcal{B} representing v_1, v_2 accordingly such that $\mathcal{A} \cap \mathcal{B} \neq \emptyset$.

The graph FB_N comes equipped with a natural $\text{Out}(F_N)$ action by simplicial automorphisms.

Proposition 4.3. *Let $N \geq 3$. Then the following holds.*

- (1) *The graph FB_N is connected.*
- (2) *For each vertex $v = [\mathcal{A}]$ of FB_N choose some $a_v \in \mathcal{A}$. Consider the map*

$$h: V(\text{FB}_N) \rightarrow V(\text{FF}_N)$$

defined as $h(v) = [\{a_v\}]$ for every vertex v of FB_N . Extend h to a graph-map

$$h: \text{FB}_N \rightarrow \text{FF}_N$$

by sending every edge e of FB_N with endpoints v, v' to a geodesic path $[h(v), h(v')]$ in $\text{FF}_N^{(1)}$. Then:

- (a) *The map h is a quasi-isometry. In particular, the complexes FB_N and FF_N are quasi-isometric.*
- (b) *The set $h(V(\text{FB}_N))$ is 3-dense in $\text{FF}_N^{(1)}$*

Proof. First we will show that h is 4-Lipschitz. Since $h: \text{FB}_N \rightarrow \text{FF}_N^{(1)}$ is a graph-map, it suffices to check that for any two adjacent vertices v, v' of FB_B we have $d_{\text{FF}_N}(h(v), h(v')) \leq 4$.

Let $v = [\mathcal{A}]$ and $v' = [\mathcal{B}]$ be two adjacent vertices of FB_N . Hence we may choose free bases \mathcal{A} representing v and \mathcal{B} representing v' such that $\mathcal{A} \cap \mathcal{B} \neq \emptyset$. Up to re-ordering these bases, we may assume that $\mathcal{A} = \{a_1, \dots, a_N\}$, $\mathcal{B} = \{b_1, \dots, b_N\}$

and that $a_1 = b_1$. Then $a_v = a_i$ and $a_{v'} = b_j$ for some $1 \leq i, j \leq N$ and thus, by definition of h , we have $h(v) = [\langle a_i \rangle]$, $h(v') = [\langle b_j \rangle]$.

We will assume that $i > 1$ and $j > 1$ as the cases where $i = 1$ or $j = 1$ are easier. Then in FF_N we have

$$d_{\text{FF}_N}([\langle a_i \rangle], [\langle a_i, a_1 \rangle]) = d_{\text{FF}_N}([\langle a_i, a_1 \rangle], [\langle a_1 \rangle]) = 1$$

and

$$d_{\text{FF}_N}([\langle b_j \rangle], [\langle b_j, b_1 \rangle]) = d_{\text{FF}_N}([\langle b_j, b_1 \rangle], [\langle b_1 \rangle]) = 1.$$

Since $a_1 = b_1$, by the triangle inequality we conclude that

$$d_{\text{FF}_N}(h(v), h(v')) = d_{\text{FF}_N}([\langle a_i \rangle], [\langle b_j \rangle]) \leq 4.$$

Thus the map h is 4-Lipschitz, as claimed.

To show that h is a quasi-isometry we will construct a “quasi-inverse”, that is a Lipschitz map $q: \text{FF}_N^{(1)} \rightarrow \text{FB}_N$ such that there exists $C \geq 0$ with the property that for every vertex v of FB_N , $d_{\text{FB}_N}(v, q(h(v))) \leq C$ and that for every vertex u of FF_N , $d_{\text{FF}_N}(u, h(q(u))) \leq C$.

We define q on $V(\text{FF}_N)$ and then extend q to edges in a natural way, by sending every edge to a geodesic joining the images of its end-vertices.

Let $u = [K]$ be an arbitrary vertex of FF_N (so that K is a proper free factor of F_N). We choose a free basis \mathcal{B}_K of K and then a free basis \mathcal{A}_K of F_N such that $\mathcal{B}_K \subseteq \mathcal{A}_K$. Put $q(u) = [\mathcal{A}_K]$.

First we check that q is Lipschitz. Let $u = [K]$ and $u' = [K']$ be adjacent vertices of FF_N , where K, K' are proper free factors of F_N . We may assume that $K \leq K'$ is a proper free factor of K' . Since $K' \neq F_N$, there exists $t \in \mathcal{A}_{K'} \setminus \mathcal{B}_{K'}$. Since K is a free factor of K' , we can find a free basis \mathcal{A} of F_N such that $t \in \mathcal{A}$ and $\mathcal{B}_K \subseteq \mathcal{A}$. Since $t \in \mathcal{A}_{K'} \cap \mathcal{A}$, we have $d([\mathcal{A}_{K'}], [\mathcal{A}]) \leq 1$ in FB_N . Since $\mathcal{B}_K \subset \mathcal{A} \cap \mathcal{A}_K$, it follows that $d([\mathcal{A}_K], [\mathcal{A}]) \leq 1$ in FB_N . Therefore

$$d_{\text{FB}_N}(q(u), q(u')) = d_{\text{FB}_N}([\mathcal{A}_K], [\mathcal{A}_{K'}]) \leq 2.$$

Hence q is 2-Lipschitz.

For a vertex $v = [\mathcal{A}]$ of FB_N let us now estimate $d_{\text{FB}_N}(v, q(h(v)))$. We have $h(v) = [\langle a_v \rangle]$ for some $a_v \in \mathcal{A}$. The group $K = \langle a_v \rangle$ is infinite cyclic (that is free of rank 1). Therefore this group has only two possible free bases, $\{a_v\}$ and $\{a_v^{-1}\}$. We will assume that $\mathcal{B}_K = \{a_v\}$ as the case $\mathcal{B}_K = \{a_v^{-1}\}$ is similar. Then, by definition, \mathcal{A}_K is a free basis of F_N containing a_v and $q(h(v)) = q([K]) = [\mathcal{A}_K]$. Thus $a_v \in \mathcal{A} \cap \mathcal{A}_K$ and hence $d_{\text{FB}_N}(v, q(h(v))) \leq 1$ in FB_N .

Now let $u = [K]$ be an arbitrary vertex of FF_N . We need to estimate the distance $d_{\text{FF}_N}(u, h(q(u)))$. By definition, $v := h(u) = [\mathcal{A}_K]$ where \mathcal{A}_K is a free basis of F_N containing as a (proper) subset a free basis \mathcal{B}_K of K . Then $a_v \in \mathcal{A}_K$ and

$h(q(u)) = h(v) = [\langle a_v \rangle]$. Choose an element $b \in \mathcal{B}_K$. It may happen that $b = a_v$, but in any case $K' := \langle b, a_v \rangle$ is a proper free factor of F_N . Then

$$d_{\text{FF}_N}([K], [\langle b \rangle]) \leq 1, \quad d_{\text{FF}_N}([\langle b \rangle], [K']) \leq 1 \quad \text{and} \quad d_{\text{FF}_N}([K'], [\langle a_v \rangle]) \leq 1.$$

Therefore

$$d_{\text{FF}_N}(u, h(q(u))) = d_{\text{FF}_N}([K], [\langle a_v \rangle]) \leq 3.$$

Thus indeed q is a quasi-inverse for h , and hence h is a quasi-isometry, as required.

We next show that $h(V(\text{FB}_N))$ is 3-dense in $\text{FF}_N^{(1)}$. Indeed, let $K \leq F_N$ be an arbitrary proper free factor of F_N . Let a_1, \dots, a_m (where $1 \leq m < N$) be a free basis of K and choose a_{m+1}, \dots, a_N such that $\mathcal{A} = \{a_1, \dots, a_N\}$ is a free basis of F_N . Then $h([\mathcal{A}]) = [\langle a_i \rangle]$ for some $1 \leq i \leq N$. In FF_N we have

$$d_{\text{FF}_N}([K], [\langle a_1 \rangle]) \leq 1,$$

$$d_{\text{FF}_N}([\langle a_1 \rangle], [\langle a_1, a_i \rangle]) \leq 1$$

and

$$d_{\text{FF}_N}([\langle a_1, a_i \rangle], [\langle a_i \rangle]) \leq 1.$$

Since $h([\mathcal{A}]) = [\langle a_i \rangle]$, it follows that $d_{\text{FF}_N}([K], h(v)) \leq 3$. Thus indeed $h(V(\text{FB}_N))$ is 3-dense in $\text{FF}_N^{(1)}$, as claimed. \square

Definition 4.4 (Free basis defined by a marking). If $\alpha: F_N \rightarrow \pi_1(\Gamma, v)$ is a marking, and $T \subseteq \Gamma$ is a maximal tree in Γ , there is a naturally associated free basis $\mathcal{B}(\alpha, T)$ (which we will also sometimes denote $\mathcal{B}(\Gamma, T)$) of F_N . Namely, in this case $\Gamma - T$ consists of N topological edges. Choose oriented edges $e_1, \dots, e_N \in E(\Gamma - T)$ so that $E(\Gamma - T) = \{e_1^{\pm 1}, \dots, e_N^{\pm 1}\}$. For $j = 1, \dots, N$ put

$$\gamma_j = [v, o(e_j)]_T e_j [t(e_j), v]_T,$$

where $[u, u']_T$ denotes the (unique) geodesic in the tree T from u to u' for $u, u' \in V(\Gamma) = V(T)$.

Then $\gamma_1, \dots, \gamma_N$ is a free basis of $\pi_1(\Gamma, v)$. Put

$$\mathcal{B}(\alpha, T) := \{\alpha^{-1}(\gamma_1), \dots, \alpha^{-1}(\gamma_N)\}.$$

Remark 4.5. One can show that there is a constant $C = C(N) > 0$ such that if $\alpha: F_N \rightarrow \pi_1(\Gamma, v)$ and $\alpha': F_N \rightarrow \pi_1(\Gamma', v')$ are equivalent markings and $T \subseteq \Gamma$, $T' \subseteq \Gamma'$ are maximal trees, then

$$d_{\text{FB}_N}([\mathcal{B}(\alpha, T)], [\mathcal{B}(\alpha', T')]) \leq C.$$

This can be shown, for example, using the quasi-isometry $q: \text{FB}_N \rightarrow \text{FF}_N$ constructed in Proposition 4.3. Thus the definitions imply that if T is a maximal tree in Γ

and e is an edge of $\Gamma \setminus T$, then $q([\mathcal{B}(\alpha, T)])$ is a bounded distance away in FF_N from the free factor of F_N corresponding to any of the vertex groups in the graph of groups Γ_e obtained by collapsing $\Gamma \setminus e$. On the other hand, for any two edges e_1, e_2 of Γ the free splittings Γ_{e_1} and Γ_{e_2} are adjacent vertices of FS_N and therefore (e.g. using the Lipschitz map $\tau : \text{FS}_N \rightarrow \text{FF}_N$ from the Introduction), any two vertex groups A_1 and A_2 from these splittings are bounded distance away in FF_N .

5. \mathcal{A} -graphs and Stallings folds

We briefly discuss here the language and machinery of Stallings foldings, introduced by Stallings in a seminal paper [31]. We refer the reader to [21] for detailed background on the topic.

If Γ is a finite connected non-contractible graph, we denote by $\text{Core}(\Gamma)$ the unique minimal subgraph of Γ such that the inclusion $\text{Core}(\Gamma) \subseteq \Gamma$ is a homotopy equivalence. Thus $\text{Core}(\Gamma)$ carries $\pi_1(\Gamma)$ and we can obtain Γ from $\text{Core}(\Gamma)$ by attaching finitely many trees.

Definition 5.1 (\mathcal{A} -graph). Let \mathcal{A} be a free basis of F_N and let $R_{\mathcal{A}}$ be the corresponding rose marking. An \mathcal{A} -graph is a graph Γ with a labelling function $\mu : E\Gamma \rightarrow \mathcal{A}^{\pm 1}$ (where $E\Gamma$ is the set of oriented edges of Γ) such that for every oriented edge $e \in E\Gamma$ we have $\mu(e^{-1}) = \mu(e)^{-1}$.

Note that there is an obvious way to view the rose $R_{\mathcal{A}}$ as an \mathcal{A} -graph. Any \mathcal{A} -graph Γ comes equipped with a canonical label-preserving graph-map $p : \Gamma \rightarrow R_{\mathcal{A}}$ which sends all vertices of Γ to the (unique) vertex of $R_{\mathcal{A}}$ and which sends every oriented edge of Γ to the oriented edge of the rose $R_{\mathcal{A}}$ with the same label. We call p the *natural projection*.

Let Γ be a finite connected \mathcal{A} -graph containing at least one vertex of degree ≥ 3 . Following Handel–Mosher [16], we call vertices of Γ that have degree ≥ 3 *natural vertices*. The complement of the set of natural vertices in Γ consists of a disjoint union of intervals whose closures, again following [16], we call *natural edges*.

Recall that in the definition of a marking on F_N the graph appearing in that definition had no degree-one and degree-two vertices.

Remark 5.2. Suppose that Γ is a connected \mathcal{A} -graph such that the natural projection $\Gamma \rightarrow R_{\mathcal{A}}$ is a homotopy equivalence. Then the projection $p : \text{Core}(\Gamma) \rightarrow R_{\mathcal{A}}$ is a homotopy equivalence.

Then, via using the homotopy inverse of p and making inverse subdivisions in $\text{Core}(\Gamma)$ to erase all the degree-2 vertices, we get an actual marking of F_N , $\alpha : F_N \rightarrow \pi_1(\bar{\Gamma})$. Here $\bar{\Gamma}$ is the graph obtained from $\text{Core}(\Gamma)$ by doing inverse edge-subdivisions to erase all degree-two vertices. In this case we call α the *marking associated with Γ* and denote α by α_{Γ} , or, sometimes just by $\bar{\Gamma}$.

Definition 5.3 (Folded graphs and Stallings folds). Let Γ be an \mathcal{A} -graph. We say that Γ is *folded* if there does not exist a vertex v of Γ and two distinct oriented edges e_1, e_2 with $o(e_1) = o(e_2) = v$ such that $\mu(e_1) = \mu(e_2)$. Otherwise we say that Γ is *non-folded*.

Let Γ be a non-folded \mathcal{A} -graph, let e_1, e_2 be two distinct oriented edges of Γ such that $o(e_1) = o(e_2) = v \in V(\Gamma)$ and such that $\mu(e_1) = \mu(e_2) = a \in \mathcal{A}^{\pm 1}$. Construct an \mathcal{A} -graph Γ' by identifying the edges e_1 and e_2 into a single edge e with label $\mu(e) = a$. We say that Γ' is obtained from Γ by a *Stallings fold*. In this case there is also a natural label-preserving *fold map* $f: \Gamma \rightarrow \Gamma'$. It is easy to see that the fold map f is a homotopy equivalence if and only if $t(e_1) \neq t(e_2)$ in Γ . If $t(e_1) \neq t(e_2)$ in Γ , we say that f is a *type-I Stallings fold*. If $t(e_1) = t(e_2)$ in Γ , we say that f is a *type-II Stallings fold*.

Note that if Γ is a finite connected \mathcal{A} -graph such that the natural projection $\Gamma \rightarrow R_{\mathcal{A}}$ is a homotopy equivalence, and if Γ' is obtained from Γ by a Stallings fold f , then f is necessarily a type-I fold, and hence the natural projection $\Gamma' \rightarrow R_{\mathcal{A}}$ is again a homotopy equivalence.

Definition 5.4 (Maximal fold). Let Γ be a non-folded finite connected \mathcal{A} -graph, let $v \in VA$ be a natural vertex, let e_1, e_2 be two distinct oriented edges of Γ such that $o(e_1) = o(e_2) = v$ and such that $\mu(e_1) = \mu(e_2) = a \in \mathcal{A}^{\pm 1}$. Let \widehat{e}_1 and \widehat{e}_2 be the natural edges in Γ that begin with e_1, e_2 accordingly. Let z_1, z_2 be maximal initial segments of \widehat{e}_1 and \widehat{e}_2 such that the label $\mu(z_1)$ is graphically equal, as a word over $\mathcal{A}^{\pm 1}$, to the label $\mu(z_2)$. Thus z_1 starts with e_1 and z_2 starts with e_2 . Let Γ' be obtained from Γ by a chain of Stallings folds that fold z_1 and z_2 together. We say that Γ' is obtained from Γ by a *maximal fold*. Being a composition of several Stallings folds, a maximal fold also comes equipped with a *fold map* $f: \Gamma \rightarrow \Gamma'$.

Remark 5.5. Let Γ be a connected \mathcal{A} -graph such that $\Gamma = \text{Core}(\Gamma)$ and such that the natural projection $p: \Gamma \rightarrow R_{\mathcal{A}}$ is a homotopy equivalence. Let $\alpha: F_N \rightarrow \pi_1(\Gamma, v)$ be an associated marking. Let $T \subseteq \Gamma$ be a maximal tree.

Recall that according to Definition 4.4, we have an associated free basis $\mathcal{B}(\Gamma, T)$ of F_N . In this case $\mathcal{B}(\Gamma, T)$ can be described more explicitly as follows. Choose oriented edges $e_1, \dots, e_N \in E(\Gamma - T)$ so that $E(\Gamma - T) = \{e_1^{\pm 1}, \dots, e_N^{\pm 1}\}$. Let w_j be the label (i.e. a word over \mathcal{A}) of the path $[v, o(e_j)]_T e_j [t(e_j), v]_T, j = 1, \dots, N$. Then $\mathcal{B}(\Gamma, T) = \{w_1, \dots, w_N\}$.

We need the following technical notion which is a variant of the notion of a foldable map from the paper of Handel–Mosher [16].

Definition 5.6 (Foldable maps). Let Γ be a finite connected \mathcal{A} -graph such that the natural projection $p: \Gamma \rightarrow R_{\mathcal{A}}$ is a homotopy equivalence and such that $\Gamma = \text{Core}(\Gamma)$.

We say that the natural projection $p: \Gamma \rightarrow R_{\mathcal{A}}$ is *foldable* if the following conditions hold:

- (1) If v is a vertex of degree 2 in Γ and e_1, e_2 are the two distinct edges in Γ with $o(e_1) = o(e_2) = v$ then $\mu(e_1) \neq \mu(e_2)$.
- (2) If $\deg(v) \geq 3$ in Γ then there exist three (oriented) edges e_1, e_2, e_3 in Γ such that $o(e_1) = o(e_2) = o(e_3) = v$ and such that $\mu(e_1), \mu(e_2), \mu(e_3)$ are three distinct elements in $\mathcal{A}^{\pm 1}$.

If the natural projection $p: \Gamma \rightarrow R_{\mathcal{A}}$ is foldable, we will also sometimes say that the \mathcal{A} -graph Γ is *foldable*.

Remark 5.7. Let Γ be a foldable \mathcal{A} -graph and let Γ' be obtained from Γ by a maximal fold.

(1) One can check that Γ' is again foldable. Note, however, that a single Stallings fold on a foldable \mathcal{A} -graph may introduce a vertex of degree three where condition (2) of Definition 5.6 fails, so that the resulting graph is not foldable. Performing maximal folds instead of single Stallings folds avoids this problem.

(2) Lemma 2.5 in [16] implies that

$$d_{\text{FS}'_N}(z(\bar{\Gamma}), z(\bar{\Gamma}')) \leq 2.$$

Recall that the marking $\bar{\Gamma}$ was defined in Remark 5.2.

(3) As noted above, in [16] Handel and Mosher introduce the notion of a “foldable” F_N -equivariant map between trees corresponding to arbitrary minimal splittings of F_N as the fundamental group of a finite graph of groups with trivial edge groups. They also prove the existence of such “foldable maps” in that setting. The general definition and construction of foldable maps are fairly complicated, but in the context of \mathcal{A} -graphs corresponding to markings on F_N they become much easier. In particular, we will only need the following basic fact that follows directly from comparing Definition 5.6 with the Handel–Mosher definition of a foldable map:

Let Γ be a finite connected \mathcal{A} -graph such that the natural projection $p: \Gamma \rightarrow R_{\mathcal{A}}$ is a homotopy equivalence and such that $\Gamma = \text{Core}(\Gamma)$. Suppose that p is foldable in the sense of Definition 5.6 above. Then there exists a foldable (in the sense of Handel–Mosher) map $\tilde{\Gamma} \rightarrow \tilde{R}_{\mathcal{A}}$.

Handel and Mosher use foldable maps as a starting point in constructing folding paths between vertices of FS'_N , and we will need the above fact in the proof of the main result in Section 6.

6. Proof of the main result

Before giving a proof of the main result, we illustrate the relationship and the maps between $\text{FS}_N, \text{FS}'_N, \text{FF}_N$ and FB_N in the following diagram, provided by the referee:




Figure 1. A diagram showing the maps between the various curve complex analogs considered in this paper. Here $F_N = A*$ represents an HNN-extension splitting of F_N with the trivial associated subgroups and with the base group A . The map ψ is the quasi-isometry between FS'_N and FS_N given by the inclusion map.

Theorem 6.1. *Let $N \geq 3$. Then the free factor complex FF_N is Gromov-hyperbolic. Moreover, there exists a constant $H > 0$ such that for any two vertices x, y of FS_N and any geodesic $[x, y]$ in $FS_N^{(1)}$ the path $\tau([x, y])$ is H -Hausdorff close to a geodesic $[\tau(x), \tau(y)]$ in $FF_N^{(1)}$.*

Proof. Recall that FS'_N is the barycentric subdivision of the free splitting graph FS_N , so that the inclusion map $FS_N \subseteq FS'_N$ is a quasi-isometry.

Recall also that for any free basis \mathcal{A} of F_N the rose $R_{\mathcal{A}}$ defines an $(N - 1)$ -simplex in FS_N (via the canonical marking $F_N \rightarrow \pi_1(R_{\mathcal{A}})$ sending the elements of \mathcal{A} to the corresponding loop-edges of $R_{\mathcal{A}}$) and that, as in Convention 3.4, $z(R_{\mathcal{A}}) \in V(FS'_N)$ is the barycenter of that simplex. Note that by definition, if $[\mathcal{A}] = [\mathcal{B}]$ then $z(R_{\mathcal{A}}) = z(R_{\mathcal{B}})$. Put

$$S = \{z(R_{\mathcal{A}}) \mid \mathcal{A} \text{ is a free basis of } F_N\}.$$

Thus $S \subseteq V(FS'_N)$ and we may think of S as a copy of $V(FB_N)$ in $V(FS'_N)$.

For every $x, y \in S$ let $g_{x,y}$ be the path from x to y in FS'_N given by the Handel–Mosher folding line [16]. Recall that, as proved in [16], FS'_N is Gromov-hyperbolic and $g_{x,y}$ is a re-parameterized uniform quasigeodesic. Hence $g_{x,y}$ is uniformly Hausdorff close to any geodesic $[x, y]$ in FS'_N .

Consider the following map $f : V(FS'_N) \rightarrow V(FB_N)$. For every vertex u of FS'_N , which may be viewed as a splitting of F_N as the fundamental group of a graph of groups with trivial edge-groups, choose an edge e of that splitting, collapse the rest of u to a single-edge splitting corresponding to u and let $A(u)$ be a vertex group of that collapsed splitting. Thus $A(u)$ is a proper free factor of F_N and hence $[A(u)]$ is a vertex of FF_N . Then choose a vertex v of FB_N with $d(h(v), [A(u)]) \leq 3$ (such v exists by Proposition 4.3). Put $f(u) := v$. We can make the above choices to make sure that for every free basis \mathcal{A} of F_N we have $f(z(R_{\mathcal{A}})) = [\mathcal{A}]$ (note that

$d_{\text{FF}_N}(h([\mathcal{A}], A(z(R_{\mathcal{A}}))) \leq 2)$. With the above mentioned identification of S and $V(\text{FB}_N)$ we may in fact informally think that $f|_S = \text{Id}_S$. Moreover, if Γ is a foldable \mathcal{A} -graph (which therefore defines a marking $\bar{\Gamma}$) and $T \subseteq \Gamma$ is a maximal tree, then we have an associated free basis $\mathcal{B}(\Gamma, T)$ described in Remark 5.5 above. One can check that $d(f(\bar{\Gamma}), [\mathcal{B}(\Gamma, T)]) \leq B$ for some constant $B = B(N) > 0$ independent of \mathcal{A}, Γ, T .

We have defined a map $f : V(\text{FS}'_N) \rightarrow V(\text{FB}_N)$. We then extend this map to a graph-map $f : \text{FS}'_N \rightarrow \text{FB}_N$ by sending an arbitrary edge e of FS'_N with endpoints u_1, u_2 to a geodesic edge-path $[f(u_1), f(u_2)]$ in FB_N . The graph-map $f : \text{FS}'_N \rightarrow \text{FB}_N$ is L -Lipschitz for some $L \geq 0$, since τ is Lipschitz and the maps $V(\text{FS}'_N) \rightarrow \text{FF}_N^{(1)}$, given by $u \mapsto [A(u)]$ and $u \mapsto \tau(u)$, are bounded distance away from each other.

We claim that all the assumptions of Proposition 2.5 are satisfied for the map $f : \text{FS}'_N \rightarrow \text{FB}_N$ and the set S .

Condition (1) of Proposition 2.5 holds, since by assumption $f(S) = V(\text{FB}_N)$. Also, as noted above, $f : \text{FS}'_N \rightarrow \text{FB}_N$ is L -Lipschitz, and it is easy to see that S is D -dense in FS'_N , for some $D > 0$. Our task is to verify condition (3) of Proposition 2.5.

If $x', y' \in V(\text{FB}_N)$ have $d(x', y') \leq 1$ then there exist free bases \mathcal{A}, \mathcal{B} of F_N such that $x' = [\mathcal{B}]$, $y' = [\mathcal{A}]$ and such that there exists $a \in \mathcal{A} \cap \mathcal{B}$. Without loss of generality, we may assume that $\mathcal{A} = \{a_1, \dots, a_N\}$, $\mathcal{B} = \{b_1, \dots, b_N\}$ and that $a_1 = b_1 = a$. Put $x = z(R_{\mathcal{B}})$ and $y = z(\mathcal{A})$, so that $f(x) = x'$ and $f(y) = y'$.

In [16] Handel–Mosher [16], given any ordered pair of vertices x, y of $V(\text{FS}'_N)$, construct an edge-path $g_{x,y}$ from x to y in FS'_N , which we will sometimes call the *Handel–Mosher folding path*. The general definition of $g_{x,y}$ in [16] is fairly complicated. However, we only need to use this definition for the case where $x, y \in S$, in which case it becomes much simpler, and which we will now describe in greater detail for the vertices $x = z(R_{\mathcal{B}})$ and $y = z(\mathcal{A})$ defined above.

Consider an \mathcal{A} -graph Γ_0 which is a wedge of N simple loops at a common base-vertex v_0 , where the i -th loop is labeled by the freely reduced word over \mathcal{A} that is equal to b_i in F_N . Note that the first loop is just a loop-edge labelled by a_1 , since by assumption $b_1 = a_1$. The natural projection $p : \Gamma_0 \rightarrow R_{\mathcal{A}}$ is a homotopy equivalence and we also have $\Gamma_0 = \text{Core}(\Gamma_0)$. Condition (1) of Definition 5.6 holds for Γ_0 by construction. However, $p : \Gamma_0 \rightarrow R_{\mathcal{A}}$ is not necessarily foldable since Condition (2) of Definition 5.6 may fail. This happens exactly when there exists $\varepsilon \in \{1, -1\}$ such that for all $i = 2, \dots, N$ the freely reduced word over \mathcal{A} representing b_i begins with a_1^ε and ends with $a_1^{-\varepsilon}$. However, after possibly replacing \mathcal{B} by an equivalent free basis of the form $a_1^m \mathcal{B} a_1^{-m}$, for the graph Γ_0 defined as above the natural projection $p : \Gamma_0 \rightarrow R_{\mathcal{A}}$ is foldable in the sense of Definition 5.6. Note that conjugation by a_1^m fixes the element $b_1 = a_1$, so that even after the above modification of \mathcal{B} it will still be true that Γ_0 contains a loop-edge at v_0 with label a_1 .

As noted in Remark 5.7 above, as the initial input for constructing $g_{x,y}$, Handel

and Mosher need a “foldable” (in the sense of [16]) F_N -equivariant map $\widetilde{R}_{\mathcal{B}} \rightarrow \widetilde{R}_{\mathcal{A}}$. Again, as observed in Remark 5.7, such a map exists since we have arranged for the \mathcal{A} -graph Γ_0 to be foldable in the sense of Definition 5.6.

Note that by construction, the marking $\overline{\Gamma}_0$ corresponding to Γ_0 is exactly the vertex $x = z(R_{\mathcal{B}}')$ of FS'_N .

The remainder of the Handel–Mosher construction of $g_{x,y}$ in this case works as follows. Since $p: \Gamma_0 \rightarrow R_{\mathcal{A}}$ is a homotopy equivalence, there exists a finite sequence of \mathcal{A} -graphs

$$\Gamma_0, \Gamma_1, \dots, \Gamma_n = R_{\mathcal{A}}$$

where for $i = 1, \dots, n$ Γ_i is obtained from Γ_{i-1} by a maximal fold. Then the associated markings $\overline{\Gamma}_i$ are vertices of FS'_N . As observed in Remark 5.7, we have $d(\overline{\Gamma}_{i-1}, \overline{\Gamma}_i) \leq 2$ in FS'_N . Joining each consecutive pair $\overline{\Gamma}_{i-1}, \overline{\Gamma}_i$ by a geodesic path of length ≤ 2 in FS'_N produces the path $g_{x,y}$ from x to y in FS'_N . Note that each Γ_i has a base-vertex v_i which is the image of the base-vertex v_0 of Γ_0 under the sequence of folds that takes Γ_0 to Γ_i .

A crucial feature of the above construction is that every Γ_i will have a loop-edge (at the base-vertex v_i of Γ_i) with label a_1 . Since the map $f: FS'_N \rightarrow FB_N$ is L -Lipschitz, this implies that $f(g_{x,y})$ has diameter bounded by some constant M_0 independent of x, y . Indeed, Since Γ_i has a loop-edge at its base-vertex with label a_1 , there exists a free basis $\gamma_1, \dots, \gamma_N$ of $\pi_1(\Gamma_i, v_i)$ (e.g. coming from a choice of a maximal tree in Γ_i , as in Definition 4.4 and Remark 5.5) such that $\mu(\gamma_1) = a_1$ and such that $\mathcal{B}_i = \{\mu(\gamma_1), \dots, \mu(\gamma_N)\}$ is a free basis of F_N . Since $a_1 \in \mathcal{B}_i$, we have $d([\mathcal{B}_i], [\mathcal{A}]) \leq 1$ in FB_N for each i . Unpacking the definition of the map f we see that $d(f(\overline{\Gamma}_{i-1}), [\mathcal{B}_i]) \leq C$ in FB_N for some constant $C \geq 0$. Hence $d(f(\overline{\Gamma}_{i-1}), [\mathcal{A}]) \leq C + 1$. Recall that $g_{x,y}$ is a quasi-geodesic in a hyperbolic graph FS'_N and hence $g_{x,y}$ is uniformly Hausdorff-close to a geodesic $[x, y]$. Since f is L -Lipschitz, it follows that $f([x, y])$ has diameter bounded by some constant M independent of x, y . Thus condition (3) of Proposition 2.5 holds.

Therefore, by Proposition 2.5, the graph FB_N is Gromov-hyperbolic, and, moreover, for any vertices x, y of FS_N , the path $f([x, y])$ is uniformly Hausdorff-close to a geodesic $[f(x), f(y)]$.

Recall that in Proposition 4.3 we constructed an explicit quasi-isometry $h: FB_N \rightarrow FF_N$. Since FB_N is hyperbolic, it follows that FF_N is Gromov-hyperbolic as well. Moreover, the map $\tau: FS_N \rightarrow FF_N$ from the statement of Theorem 1.1, and the map $h \circ f: FS_N \rightarrow FF_N$ are bounded distance from each other. This implies that there exists a constant $H > 0$ such that for any two vertices x, y of FS_N and any geodesic $[x, y]$ in $FS_N^{(1)}$ the path $\tau([x, y])$ is H -Hausdorff close to a geodesic $[\tau(x), \tau(y)]$ in $FF_N^{(1)}$. □

Remark 6.2. The above proof implies a reasonably explicit description of certain reparameterized quasigeodesics in FB_N between two arbitrary vertices of FB_N in terms of Stallings folds. Let $\mathcal{A} = \{a_1, \dots, a_N\}$ and $\mathcal{B} = \{b_1, \dots, b_N\}$ be free bases

of F_N . Let Γ_0 be an \mathcal{A} -graph corresponding to \mathcal{B} constructed in a similar way to the way Γ_0 was constructed in the above proof. That is, let Γ_0 be a wedge of N simple loops at a common base-vertex v_0 , where the i -th loop is labeled by the freely reduced word over \mathcal{A} that is equal to b_i in F_N . Suppose that Γ_0 is such that the natural projection $p: \Gamma_0 \rightarrow R_{\mathcal{A}}$ is foldable in the sense of Definition 5.6. (Note that this assumption does not always hold; however, it may always be ensured after replacing \mathcal{B} by an equivalent free basis).

Let $\Gamma_0, \Gamma_1, \dots, \Gamma_n = R_{\mathcal{A}}$ be \mathcal{A} -graphs such that for $i = 1, \dots, n$ Γ_i is obtained from Γ_{i-1} by a maximal fold. Note that each Γ_i has a distinguished base-vertex v_i , which is the image of the base-vertex v_0 of Γ_0 under the foldings transforming Γ_0 to Γ_i .

For each $1 \leq i < n$ choose a maximal subtree T_i in Γ_i . Let $\mathcal{A}_i = \mathcal{B}(\Gamma_i, T_i)$ be the associated free basis of F_N (see Remark 5.5 above for its detailed description). Put $\mathcal{A}_0 = \mathcal{B}$ and $\mathcal{A}_n = \mathcal{A}$.

It is not hard to check that $d([\mathcal{A}_i], f(\overline{\Gamma}_i)) \leq C$ in FB_N for some constant $C = C(N) > 0$ independent of \mathcal{A}, \mathcal{B} . Since, as noted in the proof of Theorem 1.1 above, the sequence $\overline{\Gamma}_0, \dots, \overline{\Gamma}_n$ defines a (reparameterized) uniform quasigeodesic in FS'_N , it now follows from the proof of Theorem 1.1 that the set $\{[\mathcal{A}_0], [\mathcal{A}_1], \dots, [\mathcal{A}_n]\}$ is uniformly Hausdorff-close to a geodesic joining $[\mathcal{B}]$ and $[\mathcal{A}]$ in FB_N . This fact can also be derived from a careful analysis of the Bestvina–Feighn proof [4] of hyperbolicity of FF_N .

References

- [1] J. Aramayona and J. Souto, Automorphisms of the graph of free splittings. *Michigan Math. J.* **60** (2011), no. 3, 483–493. [Zbl 1242.05117](#) [MR 2861084](#)
- [2] J. Behrstock, M. Bestvina, and M. Clay, Growth of intersection numbers for free group automorphisms. *J. Topol.* **3** (2010), no. 2, 280–310. [Zbl 1209.20031](#) [MR 2651361](#)
- [3] M. Bestvina, and M. Feighn, A hyperbolic $\text{Out}(F_n)$ -complex. *Groups Geom. Dyn.* **4** (2010), no. 1, 31–58. [Zbl 1190.20017](#) [MR 2566300](#)
- [4] M. Bestvina, and M. Feighn, Hyperbolicity of the complex of free factors. *Adv. Math.* **256** (2014), 104–155; corrigendum, *ibid.* **259** (2014), 843. [Zbl 06284922](#) [MR 3177291](#)
- [5] M. Bestvina, and M. Feighn, Subfactor projections. Preprint, [arXiv:1211.1730](#).
- [6] M. Bestvina and P. Reynolds, The boundary of the complex of free factors. Preprint, [arXiv:1211.3608](#).
- [7] B. Bowditch, Relatively hyperbolic groups. *Internat. J. Algebra Comput.* **22** (2012), no. 3, 1250016, 66 pp. [Zbl 1259.20052](#) [MR 2922380](#)
- [8] B. Bowditch, Intersection numbers and the hyperbolicity of the curve complex. *J. Reine Angew. Math.* **598** (2006), 105–129. [Zbl 1119.32006](#) [MR 2270568](#)
- [9] B. Bowditch, Uniform hyperbolicity of the curve graphs. *Pacific J. Math.*, to appear.

- [10] J. Brock, R. Canary, and Y. Minsky, The classification of Kleinian surface groups, II: The ending lamination conjecture. *Ann. of Math. (2)* **176** (2012), no. 1, 1–149. [Zbl 1253.57009](#) [MR 2925381](#)
- [11] M. Culler and K. Vogtmann, Moduli of graphs and automorphisms of free groups. *Invent. Math.* **84** (1986), no. 1, 91–119 [Zbl 0589.20022](#) [MR 0830040](#)
- [12] F. Dahmani and V. Guirardel, Rotating families, and the structure of some normal subgroups in groups acting on hyperbolic spaces. Preprint, <http://perso.univ-rennes1.fr/vincent.guirardel/papiers/rotating.pdf>.
- [13] B. Farb, Relatively hyperbolic groups. *Geom. Funct. Anal.* **8** (1998), no. 5, 810–840. [Zbl 0985.20027](#) [MR 1650094](#)
- [14] U. Hamenstädt, Geometry of the complex of curves and of Teichmüller space. In *Handbook of Teichmüller Theory*, Vol. 1, A. Papadopoulos, ed., European Math. Soc., Zürich 2007, 447–467. [Zbl 1162.32010](#) [MR 2349677](#)
- [15] U. Hamenstädt, The boundary of the free splitting graph and the free factor graph. Preprint, [arXiv:1211.1630](#).
- [16] M. Handel and L. Mosher, The free splitting complex of a free group I: Hyperbolicity. *Geom. Topol.* **17** (2013), no. 3, 1581–1672. [Zbl 1278.20053](#) [MR 3073931](#)
- [17] W. J. Harvey, Boundary structure of the modular group. In *Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference* (State Univ. New York, Stony Brook, N. Y., 1978), Ann. of Math. Stud. 97, Princeton University Press, Princeton, N.J., 1981, 245–251. [Zbl 0461.30036](#) [MR 0624817](#)
- [18] A. Hatcher and K. Vogtmann, The complex of free factors of a free group. *Quart. J. Math. Oxford Ser. (2)* **49** (1998), no. 196, 459–468. [Zbl 0935.20015](#) [MR 1660045](#)
- [19] A. Hilion and C. Horbez, The hyperbolicity of the sphere complex via surgery paths. Preprint, [arXiv:1210.6183](#).
- [20] I. Kapovich and M. Lustig, Geometric intersection number and analogues of the curve complex for free groups. *Geom. Topol.* **13** (2009), no. 3, 1805–1833. [Zbl 1194.20046](#) [MR 2496058](#)
- [21] I. Kapovich and A. Myasnikov, Stallings foldings and subgroups of free groups. *J. Algebra* **248** (2002), no. 2, 608–668. [Zbl 1001.20015](#) [MR 1882114](#)
- [22] J. Maher and S. Schleimer, The graph of handlebodies. In preparation.
- [23] H. Masur, and Y. Minsky, Geometry of the complex of curves. I. Hyperbolicity. *Invent. Math.* **138** (1999), no. 1, 103–149. [Zbl 0941.32012](#) [MR 1714338](#)
- [24] H. Masur, and Y. Minsky, Geometry of the complex of curves. II. Hierarchical structure. *Geom. Funct. Anal.* **10** (2000), no. 4, 902–974. [Zbl 0972.32011](#) [MR 1791145](#)
- [25] H. Masur and S. Schleimer, The geometry of the disk complex. *J. Amer. Math. Soc.* **26** (2013), no. 1, 1–62. [Zbl 1272.57015](#) [MR 2983005](#)
- [26] Y. Minsky, End invariants and the classification of hyperbolic 3-manifolds. In *Current developments in mathematics, 2002*, International Press, Somerville, Mass., 2003, 181–217. [Zbl 1049.57010](#) [MR 2062319](#)
- [27] Y. Minsky, The classification of Kleinian surface groups. I. Models and bounds. *Ann. of Math. (2)* **171** (2010), no. 1, 1–107. [Zbl 1193.30063](#) [MR 2630036](#)

- [28] M. Mj and L. Reeves, A combination theorem for strong relative hyperbolicity. *Geom. Topol.* **12** (2008), no. 3, 1777–1798. [Zbl 1192.20027](#) [MR 2421140](#)
- [29] L. Sabalka, and D. Savchuk, On the geometry of a proposed curve complex analogue for $\text{Out}(F_n)$. Preprint, [arXiv:1007.1998](#).
- [30] L. Sabalka, and D. Savchuk, Submanifold projection. Preprint, [arXiv:1211.3111](#).
- [31] J. R. Stallings, Topology of finite graphs. *Invent. Math.* **71** (1983), no. 3, 551–565 [Zbl 0521.20013](#) [MR 0695906](#)

Received September 11, 2012; revised November 30, 2012

I. Kapovich, Department of Mathematics, University of Illinois at Urbana-Champaign,
1409 West Green Street, Urbana, IL 61801, U.S.A.

E-mail: kapovich@math.uiuc.edu

K. Rafi, Department of Mathematics, University of Oklahoma, Norman, OK 73019, U.S.A.

E-mail: rafi@math.ou.edu