

Coxeter group in Hilbert geometry

Ludovic Marquis

Abstract. A theorem of Tits and Vinberg allows to build an action of a Coxeter group Γ on a properly convex open set Ω of the real projective space, thanks to the data P of a polytope and reflection across its facets. We give sufficient conditions for such action to be of finite covolume, convex-cocompact or geometrically finite. We describe a hypothesis that makes those conditions necessary.

Under this hypothesis, we describe the Zariski closure of Γ , find the maximal Γ -invariant convex set, when there is a unique Γ -invariant convex set, when the convex set Ω is strictly convex, when we can find a Γ -invariant convex set Ω' which is strictly convex.

Mathematics Subject Classification (2010). 22E40, 20F55, 20F67, 53C60.

Keywords. Coxeter group, Hilbert geometry, Discrete subgroup of Lie group, convex projective structure on manifold and orbifold, geometric group theory.

Contents

1 Preliminaries	825
2 The Theorem of Tits and Vinberg and the Theorems of Vinberg	832
3 The setting	840
4 The lemmas	842
5 Degenerate 2-perfect Coxeter polytopes	853
6 Geometry of the action	854
7 Zariski closure of Γ_P	856
8 About the convex set	865
References	873

Introduction

General Framework. The study of groups acting on Hilbert geometry or convex projective structures on manifold starts with the pioneering work of Kuiper [Kui53] in the 50's. After came the works of Benzécri [Bnz60], Vinberg [Vin63, Vin65, Vin71], Kac and Vinberg [KV67], Koszul [Kos68], and Vey [VEY70] in

the 60's. Then the field took a deep breath, and came back in the 90's with Goldman [GOL90], followed by Suhyoung Choi, Labourie, Loftin, In Kang Kim and a long series of articles by Benoist in the 2000's. The recent works of Jaejeong Lee, Misha Kapovich, Cooper, Long, Tillmann, Thistlethwaite, Ballas, Gye-Seon Lee, Suhyoung Choi, Nie, Crampon and the author show a growing interest for this field.

We want to study the action of discrete groups Γ of $\mathrm{SL}_{d+1}^{\pm}(\mathbb{R})$ on a properly¹ convex open set Ω of the projective sphere $\mathbb{S}^d = \mathbb{S}(\mathbb{R}^{d+1}) = \{\text{Half-line of } \mathbb{R}^{d+1}\}$. Note that on every properly convex open set Ω of \mathbb{S}^d there is a distance d_{Ω} and a measure μ_{Ω} invariant by the group $\mathrm{Aut}(\Omega) = \{\gamma \in \mathrm{SL}_{d+1}^{\pm}(\mathbb{R}) \mid \gamma(\Omega) = \Omega\}$ of automorphisms of Ω .

At this moment, *divisible convex sets*, the convex sets Ω for which there exists a discrete subgroup Γ of $\mathrm{Aut}(\Omega)$ such that Ω/Γ is compact, have received almost all the attention. The *quasi-divisible convex sets*, the one for which there exist a discrete subgroup Γ of $\mathrm{Aut}(\Omega)$ such that Ω/Γ is of finite volume, are starting to be studied, see [CLT15, BAL14, BAL15, CMI14, MAR12A, MAR12B].

There is at least four ways to say that the action of Γ on Ω is “cofinite.” The first two ways are the following: the action of Γ on Ω is *cocompact* (resp. of *cofinite volume*) when the quotient orbifold Ω/Γ is compact (resp. of finite volume for the measure induced by μ_{Ω}).

If we assume moreover that the action of Γ on \mathbb{R}^{d+1} is strongly irreducible,² Benoist shows in [BEN00] that there exists a smallest closed Γ -invariant subset Λ_{Γ} of the real projective space $\mathbb{P}(\mathbb{R}^{d+1}) = \mathbb{P}^d(\mathbb{R}) = \mathbb{P}^d$. We still denote Λ_{Γ} the one of the two preimages of Λ_{Γ} in \mathbb{S}^d which is included in $\partial\Omega$. We denote by $\bar{C}(\Lambda_{\Gamma})$ the convex hull³ of Λ_{Γ} in Ω . We remark that $\bar{C}(\Lambda_{\Gamma})$ is a closed subset of Ω which has a non empty interior since the action of Γ on \mathbb{R}^{d+1} is strongly irreducible.

We will say that the action of Γ on Ω is *convex-cocompact* (resp. *geometrically finite*) when the quotient $\bar{C}(\Lambda_{\Gamma})/\Gamma$ is compact (resp. of finite volume for the measure induced by μ_{Ω}).

The definition of cocompact, finite volume or convex-cocompact action make no doubt, but the definition of geometrical finiteness deserves a detailed comment that will be done in Section 6.5.

The theory of Coxeter groups has two benefits for us. First, it gives a simple and explicit recipe to build a lot of groups with different behaviours from the point of view of geometric group theory. Second, the Theorem of Tits and Vinberg gives the hope to build a lot of interesting actions of Coxeter groups on Hilbert geometry. So, we will focus on actions of Coxeter groups W on convex subsets of the projective sphere \mathbb{S}^d .

¹ A bounded convex subset of an affine chart.

² The action of any finite index subgroup of Γ on \mathbb{R}^{d+1} is irreducible.

³ The smallest closed convex subset of Ω containing Λ_{Γ} in its closure in \mathbb{S}^d .

We point out, for the reader not familiar with Hilbert geometry, that Hilbert geometries can also be very different. For example, if Ω is the round ball of an affine chart \mathbb{R}^d of \mathbb{S}^d then (Ω, d_Ω) is isometric to the real hyperbolic space of dimension d and if Ω is a triangle then (Ω, d_Ω) is bi-Lipschitz equivalent to the euclidean plane. In particular, our discussion includes the context of hyperbolic geometry.

Precise Framework. In order to make a Coxeter group act on the projective sphere, one can take a projective polytope P of \mathbb{S}^d , and choose a projective reflection σ_s across each facet s of P .⁴ We want to consider the subgroup $\Gamma = \Gamma_P$ of $\mathrm{SL}_{d+1}^\pm(\mathbb{R})$ generated by the reflections $(\sigma_s)_{s \in S}$, where S is the set of facets of P . In order to get a discrete subgroup of $\mathrm{SL}_{d+1}^\pm(\mathbb{R})$, we need some hypothesis on the set of reflections $(\sigma_s)_{s \in S}$. Roughly speaking, the hypothesis will be that if s and t are two facets of P such that $s \cap t$ is of codimension 2 then the product $\sigma_s \sigma_t$ is conjugate to a rotation of angle $\frac{\pi}{m}$, where m is an integer. We also authorize the case $m = \infty$, and for this case a special condition is needed.

The precise definition is Definition 1.8. Such a polytope will be called a *Coxeter polytope*. Given a Coxeter polytope P , we can consider the set $\mathcal{C} = \mathcal{C}_P = \bigcup_{\gamma \in \Gamma} \gamma(P)$. The Theorem of Tits and Vinberg (Theorem 2.2) tells us that Γ is discrete and \mathcal{C} is a convex subset of \mathbb{S}^d . This theorem provides a huge amount of examples with drastically different behaviours.

The goal of this article is to tackle the following questions. Let P be a Coxeter polytope of \mathbb{S}^d , let Γ be the discrete subgroup generated by the reflections $(\sigma_s)_{s \in S}$ and $\mathcal{C} = \bigcup_{\gamma \in \Gamma} \gamma(P)$. In order to get nice irreducible examples, we assume that the action of Γ on \mathbb{R}^{d+1} is strongly irreducible, so \mathcal{C} has to be properly convex. Let Ω be the interior of \mathcal{C} . When is the action of Γ on Ω cocompact?⁵ of finite covolume? convex cocompact? geometrically finite?

Our goal is also to answer questions about the Zariski closure of Γ , about the convex set Ω and about the other possible convex set preserved by Γ .

- (1) What are the possible Zariski closures for Γ ?
- (2) Is the convex set Ω the largest properly convex open set preserved?
- (3) When does the action of Γ on \mathbb{S}^d preserve a unique properly convex open subset?
- (4) When is the convex set Ω the smallest properly convex open set preserved?
- (5) When is the convex set Ω strictly convex? with \mathcal{C}^1 boundary? both?
- (6) When does the action of Γ on \mathbb{S}^d preserve a strictly convex open set? a properly convex open set with \mathcal{C}^1 boundary? a strictly convex open set with \mathcal{C}^1 boundary?

⁴ Note that in projective geometry there are many reflections across a given hyperplane.

⁵ Already answered by Vinberg, see Theorem 2 of [Vin71] or corollary 2.3.

If we don't make any hypothesis on the Coxeter polytope P , the behaviour of the action $\Gamma \curvearrowright \Omega$ can be very complicated. So, we will make a non-trivial hypothesis along this text. The author thinks that this hypothesis is relevant and offers an access to a wide family of examples. For example, this hypothesis is satisfied by every Coxeter polygon and every Coxeter polyhedron whose dihedral angles are non-zero.

Now, we briefly explain the hypothesis that we will make most of the time along this text. A nice way to get information about a polytope is to look around a vertex. The link of a Coxeter polytope P at a vertex p is a Coxeter polytope P_p of one dimension less than P and which is “ P seen from p .” In the context of hyperbolic geometry, it is the intersection of P with a small sphere centered at p .

Vinberg introduces the following terminology in [VIN71]. A Coxeter polytope P is *perfect* when the action of Γ on Ω is cocompact. We will mainly assume that P is *2-perfect*, which means that the link of every vertex of P is perfect or equivalently that $P \cap \partial\Omega$ is contained in the set of vertices of P . See Proposition 3.1 for a precise statement.

Vinberg shows in [VIN71] that perfect Coxeter polytopes come from three different families:⁶

- P is *elliptic*, i.e. Γ is finite;
- P is *parabolic*, i.e. Ω is an affine chart;
- Otherwise, Ω is properly convex. In that case, we say that P is *loxodromic*.

In particular, if P is 2-perfect, then the link at any vertex is either elliptic, parabolic or loxodromic. We can now state our results.

Theorem A (Theorems 6.2, 6.3 and 6.4). *Let P be a 2-perfect Coxeter polytope. Let $\Gamma = \Gamma_P$ be the subgroup of $\mathrm{SL}_{d+1}(\mathbb{R})$ generated by the reflections around the facets of P . Let $\Omega = \Omega_P$ be the interior of the Γ -orbit of P . Suppose that the action of Γ on \mathbb{R}^{d+1} is strongly irreducible.*

- *The action $\Gamma \curvearrowright \Omega$ is geometrically finite.*
- *Moreover, the action $\Gamma \curvearrowright \Omega$ is of finite covolume if and only if the link P_p of every vertex p of P is elliptic or parabolic.*
- *Finally, the action $\Gamma \curvearrowright \Omega$ is convex cocompact if and only if the link P_p of every vertex p of P is elliptic or loxodromic.*

We keep the same notation and hypothesis for the following theorems.

Theorem B (Theorem 7.11). *The Zariski closure of Γ is either conjugate to $\mathrm{SO}_{d,1}^\circ(\mathbb{R})$ or is equal to $\mathrm{SL}_{d+1}(\mathbb{R})$.*

⁶ See definition 2.13.

Theorem C (Theorem 8.1). *Every properly convex open set preserved by Γ is included in Ω .*

Theorem D (Theorem 8.2). *The convex set Ω is the smallest properly convex open set preserved by Γ if and only if the action $\Gamma \curvearrowright \Omega$ is of finite covolume.*

By using one of the results of [BEN04, CLT15], we can also show:

Theorem E (Theorem 8.7). *The following are equivalent:*

- (1) *the properly convex open set Ω is strictly convex;*
- (2) *the boundary $\partial\Omega$ of Ω is of class \mathcal{C}^1 ;*
- (3) *the action $\Gamma \curvearrowright \Omega$ is of finite covolume and the group Γ is relatively hyperbolic relatively to the links P_p for which P_p is parabolic.*

In that case, the metric space (Ω, d_Ω) is Gromov-hyperbolic.

Thanks to the moduli space computed in [MAR10], we will easily get the following theorem as a corollary of Theorem E.

Theorem F. *In dimension 3, there exists an indecomposable⁷ quasi-divisible properly convex open set which is not divisible nor strictly convex.*

We recall that one cannot find such an example in dimension 2, thanks to [BNZ60, MAR12A]. A construction in any dimension is an open question in the divisible or the quasi-divisible context.

Theorem G (Theorem 8.11). *If moreover all the loxodromic vertices are simple,⁸ the following are equivalent:*

- (1) *there exists a strictly convex open set Ω' preserved by Γ ;*
- (2) *there exists a properly convex open set Ω' with \mathcal{C}^1 -boundary preserved by Γ ;*
- (3) *the group Γ is relatively hyperbolic relatively to the links P_p for which P_p is parabolic.*

Along the way, we will study a nice procedure: truncation which allows to build a new polytope from a starting one by cutting a simple vertex (See Subsection 4.5). This procedure is present in a survey of Vinberg [VIN85] in the context of hyperbolic geometry, it has also been used by the author in [MAR10], this time in the context of projective geometry. The approach in this text will be less computational and more geometrical than in [MAR10]. We think this procedure is interesting in its own right. Moreover, the introduction of this procedure gives nicer statements of the previously quoted theorems.

⁷ A convex set that is not the join of two convex sets of smaller dimension.

⁸ A vertex is *simple* when its link is a simplex.

Others works around the subject. The starting point and main inspiration for this article, is the article [VIN71] of Vinberg, which presents the notion of Coxeter polytope⁹ and studies the first properties. Cocompact actions are studied in Vinberg’s text but actions of cofinite volume, convex-cocompact or geometrically finite action are not. There are also lecture notes by Benoist [BEN09] which present a proof of the theorem of Tits and Vinberg. The examples of the article [BEN06A, BEN06B] of Benoist are built thanks to the theorem of Tits and Vinberg.

One can also study the moduli space of a Coxeter polytope but we will not do it in this text. Suhyoung Choi with Gye-Seon Lee, Craig Hodgson and the author have works on this problem [CHO06, CHL12, CLI5, MAR10]. We will devote several forthcoming articles with Suhyoung Choi, Gye-Seon Lee and/or Ryan Greene to the problem of moduli space.

The study of geometrically finite actions was started in [CMI4] of M. Crampon and the author. We stress that in the last article the authors made the hypothesis that the convex set Ω on which the group Γ acts is strictly convex with \mathcal{C}^1 -boundary. The study of actions of cofinite volume is the main purpose of the articles [CLT15] of Cooper, Long and Tillmann, [MAR12A] and [MAR12B] of the author. We stress that the hypothesis of strict convexity of Ω is central in [CMI4, MAR12B]. This hypothesis is absent from [MAR12A] and is not always present in [CLT15]. There is also a paper of Suhyoung Choi about geometrically finite actions [CHO10].

We point out that in this text, we did not make any assumption about the regularity of the boundary of Ω . One of the goals is actually to build examples where Ω is not strictly convex and the action is cocompact or of finite covolume.

Plan of the article. The first part of the article contains preliminaries about convexity, Hilbert geometry, Coxeter groups and Coxeter polytopes. The second part is a recalling of the theorem of Tits and Vinberg and of important results of Vinberg coming from the article [VIN71]. The third part presents the definition of link of a polytope and makes precise the hypothesis “ P is 2-perfect.”

The fourth part presents the lemmas for the study of the geometry around a vertex. The fifth part is a classification of degenerate 2-perfect polytopes. The sixth part is devoted to the proof of Theorem A. The seventh part studies the Zariski closure of Γ , and it contains the proof of Theorem B. The eighth part contains the proof of Theorems C, D, E, and G.

Acknowledgements. The author thanks Yves Benoist for a couple of dense discussion about this text. We thank Érnest Vinberg which is a major source of inspiration for this article. We thank the referee for the comments and corrections he suggests. Finally, we warmly thank Gye-Seon Lee who found a lot of errors in a previous version.

The author thanks the ANR facets of discrete groups and ANR Finsler geometry for their supports.

⁹ Note that Vinberg prefers to work with Γ than with P . Vinberg called such a Γ a *linear Coxeter group*.

1. Preliminaries

1.1. Convexity in the projective sphere. Let V be a real vector space. A convex cone \mathcal{C} is *sharp* when \mathcal{C} does not contain any affine line. Consider the *projective sphere* $\mathbb{S}(V) = \{\text{Half-lines of } V\} = V \setminus \{0\} / \sim$ where \sim is the equivalence relation induced by the action of \mathbb{R}_+^* by homothety on V . Of course, $\mathbb{S}(V)$ is the 2-fold cover of the real projective space $\mathbb{P}(V)$. The notion of convexity is nicer in $\mathbb{S}(V)$ than in $\mathbb{P}(V)$. We will denote $\mathbb{S}: V \setminus \{0\} \rightarrow \mathbb{S}(V)$ the natural projection.

A subset C of $\mathbb{S}(V)$ is *convex* (resp. *properly convex*) when the set $\mathbb{S}^{-1}(C)$ is a convex cone (resp. sharp convex cone) of V . Given a hyperplane H of $\mathbb{S}(V)$, the two connected components of $\mathbb{S}(V) \setminus H$ are called *affine charts*. An open set $\Omega \neq \mathbb{S}(V)$ of $\mathbb{S}(V)$ is convex (resp. properly convex) if and only if there exists an affine chart \mathbb{A} such that $\Omega \subset \mathbb{A}$ (resp. $\bar{\Omega} \subset \mathbb{A}$) and Ω is convex in the usual sense in \mathbb{A} .

1.2. Hilbert geometry. On every properly convex open set Ω of \mathbb{S}^d there is a distance d_Ω defined thanks to the cross-ratio, in the following way. Take any two points $x \neq y \in \Omega$ and draw the line between them. This line intersects the boundary $\partial\Omega$ of Ω in two points p and q . We assume that x is between p and y . Then the following formula defines a distance (see Figure 1):

$$d_\Omega(x, y) = \frac{1}{2} \ln([p : x : y : q]).$$

This distance gives to Ω the same topology than the one inherited from $\mathbb{S}(V)$. The metric space (Ω, d_Ω) is complete, the closed ball are compact, the group $\text{Aut}(\Omega)$ acts by isometries on Ω , and therefore acts properly.

This distance is called the *Hilbert distance* and has the good taste to came from a Finsler metric on Ω defined by a very simple formula. Let x be a point of Ω and v a vector of the tangent space $T_x\Omega$ of Ω at x . The quantity $\left. \frac{d}{dt} \right|_{t=0} d_\Omega(x, x + tv)$ defines a Finsler metric $F_\Omega(x, v)$ on Ω . Moreover, if we choose an affine chart \mathbb{A} containing Ω and a euclidean norm $|\cdot|$ on \mathbb{A} , we get

$$F_\Omega(x, v) = \left. \frac{d}{dt} \right|_{t=0} d_\Omega(x, x + tv) = \frac{|v|}{2} \left(\frac{1}{|xp^-|} + \frac{1}{|xp^+|} \right),$$

where p^- and p^+ are the intersection points of the half-line starting at p with direction $-v$ and v with $\partial\Omega$ and $|ab|$ is the distance between points a, b of \mathbb{A} for the euclidean norm $|\cdot|$ (see Figure 1). The regularity of this Finsler metric is the regularity of the boundary $\partial\Omega$ of Ω , and the Finsler structure gives rise to an absolutely continuous measure μ_Ω with respect to Lebesgue measure. We will not need any explicit formula for this measure; we will only use the following proposition which is straightforward and explained in [VER05].




Figure 1. Hilbert distance.

Proposition 1.1. Let $\Omega_1 \subset \Omega_2$ be two properly convex open sets; then for any Borel set A of Ω_1 , we have $\mu_{\Omega_2}(A) \leq \mu_{\Omega_1}(A)$.

1.3. Coxeter Group. Coxeter groups are going to be the main object of this paper, so we take the time to recall some basic facts.

Definition 1.2. A Coxeter system is the data of a finite set S and a symmetric matrix $M = (M_{st})_{s,t \in S}$ such that the diagonal coefficients $M_{ss} = 1$ and the other coefficients $M_{st} \in \{2, 3, \dots, n, \dots, \infty\}$. The cardinality of S is called the rank of the Coxeter system (S, M) . With a Coxeter system, one can build a Coxeter group W_S ; it is a group defined by generators and relations. The generators are the elements of S and we impose the relations $(st)^{M_{st}} = 1$ for all $s, t \in S$ such that $M_{st} \neq \infty$.

There are two basic objects associated to a Coxeter system or a Coxeter group: its Coxeter diagram and its Gram matrix. We recall the definition of these two objects and the basic consequences.

One can associate to W a labelled graph, also denoted by W , called the Coxeter diagram of W . The vertices of W are the elements of S . Two vertices $s, t \in S$ are linked by an edge if and only if $M_{st} \neq 2$. The label of an edge linking s to t in W is the number $M_{st} > 2$. A Coxeter group is irreducible when its Coxeter graph is connected. Of course, any Coxeter group is the direct product of the Coxeter groups associated to the connected components of its Coxeter graph.

One also associates to W a symmetric matrix of size the cardinality of S , namely its Gram matrix $\text{Cos}(W)$, defined by the formula

$$(\text{Cos}(W))_{st} = -2 \cos\left(\frac{\pi}{M_{st}}\right), \quad \text{for all } s, t \in S.$$

An irreducible Coxeter group W is a *spherical Coxeter group* (resp. *affine Coxeter group*) if its Gram Matrix is positive definite (resp. positive but not definite). Vinberg and Margulis showed that an irreducible Coxeter group which is not spherical nor affine is large¹⁰ in [MV00]. Therefore an irreducible Coxeter group is either spherical, affine or large.

More generally a Coxeter group is *spherical* (resp. *affine* resp. *euclidean*) when all its connected components are irreducible spherical Coxeter group (resp. affine resp. affine or spherical).

The irreducible spherical and affine Coxeter groups have been classified (by Coxeter in [Cox34] for the spherical case). We reproduce the list of those Coxeter diagrams in Figures 2 and 3. We use the usual convention that an edge that should be labelled 3 has in fact no label. We stress that among them the only ones which are not trees or have an edge labelled ∞ are the affine Coxeter diagram named \tilde{A}_n for $n \geq 1$. As already remarked by Vinberg, those Coxeter groups play a special role in this context.

1.4. The face of a properly convex closed (or open) set. Let C be a properly convex closed subset of \mathbb{S}^d . We introduce the following equivalence relation on C ; $x \sim y$ when the segment $[x, y]$ can be extended beyond x and y . The equivalence classes of \sim are called *the open faces of C* , the closure of an open face is a *face of C* . The *support* of a face or an open face is the smallest projective space containing it. The *dimension of a face* is the dimension of its support. For properly convex open subsets, we just apply this definition to their closure, so a face of Ω is a subset of $\bar{\Omega}$.

The interior of a face F in its support (i.e its relative interior) is equal to the unique open face f such that $\bar{f} = F$. Finally, one should remark that if f is an open face of C then f is a properly convex open set in its support. The only face of dimension d is C . A face of dimension $d - 1$ is called a *facet*, a face of dimension 0 a *vertex*, a face of dimension 1 an *edge* and a face of dimension $d - 2$ a *ridge*.

1.5. Mirror polytope. A *projective polytope* is a properly convex closed set P of $\mathbb{S}(V)$ with non-empty interior such that there exists a finite number of linear form $\alpha_1, \dots, \alpha_r$ on V such that $P = \mathbb{S}(\{x \in V \setminus \{0\} \mid \alpha_i(x) \leq 0, i = 1, \dots, r\})$.

A *projective reflection* is an element of $\mathrm{SL}^\pm(V)$ of order 2 which is the identity on a hyperplane. Each projective reflection σ can be written as $\sigma = \mathrm{Id} - \alpha \otimes v$ where α is a linear form and v a vector such that $\alpha(v) = 2$. This notation means that $\sigma(x) = x - \alpha(x)v$.

A *projective rotation* is an element of $\mathrm{SL}(V)$ which is the identity on a codimension 2 subspace H and conjugate to the 2×2 matrix $\begin{pmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{pmatrix}$ on a plan Π such that $H \oplus \Pi = V$. The two following lemmas are easy but essential.

¹⁰ Admits a finite index subgroup which admits an onto morphism on a non-abelian free group.




Figure 2. Irreducible spherical diagram.




Figure 3. Irreducible affine diagram.

Lemma 1.3 (Vinberg, Proposition 6 of [Vin71]). *Let $\sigma_s = \text{Id} - \alpha_s \otimes v_s$ and $\sigma_t = \text{Id} - \alpha_t \otimes v_t$ be two reflections of \mathbb{R}^2 . Let Γ be the group generated by σ_s and σ_t . Let \mathcal{C} be the cone $\{x \in \mathbb{R}^2 \mid \alpha_s(x) \leq 0 \text{ and } \alpha_t(x) \leq 0\}$. If the sets $(\gamma(\mathcal{C}))_{\gamma \in \Gamma}$ have disjoint interiors, then*

$$\left. \begin{aligned} & \alpha_s(v_t) \leq 0 \text{ and } \alpha_t(v_s) \leq 0 \quad (1) \\ \text{and} & \\ & \alpha_s(v_t) = 0 \iff \alpha_t(v_s) = 0. \quad (2) \end{aligned} \right\} \quad (C)$$

Lemma 1.4 (Vinberg, Propositions 6 and 7 of [VIN71]). *With the same notation, if condition (C) is satisfied then the group Γ preserves a symmetric bilinear form b on \mathbb{R}^2 .*

- (1) *If $\alpha_s(v_t)\alpha_t(v_s) < 4$ then b is positive definite and the element $\sigma_s\sigma_t$ is a rotation of angle $2\theta_{st}$ where $\alpha_s(v_t)\alpha_t(v_s) = 4\cos^2(\theta_{st})$. In particular, the group Γ is discrete if and only if the number $m_{st} = \frac{\pi}{\theta_{st}}$ is an integer.*
- (2) *If $\alpha_s(v_t)\alpha_t(v_s) > 4$ then b is of signature $(1, 1)$, the element $\sigma_s\sigma_t$ is loxodromic,¹¹ the action on \mathbb{P}^1 preserves a unique properly convex open Ω set, and the action on Ω is cocompact.*
- (3) *Otherwise $\alpha_s(v_t)\alpha_t(v_s) = 4$, b is positive and degenerate, the element $\sigma_s\sigma_t$ is unipotent,¹² the action on \mathbb{P}^1 preserves a unique affine chart \mathbb{A}^1 , and the action on \mathbb{A}^1 is cocompact.*

These actions on \mathbb{R}^2 are described by Figure 4.




Figure 4

¹¹ Here, this means that $\sigma_s\sigma_t$ is diagonalizable over \mathbb{R} .

¹² Here, this means that $(\sigma_s\sigma_t - \text{Id})^2 = 0$.

The two previous lemmas motivate the following definition:

Definition 1.5. A *mirror polytope* is a convex projective polytope P with the data of a projective reflection σ_s across each facet s of P , such that for any two facets s and t of P such that $s \cap t$ is a ridge of P , the pair $\{\sigma_s, \sigma_t\}$ satisfies condition (C). We say that *the dihedral angle between the facets s and t is θ_{st}* when we have $\alpha_s(v_t)\alpha_t(v_s) = 4 \cos^2(\theta_{st})$. Otherwise, we say that the angle is 0. Two mirror polytopes are *isomorphic* if one can find an isomorphism of vector spaces which sends the first polytope to the second, and sends the reflections of the first to the reflections of the second. When P and Q are isomorphic, we will write $P \simeq Q$.

Notation 1.6. The following notation will be used along this text. Let P be a mirror polytope; the symbol S will denote the set of facets of P . We can always write $P = \mathbb{S}(\{x \in V \setminus \{0\} \mid \alpha_s(x) \leq 0, s \in S\})$. For each facet $s \in S$, we denote by σ_s the reflection of P which fixes each point of s . We can write it $\sigma_s = \text{Id} - \alpha_s \otimes v_s$ with $v_s \in V$ and $\alpha_s(v_s) = 2$. Be careful that the couple (α_s, v_s) is unique up to a multiplicative positive constant,¹³ but nothing will depend on this choice. The point $[v_s] \in \mathbb{S}(V)$, which is unique, is called the *polar* of the facet s (or of σ_s) and the hyperplane $\{x \in \mathbb{S}^d \mid \alpha_s(x) = 0\}$ is called the *support* of the facet s or of σ_s . We will denote by the symbol Γ_P or simply Γ the group generated by the reflections σ_s for $s \in S$.

Corollary 1.7. Let P be a mirror polytope. If the sets $\gamma(\overset{\circ}{P})$ are disjoint for $\gamma \in \Gamma_P$, then the family $(\alpha_s(v_t))_{s,t \in S}$ verifies condition (C) and

$$\left. \begin{aligned} & \alpha_s(v_t)\alpha_t(v_s) = 4 \cos^2(\theta_{st}), & (1) \\ & \text{and the number } m_{st} = \frac{\pi}{\theta_{st}} \\ & \text{is an integer greater or equal to 2} & (D) \\ \text{or} & \\ & \alpha_s(v_t)\alpha_t(v_s) \geq 4. & (2) \end{aligned} \right\}$$

Definition 1.8. A mirror polytope P is a *Coxeter polytope* when all its dihedral angles are sub-multiples¹⁴ of π .

If P is a Coxeter polytope, the *Coxeter system associated to P* is the Coxeter system (S, M) , where S is the set of facets of P and where for all $s, t \in S$, we have $M_{st} = m_{st}$ if the facets $s, t \in S$ are such that $s \cap t$ is a ridge of P and $\theta_{st} = \frac{\pi}{m_{st}}$, otherwise $M_{st} = \infty$. We will denote by the letter W_P or simply W the *Coxeter group associated to the system (S, M)* .

¹³ By the action $\lambda \cdot (\alpha_s, v_s) = (\lambda\alpha_s, \lambda^{-1}v_s)$.

¹⁴ Precisely, $\theta = \frac{\pi}{m}$ with m an integer greater than or equal to 2 OR $m = \infty$.




Figure 5. Illustration of equation (C).

Remark 1.9. Figure 5 shows a pentagon P . Any mirror structure on this polygon verifies that the polar $[v_i]$ of the facet are in the grey triangle given by the facet i . This is a consequence of the inequalities in (C). We will see that these inequalities have usefull implications.

1.6. The limit set of positively proximal subgroup of $SL_{d+1}(\mathbb{R})$. In this section, we just state a theorem of existence of limit set. We will give a more detailed discussion in paragraph 7.4.

1.6.1. Strongly irreducible case

Theorem 1.10 (Benoist, Lemma 2.9 and 3.3 of [BEN00]). *Let Γ be a strongly irreducible subgroup of $SL_{d+1}(\mathbb{R})$ preserving a properly convex open set. There exists a smallest closed Γ -invariant subset Λ_Γ for the action of Γ on \mathbb{P}^d . This closed subset is called the **limit set** of Γ .*

Corollary 1.11. *Let Γ be a strongly irreducible subgroup of $SL_{d+1}(\mathbb{R})$ preserving a properly convex open set. There exists a smallest and a largest Γ -invariant convex open subset for the action of Γ on \mathbb{P}^d .*

1.6.2. Irreducible case

Lemma 1.12 (Benoist, Lemma 2.9 and 3.3 of [BEN00]). *Let Γ be an irreducible subgroup of $\mathrm{SL}_{d+1}(\mathbb{R})$ preserving a properly convex open set Ω . Let Γ_0 be the Zariski connected component of Γ . There exists a decomposition $\mathbb{R}^{d+1} = \bigoplus_{i=1,\dots,r} E_i$ in strongly irreducible Γ_0 -sub-modules such that the action of Γ_0 on each factor preserves a properly convex open cone. The **limit set** of Γ is the union of the limit set of Γ_0 in $\mathbb{P}(E_i)$.*

2. The Theorem of Tits and Vinberg and the Theorems of Vinberg

In this section, we recall the Theorem of Tits and Vinberg and the Theorems of Vinberg.

2.1. Tiling theorem. To avoid any confusion, we recall a general definition of a tiling.

Definition 2.1. A family $(E_i)_{i \in I}$ of closed set *tiles* a topological set X when we have the following three conditions. For all $i \in I$, the interior of E_i is dense in E_i , the union of the E_i is X and for all $i \neq j$ in I , the intersection of the interiors of E_i and E_j is empty.

If (S, M) is a Coxeter system then for every subset S' of S , one can consider the Coxeter group $W_{S'}$ associated to the Coxeter system (S', M') , where M' is the restriction of M to S' . Theorem 2.2 shows that the natural morphism $W_{S'} \rightarrow W_S$ is injective. Therefore, $W_{S'}$ may be identified with the subgroup of W_S generated by the subset S' .

If P is a Coxeter polytope and f is a face (or an open face) of P , and $\bar{f} \neq P$, then we will write $S_f = \{s \in S \mid f \subset s\}$ and $W_f = W_{S_f}$.

Let (S, M) be a Coxeter system. A *standard parabolic subgroup* of the Coxeter group W_S is a subgroup generated by some elements of S . A *parabolic subgroup* of W_S is conjugate of a standard parabolic subgroup.

Theorem 2.2 (Tits, see Chapter V [Bou68] for the Tits's simplex or Vinberg [Vin71]). *Let P be a Coxeter polytope of $\mathbb{S}(V)$, W_P be the associated Coxeter group and Γ_P the group generated by the projective reflections $(\sigma_s)_{s \in S}$. Then,*

- (1) *the morphism $\sigma: W_P \rightarrow \Gamma_P$ defined by $\sigma(s) = \sigma_s$ is an isomorphism;*
- (2) *the polytopes $(\gamma(P))_{\gamma \in \Gamma_P}$ tile a convex set \mathcal{C}_P of $\mathbb{S}(V)$;*
- (3) *the group Γ_P acts properly on $\Omega_P = \overset{\circ}{\mathcal{C}}_P$, the interior of \mathcal{C}_P ;*
- (4) *the group Γ_P is a discrete subgroup of $\mathrm{SL}^\pm(V)$;*
- (5) *an open face f of P lies in Ω_P if and only if the Coxeter group W_f is finite;*
- (6) *for every parabolic subgroup U of W_P , the union $\bigcup_{\gamma \in U} \gamma(P)$ is convex.*

Corollary 2.3. *The convex set \mathcal{C}_P is open if and only if the action of Γ_P on Ω_P is cocompact if and only if for every vertex p of P the Coxeter group W_p is finite. Following Vinberg, we will say that in this case, P is **perfect**.¹⁵*

The following theorem can give the impression to be a corollary, but in fact Vinberg uses it to conclude the proof of his theorem (see Lemma 10 of [VIN71]).

Theorem 2.4 (Coxeter [Cox34]). *The convex set \mathcal{C}_P is the projective sphere $\mathbb{S}(V)$ if and only if the group W_P is finite if and only if the Coxeter group W_P is spherical.*

Remark 2.5. The sixth point of Theorem 2.2 is not explicit in Vinberg's article but it is an easy consequence of the techniques he develops.

2.2. The Cartan Matrix of a Coxeter polytope

Definition 2.6. A matrix A of $M_m(\mathbb{R})$ is a *Cartan matrix* when

- $a_{ii} = 2$ for all $i = 1, \dots, m$,
- $a_{ij} = 0 \iff a_{ji} = 0$ for all $i, j = 1, \dots, m$,
- all non-diagonal coefficients of A are negative or null.

A matrix is *reducible* if after a simultaneous permutation of the rows and the columns, one as a non trivial diagonal bloc matrix. A matrix is *irreducible* if and only if it is not reducible.

The theorem of Perron-Frobenius shows that *the spectral radius of an irreducible matrix with positive or null coefficients is a simple eigenvalue*. Hence, an irreducible Cartan matrix A has a unique eigenvalue λ_A of minimal modulus. We will say that A is of *positive type*, *zero type* or *negative type* when $\lambda_A > 0$, $\lambda_A = 0$ or $\lambda_A < 0$.

Given a Coxeter polytope P , one can define the matrix A where $A_{ij} = \alpha_i(v_j)$. By definition of a Coxeter polytope, A is a Cartan matrix; we will call it the *Cartan matrix associated to the Coxeter polytope P* and denoted it A_P .

Of course, the Coxeter group W_P is irreducible if and only if the Cartan matrix A_P is irreducible. In that case, we say that P is of *positive type* (resp. *zero type* or *negative type*) according to the type of A_P .

If the Coxeter group W_P is not irreducible, then the Cartan matrix A_P is the sum of its irreducible components, we say that P is of *positive type*, (resp. *zero type*, or *negative type*) if all the irreducible components are of *positive type*, (resp. *zero type* or *negative type*). It is easy to find a Coxeter polytope P , such that the components of A_P do not have not the same type.

¹⁵ Definition 8 of [VIN71].

2.3. Tits's simplex. To each Coxeter group W , we can associate a Coxeter polytope. The polytope will be a simplex of dimension the rank of W minus 1. The construction¹⁶ is the following.

Suppose that W arises from the Coxeter system $M = (M_{st})_{s,t \in S}$. Consider the vector space $V = (\mathbb{R}^S)^*$, and denote by $(e_s)_{s \in S}$ the canonical basis of \mathbb{R}^S . We consider the simplicial cone $\mathcal{C} = \{\varphi \in (\mathbb{R}^S)^* \mid \varphi(e_s) \leq 0 \text{ for all } s \in S\}$; the simplex we want is $P = \mathcal{S}(\mathcal{C})$. The reflection associated to the element $s \in S$ is the reflection across the facet $\mathcal{S}(\{\varphi \mid \varphi(e_s) = 0\}) \cap P$, and given by the formula $\sigma_s(\varphi) = \varphi - 2\varphi(e_s)B_W(e_s, \cdot)$, where B_W is the symmetric bilinear form given by $B_W(e_s, e_t) = -\cos(\frac{\pi}{M_{st}})$.

The resulting Coxeter polytope will be called the *Tits simplex associated to W* and denoted by Δ_W . The polar of the facet s is the point $[2B_W(e_s, \cdot)]$ of $\mathcal{S}((\mathbb{R}^S)^*)$. We stress that the group Γ_{Δ_W} preserves the symmetric bilinear form B_W .

2.4. Proper convexity of Ω_P

Theorem 2.7 (Vinberg, Lemma 15 and Proposition 25 in [VIN71]). *Let P be a Coxeter polytope of \mathbb{S}^d . The convex set Ω_P is properly convex if and only if the Cartan matrix A_P of P is of negative type.*

Remark 2.8. The terminology in [VIN71] and the terminology we use can be in opposition. A cone C is strictly convex for [VIN71] when it is properly convex for us. Vinberg prefer to speak about a reduced linear Coxeter group while we prefer to talk about a Coxeter polytope.

2.5. Irreducible Coxeter polytope. The following proposition gives the shape of Ω_P via the type of A_P .

Proposition 2.9 (Vinberg [VIN71]). *Let P be an irreducible Coxeter polytope of \mathbb{S}^d . Let W be the Coxeter group associated to P . We are in exactly one the following five cases.*

- (1) *The Coxeter group W is spherical; in that case:*
 - A_P is of positive type and of rank $d + 1$,
 - $\Omega_P = \mathbb{S}^d$,
 - in fact, $P \simeq \Delta_W$.
- (2) *The Coxeter group W is affine but not of type \tilde{A}_n ; in that case:*
 - A_P is of zero type and of rank d ,
 - Ω_P is an affine chart,
 - in fact, $P \simeq \Delta_W$,
 - the action of Γ_P on Ω_P is cocompact and preserves a euclidean metric.

¹⁶ In order to get a Coxeter polytope, one has to take the dual of the standard representation introduced by Tits.

(3) *The Coxeter group W is affine of type \tilde{A}_n , and A_P is of zero type; in that case:*

- A_P is of rank d ,
- Ω_P is an affine chart,
- in fact, $P \simeq \Delta_W$,
- the action of Γ_P on Ω_P is cocompact and preserves a euclidean metric.

(4) *The Coxeter group W is affine of type \tilde{A}_n , and A_P is of negative type; in that case:*

- A_P is of rank $d + 1$,
- Ω_P is a simplex; in particular Ω_P is a properly convex open set,
- the action of Γ_P on Ω_P is cocompact.

(5) *The Coxeter group W is large; in that case:*

- A_P is of rank $r \leq d + 1$,
- Ω_P is a properly convex open set (which is not a simplex).

Explanation of proof. The first point is given by the Proposition 22 of [VIN71], the second and third points are given by Proposition 23 of [VIN71]. Theorem 2.7 shows that in the fourth and fifth point the convex set Ω_P is properly convex. Hence, the only thing left to explain is that in the fourth point the convex set Ω_P has to be a simplex. This is Lemma 8 in [MV00] of Margulis and Vinberg. \square

2.6. Product of Coxeter polytopes

2.6.1. Spherical projective completion. If V is a vector space, then V is an affine chart of $\mathbb{S}(V \oplus \mathbb{R})$. The space $\mathbb{S}(V)$ is a projective hyperplane of $\mathbb{S}(V \oplus \mathbb{R})$. Finally, $\mathbb{S}(V \oplus \mathbb{R}) \setminus \mathbb{S}(V)$ has two connected components, each isomorphic to the affine space V . Hence, $\mathbb{S}(V)$ is the *hyperplane at infinity of V* and $\mathbb{S}(V \oplus \mathbb{R})$ is the *spherical projective completion of V* .

2.6.2. The Coxeter cone above a Coxeter polytope. Let P be a Coxeter polytope of \mathbb{S}^d , then $\mathbb{S}^{-1}(P)$ is a convex cone of \mathbb{R}^{d+1} . The affine space \mathbb{R}^{d+1} is an affine chart of its spherical projective completion $\mathbb{S}^{d+1} = \mathbb{S}(\mathbb{R}^{d+1} \oplus \mathbb{R})$. We denote by H_∞ the projective subspace \mathbb{S}^d in \mathbb{S}^{d+1} , i.e. the hyperplane at infinity of \mathbb{R}^{d+1} .

The closure $\overline{\mathbb{S}^{-1}(P)}$ of $\mathbb{S}^{-1}(P)$ in \mathbb{S}^{d+1} is a polytope, each facet of $\mathbb{S}^{-1}(P)$ has a reflection coming from P , except the facet $H_\infty \cap \overline{\mathbb{S}^{-1}(P)}$ to which we associate the reflection across H_∞ with polar the origin of the affine chart defined by H_∞ which does not contain $\mathbb{S}^{-1}(P)$.

This Coxeter polytope associated to P will be called *the Coxeter cone above P* and denoted by the symbol $P \otimes \cdot$, it is a Coxeter polytope. One should remark that the polytope $P \otimes \cdot$ has one facet f_∞ more than P , all the ridges included in the facet f_∞ have dihedral angle $\frac{\pi}{2}$, so $W_{P \otimes \cdot} = W_P \times \mathbb{Z}/2\mathbb{Z}$ where the factor $\mathbb{Z}/2\mathbb{Z}$ is given by the reflection σ_∞ across H_∞ . Finally, one should remark that $(P \otimes \cdot) \cap H_\infty$ is the Coxeter polytope P , and that the convex set $\Omega_{P \otimes \cdot}$ is the convex hull of $\Omega_P \subset H_\infty, 0$ and $\sigma_\infty(0)$. Figure 6 may help to understand the situation.




Figure 6. The Coxeter cone above a Coxeter polytope.

In particular, the convex set $\Omega_{P \otimes \cdot}$ is never properly convex and if P is not elliptic, the action of $W_{P \otimes \cdot}$ on $\Omega_{P \otimes \cdot}$ is never cocompact.

2.6.3. The product of two convex sets. A sharp convex cone \mathcal{C} of a vector space V is *decomposable* if we can find a decomposition $V = V_1 \oplus V_2$ of V such that this decomposition induces a decomposition of \mathcal{C} (i.e. $\mathcal{C}_i = V_i \cap \mathcal{C}$ and $\mathcal{C} = \mathcal{C}_1 \times \mathcal{C}_2$). A sharp convex cone is *indecomposable* if it is not decomposable.

We induce this definition to properly convex open set. A properly convex open set Ω is *indecomposable* if the cone $\mathbb{S}^{-1}(\Omega)$ above Ω is indecomposable.

This definition suggests a definition of a *product* of two properly convex open sets which is not the Cartesian product. Given two properly convex open sets Ω_1 and Ω_2 of the spherical projective spaces $\mathbb{S}(V_1)$ and $\mathbb{S}(V_2)$, we define a new

properly convex open set $\Omega_1 \otimes \Omega_2$ of the spherical projective space $\mathbb{S}(V_1 \times V_2)$ by the following formula: if \mathcal{C}_i is the cone $\mathbb{S}^{-1}(\Omega_i)$ then $\Omega_1 \otimes \Omega_2 = \mathbb{S}(\mathcal{C}_1 \times \mathcal{C}_2)$.

It is important to note that if Ω_i is of dimension d_i then $\Omega_1 \otimes \Omega_2$ is of dimension $d_1 + d_2 + 1$. Here is a more pragmatic way to see this product. Take two properly convex subsets ω_i of a spherical projective space $\mathbb{S}(V)$ with support in direct sum, the ω_i are not open but we assume that they are open in their supports; assume also that there exists an affine chart containing both ω_i . Then the convex hull in such an affine chart of $\omega_1 \cup \omega_2$ is $\omega_1 \otimes \omega_2$. Some call $\omega_1 \otimes \omega_2$ the *join* of ω_1 and ω_2 .

Just to be clear, we give the definition of a cone in the projective context. A properly convex open set Ω is a *cone* when there exist two open faces ω_1 and ω_2 of Ω such that ω_1 is a singleton, ω_2 is of dimension $d - 1$ and $\Omega = \omega_1 \otimes \omega_2$. The face ω_1 is called the *summit* of the cone and ω_2 is called the *basis* of the cone.

2.6.4. The product of two Coxeter polytopes. Let P and Q be two Coxeter polytopes of \mathbb{S}^d and \mathbb{S}^e . Then $\mathbb{S}^{-1}(P)$ and $\mathbb{S}^{-1}(Q)$ are convex cones of \mathbb{R}^{d+1} and \mathbb{R}^{e+1} . We can take the Cartesian product of these two cones to get a convex cone $\mathcal{C}_{P,Q}$ of \mathbb{R}^{d+e+2} and then project this cone to \mathbb{S}^{d+e+1} to get a polytope $P \otimes Q$ of dimension $d + e + 1$.

The facet of $P \otimes Q$ are in correspondence with the facets of P union the facets of Q . By extending trivially each reflection from \mathbb{R}^{d+1} (or \mathbb{R}^{e+1}) to \mathbb{R}^{d+e+2} , we get a Coxeter polytope whose Coxeter group is $W_P \times W_Q$ and we get $\Omega_{P \otimes Q} = \Omega_P \otimes \Omega_Q$.

2.6.5. Return to the cone. One can remark that the sphere \mathbb{S}^0 of dimension 0 is just two points and is tiled by the Coxeter group $\mathbb{Z}/2\mathbb{Z}$ via the Coxeter polytope of dimension 0 i.e. a point, hence the Coxeter cone $P \otimes \cdot$ above P is the product of P with the Coxeter polytope of dimension 0. This explains our notation.

2.6.6. Decomposability

Definition 2.10. A Coxeter polytope P is *decomposable* if one can find two Coxeter polytopes such that $P = Q \otimes R$, otherwise P is *indecomposable*.

Remark 2.11. If a Coxeter polytope P is decomposable then the Coxeter group W_P is reducible. The converse is false, think of the right angled square, this Coxeter polygon is indecomposable but the associated Coxeter group is reducible.

2.6.7. Theorem of decomposability of Vinberg

Theorem 2.12 (Corollary 4 of [VIN71]). *Let P be a Coxeter polytope of \mathbb{S}^d . We denote by W the Coxeter group associated to P . Suppose that W is reducible. If $\text{Rank}(A_P) = d + 1$ or P is a simplex then P is decomposable.*

2.7. Elliptic, parabolic, loxodromic Coxeter polytopes

Definition 2.13. A Coxeter polytope P of \mathbb{S}^d is

- (1) *elliptic* when A_P is of positive type,
- (2) *parabolic* when A_P is of zero type and of rank d ,
- (3) *loxodromic* when A_P is of negative type and of rank $d + 1$.

Remark 2.14. Let P be a Coxeter polytope. If P is elliptic then the rank of A_P is necessarily $d + 1$. If A_P is of zero type then the rank of A_P cannot be $d + 1$, but it can be strictly less than d . If A_P is of negative type then the rank of A_P can be strictly less than $d + 1$.

Remark 2.15. We recall that for Vinberg, a Coxeter polytope P is *hyperbolic* if P is loxodromic and Γ_P preserves an ellipsoid (i.e. Γ_P is a subgroup of a conjugate of $\mathrm{SO}_{d,1}^\circ(\mathbb{R})$).

2.8. About irreducibility

2.8.1. Characterisation of the irreducibility of Γ_P

Theorem 2.16 (Vinberg, Proposition 18 and Corollary of Proposition 19 of [VIN71]). *Let P be a Coxeter polytope of \mathbb{S}^d . Then the following assertions are equivalent:*

- (1) *the representation $\rho : W_P \rightarrow \mathrm{SL}_{d+1}^\pm(\mathbb{R})$ is irreducible;*
- (2) *the Coxeter group W_P is irreducible and the family $(v_s)_{s \in S}$ generates \mathbb{R}^{d+1} ;*
- (3) *the Coxeter group W_P is irreducible and the Cartan Matrix A_P of P is of rank $d + 1$.*

In particular, if W_P is infinite then ρ is irreducible if and only if the Coxeter polytope P is irreducible and loxodromic.

Remark 2.17. Let P be a Coxeter polytope. From Theorem 1.10, we learn that we can define a limit set for the group Γ_P as soon as the group Γ_P is irreducible. Hence, Theorem 2.16 shows that the limit set of Γ_P is defined as soon as P is irreducible and loxodromic. We will denote the limit set of Γ_P by the symbol Λ_P or Λ_Γ . The limit set is a crucial object for us. Its definition is easier to handle when the group Γ_P is strongly irreducible. The next theorem shows that if P is irreducible and loxodromic then Γ_P is strongly irreducible except if W_P is affine.

2.8.2. From irreducible to strongly irreducible

Theorem 2.18 (folklore). *Let P be a Coxeter polytope of \mathbb{S}^d . Suppose that the representation $\rho: W_P \rightarrow \mathrm{SL}_{d+1}^{\pm}(\mathbb{R})$ is irreducible. Then we have the following exclusive trichotomy:*

- (1) *the Coxeter group W_P is spherical and $\Omega_P = \mathbb{S}^d$;*
- (2) *the Coxeter group W_P is affine of type \tilde{A}_n and Ω_P is a simplex;*
- (3) *the Coxeter group W_P is large, Ω_P is a properly convex open set and the linear group Γ_P is strongly irreducible.*

We give a short explanation for this theorem since we did not find any proof of it in the literature, although the result is surely known.

Proposition 2.19. *Let Γ be an infinite group of $\mathrm{SL}_{d+1}(\mathbb{R})$ acting properly on a convex set Ω of \mathbb{S}^d . If Γ is an irreducible subgroup of $\mathrm{SL}_{d+1}(\mathbb{R})$, then Ω is a properly convex open set.*

Proof. The vector space generated by $\bar{\Omega} \cap -\bar{\Omega}$ is preserved by Γ , therefore either $\Omega = -\Omega$ or $\bar{\Omega} \cap -\bar{\Omega} = \emptyset$. In the first case, $\Omega = \mathbb{S}^d$ and Γ has to be finite since the action is proper. The second condition means that Ω is properly convex. \square

Proof of Theorem 2.18. From Theorem 2.16, we know that $W = W_P$ has to be an irreducible Coxeter group, therefore we have three cases: W can be spherical, affine, or large. If W is spherical then Theorem 2.2 shows that $\Omega = \Omega_P = \mathbb{S}^d$. If W is not spherical then W is infinite and Proposition 2.19 shows that Ω is properly convex. If W is affine then Proposition 2.9 shows that W is of type \tilde{A}_n and Ω is a simplex. Of course, in that case, the linear group $\Gamma = \Gamma_P$ is not strongly irreducible since the vertices of Ω have to be fixed by a finite-index subgroup of Γ .

If W is large, it remains to show that Γ is a strongly irreducible subgroup of $\mathrm{SL}_{d+1}(\mathbb{R})$. Suppose the representation is not strongly irreducible, then consider Γ_0 the Zariski connected component of Γ . The subgroup Γ_0 of Γ is of finite index. The vector space \mathbb{R}^{d+1} is the sum of the strongly irreducible Γ_0 -submodules $\mathbb{R}^{d+1} = \bigoplus_{i \in I} E_i$. In particular, Γ_0 splits as a non-trivial direct product and this is absurd by Theorem 2.20 below. \square

Theorem 2.20 (Paris [PAR07] or Proposition 8 of de Cornulier and de la Harpe in [CH07]). *No finite index subgroup of a large irreducible Coxeter group splits as a non-trivial direct product.*

3. The setting

The study of the geometry around a vertex will be crucial in the sequel, so we introduce some definitions.

3.1. Link of a Coxeter polytope. Let P be a Coxeter polytope of \mathbb{S}^d and p be vertex of P . The *link* P_p of P at p is the set of half-lines starting at p intersecting P . It is a Coxeter polytope included in the projective space $\mathbb{S}(\mathbb{R}^{d+1}/p^2) = \mathbb{S}_p^{d-1}$ where p^2 is the line generated by the half-line p .

To avoid confusion, we get P_p by the following procedure:

- (1) forget all the facets of P not containing p , so that you get a convex cone whose summit is p ;
- (2) forget at the same time all the reflections around facets of P not containing p , so that you get a Coxeter convex cone whose summit is p ;
- (3) consider the set P_p of half-line starting at p intersecting P , look at it in the projective sphere \mathbb{S}_p^{d-1} ; it is a convex subset, better it is a polytope.
- (4) the reflections around the facets containing p fix p , so they pass to the quotient \mathbb{R}^{d+1}/p^2 and acts as reflection around the facets of P_p .

Since P_p is a Coxeter polytope of \mathbb{S}_p^{d-1} we can apply to it the Vinberg-Tits Theorem 2.2 to get a convex subset $\Omega_p := \Omega_{P_p}$ of \mathbb{S}_p^{d-1} . We shall concentrate on a special class of Coxeter polytope for which this procedure gives a lot of information.

3.2. Quasi-perfect, 2-perfect Coxeter polytopes

Proposition 3.1. *Let P be a Coxeter polytope. Then the following are equivalent:*

- (1) *the intersection $P \cap \partial\Omega_P$ is finite;*
- (2) *the intersection $P \cap \partial\Omega_P$ is included in the set of vertices of P ;*
- (3) *for every edge e of P , the Coxeter group W_e is finite;*
- (4) *for every vertex p of P , the Coxeter polytope P_p is perfect.*

*In that case, we say that the Coxeter polytope P is **2-perfect**.*

Proof. Since the convex polytope P is included in the closed convex set $\overline{\Omega_P}$, the relative interior of an edge of P intersects the boundary $\partial\Omega_P$ if and only if it is included in the boundary, so (1) \iff (2). The implication (2) \iff (3) is a direct consequence of Theorem 2.2 (5). For (3) \iff (4), there is a natural correspondence between the edges of P and the vertices of the link $(P_p)_{p \in \mathcal{V}}$, where \mathcal{V} is the set of vertices of P , by definition of P_p . The equivalence is then a consequence of Corollary 2.3. \square

Remark 3.2. Every Coxeter polygon is 2-perfect and a Coxeter polytope of dimension 3 is 2-perfect if and only if none of its dihedral angle is 0.

In [VIN71], Vinberg introduces the notion of quasi-perfect Coxeter polytope.

Definition 3.3. A Coxeter polytope P is *quasi-perfect* when P is 2-perfect and for every vertex p of P , the Coxeter polytope P_p is either elliptic or parabolic.

Remark 3.4. Let (S, M) be a Coxeter system and W the corresponding Coxeter group. The Tits simplex Δ_W is perfect if and only if for every subsystem S' such that $\text{Card}(S \setminus S') = 1$ we have $W_{S'}$ finite. A large irreducible Coxeter group such that Δ_W is perfect is usually called a *Lannér* Coxeter group. These groups have been classified by Lannér in [LAN50].

The Tits simplex Δ_W is quasi-perfect if and only if for every subsystem S' such that $\text{Card}(S \setminus S') = 1$ we have $W_{S'}$ finite or irreducible affine. A large irreducible Coxeter group such that Δ_W is quasi-perfect is usually called a *quasi-Lannér* Coxeter group (sometimes Koszul Coxeter group). They have been classified by Koszul and Chein in [KOS67, CHE69].

Finally, the Tits simplex Δ_W is 2-perfect if and only if for every subsystem S' such that $\text{Card}(S \setminus S') = 2$ we have $W_{S'}$ finite. They are sometimes called *Lorentzian* Coxeter groups. They have been classified by Maxwell in [MAX82], the complete list have been published by Chen and Labbé in [CL15].

3.3. A geometric quadrichotomy for quasi-perfect Coxeter polytope. Vinberg in [VIN71] arranges quasi-perfect polytope into four families.

Theorem 3.5 (Vinberg, Proposition 26 of [VIN71]). *Let P be a quasi-perfect Coxeter polytope; then P is in one of the following four exclusive cases:*

- (1) *elliptic,*
- (2) *parabolic,*
- (3) *loxodromic and irreducible, or*
- (4) *decomposable and not elliptic; in fact P is of the form $P = Q \otimes \cdot$, where Q is parabolic.*

Remark 3.6. We stress that in the last case of Theorem 3.5, the Coxeter polytope is not perfect. Hence, if P is perfect and decomposable then P is elliptic.

Remark 3.7. If a Coxeter polytope P is parabolic then P is indecomposable even if W_P is not irreducible. Indeed, if P were decomposable then $P = P_1 \otimes \cdots \otimes P_r$ where P_i is of dimension d'_i . Each Γ_{P_i} is virtually isomorphic to $\mathbb{Z}^{d'_i}$ with $d'_i \geq d_i$ since Γ_{P_i} acts properly on Ω_{P_i} . But, the group Γ_P acts cocompactly on an affine chart of $\mathbb{S}^{d'_1 + \cdots + d'_r + r - 1}$, so $\mathbb{Z}^{d'_1 + \cdots + d'_r}$ acts cocompactly hence $r = 1$ and $d'_i = d_i$.

3.4. The final context. In the case where P is a 2-perfect Coxeter polytope of \mathbb{S}^d , all the link P_p are perfect so the convex set Ω_p is either the whole space \mathbb{S}_p^{d-1} , an affine chart or a properly convex open set from Theorem 3.5. We want to understand the geometry of the action of Γ_P on Ω_P by mean of the shape of the $(\Omega_p)_{p \in \mathcal{V}}$, where \mathcal{V} is the set of vertices of P .

For any Coxeter polytope, we will say that a vertex p is *elliptic* (resp. *parabolic* or *loxodromic*) when the Coxeter polytope P_p is elliptic (resp. *parabolic* or *loxodromic*). For a 2-perfect Coxeter polytope, we have a nice trichotomy: every vertex p of P has to be either elliptic or parabolic or loxodromic.

Remark 3.8. We stress that the word spherical, affine, large, euclidean, irreducible refer to properties of Coxeter groups and the word elliptic, parabolic, loxodromic, indecomposable to properties of Coxeter polytope, and so of *linear* Coxeter groups.

4. The lemmas

4.1. The shape of a convex set around a point of its boundary. Let Ω be a properly convex open set and p be a point of $\partial\Omega$. We say that p is \mathcal{C}^1 when the hypersurface $\partial\Omega$ is differentiable at p (if and only if Ω admits a unique supporting hyperplane at p). We say that p is *not strictly convex* when there exists a non-trivial segment $s \subset \partial\Omega$ such that $p \in s$. When p is of class \mathcal{C}^1 and strictly convex, we say that p is *round*. A properly convex open set is *round* when every point of its boundary is round.

To study the boundary around a point, the following spaces are very useful. We denote by $\mathcal{D}_p(\Omega)$ (resp. $\mathcal{D}_p(\bar{\Omega})$) the space of half-lines starting at p and meeting Ω (resp. $\bar{\Omega}$). These two spaces are convex subsets of \mathbb{S}_p^{d-1} . We also a map between these spaces $\mathbb{S}_p: \partial\Omega \setminus \{p\} \rightarrow \mathcal{D}_p(\bar{\Omega})$ given by $\mathbb{S}_p(q) = [pq]$. The point p is \mathcal{C}^1 if and only if $\mathcal{D}_p(\Omega)$ is an affine chart of \mathbb{S}_p^{d-1} . The point p is strictly convex if and only if \mathbb{S}_p is injective. One should remark that \mathbb{S}_p is always onto.

Remark 4.1. Let P be a Coxeter polytope and p a vertex of P , then we have $\mathcal{D}_p(\Omega_P) = \Omega_{P_p} = \Omega_p$.

4.2. Consequence of ellipticity

Lemma 4.2. *Let P be a Coxeter polytope of \mathbb{S}^d . The action of Γ_P on \mathbb{S}^d has no global fixed point.*

Proof. The only fixed point of a reflection σ are the points of the support of σ . But, the intersection of the support of all the facets of P is empty. \square

Remark 4.3. The last lemma is false in the context of \mathbb{P}^d . The simplest example is the Coxeter triangle with dihedral angle $(\frac{\pi}{2}, \frac{\pi}{2}, \frac{\pi}{m})$. The reason for this difference is that in \mathbb{P}^d , a reflection fixes the point of its support and its polar.

Proposition 4.4. *Let P be an irreducible loxodromic Coxeter polytope and p a vertex of P . The vertex p is elliptic if and only if $p \in \Omega_P$, and in that case $p \in C(\Lambda_P)$, the open convex hull¹⁷ of Λ_P .*

Proof. Theorem 2.2 shows that $p \in \Omega_P$ if and only if W_p is finite and Theorem 2.4 shows that W_p is finite if and only if $\Omega_p = \mathbb{S}_p^{d-1}$ if and only if P_p is elliptic. So, we only have to prove that $p \in C(\Lambda_P)$. The point p is the unique fixed point of the finite group W_p acting on Ω_P since the action of W_p on \mathbb{S}_p^{d-1} has no global fixed point from Lemma 4.2. Hence, the point p belongs to $C(\Lambda_P)$ since the center of mass¹⁸ of any orbit of a finite group acting on a properly convex open set is a fixed point. \square

4.3. Consequence of parabolicity. We introduce formally the trick to show that a convex projective manifold is of finite volume. This trick has been used in [MAR12A, MAR12B, CLT15, CM14].

Definition 4.5. Let Ω be a properly convex open subset of \mathbb{S}^d and $p \in \partial\Omega$. We say that Ω admits *two ellipsoids of security at p* when there exist two ellipsoids \mathcal{E}^{int} and \mathcal{E}^{ext} such that $\mathcal{E}^{\text{int}} \subset \Omega \subset \mathcal{E}^{\text{ext}}$ and $\partial\mathcal{E}^{\text{int}} \cap \partial\Omega = \partial\mathcal{E}^{\text{ext}} \cap \partial\Omega = \{p\}$. (See Figure 7.)

Proposition 4.6. *Let Ω be a properly convex open set and $p \in \partial\Omega$. Let K be a compact subset of $\partial\Omega \setminus \{p\}$ and denote by $\mathcal{C}_{K,p}$ the convex hull of $K \cup \{p\}$ in Ω . Suppose that Ω admits two ellipsoids of security at p . Then p is a round point of $\partial\Omega_P$ and for a sufficiently small neighbourhood \mathcal{U} of p in \mathbb{S}^d we have $\mu_\Omega(\mathcal{C}_{K,p} \cap \mathcal{U}) < \infty$.*

The goal of this paragraph is to show the following proposition.

Proposition 4.7. *Let P be an irreducible loxodromic Coxeter polytope and p be a vertex of P . If the vertex p is parabolic then*

- (1) *the point p is a round point of $\partial\Omega_P$;*
- (2) *the convex set Ω_P admits two ellipsoids of security at p , which are preserved by Γ_P ;*
- (3) *there exists a neighbourhood \mathcal{U} of p in \mathbb{S}^d such that $\mu_{\Omega_P}(P \cap \mathcal{U}) < \infty$;*
- (4) *$p \in \Lambda_P$.*

¹⁷ The smallest convex open set that contains Λ_P in its closure.

¹⁸ For the existence of a center of mass for every bounded subset of a properly convex subset, we refer the reader to [MAR13] lemma 4.2.




Figure 7. Two ellipsoids of security.

Proof of Proposition 4.6. The roundness is obvious. For the finiteness of the volume, the claim is true and obvious when Ω is an ellipsoid since an ellipsoid endowed with its Hilbert metric is the projective model of hyperbolic geometry. Therefore the existence of \mathcal{E}^{int} via Proposition 1.1 implies the proposition. \square

The following lemma is due to Vinberg.

Lemma 4.8 (Vinberg, Proposition 23 of [VIN71]). *Let P be a parabolic Coxeter polytope of \mathbb{S}^d . Then Γ_P acts by euclidean transformation on the affine chart Ω_P (i.e. there exists a positive definite scalar product on Ω_P preserved by Γ_P).*

An avatar of the following lemma can be find in [CLT15, MAR12A, MAR12B, CMI4]. In fact in [CLT15] and [CMI4], the reader can find a proof without the second hypothesis. We give a proof with this hypothesis for the convenience of the reader.

Lemma 4.9. *Let Γ be a discrete subgroup of $SL_{d+1}(\mathbb{R})$ preserving a properly convex open set Ω of \mathbb{S}^d . Let p be a point of $\partial\Omega$ and let $\Gamma_p = \{\gamma \in \Gamma \mid \gamma(p) = p\}$. Suppose that*

- (1) *the subgroup Γ_p acts cocompactly on an affine chart \mathbb{A}^{d-1} of \mathbb{S}_p^{d-1} and*
- (2) *the action of Γ_p on \mathbb{A}^{d-1} is by euclidean transformation.*

Then Ω admits two ellipsoids of security at p which are preserved by Γ_p .

It will be convenient for the proof to call the boundary of an ellipsoid an *ellisphere*.

Proof. The action of Γ_p on \mathbb{A}^{d-1} is by euclidean transformation, therefore the action of Γ_p on \mathbb{R}^{d+1} preserves a quadratic form of signature $(d, 1)$; in other words preserves an ellipsoid \mathcal{E} such that $p \in \partial\mathcal{E}$. Also, there exists a convex compact fundamental domain D for the action of Γ_p on the affine chart \mathbb{A}^{d-1} . We denote by \mathcal{C}_p the cone of vertex p and basis D (see Figure 8).




Figure 8. The action of a parabolic subgroup.

Since, Γ_p acts cocompactly on an affine chart \mathbb{A}^{d-1} , we get that $\mathcal{D}_p(\Omega) = \mathbb{A}^{d-1}$, hence Ω has a unique supporting hyperplane H_p at p . The pencil \mathcal{F} of ellipsoids generated by $\partial\mathcal{E}$ and H_p gives a one parameter family of ellipsoids preserved by Γ_p . Moreover the intersection of any ellipsoid of the pencil \mathcal{F} with \mathcal{C}_p is compact since Γ_p acts cocompactly on $\mathcal{D}_p(\Omega)$.

Therefore to find an ellipsoid \mathcal{E}^{int} (resp. \mathcal{E}^{ext}) inside (resp. outside) Ω preserved by Γ_p , it is sufficient to see that any ellipsoid $\partial\mathcal{E}'$ of the pencil \mathcal{F} which is sufficiently close (resp. far) from p verifies $\mathcal{E}' \cap \mathcal{C}_p \subset \Omega$ (resp. $\Omega \cap \mathcal{C}_p \subset \mathcal{E}'$).

Hence, Ω admits two ellipsoids of security at p which are preserved by Γ_p . Thanks to Proposition 4.6, the point p is round. □

Proof of Proposition 4.7. The point p is parabolic, so the Coxeter polytope P_p is perfect, preserves an affine chart of \mathbb{S}_p^{d-1} and acts compactly by euclidean transformations on it by Lemma 4.8. Hence, Lemma 4.9 shows the second point.

Proposition 4.6 shows that the second point implies the first and third one. The last point is a trivial consequence of the fact that for any infinite order element γ of Γ_p and for all point $x \in \underline{\Omega}_p$, we have $\gamma^n(x) \rightarrow p$. □

4.4. A lemma about negative type Coxeter polytope

Lemma 4.10. *Let P be Coxeter polyhedron of \mathbb{S}^d . If P is of negative type then there exists an affine chart of \mathbb{S}^d containing P and all its polars.*

Proof. We can assume that P is indecomposable. By Perron-Frobenius theorem and the definition of being of negative type there exists a strictly positive vector $\mu \in (\mathbb{R}_+)^S$ and a real $\lambda < 0$ such that $A_P \mu = \lambda \mu$. So, if we take $\alpha = \sum_{s \in S} \mu_s \alpha_s$, then for each $t \in S$, we get that $\alpha(v_t) = \lambda \mu_t < 0$. This implies that the affine chart $\mathbb{A} := \{x \in \mathbb{S} \mid \alpha(x) < 0\}$ contains all the polar of P in its closure. Moreover, since $P = \{x \in P \mid \text{for all } s \in S, \alpha_s(x) \leq 0\}$, we get that $P \subset \mathbb{A}$. \square

4.5. Truncability

4.5.1. Definition of truncability

Definition 4.11. Let p be a vertex of a Coxeter polytope P of \mathbb{S}^d and S_p the set of facets of P containing p . The vertex p of a Coxeter polytope is *truncable* when the projective subspace Π_p spanned by the $[v_s]$ for $s \in S_p$:

- (1) is a hyperplane,
- (2) meets the interior of P ,
- (3) a ridge e of P verifies $e \cap \Pi_p \neq \emptyset$ if and only if $p \in e$ and $e \cap \Pi_p$ have to be included in the relative interior of e .

We will denote by Π_p^+ (resp. Π_p^-) the connected component of $\mathbb{S}^d \setminus \Pi_p$ which does not contain p (resp. containing p). We stress that Π_p^+ and Π_p^- are affine charts. See Figure 9.

Remark 4.12 (consequence). Suppose P is a Coxeter polytope and p is a truncable vertex. We can define a new polytope $P^{\dagger p}$. The facets of $P^{\dagger p}$ are the facets of P (we call them the *old facets*) plus one extra facet defined by the hyperplane Π_p (called the *new facet*). The polar of the old facets are unchanged and the polar of the new facet is p . Therefore, it is easy to check that the polytope $P^{\dagger p}$ has the following property:

- the dihedral angles of the new ridges are $\frac{\pi}{2}$;
- the vertices of the new facet are all elliptic if and only if P_p is perfect;
- the hyperplane Π_p is preserved by the reflection across the facets containing p . The intersection $P \cap \Pi_p$ is a Coxeter polytope of Π_p isomorphic to P_p .




Figure 9. Illustration of Truncability of p .

Remark 4.13. This construction was already known in the hyperbolic space. See for example the survey [VIN85] of Vinberg, Proposition 4.4. This construction in the projective context and in dimension 3 is the main ingredient of [MAR10].

4.5.2. Simple perfect loxodromic vertex are truncable

Proposition 4.14. *Let P be a Coxeter polytope of \mathbb{S}^d and p be a vertex of P . If the vertex p is simple perfect and loxodromic then P is truncable at p except if P is isomorphic to $P_p \otimes \cdot$.*

Proof. Thanks to Lemma 4.10, we can think of everything inside an affine chart that contains P and its polars. The simplicity (resp. loxodromicness) of p implies that the projective space Π_p is of dimension at most (resp. at least since the rank of A_p is d) $d - 1$.

We first look at the half-line $]pv_i)$, for $i \in S_p$. Lemma 4.10 applied to P_p gives the existence of a hyperplane H_p of \mathbb{S}^d that contains p and such that for each $i \in S_p$, the open segment $]pv_i[$ is included in the connected component H_p^- of $\mathbb{S}^d \setminus H_p$ that contains the interior of P .

Next, we look at the position of this half-line around Ω_P . Lemma 4.15 shows that Ω_p (and so P_p also) is included in the convex hull of the $S_p(v_i)$ for $i \in S_p$. Hence, the convex set Ω_P is included in the convex hull of this half-lines. Roughly speaking, the v_i for $i \in S_p$ are “all around” P and Ω_P .

Finally, we show that the v_i , for $i \in S_p$ cannot be too “far” from p . Let F_t^- denote the component of $S^d \setminus F_t$ that contains p , where F_t is the hyperplane generated by the facet $t \notin S_p$. Geometrically, the inequalities in (C) mean that $v_i \in F_t^-$ except when the angle between the face i and t is $\frac{\pi}{2}$; in that case, $v_i \in F_t$. So we know that for each facet $t \notin S_p$ and for each $i \in S_p$ the point v_i is on the segment $[pv_i) \cap \overline{F_t^-}$.

So $\Pi_p \cap P \neq \emptyset$ and $p \notin \Pi_p$. For the sake of clarity, we need to distinguish two cases: a) P is not a cone of summit p or b) it is not. If we are in the case b), the inequalities in (C) (using the several facets of P not in S_p) show that any facet f of P which intersects Π_p has to contain p , and the intersection $f \cap \Pi_p$ is included in the relative interior of f . Now, if we are in case a) then the inequalities in (C) show that either Π_p is the support of the face of P not in S_p , in which case P is isomorphic to $P_p \otimes \cdot$, or any facet f of P which intersects Π_p has to contain p , and the intersection $f \cap \Pi_p$ is included in the relative interior of f . \square

Lemma 4.15 (Nie, Lemma 3 of [Nie15]). *Let P be a perfect loxodromic simplex. The convex set Ω_P is included in the convex hull of its polar.*

Remark 4.16. A more careful analysis of the situation would show that if P is an indecomposable Coxeter polytope of S^d and p is a simple perfect vertex of P , then, p is elliptic if and only if $\Omega_P \cap \Pi_p = \emptyset$, p is parabolic if and only if $\Omega_P \cap \Pi_p = \{p\}$ and p is loxodromic if and only if p is truncable.

4.5.3. Iteration of truncation

Lemma 4.17. *Let P be a loxodromic Coxeter polytope and p, q two vertices of P . Suppose that p and q are perfect simple loxodromic vertices, denote by f_p (resp. f_q) the facets obtained by truncation of P at p (resp. q), then the facets f_p and f_q do not meet.*

Proof. Let π_p (resp. π_q) be the intersection of the projective space spanned by f_p (resp. f_q) and $\overline{\Omega_P}$. Since $f_p \subset \pi_p$ and $f_q \subset \pi_q$, this lemma is a consequence of the fact that $\pi_p \cap \pi_q$ is included in $\partial\Omega_P$. Since p is perfect, the f_p is included in the relative interior of π_p (by Corollary 2.3).

Let us now prove this fact (Figure 10 can be useful). Choose an affine chart \mathbb{A} containing $\overline{\Omega_P}$, denote by \mathcal{C}_p (resp. \mathcal{C}_q) the cone of summit p (resp. q) and basis π_p (resp. π_q) and by $\widehat{\mathcal{C}}_p$ (resp. $\widehat{\mathcal{C}}_q$) the cone of summit p generated by π_p (resp. π_q) in the affine chart \mathbb{A} . We remark that $\overline{\Omega_P}$ contains the cones \mathcal{C}_p and \mathcal{C}_q and is contained in $\widehat{\mathcal{C}}_p \cap \widehat{\mathcal{C}}_q$. Since Ω_P is convex, this is possible only when $\pi_p \cap \pi_q$ is included in $\partial\Omega_P$. \square




Figure 10. A possible situation and an impossible situation.

The last lemma shows that given a loxodromic Coxeter polytope, if we denote by \mathcal{L}^{sp} the set of simple perfect loxodromic vertices, then we can define a new Coxeter polytope P^\dagger which is obtained from P by doing the truncation around every vertex $p \in \mathcal{L}^{sp}$. We will call it the *truncated Coxeter polytope* of P and we will use the notation P^\dagger to represent it. We will call *old* (resp. *new*) the vertices, edges, facets, ridges of P^\dagger that were (resp. were not) in P . Figure 11 illustrates the situation.

The following lemma gives the main properties of P^\dagger . To stay it, the notion of (Γ, Γ') -precisely invariant region is useful. If Γ acts on Ω and Γ' is a subgroup of Γ then a subset A of Ω is (Γ, Γ') -precisely invariant when for every $\gamma \in \Gamma \setminus \Gamma'$, we have $\gamma(A) \cap A = \emptyset$ and for every $\gamma \in \Gamma'$, we have $\gamma(A) = A$.




Figure 11. The starting of the tiling given here is obtained thanks to a square with three right angles and one loxodromic vertex: p . The convex set Ω_P is the union of the convex set $\Omega_{P^\dagger p}$ and the Γ_P -orbits of the hatching zone. The limit set is the boundary of $\Omega_{P^\dagger p}$ minus the interior of the Γ_P -orbits of the hatching zone.

Lemma 4.18. *Let P be an irreducible 2-perfect loxodromic Coxeter polytope whose loxodromic vertices are simple. Consider the truncated Coxeter polytope P^\dagger of P . For each loxodromic vertex p , we denote by \mathcal{C}_p the cone of summit p and basis the intersection of Ω_P with the hyperplane generated by the polars of the facets containing p . We have the following:*

- (1) $P^\dagger \subset P$ and $\Gamma_P \subset \Gamma_{P^\dagger}$;
- (2) for every loxodromic vertex p of P , the cone \mathcal{C}_p is (Γ_P, Γ_p) -precisely invariant;
- (3) $P \cap \bar{C}(\Lambda_P) = P^\dagger$;
- (4) $\Omega_{P^\dagger} \subset \Omega_P$;
- (5) the Coxeter polytope P^\dagger is a quasi-perfect Coxeter polytope.

Proof. The first statement is trivial. The second statement is a consequence of the fact that the action of Γ_P on the d -cell of the tiling of Ω_P is free. The main interest of the second statement is that we found a (Γ_P, Γ_p) -precisely-invariant region C_p which is convex and such $\Omega_P \setminus C_p$ is also convex. Hence the set $\Omega' = \bigcup_{\gamma \in \Gamma_P} \gamma(P^\dagger)$ is convex and Γ_P -invariant. Hence $\Lambda_P \subset \partial\Omega'$, and so $C(\Lambda_P) \subset \Omega'$. Moreover, the limit set of Γ_p is included in the intersection π_p of Ω_P with the hyperplane generated by the polars of the facets containing p . So $P \cap \bar{C}(\Lambda_P) = P^\dagger$.

By definition, Ω_{P^\dagger} is the union of the orbits of P^\dagger under Γ_{P^\dagger} . Let p be a loxodromic vertex and let f be the new facet associated to the truncation of p . The set $\Omega_P \setminus \pi_p$ has two connected components, the cone \mathcal{C}_p , and a convex set Ω^+ which contains the interior of P^\dagger and satisfies $\sigma_f(\Omega^+) \subset \mathcal{C}_p \subset \Omega_P$. So we have $\Omega_{P^\dagger} \subset \Omega_P$.

The last point is trivial, the truncation eliminates all the old loxodromic vertices. Moreover, since P is 2-perfect the truncation process creates only elliptic vertices, so the Coxeter polytope P^\dagger has only elliptic and parabolic vertices. \square

4.6. Consequence of loxodromicness for non-simple vertices. Before starting the proof, we make an important remark.

Remark 4.19 (structure of the tiling). The tiling given by Coxeter group has a special feature, roughly speaking, “face extend to subspace.” More precisely let P be a Coxeter polytope. The union $\bigcup_{\gamma \in \Gamma_P} \gamma(\partial P)$ is contained in a union of hyperplanes, in other words, every facet of the tiling extends to a hyperplane of the tiling. Even better, the k -skeleton of the tiling is a union of k -subspaces of $\overline{\Omega_P}$ (i.e. intersections of k -planes with $\overline{\Omega_P}$).

Proposition 4.20. *Let P be an irreducible loxodromic Coxeter polytope of \mathbb{S}^d and p a vertex of P . If the vertex p is perfect loxodromic then $p \notin \Lambda_P$.*

Proof. Suppose that $p \in \Lambda_P$; then there exists a sequence of distinct elements $\gamma_n \in \Gamma_P \setminus \Gamma_p$ such that $q_n := \gamma_n(p) \rightarrow p$. We choose an affine chart \mathbb{A} containing $\overline{\Omega_P} = \overline{\Omega}$. Let K_p be the cone of summit p generated by P in \mathbb{A} intersected with Ω . Define $K_{q_n} := \gamma_n(K_p)$ and $Q_n = \gamma_n(P)$. Since $\Sigma = \bigcup_{\gamma \in \Gamma_P} \gamma(\partial P \cap \Omega)$ is contained ∂K_p , we must have $p \in K_{q_n}$ for n big enough. We claim that $p \in \partial K_{q_n}$ for n big enough. Indeed, if not, then, $\mathcal{D}_p(\Omega) = \lim_n \mathcal{D}_{q_n}(\Omega)$ is an affine chart contradicting the fact that p is loxodromic. By symmetry, we get that $q_n \in \partial K_p$ for n big enough. Hence, q_n is on the hyperplane generated by a facet of P for n big enough, this is in contradiction with the fact that q_n converges to p . \square

Remark 4.21. Choi proves a similar statement for the action of a discrete group Γ on a properly convex open set Ω with the hypothesis that Γ_p is Gromov-hyperbolic and also a technical condition on the eigenvalue of Γ_p (See Theorem 6.4 of [Choi13]). Here we do not assume Γ_p Gromov-hyperbolic but we assume we are in a “Coxeter situation.”

The following definition is ad hoc but it will be useful. A *nicely embedded cone* \mathcal{C} in a properly convex open set Ω is a properly convex open cone \mathcal{C} such that $\mathcal{C} \subset \Omega$ and $\partial \mathcal{C} \cap \Omega$ is the relative interior \mathcal{B} of the basis of \mathcal{C} . Hence, we have $\partial \mathcal{C} \setminus \mathcal{B} \subset \partial \Omega$.

Remark 4.22. Let P be a Coxeter polytope and p a vertex of P . When p is truncable there is a *canonical* properly embedded cone which is (Γ_P, Γ_p) -precisely invariant: $\mathcal{C}_p = \Pi_p^- \cap \Omega_P$.

Corollary 4.23. Let P be an irreducible loxodromic Coxeter polytope of \mathbb{S}^d and p be a vertex of P . If the vertex p is perfect loxodromic then there exists a properly embedded cone which is (Γ_P, Γ_p) -precisely invariant.

Proof. Since $p \notin \Lambda_P$, we get that $p \notin \bar{C}(\Lambda_P)$, so one can choose an hyperplane H such that $H \cap \Omega_P \neq \emptyset$ and one connected component $\mathbb{S}^d \setminus H$ contains $\bar{C}(\Lambda_P)$ while the other H^- contains p . The cone $\mathcal{C}_p = H^- \cap \Omega_P$ does the job. \square

Proposition 4.24. Let P be an irreducible loxodromic Coxeter polytope of \mathbb{S}^d and \mathcal{L} be set of perfect loxodromic vertices of P . For each $p \in \mathcal{L}$, we choose a nicely embedded cone \mathcal{C}_p which is (Γ_P, Γ_p) -precisely invariant and let $\Omega' = \Omega_P \setminus \bigcup_{p \in \mathcal{L}} \Gamma_P(\bar{\mathcal{C}})$. Then

- (1) the open set Ω' is a Γ_P -invariant properly convex;
- (2) $C(\Lambda_P) \subset \Omega'$;
- (3) for every $p \in \mathcal{L}$, the point $p \in \partial\Omega_P$ is neither strictly convex nor with \mathcal{C}^1 boundary;
- (4) the point p is an extremal point of $\partial\Omega_P$;
- (5) for every neighbourhood \mathcal{U} of p in \mathbb{S}^d we have $\mu_{\Omega_P}(\mathcal{U} \cap P) = \infty$.

Proof. The existence of such a \mathcal{C}_p is a consequence of Propositions 4.14 and 4.20. The first, third and fourth points are direct consequences of the (Γ_P, Γ_p) -precise invariance of the nicely embedded cone \mathcal{C}_p . The second point is a consequence of the fact that Λ_P is the smallest closed subset of \mathbb{P}^d that is Γ_P -invariant. For the last point, since $\mathcal{D}_p(\Omega_P)$ is properly convex, we can find a cone ω_p of summit p that contains Ω_P and such that $\partial\Omega_P \cap \partial\omega_p = \{p\}$. Proposition 1.1 shows that $\mu_{\omega_p}(P) \leq \mu_{\Omega_P}(P)$ and Lemma 4.25 below shows that $\mu_{\omega_p}(P) = \infty$. \square

Lemma 4.25. Let Ω be a properly convex open set. Suppose that Ω is a cone. Let p be the summit of Ω and P a convex subset of Ω such that $\bar{P} \cap \partial\Omega = \{p\}$; then $\mu_\Omega(P) = \infty$.

Proof of Lemma 4.25. Consider the affine chart \mathbb{A} whose hyperplane at infinity is the hyperplane generated by the basis of Ω . The automorphism group of Ω contains the homothety h of the affine chart \mathbb{A} of ratio $\frac{1}{2}$ fixing p and $h(P) \subset P$. Of course, as h is an automorphism of Ω , we have $\mu_\Omega(h(P)) = \mu_\Omega(P)$, it follows that $\mu_\Omega(P) = \infty$. \square

5. Degenerate 2-perfect Coxeter polytopes

The reader has probably noticed that the quadrotomy of Theorem 3.5 is very useful. Hence, we believe that the study of the similar question for 2-perfect Coxeter polytopes can be useful even if we will not use it.

Proposition 5.1. *Let P be a 2-perfect Coxeter polytope of \mathbb{S}^d . Then one of the following assertions is true:*

- (1) P is elliptic;
- (2) P is parabolic;
- (3) P is loxodromic and irreducible;
- (4) P is decomposable; in fact $P = Q \otimes \cdot$ where Q is parabolic or loxodromic perfect.

Remark 5.2. So, a loxodromic 2-perfect Coxeter polyhedron has to be irreducible.

Proof. Consider the Cartan matrix A_P of P ; we will distinguish four cases:

- (1) $\text{Rank}(A_P) = d + 1$ and W_P is irreducible;
- (2) $\text{Rank}(A_P) = d + 1$ and W_P is not irreducible;
- (3) P has a loxodromic vertex;
- (4) $\text{Rank}(A_P) < d + 1$ and no loxodromic vertices.

Suppose we are in the first case; then A_P is either of positive type or of negative type, hence P is either elliptic or irreducible loxodromic. Suppose we are in the second case; since $\text{Rank}(A_P) = d + 1$, P is decomposable by Theorem 2.12 and Lemma 5.3 takes care of this case.

Suppose we are in the third case. Let p be a loxodromic vertex of P . Consider the projective space π_p spanned by the polar $[v_s]$ for $s \in S_p$. If $\Pi_p = \mathbb{S}^d$ or p is not simple then we must have $\text{Rank}(A_P) = d + 1$ and we are back to the previous case. If not, then Π_p is a hyperplane and p is simple. A) If P is indecomposable then Proposition 4.14 shows that P is truncable at p , hence there exists a polar of P not in Π_p and so $\text{Rank}(A_P) = d + 1$, and we are back to the previous case again. B) If P is decomposable then Lemma 5.3 takes care of this case.

Suppose we are in the fourth case. Since P is 2-perfect, it has only elliptic or parabolic vertices then Lemma 5.4 of Vinberg concludes. \square

This lemma is a direct adaptation of Vinberg's analogous lemma for the proof of Theorem 3.5.

Lemma 5.3. *Let P be a 2-perfect Coxeter polytope of \mathbb{S}^d . Suppose that P is the product $P_1 \otimes P_2$ of two Coxeter polytopes P_1 and P_2 , then either*

- (1) *both are elliptic or*
- (2) *one is parabolic and the other one is the point Coxeter polytope or*
- (3) *one is loxodromic and the other one is the point Coxeter polytope.*

Proof. Suppose P_1 is not elliptic. A vertex p of P_2 defines a vertex \tilde{p} of $P_1 \otimes P_2$ and the link $P_{\tilde{p}}$ of $P = P_1 \otimes P_2$ at \tilde{p} verifies $P_{\tilde{p}} = P_1 \otimes P_{2p}$. The Coxeter polytope $P_{\tilde{p}} = P_1 \otimes P_{2p}$ is perfect hence elliptic, parabolic or loxodromic (Theorem 3.5). The first case is impossible since P_1 is not elliptic.

So $P_{\tilde{p}}$ is perfect but not elliptic. Then by Theorem 3.5, $P_{\tilde{p}}$ is indecomposable, hence $P_{2p} = \emptyset$, which means that P_2 is a point and $P = P_1 \otimes \cdot$. \square

Lemma 5.4 (Vinberg [VIN71], proof of Proposition 26). *Let P be a Coxeter polytope such that $\text{Rank}(A_P) < d + 1$.*

- (1) *If P has an elliptic vertex then P is either parabolic or decomposable.*
- (2) *If P has a parabolic vertex then P is parabolic or $P = Q \otimes \cdot$ where Q is a parabolic.*

6. Geometry of the action

In this part, we prove Theorem A.

6.1. Cocompact action. We rephrase Corollary 2.3 in our language to get used to it.

Theorem 6.1 (Vinberg [VIN71]). *Let P be a Coxeter polytope. The action of Γ_P on Ω_P is cocompact if and only if all the vertices of P are elliptic (i.e. P is perfect).*

6.2. Geometrically finite action

Theorem 6.2. *Let P be a loxodromic 2-perfect Coxeter polytope. Then we always have*

$$\mu_{\Omega_P}(C(\Lambda_P) \cap P) < \infty.$$

In other word, the action of Γ_P on Ω_P is always geometrically finite.

Proof. Let \mathcal{L} be the set of loxodromic vertices of P . Proposition 4.24 shows that for each vertex $p \in \mathcal{L}$, one can find a (Γ_P, Γ_p) -precisely invariant nicely embedded cone \mathcal{C}_p . One can suppose these cones disjoint by taking them smaller. By removing the Γ_P -orbits of all the \mathcal{C}_p , for $p \in \mathcal{L}$, one obtains a Γ_P -invariant

properly convex open set Ω' such that $P \cap \partial\Omega'$ is exactly the set of parabolic points of P . Now, Proposition 4.7 shows that there exists a neighbourhood \mathcal{U}_p of p in \mathbb{S}^d such that $\mu_{\Omega_P}(P \cap \mathcal{U}_p) < \infty$. Since P has only a finite number of vertices, we get that $\mu_{\Omega_P}(P \cap \Omega') < \infty$. Since $C(\Lambda_P) \subset \Omega'$, we have $\mu_{\Omega_P}(C(\Lambda_P) \cap P) < \infty$. Hence, the action of Γ_P on Ω_P is geometrically finite. \square

6.3. Finite volume case

Theorem 6.3. *Let P be a loxodromic 2-perfect Coxeter polytope. The action of Γ_P on Ω_P is of finite covolume if and only if all the vertices of P are elliptic or parabolic (i.e. P is quasi-perfect).*

Proof. Suppose the action of Γ_P on Ω_P is of finite covolume. Assume one of the vertices p of P is loxodromic. Then the fifth point of Proposition 4.24 shows that $\mu_{\Omega_P}(P) = \infty$. This is absurd, so every vertex of P is either elliptic or parabolic.

Suppose all the vertices of P are elliptic or parabolic. We know from Theorem 6.2 that the action is geometrically finite, but since there is no loxodromic vertices, we have $\Omega' = \Omega_P$ in the proof of 6.2 and we get that $\mu_{\Omega_P}(P) < \infty$. \square

6.4. Convex-cocompact action

Theorem 6.4. *Let P be a loxodromic 2-perfect Coxeter polytope. The action of Γ_P on Ω_P is convex-cocompact if and only if all the vertices of P are elliptic or loxodromic.*

The following corollary is immediate, thanks to Proposition 4.14.

Corollary 6.5. *Let P be a loxodromic 2-perfect Coxeter polytope whose loxodromic vertices are simple. Then, the action of Γ_P on Ω_P is convex-cocompact if and only if the truncated Coxeter polytope P^\dagger of P is perfect.*

Proof of Theorem 6.4. Suppose the action of Γ_P on Ω_P is convex-cocompact. Let p be a vertex of P . We claim that $p \notin \Lambda_P$. Indeed, first $p \in \partial\Omega_P$ if and only if p is not elliptic; second if $p \in \partial\Omega_P$, consider the ray of Ω_P from any point $x_0 \in P$ to p . The projection r of this ray leaves every compact of Ω_P/Γ_P since P is a convex fundamental domain. In particular, the ray r leaves the compact set $\bar{C}(\Lambda_P)/\Gamma_P$, thereby p is not in Λ_P . So, p is not parabolic by Proposition 4.7(4).

Suppose all the vertices of P are elliptic or loxodromic. We know from Theorem 6.2 that the action is geometrically finite, but since there is no parabolic vertices, we get that $P \cap \Omega'$ is bounded in (Ω, d_Ω) in the proof of 6.2 and so the action of Γ_P on Ω_P is convex-cocompact. \square

6.5. Geometrical definition of geometrical finiteness vs the topological one.

In this paragraph, we motivated our definition of geometrical finiteness by comparing it to the definitions in pinched negative curvature and explaining why the definition we choose implies the other classical definitions.

It is classical that if X is a simply connected pinched negatively curved Riemannian manifold (i.e. a Hadamard manifold), then for every irreducible discrete group Γ of isometries of X , the thick part of the convex core is compact if and only if the volume of the convex core is finite and the group Γ is of finite type (thanks to [Bow95]).

When X is a properly convex open subset of \mathbb{P}^d which is strictly convex with \mathcal{C}^1 -boundary then this equivalence remains true ([CM14]). We stress that Margulis's lemma holds in any Hilbert geometry ([CLT15] or [CM13]).

But there is also a topological version of geometrical finiteness. The action of Γ on X is *geometrically finite* if all the points of the limit set of Γ are *conical limit points* or *bounded parabolic fixed points*. See [Bow95] for the definition. We only stress that these definitions are purely topological.

When X is a Hadamard manifold and Γ an irreducible group of isometry of X then the topological definition of geometrically finite action is equivalent to the geometrical definition by [Bow95]. But, when X is a properly convex open subset of \mathbb{P}^d which is strictly convex with \mathcal{C}^1 -boundary, this is no longer true. We only have that the geometrical definition implies the topological one, see [CM14] for the implication and a counterexample in dimension 4.

Maybe even worst, if X is a properly convex open subset of \mathbb{P}^d , which is not supposed strictly convex nor with \mathcal{C}^1 -boundary then *if the volume of the convex core is finite and the group Γ is of finite type then the thick part of the convex core is compact*. But, the author does not know if the converse is true. This implication is just a consequence of the fact that the volume of balls of radius $r > 0$ in Hilbert geometry are bounded from below by a universal constant depending only on the dimension d (thanks to Benzécri's theorem, see [CM14] for the details).

7. Zariski closure of Γ_P

7.1. Notations. Let us introduce some notation for the sake of clarity. We will denote by Trans_d the subgroup of $\text{SL}_{d+1}(\mathbb{R})$ which is the group of translations in the standard affine chart. In term of matrices, it is the group

$$\text{Trans}_d = \left\{ \left(\begin{array}{ccc|c} 1 & & 0 & u_1 \\ & \ddots & & \vdots \\ 0 & & 1 & u_d \\ 0 & \cdots & 0 & 1 \end{array} \right) \middle| (u_1, \dots, u_d) \in \mathbb{R}^d \right\}$$

and by Diag_d the subgroup of $\text{SL}_{d+1}(\mathbb{R})$ of diagonal matrices with positive entries:

$$\text{Diag}_d = \left\{ \begin{pmatrix} \lambda_1 & & 0 \\ & \ddots & \\ 0 & & \lambda_{d+1} \end{pmatrix} \middle| \lambda_1, \dots, \lambda_{d+1} \in \mathbb{R}_+^* \text{ such that } \lambda_1 \cdots \lambda_{d+1} = 1 \right\}.$$

These two groups are isomorphic as abstract groups.

Notation 7.1. If P is a Coxeter polytope, we will denote by G_P the connected component of the Zariski closure of the discrete subgroup Γ_P of $\text{SL}_{d+1}(\mathbb{R})$.

7.2. The perfect case. In the perfect case, all the arguments are in the literature, we just put them together.

7.2.1. The easy case

Theorem 7.2. *Let P be a perfect Coxeter polytope. Let G_P be the connected component of the Zariski closure of Γ_P in $\text{SL}_{d+1}(\mathbb{R})$.*

- (1) *If P is elliptic, then $G_P = \{1\}$.*
- (2) *If P is parabolic, then G_P is conjugate to the group Trans_d .*
- (3) *If P is loxodromic and W_P is affine, then Ω_P is a simplex, the Coxeter group W_P is affine irreducible of type \tilde{A}_n and the group G_P is conjugate to Diag_d .*

Proof. In the first case, the group Γ_P is finite so $G_P = \{1\}$. In the second case, Γ_P is a lattice of a conjugate of $\text{Trans}_d \rtimes \text{SO}_d$ and the image of Γ_P in SO_d is finite, therefore G_P is conjugate to Trans_d . In the third case, by Proposition 2.9 (4), Γ_P is a lattice of $\text{Aut}(\Omega_P) = \text{Diag}_d \rtimes \mathfrak{S}_{d+1}$, the conclusion follows, where \mathfrak{S}_{d+1} is the symmetric group on $\{1, \dots, d + 1\}$ acting canonically on \mathbb{R}^{d+1} . \square

7.2.2. The interesting case

Remark 7.3. From Theorem 3.5, we learn that if P is a perfect polytope then P is either elliptic, parabolic, loxodromic with W_P affine irreducible or loxodromic with W_P large irreducible.

Theorem 7.4 (Benoist + folklore). *Let P be a perfect loxodromic Coxeter polytope with W_P large irreducible. Then we have the following alternative:*

- Ω_P is an ellipsoid and G_P is conjugate to $\text{SO}_{d,1}^\circ(\mathbb{R})$ or
- Ω_P is not an ellipsoid and $G_P = \text{SL}_{d+1}(\mathbb{R})$.

Proof. First, from Theorem 2.18, we know that Γ_P is strongly irreducible and so Ω_P is indecomposable. We need to distinguish three cases: Ω_P is an ellipsoid, Ω_P is symmetric¹⁹ but not an ellipsoid, or Ω_P is not symmetric.

If Ω_P is an ellipsoid, then $\text{Aut}(\Omega_P)$ is conjugate to $\text{SO}_{d,1}^\circ(\mathbb{R})$ and Γ_P is a cocompact lattice of $\text{Aut}(\Omega_P)$. The conclusion follows from Borel density Theorem 7.5.

Assume Ω_P is symmetric but not an ellipsoid. Then $\text{Aut}(\Omega_P)$ has property (T), from Theorem 7.6 and Theorem 7.7 below. So, Γ_P has property (T)²⁰ too since it is a lattice of $\text{Aut}(\Omega_P)$ by Theorem 7.8. But an infinite Coxeter group does not have property (T) by Theorem 7.9. So this case is absurd.

If Ω_P is not symmetric, then $G_P = \text{SL}_{d+1}(\mathbb{R})$ by Theorem 7.10. □

Theorem 7.5 (Borel density theorem, [BOR60]). *A lattice of a semi-simple lie group without compact factor is Zariski-dense.*

Theorem 7.6 (Koecher and Vinberg, see [VIN63], [FK94], or [KOE99]). *Let Ω be an indecomposable symmetric properly convex open subset of \mathbb{P}^d . Then Ω is the symmetric space associated to $\text{SO}_{d,1}^\circ(\mathbb{R})$ or $\text{SL}_m(\mathbb{K})$ where $\mathbb{K} = \mathbb{R}, \mathbb{C}, \mathbb{H}$ and $m \geq 3$ or to the exceptional group $E_{6(-26)}$. In particular, the automorphism group of an indecomposable symmetric properly convex open set which is not an ellipsoid is a quasi-simple²¹ Lie group of real rank²² at least two.*

Theorem 7.7 (Kazhdan, Delaroché, and Kirillov, Vasertein, and Wang, see Theorem 1.6.1 in [BHV08]). *A quasi-simple Lie group of real rank at least two has property (T).*

Theorem 7.8 (Kazhdan, see Theorem 1.7.1 of [BHV08]). *A lattice Γ of a locally compact group G has property (T) if and only if G has property (T).*

Theorem 7.9 (Bozejko, Januszkiewicz, and Spatzier [BJS88]). *Let W be a Coxeter group. If W has property (T) then, W is finite.*

Theorem 7.10 (Benoist [BEN03]). *Let Γ be a discrete group of $\text{SL}_{d+1}(\mathbb{R})$ acting cocompactly on a properly convex open set Ω . If the group Γ is strongly irreducible and Ω is not symmetric, then Γ is Zariski-dense in $\text{SL}_{d+1}(\mathbb{R})$.*

¹⁹ A properly convex open set is *symmetric* if for every point $x \in \Omega$ there exists an isometry γ of (Ω, d_Ω) which fixes x and whose differential at x is $-\text{Id}$.

²⁰ We don't give the definition of property (T), since we don't need the definition for our purpose. The reader is referred to the book [BHV08] for a definition and all the proof of the theorem we will use in the sequel.

²¹ A Lie group is *quasi-simple* when its Lie algebra is simple or equivalently when all its normal subgroups are discrete.

²² The real rank of a semi-simple Lie group is the common dimension of all the maximal splitted connected abelian subgroups, e.g maximal splitted tori.

7.3. Non-degenerate 2-perfect case

Theorem 7.11. *Let P be a loxodromic 2-perfect Coxeter polytope of \mathbb{S}^d which is not perfect. Then either G_P is conjugate to $\mathrm{SO}_{d,1}^{\circ}(\mathbb{R})$, or $G_P = \mathrm{SL}_{d+1}(\mathbb{R})$.*

A nice corollary is the following.

Corollary 7.12. *Let P be a loxodromic 2-perfect Coxeter polytope of \mathbb{S}^d which is not perfect and whose loxodromic vertices are truncable. Let P^{\dagger} be the truncated Coxeter polytope associated to P then either $\Omega_{P^{\dagger}}$ is an ellipsoid, or $G_P = \mathrm{SL}_{d+1}(\mathbb{R})$. In particular, in both cases, $G_P = G_{P^{\dagger}}$.*

We will show the Theorem 7.11 and the Corollary 7.12 at paragraph 7.6.

7.4. Proximality, limit sets and Zariski closure. The following paragraph presents basic facts about proximal action on the projective space. We have included the facts we will need and some arguments to make the reading easier to the reader not familiar to the theory. We have tried to give references when we thought an argument would divert the reader's attention or be too long. This paragraph has nothing original, we borrow a lot from [AMS95, GG96, BEN00].

7.4.1. Proximal elements and proximal subgroups. An element γ of $\mathrm{SL}_{d+1}(\mathbb{R})$ is *proximal* when the eigenvalue of maximal modulus is a simple eigenvalue. In that case, the eigenvalue of maximal modulus has to be real, it has to be the spectral radius ρ or its opposite $-\rho$. The corresponding eigenspace is a line, so a point x_{γ}^{+} of \mathbb{P}^d . This point is called the *attractive fixed point* of γ . Indeed, it is easy to see that outside a projective hyperplane H , for every point $x \in \mathbb{P}^d \setminus H$, we have $\gamma^n(x) \rightarrow x_{\gamma}^{+}$ when $n \rightarrow +\infty$.

A subgroup G of \mathbb{P}^d is *proximal* when it contains a proximal element.

7.4.2. Proximal action. The action of a group G on \mathbb{P}^d is *proximal* when for every two points $x, y \in \mathbb{P}^d$ there exists a sequence $(g_k)_{k \in \mathbb{N}}$ in G such that the sequences $(g_k(x))_{k \in \mathbb{N}}$ and $(g_k(y))_{k \in \mathbb{N}}$ converge to the same point.

The link between the notion of proximal group and proximal action is given by the following equivalence. If G is a subgroup of $\mathrm{SL}_{d+1}(\mathbb{R})$ then “ G is irreducible and the action of G on \mathbb{P}^d is proximal” if and only if “the group G is strongly irreducible and proximal” (Theorem 2.9 of [GG96]).

7.4.3. Limit set. Suppose G is strongly irreducible and proximal. Then one can show that the closure Λ_G of all the attractive fixed points of all the proximal elements of G is the smallest²³ closed G -invariant subset of \mathbb{P}^d (see Theorem 2.3 of [GG96]). So in particular, the action of G on Λ_G is minimal.²⁴ This closed subset Λ_G is called the *limit set* of G .

²³ Every closed G -invariant subset contains Λ_G .

²⁴ Every orbit is dense.

7.4.4. The case of an algebraic group. If we assume moreover that G is a Zariski closed subgroup of $\mathrm{SL}_{d+1}(\mathbb{R})$. Then Λ_G is the unique closed orbit of the action of G on \mathbb{P}^d . In fact, Λ_G is even Zariski closed. In particular, Λ_G is a smooth algebraic sub-manifold of \mathbb{P}^d . This is due to the following fact.

The action of a Zariski closed subgroup G of $\mathrm{SL}_{d+1}(\mathbb{R})$ on \mathbb{P}^d is algebraic, so in particular every orbit is locally closed for the Zariski topology, i.e. every orbit is Zariski-open in its Zariski-closure. First, the limit set is closed for the Zariski topology. Indeed, take a point $x \in \Lambda_G$, the orbit $G \cdot x$ is open in its Zariski closure $\overline{G \cdot x}^{\mathrm{Zar}}$, hence $\overline{G \cdot x}^{\mathrm{Zar}} \setminus G \cdot x$ is Zariski closed, hence closed also in the usual sense. But Λ_G is the smallest closed invariant set, hence $G \cdot x = \Lambda_G$ and is Zariski closed. The fact that the orbit of a point outside Λ_G is not closed is a consequence of the definition of Λ_G . Finally, the limit set Λ_G is a smooth algebraic manifold since there exists a transitive action on it.

7.4.5. The point of view of semi-simple group's representation theory. Even better, since G is a Zariski closed subgroup of $\mathrm{SL}_{d+1}(\mathbb{R})$, it is a Lie group. Let G_0 be the connected component of G . Since the action of G on \mathbb{R}^{d+1} is strongly irreducible, the action of G_0 on \mathbb{R}^{d+1} is also strongly irreducible (an algebraic variety can only have a finite number of connected components, so the index of G_0 in G is finite). Much better, the Lie group G_0 is semi-simple since the group G is proximal.²⁵

Hence, the representation $\rho_0: G_0 \rightarrow \mathrm{SL}_{d+1}(\mathbb{R})$ is an irreducible representation of the semi-simple group G_0 . Let $KAN = G_0$ be an Iwasawa decomposition of G_0 where K is a maximal compact subgroup of G , A a maximal abelian connected and diagonalizable over \mathbb{R} subgroup and N a maximal unipotent subgroup.

A representation ρ of a connected semi-simple group with finite center G_0 is proximal when the subspace $\mathrm{Fix}(N) = \{x \in \mathbb{R}^{d+1} \mid \text{for all } n \in N, n(x) = x\}$ is a line. In [AMS95] Abels, Margulis, and Soifer show that an irreducible representation $\rho: G_0 \rightarrow \mathrm{SL}_{d+1}(\mathbb{R})$ is proximal if and only if the group $\rho(G_0)$ is proximal (Theorem 6.3). In particular, the representation ρ_0 is proximal.

Since the subspace $\mathrm{Fix}(N)$ is a line, it is a point x_N of \mathbb{P}^d . The orbit of x_N under the group G_0 is equal to the orbit of x_N under the compact group K (since N is normal in AN), hence it is closed, thereby it is the unique closed orbit of G on \mathbb{P}^d , i.e. the limit set Λ_G .

7.4.6. Zariski closure. This procedure is particularly interesting when one starts with a discrete subgroup Γ of $\mathrm{SL}_{d+1}(\mathbb{R})$. Then one can consider G_0 the connected component of the Zariski closure of Γ . The action of Γ on \mathbb{R}^{d+1} is strongly

²⁵The group G_0 is a reductive group (i.e. its unipotent radical is trivial) because G_0 is irreducible. So, to show that G_0 is semi-simple, one just has to show that the center of G_0 is discrete. Now, any element of the center has to preserve the eigenspaces of all elements of G_0 , in particular the proximal one, hence the center is composed only of a homothety of determinant one.

irreducible if and only if the action of G_0 on \mathbb{R}^{d+1} is irreducible. Moreover, in that case, the action of Γ on \mathbb{P}^d is proximal if and only if the action of G_0 on \mathbb{P}^d is proximal (Theorem 6.3 of [GM89]).

Hence, if one starts with a strongly irreducible and proximal subgroup Γ of $\mathrm{SL}_{d+1}(\mathbb{R})$, this procedure gives a connected semi-simple group with finite center G_0 , an irreducible representation $\rho: G_0 \rightarrow \mathrm{SL}_{d+1}(\mathbb{R})$ and two closed subsets of \mathbb{P}^d : $\Lambda_\Gamma \subset \Lambda_{G_0}$.

7.5. Positive proximality and Zariski closure. In this article, we are interested in discrete subgroups of $\mathrm{SL}_{d+1}(\mathbb{R})$ which preserve a properly convex open subset of \mathbb{P}^d . In this context, the notion of positive proximality is interesting.

7.5.1. Positively proximal element and positively proximal group. A proximal element γ of $\mathrm{SL}_{d+1}(\mathbb{R})$ is *positively proximal* when its spectral radius ρ is an eigenvalue. A proximal subgroup G of $\mathrm{SL}_{d+1}(\mathbb{R})$ is *positively proximal* when all its proximal elements are positively proximal.

A theorem of Benoist makes a bridge between being positive proximal and preserving a properly convex open set. Suppose Γ is strongly irreducible. *Then the group Γ preserves a properly convex open set if and only if the group Γ is positively proximal* (Proposition 1.1 of [BEN00]). In particular, the group Γ is proximal, and the construction explained in the previous paragraph applied.

7.5.2. A key lemma of Benoist

Lemma 7.13 (Benoist [BEN00]). *Let Γ be a strongly irreducible subgroup of $\mathrm{SL}_{d+1}(\mathbb{R})$ preserving a properly convex open subset. The connected component G of the Zariski closure of Γ is a semi-simple Lie group and the action of G on \mathbb{P}^d is proximal. Moreover, we can be more precise in two extremal cases:*

- (1) *if the limit set Λ_G of G is the boundary of a properly convex open subset of \mathbb{P}^d , then Λ_G is an ellipsoid and G is conjugate to $\mathrm{SO}_{d,1}^\circ(\mathbb{R})$;*
- (2) *if $\Lambda_G = \mathbb{P}^d$ then $G = \mathrm{SL}_{d+1}(\mathbb{R})$.*

The following lemma is an easy consequence of Lemma 7.13. We state it to clarify our strategy to find the Zariski closure of Γ_P .

Lemma 7.14. *Let Γ be a strongly irreducible subgroup of $\mathrm{SL}_{d+1}(\mathbb{R})$ preserving a properly convex open set. Let G be the connected component of the Zariski closure of Γ . Suppose there exists a point $x \in \Lambda_G$ and a Zariski closed subgroup H of G such that the orbit $H \cdot x$ is a sub-manifold of \mathbb{P}^d of dimension $d - 1$. Then G is conjugate to $\mathrm{SO}_{d,1}^\circ(\mathbb{R})$ or $G = \mathrm{SL}_{d+1}(\mathbb{R})$.*

7.5.3. A useful remark. The following remark gives a description of the maximal properly convex open set preserved by a strongly irreducible positively proximal discrete subgroup Γ of $SL_{d+1}(\mathbb{R})$.

An element $\gamma \in SL_{d+1}(\mathbb{R})$ is bi-proximal if γ and γ^{-1} are proximal. We introduce the following notation. If γ is a bi-proximal element of $SL_{d+1}(\mathbb{R})$, then γ^+ is the eigenvalue corresponding to the spectral radius, H_γ is the projective subspace spanned by all the eigenvectors except the one corresponding to the smallest (in modulus) eigenvalue and H_γ^+ is the affine chart $\mathbb{P}^d \setminus H_\gamma$. Hence, γ^+ is the unique attractive fixed point of $\gamma \curvearrowright \mathbb{P}^d$ and H_γ is the unique attractive fixed point of $\gamma \curvearrowright \mathbb{P}^{d*}$, where \mathbb{P}^{d*} is the dual of \mathbb{P}^d .

Remark 7.15. The smallest properly convex open set Ω_{\min} preserved by Γ is the convex hull of the limit set Λ_Γ . The largest Ω_{\max} is the dual of the convex hull $\Omega_{\min,*}$ of the limit set $\Lambda_{\Gamma,*}$ of the dual action of Γ on \mathbb{P}^{d*} .

Let $\text{AFP}(\Gamma)$ (resp. $\text{AFP}(\Gamma^*)$) be the set of attractive fixed points of proximal elements of Γ (resp. Γ^*). We know that $\text{AFP}(\Gamma)$ is dense in Λ_Γ , so we get Ω_{\min} is the convex hull of $\text{AFP}(\Gamma)$. Now, since Ω_{\max} is the dual of $\Omega_{\min,*} = C(\text{AFP}(\Gamma^*))$, we get $\Omega_{\max} = \bigcap_{\gamma \in \Gamma^{\text{prox}}} H_\gamma^+$, where Γ^{prox} is the set of bi-proximal elements of Γ .

7.6. Proof of Theorem 7.11

7.6.1. The action of \mathcal{U}_{d-1} on \mathbb{P}^d . We define a subgroup of $SL_{d+1}(\mathbb{R})$:

$$\mathcal{U}_{d-1} = \left\{ \begin{pmatrix} 1 & u_1 & \cdots & u_{d-1} & \frac{1}{2}(u_1^2 + \cdots + u_{d-1}^2) \\ & 1 & & 0 & u_1 \\ & & \ddots & & \vdots \\ 0 & & & 1 & u_{d-1} \\ 0 & \cdots & & 0 & 1 \end{pmatrix} \middle| (u_1, \dots, u_{d-1}) \in \mathbb{R}^{d-1} \right\}.$$

The group \mathcal{U}_{d-1} preserves an ellipsoid \mathcal{E} , fixes a point $p \in \partial\mathcal{E}$ and fixes every horosphere of \mathcal{E} centered at p . In other words, \mathcal{U}_{d-1} is included in the stabilizer of a horosphere in the hyperbolic space $(\mathcal{E}, d_\mathcal{E})$. More precisely, \mathcal{U}_{d-1} is the subgroup composed of the non-screw parabolic elements fixing p of the hyperbolic space $(\mathcal{E}, d_\mathcal{E})$. In particular, \mathcal{U}_{d-1} is isomorphic to \mathbb{R}^{d-1} . Moreover, if x is not in the tangent space to $\partial\mathcal{E}$ at p then the space $\mathcal{U}_{d-1} \cdot x \cup \{p\}$ is an ellisphere.

The following lemma is then a direct corollary of Proposition 4.7:

Lemma 7.16. *Let P be an irreducible loxodromic Coxeter polytope of \mathbb{S}^d and let p be a parabolic vertex of P . Then the connected component G_p of the Zariski closure of Γ_p is conjugate to \mathcal{U}_{d-1} .*




Figure 12. The orbits of the action of U_{d-1} on \mathbb{P}^d .

7.6.2. The action of $SO_{d-1,1}^\circ(\mathbb{R})$ on \mathbb{P}^d . The action of $SO_{d-1,1}^\circ(\mathbb{R})$ on \mathbb{P}^d has seven types of orbits. To see this, one should think of \mathbb{P}^d as the projective space $\mathbb{P}(\mathbb{R}^d \oplus \mathbb{R})$ i.e. the projective completion of \mathbb{R}^d . The action $SO_{d-1,1}^\circ(\mathbb{R})$ on \mathbb{P}^d preserves the hyperplane at infinity H_∞ of the affine chart \mathbb{R}^d of $\mathbb{P}^d = \mathbb{P}(\mathbb{R}^d \oplus \mathbb{R})$. The action of $SO_{d-1,1}^\circ(\mathbb{R})$ on $H_\infty = \mathbb{P}^{d-1}$ has three orbits, the limit set for the proximal action on H_∞ which is an ellisphere of dimension $d - 2$ and the two connected components of $H_\infty \setminus \mathcal{E}$, one of them being a ball.

For the action on the affine chart $\mathbb{P}^d \setminus H_\infty$, the origin is fixed, there is a cone $\mathcal{C}_{\text{light}}$ which gives two orbits. Finally, the orbit of an element inside the cone is one sheet of a hyperboloid with two sheets and the orbit of an element outside the cone is a hyperboloid with one sheet. The space $\mathbb{P}^d \setminus (H_\infty \cup \mathcal{C}_{\text{light}})$ has three connected components, two of them are balls, these are the inside of the cone, the remaining one is the outside.

7.6.3. Action of Diag_{d-1} on \mathbb{P}^d . We define the group

$$\text{Diag}_{d-1} = \left\{ \left(\begin{array}{cccc} \lambda_1 & & & 0 \\ & \ddots & & \\ & & \lambda_d & \\ 0 & & & 1 \end{array} \right) \mid \lambda_1, \dots, \lambda_d \in \mathbb{R}_+^* \text{ such that } \lambda_1 \cdots \lambda_d = 1 \right\}.$$

The action of Diag_{d-1} on \mathbb{P}^d has exactly $d + 1$ fixed points which are in generic position. This action preserves $d + 1$ hyperplanes $(H_i)_{i=1, \dots, d+1}$ each of them generated by d fixed points. The orbit of any point $x \in \mathbb{P}^d$ which is not in one of the $(H_i)_{i=1, \dots, d+1}$ is a convex hypersurface of \mathbb{P}^d i.e. an open subset of the boundary of a properly convex open subset.

7.6.4. Conclusion

Lemma 7.17. *Let P be an irreducible loxodromic Coxeter polytope of \mathbb{S}^d . If P has a perfect non-elliptic vertex then G_P is conjugate to $\mathrm{SO}_{d,1}^\circ(\mathbb{R})$ or equal to $\mathrm{SL}_{d+1}(\mathbb{R})$.*

Proof. To simplify notation, we denote G_P by G . Since P is irreducible and loxodromic, we know from Theorem 7.13 and Proposition 2.9 that G is a semi-simple proximal Lie subgroup of $\mathrm{SL}_{d+1}(\mathbb{R})$. Hence the limit set Λ_G of G is the unique closed orbit of the action of G on \mathbb{P}^d and a smooth Zariski closed submanifold of \mathbb{P}^d .

If P admits a parabolic vertex p , then the Zariski closure of Γ_p is conjugate to \mathcal{U}_{d-1} (Lemma 7.16). Apart from the points on a unique hyperplane H_p containing p , for every $x \notin H_p$, the space $\mathcal{U}_{d-1} \cdot x \cup \{p\}$ is an ellipsoid \mathcal{E}_x . Since G is irreducible, we can find a point $x \in \Lambda_G$ but not in H_p . Thus, the limit set Λ_G must contain an ellipsoid, and so Λ_G is an ellipsoid or the all \mathbb{P}^d . Lemma 7.14 concludes.

If P admits a loxodromic vertex p such that W_p is not affine, then the connected component of the Zariski closure G_p of Γ_p is conjugate to $\mathrm{SL}_d(\mathbb{R})$ or $\mathrm{SO}_{d-1,1}^\circ(\mathbb{R})$, thanks to Theorem 7.4. If P admits a loxodromic vertex p such that W_p is affine, then the connected component of the Zariski closure G_p of Γ_p is conjugate to Diag_{d-1} thanks to Theorem 7.2. We again apply the idea of Lemma 7.14. In all these three cases, since the action of G is irreducible, we can find a point $x \in \Lambda_G$ such that the orbit of x under G_p is of dimension $d - 1$. Hence, Lemma 7.14 concludes. \square

Proof of Theorem 7.11. We assume P is not perfect, so P admits a perfect non-elliptic vertex and Lemma 7.17 concludes. \square

Proof of Corollary 7.12. Thanks to Theorems 7.4 and 7.11 which can be applied to P or P^\dagger and the fact that $\Gamma_P \subset \Gamma_{P^\dagger}$, we just have to prove that if $G_P = \mathrm{SO}_{d,1}^\circ(\mathbb{R})$ then $\Gamma_{P^\dagger} \subset \mathrm{SO}_{d,1}^\circ(\mathbb{R})$. In that case, Γ_P preserves a unique ellipsoid \mathcal{E} , any loxodromic vertex p is outside $\bar{\mathcal{E}}$. Let Π_p be the hyperplane spanned by the polar $[v_s]$ for s facets of P containing p . The hyperplane Π_p is the hyperplane p^\perp for the quadratic form defined by \mathcal{E} . Hence, the group $\Gamma_{P^\dagger} \subset \mathrm{SO}_{d,1}^\circ(\mathbb{R})$. \square

7.7. Degenerate 2-perfect case. We just give the statement for the degenerate 2-perfect case without proof since the proof are similar and easier. The subgroups Trans_{d-1} , $\mathrm{SO}_{d-1,1}^\circ(\mathbb{R})$ and $\mathrm{SL}_d(\mathbb{R})$ of $\mathrm{SL}_d(\mathbb{R})$ can be embedded in $\mathrm{SL}_{d+1}(\mathbb{R})$ in the upper-left corner. We make the abuse of notation of identifying these subgroups of $\mathrm{SL}_d(\mathbb{R})$ with their images in $\mathrm{SL}_{d+1}(\mathbb{R})$.

Proposition 7.18. *Let P be a 2-perfect Coxeter polytope of \mathbb{S}^d which is not perfect. If P is decomposable then G_P is conjugate to Trans_{d-1} , $\mathrm{SO}_{d-1,1}^\circ(\mathbb{R})$ or $\mathrm{SL}_d(\mathbb{R})$.*

8. About the convex set

In this section, we prove Theorems C, D, E, and G.

8.1. The convex set Ω_P is the largest convex open subset of \mathbb{P}^d preserved by Γ_P . We start with Theorem C.

Theorem 8.1. *Let P be a loxodromic 2-perfect Coxeter polytope. Then Ω_P is the largest convex open subset preserved by Γ_P .*

Proof. Remark 7.15 shows that $\Omega_{\max} = \bigcap_{\gamma \in \Gamma^{\text{prox}}} H_\gamma^+$. Proposition 4.24 shows that $\Omega_P \setminus \Omega_{\min}$ modulo Γ is a finite union of sets each containing a (Γ_P, Γ_p) -precisely invariant nicely embedded cone \mathcal{C}_p , for p running over the set of loxodromic vertices of P and $\mathcal{D}_p(\mathcal{C}_p) = \Omega_p$.

The closure of the set F_p^{prox} of attractive bi-proximal fixed points of Γ_p is the limit set Λ_p (by Benoist [BEN00]), and Vey shows that since the action of Γ_p on Ω_p is cocompact, we also have $C(F_p^{\text{prox}}) = \Omega_p$ ([VEY70]). We stress that $p \in H_\gamma$ for every $\gamma \in \Gamma_p^{\text{prox}}$.

Hence, the convex set $\Omega_{\max} = \bigcap_{\gamma \in \Gamma^{\text{prox}}} H_\gamma^+$ contains in its boundary any loxodromic vertex p of P , and we have $\mathcal{D}_p(\Omega_{\max}) = \Omega_p$. Let \mathbb{A} be an affine chart containing $\overline{\Omega_{\max}}$. The convex set $\Omega'_p = \bigcap_{\gamma \in \Gamma_p^{\text{prox}}} H_\gamma^+ \cap \mathbb{A}$ is a cone of summit p such that $\mathcal{D}_p(\Omega'_p) = \Omega_p$, so $\Omega_{\max} \subset \Omega_P$. □

8.2. When is Ω_P the smallest convex open subset of \mathbb{P}^d preserved by Γ_P ? We are ready to prove Theorem D.

Theorem 8.2. *Let P be a loxodromic 2-perfect Coxeter polytope. The convex set Ω_P is the smallest convex open subset of \mathbb{P}^d preserved by Γ_P if and only if the action of Γ_P on Ω_P is of finite covolume. In that case, the convex set Ω_P is the unique properly convex open set preserved by Γ_P .*

Proof. Thanks to Theorem 6.3, we only have to show that the convex set Ω_P is the smallest convex open subset of \mathbb{P}^d preserved by Γ_P if and only if every vertex of P is elliptic or parabolic. Suppose one vertex p of P is loxodromic, Proposition 4.24 builds a convex set Ω' preserved by Γ_P which is strictly included in Ω_P .

Suppose every vertex of P is elliptic or parabolic. The parabolic vertices of P are in Λ_P by Proposition 4.7 and the elliptic vertices are in $C(\Lambda_P)$ by Proposition 4.4. Thereby, the vertices of P are in $\overline{C(\Lambda_P)}$, so $P \cap \Omega_P \subset C(\Lambda_P)$. This implies $\Omega_P \subset C(\Lambda_P)$ by definition of Ω_P and so $\Omega_P = C(\Lambda_P)$. Hence, Ω_P is the smallest properly convex open set preserved by Γ_P . □

8.3. Strict convexity of Ω_P . Here we show Theorem E. The word parabolic can cover different things in geometry. We need to recall some definitions to be precise.

8.3.1. Parabolic automorphism. An automorphism γ of a properly convex open set Ω is *parabolic* when the quantity $\inf_{x \in \Omega} d_\Omega(x, \gamma \cdot x) = 0$ and the infimum is not achieved. One can show that such an element has spectral radius 1 and fixes every point of a unique face of Ω (see [CLT15]).

An isometry γ of a Gromov-hyperbolic space X is *parabolic* when the quantity $\inf_{x \in \Omega} d_\Omega(x, \gamma \cdot x) = 0$ and the infimum is not achieved. Such an isometry has a unique fixed point in the boundary ∂X of X . Every point fixed by a parabolic element of a group Γ acting on ∂X is a *parabolic fixed point*.

8.3.2. Projective structure and holonomy. A *convex projective manifold* M is a quotient Ω/Γ of a properly convex open set Ω by a torsion-free discrete subgroup Γ of $\text{Aut}(\Omega)$. The *holonomy* of an element $\gamma \in \pi_1(M)$ is the corresponding element in Γ . We say that an element $\gamma \in \pi_1(M)$ has *parabolic holonomy* when the corresponding element in Γ is parabolic. Every point of $\partial\Omega$ fixed by a parabolic element is called a *parabolic fixed point*.

8.3.3. The notion of relative hyperbolicity

Definition 8.3. Let Γ be a discrete group and $(\mathcal{P}_i)_{i \in I}$ a finite family of subgroups of Γ . The group Γ is *relatively hyperbolic relatively to the family* $(\mathcal{P}_i)_{i \in I}$ if and only if there exists a proper Gromov-hyperbolic space X and a geometrically finite action²⁶ of Γ on X such that the stabilizer of any parabolic fixed point is conjugate to one of the $(\mathcal{P}_i)_{i \in I}$.

8.3.4. The statement

Theorem 8.4 (compact case by Benoist [BEN04] and Cooper, Long, and Tillmann [CLT15]). *Let Γ be a torsion free discrete subgroup of $\text{SL}_{d+1}(\mathbb{R})$ acting on a properly convex open set Ω . Suppose the action is of finite covolume, the manifold Ω/Γ is the interior of a compact manifold N with boundary and the holonomy of every component of ∂N is parabolic. Then the following are equivalent:*

- (1) *the metric space (Ω, d_Ω) is Gromov-hyperbolic;*
- (2) *the properly convex open set Ω is strictly convex;*
- (3) *the boundary $\partial\Omega$ of Ω is \mathcal{C}^1 ;*
- (4) *the group Γ is relatively hyperbolic relatively to the stabilizer of its parabolic fixed points.*

Remark 8.5. Without any action of a group, one can show that a properly convex open set Ω such that (Ω, d_Ω) is Gromov-hyperbolic has to be strictly convex (Benoist [BEN04]) and with \mathcal{C}^1 -boundary (Karlsson and Noskov [KN02]).

²⁶ See paragraph 6.5 for a definition.

Remark 8.6. One can find avatars of this theorem in the literature, one by Choi in [Cho10] and the implication (1) \implies (4) by M. Crampon and the author in [CM14] in the context of geometrically finite actions. For the next theorem we will need the version quoted previously.

Let P be a Coxeter polytope. If p is a parabolic vertex of P we say the subgroup Γ_p is a *geometric parabolic subgroup* of Γ_P .

Theorem 8.7. *Let P be a loxodromic Coxeter polytope. The following are equivalent:*

- (1) *the properly convex open set Ω_P is strictly convex;*
- (2) *the Coxeter polytope P is 2-perfect and the boundary $\partial\Omega_P$ of Ω_P is \mathcal{C}^1 ;*
- (3) *the Coxeter polytope P is quasi-perfect and the group Γ_P is relatively hyperbolic relatively to its geometrical parabolic subgroups.*

In that case, the metric space (Ω_P, d_{Ω_P}) is Gromov-hyperbolic and the action is of finite covolume.

Proof. Suppose we have (3) and let show (1) and (2). Theorem 6.3 shows that the action of Γ_P on Ω_P is of finite covolume. Since Γ_P is of finite type by Selberg's lemma we can find a finite index subgroup Γ' of Γ_P which is torsion free. The quotient manifold Ω_P/Γ' is of finite volume, it is the interior of a compact manifold N and the holonomy of each component of ∂N is parabolic since P is quasi-perfect. Hence, Theorem 8.4 shows that (Ω_P, d_{Ω_P}) is Gromov-hyperbolic, therefore strictly convex with \mathcal{C}^1 -boundary by Remark 8.5.

We first show that P has to be 2-perfect if we assume 1). If P is not 2-perfect then it exists an edge e of P such that the group W_e is infinite (Proposition 3.1). This implies $e \subset \partial\Omega_P$ by Theorem 2.2 (5). In particular, Ω_P is not strictly convex.

Suppose we have (1) or (2) and P is 2-perfect. First remark that no vertex can be loxodromic from Proposition 4.24 (3). Thereby, every vertex of P is either elliptic or parabolic, hence P is quasi-perfect, so Theorem 6.3 shows that $\mu_{\Omega_P}(P) < \infty$, and we have the first part of the assertion. For the same reason than in the first paragraph of this proof we can use Theorem 8.4 which shows that Γ_P is relatively hyperbolic relatively to the stabiliser of its parabolic fixed points (i.e. its geometrical parabolic subgroups). \square

The following statement is a straightforward corollary of Theorem 8.7 which does not use Theorem 8.4.

Corollary 8.8. *Let W be a Coxeter group. The Tits convex set Ω_{Δ_W} is strictly convex if and only if W is quasi-Lannér. In particular, in that case, Ω_{Δ_W} is an ellipsoid.*

Proof. If W is quasi-Lannér, then Ω_{Δ_W} is an ellipsoid and the action of W on Ω_{Δ_W} is of finite covolume. Now, suppose Ω_{Δ_W} is strictly convex. Then Theorem 8.7 shows that Δ_W is quasi-perfect.²⁷ This means by Remark 3.4 that W is quasi-Lannér. \square




Figure 13. An indecomposable quasi-divisible prism which gives a non-strictly convex set.

Proof of Theorem F. Consider the prism \mathfrak{G} given by Figure 13. The main result of [MAR10] shows that the space of finite covolume Coxeter prisms P such that the dihedral angle of P are the one given by the label of the edges of \mathfrak{G} is homeomorphic to \mathbb{R}^* , so in particular not empty. The group Γ_P is not relatively hyperbolic relatively to the unique parabolic vertex (intersection of the faces 1-3-5) because the subgroup generated by $\sigma_1, \sigma_2, \sigma_3$ is virtually \mathbb{Z}^2 . Theorem 2.18 shows that Γ_P is strongly irreducible, hence Ω_P is indecomposable. Theorem 6.3 shows that $\mu_{\Omega_P}(P) < \infty$. Theorem 8.7 concludes that Ω_P is not strictly convex. \square

8.4. Existence of a strictly convex open set preserved. We now show Theorem G.

8.4.1. The statement. Two standard Coxeter subsystems T and U of (S, M) are *orthogonal* when for every $t \in T, u \in U, m_{tu} = 2$. We denote by T^\perp the maximal subsystem orthogonal to T .

Theorem 8.9 (Moussong [MOU88] in the hyperbolic case, Caprace [CAP09, CAPERR]). *For every Coxeter system (S, M) , and every collection \mathcal{P} of standard parabolic subgroups of W_S , the group W_S is relatively hyperbolic relatively to the W_T for $T \in \mathcal{P}$ if and only if the following three conditions are satisfied.*

- (1) *Each affine sub-system of rank at least 3 of (S, M) is included in one $T \in \mathcal{P}$. For each pair S_1, S_2 of irreducible infinite subsystem which are orthogonal, there exists a $T \in \mathcal{P}$ such that $S_1 \cup S_2 \subset T$.*
- (2) *For all $T \neq T' \in \mathcal{P}$, $T \cap T'$ is a spherical Coxeter system.*
- (3) *For each $T \in \mathcal{P}$, for each irreducible infinite subsystem U of T , we have $U^\perp \subset T$.*

²⁷ We don't need to know that W is relatively hyperbolic since every quasi-Lannér Coxeter group is relatively hyperbolic.

We are going to make the abuse of taking together parabolic vertices and the associated Coxeter parabolic subgroup.

Proposition 8.10. *Let P be a loxodromic 2-perfect Coxeter polytope and \mathcal{P} the set of parabolic vertices of P . Then the pair (W_P, \mathcal{P}) satisfies both point (2) and point (3) of Theorem 8.9.*

Proof. We begin by the second point. Given two vertices p, q of P , the segment $[p, q]$ is included in a unique face f of P of minimal dimension and $W_p \cap W_q = W_f$, which is spherical since $\dim(f) \geq 1$ and P is 2-perfect.

For the third point, for each $p \in \mathcal{P}$, the Coxeter group W_p is affine, hence a direct product of irreducible affine Coxeter group W_{a_1}, \dots, W_{a_r} . Let a be the union of some a_i and b the union of the others, so that $W_a \times W_b = W_p$.

Suppose there exists a facet f of P in $a^\perp \setminus b$. If s is a facet of P , then F_s denotes its support and \mathbb{A}_s the affine chart $\mathbb{S}^d \setminus F_s$ not containing p . Let l be the intersection $l = \bigcap_{s \in a} F_s$ (if $r = 1$ then $l = \{p, -p\}$).

The polar v_f of f belongs to l . Moreover, $\alpha_f(v_f) = 2$ and $\alpha_f(p) < 0$ (since $f \notin a \cup b$), hence $v_f \in \mathbb{A}_f \cap l$ (if $r = 1$ then we get $v_f = -p$). So, there cannot exist an affine chart containing P and its polars, contradicting Lemma 4.10. \square

When P is a 2-perfect Coxeter polytope and p a loxodromic vertex, we will call Γ_p a *geometrical loxodromic subgroup* of Γ_P .

Corollary 8.11. *Let P be a loxodromic 2-perfect Coxeter polytope whose loxodromic vertices are simple. The following are equivalent:*

- (1) *the convex set Ω_{P^\dagger} is strictly convex;*
- (2) *the boundary of Ω_{P^\dagger} is \mathcal{C}^1 ;*
- (3) *there exists a strictly convex open set Ω' preserved by Γ_P ;*
- (4) *there exists a properly convex open set Ω' with \mathcal{C}^1 -boundary preserved by Γ_P ;*
- (5) *the group Γ_P is relatively hyperbolic relatively to its geometric parabolic subgroups;*
- (6) *the group Γ_{P^\dagger} is relatively hyperbolic relatively to its geometric parabolic subgroups.*

In this case, the metric space $(\Omega_{P^\dagger}, d_{\Omega_{P^\dagger}})$ is Gromov-hyperbolic, hence Ω_{P^\dagger} is strictly convex with \mathcal{C}^1 -boundary.

Remark 8.12. If the group Γ_P is relatively hyperbolic relatively to its geometric parabolic subgroups then its loxodromic subgroups are Gromov-hyperbolic since for every ridge r the group Γ_r is finite.

8.4.2. A lemma about just-infinite subsystems

Definition 8.13. Let W be a Coxeter group given by the Coxeter system (S, M) . A subsystem U of S is *just infinite* when the Coxeter group W_U is infinite and for every element $u \in U$, the Coxeter group $W_{U \setminus \{u\}}$ is finite.

An infinite Coxeter group W always contains a just infinite subsystem. A Coxeter group W is just infinite if and only if W is irreducible affine or Lannér.

Definition 8.14. Let P be a Coxeter polytope. Let U be a set of facets of P . We say U *bounds a right angle facet* when there exists a facet f of P such that every ridge of f is also a ridge of a facet of U , and all the ridges of f are right angle.

If U bounds a right angle facet f then the projective subspace Π_U spanned by the polar of the facets of U is included in the support of f .

Definition 8.15. Let P be a Coxeter polytope. Let U be a set of facets of P of cardinal r . The projective subspace Π_U *meets nicely* P when

- (1) Π_U is of dimension $r - 1$,
- (2) $\Pi_U \cap P \neq \emptyset$,
- (3) U bounds a right angle facet, or
- (3)' the only facets of P met by Π_U are the facets of U , and the ridges of P met by Π_U are met in their relative interior.

Remark 8.16. Let P be a Coxeter polytope. Let U be a subsystem of facets of P such that the projective space Π_U meets nicely P . Then $P \cap \Pi_U$ is a polytope and its facets are in correspondence with U , hence P induces a Coxeter structure on $P \cap \Pi_U$, and $P \cap \Pi_U$ is tiling the convex set $\Omega_P \cap \Pi_U$. Roughly speaking, we find a sub-Coxeter-polytope of P .

Remark 8.17. Let P be an irreducible loxodromic Coxeter polytope. Let p be a vertex of P , and S_p be the set of facets containing p . We saw at Proposition 4.14 that the projective space Π_{S_p} meets nicely P if p is loxodromic, perfect and simple.

Remark 8.18. Let P be an irreducible loxodromic Coxeter polytope. Let U be the union of two facets which do not intersect. Then the projective space $l = \Pi_U$ is a line that intersects P nicely thanks to the inequalities in (C). Hence, $P \cap l$ is a Coxeter segment which is tiling the segment $\Omega_P \cap l$.

Lemma 8.19. *Let P be an irreducible loxodromic Coxeter polytope. Let U be a just-infinite set of facets of P such that $U \not\subset S_p$ for every parabolic or loxodromic vertex p of P . Then the projective space Π_U meets nicely P , the Coxeter polytope $\Delta = \Pi_U \cap P$ is a simplex, and verifies $A_\Delta = A_U$ and $W_\Delta = W_U$.*

In particular, the group Γ_U acts cocompactly on $\Pi_U \cap \Omega_P$. In particular, Γ_U contains a bi-proximal element.

Proof. First, we show that Π_U is of dimension the rank r of U minus 1. If W_U is a Lannér Coxeter group then A_U is the Cartan matrix of a perfect loxodromic simplex so of strictly negative determinant hence of full rank qed. If W_U is irreducible affine then either A_U is the Cartan matrix of a perfect loxodromic simplex (and $W_U = \tilde{A}_{r-1}$) and we conclude by the same argument. Otherwise, A_U is the Cartan matrix of a parabolic simplex and Lemma 8.20 shows that there exists a vertex p of P such that $U = S_p$. We assume that this is not the case.

We denote by S the set of facets of P and by T the complement $S \setminus U$ of U . If $t \in T$, let F_t be the hyperplane spanned by t . We denote by \mathbb{A}_t the connected component of $\mathbb{S}^d \setminus F_t$ that contains the interior of P . Finally, let $C_T = \bigcap_{t \in T} \overline{\mathbb{A}_t}$. The convex set C_T is not necessarily properly convex. The inequalities (C) show that for every $u \in U$, the polar $v_u \in C_T$.

Let U' be any proper subset of U . Since U is just-infinite, U' is spherical and Lemma 8.20 shows that the intersection $f_{U'} = \bigcap_{u \in U'} u$ is a face of P . The intersection $f_U = \bigcap_{u \in U} u$ is not a face of P because otherwise we would have $U \subset S_p$ for some vertex p of P . So there exists a set V of $d - r + 2$ facets of P not in U such that the polytope Q obtained from P by keeping only the facets in $U \cup V$ is a polytope of dimension d with $d + 2$ facets. The combinatorics of such a polytope is well-known, they are product of two simplices, or cone over a polytope of dimension $d - 1$ with $(d - 1) + 2$ facets.

The polytope Q is not a cone, since the intersection of any two facets of U is a ridge of Q , thanks to Lemma 8.20 that can be applied because U is just-infinite. So Q is the product of two simplices. Finally, any proper subset of facets of U intersect to give a face of Q , so Q is the product of a $(r - 1)$ -simplex by a $(d - r + 1)$ -simplex.

Let $C_V = \bigcap_{t \in V} \overline{\mathbb{A}_t}$. We have $P \cap \Pi_U \subset C_V$, thanks to the inequalities in (C). Now since Q is the product of a $(r - 1)$ -simplex by a $(d - r + 1)$ -simplex, we get that $P \cap \Pi_U \neq \emptyset$.

If U bound a right angle facet. Then Π_U meets nicely P by definition. Suppose U does not bound a right angle facet. Then $P \cap \Pi_U$ is included in the interior of C_T and so a facet f of P such that $f \cap \Pi_U \neq \emptyset$ is a facet of U and the ridges of P met by Π_U are met on their interior. The polytope $P \cap \Pi_U$ is a simplex since Lemma 8.20 shows that any proper set of facets of $P \cap \Pi_U$ meets. \square

Lemma 8.20 (Vinberg [Vin71], Theorem 7). *Let P be a Coxeter polytope. Let (S, M) be the Coxeter system associated to P and W the corresponding Coxeter group. Let S' be a subsystem.*

- *If $W_{S'}$ is finite, then there exists a face f of P such that $S' = S_f = \{s \in S \mid s \supset f\}$.*
- *If $A_{S'}$ is the Cartan matrix of a parabolic simplex then there exists a vertex p of P such that $S' = S_p$.*

8.4.3. The proof of Theorem 8.11

Proof of Theorem 8.11. We begin by (6) \iff (1) \iff (2). Since P^\dagger is quasi-perfect, the conclusion follows from Theorem 8.7. The implication (1) \implies (3) and (2) \iff (4) are obvious since the convex set Ω_{P^\dagger} is preserved by Γ_P and $\Gamma_P \subset \Gamma_{P^\dagger}$. Theorem 8.9 shows (5) \iff (6).

\neg (5) \implies (\neg (3) & \neg (4)). Let Ω' be a properly convex open set preserved by Γ_P . By Theorem 8.9 and Proposition 8.10, we only have to distinguish the cases, A) there exists a loxodromic vertex p such that W_p is not Gromov-hyperbolic, B) there exists an affine sub-system U of rank at least 3 which is not included in a geometric parabolic or loxodromic subgroup of Γ_P , and C) there exist two infinite sub-systems U_1 and U_2 which are orthogonal and $U_1 \cup U_2$ is not included in a geometric parabolic or loxodromic subgroup of Γ_P .

Suppose we are in case A). We have to show that Ω' is not strictly convex nor with \mathcal{C}^1 -boundary. Consider the projective space $\Pi_p = \Pi_{S_p}$ where S_p is the set of facets containing p . Since p is simple loxodromic, the projective space spanned by the limit set Λ_p of Γ_p is Π_p , and we know from Proposition 4.14 that Π_p is of dimension $d - 1$. So the convex set $\Pi_p \cap \Omega'$ is of dimension $d - 1$ and the action of Γ_P on it is cocompact since P is 2-perfect, hence by Theorem 8.4 (cocompact case) the convex set $\Omega' \cap \Pi_p$ is not strictly convex nor with \mathcal{C}^1 -boundary since $\Gamma_p = W_p$ is not Gromov-hyperbolic. Hence, the same is true for Ω' .

Suppose we are in case B) or C). we claim that in this case, the group Γ_P contains a subgroup isomorphic to \mathbb{Z}^2 generated by two bi-proximal elements hence Lemma 8.21 below shows that Ω_P cannot be strictly convex nor with \mathcal{C}^1 -boundary.

Suppose we are in case B). We can assume U is just-infinite. Since $U \not\subset S_p$ for any loxodromic or parabolic vertex p of P , by Lemma 8.19, the projective space Π_U meets nicely P , hence Γ_U acts cocompactly on $\Omega_P \cap \Pi_U$. Since, W_U is an irreducible affine Coxeter group, we know that W_U has to be of type \tilde{A}_n with $n \geq 2$ and by Proposition 2.9 $\Omega_P \cap \Pi_U$ is a simplex. Hence, Γ_U contains two bi-proximal elements which generate a \mathbb{Z}^2 and Lemma 8.21 concludes.

Suppose we are in case C). We can assume U_1 and U_2 are just-infinite sub-systems. We need to distinguish two cases before concluding. a) $U_1 \subset S_p$ for some vertex p of P . In that case, $U_1 \not\subset (S_p)_q$ for any parabolic or loxodromic

vertex q of P_p since P_p is perfect. Hence, Lemma 8.19 applied to P_p shows Γ_{U_1} contains a bi-proximal element for its action on \mathbb{S}_p^{d-1} . But the eigenvalue at p for any element of Γ_p is one, hence Γ_{U_1} as a subgroup of $\mathrm{SL}_{d+1}(\mathbb{R})$, has a bi-proximal element. b) $U_1 \not\subset S_p$, for any vertex p of P . Then Lemma 8.19 applied to P shows Γ_{U_1} contains a bi-proximal element.

So in any situation, the groups Γ_{U_1} and Γ_{U_2} contain a bi-proximal element. Since these two groups commute, we get that Γ_P contains two elements which are bi-proximal and generate a \mathbb{Z}^2 . Lemma 8.21 concludes. \square

Lemma 8.21. *Let Ω be a properly convex open set. Suppose $\mathrm{Aut}(\Omega)$ contains two bi-proximal elements γ, δ which generate a \mathbb{Z}^2 . Then Ω is not strictly convex nor with \mathcal{C}^1 -boundary.*

Proof. Let $p_\gamma^+, p_\gamma^-, p_\delta^+, p_\delta^-$ be the attractive and repulsive fixed points of γ and δ . Let Γ be the group generated by γ and δ . We claim that the set $F = \{p_\gamma^+, p_\gamma^-, p_\delta^+, p_\delta^-\}$ is of cardinality 3. Indeed, if F is of cardinality 2 then the group Γ acts properly on the segment joining the two points of F include in Ω , hence Γ is cyclic. If F is of cardinality 4, then a ping-pong argument shows Γ contains a free subgroup of rank 2.

We call p^0 the point p_δ^+ or p_δ^- different from p_γ^+, p_γ^- . Hence, the plane Π generated by $p^0, p_\delta^+, p_\delta^-$ is preserved by γ and we are in a dimension 2 situation. It is then easy to see that the segments $[p^0, p_\delta^+]$ and $[p^0, p_\delta^-]$ are included in $\partial\Omega \cap \Pi$. Thereby, Ω is not strictly convex nor with \mathcal{C}^1 -boundary. See [MAR12A] for more details. \square

Remark 8.22. If we do not assume that the loxodromic vertices are simple then the statements (1), (2), and (6) of Theorem 8.11 do not make sense any more. But, we still have (3) or (4) \implies (5). But, the author does not know how to build a strictly convex invariant open set (or with \mathcal{C}^1 boundary) assuming (5).

References

- [AMS95] H. Abels, G. Margulis, and G. Soifer, Semigroups containing proximal linear maps. *Israel J. Math.* **91** (1995), no. 1-3, 1–30. [Zbl 0845.22004](#) [MR 1348303](#)
- [Bal14] S. Ballas, Deformations of noncompact projective manifolds. *Algebr. Geom. Topol.* **14** (2014), no. 5, 2595–2625. [Zbl 1309.57014](#) [MR 3276842](#)
- [Bal15] S. Ballas, Finite volume properly convex deformations of the figure-eight knot. *Geom. Dedicata* **178** (2015), 49–73. [Zbl 1329.57004](#) [MR 3397481](#)
- [BHV08] B. Bekka, P. de la Harpe, and A. Valette, *Kazhdan's property (T)*. New Mathematical Monographs, 11. Cambridge University Press, Cambridge, 2008. [Zbl 1146.22009](#) [MR 2415834](#)

- [Ben00] Y. Benoist, Automorphismes des Cônes convexes. *Invent. Math.* **141** (2000), no. 1, 149–193. [Zbl 0957.22008](#) [MR 1767272](#)
- [Ben03] Y. Benoist, Convexes divisibles. II. *Duke Math. J.* **120** (2003), no. 1, 97–120. [Zbl 1037.22022](#) [MR 2010735](#)
- [Ben04] Y. Benoist, Convexes divisibles. I. In S. G. Dani and G. Prasad (eds.), *Algebraic groups and arithmetic*. (Mumbai, 2001.) Published for the Tata Institute of Fundamental Research by Narosa Publishing House, New Delhi, 2004, 339–374. [Zbl 1084.37026](#) [MR 2094116](#)
- [Ben06a] Y. Benoist, Convexes divisibles. IV. Structure du bord en dimension 3. *Invent. Math.* **164** (2006), no. 2, 249–278. [Zbl 1107.22006](#) [MR 2218481](#)
- [Ben06b] Y. Benoist, Convexes hyperboliques et quasiisométries. *Geom. Dedicata* **122** (2006), 109–134. [Zbl 1122.20020](#) [MR 2295544](#)
- [Ben09] Y. Benoist, Five lectures on lattices in semisimple Lie groups. In L. Bessières, A. Parreau, and B. Rémy (eds.), *Géométries à courbure négative ou nulle, groupes discrets et rigidités*. (Grenoble, 2004.) Séminaires et Congrès, 18. Société Mathématique de France, Paris, 2009, 117–176. [Zbl 1252.22006](#) [MR 2655311](#)
- [Bnz60] J.-P. Benzécri, Sur les variétés localement affines et localement projectives. *Bull. Soc. Math. France* **88** (1960), 229–332. [Zbl 0098.35204](#) [MR 0124005](#)
- [Bor60] A. Borel, Density properties for certain subgroups of semi-simple groups without compact components. *Ann. of Math. (2)* **72** (1960) 179–188. [Zbl 0094.24901](#) [MR 0123639](#)
- [Bou68] N. Bourbaki, *Éléments de mathématique*. Fasc. XXXIV. Groupes et algèbres de Lie. Chapitres IV–VI. Actualités Scientifiques et Industrielles, 1337. Hermann, Paris, 1968. [Zbl 0186.33001](#) [MR 0240238](#)
- [Bow95] B. Bowditch, Geometrical finiteness with variable negative curvature. *Duke Math. J.* **77** (1995), no. 1, 229–274. [Zbl 0877.57018](#) [MR 1317633](#)
- [BJS88] M. Bożejko, T. Januszkiewicz, and R. Spatzier, Infinite Coxeter groups do not have Kazhdan’s property. *J. Operator Theory* **19** (1988), no. 1, 63–67. [Zbl 0662.20040](#) [MR 0950825](#)
- [Cap09] P.-E. Caprace, Buildings with isolated subspaces and relatively hyperbolic Coxeter groups. *Innov. Incidence Geom.* **10** (2009), 15–31. [Zbl 1254.20036](#) [MR 2665193](#)
- [CapErr] P.-E. Caprace, Erratum to “Buildings with isolated subspaces and relatively hyperbolic Coxeter groups.” *Innov. Incidence Geom.* **14** (2015), 77–79. [Zbl 1336.20044](#) [MR 3450952](#)
- [Che69] M. Chein, Recherche des graphes des matrices de Coxeter hyperboliques d’ordre ≤ 10 . *Rev. Française Informat. Recherche Opérationnelle* **3** (1969), Sér. R-3, 3–16. [Zbl 0276.20043](#) [MR 0294181](#)
- [CL15] H. Chen and J.-Ph. Labbé, Lorentzian Coxeter systems and Boyd–Maxwell ball packings. *Geom. Dedicata* **174** (2015), 43–73. [Zbl 1312.52012](#) [MR 3303040](#)

- [Cho06] S. Choi, The deformation spaces of projective structures on 3-dimensional Coxeter orbifolds. *Geom. Dedicata* **119** (2006), 69–90. [Zbl 1103.57013](#) [MR 2247648](#)
- [Cho10] S. Choi, The convex real projective manifolds and orbifolds with radial ends: the openness of deformations. Preprint 2010. [arXiv:1011.1060](#) [math.GT]
- [Cho13] S. Choi, A classification of radial and totally geodesic ends of properly convex real projective orbifolds. Preprint 2013. [arXiv:1304.1605](#) [math.GT]
- [CHL12] S. Choi, C. Hodgson, and G.-S. Lee, Projective deformations of hyperbolic Coxeter 3-orbifolds. *Geom. Dedicata* **159** (2012), 125–167. [Zbl 1266.57012](#) [MR 2944525](#)
- [CL15] S. Choi and G.-S. Lee, Projective deformations of weakly orderable hyperbolic Coxeter orbifolds. *Geom. Topol.* **19** (2015), no. 4, 1777–1828. [Zbl 1333.57028](#) [MR 3375519](#)
- [CLT15] D. Cooper, D. Long, and S. Tillmann, On Convex Projective Manifolds and Cusps. *Adv. Math.* **277** (2015), 181–251. [Zbl 06431971](#) [MR 3336086](#)
- [CH07] Y. de Cornulier and P. de la Harpe, Décompositions de groupes par produit direct et groupes de Coxeter. In G. N. Arzhantseva, L. Bartholdi, J. Burillo, and E. Ventura, *Geometric group theory*. (Geneva and Barcellona, 2005.) Trends in Mathematics. Birkhäuser Verlag, Basel, 2007, 75–102. [Zbl 1147.20023](#) [MR 2395791](#)
- [Cox34] H. Coxeter, Discrete groups generated by reflections. *Ann. of Math. (2)* **35** (1934), no. 3, 588–621. [JFM 60.0898.02](#) [Zbl 0010.01101](#) [MR 1503182](#)
- [CM13] M. Crampon and L. Marquis, Un lemme de Kazhdan–Margulis–Zassenhaus pour les géométries de Hilbert. *Ann. Math. Blaise Pascal* **20** (2013), no. 2, 363–376. [Zbl 1282.22007](#) [MR 3138033](#)
- [CM14] M. Crampon and L. Marquis, Finitude géométrique en géométrie de Hilbert. *Ann. Inst. Fourier (Grenoble)* **64** (2014), no. 6, 2299–2377. [Zbl 1306.22005](#) [MR 3331168](#)
- [FK94] J. Faraut and A. Korányi, *Analysis on symmetric cones*. Oxford Mathematical Monographs. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1994. [Zbl 0841.43002](#) [MR 1446489](#)
- [Gol90] W. Goldman, Convex real projective structures on compact surfaces. *J. Differential Geom.* **31** (1990), no. 3, 791–845. [Zbl 0711.53033](#) [MR 1053346](#)
- [GG96] I. Gol'dsheĭd and Y. Guivarc'h, Zariski closure and the dimension of the Gaussian law of the product of random matrices. I. *Probab. Theory Related Fields* **105** (1996), no. 1, 109–142. [Zbl 0854.60006](#) [MR 1389734](#)
- [GM89] I. Gol'dsheĭd and G. Margulis, Lyapunov exponents of a product of random matrices. *Uspekhi Mat. Nauk* **44** (1989), no. 5(269), 13–60. In Russian. English translation, *Russian Math. Surveys* **44** (1989), no. 5, 11–71. [Zbl 0705.60012](#) [MR 1040268](#)
- [KV67] V. Kac and É. Vinberg, Quasi-homogeneous cones. *Mat. Zametki* **1** (1967), 347–354. In Russian. English translation, *Math. Notes* **1** (1967), 231–235. [Zbl 0163.16902](#) [MR 0208470](#)

- [KN02] A. Karlsson and G. Noskov, The Hilbert metric and Gromov hyperbolicity. *Enseign. Math. (2)* **48** (2002), no. 1-2, 73–89. [Zbl 1046.53026](#) [MR 1923418](#)
- [Koe99] M. Koecher, *The Minnesota notes on Jordan algebras and their applications*. Edited, annotated and with a preface by A. Krieg and S. Walcher. Lecture Notes in Mathematics, 1710. Springer-Verlag, Berlin, 1999. [Zbl 1072.17513](#) [MR 1718170](#)
- [Kos67] J.-L. Koszul, *Lectures on hyperbolic Coxeter groups*. University of Notre Dame, Notre Dame, IN, 1967.
- [Kos68] J.-L. Koszul, Déformations de connexions localement plates. *Ann. Inst. Fourier (Grenoble)* **18** (1968), fasc. 1, 103–114. [Zbl 0167.50103](#) [MR 0239529](#)
- [Kui53] N. Kuiper, On convex locally-projective spaces. In *Convegno Internazionale di Geometria Differenziale*. (1953) Edizioni Cremonese, Roma, 1954, 200–213. [Zbl 0057.14303](#) [MR 0063115](#)
- [Lan50] F. Lannér, On complexes with transitive groups of automorphisms. *Comm. Sémin., Math. Univ. Lund [Medd. Lunds Univ. Mat. Sem.]* **11**, (1950). 71 pp. [Zbl 0037.39802](#) [MR 0042129](#)
- [Mar10] L. Marquis, Espace des modules de certains polyèdres projectifs miroirs. *Geom. Dedicata* **147** (2010), 47–86. [Zbl 1243.52015](#) [MR 2660566](#)
- [Mar12a] L. Marquis, Surface projective convexe de volume fini. *Ann. Inst. Fourier (Grenoble)* **62** (2012), no. 1, 325–392. [Zbl 1254.57015](#) [MR 2986273](#)
- [Mar12b] L. Marquis, Exemples de variétés projectives strictement convexes de volume fini en dimension quelconque. *Enseign. Math. (2)* **58** (2012), no. 1-2, 3–47. [Zbl 1284.57021](#) [MR 2985008](#)
- [Mar13] L. Marquis, Around groups in Hilbert geometry. In A. Papadopoulos and M. Troyanov (eds.), *Handbook of Hilbert geometry*. IRMA Lectures in Mathematics and Theoretical Physics, 22. European Mathematical Society (EMS), Zürich, 2014, 07–261. [MR 3329882](#)
- [MV00] G. Margulis and É. Vinberg, Some linear groups virtually having a free quotient. *J. Lie Theory* **10** (2000), no. 1, 171–180. [Zbl 0958.22008](#) [MR 1748082](#)
- [Max82] G. Maxwell, Sphere packings and hyperbolic reflection groups. *J. Algebra* **79** (1982), no. 1, 78–97. [Zbl 0556.51003](#) [MR 0679972](#)
- [Mou88] G. Moussong, *Hyperbolic Coxeter groups*. Ph. D. Thesis. Ohio State University, Columbus, OH, 1988. [MR 2636665](#)
- [Nie15] X. Nie, On the Hilbert geometry of simplicial Tits sets. *Ann. Inst. Fourier (Grenoble)* **65** (2015), no. 3, 1005–1030. [Zbl 1341.37018](#) [MR 3449173](#)
- [Par07] L. Paris, Irreducible Coxeter groups. *Internat. J. Algebra Comput.* **17** (2007), no. 3, 427–447. [Zbl 1134.20046](#) [MR 2333366](#)
- [Ver05] C. Vernicos, Introduction aux géométries de Hilbert. In *Actes de Séminaire de Théorie Spectrale et Géométrie*. Vol. 23. Année 2004–2005. Université de Grenoble I, Institut Fourier, Saint-Martin-d’Hères, 2005, 145–168. [Zbl 1100.53031](#) [MR 2270228](#)

- [Vey70] J. Vey, Sur les automorphismes affines des ouverts convexes saillants. *Ann. Scuola Norm. Sup. Pisa* (3) **24** (1970), 641–665. [Zbl 0206.51302](#) [MR 0283720](#)
- [Vin63] É. Vinberg, The theory of homogeneous convex cones. *Trudy Moskov. Mat. Obšč.* **12** (1963), 303–358. In Russian. English translation, *Trans. Mosc. Math. Soc.* **12** (1963), 340–403. [Zbl 0138.43301](#) [MR 0158414](#)
- [Vin65] É. Vinberg, Structure of the group of automorphisms of a homogeneous convex cone. *Trudy Moskov. Mat. Obšč.* **13** (1965), 56–83. [Zbl 0224.17010](#) [MR 0201575](#)
- [Vin71] É. Vinberg, Discrete linear groups that are generated by reflections. *Izv. Akad. Nauk SSSR Ser. Mat.* **35** (1971), 1072–1112. [Zbl 0247.20054](#) [MR 0302779](#)
- [Vin85] É. Vinberg, Hyperbolic groups of reflections. *Uspekhi Mat. Nauk* **40** (1985), no. 1(241), 29–66, 255. In Russian. English translation, *Russ. Math. Surv.* **40** (1985), no. 1, 31–75. [Zbl 0579.51015](#) [MR 0783604](#)

Received June 1, 2015

Ludovic Marquis, IRMAR, UMR CNRS 6625, UFR Mathématiques,
Campus Scientifique de Beaulieu, 263 Avenue du général Leclerc – CS 74205,
35042 Rennes Cedex, France

e-mail: ludovic.marquis@univ-rennes1.fr