

## Almost commuting matrices with respect to the rank metric

Gábor Elek<sup>1</sup> and Łukasz Grabowski<sup>2</sup>

**Abstract.** We show that if  $A_1, A_2, \dots, A_n$  are square matrices, each of them is either unitary or self-adjoint, and they almost commute with respect to the rank metric, then one can find commuting matrices  $B_1, B_2, \dots, B_n$  that are close to the matrices  $A_i$  in the rank metric.

**Mathematics Subject Classification (2010).** 13A99, 15B57.

**Keywords.** Rank metric, almost-commuting matrices, stability problems.

### 1. Introduction

Recently there has been a considerable amount of research devoted to the following family of questions: suppose that square matrices  $A$  and  $B$  fulfil some relation “approximately”. Can we then perturb  $A$  and  $B$  so that the resulting matrices  $A'$  and  $B'$  actually fulfil the relation in question? Let us make it more precise by reviewing some historical and more recent examples.

We start with the most famous one. Paul Halmos [12] posed the following problem, known since as the *Halmos problem*. Let  $\delta > 0$ , and suppose that  $A$  and  $B$  are self-adjoint matrices of norm 1. Can we find  $\varepsilon > 0$  such that if the operator norm of  $AB - BA$  is at most  $\varepsilon$  then there exist self-adjoint matrices  $A'$  and  $B'$  such that  $A'B' = B'A'$  and such that the operator norms of  $A' - A$  and of  $B' - B$  are at most  $\delta$ ?

An affirmative answer to this question was given by Huaxin Lin [15] (see also [9] and [13]). On the other hand, Voiculescu proved that for integers  $d \geq 7$  there exist  $d \times d$  unitary matrices  $U_d, V_d$  such that

---

<sup>1</sup> Gábor Elek was partially supported by the ERC Consolidator Grant “Asymptotic invariants of discrete groups, sparse graphs and locally symmetric spaces” No. 648017.

<sup>2</sup> Łukasz Grabowski was partially supported by the ERC Starting Grant “Limits of Structures in Algebra and Combinatorics” No. 805495.

- $\|U_d V_d - V_d U_d\| = |1 - e^{2\pi i/d}|$ , and
- for any pair  $A_d, B_d$  of commuting  $d \times d$  matrices we have

$$\|U_d - A_d\| + \|V_d - B_d\| \geq \sqrt{2 - |1 - e^{2\pi i/d}|} - 1.$$

In other words, in the original Halmos problem, if we replace the assumption that  $A$  and  $B$  are self-adjoint with the assumption that  $A$  and  $B$  are unitary, then the answer is negative, even if we do not demand that the nearby commuting matrices  $A'$  and  $B'$  should be unitary. Furthermore, counterexamples were found by Davidson [6] if we ask about three or more almost commuting self-adjoint matrices.

A similar question had previously been asked by Rosenthal [19], where the “closeness” and “almost commutativity” of the matrices were defined using the normalised Hilbert-Schmidt norm in place of the operator norm. Affirmative answers to this version of the Halmos Problem were given for arbitrarily large finite families of normal operators by various authors [11], [8], and [10].

More recently the analogous question was studied in [2] for permutations and the Hamming distance. Arzhantseva and Paunescu showed the following result, which was a direct motivation for the investigations presented in this article. For every  $\delta > 0$  there exists  $\varepsilon > 0$  such that if  $A$  and  $B$  are permutations such that the normalised Hamming distance between  $AB$  and  $BA$  is at most  $\varepsilon$  then we can find permutations  $A'$  and  $B'$  such that  $A'B' = B'A'$  and the normalised Hamming distances between  $A$  and  $A'$ , as well as  $B$  and  $B'$ , are both bounded by  $\delta$ . The corresponding result is true also for an arbitrary finite number of permutations.

In this paper we study the analogous question for the *rank metric*. We refer to [3] and the references therein for the background and motivation for studying rank metric, and here we only state the definitions. The set of natural numbers is  $\mathbb{N} := \{0, 1, \dots\}$  and we let  $\mathbb{N}_+ := \{1, 2, \dots\}$ . For  $d \in \mathbb{N}_+$  let  $\text{Mat}(d)$  be the set of all  $d \times d$  square matrices with complex coefficients. Finally, for  $A \in \text{Mat}(d)$  we let  $\text{rank}(A) := \frac{\dim_{\mathbb{C}}(\text{im}(A))}{d}$ . This norm defines a metric on  $\text{Mat}(d)$  in a usual way, i.e.  $d_{\text{rank}}(A, B) := \text{rank}(A - B)$ .

Our main aim in this note is to show the following theorem.

**Theorem 1.** *For every  $\varepsilon > 0$  and  $n \in \mathbb{N}_+$  there exists  $\delta > 0$  such that for all  $d \in \mathbb{N}_+$  we have the following. If  $A_1, A_2, \dots, A_n \in \text{Mat}(d)$  are matrices, each of them is either unitary or self-adjoint, and for all  $1 \leq i, j \leq n$  we have  $\text{rank}(A_i A_j - A_j A_i) \leq \delta$ , then there exist commuting matrices  $B_1, B_2, \dots, B_n$  such that for every  $1 \leq i \leq n$  we have  $\text{rank}(A_i - B_i) \leq \varepsilon$ .*

A more general statement will be presented in Theorem 6.

**Remark 2.** It is natural to ask whether the matrices  $B_1, \dots, B_n$  can be taken to be “of the same type” as the matrices  $A_1, \dots, A_n$ , e.g. whether we can demand, say, the matrix  $B_1$  to be unitary, provided that  $A_1$  is unitary. We do not know the answer to this question.

We think that Theorem 1 likely stays true when  $A_1, \dots, A_n$  are allowed to be arbitrary normal invertible matrices. On the other hand, it would be interesting to find a counterexample when  $A_1, \dots, A_n$  are allowed to be arbitrary invertible matrices.

Becker, Lubotzky and Thom [4] generalised the results from [2] to the context of finitely presented polycyclic groups, and showed that there are significant obstacles to generalise it further. We are able to prove some analogous results in the context of the rank metric. Let us make it precise now.

Let  $\Gamma$  be a finitely presented group with presentation

$$\langle \gamma_1, \dots, \gamma_g \mid P_1(\gamma_1, \dots, \gamma_g), \dots, P_r(\gamma_1, \dots, \gamma_g) \rangle,$$

where  $P_i$  are non-commutative monomials in  $g$  variables (we allow negative exponents here).

For a  $k \times k$  matrix  $B$  we denote with  $\widehat{B}$  the operator on the vector space  $\mathbb{C}^{\oplus \mathbb{N}}$  which acts as  $B$  on the first  $k$  basis vectors and is 0 otherwise.

We say that  $\Gamma$  is *stable with respect to the rank metric* if for every  $\delta > 0$  there exists  $\varepsilon > 0$  such that the following holds. For all  $d \in \mathbb{N}$  we have that if  $A_1, \dots, A_g$  are unitary  $d \times d$  matrices with  $\text{rank}(P_i(A_1, \dots, A_g) - \text{Id}_d) \leq \delta$ , then there exist  $k \in \mathbb{N}$  and invertible  $k \times k$  matrices  $B_1, \dots, B_g$  with  $\dim(\text{im}(\widehat{A}_i - \widehat{B}_i)) \leq \varepsilon \cdot d$  and such that  $P_i(B_1, \dots, B_g) = \text{Id}_k$  for all  $i = 1, \dots, r$ .

**Remarks 3.** (1) Originally, we have not worked with  $\widehat{A}_i$  but rather with  $A_i$  in the definition above. We thank Narutaka Ozawa for pointing out that it is more natural to take  $\widehat{A}_i$ .

(2) It is not hard to check (and we use it implicitly in the discussion above) that the property of being stable with respect to the rank metric does not depend on the choice of a finite presentation of the group  $\Gamma$ .

(3) Theorem 1 implies that the groups  $\mathbb{Z}^k$ , where  $k = 1, 2, \dots$ , are stable with respect to the rank metric. We remark that there exist other natural notions of *being stable with respect to the rank metric*: for example, we could demand the matrices  $B_i$  to be unitary, or we could remove the assumption that the matrices  $A_i$  are unitary. Thus, to avoid confusion, it might be useful to talk about, say,  $(\mathcal{A}, \mathcal{B})$ -stability, where  $\mathcal{A} = ((A_1, d_1), (A_2, d_2), \dots)$  is a sequence of monoids with metrics, and  $\mathcal{B} = (B_1, B_2, \dots)$  is a sequence of groups such that  $B_i \subset A_i$ . We refrain from doing this in this paper as all our results are about the stability with respect to the rank metric, as defined above.

Perhaps the most interesting question which we cannot tackle at present is inspired by the results of [4]: are polycyclic groups stable with respect to the rank metric? However, by using some of the ideas from [4] we can show the following result.

Let  $p$  be a prime number. Recall that Abels' group  $A_p$  (see [1]) is the group of 4-by-4 matrices of the form

$$\begin{pmatrix} 1 & * & * & * \\ & p^m & * & * \\ & & p^n & * \\ & & & 1 \end{pmatrix},$$

where  $m, n \in \mathbb{Z}$ , and where the stars are arbitrary elements of the ring  $\mathbb{Z}[\frac{1}{p}]$  of rational numbers which can be written with a power of  $p$  as the denominator.

**Theorem 4.** *For any prime number  $p$  the Abels' group  $A_p$  is not stable with respect to the rank metric.*

**Remarks 5.** (1) It is not difficult to show that if a finitely presented amenable group is stable with respect to the rank metric then it is residually linear. Thus, mimicking the question posed in [2], one could ask whether every finitely presented linear amenable group is stable with respect to the rank metric. Since Abels' group is a solvable group of step 3 which is finitely presented and linear, Theorem 4 gives a negative answer to this question.

(2) Our proof of Theorem 4 is based on an argument from [4] used to show that the Abels' groups are not stable with respect to the Hamming distance. In fact, Theorem 4 is a generalisation of that particular result from [4]. We will present the proof of Theorem 4 in Section 5. It is very self-contained and can also serve a minor role as an alternative exposition of one of the results of [4] (the advantage of our proof of Theorem 4 compared with the exposition in [4] is somewhat smaller definitional overheads).

**Acknowledgment.** We would like to thank the referees for a very careful reading of this paper, and numerous helpful suggestions and corrections.

## 2. The strategy of the proof and the general statement of Theorem 1

Let us very informally discuss the strategy of the proof of Theorem 1. For simplicity let us assume that we are given two  $d \times d$  matrices  $A$  and  $B$  which are almost commuting with respect to the rank metric.

First, we need to find a large subspace  $W \subset \mathbb{C}^d$  and a decomposition  $W = \bigoplus_{i=1}^N B_i$ , such that each space  $B_i$  has the following two properties:

- (1) there exists  $R_i \in \mathbb{N}$ , an ideal  $\mathfrak{a}_i \subset \mathbb{C}[X, Y]$ , and a linear embedding  $\varphi_i: B_i \rightarrow \mathbb{C}[X, Y]/\mathfrak{a}_i$  whose image consists of all elements of degree at most  $R_i$ , and
- (2) “ $B_i$  is almost invariant for the actions of  $A$  and  $B$ .”

Most of Section 4 is devoted to finding such  $W$ , culminating in Lemma 25. This allows us to replace the original  $A$  and  $B$  with direct sums of multiplication operators in commutative algebras, restricted to “balls in the algebras,” i.e. to subspaces of polynomials with degree bounded by  $R_i$ .

The reduction of the proof of Theorem 6 to finding such  $W$  is described in Lemma 17.

The property that  $A$  and  $B$  are either self-adjoint or unitary is used in two ways. The first use is controlling the nilpotent elements in the resulting commutative algebras. This is done in Lemma 23. While controlling the nilpotent elements greatly simplifies the proof, the authors believe it is not essential.

The second, more crucial, use is making sure that the subspace  $W$  is large. Informally speaking, the assumption that  $A$  and  $B$  are either self-adjoint or unitary allows us to argue that if  $W$  is small, then we can add some extra subspaces  $B_i$  in the orthogonal complement of  $W$  (see Lemma 24). The argument is very similar to the “Ornstein–Weiss trick” (see [18]), and the assumption on  $A$  and  $B$  allows us to replace “disjointedness” with “orthogonality”.

After finding  $W$  we still need to consider the operators of multiplication by  $X$  and  $Y$  in  $\mathbb{C}[X, Y]/\mathfrak{a}_i$  restricted to polynomials of degree bounded by  $R_i$ . These two restrictions clearly almost commute, and we need to perturb them with small rank operators to obtain commuting operators.

In order to be able to carry out the Ornstein–Weiss trick in our setting, we make use of the effective Nullstellensatz (encapsulated in Theorem 9) and the Macaulay theorem on growth in graded algebras (encapsulated in Corollary 13). The final commutative algebra tool which we use is the standard Nullstellensatz (Proposition 14).

The effective Nullstellensatz (i) allows us to argue that the embeddings  $\varphi_i$  exist, i.e. reduce “the local situation to the commutative algebra,” and (ii) together with the assumption that  $A$  and  $B$  are either unitary or self-adjoint, it allows us to control the nilpotent elements in the resulting commutative algebras  $\mathbb{C}[X, Y]/\mathfrak{a}_i$ . It is used in Lemma 23.

The Macaulay theorem (i) allows us to argue that the complement of  $W$  is small, and (ii) it allows us to argue that the commuting perturbations of multiplication operators in commutative algebras which we find, are indeed small rank perturbations. It is used in Lemmas 20 and 24.

**Definitions and the general statement.** Elements of  $\text{Mat}(d)$  will be called  $d$ -matrices. Tuples of  $d$ -matrices will be called  $d$ -matrix tuples, and will be denoted with curly letters, e.g.  $\mathcal{A} = (A_1, \dots, A_n)$  and  $\mathcal{M} = (M_1, \dots, M_n)$ .

For  $a \in \mathbb{N}_+$ , the symbol  $[a]$  denotes the set  $\{1, 2, \dots, a\}$ , and we let  $[0]$  denote the empty set. We say that a matrix tuple  $\mathcal{A} = (A_1, \dots, A_n)$  is *commuting* if for all  $i, j \in [n]$  we have  $A_i A_j - A_j A_i = 0$ . More generally, for  $\varepsilon \geq 0$  we say that  $\mathcal{A}$  is  $\varepsilon$ -commuting if

$$\max_{i, j \in [n]} \text{rank}(A_i A_j - A_j A_i) \leq \varepsilon.$$

If  $d \in \mathbb{N}_+$  and  $\mathcal{A} = (A_1, \dots, A_n)$ ,  $\mathcal{B} = (B_1, \dots, B_n)$  are two  $d$ -matrix tuples, then we let

$$d_{\text{rank}}(\mathcal{A}, \mathcal{B}) := \max_{i \in [n]} \text{rank}(A_i - B_i).$$

Given a matrix  $A$ , we denote the adjoint of  $A$  by  $A^*$ . We say that a  $d$ -matrix tuple  $\mathcal{M} = (M_1, \dots, M_n)$  is *\*-closed* if for every  $i \in [n]$  there exists  $j \in [n]$  such that  $M_i^* = M_j$ .

Our general result is as follows.

**Theorem 6.** *For every  $\varepsilon \geq 0$  and  $n \in \mathbb{N}_+$  there exists  $\delta \geq 0$  such that if*

$$\mathcal{A} = (A_1, \dots, A_n)$$

*is a \*-closed  $\delta$ -commuting matrix tuple then we can find a commuting matrix tuple  $\mathcal{B}$  with*

$$d_{\text{rank}}(\mathcal{A}, \mathcal{B}) \leq \varepsilon.$$

Let us argue how to deduce Theorem 1 from Theorem 6. First, we note that if we replace the expression *a \*-closed  $\delta$ -commuting matrix tuple* in the statement of Theorem 6 by *a  $\delta$ -commuting matrix tuple such that each of the matrices  $A_1, \dots, A_n$  is either self-adjoint or unitary* then we obtain the statement of Theorem 1.

But if  $(A_1, \dots, A_n)$  is any matrix tuple, then  $(A_1, \dots, A_n, A_1^*, \dots, A_n^*)$  is a \*-closed matrix tuple. As such, in order to deduce Theorem 1 from Theorem 6 it is enough to prove the following proposition.

**Proposition 7.** *For every  $n \in \mathbb{N}_+$ , every  $\delta > 0$  and every  $d \in \mathbb{N}_+$ , we have that if  $(A_1, \dots, A_n)$  is a  $\delta$ -commuting  $d$ -matrix tuple and each of the matrices  $A_1, \dots, A_n$  is either unitary or self-adjoint, then the  $d$ -matrix tuple  $(A_1, \dots, A_n, A_1^*, \dots, A_n^*)$  is  $\delta$ -commuting as well.*

*Proof.* Using induction, it is enough to show that if  $A$  is a  $d$ -matrix and  $B$  is either a unitary or a self-adjoint  $d$ -matrix with  $\text{rank}(A, B) \leq \delta$  then also  $\text{rank}(A, B^*) \leq \delta$ .

If  $B$  is self-adjoint then there is nothing to prove. If  $B$  is unitary then we will use the fact that  $B^* = B^{-1}$ . We let

$$W := \ker(AB - BA),$$

and by assumption we have  $\dim(W) \geq 1 - \delta$ . For  $v \in B(W)$  we can write  $v = B(w)$  for some  $w \in W$ , hence we obtain that

$$B^{-1}A(v) = B^{-1}AB(w) = B^{-1}BA(w) = A(w).$$

On the other hand we can write

$$AB^{-1}(v) = AB^{-1}B(w) = A(w).$$

This shows that  $B(W) \subset \ker(AB^{-1} - B^{-1}A)$ , finishing the proof because

$$\dim(B(W)) = \dim(W) \geq 1 - \delta. \quad \square$$

**Remark 8.** With a little bit more effort we could also deal with matrix tuples whose all elements are normal matrices with spectrum contained in the union of the real line and the unit circle.

For the rest of the paper we fix a positive natural number  $n$ . From now on all matrix tuples will have length  $n$ .

### 3. Commutative algebra preliminaries

Let  $\mathbb{C}$  be the field of complex numbers. The ring  $\mathbb{C}[X_1, \dots, X_n]$  will be denoted by  $\mathbb{C}[X]$ . Recall that an ideal  $\mathfrak{a} \subset \mathbb{C}[X]$  is *radical* if for all  $m \in \mathbb{N}_+$  and  $f \in \mathbb{C}[X]$  we have that  $f^m \in \mathfrak{a}$  implies  $f \in \mathfrak{a}$ . By Hilbert’s Nullstellensatz we have that  $\mathfrak{a} = \bigcap \mathfrak{m}$ , where the intersection is over all maximal ideals which contain  $\mathfrak{a}$ .

Given an arbitrary ideal  $\mathfrak{a}$  we denote by  $\text{rad}(\mathfrak{a})$  the *radical of  $\mathfrak{a}$* , i.e. the radical ideal defined as  $\text{rad}(\mathfrak{a}) := \{f \in \mathbb{C}[X]: \text{there exists } m \in \mathbb{Z}_+ \text{ with } f^m \in \mathfrak{a}\}$ .

The next theorem follows from the effective Nullstellensatz of Grete Hermann [14] and the Rabinowitsch trick (see e.g. [5, Theorem 1 and the corollary afterwards]).

**Theorem 9.** *There exists an increasing function  $K: \mathbb{N} \rightarrow \mathbb{N}$  such that we have the following properties. Let  $f, f_1, \dots, f_k \in \mathbb{C}[X]$  be polynomials of degree at most  $R$ , and let  $\mathfrak{a}$  be the ideal generated by  $f_1, \dots, f_k$ .*

(1) *If  $f \in \mathfrak{a}$  then there exist  $h_1, \dots, h_k \in \mathbb{C}[X]$  such that*

$$h_1 f_1 + \dots + h_k f_k = f$$

*and  $\deg(h_i f_i) \leq K(R)$ .*

(2) *If  $f \in \text{rad}(\mathfrak{a})$  then we can find  $m \in \mathbb{N}$  and  $g_1, \dots, g_k \in \mathbb{C}[X]$  such that*

$$g_1 f_1 + \dots + g_k f_k = f^m$$

*and  $\deg(g_i f_i) \leq K(R)$ .*

□

In the applications of this theorem we will implicitly use that  $K(R) \geq R$ .

For the next proposition we need to recall some definitions. A *standard graded  $\mathbb{C}$ -algebra* is a  $\mathbb{C}$ -algebra  $A$  together with a family of vector spaces  $A_i$ ,  $i \in \mathbb{N}$ , such that

- (1)  $A_0 = \mathbb{C}$ ,  $A = \bigoplus_{i \in \mathbb{N}} A_i$ ,
- (2)  $A$  is generated as a  $\mathbb{C}$ -algebra by finitely many elements of  $A_1$ ,
- (3) for all  $i, j \in \mathbb{N}$  we have  $A_i A_j \subset A_{i+j}$ .

A *filtration* on a  $\mathbb{C}$ -algebra  $A$  is an ascending family  $F_0 \subset F_1 \subset \dots$  of linear subspaces of  $A$  such that

- (1)  $F_0 = \mathbb{C}$ ,  $A = \bigcup_{i \in \mathbb{N}} F_i$ ,
- (2) for all  $i, j \in \mathbb{N}$  we have  $F_i F_j \subset F_{i+j}$ .

Given an algebra  $A$  with a filtration  $F_i$ ,  $i \in \mathbb{N}$ , we can associate to it a graded algebra  $\text{gr}(A)$  as follows. As a  $\mathbb{C}$ -vector space we let  $\text{gr}(A) := F_0 \oplus \bigoplus_{i>0} F_i / F_{i-1}$ . We define the multiplication on  $\text{gr}(A)$  first on the elements of the form  $a + F_i$  and  $b + F_j$ , where  $i, j \geq 0$ ,  $a \in F_{i+1}$ ,  $b \in F_{j+1}$ , by the formula  $(a + F_i) \cdot (b + F_j) := ab + F_{i+j+1}$ . In general we extend this multiplication to all of  $\text{gr}(A)$  by  $\mathbb{C}$ -linearity.

**Remark 10.** The reason why  $\text{gr}(A)$  is not always a *standard* graded algebra is that it may happen not to be generated by the elements of  $F_1 / F_0$ . This may be the case even if  $A$  is generated by finitely many elements of  $F_1$  as a  $\mathbb{C}$ -algebra.

For example, let  $A := \mathbb{C}[X_1]$ , let  $F_0 = \mathbb{C}$ , let  $F_1$  be the vector space of polynomials of degree at most 1, and finally for  $i \geq 2$  let  $F_i$  be the vector space of polynomials of degree at most  $2i - 1$ . In this case we have  $(X_1 + F_0)^2 = X_1^2 + F_1$ , and therefore  $(X_1 + F_0)^3 = X_1^3 + F_2$ , i.e.  $(X_1 + F_0)^3$  is equal to 0 in  $\text{gr}(A)$ .

In fact, it is not difficult to construct examples where  $\text{gr}(A)$  fails to be finitely-generated, even when  $A$  is generated by finitely many elements of  $F_1$ .

We say that  $F_i$  is a *standard* filtration on  $A$  if the associated graded algebra  $\text{gr}(A)$  is standard.

The following is a consequence of Macaulay's theorem [16]. We will use the exposition from [7, Section 5].

**Proposition 11.** *Let  $A$  be a  $\mathbb{C}$ -algebra with a standard filtration  $F_i$ . Then for every  $i > 0$  we have*

$$\dim_{\mathbb{C}}(F_i / F_{i-1}) < \frac{\dim_{\mathbb{C}}(F_1)}{i} \dim_{\mathbb{C}}(F_{i-1}).$$

*Proof.* For a natural number  $k$  and a real number  $x$  we let  $\binom{x}{k}$  denote the number  $\frac{1}{k!} \cdot x(x-1) \cdots (x-k+1)$ .



Let us fix  $i > 0$ . After applying [7, Theorem 5.10] to the standard graded algebra  $\text{gr}(A)$  we obtain the following. Let  $x$  be the unique real number such that  $x \geq i - 1$  and

$$\dim_{\mathbb{C}}(F_{i-1}) = \binom{x}{i-1}.$$

Then we have that

$$\dim_{\mathbb{C}}(F_i) \leq \binom{x+1}{i}.$$

In particular, we also obtain that

$$\frac{\dim(F_i)}{\dim(F_{i-1})} \leq \frac{\binom{x+1}{i}}{\binom{x}{i-1}} = \frac{x+1}{i}. \tag{1}$$

On the other hand, we have  $\dim(F_1) = \binom{\dim(F_1)}{1}$ , and the function  $X \mapsto \binom{X}{k}$  is increasing for  $X \geq k - 1$  (see [7, Lemma 5.6]). Thus if we apply [7, Theorem 5.10]  $i - 2$  times starting with  $\dim(F_1) = \binom{\dim(F_1)}{1}$ , then we obtain

$$\dim_{\mathbb{C}}(F_{i-1}) \leq \binom{\dim(F_1) + i - 2}{i-1},$$

implying that  $x < \dim(F_1) + i - 1$ . Together with (1), this shows that

$$\frac{\dim(F_i)}{\dim(F_{i-1})} < \frac{i + \dim(F_1)}{i},$$

so the proposition follows since

$$\dim(F_i) = \dim(F_i/F_{i-1}) + \dim(F_{i-1}). \quad \square$$

**Definition 12.** Given an ideal  $\mathfrak{a} \subset \mathbb{C}[X]$ , we introduce a standard filtration  $F_i^{\mathfrak{a}}$ ,  $i \in \mathbb{N}$ , on  $\mathbb{C}[X]/\mathfrak{a}$  by defining  $F_i^{\mathfrak{a}}$  to be the space of all those elements of  $\mathbb{C}[X]/\mathfrak{a}$  which can be written as  $f + \mathfrak{a}$  with  $\deg(f) \leq i$ .

Applying Proposition 11 to the filtration  $F_i^{\mathfrak{a}}$ , we obtain the following Corollary.

**Corollary 13.** *Let  $\mathfrak{a} \subset \mathbb{C}[X]$  be an ideal. Then for any  $i > 0$  we have*

$$\dim_{\mathbb{C}}(F_i^{\mathfrak{a}}/F_{i-1}^{\mathfrak{a}}) \leq \frac{n}{i} \dim_{\mathbb{C}}(F_{i-1}^{\mathfrak{a}}).$$

We now proceed to derive some properties of multiplication operators restricted to the spaces  $F_i^{\mathfrak{a}}$ ,  $i \in \mathbb{N}$ . We start with a simple consequence of Hilbert’s Nullstellensatz. When for some  $k \in \mathbb{N}_+$  we consider the space  $\mathbb{C}^k$  as a  $\mathbb{C}$ -algebra, it is meant to be with the pointwise multiplication.

**Proposition 14.** *Let  $\mathfrak{a} \subset \mathbb{C}[X]$  be an ideal and let  $V \subset \mathbb{C}[X]$  be a finite dimensional  $\mathbb{C}$ -linear subspace with the property that  $V \cap \text{rad}(\mathfrak{a}) = \{0\}$ . Then there exists a surjective algebra homomorphism  $\sigma: \mathbb{C}[X] \rightarrow \mathbb{C}^{\dim(V)}$  which is injective on  $V$  and such that  $\mathfrak{a} \subset \ker(\sigma)$ .*

*Proof.* We prove by induction on  $\dim(V)$  the following statement: there exist distinct maximal ideals  $\mathfrak{m}_1, \dots, \mathfrak{m}_{\dim(V)}$  such that for all  $i$  we have  $\text{rad}(\mathfrak{a}) \subset \mathfrak{m}_i$  and

$$V \cap \mathfrak{m}_1 \cap \dots \cap \mathfrak{m}_{\dim(V)} = \{0\}.$$

For the case  $\dim(V) = 1$  let us first choose a non-zero element  $v \in V$ . Now since  $v \notin \text{rad}(\mathfrak{a})$  and  $\text{rad}(\mathfrak{a})$  is equal to an intersection of maximal ideals, we can find a maximal ideal  $\mathfrak{m}_1$  such that  $\text{rad}(\mathfrak{a}) \subset \mathfrak{m}_1$  and  $v \notin \mathfrak{m}_1$ .

Let us therefore assume that we know the inductive statement when  $\dim(V) = k$  for some  $k$  and let us fix  $V$  such that  $\dim(V) = k + 1$ . Let  $W \subset V$  be a  $k$ -dimensional subspace. By the inductive assumption we can find  $\mathfrak{m}_1, \dots, \mathfrak{m}_k$  such that  $W \cap \mathfrak{m}_1 \cap \dots \cap \mathfrak{m}_k = \{0\}$ . Thus the intersection  $V \cap \mathfrak{m}_1 \cap \dots \cap \mathfrak{m}_k$  is at most one-dimensional. It cannot be zero-dimensional because the composition  $V \hookrightarrow \mathbb{C}[X] \rightarrow \mathbb{C}[X]/(\mathfrak{m}_1 \cap \dots \cap \mathfrak{m}_k)$  has a non-trivial kernel, since the Chinese remainder theorem implies that the right-hand side is isomorphic to  $\mathbb{C}^k$ .

Thus the intersection  $V \cap \mathfrak{m}_1 \cap \dots \cap \mathfrak{m}_k$  is one-dimensional. Let  $v$  be a non-zero element of  $V \cap \mathfrak{m}_1 \cap \dots \cap \mathfrak{m}_k$ . Since  $v \notin \text{rad}(\mathfrak{a})$  and  $\text{rad}(\mathfrak{a})$  is equal to an intersection of maximal ideals, we can find a maximal ideal  $\mathfrak{m}_{k+1}$  such that  $\text{rad}(\mathfrak{a}) \subset \mathfrak{m}_{k+1}$  and  $v \notin \mathfrak{m}_{k+1}$ . Thus  $V \cap \mathfrak{m}_1 \cap \dots \cap \mathfrak{m}_k \cap \mathfrak{m}_{k+1} = \{0\}$ , finishing the proof of the inductive claim.

Now we can define  $\sigma$  as being the quotient map  $\mathbb{C}[X] \rightarrow \mathbb{C}[X]/(\mathfrak{m}_1 \cap \dots \cap \mathfrak{m}_{\dim(V)})$ . This finishes the proof. □

**Definition 15.** Given an ideal  $\mathfrak{a} \subset \mathbb{C}[X]$  we denote by  $\mu_i^\mathfrak{a}: \mathbb{C}[X]/\mathfrak{a} \rightarrow \mathbb{C}[X]/\mathfrak{a}$  the linear map defined by  $\mu_i^\mathfrak{a}(f) = X_i \cdot f$ .

**Corollary 16.** *Let  $\mathfrak{a} \subset \mathbb{C}[X]$  be an ideal and let  $R \in \mathbb{N}$  be such that*

$$F_R^\mathfrak{a} \cap (\text{rad}(\mathfrak{a})/\mathfrak{a}) = \{0 + \mathfrak{a}\}.$$

*Then there exist simultaneously diagonalisable linear maps*

$$M_1, \dots, M_n: F_R^\mathfrak{a} \longrightarrow F_R^\mathfrak{a}$$

*such that for  $v \in F_{R-1}^\mathfrak{a}$  and all  $i = 1, \dots, n$  we have  $M_i(v) = \mu_i^\mathfrak{a}(v)$ .*

*Proof.* Let  $d := \dim(F_R^\mathfrak{a})$ , and let  $f_1, \dots, f_d \in \mathbb{C}[X]$  be such that  $f_i + \mathfrak{a}$  is a basis of  $F_R^\mathfrak{a}$ . Let  $V \subset \mathbb{C}[X]$  be the linear span of the elements  $f_i$ . Let us observe that  $V \cap \text{rad}(\mathfrak{a}) = \{0\}$ . Indeed, if  $f \in V \cap \text{rad}(\mathfrak{a})$  then  $f + \mathfrak{a} \in F_R^\mathfrak{a} \cap \text{rad}(\mathfrak{a})/\mathfrak{a}$  and so by assumption we see that  $f \in \mathfrak{a}$ .

Hence by the previous proposition we can find a surjective algebra homomorphism

$$\sigma: \mathbb{C}[X]/\mathfrak{a} \longrightarrow \mathbb{C}^d$$

such that  $\sigma$  is injective on  $F_R^\mathfrak{a}$ . Let  $\tau: \mathbb{C}^d \rightarrow F_R^\mathfrak{a}$  be the unique linear isomorphism such that for  $v \in F_R^\mathfrak{a}$  we have  $\tau(\sigma(v)) = v$ . Since  $\sigma$  is surjective, it follows that for all  $v \in \mathbb{C}^d$  we have  $\sigma(\tau(v)) = v$ .

For  $v \in F_R^\mathfrak{a}$  let us define

$$M_i(v) := \tau(\sigma(X_i \cdot v)).$$

If  $v \in F_{R-1}^\mathfrak{a}$  then  $X_i \cdot v \in F_R^\mathfrak{a}$  and so  $M_i(v) = \tau(\sigma(X_i \cdot v)) = X_i \cdot v = \mu_i^\mathfrak{a}(v)$ . Thus in order to finish the proof we only need to check that the maps  $M_i$  are simultaneously diagonalisable.

Let  $e_1, \dots, e_d$  be the standard basis of  $\mathbb{C}^d$ . In particular  $\tau(e_1), \dots, \tau(e_d)$  is a basis of  $F_R^\mathfrak{a}$ , and we claim that for every  $i \in \{1, \dots, n\}$  we have that the vectors  $\tau(e_j)$ ,  $j = 1, \dots, d$ , are eigenvectors for  $M_i$ . Indeed, first we note that for every  $i$  and  $j$  we have that  $\sigma(X_i) \cdot e_j$  is a multiple of  $e_j$ , and so we can define numbers  $\lambda_{ij} \in \mathbb{C}$  by the formula

$$\sigma(X_i) \cdot e_j = \lambda_{ij} e_j.$$

Now we can write

$$\begin{aligned} M_i(\tau(e_j)) &= \tau(\sigma(X_i \cdot \tau(e_j))) = \tau(\sigma(X_i) \cdot \sigma(\tau(e_j))) \\ &= \tau(\sigma(X_i) \cdot e_j) = \tau(\lambda_{ij} e_j) = \lambda_{ij} \tau(e_j), \end{aligned}$$

finishing the proof. □

### 4. Proof of Theorem 6

We will first prove several lemmas. The first lemma, informally speaking, allows us to deduce Theorem 6 provided that we can construct large subspaces by “growing balls around points”. To make it precise we state a few definitions.

Given two positive natural numbers  $a, b$  we let  $\text{Map}(a, b)$  be the set of all maps from  $[a]$  to  $[b]$ , and furthermore we let  $\text{Map}_{\leq}(a, b) := \bigcup_{i=0}^a \text{Map}(i, b)$ .

Let  $W$  be a  $\mathbb{C}$ -vector space and let  $\mathcal{M} = (M_1, \dots, M_n)$  be a tuple of endomorphisms of  $W$ . Given  $R \in \mathbb{N}$  and  $\alpha \in \text{Map}(R, n)$  we let

$$\mathcal{M}_\alpha := M_{\alpha(1)} \cdots M_{\alpha(R)}.$$

Note that the unique element of  $\text{Map}(0, n)$  is the empty set. Our convention is that  $\mathcal{M}_\emptyset$  is the identity map.

Given  $w \in W$  we let  $B_{\mathcal{M}}(w, R)$  to be the linear span of the vectors  $\mathcal{M}_\alpha(w)$ , where  $\alpha \in \text{Map}_{\leq}(R, n)$ . We will call  $B_{\mathcal{M}}(w, R)$  the  $R$ -ballspace for  $\mathcal{M}$  with

root  $w$ . If  $\mathcal{M}$  is clear from the context, then we denote  $B_{\mathcal{M}}(w, R)$  simply with  $B(w, R)$ .

Recall from Definition 15 that given an ideal  $\mathfrak{a} \subset \mathbb{C}[X]$  and  $i \in [n]$ , we denote by  $\mu_i^{\mathfrak{a}}: \mathbb{C}[X]/\mathfrak{a} \rightarrow \mathbb{C}[X]/\mathfrak{a}$  the linear map defined by  $\mu_i^{\mathfrak{a}}(f + \mathfrak{a}) = (X_i \cdot f) + \mathfrak{a}$ . Note that the  $R$ -ballspace for  $(\mu_1^{\mathfrak{a}}, \dots, \mu_n^{\mathfrak{a}})$  with root  $1 + \mathfrak{a} \in \mathbb{C}[X]/\mathfrak{a}$  is equal to  $F_R^{\mathfrak{a}}$ .

In general we will say that  $B_{\mathcal{M}}(w, R)$  is *regular* if there exists an ideal  $\mathfrak{a} \subset \mathbb{C}[X]$  and a linear isomorphism  $\varphi: B_{\mathcal{M}}(w, R) \rightarrow F_R^{\mathfrak{a}}$  such that the following two conditions hold.

- (1) For every  $v \in B_{\mathcal{M}}(w, R - 1)$  and  $i \in [n]$  we have that  $\varphi(v) \in F_{R-1}^{\mathfrak{a}}$  and  $\varphi(M_i(v)) = \mu_i^{\mathfrak{a}}(\varphi(v))$ .
- (2) If for some  $f \in \mathbb{C}[X]$  and  $m \in \mathbb{N}_+$  we have  $f + \mathfrak{a} \in F_R^{\mathfrak{a}}$  and  $f^m \in \mathfrak{a}$  then  $f \in \mathfrak{a}$ . In other words we have  $F_R^{\mathfrak{a}} \cap (\text{rad}(\mathfrak{a})/\mathfrak{a}) = \{0 + \mathfrak{a}\}$ .

Let  $d \in \mathbb{N}$  and let  $(M_1, \dots, M_n)$  be a  $d$ -matrix tuple. Given  $R \in \mathbb{N}$  and a subspace  $W \subset \mathbb{C}^d$  we say that  $W$  is an  $R$ -multi-ballspace if there exist  $w_1, \dots, w_k \in W$  and natural numbers  $R_1, \dots, R_k$  with  $R_j \geq R$  such that

- (1) the ballspaces  $B(w_j, R_j)$  are regular, and
- (2)  $W$  is equal to the direct sum  $\bigoplus_{j=1}^k B(w_j, R_j)$ .

The *roots* of such  $W$  are the points  $\{w_1, \dots, w_k\}$ .

If all elements of a matrix tuple  $\mathcal{A}$  can be diagonalised simultaneously, then  $\mathcal{A}$  will be called *simultaneously diagonalisable*. Clearly, if  $\mathcal{A}$  is a simultaneously diagonalisable tuple then it is also a commuting tuple.

**Lemma 17.** *For every  $\varepsilon > 0$  there exists  $R \in \mathbb{N}$  and  $\delta > 0$  such that the following holds. Suppose that  $d \in \mathbb{N}$ , let  $\mathcal{M}$  be a  $d$ -matrix tuple, and let  $W \subset \mathbb{C}^d$  be an  $R$ -multi-ballspace with  $\dim(W) \geq (1 - \delta) \cdot d$ .*

*Then there exists a simultaneously diagonalisable  $d$ -matrix tuple  $\mathcal{A}$  such that*

$$d_{\text{rank}}(\mathcal{M}, \mathcal{A}) \leq \varepsilon.$$

*Proof.* Let  $R$  be such that  $\frac{n}{R} < \frac{\varepsilon}{2}$  and let  $\delta$  be such that  $\delta < \frac{\varepsilon}{2}$ . Let  $\mathcal{M} = (M_1, \dots, M_n)$  be the  $d$ -matrix tuple, let  $w_1, \dots, w_k \in \mathbb{C}^d$  and let  $R_1, \dots, R_k \in \mathbb{N}$  be such that  $R_i \geq R$  and such that the ballspaces  $B(w_j, R_j)$  are regular and  $W = \bigoplus_{j=1}^k B(w_j, R_j)$ .

We need to find a simultaneously diagonalisable tuple  $\mathcal{A} = (A_1, \dots, A_n)$  such that  $d_{\text{rank}}(\mathcal{M}, \mathcal{A}) \leq \varepsilon$ .

For every  $j = 1, \dots, k$ , let  $\varphi_j: B(w_j, R_j) \rightarrow F_{R_j}^{\mathfrak{a}_j}$  be the linear isomorphism witnessing the regularity of the ballspace  $B(w_j, R_j)$ .

By Corollary 16, for every  $j = 1, \dots, k$  we can find maps  $M_{ij}: F_{R_j}^{\mathfrak{a}_j} \rightarrow F_{R_j}^{\mathfrak{a}_j}$ , where  $i = 1, \dots, n$ , such that the maps  $M_{1j}, \dots, M_{nj}$  pairwise commute, are

simultaneously diagonalisable, and for  $v \in B(w_j, R_j - 1)$  we have  $M_i(v) = \varphi_j^{-1} \cdot M_{ij} \cdot \varphi_j(v)$ .

Let us fix a projection  $\pi: \mathbb{C}^d \rightarrow \bigoplus_{j=1}^k B(w_j, R_j)$ , and for  $i = 1, \dots, n$  let

$$A_i := \left( \bigoplus_j \varphi_j^{-1} \cdot M_{ij} \cdot \varphi_j \right) \cdot \pi.$$

It is clear that the maps  $A_1, \dots, A_n$  are simultaneously diagonalisable. Also for every  $v \in \bigoplus_{j=1}^k B(w_j, R_j - 1)$  we have  $A_i(v) = M_i(v)$ , so

$$d_{\text{rank}}(\mathcal{M}, \mathcal{A}) \leq \frac{1}{d} \left( \dim(\ker(\pi)) + \sum_{j=1}^k (\dim(B(w_j, R_j)) - \dim(B(w_j, R_j - 1))) \right).$$

Since the ballspaces  $B(w_j, R_j)$  are regular, by Proposition 11 we have

$$\begin{aligned} \dim(B(w_j, R_j)) - \dim(B(w_j, R_j - 1)) &\leq \frac{n}{R_j} \cdot \dim(B(w_j, R_j - 1)) \\ &\leq \frac{n}{R} \cdot \dim(B(w_j, R_j)). \end{aligned}$$

Hence we see that

$$\sum_{j=1}^k (\dim(B(w_j, R_j)) - \dim(B(w_j, R_j - 1))) \leq \frac{n}{R} \sum_{j=1}^k \dim(B(w_j, R_j)) < \frac{\varepsilon}{2} \cdot d.$$

Thus altogether we have

$$d_{\text{rank}}(\mathcal{M}, \mathcal{A}) < \frac{\varepsilon}{2} + \delta \leq \varepsilon,$$

finishing the proof. □

Given  $r, d \in \mathbb{N}$ , a  $d$ -matrix tuple  $\mathcal{M} = (M_1, \dots, M_n)$ , and  $v \in \mathbb{C}^d$ , we say that  $\mathcal{M}$  is  $r$ -commutative on  $v$  if for any  $i \leq r$ , any  $\alpha \in \text{Map}(i, n)$  and any permutation  $\sigma: [i] \rightarrow [i]$  we have

$$\mathcal{M}_\alpha(v) = \mathcal{M}_{\alpha \circ \sigma}(v).$$

If  $W \subset \mathbb{C}^d$  then we say that  $\mathcal{M}$  is  $r$ -commutative on  $W$  if for every  $v \in W$  we have that  $\mathcal{M}$  is  $r$ -commutative on  $v$ .

**Lemma 18.** *For every  $R \in \mathbb{N}$  and  $\varepsilon > 0$  there exists  $\eta > 0$  such that if  $d \in \mathbb{N}$  and  $\mathcal{M}$  is an  $\eta$ -commuting  $d$ -matrix tuple, then there exists a subspace  $W \subset \mathbb{C}^d$  such that  $\mathcal{M}$  is  $R$ -commutative on  $W$  and  $\dim(W) \geq d(1 - \varepsilon)$ .*

*Proof.* For  $k \in \mathbb{N}$  we let  $\text{Bij}(k)$  be the set of all bijections of the set  $[k]$ . Let us prove by induction on  $R$  that for every  $\varepsilon > 0$  there exists  $\eta > 0$  such that if  $d \in \mathbb{N}$  and  $\mathcal{M}$  is an  $\eta$ -commuting  $d$ -matrix tuple, then

$$\dim \left( \bigcap_{\substack{\alpha \in \text{Map}(R, n) \\ \sigma \in \text{Bij}(R)}} \ker(\mathcal{M}_\alpha - \mathcal{M}_{\alpha \circ \sigma}) \right) \geq d(1 - \varepsilon).$$

When  $R = 2$ , we can set  $\eta := \frac{\varepsilon}{n^2}$ . Indeed, if  $\mathcal{M}$  is an  $\eta$ -commuting tuple then by definition for  $i, j \in [n]$  we have  $\dim(\ker([M_i, M_j])) \geq (1 - \eta)d$ . And so we have

$$\dim \left( \bigcap_{i, j \in [n]} \ker([M_i, M_j]) \right) \geq (1 - n^2\eta)d = (1 - \varepsilon)d.$$

Thus let us assume that we have shown the inductive statement for some  $R$  and let us prove it for  $R + 1$ . Let us fix  $\varepsilon > 0$  and let  $\eta$  be given by the inductive assumption for  $\frac{\varepsilon}{n+1}$ . Thus given an  $\eta$ -commuting tuple  $\mathcal{M}$  we obtain a subspace  $W \subset \mathbb{C}^d$  such that  $\dim(W) \geq d(1 - \frac{\varepsilon}{n+1})$  and  $\mathcal{M}$  is  $R$ -commutative on  $W$ , i.e. for any  $w \in W$ , any  $\alpha \in \text{Map}(k, n)$  with  $k \leq R$  and any permutation  $\sigma: [k] \rightarrow [k]$  we have

$$\mathcal{M}_\alpha(w) = \mathcal{M}_{\alpha \circ \sigma}(w).$$

Let us define  $V := W \cap \bigcap_{i=1}^n M_i^{-1}(W)$ . Clearly  $\dim(V) \geq d(1 - \varepsilon)$ .

Now let  $\beta \in \text{Map}(R + 1, n)$ , let  $\tau: [R + 1] \rightarrow [R + 1]$  be a permutation, and let  $v \in V$ . Let  $i \in [n]$  be such that for some  $2 \leq j_1 \leq R + 1$  we have  $\beta(j_1) = i$  and for some  $2 \leq j_2 \leq R + 1$  we have  $\beta(\tau(j_2)) = i$ . We can find such  $i$  because  $R + 1 \geq 3$ .

Since in particular  $v \in W$ , we can find  $\gamma \in \text{Map}(R, n)$  and a permutation  $\rho: [R] \rightarrow [R]$  such that

$$\mathcal{M}_\beta(v) = \mathcal{M}_\gamma \cdot M_i(v)$$

and

$$\mathcal{M}_{\beta \circ \tau}(v) = \mathcal{M}_{\gamma \circ \rho} \cdot M_i(v).$$

Since  $M_i(v) \in W$ , we have

$$\mathcal{M}_{\gamma \circ \rho} \cdot M_i(v) = \mathcal{M}_\gamma \cdot M_i(v),$$

which finishes the proof. □

**Definition 19.** If  $r \in \mathbb{N}$ ,  $\varepsilon > 0$ , and  $A$  and  $B$  are subspaces of  $\mathbb{C}^d$ , then we say that  $(A, B)$  is an  $(r, \varepsilon)$ -pair for the  $d$ -tuple  $\mathcal{M} = (M_1, M_2, \dots, M_n)$  if

- (1)  $A \subset B$  and  $\dim(B/A) \leq \varepsilon \cdot d$ , and
- (2) for every  $\alpha \in \text{Map}_{\leq}(r, n)$  and  $v \in A$  we have  $\mathcal{M}_\alpha(v) \in B$ .

**Lemma 20.** *Let  $\varepsilon > 0$ , let  $r, R, d \in \mathbb{N}$  with  $n \leq r < R$ , and let  $\mathcal{M} = (M_1, \dots, M_n)$  be a  $*$ -closed  $d$ -matrix tuple. Let  $(A_1, B_1)$  be an  $(R, \varepsilon)$ -pair and let  $W = \bigoplus_{j=1}^k B(w_j, R_j)$  be an  $R$ -multi-ballspace for  $\mathcal{M}$  contained in  $B_1$ . Furthermore let us assume that the ballspace  $B(w_j, R_j + r)$  are regular for all  $j$ . Finally, let  $B_2$  be the orthogonal complement of  $W$  in  $B_1$ .*

*Then there exists a subspace  $A_2 \subset A_1 \cap B_2$  such that  $(A_2, B_2)$  is an  $(r, \varepsilon + \frac{n}{R}2^r)$ -pair.*

*Proof.* Let  $V \subset \mathbb{C}^d$  be the space spanned by the ballspace  $B(w_j, R_j + r)$ ,  $j = 1, \dots, k$ . Let  $V^\perp \subset \mathbb{C}^d$  be the space orthogonal to  $V$  and let  $A_2 = A_1 \cap V^\perp$ .

Let us show that  $\dim(B_2/A_2) \leq d(\varepsilon + \frac{n}{R}2^r)$ . By basic linear algebra, it is easy to check that  $\dim(B_2/A_2)$  is bounded from above by

$$\dim(B_1/A_1) + \dim(V) - \dim(W). \tag{2}$$

We can bound (2) by

$$\dim(B_1/A_1) + \sum_{j=1}^k (\dim(B(w_j, R_j + r)) - \dim(B(w_j, R_j))).$$

By Corollary 13, the quantity above is at most

$$\varepsilon \cdot d + \sum_{j=1}^k \dim(B(w_j, R_j)) \left( \left(1 + \frac{n}{R}\right)^r - 1 \right) \leq \varepsilon \cdot d + d \cdot \frac{n}{R} \cdot 2^r,$$

where we use the inequality  $(1 + x)^r \leq 1 + (2^r - 1)x$ , valid for  $x \in [0, 1]$  and  $r \geq 1$ . Therefore we obtain that  $\dim(B_2/A_2) \leq d(\varepsilon + \frac{n}{R}2^r)$ .

Thus to finish the proof we only need to show that for  $x \in A_2$  and  $\alpha \in \text{Map}(q, n)$  with  $q \leq r < R$  we have  $\mathcal{M}_\alpha(x) \in B_2$ . In other words, we need to show that if  $x \in A_1$  is orthogonal to  $B(w_j, R_j + r)$  then  $\mathcal{M}_\alpha(x)$  is orthogonal to  $B(w_j, R_j)$ .

Indeed, let  $w \in B(w_j, R_j)$ . Since  $\mathcal{M}$  is  $*$ -closed,  $\mathcal{M}_\alpha^*(w) \in B(w_j, R_j + r)$  and hence

$$\langle \mathcal{M}_\alpha(x), w \rangle = \langle x, \mathcal{M}_\alpha^*(w) \rangle = 0,$$

finishing the proof. □

Let  $\mathbb{C}\langle Y_1, \dots, Y_n \rangle$  be the ring of polynomials in  $n$  non-commuting variables, and let  $\pi: \mathbb{C}\langle Y_1, \dots, Y_n \rangle \rightarrow \mathbb{C}[X]$  be the algebra homomorphism such that  $\pi(Y_i) = X_i$ . Let  $c: \mathbb{C}[X] \rightarrow \mathbb{C}\langle Y_1, \dots, Y_n \rangle$  be the unique  $\mathbb{C}$ -linear map such that for any  $\alpha \in \text{Map}(d, n)$  with  $\alpha(1) \leq \alpha(2) \leq \dots \leq \alpha(d)$  we have

$$c \circ \pi(Y_{\alpha(1)} \dots Y_{\alpha(d)}) = Y_{\alpha(1)} \dots Y_{\alpha(d)}.$$

In other words, the map  $c$  allows us to treat commutative polynomials as non-commutative ones, by fixing an order on the variables. Given a matrix tuple  $\mathcal{M} = (M_1, \dots, M_n)$  and  $f \in \mathbb{C}[X]$ , we define  $f(\mathcal{M})$  to be the matrix  $c(f)(M_1, \dots, M_n)$ .

Let us define a  $*$ -operation on  $\mathbb{C}\langle Y_1, \dots, Y_n \rangle$  in the following way. For any  $\alpha \in \text{Map}(d, n)$  and  $a \in \mathbb{C}$  we define

$$(a \cdot Y_{\alpha(1)} \cdots Y_{\alpha(d)})^* := \bar{a} \cdot Y_{\alpha(d)} \cdots Y_{\alpha(1)},$$

and we extend this definition to arbitrary elements of  $\mathbb{C}\langle Y_1, \dots, Y_n \rangle$  by linearity.

For  $f \in \mathbb{C}[X]$  we define  $f^*(\mathcal{M}) := (c(f))^*(M_1, \dots, M_n)$ . The following simple observation will be used without reference.

**Lemma 21.** *For any matrix tuple  $\mathcal{M}$  and any  $f \in \mathbb{C}[X]$  we have that the matrices  $f(\mathcal{M})$  and  $f^*(\mathcal{M}^*)$  are adjoint to each other.  $\square$*

We will also need the following lemma.

**Lemma 22.** *Let  $f \in \mathbb{C}[X]$  and let  $k \in \mathbb{N}$ . Let  $\mathcal{M}$  be a  $*$ -closed  $d$ -matrix tuple and let  $x \in \mathbb{C}^d$  be such that  $\mathcal{M}$  is  $(2k \deg(f))$ -commutative at  $x$ . Then*

$$[f^*(\mathcal{M}^*)f(\mathcal{M})]^k(x) = [f^*(\mathcal{M}^*)]^k[f(\mathcal{M})]^k(x)$$

*Proof.* For  $f \in \mathbb{C}[X]$  let us define  $\bar{f}$  to be the polynomial which arises from  $f$  by conjugating the coefficients. Then the left-hand side is equal to

$$[\bar{f}(\mathcal{M}^*)f(\mathcal{M})]^k(x),$$

and the right-hand side is equal to

$$[\bar{f}(\mathcal{M}^*)]^k[f(\mathcal{M})]^k(x).$$

These two expressions are clearly equal if  $\mathcal{M}$  is  $(2k \deg(f))$ -commutative at  $x$ .  $\square$

Recall that  $K(R)$  is a function defined in Theorem 9.

**Lemma 23.** *Let  $d \in \mathbb{N}_+$  and let  $\mathcal{M} = (M_1, \dots, M_n)$  be a  $*$ -closed  $d$ -matrix tuple. Let  $R \geq 0$  and let  $v \in \mathbb{C}^d$  be such that  $\mathcal{M}$  is  $(2K(R))$ -commutative at  $v$ . Then the ballspace  $B(v, R)$  is regular.*

*Proof.* Given  $\alpha \in \text{Map}(q, n)$ , we define  $X_\alpha \in \mathbb{C}[X]$  to be the monomial

$$X_\alpha := X_{\alpha(1)} \cdots X_{\alpha(q)}.$$



Let  $P \subset \mathbb{C}[X]$  be defined as follows. We let  $f \in P$  if and only if  $\deg(f) \leq R$  and  $f(\mathcal{M})(v) = 0$ . Let  $\mathfrak{a}$  be the ideal generated by  $P$ . Let us define a map  $\varphi: B(v, R) \rightarrow \mathbb{C}[X]/\mathfrak{a}$  as follows:

$$\varphi(\mathcal{M}_\alpha(v)) := X_\alpha + \mathfrak{a}.$$

Let us check that  $\varphi$  is well-defined. For this let us assume that

$$\sum_{\alpha \in \text{Map}_{\leq}(R, n)} s_\alpha \mathcal{M}_\alpha(v) = \sum_{\alpha \in \text{Map}_{\leq}(R, n)} t_\alpha \mathcal{M}_\alpha(v),$$

where  $s_\alpha, t_\alpha \in \mathbb{C}$ . But then

$$\sum_{\alpha \in \text{Map}_{\leq}(R, n)} (s_\alpha - t_\alpha) \mathcal{M}_\alpha(v) = 0,$$

and therefore  $\sum_{\alpha \in \text{Map}_{\leq}(R, n)} (s_\alpha - t_\alpha) X_\alpha \in P$ . In particular we get that

$$\sum_{\alpha \in \text{Map}_{\leq}(R, n)} s_\alpha X_\alpha + \mathfrak{a} = \sum_{\alpha \in \text{Map}_{\leq}(R, n)} t_\alpha X_\alpha + \mathfrak{a},$$

which shows that  $\varphi$  is well-defined.

Now let us see that  $\varphi$  is injective. Indeed suppose that

$$\varphi\left(\sum_{\alpha} s_\alpha \mathcal{M}_\alpha(v)\right) = 0,$$

where  $\alpha$  runs through the elements of  $\text{Map}_{\leq}(R, n)$ , and  $s_\alpha \in \mathbb{C}$ . Then  $\sum_{\alpha} s_\alpha X_\alpha \in \mathfrak{a}$ , and so we can find  $f_i \in P$  and  $h_i \in \mathbb{C}[X]$  with  $\deg(h_i f_i) \leq K(R)$  such that

$$\sum_{i=1}^k h_i f_i = \sum_{\alpha} s_\alpha X_\alpha.$$

But since  $\mathcal{M}$  is  $K(R)$ -commutative at  $v$ , we have

$$\sum_{i=1}^k h_i(\mathcal{M}) f_i(\mathcal{M})(v) = \sum_{\alpha} s_\alpha \mathcal{M}_\alpha(v).$$

The left-hand side is equal to 0 since  $f_i \in P$ , and so we see that  $\sum_{\alpha} s_\alpha \mathcal{M}_\alpha(v) = 0$ . This finishes the proof of injectivity of  $\varphi$ .

Since clearly the image of  $\varphi$  is equal to  $F_R^{\mathfrak{a}}$ , it remains to prove that

$$F_R^{\mathfrak{a}} \cap (\text{rad}(\mathfrak{a})/\mathfrak{a}) = \{0 + \mathfrak{a}\}.$$

By Theorem 9, if  $f \in \mathbb{C}[X]$  is such that  $\deg(f) \leq R$  and  $f \in \text{rad}(\mathfrak{a})$  then we can find  $m \in \mathbb{N}$ , elements  $f_i \in P$  and  $g_i \in \mathbb{C}[X]$  with  $\deg(g_i f_i) \leq K(R)$ , such that

$$f^m = \sum_{i=1}^k g_i f_i,$$

Since  $\mathcal{M}$  is  $K(R)$ -commutative at  $v$ , we have

$$0 = \sum_{i=1}^k g_i(\mathcal{M}) f_i(\mathcal{M})(v) = f(\mathcal{M})^m(v),$$

and so by  $2K(R)$ -commutativity, and since  $R \leq K(R)$ , we also have

$$0 = f(\mathcal{M})^m(v) = f^*(\mathcal{M}^*)^m f(\mathcal{M})^m(v) = (f^*(\mathcal{M}^*) f(\mathcal{M}))^m(v).$$

In particular we can define  $t$  to be the smallest positive integer such that

$$(f^*(\mathcal{M}^*) f(\mathcal{M}))^t(v) = 0.$$

We will show  $t = 1$ . By way of contradiction, let us consider two cases: first let us assume that  $t$  is even and equal to  $2l$  for some  $l \geq 1$ .

By Lemma 22, we have

$$0 = f^*(\mathcal{M}^*)^{2l} f(\mathcal{M})^{2l}(v) = (f^*(\mathcal{M}^*) f(\mathcal{M}))^{2l}(v). \tag{3}$$

Therefore, we also have

$$\begin{aligned} 0 &= \langle (f^*(\mathcal{M}^*) f(\mathcal{M}))^{2l}(v), v \rangle \\ &= \langle (f^*(\mathcal{M}^*) f(\mathcal{M}))^l(v), (f^*(\mathcal{M}^*) f(\mathcal{M}))^l(v) \rangle. \end{aligned}$$

This shows that  $(f^*(\mathcal{M}^*) f(\mathcal{M}))^l(v) = 0$ , which contradicts the minimality of  $t$ .

In the second case, let us assume that  $t$  is odd and equal to  $2l + 1$  for some  $l \geq 1$ . We proceed in a similar fashion. By Lemma 22 we have that  $(f^*(\mathcal{M}^*) f(\mathcal{M}))^{2l+1}(v) = 0$ . Hence, we also have

$$\begin{aligned} 0 &= \langle (f^*(\mathcal{M}^*) f(\mathcal{M}))^{2l+1}(v), f^*(\mathcal{M}^*) f(\mathcal{M})(v) \rangle \\ &= \langle (f^*(\mathcal{M}^*) f(\mathcal{M}))^{l+1}(v), (f^*(\mathcal{M}^*) f(\mathcal{M}))^{l+1}(v) \rangle, \end{aligned}$$

and since  $l + 1 < t$ , we obtain a contradiction exactly as in the first case.

Thus all in all we have showed that  $f^*(\mathcal{M}^*) f(\mathcal{M})(v) = 0$ . Since  $f^*(\mathcal{M}^*)$  and  $f(\mathcal{M})$  are adjoint to each other, we also have that  $f(\mathcal{M})(v) = 0$ . This shows that  $f \in P$ , and hence  $f \in \mathfrak{a}$ , which finishes the proof.  $\square$

**Lemma 24.** *Let  $R \in \mathbb{N}$ ,  $d \in \mathbb{N}$ , let  $\mathcal{M}$  be a  $*$ -closed  $d$ -matrix tuple, let  $A \subset \mathbb{C}^d$ , and let us assume that  $\mathcal{M}$  is  $2K(2R)$ -commutative on  $A$ . Then there exist  $k \in \mathbb{N}$  and  $w_1, \dots, w_k \in A$  such that the  $R$ -ballspaces  $B_{\mathcal{M}}(w_i, R)$  are regular, pairwise orthogonal, and we have that*

$$\sum_{i=1}^k \dim(B_{\mathcal{M}}(w_i, R)) \geq \frac{1}{e^n} \cdot \dim(A), \tag{4}$$

where  $e = 2.71 \dots$

*Proof.* Note that by Lemma 23 all  $R$ -ballspaces with roots in  $A$  are regular. Let us consider the subset  $Q$  of  $A^{\oplus \mathbb{N}}$  which consists of those tuples  $(w_1, \dots, w_k)$  such that the ballspaces  $B_{\mathcal{M}}(w_1, R), \dots, B_{\mathcal{M}}(w_k, R)$  are pairwise orthogonal to each other.

Let  $(w_1, \dots, w_k) \in Q$  be a tuple for which the number

$$\sum_{i=1}^k \dim(B_{\mathcal{M}}(w_i, R))$$

is maximal. By Lemma 23, it is enough to show that  $\sum_{i=1}^k \dim(B_{\mathcal{M}}(w_i, R)) \geq \frac{1}{e^n} \cdot \dim(A)$ . Consider the vector space  $V$  spanned by the ballspaces  $B_{\mathcal{M}}(w_i, 2R)$ . By Lemma 23, the ballspaces  $B_{\mathcal{M}}(w_i, 2R)$  are regular, and so by Corollary 13 we have that

$$\begin{aligned} \dim(B_{\mathcal{M}}(w_i, 2R)) &\leq \\ &\leq \left(1 + \frac{n}{2R}\right) \left(1 + \frac{n}{2R-1}\right) \dots \left(1 + \frac{n}{R+1}\right) \dim(B_{\mathcal{M}}(w_i, R)), \end{aligned}$$

which easily implies that

$$\dim(B_{\mathcal{M}}(w_i, 2R)) \leq e^n \dim(B_{\mathcal{M}}(w_i, R)).$$

This shows that

$$\dim(V) \leq e^n \dim\left(\bigoplus_{i=1}^k B_{\mathcal{M}}(w_i, R)\right). \tag{5}$$

Let us observe that if  $x \in A$  is orthogonal to  $V$  then  $B_{\mathcal{M}}(x, R)$  is orthogonal to the space  $\bigoplus_{i=1}^k B_{\mathcal{M}}(w_i, R)$ . Indeed, since  $\mathcal{M}$  is  $*$ -closed, for any  $\alpha \in \text{Map}_{\leq}(R, n)$  and any  $w \in B_{\mathcal{M}}(w_i, R)$  we have that  $\mathcal{M}_{\alpha}^*(w) \in B_{\mathcal{M}}(w_i, 2R)$ . It follows that

$$\langle \mathcal{M}_{\alpha}(x), w \rangle = \langle x, \mathcal{M}_{\alpha}^*(w) \rangle = 0.$$

But by the maximality of  $(w_1, \dots, w_k)$ , the above shows that there are no points in  $A$  orthogonal to  $V$ , so in fact we have  $A \subset V$ . In particular, we have

$\dim(V) \geq \dim(A)$ , and hence by (5) we have

$$\dim(A) \leq e^n \dim\left(\bigoplus_{i=1}^k B_{\mathcal{M}}(w_i, R)\right),$$

finishing the proof. □

The final lemma which we need for the proof of Theorem 6 is an ‘‘Ornstein–Weiss type’’ lemma.

**Lemma 25.** *For every  $\delta > 0$  and  $r \in \mathbb{N}$  there exists  $\eta > 0$  such that if  $d \in \mathbb{N}$ , and  $\mathcal{M}$  is an  $\eta$ -commuting  $*$ -closed  $d$ -matrix tuple, then there exists an  $r$ -multiball  $W \subset \mathbb{C}^d$  for  $\mathcal{M}$ , such that  $\dim(W) \geq (1 - \delta) \cdot d$ .*

*Proof.* Let us fix  $\delta > 0$  and  $r \in \mathbb{N}$ . Let us first fix a positive integer  $k$  such that  $k(1 - \frac{1}{e^n})^k < \frac{\delta}{4}$ , and then let us choose  $\varepsilon \in (0, \min(\frac{\delta}{4}, \frac{\delta}{2k}, \frac{1}{e^n}))$  and natural numbers  $r_0 > r_1 > \dots > r_k = r$  such that for  $i = 0, \dots, k - 1$  we have

$$\varepsilon + \frac{n}{r_i} 2^{r_{i+1}} < \left(1 - \frac{1}{e^n}\right)^k.$$

By Lemma 18, we can fix  $\eta$  to be such that if  $\mathcal{M}$  is  $\eta$ -commuting then there exists a subspace  $S \subset \mathbb{C}^d$  such that  $\dim(S) \geq (1 - \varepsilon)d$  and  $\mathcal{M}$  is  $2K(2(r_0 + r_1))$ -commutative on  $S$ .

Let  $\bar{d} := (1 - \varepsilon)d$ . We will prove by induction on  $i$  the following statement: for every  $i = 1, \dots, k$  there exists  $g(i) \in \mathbb{N}$ , roots  $w_1, \dots, w_{g(i)} \in S$  and radii  $R_1, \dots, R_{g(i)}$  with  $r_i \leq R_j \leq r_0$  for all  $j = 1, \dots, g(i)$ , such that the balls  $B_{\mathcal{M}}(w_j, R_j)$  are pairwise orthogonal and

$$\sum_{j=1}^{g(i)} \dim(B_{\mathcal{M}}(w_j, R_j)) \geq \bar{d} \left(1 - i \left(1 - \frac{1}{e^n}\right)^i\right) - d \cdot \frac{i\delta}{2k}.$$

This will be enough to finish the proof because for  $i = k$  the right hand side above is equal to

$$(1 - \varepsilon)d \left(1 - k \left(1 - \frac{1}{e^n}\right)^k\right) - d \frac{\delta}{2} > \left(1 - \frac{\delta}{4}\right) \left(1 - \frac{\delta}{4}\right) \cdot d - \frac{\delta}{2}d > (1 - \delta) \cdot d$$

For  $i = 1$  the inductive claim is implied by Lemma 24. Suppose that the inductive claim holds for some  $i \in \{1, \dots, k - 1\}$  and let us prove it for  $i + 1$ .

Let  $W_i = \bigoplus_{j=1}^{g(i)} B_{\mathcal{M}}(w_j, R_j)$  and let  $W_i^\perp$  be the orthogonal complement of  $W_i$  in  $\mathbb{C}^d$ . Since  $\mathcal{M}$  is  $2K(2(r_0 + r_{i+1}))$ -commutative on  $S$ , we have that all the ballspaces

$$B_{\mathcal{M}}(w_j, R_j + r_{i+1})$$

are regular, and so we can apply Lemma 20 for the  $(r_i, \varepsilon)$ -pair  $(S, \mathbb{C}^d)$  and the  $r_i$ -multiballspace  $W_i$ .

As a result we obtain a subspace  $S_i \subset S \cap W_i^\perp$  such that  $(S_i, W_i^\perp)$  is an  $(r_{i+1}, \varepsilon + \frac{n}{r_i}2^{r_{i+1}})$ -pair.

Now, by Lemma 24 we obtain  $g(i + 1) \in \mathbb{N}$  and roots  $w_{g(i)+1}, w_{g(i)+2}, \dots, w_{g(i+1)} \in S_i$ , such that the ballspace  $B_{\mathcal{M}}(w_{g(i)+s}, r_{i+1}), s = 1, \dots, g(i + 1) - g(i)$ , are regular, pairwise orthogonal, and

$$\sum_{s=1}^{g(i+1)-g(i)} \dim(B_{\mathcal{M}}(w_{g(i)+s}, r_{i+1})) \geq \frac{1}{e^n} \dim(S_i).$$

Since  $(S_i, W_i^\perp)$  is an  $(r_{i+1}, \varepsilon + \frac{n}{r_i}2^{r_{i+1}})$ -pair, we have

$$\dim(S_i) \geq \dim(W_i^\perp) - d\left(\varepsilon + \frac{n}{r_i}2^{r_{i+1}}\right)$$

and so

$$\begin{aligned} \sum_{s=1}^{g(i+1)-g(i)} \dim(B_{\mathcal{M}}(w_{g(i)+s}, r_{i+1})) &\geq \frac{1}{e^n} \dim(W_i^\perp) - \frac{d(1 - \frac{1}{e^n})^k}{e^n} \\ &\geq \frac{1}{e^n} \dim(W_i^\perp) - \bar{d}\left(1 - \frac{1}{e^n}\right)^{i+1}. \end{aligned}$$

Therefore we have also

$$\begin{aligned} \sum_{j=1}^{g(i+1)} \dim(B_{\mathcal{M}}(w_j, R_j)) &\geq \dim(W_i) + \frac{1}{e^n} \dim(W_i^\perp) - \bar{d}\left(1 - \frac{1}{e^n}\right)^{i+1} \\ &= d - \dim(W_i^\perp) + \frac{1}{e^n} \dim(W_i^\perp) - \bar{d}\left(1 - \frac{1}{e^n}\right)^{i+1} \\ &\geq \bar{d}\left(1 - \left(1 - \frac{1}{e^n}\right)^{i+1}\right) - \left(1 - \frac{1}{e^n}\right) \dim(W_i^\perp). \end{aligned}$$

By the inductive assumption, we have  $\dim(W_i^\perp) \leq \bar{d} \cdot i(1 - \frac{1}{e^n})^i + d \cdot i \frac{\delta}{2k} + \varepsilon d$ , so altogether we have

$$\begin{aligned} \sum_{j=1}^{g(i+1)} \dim(B_{\mathcal{M}}(w_j, R_j)) &\geq \bar{d}\left(1 - \left(1 - \frac{1}{e^n}\right)^{i+1}\right) - \bar{d} \cdot i\left(1 - \frac{1}{e^n}\right)^{i+1} - d\left(i \frac{\delta}{2k} + \varepsilon\right) \\ &= \bar{d}\left(1 - (i + 1)\left(1 - \frac{1}{e^n}\right)^{i+1}\right) - d\left(i \frac{\delta}{2k} + \varepsilon\right) \\ &\geq \bar{d}\left(1 - (i + 1)\left(1 - \frac{1}{e^n}\right)^{i+1}\right) - d\left(i \frac{\delta}{2k} + \frac{\delta}{2k}\right), \end{aligned}$$

which is the inductive statement we wanted to show. Hence the lemma follows.  $\square$

We have now everything in place to prove Theorem 6.

*Proof of Theorem 6.* Let us fix  $\varepsilon > 0$ . By Lemma 17, we can fix  $R \in \mathbb{N}$  and  $\eta > 0$  such that if  $\mathcal{A}$  is a  $d$ -matrix tuple for some  $d \in \mathbb{N}$ , and  $W \subset \mathbb{C}^d$  is an  $R$ -multi-ball space for  $\mathcal{A}$  with  $\dim(W) \geq (1 - \eta)d$ , then we can find a commuting  $d$ -matrix tuple  $\mathcal{B}$  such that

$$d_{\text{rank}}(\mathcal{A}, \mathcal{B}) \leq \varepsilon.$$

However by Lemma 25, we can find  $\delta > 0$  such that if  $d \in \mathbb{N}$  and  $\mathcal{A}$  is a  $*$ -closed  $\delta$ -commuting tuple then there exists an  $R$ -multi-ball space  $W \subset \mathbb{C}^d$  for  $\mathcal{A}$  such that

$$\dim(W) \geq (1 - \eta)d.$$

This finishes the proof.  $\square$

## 5. Abels' group is not stable with respect to the rank metric

We finish the article with the following proof.

*Proof of Theorem 4.* The centre  $Z(A_p)$  of  $A_p$  is the group of matrices of the form

$$\begin{pmatrix} 1 & 0 & 0 & * \\ & 1 & 0 & 0 \\ & & 1 & 0 \\ & & & 1 \end{pmatrix},$$

isomorphic to  $\mathbb{Z}[\frac{1}{p}]$ . Consider the central subgroup  $H$  of the elements of the form

$$\begin{pmatrix} 1 & 0 & 0 & x \\ & 1 & 0 & 0 \\ & & 1 & 0 \\ & & & 1 \end{pmatrix},$$

where  $x \in \mathbb{Z}$ .

Let  $\Delta$  be the quotient group  $A_p/H$  and let  $\pi: A_p \rightarrow \Delta$  be the quotient map. Let  $\gamma_1, \dots, \gamma_g$  be generators of  $A_p$  and let  $P_1, \dots, P_r$  be noncommutative monomials (possibly with negative exponents) such that

$$\langle \gamma_1, \dots, \gamma_g | P_1(\gamma_1, \dots, \gamma_g), \dots, P_r(\gamma_1, \dots, \gamma_g) \rangle,$$

is a presentation of  $A_p$ , and let  $H$  be such that for all  $i$  we have that  $P_i$  has length at most  $H$ . Let  $F_1, F_2, \dots \subset \Delta$  be a sequence of Følner sets in  $\Delta$ . Let  $S \subset \Delta$  be the set of those elements which can be represented as words of length at most  $H$  in the elements  $\pi(\gamma_1), \dots, \pi(\gamma_g)$  and their inverses. For  $i \in \mathbb{N}_+$  let  $\text{int}(F_i)$  be the subset of those  $f \in F_i$  such that for all  $s \in S$  we have  $sf \in F_i$ .

For  $i \in \mathbb{N}_+$  and  $j = 1, \dots, g$  let  $A_i^j$  be a permutation of  $F_i$  which is equal to  $\pi(\gamma_j)$  on  $\text{int}(F_i)$  (there is in general no unique such permutation). In what follows we will think of  $A_i^j$  as permutation matrices – in particular they are unitary matrices.

Since  $F_i$  is a Følner sequence, we have, for any  $k \in \{1, \dots, r\}$  that

$$\text{rank}(P_k(A_i^1, \dots, A_i^g) - \text{Id}_{|F_i|}) \xrightarrow{i \rightarrow \infty} 0,$$

since the left-hand side is bounded from above by  $1 - \frac{|\text{int } F_i|}{|F_i|}$ .

By way of contradiction, let us assume that  $A_p$  is stable with respect to the rank metric. It follows that we can find  $g$  sequences of invertible matrices  $B_i^1, \dots, B_i^g$  with

$$\frac{1}{|F_i|} \dim(\text{im}(\widehat{A}_i - \widehat{B}_i)) \xrightarrow{i \rightarrow \infty} 0$$

and such that  $P_i(B_1, \dots, B_g) = 1$  for all  $i = 1, \dots, r$ . In particular for each  $i = 1, 2, \dots$  we get a representation  $\rho_i: A_p \rightarrow GL(n_i, \mathbb{C})$  for suitable  $n_i \in \mathbb{N}$ .

Now let  $t$  be a generator of  $H$ . Since  $t$  is a central element, we have that each eigenspace of  $\rho_i(t)$  is preserved under the action of  $A_p$ . Let  $V_i \subset \mathbb{C}^{n_i}$  be the eigenspace of  $\rho_i(t)$  corresponding to the eigenvalue 1, i.e. the set of all  $v \in \mathbb{C}^{n_i}$  such that  $\rho_i(t)(v) = v$ . Note that  $\frac{\dim(V_i)}{n_i} \xrightarrow{i \rightarrow \infty} 1$

We have representations  $\bar{\rho}_i$  of  $A_p/H$  on all the spaces  $V_i$ . Now let  $K \subset Z(A_p)$  be the subgroup of  $Z(A_p)$  of elements of the form  $\frac{n}{p}$ , where  $n \in \mathbb{Z}$ , and let  $\bar{K}$  be the image of  $K$  in  $Z(A_p)/H$ . Since  $\bar{K}$  is finite, we may assume that  $i$  is big enough so that  $\text{rank}(\bar{\rho}_i(\gamma) - 1) \geq \frac{1}{2}$  for all  $\gamma \in \bar{K} \setminus \{e\}$ .

But for every element  $\eta \in Z(A_p) \setminus H$  there exists  $n \in \mathbb{N}_+$  such that  $\eta^n \in K \setminus H$ . It follows that for every  $\eta \in Z(A_p) \setminus H$  we have that  $\rho_i(\eta)$  does not act as the identity on  $V_i$ . This shows that  $\bar{\rho}_i$  is injective on  $Z(A_p)/H$ .

But  $\Delta$  is finitely-generated, and hence  $\bar{\rho}_i(\Delta)$  is a finitely-generated linear group, which by Malcev’s theorem [17] shows that  $\Delta$  is residually finite. But the abelian group  $Z(A_p)/H \subset \Delta$  is not residually finite (see [4] for a short argument), which is a contradiction. This finishes the proof. □

### References

- [1] H. Abels, An example of a finitely presented solvable group. In C. T. C. Wall (ed.), *Homological group theory*. (Durham, 1977.) London Mathematical Society Lecture Note Series, 36. Cambridge University Press, Cambridge and New York, 1979, 205–211. [Zbl 0422.20026](#) [MR 0564423](#)
- [2] G. Arzhantseva and L. Paunescu, Almost commuting permutations are near commuting permutations. *J. Funct. Anal.* **269** (2015), no. 3, 745–757. [Zbl 1368.20025](#) [MR 3350728](#)

- [3] G. Arzhantseva and L. Paunescu, Linear sofic groups and algebras. *Trans. Amer. Math. Soc.* **369** (2017), no. 4, 2285–2310. [Zbl 1406.20038](#) [MR 3592512](#)
- [4] O. Becker, A. Lubotzky, and A. Thom, Stability and invariant random subgroups. *Duke Math. J.* **168** (2019), no. 12, 2207–2234. [Zbl 07145001](#) [MR 3999445](#)
- [5] W. D. Brownawell, Bounds for the degrees in the Nullstellensatz. *Ann. of Math. (2)* **126** (1987), no. 3, 577–591. [Zbl 0641.14001](#) [MR 0916719](#)
- [6] K. R. Davidson, Almost commuting Hermitian matrices. *Math. Scand.* **56** (1985), no. 2, 222–240. [Zbl 0563.15010](#) [MR 0813638](#)
- [7] S. Eliahou, Wilf’s conjecture and Macaulay’s theorem. *J. Eur. Math. Soc. (JEMS)* **20** (2018), no. 9, 2105–2129. [Zbl 1436.20114](#) [MR 3836842](#)
- [8] N. Filonov and Y. Safarov, On the relation between an operator and its self-commutator. *J. Funct. Anal.* **260** (2011), no. 10, 2902–2932. [Zbl 1239.47029](#) [MR 2774059](#)
- [9] P. Friis and M. Rørdam, Almost commuting matrices – A short proof of Huaxin Lin’s Theorem. *J. Reine Angew. Math.* **479** (1996), 121–131. [Zbl 0859.47018](#) [MR 1414391](#)
- [10] L. Glebsky, Almost commuting matrices with respect to normalized Hilbert–Schmidt norm. Preprint, 2010. [arXiv:1002.3082](#)
- [11] D. Hadwin and W. Li, A note on approximate liftings. *Oper. Matrices* **3** (2009), no. 1, 125–143. [Zbl 1181.46042](#) [MR 2500597](#)
- [12] P. R. Halmos, Some unsolved problems of unknown depth about operators on Hilbert space. *Proc. Roy. Soc. Edinburgh Sect. A* **76** (1976/77), no. 1, 67–76. [Zbl 0346.47001](#) [MR 0451002](#)
- [13] M. B. Hastings, Making almost commuting matrices commute. *Comm. Math. Phys.* **291** (2009), no. 2, 321–345. [Zbl 1189.15018](#) [MR 2530163](#)
- [14] G. Hermann, Die Frage der endlich vielen Schritte in der Theorie der Polynomideale. *Math. Ann.* **95** (1926), no. 1, 736–788. [MR 1512302](#)
- [15] H. Lin, Almost commuting self-adjoint matrices and applications. *Fields. Inst. Commun* **13** (1995), 193–233.
- [16] F. S. MacAulay, Some properties of enumeration in the theory of modular systems. *Proc. London Math. Soc. (2)* **26** (1927), 531–555. [JFM 53.0104.01](#) [MR 1576950](#)
- [17] A. I. Malcev, On isomorphic matrix representations of infinite groups of matrices. *Mat. Sb.* **8** (1940), 405–422. In Russian. English transl., *Amer. Math. Soc. Transl. (2)* **45** (1965), 1–18. [JFM 66.0088.03](#) [MR 0003420](#)
- [18] D. S. Ornstein and B. Weiss, Entropy and isomorphism theorems for actions of amenable groups. *J. Analyse Math.* **48** (1987), 1–141. [Zbl 0637.28015](#) [MR 0910005](#)
- [19] P. Rosenthal, Research problems: are almost commuting matrices near commuting matrices? *Amer. Math. Monthly* **76** (1969), no. 8, 925–926. [MR 1535586](#)
- [20] D. Voiculescu, Asymptotically commuting finite rank unitary operators without commuting approximants. *Acta Sci. Math. (Szeged)* **45** (1983), no. 1-4, 429–431. [Zbl 0538.47003](#) [MR 0708811](#)



Received April 2, 2019

Gábor Elek, Department of Mathematics And Statistics, Fylde College,  
Lancaster University, Lancaster, LA1 4YF, United Kingdom

e-mail: [g.elek@lancaster.ac.uk](mailto:g.elek@lancaster.ac.uk)

Łukasz Grabowski, Department of Mathematics And Statistics, Fylde College,  
Lancaster University, Lancaster, LA1 4YF, United Kingdom

e-mail: [lukasz.grabowski@lancaster.ac.uk](mailto:lukasz.grabowski@lancaster.ac.uk)