

Almost finiteness and homology of certain non-free actions

Eduard Ortega and Eduardo Scarparo

Abstract. We show that Cantor minimal $\mathbb{Z} \rtimes \mathbb{Z}_2$ -systems and essentially free amenable odometers are almost finite. We also compute the homology groups of Cantor minimal $\mathbb{Z} \rtimes \mathbb{Z}_2$ -systems and show that the associated transformation groupoids satisfy the HK conjecture if and only if the action is free.

1. Introduction

The property of almost finiteness for ample groupoids was introduced by Matui and applied to questions in groupoid homology in [13]. Furthermore, in [2, 10, 11, 20], this property was applied to problems in classification of C^* -algebras.

Kerr and Szabó in [11] and Suzuki in [20] observed that it is a consequence of work of Downarowicz and Zhang [5] that any free action of a group with subexponential growth on the Cantor set is almost finite. Moreover, free odometers arising from sequences of finite index normal subgroups of an amenable group were shown to be almost finite by Kerr in [10].

On the other hand, there exist interesting examples of non-free odometers. For example, in [19], certain non-free $\mathbb{Z} \rtimes \mathbb{Z}_2$ -odometers were shown to be counterexamples to the *HK conjecture*, which is a conjecture posted by Matui in [14] that relates the homology groups of an ample groupoid with the K-theory of its reduced C^* -algebra.

In Section 2, we investigate almost finiteness of odometers and of Cantor minimal $\mathbb{Z} \rtimes \mathbb{Z}_2$ -systems.

In particular, we prove that an amenable odometer is essentially free if and only if it is almost finite. The proof uses a result by Kar et al. [8] which shows that, given an amenable essentially free odometers, the acting group admits a Følner sequence consisting of coset representatives.

We also show that any Cantor minimal $\mathbb{Z} \rtimes \mathbb{Z}_2$ -system is almost finite. For that, we use a structure result for Cantor minimal $\mathbb{Z} \rtimes \mathbb{Z}_2$ -systems from [3].

In Section 3, we compute the homology groups of Cantor minimal $\mathbb{Z} \rtimes \mathbb{Z}_2$ -systems and conclude that these systems satisfy the HK conjecture if and only if the action is free.

It should come as no surprise that, in determining the homology groups, we are able to mainly follow the ideas introduced by Bratteli et al. in [3] and Thomsen in [22] in their computation of the K-theory of the associated crossed products since, although the HK conjecture is now known to be false, it has been verified in many cases (see the work of Proietti and Yamashita [16, Remark 3.5] and the references therein).

2. Almost finiteness

In this section, we review some terminology about étale groupoids and verify almost finiteness for certain classes of non-principal groupoids. The reader can find an introduction to étale groupoids in, e.g., [18, Chapter 3].

2.1. Almost finite groupoids

Let G be a Hausdorff étale groupoid with range and source maps denoted by r and s . A *bisection* is a subset $S \subset G$ such that $r|_S$ and $s|_S$ are injective maps.

Let $G' := \{g \in G : r(g) = s(g)\}$ and let $G^{(0)}$ be the unit space of G . Then G is said to be *principal* if $G' = G^{(0)}$ and *effective* if the interior of G' equals $G^{(0)}$. The *orbit* of a point $x \in G^{(0)}$ is the set $G(x) := r(s^{-1}(x))$. We say that G is *minimal* if the orbit of each $x \in G^{(0)}$ is dense in $G^{(0)}$. Also G is said to be *ample* if its unit space is totally disconnected.

Example 2.1. Let X be a locally compact Hausdorff space and let $\alpha : \Gamma \curvearrowright X$ be an action of a discrete group Γ on X . Given $g \in \Gamma$ and $x \in X$, we will denote $gx := \alpha(g)(x)$. As a space, the *transformation groupoid* G associated with α is $G := \Gamma \times X$ equipped with the product topology. The product of two elements $(h, y), (g, x) \in G$ is defined if and only if $y = gx$, in which case $(h, gx)(g, x) := (hg, x)$. Inversion is given by $(g, x)^{-1} := (g^{-1}, gx)$. The unit space $G^{(0)}$ is naturally identified with X . Note that G is principal if and only if α is free ($gx = x \Rightarrow g = e$). If X is totally disconnected, then G is ample.

Let G be an ample groupoid with compact unit space. A subgroupoid $K \subset G$ is said to be *elementary* if K is compact-open principal and $K^{(0)} = G^{(0)}$. The groupoid G is said to be *almost finite* if for any compact subset $C \subset G$ and $\varepsilon > 0$, there is $K \subset G$ elementary subgroupoid such that, for any $x \in G^{(0)}$, we have that

$$\frac{|CKx \setminus Kx|}{|K(x)|} < \varepsilon.$$

For compact groupoids, the following holds:

Proposition 2.2. *Let G be an ample almost finite groupoid. If G is compact, then G is principal.*

Proof. Compactness of G implies that there is a finite partition of G into compact-open bisections. Hence, there is $M > 0$ such that, for each $x \in G^{(0)}$, we have that $|G(x)| < M$.

If G is not principal, there is $g \in G \setminus G^{(0)}$ such that $r(g) = s(g)$. Then, for any elementary subgroupoid $K \subset G$, we have that $g \notin K$ and

$$\frac{|gKs(g) \setminus Ks(g)|}{|K(s(g))|} > \frac{1}{M}.$$

Therefore, G is not almost finite. ■

Given an étale groupoid G with compact unit space, denote by $M(G)$ the set of G -invariant probability measures on $G^{(0)}$.

Remark 2.3. It was observed in [13, Remark 6.6] that if G is an almost finite groupoid, then $M(G) \neq \emptyset$ and, given $\mu \in M(G)$, we have that

$$\mu(r(U \cap (G' \setminus G^{(0)}))) = 0, \quad \text{for any } U \subset G \text{ compact-open bisection.} \quad (1)$$

2.2. Almost finite actions

Let us recall the characterization of almost finiteness for transformation groupoids presented in [11, 20]. We will restrict ourselves to actions on totally disconnected spaces.

Let Γ be a group acting on a compact Hausdorff totally disconnected space X . A *clopen tower* is a pair (V, S) consisting of a clopen subset V of X and a finite subset S of Γ such that the sets sV for $s \in S$ are pairwise disjoint. The set S is said to be the *shape* of the tower. A *clopen castle* is a finite collection of clopen towers $\{(V_i, S_i)\}_{i \in I}$ such that the sets $S_i V_i = \{gV_i : g \in S_i\}$ for $i \in I$ are pairwise disjoint.

Given $\varepsilon > 0$ and $K \subset \Gamma$ finite, we say that a finite set $F \subset \Gamma$ is (K, ε) -invariant if $|KF \Delta F| < \varepsilon|F|$.

Let G be the transformation groupoid associated to the action of Γ on X . Then G is almost finite if and only if, given $K \subset \Gamma$ finite and $\varepsilon > 0$, there is a clopen castle which partitions X , and whose shapes are (K, ε) -invariant (see [20, Lemma 5.2] for a proof of this fact). In this case, we say that the *action is almost finite*.

Recall that an action of a group Γ on a locally compact Hausdorff space X is said to be *minimal* if the orbit of any $x \in X$ is dense in X . Clearly, this is equivalent to the associated transformation groupoid being minimal. It is also equivalent to every open (or closed) Γ -invariant set being trivial. If the action is minimal, then any $f \in C(X, \mathbb{Z})$ which is Γ -invariant must be constant. We will say that a homeomorphism φ on X is minimal if the \mathbb{Z} -action induced by φ is minimal. By a *Cantor minimal Γ -system*, we mean a minimal action of Γ on the Cantor set.

Given Γ a group acting on a compact Hausdorff space X , we denote by $M_\Gamma(X)$ the set of Γ -invariant probability measures on X . For $g \in \Gamma$, let $\text{Fix}_g \subset X$ be the set of points fixed by g .

If $\mu \in M_\Gamma(X)$, we say that the action of Γ on (X, μ) is *essentially free* if, for any $g \in \Gamma \setminus \{e\}$, we have $\mu(\text{Fix}_g) = 0$. The action is said to be *topologically free* if the interior of Fix_g is empty for each $g \in \Gamma \setminus \{e\}$. Notice that topological freeness is equivalent to effectiveness of the associated transformation groupoid.

Also observe that if $\Gamma \curvearrowright X$ is a minimal action, then any $\mu \in M_\Gamma(X)$ has full support. Therefore, for minimal actions, essential freeness of $\Gamma \curvearrowright (X, \mu)$ implies topological freeness of $\Gamma \curvearrowright X$.

Remark 2.4. If G is a transformation groupoid associated to an action of a group Γ on a compact Hausdorff space X and $\mu \in M(G) \approx M_\Gamma(X)$, then the condition in (1) is equivalent to the action of Γ on (X, μ) being essentially free. It follows from Remark 2.3 that if the action is almost finite, then $\Gamma \curvearrowright (X, \mu)$ is essentially free.

2.3. Odometers

Let Γ be a group and $(\Gamma_i)_{i \in \mathbb{N}}$ a sequence of finite index subgroups of Γ such that, for every $i \in \mathbb{N}$, $\Gamma_i \supsetneq \Gamma_{i+1}$.

For each $i \in \mathbb{N}$, let $p_i: \Gamma/\Gamma_{i+1} \rightarrow \Gamma/\Gamma_i$ be the surjection given by

$$p_i(\gamma\Gamma_{i+1}) := \gamma\Gamma_i, \quad \text{for } \gamma \in \Gamma. \quad (2)$$

Let $X := \varprojlim (\Gamma/\Gamma_i, p_i) = \{(x_i) \in \prod \Gamma/\Gamma_i : p_i(x_{i+1}) = x_i, \forall i \in \mathbb{N}\}$. Then X is homeomorphic to the Cantor set and Γ acts in a minimal way on X by $\gamma(x_i) := (\gamma x_i)$, for $\gamma \in \Gamma$ and $(x_i) \in X$. This action is called an *odometer*.

Given $j \geq 1$ and $g\Gamma_j \in \Gamma/\Gamma_j$, let $U(j, g\Gamma_j) := \{(x_i) \in X : x_j = g\Gamma_j\}$. Then

$$\{U(j, g\Gamma_j) : j \in \mathbb{N}, g\Gamma_j \in \Gamma/\Gamma_j\}$$

is a basis for X consisting of compact-open sets.

Notice that X admits a unique Γ -invariant probability measure. Hence, there is no ambiguity in calling an odometer action essentially free. It is known (see, for example, [7, (1.6)]) that the odometer is essentially free if and only if, for each $g \in \Gamma \setminus \{e\}$, we have

$$\lim_i \frac{|\{x \in \Gamma/\Gamma_i : gx = x\}|}{|\Gamma/\Gamma_i|} = 0.$$

If the odometer is free, then $\bigcap_i \Gamma_i = \{e\}$, but the converse does not hold in general (see, e.g., [1, Theorem 1]).

Let us now give a characterization of almost finite odometers. First recall that a Følner sequence (F_n) for Γ is a sequence of finite subsets of Γ such that for every $g \in \Gamma$ we have that $\frac{|gF_n \Delta F_n|}{|F_n|} \rightarrow 0$.

Theorem 2.5 ([8, 9]). *Let Γ be a countable amenable group and $\Gamma \curvearrowright X := \varprojlim \Gamma/\Gamma_i$ an odometer. The following conditions are equivalent:*

- (i) *the action is essentially free;*
- (ii) *there is a Følner sequence (F_n) for Γ such that each F_n is a complete set of representatives for Γ/Γ_n ;*
- (iii) *the action is almost finite;*
- (iv) *there is a unique tracial state on $C(X) \rtimes \Gamma$.*

Proof. The equivalence of (i) and (iv) is a consequence of [9, Corollary 2.8]. That (iii) implies (i) is a consequence of Remark 2.4. The implication from (i) to (ii) is the content of [8, Theorem 7]. Let us show then that (ii) implies (iii).

Notice that for each $n \in \mathbb{N}$,

$$X = \bigsqcup_{\gamma \in F_n} \gamma U(n, \Gamma_n). \quad (3)$$

By taking n sufficiently big, we can make the shapes of the castle (3) arbitrarily invariant. ■

A few remarks are in order about the result above:

Remark 2.6. (i) It follows from Theorem 2.5 and the results in [2] that crossed products associated to essentially free odometers in which the acting group is countable and amenable are classifiable by their Elliott-invariant.

(ii) It is a consequence of [12, Proposition 4.7] that if $\Gamma \curvearrowright \varprojlim \Gamma / \Gamma_i$ is an almost finite odometer, then Γ is amenable.

(iii) In [19, Proposition 2.1], it was shown that an odometer $\Gamma \curvearrowright X := \varprojlim \Gamma / \Gamma_i$ is topologically free if and only if, for every $\gamma \in \bigcap \Gamma_i \setminus \{e\}$ and $j \geq 1$, there exists $b \in \Gamma_j$ such that $b^{-1}\gamma b \notin \bigcap \Gamma_i$.

We do not know whether, for a countable amenable group Γ , topological freeness of a Γ -odometer implies essential freeness. If one drops the amenability assumption, there are several counterexamples in the literature (see, for example, [1, Theorem 1]).

Example 2.7. Recall that the *infinite dihedral group* is the semidirect product $\mathbb{Z} \rtimes \mathbb{Z}_2$ associated to the action of \mathbb{Z}_2 on \mathbb{Z} by multiplication by -1 .

Let (n_i) be a strictly increasing sequence of natural numbers such that $n_i | n_{i+1}$, for every $i \in \mathbb{N}$. Define $\Gamma := \mathbb{Z} \rtimes \mathbb{Z}_2$ and, for $i \geq 1$, $\Gamma_i := n_i \mathbb{Z} \rtimes \mathbb{Z}_2$.

By [19, Lemma 3.2] and the fact that any element of the form $(n, 1) \in \mathbb{Z} \rtimes \mathbb{Z}_2$ is conjugate to either $(0, 1)$ or $(1, 1)$, we obtain that any $g \in \Gamma \setminus \{e\}$ fixes at most finitely many points in $\varprojlim \Gamma / \Gamma_i$. Hence, this odometer is essentially free.

Let us describe a Følner sequence for $\mathbb{Z} \rtimes \mathbb{Z}_2$ satisfying condition (ii) in Theorem 2.5. Given $m \in \mathbb{N}$, let $F_m \subset \mathbb{Z} \rtimes \mathbb{Z}_2$ be defined by

$$F_m = \begin{cases} \left(\left[-\frac{m}{2}, -1 \right] \times \{1\} \right) \cup \left(\left[0, \frac{m}{2} \right] \times \{0\} \right), & \text{if } m \text{ is even,} \\ \left(\left[-\frac{m-1}{2}, -1 \right] \times \{1\} \right) \cup \left(\left[0, \frac{m-1}{2} \right] \times \{0\} \right), & \text{if } m \text{ is odd.} \end{cases} \quad (4)$$

Then $(F_m)_{m \in \mathbb{N}}$ is a Følner sequence for $\mathbb{Z} \rtimes \mathbb{Z}_2$ such that each F_m is a set of representatives for $\frac{\mathbb{Z} \rtimes \mathbb{Z}_2}{(m\mathbb{Z}) \rtimes \mathbb{Z}_2}$.

2.4. Almost finiteness of Cantor minimal $\mathbb{Z} \rtimes \mathbb{Z}_2$ -systems

Notice that any action α of $\mathbb{Z} \rtimes \mathbb{Z}_2$ on a set X is given by a pair of bijections on X (φ, σ) such that $\sigma^2 = \text{Id}_X$ and $\sigma\varphi\sigma = \varphi^{-1}$, so that $\alpha_{n,i} = \varphi^n \sigma^i$, for $(n, i) \in \mathbb{Z} \rtimes \mathbb{Z}_2$.

There is an isomorphism $\mathbb{Z} \rtimes \mathbb{Z}_2 \simeq \mathbb{Z}_2 * \mathbb{Z}_2$ which takes the canonical generators $a, b \in \mathbb{Z}_2 * \mathbb{Z}_2$ to $(1, 1), (0, 1) \in \mathbb{Z} \rtimes \mathbb{Z}_2$.

Proposition 2.8. *Let $\alpha := (\varphi, \sigma)$ be a minimal action of $\mathbb{Z} \rtimes \mathbb{Z}_2$ on the Cantor set X . The following holds.*

- (i) *The \mathbb{Z} -action induced by φ is free and α is topologically free. If α is not free, then either σ or $\varphi\sigma$ has at least one fixed point.*
- (ii) *If the \mathbb{Z} -action induced by φ is not minimal, then there exists a clopen set Y such that $Y \cap \sigma(Y) = \emptyset$, $Y \cup \sigma(Y) = X$, $\varphi(Y) = Y$, and $\varphi|_Y$ is minimal. In particular, α is free.*

Proof. (i) Suppose that the \mathbb{Z} -action induced by φ is not free. Then there is $n \in \mathbb{Z} \setminus \{0\}$ and $x \in X$ such that $\varphi^n(x) = x$; then $\varphi^n\sigma(x) = \sigma\varphi^{-n}(x) = \sigma(x)$. Hence, the α -orbit of x is finite, which contradicts the fact that α is minimal. Therefore, the \mathbb{Z} -action induced by φ is free.

Suppose that α is not topologically free. Then there is a non-empty open set $U \subset X$ and $n \in \mathbb{Z}$ such that $\varphi^n\sigma$ fixes U pointwise.

Fix $x \in U$. By minimality of α , there is $(m, i) \in \mathbb{Z} \rtimes \mathbb{Z}_2$ such that $\varphi^m\sigma^i(x) \in U$ and $\varphi^m\sigma^i(x) \neq x$. Furthermore, by multiplying (m, i) on the left by $(n, 1)$, we can assume that $i = 0$.

Since $\varphi^m(x) \in U$, we get that $\varphi^m(x) = \varphi^n\sigma\varphi^m(x) = \varphi^{-m}\varphi^n\sigma(x) = \varphi^{-m}(x)$; hence $\varphi^{2m}(x) = x$. But this contradicts the fact that the \mathbb{Z} -action induced by φ is free. Therefore, α is topologically free.

If α is not free, then there is $n \in \mathbb{Z}$ and $x \in X$ such that $\varphi^n\sigma(x) = x$. Since any element of the form $(n, 1) \in \mathbb{Z} \rtimes \mathbb{Z}_2$ is conjugate to $(0, 1)$ or $(1, 1)$, we conclude that either σ or $\varphi\sigma$ has at least one fixed point.

- (ii) This is the content of [15, Proposition 2.10] (see also [22, Lemma 4.28]). ■

The following lemma is a slight modification of [3, Lemma 1.4], and we include the proof for the sake of completeness.

Lemma 2.9. *Let (φ, σ) be a minimal action of $\mathbb{Z} \rtimes \mathbb{Z}_2$ on the Cantor set X and $Y \subset X$ a non-empty clopen set such that $\sigma(Y) = Y$.*

Then there is a partition of Y into clopen sets Y_1, \dots, Y_K and positive integers $J_1 < J_2 < \dots < J_K$ such that $\{\varphi^k(Y_i) : i = 1, \dots, K, k = 0, \dots, J_i - 1\}$ is a partition of X , and $\sigma\varphi^{J_i}(Y_i) = Y_i$ for each $i = 1, \dots, K$.

Proof. Define $\lambda(y) := \min\{n > 0 : \varphi^n(y) \in Y\}$, for $y \in Y$. From Proposition 2.8, it follows that the map λ is well defined, in the sense that for each $y \in Y$ there is $n > 0$ such that $\varphi^n(y) \in Y$.

It is easy to check that λ is continuous. Hence, it has a finite range $\{J_1, \dots, J_K\}$, where $J_1 < J_2 < \dots < J_K$. Define $Y_i = \lambda^{-1}(J_i)$ for each i . Then the sets $\{\varphi^k(Y_i) : i = 1, \dots, K, k = 0, \dots, J_i - 1\}$ are pairwise disjoint. From Proposition 2.8, it follows that

the union of these sets is φ and σ -invariant, hence minimality of the action implies that this is a partition of X .

Let us show that $\sigma\varphi^{J_i}(Y_i) = Y_i$ for each i . Observe first that $\sigma\varphi^{J_i}(Y_i) \subset Y$ since $\varphi^{J_i}(Y_i) \subset Y$ and Y is σ -invariant.

Take $y \in \sigma\varphi^{J_i}(Y_i)$. Then there is $x \in Y_i$ such that $y = \sigma\varphi^{J_i}(x)$. In particular, $\varphi^{J_i}(y) = \sigma(x) \in Y$. Therefore, $\lambda(y) \leq J_i$.

If $i = 1$, we must then have $\lambda(y) = J_1$. Therefore, $\sigma\varphi^{J_1}(Y_1) \subset Y_1$. Since $\sigma\varphi^{J_1}$ is involutive, we conclude that $\sigma\varphi^{J_1}(Y_1) = Y_1$. Arguing the same way for $i = 2, \dots, K$, the result follows. ■

Theorem 2.10. *Any minimal action of $\mathbb{Z} \rtimes \mathbb{Z}_2$ on the Cantor set is almost finite.*

Proof. Let $\alpha = (\varphi, \sigma)$ be a minimal action of $\mathbb{Z} \rtimes \mathbb{Z}_2$ on the Cantor set X . It follows from the fact that α is topologically free (Proposition 2.8) and the Baire category theorem that there is $y \in X$ such that if $(n, i), (m, j) \in \mathbb{Z} \rtimes \mathbb{Z}_2$ are distinct, then $\varphi^n\sigma^i(x) \neq \varphi^m\sigma^j(x)$.

Let Z be a clopen neighborhood of y and $Y := Z \cup \sigma(Z)$. Then $\sigma(Y) = Y$ and, given $N \in \mathbb{N}$, if we take Z sufficiently small, we can assume that the sets $\{Y, \varphi(Y), \dots, \varphi^{N-1}(Y)\}$ are disjoint. Then, by Lemma 2.9, we can partition Y into clopen sets Y_1, \dots, Y_K such that

$$X = \sqcup_{i=1}^K \sqcup_{k=0}^{J_i-1} \varphi^k(Y_i)$$

and $\sigma\varphi^{J_i}(Y_i) = Y_i$ for each $i = 1, \dots, K$. In particular, $\varphi^{-l}\sigma(Y_i) = \varphi^{J_i-l}(Y_i)$ for any l .

Consider the Følner sequence (F_m) introduced in (4), and notice that

$$X = \bigsqcup_{i=1}^K \bigsqcup_{g \in F_{J_i}} \alpha_g(Y_i). \quad (5)$$

Furthermore, each $J_i \geq N$. By taking N sufficiently big, we can make the shapes of the castle (5) arbitrarily invariant. ■

3. Homology of Cantor minimal $\mathbb{Z} \rtimes \mathbb{Z}_2$ -systems

In this section, we compute the homology groups of Cantor minimal $\mathbb{Z} \rtimes \mathbb{Z}_2$ -systems.

Given a group Γ and a Γ -module M , we denote by M_Γ the quotient of M by the subgroup generated by elements of the form $m - mg$, for $m \in M$ and $g \in \Gamma$. Recall that M_Γ is canonically isomorphic to $H_0(\Gamma, M)$. If $i: \Lambda \rightarrow \Gamma$ is an embedding, we denote the canonical map $i_*: H_*(\Lambda, M) \rightarrow H_*(\Gamma, M)$ by cor.

We will use the following result about the homology of free products, whose proof can be found in [21, Theorem 2.3].

Theorem 3.1. *Let Γ_1 and Γ_2 be groups and M a $(\Gamma_1 * \Gamma_2)$ -module. Then, for $n \geq 2$,*

$$H_n(\Gamma_1, M) \oplus H_n(\Gamma_2, M) \simeq H_n(\Gamma_1 * \Gamma_2, M)$$

and there is an exact sequence

$$\begin{aligned} 0 \rightarrow H_1(\Gamma_1, M) \oplus H_1(\Gamma_2, M) &\xrightarrow{(\text{cor}, \text{cor})} H_1(\Gamma_1 * \Gamma_2, M) \\ \rightarrow M &\xrightarrow{(\text{cor}, -\text{cor})} M_{\Gamma_1} \oplus M_{\Gamma_2} \xrightarrow{(\text{cor}, \text{cor})} M_{\Gamma_1 * \Gamma_2} \rightarrow 0. \end{aligned}$$

Given an action of a group Γ on a topological space X , then $C(X, \mathbb{Z})$ has a structure of Γ -module given by $fa(x) := f(ax)$ for every $f \in C(X, \mathbb{Z})$ and $a \in \Gamma$.

Lemma 3.2. *Let a and b be involutive homeomorphisms on the Cantor set X and suppose that the \mathbb{Z} -action induced by ab is minimal. Denote by A and B the abelian group $C(X, \mathbb{Z})$ endowed with the \mathbb{Z}_2 -action given by a and b , respectively.*

*Then $(\text{cor}, \text{cor}): H_1(\mathbb{Z}_2, A) \oplus H_1(\mathbb{Z}_2, B) \rightarrow H_1(\mathbb{Z}_2 * \mathbb{Z}_2, C(X, \mathbb{Z}))$ is an isomorphism and the following sequence is exact:*

$$0 \rightarrow C(X, \mathbb{Z}) \xrightarrow{(\text{cor}, -\text{cor})} A_{\mathbb{Z}_2} \oplus B_{\mathbb{Z}_2} \xrightarrow{(\text{cor}, \text{cor})} C(X, \mathbb{Z})_{\mathbb{Z}_2 * \mathbb{Z}_2} \rightarrow 0.$$

Proof. By Theorem 3.1, we only need to show that

$$(\text{cor}, -\text{cor}): C(X, \mathbb{Z}) \rightarrow A_{\mathbb{Z}_2} \oplus B_{\mathbb{Z}_2}$$

is injective.

Take $f \in C(X, \mathbb{Z})$ such that $(\text{cor}, -\text{cor})(f) = 0$. This implies that there exist $g, h \in C(X, \mathbb{Z})$ such that $f = g - ga = h - hb$. In particular, $fa = fb = -f$. Therefore, $fab = f$. Since the \mathbb{Z} -action induced by ab is minimal, we conclude that f is constant. Finally, as $fa = -f$, we must have $f = 0$. ■

In order to apply Theorem 3.1 and Lemma 3.2 to Cantor minimal $\mathbb{Z}_2 * \mathbb{Z}_2$ -systems, we need to compute homology groups of the form $H_*(\mathbb{Z}_2, C(X, \mathbb{Z}))$.

Lemma 3.3. *Let a be an involutive homeomorphism on a compact, Hausdorff, totally disconnected space X . Then, for $k \geq 0$, we have $H_{2k+1}(\mathbb{Z}_2, C(X, \mathbb{Z})) \simeq C(\text{Fix}_a, \mathbb{Z}_2)$.*

Proof. By [23, Theorem 6.2.2], we have that

$$H_{2k+1}(\mathbb{Z}_2, C(X, \mathbb{Z})) = \frac{\{f \in C(X, \mathbb{Z}) : f = fa\}}{\{f + fa : f \in C(X, \mathbb{Z})\}}.$$

Let $E: \frac{\{f \in C(X, \mathbb{Z}) : f = fa\}}{\{f + fa : f \in C(X, \mathbb{Z})\}} \rightarrow C(\text{Fix}_a, \mathbb{Z}_2)$ be the map given by restriction to Fix_a . Clearly, this is a well-defined homomorphism, and we will show that it is bijective.

Given $F \subset \text{Fix}_a$ clopen, take for each $x \in F$ a clopen set $U_x \subset X$ such that $x \in U_x$, $U_x \cap \text{Fix}_a \subset F$, and $a(U_x) = U_x$. Then there exist $x_1, \dots, x_n \in F$ such that

$$F = \bigcup (U_{x_i} \cap \text{Fix}_a).$$

Let $U := \bigcup U_{x_i}$. Notice that $a(U) = U$. Hence, $E([1_U]) = 1_F$. Therefore, E is surjective.

Let us show now injectivity of E . Take $f \in C(X, \mathbb{Z})$ such that $f = fa$ and $E([f]) = 0$, and we will show that $[f] = 0$. We claim that we can assume that $f|_{\text{Fix}_a} = 0$. Indeed, let U_1, \dots, U_n be a -invariant clopen subsets of X whose union cover Fix_a , and such that f is constant on each U_i . By taking differences, we can assume that these sets are disjoint. By summing f with functions of the form $2m_i 1_{U_i}$, we get our claim.

Assume then that $f|_{\text{Fix}_a} = 0$. We have

$$f = \sum_{q \in \mathbb{Z} \setminus \{0\}} q 1_{f^{-1}(q)}.$$

Since the support of f does not intersect Fix_a , we can, for each $q \in \mathbb{Z} \setminus \{0\}$, partition $f^{-1}(q)$ as $f^{-1}(q) = A_q \sqcup a(A_q)$, for some A_q clopen. Hence, $[f] = 0$. ■

Lemma 3.4. *Let a be an involutive homeomorphism on a compact, Hausdorff, totally disconnected space X and $G_a := \{f \in C(X, \mathbb{Z}) : fa = f \text{ and } f(\text{Fix}_a) \subset 2\mathbb{Z}\}$. Then*

$$\begin{aligned} \psi : C(X, \mathbb{Z})_{\mathbb{Z}_2} &\rightarrow G_a \\ [f] &\mapsto f + fa \end{aligned}$$

is an isomorphism and, for $k \geq 1$, we have $H_{2k}(\mathbb{Z}_2, C(X, \mathbb{Z})) = 0$.

Proof. Clearly, ψ is a well-defined homomorphism, and we will show that it is bijective.

Given $f \in C(X, \mathbb{Z})$, suppose that $f + fa = 0$. Then $f|_{\text{Fix}_a} = 0$. Take A clopen neighborhood of Fix_a such that $a(A) = A$ and f vanishes on A .

Then we can partition A^c as $A^c = B \sqcup a(B)$ for some clopen set B . Let $g := f 1_B$. Then $f = g - ga$. Hence, ψ is injective.

Let us now show surjectivity of ψ . Take $f \in C(X, \mathbb{Z})$ such that $fa = f$ and $f(\text{Fix}_a) \subset 2\mathbb{Z}$, and let us show that $f \in \text{Im } \psi$.

Let U_1, \dots, U_n be a -invariant clopen subsets of X whose union cover Fix_a , and such that f is constant on each U_i . By taking differences, we can assume that these sets are disjoint. By summing f with functions of the form $2m_i 1_{U_i}$, we may assume that $f|_{\text{Fix}_a} = 0$. We have $f = \sum_{q \in \mathbb{Z} \setminus \{0\}} q 1_{f^{-1}(q)}$. Since the support of f does not intersect Fix_a , we can, for each $q \in \mathbb{Z} \setminus \{0\}$, partition $f^{-1}(q)$ as $f^{-1}(q) = A_q \sqcup a(A_q)$, for some A_q clopen. Let $g := \sum_{q \in \mathbb{Z} \setminus \{0\}} q A_q$. Then $\psi([g]) = f$.

Finally, notice that by [23, Theorem 6.2.2], we have that, for $k \geq 1$,

$$H_{2k}(\mathbb{Z}_2, C(X, \mathbb{Z})) = \frac{\{f \in C(X, \mathbb{Z}) : f + fa = 0\}}{\{f - fa : f \in C(X, \mathbb{Z})\}}. \quad (6)$$

Since the right-hand side of (6) is equal to $\ker \psi$, the result follows. ■

Given a finite index subgroup Λ of a group Γ , and M a Γ -module, there is a homomorphism $\text{tr} : M_\Gamma \rightarrow M_\Lambda$ given by $\text{tr}([m]) = [\sum g_i m]$, where $\{g_1, \dots, g_{[\Gamma:\Lambda]}\}$ is a complete set of right coset representatives (this is the so-called *transfer map*, and it does not depend on the choice of representatives).

Theorem 3.5. *Let $\alpha := (\varphi, \sigma)$ be an action of $\mathbb{Z} \rtimes \mathbb{Z}_2$ on the Cantor set X such that the restricted \mathbb{Z} -action is minimal. Let*

$$\begin{aligned} \text{tr}: H_0(\mathbb{Z} \rtimes \mathbb{Z}_2, C(X, \mathbb{Z})) &\rightarrow (1 + \sigma_*)H_0(\mathbb{Z}, C(X, \mathbb{Z})) \\ [f] &\mapsto [f + f\sigma]. \end{aligned}$$

If α is not free, then tr is an isomorphism. If α is free, then $\ker \text{tr} \simeq \mathbb{Z}_2$ and it is generated by $[1_K] - [1_L]$, where K and L are clopen sets such that $X = K \sqcup \sigma(K) = L \sqcup \varphi\sigma(L)$.

Proof. Let $f \in \ker \text{tr}$. In this case, there is $h \in C(X, \mathbb{Z})$ such that

$$f + f\sigma = h - h\varphi. \quad (7)$$

This implies that $h - h\varphi = (h - h\varphi)\sigma = h\sigma - h\varphi\sigma$.

Therefore, $h + h\varphi\sigma = h\sigma + h\varphi$. Composing the right-hand side of this equation with φ^{-1} , we obtain that $h + h\varphi\sigma$ is φ -invariant. Since, φ is minimal, we conclude that there is an integer z such that

$$h + h\varphi\sigma = -z. \quad (8)$$

Furthermore, from (7), we get that $f + f\sigma - h = -h\varphi = -h\varphi - h\sigma + h\sigma = z + h\sigma$. Therefore,

$$f + f\sigma - h - h\sigma = z. \quad (9)$$

Let G_σ and $G_{\varphi\sigma}$ be as in Lemma 3.4. By Lemmas 3.2 and 3.4, there is an isomorphism

$$\psi: C(X, \mathbb{Z})_{\mathbb{Z} \rtimes \mathbb{Z}_2} \rightarrow \frac{G_\sigma \oplus G_{\varphi\sigma}}{\{(g + g\sigma, -g - g\varphi\sigma) : g \in C(X, \mathbb{Z})\}}$$

such that, for $f_1, f_2 \in C(X, \mathbb{Z})$, we have $\psi([f_1 + f_2]) = [(f_1 + f_1\sigma, f_2 + f_2\varphi\sigma)]$.

From (8) and (9), we get that $\psi([f]) = \psi([f + 0]) = [(f + f\sigma, 0)] = [(z, -z)]$.

If z is even, then clearly $[(z, -z)] = 0$. Besides, if α is not free, then it follows from (8), (9), and Proposition 2.8 that z is even.

Now suppose that α is free and take K and L clopen sets as in the statement. Clearly, $[1_K - 1_L] \in \ker \text{tr}$ and $\psi([1_K - 1_L]) = [(1, -1)]$. Finally, notice that $[(1, -1)] \neq 0$. Indeed, if there is $g \in C(X, \mathbb{Z})$ such that $1 = g + g\sigma = g + g\varphi\sigma$, then $g\sigma = g\varphi\sigma$, which implies that g is φ -invariant, hence constant. But this contradicts the fact that $1 = g + g\sigma$. ■

The next result is essentially a summary of what we have obtained so far.

Theorem 3.6. *Let $\alpha := (\varphi, \sigma)$ be a minimal action of $\mathbb{Z} \rtimes \mathbb{Z}_2$ on the Cantor set X .*

- (i) *If φ is not minimal, then there exists $Y \subset X$ clopen and φ -invariant such that $X = Y \sqcup \sigma(Y)$ and $\varphi|_Y$ is minimal. Furthermore,*

$$\begin{aligned} H_0(\mathbb{Z} \rtimes \mathbb{Z}_2, C(X, \mathbb{Z})) &\simeq H_0(\mathbb{Z}, C(Y, \mathbb{Z})), \\ H_1(\mathbb{Z} \rtimes \mathbb{Z}_2, C(X, \mathbb{Z})) &\simeq \mathbb{Z}, \\ H_n(\mathbb{Z} \rtimes \mathbb{Z}_2, C(X, \mathbb{Z})) &= 0, \quad \text{for } n \geq 2. \end{aligned}$$

(ii) If φ is minimal and α is free, then

$$\begin{aligned} H_0(\mathbb{Z} \rtimes \mathbb{Z}_2, C(X, \mathbb{Z})) &\simeq \mathbb{Z}_2 \oplus (1 + \sigma_*)H_0(\mathbb{Z}, C(X, \mathbb{Z})), \\ H_n(\mathbb{Z} \rtimes \mathbb{Z}_2, C(X, \mathbb{Z})) &= 0, \quad \text{for } n \geq 1. \end{aligned}$$

(iii) If α is not free, then

$$\begin{aligned} H_0(\mathbb{Z} \rtimes \mathbb{Z}_2, C(X, \mathbb{Z})) &\simeq (1 + \sigma_*)H_0(\mathbb{Z}, C(X, \mathbb{Z})), \\ H_{2n-1}(\mathbb{Z} \rtimes \mathbb{Z}_2, C(X, \mathbb{Z})) &\simeq C(\text{Fix}_\sigma \sqcup \text{Fix}_{\varphi\sigma}, \mathbb{Z}_2), \quad \text{for } n \geq 1, \\ H_{2n}(\mathbb{Z} \rtimes \mathbb{Z}_2, C(X, \mathbb{Z})) &= 0, \quad \text{for } n \geq 1. \end{aligned}$$

Proof. (i) The existence of Y is the content of Proposition 2.8. Notice that

$$C(X, \mathbb{Z}) \simeq \text{Ind}_{\mathbb{Z}}^{\mathbb{Z} \rtimes \mathbb{Z}_2} C(Y, \mathbb{Z}).$$

Shapiro's lemma [4, Proposition 6.2] implies then that

$$H_*(\mathbb{Z} \rtimes \mathbb{Z}_2, C(X, \mathbb{Z})) \simeq H_*(\mathbb{Z}, C(Y, \mathbb{Z})).$$

Finally, the homology groups of a Cantor minimal \mathbb{Z} -system are easy to compute.

(ii) Let us begin by showing that

$$H_0(\mathbb{Z} \rtimes \mathbb{Z}_2, C(X, \mathbb{Z})) \simeq \mathbb{Z}_2 \oplus (1 + \sigma_*)H_0(\mathbb{Z}, C(X, \mathbb{Z})). \quad (10)$$

Consider the map

$$\text{tr}: H_0(\mathbb{Z} \rtimes \mathbb{Z}_2, C(X, \mathbb{Z})) \rightarrow (1 + \sigma_*)H_0(\mathbb{Z}, C(X, \mathbb{Z}))$$

from Theorem 3.5. Since $H_0(\mathbb{Z}, C(X, \mathbb{Z}))$ is torsion-free, it follows that its subgroup $(1 + \sigma_*)H_0(\mathbb{Z}, C(X, \mathbb{Z}))$ is torsion-free as well. Since tr is surjective, we conclude from [6, Section 23 (H)] that $\ker \text{tr}$ is a pure subgroup of $H_0(\mathbb{Z} \rtimes \mathbb{Z}_2, C(X, \mathbb{Z}))$. By Theorem 3.5, we have that $\ker \text{tr} \simeq \mathbb{Z}_2$. Therefore, [6, Theorem 24.1] implies that $\ker \text{tr}$ is a direct summand of $H_0(\mathbb{Z} \rtimes \mathbb{Z}_2, C(X, \mathbb{Z}))$. This concludes the proof of (10).

The remaining computations of the homology groups in cases (ii) and (iii) are a consequence of Theorem 3.1 and Lemmas 3.2, 3.3, and 3.4. In case (iii), since the \mathbb{Z} -action induced by φ is free, notice that Fix_σ is disjoint from $\text{Fix}_{\varphi\sigma}$. ■

Let us now apply Theorem 3.6 to an example which had its K-theory computed in [3, Corollary 4.4].

Example 3.7. Fix $\theta \in (0, 1)$ an irrational number. Let \tilde{X} be the set obtained from \mathbb{R} by replacing each $t \in \mathbb{Z} + \theta\mathbb{Z}$ by two elements $\{t^-, t^+\}$, and endow \tilde{X} with the order topology. Notice that there is an action $\mathbb{Z} \curvearrowright_\alpha \tilde{X}$ by translations ($\alpha_n(x) = n + x$). We let $X := \tilde{X}/\mathbb{Z}$. Then X is homeomorphic to the Cantor set and there is a minimal homeomorphism $R_\theta: X \rightarrow X$ given by $R_\theta(x) = x + \theta$.

Furthermore, there is an involutive homeomorphism σ on X given by $\sigma(x) = -x$ and $\sigma(t^\pm) = t^\mp$. Notice that the only fixed point of σ is $\frac{1}{2}$, and the only fixed points of $R_\theta \circ \sigma$ are $\frac{1+\theta}{2}$ and $\frac{\theta}{2}$.

It was shown in [17, Theorem 2.1] that $H_0(\mathbb{Z}, C(X, \mathbb{Z})) \simeq \mathbb{Z}^2$ and the generators are $[1_{[0^+, \theta^+]})]$ and $[1_{[\theta^+, 1^+]})]$. Observe that σ_* acts trivially on these two elements.

From these observations and Theorem 3.6, it follows that, for $k \geq 1$,

$$\begin{aligned} H_0(\mathbb{Z} \rtimes \mathbb{Z}_2, C(X, \mathbb{Z})) &\simeq \mathbb{Z}^2, \\ H_{2k-1}(\mathbb{Z} \rtimes \mathbb{Z}_2, C(X, \mathbb{Z})) &\simeq \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2, \\ H_{2k}(\mathbb{Z} \rtimes \mathbb{Z}_2, C(X, \mathbb{Z})) &= 0. \end{aligned}$$

Remark 3.8. In [14], Matui conjectured that if G is an ample, effective, second countable, minimal groupoid with unit space homeomorphic to the Cantor set, then

$$K_*(C_r^*(G)) \simeq \bigoplus_{n=0}^{\infty} H_{2n+*}(G)$$

for $* = 0, 1$ (*HK conjecture*). In [19], the (transformation groupoids of the) $\mathbb{Z} \rtimes \mathbb{Z}_2$ -odometers from Example 2.7 were shown to be counterexamples to the HK conjecture.

Let α be a minimal action of $\mathbb{Z} \rtimes \mathbb{Z}_2$ on the Cantor set X . If α is not free, then the associated crossed product $C(X) \rtimes \mathbb{Z} \rtimes \mathbb{Z}_2$ is AF [3, Theorem 3.5], hence $K_1(C(X) \rtimes \mathbb{Z} \rtimes \mathbb{Z}_2) = 0$. On the other hand, it follows from Proposition 2.8 and Theorem 3.6 that $H_{2n+1}(\mathbb{Z} \rtimes \mathbb{Z}_2, C(X, \mathbb{Z})) \neq 0$. Therefore, α is a counterexample to the HK conjecture.

If α is free, it follows from [22, Theorems 4.30 and 4.42] and Theorem 3.6 that α satisfies the HK conjecture. Alternatively, this also follows from Theorem 3.6 and [16, Remark 4.7].

Remark 3.9. Given an ample groupoid G with compact unit space, the *topological full group* of G , denoted by $[[G]]$, is the group of compact-open bisections U such that $r(U) = s(U) = G^{(0)}$.

Given an effective, minimal, second countable groupoid G with compact unit space homeomorphic to the Cantor set, Matui conjectured in [14] that the index map $I: [[G]]_{\text{ab}} \rightarrow H_1(G)$ is surjective and that its kernel is a quotient of $H_0(G) \otimes \mathbb{Z}_2$ under a certain canonical map (*AH conjecture*). Assuming that G is also minimal and almost finite, Matui proved that the index map is surjective and if G is also principal, then it satisfies AH conjecture.

We have not been able to verify whether the AH conjecture holds for non-free Cantor minimal $\mathbb{Z} \rtimes \mathbb{Z}_2$ -systems in general, but note that in [19] it was shown that the $\mathbb{Z} \rtimes \mathbb{Z}_2$ -odometers from Example 2.7 satisfy it.

Acknowledgments. This work was carried out during the tenure of an ERCIM “Alain Bensoussan” Fellowship Programme. The authors would like to thank the referee for a careful reading of the paper and the helpful comments.

References

- [1] M. Abért and G. Elek, Non-abelian free groups admit non-essentially free actions on rooted trees. 2007, arXiv:[0707.0970](https://arxiv.org/abs/0707.0970)
- [2] P. Ara, C. Bönicke, J. Bosa, and K. Li, Strict comparison for C^* -algebras arising from almost finite groupoids. *Banach J. Math. Anal.* **14** (2020), no. 4, 1692–1710 Zbl [1454.46052](https://zbmath.org/?q=1454.46052) MR [4133645](https://mathscinet.org/mr/4133645)
- [3] O. Bratteli, D. E. Evans, and A. Kishimoto, Crossed products of totally disconnected spaces by $Z_2 * Z_2$. *Ergodic Theory Dynam. Systems* **13** (1993), no. 3, 445–484 Zbl [0819.46049](https://zbmath.org/?q=0819.46049) MR [1245825](https://mathscinet.org/mr/1245825)
- [4] K. S. Brown, *Cohomology of groups. Corrected reprint of the 1982 original*. Grad. Texts in Math. 87, Springer, New York, 1994 MR [1324339](https://mathscinet.org/mr/1324339)
- [5] T. Downarowicz and G. Zhang, Symbolic extensions of amenable group actions and the comparison property. 2019, arXiv:[1901.01457](https://arxiv.org/abs/1901.01457)
- [6] L. Fuchs, *Abelian groups*. Int. Ser. Monogr. Pure Appl. Math., Pergamon Press, New York, 1960 Zbl [0100.02803](https://zbmath.org/?q=0100.02803) MR [0111783](https://mathscinet.org/mr/0111783)
- [7] A. Ioana, Cocycle superrigidity for profinite actions of property (T) groups. *Duke Math. J.* **157** (2011), no. 2, 337–367 Zbl [1235.37005](https://zbmath.org/?q=1235.37005) MR [2783933](https://mathscinet.org/mr/2783933)
- [8] A. Kar, P. Kropholler, and N. Nikolov, On growth of homology torsion in amenable groups. *Math. Proc. Cambridge Philos. Soc.* **162** (2017), no. 2, 337–351 Zbl [1383.20035](https://zbmath.org/?q=1383.20035) MR [3604918](https://mathscinet.org/mr/3604918)
- [9] S. Kawamura, H. Takemoto, and J. Tomiyama, State extensions in transformation group C^* -algebras. *Acta Sci. Math. (Szeged)* **54** (1990), 191–200 Zbl [0741.46031](https://zbmath.org/?q=0741.46031) MR [1073433](https://mathscinet.org/mr/1073433)
- [10] D. Kerr, Dimension, comparison, and almost finiteness. *J. Eur. Math. Soc. (JEMS)* **22** (2020), no. 11, 3697–3745 Zbl [1465.37010](https://zbmath.org/?q=1465.37010) MR [4167017](https://mathscinet.org/mr/4167017)
- [11] D. Kerr and G. Szabó, Almost finiteness and the small boundary property. *Comm. Math. Phys.* **374** (2020), no. 1, 1–31 Zbl [1446.46040](https://zbmath.org/?q=1446.46040) MR [4066584](https://mathscinet.org/mr/4066584)
- [12] X. Li, Partial transformation groupoids attached to graphs and semigroups. *Int. Math. Res. Not. IMRN* **2017** (2017), no. 17, 5233–5259 Zbl [1405.37007](https://zbmath.org/?q=1405.37007) MR [3694599](https://mathscinet.org/mr/3694599)
- [13] H. Matui, Homology and topological full groups of étale groupoids on totally disconnected spaces. *Proc. Lond. Math. Soc. (3)* **104** (2012), no. 1, 27–56 Zbl [1325.19001](https://zbmath.org/?q=1325.19001) MR [2876963](https://mathscinet.org/mr/2876963)
- [14] H. Matui, Étale groupoids arising from products of shifts of finite type. *Adv. Math.* **303** (2016), 502–548 Zbl [1352.19003](https://zbmath.org/?q=1352.19003) MR [3552533](https://mathscinet.org/mr/3552533)
- [15] V. Nekrashevych, Palindromic subshifts and simple periodic groups of intermediate growth. *Ann. of Math. (2)* **187** (2018), no. 3, 667–719 Zbl [1437.20038](https://zbmath.org/?q=1437.20038) MR [3779956](https://mathscinet.org/mr/3779956)
- [16] V. Proietti and M. Yamashita, Homology and K -theory of dynamical systems I. Torsion-free ample groupoids. *Ergodic Theory Dynam. Systems* **42** (2022), no. 8, 2630–2660 MR [4448401](https://mathscinet.org/mr/4448401)
- [17] I. F. Putnam, The C^* -algebras associated with minimal homeomorphisms of the Cantor set. *Pacific J. Math.* **136** (1989), no. 2, 329–353 Zbl [0631.46068](https://zbmath.org/?q=0631.46068) MR [978619](https://mathscinet.org/mr/978619)
- [18] I. F. Putnam, Lecture notes on C^* -algebras. 2019, available at www.math.uvic.ca/faculty/putnam/ln/C*-algebras.pdf
- [19] E. Scarparo, Homology of odometers. *Ergodic Theory Dynam. Systems* **40** (2020), no. 9, 2541–2551 Zbl [1460.46065](https://zbmath.org/?q=1460.46065) MR [4130816](https://mathscinet.org/mr/4130816)
- [20] Y. Suzuki, Almost finiteness for general étale groupoids and its applications to stable rank of crossed products. *Int. Math. Res. Not. IMRN* **2020** (2020), no. 19, 6007–6041 Zbl [0747.1405](https://zbmath.org/?q=0747.1405) MR [4165470](https://mathscinet.org/mr/4165470)

- [21] R. G. Swan, Groups of cohomological dimension one. *J. Algebra* **12** (1969), 585–610
Zbl [0188.07001](#) MR [240177](#)
- [22] K. Thomsen, The homoclinic and heteroclinic C^* -algebras of a generalized one-dimensional solenoid. *Ergodic Theory Dynam. Systems* **30** (2010), no. 1, 263–308 Zbl [1194.37003](#)
MR [2586354](#)
- [23] C. A. Weibel, *An introduction to homological algebra*. Cambridge Stud. Adv. Math. 38, Cambridge University Press, Cambridge, 1994 Zbl [0797.18001](#) MR [1269324](#)

Received 14 July 2020.

Eduard Ortega

Department of Mathematical Sciences, Norwegian University of Science and Technology (NTNU),
7491 Trondheim, Norway; eduard.ortega@ntnu.no

Eduardo Scarparo

Center for Engineering, Federal University of Pelotas, R. Benjamin Constant 989, Pelotas, RS,
CEP 96010-020, Brazil; eduardo.scarparo@ufpel.edu.br