

Quasi-isometry invariance of relative filling functions

Sam Hughes, Eduardo Martínez-Pedroza, and Luis Jorge Sánchez Saldaña
(with an appendix by Ashot Minasyan)

Abstract. For a finitely generated group G and collection of subgroups \mathcal{P} , we prove that the relative Dehn function of a pair (G, \mathcal{P}) is invariant under quasi-isometry of pairs. Along the way, we show quasi-isometries of pairs preserve almost malnormality of the collection and fineness of the associated coned-off Cayley graphs. We also prove that for a cocompact simply connected combinatorial G -2-complex X with finite edge stabilisers, the combinatorial Dehn function is well defined if and only if the 1-skeleton of X is fine. We also show that if H is a hyperbolically embedded subgroup of a finitely presented group G , then the relative Dehn function of the pair (G, H) is well defined. In the appendix, it is shown that the Baumslag–Solitar group $BS(k, l)$ has a well-defined Dehn function with respect to the cyclic subgroup generated by the stable letter if and only if neither k divides l nor l divides k .

1. Introduction

The main objects of study in this article are pairs (G, \mathcal{P}) , where G is a *finitely generated group* with a chosen word metric dist_G , and \mathcal{P} is a *finite collection of subgroups*; note that *these assumptions will stand throughout the introduction*.

Let hdist_G denote the Hausdorff distance between subsets of G , and let G/\mathcal{P} denote the collection of left cosets gP for $g \in G$ and $P \in \mathcal{P}$.

For constants $L \geq 1$, $C \geq 0$ and $M \geq 0$, an (L, C, M) -quasi-isometry of pairs

$$q: (G, \mathcal{P}) \rightarrow (H, \mathcal{Q})$$

is an (L, C) -quasi-isometry $q: G \rightarrow H$ such that the relation

$$\{(A, B) \in G/\mathcal{P} \times H/\mathcal{Q} : \text{hdist}_H(q(A), B) < M\}$$

satisfies the condition that the projections to G/\mathcal{P} and H/\mathcal{Q} are surjective.

This article is part of the program of investigating which properties of pairs (G, \mathcal{P}) are invariant under quasi-isometry of pairs. There are recent results in this direction. For example, it is a consequence of the quasi-isometric rigidity of relative hyperbolicity [1],

that if (G, \mathcal{P}) is a relatively hyperbolic pair, \mathcal{P} is a collection of non-relatively hyperbolic groups, and (G, \mathcal{P}) and (H, \mathcal{Q}) are quasi-isometric pairs, then H is hyperbolic relative to a refinement of \mathcal{Q} as defined below. Under natural assumptions, quasi-isometries of pairs between relatively hyperbolic pairs induce canonical homeomorphisms between their Bowditch boundaries [11] and canonical isomorphisms of JSJ trees [10]. Outside the framework of relatively hyperbolic groups, it is known that quasi-isometries of pairs preserve the number of Bowditch’s filtered ends [19] and under technical hypotheses acylindrical hyperbolicity [13]. For a recent survey, we direct the reader to [14].

For a pair (G, \mathcal{P}) , Osin introduced the notions of *finite relative presentation* and *relative Dehn function* $\Delta_{G,\mathcal{P}}$ as natural generalisations of their standard counterparts for finitely generated groups, see [23]. These notions characterise relatively hyperbolic pairs (G, \mathcal{P}) as the ones which are relatively finitely presented and have relative Dehn function bounded from above by a linear function. By quasi-isometric rigidity of relative hyperbolicity, among relatively finitely presented pairs, quasi-isometries of pairs preserve having linear relative Dehn function.

The main result of this article confirms the natural expectation that among relatively finitely presented pairs, quasi-isometric pairs have equivalent relative Dehn functions. This is not an elementary statement, as we describe below.

Convention 1.1 ($\Delta_{G,\mathcal{P}}$ is well defined). By $\Delta_{G,\mathcal{P}}$ is well defined, we mean that G is finitely presented relative to \mathcal{P} and the relative Dehn function $\Delta_{G,\mathcal{P}}$ takes only finite values with respect to a finite relative presentation of G and \mathcal{P} . From here on, when we refer to a relative Dehn function, we always assume that it has been defined using a finite relative presentation.

Let \mathcal{P} be a collection of subgroups of group G . A *refinement* \mathcal{P}^* of \mathcal{P} is a set of representatives of conjugacy classes of the collection of subgroups $\{\text{Comm}_G(gPg^{-1}): P \in \mathcal{P} \text{ and } g \in G\}$, where $\text{Comm}_G(P)$ denotes the commensurator of the subgroup P in G .

Theorem A. *Let $(G, \mathcal{P}) \rightarrow (H, \mathcal{Q})$ be a quasi-isometry of pairs and let \mathcal{P}^* be a refinement of \mathcal{P} . If the relative Dehn function $\Delta_{H,\mathcal{Q}}$ is well defined, then Δ_{G,\mathcal{P}^*} is well defined and $\Delta_{G,\mathcal{P}^*} \asymp \Delta_{H,\mathcal{Q}}$.*

A phenomenon that occurs for pairs (G, \mathcal{P}) is that being relatively finitely presented does not imply that the relative Dehn function is well defined. This is in sharp contrast with the standard framework where a finitely presented group always has a well-defined Dehn function. The proof of Theorem A provides an insight into this phenomenon via the following results on which our argument relies on.

In the framework of relatively hyperbolic groups, Bowditch introduced the notion of fine graph [2]. A *circuit* in a simplicial graph is an embedded close path. A simplicial graph Γ is *fine* if for every $n \geq 0$ and every edge e in Γ , there are finitely many circuits of length less than or equal to n which contain e . This is weaker than the graph being locally finite. The relationship between this notion and isoperimetric functions was made explicit by Groves and Manning [8, Proposition 2.50, Question 2.51]. The following result can be

interpreted as a homotopical version of [17, Theorem 1.3] where an analogous statement is proved for homological Dehn functions.

Theorem B (Theorem 2.1). *Let X be a cocompact simply connected combinatorial G -2-complex with finite edge stabilisers. The combinatorial Dehn function Δ_X of X takes only finite values if and only if the 1-skeleton of X is a fine graph.*

It is obvious that being fine is not a property preserved by quasi-isometries in the class of graphs. For a pair (G, \mathcal{P}) , together with a finite generating set S of G , one can assign a connected and cocompact G -graph known as the coned-off Cayley graph $\widehat{\Gamma}(G, \mathcal{P}, S)$; a notion introduced by Farb [5], see Definition 4.6. It is an observation that the quasi-isometry type of $\widehat{\Gamma}(G, \mathcal{P}, S)$ is independent of the finite generating set S ; throughout the introduction $\widehat{\Gamma}(G, \mathcal{P})$ denotes the coned-off Cayley graph with respect to some finite generating set of G . In this framework, under some assumptions, we are able to prove that fineness is preserved under quasi-isometry of pairs in the class of coned-off Cayley graphs. A collection of subgroups \mathcal{P} of a group G is *reduced* if for any $P, Q \in \mathcal{P}$ and $g \in G$, then P and gQg^{-1} being commensurable subgroups implies $P = Q$ and $g \in P$.

Theorem C (Theorem 5.15). *Let $q: (G, \mathcal{P}) \rightarrow (H, \mathcal{Q})$ be a quasi-isometry of pairs. Suppose that \mathcal{P} and \mathcal{Q} are reduced. Then there is an induced quasi-isometry of graphs $\widehat{q}: \widehat{\Gamma}(G, \mathcal{P}) \rightarrow \widehat{\Gamma}(H, \mathcal{Q})$, and if $\widehat{\Gamma}(H, \mathcal{Q})$ is a fine graph, then $\widehat{\Gamma}(G, \mathcal{P})$ is a fine graph.*

The condition that the coned-off Cayley graph $\widehat{\Gamma}(G, \mathcal{P})$ is fine forces the collection \mathcal{P} to be almost malnormal (see Definition 6.5). It is an observation that any almost malnormal collection of infinite subgroups is reduced. We prove that the property of being almost malnormal is preserved under quasi-isometry of pairs up to taking a refinement.

Theorem D (Theorem 6.12). *Let $q: (G, \mathcal{P}) \rightarrow (H, \mathcal{Q})$ be a quasi-isometry of pairs. If \mathcal{Q} is an almost malnormal collection of infinite subgroups, then any refinement \mathcal{P}^* of \mathcal{P} is almost malnormal and $q: (G, \mathcal{P}^*) \rightarrow (H, \mathcal{Q})$ is a quasi-isometry of pairs.*

The previous results can be linked to Osin’s definition of relative Dehn function $\Delta_{G, \mathcal{P}}$ of a relatively finitely presented pair (G, \mathcal{P}) via the following result. A connected graph Γ is called *fillable* if, when considering Γ with the length metric obtained by regarding each edge as a segment of length one, there is an integer k such that the coarse isoperimetric function f_k^Γ takes only finite values, see Section 3 for definitions.

Theorem E (See Theorem 4.17). *If (G, \mathcal{P}) is a relatively finitely presented pair, then*

- (1) $\widehat{\Gamma}(G, \mathcal{P})$ is fillable.
- (2) The relative Dehn function $\Delta_{G, \mathcal{P}}$ is well defined if and only if $\widehat{\Gamma}(G, \mathcal{P})$ is fine graph.

Conversely, if $\widehat{\Gamma}(G, \mathcal{P})$ is fine and fillable, then (G, \mathcal{P}) is a relatively finitely presented pair, and hence $\Delta_{G, \mathcal{P}}$ is well defined.

The following result is a restatement of a result of Osin [23, Theorem 2.53], see Proposition 4.8. This statement allows us to translate his definition of relative Dehn function to the realm of coarse isoperimetric functions of coned-off Cayley graphs.

Theorem F (Osin). *Let G be a group and let \mathcal{P} be a collection of subgroups. Suppose that $\Delta_{G,\mathcal{P}}$ is well defined. Then $\Delta_{G,\mathcal{P}}$ is equivalent to the coarse isoperimetric function $f_N^{\widehat{\Gamma}(G,\mathcal{P})}$ of $\widehat{\Gamma}(G,\mathcal{P})$ for all sufficiently large integers N .*

Let us describe the argument proving Theorem A using the results that have been stated.

Proof of Theorem A. Let us first observe that we can assume that the collections \mathcal{P} and \mathcal{Q} contain only infinite subgroups. First note that if \mathcal{P}_∞ and \mathcal{Q}_∞ are the collections obtained by removing finite subgroups from \mathcal{P} and \mathcal{Q} , respectively, then $q: (G, \mathcal{P}_\infty) \rightarrow (H, \mathcal{Q}_\infty)$ is a quasi-isometry of pairs as well. Moreover, for an arbitrary pair (K, \mathcal{L}) , adding or removing a finite subgroup of K to \mathcal{L} preserves having a well-defined relative Dehn function, and if the functions are well defined, they are equivalent, see, for example, [23, Theorem 2.40].

Assume that \mathcal{P} and \mathcal{Q} consist only of infinite subgroups. Since (H, \mathcal{Q}) is relatively finitely presented and $\Delta_{H,\mathcal{Q}}$ is well defined, Theorem E implies that $\widehat{\Gamma}(H, \mathcal{Q})$ is fillable and fine. Since $\widehat{\Gamma}(H, \mathcal{Q})$ is a fine graph, it follows that \mathcal{Q} is an almost malnormal collection. Then Theorem D implies that \mathcal{P}^* is an almost malnormal collection. Hence, both \mathcal{Q} and \mathcal{P}^* are reduced collections, and $q: (G, \mathcal{P}^*) \rightarrow (H, \mathcal{Q})$ is a quasi-isometry of pairs. Now we can invoke Theorem C to obtain a quasi-isometry $\widehat{q}: \widehat{\Gamma}(G, \mathcal{P}^*) \rightarrow \widehat{\Gamma}(H, \mathcal{Q})$ and also obtain that $\widehat{\Gamma}(G, \mathcal{P}^*)$ is fine. It is a standard result in the literature that being fillable is a property preserved by quasi-isometry in the class of connected graphs, and any two quasi-isometric graphs have equivalent coarse isoperimetric functions (see, for instance, [3, Proposition III.H.2.2]). The quasi-isometry \widehat{q} implies that $\widehat{\Gamma}(G, \mathcal{P}^*)$ is fillable and both $\widehat{\Gamma}(G, \mathcal{P}^*)$ and $\widehat{\Gamma}(H, \mathcal{Q})$ have equivalent coarse isoperimetric inequalities. Then Theorem E implies that (G, \mathcal{P}^*) is relatively finitely presented and Δ_{G,\mathcal{P}^*} is well defined. The proof concludes by invoking Theorem F. ■

In the class of finitely generated groups, being finitely presented is a quasi-isometry invariant. We do not know the answer to the following general question.

Question 1.2. Suppose that $q: (G, \mathcal{P}) \rightarrow (H, \mathcal{Q})$ is a quasi-isometry of pairs and (H, \mathcal{Q}) is relatively finitely presented. Is (G, \mathcal{P}) relatively finitely presented?

There is a rich class of pairs (G, \mathcal{P}) with well-defined relative Dehn function. Hyperbolically embedded subgroups were introduced in [4] by Dahmani, Guirardel and Osin. Given a group G , $X \subset G$ and $H \leq G$, let $H \hookrightarrow_h (G, X)$ denote that H is a hyperbolically embedded subgroup of G with respect to X .

Theorem G (Theorem 7.2). *Let G be a finitely presented group and $H \leq G$ be a subgroup. If $H \hookrightarrow_h G$, then the relative Dehn function $\Delta_{G,H}$ is well defined.*

In the context of Theorem G, the relative Dehn function $\Delta_{G,\mathcal{P}}$ is bounded from above by a linear function if and only if G is hyperbolic relative to H , see [23]. It is well known that the class of pairs (G, H) such that $H \hookrightarrow_n G$ properly extends relative hyperbolicity, for examples, see [4].

In a preliminary version of this manuscript, we asked whether there exist pairs (G, \mathcal{P}) such that $\Delta_{G,\mathcal{P}}$ is well defined, but \mathcal{P} is not hyperbolically embedded in G . In this regard, consider the Baumslag–Solitar groups $BS(k, l) = \langle a, t \mid ta^k t^{-1} = a^l \rangle$, where $k, l \in \mathbb{Z} \setminus \{0\}$. In Example 7.6, we show that $BS(k, l)$ does not have a well-defined Dehn function with respect to the cyclic subgroup generated by the stable letter t if either k divides l or l divides k . On the other hand, in the appendix Ashot Minasyan shows that the converse holds, that is, if neither $k \mid l$ nor $l \mid k$, then $\Delta_{BS(k,l),(t)}$ is well defined.

2. Combinatorial Dehn functions and fine graphs

The goal of this section is to prove Theorem B. We use the notion of disk diagram in a combinatorial complex; for definitions see, for example, [20]. We begin by recalling the definition of a combinatorial Dehn function, then we prove each direction of Theorem B individually as Lemmas 2.3 and 2.4. Note that Lemma 2.4 does not require the hypothesis of finite edge stabilisers.

Suppose X is a combinatorial 2-complex and let $c: S^1 \rightarrow X$ be a closed path in $X^{(1)}$ that is null-homotopic in X . Then there is a disk diagram $i: D^2 \rightarrow X$ spanning c , that is, i is a combinatorial map and $i(\partial D^2) = c$. Let $\text{Area}(D)$ denote the number of faces of D and define

$$\delta_X(c) := \min\{\text{Area}(D) : D \text{ is a disk spanning } c\},$$

the *combinatorial Dehn function* Δ_X of X is defined to be

$$\Delta_X(n) := \max\{\delta_X(c) : c \text{ is a closed path in } X^{(1)}, \text{ null-homotopic in } X, \text{ with } |c| \leq n\}.$$

Unless otherwise stated, all graphs in this article are assumed to be simplicial. A *circuit* in a simplicial graph is an embedded close path. We recall the following definition due to Bowditch [2, Proposition 2.1]. A graph Γ is *fine* if for every $n \geq 0$ and every edge e in Γ , there are finitely many circuits of length less than or equal to n which contains e .

Theorem 2.1 (Theorem B). *Let X be a cocompact simply connected combinatorial G -2-complex with finite edge stabilisers. The combinatorial Dehn function Δ_X of X takes only finite values if and only if the 1-skeleton of X is a fine graph.*

The next three lemmas prove the theorem. The method is essentially a van Kampen diagram approach to the proof of [17, Theorem 1.3]. The first lemma is a triviality.

Lemma 2.2. *Let X be a cocompact simply connected combinatorial G -2-complex with finite edge stabilisers. Then each edge is contained in finitely many 2-cells.*

The next lemma proves the “only if” direction of Theorem 2.1.

Lemma 2.3. *Let X be a cocompact simply connected combinatorial G -2-complex with finite edge stabilisers. If the combinatorial Dehn function Δ_X of X is well defined, then $X^{(1)}$ is a fine graph.*

Proof. Let D be a cellular 2-disk. We say D is *golden* if D has an enumeration of its 2-cells f_1, \dots, f_k with the property that ∂f_{i+1} contains a 1-cell of the subcomplex induced by $f_1 \cup \dots \cup f_i$, and there is a cellular map $D \rightarrow X$.

Let R be a 2-cell and $f_1: R \rightarrow X$, then a simple counting argument yields there are only finitely many golden disks with at most $n \geq 0$ faces making the following diagram commute:

$$\begin{array}{ccc} R & \xrightarrow{f_1} & X \\ \downarrow & \nearrow & \\ D. & & \end{array}$$

Observe that by taking the minimal area filling for a circuit c of length n in X gives rise to a golden disk D with at most $\Delta_X(n)$ many 2-cells. Now, there are only finitely many 2-cells containing a given edge e , so by the previous paragraph, there are only finitely many golden disks D containing e with at most $\Delta_X(n)$ many 2-cells. In particular, for each $n \geq 0$, there are only finitely many circuits in X of length less than or equal n containing e . It follows that $X^{(1)}$ is a fine graph. ■

The next lemma proves the “if” direction of Theorem 2.1. Note that we can drop the hypothesis of finite edge stabilisers.

Lemma 2.4. *Let X be a cocompact simply connected combinatorial G -2-complex. If $X^{(1)}$ is a fine graph, then the combinatorial Dehn function Δ_X of X is well defined.*

Proof. Let Y_n denote the set of circuits of length less than or equal to n in X .

Claim. *The set Y_n is a G -set with finitely many orbits.*

Let $\{e_1, \dots, e_r\}$ be edges representing the orbits of the G -action on $X^{(1)}$. Every circuit in X of length less than or equal to n can be translated to contain some e_i , the claim now follows from fineness of $X^{(1)}$.

Let A_n be an upper bound for the area of a circuit of length less than or equal to n in X , and it is well defined by the previous claim. Let γ be a closed path without backtracks in X , then γ can be expressed as a concatenation of closed paths $\gamma_1 \dots \gamma_k$ such that $1 \leq k \leq \text{Len}(\gamma)$, for $i = 1, \dots, k$, we have $\text{Len}(\gamma_i) \leq \text{Len}(\gamma)$ and each γ_i is a circuit. Now, filling each γ_i , we have

$$\text{Area}(\gamma) \leq \sum_{i=1}^k \text{Area}(\gamma_i) \leq \sum_{i=1}^k A_{\text{Len}(\gamma_i)} \leq k A_{\text{Len}(\gamma)} \leq \text{Len}(\gamma) A_{\text{Len}(\gamma)}.$$

This yields a finite upper bound for $\Delta_X(\ell)$ and so we conclude that Δ_X is well defined. ■

Remark 2.5. One can define a combinatorial Dehn function of a 2-complex using circuits instead of arbitrary closed paths. Let us call this function Δ_X^{circ} . In this case,

$$\Delta_X^{\text{circ}}(n) \leq \Delta_X(n) \leq \overline{\Delta_X^{\text{circ}}}(n),$$

where $\overline{\Delta_X^{\text{circ}}}$ is the superadditive closure of Δ_X^{circ} . We do not know whether $\Delta_X^{\text{circ}}(n)$ is equivalent to $\Delta_X(n)$. This resembles a conjecture of Mark Sapir of whether the Dehn function of a finitely presented group is equivalent to a superadditive function (see [9]).

3. Coarse isoperimetric functions

To prove quasi-isometry invariance, we will use the less general version of ε -fillings for graphs and 2-complexes defined in [23]. The original definition, set up for essentially arbitrary metric spaces, can be found in [3, Chapter III.H.2]. The main result of this section is Proposition 3.2 – a generalisation of a result of Osin [23, Theorem 2.53] alluded to in the introduction.

Let X be a 2-complex. A *singular combinatorial loop* $c: S^1 \rightarrow X$ is a combinatorial structure on S^1 and a continuous map such that for every open cell of S^1 , either $f|_e$ is a homeomorphism onto an open cell of X , or else $f(e)$ is contained in the 0-skeleton of X .

Let c be a combinatorial cycle in X . An ε -*filling* of c is a pair (P, Φ) consisting of a triangulation P of a 2-disk D^2 and a singular combinatorial map $\Phi: P^{(1)} \rightarrow X^{(1)}$, such that $\Phi|_{S^1} = c$ and the image under Φ of each face of P is a set of diameter at most ε . Define $|\Phi|$ to be the number of faces of Φ and

$$\text{Area}_\varepsilon(c) := \min\{|\Phi|: (P, \Phi) \text{ an } \varepsilon\text{-filling of } c\}.$$

The *coarse isoperimetric function* of X is then defined to be

$$f_\varepsilon^X(\ell) := \sup\{\text{Area}_\varepsilon(c): \text{Len}(c) \leq \ell\}.$$

Definition 3.1. For two functions $f, g: \mathbb{N} \rightarrow \mathbb{N}$, we say that f is *asymptotically less than* g , and we write $f \preceq g$ if there exist constants $C, K, L \in \mathbb{N}$ such that

$$f(n) \leq Cg(Kn) + Ln.$$

Further, we say f is *asymptotically equivalent* to g , and write $f \asymp g$ if $f \preceq g$ and $g \preceq f$.

Proposition 3.2. *Let X be a cocompact simply connected combinatorial G -2-complex. If Δ_X takes only finite values, then for $N \in \mathbb{N}$ large enough, f_N^X takes only finite values and $f_N^X \asymp \Delta_X$.*

Proof. Since X is a cocompact G -2-complex, there are only finitely many G -orbits of 2-cells in X . Let $\{D_1, \dots, D_n\}$ denote a representative set of orbits and let N be an integer greater than the maximum diameter of each disk D_i for $i = 1, \dots, n$.

First, we will show $f_N^X \leq \Delta_X$. Let $c: S^1 \rightarrow X$ be a singular combinatorial loop. Let $\Phi: D \rightarrow X$ be a disk diagram of minimal area that fills $c: S^1 \rightarrow X$. Barycentric subdividing D twice to obtain D'' yields a simplicial disk such that the image of each face in X has diameter less than N , i.e., (D'', Φ) is an N -filling of c . It follows that $\text{Area}_\varepsilon(c) \leq 12N\Delta_X(\text{Len}(c))$. In particular, $f_N^X \leq \Delta_X$.

It remains to show that $\Delta_X \leq f_N^X$. Consider an N -filling (P, Φ) of a combinatorial loop c in $X^{(1)}$. Considering (P, Φ) as a $3N$ -filling, we may assume that each 0-cell of P maps to a 0-cell of X and each 1-cell of P maps to an edge path in X of length at most N . Thus, after subdividing P at most N times, we may assume that Φ is cellular on $P^{(1)}$. For each 2-cell of the subdivided P , its boundary map determines a cellular loop in X with length bounded by $3N$. Now, we fill each such loop with some disk diagram $D \rightarrow X$ to obtain a diagram for c in X which has area at most $\Delta_X(3N)f_{3N}^X(\text{Len}(c))$. In particular, we conclude that $\Delta_X(\text{Len}(c)) \leq \Delta_X(3N)f_{3N}^X(\text{Len}(c))$. ■

A connected graph Γ is *fillable* if, when considering Γ with the length metric obtained by regarding each edge as a segment of length one, there is an integer k such that the coarse isoperimetric function f_k^Γ takes only finite values.

Proposition 3.3 ([3, Proposition III.H.2.2]). *If Γ and Γ' are quasi-isometric connected graphs such that Γ is fillable, then Γ' is fillable and $f_k^\Gamma \asymp f_k^{\Gamma'}$ for large enough k .*

Remark 3.4. If a connected graph Γ is fillable, then there is a positive integer m such that the complex obtained by attaching 2-cells to all circuits of length less than or equal to m is simply connected.

4. Relative Dehn functions of groups

Definition 4.1 (Finite relative presentation). Let G be a group, \mathcal{P} an arbitrary collection of subgroups of G , and let S be a subset of G . We say that G is *generated by S relative to \mathcal{P}* if G is generated by the set $\mathcal{S} = S \sqcup \bigsqcup_{P \in \mathcal{P}} (P - \{1\})$, equivalently, the natural homomorphism

$$F = F(S) * \bigstar_{P \in \mathcal{P}} P \rightarrow G \tag{4.1}$$

is surjective. In the case where S is finite, G is relatively finitely generated with respect to \mathcal{P} .

Let $R \subseteq F$ be a set that normally generates the kernel of the above homomorphism, then we say

$$G = \langle S, \mathcal{P} \mid R \rangle \tag{4.2}$$

is a *presentation of G relative to \mathcal{P}* . If both S and R are finite, we say G is *relatively finitely presented with respect to \mathcal{P}* , or just, *relatively finitely presented* if the collection \mathcal{P} is clear from the context, and (4.1) is a *relative finite presentation*.

Definition 4.2 (Relative Dehn function of a relative presentation). Let $G = \langle S, \mathcal{P} \mid R \rangle$ be a relative presentation. For a word W over the alphabet $\mathcal{S} = S \sqcup \bigsqcup_{P \in \mathcal{P}} (P - \{1\})$ representing the trivial element in G , there is an expression

$$W = \prod_{i=1}^k f_i^{-1} R_i f_i, \tag{4.3}$$

where $R_i \in R$ and $f_i \in F$.

We say a function $f: \mathbb{N} \rightarrow \mathbb{N}$ is a *relative isoperimetric function* of the presentation $G = \langle S, \mathcal{P} \mid R \rangle$ if, for any $n \in \mathbb{N}$, and any word W as above of length $\leq n$, one can write W as in (4.3) with $k \leq f(n)$. The smallest relative isoperimetric function of $G = \langle S, \mathcal{P} \mid R \rangle$ is called the *relative Dehn function of G with respect to \mathcal{P}* , and it is denoted $\Delta_{G, \mathcal{P}}$.

Definition 3.1 and Theorem 4.3 below justify the notation $\Delta_{G, \mathcal{P}}$ for the relative Dehn function of G with respect to \mathcal{P} .

Theorem 4.3 ([23, Theorem 2.34]). *Let G be a finitely presented group relative to \mathcal{P} . Let Δ_1 and Δ_2 be the relative Dehn functions associated to two finite relative presentations. If Δ_1 takes only finite values, then Δ_2 takes only finite values, and $\Delta_1 \asymp \Delta_2$.*

Definition 4.4 (Osin–Cayley graph and Osin–Cayley complex). Assume G has a relative presentation as in (4.2). We call the Cayley graph $\Gamma(G, \mathcal{S})$ with $\mathcal{S} = S \sqcup \bigsqcup_{P \in \mathcal{P}} (P - \{1\})$ the *Osin–Cayley graph* and denote it by $\bar{\Gamma}(G, \mathcal{P}, S)$. Note that in general this graph is not simplicial.

For each $P \in \mathcal{P}$, denote by R_P the set of all words in the alphabet $P - \{1\}$ that represent the identity in P , that is, we have the presentation $P = \langle P - \{1\} \mid R_P \rangle$. We also have the following presentation:

$$F = \left\langle S, \bigsqcup_{P \in \mathcal{P}} (P - \{1\}) \mid \bigsqcup_{P \in \mathcal{P}} R_P \right\rangle.$$

The Osin–Cayley complex $\bar{X}(G, \mathcal{P}, S)$ is the 2-complex with 1-skeleton $\bar{\Gamma}(G, \mathcal{P}, S)$, and we attach

- One 2-cell for each loop labelled with a word in R , which we call from now on *R-cells*.
- One 2-cell for each loop labelled by a word in $\bigsqcup_{P \in \mathcal{P}} R_P$, which we call from now on *\mathcal{P} -cells*.

Remark 4.5. By [23, Definition 2.31], the relative Dehn function $\Delta_{G, \mathcal{P}}$ can be described as follows. For any combinatorial loop $\gamma: S^1 \rightarrow \bar{X}(G, \mathcal{P}, S)$, the relative area $\text{Area}^{\text{rel}}(\gamma)$ of γ is the number of R -cells in a minimal disk diagram for γ , where minimality is with respect to the number of R -cells. Then

$$\Delta_{G, \mathcal{P}}(n) = \max\{\text{Area}^{\text{rel}}(\gamma) : \gamma \text{ is a loop in } \bar{X}(G, \mathcal{P}, S) \text{ of length at most } n\}.$$

Definition 4.6 (Coned-off Cayley graph). Let G be a group, let \mathcal{P} be an arbitrary collection of subgroups of G , and let S be a generating set of G . Denote by G/\mathcal{P} the set of all cosets gP with $g \in G$ and $P \in \mathcal{P}$. The *coned-off Cayley graph of G with respect to \mathcal{P}* is the graph $\widehat{\Gamma}(G, \mathcal{P}, S)$ with the vertex set $G \cup G/\mathcal{P}$ and edges are of the following type:

- $\{g, gs\}$ for $s \in S$,
- $\{x, gP\}$ for $g \in G, P \in \mathcal{P}$ and $x \in gP$.

We call vertices of the form gP *cone points*.

Note that $\widehat{\Gamma}(G, \mathcal{P}, S)$ contains the Cayley graph of G with respect to the generating set S , and the quasi-isometry type of $\widehat{\Gamma}(G, \mathcal{P}, S)$ is independent of the finite generating set S of G . This justifies the notation $\widehat{\Gamma}(G, \mathcal{P})$ that we use throughout the article.

Definition 4.7 (A natural quasi-isometry between $\overline{\Gamma}(G, \mathcal{P}, S)$ and $\widehat{\Gamma}(G, \mathcal{P}, S)$). Assume G is generated by S relatively to \mathcal{P} . Let

$$\varphi: \overline{\Gamma}(G, \mathcal{P}, S) \rightarrow \widehat{\Gamma}(G, \mathcal{P}, S)$$

be the map defined as follows. Add a vertex at the midpoint of each edge $e = \{g, gh\}$ of $\overline{\Gamma}(G, \mathcal{P}, S)$ with $h \in P, p \in \mathcal{P}$, and label in P . Consider the inclusion of the vertex set of $\overline{\Gamma}(G, \mathcal{P}, S)$ into the vertex set of $\widehat{\Gamma} = \widehat{\Gamma}(G, \mathcal{P}, S)$. Observe that this map extends to a G -equivariant cellular map between $\overline{\Gamma}(G, \mathcal{P}, S)$ and $\widehat{\Gamma}(G, \mathcal{P}, S)$. Specifically, for an edge $e = \{g, gh\}$ with $h \in P$ and label in P of $\overline{\Gamma}(G, \mathcal{P}, S)$, the midpoint of e maps to the vertex gP ; an edge $\{g, gs\}$ with label in S is an edge that is common to both $\overline{\Gamma}(G, \mathcal{P}, S)$ and $\widehat{\Gamma}(G, \mathcal{P}, S)$. Observe that the map $\varphi: \overline{\Gamma}(G, \mathcal{P}, S) \rightarrow \widehat{\Gamma}(G, \mathcal{P}, S)$ is indeed a $(1, 1)$ -quasi-isometry.

Proposition 4.8. *Let G be a group and let \mathcal{P} be a collection of subgroups. If $\Delta_{G, \mathcal{P}}$ is well defined, then $\Delta_{G, \mathcal{P}} \simeq f_N^{\widehat{\Gamma}(G, \mathcal{P})}$ for all sufficiently large integers N .*

Proof. This is a restatement of Osin’s result [23, Theorem 2.53] modulo the fact that $\widehat{\Gamma}(G, \mathcal{P}, S)$ and the Cayley graph $\overline{\Gamma}(G, \mathcal{P}, S)$ are quasi-isometric graphs, see Definition 4.7. ■

Proposition 4.9. *Let G be a group, \mathcal{P} be a collection of subgroups, and S a relative generating set. Let \widetilde{G} be the free product $F(S) * \ast_{P \in \mathcal{P}} P$, and consider the short exact sequence,*

$$1 \rightarrow N \hookrightarrow \widetilde{G} \xrightarrow{\varphi} G \rightarrow 1,$$

where φ is the homomorphism induced by the inclusion $S \cup \bigcup_{i=1}^n P_i$ into G , and N is the kernel of φ . Then the coned-off Cayley graph $\widehat{\Gamma} = \widehat{\Gamma}(G, \mathcal{P}, S)$ is connected and for any vertex x_0 of $\widehat{\Gamma}$, there is a group isomorphism

$$N \rightarrow \pi_1(\widehat{\Gamma}, x_0), \quad g \mapsto [\gamma_g],$$

where γ_g is a combinatorial closed path in $\widehat{\Gamma}$ based at x_0 .

Proof. Consider the splitting of \tilde{G} as the fundamental group of the graph of groups \mathbf{Y} that consists of a vertex v labelled with the trivial group, one vertex v_P labelled with each $P \in \mathcal{P}$, respectively, one edge e_P that joins v with v_P for each $P \in \mathcal{P}$ labelled with the trivial group, and one edge loop e_s based at v for each $s \in S$ labelled with the trivial group.

Let \mathcal{T} be the Bass–Serre tree of \mathbf{Y} , see [24]. Since each subgroup P of \tilde{G} survives in the quotient G , we have that the subgroup N acts freely on \mathcal{T} , and the quotient map $\rho: \mathcal{T} \rightarrow \mathcal{T}/N$ is a covering map. Moreover, G acts on the quotient \mathcal{T}/N . We leave the reader to verify that the quotient \mathcal{T}/N is G -homeomorphic to the coned-off Cayley graph $\hat{\Gamma}(G, \mathcal{P}, S)$.

Fix a vertex \tilde{x}_0 of \mathcal{T} such that $\rho(\tilde{x}_0) = x_0$. Then any element g of N induces a unique embedded path α_g from \tilde{x}_0 to $g\tilde{x}_0$. Let $\gamma_g = \rho \circ \alpha_g$ and note it is a closed combinatorial path in $\hat{\Gamma}$ based at x_0 . Since \mathcal{T} is simply connected, standard covering space theory implies that the map $N \rightarrow \pi_1(\hat{\Gamma}, \rho(x_0))$ given by $g \mapsto [\gamma_g]$ is a group isomorphism. ■

Definition 4.10 (Coned-off Cayley complex $\hat{X}(G, \mathcal{P}, S)$). Consider a finite relative presentation $G = \langle S, \mathcal{P} \mid R \rangle$. The *coned-off Cayley complex* $\hat{X}(G, \mathcal{P}, S)$ of G is a 2-dimensional G -complex with 1-skeleton the coned-off Cayley graph $\hat{\Gamma}(G, \mathcal{P}, S)$ defined as follows.

We use the setup of Proposition 4.9. In particular, N is the normal subgroup of \tilde{G} generated by R , we have fixed a vertex x_0 of $\hat{\Gamma}$, and we have a group isomorphism $N \rightarrow \pi_1(\hat{\Gamma}, x_0)$ given by $g \mapsto [\gamma_g]$, where γ_g is a combinatorial closed path based at x_0 .

For $g \in G$ and $r \in R$, let $g \cdot \gamma_r$ be the translated closed path in $\hat{\Gamma}$ without an initial point, i.e., these are cellular maps from $S^1 \rightarrow \hat{\Gamma}$. Consider the G -set $\Omega = \{g \cdot \gamma_r : r \in R, g \in G\}$ of closed paths in $\hat{\Gamma}$. The complex \hat{X} is then obtained by attaching a 2-cell to $\hat{\Gamma}$ for every closed path in Ω . In particular, the pointwise G -stabiliser of a 2-cell of \hat{X} coincides with the pointwise G -stabiliser of its boundary path. The natural isomorphism from N to $\pi_1(\hat{\Gamma}, \rho(x_0))$ implies that \hat{X} is simply connected. Moreover, the G -action is cocompact since R is finite.

Definition 4.11 (A natural map between $\bar{X}(G, \mathcal{P}, S)$ and $\hat{X}(G, \mathcal{P}, S)$). There exists a G -map $\varphi: \bar{X}(G, \mathcal{P}, S) \rightarrow \hat{X}(G, \mathcal{P}, S)$ that extends the natural quasi-isometry

$$\varphi: \bar{\Gamma}(G, \mathcal{P}, S) \rightarrow \hat{\Gamma}(G, \mathcal{P}, S).$$

In particular, we have a commutative diagram

$$\begin{array}{ccc} \bar{\Gamma}(G, \mathcal{P}, S) & \xrightarrow{\varphi} & \hat{\Gamma}(G, \mathcal{P}, S) \\ \downarrow & & \downarrow \\ \bar{X}(G, \mathcal{P}, S) & \xrightarrow{\varphi} & \hat{X}(G, \mathcal{P}, S). \end{array}$$

Specifically, every R -cell in $\bar{X}(G, \mathcal{P}, S)$ is sent homeomorphically to the corresponding 2-cell in $\hat{X}(G, \mathcal{P}, S)$, while every \mathcal{P} -cell in $\bar{X}(G, \mathcal{P}, S)$ is collapsed to a star-like 1-complex as we see in Figure 1.






Figure 1. The image of the boundary of a \mathcal{P} -cell on $\bar{\Gamma}(G, \mathcal{P}, S)$ under the quasi-isometry φ .

Remark 4.12. The following statements are straightforward to verify from the definition of $\varphi: \bar{X}(G, \mathcal{P}, S) \rightarrow \hat{X}(G, \mathcal{P}, S)$ and Figure 1. Denote by $\Delta_{\hat{X}}$ the combinatorial Dehn function of $\hat{X}(G, \mathcal{P}, S)$.


- (1) Let $\hat{\gamma}: S^1 \rightarrow \hat{X}(G, \mathcal{P}, S)$ be a loop with no backtracks in the coned-off Cayley complex. Then we can pull back $\hat{\gamma}$ to a loop $\gamma: S^1 \rightarrow \bar{X}(G, \mathcal{P}, S)$ in such a way that the following diagram commutes:



Let $D \rightarrow \bar{X}(G, \mathcal{P}, S)$ be a disk diagram filling a combinatorial loop $\gamma: S^1 \rightarrow \bar{X}(G, \mathcal{P}, S)$. Then there exists a disk diagram $\hat{D} \rightarrow \hat{X}(G, \mathcal{P}, S)$ so that the following diagram commutes:



- (2) Let $\gamma: S^1 \rightarrow \bar{X}(G, \mathcal{P}, S)$ be a combinatorial loop of length n , then we can push it to a loop $\hat{\gamma} = \varphi \circ \gamma: S^1 \rightarrow \hat{X}(G, \mathcal{P}, S)$ of length at most $2n$, that is, we have the following commutative diagram:



Let $\widehat{D} \rightarrow \widehat{X}(G, \mathcal{P}, S)$ be a disk diagram filling the cycle $\widehat{\gamma} = \varphi \circ \gamma$. Then there exists a disk diagram $D \rightarrow \bar{X}$ such that the following diagram commutes:

$$\begin{array}{ccccc}
 \bar{X}(G, \mathcal{P}, S) & \xrightarrow{\varphi} & \widehat{X}(G, \mathcal{P}, S) & \xleftarrow{\widehat{\gamma}} & S^1 \\
 \uparrow \text{---} & & \uparrow & & \parallel \\
 D & \text{---} & \widehat{D} & \xleftarrow{\quad} & \partial \widehat{D}
 \end{array}$$

(3) In both items above, $\text{Area}^{\text{rel}}(D) = \text{Area}(\widehat{D})$.

Proposition 4.13. *Let $G = \langle S, \mathcal{P} \mid R \rangle$ be a finite relative presentation, and let $\Delta_{G, \mathcal{P}}$ and \widehat{X} be the corresponding relative Dehn function and coned-off Cayley complex, respectively. Then $\Delta_{G, \mathcal{P}}(n) \asymp \Delta_{\widehat{X}}(n)$ for every $n \in \mathbb{N}$.*

Proof. Let $\widehat{\gamma}: S^1 \rightarrow \widehat{X}(G, \mathcal{P}, S)$ be a loop of length n with no backtracks in the coned-off Cayley complex. By the first item of Remark 4.12 and considering a minimal relative area disk diagram $D \rightarrow \bar{X}(G, \mathcal{P}, S)$ filling a pullback cycle $\widehat{\gamma}: S^1 \rightarrow \widehat{\Gamma}(G, \mathcal{P}, S)$ of γ , it follows that

$$\Delta_{G, \mathcal{P}}(|\widehat{\gamma}|) \geq \Delta_{G, \mathcal{P}}(|\gamma|) \geq \text{Area}^{\text{rel}}(D) = \text{Area}(\widehat{D}) \geq \text{Area}(\widehat{\gamma}),$$

where the equality comes from the third item of Remark 4.12. Therefore, $\Delta_{G, \mathcal{P}}(n) \geq \Delta_{\widehat{X}}(n)$ for all $n \in \mathbb{N}$. Analogously, let $\gamma: S^1 \rightarrow \bar{X}(G, \mathcal{P}, S)$ be a combinatorial loop. By the second item of Remark 4.12 and considering a minimal area disk diagram $\widehat{D} \rightarrow \widehat{X}(G, \mathcal{P}, S)$ filling $\widehat{\gamma} = \varphi \circ \gamma$, it follows that

$$\text{Area}^{\text{rel}}(\gamma) \leq \text{Area}^{\text{rel}}(D) = \text{Area}(\widehat{D}) \leq \Delta_{\widehat{X}}(|\widehat{\gamma}|) \leq \Delta_{\widehat{X}}(2|\gamma|),$$

and hence $\Delta_{G, \mathcal{P}}(n) \leq \Delta_{\widehat{X}}(2n)$ for all $n \in \mathbb{N}$. ■

The following corollary is a direct consequence of Theorem 2.1 and Proposition 4.13.

Corollary 4.14. *Let G be finitely presented relative to a collection of subgroups \mathcal{P} . The following statements are equivalent:*

- (1) *The relative Dehn function $\Delta_{G, \mathcal{P}}$ takes only finite values.*
- (2) *The graph $\widehat{\Gamma}(G, \mathcal{P})$ is fine.*

The proof of [8, Proposition 2.50] contains an argument proving (1) implies (2) of the above corollary.

The following corollary is a straightforward consequence of Propositions 4.13 and 3.2.

Corollary 4.15. *Let G be a group finitely presented relative to a finite collection of subgroups \mathcal{P} . If $\Delta_{G, \mathcal{P}}$ takes only finite values, then $\widehat{\Gamma}(G, \mathcal{P})$ is fillable for some integer m .*

Proposition 4.16. *Let G be a group finitely generated by S with respect to \mathcal{P} . If the graph $\widehat{\Gamma}(G, \mathcal{P}, S)$ is connected, fine, cocompact, and k -fillable, then G is finitely presented relative to \mathcal{P} .*

Proof. We use the setup of Proposition 4.9. In particular, N is the normal subgroup of \tilde{G} generated by R , we fix a vertex x_0 of $\hat{\Gamma} = \hat{\Gamma}(G, \mathcal{P}, S)$, and we have a group isomorphism $\psi: N \rightarrow \pi_1(\hat{\Gamma}, x_0)$.

Since $\hat{\Gamma}$ is k -fillable, there is an integer m such that the complex \hat{X} obtained by attaching 2-cells with boundary paths the circuits of length at most m is simply connected, see Remark 3.4.

Since $\hat{\Gamma}$ is fine and there are finitely many G -orbits of edges, there are finitely many G -orbits of circuits of length at most m . Let $\{\gamma_1, \dots, \gamma_\ell\}$ be a collection of representatives of circuits of length m , and after translations assume that each γ_i contains the vertex x_0 corresponding to the identity element of G . Then each γ_i defines an element of the fundamental group $\pi_1(\hat{\Gamma}, x_0)$. Let $r_i \in N$ be defined by $\psi(r_i) = \gamma_i$.

Since \hat{X} is simply connected, we have that $\pi_1(\hat{\Gamma}, x_0)$ is generated by the closed paths arising as concatenations of the form $\alpha_g \cdot \gamma_i \cdot \bar{\alpha}_g$ for $g \in \tilde{G}$, where α_g is the projection via ρ of the unique path from \tilde{x}_0 to $g.\tilde{x}_0$. Equivalently, N is generated by the elements $g r_i g^{-1}$ for $g \in \tilde{G}$. We have shown that N is normally generated by $\mathcal{R} = \{r_1, \dots, r_\ell\}$.

Since $\hat{\Gamma} = \hat{\Gamma}(G, \mathcal{P})$ is cocompact, the collection \mathcal{P} is finite. Therefore, $\langle S, \mathcal{P} \mid \mathcal{R} \rangle$ is a finite relative presentation of G . ■

Summarising the results of this section, we obtain Theorem 4.17 below.

Theorem 4.17 (Theorem E). *Let G be a group finitely generated relative to a finite collection of subgroups \mathcal{P} . If G is finitely presented relative to \mathcal{P} , then*

- (1) $\hat{\Gamma}(G, \mathcal{P})$ is fillable.
- (2) The relative Dehn function $\Delta_{G, \mathcal{P}}$ is well defined if and only if $\hat{\Gamma}(G, \mathcal{P})$ is fine graph.

Conversely, if $\hat{\Gamma}(G, \mathcal{P})$ is fine and fillable, then G is finitely presented relative to \mathcal{P} and hence $\Delta_{G, \mathcal{P}}$ is well defined.

Proof. This follows from Corollaries 4.14, 4.15 and Proposition 4.16. ■

Note that Theorem E from the introduction is a particular case Theorem 4.17.

5. Fineness and quasi-isometries of pairs

In this section, we will prove Theorem C from the introduction. The heart of the argument is establishing Proposition 5.6 which gives conditions on a quasi-isometry of pairs $q: (G, \mathcal{P}) \rightarrow (H, \mathcal{Q})$ to induce a quasi-isometry of coned-off Cayley graphs. The remainder of the section then works towards replacing the geometric-set-theoretic conditions on q with algebraic conditions on \mathcal{P} and \mathcal{Q} . This yields Proposition 5.12. Finally, we give a proof of Theorem C.

Another equivalent definition of Bowditch’s fine graphs is used in this section [2, Proposition 2.1].

Definition 5.1 (Fine). Let Γ be a graph and let v be a vertex of Γ . Let

$$T_v\Gamma = \{w \in V(\Gamma) : \{v, w\} \in E(\Gamma)\}$$

denote the set of the vertices adjacent to v . For $x, y \in T_v\Gamma$, the *angle metric* $\angle_v(x, y)$ is the length of the shortest path in the graph $\Gamma \setminus \{v\}$ between x and y , with $\angle_v(x, y) = \infty$ if there is no such path. The graph Γ is *fine at v* if $(T_v\Gamma, \angle_v)$ is a locally finite metric space. The graph Γ is *fine at $C \subseteq V(\Gamma)$* if Γ is fine at v for all $v \in C$. The graph Γ is a *fine graph* if it is fine at every vertex.

Definition 5.2 (Quasi-isometry of pairs). Consider two pairs (G, \mathcal{P}) and (H, \mathcal{Q}) , where G and H are finitely generated groups with chosen word metrics dist_G and dist_H with respect to some finite generating sets. Denote the Hausdorff distance between subsets of H by hdist_H . An (L, C) -quasi-isometry $q: G \rightarrow H$ is an (L, C, M) -quasi-isometry of pairs $q: (G, \mathcal{P}) \rightarrow (H, \mathcal{Q})$ if the relation

$$\dot{q} = \{(A, B) \in G/\mathcal{P} \times H/\mathcal{Q} : \text{hdist}_H(q(A), B) < M\}$$

satisfies that the projections into G/\mathcal{P} and H/\mathcal{Q} are surjective.

In this section, we explicitly use the relational approach of the notion of a function between sets: a function f from A to B is a subset of $A \times B$ so that for every $a \in A$, there is a unique $b \in B$ such that $(a, b) \in f$.

The proof of Theorem C, the main objective of this section, relies on the study of the relation \dot{q} defined by a quasi-isometry of pairs $q: (G, \mathcal{P}) \rightarrow (H, \mathcal{Q})$. We will show that in the case where \dot{q} defines a bijection $G/\mathcal{P} \rightarrow H/\mathcal{Q}$, the coned-off Cayley graphs $\widehat{\Gamma}(G, \mathcal{P})$ and $\widehat{\Gamma}(H, \mathcal{Q})$ share global a local geometric conditions, see Proposition 5.6. In the second part of the section, we provide algebraic conditions guaranteeing that the relation \dot{q} is a bijection, see Proposition 5.12.

Remark 5.3. Note that in Definition 5.2, the notion of a quasi-isometry of pairs is independent of the chosen finite generating sets for G and H . In the case where we want to keep track of specific generating sets we use the following notation. If G and H are groups generated by finite generating sets S_0 and T_0 , respectively, by a quasi-isometry of pairs $(G, \mathcal{P}, S_0) \rightarrow (H, \mathcal{Q}, T_0)$, we mean a quasi-isometry of pairs $(G, \mathcal{P}) \rightarrow (H, \mathcal{Q})$ with respect to the word metrics induced by S_0 and T_0 .

Remark 5.4. If \mathcal{P} is a finite collection, then the metric space $(G/\mathcal{P}, \text{hdist})$ is locally finite. Indeed, when fixing $P \in \mathcal{P}$ and $r > 0$, there are finitely many left cosets in G/\mathcal{P} such that

$$\text{hdist}(P, gP) < r.$$

Moreover, the left G -action on G/\mathcal{P} by multiplication on the left preserves the Hausdorff distance hdist between subsets of G and hence it is an action by isometries.

Remark 5.5. If $q: (G, \mathcal{P}) \rightarrow (H, \mathcal{Q})$ is an (L, C, M) -quasi-isometry of pairs, and \dot{q} is a function $G/\mathcal{P} \rightarrow H/\mathcal{Q}$, then

$$\frac{1}{L} \text{hdist}(A, B) - C - 2M \leq \text{hdist}(\dot{q}(A), \dot{q}(B)) \leq L \text{hdist}(A, B) + C + 2M.$$

In particular, $\dot{q}: (G/\mathcal{P}, \text{hdist}) \rightarrow (H/\mathcal{Q}, \text{hdist})$ is a quasi-isometry.

The main technical result of this section is the following proposition. Note that given a connected graph Γ , we consider the vertex set as a metric space with metric induced by the path metric. In particular, a quasi-isometry between graphs is a function of the vertex sets satisfying the usual axioms.

Proposition 5.6. *Let G and H be groups, let $S \subset G$ and $T \subset H$, and let $S_0 \subset S$ and $T_0 \subset T$ be finite generating sets of G and H , respectively. Consider collections \mathcal{P} and \mathcal{Q} of subgroups of G and H , respectively. Let $q: G \rightarrow H$ be a function.*

Suppose q is a quasi-isometry $\Gamma(G, S) \rightarrow \Gamma(H, T)$, \hat{q} is a quasi-isometry of pairs $(G, \mathcal{P}, S_0) \rightarrow (H, \mathcal{Q}, T_0)$, and \dot{q} is a bijection $G/\mathcal{P} \rightarrow H/\mathcal{Q}$.

- (1) *If $\hat{q} = q \cup \dot{q}$, then \hat{q} is a quasi-isometry $\hat{\Gamma}(G, \mathcal{P}, S) \rightarrow \hat{\Gamma}(H, \mathcal{Q}, T)$.*
- (2) *If $\hat{\Gamma}(H, \mathcal{Q}, T)$ is fine at cone vertices, then $\hat{\Gamma}(G, \mathcal{P}, S)$ is fine at cone vertices.*

Remark 5.7. There are algebraic conditions on \mathcal{P} and \mathcal{Q} that imply that \dot{q} is a bijection, see Proposition 5.12.

Corollary 5.8. *Suppose that $q: (G, \mathcal{P}) \rightarrow (H, \mathcal{Q})$ is a quasi-isometry of pairs and \dot{q} is a bijection. Then $\hat{q}: \hat{\Gamma}(G, \mathcal{P}) \rightarrow \hat{\Gamma}(H, \mathcal{Q})$ is a quasi-isometry, and if $\hat{\Gamma}(H, \mathcal{Q})$ is a fine graph, then $\hat{\Gamma}(G, \mathcal{P})$ is a fine graph.*

The following argument is patterned from the proof of [18, Proposition 5.4].

Proof of Proposition 5.6. Suppose $q: \Gamma(G, S) \rightarrow \Gamma(H, T)$ is a (\bar{L}, \bar{C}) -quasi-isometry and $q: (G, \mathcal{P}, S_0) \rightarrow (H, \mathcal{Q}, T_0)$ is a (L, C, M) -quasi-isometry of pairs.

For any path $\alpha = [v_0, v_1, \dots, v_\ell]$ in $\hat{\Gamma}(G, \mathcal{P}, S)$, let $\hat{q}(\alpha)$ denote a path in $\hat{\Gamma}(H, \mathcal{Q}, T)$ from $\hat{q}(v_0)$ to $\hat{q}(v_\ell)$ obtained as the concatenation of paths $\beta_0, \dots, \beta_{\ell-1}$, where β_i is a path from $\hat{q}(v_i)$ to $\hat{q}(v_{i+1})$ defined as follows:

- (1) If v_i and v_{i+1} are elements of G , then β_i is a geodesic in $\Gamma(H, T)$ from $q(v_i)$ to $q(v_{i+1})$. Since $q: \Gamma(G, S) \rightarrow \Gamma(H, T)$ is a (\bar{L}, \bar{C}) -quasi-isometry, β_i has length bounded by $\bar{L} + \bar{C}$.
- (2) Suppose $v_i \in G$ and $v_{i+1} \in G/\mathcal{P}$. Observe that v_i is an element of the left coset v_{i+1} . Since $q: (G, \mathcal{P}, S_0) \rightarrow (H, \mathcal{Q}, T_0)$ is an (L, C, M) -quasi-isometry of pairs, there is a geodesic of length at most M in $\Gamma(H, T_0)$ from $q(v_i)$ to an element w of the left coset $\dot{q}(v_{i+1})$. Let β_i be the concatenation of this geodesic in $\Gamma(H, T_0)$ followed by the edge between w and the cone vertex $\dot{q}(v_{i+1})$. Observe that β_i is a path of length at most $M + 1$ in $\hat{\Gamma}(H, \mathcal{Q}, T)$.

- (3) If $v_i \in G/\mathcal{P}$ and $v_{i+1} \in G$, then β_i is defined in an analogous way as in the previous case, and also has length at most $M + 1$.

Observe that $|\hat{q}(\alpha)| \leq (\bar{L} + \bar{C} + M + 1)|\alpha|$. The above inequality applied in the case where α is a geodesic between vertices x and y of $\hat{\Gamma}(G, \mathcal{P}, S)$ implies that

$$\text{dist}_{\hat{\Gamma}(H, \mathcal{Q}, T)}(\hat{q}(x), \hat{q}(y)) \leq (\bar{L} + \bar{C} + M + 1) \text{dist}_{\hat{\Gamma}(G, \mathcal{P}, S)}(x, y)$$

for any pair of vertices x, y of $\hat{\Gamma}(G, \mathcal{P}, S)$. By symmetry, an analogous inequality holds for vertices of $\hat{\Gamma}(H, \mathcal{Q})$. Since \hat{q} is a bijection, the definition of $\hat{q}(\alpha)$ shows that α passes through a cone vertex A if and only if $\hat{q}(\alpha)$ passes through the cone vertex $\hat{q}(A)$. We summarise this discussion in the following lemma.

Lemma 5.9. *There are constants $\hat{L} \geq 1$ and $\hat{C} \geq 0$ such that*

- (1) *The function \hat{q} is a (\hat{L}, \hat{C}) -quasi-isometry from $\hat{\Gamma}(G, \mathcal{P}, S)$ to $\hat{\Gamma}(H, \mathcal{Q}, T)$.*
- (2) *Let α be a path in $\hat{\Gamma}(G, \mathcal{P}, S)$.*
 - (a) *For any $A \in G/\mathcal{P}$, α passes through the cone vertex A if and only if $\hat{q}(\alpha)$ passes through the cone vertex $\hat{q}(A)$.*
 - (b) *$|\hat{q}(\alpha)| \leq \hat{L} |\alpha|$.*

We prove the contrapositive of the second statement of the proposition. See Figure 2 for a schematic of the argument that follows. Suppose that $\hat{\Gamma}(G, \mathcal{P}, S)$ is not fine at cone vertices. Then there is $P \in \mathcal{P}$ such that $(T_P \hat{\Gamma}, \angle_P)$ is not locally-finite. Let $r > 0$ and let $\{g_i\} \subseteq P$ be an infinite subset such that $\angle_P(g_i, g_j) \leq r$ for every i, j . Let $\alpha_{i,j}$ be a path in $\hat{\Gamma}(G, \mathcal{P}, S)$ from g_i to g_j of length at most r that does not contain the cone vertex P . Let Q denote the left coset $\hat{q}(P)$. Let γ_i be a geodesic in $\Gamma(H, T_0)$ from an element h_i of Q to $q(g_i)$ such that $\text{dist}_H(h_i, q(g_i)) = \text{dist}_H(Q, q(g_i))$. Since q is a (L, C, M) quasi-isometry of pairs, each γ_i has length at most M .

Let us prove that the set $\{h_i\}$ is infinite. Suppose, for contradiction, that $\{h_i\}$ is a finite set. Since T_0 is a finite generating set, $\Gamma(H, T_0)$ is a locally finite graph and hence it admits only finitely many paths of length at most M with initial vertex in $\{h_i\}$. Since each γ_i has length at most M with initial vertex in $\{h_i\}$, it follows that the set $\{q(g_i)\}$ is finite and, in particular, bounded. Since q is a quasi-isometry $\Gamma(G, S_0) \rightarrow \Gamma(G, T_0)$, it follows that the set $\{g_i\}$ is a bounded subset of vertices in the locally finite graph $\Gamma(G, S_0)$, hence the set $\{g_i\}$ is finite, a contradiction.

To conclude the proof, we show that $\hat{\Gamma}(H, \mathcal{Q}, T)$ is not fine at the cone vertex Q . Since $\{h_i\}$ is an infinite subset of Q , it is enough to show that $\angle_Q(h_i, h_j) \leq r\hat{L} + M$ for any i, j . Consider the path $\beta_{i,j}$ from h_i to h_j obtained as the concatenation of the path γ_i from h_i to $\hat{q}(g_i)$, followed by the path $\hat{q}(\alpha_{i,j})$ from $\hat{q}(g_i)$ to $\hat{q}(g_j)$, and then the path $\bar{\gamma}_j$ from $\hat{q}(g_j)$ to h_j . The paths γ_i and $\bar{\gamma}_j$ have length bounded by M , and they do not contain the cone vertex Q as they are paths in $\Gamma(H, T_0)$; the path $\hat{q}(\alpha_{i,j})$ has length at most $r\hat{L}$ and does not contain the cone vertex Q by Lemma 5.9. Therefore, $\angle_Q(h_i, h_j) \leq |\gamma_i| + |\hat{q}(\alpha_{i,j})| + |\bar{\gamma}_j| \leq 2M + r\hat{L}$ as desired. This completes the proof of Proposition 5.6. ■




Figure 2. Illustration of the proof of Proposition 5.6.

The goal for the remainder of this section is to give algebraic conditions on \mathcal{P} and \mathcal{Q} to ensure \hat{q} is a bijection. The following key definition will provide such criteria.

Definition 5.10 (Reduced collection). A collection of subgroups \mathcal{P} of a group G is *reduced* if for any $P, Q \in \mathcal{P}$ and $g \in G$, then P and gQg^{-1} being commensurable subgroups implies $P = Q$ and $g \in P$.

Remark 5.11. If \mathcal{P} is a reduced collection of subgroups of a group G , then

$$P = \text{Comm}_G(P) \quad \text{for any } P \in \mathcal{P}.$$

Proposition 5.12. Let $q: (G, \mathcal{P}) \rightarrow (H, \mathcal{Q})$ be a (L, C, M) -quasi-isometry of pairs. Then

- (1) \hat{q} is a surjective function $G/\mathcal{P} \rightarrow H/\mathcal{Q}$ if \mathcal{Q} is reduced.
- (2) \hat{q} is a bijection $G/\mathcal{P} \rightarrow H/\mathcal{Q}$ if \mathcal{P} and \mathcal{Q} are reduced.

There are different versions of the following lemma in the literature: [21, Lemma 2.2], [16, Lemma 4.7] and [12, Proposition 9.4], the statement below is taken from the later reference. For $A \subset G$, $\mathcal{N}_k(A)$ denotes the closed neighbourhood of A in (G, dist_G) .

Lemma 5.13. Let G be a finitely generated group with word metric dist_G . Let gP and fQ be arbitrary left cosets of subgroups of G . Then for any $k > 0$ there is $M > 0$ such that

$$\mathcal{N}_k(gP) \cap \mathcal{N}_k(fQ) \subseteq \mathcal{N}_M(gPg^{-1} \cap fQf^{-1}).$$

Lemma 5.14. Let G be a finitely generated group with a word metric dist_G , let P and Q be subgroups, and let $g \in G$. Then P and gQg^{-1} are commensurable subgroups if and only if $\text{hdist}_G(P, gQ) < \infty$.

Proof. Suppose K is a finite index subgroup of P and gQg^{-1} . Then $\text{hdist}(K, P) < \infty$ and $\text{hdist}(K, gQg^{-1})$ are finite. Since $\text{hdist}(gQg^{-1}, gQ) \leq \text{dist}(1, g) < \infty$, it follows that

$$\text{hdist}(P, gQ) \leq \text{hdist}(P, K) + \text{hdist}(K, gQg^{-1}) + \text{hdist}(gQg^{-1}, gQ) < \infty.$$

Conversely, suppose $\text{hdist}(P, gQ)$ is finite. Then $P \subset P \cap \mathcal{N}_k(gQ)$ for some k , and therefore Lemma 5.13 implies that $P \subseteq \mathcal{N}_M(P \cap gQg^{-1})$ for some M . It follows that $P \cap gQg^{-1}$ is a finite index subgroup of P . In an analogous way, one shows that $P \cap gQg^{-1}$ is a finite index subgroup of gQg^{-1} . Whence, P and gQg^{-1} are commensurable subgroups. ■

Proof of Proposition 5.12. To prove the first statement, we only need to show that the relation \dot{q} is a function. Suppose that \mathcal{Q} is reduced and the pairs (A, h_1Q_1) and (A, h_2Q_2) belong to \dot{q} . Then $h_1Q_1, h_2Q_2 \in H/\mathcal{Q}$ and $\text{hdist}_H(h_1Q_1, h_2Q_2) < \infty$. Lemma 5.14 implies that $h_1Q_1h_1^{-1}$ and $h_2Q_2h_2^{-1}$ are commensurable subgroups. Since \mathcal{Q} is reduced, it follows that $Q_1 = Q_2$ and $h_2 \in h_1Q_1$. In particular, $h_1Q_1 = h_2Q_2$ and hence \dot{q} is a function. The second statement of the lemma follows from the first one. ■

We are now ready to prove Theorem C from the introduction.

Theorem 5.15 (Theorem C). *Let $q: (G, \mathcal{P}) \rightarrow (H, \mathcal{Q})$ be a quasi-isometry of pairs. Suppose \mathcal{P} and \mathcal{Q} are reduced finite collections. Then there is an induced quasi-isometry of graphs $\hat{q}: \hat{\Gamma}(G, \mathcal{P}) \rightarrow \hat{\Gamma}(H, \mathcal{Q})$, and if $\hat{\Gamma}(H, \mathcal{Q})$ is a fine graph, then $\hat{\Gamma}(G, \mathcal{P})$ is a fine graph.*

Proof. The result follows from applying Proposition 5.12 to Corollary 5.8. ■

6. Almost malnormal collections and quasi-isometries of pairs

In this section, we will prove Theorem D from the introduction. First, we introduce a refinement \mathcal{P}^* of a collection \mathcal{P} . In Proposition 6.3, we show under a mild hypothesis that (G, \mathcal{P}) and (G, \mathcal{P}^*) are quasi-isometric pairs under the identity map.

Definition 6.1. Let \mathcal{P} be a collection of subgroups of a group G . A refinement \mathcal{P}^* of \mathcal{P} is a set of representatives of conjugacy classes of the collection of subgroups

$$\{\text{Comm}_G(gPg^{-1}): P \in \mathcal{P} \text{ and } g \in G\}.$$

Remark 6.2. Observe that for a collection of subgroups \mathcal{P} of a group G , there is a refinement \mathcal{P}^* such that each of its elements are of the form $\text{Comm}_G(P)$ for some $P \in \mathcal{P}$. This is a consequence of $\text{Comm}_G(gPg^{-1}) = g \text{Comm}_G(P)g^{-1}$ for each subgroup P of G .

Proposition 6.3. *Let \mathcal{P}^* be a refinement of a finite collection of subgroups \mathcal{P} of a finitely generated group G . If P is a finite index subgroup of $\text{Comm}_G(P)$ for every $P \in \mathcal{P}$, then (G, \mathcal{P}) and (G, \mathcal{P}^*) are quasi-isometric pairs via the identity map on G .*

Proof. Let $\mathcal{P} = \{P_1, \dots, P_k\}$. By the previous remark, we may assume that every subgroup in \mathcal{P}^* is of the form $\text{Comm}_G(P)$ for some $P \in \mathcal{P}$. Let $q: G \rightarrow G$ be the identity map. Since q is a $(1, 0)$ -quasi-isometry, it is enough to show that there is $M > 0$ such that the relation

$$\dot{q} = \{(A, B) \in G/\mathcal{P} \times G/\mathcal{P}^* : \text{hdist}(A, B) < M\}$$

satisfies that it projects surjectively on G/\mathcal{P} and on G/\mathcal{P}^* .

For any $P_i \in \mathcal{P}$, note that $\text{hdist}(P_i, \text{Comm}_G(P_i)) < \infty$ since P_i has finite index in $\text{Comm}_G(P_i)$. Let

$$M_1 = \max\{\text{hdist}(P_i, \text{Comm}_G(P_i)) : 1 \leq i \leq k\}.$$

By definition of \mathcal{P}^* , for any P_i , there is $Q_i \in \mathcal{P}^*$ and $g_i \in G$ such that $\text{Comm}_G(P_i) = g_i Q_i g_i^{-1}$. In particular, $\text{hdist}(\text{Comm}_G(P_i), g_i Q_i)$ is finite. Let

$$M_2 = \max\{\text{hdist}(\text{Comm}_G(P_i), g_i Q_i) : 1 \leq i \leq k\}.$$

Let $M > M_1 + M_2$. Then for any $gP_i \in G/\mathcal{P}$, $(gP_i, g g_i Q_i) \in \dot{q}$. On the other hand, if $gQ \in G/\mathcal{P}^*$, then $Q = \text{Comm}_G(P)$ for some $P \in \mathcal{P}$ and hence $(gP, gQ) \in \dot{q}$. ■

Remark 6.4. Note that in the previous proposition if \mathcal{P} is infinite, the map $\dot{q}: G/\mathcal{P} \rightarrow G/\mathcal{P}^*$ must be finite-to-one. Otherwise after conjugating, there will be a sequence of subgroups $P_i \leq \text{Comm}_G(P_0)$ such that $|\text{Comm}_G(P_0) : P_i| \rightarrow \infty$, in particular, the sequence of Hausdorff distances $\text{hdist}(\text{Comm}_G(P_0), P_i)$ is not bounded.

Definition 6.5. A collection of subgroups \mathcal{P} of a group G is *almost malnormal* if for any $P, P' \in \mathcal{P}$ and $g \in G$, either $gPg^{-1} \cap P'$ is finite, or $P = P'$ and $g \in P$.

Remark 6.6. If \mathcal{P} is an almost malnormal collection of infinite subgroups of a group G , then \mathcal{P} is reduced.

Remark 6.7. If a group G acts by automorphisms on a fine graph Γ such that edge stabilisers are finite and \mathcal{P} is a collection of representatives of conjugacy classes of vertex stabilisers, then \mathcal{P} is an almost malnormal collection.

Proposition 6.8. *Let $q: (G, \mathcal{P}) \rightarrow (H, \mathcal{Q})$ be a quasi-isometry of pairs. If \mathcal{Q} is an almost malnormal finite collection of infinite subgroups and \mathcal{P} is a finite collection, then any refinement \mathcal{P}^* of \mathcal{P} is almost malnormal.*

The proof of Proposition 6.8 relies on the following lemmas.

Lemma 6.9. *Let \mathcal{P} be a collection of subgroups of a group G . Suppose P is a finite index subgroup of $\text{Comm}_G(P)$ for every $P \in \mathcal{P}$. Then any refinement \mathcal{P}^* of \mathcal{P} is a reduced collection.*

Proof. Since commensurable subgroups have equal commensurator,

$$\text{Comm}_G(\text{Comm}_G(P)) = \text{Comm}_G(P)$$

for every $P \in \mathcal{P}$. Let $P_1, P_2 \in \mathcal{P}$ such that $\text{Comm}_G(P_1)$ and $\text{Comm}_G(P_2)$ are in \mathcal{P}^* , and let $g \in G$. Suppose $\text{Comm}_G(P_1)$ and $g \text{Comm}_G(P_2)g^{-1}$ are commensurable subgroups. Then

$$\begin{aligned} \text{Comm}_G(P_1) &= \text{Comm}_G(\text{Comm}_G(P_1)) = \text{Comm}_G(g \text{Comm}_G(P_2)g^{-1}) \\ &= \text{Comm}_G(\text{Comm}_G(gP_2g^{-1})) = \text{Comm}_G(gP_2g^{-1}) \\ &= g \text{Comm}_G(P_2)g^{-1}. \end{aligned}$$

Since, by definition, \mathcal{P}^* does not have two subgroups that are conjugate to each other, it follows that $\text{Comm}_G(P_1) = \text{Comm}_G(P_2)$ and $g \in \text{Comm}_G(P_1)$. Therefore, \mathcal{P}^* is reduced. ■

Lemma 6.10. *Let \mathcal{P} be a finite collection of infinite subgroups of a finitely generated group G . Then \mathcal{P} is almost malnormal if and only if for any $A, B \in G/\mathcal{P}$, either $A = B$ or $\mathcal{N}_n(A) \cap \mathcal{N}_n(B)$ is a finite subset of G for every n .*

Proof. Suppose that \mathcal{P} is an almost malnormal collection of infinite subgroups. Let $g_1P_1, g_2P_2 \in G/\mathcal{P}$ and suppose that $\mathcal{N}_n(g_1P_1) \cap \mathcal{N}_n(g_2P_2)$ is an infinite (and hence unbounded) subset of G for some integer n . By Lemma 5.13, there is an integer m such that $\mathcal{N}_n(g_1P_1) \cap \mathcal{N}_n(g_2P_2) \subset \mathcal{N}_m(g_1P_1g_1^{-1} \cap g_2P_2g_2^{-1})$. It follows that $g_1P_1g_1^{-1} \cap g_2P_2g_2^{-1}$ is an infinite subgroup and hence $P_1 = P_2$ and $g_1^{-1}g_2 \in P_1$ by almost malnormality. Therefore, $g_1P_1 = g_2P_2$.

Conversely, suppose that for any $A, B \in G/\mathcal{P}$, either $A = B$ or $\mathcal{N}_n(A) \cap \mathcal{N}_n(B)$ is a finite set for every n . Let $P, P' \in \mathcal{P}$ and $g \in G$ and suppose that $gPg^{-1} \cap P'$ is an infinite subgroup. It follows that there is $n > 0$ such that $\mathcal{N}_n(gP) \cap \mathcal{N}_n(P')$ is an infinite subset of G . Hence $gP = P'$ and, in particular, $P = P'$ and $g \in P$. ■

Lemma 6.11. *Let $q: (G, \mathcal{P}) \rightarrow (H, \mathcal{Q})$ be a quasi-isometry of pairs. Suppose that \mathcal{P} and \mathcal{Q} are finite collections, and \mathcal{Q} is reduced. If Q is of finite index in $\text{Comm}_H(Q)$ for every $Q \in \mathcal{Q}$, then P is of finite index in $\text{Comm}_G(P)$ for every $P \in \mathcal{P}$.*

Proof. Since \mathcal{Q} is reduced, \dot{q} is a function from $G/\mathcal{P} \rightarrow G/\mathcal{Q}$. Since both \mathcal{P} and \mathcal{Q} are finite collections, it follows that $\dot{q}: (G/\mathcal{P}, \text{hdist}) \rightarrow (H/\mathcal{Q}, \text{hdist})$ is a quasi-isometry between locally finite metric spaces. Suppose that $P \in \mathcal{P}$ has infinite index in $\text{Comm}_G(P)$. Lemma 5.14 implies that there is an infinite collection of left cosets $\mathcal{A} = \{g_iP : i \in I\}$ such that $\text{hdist}(g_iP, g_jP) < \infty$ for any $i, j \in I$. By local finiteness of $(G/\mathcal{P}, \text{hdist})$, the collection \mathcal{A} is an unbounded subset of G/\mathcal{P} . It follows that $\mathcal{B} = \{\dot{q}(g_iP) : i \in I\}$ is an unbounded subset of H/\mathcal{Q} . Since \mathcal{Q} is a finite collection, and $\dot{q}(g_iP) = h_iQ_i$ for some $h_i \in H$ and $Q_i \in \mathcal{Q}$, the pigeonhole principle implies that we can assume that all Q_i 's are a fixed $Q \in \mathcal{Q}$. By Lemma 5.14, the subgroup Q has infinite index in $\text{Comm}_H(Q)$. ■

Proof of Proposition 6.8. Suppose that $q: (G, \mathcal{P}) \rightarrow (G, \mathcal{Q})$ is a quasi-isometry of pairs. Since \mathcal{Q} is an almost malnormal collection of infinite subgroups, it is a reduced collection and every element of \mathcal{Q} has a finite index in its commensurator. Since \mathcal{P} and \mathcal{Q} are finite

collections, Lemma 6.11 implies that every element of \mathcal{P} has a finite index in its commensurator. Let \mathcal{P}^* be a refinement of \mathcal{P} in G . By Proposition 6.3, there is a quasi-isometry of pairs $p: (G, \mathcal{P}^*) \rightarrow (G, \mathcal{P})$. Then the composition $r = p \circ q$ is an (L, C, M) -quasi-isometry of pairs $(G, \mathcal{P}^*) \rightarrow (H, \mathcal{Q})$. Lemma 6.9 implies that \mathcal{P}^* is a reduced collection. Therefore, \dot{r} is a bijection $G/\mathcal{P}^* \rightarrow H/\mathcal{Q}$ by Proposition 5.12. To conclude that \mathcal{P}^* is an almost malnormal, we verify the hypothesis of Lemma 6.10.

Claim. *The collection \mathcal{P}^* is a finite collection of infinite subgroups.*

Since \mathcal{P} is finite, then \mathcal{P}^* is finite. Every element of \mathcal{P}^* is a conjugate of a subgroup of the form $\text{Comm}_G(P)$ for some $P \in \mathcal{P}$, hence it is enough to show that \mathcal{P} contains only infinite subgroups. Observe that any $P \in \mathcal{P}$ is an infinite subgroup since $\text{hdist}(\dot{q}(P), Q) < \infty$ for some $Q \in H/\mathcal{Q}$ and every subgroup in \mathcal{Q} is infinite.

Claim. *For any $A, B \in G/\mathcal{P}^*$, either $A = B$ or $\mathcal{N}_n(A) \cap \mathcal{N}_n(B)$ is a finite subset of G for every n .*

Let $A, B \in G/\mathcal{P}^*$ and suppose that $A \neq B$. Since $\dot{r}: G/\mathcal{P}^* \rightarrow H/\mathcal{Q}$ is a bijection, it follows that $\dot{r}(A)$ and $\dot{r}(B)$ are distinct elements of H/\mathcal{Q} . Since \mathcal{Q} is an almost malnormal collection, Lemma 6.10 implies that for any integer m , the intersection $\mathcal{N}_m(\dot{r}(A)) \cap \mathcal{N}_m(\dot{r}(B))$ is a finite (and hence bounded) subset of H . Since $r: G \rightarrow H$ is a quasi-isometry, it follows that for every n , the intersection $\mathcal{N}_n(A) \cap \mathcal{N}_n(B)$ is a bounded (and hence finite) subset of G . ■

Theorem 6.12 (Theorem D). *Let $q: (G, \mathcal{P}) \rightarrow (H, \mathcal{Q})$ be a quasi-isometry of pairs. If \mathcal{Q} is an almost malnormal finite collection of infinite subgroups and \mathcal{P} is a finite collection, then any refinement \mathcal{P}^* of \mathcal{P} is almost malnormal and $q: (G, \mathcal{P}^*) \rightarrow (H, \mathcal{Q})$ is a quasi-isometry of pairs.*

Proof. The result follows from Propositions 6.3 and 6.8. ■

7. Examples

In this section, we show that there are examples of pairs (G, H) with a well-defined relative Dehn function outside of the context of relatively hyperbolic groups. Hyperbolically embedded subgroups were introduced in [4] by Dahmani, Guirardel and Osin. Given a group G , $X \subset G$ and $H \leq G$, let $H \hookrightarrow_h (G, X)$ denote that H is a hyperbolically embedded subgroup of G with respect to X . There is a characterisation in [18] of H being hyperbolically embedded into G that fits into the context of our Theorem E, namely, in terms of fine vertices in coned-off Cayley graphs (see Definition 5.1).

Proposition 7.1 ([18, Proposition 1.4]). *Let G be a group, $X \subset G$ and $H \leq G$. Then $H \hookrightarrow_h (G, X)$ if and only if $\hat{\Gamma}(G, H, X)$ is connected, hyperbolic, and fine at cone vertices.*

The following theorem provides our examples.

Theorem 7.2 (Theorem G). *Let G be a finitely presented group and $H \leq G$ be a subgroup. If $H \hookrightarrow_h G$, then the relative Dehn function $\Delta_{G,H}$ is well defined.*

The proof of the theorem is discussed after the following lemma.

Lemma 7.3. *Let G be a finitely generated group and H a finitely presented subgroup. Then G is finitely presented if and only if G is finitely presented relative to H .*

Proof. Suppose that G has a finite presentation $\langle A \mid R \rangle$. Let R_H be the collection of all relations in H over the generating set $H - \{1\}$, that is, $H = \langle H - \{1\} \mid R_H \rangle$. Let $\{h_1, \dots, h_k\}$ be a finite generating set of H . Then, there is a word w_i over the alphabet A that represents h_i . Observe that

$$\langle A \sqcup (H - \{1\}) \mid R, R_H, h_1 = w_1, \dots, h_k = w_k \rangle$$

yields a finite relative presentation of G with respect to H .

Conversely, suppose that $\langle A, H \mid R \rangle$ is a finite relative presentation of G with respect to H , and let $\langle B \mid T \rangle$ be a finite presentation of H . Then $\langle A \sqcup B, H \mid R \sqcup T, h_1 = w_1, \dots, h_k = w_k \rangle$ is a finite relative presentation of G with respect to H , where $\{h_1, \dots, h_k\} \subset H$ is a finite generating set of H and w_i is a word over B that represents the element h_i (after choosing an isomorphism $F(B)/\langle\langle T \rangle\rangle \rightarrow H$). This relative presentation yields a standard presentation $\langle A \sqcup B \sqcup (H - \{1\}) \mid R \sqcup T \sqcup R_H \sqcup \{h_1 = w_1, \dots, h_k = w_k\} \rangle$ of G , where R_H is the collection of all relations in H over the generating set $H - \{1\}$. Since the $\{h_1, \dots, h_k\}$ generate H , using Tietze transformations one obtains that $\langle A \sqcup B \mid R \sqcup T \sqcup \{h_1 = w_1, \dots, h_k = w_k\} \rangle$ is a presentation of G which is finite. ■

Proof of Theorem 7.2. First, note that the theorem is trivial in the case where H is a finite subgroup of G . Indeed, any finite subgroup is hyperbolically embedded by definition and a finite relative presentation of a group with respect to a finite subgroup is in fact a finite presentation. In particular, the relative Dehn function coincides with the Dehn function and the Dehn function of a finitely presented group is always well defined.

Since G is finitely presented and $H \hookrightarrow_h G$, it follows from [4, Corollary 4.32] that H is finitely presented. Hence, by Lemma 7.3, G is finitely presented relative to H .

Let S be a finite generating set of G . In view of Theorem E(2), to conclude that $\Delta_{G,H}$ is well defined, it is enough to prove that $\widehat{\Gamma}(G, \mathcal{P}, S)$ is a fine graph.

Suppose $H \hookrightarrow_h (G, X)$ for some $X \subset G$. Without loss of generality, assume that X contains the finite generating set S , see [4, Corollary 4.27]. It follows that $\widehat{\Gamma}(G, H, S)$ is a subgraph of $\widehat{\Gamma}(G, H, X)$. Since S is finite, observe that every vertex of $\widehat{\Gamma}(G, H, S)$ has either finite degree or is cone-vertex. By Proposition 7.1, the graph $\widehat{\Gamma}(G, H, X)$ is fine at every cone vertex, and hence so is $\widehat{\Gamma}(G, H, S)$. Therefore, $\widehat{\Gamma}(G, H, S)$ is a fine graph. ■

Example 7.4. In [6], the author shows that amongst graph products of finite groups various *eccentric subgroups* (see loc. cit. for a definition) are quasi-isometrically rigid in the

sense of [19]. Let G be a graph product of finite groups that is not virtually cyclic or a direct product of two infinite groups, then G is acylindrically hyperbolic. Suppose H is an eccentric subgroup, then $H \hookrightarrow_h G$ if and only if H is almost malnormal. In particular, if H is almost malnormal, then by Theorem 7.2, we see that $\Delta_{G,H}$ is well defined. Moreover, for any graph product of finite groups G' quasi-isometric to G , there exists a subgroup $H' < G'$, such that $\Delta_{G',H'} \asymp \Delta_{G,H}$.

The following example demonstrates that $\Delta_{G,\mathcal{P}}$ being well defined is not implied by \mathcal{P} being a qi-characteristic collection in the sense of [19].

Example 7.5. Let F be a finite group and let H be a finitely presented one-ended group. Consider the wreath product $G = F \wr H$. In the work of Genevois and Tessera [7, Proof of Theorem 7.1], they show that a quasi-isometry of $q: G \rightarrow G$ is a quasi-isometry of pairs $q: (G, H) \rightarrow (G, H)$. Moreover, H is an almost malnormal subgroup and, in fact, is qi-characteristic in the sense of [19], see [7, Theorem 1.18]. However, the coned-off Cayley graph of G with respect to H is not fine, so the group G cannot have a well-defined Dehn function by Theorem 4.17. To prove this, suppose F is the group with two elements and let H be a group with an element of infinite order a . Consider the wreath product $G = F \wr H$. If F has non-trivial element x , then G has a relative presentation

$$\langle x, H \mid x^2, [x, gxg^{-1}] \text{ for all } G - \{e\} \rangle.$$

Let us observe that the coned-off Cayley graph $\widehat{\Gamma}(G, H, \{x\})$ is not fine. Consider the edge $\{e, H\}$. We will show that there are infinitely many circuits of length twelve that contain this edge, each of them induced by a word

$$w_n = xa^n xa^{-n} xa^n xa^{-n}$$

which represents the identity. For an arbitrary integer $n > 0$, the sequence of vertices

$$\gamma_n = [H, e, x, xH, xa^n, xa^n x, xa^n xH, xa^n xa^{-n}, a^n xa^{-n}, a^n xH, a^n x, a^n, H]$$

is a closed path of length twelve in $\widehat{\Gamma}$ containing the edge $\{e, H\}$; the only non-trivial adjacency follows from $xa^n xa^{-n} = a^n xa^{-n}$. It follows that γ_n is a circuit since one can show that the left cosets $H, xH, xa^n xH, a^n xH$ are all distinct. On the other hand, $a^n xH = a^m xH$ if and only if $n = m$, and therefore $\gamma_n \neq \gamma_m$ if $m \neq n$. Note, we do not know the existence of a finite relative presentation for G with respect to H , but observe that we do not use this in the remark.

Finally, we will show the relative Dehn function of $\text{BS}(k, l)$ with respect to the stable letter is not well defined if either k or l divides the other one.

Example 7.6. Let $G = \text{BS}(k, l) = \langle a, t \mid ta^k t^{-1} = a^l \rangle$. We claim that if $k \mid l$ or $l \mid k$, then $\Delta_{G,\langle t \rangle}$ is not well defined. As in the previous example, we will show that the coned-off Cayley graph $\widehat{\Gamma}(G, \langle t \rangle, \{a, t\})$ is not fine and apply Theorem 4.17.

Without loss of generality, let $\ell = km$ and consider $w_n = t^n a^k t^{-n} a t^n a^{-k} t^{-n} a^{-1}$. Observe that $w_n = 1_G$ since $t^n a^k t^{-n} = a^{k\ell^n}$ and $t^n a^{-k} t^{-n} = a^{-k\ell^n}$. The word w_n describes a circuit of length $2k + 6$ in $\widehat{\Gamma}(G, \langle t \rangle, \{a, t\})$ because the four left cosets $\langle t \rangle$, $t^n a^k \langle t \rangle = a^{k\ell^n} \langle t \rangle$, $t^n a^k t^{-n} a \langle t \rangle = a^{k\ell^n + 1} \langle t \rangle$, and $t^n a^k t^{-n} a^{-k} \langle t \rangle = a \langle t \rangle$ are all distinct. In particular, the coned-off Cayley graph $\widehat{\Gamma}(G, \langle t \rangle, \{a, t\})$ is not fine.

A. Relative Dehn functions of Baumslag–Solitar groups (by Ashot Minasyan)

For two non-zero integers k and l , we define the Baumslag–Solitar group $BS(k, l)$ by the presentation

$$BS(k, l) = \langle a, t \mid ta^k t^{-1} = a^l \rangle.$$

Evidently, $BS(k, l)$ is finitely presented relative to its cyclic subgroup $\langle t \rangle$ and we can consider the relative presentation

$$BS(k, l) = \langle a, \langle t \rangle \mid \mathcal{R} \rangle, \tag{A.1}$$

where \mathcal{R} consists of all cyclic permutations of the relator $ta^k t^{-1} a^{-l}$ and its inverse.

Let $F = F(a, t)$ be the free group freely generated by $\{a, t\}$. The generating set $\{a\} \cup \langle t \rangle$ of F gives rise to the relative word length $\|\cdot\|_{\{a\} \cup \langle t \rangle}$ for words over the alphabet $\{a\}^{\pm 1} \cup \langle t \rangle$.

The goal of this appendix is to provide a characterisation for the Dehn function of $BS(k, l)$ with respect to $\langle t \rangle$ to be well defined (we shall use the definitions of the relative area and relative Dehn functions from Remark 4.5).

Theorem A.1. *Let $G = BS(k, l)$, for some non-zero integers k, l . The relative Dehn function $\Delta_{G, \langle t \rangle}$ is well defined if and only if k does not divide l and l does not divide k .*

Remark A.2. Theorem A.1 implies that the relative Dehn function of the group $G = BS(2, 3)$ with respect to the cyclic subgroup $\langle t \rangle$ is well defined. However, we note that $\langle t \rangle \not\hookrightarrow_h G$, so the converse of Theorem G is false. In fact, G does not contain any proper infinite hyperbolically embedded subgroups: see [22, Theorem 1.2 and Example 7.4].

Proof of Theorem A.1. The necessity has already been proved in Example 7.6, using Theorem 4.17. Below we give a different argument, based on the results of Osin [23].

Throughout the argument, we will use the following well-known elementary facts about $G = BS(k, l)$: the elements a and t have infinite order and $\langle a \rangle \cap \langle t \rangle = \{1\}$ in G .

Assume, without loss of generality, that k divides l , so that $l = km$ for some $m \in \mathbb{Z} \setminus \{0\}$. Arguing by contradiction, suppose that the Dehn function $\Delta_{G, \langle t \rangle}$ is well defined. Then, in accordance with [23, Proposition 2.36], $\langle t \rangle$ is a malnormal subgroup of G (i.e., $g \langle t \rangle g^{-1} \cap \langle t \rangle = \{1\}$ for any $g \in G \setminus \langle t \rangle$).

If $l = \pm k$, then $t^2 a^k t^{-2} = a^k$, so that $a^{-k} t^2 a^k = t^2 \in a^{-k} \langle t \rangle a^k \cap \langle t \rangle = \{1\}$, contradicting to the fact that t has infinite order in G . Therefore, we can further assume that $|k| \neq |l|$, so that $|m| > 1$.

For any $s \in \mathbb{N}$, we have $t^s a^k t^{-s} = a^{m^s k}$ in G , whence the commutator word

$$W_s = [t^s a^k t^{-s}, a] = t^s a^k t^{-s} a t^s a^{-k} t^{-s} a^{-1}$$

represents the trivial element of G . Note that $\|W_s\|_{\{a\} \cup \langle t \rangle} = 2k + 6$, so, since the Dehn function $\Delta_{G, \langle t \rangle}$ is well defined, there exists a constant $C \geq 0$ such that

$$\text{Area}^{\text{rel}}(W_s) \leq C \quad \text{for all } s \in \mathbb{N}.$$

For each $s \in \mathbb{N}$, let q_s be the cycle in the Cayley graph $\Gamma(G, \{a\} \cup \langle t \rangle \setminus \{1\})$ based at the identity element and labelled by the word W_s . By the definition of W_s , q_s is a concatenation of eight subpaths p_1, p_2, \dots, p_8 , where p_1 is the edge labelled by t^k , p_2 has length $|k|$ and is labelled by a^k , and so on: see Figure 3.




Figure 3. The cycle q_s (markings inside the polygon represent the labels of the subpaths p_1, \dots, p_8).

Using Osin’s terminology from [23, Section 2.2], we see that p_1, p_3, p_5 and p_7 is the list of $\langle t \rangle$ -components of q . Let us show that p_1 is an isolated component of q_s . Indeed, if p_1 is connected to p_3 , then the label of p_2, a^k , must represent an element of $\langle t \rangle$ in G . The latter is impossible since $\langle a \rangle \cap \langle t \rangle = \{1\}$ in G and $a^k \neq 1$. Similarly, p_1 cannot be connected to p_7 . Finally, if p_1 is connected to p_5 , then the label of the path $p_1 p_2 p_3 p_4$ must represent an element of $\langle t \rangle$ in G . However, this label is equal to $t^s a^k t^{-s} a$, which simplifies to $a^{m^s k + 1}$ in G . This again yields a contradiction because $m^s k + 1 \neq 0$ (which is true as $|m| > 1$ and $s \in \mathbb{N}$).

Therefore, we can apply [23, Lemma 2.27] to the cycle q_s , claiming that

$$|t^s|_\Omega \leq M \text{Area}^{\text{rel}}(W_s),$$

where $\Omega = \{t, t^{-1}\}$ and $M = \max\{\|R\|_{\{a\} \cup \langle t \rangle} \mid R \in \mathcal{R}\} = |k| + |l| + 2$. It follows that $s \leq MC$ for all $s \in \mathbb{N}$. This contradiction shows that the Dehn function $\Delta_{G, \langle t \rangle}$ is not well defined, so the necessity statement of the theorem has been proved.

The proof of the sufficiency occupies the rest of the appendix and will be completed in Theorem A.8 below. ■

A.1. Notation

We will use \mathbb{Z} to denote the set of all integers, $\mathbb{N} = \{1, 2, \dots\}$ – the set of natural numbers and $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. Given a prime p and an integer $n \in \mathbb{Z} \setminus \{0\}$, we will write

$$v_p(n) = \max\{s \in \mathbb{N}_0: p^s \text{ divides } n\} \in \mathbb{N}_0.$$

Evidently, $v_p(n) \leq \log_p(|n|)$ and $v_p(mn) = v_p(m) + v_p(n)$ for all $m, n \in \mathbb{Z} \setminus \{0\}$.

Further on k, l will be some fixed non-zero integers and G will be the Baumslag–Solitar group $BS(k, l)$, equipped with the relative presentation (A.1). For two words W and W' over the alphabet $\{a\} \cup \langle t \rangle$, we will write $W \stackrel{G}{=} W'$ if W and W' represent the same element of G .

A.2. Some terminology

By the normal form theorem for free products, we know that any word W over the alphabet $\{a\}^{\pm 1} \cup \langle t \rangle$ is equal in $F = F(a, t)$ to a unique *freely reduced word*, which has the form

$$a^{u_0} t^{v_1} a^{u_1} \dots t^{v_m} a^{u_m}, \quad \text{where } m \geq 0, u_0, u_m \in \mathbb{Z},$$

$$u_1, \dots, u_{m-1}, v_1, \dots, v_m \in \mathbb{Z} \setminus \{0\}, \tag{A.2}$$

and $t^{v_1}, \dots, t^{v_m} \in \langle t \rangle \setminus \{1\}$ are treated as single letters from the alphabet $\{a\}^{\pm 1} \cup \langle t \rangle$. We will call the number m the *syllable length of W* and will denote it $sl(W)$. Observe that $sl(W) \leq \|W\|_{\{a\} \cup \langle t \rangle}$ for any freely reduced word W .

Definition A.3 (Reduction of the first type). Suppose that W is a word of form (A.2). If for some $i \in \{1, \dots, m - 1\}$, we have $v_i > 0, v_{i+1} < 0$ and $u_i \in k\mathbb{Z} \setminus \{0\}$, then we can perform a reduction of the first type on W as follows.

Set $s = u_i/k \in \mathbb{Z}$ and observe that, by applying a defining relation from presentation (A.1) $|s|$ times, we get

$$t a^{u_i} t^{-1} \stackrel{G}{=} (t a^k t^{-1})^s \stackrel{G}{=} a^{ls} \quad \text{in } G.$$

Therefore, W is equal in G to the word

$$W' = a^{u_0} t^{v_1} a^{u_1} \dots a^{u_{i-1}} t^{v_i-1} a^{ls} t^{v_{i+1}+1} a^{u_{i+1}} \dots t^{v_m} a^{u_m}. \tag{A.3}$$

We will say that W' has been obtained from W by applying a *reduction of the first type at place i* , writing $W \xrightarrow{i} W'$.

We can similarly define basic reductions of the second type.

Definition A.4 (Reduction of the second type). Suppose that W is a word of form (A.2). If for some $i \in \{1, \dots, m - 1\}$, we have $v_i < 0$, $v_{i+1} > 0$ and $u_i \in \mathbb{Z} \setminus \{0\}$, then we can perform a reduction of the second type on W as follows.

Set $s = u_i/l \in \mathbb{Z}$ and observe that, by applying a defining relation from presentation (A.1) $|s|$ times, we get

$$t^{-1} a^{u_i} t \stackrel{G}{=} (t^{-1} a^l t)^s \stackrel{G}{=} a^{ks} \quad \text{in } G.$$

Therefore, W is equal in G to the word

$$W' = a^{u_0} t^{v_1} a^{u_1} \dots a^{u_{i-1}} t^{v_i+1} a^{ls} t^{v_{i+1}-1} a^{u_{i+1}} \dots t^{v_m} a^{u_m}. \tag{A.4}$$

We will say that W' has been obtained from W by applying a *reduction of the second type at place i* , writing $W \xrightarrow[i]{2} W'$.

When the type of the reduction does not matter, we will simply write $W \xrightarrow[i]{} W'$. Note that after applying a reduction (of any type) to a freely reduced word W the resulting word W' satisfies $\text{sl}(W') \leq \text{sl}(W)$. Moreover, if $\text{sl}(W') = \text{sl}(W)$ then the word W' (from (A.3) or (A.4)) is again freely reduced in the above sense.

Definition A.5 (Trimming chain). Let W be a freely reduced word over the alphabet $\{a\}^{\pm 1} \cup \{t\}$ and $i \in \{1, \dots, \text{sl}(W) - 1\}$. For any $\ell \in \mathbb{N}$, a *trimming chain of the first type at place i of length ℓ* is a sequence of reductions

$$W = W_0 \xrightarrow[i]{1} W_1 \xrightarrow[i]{1} \dots \xrightarrow[i]{1} W_{\ell-1} \xrightarrow[i]{1} W_\ell,$$

where $\text{sl}(W) = \text{sl}(W_1) = \dots = \text{sl}(W_{\ell-1})$ and $\text{sl}(W_\ell) < \text{sl}(W)$.

A *trimming chain of the second type at place i of length ℓ* ,

$$W = W_0 \xrightarrow[i]{2} W_1 \xrightarrow[i]{2} \dots \xrightarrow[i]{2} W_{\ell-1} \xrightarrow[i]{2} W_\ell,$$

is defined similarly.

A.3. Technical lemmas

From now on, we assume that $k \nmid l$ and $l \nmid k$. In this case, we can choose some primes $p, q \in \mathbb{N}$ such that $v_p(k) > v_p(l)$ and $v_q(l) > v_q(k)$.

Lemma A.6. *Let W be the word given by (A.2) with $\text{sl}(W) = m > 0$. If W represents the trivial element of G , then there is $i \in \{1, \dots, m - 1\}$ such that either W admits a trimming chain of the first type at place i of length at most $v_p(u_i)$ or it admits a trimming chain of the second type at place i of length at most $v_q(u_i)$.*

Proof. We will prove the statement by induction on the total number of t 's occurring in W , i.e., on the number $v(W) = \sum_{r=1}^m |v_r|$.

Since $W \stackrel{G}{=} 1$, by Britton’s lemma (see [15, Section IV.2]), the number $v(W)$ must be at least 2, and if $v(W) = 2$, then $\text{sl}(W) = m = 2$ and either $v_1 = 1, v_2 = -1$ and $u_1 \in k\mathbb{Z} \setminus \{0\}$ (i.e., W admits a reduction of the first type) or $v_1 = -1, v_2 = 1$ and $u_1 \in l\mathbb{Z} \setminus \{0\}$ (i.e., W admits a reduction of the second type). Without loss of generality, let us assume that we are in the former case. Applying a reduction of the first type to W , we obtain a word W' with $\text{sl}(W') = 0 < \text{sl}(W)$, so $W \xrightarrow{i} W'$ is a trimming chain of the first type at place 1 of length 1. Moreover, $1 \leq v_p(u_1)$, as $p \mid k \mid u_1$, so the base of induction has been established.

Suppose now that $v(W) > 2$. By Britton’s lemma, W admits a reduction (of some type) at some place $i \in \{1, \dots, \text{sl}(W) - 1\}$, and again, without loss of generality, we assume that it is a reduction of the first type (the other case is similar). Let W' be the word (A.3) resulting in this reduction.

If $\text{sl}(W') < \text{sl}(W)$, then $W \xrightarrow{i} W'$ is a trimming chain of the first type of length $1 \leq v_p(u_i)$, as required. So we can further assume that $\text{sl}(W') = \text{sl}(W) = m$, whence W' is again freely reduced and $v(W') = v(W) - 2 < v(W)$. By induction, W' must admit a trimming chain (of one of the two types) at some place $j \in \{1, \dots, m - 1\}$ of length $\ell \in \mathbb{N}$. If $j \neq i$, then we can perform the same trimming chain on W since u_j is not affected by the original reduction $W \xrightarrow{i} W'$ and $\text{sl}(W') = \text{sl}(W)$. The desired inequality on ℓ will then follow by induction.

Now let us suppose that $j = i$. Since $\text{sl}(W') = \text{sl}(W)$, the trimming chain at place i for W' must have the same type as the original reduction from W to W' , thus we have a trimming chain

$$W' = W_0 \xrightarrow{i} W_1 \xrightarrow{i} \dots \xrightarrow{i} W_{\ell-1} \xrightarrow{i} W_\ell.$$

By precomposing this trimming chain with the original reduction $W \xrightarrow{i} W'$, we obtain a trimming chain of the first type at place i of length $\ell + 1$ for W . By induction and the construction of W' (see (A.3)), we have $\ell \leq v_p(ls)$, where $s = u_i/k \in \mathbb{Z} \setminus \{0\}$. Since $v_p(k) \geq v_p(l) + 1$, we can conclude that

$$\ell + 1 \leq v_p(ls) + 1 = v_p(l) + v_p(s) + 1 \leq v_p(k) + v_p(s) = v_p(ks) = v_p(u_i).$$

Thus we have established the step of induction, and so the statement is proved. ■

Denote

$$\alpha = \max\left\{\left|\frac{l}{k}\right|, \left|\frac{k}{l}\right|\right\} > 1. \tag{A.5}$$

Lemma A.7. *Let W be a word of form (A.2), representing the trivial element of G . Suppose that*

$$W = W_0 \xrightarrow{i} W_1 \xrightarrow{i} \dots \xrightarrow{i} W_{\ell-1} \xrightarrow{i} W_\ell \tag{A.6}$$

is a sequence of reductions (of the same type) at place $i \in \{1, \dots, \text{sl}(W) - 1\}$, where $\ell \in \mathbb{N}$ and $\text{sl}(W) = \text{sl}(W_1) = \dots = \text{sl}(W_{\ell-1})$. Denote $n = \|W\|_{\{a\} \cup \langle t \rangle} \in \mathbb{N}$, then

$$\|W_\ell\|_{\{a\} \cup \langle t \rangle} \leq \alpha^\ell n \quad \text{and} \quad \text{Area}^{\text{rel}}(W) \leq \text{Area}^{\text{rel}}(W_\ell) + \frac{\alpha^\ell - 1}{\alpha - 1} n. \tag{A.7}$$

Proof. Without loss of generality, we will assume that all of the reductions in the sequence (A.6) are of the first type. We will argue by induction on ℓ .

Suppose, first, $\ell = 1$ and $W_1 = a^{u_0} t^{v_1} a^{u_1} \dots a^{u_{i-1}} t^{v_i-1} a^{ls} t^{v_{i+1}+1} a^{u_{i+1}} \dots t^{v_m} a^{u_m}$, where $s = u_i/k$. Then

$$n = \|W\|_{\{a\} \cup \langle t \rangle} = m + \sum_{r=0}^m |u_r| \quad \text{and} \quad \|W_1\|_{\{a\} \cup \langle t \rangle} \leq m + \sum_{r=0, r \neq i}^m |u_r| + |ls|.$$

Since $|ls|/|u_i| = |l/k| \leq \alpha$ and $\alpha > 1$, we see that $\|W_1\|_{\{a\} \cup \langle t \rangle} \leq \alpha n$. The word W_1 can be obtained from the word W by applying a defining relation from presentation (A.1) $|s|$ times, so, since $|s| \leq |u_i| \leq n$, we have

$$\text{Area}^{\text{rel}}(W) \leq \text{Area}^{\text{rel}}(W_1) + |s| \leq \text{Area}^{\text{rel}}(W_1) + n.$$

Now assume that $\ell \geq 2$ and set $n_1 = \|W_1\|_{\{a\} \cup \langle t \rangle}$. By induction, we know that

$$\|W_\ell\|_{\{a\} \cup \langle t \rangle} \leq \alpha^{\ell-1} n_1 \quad \text{and} \quad \text{Area}^{\text{rel}}(W_1) \leq \text{Area}^{\text{rel}}(W_\ell) + \frac{\alpha^{\ell-1} - 1}{\alpha - 1} n_1. \tag{A.8}$$

We have shown above that $n_1 \leq \alpha n$ and $\text{Area}^{\text{rel}}(W) \leq \text{Area}^{\text{rel}}(W_1) + n$. Combining this with inequalities (A.8), we obtain (A.7). ■

A.4. Proof of the sufficiency in Theorem A.1

Theorem A.8. *Let G be the Baumslag–Solitar group $\text{BS}(k, l)$ for some $k, l \in \mathbb{Z} \setminus \{0\}$. If neither of k, l divides the other one, then the relative Dehn function $\Delta_{G, \langle t \rangle}$ is well defined.*

Proof. Choose primes $p, q \in \mathbb{N}$ as in the beginning of Section A.3 and let $\alpha > 1$ be defined by (A.5).

To prove that $\Delta_{G, \langle t \rangle}$ is well defined, it is sufficient to show that there is a function $h: \mathbb{N}_0 \times \mathbb{N}_0 \rightarrow \mathbb{N}_0$ such that for all $m, n \in \mathbb{N}_0$ if W is a freely reduced word over the alphabet $\{a\}^{\pm 1} \cup \langle t \rangle$, representing the trivial element of G and satisfying $\text{sl}(W) = m$ and $\|W\|_{\{a\} \cup \langle t \rangle} = n$, then

$$\text{Area}^{\text{rel}}(W) \leq h(m, n).$$

(Since $\text{sl}(W) \leq \|W\|_{\{a\} \cup \langle t \rangle}$, the function $f: \mathbb{N}_0 \rightarrow \mathbb{N}_0$, $f(n) = \max\{h(m', n'): 0 \leq m', n' \leq n\}$ will serve as a relative isoperimetric function of G with respect to $\langle t \rangle$.)

The proof will use induction on m . By Britton’s lemma, a freely reduced word W of syllable length at most 1 cannot represent the trivial element of G , hence we can define $h(0, n) = h(1, n) = 0$ for all $n \in \mathbb{N}_0$.

Now suppose that $m \geq 2$ and the values of the desired function $h(s, n)$ have been found for all $s \in \{0, \dots, m - 1\}$ and all $n \in \mathbb{N}_0$. Take any $n \in \mathbb{N}_0$. If there are no freely reduced words W such that $\text{sl}(W) = m$, $\|W\|_{\{a\} \cup \{t\}} = n$ and $W \stackrel{G}{=} 1$ in G , then we set $h(m, n) = 0$. Otherwise, let W be such a word (in particular, $n \geq m \geq 2$).

If W is given by (A.2), then according to Lemma A.6, W admits a trimming chain

$$W = W_0 \xrightarrow{i} W_1 \xrightarrow{i} \dots \xrightarrow{i} W_{\ell-1} \xrightarrow{i} W_\ell$$

at some place $i \in \{1, \dots, m - 1\}$ of length $\ell \in \mathbb{N}$, where $\ell \leq \max\{v_p(u_i), v_q(u_i)\}$. Since $|u_i| \leq n$, we see that $\ell \leq \max\{\log_p(n), \log_q(n)\} = \log_r(n)$, where $r = \min\{p, q\}$.

Define $\beta = \log_r(\alpha) + 1$ and observe that

$$\alpha^\ell n \leq \alpha^{\log_r(n)} n = r^{\log_r(\alpha) \log_r(n)} n = n^{\log_r(\alpha)+1} = n^\beta,$$

and

$$\frac{\alpha^\ell - 1}{\alpha - 1} n \leq \frac{1}{\alpha - 1} \alpha^\ell n \leq \frac{1}{\alpha - 1} n^\beta.$$

Thus inequalities (A.7), given by Lemma A.7, imply that

$$\|W_\ell\|_{\{a\} \cup \{t\}} \leq n^\beta \quad \text{and} \quad \text{Area}^{\text{rel}}(W) \leq \text{Area}^{\text{rel}}(W_\ell) + \frac{1}{\alpha - 1} n^\beta. \tag{A.9}$$

Since $m' = \text{sl}(W_\ell) < \text{sl}(W)$, by induction we have $\text{Area}^{\text{rel}}(W_\ell) \leq h(m', n')$, where $n' = \|W_\ell\|_{\{a\} \cup \{t\}}$. In view of (A.9), after defining

$$h(m, n) = \max \left\{ h(m', n') + \left\lfloor \frac{1}{\alpha - 1} n^\beta \right\rfloor \mid 0 \leq m' \leq m - 1, 0 \leq n' \leq n^\beta \right\} \in \mathbb{N}_0,$$

we shall have $\text{Area}^{\text{rel}}(W) \leq h(m, n)$.

Thus we have found the required function $h: \mathbb{N}_0 \times \mathbb{N}_0 \rightarrow \mathbb{N}_0$, so the proof of the theorem is complete. ■

Remark A.9. The argument from the proof of Theorem A.8 gives a double exponential upper bound $\Delta_{G, \{t\}}(n) \leq n^{\beta n}$ for all $n \in \mathbb{N}_0$, where $\beta > 1$ is the constant from that proof.

Acknowledgements. The authors thank Ashot Minasyan for comments on an earlier version of the manuscript, and for pointing out a necessary correction in the proof of Theorem G. The authors also thank Anthony Genevois for comments and pointing us to his work with Tessera, see Example 7.5. The first author would like to thank his PhD supervisor Professor Ian Leary. The second author thanks Noel Brady for discussions on the topics of the article. All three authors would like to thank the organisers of the online seminar ‘‘Algebra at Bicocca’’, without which this collaboration would not have happened. We thank the anonymous referee for a number of helpful comments.

Funding. The first author was supported by the Engineering and Physical Sciences Research Council grant number 2127970. The second author acknowledges funding by the Natural Sciences and Engineering Research Council of Canada, NSERC. The third author was supported by grant PAPIIT-IA101221.

References

- [1] J. Behrstock, C. Druţu, and L. Mosher, [Thick metric spaces, relative hyperbolicity, and quasi-isometric rigidity](#). *Math. Ann.* **344** (2009), no. 3, 543–595 Zbl 1220.20037 MR 2501302
- [2] B. H. Bowditch, [Relatively hyperbolic groups](#). *Internat. J. Algebra Comput.* **22** (2012), no. 3, article no. 1250016 Zbl 1259.20052 MR 2922380
- [3] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*. Grundlehren Math. Wiss. 319, Springer, Berlin, 1999 Zbl 0988.53001 MR 1744486
- [4] F. Dahmani, V. Guirardel, and D. Osin, [Hyperbolically embedded subgroups and rotating families in groups acting on hyperbolic spaces](#). *Mem. Amer. Math. Soc.* **245** (2016), no. 1156, v+152 pp. Zbl 1396.20041 MR 3589159
- [5] B. Farb, [Relatively hyperbolic groups](#). *Geom. Funct. Anal.* **8** (1998), no. 5, 810–840 Zbl 0985.20027 MR 1650094
- [6] A. Genevois, [Quasi-isometrically rigid subgroups in right-angled Coxeter groups](#). *Algebr. Geom. Topol.* **22** (2022), no. 2, 657–708 Zbl 07570603 MR 4464462
- [7] A. Genevois and R. Tessera, [Asymptotic geometry of lamplighters over one-ended groups](#). 2021, arXiv:2105.04878
- [8] D. Groves and J. F. Manning, [Dehn filling in relatively hyperbolic groups](#). *Israel J. Math.* **168** (2008), 317–429 Zbl 1211.20038 MR 2448064
- [9] V. S. Guba and M. V. Sapir, [On Dehn functions of free products of groups](#). *Proc. Amer. Math. Soc.* **127** (1999), no. 7, 1885–1891 Zbl 0942.20023 MR 1469408
- [10] M. Haulmark and G. C. Hruska, [On canonical splittings of relatively hyperbolic groups](#). *Israel J. Math.* (2023)
- [11] B. B. Healy and G. C. Hruska, [Cusped spaces and quasi-isometries of relatively hyperbolic groups](#). 2021, arXiv:2010.09876
- [12] G. C. Hruska, [Relative hyperbolicity and relative quasiconvexity for countable groups](#). *Algebr. Geom. Topol.* **10** (2010), no. 3, 1807–1856 Zbl 1202.20046 MR 2684983
- [13] S. Hughes and E. Martínez-Pedroza, [Hyperbolically embedded subgroups and quasi-isometries of pairs](#). *Canad. Math. Bull.* (2023)
- [14] S. Hughes, E. Martínez-Pedroza, and L. J. Sánchez Saldaña, [A survey on quasi-isometries of pairs: invariants and rigidity](#). 2021, arXiv:2112.15046
- [15] R. C. Lyndon and P. E. Schupp, *Combinatorial group theory*. Classics Math., Springer, Berlin, 2001 Zbl 0997.20037 MR 1812024
- [16] E. Martínez-Pedroza, [Combination of quasiconvex subgroups of relatively hyperbolic groups](#). *Groups Geom. Dyn.* **3** (2009), no. 2, 317–342 Zbl 1186.20029 MR 2486802
- [17] E. Martínez-Pedroza, [A note on fine graphs and homological isoperimetric inequalities](#). *Canad. Math. Bull.* **59** (2016), no. 1, 170–181 Zbl 1360.20037 MR 3451908
- [18] E. Martínez-Pedroza and F. Rashid, [A note on hyperbolically embedded subgroups](#). *Comm. Algebra* **50** (2022), no. 4, 1459–1468 Zbl 07488675 MR 4391500

- [19] E. Martínez-Pedroza and L. J. Sánchez Saldaña, [Quasi-isometric rigidity of subgroups and filtered ends](#). *Algebr. Geom. Topol.* **22** (2022), no. 6, 3023–3057 Zbl [1510.20039](#) MR [4520684](#)
- [20] J. P. McCammond and D. T. Wise, [Fans and ladders in small cancellation theory](#). *Proc. London Math. Soc. (3)* **84** (2002), no. 3, 599–644 Zbl [1022.20012](#) MR [1888425](#)
- [21] L. Mosher, M. Sageev, and K. Whyte, [Quasi-actions on trees II: Finite depth Bass–Serre trees](#). *Mem. Amer. Math. Soc.* **214** (2011), no. 1008, vi+105 pp. Zbl [1234.20034](#) MR [2867450](#)
- [22] D. Osin, [Acylindrically hyperbolic groups](#). *Trans. Amer. Math. Soc.* **368** (2016), no. 2, 851–888 Zbl [1380.20048](#) MR [3430352](#)
- [23] D. V. Osin, [Relatively hyperbolic groups: intrinsic geometry, algebraic properties, and algorithmic problems](#). *Mem. Amer. Math. Soc.* **179** (2006), no. 843, vi+100 pp. Zbl [1093.20025](#) MR [2182268](#)
- [24] J.-P. Serre, *Trees*. Springer Monogr. Math., Springer, Berlin, 2003 Zbl [1013.20001](#) MR [1954121](#)

Received 22 September 2021; revised 25 January 2022.

Sam Hughes

Mathematical Institute, University of Oxford, Andrew Wiles Building, Observatory Quarter, OX2 6GG Oxford, UK; sam.hughes@maths.ox.ac.uk

Eduardo Martínez-Pedroza

Department of Mathematics and Statistics, Memorial University of Newfoundland, St. John's, NL, A1C 5S7, Canada; emartinezped@mun.ca

Luis Jorge Sánchez Saldaña

Universidad Nacional Autónoma de México, Av. Universidad 3000, 04510 México, Mexico; luisjorge@ciencias.unam.mx

Ashot Minasyan

School of Mathematical Sciences, University of Southampton, Highfield, SO17 1BJ Southampton, UK; aminasyan@gmail.com