

The derived non-commutative Poisson bracket on Koszul Calabi–Yau algebras

Xiaojun Chen, Alimjon Eshmatov, Farkhod Eshmatov and Song Yang

Abstract. Let A be a Koszul (or more generally, N -Koszul) Calabi–Yau algebra. Inspired by the works of Kontsevich, Ginzburg and Van den Bergh, we show that there is a derived non-commutative Poisson structure on A , which induces a graded Lie algebra structure on the cyclic homology of A ; moreover, we show that the Hochschild homology of A is a Lie module over the cyclic homology and the Connes long exact sequence is in fact a sequence of Lie modules. Finally, we show that the Leibniz–Loday bracket associated to the derived non-commutative Poisson structure on A is naturally mapped to the Gerstenhaber bracket on the Hochschild cohomology of its Koszul dual algebra and hence on that of A itself. Relations with some other brackets in literature are also discussed and several examples are given in detail.

Mathematics Subject Classification (2010). 14A22; 16S38.

Keywords. Noncommutative Poisson structure, Calabi–Yau algebra, cyclic homology.

1. Introduction

The notion of Calabi–Yau algebras is introduced by Ginzburg [23], and has been intensively studied in recent years. They are associative algebras with some additional properties, and may be viewed as non-commutative generalization of affine Calabi–Yau varieties. It turns out that they are related to representation theory, non-commutative symplectic/algebraic geometry, mirror symmetry, and much more. For more details, see, for example, [7, 8, 13, 23, 32] and references therein.

In this paper, we study the *derived non-commutative Poisson structure* on Koszul (or more generally, N -Koszul in the sense of Berger [4]) Calabi–Yau algebras, continuing the work of Berest, Chen, Eshmatov and Ramadoss [1]. Let us start with some background.

1.1. Derived non-commutative Poisson structures. Let k be an algebraically closed field of characteristic zero. In 2005 Crawley-Boevey [12] introduced for associative algebras the notion of H_0 -Poisson structure. Suppose A is an associative algebra over k , then an H_0 -Poisson structure on A is a Lie bracket on $A/[A, A]$ such

that

$$[\bar{a}, -] : A/[A, A] \rightarrow A/[A, A]$$

is introduced by a derivation $d_{\bar{a}} : A \rightarrow A$, for all $\bar{a} \in A/[A, A]$. Such a notion perfectly fits a principle raised by Kontsevich and Rosenberg [34] in the study of non-commutative geometry, that is, any non-commutative geometric structure (such as the non-commutative symplectic, non-commutative Poisson, etc.) on a non-commutative space (here we mean an associative algebra) should induce its classical counterpart on the moduli space of its representations, i.e. on its representation scheme. Recall that for an associative algebra A , $A/[A, A]$ is always considered as the space of functions on A , and there is the canonical *trace map*

$$\begin{aligned} \text{Tr} : A/[A, A] &\longrightarrow k[\text{Rep}_V(A)] \\ \bar{a} &\longmapsto \{\rho \mapsto \text{trace}(\rho(a))\} \end{aligned} \quad (1.1)$$

from the functions on A to the functions on the representation scheme of A in a k -vector space V . The trace map is $\text{GL}(V)$ -invariant, and Crawley-Boevey showed that if A admits an H_0 -Poisson structure, then it naturally induces via the trace map a unique Poisson structure on $\text{Rep}_V(A)//\text{GL}(V)$ for all $n \in \mathbb{N}$ such that Tr is a map of Lie algebras. The notation “ H_0 ” means the *zero-th homology*, since $A/[A, A]$ is the zero-th Hochschild/cyclic homology of A .

In 2012 the H_0 -Poisson structure was generalized to the higher degree case in [1], where all cyclic homology groups are taken into account. The starting point is that the trace map (1.1) is not perfect in the sense that in very rare cases $\text{Rep}_V(A)$ is a smooth variety (cf. [14] for further studies), and there are obstructions for $\text{Rep}_V(A)$ to have the desired geometric property. The work [3] shows that instead one has to consider the homotopy category (in the sense of Quillen) of DG associative algebras. The main idea is to replace the associative algebra A by its cofibrant resolution QA , and then consider the DG representations of QA . It turns out that: (i) there is a surjective map from the cyclic homology $\text{HC}_\bullet(A)$ to the homology of the commutator quotient space $QA/[QA, QA]$; (ii) the n -dimensional DG representation scheme of QA , which up to homotopy is denoted by $\text{DRep}_V(A)$, is smooth in the differential graded sense. By passing to the homotopy category one obtains a natural map (called the *derived trace map*)

$$\text{HC}_\bullet(A) \longrightarrow \text{H}_\bullet(\text{DRep}_V(A)) \quad (1.2)$$

from the cyclic homology of A to the homology of the *derived representation scheme* of A . For more details of $\text{DRep}_V(A)$, one may refer to [2, 3].

The work [1] may be viewed as an application of the general result of [2, 3]. In that paper, an algebra A is called to admit a *derived non-commutative Poisson* structure if there is a DG non-commutative Poisson structure in the sense of Crawley-Boevey on its cofibrant resolution. It is proved in [1] that if A admits a derived non-commutative Poisson structure, then (1.2) induces a unique graded Poisson structure on the derived representation schemes.

As an important example, it is shown in [1] that the cobar construction $\Omega(C)$ of a cyclic coalgebra C (“cyclic” here means the dual space of C is a cyclic associative algebra) admits a derived non-commutative Poisson structure. Since $\Omega(C)$ is always a quasi-free (and hence cofibrant) DG algebra, from the above argument one obtains that any algebra A which is quasi-isomorphic to $\Omega(C)$ admits a derived non-commutative Poisson structure as well. It is exactly at this point that Koszul Calabi–Yau algebras come in.

1.2. Koszul Calabi–Yau algebras. According to Ginzburg [23], an associative algebra A is called *Calabi–Yau of dimension d* (or *d -Calabi–Yau* for short) if

- A is homologically smooth, that is, it has a finite resolution of finitely generated projective $A \otimes A^{\text{op}}$ modules;
- there exists an isomorphism

$$\text{RHom}_{A \otimes A^{\text{op}}}(A, A \otimes A) \simeq A[-d]$$

in the derived category of $A \otimes A^{\text{op}}$ modules.

Ginzburg in *loc. cit.* also showed that if a Calabi–Yau algebra A is Koszul, then its Koszul dual algebra, denoted by $A^!$, is cyclic. Dually, the Koszul dual coalgebra of A , denoted by A^i , is a cyclic coalgebra. From Koszul duality theory there is a quasi-isomorphism

$$\Omega(A^i) \xrightarrow{\simeq} A,$$

and since $\Omega(A^i)$ is cofibrant, by the work [1] sketched above, we thus obtain a derived non-commutative Poisson structure on A .

In fact, a slightly more general class of Calabi–Yau algebras has the above property. In [4] Berger introduced the notion of N -Koszul algebras, where 2-Koszul is Koszul in the usual sense. If a Calabi–Yau algebra A is N -Koszul, then its Koszul dual coalgebra A^i is a cyclic A_∞ coalgebra. Denote the cobar construction of A^i by $\Omega_\infty(A^i)$; we still have $A \simeq \Omega_\infty(A^i)$. There are many examples of N -Koszul Calabi–Yau algebras, such as the Sklyanin algebras, universal enveloping algebra of semi-simple Lie algebras, and Yang–Mills algebras, etc. The theorem below studies the derived non-commutative Poisson structure on N -Koszul Calabi–Yau algebras in this general setting:

Theorem 1.1. *Let A be an N -Koszul d -Calabi–Yau algebra. Then*

- (1) *there is a degree $2 - d$ derived non-commutative Poisson structure on A , which induces a degree $2 - d$ graded Lie algebra structure on the cyclic homology $\text{HC}_\bullet(A)$ of A ; and*
- (2) *there is a degree $2 - d$ Lie module structure on the Hochschild homology $\text{HH}_\bullet(A)$ over $\text{HC}_\bullet(A)$.*

The first statement in the theorem may be viewed as an application of [1, Lemma 11] to the Koszul Calabi–Yau case, and it also answers a question raised in the last paragraph in *loc. cit.* §5.4; the second statement is new.

1.3. The Connes long exact sequence of Lie modules. The key ingredient in the above theorem is the structure of a differential graded version of the *double Poisson bracket* in the sense of Van den Bergh [45] on $\Omega_\infty(\tilde{A}^i)$, where $\tilde{A}^i := k \oplus A^i$ is the co-augmentation of A^i . According to Van den Bergh, a double Poisson bracket on an associative algebra, say R , is a bilinear map

$$\{\{-, -\}\} : R \times R \rightarrow R \otimes R$$

satisfying some additional conditions. If R admits a double Poisson structure, then the commutator quotient space $R_{\natural} = R/[R, R]$ naturally admits a Lie algebra structure satisfying the criterion of Crawley-Boevey. From the famous result of Feigin and Tsygan [20], for a Koszul algebra A , the homology of $\Omega_\infty(\tilde{A}^i)_{\natural}$ is exactly $\mathrm{HC}_\bullet(A)$ and Theorem 1.1 part (1) follows.

Moreover, we recall that for an associative algebra A there is a well-known long exact sequence due to Connes, relating the cyclic and Hochschild homologies:

$$\cdots \xrightarrow{B} \mathrm{HH}_\bullet(A) \xrightarrow{I} \mathrm{HC}_\bullet(A) \xrightarrow{S} \mathrm{HC}_{\bullet-2}(A) \xrightarrow{B} \mathrm{HH}_{\bullet-1}(A) \xrightarrow{I} \cdots$$

By using an elegant interpretation of the Hochschild and cyclic homology of an algebra in terms of its bar construction, which is due to Quillen [41], and by Theorem 1.1 part (1), we in fact obtain that $\mathrm{HH}_\bullet(A)$ and $\mathrm{HC}_\bullet(A)$ are Lie modules over the Lie algebra $\mathrm{HC}_\bullet(A)$, from which Theorem 1.1 part (2) follows, and we now claim that

Theorem 1.2. *The maps B , I and S are morphisms of Lie modules of degree $2 - d$.*

1.4. The Poisson structure on derived representation schemes. The significance of the above two theorems is that the Lie module morphism naturally induces a Lie module morphism on the (derived) representation scheme of the Calabi–Yau algebra. As we mentioned above, there is a derived trace map

$$\mathrm{Tr} : \mathrm{HC}_\bullet(A) \rightarrow \mathrm{H}_\bullet(\mathrm{DRep}_V(A)).$$

The images are $\mathrm{GL}(V)$ -invariant, and in [3, Theorem 5.2] this map is extended to Hochschild homology and there is in fact a commutative diagram

$$\begin{array}{ccc} \mathrm{HC}_\bullet(A) & \xrightarrow{B} & \mathrm{HH}_{\bullet+1}(A) \\ \downarrow \mathrm{Tr} & & \downarrow \mathrm{Tr} \\ \mathrm{H}_\bullet(\mathrm{DRep}_V(A)^{\mathrm{GL}(V)}) & \xrightarrow{B_V} & \mathrm{H}_\bullet(\Omega^1(\mathrm{DRep}_V(A)^{\mathrm{GL}(V)})), \end{array} \quad (1.3)$$

where B in the upper line is the Connes differential, B_V in the bottom line is the de Rham differential and $(-)^{\mathrm{GL}(V)}$ means the $\mathrm{GL}(V)$ -invariant space.

Now recall that in classical Poisson geometry, if a manifold M has a Poisson structure, then the space of differential forms $\Omega^*(M)$ on M is a Lie module over the space of functions $\Omega^0(M)$, where the Lie action is given as follows: for $f \in \Omega^0(M)$, $\omega \in \Omega^*(M)$,

$$[f, \omega] := \mathcal{L}_{X_f} \omega,$$

where \mathcal{L}_{X_f} is the Lie derivative of the vector field X_f associated to f . In derived non-commutative geometry, we have similar results, and claim that

Theorem 1.3 (to appear in [9]). *Let A be a Koszul Calabi–Yau algebra. Then the diagram (1.3) is a commutative diagram of Lie module morphisms.*

To prove this theorem, we shall have to discuss the Lie derivative on the derived representation schemes, which is very much involved. We decide to give a complete proof in a separate paper; however, for reader’s convenience we give enough backgrounds in §7.

1.5. Relations to the Gerstenhaber and the de Völcsey–Van den Bergh brackets.

The construction of the double Poisson bracket on $\Omega_\infty(\tilde{A}^i)$ is also inspired by the Kontsevich bracket in non-commutative symplectic geometry (see Kontsevich [33] as well as Ginzburg [22] and Van den Bergh [45] for further discussions). It is direct to check that if $\{\{-, -\}\}$ is a double Poisson bracket on R then

$$\{-, -\} = \mu \circ \{\{-, -\}\} : R \times R \rightarrow R$$

defines a Leibniz–Loday bracket on R , where μ is the multiplication. For more details of the Leibniz–Loday bracket, see §3. It has been interesting for a long time to explore the relationships among the brackets such as the Gerstenhaber bracket, the Leibniz–Loday bracket and the non-commutative Poisson bracket of Kontsevich and Van den Bergh, etc. In this paper we also study this problem in the case of Koszul Calabi–Yau algebras with some detail.

First, we observe that there is a natural quasi-isomorphism (a version of non-commutative Poincaré duality originally due to Tradler [44])

$$\Phi : \text{CH}_\bullet(A^i) \longrightarrow \text{CH}^\bullet(A^1)$$

from the Hochschild chain complex of A^i to the Hochschild cochain complex of A^1 . Second, we observe that $\Omega(A^i)$ naturally embeds into $\text{CH}_\bullet(A^i)$ as chain complexes. By combining these two observations, we obtain the following theorem:

Theorem 1.4. *Let A be an N -Koszul d -Calabi–Yau algebra. Denote by A^i its Koszul dual coalgebra. Denote by $\{-, -\}_{\text{DNCP}}$ the Leibniz–Loday bracket associated to the derived non-commutative Poisson structure on A and by $\{-, -\}_{\text{G}}$ the Gerstenhaber bracket on $\text{CH}^\bullet(A^1)$, respectively. Then we have*

$$\Phi \circ B\{u, v\}_{\text{DNCP}} = \{\Phi \circ B(u), \Phi \circ B(v)\}_{\text{G}},$$

for any $u, v \in \Omega_\infty(A^i)$, where B is the Connes cyclic operator.

A classical result of Keller [31, Theorem 3.5] says that if A is Koszul, then the Hochschild cohomology of A and of $A^!$ are isomorphic as Gerstenhaber algebras. His theorem holds for N -Koszul algebras, too. Therefore, Theorem 1.4 implies that $\Phi \circ B$, composed with the isomorphism of Keller, maps the Leibniz–Loday bracket of A to the Gerstenhaber bracket on the Hochschild cohomology of A itself.

The above result allows us to give an explicit formula for the Lie bracket on the cyclic homology and the Lie module structure on the Hochschild homology. First, we recall that $(\mathrm{HH}^\bullet(A), \cup)$ is a graded commutative algebra and $(\mathrm{HH}_\bullet(A), \cap)$ is a graded module

$$\cap : \mathrm{HH}_m(A) \otimes \mathrm{HH}^n(A) \rightarrow \mathrm{HH}_{m-n}(A), \alpha \otimes f \mapsto \alpha \cap f$$

and we denote $\iota_f(\alpha) := \alpha \cap f$. Now if A is d -Calabi–Yau, then there in fact exists an element $\omega \in \mathrm{HH}_d(A)$ such that the cap product with ω :

$$\begin{aligned} \Psi : \mathrm{HH}^\bullet(A) &\longrightarrow \mathrm{HH}_{d-\bullet}(A) \\ f &\longmapsto \iota_f \omega \end{aligned}$$

is an isomorphism, which is called the *non-commutative Van den Bergh–Poincaré duality* for A . We can show

Corollary 1.5. *The Lie bracket of Theorem 1.1 on $\mathrm{HC}_\bullet(A)$ is given by*

$$\{\alpha, \beta\}_{\mathrm{DNCP}} = (-1)^{(d-|\alpha|-1)} \iota_{\Psi^{-1}(B(\alpha))} B(\beta) \quad (1.4)$$

for $\alpha, \beta \in \mathrm{HC}_\bullet(A)$, where $B : \mathrm{HC}_\bullet(A) \rightarrow \mathrm{HH}_{\bullet+1}(A)$ is the Connes operator. The Lie module structure on $\mathrm{HH}_\bullet(A)$ is also given by the bracket (1.4) for $\beta \in \mathrm{HH}_\bullet(A)$.

Another consequence of Theorem 1.4 is that it relates the Lie bracket $\{-, -\}_{\mathrm{DNCP}}$ of Theorem 1.1 on $\mathrm{HC}_\bullet(A)$ with the Lie bracket $\{-, -\}_{\mathrm{dVV}}$ of de Völcsey–Van den Bergh on the negative cyclic homology $\mathrm{HC}_\bullet^-(A)$ introduced in [17]. Let us briefly recall their construction. Let $\mathrm{CC}_\bullet^-(A)$ be the negative cyclic complex and let $\pi : \mathrm{CC}_\bullet^-(A) \rightarrow \mathrm{CH}_\bullet(A)$ be the natural projection in to the last two columns (see e.g. [35, Section 5.1.4.1]). Next, let \cup be the cup product on $\mathrm{HH}^\bullet(A)$. Then

$$\begin{aligned} \{-, -\}_{\mathrm{dVV}} : \mathrm{HC}_n^-(A) \times \mathrm{HC}_m^-(A) &\longrightarrow \mathrm{HC}_{n+m-d+1}^-(A) \\ (\eta_1, \eta_2) &\longmapsto (-1)^{|\eta_1|+d} B \circ \Psi(\Psi^{-1}(\pi(\eta_1)) \cup \Psi^{-1}(\pi(\eta_2))) \end{aligned} \quad (1.5)$$

where $B : \mathrm{HH}_q(A) \rightarrow \mathrm{HC}_{q+1}^-(A)$ is the map induced from the Connes operator and Ψ is the non-commutative Poincaré duality. From the following commutative diagram (see [35, Proposition 5.1.5])

$$\begin{array}{ccc} \mathrm{HC}_\bullet(A) & \xrightarrow{B} & \mathrm{HC}_{\bullet+1}^-(A) \\ \downarrow id & & \downarrow \pi \\ \mathrm{HC}_\bullet(A) & \xrightarrow{B} & \mathrm{HH}_{\bullet+1}(A), \end{array} \quad (1.6)$$

we have

Theorem 1.6. *Let A be an N -Koszul d -Calabi–Yau algebra. Then*

$$B\{\alpha, \beta\}_{\text{DNCP}} = \{B(\alpha), B(\beta)\}_{d\text{VV}}$$

for any $\alpha, \beta \in \text{HC}_\bullet(A)$.

The rest of the paper is devoted to the proof of the above theorems. It is organized as follows: in §2 we recollect several basic notions such as the Hochschild and cyclic homology of algebras and coalgebras; in §3 we recall the definition of double Poisson algebras in the sense of Van den Bergh and their bimodules; in §4 we show that for a class of coalgebras there is a double Poisson bracket on their cobar construction; in §5 we continue to show that, for the cobar construction, when viewing it as a DG algebra, its non-commutative differential 1-forms admit a double Poisson bimodule structure; in §6 we show N -Koszul Calabi–Yau algebras have a derived non-commutative Poisson structure; in §7 we give a brief introduction to derived representation schemes and derived non-commutative Poisson structures; in §8 we prove the main theorems listed in §1; and in §9 we give by explicit formulas several brackets on the space of polynomials of several variables.

Acknowledgements. We would like to thank Yuri Berest for helpful communications and NSFC (No. 11271269) for partial support.

2. Hochschild and cyclic homologies of coalgebras

2.1. Bar and cobar constructions. Let k be a field of characteristic zero. An associative k -algebra A is said to be *augmented* if there is an algebra homomorphism $\epsilon : A \rightarrow k$, called the *augmentation map*. In particular, A is canonically isomorphic, as a vector space, to $k \oplus \bar{A}$, where $\bar{A} = \text{Ker}(\epsilon)$.

The *bar construction* of an augmented algebra A , denoted by $\mathbf{B}(A)$, is a DG coalgebra defined as follows. First, recall that the *suspension* of a graded vector space V is the graded vector space sV such that $(sV)_i = V_{i-1}$. Similarly, the *desuspension* is $s^{-1}V$ such that $(s^{-1}V)_i = V_{i+1}$. As a coalgebra $\mathbf{B}(A)$ is the tensor coalgebra $T(s\bar{A})$ of the underlying vector space of \bar{A} located in degree one. The coproduct is

$$\Delta(a_1, \dots, a_n) = \sum_{i=0}^n (a_1, \dots, a_i) \otimes (a_{i+1}, \dots, a_n)$$

while the differential is

$$b'(a_1, \dots, a_n) = \sum_{i=1}^{n-1} (-1)^{i-1} (a_1, \dots, a_i \cdot a_{i+1}, \dots, a_n).$$

The counit $\eta : \mathbf{B}(A) \rightarrow k$ is the projection onto $A^{\otimes 0} = k$ and we will denote by $\bar{\mathbf{B}}(A)$ the $\text{Ker}(\eta)$.

A coassociative coalgebra (C, Δ) is a *coaugmented* coalgebra if there is coalgebra homomorphism $u : k \rightarrow C$. Then C can be identified as a vector space to $k \oplus \bar{C}$, where \bar{C} is the $\text{Coker}(u)$. Define the *reduced coproduct* $\bar{\Delta} : \bar{C} \rightarrow \bar{C} \otimes \bar{C}$ as the composite

$$\bar{C} \xrightarrow{i} C \xrightarrow{\Delta} C \otimes C \xrightarrow{\pi \otimes \pi} \bar{C} \otimes \bar{C}$$

where i and π are canonical inclusion and projection. Then *cobar construction* of C is a DG algebra $\mathbf{\Omega}(C)$ ¹ defined as follows. As an algebra it is the tensor algebra $T(s^{-1}\bar{C})$ and the differential is

$$b'(c_1, \dots, c_n) = \sum_{i=1}^n \sum_{(c_i)} (-1)^{i-1} (c_1, \dots, c'_i, c''_i, \dots, c_n)$$

where $\bar{\Delta}(c_i) = \sum_{(c_i)} (c'_i, c''_i)$. The unit map $\epsilon : k \rightarrow \mathbf{\Omega}(C)$ is the inclusion into $\bar{C}^{\otimes 0} = k$, and we will write $\bar{\mathbf{\Omega}}(C)$ for $\text{Coker}(\epsilon)$, which is the *reduced cobar construction* of C .

2.2. The cyclic bicomplex. Let C be a coalgebra over k . We write $\Delta(c) = \sum_{(c)} c' \otimes c''$ for the coproduct in C . Then we consider the following double complex which is obtained by reversing the arrows in the standard (Tsygan) double complex of an algebra:

$$\begin{array}{ccccccc}
 & & \uparrow & & \uparrow & & \uparrow \\
 & \xrightarrow{N} & C^{\otimes 3} & \xrightarrow{1-T} & C^{\otimes 3} & \xrightarrow{N} & C^{\otimes 3} & \xrightarrow{1-T} & \\
 & & \uparrow & & \uparrow & & \uparrow & & \\
 & \xrightarrow{N} & C^{\otimes 2} & \xrightarrow{1-T} & C^{\otimes 2} & \xrightarrow{N} & C^{\otimes 2} & \xrightarrow{1-T} & \\
 & & \uparrow & & \uparrow & & \uparrow & & \\
 & \xrightarrow{N} & C & \xrightarrow{1-T} & C & \xrightarrow{N} & C & \xrightarrow{1-T} & \\
 & & \uparrow & & \uparrow & & \uparrow & & \\
 & & 0 & & 0 & & 0 & &
 \end{array}$$

¹In this paper the cobar construction is denoted by the bold face $\mathbf{\Omega}$ while the standard Ω means the non-commutative differential forms.

This double complex is 2-periodic in horizontal direction, with operators b', b, T, N given by

$$\begin{aligned} b'(c_1, \dots, c_n) &:= \sum_{i=1}^n \sum_{(c_i)} (-1)^{i-1} (c_1, \dots, c'_i, c''_i, \dots, c_n), \\ b(c_1, \dots, c_n) &:= b'(c_1, \dots, c_n) + \sum (-1)^n (c''_1, c_2, \dots, c_n, c'_1), \\ T(c_1, \dots, c_n) &:= (-1)^{n-1} (c_2, \dots, c_n, c_1), \\ N &:= \sum_{i=0}^{n-1} T^i. \end{aligned}$$

The b -column is called the Hochschild chain complex $\text{CH}_\bullet(C)$ of C : it defines the Hochschild homology $\text{HH}_\bullet(C)$. The b' -column is the reduced cobar construction of co-augmented coalgebra $\tilde{C} = k \oplus C$ shifted by degree one; here \tilde{C} is the co-augmentation of C , which, as a vector space, is $k \oplus C$, with k being the co-unit. The kernel of $1 - T$ from the b -complex to the b' -complex is called the cyclic complex $\text{CC}_\bullet(C)$: by definition, its homology is the cyclic homology HC_\bullet of C .

In practice, to compute the Hochschild and cyclic homologies, one usually considers the normalized Hochschild complex. For a co-augmented coalgebra C , the normalized Hochschild complex

$$\overline{\text{CH}}_n(C) := C \otimes (\tilde{C})^{\otimes n}$$

with the differential induced from b . Similarly to the algebra case, $\overline{\text{CH}}_\bullet(C)$ and $\text{CH}_\bullet(C)$ are quasi-isomorphic; however, $\overline{\text{CH}}_\bullet(C)$ may be viewed as a tensor product $C \otimes \Omega(C)$ with a twisted differential given by $id \otimes b' + \tau_L + \tau_R$, where b' is the differential in the cobar construction, and τ_L and τ_R are given by

$$\begin{aligned} \tau_L(c_0, c_1, \dots, c_n) &:= \sum_{(c_0)} (c'_0, c''_0, c_1, \dots, c_n), \\ \tau_R(c_0, c_1, \dots, c_n) &:= \sum_{(c_0)} (-1)^n (c''_0, c_1, \dots, c_n, c'_0). \end{aligned}$$

Under this identification, one sees that the cobar construction embeds into the Hochschild complex as complexes

$$\begin{aligned} \Omega(C) &\longrightarrow \text{CH}_\bullet(C) \\ (c_1, \dots, c_n) &\longmapsto 1 \otimes (c_1, \dots, c_n). \end{aligned} \tag{2.1}$$

We now recall some facts about the cyclic bicomplex from Quillen [41, Section 1.3]. Let A be an associative algebra. The commutator subspace of A is $[A, A]$ which is the image of $\mu - \mu\sigma : A \otimes A \rightarrow A$ where μ is the product in A and σ is the switching operator, and the commutator quotient space is

$$A_{\text{q}} := A/[A, A] = \text{Coker}(\mu - \mu\sigma).$$

Dually, for a coassociative coalgebra C the cocommutator subspace of C is

$$C^{\natural} := \text{Ker}\{\Delta - \sigma\Delta : C \rightarrow C \otimes C\}.$$

Let $\tilde{A} = k \oplus A$ be the augmentation of A (recall that the augmentation of a not-necessarily unital algebra A is the algebra $\tilde{A} = k \oplus A$ where k plays the role of the unit). Let $\tilde{\mathbf{B}}$ be the *reduced bar construction* of \tilde{A} . The following lemma is [41, Lemma 1.2]:

Lemma 2.1. *The space $\tilde{\mathbf{B}}_n^{\natural}$ is the kernel of $1 - T$ acting on $A^{\otimes n}$. Hence, we have the isomorphism of complexes $\text{CC}_{\bullet-1}(A) \cong \tilde{\mathbf{B}}^{\natural}$.*

Remark 2.2. For the convenience of later discussions, from now on we shall shift the degrees of $\text{CC}_{\bullet}(A)$ up by one, and just write the above identity and alike as $\text{CC}_{\bullet}(A) \cong \tilde{\mathbf{B}}^{\natural}$.

Dually, the space $(\mathbf{\Omega}(\tilde{C})_{\natural})_n$ is the cokernel of $1 - T$ acting on $C^{\otimes n}$. Thus, by the isomorphisms

$$\begin{aligned} \text{CC}_{\bullet}(A) &= \text{Coker}(1 - T) \cong \text{Ker}(1 - T), \\ \text{CC}_{\bullet}(C) &= \text{Ker}(1 - T) \cong \text{Coker}(1 - T), \end{aligned} \tag{2.2}$$

we have the following lemma

Lemma 2.3. *As complexes of k -vector spaces $\overline{\mathbf{\Omega}}(\tilde{C})_{\natural} \cong \text{CC}_{\bullet}(C)$. Hence,*

$$\text{HC}_{\bullet}(C) \cong \text{H}_{\bullet}[\overline{\mathbf{\Omega}}(\tilde{C})_{\natural}].$$

There is an efficient way to compute the cyclic homology. Let us recall that (cf. [35]) for a unital and augmented algebra $A = k \oplus \tilde{A}$, its reduced cyclic chain complex

$$\overline{\text{CC}}_{\bullet}(A) := \text{Coker}\{1 - T : \tilde{C}^{\otimes n} \rightarrow \tilde{C}^{\otimes n}\}.$$

The associated homology is the reduced cyclic homology of A , and is denoted by $\overline{\text{HC}}_{\bullet}(A)$, and there is in fact a decomposition

$$\text{HC}_{\bullet}(A) \cong \text{HC}_{\bullet}(k) \oplus \overline{\text{HC}}_{\bullet}(A).$$

The following is originally due to Feigin and Tsygan, and is now well-known (see e.g. [3, Proposition 4.2]):

Proposition 2.4. *Let $R \xrightarrow{\cong} A$ be a quasi-free resolution of an algebra A . Then there is quasi-isomorphism $\overline{\text{CC}}_{\bullet}(A) \rightarrow \overline{R}_{\natural}$ inducing isomorphism of homologies $\overline{\text{HC}}_{\bullet}(A) \simeq \text{H}_{\bullet}(\overline{R}_{\natural})$, where $\overline{\text{CC}}_{\bullet}(-)$ and $\overline{\text{HC}}_{\bullet}(-)$ are the reduced cyclic chain complex and reduced cyclic homology respectively.*

Thus, combining the above two results, one obtains

Corollary 2.5. *Let C be a coaugmented coalgebra such that $\Omega(C) \xrightarrow{\cong} A$ is a quasi-free resolution of an algebra A . Then $\overline{\mathrm{HC}}_{\bullet}(A) \cong \overline{\mathrm{HC}}_{\bullet}(C)$. Moreover, if A is an augmented algebra then $\mathrm{HC}_{\bullet}(A) \cong \mathrm{HC}_{\bullet}(C)$.*

Proof. First, by Proposition 2.4, $\overline{\mathrm{HC}}_{\bullet}(A) \simeq \mathrm{H}_{\bullet}[\overline{\Omega}(C)_{\mathfrak{h}}]$. The later by definition is isomorphic to $\mathrm{HC}_{\bullet}(\overline{C})$. Since C is coaugmented, using arguments similar to that in [35, Proposition 2.2.16], we obtain $\mathrm{HC}_{\bullet}(\overline{C}) \cong \overline{\mathrm{HC}}_{\bullet}(C)$.

The second part follows from the first one, since in this case $\mathrm{HC}_{\bullet}(A) = \mathrm{HC}_{\bullet}(k) \oplus \overline{\mathrm{HC}}_{\bullet}(A)$ and $\mathrm{HC}_{\bullet}(C) = \mathrm{HC}_{\bullet}(k) \oplus \overline{\mathrm{HC}}_{\bullet}(C)$. \square

2.3. Non-commutative differential forms on algebras and coalgebras. The most of the material in this section is either taken from [15, 41] or merely stating the dual version of those results.

For a DG algebra R , we denote by Ω_R^1 the kernel of the multiplication map $R \otimes R \rightarrow R$. It is a DG bimodule induced from outer bimodule structure on $R \otimes R$ and it represents $\underline{\mathrm{Der}}(R, -)$, the complex of k -linear graded derivations. Thus, for any R -bimodule M , we have

$$\underline{\mathrm{Der}}(R, M) \cong \underline{\mathrm{Hom}}_{R^e}(\Omega_R^1, M).$$

If $M = \Omega_R^1$, the derivation $\partial : R \rightarrow \Omega_R^1$ then corresponding to the identity map under this isomorphism, is a universal derivation.

Let V be a k -linear vector space and R be the free algebra $\bigoplus_{n \geq 0} V^{\otimes n}$. Then Ω_R^1 can be identified with $R \otimes V \otimes R$. Indeed, there is a bijection $I : R \otimes V \otimes R \rightarrow \Omega_R^1$ (see [41, Example 3.10])

$$\begin{aligned} I\{(v_1 \cdots v_{p-1}) \otimes v_p \otimes (v_{p+1} \cdots v_m)\} \\ = (v_1 \cdots v_p) \otimes (v_{p+1} \cdots v_m) - (v_1 \cdots v_{p-1}) \otimes (v_p \cdots v_m) \end{aligned} \quad (2.3)$$

and the universal derivation $\partial : R \rightarrow \Omega_R^1$ is given by

$$\partial(v_1 v_2 \cdots v_m) = \sum_{i=1}^m (v_1 \cdots v_{i-1}) \otimes v_i \otimes (v_{i+1} \cdots v_m).$$

A simple computation shows

$$I(\partial(v_1 v_2 \cdots v_m)) = (v_1 v_2 \cdots v_m) \otimes 1 - 1 \otimes (v_1 v_2 \cdots v_m). \quad (2.4)$$

Next, for an R -bimodule M , we define $M_{\mathfrak{h}} := M/[R, M]$. Then $\Omega_{R, \mathfrak{h}}^1$ is isomorphic to $R \otimes V$ and the map $\bar{\partial} : R \rightarrow \Omega_{R, \mathfrak{h}}^1$ induced by ∂ is defined as

$$\bar{\partial}(v_1 v_2 \cdots v_m) = \sum_{i=1}^m (-1)^{(|v_1| + \cdots + |v_i|)(|v_{i+1}| + \cdots + |v_m|)} (v_{i+1} \cdots v_m v_1 \cdots v_{i-1}) \otimes v_i. \quad (2.5)$$

One can also define a graded map $\beta : \Omega^1(R)_{\mathfrak{h}} \rightarrow R$ given by

$$\begin{aligned} & \beta((v_1 \cdots v_m) \otimes v_{m+1}) \\ &= (v_1 \cdots v_m v_{m+1}) - (-1)^{|v_{m+1}|(|v_1| + \cdots + |v_m|)} (v_{m+1} v_1 \cdots v_m). \end{aligned} \quad (2.6)$$

It is easy to check that $\beta \bar{\partial} = \bar{\partial} \beta = 0$ (see [41, Proposition 3.8]).

The above maps give a commutative diagram

$$\begin{array}{ccc} \Omega_R^1 \xrightarrow{C} & R \otimes R & \\ \mathfrak{h} \downarrow & & \downarrow \mu \circ \sigma \\ \Omega_{R, \mathfrak{h}}^1 & \xrightarrow{\beta} & R \end{array} \quad (2.7)$$

where $\sigma : R \otimes R \rightarrow R \otimes R, r \otimes q \mapsto (-1)^{|r||q|} q \otimes r$ the graded switching operator.

Now let (C, Δ) be a coalgebra. Then M is a *bicomodule* over C is a vector space equipped with left and right coproducts $\Delta_l : M \rightarrow C \otimes M, \Delta_r : M \rightarrow M \otimes C$ defining left and right comodule structures which commute: $(\Delta_l \otimes 1) \circ \Delta_r = (1 \otimes \Delta_r) \circ \Delta_l$. The cocommutator subspace of M is

$$M^{\mathfrak{h}} := \text{Ker}\{\Delta_l - \sigma \Delta_r : M \rightarrow C \otimes M\}.$$

We let Ω^C , the *non-commutative differentials one-forms* on C , be the bicomodule $\text{Coker}(\Delta)$. The cocommutator subspace we denote by $\Omega^{C, \mathfrak{h}}$. We can introduce the maps $\beta : C \rightarrow \Omega^{C, \mathfrak{h}}$ and $\bar{\partial} : \Omega^{C, \mathfrak{h}} \rightarrow C$.

Recall that \mathbf{B} is the bar construction of \tilde{A} . The following is proved in [41, Theorem 4]:

Theorem 2.6. *The complex $\Omega^{\mathbf{B}, \mathfrak{h}}$ is canonically isomorphic to $\text{CH}_\bullet(A)$, the Hochschild complex of A . Under this identification $\beta = 1 - T$ and $\bar{\partial} = N$.*

Dually we can show

Theorem 2.7. *Let C be a coalgebra and let $R = \Omega(\tilde{C})$, where $\tilde{C} := k \oplus C$. Then the complex $\Omega_{R, \mathfrak{h}}^1$ is isomorphic to $\text{CH}_\bullet(C)$. Under this identification $\beta = 1 - T$ and $\bar{\partial} = N$.*

It has been pointed out in [41, Remark 5.14] that the cyclic bicomplex for the algebra A can be identified with the periodic sequence of complexes

$$\xrightarrow{\bar{\partial}} \bar{\mathbf{B}} \xrightarrow{-\beta} \Omega^{\mathbf{B}, \mathfrak{h}} \xrightarrow{\bar{\partial}} \bar{\mathbf{B}} \xrightarrow{-\beta} \xrightarrow{\quad} \quad (2.8)$$

Similarly, in view of Lemma 2.3 and Theorem 2.7 one has

Proposition 2.8. *The cyclic bicomplex for the coalgebra C can be identified with*

$$\xrightarrow{\bar{\partial}} \Omega_{R, \mathfrak{h}}^1 \xrightarrow{-\beta} \bar{\Omega}(\tilde{C}) \xrightarrow{\bar{\partial}} \Omega_{R, \mathfrak{h}}^1 \xrightarrow{-\beta} \xrightarrow{\quad} \quad (2.9)$$

The following result is essentially established in [30] (see also [10]):

Lemma 2.9. *Let C be a coaugmented coalgebra such that $\Omega(C) \xrightarrow{\cong} A$ is a quasi-free resolution of an algebra A . Then $\mathrm{HH}_\bullet(A) \cong \mathrm{HH}_\bullet(C)$.*

Combining the above lemma with Theorem 2.7 one obtains

Corollary 2.10. *If $\Omega(C) \xrightarrow{\cong} A$ is a quasi-free resolution of A and $R = \Omega(\tilde{C})$ then*

$$\mathrm{HH}_\bullet(A) \cong \mathrm{H}_\bullet(\Omega_{R, \natural}^1). \tag{2.10}$$

2.4. The A_∞ algebra and coalgebra case. The advantage of rephrasing the Hochschild and cyclic homology groups in the proceeding subsections is that they can be easily generalized to the A_∞ algebra and coalgebra case.

Definition 2.11 (A_∞ algebra). Let A be a graded vector space. An A_∞ algebra structure on A is a sequence of linear operators

$$m_n : A^{\otimes n} \longrightarrow A, \quad n = 1, 2, \dots \tag{2.11}$$

of degree $n - 2$ such that

$$\sum_{r+s+t=n} (-1)^{r+st} m_{r+1+t}(\mathrm{id}^{\otimes r} \otimes m_s \otimes \mathrm{id}^{\otimes t}) = 0. \tag{2.12}$$

An A_∞ algebra A is called *unital*, if there exists a map $k \rightarrow A$ which maps 1 to $\mathbf{1}$, such that

$$\begin{aligned} \mu_1(\mathbf{1}) &= 0, \quad \mu_2(a, \mathbf{1}) = \mu_2(\mathbf{1}, a) = a, \\ \text{and } \mu_n(a_1, \dots, a_{i-1}, \mathbf{1}, a_{i+1}, a_n) &= 0, \quad \text{for all } n \geq 3, \end{aligned}$$

where μ_1, μ_2, \dots are the A_∞ operators. It is called *unital and augmented* if furthermore there is a map $A \rightarrow k$ such that the composition

$$k \rightarrow A \rightarrow k$$

is the identity. In this case, A is decomposed into direct sum $k \cdot \mathbf{1} \oplus \bar{A}$ of A_∞ algebras.

Definition 2.12 (A_∞ coalgebras). Let C be a graded vector space. An A_∞ coalgebra structure on C is a sequence of linear operators

$$\Delta_n : C \longrightarrow C^{\otimes n}, \quad n = 1, 2, \dots \tag{2.13}$$

of degree $n - 2$ such that

$$\sum_{r+s+t=n} (-1)^{r+st} (\mathrm{id}^{\otimes r} \otimes \Delta_s \otimes \mathrm{id}^{\otimes t}) \Delta_{r+1+t} = 0. \tag{2.14}$$

An A_∞ coalgebra C is said to be *co-unital* if there is a map $\eta : C \rightarrow k$ such that for all $c \in C$,

$$\begin{aligned} \eta \circ \Delta_1(c) &= 0, \\ (\eta \otimes \text{id} + \text{id} \otimes \eta) \circ \Delta_2(c) &= 1 \otimes c + c \otimes 1, \\ \text{and } \left(\sum_{j+k+1=n} \text{id}^{\otimes j} \otimes \eta \otimes \text{id}^{\otimes k} \right) \circ \Delta_n(c) &= 0, \quad \text{for all } n \geq 3. \end{aligned}$$

It is called *co-unital and co-augmented* if furthermore there is a map $k \rightarrow C$ such that the composition $k \rightarrow C \rightarrow k$ is the identity.

Assumption 2.13. *From now on we shall assume² that for an A_∞ -coalgebra C , in each grading C_i is finite dimensional (in literature C is called locally finite dimensional), and that all but finitely many $\Delta_n(v)$ vanish, for each $v \in C$. In particular, the Koszul dual A_∞ coalgebra A^\natural of an N -Koszul algebra (to be studied later), which only has Δ_2 and Δ_N , satisfies this assumption.*

Definition 2.14 (Bar and cobar constructions). Suppose A is a unital and augmented A_∞ algebra, and C is a co-unital and co-augmented A_∞ coalgebra (satisfying Assumption 2.13 above). The *bar construction* of A , denoted by $\mathbf{B}_\infty(A)$, is the quasi-free graded coalgebra $T(s\bar{A})$ generated by $s\bar{A}$, with differential $d = d_1 + d_2 + \dots$, where d_n is defined on the co-generators by

$$\begin{aligned} d_n : (s\bar{A})^{\otimes n} &\longrightarrow s\bar{A} \subset T(s\bar{A}) \\ (s\bar{a}_1, \dots, s\bar{a}_n) &\longmapsto (-1)^{(n-1)|a_1| + (n-2)|a_2| + \dots + |a_{n-1}|} s \circ \pi \circ \mu_n(a_1, \dots, a_n) \end{aligned}$$

and is extended to $T(s\bar{A})$ by derivation, where $\pi : A \rightarrow \bar{A}$ is the projection.

Similarly, the *cobar construction* of C , denoted by $\mathbf{\Omega}_\infty(C)$, is the quasi-free graded algebra $T(s^{-1}\bar{C})$ generated by $s^{-1}\bar{C}$ with differential d defined on the generators by

$$\begin{aligned} s^{-1}\bar{c} &\longrightarrow T(s^{-1}\bar{C}) \\ s^{-1}\bar{c} &\longmapsto (s^{-1} \circ \pi \circ \Delta_1 + (s^{-1})^{\otimes 2} \circ (\pi^{\otimes 2}) \circ \Delta_2 + \dots)(c), \end{aligned}$$

which extends to $T(s^{-1}(A))$ by derivation, where in the above expression, $\pi : C \rightarrow \bar{C}$ is the projection.

Definition 2.15 (Hochschild and cyclic homology of A_∞ algebras and coalgebras).

(1) For an A_∞ algebra A , define its *Hochschild homology* $\text{HH}_\bullet(A)$ to be $\text{H}_\bullet(\Omega^{\mathbf{B}_\infty(\bar{A}), \natural})$ as in Theorem 2.6, and its *cyclic homology* $\text{HC}_\bullet(A)$ to be $\text{H}_\bullet(\bar{\mathbf{B}}_\infty(\bar{A})^\natural)$ as in Lemma 2.1.

²This assumption is also used by Prouté in his definition of A_∞ coalgebras; see [39, Définition 3.2].

(2) For an A_∞ coalgebra C , define its *Hochschild homology* $\mathrm{HH}_\bullet(C)$ to be $\mathrm{H}_\bullet(\Omega^1_{\mathfrak{Q}_\infty(\tilde{C}), \mathfrak{H}})$ as in Theorem 2.7, and its *cyclic homology* $\mathrm{HC}_\bullet(C)$ to be $\mathrm{H}_\bullet(\overline{\mathfrak{Q}}_\infty(\tilde{C})_{\mathfrak{H}})$ as in Lemma 2.3.

In the above definition, \tilde{A} and \tilde{C} are the augmentation of A and the co-augmentation of C , respectively, which are defined to be the same as for algebras and coalgebras.

Remark 2.16. As has been shown above, for DG algebras and coalgebras, the definitions given above coincide with the standard ones (cf. Loday [35]).

Proposition 2.17. *Throughout Lemma 2.1–Corollary 2.10, the statements remain true when the algebra A is replaced by an A_∞ algebra A and respectively the coalgebra C is replaced by an A_∞ coalgebra C .*

Proof. In the proofs of these statements, the only fact that is used is that $\mathbf{B}(\tilde{A})$ is a quasi-free DG coalgebra and $\mathfrak{Q}(\tilde{C})$ is a quasi-free DG algebra, which is also true for $\mathbf{B}_\infty(\tilde{A})$ and $\mathfrak{Q}_\infty(\tilde{C})$ respectively. \square

Finally, we show that for unital and augmented A_∞ algebras and coalgebras, their Hochschild chain complex is quasi-isomorphic to their *normalized Hochschild chain complex*.

Proposition 2.18. *Suppose A is a unital and augmented A_∞ algebra and C is a co-unital and co-augmented A_∞ coalgebra. The the following are quasi-isomorphic*

$$\mathrm{CH}_\bullet(A) \simeq \overline{\mathrm{CH}}_\bullet(A), \quad \mathrm{CH}_\bullet(C) \simeq \overline{\mathrm{CH}}_\bullet(C).$$

Proof. The same argument for algebras (see Loday [35, Proposition 1.6.5]) remains to hold for the A_∞ case. \square

As an application, we see that

$$\overline{\mathrm{CH}}_\bullet(A) \simeq A \otimes \mathbf{B}_\infty(A), \quad \overline{\mathrm{CH}}_\bullet(C) \simeq C \otimes \mathfrak{Q}_\infty(C),$$

with the differential properly defined. In particular, there is an embedding

$$\begin{aligned} \mathfrak{Q}_\infty(C) &\longrightarrow \overline{\mathrm{CH}}_\bullet(C) \\ (a_1, \dots, a_n) &\longmapsto (1, a_1, \dots, a_n) \end{aligned} \tag{2.15}$$

of chain complexes similar to the coalgebra case (compare with (2.1)).

3. Double Poisson algebras and bimodules

In this section we remind the definition of a double Poisson algebra A , and introduce the notion of a double Poisson bimodule M . After that, we discuss the construction of n -Poisson structures.

3.1. Double Poisson algebras.

Definition 3.1. Suppose A is a unital, associative algebra over a field k . A *double bracket* on A is a bilinear map $\{\{-, -\}\} : A \times A \rightarrow A \otimes A$ which satisfies

$$\{\{a, b\}\} = -\{\{b, a\}\}^\circ, \quad (3.1)$$

$$\{\{a, bc\}\} = b\{\{a, c\}\} + \{\{a, b\}\}c, \quad (3.2)$$

where $(u \otimes v)^\circ = v \otimes u$. Here the action of b and c is given via outer bimodule structure on $A \otimes A$, that is, $b(a_1 \otimes a_2)c := ba_1 \otimes a_2c$. We recall that the *inner* bimodule structure on $A \otimes A$ is given by

$$b * (a_1 \otimes a_2) * c := a_1c \otimes ba_2. \quad (3.3)$$

The formulas (3.1) and (3.2) imply that $\{\{-, -\}\}$ is a derivation on its first argument for the inner bimodule structure

$$\{\{ab, c\}\} = a * \{\{b, c\}\} + \{\{a, c\}\} * b. \quad (3.4)$$

Suppose that $\{\{-, -\}\}$ is a double bracket on A . For $a, b_1, \dots, b_n \in A$, let

$$\{\{a, b_1 \otimes \dots \otimes b_n\}\}_L := \{\{a, b_1\}\} \otimes b_2 \otimes \dots \otimes b_n,$$

and let

$$\sigma_s(b_1 \otimes \dots \otimes b_n) := b_{s^{-1}(1)} \otimes \dots \otimes b_{s^{-1}(n)},$$

where s is a permutation of $\{1, 2, \dots, n\}$. If furthermore A satisfies the following *double Jacobi identity*

$$\{\{a, \{\{b, c\}\}\}\}_L + \sigma_{(123)}\{\{b, \{\{c, a\}\}\}\}_L + \sigma_{(132)}\{\{c, \{\{a, b\}\}\}\}_L = 0, \quad (3.5)$$

then A is called a *double Poisson algebra*.

Let $\mu : A \otimes A \rightarrow A$ denote the multiplication on A , and let $\{-, -\} := \mu \circ \{\{-, -\}\} : A \otimes A \rightarrow A$. Then $\{-, -\}$ induces well-defined maps $A_{\mathfrak{h}} \times A \rightarrow A$ and $A_{\mathfrak{h}} \times A_{\mathfrak{h}} \rightarrow A_{\mathfrak{h}}$ (see [45, Lemma 2.4.1]). Furthermore, the latter bracket is anti-symmetric.

Definition 3.2. A left *Leibniz–Loday algebra*³ is a vector space L with a bilinear operation $[-, -]$ such that it satisfies

$$[a, [b, c]] = [[a, b], c] + [b, [a, c]].$$

From the definition one immediately sees that

Lemma 3.3. $(A, \{-, -\})$ is a left *Leibniz–Loday algebra*.

As a consequence of Lemma 3.3 and (3.1), we have

Corollary 3.4. If A is a double Poisson algebra, then $\{-, -\}$ makes $A_{\mathfrak{h}} = A/[A, A]$ into a Lie algebra and A into a Lie module over $A_{\mathfrak{h}}$.

Proof. See Van den Bergh [45, Lemmas 2.4.2 and 2.6.2]. \square

³This algebraic structure is introduced by Loday, which he calls *Leibniz algebra*, while some other authors, for example, Van den Bergh [45], call it *Loday algebra*; we here combine these two terminologies together.

3.2. Double Poisson bimodules.

Definition 3.5. Let A be a double Poisson algebra with the double bracket $\{\{-, -\}\}$ and let M be an A -bimodule. Then a *double Poisson bracket* on M is a bilinear product $\{\{-, -\}\}_M : A \times M \rightarrow (A \otimes M) \oplus (M \otimes A)$ such that the following axioms hold for all $a, b \in A$ and all $m, n \in M$:

- (i) $\{\{a, bm\}\}_M = \{\{a, b\}\}m + b\{\{a, m\}\}_M$,
 $\{\{a, mb\}\}_M = \{\{a, m\}\}_M b + m\{\{a, b\}\}_M$;
- (ii) $\{\{ab, m\}\}_M = a * \{\{b, m\}\}_M + \{\{a, m\}\}_M * b$.

Remark 3.6. Expressions $\{\{a, b\}\}m$ and $m\{\{a, b\}\}_M$ should be understood as follows. If $\{\{a, b\}\} = \{\{a, b\}'\} \otimes \{\{a, b\}''\}$, then $\{\{a, b\}\}m \in A \otimes M$ via the left action of $\{\{a, b\}'\}$ on m and $m\{\{a, b\}\}_M \in M \otimes A$ via the right action of $\{\{a, b\}''\}$ on m . Also, $*$ in $a * \{\{b, m\}\}_M$ and $\{\{a, m\}\}_M * b$ is the action of A on $A \otimes M$ and $M \otimes A$ as inner bimodules (compare to (3.3)).

Our next is to introduce the Jacobi identity for the double bracket $\{\{-, -\}\}_M$. For this we need to define the following expressions: $\{\{a, \{\{b, m\}\}_M\}\}_L$, $\{\{b, \{\{m, a\}\}_M\}\}_L$, $\{\{m, \{\{b, a\}\}\}\}_L$. First, we define

$$\{\{m, b\}\}_M := -(\{\{b, m\}\}_M)^\circ, \quad (3.6)$$

that is, if $\{\{b, m\}\}_M = (b_1 \otimes m_1) \oplus (m_2 \otimes b_2)$, then $\{\{m, b\}\}_M = -(b_2 \otimes m_2) \oplus (m_1 \otimes b_1)$. Then

$$\{\{a, \{\{b, m\}\}_M\}\}_L := (\{\{a, b_1\}\} \otimes m_1) \oplus (\{\{a, m_2\}\}_M \otimes b_2), \quad (3.7)$$

which is in $(A \otimes A \otimes M) \oplus (A \otimes M \otimes A)$. Using (3.6) and (3.7), we can define $\{\{b, \{\{m, a\}\}_M\}\}_L$. Finally, if $\{\{a, b\}\} = \{\{a, b\}'\} \otimes \{\{a, b\}''\}$, then

$$\{\{m, \{\{a, b\}\}\}\}_L := \{\{m, \{\{a, b\}'\}\}\}_M \otimes \{\{a, b\}''\} \in (A \otimes M \otimes A) \oplus (M \otimes A \otimes A).$$

Definition 3.7. Let A be a double Poisson algebra with the double bracket $\{\{-, -\}\}$ and let M be an A -bimodule with a double bracket $\{\{-, -\}\}_M$. Then we say that M is a *double Poisson A -bimodule* if

$$(iii) \quad \{\{a, \{\{b, m\}\}_M\}\}_L + \sigma_{(123)} \{\{b, \{\{m, a\}\}_M\}\}_L + \sigma_{(132)} \{\{m, \{\{b, a\}\}\}\}_L = 0, \quad (3.8)$$

for all $a, b \in A$ and $m, n \in M$.

Remark 3.8. It is clear that A itself a double Poisson A -bimodule.

Let $(M, \{\{-, -\}\}_M)$ be a double Poisson A -bimodule. Then we define

$$\{-, -\}_M = \mu_M \circ \{\{-, -\}\}_M : A \times M \rightarrow M,$$

where $\mu_M : (A \otimes M) \oplus (M \otimes A) \rightarrow M$ is the bimodule action map. And one can prove the following.

Proposition 3.9. $\{-, -\}_M$ induces well-defined maps $A_{\natural} \times M \rightarrow M$ and $A_{\natural} \times M_{\natural} \rightarrow M_{\natural}$.

Proof. (1) Let $a, b \in A$ and $m \in M$. Then

$$\begin{aligned} \{ab - ba, m\}_M &= \mu_M \{\{ab, m\}\}_M - \mu_M \{\{ba, m\}\}_M \\ &= \mu_M (a * \{\{b, m\}\}_M + \{\{a, m\}\}_M * b) \\ &\quad - \mu_M (b * \{\{a, m\}\}_M + \{\{b, m\}\}_M * a) \\ &= \{\{b, m\}\}'_M a \{\{b, m\}\}''_M + \{\{a, m\}\}'_M b \{\{a, m\}\}''_M \\ &\quad - \{\{a, m\}\}'_M b \{\{a, m\}\}''_M - \{\{b, m\}\}'_M a \{\{b, m\}\}''_M = 0. \end{aligned}$$

(2) We let

$$\{-, -\}_{M, \natural} := \natural \circ \{-, -\}_M : A \times M \rightarrow M_{\natural}, \quad (3.9)$$

where $\natural : M \rightarrow M_{\natural}$ is the projection map. Then

$$\{a, cm - mc\}_{M, \natural} = \natural(\{a, c\}m - m\{a, c\}) + \natural(c\{a, m\}_M - \{a, m\}_M c) = 0,$$

and hence by (1) we have $\{-, -\}_{M, \natural} : A_{\natural} \times M_{\natural} \rightarrow M_{\natural}$. \square

Next, we establish the following statement

Proposition 3.10. The brackets $\{-, -\}_M$ and $\{-, -\}_{M, \natural}$ define on M and M_{\natural} respectively Lie module structures over the Lie algebra A_{\natural} .

Proof. We need to show that

$$\{\{a, b\}, m\}_M = \{a, \{b, m\}_M\}_M - \{b, \{a, m\}_M\}_M. \quad (3.10)$$

Indeed, we have

$$\begin{aligned} \{\{a, b\}, m\}_M &= \mu_M [\{\{\{a, b\}\}' \cdot \{\{a, b\}\}''\}_M] \\ &= \mu_M [\{\{a, b\}\}' * \{\{\{a, b\}\}''\}_M + \{\{\{a, b\}\}'\}_M * \{\{a, b\}\}''] \\ &= \mu_M [(1 \otimes \mu_M) \circ \sigma_{(132)} \{\{m, \{\{b, a\}\}\}_L \\ &\quad - (\mu_M \otimes 1) \circ \sigma_{(132)} \{\{m, \{\{a, b\}\}\}_L], \\ \{a, \{b, m\}_M\}_M &= \mu_M [\{\{a, \{\{b, m\}\}' \cdot \{\{b, m\}\}''\}\}_M] \\ &= \mu_M [\{\{a, \{\{b, m\}\}'\}\} \cdot \{\{b, m\}\}'' + \{\{b, m\}\}' \cdot \{\{a, \{\{b, m\}\}''\}\}] \\ &= \mu_M [(\mu_M \otimes 1) \{\{a, \{\{b, m\}\}\}\}_L \\ &\quad - (1 \otimes \mu_M) \circ \sigma_{(123)} \{\{a, \{\{m, b\}\}\}_L], \\ \{b, \{a, m\}_M\}_M &= \mu_M [\{\{b, \{\{a, m\}\}' \cdot \{\{a, m\}\}''\}\}_M] \\ &= \mu_M [\{\{b, \{\{a, m\}\}'\}\} \cdot \{\{a, m\}\}'' + \{\{a, m\}\}' \cdot \{\{b, \{\{a, m\}\}''\}\}] \\ &= \mu_M [(1 \otimes \mu_M) \{\{b, \{\{a, m\}\}\}\}_L \\ &\quad - (\mu_M \otimes 1) \circ \sigma_{(123)} \{\{b, \{\{m, a\}\}\}\}_L]. \end{aligned}$$

Using these identities and (3.8) we obtain (3.10). \square

3.3. Double n -Poisson structures. One can easily generalize the above notions to the n -graded case. Indeed, let $A = \bigoplus_{i \in \mathbb{Z}} A_i$ be a \mathbb{Z} -graded vector space. We denote by $A[n]$ the graded vector space with degree shifted by n , explicitly, $A[n] = \bigoplus (A[n])_i$ with $(A[n])_i = A_{i-n}$. Then a *double bracket* of degree n on A is a bilinear map $\{\{-, -\}\} : A \times A \rightarrow (A \otimes A)[-n]$ satisfying

$$\{\{a, b\}\} = -(-1)^{(|a|+n)(|b|+n)} \{\{b, a\}\}^\circ, \tag{3.11}$$

$$\{\{a, b \cdot c\}\} = \{\{a, b\}\} \cdot c + (-1)^{|b|(|a|+n)} b \cdot \{\{a, c\}\}. \tag{3.12}$$

The pair $(A, \{\{-, -\}\})$ is a *double n -Poisson algebra* if in addition it satisfies the n -graded version of the Jacobi identity

$$\begin{aligned} \{\{a, \{\{b, c\}\}\}\}_L + (-1)^{(|a|+n)(|b|+|c|)} \sigma_{(123)} \{\{b, \{\{c, a\}\}\}\}_L \\ + (-1)^{(|c|+n)(|a|+|b|)} \sigma_{(132)} \{\{c, \{\{a, b\}\}\}\}_L = 0. \end{aligned}$$

Similarly, we can define the notion of a double n -Poisson bimodule and n -graded version of results of previous two sections can be summarized in the following proposition.

Proposition 3.11. *Let A be a double n -Poisson algebra and M be a double n -Poisson A -bimodule. Then*

- (i) *the bracket $\{-, -\}$ makes $A_{\mathfrak{h}}$ into an n -Lie algebra and A into n -Lie module;*
- (ii) *the brackets $\{-, -\}_M$ and $\{-, -\}_{M, \mathfrak{h}}$ make M and $M_{\mathfrak{h}}$ respectively into n -Lie module over $A_{\mathfrak{h}}$.*

4. Double Poisson structure on the cobar construction

In this section, we will define a natural double bracket on the DG algebra R which is the cobar construction of a cyclic A_∞ coalgebra.

4.1. Cyclic algebras and coalgebras. The notion of *cyclic A_∞ algebras* is first introduced by Kontsevich [33]. Recall that a *symmetric bilinear form of degree $-d$* on a graded vector space V is a bilinear pairing $\langle -, - \rangle : V \otimes V \rightarrow k[d]$ such that

$$\langle v, w \rangle = (-1)^{|v||w|} \langle w, v \rangle \quad \text{for all } v, w \in V. \tag{4.1}$$

A cyclic A_∞ algebra A is an A_∞ algebra with a non-degenerate symmetric bilinear form of degree $-d$

$$\langle -, - \rangle : A \otimes A \rightarrow k[d]$$

such that

$$\begin{aligned} \langle \mu_n(a_1, a_2, \dots, a_n), a_{n+1} \rangle \\ = (-1)^{n+|a_{n+1}|(|a_1|+\dots+|a_n|)} \langle \mu_n(a_{n+1}, a_1, \dots, a_{n-1}), a_n \rangle \end{aligned} \tag{4.2}$$

for all $n \in \mathbb{N}$ and all $a_0, a_1, \dots, a_n \in A$.

Similarly, one can define *cyclic A_∞ coalgebras*:

Definition 4.1 (Cyclic A_∞ coalgebra). Suppose $(C, \{\Delta_k\})$ is an A_∞ coalgebra. C is called *cyclic of degree $-d$* if there is a non-degenerate symmetric bilinear form of degree $-d$

$$\langle -, - \rangle : C \otimes C \rightarrow k[d]$$

such that for any $a, b \in C$,

$$\langle a, b^1 \rangle \cdot b^2 \cdots b^r = (-1)^{r+|b^1|(|a|+r)} \langle b, a^r \rangle \cdot a^1 \cdots a^{r-1} \in C^{\otimes(r-1)}, \quad (4.3)$$

where we write $\Delta_r(a) = a^1 a^2 \cdots a^r$ and $\Delta_r(b) = b^1 b^2 \cdots b^r$.

The notion of cyclic A_∞ coalgebras will be useful for our next discussion. We have the following lemma:

Lemma 4.2. *Suppose C is of finite dimension. Then C is a cyclic A_∞ coalgebra if and only if $A := C^* = \text{Hom}_k(C, k)$ is a cyclic A_∞ algebra.*

Proof. Denote the A_∞ operators of C and A by $\Delta_1, \Delta_2, \dots$ and μ_1, μ_2, \dots , respectively. Under the isomorphism of k -vector spaces

$$\begin{aligned} C &\xrightarrow{\cong} A \\ a &\mapsto a^* := \langle -, a \rangle, \end{aligned}$$

consider the evaluation of both sides of (4.3) on any $(x_1, x_2, \dots, x_{r-1}) \in A^{\otimes(r-1)}$. We have:

$$\begin{aligned} (x_1, x_2, \dots, x_{r-1})(\langle b, a^r \rangle (a^1 \cdots a^{r-1})) &= \langle b, a^r \rangle \cdot (x_1, x_2, \dots, x_{r-1})(a^1 \cdots a^{r-1}) \\ &= (x_1, x_2, \dots, x_{r-1}, b^*) (a^1 a^2 \cdots a^r) \\ &= (x_1, x_2, \dots, x_{r-1}, b^*) \Delta_r(a) \\ &= (\mu_r(x_1, x_2, \dots, x_{r-1}, b^*)(a)) \\ &= \langle \mu_r(x_1, x_2, \dots, x_{r-1}, b^*), a^* \rangle. \end{aligned} \quad (4.4)$$

Similarly, the evaluation

$$\begin{aligned} (x_1, x_2, \dots, x_{r-1})(\langle a, b^1 \rangle (b^2 \cdots b^r)) &= (a^*, x_1, x_2, \dots, x_{r-1})(b^1 \cdots b^r) \\ &= (a^*, x_1, x_2, \dots, x_{r-1}) \Delta_r(b) \\ &= \mu_r(a^*, x_1, x_2, \dots, x_{r-1})(b) \\ &= \langle \mu_r(a^*, x_1, x_2, \dots, x_{r-1}), b^* \rangle. \end{aligned} \quad (4.5)$$

The most right hand side of (4.4) equals the most right hand side of (4.5) with sign counted if and only if (4.2) holds, and thus the lemma follows. \square

Remark 4.3. By repeatedly applying (4.2) to (4.4) or to (4.5) one gets more identities like

$$\langle a^m, b \rangle \cdot a^{m+1} \cdots a^r a^1 \cdots a^{r-1} = \pm \langle a, b^\ell \rangle \cdot b^1 \cdots b^{\ell-1} b^{\ell+1} \cdots b^r. \quad (4.6)$$

We leave the check to the interested reader.

4.2. Double bracket on the cobar construction. The following result has already appeared in [1, Theorem 15] when C is a cyclic coalgebra (not A_∞).

Lemma 4.4. *Let C be a cyclic A_∞ coalgebra of degree $-d$ and let $R = \Omega_\infty(\tilde{C})$. Then we define $\{\{-, -\}\} : \bar{R} \times \bar{R} \rightarrow \bar{R} \otimes \bar{R}$ by*

$$\{\{r, q\}\} := \sum_{\substack{(i=1, \dots, k) \\ (j=1, \dots, m)}} (-1)^{|v_i|+\epsilon} \langle v_i, w_j \rangle \cdot (s^{-1}w_1 \cdots s^{-1}w_{j-1}s^{-1}v_{i+1} \cdots s^{-1}v_k) \\ \otimes (s^{-1}v_1 \cdots s^{-1}v_{i-1}s^{-1}w_{j+1} \cdots s^{-1}w_m), \quad (4.7)$$

for $r = (s^{-1}v_1s^{-1}v_2 \cdots s^{-1}v_k)$ and $q = (s^{-1}w_1s^{-1}w_2 \cdots s^{-1}w_m)$, where ϵ is

$$(|r| + d)(|s^{-1}w_1| + \cdots + |s^{-1}w_{j-1}|) \\ + (|s^{-1}v_1| + \cdots + |s^{-1}v_{i-1}| + |s^{-1}w_j| + d)(|s^{-1}v_{i+1}| + \cdots + |s^{-1}v_k|).$$

Then the bracket (4.7) gives a DG double n -Poisson structure on R , where $n = 2 - d$.

Proof. The proof is essentially the same as in [1, Theorem 15]. We only need to show that the bracket commutes with the differential, which involves the A_∞ operators.

From the definition of the double bracket,

$$\{\{d(a_1a_2 \cdots a_m), b_1b_2 \cdots b_n\}\} + (-1)^{|a|} \{\{a_1a_2 \cdots a_m, d(b_1b_2 \cdots b_n)\}\} \\ - d(\{\{a_1a_2 \cdots a_m, b_1b_2 \cdots b_n\}\})$$

contains summands whose coefficient are pairings of components of $\Delta(a)$ with b , or pairings of a with components of $\Delta(b)$, namely,

$$\sum_{i,j} \sum_k \langle a_i^k, b_j \rangle b_1b_2 \cdots b_{j-1}a_i^{k+1} \cdots a_i^r a_{i+1} \cdots a_m \\ \otimes a_1a_2 \cdots a_{i-1}a_i^1 \cdots a_i^{k-1} b_{j+1} \cdots b_n \\ - \sum_{i,j} \sum_\ell \langle a_i, b_j^\ell \rangle b_1b_2 \cdots b_{j-1}b_j^1 \cdots b_j^{\ell-1} a_{i+1} \cdots a_m \\ \otimes a_1a_2 \cdots a_{i-1}b_j^{\ell+1} \cdots b_j^s b_{j+1} \cdots b_n,$$

if we write $\Delta_k(a_i) = a_i^1a_i^2 \cdots a_i^k$ and $\Delta_\ell(b_j) = b_j^1b_j^2 \cdots b_j^\ell$. However, from the cyclicity of the pairing (see (4.3) or its general form (4.6)), these two types of terms exactly cancel with each other. Thus the double bracket commutes with the differential. \square

Remark 4.5. One can easily extend $\{\{-, -\}\}$ to R by taking $\{\{r, 1\}\} = 0$.

5. Brackets on the bimodule of one-forms

Let C be a cyclic A_∞ coalgebra and let $R := \Omega_\infty(\tilde{C})$. Using the double bracket on R defined in Lemma 4.4, we introduce a double bracket on Ω_R^1 and the induced bracket on $\Omega_{R, \natural}^1$. Then we consider the corresponding Lie brackets on Ω_R^1 and $\Omega_{R, \natural}^1$.

5.1. Double bracket on Ω_R^1 . First we define a double bracket

$$\{\{-, -\}\}_{R \otimes R} : R \times (R \otimes R) \rightarrow R \otimes (R \otimes R) \oplus (R \otimes R) \otimes R$$

as follows

$$\{\{r, p \otimes q\}\}_{R \otimes R} := \{\{r, p\}\} \otimes q + (-1)^{|p|(|r|+n)} p \otimes \{\{r, q\}\}, \quad (5.1)$$

where $\{\{-, -\}\}$ is a double bracket on R defined in Lemma 4.4, and in the right-hand side of (5.1), the first summand lies in $R \otimes (R \otimes R)$ and the second lies in $(R \otimes R) \otimes R$. Then one can show

Lemma 5.1. $(R \otimes R, \{\{-, -\}\}_{R \otimes R})$ is a DG double n -Poisson R -bimodule.

Proof. We need to verify axioms (i)–(iii) for the double bimodule. We first verify

$$(i) \quad \{\{r, q \cdot (p_1 \otimes p_2)\}\}_{R \otimes R} = \{\{r, q\}\}(p_1 \otimes p_2) + (-1)^{|q|(|r|+n)} q \{\{r, p_1 \otimes p_2\}\}_{R \otimes R}.$$

By definition the left hand side (LHS) is

$$\begin{aligned} \{\{r, q \cdot p_1 \otimes p_2\}\}_{R \otimes R} &= \{\{r, q\}\} \cdot (p_1 \otimes p_2) + (-1)^{|q|(|r|+n)} q \cdot \{\{r, p_1\}\} \otimes p_2 \\ &\quad + (-1)^{(|q|+|p_1|)(|r|+n)} q \cdot p_1 \otimes \{\{r, p_2\}\} \end{aligned}$$

which is identically equal to the right hand side (RHS). Next we verify

$$(ii) \quad \{\{r \cdot q, p_1 \otimes p_2\}\}_{R \otimes R} = r * \{\{q, p_1 \otimes p_2\}\}_{R \otimes R} + (-1)^{|q|(|p_1|+|p_2|+n)} \{\{r, p_1 \otimes p_2\}\}_{R \otimes R} * q.$$

Indeed, the LHS is equal to

$$\begin{aligned} \{\{r \cdot q, p_1\}\} \otimes p_2 + (-1)^{|p_1|(|r|+|q|+n)} p_1 \otimes \{\{r \cdot q, p_2\}\} \\ = r * \{\{q, p_1\}\} \otimes p_2 + (-1)^{|q|(|p_1|+n)} \{\{r, p_1\}\} * q \otimes p_2 \\ + (-1)^{|p_1|(|r|+|q|+n)} p_1 \otimes (r * \{\{q, p_2\}\} + (-1)^{|q|(|p_2|+n)} \{\{r, p_2\}\} * q) \end{aligned}$$

and the RHS is

$$\begin{aligned} r * \{\{q, p_1\}\} \otimes p_2 + (-1)^{|p_1|(|r|+|q|+n)} p_1 \otimes r * \{\{q, p_2\}\} \\ + (-1)^{|q|(|p_1|+|p_2|+n)+|q||p_2|} \{\{r, p_1\}\} * q \otimes p_2 \\ + (-1)^{|q|(|p_1|+|p_2|+n)+|p_1|(|r|+n)} p_1 \otimes \{\{r, p_2\}\} * q \end{aligned}$$

and hence they are equal. Similarly, we can prove the Jacobi identity (iii).

Next we show that $\{\{-, -\}\}_{R \otimes R}$ commutes with the differential. Recall that $d_{R \otimes R} = d_R \otimes 1 + 1 \otimes d_R$. So we need to prove

$$d_R \{\{r, p \otimes q\}\}_{R \otimes R} = \{\{d_R(r), p \otimes q\}\}_{R \otimes R} + (-1)^{|r|+n} \{\{r, d_R(p \otimes q)\}\}_{R \otimes R}.$$

Indeed, using (5.1) one has

$$\begin{aligned}
d_R\{\{r, p \otimes q\}\}_{R \otimes R} &= d_R\{\{r, p\}\} \otimes q + (-1)^{|r|+|p|+n}\{\{r, p\}\} \otimes d_R(q) \\
&\quad + (-1)^{|p|(|r|+n)}d_R(p) \otimes \{\{r, q\}\} + (-1)^{|p|(|r|+n+1)}p \otimes d_R\{\{r, q\}\}; \\
\{\{d_R(r), p \otimes q\}\}_{R \otimes R} &= \{\{d_R(r), p\}\} \otimes q + (-1)^{|p|(|r|+n+1)}p \otimes \{\{d_R(r), q\}\}; \\
\{\{r, d_R(p \otimes q)\}\}_{R \otimes R} &= \{\{r, d_R(p) \otimes q\}\} + (-1)^{|p|}\{\{r, p \otimes d_R(q)\}\} \\
&= \{\{r, d_R(p)\}\} \otimes q + (-1)^{(|p|+1)(|r|+n)}d_R(p) \otimes \{\{r, q\}\} \\
&\quad + (-1)^{|p|}\{\{r, p\}\} \otimes d_R(q) + (-1)^{|p|(|r|+n+1)}p \otimes \{\{r, d_R(q)\}\}.
\end{aligned}$$

Combining the RHS of these identities we get

$$\begin{aligned}
&(d_R\{\{r, p\}\} - \{\{d_R(r), p\}\} - (-1)^{|r|+n}\{\{r, d_R(p)\}\}) \otimes q \\
&+ (-1)^{|p|(|r|+n+1)}p \otimes (d_R\{\{r, q\}\} - \{\{d_R(r), q\}\} - (-1)^{|r|+n}\{\{r, d_R(q)\}\}) \quad (5.2)
\end{aligned}$$

and this expression is equal to 0, since by Lemma 4.4 $\{\{-, -\}$ commutes with d_R . This finishes our proof. \square

We claim that the bracket in (5.1) can be restricted to Ω_R^1 . Recall that $\Omega_R^1 \cong R \otimes s^{-1}C \otimes R$ and $\Omega_{R, \mathfrak{q}}^1 \cong s^{-1}C \otimes R$, where identifications are given by map I defined in (2.3). Indeed, let

$$\omega = (s^{-1}v_1 \cdots s^{-1}v_{p-1}) \otimes s^{-1}v_p \otimes (s^{-1}v_{p+1} \cdots s^{-1}v_m) \in \Omega_R^1.$$

Then $I(\omega) = b \cdot s^{-1}v_p \otimes c - b \otimes s^{-1}v_p \cdot c$, where $b = (s^{-1}v_1 \cdots s^{-1}v_{p-1})$ and $c = (s^{-1}v_{p+1} \cdots s^{-1}v_m)$ and one has

$$\{\{a, I(\omega)\}\}_{R \otimes R} = \{\{a, b \cdot s^{-1}v_p\}\} \otimes c + (-1)^{(|b|+|s^{-1}v_p|)(|a|+n)}b \cdot s^{-1}v_p \otimes \{\{a, c\}\}$$

$$- \{\{a, b\}\} \otimes s^{-1}v_p \cdot c - (-1)^{|b|(|a|+n)}b \otimes \{\{a, s^{-1}v_p \cdot c\}\}$$

$$= \{\{a, b\}\}^{(1)} \otimes \{\{a, b\}\}^{(2)} \cdot s^{-1}v_p \otimes c \quad (5.3)$$

$$+ (-1)^{|b|(|a|+n)}b \cdot \{\{a, s^{-1}v_p\}\}^{(1)} \otimes \{\{a, s^{-1}v_p\}\}^{(2)} \otimes c \quad (5.4)$$

$$+ (-1)^{(|b|+|s^{-1}v_p|)(|a|+n)}b \cdot s^{-1}v_p \otimes \{\{a, c\}\}^{(1)} \otimes \{\{a, c\}\}^{(2)} \quad (5.5)$$

$$- \{\{a, b\}\}^{(1)} \otimes \{\{a, b\}\}^{(2)} \otimes s^{-1}v_p \cdot c \quad (5.6)$$

$$- (-1)^{|b|(|a|+n)}b \otimes \{\{a, s^{-1}v_p\}\}^{(1)} \otimes \{\{a, s^{-1}v_p\}\}^{(2)} \cdot c \quad (5.7)$$

$$- (-1)^{(|b|+|s^{-1}v_p|)(|a|+n)}b \otimes s^{-1}v_p \cdot \{\{a, c\}\}^{(1)} \otimes \{\{a, c\}\}^{(2)}. \quad (5.8)$$

Now (5.3) + (5.6) $\in R \otimes \Omega_R^1$ and (5.5) + (5.8) $\in \Omega_R^1 \otimes R$. On the other hand

$$(5.4) - (-1)^{|b|(|a|+n)} b \otimes \{\{a, s^{-1}v_p\}\}^{(1)} \cdot \{\{a, s^{-1}v_p\}\}^{(2)} \otimes c \in \Omega_R^1 \otimes R, \quad (5.9)$$

$$(5.7) + (-1)^{|b|(|a|+n)} b \otimes \{\{a, s^{-1}v_p\}\}^{(1)} \cdot \{\{a, s^{-1}v_p\}\}^{(2)} \otimes c \in R \otimes \Omega_R^1. \quad (5.10)$$

These two formulas hold since, for example in (5.9), by taking out the third common component c in the tensor, the first two components exactly lie in Ω_R^1 . Thus, we define $\{\{-, -\}\}_{\Omega_R^1} : R \times \Omega_R^1 \rightarrow (R \otimes \Omega_R^1) \oplus (\Omega_R^1 \otimes R)$ as

$$\{\{a, \omega\}\}_{\Omega_R^1} := \{\{a, I(\omega)\}\}_{R \otimes R} \quad (5.11)$$

and we have proved

Corollary 5.2. $(\Omega_R^1, \{\{-, -\}\}_{\Omega_R^1})$ is a DG double n -Poisson sub-bimodule of $R \otimes R$.

5.2. Lie brackets on Ω_R^1 and $\Omega_{R, \mathfrak{h}}^1$. Let us recall some notations from the previous section. First, $\{-, -\} := \mu_R \circ \{\{-, -\}\}$, where μ_R is the multiplication map on R . Second,

$$\{-, -\}_{\Omega_R^1} := \mu_{\Omega_R^1} \circ \{\{-, -\}\}_{\Omega_R^1}, \quad \{-, -\}_{\Omega_{R, \mathfrak{h}}^1} := \mathfrak{h} \circ \{-, -\}_{\Omega_R^1},$$

where $\mu_{\Omega_R^1}$ is the bimodule action map.

Theorem 5.3. Let C be a cyclic A_∞ coalgebra of degree $-d$ and let $R = \overline{\Omega}_\infty(\tilde{C})$. Let $n := 2 - d$. Then

- (a) $\{-, -\}$ induces a DG n -Lie algebra structure on $R_{\mathfrak{h}}$ and a DG n -Lie module on R .
- (b) $\{-, -\}_{\Omega_R^1}$ and $\{-, -\}_{\Omega_{R, \mathfrak{h}}^1}$ make Ω_R^1 and $\Omega_{R, \mathfrak{h}}^1$ into DG n -Lie modules over $R_{\mathfrak{h}}$.

In particular, $H_\bullet(R_{\mathfrak{h}})$ is a graded n -Lie algebra and $H_\bullet(R)$, $H_\bullet(\Omega_R^1)$ and $H_\bullet(\Omega_{R, \mathfrak{h}}^1)$ are n -Lie modules.

Proof. Follows from Proposition 3.11 for the double bracket on R defined in Lemma 4.4 and the double bracket on Ω_R^1 defined in Corollary 5.2. \square

Now we turn our attention to the 2-periodic sequence of complexes

$$\xrightarrow{\bar{\partial}} \Omega_{R, \mathfrak{h}}^1 \xrightarrow{-\beta} \overline{R} \xrightarrow{\bar{\partial}} \Omega_{R, \mathfrak{h}}^1 \xrightarrow{-\beta} \rightarrow$$

which by Proposition 2.8 is identical to the cyclic bicomplex of C . Our claim is

Theorem 5.4. $\bar{\partial}$ and β are morphisms of Lie modules, that is,

$$I(\bar{\partial}\{r, q\}) = \{r, \bar{\partial}(q)\}_{\Omega_{R, \mathfrak{h}}^1}, \quad \beta\{r, \omega\}_{\mathfrak{h}} = \{r, \beta(\omega)\}. \quad (5.12)$$

We need the following auxiliary statement.

Lemma 5.5. *Let $r \in R$ and $\omega = q \otimes s^{-1}v \in \Omega_{R, \natural}^1$. Then*

$$\begin{aligned} \{r, \omega\}_{\Omega_{R, \natural}^1} &= \natural \circ \left[\{r, q\}s^{-1}v \otimes 1 - \{r, q\} \otimes s^{-1}v \right. \\ &\quad \left. + (-1)^{|q|(|r|+n)}(q \cdot \{r, s^{-1}v\} \otimes 1 - q \otimes \{r, s^{-1}v\}) \right]. \end{aligned} \quad (5.13)$$

Proof. It follows directly from the definition of the double bracket on Ω_R^1 . Indeed, we have

$$\begin{aligned} \{r, \omega\}_{\Omega_R^1} &= \mu \circ \{\{r, \mathbf{I}(\omega)\}\}_{R \otimes R} \\ &= \mu_{\Omega_R^1} \circ [\{\{r, q \cdot s^{-1}v \otimes 1\}\}_{R \otimes R} - \{\{r, q \otimes s^{-1}v\}\}_{R \otimes R}] \\ &= \mu_{\Omega_R^1} \circ [\{\{r, q\}\} \cdot s^{-1}v \otimes 1 - \{\{r, q\}\} \otimes s^{-1}v] \\ &\quad + (-1)^{|q|(|r|+n)} \mu_{\Omega_R^1} \circ [q \cdot \{\{r, s^{-1}v\}\} \otimes 1 - q \otimes \{\{r, s^{-1}v\}\}] \\ &= \{r, q\}s^{-1}v \otimes 1 - \{r, q\} \otimes s^{-1}v \\ &\quad + (-1)^{|q|(|r|+n)} \mu_{\Omega_R^1} \circ [q \cdot \{\{r, s^{-1}v\}\}^{(1)} \otimes \{\{r, s^{-1}v\}\}^{(2)} \otimes 1 - q \otimes \{r, s^{-1}v\} \otimes 1] \\ &\quad + (-1)^{|q|(|r|+n)} \mu_{\Omega_R^1} \circ [q \otimes \{r, s^{-1}v\} \otimes 1 - q \otimes \{\{r, s^{-1}v\}\}^{(1)} \otimes \{\{r, s^{-1}v\}\}^{(2)}] \\ &= \{r, q\}s^{-1}v \otimes 1 - \{r, q\} \otimes s^{-1}v \\ &\quad + (-1)^{|q|(|r|+n)}(q \cdot \{\{r, s^{-1}v\}\}^{(1)} \otimes \{\{r, s^{-1}v\}\}^{(2)} - q \otimes \{r, s^{-1}v\}) \\ &\quad + (-1)^{|q|(|r|+n)}(q \cdot \{r, s^{-1}v\} \otimes 1 - q \cdot \{\{r, s^{-1}v\}\}^{(1)} \otimes \{\{r, s^{-1}v\}\}^{(2)}) \\ &= \{r, q\} \cdot s^{-1}v \otimes 1 - \{r, q\} \otimes s^{-1}v \\ &\quad + (-1)^{|q|(|r|+n)}(q \cdot \{r, s^{-1}v\} \otimes 1 - q \otimes \{r, s^{-1}v\}). \quad \square \end{aligned}$$

Proof of Theorem 5.4. To prove the first identity of (5.12), it suffices to show

$$\mathbf{I}(\partial\{r, q\}) = \{r, \partial(q)\}_{\Omega_R^1}.$$

By (2.4), we have $\mathbf{I}(\partial\{r, q\}) = \{r, q\} \otimes 1 - 1 \otimes \{r, q\}$. On the other hand

$$\begin{aligned} \{r, \partial(q)\}_{\Omega_R^1} &= \mu_{\Omega_R^1} \{\{r, \mathbf{I}(\partial(q))\}\}_{R \otimes R} = \mu_{\Omega_R^1} \{\{r, q \otimes 1 - 1 \otimes q\}\}_{R \otimes R} \\ &= \mu_{\Omega_R^1} [\{\{r, q\}\} \otimes 1 - 1 \otimes \{\{r, q\}\}] = \{r, q\} \otimes 1 - 1 \otimes \{r, q\}. \end{aligned}$$

Let $\omega = q \otimes s^{-1}v$ for some $q \in R$ and $v \in C$. Then

$$\beta(\omega) = q \cdot s^{-1}v - (-1)^{|s^{-1}v||q|}s^{-1}v \cdot q, \quad \mathbf{I}(\omega) = q \cdot s^{-1}v \otimes 1 - q \otimes s^{-1}v,$$

and

$$\begin{aligned}
\{r, \beta(\omega)\} &= \{r, q \cdot s^{-1}v\} - (-1)^{|s^{-1}v||q|} \{r, s^{-1}v \cdot q\} \\
&= \{r, q\} \cdot s^{-1}v + (-1)^{|q|(|r|+n)} q \cdot \{r, s^{-1}v\} \\
&\quad - (-1)^{|s^{-1}v||q|} \{r, s^{-1}v\} \cdot q - (-1)^{|s^{-1}v|(|q|+|r|+n)} s^{-1}v \cdot \{r, q\}.
\end{aligned} \tag{5.14}$$

On the other hand, by the commutative diagram (2.7) for R , we have

$$\beta(\{r, \omega\}_{\mathfrak{h}}) = \mu \circ \sigma(\mathbf{I}\{r, \omega\}_{\Omega_R^1}). \tag{5.15}$$

Using (5.13), one has

$$\begin{aligned}
\beta(\{r, \omega\}_{\mathfrak{h}}) &= \{r, q\} \cdot s^{-1}v - (-1)^{|s^{-1}v|(|r|+|q|+n)} s^{-1}v \cdot \{r, q\} \\
&\quad + (-1)^{|q|(|r|+n)} q \cdot \{r, s^{-1}v\} - (-1)^{|s^{-1}v||q|} \{r, s^{-1}v\} \cdot q.
\end{aligned} \tag{5.16}$$

We finish our proof by comparing (5.14) and (5.16). \square

6. N -Koszul Calabi–Yau algebras

Definition 6.1 (Ginzburg [23]). An associative algebra A is said to be a *Calabi–Yau algebra* of dimension d if A is homologically smooth and there exists an isomorphism

$$\eta : \mathrm{RHom}_{A \otimes A^{\mathrm{op}}}(A, A \otimes A) \rightarrow A[-d]$$

in the derived category of $A \otimes A^{\mathrm{op}}$ -modules.

In the above definition, an algebra A is said to be *homologically smooth* if A is a perfect $A \otimes A^{\mathrm{op}}$ -module, i.e. it has a finitely-generated projective resolution of finite length.

Ever since they are first introduced by Ginzburg, Calabi–Yau algebras have been widely studied by mathematicians from various fields.

So far, most of Calabi–Yau algebras appeared in literature are Koszul or N -Koszul, (the most general case is due to Van den Bergh [46]), and hence they are all of the form

$$\Omega_{\infty}(A^i) \xrightarrow{\cong} A,$$

where A^i is the Koszul dual A_{∞} coalgebra of A , which is a cyclic A_{∞} coalgebra.

We now recall the definition of N -Koszul algebras. Let V be a finite dimensional vector space over k , and S be a subspace of $V^{\otimes N}$, where $N \geq 2$ is an integer. Let TV be the tensor algebra of V , and $\langle S \rangle$ be the two-sided ideal of TV generated by S . The quotient algebra $A := TV/\langle S \rangle$ is called an *N -homogeneous algebra*, and is denoted by $A = A(V, S)$. The N -homogenous algebra $A^{\vee} := TV^*/\langle S^{\perp} \rangle$ is called

N-homogeneous dual algebra, where V^* is the dual space of V and $S^\perp \subset (V^*)^{\otimes N}$ is the orthogonal complement of S in $(V^*)^{\otimes N}$. The *N*-Koszul dual algebra $A^!$ of the *N*-homogenous algebra A is defined as follows: set

$$N(i) = \begin{cases} Nj, & \text{if } i = 2j, \\ Nj + 1, & \text{if } i = 2j + 1, \end{cases}$$

and define $A^! := \bigoplus_i A_{N(i)}^\vee$. We also denote $A^i := (A^!)^* \subset TV$, which is called *N*-Koszul dual coalgebra of A . Here we use the convention that the dual of a graded space with finite-dimensional component is the direct sum of component-wise duals.

Definition 6.2 (Berger [4], *N*-Koszul algebra). An *N*-homogeneous algebra A is (left) *N*-Koszul if the trivial left A -module ${}_A k$ admits a linear projective resolution

$$\dots \rightarrow P_i \xrightarrow{b} P_{i-1} \xrightarrow{b} \dots \rightarrow P_1 \xrightarrow{b} P_0 \xrightarrow{b} {}_A k$$

with $P_i = A \otimes_k A_i^!$, and if we choose a basis $\{e_i\}$ for V and let $\{e^*\}$ be the dual basis, then the differential $b = \sum_i^{\dim V} e_i \otimes e_i^*$.

One can define the right *N*-Koszul algebra similarly. In [4], it is proved that an *N*-homogeneous algebra is left *N*-Koszul if and only if it is right *N*-Koszul. Note that 2-Koszul algebras are Koszul algebras in the usual sense. For a 2-Koszul algebra A , it is known that the Yoneda algebra $\text{Ext}_A^\bullet(k, k)$ is an associative algebra. The A_∞ algebra structure on $\text{Ext}_A^\bullet(k, k)$ is started by Lu, Palmieri, Wu and Zhang in papers [37, 38].

Proposition 6.3 (Berger–Marconnet; He–Lu). *Let A be an N -Koszul algebra. Then its Yoneda algebra $\text{Ext}_A^\bullet(k, k)$ is isomorphic to its N -Koszul dual algebra $A^!$ as unital, augmented A_∞ algebras.*

Proof. See Berger–Marconnet [5, Proposition 3.1], or He–Lu [25, Theorem 6.5]. \square

More precisely, the authors cited above showed that for *N*-Koszul algebras, all the A_∞ operators but m_2, m_N vanish. Now if A is an *N*-Koszul algebra, then A^i is a co-unital, co-augmented A_∞ coalgebra.

Theorem 6.4 (Dotsenko–Vallette). *Let A be an N -Koszul algebra, and A^i be its N -Koszul dual A_∞ coalgebra. Then the morphism of DG algebras $\Omega_\infty(A^i) \xrightarrow{\cong} A$ is a quasi-isomorphism.*

Proof. The $N = 2$ case is standard, see, for example, Loday–Vallette [36, Theorem 3.4.6]. For the *N*-Koszul case, see Dotsenko–Vallette [18, Theorem 5.1]. \square

Proposition 6.5 (He–Van Oystaeyen–Zhang; Wu–Zhu). *Let A be an N -Koszul algebra ($N \geq 3$). Then the following conditions are equivalent:*

- (1) A is a d -Calabi–Yau algebra;
- (2) $\text{Ext}_A^\bullet(k, k)$ is a unital and augmented cyclic A_∞ algebra of degree $-d$.

Proof. See Wu–Zhu [48, Corollary 4.11] and He–Van Oystaeyen–Zhang [27, Proposition 3.3]. \square

In view of Lemma 4.2, we can summarize the above three results in terms of A_∞ coalgebras in the following form (due to the works of Berger, Dotsenko–Vallette, He–Lu, He–Van Oystaeyen–Zhang, Van den Bergh and Wu–Zhu cited above):

Theorem 6.6. *Let A be an N -Koszul algebra. Then A is d -Calabi–Yau if and only if the Koszul dual A_∞ coalgebra A^i is cyclic of degree $-d$. Moreover, in this case $\Omega_\infty(A^i)$ is a cofibrant resolution of A .*

The most general form of this theorem is due to Van den Bergh given in [46, cf. Theorem 11.1]. Van den Bergh does not use the terminology “ N -Koszul”, but it is clear that all N -Koszul algebras are Koszul in the sense of [46], where the latter is the linear dual of the bar construction. Since the finite dimensionality $\text{Ext}_A^\bullet(k, k)$ is essentially used in the following construction, we focus only on the N -Koszul case.

6.1. Examples. In this subsection, we list several known examples of N -Koszul Calabi–Yau algebras, and therefore they all admit a derived non-commutative Poisson structure. Note that by the above theorem, we only need to describe the cyclic A_∞ algebra structure on their Koszul dual.

First let us remind that all graded 2- and 3-Calabi–Yau algebras are N -Koszul. This fact is proved by Berger–Marconnet in [5, Proposition 5.2]. Being graded is important here, since from Davison’s result there exist 3-Calabi–Yau algebras which are not superpotential algebras and hence not N -Koszul [16].

6.1.1. Three dimensional Sklyanin algebras. Let $a, b, c \in k$. The three dimensional Sklyanin algebra $A = A(a, b, c)$ is the graded k -algebra with generators x, y, z of degree one, and relations

$$\begin{aligned} f_1 &= cx^2 + bzy + ayz = 0, \\ f_2 &= azx + cy^2 + bxz = 0, \\ f_3 &= byx + axy + cz^2 = 0. \end{aligned}$$

This Sklyanin algebra is one of the important examples of Ginzburg’s Calabi–Yau algebras (see [23, Example 1.3.8]).

Smith showed in [42, Example 10.1] that A is Koszul, whose dual algebra $A^!$ is generated by ξ_1, ξ_2, ξ_3 with relations

$$\begin{aligned} c\xi_2\xi_3 - b\xi_3\xi_2, & \quad b\xi_1^2 - a\xi_2\xi_3, \\ c\xi_3\xi_1 - b\xi_1\xi_3, & \quad b\xi_2^2 - a\xi_3\xi_1, \\ c\xi_1\xi_2 - b\xi_2\xi_1, & \quad b\xi_3^2 - a\xi_1\xi_2. \end{aligned}$$

To describe the non-degenerate pairing, choose a basis for $A^!$:

$$\begin{aligned} A_0^! &: 1 \\ A_1^! &: \xi_1, \xi_2, \xi_3 \\ A_2^! &: \xi_1^2, \xi_2^2, \xi_3^2 \\ A_3^! &: \xi_1 \xi_2 \xi_3 \end{aligned}$$

The pairing $\langle u, v \rangle$ for homogeneous $u, v \in A^!$ is given by the scalar of uv with respect to $\xi_1 \xi_2 \xi_3$. This pairing is cyclically invariant, and therefore we obtain a derived non-commutative Poisson structure on A .

6.1.2. Four dimensional Sklyanin algebras. Let $\alpha, \beta, \gamma \in k$ such that

$$\alpha + \beta + \gamma + \alpha\beta\gamma = 0, \quad \{\alpha, \beta, \gamma\} \cap \{0, \pm 1\} = \emptyset.$$

The four dimensional Sklyanin algebra $A = A(\alpha, \beta, \gamma)$ is the graded k -algebra with generators x_0, x_1, x_2, x_3 of degree one, and relations $f_i = 0$, where

$$\begin{aligned} f_1 &= x_0 x_1 - x_1 x_0 - \alpha(x_2 x_3 + x_3 x_2), & f_2 &= x_0 x_1 + x_1 x_0 - (x_2 x_3 - x_3 x_2), \\ f_3 &= x_0 x_2 - x_2 x_0 - \beta(x_3 x_1 + x_1 x_3), & f_4 &= x_0 x_2 + x_2 x_0 - (x_3 x_1 - x_1 x_3), \\ f_5 &= x_0 x_3 - x_3 x_0 - \gamma(x_1 x_2 + x_2 x_1), & f_6 &= x_0 x_3 + x_3 x_0 - (x_1 x_2 - x_2 x_1). \end{aligned}$$

As proved by Smith and Stafford [43, Propositions 4.3–4.9], A is Koszul, whose Koszul dual algebra $A^!$ is generated by $\xi_0, \xi_1, \xi_2, \xi_3$ with the following relations:

$$\begin{aligned} \xi_0^2 &= \xi_1^2 = \xi_2^2 = \xi_3^2 = 0, \\ 2\xi_2 \xi_3 + (\alpha + 1)\xi_0 \xi_1 - (\alpha - 1)\xi_1 \xi_0 &= 0, \\ 2\xi_3 \xi_2 + (\alpha - 1)\xi_0 \xi_1 - (\alpha + 1)\xi_1 \xi_0 &= 0, \\ 2\xi_3 \xi_1 + (\beta + 1)\xi_0 \xi_2 - (\beta - 1)\xi_2 \xi_0 &= 0, \\ 2\xi_1 \xi_3 + (\beta - 1)\xi_0 \xi_2 - (\beta + 1)\xi_2 \xi_0 &= 0, \\ 2\xi_1 \xi_2 + (\gamma + 1)\xi_0 \xi_3 - (\gamma - 1)\xi_3 \xi_0 &= 0, \\ 2\xi_2 \xi_1 + (\gamma - 1)\xi_0 \xi_3 - (\gamma + 1)\xi_3 \xi_0 &= 0. \end{aligned}$$

Smith and Stafford also showed that $A^!$ admits a non-degenerate symmetric pairing. To see this, $A^!$ is spanned by the following elements:

$$\begin{aligned} A_0^! &: 1 \\ A_1^! &: \xi_0, \xi_1, \xi_2, \xi_3 \\ A_2^! &: \xi_0 \xi_1, \xi_0 \xi_2, \xi_0 \xi_3, \xi_1 \xi_0, \xi_2 \xi_0, \xi_3 \xi_0 \\ A_3^! &: \xi_0 \xi_1 \xi_0, \xi_0 \xi_2 \xi_0, \xi_0 \xi_3 \xi_0, \xi_1 \xi_0 \xi_1 \\ A_4^! &: \xi_0 \xi_1 \xi_0 \xi_1 \end{aligned}$$

and all other degree components are zero. Also, in degree 4, we have the following identities

$$\begin{aligned}\xi_0 \xi_j \xi_0 \xi_j &= -\xi_j \xi_0 \xi_j \xi_0 \quad \text{for } 1 \leq j \leq 3, & \xi_0 \xi_i \xi_0 \xi_j &= 0 \quad \text{for } i \neq j, \\ \xi_0 \xi_3 \xi_0 \xi_3 &= \frac{1+\alpha}{1-\gamma} \xi_0 \xi_1 \xi_0 \xi_1, \\ \xi_0 \xi_2 \xi_0 \xi_2 &= \frac{1+\gamma}{1-\beta} \frac{1+\alpha}{1-\gamma} \xi_0 \xi_1 \xi_0 \xi_1.\end{aligned}$$

The pairing $\langle a, b \rangle$ for homogeneous $a, b \in A^1$ is defined to be the scalar of ab with respect to $\xi_0 \xi_1 \xi_0 \xi_1$. One easily sees the pairing such defined is graded symmetric and is cyclic. Therefore $A(\alpha, \beta, \gamma)$ is Koszul Calabi–Yau of dimension 4, and there is derived non-commutative Poisson structure on it.

6.1.3. Universal enveloping algebras. Let us first say a bit about *linear-quadratic Koszul algebras*. Suppose V is a finite dimensional vector space. A *linear quadratic relation* is a subspace $S \subset V \oplus V^{\otimes 2}$. And we may define the linear quadratic algebra $A(V, S)$ as before.

In this subsection, we assume S satisfies the following two conditions:

$$S \cap V = 0, \tag{6.1}$$

$$\{S \otimes V + V \otimes S\} \cap V^{\otimes 2} = S \cap V^{\otimes 2}. \tag{6.2}$$

Let $qS : S \rightarrow V^{\otimes 2}$ be the projection.

Definition 6.7 (Linear quadratic Koszul algebra). A linear quadratic algebra $A = A(V, S)$ is said to be *Koszul* if it satisfies conditions (6.1) and (6.2) and if the quadratic algebra $A(V, qS)$ is Koszul.

Since qS is the image of the projection of S to $V^{\otimes 2}$, we in fact have a map

$$\phi : qS \rightarrow V$$

such that $S = \{X - \phi(X) \mid X \in qS\}$. Denote by $(qA)^i$ the quadratic dual coalgebra of $T(V)/(qS)$, then this ϕ gives a map

$$d_\phi : (qA)^i \rightarrow qS \rightarrow V,$$

which extends to a coderivation $d_\phi : (qA)^i \rightarrow T(V)$.

Now if

$$\{S \otimes V + V \otimes S\} \cap S^{\otimes 2} \subset qS,$$

then the images of d_ϕ lie in $(qA)^i$. We in fact get a co-derivation

$$d_\phi : (qA)^i \rightarrow (qA)^i.$$

And if furthermore,

$$\{S \otimes V + V \otimes S\} \cap S^{\otimes 2} \subset S \otimes V^{\otimes 2}, \tag{6.3}$$

then $(d_\phi)^2 = 0$, and we obtain a differential graded coalgebra $((qA)^i, d_\phi)$, which is the Koszul dual coalgebra of $A(V, S)$, and is denoted by $(A(V, S)^i, d)$. Its linear dual is a differential graded algebra, and is denoted by $A(V, S)^!$. For more details, see [36, §3.6].

Lemma 6.8. *Suppose $A(V, qS)$ is a Calabi–Yau algebra of dimension n . Then $A(V, S)$ is a Calabi–Yau algebra of the same dimension if and only if any nonzero element in $A(V, S)_n^!$ is a cycle.*

Proof. As underlying spaces, $A(V, S)^!$ is isomorphic to $A(V, qS)^!$. Since $A(V, qS)$ is Calabi–Yau, there is a cyclically invariant non-degenerate pairing on $A(V, qS)^!$, which then gives the same pairing on $A(V, S)^!$. Thus to show $A(V, S)$ is Calabi–Yau, it is equivalent to show such pairing respects the differential, which is again equivalent to show the fundamental chain (the top chain in $A(V, S)^i$) is a cycle. This completes the proof. \square

Suppose \mathfrak{g} is a Lie algebra, then the universal enveloping algebra $U(\mathfrak{g})$ has Koszul dual differential graded coalgebra which is exactly the Chevalley–Eilenberg complex $(CE_\bullet(\mathfrak{g}), d)$. It is also known (cf. [24, Chapter V]) that the top chain of the Chevalley–Eilenberg complex is a cycle if and only if $\text{Tr}(\text{ad}(g)) = 0$ for all $g \in \mathfrak{g}$. A Lie algebra satisfying this property is called *unimodular*. Examples of unimodular Lie algebras are abelian Lie algebras, semi-simple Lie algebras, Heisenberg Lie algebras, and the Lie algebra of compact groups. Thus as a corollary to Lemma 6.8, we have the following statement, which is due to He, Van Oystaeyen, and Zhang [26, Theorem 5.3].

Theorem 6.9 (He, Van Oystaeyen, and Zhang). *Let \mathfrak{g} be a Lie algebra of dimension n . Then the universal enveloping algebra $U(\mathfrak{g})$ is n -Calabi–Yau if and only if \mathfrak{g} is unimodular.*

The derived non-commutative Poisson structure on $U(\mathfrak{g})$ is highly nontrivial, even in the polynomial case (i.e. the case where \mathfrak{g} is abelian); for more details see §9.

6.1.4. Yang–Mills algebras. Yang–Mills algebras are introduced by Connes and Dubois-Violette [11]. An algebra A is called a *Yang–Mills algebra* if A is generated by the elements x_i ($i \in \{1, \dots, n\}$) with the following relations:

$$g^{ij}[x_i, [x_j, x_l]] = 0, \quad l \in \{1, 2, \dots, n\},$$

where (g^{ij}) is a symmetric invertible $n \times n$ -matrix. Equivalently, $A = A(V, S)$ is a 3-homogenous algebra, with $V := \bigoplus_i kx_i$, and $S \subset V^{\otimes 3}$ spanned by

$$g^{ij}(x_i \otimes x_j \otimes x_l + x_l \otimes x_i \otimes x_i - 2x_i \otimes x_l \otimes x_j).$$

In literature, the above A is also denoted by $YM(n)$.

Yang–Mills algebras are 3-Koszul 3-Calabi–Yau. In fact Connes and Dubois-Violette [11, Theorem 1] proved that Yang–Mills algebras are 3-Koszul, and Berger–Taillefer [6, Proposition 4.5] proved they are 3-Calabi–Yau.

We give a brief description of the A_∞ operators and the pairing on $\text{YM}(n)^\dagger$. For simplicity, we take (g^{ij}) to be the identity matrix. Denote $V = \text{Span}_k \{x_1, x_2, \dots, x_n\}$, and

$$S = \text{Span}_k \left\{ \sum_{i=1}^n [x_i, [x_i, x_j]] : 1 \leq j \leq n \right\} \subset V^{\otimes 3}.$$

Then

$$\text{YM}(n) = TV / \langle S \rangle.$$

And the homogeneous dual algebra of $\text{YM}(n)$ is given as follows (see Connes and Dubois-Violette [11, Proposition 1] and also Herscovich and Solotar [29, Proposition 2.3]):

$$\begin{aligned} \text{YM}(n)_0^\vee &= k1, & \text{YM}(n)_1^\vee &= V^*, & \text{YM}(n)_2^\vee &= \bigoplus_{i,j=1}^n kx_i^*x_j^*, \\ \text{YM}(n)_3^\vee &= \bigoplus_{i=1}^n kx_i^*z, & \text{YM}(n)_4^\vee &= kz^2, & \text{YM}(n)_i^\vee &= 0 \quad (i > 4), \end{aligned}$$

where $z = \sum_{i=1}^n (x_i^*)^2$. The Koszul dual algebra of $\text{YM}(n)$ is thus given taking

$$\begin{aligned} \text{YM}(n)_0^\dagger &= \text{YM}(n)_0^\vee, & \text{YM}(n)_1^\dagger &= \text{YM}(n)_1^\vee, \\ \text{YM}(n)_2^\dagger &= \text{YM}(n)_3^\vee, & \text{YM}(n)_3^\dagger &= \text{YM}(n)_4^\vee, \end{aligned}$$

and the A_∞ operators (recall that we only have two nontrivial operators μ_2 and μ_3 in this case) are given as follows: denoting the above identification by

$$\phi : \text{YM}(n)_i^\dagger \xrightarrow{\cong} \text{YM}(n)_j^\vee,$$

then for $f_1, f_2, f_3 \in \text{YM}(n)^\dagger$,

$$\mu_2(f_1, f_2) = \begin{cases} \phi^{-1}(\phi(f_1) \cdot \phi(f_2)), & \text{if } \phi(f_1) \cdot \phi(f_2) \notin \text{YM}(n)_2^\vee, \\ 0, & \text{otherwise,} \end{cases} \quad (6.4)$$

and

$$\mu_3(f_1, f_2, f_3) = \begin{cases} \phi^{-1}(\phi(f_1) \cdot \phi(f_2) \cdot \phi(f_3)), & \text{if } \phi(f_1) \cdot \phi(f_2) \cdot \phi(f_3) \notin \text{YM}(n)_2^\vee, \\ 0, & \text{otherwise.} \end{cases} \quad (6.5)$$

There is a pairing on $\text{YM}(n)^\vee$, which is again the scalar of the corresponding product with respect to z^2 , and hence induces the pairing on $\text{YM}(n)^\dagger$ via ϕ .

7. Derived representation schemes

As we mentioned in §1, according to the Kontsevich–Rosenberg Principle, any meaningful non-commutative geometric structure on an associative algebra A should naturally induce its classical counterpart on the representative scheme $\text{Rep}_V(A)$, for all k -vector space V . Cuntz and Quillen in [14] first studied the smoothness problem for associative algebras, and introduced the notion of *smooth algebras*. If an algebra is smooth (e.g. cofibrant), then its representation scheme is smooth, too. However, in practice, an algebra is rarely smooth, and therefore it is really difficult to study the geometry on the representation scheme.

In the following, we first briefly recall the derived representation scheme of a DG algebra, and then discuss its relations to cyclic homology; most materials are taken from [2, 3]. After that, we recall several results on derived non-commutative Poisson structures from [1].

7.1. Derived representation scheme and cyclic homology. We start with the representation scheme of associative algebras. Suppose A is an associative algebra, and V is a k -vector space. Consider the following functor

$$\text{Rep}_V(A) : \text{CommAlg}_k \rightarrow \text{Sets}, \quad B \mapsto \text{Hom}_{\text{Alg}_k}(A, \text{End}(V) \otimes B),$$

where CommAlg_k is the category of commutative, unital k -algebras and Alg_k is the category of unital k -algebras. This functor is representable, which means there exists a commutative algebra, which we denote by $k[\text{Rep}_V(A)]$, such that

$$\text{Hom}_{\text{Alg}_k}(A, \text{End}(V) \otimes B) \cong \text{Hom}_{\text{CommAlg}_k}(k[\text{Rep}_V(A)], B).$$

Keeping V fixed, we in fact get a functor

$$(-)_V : \text{Alg}_k \rightarrow \text{CommAlg}_k, \quad A \mapsto k[\text{Rep}_V(A)]$$

which we call the *representation functor* in V , and the corresponding scheme is called the *representation scheme*. The representation functor can be extended to the category of DG algebras, DGA_k , which has a model structure in the sense of Quillen [40]. In [3] the authors showed that $(-)_V$ defines a left Quillen functor on DGA_k and therefore it has a total derived functor

$$\mathbf{L}(-)_V : \text{Ho}(\text{DGA}_k) \rightarrow \text{Ho}(\text{CDGA}_k)$$

from the homotopy category of DG algebras to the homotopy category of commutative DG algebras. When applied to A , $\mathbf{L}(A)_V$ is called the *derived representation scheme* of A , and is represented by a commutative DG algebra, which we denote by $\text{DRep}_V(A)$. The homology of $\text{DRep}_V(A)$, namely $H_\bullet(\text{DRep}_V(A))$, only depends on A and V , and is called the *representation homology* of A . We have that $H_0(\text{DRep}_V(A))$ is exactly $k[\text{Rep}_V(A)]$.

Now let us remind some results in model category theory (we recommend the excellent survey of Dwyer and Spalinski [19] for an introduction to model categories). Suppose \mathcal{A} is a model category, then the homotopy category $\mathrm{Ho}(\mathcal{A})$ is a category where the objects remain the same as \mathcal{A} , and the morphisms for two objects, say A and B , are given by

$$\mathrm{Hom}_{\mathrm{Ho}(\mathcal{A})}(A, B) := \mathrm{Hom}_{\mathcal{A}}(QA, QB)/\text{quasi-equivalences},$$

where QA and QB are the *cofibrant resolutions* of A and B respectively (for more details see [19, §5]). In the category of non-negatively (or non-positively) graded DG algebras, any quasi-free resolution is a cofibrant resolution.

Now suppose A is a DG algebra, and R is its cofibrant resolution. Then by definition $\mathbf{L}(A)_V$ can be represented by $R \mapsto \mathrm{DRep}_V(A)$. A key result in [3] is the construction of the higher trace map

$$\mathrm{Tr} : \mathbf{H}_\bullet(R_{\mathfrak{q}}) \rightarrow \mathbf{H}_\bullet(\mathrm{DRep}_V(A)).$$

Via Feigin–Tsygan’s result $\mathbf{H}_\bullet(\bar{R}_{\mathfrak{q}}) \cong \overline{\mathrm{HC}}_\bullet(A)$ (see [20] and also §2, Proposition 2.4) and by the fact there is a surjective map $\mathrm{HC}_\bullet(A) \rightarrow \overline{\mathrm{HC}}_\bullet(A)$, we in fact get a map

$$\mathrm{Tr} : \mathrm{HC}_\bullet(A) \rightarrow \mathbf{H}_\bullet(\mathrm{DRep}_V(A)), \quad (7.1)$$

called the *derived trace map*, which, when restricting to degree zero, is exactly the usual trace map $A_{\mathfrak{q}} = A/[A, A] \rightarrow k[\mathrm{Rep}_V(A)]$.

For an associative algebra A , $\mathrm{Rep}_V(A)$ has a natural $\mathrm{GL}(V)$ action, and the covariant space $\mathrm{Rep}_V(A)^{\mathrm{GL}(V)}$ classifies the isomorphism classes of representations of A on V . In the derived representation case, there is an analogous result, and moreover, the authors of [3] showed that there is an isomorphism

$$\mathbf{H}_\bullet(\mathrm{DRep}_V(A)^{\mathrm{GL}(V)}) \cong \mathbf{H}_\bullet(\mathrm{DRep}_V(A))^{\mathrm{GL}(V)}$$

and the derived trace map (7.1) is $\mathrm{GL}(V)$ invariant, and hence is a map

$$\mathrm{Tr} : \mathrm{HC}_\bullet(A) \rightarrow \mathbf{H}_\bullet(\mathrm{DRep}_V(A))^{\mathrm{GL}(V)}.$$

7.2. Derived non-commutative Poisson structures. The above general setting is successfully applied to the study of the derived non-commutative Poisson structures [1].

Suppose A is an associative algebra, or more generally a DG algebra. A *derived non-commutative Poisson structure* on A , following the Kontsevich–Rosenberg Principle in a naive way, is such a structure on its cofibrant resolution QA that naturally induces a DG Poisson structure on its derived representation scheme $\mathrm{DRep}_V(A)$ (or more precisely on $\mathrm{DRep}_V(A)^{\mathrm{GL}(V)}$), for all V . In [1], the authors proposed the following:

Definition 7.1 (Derived non-commutative Poisson structure; [1]). Let A be a DG algebra. A *derived non-commutative Poisson bracket* of degree n on A is a DG non-commutative Poisson bracket of degree n in the sense of Crawley-Boevey on its cofibrant resolution QA , namely, it is a differential graded Lie bracket of degree n on $(QA)_{\mathfrak{h}}$ such that

$$[\bar{a}, -] : (QA)_{\mathfrak{h}} \rightarrow (QA)_{\mathfrak{h}}$$

is induced by a derivation $d_a : QA \rightarrow QA$, which commutes with the differential.

The derived non-commutative Poisson bracket does not depend on the choice of resolutions QA up to homotopy, and hence is well defined on the homotopy category of DG algebras. The following is the main result proved in [1]:

Theorem 7.2 ([1, Theorem 9]). *Suppose A is an associative (DG) algebra equipped with a derived n -Poisson structure. Then there exists a unique graded n -Poisson algebra structure on $H_{\bullet}(\text{DRep}_V(A))^{\text{GL}(V)}$ for all V such that the derived trace map is a map of graded Lie algebras.*

8. Proof of main theorems

8.1. Proof of Theorems 1.1 and 1.2. Let A be an N -Koszul d -Calabi–Yau algebra. Then by Theorem 6.6 there is a cofibrant resolution $p : QA \xrightarrow{\cong} A$, where $QA = \Omega_{\infty}(A^i)$ and A^i is a cyclic A_{∞} coalgebra of degree $-d$. Next, by Lemma 2.3 and Corollary 2.5, we get

$$\text{HC}_{\bullet}(A) \cong \text{HC}_{\bullet}(A^i) \cong H_{\bullet}[\bar{R}_{\mathfrak{h}}],$$

where $R = \Omega_{\infty}(\tilde{A}^i)$. And by Corollary 2.10, one has

$$\text{HH}_{\bullet}(A) \cong H_{\bullet}[\Omega_{R, \mathfrak{h}}^1].$$

Proof of Theorem 1.1. By Theorem 5.3, there is a bracket $\{-, -\}$ making $H_{\bullet}[\bar{R}_{\mathfrak{h}}]$ a $(2-d)$ -Lie algebra and a bracket $\{-, -\}_{\Omega_{R, \mathfrak{h}}^1}$ making $H_{\bullet}[\Omega_{R, \mathfrak{h}}^1]$ a Lie module over it. Since, the first one can be identified with $\text{HC}_{\bullet}(A)$ and the second one with $\text{HH}_{\bullet}(A)$, we have proved this theorem. \square

Proof of Theorem 1.2. We recall (see (2.2)) that

$$\text{CC}_{\bullet}(C) = \text{Ker}(1 - T) \cong \text{Coker}(1 - T).$$

Hence from the 2-periodic complex (2.9) one obtains the following exact sequence of complexes

$$0 \longrightarrow \text{CC}_{\bullet}(C) \xrightarrow{i} \Omega_{R, \mathfrak{h}}^1 \xrightarrow{-\beta} \bar{R} \xrightarrow{\pi} \text{CC}_{\bullet}(C) \longrightarrow 0, \quad (8.1)$$

since $-\beta = 1 - T$. Note we can identify $CC_\bullet(C)$ with \bar{R}_1 which is a Lie module and the inclusion i is a DG Lie module homomorphism. β is a DG Lie module homomorphism by Theorem 5.4 and so is the projection π .

From (8.1) we get the Connes long exact sequence for the coalgebra C

$$\longrightarrow HH_\bullet(C) \longrightarrow HC_\bullet(C) \longrightarrow HC_{\bullet-2}(C) \longrightarrow HH_{\bullet-1}(C) \longrightarrow$$

By the above discussion the maps between homologies are Lie module homomorphisms. Finally, by Corollary 2.5 and Lemma 2.9, this sequence can be identified with the Connes long sequence for the algebra A . \square

8.2. Proof of Theorem 1.4. Let us first remind the definition of Hochschild cohomology groups.

Definition 8.1 (Hochschild cohomology). Let A be an associative algebra, and M be an A -bimodule. The *Hochschild cochain complex* $CH^\bullet(A; M)$ of A with value in M is the complex whose underlying space is

$$\bigoplus_{n \geq 0} \text{Hom}(A^{\otimes n}, M)$$

with coboundary $\delta : \text{Hom}(A^{\otimes n}, M) \rightarrow \text{Hom}(A^{\otimes n+1}, M)$ defined by

$$\begin{aligned} (\delta f)(a_0, a_1, a_2, \dots, a_n) &= a_0 f(a_1, \dots, a_n) \\ &+ \sum_{i=0}^{n-1} (-1)^{i+1} f(a_0, \dots, a_i a_{i+1}, \dots, a_n) + (-1)^n f(a_0, \dots, a_{n-1}) a_n. \end{aligned} \quad (8.2)$$

The associated cohomology is called the *Hochschild cohomology* of A with value in M , and is denoted by $HH^\bullet(A; M)$. In particular, if $M = A$, then $HH^\bullet(A; A)$ is called the *Hochschild cohomology* of A .

Definition 8.2. Let A be an associative algebra and let $CH^\bullet(A; A)$ be its Hochschild cochain complex.

- (1) The *Gerstenhaber cup product* on $CH^\bullet(A; A)$ is defined as follows:
for any $f \in CH^n(A; A)$, $g \in CH^m(A; A)$, and $a_1, \dots, a_{n+m} \in A$,

$$f \cup g(a_1, \dots, a_{n+m}) := (-1)^{nm} f(a_1, \dots, a_n) g(a_{n+1}, \dots, a_{n+m}).$$

- (2) For any $f \in CH^n(A; A)$, $g \in CH^m(A; A)$, and $a_1, \dots, a_{n+m-1} \in A$, let

$$\begin{aligned} f \circ g(a_1, \dots, a_{n+m-1}) &:= \sum_{i=1}^{n-1} (-1)^{(m-1)(i-1)} \\ &\cdot f(a_1, \dots, a_{i-1}, g(a_i, \dots, a_{i+m-1}), a_{i+m}, \dots, a_{n+m-1}). \end{aligned}$$

The Gerstenhaber bracket on $\text{CH}^\bullet(A; A)$ is defined to be

$$\{f, g\}_G := f \circ g - (-1)^{(n-1)(m-1)} g \circ f.$$

- (3) For any homogeneous elements $f \in \text{CH}^n(A; A)$ and $\alpha = (a_0, a_1, \dots, a_m) \in \text{CH}_m(A, A)$, define the cap product

$$\cap : \text{CH}_m(A; A) \times \text{CH}^n(A; A) \rightarrow \text{CH}_{m-n}(A; A)$$

by

$$\alpha \cap f := (a_0 f(a_1, \dots, a_n), a_{n+1}, \dots, a_m),$$

for $m \geq n$, and zero otherwise.

Both the Gerstenhaber product and the Gerstenhaber bracket induce a well-defined product and bracket on Hochschild cohomology $\text{HH}^\bullet(A; A)$, which makes $\text{HH}^\bullet(A; A)$ into a Gerstenhaber algebra, and the cap product makes $\text{HH}_\bullet(A)$ into an $(\text{HH}^\bullet(A), \cup)$ module. Recall that a Gerstenhaber algebra is a graded commutative algebra together with a degree -1 Lie bracket $\{-, -\}$ such that

$$a \mapsto \{a, b\}$$

are derivations with respect to the product.

Theorem 8.3 (Gerstenhaber). *Let A be an algebra. Then the Hochschild cohomology $\text{HH}^\bullet(A; A)$ of A equipped with the Gerstenhaber cup product and the Gerstenhaber bracket forms a Gerstenhaber algebra.*

Proof. For a proof, see Gerstenhaber [21, Theorems 3–5]. □

The Gerstenhaber algebra structure is even more interesting in the case of cyclic A_∞ algebra case. Tradler, in his paper [44], showed that for A_∞ algebras, one may similarly define the Gerstenhaber product and bracket on the Hochschild cochain complex, where the Gerstenhaber product is associative up to homotopy, and hence is well-defined on the cohomology level. Moreover he showed the following (see [44, Theorem 2]):

Lemma 8.4 (Tradler). *Suppose C is a cyclic A_∞ coalgebra of degree $-d$, and denote by A its dual A_∞ algebra. Then there is an isomorphism of graded vector spaces*

$$\Phi : \text{HH}_\bullet(C) \cong \text{HH}^{d-\bullet}(A).$$

Proof. Since C and hence A are locally finite dimensional, we have an isomorphism

$$\begin{aligned} \phi : C &\longrightarrow A[d] \\ c &\longmapsto \langle c, - \rangle \end{aligned}$$

of A -bimodules, from which we obtain an isomorphism of chain complexes

$$\begin{aligned} \Phi : \quad \text{CH}_\bullet(C) &\longrightarrow \text{CH}^\bullet(A) \\ (c_1, c_1, \dots, c_{n+1}) &\longmapsto \{(a_1, \dots, a_n) \mapsto (c_1, a_1) \cdots (c_n, a_n) \cdot \phi(c_{n+1})\}. \end{aligned}$$

In other words the isomorphism is given by

$$\Phi : (c_1, c_2, \dots, c_{n+1}) \longmapsto (c_1, c_2, \dots, c_n) \otimes \phi(c_n)$$

where the right hand side is viewed as an element in $\text{CH}^n(A)$ via the isomorphism

$$\text{Hom}(A^{\otimes n}, A) = (A^{\otimes n})^* \otimes A = (A^*)^{\otimes n} \otimes A = C^{\otimes n} \otimes A. \quad \square$$

Proof of Theorem 1.4. Since A^i is finite dimensional, from the above lemma any homogeneous element $f \in \text{CH}^\bullet(A^1)$ is a linear combination of elements in the form

$$(u_1, u_2, \dots, u_n) \otimes \bar{u}, \quad \text{where } u_i \in A^i, \bar{u} \in A^1.$$

For two elements, say $f = (u_1, u_2, \dots, u_n) \otimes \bar{u}$, $g = (v_1, v_2, \dots, v_m) \otimes \bar{v}$,

$$\begin{aligned} &\{f, g\}_G(x_1, x_2, \dots, x_{n+m-1}) \\ &= \sum_i (-1)^{\eta_i} f(x_1, \dots, x_{i-1}, g(x_i, \dots, x_{i+m-1}), x_{i+m}, \dots, x_{n+m-1}) \\ &- \sum_j (-1)^{(|f|-1)(|g|-1)+\eta_j} g(x_1, \dots, x_{j-1}, f(x_j, \dots, x_{j+n-1}), \\ &\hspace{15em} x_{j+n}, \dots, x_{n+m-1}) \\ &= \sum_i (-1)^{\eta_i} u_1(x_1) \cdots u_{i-1}(x_{i-1}) v_1(x_i) \cdots v_m(x_{i+m-1}) u_i(\bar{v}) u_{i+1}(x_{i+m}) \cdots \\ &\hspace{15em} \cdots u_n(x_{n+m-1}) \bar{u} \\ &- \sum_j (-1)^{\eta+\eta_j} v_1(x_1) \cdots v_{j-1}(x_{j-1}) u_1(x_j) \cdots u_n(x_{j+n-1}) v_j(\bar{u}) v_{j+1}(x_{j+n}) \cdots \\ &\hspace{15em} \cdots v_m(x_{n+m-1}) \bar{v}, \end{aligned}$$

for any homogeneous element $(x_1, x_2, \dots, x_{n+m-1}) \in (A^1)^{\otimes n+m-1}$, where $\eta_i = |g|(|x_1| + \cdots + |x_{i-1}|)$, $\eta_j = |f|(|x_1| + \cdots + |x_{j-1}|)$ and $\eta = (|f|-1)(|g|-1)$. In other words,

$$\begin{aligned} \{f, g\}_G &= \sum_i (-1)^{\eta_i} u_i(\bar{v}) (u_1, \dots, u_{i-1}, v_1, \dots, v_m, u_{i+1}, \dots, u_n) \otimes \bar{u} \\ &- \sum_j (-1)^{\eta+\eta_j} v_j(\bar{u}) (v_1, \dots, v_{j-1}, u_1, \dots, u_n, v_{j+1}, \dots, v_m) \otimes \bar{v}. \quad (8.3) \end{aligned}$$

Now let $u = [u_1|u_2|\cdots|u_n]$, $v = [v_1|v_2|\cdots|v_m] \in \mathbf{\Omega}_\infty(A^i)$ be two homogeneous elements where u_i 's and v_j 's are generically different from each other.

Viewing them as elements in $\text{CH}_\bullet(A^i)$ via the embedding (2.1) or (2.15), we have

$$B(u) = \sum_{i=1}^n (-1)^{\varepsilon_i} (u_{i+1}, \dots, u_n, u_1, \dots, u_i),$$

$$B(v) = \sum_{j=1}^m (-1)^{\varepsilon_j} (v_{j+1}, \dots, v_m, v_1, \dots, v_j),$$

where $\varepsilon_i = (|u_1| + \dots + |u_i|)(|u_{i+1}| + \dots + |u_n|)$ and $\varepsilon_j = (|v_1| + \dots + |v_j|)(|v_{j+1}| + \dots + |v_m|)$. Applying the above lemma to A^i , we have

$$\Phi \circ B(u) = \sum_{i=1}^n (-1)^{\varepsilon_i} (u_{i+1}, \dots, u_n, u_1, \dots, u_{i-1}) \otimes \bar{u}_i, \tag{8.4}$$

$$\Phi \circ B(v) = \sum_{j=1}^m (-1)^{\varepsilon_j} (v_{j+1}, \dots, v_m, v_1, \dots, v_{j-1}) \otimes \bar{v}_j, \tag{8.5}$$

where $\bar{u}_i = \phi(u_i)$ and $\bar{v}_j = \phi(v_j)$. Thus for two summands in $\Phi \circ B(u)$ and $\Phi \circ B(v)$ respectively, say

$$(u_1, \dots, u_{n-1}) \otimes \bar{u}_n, \quad (v_1, \dots, v_{m-1}) \otimes \bar{v}_m,$$

by (8.3) we have

$$\begin{aligned} & \{(u_1, \dots, u_{n-1}) \otimes \bar{u}_n, (v_1, \dots, v_{m-1}) \otimes \bar{v}_m\}_G \tag{8.6} \\ &= \sum_i (-1)^{\mu_i} \langle sv_m, su_i \rangle (u_1, \dots, u_{i-1}, v_1, \dots, v_{m-1}, u_{i+1}, \dots, u_{n-1}) \otimes \bar{u}_n \\ & \quad - \sum_j (-1)^{\nu_j} \langle su_n, sv_j \rangle (v_1, \dots, v_{j-1}, u_1, \dots, u_{n-1}, v_{j+1}, \dots, v_{m-1}) \otimes \bar{v}_m, \end{aligned}$$

where $\mu_i = |v_m| + (|v| + d)(|u_{i+1}| + \dots + |u_n| + d)$, $\nu_j = |u_n| + (|u| + d)(|v_1| + \dots + |v_{j-1}|)$, and $\langle -, - \rangle$ is the graded symmetric pairing on C .

On the other hand, by (4.7) we have

$$\begin{aligned} \Phi \circ B\{(u_1, \dots, u_n), (v_1, \dots, v_m)\}_{\text{DNCP}} &= \Phi \circ B \left(\sum_{i,j} (-1)^{|u_i| + \epsilon} \langle su_i, sv_j \rangle \right. \\ & \quad \left. \cdot (v_1, \dots, v_{j-1}, u_{i+1}, \dots, u_n, u_1, \dots, u_{i-1}, v_{j+1}, \dots, v_m) \right) \tag{8.7} \end{aligned}$$

where ϵ is given as in (4.7). From this one sees that (8.6) are summands in (8.7). For other summands in $\Phi \circ B(u)$ and $\Phi \circ B(v)$, by the same argument, their Gerstenhaber brackets also lie in (8.7) as summands.

Also, from (8.6) one sees that in the expression

$$\{\Phi \circ B(u), \Phi \circ B(v)\}_G$$

none of the summands literally cancels with any other, and therefore to show

$$\{\Phi \circ B(u), \Phi \circ B(v)\}_G = \Phi \circ B\{u, v\}_{\text{DNCP}},$$

it is enough to show both sides have the same number of summands.

In fact, on one hand, from (8.4) and (8.5) we see that $\Phi \circ B(u)$ and $\Phi \circ B(v)$ have n and m summands respectively, and therefore

$$\{\Phi \circ B(u), \Phi \circ B(v)\}_G$$

has $mn(m+n-2)$ summands. On the other hand, from (8.7) we know that $\{u, v\}_{\text{DNCP}}$ has mn summands, and each summand contains $m+n-2$ components, and therefore $\Phi \circ B\{u, v\}_{\text{DNCP}}$ has $mn(m+n-2)$ summands. The numbers of summands are equal, which proves the theorem. \square

Let us now recall a theorem of Keller [31] (see also Herscovich [28]):

Theorem 8.5 (Keller). *Let A be an N -Koszul algebra, and denote by $A^!$ its Koszul dual algebra. Then we have isomorphism*

$$\Xi : \text{HH}^\bullet(A^!) \rightarrow \text{HH}^\bullet(A)$$

of Gerstenhaber algebras.

Proof. For $N = 2$, this is proved in Keller [31, Theorem 3.5]. For the general case, it is proved by Herscovich in [28]. \square

Corollary 8.6. *Let A be an N -Koszul Calabi–Yau algebra. Then $\Xi \circ \Phi \circ B$ maps the Leibniz–Loday bracket $\{-, -\}_{\text{DNCP}}$ on $\tilde{R} = \Omega_\infty(\tilde{A}^!)$ to the Gerstenhaber bracket $\{-, -\}_G$ on $\text{HH}^\bullet(A)$.*

Proof. The claim follows from a combination of Theorem 1.4 and Theorem 8.5 of Keller above. \square

Roughly speaking, the above corollary says that the derived non-commutative Poisson bracket of a Koszul Calabi–Yau algebra is naturally mapped to the Gerstenhaber bracket on its Hochschild cohomology.

8.3. Proof of Theorem 1.6. Before we proceed to the proof of Theorem 1.6, let us recall a Batalin–Vilkovisky algebra structure on the Hochschild cohomology $\mathrm{HH}^\bullet(A)$. In order to do that, let us first observe that, for an associative algebra A , we have the following two structures:

- there is the Connes cyclic operator $B : \mathrm{HH}_\bullet(A) \rightarrow \mathrm{HH}_{\bullet+1}(A)$ whose square is zero; and
- there is a Gerstenhaber algebra structure on $(\mathrm{HH}^\bullet(A), \cup)$, and $\mathrm{HH}_\bullet(A)$ is a module over $\mathrm{HH}^\bullet(A)$. Denote the action by \cap .

For a d -Calabi–Yau algebra A , these two structures are perfectly matched. Note that there is an element $\omega \in \mathrm{HH}_d(A)$ such that the following map

$$\Psi : \begin{array}{ccc} \mathrm{HH}^\bullet(A) & \xrightarrow{\cong} & \mathrm{HH}_{d-\bullet}(A) \\ f & \longmapsto & \iota_f \omega \end{array} \quad (8.8)$$

is an isomorphism, called the *non-commutative Van den Bergh–Poincaré duality*; for a proof of this result see [17, Proposition 5.5].

Definition 8.7 (Batalin–Vilkovisky algebra). Let (V, \bullet) be a graded commutative algebra. A *Batalin–Vilkovisky operator* on V is a degree -1 differential operator

$$\Delta : V_i \rightarrow V_{i-1}, \quad \text{for all } i$$

such that the *deviation from being a derivation*

$$\{a, b\} := (-1)^{|a|}(\Delta(a \bullet b) - \Delta(a) \bullet b - (-1)^{|a|} a \bullet \Delta(b)), \quad \text{for all } a, b \in V \quad (8.9)$$

is a derivation for each component, i.e.

$$\{a, b \bullet c\} = \{a, b\} \bullet c + (-1)^{|b|(|a|-1)} b \bullet \{a, c\}, \quad \text{for all } a, b, c \in V.$$

The triple (V, \bullet, Δ) is called a *Batalin–Vilkovisky algebra*.

If (V, \bullet, Δ) is a Batalin–Vilkovisky algebra, then by definition $(V, \bullet, \{-, -\})$, where $\{-, -\}$ is given by (8.9), is a Gerstenhaber algebra. In other words, a Batalin–Vilkovisky algebra is a special class of Gerstenhaber algebras.

Let A be a d -Calabi–Yau algebra and let

$$\Delta := \Psi^{-1} \circ B \circ \Psi : \mathrm{HH}^\bullet(A) \rightarrow \mathrm{HH}^{\bullet-1}(A),$$

we have

Theorem 8.8 (Ginzburg [23, Theorem 3.4.3]). *Suppose that A is an d -Calabi–Yau algebra. Then $(\mathrm{HH}^\bullet(A; A), \cup, \Delta)$ is a Batalin–Vilkovisky algebra. Moreover, the bracket (8.9) for this triple is exactly the classical Gerstenhaber bracket $\{-, -\}_G$.*

Recently the first author with the fourth author and G. D. Zhou proved in [10, Theorem A] that for Koszul (and more generally, N -Koszul) Calabi–Yau algebras, the isomorphism in Theorem 8.5 is in fact an isomorphism of Batalin–Vilkovisky algebras. The key point in the proof is that there is a canonical complex (the Koszul complex equipped with an appropriate differential) which computes both $\mathrm{HH}_\bullet(A)$ and $\mathrm{HH}_\bullet(A^i)$, and we have the following commutative diagram

$$\begin{array}{ccc} \mathrm{HH}^\bullet(A^i) & \xrightarrow{\Xi} & \mathrm{HH}^\bullet(A) \\ \uparrow \Phi & & \downarrow \Psi \\ \mathrm{HH}_{d-\bullet}(A^i) & \xlongequal{\quad} & \mathrm{HH}_{d-\bullet}(A). \end{array} \quad (8.10)$$

Proof of Theorem 1.6. Let $\alpha, \beta \in \mathrm{HC}_\bullet(A)$. Then

$$\begin{aligned} \{B(\alpha), B(\beta)\}_{\mathrm{dVV}} &= (-1)^{|\alpha|+d+1} B \circ \Psi(\Psi^{-1}(\pi(B(\alpha))) \cup \Psi^{-1}(\pi(B(\beta)))) \quad (8.11) \\ &= (-1)^{|\alpha|+d+1} \Psi \circ \Delta(\Psi^{-1}(\pi(B(\alpha))) \cup \Psi^{-1}(\pi(B(\beta)))) \\ &= (-1)^{|\alpha|+d+1} \Psi \circ \Delta(\Delta(\Psi^{-1}(\alpha)) \cup \Delta(\Psi^{-1}(\beta))) \\ &= (-1)^{|\alpha|+d+1+d-|\alpha|-1} \Psi \{\Delta(\Psi^{-1}(\alpha)), \Delta(\Psi^{-1}(\beta))\}_{\mathrm{G}} \\ &= \Psi \{\Psi^{-1}(B(\alpha)), \Psi^{-1}(B(\beta))\}_{\mathrm{G}} \\ &= \Psi \Xi \{\Xi^{-1}(\Psi^{-1}(B(\alpha))), \Xi^{-1}(\Psi^{-1}(B(\beta)))\}_{\mathrm{G}} \\ &= \Phi^{-1} \{\Phi \circ B(\alpha), \Phi \circ B(\beta)\}_{\mathrm{G}} \\ &= B \{\alpha, \beta\}_{\mathrm{DNCP}}, \end{aligned}$$

where the first two equalities follow definitions of $\{-, -\}_{\mathrm{dVV}}$ and Δ respectively, the third one follows from (1.6) and the definition of Δ , the fourth equality follows from applying (8.9), the fifth equality follows again from the definition of Δ , the sixth equality holds due to Theorem 8.5, the seventh equality follows from (8.10), and finally the eighth equality follows from Theorem 1.4. \square

Proof of Corollary 1.5. It follows from (8.11) that

$$B \{\alpha, \beta\}_{\mathrm{DNCP}} = (-1)^{|\alpha|+d+1} B \circ \Psi(\Psi^{-1}(B(\alpha)) \cup \Psi^{-1}(B(\beta))).$$

Since A is a graded algebra the Connes operator B is injective (see e.g. [47, Theorem 9.9.1]) and hence

$$\{\alpha, \beta\}_{\mathrm{DNCP}} = (-1)^{|\alpha|+d+1} \Psi(\Psi^{-1}(B(\alpha)) \cup \Psi^{-1}(B(\beta))).$$

Note that by (8.8) the RHS of the above identity

$$\begin{aligned} \Psi(\Psi^{-1}(B(\alpha)) \cup \Psi^{-1}(B(\beta))) &= \iota_{\Psi^{-1}(B(\alpha)) \cup \Psi^{-1}(B(\beta))} \omega \\ &= \iota_{\Psi^{-1}(B(\alpha))} \iota_{\Psi^{-1}(B(\beta))} \omega \\ &= \iota_{\Psi^{-1}(B(\alpha))} B(\beta), \end{aligned}$$

and the result follows. \square

9. Example: the polynomial algebra

Let A be the symmetric algebra $\text{Sym}(V)$ where V is a vector space of dimension $m \geq 1$ concentrated in homological degree 0. This is a quadratic Koszul, Calabi–Yau algebra of dimension m . By the well-known result of Hochschild–Kostant–Rosenberg, we can identify $\text{HH}_\bullet(A) \cong \Omega^\bullet(V)$ and $\text{HH}^\bullet(A) \cong \Theta_\bullet(V)$, where $\Omega^\bullet(V)$ is the vector spaces of differential forms and $\Theta_\bullet(V)$ is that of polyvector fields on V . Moreover, $\text{HC}_\bullet(A)$ can be identified with $\bigoplus_n \Omega^n(V)/d(\Omega^{n-1}(V))$, where d is the de Rham differential. Using Corollary 8.6 we will give an geometric description of $\{-, -\}_{\text{DNCP}}$ (to simplify the notation in the following we write $\{-, -\}_{\text{DNCP}}$ as $\{-, -\}$).

9.1. Koszul dual coalgebra. We recall that A is a quadratic Koszul algebra defined by the quadratic data (V, S) with $S \subset V \otimes V$ spanned by the vectors of the form $v \otimes u - u \otimes v$, which has a minimal semi-free resolution $R := \Omega(C)$ given by the cobar construction of the Koszul dual coalgebra $C := A^i = C(sV, s^2S)$. The coalgebra C is an exterior coalgebra whose degree n elements we denote by

$$\mu(v_1, v_2, \dots, v_n) := (sv_1 \wedge sv_2 \wedge \dots \wedge sv_n).$$

We can always assume that v_1, \dots, v_n are linearly independent. The coproduct $\Delta : C \rightarrow C \otimes C$ is given by

$$\begin{aligned} \Delta(\mu(v_1, v_2, \dots, v_n)) \\ := \sum_{p=0}^n \sum_{\sigma \in Sh_{p, n-p}} \text{sgn}(\sigma) \mu(v_{\sigma(1)}, \dots, v_{\sigma(p)}) \otimes \mu(v_{\sigma(p+1)}, \dots, v_{\sigma(n)}), \end{aligned}$$

where $Sh_{p, n-p} \subset S_n$ is the subset of $(p, n - p)$ shuffles. We denote counit in C by e . For example, we have

$$\Delta(\mu(v_1, v_2)) = e \otimes \mu(v_1, v_2) + \mu(v_1, v_2) \otimes e + \mu(v_1) \otimes \mu(v_2) - \mu(v_2) \otimes \mu(v_1).$$

We recall that a symmetric bilinear form of degree $-d$ on C is a pairing $\langle, \rangle : C \otimes C \rightarrow k[d]$ such that

$$\langle a, b \rangle = (-1)^{|a||b|} \langle b, a \rangle, \quad \text{for all } a, b \in C. \tag{9.1}$$

It is *cyclic* if

$$\langle a, b^2 \rangle \cdot b^1 = \langle a^1, b \rangle \cdot a^2, \quad \text{for all } a, b \in C, \tag{9.2}$$

where $\Delta(a) = \sum a^1 \otimes a^2$ and $\Delta(b) = \sum b^1 \otimes b^2$ using Sweedler’s notation.

9.2. Cyclic forms on C .

First we show

Proposition 9.1. *Let $\langle -, - \rangle$ be a cyclic form on C . Then*

- (1) $\langle e, \mu(v_1, v_2, \dots, v_n) \rangle = 0$ for $n \leq m - 1$.
- (2) $\langle \mu(v_1, v_2, \dots, v_n), \mu(w_1, \dots, w_l) \rangle = 0$ for $l \leq m - 1$,
if each $v_i \in \text{Span}\{w_1, \dots, w_l\}$.

Proof. (1) Since $n \leq m - 1$ we can choose $v \in V$ such that v is not in $\text{Span}\{v_1, \dots, v_n\}$. Let $a = \mu(v)$ and $b = \mu(v_1, v_2, \dots, v_n)$. Then

$$\begin{aligned}\Delta(a) &= e \otimes \mu(v) + \mu(v) \otimes e, \\ \Delta(b) &= e \otimes \mu(v_1, \dots, v_n) + \mu(v_1, \dots, v_n) \otimes e + \dots.\end{aligned}$$

Using (9.2) we get

$$\begin{aligned}\langle \mu(v), \mu(v_1, \dots, v_n) \rangle \cdot e + \langle \mu(v), e \rangle \cdot \mu(v_1, \dots, v_n) + \dots \\ = \langle \mu(v), \mu(v_1, \dots, v_n) \rangle \cdot e + \langle e, \mu(v_1, \dots, v_n) \rangle \cdot \mu(v).\end{aligned}$$

Since the left hand side does not contain the term with $\mu(v)$ we can conclude this statement.

(2) Due to linearity of the form it suffices to show it in the case when each v_i is of one w_j 's and after reordering we can also assume that $v_1 = w_1, \dots, v_n = w_n$. So we only need to show

$$\langle \mu(w_1, w_2, \dots, w_n), \mu(w_1, \dots, w_l) \rangle = 0.$$

First, we choose $v \in V$ such that $v \notin \text{Span}\{w_1, \dots, w_l\}$. Second, let $a = \mu(w_1, \dots, w_n)$ and $b = \mu(w_1, \dots, w_l, v)$. Then

$$\begin{aligned}\Delta(a) &= e \otimes \mu(w_1, \dots, w_n) + \mu(w_1, \dots, w_n) \otimes e + \dots, \\ \Delta(b) &= \dots + \mu(w_1, \dots, w_l) \otimes \mu(v) + (-1)^l \mu(v) \otimes \mu(w_1, \dots, w_l) + \dots.\end{aligned}$$

Now using (9.2) we get

$$\begin{aligned}(-1)^l \langle \mu(w_1, \dots, w_n), \mu(v) \rangle \cdot \mu(w_1, \dots, w_l) \\ + \langle \mu(w_1, \dots, w_n), \mu(w_1, \dots, w_l) \rangle \cdot \mu(v) + \dots \\ = \langle e, \mu(w_1, \dots, w_l, v) \rangle \cdot \mu(w_1, \dots, w_n) \\ + \langle \mu(w_1, \dots, w_n), \mu(w_1, \dots, w_l, v) \rangle \cdot e + \dots\end{aligned}$$

Left side has no term with $\mu(v)$ while right side has only one term with $\mu(v)$ and therefore coefficient in front of this term must vanish. \square

Next, we prove the following statement.

Proposition 9.2. *If $v \notin \text{Span}\{v_1, \dots, v_n, w_1, \dots, w_l\}$ then*

$$\langle \mu(v_1, v_2, \dots, v_n, v), \mu(w_1, \dots, w_l) \rangle = \langle \mu(v_1, v_2, \dots, v_n), \mu(v, w_1, \dots, w_l) \rangle.$$

Proof. For $a = \mu(v_1, v_2, \dots, v_n, v)$ and $b = \mu(v, w_1, \dots, w_l)$:

$$\begin{aligned} \Delta(a) &= \mu(v_1, \dots, v_n) \otimes \mu(v) + (-1)^n \mu(v) \otimes \mu(v_1, \dots, v_n) + \dots \\ \Delta(b) &= \mu(v) \otimes \mu(w_1, \dots, w_l) + (-1)^l \mu(w_1, \dots, w_l) \otimes \mu(v) + \dots \end{aligned}$$

Since $v \notin \text{Span}\{v_1, \dots, v_n, w_1, \dots, w_l\}$, the above terms in coproducts of a and b are the only terms containing $\mu(v)$. By (9.2) we have

$$\begin{aligned} \langle \mu(v_1, v_2, \dots, v_n, v), \mu(w_1, \dots, w_l) \rangle \mu(v) + \dots \\ = \langle \mu(v_1, v_2, \dots, v_n), \mu(v, w_1, \dots, w_l) \rangle \mu(v) + \dots \end{aligned}$$

This finishes our proof. \square

The next corollary follows immediately from Propositions 9.1 and 9.2.

Corollary 9.3. *We have*

$$\langle \mu(v_1, v_2, \dots, v_n), \mu(w_1, \dots, w_l) \rangle = 0$$

unless $W := \text{Span}\{v_1, v_2, \dots, v_n, w_1, \dots, w_l\} = V$.

Proof. Let $W \neq V$. First, we consider the case when none of v_i 's in W . Then using Proposition 9.2 we obtain

$$\langle \mu(v_1, v_2, \dots, v_n), \mu(w_1, \dots, w_l) \rangle = \langle e, \mu(v_1, v_2, \dots, v_n, w_1, \dots, w_l) \rangle.$$

The RHS is 0 by Proposition 9.1 part (1).

Second, we can assume that $v_i \in W$ for all $i = 1, \dots, n$. Otherwise, we can repeatedly use Proposition 9.2 to move those v_i 's which are not in W to the right. Finally, by Proposition 9.1 part (2), we have vanishing of the bilinear form in this case. \square

The above results can be summarized into the following statement

Proposition 9.4. *Let V be an m -dimensional vector space and let $A := \text{Sym}(V)$. Then the Koszul dual coalgebra C has a unique cyclic form and it is of degree $-m$. This form is completely determined by its value $\langle e, \mu(v_1, v_2, \dots, v_m) \rangle$, where $\{v_1, v_2, \dots, v_m\}$ is a basis for V .*

9.3. Lie bracket on $\mathbf{HC}_\bullet(A)$. We have an isomorphism $\Psi : \Theta_k(V) \rightarrow \Omega^{m-k}(V)$ such that $\Psi(\xi) = \iota_\xi \omega$. The space $\Theta_*(V)$ equipped with BV operator $\Delta : \Theta_*(V) \rightarrow \Theta_{*-1}(V)$ such that the following diagram commutes:

$$\begin{array}{ccc} \Theta_k(V) & \xrightarrow{\Delta} & \Theta_{k-1}(V) \\ \downarrow \Psi & & \downarrow \Psi \\ \Omega^{n-k}(V) & \xrightarrow{d} & \Omega^{m-k+1}(V) \end{array}$$

By Corollary 8.6, the bracket $\{-, -\}$ on $\Omega^*(V)$ is given by

$$d\{\alpha, \beta\} = \Psi\{\Psi^{-1}(d\alpha), \Psi^{-1}(d\beta)\}_G$$

where $\alpha, \beta \in \Omega^*(V)$ and $\{-, -\}_G$ is the Gerstenhaber bracket of poly-vector fields.

Lemma 9.5. For given forms α and β , their bracket is given by

$$d\{\alpha, \beta\} = (-1)^{m-|\alpha|-1} d\iota_\xi d\beta = (-1)^{(m-|\alpha|-1)(m-|\beta|)} d\iota_\eta d\alpha$$

where $\Psi^{-1}(d\alpha) = \xi$ and $\Psi^{-1}(d\beta) = \eta$.

Proof. Recall for $a, b \in \Theta_*(V)$ their Gerstenhaber bracket can be expressed in terms of Δ

$$\{a, b\}_G = (-1)^{|a|} (\Delta(a \wedge b) - \Delta(a) \wedge b - (-1)^{|a|} a \wedge \Delta(b)).$$

Hence using commutativity above diagram we have

$$\{\Psi^{-1}(d\alpha), \Psi^{-1}(d\beta)\}_G = (-1)^{m-|\alpha|-1} \Delta(\Psi^{-1}(d\alpha) \wedge \Psi^{-1}(d\beta)),$$

and applying Ψ we have

$$\begin{aligned} \Psi\{\Psi^{-1}(d\alpha), \Psi^{-1}(d\beta)\}_G &= (-1)^{m-|\alpha|-1} \Psi \Delta(\Psi^{-1}(d\alpha) \wedge \Psi^{-1}(d\beta)) \\ &= (-1)^{m-|\alpha|-1} d \circ \Psi(\Psi^{-1}(d\alpha) \wedge \Psi^{-1}(d\beta)). \end{aligned}$$

Now using the well-known formula

$$\iota_{\xi \wedge \eta} \omega = \iota_\xi (\iota_\eta \omega) = (-1)^{|\xi||\eta|} \iota_{\eta \wedge \xi} \omega$$

we obtain

$$\begin{aligned} \Psi(\Psi^{-1}(d\alpha) \wedge \Psi^{-1}(d\beta)) &= \iota_{\Psi^{-1}(d\alpha) \wedge \Psi^{-1}(d\beta)} \omega \\ &= \iota_{\Psi^{-1}(d\alpha)} (\iota_{\Psi^{-1}(d\beta)} \omega) = \iota_{\Psi^{-1}(d\alpha)} d\beta, \end{aligned}$$

from which the result immediately follows. \square

Since $\mathrm{HH}_\bullet(A) \cong \Omega^\bullet(V)$ and $\mathrm{HC}_\bullet(A) \cong \bigoplus_n \Omega^n(V)/d(\Omega^{n-1}(V))$, we have established

Corollary 9.6. *The Lie bracket on $\mathrm{HC}_\bullet(A)$ and the Lie module structure on $\mathrm{HH}_\bullet(A)$ in Theorem 1.1 for $A = \mathrm{Sym}(V)$ is given as follows. For $\alpha, \beta \in \mathrm{HC}_\bullet(A)$ or $\beta \in \mathrm{HH}_\bullet(A)$, we have*

$$\{\alpha, \beta\} = (-1)^{(m-|\alpha|-1)(m-|\beta|)} \iota_\eta d\alpha,$$

where $\eta = \Psi^{-1}(d\beta)$.

Remark 9.7. One can directly check that $\{-, -\}$ is not a Lie bracket on $\Omega^*(V)$, since the Jacobi identity fails. Thus $\{-, -\}$ defines only a Lie module structure on $\mathrm{HH}_\bullet(A)$ not Lie algebra.

More precisely, let

$$\alpha = x^2 y z dx, \quad \beta = x y z dy, \quad \gamma = x z dz,$$

and $\omega = dx \wedge dy \wedge dz$, then

$$\begin{aligned} \Psi^{-1}(d\alpha) &= -x^2 z \frac{\partial}{\partial z} + x^2 y \frac{\partial}{\partial y}, & \Psi^{-1}(d\beta) &= y z \frac{\partial}{\partial z} - x y \frac{\partial}{\partial x}, \\ \Psi^{-1}(d\gamma) &= -z \frac{\partial}{\partial y}, \end{aligned}$$

and we have

$$\begin{aligned} \{\{\alpha, \beta\}, \gamma\} &= x^2 y z^2 dx - x^3 y z dz, \\ \{\alpha, \{\beta, \gamma\}\} &= -x^3 z^2 dy - x^3 y z dz, \\ \{\beta, \{\alpha, \gamma\}\} &= 2x^2 y z^2 dx + 2x^3 y z dz. \end{aligned}$$

Thus we have

$$\begin{aligned} \{\{\alpha, \beta\}, \gamma\} - \{\alpha, \{\beta, \gamma\}\} + \{\beta, \{\alpha, \gamma\}\} &= 3x^2 y z^2 dx + x^3 z^2 dy + 2x^3 y z dz \\ &= d(x^3 y z^2), \end{aligned}$$

which is not zero, and therefore $\mathrm{HH}_\bullet(A)$ with $\{-, -\}$ above does not form a Lie algebra.

References

- [1] Y. Berest, X. Chen, F. Eshmatov, and A. Ramadoss, Non-commutative Poisson structures, derived representation schemes and Calabi–Yau algebras, in *Mathematical aspects of quantization*, 219–246, Contemp. Math., 583, Amer. Math. Soc., Providence, RI, 2012. Zbl 1316.17015 MR 3013096

- [2] Y. Berest, G. Felder, and A. Ramadoss, Derived representation schemes and non-commutative geometry, in *Expository lectures on representation theory*, 113–162, Contemp. Math., 607, Amer. Math. Soc., Providence, RI, 2014. Zbl 1333.16014 MR 3204869
- [3] Y. Berest, G. Khachatryan, and A. Ramadoss, Derived representation schemes and cyclic homology, *Adv. Math.*, **245** (2013), 625–689. Zbl 1291.14006 MR 3084440
- [4] R. Berger, Koszulity for nonquadratic algebras, *J. Algebra*, **239** (2001), 705–734. Zbl 1035.16023 MR 1832913
- [5] R. Berger and N. Marconnet, Koszul and Gorenstein properties for homogeneous algebras, *Algebr. Represent. Theory*, **9** (2006), 67–97. Zbl 1125.16017 MR 2233117
- [6] R. Berger and R. Taillefer, Poincaré-Birkhoff-Witt deformations of Calabi–Yau algebras, *J. Noncommut. Geom.*, **1** (2007), 241–270. Zbl 1161.16022 MR 2308306
- [7] R. Bocklandt, Graded Calabi–Yau algebras of dimension 3, with an appendix “The signs of Serre functor” by M. Van den Bergh, *J. Pure Appl. Algebra*, **212** (2008), 14–32. Zbl 1132.16017 MR 2355031
- [8] T. Bridgeland and D. Stern, Helices on del Pezzo surfaces and tilting Calabi–Yau algebras, *Adv. Math.*, **224** (2010), 1672–1716. Zbl 1193.14022 MR 2646308
- [9] X. Chen, Y. Chen, A. Eshmatov, and F. Eshmatov, The shifted Poisson structure on derived representation schemes of Koszul Calabi–Yau algebras, *in preparation*.
- [10] X. Chen, S. Yang and G. D. Zhou, Batalin–Vilkovisky algebras and the non-commutative Poincaré duality of Koszul Calabi–Yau algebras, *J. Pure Appl. Algebra*, **220** (2016), no. 7, 2500–2532. Zbl 06546716 MR 3457981
- [11] A. Connes and M. Dubois-Violette, Yang–Mills Algebra, *Lett. Math. Phys.*, **61** (2002), 149–158. Zbl 1028.53025 MR 1936574
- [12] W. Crawley-Boevey, Poisson structures on moduli spaces of representations, *J. Algebra*, **325** (2011), 205–215. Zbl 1255.17012 MR 2745537
- [13] W. Crawley-Boevey, P. Etingof, and V. Ginzburg, Non-commutative geometry and quiver algebras, *Adv. Math.*, **209** (2007), 274–336. Zbl 1111.53066 MR 2294224
- [14] J. Cuntz and D. Quillen, Algebra extensions and nonsingularity, *J. Amer. Math. Soc.*, **8** (1995), 251–289. Zbl 0838.19001 MR 1303029
- [15] J. Cuntz and D. Quillen, Operators on non-commutative differential forms and cyclic homology, in *Geometry, topology, & physics for Raoul Bott*, 77–111, Conf. Proc. Lecture Notes Geom. Topology, IV, Int. Press, Cambridge, MA, 1995. Zbl 0865.18009 MR 1358613
- [16] B. Davison, Superpotential algebras and manifolds, *Adv. Math.*, **231** (2012), 879–912. Zbl 1278.53083 MR 2955196
- [17] L. de Thanhoffer de Völcsey and M. Van den Bergh, Calabi–Yau Deformations and Negative Cyclic Homology, to appear in *J. Noncommut. Geom.* arXiv:1201.1520
- [18] V. Dotsenko and B. Vallette, Higher Koszul duality for associative algebras, *Glasg. Math. J.*, **55** (2013), 55–74. Zbl 1284.18026 MR 3110804
- [19] W. G. Dwyer and J. Spalinski, Homotopy theories and model categories, in *Handbook of Algebraic Topology*, 73–126, Elsevier, 1995. Zbl 0869.55018 MR 1361887

- [20] B. Feigin and B. Tsygan, Additive K-theory and crystalline cohomology, *Funct. Anal. Appl.*, **19** (1985), 124–132. Zbl 0585.18007 MR 800920
- [21] M. Gerstenhaber, The cohomology structure of an associative ring, *Ann. of Math.*, **78** (1963), 267–288. Zbl 0131.27302 MR 161898
- [22] V. Ginzburg, Non-commutative symplectic geometry, quiver varieties, and operads, *Math. Res. Lett.*, **8** (2001), 377–400. Zbl 1113.17306 MR 1839485
- [23] V. Ginzburg, Calabi–Yau algebras, *preprint*, 2006. arXiv:math/0612139
- [24] W. Greub, S. Halperin, and R. Vanstone, *Connections, curvature, and cohomology*, Vol. III, Academic Press, 1976. Zbl 0372.57001 MR 400275
- [25] J.-W. He and D.-M. Lu, Higher Koszul algebras and A-infinity algebras, *J. Algebra*, **293** (2005), 335–362. Zbl 1143.16027 MR 2172343
- [26] J.-W. He, F. Van Oystaeyen, and Y. Zhang, Cocommutative Calabi–Yau Hopf algebras and deformations, *J. Algebra*, **324** (2010), 1921–1939. Zbl 1213.16026 MR 2678828
- [27] J.-W. He, F. Van Oystaeyen, and Y. Zhang, Hopf algebra actions on differential graded algebras and applications, *Bulletin of Belgian Math. Soc.*, **18** (2011), 99–111. Zbl 1228.16011 MR 2809906
- [28] E. Herscovich, Hochschild (co)homology and Koszul duality, 2015. arXiv:1405.2247
- [29] E. Herscovich and A. Solotar, Hochschild and cyclic homology of Yang–Mills algebras, *J. Reine Angew. Math.*, **665** (2012), 73–156. Zbl 1305.17018 MR 2908741
- [30] J. D. S. Jones and J. McCleary, Hochschild homology, cyclic homology, and the cobar construction, in *Adams Memorial Symposium on Algebraic Topology, I (Manchester, 1990)*, 53–65, London Math. Soc. Lecture Note Ser., 175, Cambridge Univ. Press, Cambridge, 1992. Zbl 0752.55003 MR 1170570
- [31] B. Keller, *Derived invariance of higher structures on the Hochschild complex*. Available at <http://www.math.jussieu.fr/~keller/publ/index.html>
- [32] B. Keller, Deformed Calabi–Yau completions. With an appendix by Michel Van den Bergh, *J. Reine Angew. Math.*, **654** (2011), 125–180. Zbl 1220.18012 MR 2795754
- [33] M. Kontsevich, Feynman diagrams and low-dimensional topology, in *First European Congress of Mathematics, Vol. II (Paris, 1992)*, 97–121, Progr. Math., 120, Birkhäuser, Basel, 1994. Zbl 0872.57001 MR 1341841
- [34] M. Kontsevich and A. Rosenberg, Non-commutative smooth spaces, in *The Gelfand Mathematical Seminars 1996–1999*, 85–108, Gelfand Math. Sem., Birkhäuser Boston, Boston, MA, 2000. Zbl 1003.14001 MR 1731635
- [35] J.-L. Loday, *Cyclic homology*, 2nd edition, Grundle. Math. Wiss, 301, Springer-Verlag, Berlin, 1998. Zbl 0885.18007 MR 1600246
- [36] J.-L. Loday and B. Vallette, *Algebraic operads*, Grundle. Math. Wiss, 346, Springer-Verlag, 2012. Zbl 1260.18001 MR 2954392
- [37] D.-M. Lu, J. H. Palmieri, Q.-S. Wu, and J. J. Zhang, Regular algebras of dimension 4 and their A_∞ -Ext-algebras, *Duke Math. J.*, **137** (2007), 537–584. Zbl 1193.16014 MR 2309153
- [38] D.-M. Lu, J. H. Palmieri, Q.-S. Wu, and J. J. Zhang, A_∞ structure on Ext-algebras, *J. Pure Appl. Algebra*, **213** (2009), 2017–2037. Zbl 1231.16008 MR 2533303

- [39] A. Prouté, A_∞ -structures: Modèles Minimaux de Baues-Lemaire et Kadeishvili et Homologie des Fibrations, reprint of the 1986 original, with a preface to the reprint by Jean-Louis Loday, *Repr. Theory Appl. Categ.*, (2011), no. 21, 1–99. Zbl 1245.55007 MR 2844537
- [40] D. Quillen, *Homotopical algebra*, Lecture Notes in Mathematics, 43, Springer-Verlag, Berlin-New York, 1967. Zbl 0168.20903 MR 223432
- [41] D. Quillen, Algebra cochains and cyclic cohomology, *Inst. Hautes Études Sci. Publ. Math.*, **68** (1989), 139–174. Zbl 0696.16021 MR 1001452
- [42] S. P. Smith, Some finite dimensional algebras related to elliptic curves, in *Representation Theory of Algebras and Related Topics (Mexico City, 1994)*, 315–348, CMS Conf. Proc., 19, Amer. Math. Soc., Providence, 1996. Zbl 0856.16009 MR 1388568
- [43] S. P. Smith and J. T. Stafford, Regularity of the four dimensional Sklyanin algebra, *Compositio Math.*, **83** (1992), 259–289. Zbl 0758.16001 MR 1175941
- [44] T. Tradler, The Batalin–Vilkovisky algebra on Hochschild cohomology induced by infinity inner products, *Ann. Inst. Fourier (Grenoble)*, **58** (2008), 2351–2379. Zbl 1218.16004 MR 2498354
- [45] M. Van den Bergh, Double Poisson algebras, *Trans. Amer. Math. Soc.*, **360** (2008), 5711–5769. Zbl 1157.53046 MR 2425689
- [46] M. Van den Bergh, Calabi–Yau algebras and superpotentials, *Sel. Math. New Ser.*, **21** (2015), no. 2, 555–603. Zbl 06441727 MR 3338683
- [47] C. Weibel, *An introduction to homological algebra*, Cambridge Studies in Adv. Math., 38, Cambridge University Press, 1994. Zbl 0797.18001 MR 1269324
- [48] Q.-S. Wu and C. Zhu, Skew group algebras of Calabi–Yau algebras, *J. Algebra*, **340** (2011), 53–76. Zbl 1248.16010 MR 2813562

Received 28 May, 2015

X. Chen, Department of Mathematics, Sichuan University,
Chengdu, Sichuan Province 610064, P. R. China

E-mail: xjchen@scu.edu.cn

A. Eshmatov, Department of Mathematics, University of Western Ontario,
London, Ontario N6A 5B7, Canada

E-mail: aeshmato@uwo.ca

F. Eshmatov, Department of Mathematics, Sichuan University,
Chengdu, Sichuan Province 610064, P. R. China

E-mail: olimjon55@hotmail.com

S. Yang, Department of Mathematics, Sichuan University,
Chengdu, Sichuan Province 610064, P. R. China

E-mail: syang.math@gmail.com