

An equivariant Poincaré duality for proper cocompact actions by matrix groups

Hao Guo and Varghese Mathai

Abstract. Let G be a linear Lie group acting properly on a G - spin^c manifold M with compact quotient. We give a short proof that Poincaré duality holds between G -equivariant K -theory of M , defined using finite-dimensional G -vector bundles, and G -equivariant K -homology of M , defined through the geometric model of Baum and Douglas.

1. Introduction

Poincaré duality in K -theory asserts that the K -theory group of a closed spin^c manifold is naturally isomorphic to its K -homology group via cap product with the fundamental class in K -homology. This class can be represented geometrically by the spin^c -Dirac operator. More generally, if a compact Lie group acts on the manifold preserving the spin^c structure, the analogous map implements Poincaré duality between the equivariant versions of K -theory and K -homology. In the case when the Lie group is non-compact but has finite component group, induction on K -theory and K -homology allows one to establish the analogous result [7] for proper actions. The observation underlying Poincaré duality in all of these cases is that there exist enough equivariant vector bundles with which to pair the fundamental class.

In contrast, Phillips [14] showed through a counter-example with a non-linear group that, for proper actions by a general Lie group G on a manifold X with compact quotient space, finite-dimensional vector bundles do not exhaust the G -equivariant K -theory of X , and that it is necessary to introduce infinite-dimensional vector bundles into the description of K -theory (see also [10]).

One case in which finite-dimensional bundles are sufficient is when the group G is linear (see [13]), owing to the key fact that in this case every G -equivariant vector bundle over X is a direct summand of a G -equivariantly trivial bundle, i.e. one that is isomorphic to $X \times V$ for some finite-dimensional representation of G on V .

Motivated by this, we give a short proof of Poincaré duality in this setting. That is, we show that the natural map from G -equivariant K -theory to G -equivariant K -homology (which for us means Baum–Douglas’ geometric K -homology [2]), given by pairing with

the fundamental class, is an isomorphism for proper cocompact G -spin^c manifolds where G is a matrix group.

Theorem 1.1. *Suppose that a linear Lie group G acts properly and cocompactly on a G -equivariantly spin^c manifold X . Then there is a natural isomorphism*

$$\psi: K_G^*(X) \rightarrow K_*^G(X), \tag{1.1}$$

where the left- and right-hand sides denote G -equivariant K -theory and geometric K -homology, respectively.

For compact Lie group actions, Theorem 1.1 is implied by the work of [5] on the isomorphism between the equivariant geometric and analytic models of K -homology. Meanwhile, by the Peter–Weyl theorem, such groups form a subclass of linear Lie groups. Consequently, Theorem 1.1 provides another approach to some of the results in [5].

While Theorem 1.1 makes no reference to the *analytic* model of K -homology [1, 9], we note that (1.1) still holds when the right-hand side is replaced by the analytic K -homology group $KK_*^G(C_0(X), \mathbb{C})$. Indeed, as a special case of Emerson–Meyer’s general second duality result in [6, Section 6], (in particular, see the first display after (1.5) in [6]), we have

$$KK_*^G(C_0(X), \mathbb{C}) \cong K_G^*(TX).$$

By the Thom isomorphism from [14, Theorem 8.11] (see also Theorem 2.6 below), we have

$$K_G^*(TX) \cong K_G^*(X).$$

Putting this together gives $K_G^*(X) \cong KK_*^G(C_0(X), \mathbb{C})$. Combined with Theorem 1.1, this gives the following corollary.

Corollary 1.2. *Suppose that a linear Lie group G acts properly and cocompactly on a G -equivariantly spin^c manifold X . Then we have an isomorphism between the G -equivariant geometric and analytic K -homology groups of X :*

$$K_*^G(X) \cong KK_*^G(C_0(X), \mathbb{C}). \tag{1.2}$$

In particular, the geometric K -homology groups K_^G are part of a G -equivariant extraordinary homology theory.*

The natural map realizing this isomorphism (1.2) is the Baum–Douglas map [2, 7], which can be described as follows. Let (M, E, f) be a G -equivariant geometric K -cycle for $K_*^G(X)$ (see Definition 2.8 below). Then M is a G -equivariantly spin^c manifold with Dirac operator D_M acting on sections of a spinor bundle S_M , E is a G -equivariant vector bundle over M , and $f: M \rightarrow X$ is a G -equivariant continuous map. The Baum–Douglas map takes

$$[M, E, f] \mapsto f_*[L^2(S_M \otimes E), \varphi, D_E(1 + D_E^2)^{-1/2}],$$

where φ is the $*$ -representation of $C_0(M)$ on $\mathcal{B}(L^2(E))$ given by pointwise multiplication, and the right-hand side is the pushforward under f of a class in $KK_*^G(C_0(M), \mathbb{C})$. Corollary 1.2 then implies that the following holds.

Corollary 1.3. *Suppose that a linear Lie group G acts properly and cocompactly on a G -equivariantly spin^c manifold X . Then for any $e \in KK_*^G(C_0(X), \mathbb{C})$, there exist a G -equivariantly cocompact spin^c manifold M and a G -equivariant continuous map $f: M \rightarrow X$ such that e is the pushforward under f of the class of a Dirac-type operator on M in $KK_*^G(C_0(M), \mathbb{C})$. More precisely, there exists a vector bundle $E \rightarrow M$ such that*

$$e = f_*[L^2(S_M \otimes E), \varphi, D_E(1 + D_E^2)^{-1/2}].$$

2. Preliminaries

We begin by recalling the definitions and facts we will need. Unless specified otherwise, G will always denote a closed subgroup of $GL(n, \mathbb{R})$ for some n . All vector bundles will be complex. For this section, let X be a locally compact proper G -space.

2.1. Equivariant K -theory

In [13], Phillips showed that the G -equivariant K -theory of the space X with G -cocompact supports can be defined in a such a way that is directly analogous to non-equivariant, compactly supported K -theory.

Definition 2.1 ([13, Definition 1.1]). A G -equivariant K -cocycle for X is a triple (E, F, t) consisting of two finite-dimensional complex G -vector bundles E and F over X and a G -equivariant bundle bundle map $t: E \rightarrow F$ whose restriction to the complement of some G -cocompact subset of X is an isomorphism. Two K -cocycles (E, F, t) and (E', F', t') are said to be equivalent if there exist finite-dimensional G -vector bundles H and H' and G -equivariant isomorphisms

$$a: E \oplus H \rightarrow E' \oplus H, \quad b: F \oplus H \rightarrow F' \oplus H'$$

such that $b_x^{-1}(t'_x \oplus \text{id})a_x = t_x \oplus \text{id}$ for all x in the complement of a G -cocompact subset of X . The set of equivalence classes $[E, F, t]$ of K -cocycles forms a semigroup under the direct sum operation, and we define the group $K_G^0(X)$ as the Grothendieck completion of the semigroup of finite-dimensional complex G -vector bundles over X .

Remark 2.2. When it is clear from context, or when X is G -cocompact, we will omit the map t from the cycle, and simply denote a class in $K_G^0(X)$ by $[E] - [F]$.

Remark 2.3. For general locally compact groups, Definition 2.1 needs to be modified to include infinite-dimensional bundles [14, Chapter 3]. For G linear, this is not necessary [13, Theorem 2.3].

Definition 2.4. For each non-negative integer i , let

$$K_G^i(X) = K_G^0(X \times \mathbb{R}^i),$$

where G acts trivially on \mathbb{R}^i .

By [13, Lemma 2.2], K_G^i satisfies Bott periodicity, so that we have a natural isomorphism $K_G^i(X) \cong K_G^{i+2}(X)$ for each i . We will use the notation

$$K_G^*(X) = K_G^0(X) \oplus K_G^1(X).$$

In addition, K_G^i are contravariant functors from the category of proper G -spaces and proper G -equivariant maps to the category of abelian groups, and form an equivariant extraordinary cohomology theory with a continuity property. In particular, Bott periodicity implies that for any G -invariant open subset $U \subseteq X$, there is a six-term exact sequence of abelian groups

$$\begin{CD} K_G^0(U) @>>> K_G^0(X) @>>> K_G^0(X \setminus U) \\ @. @. @VV \partial V \\ @. @. @VV \partial V \\ K_G^1(X \setminus U) @<<< K_G^1(X) @<<< K_G^1(U). \end{CD} \tag{2.1}$$

Here, the map $K_G^i(U) \rightarrow K_G^i(X)$ is induced by the extension-by-zero homomorphism – see Remark 2.5 below, and the boundary maps ∂ are defined as in equivariant K -theory for compact group actions [15].

Remark 2.5 (Extension-by-zero). Any inclusion of G -invariant open subsets $U_1 \hookrightarrow U_2$ induces in the obvious way an extension-by-zero $*$ -homomorphism $C_0(U_1) \rightarrow C_0(U_2)$. This extends to a $*$ -homomorphism $C_0(U_1) \rtimes G \rightarrow C_0(U_2) \rtimes G$ between crossed products. The induced map on operator K -theory, together with the identification

$$K_G^i(U_j) \cong K_i(C_0(U_j) \rtimes G),$$

gives a map

$$K_G^i(U_1) \rightarrow K_G^i(U_2).$$

To prove the Poincaré duality, we will make use of the following Thom isomorphism theorem for G -spin^c bundles:

Theorem 2.6 ([14, Theorem 8.11]). *Let E be a finite-dimensional G -equivariant spin^c vector bundle over X . Then there is a natural isomorphism*

$$T_G: K_G^i(X) \xrightarrow{\cong} K_G^{i+\dim E}(E)$$

for $i = 0, 1$, where $\dim E$ is the real dimension of E and $i + \dim E$ is taken mod 2.

2.2. The Gysin homomorphism

Theorem 2.6 can be used to give an explicit geometric description of the Gysin (pushforward) homomorphism in G -equivariant K -theory, which we will need later.

Let Y_1 and Y_2 be two G -cocompact G -spin^c manifolds and $f: Y_1 \rightarrow Y_2$ a G -equivariant continuous map. By cocompactness, both Y_1 and Y_2 contain only finitely many orbit types. Together with the fact that G is a linear group, this implies, by [11, Theorem 4.4.3], that there exists a G -equivariant embedding

$$j_{Y_1}: Y_1 \rightarrow \mathbb{R}^{2n}$$

for some n , where G is considered as a subgroup of $\text{GL}(2n, \mathbb{R})$. Let

$$i_{Y_2}: Y_2 \rightarrow Y_2 \times \mathbb{R}^{2n}$$

denote the zero section, and define the G -equivariant embedding

$$\begin{aligned} i_{Y_1}: Y_1 &\rightarrow Y_2 \times \mathbb{R}^{2n}, \\ y &\mapsto (f(y), j_{Y_1}(y)). \end{aligned}$$

Let ν_1 be the normal bundle of i_{Y_1} , which we identify with a G -invariant tubular neighbourhood U_1 of its image. Note that it follows from the two-out-of-three lemma for G -equivariant spin^c-structures (see [12, Section 3.1] and [8, Remark 2.6]), together with the assumption that Y_1 and Y_2 are G -spin^c, that ν_1 has a G -spin^c structure, and so Theorem 2.6 applies. Identifying the normal bundle with U_1 , we have the Thom isomorphism

$$T_G: K_G^i(Y_1) \xrightarrow{\cong} K_G^{i+\dim \nu_1}(U_1), \tag{2.2}$$

for $i = 0, 1$. The Gysin homomorphism

$$f_!: K_G^i(Y_1) \rightarrow K_G^{i+\dim Y_2 - \dim Y_1}(Y_2) \tag{2.3}$$

associated to f is then the composition

$$\begin{array}{ccc} K_G^i(Y_1) & \xrightarrow{T_G} & K_G^{i+\dim Y_2+2n-\dim Y_1}(U_1) \\ & & \downarrow \lambda \\ & & K_G^{i+\dim Y_2+2n-\dim Y_1}(Y_2 \times \mathbb{R}^{2n}) \xrightarrow{\cong} K_G^{i+\dim Y_2-\dim Y_1}(Y_2), \end{array}$$

where λ is induced by the extension-by-zero map associated to the inclusion of U_1 into $Y_2 \times \mathbb{R}^{2n}$ (see Remark 2.5 below), and the right horizontal isomorphism is due to Bott periodicity.

Remark 2.7. It can be seen from the above that $f_!$ depends only on the G -homotopy class of f and that the Gysin map is functorial under compositions.

2.3. Equivariant geometric K -homology

We briefly review the equivariant version of Baum and Douglas’ geometric definition of K -homology [2]; see [3–5], or [7] for more details. As before, X is a locally compact proper G -space.

Definition 2.8. A G -equivariant *geometric K -cycle* for X is a triple (M, E, f) , where

- M is a proper G -cocompact manifold with a G -equivariant spin^c -structure,
- E is a smooth G -equivariant Hermitian vector bundle over M ,
- $f: M \rightarrow X$ is a G -equivariant continuous map.

For $i = 0$ or 1 , the G -equivariant *geometric K -homology group* $K_i^G(X)$ is the abelian group generated by geometric K -cycles (M, E, f) , where $\dim M = i \pmod 2$, subject to the equivalence relation generated by the following three elementary relations.

- (i) (Direct sum – disjoint union) For two G -equivariant Hermitian vector bundles E_1 and E_2 over M and a G -equivariant continuous map $f: M \rightarrow X$,

$$(M \sqcup M, E_1 \sqcup E_2, f \sqcup f) \sim (M, E_1 \oplus E_2, f).$$

- (ii) (Bordism) Suppose two cycles (M_1, E_1, f_1) and (M_2, E_2, f_2) are *bordant*, so that there exists a G -cocompact proper G - spin^c manifold W with boundary, a smooth G -equivariant Hermitian vector bundle $E \rightarrow W$ and a continuous G -equivariant map $f: W \rightarrow X$ such that $(\partial W, E|_{\partial W}, f|_{\partial W})$ is isomorphic to

$$(M_1, E_1, f_1) \sqcup (-M_2, E_2, f_2),$$

where $-M_2$ denotes M_2 with the opposite G - spin^c structure. Then

$$(M_1, E_1, f_1) \sim (M_2, E_2, f_2).$$

- (iii) (Vector bundle modification) Let V be a G - spin^c vector bundle of real rank $2k$ over M . Upon choosing a G -invariant metric on V , let \widehat{M} be the unit sphere bundle of $(M \times \mathbb{R}) \oplus V$, where the bundle $M \times \mathbb{R}$ is equipped with the trivial G -action. Let F be the *Bott bundle* over \widehat{M} , which is fibrewise the non-trivial generator of $K^0(S^{2k})$. (See [4, Section 3] for a more detailed description.) Then

$$(M, E, f) \sim (\widehat{M}, F \otimes \pi^*(E), f \circ \pi),$$

where $\pi: \widehat{M} \rightarrow M$ is the canonical projection.

Addition in $K_i^{\text{geo}, G}(X)$ is given by

$$[M_1, E_1, f_1] + [M_2, E_2, f_2] = [M_1 \sqcup M_2, E_1 \sqcup E_2, f_1 \sqcup f_2],$$

the additive inverse of $[M, E, f]$ is its opposite $[-M, E, f]$, while the additive identity is given by the empty cycle, where $M = \emptyset$.

Remark 2.9. The above definition of classes $[M, E, f]$ continues to make sense if we replace the bundle E by a K -theory class. Indeed, if

$$e = [E_1] - [E_2] = [E'_1] - [E'_2] \in K_G^0(M),$$

then there exists a G -vector bundle F over X such that

$$E_1 \oplus E'_2 \oplus F \cong E'_1 \oplus E_2 \oplus F.$$

By Definition 2.8 (i), this means that

$$[M, E_1, f] + [M, E'_2, f] + [M, F, f] = [M, E'_1, f] + [M, E_2, f] + [M, F, f].$$

Adding the inverse of $[M, F, f]$ to both sides and rearranging shows that the class

$$[M, e, f] := [M, E_1, f] - [M, E_2, \text{id}] = [M, E'_1, \text{id}] - [M, E'_2, \text{id}]$$

is well defined.

Finally, we can describe vector bundle modification using the Gysin homomorphism (2.3). To do this, let \widehat{M} be the manifold underlying the vector bundle modification of a cycle (M, E, f) by a bundle V , as in Definition 2.8 (iii). Then \widehat{M} is the unit sphere bundle of $(M \times \mathbb{R}) \oplus V$. We will refer to the G -equivariant embedding

$$\begin{aligned} s: M &\rightarrow \widehat{M} \subseteq (M \times \mathbb{R}) \oplus V, \\ m &\mapsto (m, 1, 0) \end{aligned}$$

as the *north pole section*.

Lemma 2.10. *Let (M, E, f) be a geometric cycle for X . Let $(\widehat{M}, F \otimes \pi^*(E), f \circ \pi)$ be its modification by a G -spin^c vector bundle V of even real rank, and let $\pi: \widehat{M} \rightarrow M$ be the projection. Let $s: M \rightarrow \widehat{M}$ be the north pole section. Then*

$$(\widehat{M}, F \otimes \pi^*(E), f \circ \pi) \sim (\widehat{M}, s_! [E], f \circ \pi).$$

Proof. The proof we give is similar to that of [5, Lemma 3.5] concerning the case of compact Lie group actions; compare also the discussion following [4, Definition 6.9]. To begin, observe that the total space of V can be identified G -equivariantly with a G -invariant tubular neighbourhood U of the embedding $s: M \rightarrow \widehat{M}$. The Gysin map $s_!$ is then the composition

$$K_G^*(M) \xrightarrow{T_G} K_G^*(U) \xrightarrow{\lambda} K_G^*(\widehat{M}), \tag{2.4}$$

where T_G is the Thom isomorphism in the form (2.2), while λ is the homomorphism induced by the extension-by-zero map $C_0(U) \rightarrow C_0(\widehat{M})$. Note that T_G is essentially given via tensor product with a “Bott element”, and (2.4) admits the following geometric description. Let F be the Bott bundle over \widehat{M} , and let F_0 be the bundle defined by pulling

back the restriction $F|_M$ along π . The composition (2.4) is then given by pulling back a vector bundle over M along π and tensoring with the class $[F] - [F_0]$. On the other hand, since \widehat{M} is the boundary of the unit sphere bundle of $(M \times \mathbb{R}) \oplus W$, and the bundle F_0 is pulled back from M , the cycle $(\widehat{M}, F_0 \otimes \pi^*(E), f \circ \pi)$ is bordant to the empty cycle. Thus we have a bordism of cycles

$$(\widehat{M}, F \otimes \pi^*(E), f \circ \pi) \sim (\widehat{M}, ([F] - [F_0]) \otimes \pi^*(E), f \circ \pi),$$

whence the right-hand side is equal to $(\widehat{M}, s![E], f \circ \pi)$ by the description of the composition (2.4) given above. ■

3. Poincaré duality for geometric K -homology

In this section, we prove Theorem 1.1. For the rest of this section, let X be a proper G -spin^c manifold with X/G being compact.

We can define the following natural map between $K_G^*(X)$ and $K_*^G(X)$, which can be thought of as cap product with the fundamental K -homology class on X . For this, let S^1 be the unit circle in \mathbb{C} , and define the map

$$\begin{aligned} c: X &\rightarrow X \times S^1 \\ x &\mapsto (x, 1). \end{aligned} \tag{3.1}$$

Definition 3.1. Define $\phi: K_G^*(X) \rightarrow K_*^G(X)$ by

$$\begin{aligned} \phi: K_G^i(X) &\rightarrow K_{i+\dim X}^G(X), \\ x &\mapsto \begin{cases} [X, e, \text{id}], & \text{if } i = 0, \\ [X \times S^1, c_!(e), \text{pr}_1], & \text{if } i = 1, \end{cases} \end{aligned}$$

where $\text{pr}_1: X \times S^1$ is the projection onto the first factor, and we have used the notation from Remark 2.9.

We now show that ϕ is an isomorphism by defining explicitly a map ψ that will turn out to be its inverse.

Definition 3.2. Define the map $\psi: K_*^G(X) \rightarrow K_G^*(X)$ by

$$\begin{aligned} \psi: K_{i+\dim X}^G(X) &\rightarrow K_G^i(X), \\ [M, E, f] &\mapsto f_![E], \end{aligned}$$

for $i = 0, 1$, where $f_!$ is the Gysin homomorphism from (2.3).

Remark 3.3. Note that

$$f_![E] \in f_!(K_G^0(M)) \subseteq K_G^{\dim X - \dim M}(X) = K_G^{\dim X - (i + \dim X)}(X) = K_G^i(X),$$

so the degrees make sense.

We first need to show that ψ is well defined. For this, we will use the following lemma.

Lemma 3.4. *Let W be a G -cocompact, G -spin^c manifold-with-boundary, and let X be a G -cocompact G -spin^c manifold. Let $h: W \rightarrow X$ be a G -equivariant map, and let $i: \partial W \hookrightarrow W$ be the natural inclusion. Then the composition*

$$K_G^*(W) \xrightarrow{i^*} K_G^*(\partial W) \xrightarrow{(h|_{\partial W})!} K_G^*(X)$$

is the zero map.

Proof. We give the proof when $\dim \partial W = \dim X \pmod 2$ and show that

$$K_G^0(W) \xrightarrow{i^*} K_G^0(\partial W) \xrightarrow{(h|_{\partial W})!} K_G^0(X) \tag{3.2}$$

is the zero map; the proofs for the other cases are similar.

Let us consider the composition (3.2) upon applying the Thom isomorphism, Theorem 2.6, in the form of (2.2), and use the description of the Gysin map from (2.3).

Let j_W be a G -equivariant embedding of W into \mathbb{R}^{2n} for some n , where G is realized as a subgroup of $\text{GL}(2n, \mathbb{R})$; note that this is possible because G is assumed to be linear. Let $j_{\partial W}$ denote the restriction of j_W to ∂W . Let $i_X: X \rightarrow X \times \mathbb{R}^{2n}$ be the zero section. Define the embedding

$$\begin{aligned} i_W: W &\rightarrow X \times \mathbb{R}^{2n}, \\ w &\mapsto (h(w), j_W(w)), \end{aligned}$$

and let $i_{\partial W}$ be the restriction of i_W to ∂W . Let ν_W and $\nu_{\partial W}$ denote the respective normal bundles of the embeddings i_W and $i_{\partial W}$. We may identify these normal bundles with G -invariant tubular neighbourhoods U_W and $U_{\partial W}$ in $X \times \mathbb{R}^{2n}$, noting that in general U_W has boundary. Since the normal bundle of ∂W in W is trivial and one-dimensional, there is a natural G -equivariant identification

$$U_{\partial W} \cong \partial U_W \times (-\varepsilon, \varepsilon) \tag{3.3}$$

for some $\varepsilon > 0$. It follows from the two-out-of-three lemma for G -equivariant spin^c-structures (see [12, Section 3.1] and [8, Remark 2.6]), together with the fact that W and X are G -spin^c, that ν_W and $\nu_{\partial W}$ are G -spin^c vector bundles, and hence Theorem 2.6 applies. The resulting Thom isomorphisms for W , ∂W , and X (in the notation of (2.2)) are shown as vertical arrows in the following commutative diagram:

$$\begin{CD} K_G^0(W) @>i^*>> K_G^0(\partial W) @>(h|_{\partial W})!>> K_G^0(X) \\ @VV T_G V @VV T_G V @VV T_G V \\ K_G^1(U_W) @>T_G i^*>> K_G^0(U_{\partial W}) @>\lambda>> K_G^0(X \times \mathbb{R}^{2n}), \end{CD} \tag{3.4}$$

Proposition 3.6. *The map ϕ is injective.*

Proof. For any $e \in K_G^i(X)$, we have

$$\psi \circ \phi(e) = \begin{cases} \psi[X, e, \text{id}] = \text{id}_!(e), & \text{if } i = 0, \\ \psi[X \times S^1, c_!(e), \text{pr}_1] = (\text{pr}_1)_!(\text{id}_!(e)) = (pr_1 \circ \text{id})_!(e), & \text{if } i = 1, \end{cases}$$

which are both equal to e , where we have used functoriality of the Gysin map. Hence $\psi \circ \phi = \text{id}$, so ϕ is injective. ■

For surjectivity, we will use the Gysin homomorphism from Subsection 2.2, together with the following result, which is a special case of [5, Theorem 4.1] but applied to linear instead of compact G .

Lemma 3.7. *Let M, N, X be three G -cocompact G -spin^c manifolds, $g: N \rightarrow X$ a G -equivariant continuous map, and $f: M \rightarrow N$ a G -equivariant embedding with even codimension. Then for any G -vector bundle $E \rightarrow M$, we have*

$$[M, E, g \circ f] = [N, f![E], g] \in K_*^G(X).$$

Proof. The proof of Theorem 4.1 in [5], which was stated for compact Lie groups, goes through with no changes to our setting. ■

Proposition 3.8. *The map ϕ is surjective.*

Proof. Examining Definitions 3.1 and 3.2, one sees that $\phi \circ \psi$ is given, at the level of geometric cycles, by

$$[M, E, f] \mapsto f_![E] \mapsto \begin{cases} [X, f_![E], \text{id}], & \text{if } \dim M = \dim X \pmod 2, \\ [X \times S^1, c_!(f_![E]), \text{pr}_1], & \text{otherwise,} \end{cases}$$

where the map c was defined in (3.1). Let $i: M \rightarrow \mathbb{R}^{2n}$ be a G -equivariant embedding for some n , and let $j: X \rightarrow \mathbb{R}^{2n} \times X$ be the zero section. Upon compactifying \mathbb{R}^{2n} , f factors as

$$\begin{array}{ccc} & & S^{2n} \times X \\ & \nearrow^{i \times f} & \downarrow \text{pr}_2 \\ M & \xrightarrow{f} & X, \end{array} \tag{3.7}$$

where pr_2 is the projection onto the second factor, and j becomes an embedding $X \rightarrow S^{2n} \times X$.

Suppose first that $\dim M = \dim X \pmod 2$. Then it suffices to prove that any geometric cycle of the form (M, E, f) is equivalent to $(X, f_![E], \text{id})$. By Lemma 3.7 applied to the embedding $i \times f$, we have

$$[M, E, f] = [S^{2n} \times X, (i \times f)_![E], \text{pr}_2] \in K_*^G(X). \tag{3.8}$$

Meanwhile, Lemma 3.7 applied to j , together with functoriality of the Gysin map, yields

$$\begin{aligned} [X, f_! [E], \text{id}] &= [X, f_! [E], \text{pr}_2 \circ j] \\ &= [S^{2n} \times X, j_! (f_! [E]), \text{pr}_2] \\ &= [S^{2n} \times X, (j \circ f)_! [E], \text{pr}_2]. \end{aligned} \tag{3.9}$$

Finally, the maps $i \times f$ and $j \circ f$ are G -homotopic through

$$\begin{aligned} F: M \times I &\rightarrow \mathbb{R}^{2n} \times X \hookrightarrow S^{2n} \times X, \\ (m, t) &\mapsto ((1 - t)i(m), f(m)). \end{aligned}$$

Invariance of the Gysin map under G -homotopy now implies that (3.8) and (3.9) are equal, and we conclude.

The case of $\dim M \not\equiv \dim X \pmod{2}$ proceeds analogously as follows. Here, we need to show that any geometric cycle of the form (M, E, f) is equivalent to $(X \times S^1, c_!(f_! [E]), \text{pr}_1)$, where c was defined in (3.1). Similar to (3.8), we have

$$[M, E, f] = [S^{2n} \times X \times S^1, (i \times (c \circ f))_! [E], \text{pr}_2] \in K_*^G(X), \tag{3.10}$$

Define

$$\begin{aligned} \tilde{j}: X \times S^1 &\rightarrow S^{2n} \times X \times S^1, \\ (x, s) &\mapsto (j(x), s). \end{aligned}$$

Then

$$\begin{aligned} [X \times S^1, c_!(f_! [E]), \text{pr}_1] &= [X \times S^1, c_!(f_! [E]), \text{pr}_2 \circ \tilde{j}] \\ &= [S^{2n} \times X \times S^1, \tilde{j}_!(c_!(f_! [E])), \text{pr}_2] \\ &= [S^{2n} \times X, (\tilde{j} \circ c \circ f)_! [E], \text{pr}_2]. \end{aligned} \tag{3.11}$$

The maps $i \times (c \circ f)$ and $\tilde{j} \circ c \circ f$ are G -homotopic through

$$\begin{aligned} F: M \times I &\rightarrow \mathbb{R}^{2n} \times X \times S^1 \hookrightarrow S^{2n} \times X \times S^1, \\ (m, t) &\mapsto ((1 - t)i(m), f(m), 1), \end{aligned}$$

and the claim follows from G -homotopy invariance of the Gysin map. ■

Proposition 3.8 and equation (3.6) together imply that $\phi: K_G^i(X) \rightarrow K_{i+\dim X}^G(X)$ is an isomorphism for $i = 0, 1$, which establishes Theorem 1.1.

Acknowledgments. The authors would like to thank Peter Hochs and Hang Wang for their useful feedback on this paper.

Funding. Hao Guo was partially supported by funding from the Australian Research Council through the Discovery Project DP200100729, and partially by the National Science Foundation through the NSF DMS-2000082. Varghese Mathai was partially supported by funding from the Australian Research Council, through the Australian Laureate Fellowship FL170100020.

References

- [1] P. Baum, A. Connes, and N. Higson, Classifying space for proper actions and K -theory of group C^* -algebras. In *C^* -algebras: 1943–1993 (San Antonio, TX, 1993)*, pp. 240–291, Contemp. Math. 167, Amer. Math. Soc., Providence, RI, 1994 Zbl [0830.46061](#) MR [1292018](#)
- [2] P. Baum and R. G. Douglas, Index theory, bordism, and K -homology. In *Operator algebras and K -theory (San Francisco, Calif., 1981)*, pp. 1–31, Contemp. Math. 10, Amer. Math. Soc., Providence, RI, 1982 Zbl [0507.55004](#) MR [658506](#)
- [3] P. Baum, N. Higson, and T. Schick, On the equivalence of geometric and analytic K -homology. *Pure Appl. Math. Q.* **3** (2007), no. 1 (part 3), 1–24 Zbl [1146.19004](#) MR [2330153](#)
- [4] P. Baum, N. Higson, and T. Schick, A geometric description of equivariant K -homology for proper actions. In *Quanta of maths*, pp. 1–22, Clay Math. Proc. 11, Amer. Math. Soc., Providence, RI, 2010 Zbl [1216.19006](#) MR [2732043](#)
- [5] P. Baum, H. Oyono-Oyono, T. Schick, and M. Walter, Equivariant geometric K -homology for compact Lie group actions. *Abh. Math. Semin. Univ. Hambg.* **80** (2010), no. 2, 149–173 Zbl [1242.19006](#) MR [2734682](#)
- [6] H. Emerson and R. Meyer, Dualities in equivariant Kasparov theory. *New York J. Math.* **16** (2010), 245–313 Zbl [1244.19004](#) MR [2740579](#)
- [7] H. Guo, V. Mathai, and H. Wang, Positive scalar curvature and Poincaré duality for proper actions. *J. Noncommut. Geom.* **13** (2019), no. 4, 1381–1433 Zbl [1443.53031](#) MR [4059824](#)
- [8] P. Hochs and V. Mathai, Geometric quantization and families of inner products. *Adv. Math.* **282** (2015), 362–426 Zbl [1325.53115](#) MR [3374530](#)
- [9] G. G. Kasparov, Equivariant KK -theory and the Novikov conjecture. *Invent. Math.* **91** (1988), no. 1, 147–201 Zbl [0647.46053](#) MR [918241](#)
- [10] W. Lück and B. Oliver, The completion theorem in K -theory for proper actions of a discrete group. *Topology* **40** (2001), no. 3, 585–616 Zbl [0981.55002](#) MR [1838997](#)
- [11] R. S. Palais, On the existence of slices for actions of non-compact Lie groups. *Ann. of Math.* (2) **73** (1961), 295–323 Zbl [0103.01802](#) MR [126506](#)
- [12] M. G. Penington and R. J. Plymen, The Dirac operator and the principal series for complex semisimple Lie groups. *J. Funct. Anal.* **53** (1983), no. 3, 269–286 Zbl [0542.22013](#) MR [724030](#)
- [13] N. C. Phillips, Equivariant K -theory for proper actions and C^* -algebras. In *Index theory of elliptic operators, foliations, and operator algebras (New Orleans, LA/Indianapolis, IN, 1986)*, pp. 175–204, Contemp. Math. 70, Amer. Math. Soc., Providence, RI, 1988 Zbl [0727.46047](#) MR [948693](#)
- [14] N. C. Phillips, *Equivariant K -theory for proper actions*. Pitman Res. Notes Math. Ser. 178, Longman Scientific & Technical, Harlow; copublished in the United States with John Wiley & Sons, New York, 1989 Zbl [0684.55018](#) MR [991566](#)
- [15] G. Segal, Equivariant K -theory. *Inst. Hautes Études Sci. Publ. Math.* (1968), no. 34, 129–151 Zbl [0199.26202](#) MR [234452](#)

Received 14 December 2020.

Hao Guo

Department of Mathematics, Texas A&M University, College Station, TX 77840, USA;
haoguo@math.tamu.edu

Varghese Mathai

School of Mathematical Sciences, University of Adelaide, Adelaide, SA 5000, Australia;
mathai.varghese@adelaide.edu.au