

Non-embeddable II_1 factors resembling the hyperfinite II_1 factor

Isaac Goldbring

Abstract. We consider various statements that characterize the hyperfinite II_1 factors amongst embeddable II_1 factors in the non-embeddable situation. In particular, we show that “generically” a II_1 factor has the Jung property (which states that every embedding of itself into its ultrapower is unitarily conjugate to the diagonal embedding) if and only if it is self-tracially stable (which says that every such embedding has an approximate lifting). We prove that the enforceable factor, should it exist, has these equivalent properties. Our techniques are model-theoretic in nature. We also show how these techniques can be used to give new proofs that the hyperfinite II_1 factor has the aforementioned properties.

1. Introduction

In [10], Murray and von Neumann proved that there exists a unique (up to isomorphism) separable hyperfinite II_1 factor. This unique factor, henceforth denoted by \mathcal{R} , plays a crucial role in the theory of finite von Neumann algebras. By Connes’ seminal work in [4], we know that \mathcal{R} is also the unique separable II_1 factor possessing any of the following properties: injectivity, semidiscreteness, and amenability.

In this article, our focus will be on some statements that characterize \mathcal{R} amongst the class of separable *embeddable* II_1 factors, where a separable tracial von Neumann algebra is embeddable if it embeds into some (equivalently, any) ultrapower $\mathcal{R}^{\mathcal{U}}$ of \mathcal{R} with \mathcal{U} a nonprincipal ultrafilter on \mathbb{N} . For example, as proven by Jung in [9], any embedding of \mathcal{R} into $\mathcal{R}^{\mathcal{U}}$ is unitarily conjugate to the diagonal embedding. In [2], the authors say that a separable II_1 factor M has the *Jung property* if and only if any embedding of M into $M^{\mathcal{U}}$ is unitarily conjugate to the diagonal embedding. In [3] (see also [2, Theorem 3.1.3]), the authors show that \mathcal{R} is the unique separable embeddable II_1 factor with the Jung property.

In [1], the author defines a separable tracial von Neumann algebra M to be *self-tracially stable* if any embedding of M into $M^{\mathcal{U}}$ has an “approximate lifting.” (See the next section for a precise definition.) It is easy to see that any II_1 factor with the Jung property is self-tracially stable (see [2, Proposition 3.3.14] for a proof). It follows that \mathcal{R} is self-tracially stable. The fact that \mathcal{R} is the unique separable embeddable self-tracially stable II_1 factor is the content of [3, Theorem 2.4].

Recall that the *Connes Embedding Problem* (CEP) asks whether or not every separable¹ tracial von Neumann algebra is embeddable. As announced in the recent landmark paper [8], the Connes Embedding Problem has a negative answer. It thus makes sense to ask whether or not there are separable non-embeddable II_1 factors that have the Jung property or are self-tracially stable. (See [2, Question 3.3.12] and [1, Question 1.8] respectively.)²

Our first main result is that “generically” these are the same question. To explain this, recall that a tracial von Neumann algebra M is *existentially closed* (or e.c. for short) if the following holds: whenever M is contained in the tracial von Neumann algebra N , then there is an embedding of N into $M^{\mathcal{U}}$ that restricts to the diagonal embedding of M .³ The notion of e.c. tracial von Neumann algebras comes from model theory and has proven useful in applications of model theory to operator algebras. Much is known about the class of e.c. tracial von Neumann algebras: they must be McDuff II_1 factors with only approximately inner automorphisms (see [5] for more on this class). The generic separable tracial von Neumann algebra is e.c. in the sense that in a natural Polish topology on the space of separable tracial von Neumann algebras, the e.c. algebras form a comeager set. The notion of e.c. factor can be relativized to the class of embeddable factors, in which case \mathcal{R} is an e.c. embeddable factor.⁴

We show the following:

Theorem 1. *If M is a separable e.c. factor, then any embedding of M into itself is approximately inner.*

From this theorem, it follows fairly quickly that if M is separable, e.c., and self-tracially stable, then M has the Jung property; see Proposition 6 below. In connection with the previous theorem, it is worth noting that recently Popa and Vaes constructed a family of examples of embeddable factors all of whose endomorphisms are approximately inner [11, Theorem D].

We next turn to two model theoretic characterizations of \mathcal{R} amongst embeddable factors. We call an e.c. (embeddable) factor *enforceable* if it embeds into all other e.c. (embeddable) factors.⁵ Should the enforceable (embeddable) factor exist, it is automatically unique. In [7, Theorems 5.1 and 5.2], it is shown that \mathcal{R} is the enforceable embed-

¹Separability is not an issue here if one allows ultrafilters over larger index sets.

²We should mention that in [2, Theorem 3.2.5] it was shown that \mathcal{R} is the unique separable embeddable II_1 factor with the *generalized Jung property*, meaning that any two embeddings of itself into its ultrapower are conjugate by some (not necessarily inner) automorphism of the ultrapower; [2, Theorem 3.3.1] shows that there are non-embeddable factors with this property.

³Again, this definition makes sense for not necessarily separable factors using ultrafilters on larger index sets. Alternatively, one can give a purely syntactical, model-theoretic, definition which makes it clear that density character is irrelevant.

⁴This follows immediately from the fact that \mathcal{R} has the Jung property, but we will discuss another proof at the end of this paper.

⁵This is not the original definition given in [7], but is equivalent by the results in Section 6 of that paper.

dable factor and that the CEP is equivalent to \mathcal{R} being the enforceable factor. Due to the negative solution of the CEP, we see that \mathcal{R} is not the enforceable factor. This does not, however, preclude the existence of the enforceable factor. We view the problem of the existence of the enforceable factor to be one of the central problems in the model theory of II_1 factors, for if the enforceable factor exists, then it is a canonical object deserving of further study, whereas any proof that it does not exist yields a stronger refutation of CEP.

In this paper, we prove the following:

Theorem 2. *If the enforceable factor exists, then it has the Jung property.*

It is worth noting that by [6, Theorem 2.14], if M is a II_1 factor with the Jung property and M is *elementarily equivalent* to $M \otimes M$, then $M \cong \mathcal{R}$.⁶ Consequently, if the enforceable factor \mathcal{E} exists, then we have that \mathcal{E} is not elementarily equivalent to $\mathcal{E} \otimes \mathcal{E}$.⁷

Our final result concerns the *finite forcing companion*. A *finitely generic factor* is a particular kind of e.c. II_1 factor with the generalized Jung property. (See [7, Definition 5.3] for a precise definition. Alternatively, [7, Proposition 3.10] presents a more workable version of the notion.) These factors always exist and any two of them are elementarily equivalent; the common first-order theory of the finitely generic factors is known as the finite forcing companion, denoted by T^f . In the embeddable situation, \mathcal{R} is a finitely generic embeddable factor (see [7, Corollary 3.14]), whence the finite forcing companion is simply the complete theory of \mathcal{R} . Since \mathcal{R} embeds in any model of its theory (due to the axiomatizability of being McDuff), the fact that \mathcal{R} has the generalized Jung property implies that it is the *prime model* of its theory.⁸

It is currently unknown whether or not T^f has a prime model. However, if it does, then it is a non-embeddable factor with the Jung property:

Theorem 3. *If T^f has a prime model M , then M has the Jung property.*

In the final section, we revisit the embeddable situation and give model-theoretic proofs that \mathcal{R} has the Jung property and is self-tracially stable that might be of independent interest.

In order to keep this note fairly short, we will freely use model-theoretic language when necessary. The reader is advised to consult [2, Section 2] for a more thorough introduction.

⁶Here, two II_1 factors are elementarily equivalent if they have the same first-order theory. By the Keisler–Shelah theorem, we can equivalently say that they have isomorphic ultrapowers. The reference [6] actually only deals with strongly self-absorbing C^* -algebras, but the proof there implies the result that we mention above.

⁷By the work in [7, Remark 5.8], the failure of CEP already implies that \mathcal{E} , should it exist, could not be isomorphic to $\mathcal{E} \otimes \mathcal{E}$.

⁸In general, the prime model of a theory is a model which elementarily embeds into any other model of the theory and, if it exists, it is automatically unique.

2. Proofs of theorems

We first prove Theorem 1. In fact, the following yields an even stronger result:

Theorem 4. *Suppose that M is an e.c. factor with subalgebra N . Then any embedding of N in M is approximately unitarily conjugate to the inclusion map.*

Proof. Let $f : N \rightarrow M$ be an embedding. Let P be the HNN extension obtained from M and f ; we note that P is finite and there is a trace on P such that the inclusion $M \subseteq P$ is trace-preserving (see [12, Corollary 4.2]). In particular, there is a unitary $u \in P$ such that $uf(x)u^* = x$ for all $x \in N$. Since M is e.c., this implies that for any finite $F \subseteq N$ and $\varepsilon > 0$, there is a unitary $v \in M$ such that $\|vf(x)v^* - x\|_2 < \varepsilon$ for all $x \in F$, as desired. ■

Remark 5 (For the model theorists). Theorem 4 implies that, in any e.c. II_1 factor, the quantifier-free type of a tuple implies its complete type. It would be interesting to see if one could leverage this fact to gain any further insight into the class of e.c. factors.

In connection with Theorem 1, we say that a II_1 factor M has the *weak Jung property* if every endomorphism of M is approximately inner.

Proposition 6. *A separable II_1 factor has the Jung property if and only if it has the weak Jung property and is self-tracially stable.*

Proof. First suppose that M has the Jung property and that $f : M \rightarrow M$ is an endomorphism. By viewing f as taking values in $M^{\mathcal{U}}$, there is a unitary $u \in M^{\mathcal{U}}$ such that $uf(x)u^* = x$ for all $x \in M$. In particular, given any finite $F \subseteq M$ and $\varepsilon > 0$, there is a unitary $v \in M$ such that $\|vf(x)v^* - x\|_2 < \varepsilon$ for all $x \in F$, whence f is approximately inner. As mentioned in the introduction, any II_1 factor with the Jung property is self-tracially stable.

The converse is clear. ■

Given the fact that \mathcal{R} is the unique separable embeddable factor with either the Jung property or the property of being self-tracially stable, the following question seems natural⁹:

Question 7. Must there be at most one non-embeddable factor with the Jung property? That is self-tracially stable?

We now move on to Theorem 2, which will follow from an alternative characterization of the enforceable factor. First, recall from [1] that if \mathcal{C} is a class of tracial von Neumann algebras, then a tracial von Neumann algebra M is said to be \mathcal{C} -tracially stable if whenever $f : M \rightarrow \prod_{\mathcal{U}} N_i$ is an embedding with \mathcal{U} a nonprincipal ultrafilter on \mathbb{N} and each N_i belongs to \mathcal{C} , then there are *-homomorphisms $f_i : M \rightarrow N_i$ such that $f(x) = (f_i(x))_{\mathcal{U}}$ for all $x \in M$. (We refer to the sequence $(f_i)_{i \in \mathbb{N}}$ as an “approximate lifting” of f .) In particular, M is self-tracially stable if and only if M is $\{M\}$ -tracially stable.

⁹On the other hand, there may be more than one non-embeddable factor with the generalized Jung property; see [2, Corollary 3.3.5].

We let \mathfrak{E} denote the class of e.c. factors. The following theorem immediately implies Theorem 2:

Theorem 8. *The II_1 factor M is the enforceable factor if and only if it is e.c. and \mathfrak{E} -tracially stable.*

In order to prove Theorem 8, we need to recall a few model-theoretic facts from [7]. First, if M is an e.c. factor and a is a tuple from M , the *existential type of a in M* , denoted by $\text{etp}^M(a)$, is the collection of existential formulae $\varphi(x)$ such that $\varphi^M(a) = 0$. Such an existential type is called *isolated* if, given any $\varepsilon > 0$, there is an existential formula $\varphi(x)$ and $\delta > 0$ such that $\varphi^M(a) = 0$ and whenever N is an e.c. factor with a tuple $b \in N$ such that $\varphi^N(b) < \delta$, then there is $c \in N$ such that $\|b - c\|_2 < \varepsilon$ and for which $\text{etp}^M(a) = \text{etp}^N(c)$. The e.c. factor M is called *e-atomic* if the existential types of all finite tuples are isolated. It is shown in [7, Section 6] that an e.c. factor M is e-atomic if and only if it is the enforceable factor (in which case it is unique).

Proof of Theorem 8. First suppose that M is the enforceable factor. We must show that M is \mathfrak{E} -tracially stable. Towards this end, fix an embedding $f : M \rightarrow \prod_{\mathcal{U}} N_i$ with each N_i e.c. Since M is e.c., it is McDuff, whence singly generated. Fix a generator a of M and write $f(a) = (a_i)_{\mathcal{U}}$. Fix $\varepsilon > 0$ and let $\varphi(x)$ and $\delta > 0$ be as in the definition of isolated existential type for a and ε . Since M is e.c., f is an existential embedding, meaning that $\varphi^{\prod_{\mathcal{U}} N_i}(f(a)) = 0$ and thus $\varphi^{N_i}(a_i) < \delta$ for \mathcal{U} -almost all i . For these i , there is $b_i \in N_i$ such that $\|a_i - b_i\|_2 < \varepsilon$ and for which the map $a \mapsto b_i$ extends to an isomorphism between M and the subalgebra of N_i generated by b_i . Thus, f has an approximate lifting.

Conversely, suppose that M is e.c. and \mathfrak{E} -tracially stable. It follows that M embeds into every e.c. II_1 factor, whence M is enforceable. ■

Finally, we prove Theorem 3.

Proof of Theorem 3. Let M be the prime model of T^f . To show that M has the Jung property, we show that M has the weak Jung property and is self-tracially stable.

Fix a finitely generic factor N . Since M is the prime model of T^f , we have that M embeds elementarily in N . Thus, by [7, Corollary 3.12], M itself is finitely generic. In particular, M is e.c. and thus has the weak Jung property.

It remains to show that M is self-tracially stable. The argument for showing this is similar to that showing that the enforceable factor is self-tracially stable. Indeed, fix an embedding $f : M \rightarrow M^{\mathcal{U}}$. This time, given any $a \in M$, the complete type of a in M , denoted by $\text{tp}^M(a)$, is isolated¹⁰. Fix a generator a of M and write $f(a) = (a_i)_{\mathcal{U}}$. Given $\varepsilon > 0$, there is some formula $\varphi(x)$ and $\delta > 0$ such that $\varphi^M(a) = 0$ and such that, given any model N of T^f and any $b \in N$ with $\varphi^N(b) < \delta$, there is $c \in N$ such that $\text{tp}^M(a) = \text{tp}^N(c)$ and $\|b - c\|_2 < \varepsilon$. Since M is finitely generic, f is an elementary map, whence $\varphi^{M^{\mathcal{U}}}(f(a)) = 0$ and thus $\varphi^M(a_i) < \delta$ for \mathcal{U} -almost all i . As before, for these i , this

¹⁰This follows from the fact that prime models of theories are atomic models.

guarantees the existence of $b_i \in M$ such that $\|a_i - b_i\|_2 < \varepsilon$ and such that the map $a \mapsto b_i$ extends to an embedding of M into itself. Thus, f has an approximate lift. ■

Remark 9. It is not clear if there is any relationship between the existence of the enforceable factor and the existence of the prime model of T^f .

3. Revisiting the embeddable situation

In this section, we show how our techniques from above can yield different proofs that \mathcal{R} has the Jung property and is self-tracially similar.

Recall from the introduction that \mathcal{R} is the enforceable embeddable factor. Besides the model theory behind building models by games, the two main operator-algebraic ingredients in the proof are as follows:

- being hyperfinite is $\forall \bigvee \exists$ -axiomatizable¹¹, whence being hyperfinite is an enforceable property for embeddable factors;
- \mathcal{R} is the unique separable hyperfinite factor.

Noting that our proof from the previous section that the enforceable factor (should it exist) is self-tracially stable relativizes immediately to the embeddable situation, we obtain the fact that \mathcal{R} is self-tracially stable, *without resorting to the fact that \mathcal{R} has the Jung property*.

Unfortunately, our proof in the previous section that the enforceable factor (should it exist) has the weak Jung property does not necessarily relativize to the embeddable situation as the following seems to be an open question:

Question 10. Is the class of embeddable tracial von Neumann algebras closed under HNN extensions?

If the answer to the previous question is positive, then we learn that all e.c. embeddable factors (and thus, in particular, \mathcal{R} itself) have the weak Jung property.

Nevertheless, we can give a proof that is similar in spirit that does relativize to the embeddable situation. Indeed, fix an endomorphism $f : \mathcal{R} \rightarrow \mathcal{R}$; we show that f is approximately inner. Let a be a generator of \mathcal{R} . Since \mathcal{R} is a finitely generic embeddable factor¹², we have $\text{tp}^{\mathcal{R}}(a) = \text{tp}^{\mathcal{R}}(f(a))$. Consequently, there is an elementary extension N of \mathcal{R} and an automorphism σ of N such that $\sigma(f(a)) = a$. Using that $\mathcal{R} \subseteq N \rtimes_{\sigma} \mathbb{Z}$ and the class of embeddable factors is closed under crossed products by \mathbb{Z} (and, more generally, by any amenable group), we have that, given any $\varepsilon > 0$, there is a unitary $u \in \mathcal{R}$ such that $\|uf(a)u^* - a\|_2 < \varepsilon$. Consequently, f is approximately inner.

Combining these proofs gives a new proof that \mathcal{R} has the Jung property.

¹¹Morally speaking, one just axiomatizes the property that any finite tuple is within any positive tolerance of a copy of some matrix algebra.

¹²This follows from being the enforceable factor.

Acknowledgments. We would like to thank Scott Atkinson and Srivatsav Kunnawalkam Elayavalli for helpful discussions in preparing this paper.

Funding. The author was partially supported by NSF CAREER grant DMS-1349399.

References

- [1] S. Atkinson, Some results on tracial stability and graph products. *Indiana Univ. Math. J.* **70** (2021), no. 3, 1167–1187 Zbl 1484.46050 MR 4284110
- [2] S. Atkinson, I. Goldbring, and S. Kunnawalkam Elayavalli, Factorial relative commutants and the generalized Jung property for II_1 factors. *Adv. Math.* **396** (2022), Paper No. 108107, 53 Zbl 1487.46068 MR 4370469
- [3] S. Atkinson and S. Kunnawalkam Elayavalli, On ultraproduct embeddings and amenability for tracial von Neumann algebras. *Int. Math. Res. Not. IMRN* **2021** (2021), no. 4, 2882–2918 Zbl 1481.46060 MR 4218341
- [4] A. Connes, Classification of injective factors. Cases II_1 , II_∞ , III_λ , $\lambda \neq 1$. *Ann. of Math. (2)* **104** (1976), no. 1, 73–115 Zbl 0343.46042 MR 454659
- [5] I. Farah, I. Goldbring, B. Hart, and D. Sherman, Existentially closed II_1 factors. *Fund. Math.* **233** (2016), no. 2, 173–196 Zbl 1436.03213 MR 3466004
- [6] I. Farah, B. Hart, M. Rørdam, and A. Tikuisis, Relative commutants of strongly self-absorbing C^* -algebras. *Selecta Math. (N.S.)* **23** (2017), no. 1, 363–387 Zbl 1373.03064 MR 3595896
- [7] I. Goldbring, Enforceable operator algebras. *J. Inst. Math. Jussieu* **20** (2021), no. 1, 31–63 Zbl 1460.03009 MR 4205777
- [8] Z. Ji, A. Natarajan, T. Vidick, J. Wright, and H. Yuen, $MIP^* = RE$. 2022, arXiv:2001.04383
- [9] K. Jung, Amenability, tubularity, and embeddings into \mathcal{R}^ω . *Math. Ann.* **338** (2007), no. 1, 241–248 Zbl 1121.46052 MR 2295511
- [10] F. J. Murray and J. von Neumann, On rings of operators. IV. *Ann. of Math. (2)* **44** (1943), 716–808 Zbl 0060.26903 MR 9096
- [11] S. Popa and S. Vaes, W^* -rigidity paradigms for embeddings of II_1 factors. *Comm. Math. Phys.* **395** (2022), no. 2, 907–961 Zbl 07593346 MR 4487528
- [12] Y. Ueda, HNN extensions of von Neumann algebras. *J. Funct. Anal.* **225** (2005), no. 2, 383–426 Zbl 1088.46034 MR 2152505

Received 3 February 2021.

Isaac Goldbring

Department of Mathematics, University of California, 340 Rowland Hall, Irvine, CA 92697, USA;
isaac@math.uci.edu

