

# Symmetries of simple $A\mathbb{T}$ -algebras

Yuanhang Zhang

**Abstract.** Let  $A$  be a unital simple  $A\mathbb{T}$ -algebra of real rank zero. Given an order two automorphism  $h : K_1(A) \rightarrow K_1(A)$ , we show that there is an order two automorphism  $\alpha : A \rightarrow A$  such that  $\alpha_{*0} = \text{id}$ ,  $\alpha_{*1} = h$  and the action of  $\mathbb{Z}_2$  generated by  $\alpha$  has the tracial Rokhlin property. Consequently,  $C^*(A, \mathbb{Z}_2, \alpha)$  is a simple unital AH-algebra with no dimension growth, and with tracial rank zero. Thus, our main result can be considered the  $\mathbb{Z}_2$ -action analogue of the Lin–Osaka theorem. As a consequence, a positive answer to a lifting problem of Blackadar is also given for certain split case.

## 1. Introduction

It has been an important issue to find and classify all (or some particular) finite-order automorphisms of a given  $C^*$ -algebra. Historically, partly because of their intrinsic interest and partly because of their applications in  $C^*$ -dynamical systems, these kinds of problems have attracted considerable attention in the literature (see [4, 7, 16, 17, 21–23, 25, 26, 28, 29, 34, 41]). One landmark among them is Blackadar’s famous construction of symmetries (automorphisms of order 2) on the CAR algebra whose fixed point algebras have nontrivial  $K_1$ -group [4], hence giving a negative answer to one of two questions about AF-algebras posed by him in [3, 10.11.3]. The other one is a lifting question, which is as follows.

**Question 1.1.** Let  $A$  be an AF-algebra and  $\sigma$  an automorphism of the scaled ordered group  $K_0(A)$  with  $\sigma^n = \text{id}$ . Is there an automorphism  $\alpha$  of  $A$  with  $\alpha_{*0} = \sigma$  and  $\alpha^n = \text{id}$ ?

More generally, if a certain class of  $C^*$ -algebras is well understood, one could seek whether every finite group action on the level of  $K$ -theory of a  $C^*$ -algebra in this class can be lifted to a group action on the  $C^*$ -algebra. To be precise, one can consider the following folklore question [1, 39].

**Question 1.2.** If  $A$  belongs to a class of unital simple  $C^*$ -algebras that is classifiable by Elliott invariant and  $\sigma : G \rightarrow \text{Ell}(A)$  is an action of a finite group on the Elliott invariant of  $A$ , does there exist an action  $\alpha : G \rightarrow A$  with  $\text{Ell}(\alpha) = \sigma$ ?

For the case that  $A$  is a unital universal coefficient theorem (UCT) Kirchberg algebra, satisfactory answers to this question have been given. Firstly, Benson, Kumjian, and Phillips solved this question affirmatively for  $G = \mathbb{Z}_2$  and for unital UCT Kirchberg alge-

bras  $A$  in the Cuntz standard form [2]. Later, this result was extended by Spielberg who showed this question has an affirmative answer for  $G = \mathbb{Z}_p$ , where  $p$  is a prime number, and for an arbitrary unital UCT Kirchberg algebra [38]. Finally, in [27], this was further extended by Katsura to actions of finite groups whose Sylow subgroups are cyclic.

Within the setting of  $A$  being a unital simple stably finite  $C^*$ -algebra, compared to the recent progress in the Elliott classification programme (see, e.g., [19, 20, 40]), this question is still at an early stage, and there is much to do. We note that even Question 1.1 appears to be still open. Recently, Barlak and Szabó showed that if  $A$  is a separable, unital, simple and nuclear  $C^*$ -algebra with tracial rank zero which satisfies the UCT, then any action of a finite group  $G$ -action on the Elliott invariant could be lifted to a Rokhlin action of  $G$  on  $A$ , provided that  $A$  absorbs the UHF-algebra  $M_{|G|^\infty}$  [1, Corollary 2.13].

Our main goal in the present article is to examine Question 1.2 with the setting of  $G = \mathbb{Z}_2$  and  $A$  being a unital, simple  $A\mathbb{T}$ -algebra of real rank zero. Recall that an  $A\mathbb{T}$ -algebra is a  $C^*$ -algebra which is an inductive limit of  $C^*$ -algebras that are finite direct sums of matrix algebras over continuous functions on the circle  $\mathbb{T}$ . Unital simple  $A\mathbb{T}$ -algebras of real rank zero are classified by Elliott using scaled ordered  $K_0$ -groups and  $K_1$ -groups in [13]. Many  $C^*$ -algebras of interest (e.g., [6, 8, 35]), including irrational rotation algebras [14], are in this class.

As we shall see below (Theorem 3.3), for  $A$  being a unital simple  $A\mathbb{T}$ -algebra of real rank zero, given an order two automorphism  $h : K_1(A) \rightarrow K_1(A)$ , we show that there is a symmetry  $\alpha : A \rightarrow A$  such that  $\alpha_{*0} = \text{id}$ ,  $\alpha_{*1} = h$  and the action of  $\mathbb{Z}_2$  generated by  $\alpha$  has the tracial Rokhlin property. Consequently,  $C^*(A, \mathbb{Z}_2, \alpha)$  is a simple unital AH-algebra with no dimension growth, and with tracial rank zero. In fact, in the  $\mathbb{Z}$ -action setting, Lin and Osaka show that any unital simple  $A\mathbb{T}$ -algebra  $A$  admits an automorphism  $\alpha$  with the tracial (cyclic) Rokhlin property such that the induced homomorphism  $\alpha_{*1}$  on  $K_1(A)$  is equal to any given isomorphism of  $K_1(A)$ , and the induced homomorphism  $\alpha_{*0}$  on  $K_0(A)$  is the identity [32, Theorem 3.5]. Therefore, our aforementioned result could be viewed as a  $\mathbb{Z}_2$ -action analogue of the Lin–Osaka theorem.

In Section 4, as a variation of Theorem 3.3, we obtain the following: Let  $\mathfrak{A}$  be the AF-algebra whose scaled ordered group  $K_0(\mathfrak{A})$  is  $(G \oplus H, (G_+ \setminus \{0\}) \oplus H \cup \{(0, 0)\}, \tilde{g} \oplus \tilde{h})$ , where  $(G, G_+, \tilde{g})$  is the scaled ordered group  $K_0(B)$  of a unital simple AF-algebra  $B$ , and  $H$  is a countable torsion-free abelian group,  $\tilde{h} \in H$ . Let  $\sigma$  be an order two automorphism of  $K_0(\mathfrak{A})$ , defined by  $\sigma(g \oplus h) = g \oplus \eta(h)$ , where  $g \oplus h \in G \oplus H$ , and  $\eta$  is an order two automorphism of  $H$ . Then, there is a symmetry  $\alpha$  of  $\mathfrak{A}$  such that  $\alpha_{*0} = \sigma$  and the action of  $\mathbb{Z}_2$  generated by  $\alpha$  has the tracial Rokhlin property (Theorem 4.1). It is a generalization of [43, Theorem 4.1], where  $\eta$  is further assumed to be of type I and hence provides a partial affirmative answer to Question 1.1.

## 2. Preliminaries

In this section, we will review definitions, elementary facts and important results which we need in later sections.

We use the notation  $\mathbb{Z}_2$  for  $\mathbb{Z}/2\mathbb{Z}$ . If  $A$  is a  $C^*$ -algebra and  $\alpha : A \rightarrow A$  is an automorphism of order two, then we write  $C^*(\mathbb{Z}_2, A, \alpha)$ ,  $A^\alpha$  for the crossed product and the fixed point subalgebra of  $A$  by the action of  $\mathbb{Z}_2$  generated by  $\alpha$ , respectively. Given a  $C^*$ -algebra  $A$ , and a unitary  $u \in A$ , we denote by  $\text{Ad } u : A \rightarrow A$  the continuous linear map  $\text{Ad } u(a) = u^*au$ . Let  $\text{RR}$  denote the real rank. We take  $\mathbb{N} = \{n \in \mathbb{Z} : n > 0\}$ . Throughout this paper, an AF-algebra is always assumed to be a non-elementary one.

We shall assume that the reader is familiar with the notions and fundamental properties of the inductive limits of  $C^*$ -algebras and those of abelian groups. We shall also assume that the reader is familiar with  $K$ -theory, especially the functionalities of  $K_0$  and  $K_1$ , as found in [30, 36]. We also assume that the reader is familiar with approximately finite algebras, or AF-algebras, as inductive limits of finite-dimensional  $C^*$ -algebras and their classification [12] in terms of  $K$ -theory. The reader may refer to [30] for more details if required.

**2.1.** We shall use  $a^{\sim k}$  to denote  $\overbrace{a, \dots, a}^{k \text{ copies}}$  as used in [18, 1.1.7 (b)]. For example,

$$\{a^{\sim 2}, b^{\sim 3}\} = \{a, a, b, b, b\}.$$

**2.2.** Under the canonical bases of  $\mathbb{Z}^n$  and  $\mathbb{Z}^m$ , we shall identify a group homomorphism  $T$  from  $\mathbb{Z}^n \rightarrow \mathbb{Z}^m$  with its matrix representation  $T = (t_{i,j}) \in M_{m \times n}(\mathbb{Z})$ . Set

$$|T|_{\max} = \max \{|t_{i,j}|, 1 \leq i \leq m, 1 \leq j \leq n\},$$

and

$$|T|_{\min} = \min \{|t_{i,j}|, 1 \leq i \leq m, 1 \leq j \leq n\}.$$

**2.3.** Define

$$v(t) = \begin{cases} 0, & t > 0; \\ 1, & t < 0. \end{cases}$$

Let  $\lambda$  be a homeomorphism of  $X$ . For  $n \geq 0$ , define  $\lambda^n$  as the power  $n$  iteration of  $\lambda$ ; in particular,  $\lambda^0 = \text{id}_X$ .

**2.4.** Let  $A, B$  be unital  $C^*$ -algebras and  $\Phi : A \rightarrow B$  a unital homomorphism. Let us denote by  $\Phi_{*i} : K_i(A) \rightarrow K_i(B)$  the map induced by  $\Phi$ ,  $i = 0, 1$ .

**2.5.** Here are some basic  $K$ -theory properties of the reflection maps of  $S^n$ ,  $n = 1, 2$ .

(1) Let  $(w_1, w_2) \in S^1$ , and let  $\lambda$  be the reflection map defined by

$$\lambda(w_1, w_2) = (w_1, -w_2).$$

It is well known that  $K_0(C(S^1)) = \mathbb{Z}$ ,  $K_1(C(S^1)) = \mathbb{Z}$ , and  $\lambda_{*0}(m) = m$ ,  $\lambda_{*1}(n) = -n$ , for  $m \in K_0(C(S^1))$ ,  $n \in K_1(C(S^1))$ .

(2) Let  $(w_1, w_2, w_3) \in S^2$ , and let  $\lambda$  be the reflection map defined by

$$\lambda(w_1, w_2, w_3) = (w_1, w_2, -w_3).$$

It is well known that  $K_0(C(S^2)) = \mathbb{Z} \oplus \mathbb{Z}$ , and  $\lambda_{*0}(m, n) = (m, -n)$ , for  $(m, n) \in K_0(C(S^2))$ , where the first coordinate of  $\mathbb{Z} \oplus \mathbb{Z}$  denotes the rank part.

**2.6.** Let  $A = \lim_{k \rightarrow \infty} (A_k, \Phi_k)$  be an inductive limit of  $C^*$ -algebras. Here,  $\Phi_k$  is a homomorphism from  $A_k$  to  $A_{k+1}$ . We will use  $\Phi_{k,\infty} : A_k \rightarrow A$  to denote the homomorphism induced by the inductive limit system. Similarly, the notions could be defined *mutatis mutandis* to the setting of an inductive limit of abelian groups.

**Definition 2.7.** Let  $G_1 = \bigoplus_{i=1}^{m_1} G_{1,i}$  and  $G_2 = \bigoplus_{j=1}^{m_2} G_{2,j}$ , where  $G_{1,i} \cong G_{2,j} \cong \mathbb{Z}$ . Let  $\pi_j : G_2 \rightarrow G_{2,j}$  be the quotient maps. Suppose that  $\varphi : G_1 \rightarrow G_2$ . A partial map of  $\varphi$  from  $G_{1,i}$  to  $G_{2,j}$  is the map  $\varphi^{(i,j)} = \pi_j \circ \varphi|_{G_{1,i}}$  induced by  $\varphi$ . If  $\varphi^{(i,j)}(1) = l$ , then we say the multiplicity of partial map  $\varphi^{(i,j)}$ , denoted by  $|\varphi^{(i,j)}|$ , is  $l$ .

**Definition 2.8.** Let  $\Phi$  be a unital homomorphism from  $A_1 = \bigoplus_{i=1}^{m_1} M_{l(i,i)}(C(X_i))$  to  $A_2 = \bigoplus_{j=1}^{m_2} M_{l(2,j)}(C(Y_j))$ . For any  $i, j$ , if the partial map  $\Phi^{(i,j)}$ , the restriction of the map  $\Phi$  to any direct summands  $M_{l(1,i)}(C(X_i))$  and  $M_{l(2,j)}(C(Y_j))$ , has the form

$$f \mapsto \begin{bmatrix} f \circ \lambda_1 & & & \\ & \ddots & & \\ & & \ddots & \\ & & & f \circ \lambda_{d(i,j)} \end{bmatrix}$$

for some positive integer  $d(i, j)$  and some continuous maps  $\lambda_1, \dots, \lambda_{d(i,j)} : Y_j \rightarrow X_i$ , then  $\Phi$  is called a diagonal map, and  $|\Phi^{(i,j)}| := d(i, j)$  is called the multiplicity of  $\Phi^{(i,j)}$ .

The following is the Elliott–Gong classification theorem.

**Theorem 2.9** ([15]). *Let  $A$  and  $B$  be two unital simple AH-algebras with slow dimension growth and with real rank zero. Then,  $A \cong B$  if and only if*

$$(K_0(A), K_0(A)_+, [1_A], K_1(A)) \cong (K_0(B), K_0(B)_+, [1_B], K_1(B)).$$

Lastly, we introduce a special case of a useful criterion for an action of  $\mathbb{Z}_2$  to have the tracial Rokhlin property obtained by Phillips. The reader is referred to Phillips' seminal paper [33] for details and more background information about tracial Rokhlin property.

**Lemma 2.10** ([34, Lemma 1.8]). *Let  $A$  be a separable infinite-dimensional simple unital  $C^*$ -algebra with tracial rank zero. Let  $\alpha \in \text{Aut}(A)$  satisfy  $\alpha^2 = \text{id}_A$ . Suppose that for every finite set  $F \subset A$  and every  $\varepsilon > 0$ , there are mutually orthogonal projections  $e_0, e_1$  such that*

- (1)  $\|\alpha(e_0) - e_1\| < \varepsilon$ ;
- (2)  $\|e_j a - a e_j\| < \varepsilon$  for all  $a \in F$  and  $j = 0, 1$ ;
- (3) with  $e = e_0 + e_1$ ,  $\tau(1 - e) < \varepsilon$  for each tracial state  $\tau$  on  $A$ .

The action of  $\mathbb{Z}_2$  generated by  $\alpha$  has the tracial Rokhlin property.

### 3. $\mathbb{Z}_2$ -action analogue of the Lin–Osaka theorem

The purpose of this section is to present Theorem 3.3. We will start with a construction about the equivalence of two Bratteli diagrams, which proves very useful in our later constructions.

**Lemma 3.1.** *Let  $B$  be a unital simple AF-algebra. Let  $\{n_k\}, \{c_k\}$  be two increasing sequences of positive integers. Then,  $B$  could be written as*

$$B = \lim_{k \rightarrow \infty} (B_k, \Psi_k)$$

such that for  $k \in \mathbb{N}$ ,

- (1)  $B_k = M_{l(k,1)} \oplus \cdots \oplus M_{l(k,m_k)}$ , where  $m_k \geq 3n_k$ ;
- (2)  $\Psi_k$  is diagonal, and each partial map of  $\Psi_k$  has a positive multiplicity of at least  $(2k+3)c_k$ ;
- (3) for  $1 \leq i \leq n_k$ ,  $1 \leq j \leq m_{k+1}$ ,  $|\Psi_k^{(i,j)}| = (2k+3)c_k$ ; for  $n_k+1 \leq i \leq m_k$ ,  $1 \leq j_1, j_2 \leq n_{k+1}$ ,  $|\Psi_k^{(i,j_1)}| = |\Psi_k^{(i,j_2)}|$ ;
- (4)  $l(k,1) = \cdots = l(k,n_k)$ .

*Proof.* By [30, Proposition 4.7.2 and Lemma 4.7.3],  $B$  can be written so that

$$B = \lim_{k \rightarrow \infty} (C_k, \Phi_k),$$

where  $C_k = \bigoplus_{r=1}^{l_k} M_{h(k,r)}$ ,  $G'_k := K_0(C_k)$  is a finite direct sum of  $l_k$  copies of  $\mathbb{Z}$  with  $l_k \geq 2n_k$  and each partial map of  $\Phi_k$  has a positive multiplicity of at least 2,  $k \in \mathbb{N}$ . Without loss of generality, we may assume that the connecting map  $\Phi_k$  is a diagonal map,  $k \in \mathbb{N}$ . Since  $B$  is simple, by passing to a subsequence if necessary, we may further assume that

$$|\Phi_k^{(r,s)}| \geq 2n_k(2k+3)c_k,$$

where  $1 \leq r \leq l_k$ ,  $1 \leq s \leq l_{k+1}$ . For  $k \in \mathbb{N}$ , set  $\varphi_k = (\Phi_k)_*0$ .

We will first microscope the Bratteli diagram in a suitable way and then telescope it. To have a quick understanding about the construction, it is useful to draw the corresponding Bratteli diagrams.

Fix  $k \in \mathbb{N}$ , and let  $G_k = \mathbb{Z}^{m_k}$ , where  $m_k := l_k + n_k \geq 3n_k$ .

- (a) For  $r = 1$ , define  $\delta_k^{(1,i)} = 1$ , for  $1 \leq i \leq n_k + 1$ ; define  $\delta_k^{(1,i)} = 0$ , for  $2 + n_k \leq i \leq m_k$ . For each  $2 \leq r \leq l_k$ , define  $\delta_k^{(r,i)} = 1$ , for  $i = r + n_k$ ; define  $\delta_k^{(r,i)} = 0$ , otherwise. Hence, we define a homomorphism  $\delta_k$  from  $G'_k$  to  $G_k$ .
- (b) For  $1 \leq i \leq n_k$ , for  $1 \leq s \leq l_{k+1}$ , define  $\theta_k^{(i,s)} = (2k+3)c_k$ . For  $i = n_k + 1$ , for  $1 \leq s \leq l_{k+1}$ , define  $\theta_k^{(i,s)} = |\Phi_k^{(1,s)}| - n_k(2k+3)c_k$ . For  $n_k + 2 \leq i \leq m_k$ , for  $1 \leq s \leq l_{k+1}$ , define  $\theta_k^{(i,s)} = |\Phi_k^{(i-n_k,s)}|$ . Consequently, we define a homomorphism  $\theta_k$  from  $G_k$  to  $G'_{k+1}$  similarly.

Fix  $k \in \mathbb{N}$ . It is routine to verify that  $\varphi_k = \theta_k \circ \delta_k$ . In fact, for  $r = 1$ ,  $1 \leq s \leq l_{k+1}$ ,

$$\begin{aligned} |(\theta_k \circ \delta_k)^{(1,s)}| &= \sum_{i=1}^{m_k} \delta_k^{(1,i)} \theta_k^{(i,s)} = \sum_{i=1}^{n_k+1} \theta_k^{(i,s)} \\ &= n_k(2k+3)c_k + [|\Phi_k^{(1,s)}| - n_k(2k+3)c_k] = |\Phi_k^{(1,s)}| = |\varphi_k^{(1,s)}|; \end{aligned}$$

for  $2 \leq r \leq l_k, 1 \leq s \leq l_{k+1}$ ,

$$\begin{aligned} |(\theta_k \circ \delta_k)^{(r,s)}| &= \sum_{i=1}^{m_k} \delta_k^{(r,i)} \theta_k^{(i,s)} = \delta_k^{(r,r+n_k)} \theta_k^{(r+n_k,s)} \\ &= |\Phi_k^{(r+n_k-n_k,s)}| = |\Phi_k^{(r,s)}| = |\varphi_k^{(r,s)}|. \end{aligned}$$

Therefore,

$$G'_1 \xrightarrow{\delta_1} G_1 \xrightarrow{\theta_1} G'_2 \xrightarrow{\delta_2} G_2 \xrightarrow{\theta_2} \cdots G'_k \xrightarrow{\delta_k} G_k \xrightarrow{\theta_k} G'_{k+1} \xrightarrow{\delta_{k+1}} \cdots \rightarrow K_0(B).$$

For  $k \in \mathbb{N}$ , define  $\psi_k := \delta_{k+1} \circ \theta_k$  as a homomorphism from  $G_k$  to  $G_{k+1}$ . For  $1 \leq i \leq n_k, 1 \leq j \leq n_{k+1} + 1$ ,

$$|\psi_k^{(i,j)}| = \sum_{s=1}^{l_{k+1}} \theta_k^{(i,s)} \delta_{k+1}^{(s,j)} = (2k+3)c_k \sum_{s=1}^{l_{k+1}} \delta_{k+1}^{(s,j)} = (2k+3)c_k \delta_{k+1}^{(1,j)} = (2k+3)c_k;$$

for  $1 \leq i \leq n_k, n_{k+1} + 2 \leq j \leq m_{k+1}$ ,

$$|\psi_k^{(i,j)}| = \sum_{s=1}^{l_{k+1}} \theta_k^{(i,s)} \delta_{k+1}^{(s,j)} = (2k+3)c_k \sum_{s=1}^{l_{k+1}} \delta_{k+1}^{(s,j)} = (2k+3)c_k \delta_{k+1}^{(j-n_{k+1},j)} = (2k+3)c_k.$$

For  $i = n_k + 1, 1 \leq j \leq n_{k+1} + 1$ ,

$$|\psi_k^{(i,j)}| = \sum_{s=1}^{l_{k+1}} \theta_k^{(i,s)} \delta_{k+1}^{(s,j)} = \sum_{s=1}^{l_{k+1}} [|\Phi_k^{(1,s)}| - n_k(2k+3)c_k] \delta_{k+1}^{(s,j)} = |\Phi_k^{(1,1)}| - n_k(2k+3)c_k,$$

which is independent of  $j$ ; for  $i = n_k + 1, n_{k+1} + 2 \leq j \leq m_{k+1}$ ,

$$\begin{aligned} |\psi_k^{(i,j)}| &= \sum_{s=1}^{l_{k+1}} \theta_k^{(i,s)} \delta_{k+1}^{(s,j)} = \sum_{s=1}^{l_{k+1}} [|\Phi_k^{(1,s)}| - n_k(2k+3)c_k] \delta_{k+1}^{(s,j)} \\ &= |\Phi_k^{(1,j-n_{k+1})}| - n_k(2k+3)c_k. \end{aligned}$$

For  $n_k + 2 \leq i \leq m_k, 1 \leq j \leq n_{k+1} + 1$ ,

$$|\psi_k^{(i,j)}| = \sum_{s=1}^{l_{k+1}} \theta_k^{(i,s)} \delta_{k+1}^{(s,j)} = \sum_{s=1}^{l_{k+1}} |\Phi_k^{(i-n_k,s)}| \delta_{k+1}^{(s,j)} = |\Phi_k^{(i-n_k,1)}|,$$

which is independent of  $j$ ; for  $n_k + 2 \leq i \leq m_k, n_{k+1} + 2 \leq j \leq m_{k+1}$ ,

$$|\psi_k^{(i,j)}| = \sum_{s=1}^{l_{k+1}} \theta_k^{(i,s)} \delta_{k+1}^{(s,j)} = \sum_{s=1}^{l_{k+1}} |\Phi_k^{(i-n_k,s)}| \delta_{k+1}^{(s,j)} = |\Phi_k^{(i-n_k,j-n_{k+1})}|.$$

Summarizing, for  $1 \leq i \leq n_k$ ,  $1 \leq j \leq m_{k+1}$ ,  $|\psi_k^{(i,j)}| = (2k+3)c_k$ ; for  $n_k+1 \leq i \leq m_k$ ,  $1 \leq j_1, j_2 \leq n_{k+1}$ ,  $|\psi_k^{(i,j_1)}| = |\psi_k^{(i,j_2)}|$ . Since the multiplicity of each part map of  $\Phi_k$  is at least  $2n_k(2k+3)c_k$ , it follows that

$$|\psi_k^{(i,j)}| \geq (2k+3)c_k, \quad 1 \leq i \leq m_k, \quad 1 \leq j \leq m_{k+1}, \quad k \in \mathbb{N}.$$

Let  $B_1 = \bigoplus_{i=1}^{m_1} M_{l(1,i)}$ , where  $l(1,i) = h(1,1)$ , for  $1 \leq i \leq n_1$ , and  $l(1,i) = h(1, i - n_1)$ , for  $1 + n_1 \leq i \leq m_1$ . Define

$$l(k+1, j) = \sum_{i=1}^{m_k} |\psi_k^{(i,j)}| l(k, i)$$

inductively. Since for  $1 \leq j_1, j_2 \leq n_{k+1}$ ,

$$|\psi_k^{(i,j_1)}| = |\psi_k^{(i,j_2)}|, \quad \forall 1 \leq i \leq m_k,$$

it follows that

$$l(k+1, j_1) = \sum_{i=1}^{m_k} |\psi_k^{(i,j_1)}| l(k, i) = \sum_{i=1}^{m_k} |\psi_k^{(i,j_2)}| l(k, i) = l(k+1, j_2).$$

Set  $B_k = \bigoplus_{i=1}^{m_k} M_{l(k,i)}$ ,  $k \geq 2$ . Fix  $k \in \mathbb{N}$ , and define

$$\Delta_k^{(r,i)}(a) = \text{diag}\{a \sim \delta_k^{(r,i)}\},$$

where  $a \in M_{h(k,r)}$ ,  $1 \leq r \leq l_k$ ,  $1 \leq i \leq m_k$ . Set  $\Delta_k = \bigoplus_{r,i} \Delta_k^{(r,i)} : C_k \rightarrow B_k$ . Similarly, define

$$\Theta_k^{(i,s)}(a) = \text{diag}\{a \sim \theta_k^{(i,s)}\},$$

where  $a \in M_{l(k,i)}$ ,  $1 \leq i \leq m_k$ ,  $1 \leq s \leq l_{k+1}$ . Set  $\Theta_k = \bigoplus_{i,s} \Theta_k^{(i,s)} : B_k \rightarrow C_{k+1}$ .

Then, it is standard to check  $\Phi_k = \Theta_k \circ \Delta_k$ ,  $k \in \mathbb{N}$ ; hence,

$$C_1 \xrightarrow{\Delta_1} B_1 \xrightarrow{\Theta_1} C_2 \xrightarrow{\Delta_2} B_2 \xrightarrow{\Theta_2} \cdots \xrightarrow{\Delta_k} B_k \xrightarrow{\Theta_k} C_{k+1} \xrightarrow{\Delta_{k+1}} B_{k+1} \cdots \rightarrow B.$$

For  $k \in \mathbb{N}$ , define

$$\Psi_k = \Delta_{k+1} \circ \Theta_k : B_k \rightarrow B_{k+1}.$$

Then, it is obvious that

$$B = \lim_{k \rightarrow \infty} (B_k, \Psi_k).$$

Finally, the lemma now follows from the constructions.  $\blacksquare$

The following proposition could be found in [43]. It states that any order two automorphism of a countable torsion-free abelian group is actually an inductive limit action. For the reader's convenience, we give a detailed proof here (comparing with the original proof in [43]).

**Proposition 3.2.** *Let  $H$  be a nonzero countable torsion-free abelian group and  $\eta$  an order two automorphism of  $H$ . Then, there are a nondecreasing sequence of positive integers  $\{n_k : k \in \mathbb{N}\}$  and monomorphisms  $\beta_k : \mathbb{Z}^{n_k} \rightarrow \mathbb{Z}^{n_{k+1}}$ , order two automorphisms  $\eta_k : \mathbb{Z}^{n_k} \rightarrow \mathbb{Z}^{n_k}$  such that the following diagram commutes:*

$$\begin{array}{ccccccc} \mathbb{Z}^{n_1} & \xrightarrow{\beta_1} & \mathbb{Z}^{n_2} & \xrightarrow{\beta_2} & \mathbb{Z}^{n_3} & \xrightarrow{\beta_3} & \dots \longrightarrow H \\ \downarrow \eta_1 & & \downarrow \eta_2 & & \downarrow \eta_3 & & \downarrow \eta \\ \mathbb{Z}^{n_1} & \xrightarrow{\beta_1} & \mathbb{Z}^{n_2} & \xrightarrow{\beta_2} & \mathbb{Z}^{n_3} & \xrightarrow{\beta_3} & \dots \longrightarrow H \end{array}$$

and hence  $H = \lim_{k \rightarrow \infty} (\mathbb{Z}^{n_k}, \beta_k)$ ,  $\eta = \lim_{k \rightarrow \infty} \eta_k$ . Moreover, it can be required that under the canonical basis of  $\mathbb{Z}^{n_k}$ ,  $\eta_k$  has the form

$$\eta_k = \text{diag} \left\{ \overbrace{1, \dots, 1}^{p_k}; \overbrace{-1, \dots, -1}^{q_k}; \overbrace{\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \dots, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}}^{r_k} \right\}$$

for suitable nonnegative integers  $p_k, q_k, r_k$  such that  $p_k + q_k + 2r_k = n_k$ ,  $k \in \mathbb{N}$ .

*Proof.* Since  $H$  is countable, we can write  $H$  as  $H = \{e_1, e_2, \dots\}$ . Define

$$H_k = \mathbb{Z}[e_1, \dots, e_k; \eta(e_1), \dots, \eta(e_k)],$$

so for  $k \in \mathbb{N}$ ,  $\eta(H_k) = H_k$ ,  $H = \lim_{k \rightarrow \infty} (H_k, \iota_k)$ , where  $\iota_k$  is the embedding map from  $H_k$  to  $H_{k+1}$ . Since  $H_k$  is finitely generated and torsion-free, there exist a positive integer  $n_k$  and an isomorphism  $\chi_k$  such that  $\chi_k(\mathbb{Z}^{n_k}) = H_k$ . Since  $H_k \subset H_{k+1}$ , we have  $n_k \leq n_{k+1}$ . For  $k \in \mathbb{N}$ , define  $\psi_k = \chi_{k+1}^{-1} \circ \iota_k \circ \chi_k$ . Thus, it is easy to check that the following diagram commutes:

$$\begin{array}{ccccccc} \mathbb{Z}^{n_1} & \xrightarrow{\psi_1} & \mathbb{Z}^{n_2} & \xrightarrow{\psi_2} & \mathbb{Z}^{n_3} & \xrightarrow{\psi_3} & \dots \longrightarrow \lim_{k \rightarrow \infty} (\mathbb{Z}^{n_k}, \psi_k) \\ \downarrow \chi_1 & & \downarrow \chi_2 & & \downarrow \chi_3 & & \\ H_1 & \xrightarrow{\iota_1} & H_2 & \xrightarrow{\iota_2} & H_3 & \xrightarrow{\iota_3} & \dots \longrightarrow H \\ \downarrow \eta & & \downarrow \eta & & \downarrow \eta & & \\ H_1 & \xrightarrow{\iota_1} & H_2 & \xrightarrow{\iota_2} & H_3 & \xrightarrow{\iota_3} & \dots \longrightarrow H \\ \downarrow \chi_1^{-1} & & \downarrow \chi_2^{-1} & & \downarrow \chi_3^{-1} & & \\ \mathbb{Z}^{n_1} & \xrightarrow{\psi_1} & \mathbb{Z}^{n_2} & \xrightarrow{\psi_2} & \mathbb{Z}^{n_3} & \xrightarrow{\psi_3} & \dots \longrightarrow \lim_{k \rightarrow \infty} (\mathbb{Z}^{n_k}, \psi_k). \end{array}$$

Thus,

$$\lim_{k \rightarrow \infty} (\mathbb{Z}^{n_k}, \psi_k) = H.$$

Moreover,  $\theta_k := \chi_k^{-1} \circ \eta \circ \chi_k$  is an order two automorphism of  $\mathbb{Z}^{n_k}$ ,  $k \in \mathbb{N}$ , or equivalently,  $\theta_k$  is an involution matrix in  $M_{n_k}(\mathbb{Z})$ . By [24, Lemma 1] (or [2, Lemma 2.1]), for

$k \in \mathbb{N}$ , there are an invertible matrix  $S \in M_{n_k}(\mathbb{Z})$  and nonnegative integers  $p_k, q_k, r_k$  with

$$p_k + q_k + 2r_k = n_k$$

such that

$$\theta_k = S_k^{-1} \eta_k S_k,$$

where

$$\eta_k := \text{diag} \left\{ \overbrace{1, \dots, 1}^{p_k}; \overbrace{-1, \dots, -1}^{q_k}; \overbrace{\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \dots, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}}^{r_k} \right\}.$$

For  $k \in \mathbb{N}$ , set  $\beta_k = S_{k+1} \psi_k S_k^{-1}$ . Thus, it is easy to check that the following diagram commutes:

$$\begin{array}{ccccccc} \mathbb{Z}^{n_1} & \xrightarrow{\beta_1} & \mathbb{Z}^{n_2} & \xrightarrow{\beta_2} & \mathbb{Z}^{n_3} & \xrightarrow{\beta_3} & \dots \longrightarrow \lim_{k \rightarrow \infty} (\mathbb{Z}^{n_k}, \beta_k) \\ \downarrow S_1^{-1} & & \downarrow S_2^{-1} & & \downarrow S_3^{-1} & & \\ \mathbb{Z}^{n_1} & \xrightarrow{\psi_1} & \mathbb{Z}^{n_2} & \xrightarrow{\psi_2} & \mathbb{Z}^{n_3} & \xrightarrow{\psi_3} & \dots \longrightarrow H \\ \downarrow \theta_1 & & \downarrow \theta_2 & & \downarrow \theta_3 & & \\ \mathbb{Z}^{n_1} & \xrightarrow{\psi_1} & \mathbb{Z}^{n_2} & \xrightarrow{\psi_2} & \mathbb{Z}^{n_3} & \xrightarrow{\psi_3} & \dots \longrightarrow H \\ \downarrow S_1 & & \downarrow S_2 & & \downarrow S_3 & & \\ \mathbb{Z}^{n_1} & \xrightarrow{\beta_1} & \mathbb{Z}^{n_2} & \xrightarrow{\beta_2} & \mathbb{Z}^{n_3} & \xrightarrow{\beta_3} & \dots \longrightarrow \lim_{k \rightarrow \infty} (\mathbb{Z}^{n_k}, \beta_k). \end{array}$$

Therefore,

$$H = \lim_{k \rightarrow \infty} (\mathbb{Z}^{n_k}, \beta_k).$$

Note that  $\iota_k$  is injective, so is  $\psi_k$ , and then, so is  $\beta_k$ . The proposition follows immediately from the constructions.  $\blacksquare$

We are now in a position to prove the main result.

**Theorem 3.3.** *Let  $A$  be a unital simple  $A\mathbb{T}$ -algebra with real rank zero. Let  $h : K_1(A) \rightarrow K_1(A)$  be an automorphism with  $h^2 = \text{id}_{K_1(A)}$ . Then, there exists an automorphism  $\alpha : A \rightarrow A$  such that  $\alpha^2 = \text{id}$ ,  $\alpha_{*0} = \text{id}$ ,  $\alpha_{*1} = h$ , and the action of  $\mathbb{Z}_2$  generated by  $\alpha$  has the tracial Rokhlin property. In this case,  $C^*(\mathbb{Z}_2, A, \alpha)$  is a unital simple  $AH$ -algebra with no dimension growth, and with tracial rank zero.*

*Proof.* Set  $X = S^1$ , and let  $\lambda$  be the homeomorphism of  $X$  defined by  $\lambda(w_1, w_2) = (w_1, -w_2)$ , where  $(w_1, w_2) \in S^1$ . It is evident that  $\lambda^2 = \text{id}$ . Let  $z_0$  be a fixed point of  $\lambda$  and  $\{x_i : i \in \mathbb{N}\}$  a dense set of  $X$ . Suppose that  $K_1(A) \neq 0$ . (We will make a short remark for the trivial case  $K_1(A) = 0$  afterwards.)

We divide the proof into four steps.

**Step 1.** By Proposition 3.2, there are an increasing sequence of positive integers  $\{n_k : k \in \mathbb{N}\}$  and monomorphisms  $\beta_k : \mathbb{Z}^{n_k} \rightarrow \mathbb{Z}^{n_{k+1}}$ , order two automorphisms  $\eta_k : \mathbb{Z}^{n_k} \rightarrow \mathbb{Z}^{n_k}$  such that the following diagram is commutative:

$$\begin{array}{ccccccc} \mathbb{Z}^{n_1} & \xrightarrow{\beta_1} & \mathbb{Z}^{n_2} & \xrightarrow{\beta_2} & \mathbb{Z}^{n_3} & \xrightarrow{\beta_3} & \dots \longrightarrow K_1(A) \\ \downarrow \eta_1 & & \downarrow \eta_2 & & \downarrow \eta_3 & & \downarrow \\ \mathbb{Z}^{n_1} & \xrightarrow{\beta_1} & \mathbb{Z}^{n_2} & \xrightarrow{\beta_2} & \mathbb{Z}^{n_3} & \xrightarrow{\beta_3} & \dots \longrightarrow K_1(A) \end{array}$$

and

$$K_1(A) = \lim_{k \rightarrow \infty} (\mathbb{Z}^{n_k}, \beta_k), \quad h = \lim_{k \rightarrow \infty} \eta_k.$$

Moreover, under the canonical basis of  $\mathbb{Z}^{n_k}$ ,  $\eta_k$  has the form

$$\eta_k = \text{diag} \left\{ \overbrace{1, \dots, 1}^{p_k}; \overbrace{-1, \dots, -1}^{q_k}; \overbrace{\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \dots, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}}^{r_k} \right\}$$

for suitable nonnegative integers  $p_k, q_k, r_k$  such that  $p_k + q_k + 2r_k = n_k, k \in \mathbb{N}$ .

Since  $\beta_k, \gamma_k := \beta_k \circ \eta_k = \eta_{k+1} \circ \beta_k$  are monomorphisms from  $\mathbb{Z}^{n_k}$  to  $\mathbb{Z}^{n_{k+1}}$ , write  $\beta_k$  and  $\gamma_k$  as

$$\beta_k = \begin{bmatrix} b_{1,1}^{(k)} & b_{1,2}^{(k)} & \dots & b_{1,n_k}^{(k)} \\ b_{2,1}^{(k)} & b_{2,2}^{(k)} & \dots & b_{2,n_k}^{(k)} \\ \vdots & \vdots & \ddots & \vdots \\ b_{n_{k+1},1}^{(k)} & b_{n_{k+1},2}^{(k)} & \dots & b_{n_{k+1},n_k}^{(k)} \end{bmatrix}, \quad \gamma_k = \begin{bmatrix} r_{1,1}^{(k)} & r_{1,2}^{(k)} & \dots & r_{1,n_k}^{(k)} \\ r_{2,1}^{(k)} & r_{2,2}^{(k)} & \dots & r_{2,n_k}^{(k)} \\ \vdots & \vdots & \ddots & \vdots \\ r_{n_{k+1},1}^{(k)} & r_{n_{k+1},2}^{(k)} & \dots & r_{n_{k+1},n_k}^{(k)} \end{bmatrix}.$$

Note that  $A$  is a unital simple  $A\mathbb{T}$ -algebra. Let  $B$  be a unital simple  $AF$ -algebra such that

$$(K_0(B), K_0(B)_+, [1_B]) = (K_0(A), K_0(A)_+, [1_A]).$$

Set  $c_k = |\beta_k|_{\max} + k, k \in \mathbb{N}$ . By Lemma 3.1, we may assume that

$$B = \lim_{k \rightarrow \infty} (B_k, \Psi_k),$$

where  $B_k = M_{l(k,1)} \oplus \dots \oplus M_{l(k,m_k)}$  is a finite-dimensional  $C^*$ -algebra such that

- (1)  $m_k \geq 3n_k, l(k, 1) = \dots = l(k, n_k)$ ;
- (2)  $|\psi_k|_{\min} \geq (2k + 3)c_k$ , where  $\psi_k := (\Psi_k)_{*0}$ ;
- (3) for  $1 \leq i \leq n_k, 1 \leq j \leq m_{k+1}, |\Psi_k^{(i,j)}| = (2k + 3)c_k$ ; for  $n_k + 1 \leq i \leq m_k, 1 \leq j_1, j_2 \leq n_{k+1}, |\Psi_k^{(i,j_1)}| = |\Psi_k^{(i,j_2)}|$ .

Similarly, write  $\psi_k$  as

$$\psi_k = \begin{bmatrix} s_{1,1}^{(k)} & s_{1,2}^{(k)} & \cdots & s_{1,n_k}^{(k)} & s_{1,n_k+1}^{(k)} & \cdots & s_{1,m_k}^{(k)} \\ s_{2,1}^{(k)} & s_{2,2}^{(k)} & \cdots & s_{2,n_k}^{(k)} & s_{2,n_k+1}^{(k)} & \cdots & s_{2,m_k}^{(k)} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ s_{n_{k+1},1}^{(k)} & s_{n_{k+1},2}^{(k)} & \cdots & s_{n_{k+1},n_k}^{(k)} & s_{n_{k+1},n_k+1}^{(k)} & \cdots & s_{n_{k+1},m_k}^{(k)} \\ s_{n_{k+1}+1,1}^{(k)} & s_{n_{k+1}+1,2}^{(k)} & \cdots & s_{n_{k+1}+1,n_k}^{(k)} & s_{n_{k+1}+1,n_k+1}^{(k)} & \cdots & s_{n_{k+1}+1,m_k}^{(k)} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ s_{m_{k+1},1}^{(k)} & s_{m_{k+1},2}^{(k)} & \cdots & s_{m_{k+1},n_k}^{(k)} & s_{m_{k+1},n_k+1}^{(k)} & \cdots & s_{m_{k+1},m_k}^{(k)} \end{bmatrix}.$$

Then, it follows that

- (1) for  $1 \leq i \leq n_k$ ,  $1 \leq j \leq m_{k+1}$ ,  $s_{j,i}^{(k)} = (2k+3)c_k$ ;
- (2) for  $n_k+1 \leq i \leq m_k$ ,  $1 \leq j_1, j_2 \leq n_{k+1}$ ,  $s_{j_1,i}^{(k)} = s_{j_2,i}^{(k)}$ ;
- (3)  $|\psi_k|_{\min} \geq (2k+3)c_k$ .

These will be used again and again.

**Step 2.** For  $k \in \mathbb{N}$ , define

$$A_k = (M_{l(k,1)}(C(X)) \oplus \cdots \oplus M_{l(k,n_k)}(C(X))) \oplus (M_{l(k,n_k+1)} \oplus \cdots \oplus M_{l(k,m_k)}).$$

Then,

$$K_0(A_k) = \mathbb{Z}^{m_k}, \quad K_0(A_k)_+ = \mathbb{Z}_+^{m_k}, \quad K_1(A_k) = \mathbb{Z}^{n_k},$$

where

$$\mathbb{Z}_+^{m_k} := \{(\lambda_1, \dots, \lambda_{m_k}) \in \mathbb{Z}^{m_k} : \lambda_i \geq 0, 1 \leq i \leq m_k\}.$$

We next define two unital monomorphisms  $\Phi_k, \Theta_k : A_k \rightarrow A_{k+1}$ . Here, we may recall the notation  $\nu(t)$  defined in 2.3.

- (1) For  $1 \leq i \leq n_k$ , and  $1 \leq j \leq n_{k+1}$ , if  $b_{j,i}^{(k)} \neq 0$ , define

$$\Phi_k^{(i,j)}(f) = \text{diag} \left\{ (f \circ \lambda^{(\nu(b_{j,i}^{(k)})}))^{\sim |b_{j,i}^{(k)}|}; f(z_0)^{\sim (c_k - |b_{j,i}^{(k)}|) + 2kc_k}; f(x_1), f(\lambda(x_1)); \dots; f(x_{c_k}), f(\lambda(x_{c_k})) \right\};$$

if  $b_{j,i}^{(k)} = 0$ , define

$$\Phi_k^{(i,j)}(f) = \text{diag} \left\{ f(z_0)^{\sim c_k}, f(z_0)^{\sim 2kc_k}; f(x_1), f(\lambda(x_1)); \dots; f(x_{c_k}), f(\lambda(x_{c_k})) \right\};$$

similarly, if  $r_{j,i}^{(k)} \neq 0$ , define

$$\Theta_k^{(i,j)}(f) = \text{diag} \left\{ (f \circ \lambda^{(\nu(r_{j,i}^{(k)})}))^{\sim |r_{j,i}^{(k)}|}; f(z_0)^{\sim (c_k - |r_{j,i}^{(k)}|) + 2kc_k}; f(x_1), f(\lambda(x_1)); \dots; f(x_{c_k}), f(\lambda(x_{c_k})) \right\};$$

if  $r_{j,i}^{(k)} = 0$ , define

$$\Theta_k^{(i,j)}(f) = \text{diag} \{f(z_0)^{\sim c_k}, f(z_0)^{\sim 2kc_k}; f(x_1), f(\lambda(x_1)); \dots; f(x_{c_k}), f(\lambda(x_{c_k}))\}.$$

(2) For  $1 \leq i \leq n_k$ , and  $n_{k+1} + 1 \leq j \leq m_{k+1}$ , define

$$\Phi_k^{(i,j)}(f) = \Theta_k^{(i,j)}(f) = \text{diag} \{f(z_0)^{\sim c_k}; f(z_0)^{\sim 2kc_k}; f(z_0)^{\sim 2c_k}\}.$$

(3) For  $n_k + 1 \leq i \leq m_k$ , and  $1 \leq j \leq m_{k+1}$ , define

$$\Phi_k^{(i,j)}(a) = \Theta_k^{(i,j)}(a) = \text{diag} \{a^{\sim \gamma(k,i,j)}; a^{\sim 2\xi(k,i,j)}\},$$

where

$$\begin{aligned} \gamma(k,i,j) &= 1, \quad \xi(k,i,j) = \frac{s_{j,i}^{(k)} - 1}{2}, \quad \text{if } s_{j,i}^{(k)} \text{ is odd;} \\ \gamma(k,i,j) &= 2, \quad \xi(k,i,j) = \frac{s_{j,i}^{(k)} - 2}{2}, \quad \text{if } s_{j,i}^{(k)} \text{ is even.} \end{aligned}$$

Since for  $n_k + 1 \leq i \leq m_k$ ,  $1 \leq j_1, j_2 \leq n_{k+1}$ ,  $s_{j_1,i}^{(k)} = s_{j_2,i}^{(k)}$ , one could correspondingly show that  $\Phi_k^{(i,j_1)} = \Phi_k^{(i,j_2)}$ .

Set  $\Phi_k = \bigoplus_{1 \leq i \leq m_k, 1 \leq j \leq m_{k+1}} \Phi_k^{(i,j)}$ ,  $\Theta_k = \bigoplus_{1 \leq i \leq m_k, 1 \leq j \leq m_{k+1}} \Theta_k^{(i,j)}$ , where  $\Theta_k$  will be used as an auxiliary homomorphism in Step 3. Set  $C = \lim_{k \rightarrow \infty} (A_k, \Phi_k)$ . It is a standard matter to check that

$$\varphi_k := (\Phi_k)_{*0} = \psi_k, \quad (\Phi_k)_{*1} = \beta_k.$$

Hence,

$$(K_0(C), K_0(C)_+, [1_C], K_1(C)) \cong (K_0(A), K_0(A)_+, [1_A], K_1(A)).$$

Define

$$\pi_i^{(k)} : A_k \rightarrow M_{l(k,i)}(C(X)) \quad \text{for } 1 \leq i \leq n_k$$

and

$$\pi_i^{(k)} : A_k \rightarrow M_{l(k,i)} \quad \text{for } n_k + 1 \leq i \leq m_k$$

to be the quotient maps.

We next show that  $C$  is simple. Fix  $j \in \mathbb{N}$ , for a nonzero element  $f \in A_j$ .

*Case 1:*  $\pi_i^{(j)}(f) \neq 0$  for some  $i \geq n_j + 1$ . As each partial map of  $\Phi_j$  has positive multiplicity,  $\Phi_j(f)$  is full in  $A_{j+1}$ .

*Case 2:*  $\pi_i^{(j)}(f) \neq 0$  for some  $1 \leq i \leq n_j$ . Choose  $x^* \in X$  such that

$$\pi_i^{(j)}(f)(x^*) \neq 0.$$

Since  $\pi_i^{(j)}(f)$  is a continuous map on the compact metric space  $X$ , there exists a  $\delta > 0$  such that

$$\|\pi_i^{(j)}(f)(x) - \pi_i^{(j)}(f)(x^*)\| < \frac{\|\pi_i^{(j)}(f)(x^*)\|}{2}$$

provided that  $\text{dist}(x, x^*) < \delta$ , where  $x \in X$ . Choose  $k \in \mathbb{N}$ ,  $k > j$ , such that

$$\text{dist}(x^*, \{x_l : 1 \leq l \leq c_k\}) < \frac{\delta}{2}.$$

Now,  $\Phi_{j,k+2}(f)$  is full in  $A_{k+2}$ , where  $\Phi_{j,k+2} := \Phi_{k+1} \circ \cdots \circ \Phi_j$ .

By [9, Proposition 2.1 (iii)], it follows that  $C$  is simple.

Next, we will show that  $\text{RR}(C) = 0$ . Fix  $j \in \mathbb{N}$ , and a unit norm element  $f \in A_j$ . For  $\varepsilon > 0$ , choose  $k \in \mathbb{N}$ ,  $k > j + 1$ , such that

$$\frac{1}{k} < \frac{\varepsilon}{2}.$$

Thus, for any  $y_1, y_2 \in X$ ,

$$|\text{tr}(\Phi_{j,k}(f)(y_1)) - \text{tr}(\Phi_{j,k}(f)(y_2))| \leq \frac{2}{2k+3} < \varepsilon.$$

Therefore, by [36, Proposition 3.1.4],  $\text{RR}(C) = 0$ . By Theorem 2.9,

$$C \cong A.$$

**Step 3.** For  $k \in \mathbb{N}$ , recalling that  $l(k, 1) = \cdots = l(k, n_k)$ , we could define an order two automorphism  $\rho_k : A_k \rightarrow A_k$  by

$$\rho_k(f) = (f_1, \dots, f_{p_k}; g_1 \circ \lambda, \dots, g_{q_k} \circ \lambda; \hat{h}_1, h_1, \dots, \hat{h}_{r_k}, h_{r_k}; a_{n_k+1}, \dots, a_{m_k}),$$

where

$$f = (f_1, \dots, f_{p_k}; g_1, \dots, g_{q_k}; h_1, \hat{h}_1, \dots, h_{r_k}, \hat{h}_{r_k}; a_{n_k+1}, \dots, a_{m_k}) \in A_k.$$

In fact, under the canonical basis of  $\mathbb{Z}^{m_k}$ , it is routine – if tedious – to verify that

$$(\rho_k)_{*0} = \text{diag} \left\{ \overbrace{1, \dots, 1}^{p_k+q_k}; \overbrace{\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \dots, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}}^{r_k}; \overbrace{1, \dots, 1}^{m_k-n_k} \right\} := \sigma_k,$$

and

$$(\rho_k)_{*1} = \eta_k, \quad k \in \mathbb{N}.$$

Fix  $k \in \mathbb{N}$ . We claim that

$$\sigma_{k+1} \circ \psi_k = \psi_k,$$

or equivalently,

$$(\rho_{k+1})_{*0} |_{(\Phi_k)_{*0}(K_0(A_k))} = \text{id} |_{(\Phi_k)_{*0}(K_0(A_k))}.$$

This essentially follows from the fact that the first  $n_{k+1}$  rows of the matrix  $\psi_k$  are the same. More precisely, fix

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_{n_k} \\ x_{n_k+1} \\ \vdots \\ x_{m_k} \end{bmatrix} \in K_0(A_k) = \mathbb{Z}^{m_k}.$$

Define

$$y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_{n_{k+1}} \\ y_{n_{k+1}+1} \\ \vdots \\ y_{m_{k+1}} \end{bmatrix} := \varphi_k(x) = \psi_k(x) \in K_0(A_{k+1}) = \mathbb{Z}^{m_{k+1}}.$$

One has that

$$\begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_{n_{k+1}} \\ y_{n_{k+1}+1} \\ \vdots \\ y_{m_{k+1}} \end{bmatrix} = \psi_k \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_{n_k} \\ x_{n_k+1} \\ \vdots \\ x_{m_k} \end{bmatrix} = \begin{bmatrix} \sum_{i=1}^{m_k} s_{1,i}^{(k)} x_i \\ \sum_{i=1}^{m_k} s_{2,i}^{(k)} x_i \\ \vdots \\ \sum_{i=1}^{m_k} s_{n_{k+1},i}^{(k)} x_i \\ \sum_{i=1}^{m_k} s_{n_{k+1}+1,i}^{(k)} x_i \\ \vdots \\ \sum_{i=1}^{m_k} s_{m_{k+1},i}^{(k)} x_i \end{bmatrix}.$$

Recall that for  $1 \leq i \leq n_k$ ,  $1 \leq j \leq m_{k+1}$ ,

$$s_{j,i}^{(k)} = (2k+3)c_k,$$

and for  $n_k+1 \leq i \leq m_k$ ,  $1 \leq j_1, j_2 \leq n_{k+1}$ ,

$$s_{j_1,i}^{(k)} = s_{j_2,i}^{(k)}.$$

It follows that

$$y_1 = \cdots = y_{n_{k+1}},$$

whence

$$\sigma_{k+1}(y) = y.$$

Thus,

$$\begin{aligned}(\rho_{k+1})_{*0}((\Phi_k)_{*0}(x)) &= \sigma_{k+1}(\varphi_k(x)) = \sigma_{k+1}(\psi_k(x)) = \sigma_{k+1}(y) = y \\ &= \varphi_k(x) = (\Phi_k)_{*0}(x).\end{aligned}$$

Similarly, by observing that the first  $n_k$  columns of the matrix  $\psi_k$  are the same, one could show that

$$\psi_k \circ \sigma_k = \psi_k, \quad k \in \mathbb{N}.$$

Consider that  $\rho_{k+1} \circ \Phi_k$ . Recall that  $\gamma_k = \eta_{k+1} \circ \beta_k$ .

(a) For  $1 \leq i \leq n_k$ ,  $1 \leq j \leq p_{k+1}$ , as  $b_{j,i}^{(k)} = r_{j,i}^{(k)}$ ,

$$(\rho_{k+1} \circ \Phi_k)^{(i,j)} = \Phi_k^{(i,j)}, \quad \Theta_k^{(i,j)} = \Phi_k^{(i,j)}.$$

(b) For  $1 \leq i \leq n_k$ ,  $p_{k+1} + 1 \leq j \leq p_{k+1} + q_{k+1}$ . If  $b_{j,i}^{(k)} \neq 0$ , as  $-b_{j,i}^{(k)} = r_{j,i}^{(k)}$ ,

$$\begin{aligned}\Theta_k^{(i,j)}(f) &= \text{diag} \left\{ (f \circ \lambda^{(v(r_{j,i}^{(k)}))})^{\sim|r_{j,i}^{(k)}|}; f(z_0)^{\sim(c_k - |r_{j,i}^{(k)}|) + 2kc_k}; \right. \\ &\quad \left. f(x_1), f(\lambda(x_1)); \dots; f(x_{c_k}), f(\lambda(x_{c_k})) \right\} \\ &= \text{diag} \left\{ (f \circ \lambda^{(v(-b_{j,i}^{(k)}))})^{\sim|b_{j,i}^{(k)}|}; f(z_0)^{\sim(c_k - |b_{j,i}^{(k)}|) + 2kc_k}; \right. \\ &\quad \left. f(x_1), f(\lambda(x_1)); \dots; f(x_{c_k}), f(\lambda(x_{c_k})) \right\} \\ &= \text{diag} \left\{ (f \circ \lambda^{(v(b_{j,i}^{(k)} + 1))})^{\sim|b_{j,i}^{(k)}|}; f(z_0)^{\sim(c_k - |b_{j,i}^{(k)}|) + 2kc_k}; \right. \\ &\quad \left. f(x_1), f(\lambda(x_1)); \dots; f(x_{c_k}), f(\lambda(x_{c_k})) \right\} \\ &= (\rho_{k+1} \circ \Phi_k)^{(i,j)}(f).\end{aligned}$$

If  $b_{j,i}^{(k)} = 0$ , then  $r_{j,i}^{(k)} = -b_{j,i}^{(k)} = 0$ ,

$$\begin{aligned}(\rho_{k+1} \circ \Phi_k)^{(i,j)}(f) &= \text{diag} \left\{ f(z_0)^{\sim c_k}, f(z_0)^{\sim 2kc_k}; f(x_1), f(\lambda(x_1)); \dots; f(x_{c_k}), f(\lambda(x_{c_k})) \right\} \\ &= \Theta_k^{(i,j)}(f).\end{aligned}$$

(c) For  $1 \leq i \leq n_k$ ,  $p_{k+1} + q_{k+1} + 1 \leq j \leq n_{k+1}$ . If  $2 \nmid (j - p_{k+1} - q_{k+1})$ , then  $r_{j,i}^{(k)} = b_{j+1,i}^{(k)}$ ; hence,

$$(\rho_{k+1} \circ \Phi_k)^{(i,j)} = \Phi_k^{(i,j+1)} = \Theta_k^{(i,j)};$$

if  $2 \mid (j - p_{k+1} - q_{k+1})$ , then  $r_{j,i}^{(k)} = b_{j-1,i}^{(k)}$ ; hence,

$$(\rho_{k+1} \circ \Phi_k)^{(i,j)} = \Phi_k^{(i,j-1)} = \Theta_k^{(i,j)}.$$

(d) For  $1 \leq i \leq n_k$ ,  $n_{k+1} + 1 \leq j \leq m_{k+1}$ ,

$$(\rho_{k+1} \circ \Phi_k)^{(i,j)} = \Phi_k^{(i,j)} = \Theta_k^{(i,j)}.$$

(e) For  $n_k + 1 \leq i \leq m_k$ ,  $1 \leq j \leq m_{k+1}$ ,

$$(\rho_{k+1} \circ \Phi_k)^{(i,j)} = \text{diag}\{a^{\sim\gamma(k,i,j)}; a^{\sim 2\xi(k,i,j)}\} = \Theta_k^{(i,j)}.$$

In summary,

$$\rho_{k+1} \circ \Phi_k = \Theta_k.$$

In a similar manner, consider that  $\Phi_k \circ \rho_k$ . Recall that  $\gamma_k = \beta_k \circ \eta_k$ .

(a') For  $1 \leq i \leq p_k$ ,  $1 \leq j \leq n_{k+1}$ , as  $b_{j,i}^{(k)} = r_{j,i}^{(k)}$ ,

$$(\Phi_k \circ \rho_k)^{(i,j)} = \Phi_k^{(i,j)}, \quad \Theta_k^{(i,j)} = \Phi_k^{(i,j)}.$$

(b') For  $p_k + 1 \leq i \leq p_k + q_k$ ,  $1 \leq j \leq n_{k+1}$ . If  $b_{j,i}^{(k)} \neq 0$ , as  $-b_{j,i}^{(k)} = r_{j,i}^{(k)}$ ,

$$\begin{aligned} (\Phi_k \circ \rho_k)^{(i,j)}(f) &= \text{diag} \left\{ (f \circ \lambda^{(v(b_{j,i}^{(k)})+1)})^{\sim|b_{j,i}^{(k)}|}; f(z_0)^{\sim(c_k - |b_{j,i}^{(k)}|) + 2kc_k}; \right. \\ &\quad \left. f(\lambda(x_1)), f(x_1); \dots; f(\lambda(x_{c_k})), f(x_{c_k}) \right\}, \end{aligned}$$

and

$$\begin{aligned} \Theta_k^{(i,j)}(f) &= \text{diag} \left\{ (f \circ \lambda^{(v(r_{j,i}^{(k)}))})^{\sim|r_{j,i}^{(k)}|}; f(z_0)^{\sim(c_k - |r_{j,i}^{(k)}|) + 2kc_k}; \right. \\ &\quad \left. f(x_1), f(\lambda(x_1)); \dots; f(x_{c_k}), f(\lambda(x_{c_k})) \right\} \\ &= \text{diag} \left\{ (f \circ \lambda^{(v(-b_{j,i}^{(k)}))})^{\sim|b_{j,i}^{(k)}|}; f(z_0)^{\sim(c_k - |b_{j,i}^{(k)}|) + 2kc_k}; \right. \\ &\quad \left. f(x_1), f(\lambda(x_1)); \dots; f(x_{c_k}), f(\lambda(x_{c_k})) \right\} \\ &= \text{diag} \left\{ (f \circ \lambda^{(v(b_{j,i}^{(k)})+1)})^{\sim|b_{j,i}^{(k)}|}; f(z_0)^{\sim(c_k - |b_{j,i}^{(k)}|) + 2kc_k}; \right. \\ &\quad \left. f(x_1), f(\lambda(x_1)); \dots; f(x_{c_k}), f(\lambda(x_{c_k})) \right\}. \end{aligned}$$

If  $b_{j,i}^{(k)} = 0$ , then  $r_{j,i}^{(k)} = -b_{j,i}^{(k)} = 0$ ,

$$\begin{aligned} &(\Phi_k \circ \rho_k)^{(i,j)}(f) \\ &= \text{diag} \left\{ f(z_0)^{\sim c_k}, f(z_0)^{\sim 2kc_k}; f(\lambda(x_1)), f(x_1); \dots; f(\lambda(x_{c_k})), f(x_{c_k}) \right\}, \end{aligned}$$

while

$$\begin{aligned} &\Theta_k^{(i,j)}(f) \\ &= \text{diag} \left\{ f(z_0)^{\sim c_k}, f(z_0)^{\sim 2kc_k}; f(x_1), f(\lambda(x_1)); \dots; f(x_{c_k}), f(\lambda(x_{c_k})) \right\}. \end{aligned}$$

(c') For  $p_k + q_k + 1 \leq i \leq n_k$ ,  $1 \leq j \leq n_{k+1}$ . If  $2 \nmid (i - p_k - q_k)$ , as  $r_{j,i}^{(k)} = b_{j,i+1}^{(k)}$ ,

$$(\Phi_k \circ \rho_k)^{(i,j)} = \Phi_k^{(i+1,j)} = \Theta_k^{(i,j)};$$

if  $2 \mid (i - p_k - q_k)$ , as  $r_{j,i}^{(k)} = b_{j,i-1}^{(k)}$ ,

$$(\Phi_k \circ \rho_k)^{(i,j)} = \Phi_k^{(i-1,j)} = \Theta_k^{(i,j)}.$$

(d') For  $1 \leq i \leq n_k, n_{k+1} + 1 \leq j \leq m_{k+1}$ ,

$$(\Phi_k \circ \rho_k)^{(i,j)} = \text{diag} \{ f(z_0)^{\sim c_k}; f(z_0)^{\sim 2kc_k}; f(z_0)^{\sim 2c_k} \} = \Theta_k^{(i,j)}.$$

(e') For  $n_k + 1 \leq i \leq m_k, 1 \leq j \leq m_{k+1}$ ,

$$(\Phi_k \circ \rho_k)^{(i,j)} = \Phi_k^{(i,j)} = \Theta_k^{(i,j)}.$$

Summarizing, except for the case (b'),

$$(\Phi_k \circ \rho_k)^{(i,j)} = \Theta_k^{(i,j)}.$$

Set  $u_1 = 1_{A_1}$ . For  $1 \leq i \leq p_k$ , and  $p_k + q_k + 1 \leq i \leq n_k, 1 \leq j \leq m_{k+1}$ , define

$$u_{k+1,i,j} = \text{diag} \left\{ 1^{\sim c_k}; \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}^{\sim kc_k}; 1^{\sim 2c_k} \right\};$$

for  $p_k + 1 \leq i \leq p_k + q_k, 1 \leq j \leq n_{k+1}$ , define

$$u_{k+1,i,j} = \text{diag} \left\{ 1^{\sim c_k}; \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}^{\sim kc_k}; \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}^{\sim c_k} \right\};$$

for  $p_k + 1 \leq i \leq p_k + q_k, n_{k+1} + 1 \leq j \leq m_{k+1}$ , define

$$u_{k+1,i,j} = \text{diag} \left\{ 1^{\sim c_k}; \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}^{\sim kc_k}; 1^{\sim 2c_k} \right\};$$

for  $n_k + 1 \leq i \leq m_k, 1 \leq j \leq m_{k+1}$ ,

$$u_{k+1,i,j} = \text{diag} \left\{ 1^{\sim \gamma(k,i,j)}; \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}^{\sim \xi(k,i,j)} \right\}.$$

Next, define

$$u_{k+1} = \left( \bigoplus_{j=1}^{n_{k+1}} \left( \bigoplus_{i=1}^{m_k} u_{k+1,i,j} \otimes I_{l(k,i)} \right) \otimes 1_{C(X)} \right) \\ \oplus \left( \bigoplus_{j=n_{k+1}+1}^{m_{k+1}} \left( \bigoplus_{i=1}^{m_k} u_{k+1,i,j} \otimes I_{l(k,i)} \right) \right) \in A_{k+1}, \quad k \in \mathbb{N}.$$

Then,

$$u_{k+1}^* = u_{k+1}, \quad u_{k+1}^2 = 1_{A_{k+1}}, \quad \rho_{k+1}(u_{k+1}) = u_{k+1}.$$

By comparing (a)  $\sim$  (e) and (a')  $\sim$  (e'), it is routine – if tedious – to verify that

$$\Phi_k \circ \rho_k = \text{Ad } u_{k+1} \circ \Theta_k = \text{Ad } u_{k+1} \circ (\rho_{k+1} \circ \Phi_k), \quad k \in \mathbb{N}.$$

Set  $v_1 = 1_{A_1}$ , and define  $v_{k+1} = u_{k+1} \Phi_k(v_k) \in A_{k+1}$  inductively,  $k = 1, 2, \dots$ . Define

$$\alpha_k = \text{Ad } v_k \circ \rho_k.$$

Clearly,

$$\alpha_{k+1} \circ \Phi_k = \Phi_k \circ \alpha_k, \quad k \in \mathbb{N}.$$

To check that  $\alpha_k^2 = \text{id}_{A_k}$ , since  $\rho_k^2 = \text{id}_{A_k}$ , it is sufficient to show that  $\rho_k(v_k)v_k = 1_{A_k}$ . Since  $v_1 = 1_{A_1}$ ,  $\rho_1(v_1)v_1 = 1_{A_1}$ . Assume that  $\rho_k(v_k)v_k = 1_{A_k}$ . We will show that

$$\rho_{k+1}(v_{k+1})v_{k+1} = 1_{A_{k+1}}.$$

Indeed,

$$\begin{aligned} \rho_{k+1}(v_{k+1})v_{k+1} &= \rho_{k+1}(u_{k+1}\Phi_k(v_k))u_{k+1}\Phi_k(v_k) \\ &= \rho_{k+1}(u_{k+1})\rho_{k+1}(\Phi_k(v_k))u_{k+1}\Phi_k(v_k) \\ &= u_{k+1}u_{k+1}\Phi_k(\rho_k(v_k))u_{k+1}^*u_{k+1}\Phi_k(v_k) \\ &= \Phi_k(\rho_k(v_k)v_k) \\ &= \Phi_k(1_{A_k}) \\ &= 1_{A_{k+1}}. \end{aligned}$$

Then, one can easily construct the following commutative diagram:

$$\begin{array}{ccccccc} A_1 & \xrightarrow{\Phi_1} & A_2 & \xrightarrow{\Phi_2} & A_3 & \xrightarrow{\Phi_3} & \cdots \longrightarrow A \\ \downarrow \alpha_1 & & \downarrow \alpha_2 & & \downarrow \alpha_3 & & \\ A_1 & \xrightarrow{\Phi_1} & A_2 & \xrightarrow{\Phi_2} & A_3 & \xrightarrow{\Phi_3} & \cdots \longrightarrow A. \end{array}$$

Hence, the sequence of order two automorphisms  $\alpha_k$  defines

$$\alpha : A = \lim_{k \rightarrow \infty} (A_k, \Phi_k) \rightarrow A = \lim_{k \rightarrow \infty} (A_k, \Phi_k)$$

which is also a symmetry.

Note that

$$(\alpha_k)_{*1} = (\rho_k)_{*1} = \eta_k;$$

hence,

$$\alpha_{*1} = \lim_{k \rightarrow \infty} (\alpha_k)_{*1} = \lim_{k \rightarrow \infty} \eta_k = h.$$

Since

$$(\rho_{k+1})_{*0} |_{(\Phi_k)_{*0}(K_0(A_k))} = \text{id} |_{(\Phi_k)_{*0}(K_0(A_k))}, \quad (\alpha_{k+1})_{*0} = (\rho_{k+1})_{*0},$$

it follows that

$$(\alpha_{k+1})_{*0} |_{(\Phi_k)_{*0}(K_0(A_k))} = \text{id} |_{(\Phi_k)_{*0}(K_0(A_k))}.$$

Consequently,

$$\alpha_{*0} |_{(\Phi_{k,\infty})_{*0}(K_0(A_k))} = \text{id} |_{(\Phi_{k,\infty})_{*0}(K_0(A_k))}, \quad k \in \mathbb{N}.$$

Noting that

$$\bigcup_{k=1}^{\infty} (\Phi_{k,\infty})_* 0(K_0(A_k)) = K_0(A),$$

we have

$$\alpha_* 0 = \text{id}.$$

**Step 4.** We are now in a position to show that the action of  $\mathbb{Z}_2$  generated by  $\alpha$  has the tracial Rokhlin property. To this end, we fix a finite set  $F \subset A^1 := \{a : a \in A, \|a\| \leq 1\}$ , and an  $\varepsilon > 0$ . Then, there exists a positive integer  $k$  such that

$$\text{dist}(f, \Phi_{k,\infty}(A_k)) \leq \frac{\varepsilon}{2} \quad \text{for any } f \in F,$$

and

$$\frac{3}{2k+3} < \frac{\varepsilon}{2}.$$

For  $1 \leq i \leq n_k, 1 \leq j \leq m_{k+1}$ , define

$$e_{i,j}^{(0)} = \text{diag} \left\{ 0^{\sim c_k}; \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}^{\sim k c_k}; 0^{\sim 2c_k} \right\}, \quad e_{i,j}^{(1)} = \text{diag} \left\{ 0^{\sim c_k}; \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}^{\sim k c_k}; 0^{\sim 2c_k} \right\};$$

for  $n_k + 1 \leq i \leq m_k, 1 \leq j \leq m_{k+1}$ , define

$$e_{i,j}^{(0)} = \text{diag} \left\{ 0^{\sim \gamma(k,i,j)}; \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}^{\sim \xi(k,i,j)} \right\}, \quad e_{i,j}^{(1)} = \text{diag} \left\{ 0^{\sim \gamma(k,i,j)}; \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}^{\sim \xi(k,i,j)} \right\}.$$

Define

$$e_l = \left( \bigoplus_{j=1}^{n_{k+1}} \left( \bigoplus_{i=1}^{m_k} e_{i,j}^{(l)} \otimes I_{l(k,i)} \otimes 1_{C(X)} \right) \right) \oplus \left( \bigoplus_{j=n_{k+1}+1}^{m_{k+1}} \left( \bigoplus_{i=1}^{m_k} e_{i,j}^{(l)} \otimes I_{l(k,i)} \right) \right),$$

$$l = 0, 1.$$

Then, it is easy to check that  $e_0, e_1$  are mutually orthogonal projections of  $A_{k+1}$  with

$$u_{k+1}^* e_0 u_{k+1} = e_1.$$

In addition, it is easily seen that

$$\rho_{k+1}(e_l) = e_l,$$

and  $e_l$  commutes with  $\Phi_k(A_k)$ ,  $l = 0, 1$ .

Note that

$$\begin{aligned} \alpha_{k+1}(e_0) &= v_{k+1}^* \rho_{k+1}(e_0) v_{k+1} \\ &= (u_{k+1} \Phi_k(v_k))^* \rho_{k+1}(e_0) u_{k+1} \Phi_k(v_k) \\ &= \Phi_k(v_k)^* u_{k+1}^* \rho_{k+1}(e_0) u_{k+1} \Phi_k(v_k) \\ &= \Phi_k(v_k)^* u_{k+1}^* e_0 u_{k+1} \Phi_k(v_k) \\ &= \Phi_k(v_k)^* e_1 \Phi_k(v_k) \\ &= \Phi_k(v_k)^* \Phi_k(v_k) e_1 \\ &= e_1. \end{aligned}$$

Moreover, for any tracial state  $\tau$  on  $A_{k+1}$ ,

$$\tau(1_{A_{k+1}} - e_0 - \alpha_{k+1}(e_0)) = \tau(1_{A_{k+1}} - e_0 - e_1) \leq \frac{3}{2k+3} < \frac{\varepsilon}{2}.$$

Then, by Lemma 2.10, the action of  $\mathbb{Z}_2$  generated by  $\alpha$  has the tracial Rokhlin property.

Since  $A$  is a unital simple  $A\mathbb{T}$ -algebra with  $\text{RR}(A) = 0$ , by [30, Theorem 4.3.5],  $A$  has tracial rank zero. Hence, by [33, Corollary 1.6 and Theorem 2.6],  $C^*(\mathbb{Z}_2, A, \alpha)$  is a unital simple, separable  $C^*$ -algebra with tracial rank zero. Since  $A_k$  is nuclear and  $\mathbb{Z}_2$  is amenable and compact, it follows from [42, Corollary 7.18] and [10, Proposition 6.1] that  $C^*(\mathbb{Z}_2, A_k, \alpha_k)$  is nuclear and satisfies the UCT. Hence, by [36, Proposition 2.4.7 (ii)],

$$C^*(\mathbb{Z}_2, A, \alpha) = \lim_{k \rightarrow \infty} (C^*(\mathbb{Z}_2, A_k, \alpha_k), \Phi_k)$$

is also nuclear and satisfies the UCT. Therefore, by [31, Theorem 5.2] and its proof,  $C^*(\mathbb{Z}_2, A, \alpha)$  is a unital simple AH-algebra with no dimension growth.  $\blacksquare$

**Remark 3.4.** If we do not insist on the fact that the action of  $\mathbb{Z}_2$  generated by  $\alpha$  has the tracial Rokhlin property, the unitary  $u_{k+1}$  could be constructed in a simpler form which has less flips, comparing with that we used in Step 3 of the proof of Theorem 3.3.

**Remark 3.5.** Let  $\tilde{A}$  be a unital simple  $A\mathbb{T}$ -algebra with real rank zero and  $K_1(\tilde{A}) = 0$ . Pick a *companion* algebra  $A$ , which is a unital simple  $A\mathbb{T}$ -algebra with real rank zero, and

$$(K_0(A), K_0(A)_+, [1_A], K_1(A)) \cong (K_0(\tilde{A}), K_0(\tilde{A})_+, [1_{\tilde{A}}], \mathbb{Z}).$$

Let  $h$  be the identity map of  $K_1(A)$ . By examining the proof of Theorem 3.3 carefully, it is straightforward to check that  $B \cong \tilde{A}$ , and  $B$  is a unital subalgebra of  $A$  by embedding  $B_k$  into  $A_k$ . Moreover, one could easily check that  $\alpha$  is also a symmetry of  $B$ , and  $\alpha_{*0}|_{K_0(B)} = \text{id}|_{K_0(B)}$ . Finally, exactly as that in Step 4, the action of  $\mathbb{Z}_2$  generated by  $\alpha$  has the tracial Rokhlin property, and  $C^*(\mathbb{Z}_2, B, \alpha)$  is a unital simple AH-algebra with no dimension growth, tracial rank zero (actually an AF-algebra) as desired. In other words, we finish the proof of the trivial case of Theorem 3.3.

## 4. An application to a lifting problem of Blackadar

In this section, we will modify the procedure of the proof of Theorem 3.3 to give a positive answer to a lifting problem of Blackadar for the split case.

**Theorem 4.1.** *Let  $\mathfrak{A}$  be the AF-algebra whose scaled ordered group  $K_0(\mathfrak{A})$  is (isomorphic to)*

$$(G \oplus H, (G_+ \setminus \{0\}) \oplus H \cup \{0 \oplus 0\}, \tilde{g} \oplus \tilde{h}),$$

where  $(G, G_+, \tilde{g})$  is the scaled ordered group  $K_0(B)$  of a unital simple AF-algebra  $B$ , and  $H$  is a countable torsion-free abelian group,  $\tilde{h} \in H$ . Let  $\sigma$  be an order two automorphism of  $K_0(\mathfrak{A})$ , defined by  $\sigma(g \oplus h) = g \oplus \eta(h)$ , where  $g \oplus h \in G \oplus H$ , and  $\eta$  is an order

two automorphism of  $H$ . Then, there is an automorphism  $\alpha$  of  $\mathfrak{A}$  with  $\alpha_{*0} = \sigma$ ,  $\alpha^2 = \text{id}$  and the action of  $\mathbb{Z}_2$  generated by  $\alpha$  has the tracial Rokhlin property.

Consequently,  $C^*(\mathbb{Z}_2, \mathfrak{A}, \alpha)$  is a unital simple  $AH$ -algebra with no dimension growth, and with tracial rank zero.

*Proof.* By Theorem 3.3, we assume that  $H \neq \{0\}$ . Set  $X = S^2$ , and let  $\lambda$  be the homeomorphism of  $X$  defined by  $\lambda(w_1, w_2, w_3) = (w_1, w_2, -w_3)$ , where  $(w_1, w_2, w_3) \in S^2$ . Let  $z_0$  be a fixed point of  $\lambda$  and  $\{x_i : i \in \mathbb{N}\}$  a dense set of  $X$ . Paralleled with that of Theorem 3.3, we will divide the whole proof into four steps.

**Step 1.** By Proposition 3.2, there are a sequence of positive integers  $\{n_k : k \in \mathbb{N}\}$  and monomorphisms  $\beta_k : \mathbb{Z}^{n_k} \rightarrow \mathbb{Z}^{n_{k+1}}$ , order two automorphisms  $\eta_k : \mathbb{Z}^{n_k} \rightarrow \mathbb{Z}^{n_k}$  such that the following diagram commutes:

$$\begin{array}{ccccccc} \mathbb{Z}^{n_1} & \xrightarrow{\beta_1} & \mathbb{Z}^{n_2} & \xrightarrow{\beta_2} & \mathbb{Z}^{n_3} & \xrightarrow{\beta_3} & \dots \longrightarrow H \\ \downarrow \eta_1 & & \downarrow \eta_2 & & \downarrow \eta_3 & & \downarrow \eta \\ \mathbb{Z}^{n_1} & \xrightarrow{\beta_1} & \mathbb{Z}^{n_2} & \xrightarrow{\beta_2} & \mathbb{Z}^{n_3} & \xrightarrow{\beta_3} & \dots \longrightarrow H \end{array}$$

and  $H = \lim_{k \rightarrow \infty} (\mathbb{Z}^{n_k}, \beta_k)$ ,  $\eta = \lim_{k \rightarrow \infty} \eta_k$ . Moreover, under the canonical basis of  $\mathbb{Z}^{n_k}$ ,  $\eta_k$  has the form

$$\eta_k = \text{diag} \left\{ \overbrace{1, \dots, 1}^{p_k}; \overbrace{-1, \dots, -1}^{q_k}; \overbrace{\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \dots, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}}^{r_k} \right\}$$

for suitable nonnegative integers  $p_k, q_k, r_k$  such that  $p_k + q_k + 2r_k = n_k, k \in \mathbb{N}$ .

By Lemma 3.1, we may assume that

$$B = \lim_{k \rightarrow \infty} (B_k, \Psi_k),$$

where  $B_k = M_{l(k,1)} \oplus \dots \oplus M_{l(k,m_k)}$  is a finite-dimensional  $C^*$ -algebra and (1)–(4) in Lemma 3.1 hold. Let  $\tilde{g}_k$  denote  $(l(k,1), \dots, l(k,m_k)) \in \mathbb{Z}^{m_k}, k \in \mathbb{N}$ . Then,

$$(K_0(B_k), K_0(B_k)_+, [1_{B_k}]) = (\mathbb{Z}^{m_k}, \mathbb{Z}_+^{m_k}, \tilde{g}_k).$$

Set  $\psi_k := (\Psi_k)_{*0}, k \in \mathbb{N}$ . Note that  $\psi_{k,\infty}(\tilde{g}_k) = \tilde{g}, k \in \mathbb{N}$ . Consider the ordered group  $\mathbb{Z}^{m_k} \oplus \mathbb{Z}^{n_k}$  with the positive cone

$$\left\{ (\lambda_1, \dots, \lambda_{m_k}; \mu_1, \dots, \mu_{n_k}) \in \mathbb{Z}^{m_k} \oplus \mathbb{Z}^{n_k} : \lambda_i \geq 0, 1 \leq i \leq m_k, \sum_{i=1}^{m_k} \lambda_i > 0 \right\} \\ \cup \{(0 \sim^{m_k}; 0 \sim^{n_k})\}.$$

Then, it is evident that

$$\mathbb{Z}^{m_1} \oplus \mathbb{Z}^{n_1} \xrightarrow{\psi_1 \oplus \beta_1} \mathbb{Z}^{m_1} \oplus \mathbb{Z}^{n_2} \xrightarrow{\psi_2 \oplus \beta_2} \mathbb{Z}^{m_3} \oplus \mathbb{Z}^{n_3} \xrightarrow{\psi_3 \oplus \beta_3} \dots \rightarrow G \oplus H,$$

which forms an inductive limit system of ordered groups.

**Step 2.** Define

$$A_k = (M_{2l(k,1)}(C(X)) \oplus \cdots \oplus M_{2l(k,n_k)}(C(X))) \oplus (M_{2l(k,n_k+1)} \oplus \cdots \oplus M_{2l(k,m_k)}).$$

Then,

- $K_0(A_k) = \overbrace{(\mathbb{Z} \oplus \mathbb{Z}) \oplus \cdots \oplus (\mathbb{Z} \oplus \mathbb{Z})}^{n_k} \oplus \overbrace{\mathbb{Z} \oplus \cdots \oplus \mathbb{Z}}^{m_k - n_k};$
- $K_0(A_k)_+ = \overbrace{(\mathbb{Z} \oplus \mathbb{Z})_+ \oplus \cdots \oplus (\mathbb{Z} \oplus \mathbb{Z})_+}^{n_k} \oplus \overbrace{\mathbb{Z}_+ \oplus \cdots \oplus \mathbb{Z}_+}^{m_k - n_k}$ , where
 
$$(\mathbb{Z} \oplus \mathbb{Z})_+ = \{(\lambda, \mu) : \lambda \in \mathbb{N}, \mu \in \mathbb{Z}, \text{ or } \lambda = \mu = 0\};$$
- $[1_{A_k}] = (2l(k, 1), 0, \dots, 2l(k, n_k), 0; 2l(k, n_k + 1), \dots, 2l(k, m_k)).$

Exactly as that in the proof of Theorem 3.3, we define two unital monomorphisms  $\Phi_k, \Theta_k : A_k \rightarrow A_{k+1}$ . Set  $A = \lim_{k \rightarrow \infty} (A_k, \Phi_k)$ .

For  $k \in \mathbb{N}$ , define  $\varphi_k = (\Phi_k)_{*0}$ , and

$$\omega_k : (\mathbb{Z} \oplus \mathbb{Z})^{n_k} \oplus \mathbb{Z}^{m_k - n_k} \rightarrow \mathbb{Z}^{m_k} \oplus \mathbb{Z}^{n_k}$$

by

$$\omega_k(\lambda_1, \mu_1; \dots; \lambda_{n_k}, \mu_{n_k}; \lambda_{n_k+1}, \dots, \lambda_{m_k}) = (\lambda_1, \dots, \lambda_{m_k}; \mu_1, \dots, \mu_{n_k}).$$

Then, it is routine to check that  $\omega_k$  is a positive homomorphism and bijection, while  $\omega_k^{-1}$  is not necessarily positive.

For  $k \in \mathbb{N}$ , consider

$$\chi_k := \omega_{k+1}^{-1} \circ (\psi_k \oplus \beta_k) : \mathbb{Z}^{m_k} \oplus \mathbb{Z}^{n_k} \rightarrow (\mathbb{Z} \oplus \mathbb{Z})^{n_{k+1}} \oplus \mathbb{Z}^{m_{k+1} - n_{k+1}}.$$

According to each partial map of  $\psi_k$  having positive multiplicity, it is routine to check that  $\chi_k$  is a positive homomorphism. Moreover,

$$\psi_k \oplus \beta_k = \omega_{k+1} \circ \chi_k.$$

On the other hand, by checking each basis vector of  $(\mathbb{Z} \oplus \mathbb{Z})^{n_k} \oplus \mathbb{Z}^{m_k - n_k}$ , it is straightforward – if tedious – to show that

$$\omega_{k+1} \circ \varphi_k = (\psi_k \oplus \beta_k) \circ \omega_k; \quad (\star)$$

hence,

$$\varphi_k = \chi_k \circ \omega_k, \quad k \in \mathbb{N}.$$

Therefore, we have the following commutative diagram:

$$\begin{array}{ccccccc} (\mathbb{Z} \oplus \mathbb{Z})^{(n_1)} \oplus \mathbb{Z}^{(m_1 - n_1)} & \xrightarrow{\varphi_1} & (\mathbb{Z} \oplus \mathbb{Z})^{(n_2)} \oplus \mathbb{Z}^{(m_2 - n_2)} & \xrightarrow{\varphi_2} & \cdots & \longrightarrow & K_0(A) \\ \omega_1 \downarrow & \nearrow \chi_1 & \omega_2 \downarrow & \nearrow \chi_2 & & & \omega \downarrow \\ \mathbb{Z}^{m_1} \oplus \mathbb{Z}^{n_1} & \xrightarrow{\psi_1 \oplus \beta_1} & \mathbb{Z}^{m_2} \oplus \mathbb{Z}^{n_2} & \xrightarrow{\psi_2 \oplus \beta_2} & \cdots & \longrightarrow & G \oplus H. \end{array}$$

Since each homomorphism in the above diagram is positive monomorphism, by the standard intertwining argument, it follows that, via  $\omega$ ,

$$(K_0(A), K_0(A)_+) \cong (G \oplus H, (G_+ \setminus \{0\}) \oplus H \cup \{0 \oplus 0\}) \cong (K_0(\mathfrak{A}), K_0(\mathfrak{A})_+),$$

and

$$\begin{aligned} \omega([1_A]) &= (\psi_{1,\infty} \oplus \beta_{1,\infty})(\omega_1([1_{A_1}])) = (\psi_{1,\infty} \oplus \beta_{1,\infty})(\omega_1(2\nu_k)) \\ &= (\psi_{1,\infty} \oplus \beta_{1,\infty})(2\tilde{g}_k \oplus 0_{n_k}) = 2\tilde{g} \oplus 0. \end{aligned}$$

Since  $K_1(A_k) = 0$ ,  $\forall k \in \mathbb{N}$ , one has  $K_1(A) = 0$ .

Using the same argument where appropriate in the proof of Theorem 3.3, we conclude that  $A$  is simple and  $\text{RR}(A) = 0$ . By Theorem 2.9,  $A$  is a simple AF-algebra.

**Step 3.** For  $k \in \mathbb{N}$ , define order two automorphisms  $\rho_k : A_k \rightarrow A_k$  by

$$\rho_k(f) = (f_1, \dots, f_{p_k}; g_1 \circ \lambda, \dots, g_{q_k} \circ \lambda; \hat{h}_1, h_1, \dots, \hat{h}_{r_k}, h_{r_k}; a_{n_k+1}, \dots, a_{m_k}),$$

where

$$f = (f_1, \dots, f_{p_k}; g_1, \dots, g_{q_k}; h_1, \hat{h}_1, \dots, h_{r_k}, \hat{h}_{r_k}; a_{n_k+1}, \dots, a_{m_k}) \in A_k.$$

In fact, it is routine – if tedious – to verify that

$$\begin{aligned} \varsigma_k := (\rho_k)_{*0} = \text{diag} & \left\{ \overbrace{1, \dots, 1}^{2p_k}; \overbrace{\begin{bmatrix} 1 & & & \\ & -1 & & \\ & & \ddots & \\ & & & -1 \end{bmatrix}}^{q_k}; \right. \\ & \left. \overbrace{\begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}}^{r_k}, \dots, \overbrace{\begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}}^{r_k}; \overbrace{1, \dots, 1}^{m_k - n_k} \right\}. \end{aligned}$$

For  $k \in \mathbb{N}$ , let

$$\begin{aligned} \hat{\sigma}_k := \text{diag} & \left\{ \overbrace{1, \dots, 1}^{p_k + q_k}; \overbrace{\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}}^{r_k}, \dots, \overbrace{\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}}^{r_k}; \overbrace{1, \dots, 1}^{m_k - n_k}; \overbrace{1, \dots, 1}^{p_k}; \overbrace{-1, \dots, -1}^{q_k}; \right. \\ & \left. \overbrace{\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}}^{r_k}, \dots, \overbrace{\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}}^{r_k} \right\} \\ & = \sigma_k \oplus \eta_k, \end{aligned}$$

where

$$\sigma_k := \text{diag} \left\{ \overbrace{1, \dots, 1}^{p_k+q_k}; \overbrace{\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \dots, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}}^{r_k}; \overbrace{1, \dots, 1}^{m_k-n_k} \right\}.$$

Then,  $\hat{\sigma}_k$  is an order two automorphism of the ordered group  $(\mathbb{Z}^{m_k} \oplus \mathbb{Z}^{n_k}, \mathbb{Z}_+^{m_k} \oplus \mathbb{Z}^{n_k})$ . As in Step 3 of the proof of Theorem 3.3, one could find that

$$\sigma_{k+1} \circ \psi_k = \psi_k = \psi_k \circ \sigma_k.$$

It is standard to check that

$$\hat{\sigma}_k \circ \omega_k = \omega_k \circ \zeta_k,$$

so

$$\begin{aligned} S_{k+1} \circ \chi_k &= S_{k+1} \circ \omega_{k+1}^{-1} \circ (\psi_k \oplus \beta_k) \\ &= \omega_{k+1}^{-1} \circ \hat{\sigma}_{k+1} \circ (\psi_k \oplus \beta_k) \\ &= \omega_{k+1}^{-1} \circ (\sigma_{k+1} \oplus \eta_{k+1}) \circ (\psi_k \oplus \beta_k) \\ &= \omega_{k+1}^{-1} \circ ((\sigma_{k+1} \circ \psi_k) \oplus (\eta_{k+1} \circ \beta_k)) \\ &= \omega_{k+1}^{-1} \circ ((\psi_k \circ \sigma_k) \oplus (\beta_k \circ \eta_k)) \\ &= \omega_{k+1}^{-1} \circ (\psi_k \oplus \beta_k) \circ (\sigma_k \oplus \eta_k) \\ &= \chi_k \circ \hat{\sigma}_k. \end{aligned}$$

Therefore, we have the following commutative diagram:

$$\begin{array}{ccccccc} (\mathbb{Z} \oplus \mathbb{Z})^{n_1} \oplus \mathbb{Z}^{m_1-n_1} & \xrightarrow{\omega_1} & \mathbb{Z}^{m_1} \oplus \mathbb{Z}^{n_1} & \xrightarrow{\chi_1} & (\mathbb{Z} \oplus \mathbb{Z})^{n_2} \oplus \mathbb{Z}^{m_2-n_2} & \xrightarrow{\omega_2} & \mathbb{Z}^{m_2} \oplus \mathbb{Z}^{n_2} \xrightarrow{\chi_2} \dots \\ \downarrow \zeta_1 & & \downarrow \hat{\sigma}_1 & & \downarrow \zeta_2 & & \downarrow \hat{\sigma}_2 \\ (\mathbb{Z} \oplus \mathbb{Z})^{n_1} \oplus \mathbb{Z}^{m_1-n_1} & \xrightarrow{\omega_1} & \mathbb{Z}^{m_1} \oplus \mathbb{Z}^{n_1} & \xrightarrow{\chi_1} & (\mathbb{Z} \oplus \mathbb{Z})^{n_2} \oplus \mathbb{Z}^{m_2-n_2} & \xrightarrow{\omega_2} & \mathbb{Z}^{m_2} \oplus \mathbb{Z}^{n_2} \xrightarrow{\chi_2} \dots \end{array}$$

Note that for each  $k \in \mathbb{N}$ ,

$$\omega_{k+1} \circ \chi_k = \psi_k \oplus \beta_k.$$

Thus, after telescoping the aforementioned commutative diagram, there exists an order two automorphism  $\hat{\sigma}$  of the ordered group  $G \oplus H$  such that for each  $k \in \mathbb{N}$ , the following diagram is commutative:

$$\begin{array}{ccccc} \mathbb{Z}^{m_k} \oplus \mathbb{Z}^{n_k} & \xrightarrow{\psi_k \oplus \beta_k} & \mathbb{Z}^{m_{k+1}} \oplus \mathbb{Z}^{n_{k+1}} & \xrightarrow{\psi_{k+1, \infty} \oplus \beta_{k+1, \infty}} & G \oplus H \\ \downarrow \hat{\sigma}_k & & \downarrow \hat{\sigma}_{k+1} & & \downarrow \hat{\sigma} \\ \mathbb{Z}^{m_k} \oplus \mathbb{Z}^{n_k} & \xrightarrow{\psi_k \oplus \beta_k} & \mathbb{Z}^{m_{k+1}} \oplus \mathbb{Z}^{n_{k+1}} & \xrightarrow{\psi_{k+1, \infty} \oplus \beta_{k+1, \infty}} & G \oplus H. \end{array}$$

Fix  $k \in \mathbb{N}$ . By  $(\star)$ ,

$$\begin{aligned}\widehat{\sigma}_{k+1}((\psi_k \oplus \beta_k)(g_k \oplus h_k)) &= \sigma_{k+1}(\psi_k(g_k)) \oplus \eta_{k+1}(\beta_k(h_k)) \\ &= \psi_k(g_k) \oplus \beta_k(\eta_k(h_k)).\end{aligned}$$

It follows that

$$\begin{aligned}\widehat{\sigma}(\psi_{k,\infty}(g_k) \oplus \beta_{k,\infty}(h_k)) &= \widehat{\sigma}((\psi_{k,\infty} \oplus \beta_{k,\infty})(g_k \oplus h_k)) \\ &= (\psi_{k,\infty} \oplus \beta_{k,\infty})(\widehat{\sigma}_k(g_k \oplus h_k)) \\ &= (\psi_{k+1,\infty} \oplus \beta_{k+1,\infty})((\psi_k \oplus \beta_k) \circ \widehat{\sigma}_k)(g_k \oplus h_k) \\ &= (\psi_{k+1,\infty} \oplus \beta_{k+1,\infty})((\widehat{\sigma}_{k+1} \circ (\psi_k \oplus \beta_k))(g_k \oplus h_k)) \\ &= (\psi_{k+1,\infty} \oplus \beta_{k+1,\infty})(\psi_k(g_k) \oplus \beta_k(\eta_k(h_k))) \\ &= \psi_{k,\infty}(g_k) \oplus \beta_{k,\infty}(\eta_k(h_k)) \\ &= \psi_{k,\infty}(g_k) \oplus \eta(\beta_{k,\infty}(h_k)) \\ &= \sigma(\psi_{k,\infty}(g_k) \oplus \beta_{k,\infty}(h_k)).\end{aligned}$$

Noting that

$$\bigcup_{k=1}^{\infty} \psi_{k,\infty}(G_k) \oplus \beta_{k,\infty}(H_k) = G \oplus H,$$

we have

$$\widehat{\sigma} = \sigma.$$

Set  $u_1 = \text{id}_{A_1}$ . For  $k \in \mathbb{N}$ ,  $1 \leq i \leq m_k$ ,  $1 \leq j \leq m_{k+1}$ , define  $u_{k+1,i,j}$  exactly as that in the proof of Theorem 3.3. Next, define

$$\begin{aligned}u_{k+1} &= \left( \bigoplus_{j=1}^{n_{k+1}} \left( \bigoplus_{i=1}^{m_k} u_{k+1,i,j} \otimes I_{2l(k,i)} \otimes 1_{C(X)} \right) \right) \\ &\quad \oplus \left( \bigoplus_{j=n_{k+1}+1}^{m_{k+1}} \left( \bigoplus_{i=1}^{m_k} u_{k+1,i,j} \otimes I_{2l(k,i)} \right) \right), \quad k \in \mathbb{N}.\end{aligned}$$

Exactly as that in Section 3,

$$\Phi_k \circ \rho_k = \text{Ad } u_{k+1} \circ \Theta_k = \text{Ad } u_{k+1} \circ \rho_{k+1} \circ \Phi_k \quad \text{and} \quad u_{k+1}^2 = 1_{A_{k+1}}, \quad k \in \mathbb{N}.$$

Set  $v_1 = \text{id}_{A_1}$ , and define  $v_{k+1} = u_{k+1} \Phi_k(v_k)$  inductively,  $k = 1, 2, \dots$ . Define

$$\alpha_k = \text{Ad } v_k \circ \rho_k, \quad k \in \mathbb{N}.$$

So one can easily construct the following commutative diagram:

$$\begin{array}{ccccccc} A_1 & \xrightarrow{\Phi_1} & A_2 & \xrightarrow{\Phi_2} & A_3 & \xrightarrow{\Phi_3} & \dots \longrightarrow A \\ \downarrow \alpha_1 & & \downarrow \alpha_2 & & \downarrow \alpha_3 & & \\ A_1 & \xrightarrow{\Phi_1} & A_2 & \xrightarrow{\Phi_2} & A_3 & \xrightarrow{\Phi_3} & \dots \longrightarrow A. \end{array}$$

As in the corresponding part of the proof of Theorem 3.3,  $\alpha_k^2 = \text{id}_{A_k}$ ,  $k \in \mathbb{N}$ . Hence, the automorphisms  $\alpha_k$  define

$$\alpha : A = \varinjlim_{k \rightarrow \infty} (A_k, \Phi_k) \rightarrow A = \varinjlim_{k \rightarrow \infty} (A_k, \Phi_k)$$

which is a symmetry. Moreover,

$$\alpha_{*0} = \varinjlim_{k \rightarrow \infty} (\alpha_k)_{*0} = \varinjlim_{k \rightarrow \infty} (\rho_k)_{*0} = \varinjlim_{k \rightarrow \infty} \zeta_k = \varinjlim_{k \rightarrow \infty} \hat{\sigma}_k = \hat{\sigma} = \sigma.$$

**Step 4.** Also as in Section 3, the action of  $\mathbb{Z}_2$  generated by  $\alpha$  has the tracial Rokhlin property. Note that the unit of  $\mathfrak{A}$  may not come from trivial projections in the AH inductive limit procedure. Hence, one could note that we have enlarged the algebra in each finite stage during the construction. Next, we will cut it down.

Since  $\sigma$  preserves the scale,

$$\sigma(\tilde{g} \oplus \tilde{h}) = \tilde{g} \oplus \eta(\tilde{h}) = \tilde{g} \oplus \tilde{h}.$$

Therefore,  $\eta(\tilde{h}) = \tilde{h}$ . Without loss of generality, we may assume that there exists  $h_1 \in \mathbb{Z}^{n_1}$  such that  $\beta_{1,\infty}(h_1) = \tilde{h}$ . Since  $\beta_{1,\infty}$  is injective and

$$\beta_{1,\infty}(\eta_1(\tilde{h}_1)) = \eta(\beta_{1,\infty}(\tilde{h}_1)) = \eta(\tilde{h}) = \tilde{h},$$

it follows that

$$\eta_1(\tilde{h}_1) = \tilde{h}_1.$$

Denote

$$\tilde{h}_1 = (\lambda_1, \dots, \lambda_{p_1}; \zeta_1, \dots, \zeta_{q_1}; \mu_1, \nu_1, \dots, \mu_{r_1}, \nu_{r_1}) \in \mathbb{Z}^{n_1}.$$

Since  $\eta_1(\tilde{h}_1) = \tilde{h}_1$ , we have

$$\zeta_1 = \dots = \zeta_{q_1} = 0, \quad \text{and} \quad \mu_j = \nu_j, \quad j = 1, \dots, r_1.$$

By the elementary fact of  $K$ -theory (e.g., [37, Exercise 11.2]), there exist rank one projections  $P_i \in M_2(C(X))$  such that

$$[P_{1,i}] = (1, \lambda_i) \in \mathbb{Z} \oplus \mathbb{Z} = K_0(C(X))$$

for  $1 \leq i \leq p_1$  and

$$[P_{1,i}] = (1, \mu_i) \in \mathbb{Z} \oplus \mathbb{Z} = K_0(C(X))$$

for  $i = p_1 + q_1 + 1, p_1 + q_1 + 3, \dots, p_1 + q_1 + 2r_1 - 1$ , where the first coordinate of  $\mathbb{Z} \oplus \mathbb{Z}$  denotes the rank part. For  $i = p_1 + q_1 + 2, p_1 + q_1 + 4, \dots, p_1 + q_1 + 2r_1$ , set  $P_{1,i} = P_{1,i-1}$ .

Set

$$\begin{aligned} P_1 = & \left( \bigoplus_{i=1}^{p_1} (P_{1,i} \oplus 1_{M_{l(1,i)-1}}) \right) \oplus \left( \bigoplus_{i=p_1+1}^{p_1+q_1} 1_{M_{l(1,i)}} \right) \\ & \oplus \left( \bigoplus_{i=p_1+q_1+1}^{p_1+q_1+2r_1} P_{1,i} \oplus 1_{M_{l(1,i)-1}} \right) \oplus \left( \bigoplus_{i=n_1+1}^{m_1} 1_{M_{l(1,i)}} \right). \end{aligned}$$

Then,

$$\varphi_{1,\infty}([P_1]) = (\psi_{1,\infty} \oplus \beta_{1,\infty})(\tilde{g}_1 \oplus \tilde{h}_1) = \tilde{g} \oplus \tilde{h},$$

and

$$\alpha_1(P_1) = \rho_1(P_1) = P_1.$$

Inductively, we define  $P_{k+1}$  as

$$P_{k+1} = \Phi_k(P_k), \quad k \in \mathbb{N}.$$

Noting that

$$P_1 \in A_1^{\alpha_1}, \quad \alpha_2 \circ \Phi_1 = \Phi_1 \circ \alpha_1,$$

it follows that  $P_2 \in A_2^{\alpha_2}$ ; similarly, one has  $P_k \in A_k^{\alpha_k}$ ,  $k \in \mathbb{N}$ .

For  $k \in \mathbb{N}$ , let  $\mathfrak{A}_k = P_k A_k P_k$ . Since  $P_k \in A_k^{\alpha_k}$ , it is routine to check that  $\Phi_k$  maps  $\mathfrak{A}_k$  onto  $\mathfrak{A}_{k+1}$  and  $\alpha_k$  maps  $\mathfrak{A}_k$  onto  $\mathfrak{A}_k$  and is also an order two automorphism of  $\mathfrak{A}_k$ . Let

$$\mathcal{A} = \lim_{k \rightarrow \infty} (\mathfrak{A}_k, \Phi_k).$$

Define  $P = \Phi_{1,\infty}(P_1)$ . Then, it is obvious that  $\mathcal{A} = PAP$ ; hence,  $\mathcal{A}$  is an AF-algebra.

According to the construction of  $P$ , it is easy to check that

$$(K_0(\mathcal{A}), K_0(\mathcal{A})_+, [1_{\mathcal{A}}]) = (G \oplus H, (G_+ \setminus \{0\}) \oplus H \cup \{0 \oplus 0\}, \tilde{g} \oplus \tilde{h}).$$

Therefore, by Elliott's classification theorem of AF-algebras (see, e.g., [30, Theorem 3.4.8]),  $\mathcal{A} = \mathfrak{A}$ . Again, since

$$P_1 \in A_1^{\alpha_1}, \quad \alpha \circ \Phi_{1,\infty} = \Phi_{\infty} \circ \alpha_1,$$

it follows that  $P \in A^\alpha$ . Therefore,  $\alpha$  is an order two automorphism of  $\mathfrak{A}$  with  $\alpha_{*0} = \sigma$ .

Finally, noting that  $P \in A^\alpha$ , by [33, Lemma 3.7], we deduce that the  $\mathbb{Z}_2$  action  $\alpha$  of  $\mathfrak{A}$  has the tracial Rokhlin property. Hence, by [33, Corollary 1.6 and Theorem 2.6],  $C^*(\mathbb{Z}_2, \mathfrak{A}, \alpha)$  is a unital simple, separable  $C^*$ -algebra with tracial rank zero. Since  $A_k$  is nuclear,  $\mathfrak{A}_k = P_k A_k P_k$  is a hereditary subalgebra of  $A_k$ ; hence,  $\mathfrak{A}_k$  is nuclear. Also, as the appropriate part in Step 4 of the proof of Theorem 3.3,  $C^*(\mathbb{Z}_2, \mathfrak{A}, \alpha)$  is a unital simple AH-algebra with no dimension growth. ■

**Remark 4.2.** Let  $(G, G_+)$  be a simple dimension group, that is, the ordered group of a simple AF-algebra  $B$ , and let  $H$  be a countable torsion-free abelian group. It is not hard to check that  $(G \oplus H, (G_+ \setminus \{0\}) \oplus H \cup \{(0, 0)\})$  is unperforated and satisfies the Riesz interpolation property. Hence, by the Effros–Handelman–Shen theorem [11], it is a simple dimension group.

**Remark 4.3.** Some examples which satisfy the  $K$ -theory setup of Theorem 4.1 could be found in [3, 10.11.3], [7, Section 1], [34, Examples 4.1 and 4.5].

**Corollary 4.4.** *Let  $\mathfrak{A}$  be a unital simple AF-algebra with a unique tracial state  $\tau$ , and assume that the following short exact sequence is split:*

$$0 \rightarrow \inf K_0(\mathfrak{A}) \rightarrow K_0(\mathfrak{A}) \rightarrow \tau_*(K_0(\mathfrak{A})) \rightarrow 0,$$

where  $\inf K_0(\mathfrak{A}) = \{x : x \in K_0(\mathfrak{A}), \tau_*(x) = 0\}$ . Let  $\sigma$  be an order two automorphism of the scaled ordered  $K_0(\mathfrak{A})$ . Then, there is an automorphism  $\alpha$  of  $\mathfrak{A}$  with  $\alpha_{*0} = \sigma$  and  $\alpha^2 = \text{id}$ .

*Proof.* Set

$$H := \inf K_0(\mathfrak{A}), \quad (G, G_+, \tilde{g}) := (\tau_*(K_0(\mathfrak{A})), \tau_*(K_0(\mathfrak{A})) \cap \mathbb{R}_+, 1).$$

As the short exact sequence is split,  $K_0(\mathfrak{A}) = G \oplus H$ . By [5, Theorem 3.9],

$$K_0(\mathfrak{A})_+ = \{g \oplus h : g \in G, g > 0, h \in H\} \cup \{0 \oplus 0\}.$$

For  $g \oplus h \in G \oplus H$ , set  $g_1 \oplus h_1 := \sigma(g \oplus h)$ . We claim that  $g_1 = g$ . Otherwise, since  $G$  is totally ordered, either  $g_1 > g$  or  $g > g_1$ . If  $g_1 > g$ , then  $g_1 \oplus h_1 > g \oplus h$ . Since  $\sigma$  is order preserving, it follows that

$$g \oplus h = \sigma(g_1 \oplus h_1) \geq \sigma(g \oplus h) = g_1 \oplus h_1.$$

This is a contradiction! If  $g > g_1$ , a similar argument yields the contradiction.

Set  $\eta := \sigma|_H$  which is an order two automorphism of  $H$ . Therefore, for  $g \oplus h \in G \oplus H$ ,

$$\sigma(g \oplus h) = g \oplus \eta(h).$$

Therefore, by Theorem 4.1, there is an automorphism  $\alpha$  of  $\mathfrak{A}$  with  $\alpha_{*0} = \sigma$  and  $\alpha^2 = \text{id}$ . ■

**Funding.** Yuanhang Zhang's research is supported in part by the National Natural Science Foundation of China (No. 12071174) and the Science and Technology Development Project of Jilin Province (No. 20190103028JH).

## References

- [1] S. Barlak and G. Szabó, [Rokhlin actions of finite groups on UHF-absorbing  \$C^\*\$ -algebras](#). *Trans. Amer. Math. Soc.* **369** (2017), no. 2, 833–859 Zbl [1371.46054](#) MR [3572256](#)
- [2] D. J. Benson, A. Kumjian, and N. C. Phillips, [Symmetries of Kirchberg algebras](#). *Canad. Math. Bull.* **46** (2003), no. 4, 509–528 Zbl [1079.46047](#) MR [2011390](#)
- [3] B. Blackadar, *K-theory for operator algebras*. Math. Sci. Res. Inst. Publ. 5, Springer, New York, 1986 Zbl [0597.46072](#) MR [859867](#)
- [4] B. Blackadar, [Symmetries of the CAR algebra](#). *Ann. of Math. (2)* **131** (1990), no. 3, 589–623 Zbl [0718.46024](#) MR [1053492](#)

- [5] B. E. Blackadar, [Traces on simple AF  \$C^\*\$ -algebras](#). *J. Functional Analysis* **38** (1980), no. 2, 156–168 Zbl 0443.46037 MR 587906
- [6] F. P. Boca, [The structure of higher-dimensional noncommutative tori and metric Diophantine approximation](#). *J. Reine Angew. Math.* **492** (1997), 179–219 Zbl 0884.46040 MR 1488068
- [7] O. Bratteli, G. A. Elliott, D. E. Evans, and A. Kishimoto, [Finite group actions on AF algebras obtained by folding the interval](#). *K-Theory* **8** (1994), no. 5, 443–464 Zbl 0821.46088 MR 1310287
- [8] J. W. Bunce and J. A. Deddens, [A family of simple  \$C^\*\$ -algebras related to weighted shift operators](#). *J. Functional Analysis* **19** (1975), 13–24 Zbl 0313.46047 MR 0365157
- [9] M. Dădărlat, G. Nagy, A. Némethi, and C. Pasnicu, [Reduction of topological stable rank in inductive limits of  \$C^\*\$ -algebras](#). *Pacific J. Math.* **153** (1992), no. 2, 267–276 Zbl 0809.46054 MR 1151561
- [10] S. Echterhoff, W. Lück, N. C. Phillips, and S. Walters, [The structure of crossed products of irrational rotation algebras by finite subgroups of  \$SL\_2\(\mathbb{Z}\)\$](#) . *J. Reine Angew. Math.* **639** (2010), 173–221 Zbl 1202.46081 MR 2608195
- [11] E. G. Effros, D. E. Handelman, and C. L. Shen, [Dimension groups and their affine representations](#). *Amer. J. Math.* **102** (1980), no. 2, 385–407 Zbl 0457.46047 MR 564479
- [12] G. A. Elliott, [On the classification of inductive limits of sequences of semisimple finite-dimensional algebras](#). *J. Algebra* **38** (1976), no. 1, 29–44 Zbl 0323.46063 MR 397420
- [13] G. A. Elliott, [On the classification of  \$C^\*\$ -algebras of real rank zero](#). *J. Reine Angew. Math.* **443** (1993), 179–219 Zbl 0809.46067 MR 1241132
- [14] G. A. Elliott and D. E. Evans, [The structure of the irrational rotation  \$C^\*\$ -algebra](#). *Ann. of Math. (2)* **138** (1993), no. 3, 477–501 Zbl 0847.46034 MR 1247990
- [15] G. A. Elliott and G. Gong, [On the classification of  \$C^\*\$ -algebras of real rank zero. II](#). *Ann. of Math. (2)* **144** (1996), no. 3, 497–610 Zbl 0867.46041 MR 1426886
- [16] G. A. Elliott and H. Su, [K-theoretic classification for inductive limit  \$Z\_2\$  actions on AF algebras](#). *Canad. J. Math.* **48** (1996), no. 5, 946–958 Zbl 0869.46037 MR 1414065
- [17] T. Fack and O. Maréchal, [Sur la classification des symétries des  \$C^\*\$ -algèbres UHF](#). *Canadian J. Math.* **31** (1979), no. 3, 496–523 Zbl 0361.46057 MR 536360
- [18] G. Gong, [On the classification of simple inductive limit  \$C^\*\$ -algebras. I. The reduction theorem](#). *Doc. Math.* **7** (2002), 255–461 Zbl 1024.46018 MR 2014489
- [19] G. Gong, H. Lin, and Z. Niu, [A classification of finite simple amenable  \$\mathbb{Z}\$ -stable  \$C^\*\$ -algebras, I:  \$C^\*\$ -algebras with generalized tracial rank one](#). *C. R. Math. Acad. Sci. Soc. R. Can.* **42** (2020), no. 3, 63–450 MR 4215379
- [20] G. Gong, H. Lin, and Z. Niu, [A classification of finite simple amenable  \$\mathbb{Z}\$ -stable  \$C^\*\$ -algebras, II:  \$C^\*\$ -algebras with rational generalized tracial rank one](#). *C. R. Math. Acad. Sci. Soc. R. Can.* **42** (2020), no. 4, 451–539 MR 4215380
- [21] D. Handelman and W. Rossmann, [Product type actions of finite and compact groups](#). *Indiana Univ. Math. J.* **33** (1984), no. 4, 479–509 Zbl 0559.46029 MR 749311
- [22] D. Handelman and W. Rossmann, [Actions of compact groups on AF  \$C^\*\$ -algebras](#). *Illinois J. Math.* **29** (1985), no. 1, 51–95 Zbl 0559.46028 MR 769758
- [23] R. H. Herman and V. F. R. Jones, [Period two automorphisms of UHF  \$C^\*\$ -algebras](#). *J. Functional Analysis* **45** (1982), no. 2, 169–176 Zbl 0511.46057 MR 647069
- [24] L. K. Hua and I. Reiner, [Automorphisms of the unimodular group](#). *Trans. Amer. Math. Soc.* **71** (1951), 331–348 Zbl 0045.30402 MR 43847
- [25] M. Izumi, [Finite group actions on  \$C^\*\$ -algebras with the Rohlin property. I](#). *Duke Math. J.* **122** (2004), no. 2, 233–280 Zbl 1067.46058 MR 2053753

- [26] M. Izumi, [Finite group actions on  \$C^\*\$ -algebras with the Rohlin property. II.](#) *Adv. Math.* **184** (2004), no. 1, 119–160 Zbl 1050.46049 MR 2047851
- [27] T. Katsura, [A construction of actions on Kirchberg algebras which induce given actions on their  \$K\$ -groups.](#) *J. Reine Angew. Math.* **617** (2008), 27–65 Zbl 1158.46042 MR 2400990
- [28] A. Kishimoto, [On the fixed point algebra of a UHF algebra under a periodic automorphism of product type.](#) *Publ. Res. Inst. Math. Sci.* **13** (1977/78), no. 3, 777–791 Zbl 0367.46058 MR 0500177
- [29] A. Kumjian, [An involutive automorphism of the Bunce-Deddens algebra.](#) *C. R. Math. Rep. Acad. Sci. Canada* **10** (1988), no. 5, 217–218 Zbl 0669.46028 MR 962104
- [30] H. Lin, [An introduction to the classification of amenable  \$C^\*\$ -algebras.](#) World Scientific Publishing, River Edge, NJ, 2001 Zbl 1013.46055 MR 1884366
- [31] H. Lin, [Classification of simple  \$C^\*\$ -algebras of tracial topological rank zero.](#) *Duke Math. J.* **125** (2004), no. 1, 91–119 Zbl 1068.46032 MR 2097358
- [32] H. Lin and H. Osaka, [Tracial Rokhlin property for automorphisms on simple AT-algebras.](#) *Ergodic Theory Dynam. Systems* **28** (2008), no. 4, 1215–1241 Zbl 1444.46043 MR 2437228
- [33] N. C. Phillips, [The tracial Rokhlin property for actions of finite groups on  \$C^\*\$ -algebras.](#) *Amer. J. Math.* **133** (2011), no. 3, 581–636 Zbl 1225.46049 MR 2808327
- [34] N. C. Phillips, [Finite cyclic group actions with the tracial Rokhlin property.](#) *Trans. Amer. Math. Soc.* **367** (2015), no. 8, 5271–5300 Zbl 1328.46058 MR 3347172
- [35] I. F. Putnam, [On the topological stable rank of certain transformation group  \$C^\*\$ -algebras.](#) *Ergodic Theory Dynam. Systems* **10** (1990), no. 1, 197–207 Zbl 0667.46045 MR 1053808
- [36] M. Rørdam, [Classification of nuclear, simple  \$C^\*\$ -algebras.](#) In *Classification of nuclear  $C^*$ -algebras. Entropy in operator algebras*, pp. 1–145, Encyclopaedia Math. Sci. 126, Springer, Berlin, 2002 Zbl 1016.46037 MR 1878882
- [37] M. Rørdam, F. Larsen, and N. Laustsen, [An introduction to  \$K\$ -theory for  \$C^\*\$ -algebras.](#) London Math. Soc. Stud. Texts 49, Cambridge University Press, Cambridge, 2000 Zbl 0967.19001 MR 1783408
- [38] J. Spielberg, [Non-cyclotomic presentations of modules and prime-order automorphisms of Kirchberg algebras.](#) *J. Reine Angew. Math.* **613** (2007), 211–230 Zbl 1155.46023 MR 2377136
- [39] H. Thiel, [Future targets in the classification program for amenable  \$C^\*\$ -algebras.](#) 2017, <https://www.birs.ca/workshops/2017/17w5127/files/>
- [40] A. Tikuisis, S. White, and W. Winter, [Quasidiagonality of nuclear  \$C^\*\$ -algebras.](#) *Ann. of Math. (2)* **185** (2017), no. 1, 229–284 Zbl 1367.46044 MR 3583354
- [41] S. Walters, [On the inductive limit structure of order four automorphisms of the irrational rotation algebra.](#) *Internat. J. Math.* **17** (2006), no. 1, 107–117 Zbl 1096.46034 MR 2204842
- [42] D. P. Williams, [Crossed products of  \$C^\*\$ -algebras.](#) Math. Surveys Monogr. 134, American Mathematical Society, Providence, RI, 2007 Zbl 1119.46002 MR 2288954
- [43] Y. Zhang, [On a lifting question of Blackadar.](#) *Ann. Funct. Anal.* **9** (2018), no. 4, 485–499 Zbl 1458.46060 MR 3871909

Received 23 April 2021; revised 27 June 2021.

**Yuanhang Zhang**

School of Mathematics, Jilin University, Changchun 130012, P. R. China;  
[zhangyuanhang@jlu.edu.cn](mailto:zhangyuanhang@jlu.edu.cn)