

A quantitative relative index theorem and Gromov's conjectures on positive scalar curvature

Zhizhang Xie

Abstract. In this paper, we prove a quantitative relative index theorem. It provides a conceptual framework for studying some conjectures and open questions of Gromov on positive scalar curvature. More precisely, we prove a λ -Lipschitz rigidity theorem for (possibly incomplete) Riemannian metrics on spheres with certain types of subsets removed. This λ -Lipschitz rigidity theorem is asymptotically optimal. As a consequence, we obtain an asymptotically optimal λ -Lipschitz rigidity theorem for positive scalar curvature metrics on hemispheres. These give positive answers to the corresponding open questions raised by Gromov. As another application, we prove Gromov's \square^{n-m} inequality on the bound of distances between opposite faces of spin manifolds with cube-like boundaries with a suboptimal constant. As immediate consequences, this implies Gromov's cube inequality on the bound of widths of Riemannian cubes and Gromov's conjecture on the bound of widths of Riemannian bands with suboptimal constants. Further geometric applications will be discussed in a forthcoming paper.

1. Introduction

In the past several years, Gromov has formulated an extensive list of conjectures and open questions on scalar curvature [9, 10]. This has given rise to new perspectives on scalar curvature and inspired a wave of recent activity in this area [4–6, 9–11, 13, 18, 21, 31, 36, 37]. In this paper, we develop a quantitative relative index theorem that serves as a conceptual framework for solving some of these conjectures and open questions.

For example, we answer the following conjecture of Gromov in the spin case for all dimensions with a suboptimal constant.

Conjecture 1 (Gromov's \square^{n-m} inequality, [10, Section 5.3]). *Let (X, g) be an n -dimensional compact connected orientable manifold with boundary and \underline{X}_\bullet a closed orientable manifold of dimension $n - m$. Suppose*

$$f: X \rightarrow [-1, 1]^m \times \underline{X}_\bullet$$

is a continuous map, which sends the boundary of X to the boundary of $[-1, 1]^m \times \underline{X}_\bullet$ and which has nonzero degree. Let $\partial_{j\pm}$, $j = 1, \dots, m$, be the pullbacks of the pairs of

the opposite faces of the cube $[-1, 1]^m$ under the composition of f with the projection $[-1, 1]^m \times \underline{X}_\bullet \rightarrow [-1, 1]^m$. Assume that for any m hypersurfaces $Y_j \subset X$ that separate ∂_{j-} from ∂_{j+} with $1 \leq j \leq m$, their transversal intersection $Y_{\cap} \subset X$ does not admit a metric with positive scalar curvature; furthermore, the products $Y_{\cap} \times T^k$ of Y_{\cap} and k -dimensional tori do not admit metrics with positive scalar curvature either. If $\text{Sc}(g) \geq n(n - 1)$, then the distances $\ell_j = \text{dist}(\partial_{j-}, \partial_{j+})$ satisfy the following inequality:

$$\sum_{j=1}^m \frac{1}{\ell_j^2} \geq \frac{n^2}{4\pi^2}.$$

Consequently, we have

$$\min_{1 \leq j \leq m} \text{dist}(\partial_{j-}, \partial_{j+}) \leq \sqrt{m} \frac{2\pi}{n}.$$

Here, if (X, g) is a manifold with Riemannian metric g , then $\text{Sc}(g)$ stands for the scalar curvature of g . Sometimes, we also write $\text{Sc}(X)$ for the scalar curvature of g if it is clear from the context which metric we are referring to. In [10], Gromov gave a proof of Conjecture 1 in dimension ≤ 8 by the minimal surface method [24–26]. In dimension ≥ 9 , Gromov provided an approach towards a proof of Conjecture 1 based on unpublished results of Lohkamp [20] or a generalization of Schoen–Yau’s result [26]. He stated that the above inequalities should be regarded as conjectural in dimension ≥ 9 , cf. [10, Section 5.2, page 250].

The conditions in Conjecture 1 may appear technical at the first glance. The following special case probably makes it clearer what kind of geometric problems we are dealing with here.

Conjecture 2 (Gromov’s \square^n inequality, [10, Section 3.8]). *Let g be a Riemannian metric on the cube $I^n = [0, 1]^n$. If $\text{Sc}(g) \geq n(n - 1)$, then*

$$\sum_{j=1}^n \frac{1}{\ell_j^2} \geq \frac{n^2}{4\pi^2},$$

where $\ell_j = \text{dist}(\partial_{j-}, \partial_{j+})$ is the g -distance between the pair of opposite faces ∂_{j-} and ∂_{j+} of the cube. Consequently, we have

$$\min_{1 \leq j \leq n} \text{dist}(\partial_{j-}, \partial_{j+}) \leq \frac{2\pi}{\sqrt{n}}.$$

One of the key ingredients for the proof of Conjecture 1 is the following quantitative relative index theorem.

Theorem I (cf. Theorem 3.5). *Let Z_1 and Z_2 be two closed n -dimensional Riemannian manifolds and S_j a Euclidean Cl_n -bundle¹ over Z_j for $j = 1, 2$. Suppose D_j is a Cl_n -linear Dirac-type operator acting on S_j over Z_j . Let \tilde{Z}_j be a Galois Γ -covering space of Z_j and \tilde{D}_j the lift of D_j . Let X_j be a subset of Z_j and \tilde{X}_j the preimage of X_j under*

¹Here Cl_n is the real Clifford algebra of \mathbb{R}^n . See [17, Chapter II.§7 and Chapter III.§10] for more details on Cl_n -vector bundles and the Clifford index of Cl_n -linear Dirac operators.

the covering map $\tilde{Z}_j \rightarrow Z_j$. Denote by $N_r(Z_j \setminus X_j)$ the open r -neighborhood of $Z_j \setminus X_j$. Suppose there is $r > 0$ such that all geometric data on $N_r(Z_1 \setminus X_1)$ and $N_r(Z_2 \setminus X_2)$ coincide, i.e., there is an orientation preserving Riemannian isometry $\Phi: N_r(Z_1 \setminus X_1) \rightarrow N_r(Z_2 \setminus X_2)$ such that Φ lifts to an isometric Cl_n -bundle isomorphism $\Phi: \mathcal{S}_1|_{N_r(Z_1 \setminus X_1)} \rightarrow \mathcal{S}_2|_{N_r(Z_2 \setminus X_2)}$. Assume that

(1) there exists $\sigma > 0$ such that

$$\mathcal{R}_j(x) \geq \frac{(n-1)\sigma^2}{n}$$

for all $x \in X_j$, where \mathcal{R}_j is the curvature term appearing in $D_j^2 = \nabla^* \nabla + \mathcal{R}_j$,

(2) and $D_1 = \Phi^{-1} D_2 \Phi$ on $N_r(Z_1 \setminus X_1)$.

Then there exists a universal constant $C > 0$ such that if $\sigma \cdot r > C$, then we have

$$\text{Ind}_\Gamma(\tilde{D}_1) - \text{Ind}_\Gamma(\tilde{D}_2) = 0$$

in $KO_n(C_{\max}^*(\Gamma; \mathbb{R}))$, where $\text{Ind}_\Gamma(\tilde{D}_j)$ denotes the maximal higher index of \tilde{D}_j and $C_{\max}^*(\Gamma; \mathbb{R})$ is the maximal group C^* -algebra of Γ with real coefficients.

The numerical estimates in Appendix B show that the universal constant C is ≤ 40.65 . As an application of the above quantitative relative index theorem, we solve Gromov’s \square^{n-m} inequality (Conjecture I) for all dimensions with a suboptimal constant. More precisely, we have the following theorem.

Theorem II (cf. Theorem 4.3). *Let X be an n -dimensional compact connected spin manifold with boundary. Suppose $f: X \rightarrow [-1, 1]^m$ is a continuous map that sends the boundary of X to the boundary of $[-1, 1]^m$. Let $\partial_{j\pm}$, $j = 1, \dots, m$, be the pullbacks of the pairs of the opposite faces of the cube $[-1, 1]^m$. Suppose Y_\natural is an $(n - m)$ -dimensional closed submanifold (without boundary) in X that satisfies the following conditions:*

- (1) $\pi_1(Y_\natural) \rightarrow \pi_1(X)$ is injective;
- (2) Y_\natural is the transversal intersection² of m orientable hypersurfaces $\{Y_j\}_{1 \leq j \leq m}$ of X , each of which separates ∂_{j-} from ∂_{j+} ;
- (3) the higher index $\text{Ind}_\Gamma(D_{Y_\natural})$ does not vanish in $KO_{n-m}(C_{\max}^*(\Gamma; \mathbb{R}))$, where $\Gamma = \pi_1(Y_\natural)$.

If $\text{Sc}(X) \geq n(n - 1)$, then the distances $\ell_j = \text{dist}(\partial_{j-}, \partial_{j+})$ satisfy the following inequality:

$$\sum_{j=1}^m \frac{1}{\ell_j^2} \geq \frac{n^2}{\left(\frac{8}{\sqrt{3}}C + 4\pi\right)^2},$$

where C is the universal constant from Theorem I. Consequently, we have

$$\min_{1 \leq j \leq m} \text{dist}(\partial_{j-}, \partial_{j+}) \leq \sqrt{m} \frac{\frac{8}{\sqrt{3}}C + 4\pi}{n}.$$

²In particular, this implies that the normal bundle of Y_\natural is trivial.

In a subsequent work, with Wang and Yu [30], the author proves Theorem II with the optimal constant via a different method, hence completely solves Conjecture 1 (in spin case) and Conjecture 2 for all dimensions. We point out that Cecchini [4] and Zeidler [36, 37] proved a special case of Theorem II when $m = 1$ with the optimal constant.

For spin manifolds, the assumptions on Y_{\hbar} in Theorem II above are (stably) equivalent to the assumptions in Conjecture 1, provided that the (stable) Gromov–Lawson–Rosenberg conjecture holds for $\Gamma = \pi_1(Y_{\hbar})$. See the survey paper of Rosenberg and Stolz [23] for more details. The stable Gromov–Lawson–Rosenberg conjecture for Γ follows from the strong Novikov conjecture for Γ , where the latter has been verified for a large class of groups including all word hyperbolic groups [7], all groups acting properly and isometrically on simply connected and non-positively curved manifolds [15], all subgroups of linear groups [12], and all groups that are coarsely embeddable into a Hilbert space [34].

As a special case of Theorem II, we have the following theorem, which proves Gromov’s \square^n -inequality (Conjecture 2) with a suboptimal constant.

Theorem III. *Let g be a Riemannian metric on the cube $I^n = [0, 1]^n$. If $\text{Sc}(g) \geq n(n - 1)$, then*

$$\sum_{j=1}^n \frac{1}{\ell_j^2} \geq \frac{n^2}{\left(\frac{8}{\sqrt{3}}C + 4\pi\right)^2},$$

where $\ell_j = \text{dist}(\partial_{j-}, \partial_{j+})$ is the g -distance between the pair of opposite faces ∂_{j-} and ∂_{j+} of the cube, and C is the universal constant from Theorem I. Consequently, we have

$$\min_{1 \leq j \leq n} \text{dist}(\partial_{j-}, \partial_{j+}) \leq \frac{\frac{8}{\sqrt{3}}C + 4\pi}{\sqrt{n}}.$$

Proof. Note that the higher index of the Dirac operator on a single point is a generator of $KO_0(\{e\}) = \mathbb{Z}$, hence does not vanish. If X is the cube $I^n = [0, 1]^n$ with the given Riemannian metric g , then the assumptions of Theorem II are satisfied. Hence the theorem follows from Theorem II. ■

As pointed out by Gromov in [10, Section 3.8], Theorem III has the following immediate corollary. Recall that a map $\varphi: (X, g) \rightarrow (Y, h)$ between two metric spaces is said to be λ -Lipschitz if

$$\text{dist}_h(\varphi(x_1), \varphi(x_2)) \leq \lambda \cdot \text{dist}_g(x_1, x_2)$$

for all $x_1, x_2 \in X$.

Corollary 1.1. *Let (X, g_0) be the standard unit hemisphere \mathbb{S}_+^n . If X admits a Riemannian metric g such that*

- (1) *there is a λ_n -Lipschitz homeomorphism $\varphi: (X, g) \rightarrow (X, g_0)$,*
- (2) *and $\text{Sc}(g) \geq n(n - 1) = \text{Sc}(g_0)$,*

then

$$\lambda_n \geq \frac{2 \arcsin\left(\frac{1}{\sqrt{n}}\right)}{\frac{1}{\sqrt{n}}\left(\frac{8}{\sqrt{3}}C + 4\pi\right)} > \frac{1}{\frac{4}{\sqrt{3}}C + 2\pi},$$

where C is the same universal constant from Theorem I.

As another application of our quantitative relative index theorem, we prove the following λ -Lipschitz rigidity theorem for positive scalar curvature metrics on spheres with certain subsets removed. This gives a positive answer to a corresponding open question of Gromov, cf. [9, page 687, specific problem].

Theorem IV (cf. Theorem 5.8). *Let Σ be a subset of the standard unit sphere \mathbb{S}^n contained in a geodesic ball of radius $r < \frac{\pi}{2}$. Let (X, g_0) be the standard unit sphere \mathbb{S}^n minus Σ . If a (possibly incomplete) Riemannian metric g on X satisfies that*

- (1) *there is a λ_n -Lipschitz homeomorphism $\varphi: (X, g) \rightarrow (X, g_0)$,*
- (2) *and $\text{Sc}(g) \geq n(n - 1) = \text{Sc}(g_0)$,*

then

$$\lambda_n \geq \sqrt{1 - \frac{C_r}{n^2}},$$

where³

$$C_r = \frac{8C^2}{\left(\frac{\pi}{2} - r\right)^2}$$

and C is the universal constant from Theorem I. Consequently, the lower bound for λ_n approaches 1, as $n \rightarrow \infty$.

This λ -Lipschitz rigidity theorem is asymptotically optimal in the sense that the lower bound for λ_n becomes sharp, as $n = \dim \mathbb{S}^n \rightarrow \infty$. In the case where $n = \dim \mathbb{S}^n$ is odd, an analogue of Theorem IV also holds for subsets that are contained in a pair of antipodal geodesic balls of radius $r < \frac{\pi}{6}$. We refer the reader to Theorem 5.10 for the precise details. We point out that when $\Sigma = \emptyset$, that is, when (X, g_0) is the standard unit sphere \mathbb{S}^n itself, it is a theorem of Llarul that $\lambda_n \geq 1$ for all $n \geq 2$ [19, Theorem A]. Furthermore, when Σ is either a single point or a pair of antipodal points, Gromov showed that $\lambda_n \geq 1$ when $3 \leq n \leq 8$ [10, Section 3.9].

As a consequence of Theorem IV, we have the following λ -Lipschitz rigidity result for hemispheres. This answers (asymptotically) an open question of Gromov on the sharpness of the constant λ_n for the λ -Lipschitz rigidity of positive scalar curvature metrics on hemispheres [10, Section 3.8].

³If $n = \dim \mathbb{S}^n$ is odd, our proof of Theorem IV in fact shows that we can improve C_r to be $\frac{4C^2}{\left(\frac{\pi}{2} - r\right)^2}$ instead of $\frac{8C^2}{\left(\frac{\pi}{2} - r\right)^2}$.

Theorem V (cf. Theorem 5.11). *Let (X, g_0) be the standard unit hemisphere \mathbb{S}_+^n . If a Riemannian metric g on X satisfies that*

- (1) *there is a λ_n -Lipschitz homeomorphism $\varphi: (X, g) \rightarrow (X, g_0)$,*
- (2) *and $\text{Sc}(g) \geq n(n - 1) = \text{Sc}(g_0)$,*

then

$$\lambda_n \geq \left(1 - \sin \frac{\pi}{\sqrt{n}}\right) \sqrt{1 - \frac{8C^2}{\pi^2 n}}$$

where C is the universal constant from Theorem I. Consequently, the lower bound for λ_n approaches 1, as $n \rightarrow \infty$.

The above theorem is asymptotically optimal in the sense that the lower bound for λ_n becomes sharp, as $n = \dim \mathbb{S}^n \rightarrow \infty$. In particular, it improves the lower bound for λ_n in Corollary 1.1 when $n = \dim \mathbb{S}^n$ is large.

A key geometric concept behind the proof of Theorem IV is the following notion of wrapping property for subsets of \mathbb{S}^n .

Definition 1.2. Let Z be a path-connected metric space. A subset Σ of Z is called strongly non-separating if $Z \setminus N_\varepsilon(\Sigma)$ is path-connected for all sufficiently small $\varepsilon > 0$, where $N_\varepsilon(\Sigma)$ is the open ε -neighborhood of Σ .

Definition 1.3 (Subsets with the wrapping property). A subset Σ of the standard unit sphere \mathbb{S}^n is said to have the wrapping property if Σ is strongly non-separating and furthermore there exists a smooth distance-contracting⁴ map $\Phi: \mathbb{S}^n \rightarrow \mathbb{S}^n$ such that the following are satisfied:

- (1a) if n is even, Φ equals the identity map on $N_\varepsilon(\Sigma)$;
- (1b) if n is odd, Φ equals either the identity map or the antipodal map on each of the connected components of $N_\varepsilon(\Sigma)$;
- (2) and⁵ $\deg(\Phi) \neq 1$.

Note that the conditions for satisfying the wrapping property in the odd dimensional case are slightly weaker than those in the even dimensional case. Roughly speaking, a subset $\Sigma \subset \mathbb{S}^n$ has the wrapping property if its geometric size is “relatively small”. For example, if Σ is a strongly non-separating subset of the standard unit sphere \mathbb{S}^n that is contained in a geodesic ball of radius $< \frac{\pi}{2}$, then Σ has the wrapping property (cf. Lemma 5.3). Moreover, a strongly non-separating subset of an odd dimensional sphere that is contained in a pair of antipodal geodesic balls of radius $< \frac{\pi}{6}$ also satisfies the wrapping property (cf. Lemma 5.5).

Motivated by the theorems of Llarul and Gromov and the results in the current paper, we conclude this introduction by the following open question.

⁴Recall that a smooth map $\psi: X \rightarrow Y$ between Riemannian manifolds is said to be distance-contracting if it is 1-Lipschitz, that is, $\|\psi_*(v)\| \leq \|v\|$ for all tangent vectors $v \in TX$.

⁵For example, if Φ is not surjective, then clearly $\deg(\Phi) = 0 \neq 1$.

Open Question (Rigidity for positive scalar curvature metrics on $\mathbb{S}^n \setminus \Sigma$). *Let Σ be a subset with the wrapping property in the standard unit sphere \mathbb{S}^n . Let (X, g_0) be the standard unit sphere \mathbb{S}^n minus Σ . If a (possibly incomplete) Riemannian metric g on X satisfies*

- (1) $g \geq g_0$,
- (2) and $\text{Sc}(g) \geq n(n - 1) = \text{Sc}(g_0)$,

then does it imply that $g = g_0$?

The paper is organized as follows. In Section 2, we review the construction of some standard geometric C^* -algebras and the construction of higher index. In Section 3, we prove our quantitative relative index theorem (Theorem 3.4 and Theorem 3.5). Finally, we apply the quantitative relative index theorem to prove Theorems II–V in Section 4 and Section 5.

2. Preliminaries

In this section, we review the construction of some standard geometric C^* -algebras and the construction of higher indices (cf. [32]).

Let X be a proper metric space, i.e., every closed ball in X is compact. An X -module is a Hilbert space H equipped with a $*$ -representation $\rho: C_0(X) \rightarrow \mathcal{B}(H)$ of $C_0(X)$, where $\mathcal{B}(H)$ is the algebra of all bounded linear operators on H . An X -module H is called non-degenerate if the $*$ -representation of $C_0(X)$ is non-degenerate, that is, $\rho(C_0(X))H$ is dense in H . An X -module is called ample if no nonzero function in $C_0(X)$ acts as a compact operator.

Assume that a discrete group Γ acts freely and cocompactly⁶ on X by isometries and H_X is a non-degenerate ample X -module equipped with a covariant unitary representation of Γ . If we denote by ρ and π the representations of $C_0(X)$ and Γ respectively, this means

$$\pi(\gamma)(\rho(f)v) = \rho(\gamma^* f)(\pi(\gamma)v),$$

where $f \in C_0(X)$, $\gamma \in \Gamma$, $v \in H_X$ and $\gamma^* f(x) = f(\gamma^{-1}x)$. In this case, we call (H_X, Γ, ρ) a covariant system of (X, Γ) .

Definition 2.1. Let (H_X, Γ, ρ) be a covariant system of (X, Γ) and T a Γ -equivariant bounded linear operator acting on H_X .

- (1) The propagation of T is defined to be the supremum

$$\sup\{\text{dist}(x, y) : (x, y) \in \text{supp}(T)\},$$

⁶More generally, with appropriate modifications, all constructions in this section have their obvious analogues for the case of proper and cocompact actions instead of free and cocompact actions, cf. [35, Section 2].

where $\text{supp}(T)$ is the complement of points $(x, y) \in X \times X$ for which there exist $f, g \in C_0(X)$ such that $gTf = 0$ and $f(x) \neq 0, g(y) \neq 0$;

(2) T is said to be locally compact if fT and Tf are compact for all $f \in C_0(X)$.

We recall the definition of equivariant Roe algebras.

Definition 2.2. Let X be a locally compact metric space with a free and cocompact isometric action of Γ . Let (H_X, Γ, ρ) be a covariant system. We define $\mathbb{C}[X]^\Gamma$ to be the $*$ -algebra of Γ -equivariant locally compact finite propagation operators in $\mathcal{B}(H_X)$. The equivariant Roe algebra $C_r^*(X)^\Gamma$ is defined to be the completion of $\mathbb{C}[X]^\Gamma$ in $\mathcal{B}(H_X)$ under the operator norm.

There is also a maximal version of equivariant Roe algebras.

Definition 2.3. For an operator $T \in \mathbb{C}[X]^\Gamma$, its *maximal norm* is

$$\|T\|_{\max} := \sup_{\varphi} \{ \|\varphi(T)\| : \varphi: \mathbb{C}[X]^\Gamma \rightarrow \mathcal{B}(H) \text{ is a } * \text{-representation} \}.$$

The maximal equivariant Roe algebra $C_{\max}^*(X)^\Gamma$ is defined to be the completion of $\mathbb{C}[X]^\Gamma$ with respect to $\|\cdot\|_{\max}$.

We know

$$C_r^*(X)^\Gamma \cong C_r^*(\Gamma) \otimes \mathcal{K} \quad \text{and} \quad C_{\max}^*(X)^\Gamma \cong C_{\max}^*(\Gamma) \otimes \mathcal{K},$$

where $C_r^*(\Gamma)$ (resp. $C_{\max}^*(\Gamma)$) is the reduced (resp. maximal) group C^* -algebra of Γ and \mathcal{K} is the algebra of compact operators.

Furthermore, there are also real versions of reduced and maximal equivariant Roe algebras, by using real Hilbert spaces instead of complex Hilbert spaces. We shall denote these algebras by $C_r^*(X)_{\mathbb{R}}^\Gamma$ and $C_{\max}^*(X)_{\mathbb{R}}^\Gamma$. Similarly, we have

$$C_r^*(X)_{\mathbb{R}}^\Gamma \cong C_r^*(\Gamma; \mathbb{R}) \otimes \mathcal{K}_{\mathbb{R}} \quad \text{and} \quad C_{\max}^*(X)_{\mathbb{R}}^\Gamma \cong C_{\max}^*(\Gamma; \mathbb{R}) \otimes \mathcal{K}_{\mathbb{R}},$$

where $C_r^*(\Gamma; \mathbb{R})$ (resp. $C_{\max}^*(\Gamma; \mathbb{R})$) is the reduced (resp. maximal) group C^* -algebra of Γ with real coefficients and $\mathcal{K}_{\mathbb{R}}$ is the algebra of compact operators on a real infinite dimensional Hilbert space.

Let us review the construction of the *higher index* of a first-order symmetric elliptic differential operator on a closed manifold. Suppose M is a closed Riemannian manifold. Let \tilde{M} be a Galois covering space of M whose deck transformation group is Γ . Suppose D is a symmetric elliptic differential operator acting on some vector bundle \mathcal{S} over M . In addition, if M is even dimensional, we assume \mathcal{S} to be $\mathbb{Z}/2$ -graded and D has odd-degree with respect to this $\mathbb{Z}/2$ -grading. Let \tilde{D} be the lift of D to \tilde{M} .

We choose a normalizing function χ , i.e., a continuous odd function $\chi: \mathbb{R} \rightarrow \mathbb{R}$ such that

$$\lim_{x \rightarrow \pm\infty} \chi(x) = \pm 1.$$

By the standard theory of elliptic operators on complete manifolds, \tilde{D} is essentially self-adjoint and $F = \chi(\tilde{D})$ obtained by functional calculus satisfies the condition

$$F^2 - 1 \in C_r^*(\tilde{M})^\Gamma \cong C_r^*(\Gamma) \otimes \mathcal{K}.$$

In the even dimensional case, since we assume \mathcal{S} to be $\mathbb{Z}/2$ -graded and D has odd-degree with respect to this $\mathbb{Z}/2$ -grading, we have

$$D = \begin{pmatrix} 0 & D^- \\ D^+ & 0 \end{pmatrix}.$$

In particular, it follows that

$$F = \begin{pmatrix} 0 & U \\ V & 0 \end{pmatrix}$$

for some U and V such that $UV - 1 \in C_r^*(\tilde{M})^\Gamma$ and $VU - 1 \in C_r^*(\tilde{M})^\Gamma$. Define the following invertible element:

$$W := \begin{pmatrix} 1 & U \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -V & 1 \end{pmatrix} \begin{pmatrix} 1 & U \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

and form the idempotent

$$p = W \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} W^{-1} = \begin{pmatrix} UV(2 - UV) & (2 - UV)(1 - UV)U \\ V(1 - UV) & (1 - VU)^2 \end{pmatrix}. \tag{2.1}$$

Definition 2.4. In the even dimensional case, the higher index $\text{Ind}_\Gamma(\tilde{D})$ of \tilde{D} is defined to be

$$\text{Ind}_\Gamma(\tilde{D}) := [p] - \left[\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \right] \in K_0(C_r^*(\tilde{M})^\Gamma) \cong K_0(C_r^*(\Gamma)).$$

Note that if Γ is the trivial group, then the higher index $\text{Ind}_\Gamma(\tilde{D}) \in K_0(\mathcal{K}) = \mathbb{Z}$ is simply the classical Fredholm index $\text{Ind}(D)$ of D , where the latter is defined to be

$$\text{Ind}(D) := \dim \ker(D^+) - \dim \text{coker}(D^+).$$

The construction of higher index in the odd dimensional case is similar.

Definition 2.5. In the odd dimensional case, the higher index $\text{Ind}_\Gamma(\tilde{D})$ of \tilde{D} is defined to be

$$\text{Ind}_\Gamma(\tilde{D}) := \exp\left(2\pi i \frac{\chi(\tilde{D}) + 1}{2}\right) \in K_1(C_r^*(\tilde{M})^\Gamma) \cong K_1(C_r^*(\Gamma)).$$

The higher index of \tilde{D} , as a K -theory class, is independent of the choice of the normalizing function χ . In particular, if we choose χ to be a normalizing function whose distributional Fourier transform has compact support, then $F = \chi(\tilde{D})$ has finite propagation and consequently the formula for defining $\text{Ind}_\Gamma(\tilde{D})$ produces an element of finite

propagation,⁷ that is, an element in $\mathbb{C}[\widetilde{M}]^\Gamma$, which certainly also defines a K -theory class in $K_n(C_{\max}^*(\Gamma))$. We define this class $\text{Ind}_\Gamma(\widetilde{D}) \in K_n(C_{\max}^*(\Gamma))$ to be the maximal higher index of the operator \widetilde{D} .

The higher index of an elliptic operator with real coefficients is defined the same way, and it lies in $KO_n(C_r^*(\Gamma; \mathbb{R}))$ or $KO_n(C_{\max}^*(\Gamma; \mathbb{R}))$, when the elliptic operator is appropriately graded (e.g. $\mathcal{C}\ell_n$ -graded with respect to the real Clifford algebra $\mathcal{C}\ell_n$). See [17, Chapter II. §7].

3. A quantitative relative index theorem

In this section, we prove a quantitative relative index theorem (Theorem 3.4 and Theorem 3.5), which serves a conceptual framework for studying some conjectures and open questions on Riemannian metrics of positive scalar curvature proposed by Gromov in the past several years [9, 10].

Let Y be a complete n -dimensional Riemannian manifold and \mathcal{S} a Euclidean $\mathcal{C}\ell_n$ -bundle over Y . Suppose D is a first-order symmetric elliptic $\mathcal{C}\ell_n$ -linear differential operator acting on \mathcal{S} over Y . Recall the following lemma due to Roe [22, Lemma 2.5].

Lemma 3.1. *With the same notation as above, suppose there exist $\sigma > 0$ and a subset $K \subset Y$ such that*

$$\|Df\| \geq \sigma \|f\|$$

for all $f \in C_c^\infty(Y \setminus K, \mathcal{S})$. Given $\varphi \in \mathcal{S}(\mathbb{R})$, assume the Fourier transform $\hat{\varphi}$ of φ is supported in $(-r, r)$. If $\psi \in C_0(Y)$ has support disjoint from the $2r$ -neighborhood of K , then

$$\|\varphi(D)\rho(\psi)\|_{\text{op}} \leq \|\psi\| \sup\{|\varphi(y)| : |y| \geq \sigma\}.$$

Here $\rho(\psi)$ is the bounded operator on $L^2(Y, \mathcal{S})$ given by multiplication of ψ , and $\|\psi\|$ is the supremum norm of ψ .

Roe’s proof of the above lemma makes use of the Friedrichs extension of D^2 on $Y \setminus K$ in an essential way. In Appendix A, we shall construct analogues of the Friedrichs extension in the maximal group C^* -algebra setting, which allows us to extend Roe’s lemma above to the corresponding maximal setting (by following essentially the same proof of Roe [22, Lemma 2.5]).

Let us consider the case where $n = \dim Y$ is even. Consider a normalizing function⁸ $\chi: \mathbb{R} \rightarrow \mathbb{R}$ such that the distributional Fourier transform $\hat{\chi}$ of χ is supported on $[-1, 1]$.

⁷In the odd dimensional case, one can approximate $\exp(2\pi i \frac{\chi(\widetilde{D})+1}{2})$ by a finite propagation element, since the coefficients in the power series expansion for the function $e^{2\pi i t}$ decay very fast (faster than any exponential decay, to be more precise).

⁸A normalizing function is a continuous odd function $\chi: \mathbb{R} \rightarrow \mathbb{R}$ such that $\lim_{x \rightarrow \pm\infty} \chi(x) = \pm 1$.

Let $F_t = \chi(tD)$. Since D is assumed to have odd-degree with respect to the $\mathbb{Z}/2$ -grading on \mathcal{S} , we have

$$D = \begin{pmatrix} 0 & D^- \\ D^+ & 0 \end{pmatrix}.$$

In particular, it follows that

$$F_t = \begin{pmatrix} 0 & U_t \\ V_t & 0 \end{pmatrix}$$

for some U_t and V_t . As in line (2.1), we define

$$p_t = \begin{pmatrix} U_t V_t (2 - U_t V_t) & (2 - U_t V_t)(1 - U_t V_t) U_t \\ V_t (1 - U_t V_t) & (1 - V_t U_t)^2 \end{pmatrix}. \tag{3.1}$$

The following lemma will be useful in the proof of Theorem 3.4.

Lemma 3.2. *With the same notation as above, the following hold for all $t > 0$.*

- (1) *There exists $\beta > 0$ such that $\|p_t\| \leq \beta$ for all $t > 0$.*
- (2) *The propagation $\text{prop}(p_t)$ of p_t is $\leq 5t$.*
- (3) *If there exist $\sigma > 0$ and a closed subset $K \subset Y$ such that*

$$\|Df\| \geq \sigma \|f\|$$

for all $f \in C_c^\infty(Y \setminus K, \mathcal{S})$, then there exists $\eta > 0$ such that

$$\left\| \left(p_t - \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \right) f \right\| \leq \frac{\eta}{t}$$

for all $f \in C_c^\infty(Y \setminus N_{10t}(K), \mathcal{S})$, where $N_{10t}(K)$ is the $10t$ -neighborhood of K .

Proof. Clearly, there exists $\alpha > 0$ such that $|\chi|$ is uniformly bounded by α . Hence both U_t and V_t have operator norm $\leq \alpha$. Therefore, part (1) follows from the explicit formula of p_t in line (3.1).

The distributional Fourier transform $\hat{\chi}$ of χ is supported on $[-1, 1]$. By the inverse Fourier transform formula

$$\chi(D) = \frac{1}{2\pi} \int \hat{\chi}(\xi) e^{i\xi D} d\xi$$

and the finite propagation of the wave operator $e^{i\xi D}$, we see that $\chi(D)$ has propagation no more than 1. Replacing $\chi(x)$ by $\chi(tx)$, we see that the propagation of $\chi(tD)$ is $\leq t$. In particular, U_t and V_t also have propagation $\leq t$. Hence, part (2) follows from the explicit formula of p_t in line (3.1). Furthermore, part (3) follows Lemma 3.1. This finishes the proof. ■

Let e be an element in a Banach algebra such that

$$\|e^2 - e\| < 1/4.$$

Then the spectrum of e is disjoint from the vertical line $\{\frac{1}{2} + iy : y \in \mathbb{R}\}$. Let Σ be the part of spectrum of e that is to the right of the line $\{\frac{1}{2} + iy : y \in \mathbb{R}\}$. Choose a contour γ containing Σ but disjoint from Σ and the vertical line $\{\frac{1}{2} + iy : y \in \mathbb{R}\}$. Apply the holomorphic functional calculus and define

$$\tilde{e} = \frac{1}{2\pi i} \int_{\gamma} z(z - e)^{-1}.$$

Lemma 3.3. *Let p be an idempotent in a Banach algebra with $\|p\| \leq \beta$. Let $e_s = sp + (1 - s)e$ for $s \in [0, 1]$. Suppose e is an element with $\|e\| \leq \beta + 1$ and $\|p - e\| < (\frac{1}{2\beta+2})^{\frac{1}{4}}$. Then the following hold.*

(1) We have

$$\|e_s^2 - e_s\| < \frac{1}{4}$$

for all $s \in [0, 1]$.

(2) If we define

$$\tilde{e}_s = \frac{1}{2\pi i} \int_{\gamma} z(z - e_s)^{-1}$$

as above, then $\{e_s\}_{0 \leq s \leq 1}$ is a continuous path of idempotents connecting \tilde{e}_0 and p .

Proof. Part (1) follows from the following estimate:

$$\begin{aligned} \|e_s^2 - e_s\| &= \|e_s^2 - pe_s + pe_s - p^2 + p - e_s\| \\ &\leq \|(e_s - p)e_s\| + \|p(e_s - p)\| + \|p - e_s\| \\ &\leq (\|e_s\| + \|p\| + 1)\|e_s - p\|. \end{aligned}$$

By the definition of holomorphic functional calculus, part (2) is obvious. ■

Now we are ready to prove the quantitative relative index theorem. Let us first prove a version of the quantitative relative index theorem for the reduced group C^* -algebras. The maximal version can be proved in exactly the same way, after applying the results of Appendix A.

Theorem 3.4. *Let Z_1 and Z_2 be two closed n -dimensional Riemannian manifold and \mathcal{S}_j a Euclidean Cl_n -bundle over Z_j for $j = 1, 2$. Suppose D_j is a first-order symmetric elliptic Cl_n -linear differential operators acting on \mathcal{S}_j over Z_j . Let \tilde{Z}_j be a Galois Γ -covering space of Z_j and \tilde{D}_j the lift of D_j . Let X_j be a subset of Z_j and \tilde{X}_j the preimage of X_j under the covering map $\tilde{Z}_j \rightarrow Z_j$. Denote by $N_r(Z_j \setminus X_j)$ the open r -neighborhood of $Z_j \setminus X_j$. Suppose there is $r > 0$ such that all geometric data on $N_r(Z_1 \setminus X_1)$ and $N_r(Z_2 \setminus X_2)$ coincide, i.e., there is an orientation preserving Riemannian isometry $\Phi: N_r(Z_1 \setminus X_1) \rightarrow N_r(Z_2 \setminus X_2)$ such that Φ lifts to an isometric Cl_n -bundle isomorphism $\Phi: \mathcal{S}_1|_{N_r(Z_1 \setminus X_1)} \rightarrow \mathcal{S}_2|_{N_r(Z_2 \setminus X_2)}$. Assume that*

- (1) the restriction of \tilde{D}_j on \tilde{X}_j is invertible in the following sense: there exists $\sigma > 0$ such that

$$\|\tilde{D}_j f\| \geq \sigma \|f\|$$

for all $f \in C_c^\infty(\tilde{X}_j^\circ, \tilde{\mathcal{S}}_j)$, where \tilde{X}_j° is the interior of \tilde{X}_j in \tilde{Z} ;

- (2) and $D_1 = \Phi^{-1} D_2 \Phi$ on $N_r(Z_1 \setminus X_1)$.

Then there exists a universal constant $C > 0$ such that if $\sigma \cdot r > C$, then we have

$$\text{Ind}_\Gamma(\tilde{D}_1) - \text{Ind}_\Gamma(\tilde{D}_2) = 0$$

in $KO_n(C_r^*(\Gamma; \mathbb{R}))$, where $\text{Ind}_\Gamma(\tilde{D}_j)$ denotes the maximal higher index of \tilde{D}_j and $C_r^*(\Gamma; \mathbb{R})$ is the reduced group C^* -algebra of Γ with real coefficients.

Proof. Let us prove the theorem for the case where $\dim Z_j$ is even and \mathcal{S} is a Hermitian $\mathbb{C}\ell_n$ -bundle, mainly for the reason of notational simplicity. Here $\mathbb{C}\ell_n$ is the complex Clifford algebra of \mathbb{R}^n . The proof for the real Clifford bundle case is the same. Also, the proof for the odd dimensional case is completely similar.⁹ Now, if \mathcal{S} is a Hermitian $\mathbb{C}\ell_n$ -bundle and n is even, it is equivalent to view \mathcal{S} as a Hermitian vector bundle with a $\mathbb{Z}/2$ -grading, with respect to which the operators D_1 and D_2 have odd degree.

We apply the usual higher index construction to \tilde{D}_j (cf. Section 2). Let $\chi: \mathbb{R} \rightarrow \mathbb{R}$ be a normalizing function such that its distributional Fourier transform is supported in $[-1, 1]$. Define

$$F_{1,t} = \chi(t\tilde{D}_1) \quad \text{and} \quad F_{2,t} = \chi(t\tilde{D}_2).$$

Let p_t and q_t be the idempotents constructed out of $F_{1,t}$ and $F_{2,t}$ as in line (2.1). Then for any fixed $t > 0$, the higher index $\text{Ind}_\Gamma(\tilde{D}_1) \in K_0(C_r^*(\Gamma))$ is represented by

$$[p_t] - \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix},$$

and the higher index $\text{Ind}_\Gamma(\tilde{D}_2) \in K_0(C_r^*(\Gamma))$ is represented by

$$[q_t] - \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}.$$

By assumption, there exists $\sigma > 0$ such that

$$\|\tilde{D}_j f\| \geq \sigma \|f\|$$

for all $f \in C_c^\infty(\tilde{X}_j^\circ, \tilde{\mathcal{S}}_j)$. By a standard rescaling argument, that is, by considering λD_1 and λD_2 for some appropriate $\lambda > 0$, we can without loss of generality assume $\sigma = 1$, cf.

Remark 3.6. By part (3) of Lemma 3.2, there exists $\eta > 0$ such that

$$\left\| \left(p_t - \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \right) f \right\| \leq \frac{\eta}{t}$$

⁹Alternatively, the odd dimensional case can be reduced to the even dimensional case by a standard suspension argument.

for all $f \in C_c^\infty(\tilde{Z}_1 \setminus N_{10t}(\tilde{Z}_1 \setminus \tilde{X}_1), \tilde{\mathcal{S}}_1)$. Let us define the operator P_t by setting

$$P_t(f) = \begin{cases} p_t(f) & \text{if } f \in L^2(N_{10t}(\tilde{Z}_1 \setminus \tilde{X}_1), \tilde{\mathcal{S}}_1), \\ \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} f & \text{if } f \in L^2(\tilde{Z}_1 \setminus N_{10t}(\tilde{Z}_1 \setminus \tilde{X}_1), \tilde{\mathcal{S}}_1). \end{cases}$$

In particular, by Lemma 3.3, as long as $t = t_0$ is sufficiently large, then $\|P_t - p_t\|$ is sufficiently small, which implies that

$$\|P_t^2 - P_t\| < \frac{1}{4}.$$

Furthermore, if we define

$$\tilde{P}_t = \frac{1}{2\pi i} \int_{\mathcal{Y}} z(z - P_t)^{-1}$$

then part (2) of Lemma 3.3 implies that \tilde{P}_t and p_t represent the same K -theory class, as long as $t = t_0$ is sufficiently large.

We apply the same argument above to q_t and define

$$Q_t(f) = \begin{cases} q_t(f) & \text{if } f \in L^2(N_{10t}(\tilde{Z}_2 \setminus \tilde{X}_2), \tilde{\mathcal{S}}_2), \\ \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} f & \text{if } f \in L^2(\tilde{Z}_2 \setminus N_{10t}(\tilde{Z}_2 \setminus \tilde{X}_2), \tilde{\mathcal{S}}_2). \end{cases}$$

Similarly, we define

$$\tilde{Q}_t = \frac{1}{2\pi i} \int_{\mathcal{Y}} z(z - Q_t)^{-1}$$

Then Lemma 3.3 implies that \tilde{Q}_t and q_t represent the same K -theory class, as long as $t = t_0$ is sufficiently large.

Now let us set $C = 15t_0$. Recall that we have already applied a rescaling argument to reduce the general case to the case where $\sigma = 1$. Then by assumption, we have

$$r = \sigma \cdot r > C = 15t_0.$$

It follows from the standard finite propagation of wave operators associated to \tilde{D}_1 and \tilde{D}_2 that p_{t_0} and q_{t_0} coincide as operators¹⁰ on

$$L^2(N_r(\tilde{Z}_1 \setminus \tilde{X}_1), \tilde{\mathcal{S}}_1) \cong L^2(N_r(\tilde{Z}_2 \setminus \tilde{X}_2), \tilde{\mathcal{S}}_2).$$

It follows that \tilde{P}_{t_0} and \tilde{Q}_{t_0} coincide. In particular, we have

$$\text{Ind}_\Gamma(\tilde{D}_1) = [\tilde{P}_{t_0}] - \left[\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \right] = [\tilde{Q}_{t_0}] - \left[\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \right] = \text{Ind}_\Gamma(\tilde{D}_2)$$

in $K_0(C_r^*(\Gamma; \mathbb{R}))$. This finishes the proof. ■

¹⁰As far as K -theory classes of p_{t_0} and q_{t_0} are concerned, we can simply ignore the subspaces $L^2(\tilde{Z}_1 \setminus N_r(\tilde{Z}_1 \setminus \tilde{X}_1), \tilde{\mathcal{S}}_1)$ and $L^2(\tilde{Z}_2 \setminus N_r(\tilde{Z}_2 \setminus \tilde{X}_2), \tilde{\mathcal{S}}_2)$, since p_{t_0} and q_{t_0} act on them as the trivial idempotent $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$ respectively.

By applying the results of Appendix A and the estimates in Example 3.7, the same proof for Theorem 3.4 also proves the following maximal version of the quantitative relative index theorem.

Theorem 3.5 (Theorem I). *Let Z_1 and Z_2 be two closed n -dimensional Riemannian manifolds and \mathcal{S}_j a Euclidean Cl_n -bundle over Z_j for $j = 1, 2$. Suppose D_j is a Cl_n -linear Dirac-type operator acting on \mathcal{S}_j over Z_j . Let \tilde{Z}_j be a Galois Γ -covering space of Z_j and \tilde{D}_j the lift of D_j . Let X_j be a subset of Z_j and \tilde{X}_j the preimage of X_j under the covering map $\tilde{Z}_j \rightarrow Z_j$. Denote by $N_r(Z_j \setminus X_j)$ the open r -neighborhood of $Z_j \setminus X_j$. Suppose there is $r > 0$ such that all geometric data on $N_r(Z_1 \setminus X_1)$ and $N_r(Z_2 \setminus X_2)$ coincide, i.e., there is an orientation preserving Riemannian isometry $\Phi: N_r(Z_1 \setminus X_1) \rightarrow N_r(Z_2 \setminus X_2)$ such that Φ lifts to an isometric Cl_n -bundle isomorphism $\Phi: \mathcal{S}_1|_{N_r(Z_1 \setminus X_1)} \rightarrow \mathcal{S}_2|_{N_r(Z_2 \setminus X_2)}$. Assume that*

- (1) *there exists $\sigma > 0$ such that*

$$\mathcal{R}_j(x) \geq \frac{(n - 1)\sigma^2}{n}$$

for all $x \in X_j$, where \mathcal{R}_j is the curvature term appearing in $D_j^2 = \nabla^ \nabla + \mathcal{R}_j$,*

- (2) *and $D_1 = \Phi^{-1} D_2 \Phi$ on $N_r(Z_1 \setminus X_1)$.*

Then there exists a universal constant $C > 0$ such that if $\sigma \cdot r > C$, then we have

$$\text{Ind}_\Gamma(\tilde{D}_1) - \text{Ind}_\Gamma(\tilde{D}_2) = 0$$

in $KO_n(C_{\max}^(\Gamma; \mathbb{R}))$, where $\text{Ind}_\Gamma(\tilde{D}_j)$ denotes the maximal higher index of \tilde{D}_j and $C_{\max}^*(\Gamma; \mathbb{R})$ is the maximal group C^* -algebra of Γ with real coefficients.*

The numerical estimates in Appendix B show that the universal constant C is ≤ 40.65 .

Proof. The proof follows the same strategy as that of Theorem 3.4, so we shall be brief. We claim we can assume without loss of generality that X_1 (resp. X_2) is a codimension zero submanifold with corners in Z_1 (resp. Z_2). Indeed, let $\delta > 0$ be a sufficiently small constant and $\mathcal{U} = \{U_k\}$ be an open cover of \bar{X}_j consisting of geodesically convex balls of radius $\leq \delta$. Note that \bar{X}_j is closed in Z_j , hence compact. It follows that \bar{X}_j admits a finite open cover \mathcal{V} consisting of finitely many members of \mathcal{U} . Without loss of generality, we assume

$$V \cap \bar{X}_j \neq \emptyset$$

for each member V of \mathcal{V} . Denote by W_j the union of all members of \mathcal{V} . Then the closure \bar{W}_j of W_j is contained in $N_{2\delta}(\bar{X}_j)$. By construction, W_j is an n -dimensional compact manifold with corners under the metric inherited from Z_j . If we replace X_j by W_j and replace r by $r - 2\delta$, then all assumptions of the theorem are still satisfied, as long as δ is sufficiently small. Therefore, without loss of generality, we assume that X_1 (resp. X_2) is a codimension zero submanifold with corners in Z_1 (resp. Z_2).

Let $L^2_{C^*_\max}(\tilde{Z}_j, \tilde{\mathcal{S}}_j)^\Gamma$ be the associated $C^*_\max(\Gamma; \mathbb{R})$ -Hilbert module of sections of $\tilde{\mathcal{S}}_j$ over \tilde{Z}_j (cf. line (A.5)). By assumption there exists $\sigma > 0$ such that

$$\mathcal{R}_j(x) \geq \frac{(n-1)\sigma^2}{n}$$

for all $x \in X_j$. It follows from Proposition A.5 and the proof of Roe’s lemma ([22, Lemma 2.5]) that if the Fourier transform $\hat{\varphi}$ of a function φ is supported in $(-a, a)$ and $\psi \in C_0(\tilde{Z}_j)$ has support disjoint from the $2a$ -neighborhood of \tilde{X}_j , then

$$\|\varphi(\tilde{D}_j)\rho(\psi)\| \leq \|\psi\| \sup\{|\varphi(y)| : |y| \geq \sigma\}.$$

Here $\varphi(\tilde{D}_j)$ is the bounded operator on $L^2_{C^*_\max}(\tilde{Z}_j, \tilde{\mathcal{S}}_j)^\Gamma$ obtained by applying functional calculus, and $\rho(\psi)$ is the bounded operator on $L^2_{C^*_\max}(\tilde{Z}_j, \tilde{\mathcal{S}}_j)^\Gamma$ given by multiplication of ψ and $\|\psi\|$ is supremum norm of ψ . Consequently, we also have the analogues of Lemma 3.2 and Lemma 3.3 in the current maximal setting. Now the theorem follows by the exact same proof as the one of Theorem 3.4. ■

Remark 3.6. In the proof of Theorem 3.4, we have implicitly used the fact that a Dirac-type operator has propagation speed equal to 1. Recall that the propagation speed of a first order differential operator D on a Riemannian manifold Z is defined to be

$$c_D := \sup_{x \in X} c_D(x),$$

where σ_D the principal symbol of D and

$$c_D(x) := \sup\{\|\sigma_D(x, \xi)\| : \xi \in T_x^*X, \|\xi\| = 1\}.$$

If we consider more general elliptic operators D_1 and D_2 such that both c_{D_1} and c_{D_2} are bounded by λ , then the corresponding condition $\sigma \cdot r > C$ in Theorem 3.4 should be replaced by

$$\sigma \cdot \frac{r}{\lambda} > C.$$

The following is a typical geometric setup to which Theorem 3.5 applies.

Example 3.7. Let Z be a closed n -dimensional Riemannian manifold and \mathcal{S} a Euclidean Cl_n -bundle over Z . Suppose D is a Cl_n -linear Dirac-type operator acting on \mathcal{S} over Z . Let \tilde{Z} be a Galois Γ -covering space of Z and \tilde{D} the lift of D . Let X be a subset of Z and \tilde{X} the preimage of X under the covering map $\tilde{Z} \rightarrow Z$.

By the Lichnerowicz formula, we have

$$\tilde{D}^2 = \nabla^* \nabla + \mathcal{R},$$

where \mathcal{R} is a symmetric bundle endomorphism of $\tilde{\mathcal{S}}$. If D is an actual Dirac operator, then $\mathcal{R} = \frac{\kappa}{4}$ where κ is the scalar curvature of the metric on \tilde{Z} . By the Cauchy–Schwarz inequality, we have

$$\langle \tilde{D}f, \tilde{D}f \rangle \leq n \langle \nabla f, \nabla f \rangle$$

for all $f \in C_c^\infty(\tilde{Z}, \tilde{\mathcal{S}})$ and $n = \dim Z$. Combining the two formulas above, we see that

$$\frac{n-1}{n} \langle \tilde{D}f, \tilde{D}f \rangle \geq \langle \mathcal{R}f, f \rangle$$

for all $f \in C_c^\infty(\tilde{Z}, \tilde{\mathcal{S}})$.

Let $\tilde{X}^0 = \tilde{X} - \partial\tilde{X}$ be the interior of \tilde{X} . If we assume there exists $\sigma > 0$ such that

$$\mathcal{R}(x) \geq \frac{(n-1)\sigma^2}{n}$$

for all $x \in \tilde{X}^0$, then we have

$$\frac{n-1}{n} \langle \tilde{D}f, \tilde{D}f \rangle \geq \langle \mathcal{R}f, f \rangle \geq \sigma^2 \langle f, f \rangle \tag{3.2}$$

for all $f \in C_c^\infty(\tilde{X}^0, \tilde{\mathcal{S}})$. In other words, we have

$$\|\tilde{D}f\| \geq \sigma \|f\| \tag{3.3}$$

for all $f \in C_c^\infty(\tilde{X}^0, \tilde{\mathcal{S}})$ in this case.

4. Proof of Theorem II

In this section, we apply the quantitative relative index theorem (Theorem 3.5) to prove Theorem II. In order to make our exposition more transparent, let us first prove the following special case.

Theorem 4.1 (A special case of Theorem II). *Suppose M is a closed spin manifold of dimension $n - 1$ such that the higher index of its Dirac operator does not vanish in $KO_{n-1}(C_{\max}^*(\pi_1 M; \mathbb{R}))$. If the manifold $M \times [0, 1]$ is endowed with a Riemannian metric whose scalar curvature is $\geq n(n - 1)$, then*

$$\text{width}(M \times [0, 1]) \leq \frac{\frac{8}{\sqrt{3}}C + 4\pi}{n},$$

where C is the universal constant from Theorem 3.5.

Proof. For simplicity, we shall prove the theorem for the reduced case. In fact, let us assume that the higher index of the (complexified) Dirac operator on M does not vanish in $K_{n-1}(C_r^*(\Gamma))$. The proof for the maximal case is essentially the same. For the real case, see Remark 4.2.

Let $\tilde{X} = \tilde{M} \times [0, 1]$ be the universal cover of $X = M \times [0, 1]$ and \tilde{D} the associated $\mathbb{C}\ell_n$ -linear Dirac operator on \tilde{X} . By the discussion in Example 3.7, since the scalar curvature $\text{Sc}(g) \geq n(n - 1)$, we have

$$\|\tilde{D}f\| \geq \frac{n}{2} \|f\|$$

for all $f \in C_c^\infty(\tilde{X}^0, \tilde{\mathcal{S}})$, where $\tilde{\mathcal{S}}$ is the associated spinor bundle over \tilde{X} .

We prove the theorem by contradiction. Assume to the contrary that

$$\ell := \text{width}(X) > \frac{\frac{8}{\sqrt{3}}C + 4\pi}{n}.$$

Denote by $\partial_+X = M \times \{1\}$ and $\partial_-X = M \times \{0\}$. Then for any sufficiently small $\varepsilon > 0$, there exists a hypersurface Y in X such that

$$\text{dist}(\partial_+X, Y) \geq \frac{\ell}{2} - \varepsilon \quad \text{and} \quad \text{dist}(\partial_-X, Y) \geq \frac{\ell}{2} - \varepsilon.$$

Let $\varphi: X \rightarrow \mathbb{R}$ be a real-valued smooth function such that (cf. [8, Proposition 2.1])

- (1) $\|d\varphi\| \leq 1$,
- (2) $\varphi(x) \equiv 0$ for all x between Y and ∂_-X with $\text{dist}(x, Y) \geq \frac{2\pi}{n} + \varepsilon$,
- (3) and $\varphi(x) \equiv \frac{4\pi}{n}$ for all x between Y and ∂_+X with $\text{dist}(x, Y) \geq \frac{2\pi}{n} + \varepsilon$.

From now on, let us fix a sufficiently small $\varepsilon > 0$ and denote the lift of φ from X to \tilde{X} still by φ . Define the function

$$u(x) = e^{\frac{n}{2}i\varphi(x)}$$

on \tilde{X} . We have

$$\|[\tilde{D}, u]\| = \|du\| = \frac{n}{2}\|u \cdot d\varphi\| \leq \frac{n}{2}.$$

Similarly, we also have

$$\|[\tilde{D}, u^{-1}]\| \leq \frac{n}{2}.$$

Consider the following Dirac operator on $\mathbb{S}^1 \times \tilde{X}^0$:

$$\not{D} = c \cdot \frac{d}{dt} + \tilde{D}_t \tag{4.1}$$

where c is the Clifford multiplication of the unit vector d/dt and

$$\tilde{D}_t := t\tilde{D} + (1-t)u\tilde{D}u^{-1}$$

for each $t \in [0, 1]$. Here we have chosen the parametrization $\mathbb{S}^1 = [0, 1]/\{0, 1\}$. Let $\tilde{\mathcal{S}}_{[0,1]}$ be the associated spinor bundle on $[0, 1] \times \tilde{X}^0$ and $\tilde{\mathcal{S}}_t$ its restriction on $\{t\} \times \tilde{X}^0$. Each smooth section $f \in C_c^\infty([0, 1] \times \tilde{X}^0, \tilde{\mathcal{S}}_{[0,1]})$ can be viewed as a smooth family $f(t) \in C_c^\infty(\{t\} \times \tilde{X}^0, \tilde{\mathcal{S}}_t)$. The operator \not{D} acts on the following subspace of $C_c^\infty([0, 1] \times \tilde{X}^0, \tilde{\mathcal{S}}_{[0,1]})$:

$$\{f \in C_c^\infty([0, 1] \times \tilde{X}^0, \tilde{\mathcal{S}}_{[0,1]}) : f(1) = uf(0)\}.$$

From now on, we shall simply write $C_c^\infty(\mathbb{S}^1 \times \tilde{X}^0, \tilde{\mathcal{S}})$ for the above subspace of sections.

Clearly, we have

$$\not{D}^2 = -\frac{d^2}{dt^2} + \tilde{D}_t^2 + c[\tilde{D}, u]u^{-1}.$$

By using the identity

$$\tilde{D}u\tilde{D}u^{-1} + u\tilde{D}u^{-1}\tilde{D} = [\tilde{D}, u][\tilde{D}, u^{-1}] + u\tilde{D}^2u^{-1} + \tilde{D}^2,$$

we have

$$\tilde{D}_t^2 = t\tilde{D}^2 + (1-t)u\tilde{D}^2u^{-1} + t(1-t)[\tilde{D}, u][\tilde{D}, u^{-1}]. \tag{4.2}$$

It follows from the assumption $\text{Sc}(g) \geq n(n-1)$ and the estimates in Example 3.7 that $\tilde{D}^2 \geq \frac{n^2}{4}$ on $C_c^\infty(\mathbb{S}^1 \times \tilde{X}^0, \tilde{\mathcal{S}})$, which implies also $u\tilde{D}^2u^{-1} \geq \frac{n^2}{4}$ on $C_c^\infty(\mathbb{S}^1 \times \tilde{X}^0, \tilde{\mathcal{S}})$, since u is a unitary. Therefore, we have

$$\begin{aligned} \tilde{D}_t^2 &\geq \frac{n^2}{4} - t(1-t)\|[\tilde{D}, u^{-1}][\tilde{D}, u]\| \\ &\geq \frac{n^2}{4} - \frac{n^2}{16} = \frac{3n^2}{16} \end{aligned}$$

where the second inequality uses the fact $t(1-t) \leq 1/4$ for all $t \in [0, 1]$.

Now, for each $\lambda > 0$, we define the rescaled version of \tilde{D} to be

$$\not{D}_\lambda = c \cdot \frac{d}{dt} + \lambda \tilde{D}_t \tag{4.3}$$

with $\lambda \tilde{D}_t$ in place of \tilde{D}_t . The same calculation from above shows that

$$\not{D}_\lambda^2 = -\frac{d^2}{dt^2} + \lambda^2 \tilde{D}_t^2 + \lambda c[\tilde{D}, u]u^{-1}.$$

Since $\tilde{D}_t^2 \geq \frac{3n^2}{16}$, it follows that

$$\not{D}_\lambda^2 \geq \lambda^2 \frac{3n^2}{16} - \lambda \frac{n}{2} = \frac{3n^2\lambda^2}{16} \left(1 - \frac{8}{3n\lambda}\right).$$

If we want to be explicit about the dependence of \not{D}_λ on the unitary u , we shall write $\not{D}_{\lambda,u}$ instead of \not{D}_λ .

Let $v \equiv 1$ be the trivial unitary on \tilde{X} . Define the operator

$$\not{D}_{\lambda,v} = c \frac{d}{dt} + \lambda \tilde{D}.$$

A similar (in fact simpler) calculation shows that

$$\not{D}_{\lambda,v}^2 \geq \lambda^2 \frac{n^2}{4}$$

on $C_c^\infty(\mathbb{S}^1 \times \tilde{X}^0, \tilde{\mathcal{S}})$.

Consider the doubling $\mathfrak{X} = M \times \mathbb{S}^1 = X \cup_{\partial X} (-X)$ of X , where $-X$ is a copy of X but with the opposite orientation. Extend¹¹ the Riemannian metric on X to a Riemannian

¹¹To be precise, we fix a copy of X inside of \mathfrak{X} and equip it with the Riemannian metric given by the assumption. Then we choose any Riemannian metric on \mathfrak{X} that coincides with the Riemannian metric on this chosen copy of X .

metric on \mathfrak{X} . The reader should not confuse the copy of \mathbb{S}^1 appearing in $\mathfrak{X} = M \times \mathbb{S}^1$ with the copy of \mathbb{S}^1 appearing in $\mathbb{S}^1 \times X^o = \mathbb{S}^1 \times M \times (0, 1)$. Note that the Riemannian metric on $\mathfrak{X} = M \times \mathbb{S}^1$ does *not* have positive scalar curvature everywhere in general. But \mathfrak{X} is a closed manifold, so the usual higher index theory applies. More precisely, by the construction of $u = e^{\frac{\pi}{2}i\varphi}$, it extends trivially to a unitary u on $\tilde{\mathfrak{X}} := \tilde{M} \times \mathbb{S}^1$ by setting it to be 1 in $\tilde{\mathfrak{X}} \setminus \tilde{X}$. Let $\tilde{D}^{\mathfrak{X}}$ be the Dirac operator on $\tilde{\mathfrak{X}}$. We define

$$\mathcal{D}_u^{\mathfrak{X}} = c \cdot \frac{d}{dt} + \tilde{D}_t^{\mathfrak{X}} \quad \text{where} \quad \tilde{D}_t^{\mathfrak{X}} := t\tilde{D}^{\mathfrak{X}} + (1-t)u\tilde{D}^{\mathfrak{X}}u^{-1}.$$

Similarly, let $v \equiv 1$ be the trivial unitary on $\tilde{\mathfrak{X}}$ and define

$$\mathcal{D}_v^{\mathfrak{X}} = c \cdot \frac{d}{dt} + \tilde{D}^{\mathfrak{X}}.$$

Claim. $\text{Ind}_{\Gamma}(\mathcal{D}_u^{\mathfrak{X}}) = \text{Ind}_{\Gamma}(\tilde{D}^M)$ in $K_{n-1}(C_r^*(\Gamma))$, where $\Gamma = \pi_1 M$ and \tilde{D}^M is the Dirac operator on \tilde{M} .

This can be seen as follows. The higher index $\text{Ind}_{\Gamma}(\mathcal{D}_u^{\mathfrak{X}})$ is independent of the choice of the Riemannian metric on \mathfrak{X} , since $\mathfrak{X} = M \times \mathbb{S}^1$ is a closed manifold. Furthermore, if $\{u_s\}_{0 \leq s \leq 1}$ is a continuous family of unitaries on \mathfrak{X} , then $\text{Ind}_{\Gamma}(\mathcal{D}_{u_0}^{\mathfrak{X}}) = \text{Ind}_{\Gamma}(\mathcal{D}_{u_1}^{\mathfrak{X}}) \in K_{n-1}(C_r^*(\Gamma))$. Therefore, without loss of generality, we assume the Riemannian metric on $\mathfrak{X} = M \times \mathbb{S}^1$ is given by a product metric $g_M + dx^2$ and assume¹² the unitary u on \mathfrak{X} is given by the projection map $\mathfrak{X} = M \times \mathbb{S}^1 \rightarrow \mathbb{S}^1 \subset \mathbb{C}$. In this case, the operator $\mathcal{D}_u^{\mathfrak{X}}$ becomes

$$\left(c \frac{d}{dt} + D_t^{\mathbb{S}^1}\right) \hat{\otimes} 1 + 1 \hat{\otimes} \tilde{D}_M$$

where $D_t^{\mathbb{S}^1} = tD^{\mathbb{S}^1} + (1-t)e^{2\pi i\theta}D^{\mathbb{S}^1}e^{-2\pi i\theta}$ and θ is the coordinate for the copy of \mathbb{S}^1 appearing in $\mathfrak{X} = M \times \mathbb{S}^1$. Recall that the index of the operator $c \frac{d}{dt} + D_t^{\mathbb{S}^1}$ is equal to the spectral flow of the family $\{D_t^{\mathbb{S}^1}\}_{0 \leq t \leq 1}$, which has index 1 (cf. [1, Section 7]). Therefore, it follows that

$$\text{Ind}_{\Gamma}(\mathcal{D}_u^{\mathfrak{X}}) = \text{Ind}_{\Gamma}(\tilde{D}^M) \quad \text{in } K_{n-1}(C_r^*(\Gamma)).$$

The same argument also shows that

$$\text{Ind}_{\Gamma}(\mathcal{D}_v^{\mathfrak{X}}) = 0 \quad \text{in } K_{n-1}(C_r^*(\Gamma)).$$

We conclude that

$$\text{Ind}_{\Gamma}(\mathcal{D}_u^{\mathfrak{X}}) - \text{Ind}_{\Gamma}(\mathcal{D}_v^{\mathfrak{X}}) = \text{Ind}_{\Gamma}(\tilde{D}^M) \quad \text{in } K_{n-1}(C_r^*(\Gamma)).$$

On the other hand, since we have assumed that

$$\text{width}(X) > \frac{\frac{8}{\sqrt{3}}C + 4\pi}{n},$$

¹²This can be achieved by a homotopy of unitaries on \mathfrak{X} .

the operators $\mathcal{D}_u^{\mathfrak{X}}$ and $\mathcal{D}_v^{\mathfrak{X}}$ coincide on the r -neighborhood $N_r(\mathfrak{X} \setminus X)$ of $\mathfrak{X} \setminus X$, where we have

$$r > \frac{4C}{\sqrt{3}n}$$

as long as ε chosen at the beginning of the proof is sufficiently small. In particular, we see that

$$\frac{r}{\lambda} \sqrt{\frac{3n^2\lambda^2}{16} \left(1 - \frac{8}{3n\lambda}\right)} > C,$$

as long as λ is sufficiently large. Now it follows from Theorem 3.5 and Remark 3.6 that

$$\text{Ind}_\Gamma(\mathcal{D}_u^{\mathfrak{X}}) - \text{Ind}_\Gamma(\mathcal{D}_v^{\mathfrak{X}}) = 0$$

in $K_{n-1}(C_r^*(\Gamma))$. We arrive at a contradiction, since $\text{Ind}_\Gamma(\tilde{D}^M) \neq 0$ by assumption. This finishes the proof. ■

Remark 4.2. Let us discuss how to adjust the proof of Theorem 4.1 for the real case. Roughly speaking, we replace the imaginary number $i = \sqrt{-1}$ by the matrix $\mathbf{I} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$, while viewing \mathbf{I} as a matrix acting on a 2-dimensional $\mathbb{Z}/2$ -graded real vector space. For example, multiplication by the complex number $e^{2\pi i t}$ on a 1-dimensional complex vector space is replaced by the operator $e^{2\pi t \cdot \mathbf{I}}$ acting on a 2-dimensional $\mathbb{Z}/2$ -graded real vector space. More precisely, let us describe such a modification in terms of Clifford algebras. Let $\text{Cl}_{r,s}$ be the real Clifford algebra generated by $\{e_1, e_2, \dots, e_{r+s}\}$ subject to the following relations:

$$e_j e_k + e_k e_j = \begin{cases} -2\delta_{jk} & \text{if } j \leq r, \\ +2\delta_{jk} & \text{if } j > r. \end{cases}$$

To be clear, our convention for the notation of Clifford algebras is consistent with that of [17]. In particular, $\text{Cl}_n := \text{Cl}_{n,0}$ stands for the Clifford algebra generated by $\{e_1, e_2, \dots, e_n\}$ subject to the following relations:

$$e_j^2 = -1 \quad \text{and} \quad e_j e_k + e_j e_k = 0 \quad \text{for all } 1 \leq j, k \leq n.$$

In terms of Clifford algebras, we define $\mathbf{I} = e_1 e_2$ in $\text{Cl}_{0,2}$. The operator \mathcal{D} in line (4.1) now becomes

$$\mathcal{D} = c \cdot \frac{d}{dt} + \tilde{D}_t,$$

where $c \in \text{Cl}_{1,0}$ is the Clifford multiplication of the unit vector d/dt and

$$\tilde{D}_t := t\tilde{D} + (1-t)U\tilde{D}U^{-1}$$

with $U = e^{2\pi t \mathbf{I} \varphi(x)/\ell}$. In particular, the operator \mathcal{D} is a $\text{Cl}_{n+1,2}$ -linear Dirac-type operator and its higher index lies in $KO_{n-1}(C_{\max}^*(\Gamma; \mathbb{R}))$. The same remark applies to other similar operators that appeared in the proof of Theorem 4.1. With these modifications, the proof for the real case now proceeds in the same way as the complex case.

Now we are ready to prove Theorem II.

Theorem 4.3 (Theorem II). *Let X be an n -dimensional compact connected spin manifold with boundary. Suppose $f: X \rightarrow [-1, 1]^m$ is a smooth map that sends the boundary of X to the boundary of $[-1, 1]^m$. Let $\partial_{j\pm}$, $j = 1, \dots, m$, be the pullbacks of the pairs of the opposite faces of the cube $[-1, 1]^m$. Suppose Y_{\hbar} is an $(n - m)$ -dimensional closed submanifold (without boundary) in X that satisfies the following conditions:*

- (1) $\iota: \pi_1(Y_{\hbar}) \rightarrow \pi_1(X)$ is injective, where ι is the canonical morphism on π_1 induced by the inclusion $Y_{\hbar} \hookrightarrow \pi_1(X)$;
- (2) Y_{\hbar} is the transversal intersection of m orientable hypersurfaces $Y_j \subset X$, $1 \leq j \leq m$, such that each Y_j separates ∂_{j-} from ∂_{j+} ;
- (3) the higher index $\text{Ind}_{\Gamma}(D_{Y_{\hbar}})$ does not vanish in $KO_{n-m}(C_{\max}^*(\Gamma; \mathbb{R}))$, where $\Gamma = \pi_1(Y_{\hbar})$.

If $\text{Sc}(X) \geq n(n - 1)$, then the distances $\ell_j = \text{dist}(\partial_{j-}, \partial_{j+})$ satisfy the following inequality:

$$\sum_{j=1}^m \frac{1}{\ell_j^2} \geq \frac{n^2}{\left(\frac{8}{\sqrt{3}}C + 4\pi\right)^2}.$$

Consequently, we have

$$\min_{1 \leq j \leq m} \text{dist}(\partial_{j-}, \partial_{j+}) \leq \sqrt{m} \frac{\frac{8}{\sqrt{3}}C + 4\pi}{n}.$$

Proof. For simplicity, we shall prove the theorem for the complex case, that is, complexified Dirac operators instead of $C\ell_n$ -linear Dirac operators. For the real case, see Remark 4.2.

We first show that the general case where $\iota: \pi_1(Y_{\hbar}) \rightarrow \pi_1(X)$ is injective can be reduced to the case where $\iota: \pi_1(Y_{\hbar}) \rightarrow \pi_1(X)$ is split injective.¹³ Let X_u be the universal cover of X . Since by assumption $\iota: \pi_1(Y_{\hbar}) \rightarrow \pi_1(X)$ is injective, we can view $\Gamma = \pi_1(Y_{\hbar})$ as a subgroup of $\pi_1(X)$. Let $X_{\Gamma} = X_u/\Gamma$ be the covering space of X corresponding to the subgroup $\Gamma \subset \pi_1(X)$. Then the inverse image of Y_{\hbar} under the projection $p: X_{\Gamma} \rightarrow X$ is a disjoint union of covering spaces of Y_{\hbar} , at least one of which is a diffeomorphic copy of Y_{\hbar} . Fix such a copy of Y_{\hbar} in X_{Γ} and denote it by \widehat{Y}_{\hbar} . Roughly speaking, the space X_{Γ} equipped with the lifted Riemannian metric from X could serve as a replacement of the original space X , except that X_{Γ} is not compact in general. To remedy this, we shall choose a “fundamental domain” around \widehat{Y}_{\hbar} in X_{Γ} as follows.

By assumption, $Y_{\hbar} \subset X$ is the transversal intersection of m orientable hypersurfaces $Y_j \subset X$. Let r_j be the distance function¹⁴ from ∂_{j-} , that is $r_j(x) = \text{dist}(x, \partial_{j-})$. Without

¹³We say $\iota: \pi_1(Y_{\hbar}) \rightarrow \pi_1(X)$ is split injective if there exists a group homomorphism $\varpi: \pi_1(X) \rightarrow \pi_1(Y_{\hbar})$ such that $\varpi \circ \iota = \mathbb{1}$, where $\mathbb{1}$ is the identity morphism of $\pi_1(Y_{\hbar})$.

¹⁴To be precise, let r_j be a smooth approximation of the distance function from ∂_{j-} .

loss of generality, we can assume $Y_j = r_j^{-1}(a_j)$ for some regular value $a_j \in [0, \ell_j]$. Let $Y_j^\Gamma = p^{-1}(Y_j)$ be the inverse image of Y_j in X_Γ . Denote by \bar{r}_j the lift of r_j from X to X_Γ . Let $\nabla\bar{r}_j$ be the gradient vector field associated to \bar{r}_j . A point $x \in X_\Gamma$ is said to be *permissible* if there exist a number $s \geq 0$ and a piecewise smooth curve $c: [0, s] \rightarrow X_\Gamma$ satisfying the following conditions:

- (i) $c(0) \in \widehat{Y}_\hbar$ and $c(s) = x$;
- (ii) there is a subdivision of $[0, s]$ into finitely many subintervals $\{[t_k, t_{k+1}]\}$ such that, on each subinterval $[t_k, t_{k+1}]$, the curve c is either an integral curve or a reversed integral curve¹⁵ of the gradient vector field $\nabla\bar{r}_{i_k}$ for some $1 \leq i_k \leq m$, where we require i_k ’s to be all distinct from each other;
- (iii) furthermore, when c is an integral curve of the gradient vector field $\nabla\bar{r}_{i_k}$ on the subinterval $[t_k, t_{k+1}]$, we require the length of $c|_{[t_k, t_{k+1}]}$ to be less than or equal to $(\ell_{i_k} - a_{i_k} - \frac{\varepsilon}{4})$; and when c is a reversed integral curve of the gradient vector field $\nabla\bar{r}_{i_k}$ on the subinterval $[t_k, t_{k+1}]$, we require the length of $c|_{[t_k, t_{k+1}]}$ to be less than or equal to $(a_{i_k} - \frac{\varepsilon}{4})$.

Let T be the set of all permissible points. Now T may not be a manifold with corners. To fix this, we choose an open cover $\mathcal{U} = \{U_\alpha\}_{\alpha \in \Lambda}$ of T by geodesically convex metric balls of sufficiently small radius $\delta > 0$. Now take the union of members of $\mathcal{U} = \{U_\alpha\}_{\alpha \in \Lambda}$ that do not intersect the boundary ∂T of T , and denote by Z the closure of the resulting subset. Then Z is a manifold with corners which, together with the subspace $\widehat{Y}_\hbar \subset Z$, satisfies all the conditions of the theorem, provided that ε and δ are chosen to be sufficiently small. In particular, the intersection $Y_j^\Gamma \cap Z$ of each hypersurface Y_j^Γ with Z gives a hypersurface of Z . The transversal intersection of the resulting hypersurfaces is precisely $\widehat{Y}_\hbar \subset Z$. Furthermore, note that the isomorphism $\Gamma = \pi_1(Y_\hbar^\Gamma) \rightarrow \pi_1(X^\Gamma) = \Gamma$ factors as the composition

$$\pi_1(Y_\hbar^\Gamma) \rightarrow \pi_1(Z) \rightarrow \pi_1(X^\Gamma),$$

where the morphisms $\pi_1(Y_\hbar^\Gamma) \rightarrow \pi_1(Z)$ and $\pi_1(Z) \rightarrow \pi_1(X^\Gamma)$ are induced by the obvious inclusions of spaces. It follows that $\pi_1(Y_\hbar^\Gamma) \rightarrow \pi_1(Z)$ is a split injection. Therefore, without loss of generality, it suffices to prove the theorem under the additional assumption that $\iota: \pi_1(Y_\hbar) \rightarrow \pi_1(X)$ is a split injection.

From now on, let us assume $\iota: \Gamma = \pi_1(Y_\hbar) \rightarrow \pi_1(X)$ is a split injection with a splitting morphism $\varpi: \pi_1(X) \rightarrow \pi_1(Y_\hbar) = \Gamma$. Let \tilde{X} be the Galois Γ -covering space determined by $\varpi: \pi_1(X) \rightarrow \Gamma$. In particular, the restriction of the covering map $\tilde{X} \rightarrow X$ on Y_\hbar gives the universal covering space of Y_\hbar .

¹⁵By definition, an integral curve of a vector field is a curve whose tangent vector coincides with the given vector field at every point of the curve. A reversed integral curve is an integral curve with the reversed parametrization, that is, the tangent vector field of a reserved integral curve coincides with the negative of the given vector field at every point of the curve.

Without loss of generality, assume Y_{th} is the transversal intersection of m orientable hypersurfaces $Y_j \subset X$, $1 \leq j \leq m$ such that each Y_j separates ∂_{j-} from ∂_{j+} and

$$\text{dist}(\partial_{j-}, Y_j) \geq \frac{\ell_j}{2} - \varepsilon \quad \text{and} \quad \text{dist}(\partial_{j+}, Y_j) \geq \frac{\ell_j}{2} - \varepsilon$$

for some sufficiently small $\varepsilon > 0$. Furthermore, without loss of generality, we assume

$$\ell_1 = \min_{1 \leq j \leq m} \ell_j.$$

Let us set

$$L = \left(\sum_{j=1}^m \frac{\ell_1^2}{\ell_j^2} \right)^{1/2}.$$

Assume to the contrary that

$$\sum_{j=1}^m \frac{1}{\ell_j^2} < \frac{n^2}{\left(\frac{8}{\sqrt{3}}C + 4\pi\right)^2}.$$

that is,

$$\frac{1}{\ell_1^2} \cdot L^2 < \frac{n^2}{\left(\frac{8}{\sqrt{3}}C + 4\pi\right)^2}.$$

Therefore, we have

$$\min_{1 \leq j \leq m} \ell_j = \ell_1 > \frac{L\left(\frac{8}{\sqrt{3}}C + 4\pi\right)}{n}. \tag{4.4}$$

For each $1 \leq j \leq m$, let $\varphi_j: X \rightarrow \mathbb{R}$ be a real-valued smooth function such that (cf. [8, Proposition 2.1])

- (1) $\|d\varphi_j\| \leq 1$,
- (2) $\varphi_j(x) \equiv 0$ for all x between Y_j and ∂_{j-} with $\text{dist}(x, Y) \geq \frac{2\pi L}{n} + \varepsilon$,
- (3) and $\varphi_j(x) \equiv \frac{4\pi L}{n}$ for all x between Y_j and ∂_{j+} with $\text{dist}(x, Y) \geq \frac{2\pi L}{n} + \varepsilon$.

Let us fix a sufficiently small $\varepsilon > 0$ and denote the lift of φ_j from X to \tilde{X} still by φ_j . Define the function

$$u_j(x) = \exp\left(\frac{n\ell_1}{2L\ell_j}i\varphi_j(x)\right)$$

on \tilde{X} . We have

$$\|[\tilde{D}, u_j]\| = \|du_j\| = \frac{n\ell_1}{2L\ell_j}\|u_j \cdot d\varphi_j\| \leq \frac{n\ell_1}{2L\ell_j}$$

and

$$\|[\tilde{D}, u_j^{-1}]\| \leq \frac{n\ell_1}{2L\ell_j}.$$

Let $\mathbb{T}^m = \mathbb{S}^1 \times \cdots \times \mathbb{S}^1$ be the m -dimensional torus. Consider the following differential operator on $\mathbb{T}^m \times \tilde{X}^0$:

$$\mathcal{D} = \sum_{j=1}^m c_j \frac{\partial}{\partial t_j} + \tilde{D}_{t_1, t_2, \dots, t_m}$$

where c_j is the Clifford multiplication of the unit vector $\frac{\partial}{\partial t_j}$ and $\tilde{D}_{t_1, t_2, \dots, t_m}$ is inductively defined as follows. We define

$$\tilde{D}_{t_1} = t_1 \tilde{D} + (1 - t_1)u_1 \tilde{D}u_1^{-1}$$

and

$$\tilde{D}_{t_1, t_2, \dots, t_k} := t_k (\tilde{D}_{t_1, \dots, t_{k-1}}) + (1 - t_k)u_k (\tilde{D}_{t_1, \dots, t_{k-1}})u_k^{-1}$$

for $(t_1, \dots, t_m) \in [0, 1]^m$. Here we have chosen the parametrization $\mathbb{S}^1 = [0, 1]/\{0, 1\}$.

By the assumption $\text{Sc}(X) \geq n(n - 1)$ and the estimates in Example 3.7, we have

$$\tilde{D}^2 \geq \frac{n \cdot \min_{X \in X} \text{Sc}(\tilde{X})}{4(n - 1)} \geq \frac{n^2}{4}.$$

By the calculation in the proof of Theorem 4.1, we have

$$\tilde{D}_{t_1}^2 = t_1 \tilde{D}^2 + (1 - t_1)u_1 \tilde{D}^2 u_1^{-1} + t_1(1 - t_1)[\tilde{D}, u_1^{-1}][\tilde{D}, u_1].$$

It follows that

$$\tilde{D}_{t_1}^2 \geq \frac{n^2}{4} - \frac{n^2 \ell_1^2}{16L^2 \ell_2^2}$$

Note that

$$[\tilde{D}_{t_1}, u_2] = t_1 [\tilde{D}, u_2] + (1 - t_1)u_1 [\tilde{D}, u_2]u_1^{-1},$$

which implies that

$$\|[\tilde{D}_{t_1}, u_2]\| \leq \|[\tilde{D}, u_2]\| \leq \frac{n \ell_1}{2L \ell_2}.$$

By induction, we conclude that

$$\tilde{D}_{t_1, \dots, t_k}^2 \geq \frac{n^2}{4} - \left(\sum_{j=1}^k \frac{n^2 \ell_1^2}{16L^2 \ell_j^2} \right)$$

for each $1 \leq k \leq m$. In particular,

$$\tilde{D}_{t_1, \dots, t_m}^2 \geq \frac{n^2}{4} - \left(\sum_{j=1}^m \frac{n^2 \ell_1^2}{16L^2 \ell_j^2} \right) = \frac{3n^2}{16}.$$

By applying the same rescaling argument as in line (4.3), we conclude that there exists $K > 0$ such that

$$\mathcal{D}_\lambda^2 = - \sum_{j=1}^m \frac{\partial^2}{\partial t_j^2} + \lambda^2 \tilde{D}_{t_1, \dots, t_m}^2 + \lambda \sum_{j=1}^m c_j \frac{\partial \tilde{D}_{t_1, \dots, t_m}}{\partial t_j} \geq \frac{3n^2 \lambda^2}{16} - \lambda K n$$

on $C_c^\infty(\tilde{X}^0, \tilde{\mathcal{S}})$.

Similarly, for each $1 \leq j \leq m$, we define the operator

$$\mathcal{D}_j = \sum_{i=1}^m c_i \frac{\partial}{\partial t_i} + \tilde{D}_{t_1, \dots, \hat{t}_j, \dots, t_m}$$

where $\tilde{D}_{t_1, \dots, \hat{t}_j, \dots, t_m}$ is defined the same way as $\tilde{D}_{t_1, \dots, t_j, \dots, t_m}$ except that u_j is replaced by the trivial unitary $v \equiv 1$. More generally, for each subset $\Lambda \subseteq \{1, 2, \dots, m\}$, we define the operator

$$\mathcal{D}_\Lambda = \sum_{i=1}^m c_i \frac{\partial}{\partial t_i} + \tilde{D}_\Lambda$$

where \tilde{D}_Λ is defined the same way as $\tilde{D}_{t_1, \dots, t_j, \dots, t_m}$ except that u_k is replaced by the trivial unitary $v \equiv 1$ for every $k \in \Lambda$. The same argument as above shows that

$$\mathcal{D}_{\Lambda, \lambda}^2 \geq \frac{3n^2 \lambda^2}{16} - \lambda Kn$$

for all Λ and λ .

Now we consider the doubling $\mathfrak{X} = X \cup (-X)$ of X and fix a Riemannian metric on \mathfrak{X} that extends the metric of X . Of course, this metric on \mathfrak{X} generally does *not* satisfy $\text{Sc}(\mathfrak{X}) \geq n(n-1)$. Let $\tilde{\mathfrak{X}}$ be the corresponding Galois covering of \mathfrak{X} .

We extend each unitary u_j to become a unitary u_j on $\tilde{\mathfrak{X}}$ as follows. Recall that

$$u_j(x) = \exp\left(\frac{n\ell_1}{2L\ell_j} i\varphi_j(x)\right) \quad \text{on } X.$$

Let \mathfrak{X}_j be the ‘‘partial’’ doubling of X obtained by identifying the corresponding faces $\partial_{k\pm}$ of X and $-X$ for all $1 \leq k \leq m$ except the faces $\partial_{j\pm}$. The space \mathfrak{X}_j is a manifold with corners, whose boundary consists of $\partial_+(\mathfrak{X}_j)$ and $\partial_-(\mathfrak{X}_j)$. Extend the function φ_j on the chosen copy of X to a real-valued smooth function $\check{\varphi}_j$ on \mathfrak{X}_j such that $\check{\varphi}_j(x) = 0$ in an open neighborhood of $\partial_-(\mathfrak{X}_j)$ in \mathfrak{X} and $\check{\varphi}_j(x) = \frac{4\pi L}{n}$ in an open neighborhood of $\partial_+(\mathfrak{X}_j)$. We define the unitary

$$\check{u}_j(x) = \exp\left(\frac{n\ell_1}{2L\ell_j} i\check{\varphi}_j(x)\right) \quad \text{on } \mathfrak{X}_j.$$

By construction, the unitary $\check{u}_j = 1$ near the boundary of \mathfrak{X}_j , hence actually defines a unitary¹⁶ on \mathfrak{X} , which will still be denoted by \check{u}_j . Let us denote the lift of \check{u}_j to $\tilde{\mathfrak{X}}$ by $u_j(x)$. Then u_j is a unitary on $\tilde{\mathfrak{X}}$ whose restriction on \tilde{X} is u_j .

We consider the following differential operator on $\mathbb{T}^m \times \tilde{\mathfrak{X}}$:

$$\mathcal{D}^{\mathfrak{X}} = \sum_{j=1}^m c_j \frac{\partial}{\partial t_j} + \tilde{D}_{t_1, t_2, \dots, t_m}^{\mathfrak{X}}$$

¹⁶We do not require $\|d\check{\varphi}_j\| \leq 1$ on $\mathfrak{X} \setminus X$, where the norm $\|d\check{\varphi}_j\|$ is taken with respect to the Riemannian metric on \mathfrak{X} .

where c_j is the Clifford multiplication of the unit vector $\frac{\partial}{\partial t_j}$ and $\tilde{D}_{t_1, t_2, \dots, t_m}^{\mathfrak{X}}$ is inductively defined as follows:

$$\tilde{D}_{t_1}^{\mathfrak{X}} = t_1 \tilde{D}^{\mathfrak{X}} + (1 - t_1) u_1 \tilde{D}^{\mathfrak{X}} u_1^{-1}$$

and

$$\tilde{D}_{t_1, t_2, \dots, t_k}^{\mathfrak{X}} := t_k (\tilde{D}_{t_1, \dots, t_{k-1}}^{\mathfrak{X}}) + (1 - t_k) u_k (\tilde{D}_{t_1, \dots, t_{k-1}}^{\mathfrak{X}}) u_k^{-1}$$

for $(t_1, \dots, t_m) \in [0, 1]^m$. More generally, for each subset $\Lambda \subseteq \{1, 2, \dots, m\}$, we define the operator

$$\mathcal{D}_{\Lambda}^{\mathfrak{X}} = \sum_{i=1}^m c_i \frac{\partial}{\partial t_i} + \tilde{D}_{\Lambda}^{\mathfrak{X}}$$

where $\tilde{D}_{\Lambda}^{\mathfrak{X}}$ is defined the same way as $\tilde{D}_{t_1, \dots, t_j, \dots, t_m}^{\mathfrak{X}}$ except that u_k is replaced by the trivial unitary $v \equiv 1$ for every $k \in \Lambda$. See Figure 1 for the case where $m = 2$.

Let us compute the index

$$\sum_{\Lambda \subseteq \{1, 2, \dots, m\}} (-1)^{|\Lambda|} \cdot \text{Ind}_{\Gamma}(\mathcal{D}_{\Lambda}^{\mathfrak{X}}) \tag{4.5}$$

in $KO_{n-m}(C_{\max}^*(\Gamma))$, where $|\Lambda|$ is the cardinality of the set Λ . Since \mathfrak{X} is a closed manifold, the index in line (4.5) does not change if we deform the unitaries u_j through a continuous family of unitaries. In particular, we can deform the unitaries u_j through a continuous family of unitaries so that each u_j becomes trivial (that is, equal to 1) outside a small neighborhood of the hypersurface \mathfrak{Y}_j in \mathfrak{X} , where \mathfrak{Y}_j is the doubling of Y_j . Now




Figure 1. An illustration of the indices in the $m = 2$ case where the horizontal (red) lines represent the unitary u_1 and the vertical (blue) lines represent the unitary u_2 .

we identify a small tubular neighborhood of Y_{th} in \mathfrak{X} with an open set in $Y_{\text{th}} \times \mathbb{T}^m$. By the usual relative higher index theorem for closed manifolds (cf. [3, 33]) or alternatively the proof of Theorem 3.4, we can reduce the computation to the corresponding operators on the closed manifold $Y_{\text{th}} \times \mathbb{T}^m$. Hence it remains to compute the index

$$\sum_{\Lambda \subseteq \{1, 2, \dots, m\}} (-1)^{|\Lambda|} \cdot \text{Ind}_{\Gamma} (\not{D}_{\Lambda}^{Y_{\text{th}} \times \mathbb{T}^m})$$

where $\not{D}_{\Lambda}^{Y_{\text{th}} \times \mathbb{T}^m}$ is the obvious analogue of $\not{D}_{\Lambda}^{\mathfrak{X}}$. Now to simplify the computation even further, we deform the metric on $Y_{\text{th}} \times \mathbb{T}^m$ to a product metric. In this case, the operator $\not{D}^{Y_{\text{th}} \times \mathbb{T}^m}$ becomes

$$\sum_{j=1}^m \left(c_j \frac{\partial}{\partial t_j} + u_j D^{\mathbb{S}^1} u_j^{-1} \right) \widehat{\otimes} 1 + 1 \widehat{\otimes} D^{Y_{\text{th}}}$$

on the space $\mathbb{T}^m \times Y_{\text{th}} \times \mathbb{T}^m$, where without loss of generality we can assume u_j to be the smooth function obtained by projecting to the j -component of \mathbb{T}^m ,

$$Y_{\text{th}} \times \mathbb{T}^m \rightarrow \mathbb{S}^1 \subset \mathbb{C}.$$

The operator $\sum_{j=1}^m (c_j \frac{\partial}{\partial t_j} + u_j D^{\mathbb{S}^1} u_j^{-1})$ has index 1 (cf. [1, Section 7]). Therefore, it follows that

$$\text{Ind}_{\Gamma} (\not{D}^{Y_{\text{th}} \times \mathbb{T}^m}) = \text{Ind}_{\Gamma} (D^{Y_{\text{th}}}) \in K_{n-m}(C_{\max}^*(\Gamma)).$$

Similarly, one can show that

$$\text{Ind}_{\Gamma} (\not{D}_{\Lambda}^{Y_{\text{th}} \times \mathbb{T}^m}) = 0$$

whenever Λ is a proper subset of $\{1, 2, \dots, m\}$. To summarize, we have

$$\sum_{\Lambda \subseteq \{1, 2, \dots, m\}} (-1)^{|\Lambda|} \cdot \text{Ind}_{\Gamma} (\not{D}_{\Lambda}^{\mathfrak{X}}) = \text{Ind}_{\Gamma} (D^{Y_{\text{th}}}).$$

On the other hand, we have (cf. line (4.4))

$$\min_{1 \leq j \leq m} \ell_j = \ell_1 > \frac{L(\frac{8}{\sqrt{3}}C + 4\pi)}{n}.$$

Furthermore, by appropriately choosing the metric on \mathfrak{X} that extends the metric on X , we can assume that

$$\text{supp}_{\mathfrak{X} \setminus X} (u_j - 1) \text{ and } \text{supp}_{\mathfrak{X} \setminus X} (u_k - 1) \text{ are at least } \frac{L(\frac{4}{\sqrt{3}}C + 4\pi)}{n} \text{ apart}$$

for all $j \neq k$, where $\text{supp}_{\mathfrak{X} \setminus X} (u_j - 1)$ is the support of $(u_j - 1)$ in $\mathfrak{X} \setminus X$. Now we apply the same argument as in the proof of Theorem 3.4 (and Remark 3.6) and iterate the difference construction from Lemma 4.4 below. It follows that

$$\sum_{\Lambda \subseteq \{1, 2, \dots, m\}} (-1)^{|\Lambda|} \cdot \text{Ind}_{\Gamma} (\not{D}_{\Lambda}^{\mathfrak{X}}) = 0$$

in $K_{n-m}(C_{\max}^*(\Gamma))$. We arrive at a contradiction, since $\text{Ind}_{\Gamma} (D^{Y_{\text{th}}}) \neq 0$ by assumption. This finishes the proof. ■

Lemma 4.4 (cf. [14, Section 6]). *Let p_1 and p_2 be two idempotents in a Banach algebra B . Then we have*

$$[p_1] - [p_2] = [E(p_1, p_2)] - [E_0]$$

in $K_0(B)$, where

$$E(p_1, p_2) = \begin{pmatrix} 1 + p_2(p_1 - p_2)p_2 & 0 & p_2p_1(p_1 - p_2) & 0 \\ 0 & 0 & 0 & 0 \\ (p_1 - p_2)p_1p_2 & 0 & (1 - p_2)(p_1 - p_2)(1 - p_2) & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad (4.6)$$

and

$$E_0 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

Proof. Consider the invertible element

$$U = \begin{pmatrix} p_2 & 0 & 1 - p_2 & 0 \\ 1 - p_2 & 0 & 0 & p_2 \\ 0 & 0 & p_2 & 1 - p_2 \\ 0 & 1 & 0 & 0 \end{pmatrix}$$

whose inverse is given by

$$U^{-1} = \begin{pmatrix} p_2 & 1 - p_2 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 1 - p_2 & 0 & p_2 & 0 \\ 0 & p_2 & 1 - p_2 & 0 \end{pmatrix}.$$

A direct computation shows that

$$E(p_1, p_2) = U^{-1} \begin{pmatrix} p_1 & 0 & 0 & 0 \\ 0 & 1 - p_2 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} U.$$

This proves the lemma. ■

5. Proofs of Theorems IV and V

In this section, we prove Theorems IV and V. Let us first recall the following notion of subsets with the wrapping property, which was introduced in Definition 1.3.

Definition 5.1 (Subsets with the wrapping property, cf. Definition 1.3). A subset Σ of the standard unit sphere \mathbb{S}^n is said to have *the wrapping property* if Σ is strongly non-separating (cf. Definition 1.2) and furthermore there exists a smooth distance-contracting map $\Phi: \mathbb{S}^n \rightarrow \mathbb{S}^n$ such that the following are satisfied:

- (1a) if n is even, Φ equals the identity map on $N_\varepsilon(\Sigma)$;
- (1b) if n is odd, Φ equals either the identity map or the antipodal map on each of the connected components of $N_\varepsilon(\Sigma)$;
- (2) and¹⁷ $\deg(\Phi) \neq 1$.

Loosely speaking, the class of subsets in \mathbb{S}^n with the wrapping property includes all “reasonable” geometric subsets of \mathbb{S}^n whose sizes are “relatively small”. For example, Lemma 5.3 below gives a sufficient geometric condition for a subset to satisfy the wrapping property. Let us first fix some terminology.

Definition 5.2. Consider the canonical embedding of the unit sphere \mathbb{S}^n inside the Euclidean space \mathbb{R}^{n+1} . For each unit vector $v \in \mathbb{R}^{n+1}$, denote by \mathbb{V}_v^\perp the linear subspace of \mathbb{R}^{n+1} that is orthogonal to v . We define an equator \mathbb{E} of \mathbb{S}^n to be the intersection of \mathbb{V}_v^\perp and \mathbb{S}^n for some unit vector $v \in \mathbb{R}^{n+1}$.

Lemma 5.3. *Let Σ be a strongly non-separating subset of \mathbb{S}^n . If $N_\varepsilon(\Sigma)$ is contained in a geodesic ball of radius $< \frac{\pi}{2}$ for some (hence for all) sufficiently small $\varepsilon > 0$, then Σ has the wrapping property.*

Proof. By assumption, for each sufficiently small $\varepsilon > 0$, there exists a geodesic ball B of radius $r < \frac{\pi}{2}$ that contains $N_\varepsilon(\Sigma)$. Without loss of generality, we assume that there is an equator \mathbb{E} such that B is contained in a hemisphere determined by \mathbb{E} and $\text{dist}(B, \mathbb{E}) > 2\varepsilon$. Let us denote the center of B by x_0 . Consider all geodesics in \mathbb{S}^n of length $\leq \pi$ that originate from x_0 , that is, all the shortest geodesics starting at x_0 and ending at the antipodal point of x_0 . Now we shall “wrap” the geodesics to define a distance-contracting map $\Phi: \mathbb{S}^n \rightarrow \mathbb{S}^n$ such that Φ equals the identity map on B and its image $\Phi(\mathbb{S}^n)$ lies in the hemisphere that contains B . In particular, Φ is not surjective, hence $\deg(\Phi) = 0$.

More precisely, let us first consider a smooth function $f': [-\frac{\pi}{2}, \frac{\pi}{2}] \rightarrow [-1, 1]$ such that (cf. Figure 2)

- (i) f' is odd, that is, $f'(-t) = -f'(t)$,
- (ii) $f'(t) = -1$ for all $t \in [\varepsilon, \frac{\pi}{2}]$,
- (iii) and $f'(t) \leq 0$ for all $t \in [0, \frac{\pi}{2}]$.

Define $f: [-\frac{\pi}{2}, \frac{\pi}{2}] \rightarrow \mathbb{R}$ by setting

$$f(s) = -\frac{\pi}{2} + \int_{-\frac{\pi}{2}}^s f'(t) dt.$$

¹⁷For example, if Φ is not surjective, then clearly $\deg(\Phi) = 0 \neq 1$.




Figure 2. The graph of f' .

For each shortest geodesic γ going from x_0 to its antipodal point, we parametrize γ by its arc length so that the intersection point of γ with the equator \mathbb{E} becomes the origin of the interval $[-\frac{\pi}{2}, \frac{\pi}{2}]$ and x_0 becomes $-\frac{\pi}{2}$ with respect to the parametrization. Now we define $\Phi_{\mathbb{E}}: \mathbb{S}^n \rightarrow \mathbb{S}^n$ by setting

$$\Phi_{\mathbb{E}}(\gamma(t)) = \gamma(f(t)) \tag{5.1}$$

for each $t \in [-\frac{\pi}{2}, \frac{\pi}{2}]$. For later references, let us call $\Phi_{\mathbb{E}}$ a wrapping map along the equator \mathbb{E} . For brevity, let us denote it by Φ . By construction, the wrapping map

$$\Phi: \mathbb{S}^n \rightarrow \mathbb{S}^n$$

is a smooth¹⁸ distance-contracting map such that Φ equals the identity map on B and $\Phi(\mathbb{S}^n)$ lies in a hemisphere, hence $\deg(\Phi) = 0 \neq 1$. This finishes the proof. ■

Example 5.4. By Lemma 5.3, the following subsets of \mathbb{S}^n have the wrapping property:

- (a) an open or closed geodesic ball of radius $< \frac{\pi}{2}$,
- (b) any compact simplicial complex of codimension ≥ 2 that is contained in a geodesic ball of radius $< \frac{\pi}{2}$.

For odd dimensional spheres, the following collection of subsets also satisfy the wrapping property.

Lemma 5.5. *Let Σ be a strongly non-separating subset of \mathbb{S}^{2k+1} . Let $\{\mathbb{E}_j\}_{1 \leq j \leq 2k+2}$ be a collection of $(2k + 2)$ mutually orthogonal equators of \mathbb{S}^{2k+1} so that they divide \mathbb{S}^{2k+1} into $2^{(2k+2)}$ regions. If $N_\varepsilon(\Sigma)$ is contained in an antipodal pair of such regions for some (hence for all) sufficiently small $\varepsilon > 0$, then Σ satisfies the wrapping property.*

¹⁸Due to the specific properties of f , the map Φ is smooth everywhere. In particular, Φ is smooth at the antipodal point of x_0 .

Proof. Since by assumption $N_\varepsilon(\Sigma)$ is contained in a pair of antipodal regions determined by the equators $\{\mathbb{E}_j\}_{1 \leq j \leq 2k+2}$, we can choose a wrapping map $\Phi_{\mathbb{E}_j}$ associated to \mathbb{E}_j as defined in line (5.1) for each $1 \leq j \leq 2k + 2$ such that their composition

$$\Phi := \Phi_{\mathbb{E}_1} \circ \Phi_{\mathbb{E}_2} \circ \cdots \circ \Phi_{\mathbb{E}_{2k+2}}$$

satisfies the desired properties (1b) and (2) in Definition 5.1. This finishes the proof. ■

Example 5.6. By Lemma 5.5, the following subsets of an odd dimensional sphere \mathbb{S}^{2k+1} have the wrapping property:

- (a) a pair of antipodal points on \mathbb{S}^{2k+1} ,
- (b) a pair of antipodal geodesic balls of radius $< \frac{\pi}{6}$ in \mathbb{S}^{2k+1} .
- (c) any compact simplicial complex of codimension ≥ 2 that is contained in a pair of antipodal geodesic balls of radius $< \frac{\pi}{6}$ in \mathbb{S}^{2k+1} .

For a given $\varepsilon > 0$, the geometry of the ε -neighborhood $N_\varepsilon(\Sigma)$ of Σ can be very wild, in particular at the boundary $\partial \overline{N_\varepsilon}$. However, by enlarging or shrinking $N_\varepsilon(\Sigma)$ if necessary, we can in fact always find small neighborhoods of Σ that are manifolds with boundary or manifolds with corners.

Lemma 5.7. *Let Σ be a subset of \mathbb{S}^n . Then for any sufficiently small $\varepsilon > 0$, there is a subspace $X_\varepsilon \subset \mathbb{S}^n$ with $\mathbb{S}^n \setminus N_{2\varepsilon}(\Sigma) \subset X_\varepsilon \subset \mathbb{S}^n \setminus N_\varepsilon(\Sigma)$ such that X_ε is an n -dimensional compact manifold with corners. Furthermore, if $N_\varepsilon(\Sigma)$ is non-separating for all sufficiently small $\varepsilon > 0$, then X_ε can also be chosen to be path-connected for all sufficiently small $\varepsilon > 0$.*

Proof. Let $\mathcal{U} = \{U_j\}$ be an open cover of $\overline{N_\varepsilon(\Sigma)}$ consisting of geodesically convex balls of radius $\leq \frac{\varepsilon}{2}$. Note that $\overline{N_\varepsilon(\Sigma)}$ is closed in \mathbb{S}^n , hence compact. It follows that $\overline{N_\varepsilon(\Sigma)}$ admits a finite open cover \mathcal{V} consisting of finitely many members of \mathcal{U} . Without loss of generality, we assume

$$V \cap \overline{N_\varepsilon(\Sigma)} \neq \emptyset$$

for each member V of \mathcal{V} . Denote by W the union of all members of \mathcal{V} . Then the closure \overline{W} of W is contained in $N_{2\varepsilon}(\Sigma)$.

Define X_ε to be $\mathbb{S}^n \setminus W$. By construction,¹⁹ X_ε is an n -dimensional compact manifold with corners under the metric inherited from \mathbb{S}^n such that

$$\mathbb{S}^n \setminus N_{2\varepsilon}(\Sigma) \subset X_\varepsilon \subset \mathbb{S}^n \setminus N_\varepsilon(\Sigma).$$

Furthermore, the above construction shows that if $\mathbb{S}^n \setminus N_{2\varepsilon}(\Sigma)$ is path-connected, then X_ε is path-connected. ■

¹⁹We do not rule out the possibility that X_ε could be the empty set.

Now we are ready to prove Theorem IV, which answers positively an open question of Gromov, cf. [9, page 687, specific problem] and [10, Section 3.9].

Theorem 5.8 (Theorem IV). *Let Σ be a subset of the standard unit sphere \mathbb{S}^n contained in a geodesic ball of radius $r < \frac{\pi}{2}$. Let (X, g_0) be the standard unit sphere \mathbb{S}^n minus Σ . If a (possibly incomplete) Riemannian metric g on X satisfies that*

- (1) *there is a λ_n -Lipschitz homeomorphism $\varphi: (X, g) \rightarrow (X, g_0)$,*
- (2) *and $\text{Sc}(g) \geq n(n - 1) = \text{Sc}(g_0)$,*

then

$$\lambda_n \geq \sqrt{1 - \frac{C_r}{n^2}}$$

where²⁰

$$C_r = \frac{8C^2}{(\frac{\pi}{2} - r)^2}$$

and C is a universal constant from Theorem I. Consequently, the lower bound for λ_n approaches 1, as $n \rightarrow \infty$.

Proof. We prove the theorem by contradiction. Assume to the contrary that

$$\lambda_n < \sqrt{1 - \frac{C_r}{n^2}}.$$

To avoid ambiguity, let us denote (X, g_0) by \underline{X} for the rest of the proof.

Let us first prove the even dimensional case. Recall the $\mathbb{C}\ell_n$ -Dirac bundle E_0 over \mathbb{S}^n ,

$$E_0 = P_{\text{Spin}}(\mathbb{S}^n) \times_{\ell} \mathbb{C}\ell_n \tag{5.2}$$

where $\ell: \text{Spin}_n \rightarrow \text{End}(\mathbb{C}\ell_n)$ is the representation given by left multiplication. Equip E_0 with the canonical Riemannian connection determined by the presentation $\ell: P_{\text{Spin}}(\mathbb{S}^n) \rightarrow \text{End}(\mathbb{C}\ell_n)$. Furthermore, when n is even, E_0 carries a natural $\mathbb{Z}/2$ -grading $E_0 = E_0^+ \oplus E_0^-$. By the Atiyah–Singer index theorem [2], the index of the twisted Dirac operator $D_{E_0^+}^{\mathbb{S}^n}$ is nonzero (cf. [19, equation (4.7)]).

We shall give an explicit description of the bundle E_0 as a sub-bundle of a trivial vector bundle over \mathbb{S}^n so that E_0 can be viewed a projection p in $M_k(C(\mathbb{S}^n)) = M_k(\mathbb{C}) \otimes C(\mathbb{S}^n)$, where $C(\mathbb{S}^n)$ is the C^* -algebra of continuous functions on \mathbb{S}^n . Consider the canonical embedding of the unit sphere \mathbb{S}^n inside the Euclidean space \mathbb{R}^{n+1} . Let $V = \mathbb{R}^{n+1} \times \mathbb{C}\ell_{n+1}$ be the canonical $\mathbb{C}\ell_{n+1}$ -Dirac bundle over \mathbb{R}^{n+1} . Clearly, V is a trivial vector bundle. Let us still denote by V the restriction of V on \mathbb{S}^n . Then we see that E_0 is a sub-bundle of V . Denote by v the outward unit normal vector field of \mathbb{S}^n in

²⁰If $n = \dim \mathbb{S}^n$ is odd, our proof of Theorem 5.8 in fact shows that we can improve C_r to be $\frac{4C^2}{(\frac{\pi}{2}-r)^2}$ instead of $\frac{8C^2}{(\frac{\pi}{2}-r)^2}$.

\mathbb{R}^{n+1} . Then E_0 is isomorphic to the sub-bundle of V determined by the following Bott projection²¹

$$p_n = \frac{i\bar{c}(v) + 1}{2} \tag{5.3}$$

where $\bar{c}(v)$ is the Clifford multiplication of v on $V = \mathbb{S}^n \times \mathbb{C}\ell_{n+1}$.

By assumption, Σ is contained in a geodesic ball $B_r(x_0)$ centered at x_0 of radius $r < \frac{\pi}{2}$. Let us define

$$\underline{X}_\varepsilon = \mathbb{S}^n \setminus B_{r+\frac{\varepsilon}{2}}^o(x_0)$$

where $B_{r+\frac{\varepsilon}{2}}^o(x_0)$ is the open geodesic ball centered at x_0 of radius $(r + \frac{\varepsilon}{2})$. By the proof of Lemma 5.3, there exists a smooth distance-contracting map $\Phi: \mathbb{S}^n \rightarrow \mathbb{S}^n$ such that

- (1) Φ equals the identity map on the $(\frac{\pi}{2} - r - \varepsilon)$ -neighborhood of $\mathbb{S}^n \setminus \underline{X}_\varepsilon$;
- (2) and $\text{deg}(\Phi) \neq 1$.

In order to apply the relative index theorem (Theorem 3.4), we shall view X_ε as a (topological) subset of the n -dimensional sphere. Since (X_ε, g) is an n -dimensional manifold with boundary, we can extend the Riemannian metric g on X_ε to a Riemannian metric on the sphere. Let us denote by \mathfrak{S} the resulting n -dimensional sphere with this new metric $g_\mathfrak{S}$. Of course, the metric $g_\mathfrak{S}$ generally does *not* satisfy the Lipschitz bound and scalar curvature bound on the complement of X_ε in \mathfrak{S} , when compared to the standard metric g_0 on \mathbb{S}^n . Consider the (set-theoretic) identity map

$$\mathbf{1}: \mathfrak{S} \rightarrow \mathbb{S}^n.$$

The pullback bundles of V by the map $\mathbf{1}: \mathfrak{S} \rightarrow \mathbb{S}^n$ and the map $\Phi \circ \mathbf{1}: \mathfrak{S} \rightarrow \mathbb{S}^n$ are identical, since V is a trivial vector bundle with its canonical trivial connection. We shall denote this pullback bundle on \mathfrak{S} by $W = \mathfrak{S} \times \mathbb{C}\ell_{n+1}$ from now on. Let \mathcal{S} be the spinor bundle of $(\mathfrak{S}, g_\mathfrak{S})$.

Let $p_1 = (\mathbf{1})^*(p_n)$ and $p_2 = (\Phi \circ \mathbf{1})^*(p_n)$ be the projections induced from the Bott projection p_n on \mathbb{S}^n , by the maps $\mathbf{1}$ and $\Phi \circ \mathbf{1}$ respectively. By construction, we have $p_1 = p_2$ on the $(\frac{\pi}{2} - r - \varepsilon)$ -neighborhood of $\mathfrak{S} \setminus X_\varepsilon$. The projections p_1 and p_2 can be viewed as endomorphisms of the bundle $\mathcal{S} \otimes W$. More precisely, the bundle homomorphism $1 \otimes p_j: \mathcal{S} \otimes W \rightarrow \mathcal{S} \otimes W$ satisfies that $(1 \otimes p_j)^2 = 1 \otimes p_j$ and $(1 \otimes p_j)^* = 1 \otimes p_j$, for $j = 1, 2$. Now consider the twisted Dirac operators $D_{p_j} := p_j D p_j$. Furthermore, since n is even, the bundle $p_j W$ carries a natural $\mathbb{Z}/2$ -grading inherited from the $\mathbb{Z}/2$ -grading on E_0 . We have

$$D_{p_j} = \begin{pmatrix} 0 & D_{p_j}^- \\ D_{p_j}^+ & 0 \end{pmatrix}$$

with respect to the decomposition $p_j W = (p_j W)^+ \oplus (p_j W)^-$.

The commutator $[D, p_j]$ is an endomorphism of the bundle $\mathcal{S} \otimes W$. Denote by $[D, p_j]_x: (\mathcal{S} \otimes W)_x \rightarrow (\mathcal{S} \otimes W)_x$ the endomorphism at the point $x \in \mathfrak{S}$. A key step of the proof is the following estimate for the operator norm of $[D, p_j]_x$ for every point $x \in X_\varepsilon$.

²¹To be precise, the K -theory class of p_n is a nonzero multiple of the actual Bott generator of $K^0(\mathbb{S}^n)$.

For each $x \in X_\varepsilon$, we can choose a local g_0 -orthonormal tangent frame $\{\underline{e}_1, \dots, \underline{e}_n\}$ for $T\underline{X}_\varepsilon$ and a local g -orthonormal tangent frame $\{e_1, \dots, e_n\}$ for TX_ε near x such that for each $1 \leq k \leq n$, we have

$$\mathbf{1}_*(e_k) = \mu_k \underline{e}_k$$

for some $\mu_k \geq 0$. Since $\mathbf{1}: (X_\varepsilon, g) \rightarrow (\underline{X}_\varepsilon, g_0)$ is λ_n -Lipschitz, we have $\mu_k \leq \lambda_n$ for all $1 \leq k \leq n$. If we write

$$D = \sum_{k=1}^n c(e_k) \nabla_{e_k},$$

then we have

$$\| [D, \mathfrak{p}_1]_x \| = \left\| \sum_{k=1}^n [\mu_k c(\underline{e}_k) \nabla_{\underline{e}_k}, \mathfrak{p}_n]_x \right\|.$$

A similar conclusion holds for \mathfrak{p}_2 , since $\Phi \circ \mathbf{1}: (X_\varepsilon, g) \rightarrow \mathbb{S}^n$ is also λ_n -Lipschitz.

Claim 5.9. We have

$$\| [D, \mathfrak{p}_j]_x \| \leq \frac{n \cdot \lambda_n}{2}$$

for all $x \in X_\varepsilon$ and for both $j = 1, 2$.

By the discussion above, we need to estimate

$$\left\| \sum_{k=1}^n [\mu_k c(\underline{e}_k) \nabla_{\underline{e}_k}, \mathfrak{p}_n]_x \right\| \tag{5.4}$$

for each $x \in \mathbb{S}^n$. Recall that v is the outward unit normal vector field of \mathbb{S}^n in \mathbb{R}^{n+1} . In particular, at a point $x = (x_1, x_2, \dots, x_{n+1}) \in \mathbb{S}^n \subset \mathbb{R}^{n+1}$, we have

$$\bar{c}(v)_x = \sum_{k=1}^{n+1} x_k \bar{c}_k$$

where \bar{c}_j is the Clifford multiplication of the unit vector $\frac{\partial}{\partial x_j}$ on $V = \mathbb{S}^n \times \mathbb{C}\ell_{n+1}$ from the right. Since $\text{SO}(n + 1)$ acts transitively on \mathbb{S}^n , it suffices to estimate the term in line (5.4) at the point $\underline{x} = (0, \dots, 0, 1) \in \mathbb{S}^n \subset \mathbb{R}^{n+1}$. At this point \underline{x} , after a local coordinate change if necessary, we have²²

$$\sum_{k=1}^n \mu_k c(\underline{e}_k) \nabla_{\underline{e}_k} = \sum_{k=1}^n \mu_k c_k \frac{\partial}{\partial x_k}$$

where c_k is the Clifford multiplication of the unit vector $\frac{\partial}{\partial x_j}$ on the spinor bundle of \mathbb{S}^n from the left. We conclude that

$$\sum_{k=1}^n [\mu_k c(\underline{e}_k) \nabla_{\underline{e}_k}, \mathfrak{p}_n]_{\underline{x}} = \frac{i}{2} \sum_{k=1}^n \left[\mu_k c_k \frac{\partial}{\partial x_k}, \bar{c}(v) \right]_{\underline{x}} = \frac{i}{2} \sum_{j=1}^n \mu_k c_k \otimes \bar{c}_k.$$

²²Here the term $\partial/\partial x_{n+1}$ does not appear, since it is in the normal direction.

Since $\|c_k \otimes \bar{c}_k\| = 1$ for all $1 \leq k \leq n$, it follows that

$$\left\| \sum_{k=1}^n [\mu_k c(\underline{e}_k) \nabla_{\underline{e}_k}, \mathbf{p}_n]_{\underline{x}} \right\| \leq \frac{\sum_{k=1}^n \mu_k}{2} \leq \frac{n \cdot \lambda_n}{2}.$$

This proves the claim.

For brevity, let us write \mathfrak{p} in place of \mathfrak{p}_j in the following estimation. We have

$$\begin{aligned} \langle \mathfrak{p} D \mathfrak{p} f, \mathfrak{p} D \mathfrak{p} f \rangle &= \langle \mathfrak{p} D \mathfrak{p} D \mathfrak{p} f, \mathfrak{p} f \rangle \\ &= \langle \mathfrak{p} [D, \mathfrak{p}] D \mathfrak{p} f, \mathfrak{p} f \rangle + \langle \mathfrak{p} D^2 \mathfrak{p} f, \mathfrak{p} f \rangle \\ &= -\langle D \mathfrak{p} f, [D, \mathfrak{p}] \mathfrak{p} f \rangle + \langle \mathfrak{p} D^2 \mathfrak{p} f, \mathfrak{p} f \rangle \\ &\geq -\frac{1}{2} \langle D \mathfrak{p} f, D \mathfrak{p} f \rangle - \frac{1}{2} \langle [D, \mathfrak{p}] \mathfrak{p} f, [D, \mathfrak{p}] \mathfrak{p} f \rangle + \langle \mathfrak{p} D^2 \mathfrak{p} f, \mathfrak{p} f \rangle \\ &\geq \frac{1}{2} \langle D^2 \mathfrak{p} f, \mathfrak{p} f \rangle - \frac{1}{2} \langle [D, \mathfrak{p}] \mathfrak{p} f, [D, \mathfrak{p}] \mathfrak{p} f \rangle. \end{aligned}$$

By the inequality in line (3.2), we have

$$\langle D^2 \mathfrak{p} f, \mathfrak{p} f \rangle \geq \frac{n}{n-1} \left\langle \frac{\kappa}{4} \mathfrak{p} f, \mathfrak{p} f \right\rangle,$$

where $\kappa := \text{Sc}(g)$. It follows that

$$(\mathfrak{p} D \mathfrak{p})^2 \geq \frac{1}{2} \left(\frac{n\kappa}{4(n-1)} - [D, \mathfrak{p}]^* [D, \mathfrak{p}] \right) \quad \text{on } C_c^\infty(X_\varepsilon^\circ, \mathcal{S} \otimes \mathfrak{p}W).$$

Here, $C_c^\infty(X_\varepsilon^\circ, \mathcal{S} \otimes \mathfrak{p}W)$ is the space of compactly supported smooth sections of the sub-bundle $\mathcal{S} \otimes \mathfrak{p}W \subset \mathcal{S} \otimes W$. By assumption, we have $\kappa = \text{Sc}(g) \geq n(n-1)$. It follows from Claim 5.9 that

$$\|D_{\mathfrak{p}_j} v\| \geq n \sqrt{\frac{(1-\lambda_n^2)}{8}} \|v\| \tag{5.5}$$

for all $v \in C_c^\infty(X_\varepsilon^\circ, \mathcal{S} \otimes \mathfrak{p}_j W)$ and for both $j = 1, 2$. Furthermore, the same conclusion from line (5.5) also holds for both $D_{\mathfrak{p}_j}^+$ and $D_{\mathfrak{p}_j}^-$ on $C_c^\infty(X_\varepsilon^\circ, \mathcal{S} \otimes \mathfrak{p}_j W)$.

Since we have assumed that

$$\lambda_n < \sqrt{1 - \frac{C_r}{n^2}} \quad \text{with } C_r = \frac{8C^2}{(\frac{\pi}{2} - r)^2}$$

it follows that

$$\left(\frac{\pi}{2} - r - \varepsilon\right) n \sqrt{\frac{(1-\lambda_n^2)}{8}} > C$$

for some sufficiently small $\varepsilon > 0$. By construction, the operators $D_{\mathfrak{p}_1}^+$ and $D_{\mathfrak{p}_2}^+$ coincide on the $(\frac{\pi}{2} - r - \varepsilon)$ -neighborhood of $\mathfrak{C} \setminus X_\varepsilon$. It follows from Theorem 3.4 that

$$\text{Ind}(D_{\mathfrak{p}_1}^+) - \text{Ind}(D_{\mathfrak{p}_2}^+) = 0.$$

On the other hand, by the Atiyah–Singer index theorem [2], we have

$$\text{Ind}(D_{p_1}^+) - \text{Ind}(D_{p_2}^+) = (1 - \text{deg}(\Phi)) \cdot \text{Ind}(D_{E_0^+}^{\mathbb{S}^n}) \in K_0(\mathcal{K}) = \mathbb{Z}.$$

Since $\text{Ind}(D_{E_0^+}^{\mathbb{S}^n}) \neq 0$ (cf. [19, equation (4.7)]), we conclude that

$$1 - \text{deg } \Phi = 0.$$

This contradicts the fact that $\text{deg } \Phi \neq 1$. This finishes the proof for the even dimensional case.

Now let us prove the theorem in the odd dimensional case. Since the key ideas are similar to the even dimensional case, we shall be brief. Again, consider the canonical embedding of the unit sphere \mathbb{S}^n inside the Euclidean space \mathbb{R}^{n+1} . Let $V = \mathbb{R}^{n+1} \times \mathbb{C}\ell_{n+1}$ be the canonical $\mathbb{C}\ell_{n+1}$ -Dirac bundle over \mathbb{R}^{n+1} . Denote by v the outward unit normal vector field of \mathbb{S}^n in \mathbb{R}^{n+1} . Since $n + 1$ is even in the current case, there is a canonical $\mathbb{Z}/2$ -grading of V induced by the following element:

$$\omega = i^{\frac{n+1}{2}} c_1 \cdots c_{n+1}$$

where c_j is the Clifford multiplication by the standard basis element $e_j \in \mathbb{R}^{n+1}$. Let us write $V = V^+ \oplus V^-$ for this $\mathbb{Z}/2$ -decomposition. Note that V^+ and V^- are trivial vector bundles of the same dimension. In particular, we can choose a fixed unitary matrix $U \in U_n(\mathbb{C})$ to fiberwise identify V^+ and V^- . Then a Bott element \mathbf{v}_n —a nonzero multiple of a generator of $K^1(\mathbb{S}^n) = K_1(C(\mathbb{S}^n))$ —is given by the unitary

$$\mathbf{v}_n = i\bar{c}(v) \tag{5.6}$$

where $\bar{c}(v)$ is the Clifford multiplication of v on $V^+ = \mathbb{S}^n \times \mathbb{C}\ell_{n+1}^+$.

Similar to the proof of Theorem 4.1, let us consider the following Dirac-type operator on $\mathbb{S}^1 \times \mathbb{S}^n$:

$$\not{D} = c \cdot \frac{d}{dt} + D_t$$

where c is the Clifford multiplication of the unit vector d/dt and

$$D_t := tD^{\mathbb{S}^n} + (1 - t)\mathbf{v}_n D^{\mathbb{S}^n} \mathbf{v}_n^{-1} \tag{5.7}$$

for each $t \in [0, 1]$. Here we have chosen the parametrization $\mathbb{S}^1 = [0, 1]/\{0, 1\}$. By the Atiyah–Singer index theorem [2], we have

$$\text{Ind}(\not{D}) = \int_{\mathbb{S}^n} \widehat{A}(\mathbb{S}^n) \wedge \text{ch}(\mathbf{v}_n) \neq 0$$

where $\widehat{A}(\mathbb{S}^n)$ is the \widehat{A} -form of \mathbb{S}^n and $\text{ch}(\mathbf{v}_n)$ is the odd dimensional Chern character of \mathbf{v}_n .

Let X_ε and \mathfrak{S} be as above. Also, denote by $W = \mathfrak{S} \times \mathbb{C}\ell_{n+1}^+$ the pullback bundle of the trivial bundle V^+ . Pull back the unitary \mathbf{v}_n by the maps $\mathbf{1}: \mathfrak{S} \rightarrow \mathbb{S}^n$ and $\Phi \circ \mathbf{1}: \mathfrak{S} \rightarrow \mathbb{S}^n$

and denote the resulting unitaries by $v_1 = \mathbf{1}^*(v_n)$ and $v_2 = (\Phi \circ \mathbf{1})^*(v_n)$ respectively. By construction, we have $v_1 = v_2$ on the $(\frac{\pi}{2} - r - \varepsilon)$ -neighborhood of $\mathfrak{S} \setminus X_\varepsilon$.

Consider the Dirac-type operators

$$\mathcal{D}_{v_j}^X = c \cdot \frac{d}{dt} + D_{v_j,t}$$

on $\mathbb{S}^1 \times \mathfrak{S}$, where

$$D_{v_j,t} = tD + (1-t)v_j D v_j^{-1},$$

for $j = 1, 2$. In particular, $\mathcal{D}_{v_1}^X$ and $\mathcal{D}_{v_2}^X$ coincide on the $(\frac{\pi}{2} - r - \varepsilon)$ -neighborhood of $\mathbb{S}^1 \times (\mathfrak{S} \setminus X_\varepsilon)$. Now the same calculation as in the proof of Theorem 4.1 shows that

$$\mathcal{D}_{v_j}^2 = -\frac{d^2}{dt^2} + D_{v_j,t}^2 + c[D, v_j]v_j^{-1}$$

with

$$D_{v_j,t}^2 = tD^2 + (1-t)v_j D^2 v_j^{-1} + t(1-t)[D, v_j][D, v_j^{-1}],$$

cf. line (4.2). It follows that

$$D_{v_j,t}^2 \geq \frac{n^2}{4} - \frac{n^2}{4}\lambda_n^2 = (1 - \lambda_n^2)\frac{n^2}{4} \quad \text{on } C_c^\infty(\mathbb{S}^1 \times X_\varepsilon, \mathcal{S} \otimes W).$$

Similar to the rescaling argument from (4.3), for each $\mu > 0$, we define the rescaled version of \mathcal{D}_{v_j} to be

$$\mathcal{D}_{v_j,\mu} = c \cdot \frac{d}{dt} + \mu D_{v_j,t}.$$

The same calculation as above shows that

$$(\mathcal{D}_{v_j,\mu})^2 \geq \mu^2(1 - \lambda_n^2)\frac{n^2}{4} - \mu\frac{n}{2} \quad \text{on } C_c^\infty(\mathbb{S}^1 \times X_\varepsilon, \mathcal{S} \otimes W).$$

Now, by applying the quantitative relative index theorem (Theorem 3.4), the rest of the proof for the odd dimensional case proceeds in the same way as the even dimensional case. We conclude that

$$\lambda_n \geq \sqrt{1 - \frac{4C^2}{(\frac{\pi}{2} - r)^2 n^2}}$$

where C is the universal constant from Theorem I. This completes the proof of Theorem 5.8. ■

In the case where $n = \dim \mathbb{S}^n$ is odd, the same proof for the odd dimensional case of Theorem IV in fact also proves the λ -Lipschitz rigidity of positive scalar curvature metrics on $\mathbb{S}^n \setminus \Sigma$ when Σ is a subset of \mathbb{S}^n contained in a pair of antipodal geodesic balls of radius $r < \frac{\pi}{6}$.

Theorem 5.10. *Let $\mathbb{S}^n = \mathbb{S}^{2k+1}$ be an odd dimensional standard unit sphere. Let Σ be a subset of the standard unit sphere \mathbb{S}^n contained in a pair of antipodal geodesic balls of radius $r < \frac{\pi}{6}$. Let (X, g_0) be the standard unit sphere \mathbb{S}^n minus Σ . If a (possibly incomplete) Riemannian metric g on X satisfies that*

- (1) *there is a λ_n -Lipschitz homeomorphism $\varphi: (X, g) \rightarrow (X, g_0)$,*
- (2) *and $\text{Sc}(g) \geq n(n - 1) = \text{Sc}(g_0)$,*

then

$$\lambda_n \geq \sqrt{1 - \frac{C_r}{n^2}}$$

where

$$C_r = \frac{4C^2}{(\frac{\pi}{6} - r)^2}$$

and C is the universal constant from Theorem 3.4.

Proof. By assumption, Σ is contained in a pair of antipodal geodesic balls of radius $r < \frac{\pi}{6}$. Let $\underline{X}_\varepsilon \subset \mathbb{S}^n \setminus \Sigma$ be the closed subset of \mathbb{S}^n with two antipodal open geodesic balls of radius $(r + \frac{\varepsilon}{2})$ removed. By the proof of Lemma 5.5, there exists a smooth distance-contracting map $\Phi: \mathbb{S}^n \rightarrow \mathbb{S}^n$ such that

- (1) Φ equals either the identity map or the antipodal map on each path component of the $(\frac{\pi}{6} - r - \varepsilon)$ -neighborhood of $\mathbb{S}^n \setminus \underline{X}_\varepsilon$;
- (2) and $\text{deg}(\Phi) \neq 1$.

We view X_ε as a (topological) subset of the n -dimensional sphere and extend the Riemannian metric g on X_ε to a Riemannian metric on the sphere. Let us denote by \mathfrak{C} the resulting n -dimensional sphere with this new metric $g_{\mathfrak{C}}$. Moreover, let \mathbf{v}_n be the Bott unitary defined in line (5.6).

Similar to the proof for the odd dimensional case of Theorem 5.8, we pull back the unitary \mathbf{v}_n by the maps $\mathbf{1}: \mathfrak{C} \rightarrow \mathbb{S}^n$ and $\Phi \circ \mathbf{1}: \mathfrak{C} \rightarrow \mathbb{S}^n$ and denote the resulting unitaries by $\mathbf{v}_1 = \mathbf{1}^*(\mathbf{v}_n)$ and $\mathbf{v}_2 = (\Phi \circ \mathbf{1})^*(\mathbf{v}_n)$ respectively. By construction, we have $\mathbf{v}_1 = \pm \mathbf{v}_2$ on the $(\frac{\pi}{6} - r - \varepsilon)$ -neighborhood of $\mathfrak{C} \setminus X_\varepsilon$.

Consider the Dirac-type operators

$$\mathcal{D}_{\mathbf{v}_j}^X = c \cdot \frac{d}{dt} + D_{\mathbf{v}_j,t}$$

on $\mathbb{S}^1 \times \mathfrak{C}$, where

$$D_{\mathbf{v}_j,t} = tD + (1 - t)\mathbf{v}_j D \mathbf{v}_j^{-1},$$

for $j = 1, 2$. In particular, $\mathcal{D}_{\mathbf{v}_1}^X$ and $\mathcal{D}_{\mathbf{v}_2}^X$ coincide on the $(\frac{\pi}{6} - r - \varepsilon)$ -neighborhood of $\mathbb{S}^1 \times (\mathfrak{C} \setminus X_\varepsilon)$, since $\mathbf{v}_1 = \pm \mathbf{v}_2$ on the $(\frac{\pi}{6} - r - \varepsilon)$ -neighborhood of $\mathfrak{C} \setminus X_\varepsilon$.

Now the rest of the proof proceeds the same way as the proof for the odd dimensional case of Theorem 5.8. We omit the details. This finishes the proof. ■

As a consequence of Theorem 5.8, we have the following λ -Lipschitz rigidity theorem for hemispheres. This answers (asymptotically) an open question of Gromov on the sharpness of the constant λ_n for the λ_n -Lipschitz rigidity of positive scalar curvature metrics on hemispheres [10, Section 3.8].

Theorem 5.11 (Theorem V). *Let (X, g_0) be the standard unit hemisphere \mathbb{S}_+^n . If a Riemannian metric g on X satisfies that*

- (1) *there is a λ_n -Lipschitz homeomorphism $\varphi: (X, g) \rightarrow (X, g_0)$,*
- (2) *and $\text{Sc}(g) \geq n(n - 1) = \text{Sc}(g_0)$,*

then

$$\lambda_n \geq \left(1 - \sin \frac{\pi}{\sqrt{n}}\right) \sqrt{1 - \frac{8C^2}{\pi^2 n}}$$

where C is the universal constant from Theorem I. Consequently, the lower bound for λ_n approaches 1, as $n \rightarrow \infty$.

This theorem is asymptotically optimal in the sense that the lower bound for λ_n becomes sharp, as $n = \dim \mathbb{S}^n \rightarrow \infty$. In particular, it significantly improves the lower bound for λ_n in Corollary 1.1 when $n = \dim \mathbb{S}^n$ is large.

Proof of Theorem 5.11. Let Y be the subspace of the standard unit sphere \mathbb{S}^n with an open geodesic ball of radius $(\frac{\pi}{2} - \frac{\pi}{\sqrt{n}})$ removed. It is not difficult to see there exists a $(1 - \sin \frac{\pi}{\sqrt{n}})^{-1}$ -Lipschitz homeomorphism from \mathbb{S}_+^n to Y . By composing with the map $\varphi: (X, g) \rightarrow (X, g_0)$, we obtain a $\lambda_n(1 - \sin \frac{\pi}{\sqrt{n}})^{-1}$ -Lipschitz homeomorphism $\varphi: (X, g) \rightarrow (Y, g_0)$. It follows from Theorem 5.8 that

$$\lambda_n \left(1 - \sin \frac{\pi}{\sqrt{n}}\right)^{-1} \geq \sqrt{1 - \frac{C_r}{n^2}}$$

where²³

$$C_r = \frac{8C^2}{(\frac{\pi}{2} - r)^2} \quad \text{and} \quad r = \frac{\pi}{2} - \frac{\pi}{\sqrt{n}}.$$

In other words, we have

$$\lambda_n \geq \left(1 - \sin \frac{\pi}{\sqrt{n}}\right) \sqrt{1 - \frac{8C^2}{\pi^2 n}}.$$

This finishes the proof. ■

²³If $n = \dim \mathbb{S}^n$ is odd, we can set $C_r = \frac{4C^2}{(\frac{\pi}{2} - r)^2}$ instead of $\frac{8C^2}{(\frac{\pi}{2} - r)^2}$.

A. Friedrichs extensions in the maximal group C^* -algebra setting

In this section, we show the existence of Friedrichs extensions of semibounded symmetric operators in the maximal group C^* -algebra setting.

More precisely, suppose X is an n -dimensional compact spin manifold with boundary or corners. Let \mathcal{S} be the Cl_n -Clifford bundle over X and D a Cl_n -linear Dirac-type operator acting on \mathcal{S} . Let \tilde{X} be a Galois Γ -covering space of X and $\tilde{\mathcal{S}}$ (resp. \tilde{D}) the lift of \mathcal{S} (resp. D). By the Lichnerowicz formula, we have

$$\tilde{D}^2 = \nabla^* \nabla + \mathcal{R}$$

where \mathcal{R} is a symmetric bundle endomorphism of $\tilde{\mathcal{S}}$. For the rest of this section, let us assume there exists $\sigma > 0$ such that

$$\mathcal{R}(x) \geq \frac{(n - 1)\sigma^2}{n}$$

for all $x \in \tilde{X}$.

Definition A.1. We define $H_0^1(\tilde{X}^0, \tilde{\mathcal{S}})$ to be the completion of $C_c^\infty(\tilde{X}^0, \tilde{\mathcal{S}})$ with respect to the Sobolev norm

$$\|v\|_1 = \left(\int_{\tilde{X}^0} |v|^2 + \int_{\tilde{X}^0} |\nabla v|^2 \right)^{1/2}. \tag{A.1}$$

Consider the Friedrichs extension F of \tilde{D}^2 on $L^2(\tilde{X}^0, \tilde{\mathcal{S}})$ with respect to the domain $H_0^1(\tilde{X}^0, \tilde{\mathcal{S}})$. Let us write

$$|\tilde{D}| := F^{1/2}. \tag{A.2}$$

Then it is known that the domain $\text{Dom}(|\tilde{D}|)$ of $|\tilde{D}|$ is precisely $H_0^1(\tilde{X}^0, \tilde{\mathcal{S}})$, cf. [29, Chapter 8, Proposition 1.10]. Note that the space $H_0^1(\tilde{X}^0, \tilde{\mathcal{S}})$ is the Sobolev H^1 space of sections that vanish on $\partial\tilde{X}$, i.e., sections that satisfy the Dirichlet boundary condition. It follows from a standard finite propagation argument that the wave operator $e^{it|\tilde{D}|}$ associated to $|\tilde{D}|$ has finite propagation, cf. [28, Chapter 2, Section 6].

Let \mathcal{F} be a fundamental domain of \tilde{X} under the Γ -action and ρ the characteristic function of \mathcal{F} . Define ρ_γ to be the γ -translation of ρ , that is,

$$\rho_\gamma(x) = \rho(\gamma^{-1}x).$$

For a given $a \in \mathbb{R}$, let us write $T = (|\tilde{D}| + ia)^{-1}$. We define

$$T_\gamma = \rho_\gamma \circ T \circ \rho. \tag{A.3}$$

In the following, we shall fix a length metric $l: \Gamma \rightarrow \mathbb{R}_{\geq 0}$ on Γ . Then there exist $A_\Gamma > 0$ and $B_\Gamma > 0$ such that

$$A_\Gamma^{-1} \cdot \text{dist}(\gamma\mathcal{F}, \mathcal{F}) - B_\Gamma \leq l(\gamma) \leq A_\Gamma \cdot \text{dist}(\gamma\mathcal{F}, \mathcal{F}) + B_\Gamma \tag{A.4}$$

for all $\gamma \in \Gamma$, where $\text{dist}(\gamma\mathcal{F}, \mathcal{F})$ is the distance between the two sets $\gamma\mathcal{F}$ and \mathcal{F} measured with respect to the given Riemannian metric on \tilde{X} .

Lemma A.2. *Let $T = (|\tilde{D}| + ia)^{-1}$ as above. Then there exists a constant $C_1 > 0$ such that*

$$\|T_\gamma\| \leq C_1 e^{-|a| \cdot A_\Gamma^{-1} \cdot l(\gamma)},$$

for all $\gamma \in \Gamma$, where $\|T_\gamma\|$ is the operator norm of the operator

$$T_\gamma: L^2(\tilde{X}^\circ, \tilde{\mathcal{S}}) \rightarrow L^2(\tilde{X}^\circ, \tilde{\mathcal{S}}).$$

Proof. If $a = 0$, the lemma is trivial. Without loss of generality, let us assume $a > 0$, since the case where $a < 0$ can be treated exactly the same way. The Fourier transform of $f(x) = (x + ia)^{-1}$ is

$$\hat{f}(\xi) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) e^{-i\xi x} dx = -i\sqrt{2\pi} e^{-a\xi} \theta(\xi)$$

where θ is the unit step function

$$\theta(\xi) = \begin{cases} 0 & \text{if } \xi < 0, \\ 1 & \text{if } \xi \geq 0. \end{cases}$$

In particular, \hat{f} and all of its derivatives are smooth away from $\xi = 0$ and decay exponentially as $|\xi| \rightarrow \infty$.

Let φ be a smooth function on \mathbb{R} with $0 \leq \varphi(x) \leq 1$ such that $\varphi(x) = 1$ for all $|x| \geq 2$ and $\varphi(x) = 0$ for all $|x| \leq 1$. For each $t > 0$, we define h_t to be the function on \mathbb{R} whose Fourier transform is

$$\hat{h}_t(\xi) = \varphi(t^{-1}\xi) \hat{f}(\xi).$$

For each fixed $t > 0$, we apply functional calculus to define the operator $R := h_t(|\tilde{D}|)$. We have

$$R(v) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \varphi(t^{-1}\xi) \hat{f}(\omega) e^{i\xi|\tilde{D}|} v d\xi$$

for all $v \in L^2(\tilde{X}^\circ, \tilde{\mathcal{S}})$. Define

$$R_\gamma = \rho_\gamma \circ R \circ \rho.$$

We see that there exists a constant $C_2 > 0$ such that

$$\|R_\gamma\| \leq |\rho_\gamma| \cdot \|R\| \cdot \|\rho\| \leq \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \varphi(t^{-1}\xi) |\hat{f}(\xi)| d\xi \leq C_2 e^{-at}$$

for all $\gamma \in \Gamma$. By finite propagation of the wave operator $e^{is|\tilde{D}|}$, it follows that

$$T_\gamma = R_\gamma$$

for all but finitely many $\gamma \in \Gamma$. More precisely, we have $T_\gamma = R_\gamma$ for all γ with $l(\gamma) \geq A_\Gamma \cdot t + B_\Gamma$. By varying t , it is not difficult to see that there exists a constant $C > 0$ such that

$$\|T_\gamma\| \leq C e^{-a \cdot A_\Gamma^{-1} \cdot l(\gamma)}$$

for all $\gamma \in \Gamma$. ■

For any given $a \neq 0 \in \mathbb{R}$, the range $\mathcal{R}(|\tilde{D}| + ia)$ of $|\tilde{D}| + ia$ is equal to the whole space $L^2(\tilde{X}^0, \tilde{\mathcal{S}})$, since $|\tilde{D}|$ is self-adjoint. In particular, for each $f \in C_c^\infty(\tilde{X}^0, \tilde{\mathcal{S}})$, there is $v = (|\tilde{D}| + ia)^{-1}(f) \in \text{Dom}(|\tilde{D}|)$ such that

$$(|\tilde{D}| + ia)v = f.$$

We define $\mathcal{L}_{C_{\max}^*(\Gamma; \mathbb{R})}^2$ to be the completion of $C_c^\infty(\tilde{X}^0, \tilde{\mathcal{S}})$ with respect to the following Hilbert $C_{\max}^*(\Gamma; \mathbb{R})$ -inner product

$$\langle\langle f_1, f_2 \rangle\rangle := \sum_{\gamma \in \Gamma} \langle f_1, \gamma f_2 \rangle \gamma \in C_{\max}^*(\Gamma; \mathbb{R}) \tag{A.5}$$

for all $f_1, f_2 \in C_c^\infty(\tilde{X}^0, \tilde{\mathcal{S}})$, where

$$\langle f_1, \gamma f_2 \rangle = \int_{\tilde{X}^0} \langle f_1(x), f_2(\gamma^{-1}x) \rangle dx.$$

Let us define

$$\|v\|_{\max} := \langle\langle v, v \rangle\rangle^{1/2}$$

for $v \in \mathcal{L}_{C_{\max}^*(\Gamma; \mathbb{R})}^2$.

The following lemma is a consequence of Lemma A.2.

Lemma A.3. *If $|a|$ is sufficiently large, then for every $f \in C_c^\infty(\tilde{X}^0, \tilde{\mathcal{S}})$, the element $v = (|\tilde{D}| + ia)^{-1}(f)$ lies in $\mathcal{L}_{C_{\max}^*(\Gamma; \mathbb{R})}^2$.*

Proof. Let $\{\rho_\gamma\}_{\gamma \in \Gamma}$ be the characteristic functions as above. We have

$$v = \sum_{\gamma \in \Gamma} \rho_\gamma v.$$

Clearly, each $\rho_\gamma v$ lies in $\mathcal{L}_{C_{\max}^*(\Gamma; \mathbb{R})}^2$, since $\rho_\gamma v$ is supported on a metric ball of bounded radius.

By Lemma A.2, a straightforward calculation shows that there exists a constant²⁴ $C_f > 0$ such that

$$\langle v, \beta v \rangle \leq C_f \cdot e^{-|a| \cdot A_\Gamma^{-1} \cdot l(\beta)} \cdot \|f\|_{L^2},$$

where $l(\beta)$ is the word length of β and the constant A_Γ^{-1} is defined in line (A.4). Since the group Γ has at most exponential growth, that is, there exist numbers $K_\Gamma > 0$ and C_2 such that

$$\#\{\alpha \in \Gamma : l(\alpha) \leq n\} \leq C_2 e^{K_\Gamma \cdot n}$$

for all $n \in \mathbb{N}$. It follows that

$$\|v\|_{\max}^2 = \langle\langle v, v \rangle\rangle = \sum_{\beta \in \Gamma} \langle v, \beta v \rangle \beta \in C_{\max}^*(\Gamma; \mathbb{R})$$

as long as $|a|$ is sufficiently large. This finishes the proof. ■

²⁴The constant C_f depends on f . More precisely, the constant C_f depends on the diameter of the support of f .

For each $a \in \mathbb{R}$ such that $|a|$ is sufficiently large, consider the operator

$$|\tilde{D}|_{\max} : \mathcal{L}^2_{C^*_{\max}(\Gamma; \mathbb{R})} \rightarrow \mathcal{L}^2_{C^*_{\max}(\Gamma; \mathbb{R})}$$

defined by setting $|\tilde{D}|_{\max}(v) = |\tilde{D}|(v)$ on the domain

$$\text{Dom}(|\tilde{D}|_{\max}) = C_c^\infty(\tilde{X}^0, \tilde{\mathcal{S}}) + (|\tilde{D}| + ia)^{-1}(C_c^\infty(\tilde{X}^0, \tilde{\mathcal{S}}))$$

where $(|\tilde{D}| + ia)^{-1}(C_c^\infty(\tilde{X}^0, \tilde{\mathcal{S}}))$ consists of

$$\{v \in \mathcal{L}^2_{C^*_{\max}(\Gamma; \mathbb{R})} : v = (|\tilde{D}| + ia)^{-1}f \text{ for some } f \in C_c^\infty(\tilde{X}^0, \tilde{\mathcal{S}})\}.$$

As an immediate consequence of Lemma A.3, we see that $|\tilde{D}|_{\max}$ is well defined. Moreover, $|\tilde{D}|_{\max}$ is an unbounded symmetric operator, since $|\tilde{D}|$ is symmetric with respect to the inner product from line (A.5).

Lemma A.4. *For each $a \in \mathbb{R}$ such that $|a|$ is sufficiently large, the closure of $|\tilde{D}|_{\max}$ is regular and self-adjoint.*

Proof. By construction, the operator $(|\tilde{D}|_{\max} + ia)$ has a dense range. By [16, Lemmas 9.7 and 9.8], we conclude that the closure of $|\tilde{D}|_{\max}$ is regular and self-adjoint. ■

We have the following main result of this section.

Proposition A.5. *Suppose X is an n -dimensional compact spin manifold with boundary or corners. Let \mathcal{S} be the Cl_n -Clifford bundle over X and D a Cl_n -linear Dirac-type operator acting on \mathcal{S} . Let \tilde{X} be a Galois Γ -covering space of X and $\tilde{\mathcal{S}}$ (resp. \tilde{D}) the lift of \mathcal{S} (resp. D). Let \mathcal{R} be the symmetric bundle endomorphism of $\tilde{\mathcal{S}}$ appearing in the following Lichnerowicz formula:*

$$\tilde{D}^2 = \nabla^* \nabla + \mathcal{R}.$$

Assume there exists $\sigma > 0$ such that

$$\mathcal{R}(x) \geq \frac{(n - 1)\sigma^2}{n}$$

for all $x \in \tilde{X}$. Then there exists a self-adjoint Friedrichs extension F_{\max} of \tilde{D}^2 ,

$$F_{\max} : \mathcal{L}^2_{C^*_{\max}(\Gamma; \mathbb{R})} \rightarrow \mathcal{L}^2_{C^*_{\max}(\Gamma; \mathbb{R})}$$

such that the following are satisfied:

- (1) $\|F_{\max}(f)\|_{\max} \geq \sigma^2 \|f\|_{\max}$ for all $f \in \text{Dom}(F_{\max})$,
- (2) and the wave operator $e^{is|\mathbf{D}|}$ has finite propagation, where $|\mathbf{D}| = (F_{\max})^{1/2}$.

Proof. Fix $a \in \mathbb{R}$ such that $|a|$ is sufficiently large. Let $|\tilde{D}|_{\max}$ be the operator from Lemma A.4. We denote its closure by $|\mathbf{D}|$. Let us define $F_{\max} = |\mathbf{D}|^2$. By construction, the wave operator $e^{is|\mathbf{D}|}$ has finite propagation, since $e^{is|\tilde{D}|}$ has finite propagation, where $|\tilde{D}|$ is the operator defined in line (A.2).

To prove the proposition, it suffices to verify

$$\| |\mathbf{D}|(f) \|_{\max} \geq \sigma \|f\|_{\max}$$

for all $v \in \text{Dom}(|\tilde{D}|_{\max}) = C_c^\infty(\tilde{X}^0, \tilde{\mathcal{S}}) + (|\tilde{D}| + ia)^{-1}(C_c^\infty(\tilde{X}^0, \tilde{\mathcal{S}}))$. Given

$$v = f_1 + (|\tilde{D}| + ia)^{-1} f_2$$

for some $f_1, f_2 \in C_c^\infty(\tilde{X}^0, \tilde{\mathcal{S}})$, we have

$$(|\tilde{D}|_{\max} + ia)v = (|\tilde{D}| + ia)f_1 + f_2,$$

which implies

$$\begin{aligned} |\tilde{D}|_{\max}(v) &= -iav + (|\tilde{D}| + ia)f_1 + f_2 \\ &= -iaf_1 - ia(|\tilde{D}| + ia)^{-1} f_2 + (|\tilde{D}| + ia)f_1 + f_2. \end{aligned}$$

It follows that for each $v \in \text{Dom}(|\tilde{D}|_{\max})$, we have $|\tilde{D}|_{\max}(v) \in \text{Dom}(|\tilde{D}|_{\max})$. We conclude that

$$\begin{aligned} \| |\mathbf{D}|(v) \|_{\max}^2 &= \langle |\tilde{D}|_{\max}(v), |\tilde{D}|_{\max}(v) \rangle \\ &= \langle |\tilde{D}|v, |\tilde{D}|v \rangle \\ &= \langle Fv, v \rangle \end{aligned}$$

for all $v \in \text{Dom}(|\tilde{D}|_{\max})$.

Note that $\text{Dom}(|\tilde{D}|_{\max}) = C_c^\infty(\tilde{X}^0, \tilde{\mathcal{S}}) + (|\tilde{D}| + ia)^{-1}(C_c^\infty(\tilde{X}^0, \tilde{\mathcal{S}}))$ is contained in $H_0^1(\tilde{X}^0, \tilde{\mathcal{S}})$, since $\text{Dom}(|\tilde{D}|) = H_0^1(\tilde{X}^0, \tilde{\mathcal{S}})$. It follows that

$$\langle Fv, v \rangle = \langle \tilde{D}^2 v, v \rangle = \langle \tilde{D}v, \tilde{D}v \rangle$$

for all $v \in \text{Dom}(|\tilde{D}|_{\max})$. By definition, we have

$$\tilde{D}v = \sum_{j=1}^n c(e_j) \nabla_{e_j} v$$

for all $v \in \text{Dom}(|\tilde{D}|_{\max})$, where the e_j are a local orthonormal basis of TX and $c(e_j)$ are the corresponding Clifford multiplication. By the Cauchy–Schwarz inequality, we have

$$\begin{aligned} \langle \tilde{D}v, \tilde{D}v \rangle &= \left\langle \sum_{j=1}^n c(e_j) \nabla_{e_j} v, \sum_{j=1}^n c(e_j) \nabla_{e_j} v \right\rangle \\ &\leq n \sum_{j=1}^n \langle c(e_j) \nabla_{e_j} v, c(e_j) \nabla_{e_j} v \rangle \end{aligned}$$

$$= n \sum_{j=1}^n \langle \nabla_{e_j} v, \nabla_{e_j} v \rangle = n \langle \nabla v, \nabla v \rangle$$

for all $v \in \text{Dom}(|\tilde{D}|_{\max})$. Since

$$\langle \tilde{D}v, \tilde{D}v \rangle = \langle \tilde{D}^2v, v \rangle = \langle \nabla^* \nabla v, v \rangle + \langle \mathcal{R}v, v \rangle$$

for all $v \in \text{Dom}(|\tilde{D}|_{\max})$, it follows that

$$\frac{n-1}{n} \langle \tilde{D}v, \tilde{D}v \rangle \geq \langle \mathcal{R}v, v \rangle \geq \frac{(n-1)\sigma^2}{n} \langle v, v \rangle, \tag{A.6}$$

for all $v \in \text{Dom}(|\tilde{D}|_{\max})$. To summarize, we have showed that

$$\| |\mathbf{D}|(v) \|_{\max}^2 = \langle Fv, v \rangle \geq \sigma^2 \langle v, v \rangle$$

for all $v \in \text{Dom}(|\tilde{D}|_{\max})$. This finishes the proof. ■

B. An estimate of the universal constant C in Theorem I (by Jinmin Wang and Zhizhang Xie)

In this appendix, we give a numerical estimate of the universal constant C that appeared in Theorem I. Part of the numerical computation is done with the assistance of MATLAB. We would like to thank Li Zhou for many helpful comments on the MATLAB programming. Throughout this section, we use the following convention of the Fourier transform

$$\hat{f}(\xi) = \int_{\mathbb{R}} f(x) e^{-ix\xi} dx$$

and the inverse Fourier transform

$$f(x) = \frac{1}{2\pi} \int_{\mathbb{R}} \hat{f}(\xi) e^{ix\xi} d\xi.$$

B.1. Increasing normalizing functions

Let $\chi: \mathbb{R} \rightarrow [-1, 1]$ be a normalizing function, that is, χ a continuous odd function such that $\lim_{x \rightarrow \pm\infty} \chi(x) = \pm 1$. In addition, let us assume χ is an increasing function in this subsection. This additional assumption is not necessary, but it makes the numerical estimates somewhat easier. The method in this subsection is inspired by the work of Slepian [27]. We will discuss general normalizing functions in the next subsection and carry out some estimates via a different method.

Note that given $\varepsilon > 0$, there exists $\sigma > 0$ such that

$$\sup_{|x| \geq \sigma} |1 - \chi(x)^2| < \varepsilon.$$

Here σ has the same meaning as the spectral gap σ of D on a subspace $Y \setminus K$ in Lemma 3.1 and Lemma 3.2. The key step for estimating the universal constant C from Theorem I is to answer the following minimization question.²⁵

Question B.1. *Given $\varepsilon > 0$, find the minimum value of σ among all normalizing functions χ whose Fourier transform $\hat{\chi}$ is supported on $[-2, 2]$.*

Let us set

$$f = \frac{\chi'}{2},$$

where χ' is the derivative of χ . Here the factor $1/2$ is only introduced as a normalizing constant. Since χ is assumed to be increasing, f is a positive even function such that its Fourier transform \hat{f} is supported on $[-2, 2]$ and $\hat{f}(0) = 1$. As another simplification,²⁶ let us only consider f such that $f = h^2$, where h is a function with its Fourier transform \hat{h} supported on $[-1, 1]$. Since $f = h^2$, we have $\hat{f} = \hat{h} * \hat{h}$, i.e.,

$$\hat{f}(\xi) = \int_{\mathbb{R}} \hat{h}(\xi - \eta)\hat{h}(\eta) \, d\eta.$$

It follows that

$$\hat{f}(0) = \int_{\mathbb{R}} \hat{h}(-\eta)\hat{h}(\eta) \, d\eta = \|h\|_{L^2}^2,$$

where the second equality uses the fact that h is an even function.

Observe that the minimization question (Question B.1) is essentially equivalent to the following maximization question.

Question B.2. *Given $\sigma > 0$, find the maximal value of*

$$\int_{-\sigma}^{\sigma} h^2(x) \, dx$$

among all functions $h: \mathbb{R} \rightarrow \mathbb{R}$ with $\|h\|_{L^2} = 1$ such that its Fourier transform is supported on $[-1, 1]$.

Let $c_{\sigma}(x)$ be the characteristic function of $[-\sigma, \sigma]$, that is,

$$c_{\sigma}(x) = \begin{cases} 1 & \text{if } |x| \leq \sigma, \\ 0 & \text{if } |x| > \sigma. \end{cases}$$

Then the Fourier transform of c_{σ} is given by

$$\hat{c}_{\sigma}(\xi) = \frac{2 \sin(\sigma\xi)}{\xi}.$$

²⁵Alternatively, we can fix the value of the spectral gap σ and try to minimize the propagation, that is, the support of the Fourier transform $\hat{\chi}$. The two approaches are essentially equivalent.

²⁶Such an simplification does not really affect the final estimates at the end of this subsection.

Note that

$$\int_{-\sigma}^{\sigma} f(x) \, dx = \int_{\mathbb{R}} f(x)c_{\sigma}(x) \, dx = \frac{1}{2\pi} \int_{\mathbb{R}} \hat{f}(\xi)\hat{c}_{\sigma}(\xi) \, d\xi = \int_{\mathbb{R}} \frac{\sin(\sigma\xi)}{\pi\xi} \hat{f}(\xi) \, d\xi.$$

Recall that $f = h^2$. It follows that

$$\int_{-\sigma}^{\sigma} f(x) \, dx = \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{\sin(\sigma\xi)}{\pi\xi} \hat{h}(\xi - \eta)\hat{h}(\eta) \, d\eta \, d\xi.$$

We conclude that

$$\int_{-\sigma}^{\sigma} h^2(s) \, ds = \int_{-1}^1 \int_{-1}^1 \frac{\sin(\sigma(x + y))}{\pi(x + y)} \hat{h}(x)\hat{h}(y) \, dx \, dy,$$

since \hat{h} is supported on $[-1, 1]$.

This naturally leads us to the integral operator T_{σ} on $L^2([-1, 1])$ given by

$$T_{\sigma}(\varphi)(x) = \int_{-1}^1 \frac{\sin(\sigma(x + y))}{\pi(x + y)} \varphi(y) \, dy$$

for all $\varphi \in L^2([-1, 1])$. Observe that T_{σ} is a bounded self-adjoint compact operator. Hence the maximum value of

$$\int_{-\sigma}^{\sigma} h^2$$

is equal to the largest (absolute) eigenvalue of T_{σ} , which is precisely the operator norm $\|T_{\sigma}\|$ of T_{σ} . Moreover, the maximum is achieved when h is a corresponding eigen-function of the largest (absolute) eigenvalue of T_{σ} . This eigen-function can be numerically estimated by considering the following iteration

$$\varphi_n = \|T_{\sigma}\varphi_{n-1}\|_{L^2}^{-1} \cdot T_{\sigma}(\varphi_{n-1}),$$

with $\varphi_0 \equiv 1$. In particular, φ_n converges to an eigen-function of the largest (absolute) eigenvalue of T_{σ} .

The detailed estimates can be carried out as follows.

- (1) Given any normalizing function $\chi: \mathbb{R} \rightarrow [-1, 1]$, let p_t be the idempotent from line (3.1). By Lemma 3.3, we need to estimate the operator norm $\|p_t\|$ of p_t . It amounts to estimating the operator norm of the following idempotents:

$$\mathbb{P}_a = \begin{pmatrix} a^2(2 - a^2) & (2 - a^2)(1 - a^2)a \\ a(1 - a^2) & (1 - a^2)^2 \end{pmatrix},$$

for all $a \in [-1, 1]$. A numerical estimate shows that

$$\sup_{a \in [-1, 1]} \|\mathbb{P}_a\|^2 = \sup_{a \in [-1, 1]} \{\text{eigenvalues of } \mathbb{P}_a^* \mathbb{P}_a\} \approx (1.04015)^2.$$

We conclude that

$$\|p_t\| \approx 1.04015. \tag{B.1}$$

(2) Again, by Lemma 3.3 and part (3) of Lemma 3.2, we need to estimate the operator norm

$$\left\| p_t - \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \right\|$$

on some subspace of Y . This amounts to finding $\theta > 0$ such that

$$\begin{aligned} \left\| \mathbb{P}_a - \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \right\| &= \left\| \begin{pmatrix} -(1-a^2) & (2-a^2)(1-a^2)a \\ a(1-a^2) & (1-a^2)^2 \end{pmatrix} \right\| \\ &\leq \frac{1}{(2\|\mathbb{P}_a\| + 2)} \frac{1}{4} \leq \frac{1}{(2(1.04015) + 2)} \frac{1}{4} = \frac{1}{16.3212}, \end{aligned} \tag{B.2}$$

for all a with $\theta \leq |a| \leq 1$. A numerical estimate shows that

$$\theta \approx 0.96978. \tag{B.3}$$

(3) Now the last step is to find σ such that

$$\|T_\sigma\| = \theta \approx 0.96978.$$

Again a numerical estimate shows that

$$\sigma \approx 2.86821.$$

Now let $\chi: \mathbb{R} \rightarrow [-1, 1]$ be a normalizing function that fulfills the above estimates. Since the Fourier transform $\hat{\chi}$ is supported on $[-2, 2]$, it follows from the proof of Theorem I (cf. Theorem 3.4) that we can choose $r = 2 \cdot 15 = 30$. It follows that the universal constant C in Theorem I satisfies

$$C \leq 2.86821 \cdot 2 \cdot 15 = 86.0463.$$

B.2. An improved estimate

In this subsection, we shall improve our estimate for the universal constant C by considering general normalizing functions. That is, the normalizing function χ is *not* assumed to be increasing any more.

Let sgn be the sign function, that is,

$$\text{sgn}(x) = \begin{cases} 1 & \text{if } x \geq 0, \\ -1 & \text{if } x < 0. \end{cases}$$

Its Fourier transform is given by

$$\widehat{\text{sgn}}(\xi) = \int_{\mathbb{R}} \text{sgn}(x) e^{-ix\xi} dx = \frac{2}{i\xi}.$$

Our goal is to solve the following analogue of Question B.1.

Question B.3. For each function f , let us define

$$\chi_f(x) = \frac{1}{2\pi} \int_{\mathbb{R}} f(\xi) \frac{2}{i\xi} e^{ix\xi} d\xi.$$

For given $\varepsilon_1 > 0$ and $\varepsilon_2 > 0$, find the minimum value of σ such that there exists an even function f satisfying the following conditions:

- (1) f is supported on $[-2, 2]$,
- (2) $f(0) = 1$,
- (3) $\varepsilon_1 < \chi_f(x) - 1 < \varepsilon_2$ for all $x \geq \sigma$,
- (4) and $-\varepsilon_2 < \chi_f(x) + 1 < -\varepsilon_1$ for all $x \leq -\sigma$.

Since f is an even function, it follows that χ_f is an odd function. Furthermore, we have

$$\lim_{x \rightarrow \infty} \chi_f(x) = \int_0^\infty \chi'_f(t) dt = \frac{\widehat{\chi'_f}(0)}{2} = f(0) = 1.$$

It follows that, under the above assumptions on f , the function χ_f is a normalizing function.

As a simplification, we shall only consider even functions f such that the function χ_f takes values in $[-1.2, 1.2]$. Our numerical estimates show that such a simplification essentially does not result in any loss of generality. Under the extra assumption that χ_f takes values in $[-1.2, 1.2]$, the norm estimate in line (B.1) will remain the same. Then, based on the estimate in line (B.2), we can choose

$$\varepsilon_1 \approx 1 - 0.96978 = 0.03022,$$

and

$$\varepsilon_2 \approx 1.02928 - 1 = 0.02928.$$

Now, to approximate the minimum value of σ in Question B.3, we consider the function f of the form

$$f(\xi) = \sum_{k=0}^n a_k \cos\left(\frac{k\pi\xi}{2}\right)$$

with real numbers $\{a_k\}_{0 \leq k \leq n}$ to be determined. Note that

$$\varphi_k(x) := \frac{1}{2\pi} \int_{\mathbb{R}} \cos\left(\frac{k\pi\xi}{2}\right) \frac{2}{i\xi} e^{ix\xi} d\xi = \frac{1}{2\pi} \int_{2\pi k-2x}^{2\pi k+2x} \frac{2 \sin t}{t} dt.$$

So we can write

$$\chi_f(x) = \sum_{k=0}^n a_k \varphi_k(x). \tag{B.4}$$

We see that Condition (2) in Question B.3 becomes the following linear inequality:

$$0.03022 < \sum_{k=0}^n a_k \varphi_k(x) - 1 < 0.02928$$

for each $x \geq \sigma$. This reduces Question B.3 into solving a system of linear inequalities.

If we set $n = 5$, then a numerical estimate shows that the function

$$\begin{aligned} f(\xi) \approx & 0.75382052 + 0.25425247 \cos\left(\frac{\pi\xi}{2}\right) + 0.0034679636 \cos(\pi\xi) \\ & - 0.026352193 \cos\left(\frac{3\pi\xi}{2}\right) + 0.024841712 \cos(2\pi\xi) \\ & - 0.010030481 \cos\left(\frac{5\pi\xi}{2}\right) \end{aligned} \tag{B.5}$$

satisfies the conditions in Question B.3, and we can choose

$$\sigma \approx 1.41356$$

in this case. See Figure 3 for the graph of the corresponding normalizing function χ_f . Further numerical estimates show that we can choose

$$\sigma \approx 1.3633 \quad \text{if we work with } n = 20,$$

and

$$\sigma \approx 1.355 \quad \text{if we work with } n = 50.$$

This suggests that 1.355 is rather close to the actual minimum value of σ in Question B.3. In any case, we conclude the universal constant C in Theorem I satisfies

$$C \leq 1.355 \cdot 2 \cdot 15 = 40.65.$$




Figure 3. The graph of χ_f when f is the function in line (B.5).

Acknowledgments. The author wants to thank Dean Baskin, Alex Engel, Nigel Higson, Yanli Song, Xiang Tang, Jinmin Wang, Guoliang Yu and Rudolf Zeidler for many stimulating discussions over the years. In particular, the author would like to thank Alex Engel, Jinmin Wang, Guoliang Yu and Rudolf Zeidler for helpful comments regarding some technical issues in an earlier version of the paper. The author also wants to thank the referees for constructive suggestions.

Funding. Partially supported by NSF 1800737 and NSF 1952693.

References

- [1] M. F. Atiyah, V. K. Patodi, and I. M. Singer, Spectral asymmetry and Riemannian geometry. III. *Math. Proc. Cambridge Philos. Soc.*
- [2] M. F. Atiyah and I. M. Singer, The index of elliptic operators. III. *Ann. of Math. (2)* **87** (1968), 546–604 Zbl [0164.24301](#) MR [236952](#)
- [3] U. Bunke, A K -theoretic relative index theorem and Callias-type Dirac operators. *Math. Ann.* **303** (1995), no. 2, 241–279 Zbl [0835.58035](#) MR [1348799](#)
- [4] S. Cecchini, A long neck principle for Riemannian spin manifolds with positive scalar curvature. *Geom. Funct. Anal.* **30** (2020), no. 5, 1183–1223 Zbl [1455.58008](#) MR [4181824](#)
- [5] S. Cecchini and R. Zeidler, Scalar and mean curvature comparison via the Dirac operator. 2021, arXiv:[2103.06833](#)
- [6] O. Chodosh and C. Li, Generalized soap bubbles and the topology of manifolds with positive scalar curvature. 2020, arXiv:[2008.11888](#)
- [7] A. Connes and H. Moscovici, Cyclic cohomology, the Novikov conjecture and hyperbolic groups. *Topology* **29** (1990), no. 3, 345–388 Zbl [0759.58047](#) MR [1066176](#)
- [8] R. E. Greene and H. Wu, C^∞ approximations of convex, subharmonic, and plurisubharmonic functions. *Ann. Sci. École Norm. Sup. (4)* **12** (1979), no. 1, 47–84 Zbl [0415.31001](#) MR [532376](#)
- [9] M. Gromov, Metric inequalities with scalar curvature. *Geom. Funct. Anal.* **28** (2018), no. 3, 645–726 Zbl [1396.53068](#) MR [3816521](#)
- [10] M. Gromov, Four lectures on scalar curvature. In *Perspectives on Scalar Curvature*, edited by M. L. Gromov and H. B. Lawson, Jr., pp. 1–514, World Scientific Publishing Co. Pte. Ltd., 2023
- [11] M. Gromov, No metrics with positive scalar curvatures on aspherical 5-manifolds. 2020, arXiv:[2009.05332](#)
- [12] E. Guentner, N. Higson, and S. Weinberger, The Novikov conjecture for linear groups. *Publ. Math. Inst. Hautes Études Sci.* (2005), no. 101, 243–268 Zbl [1073.19003](#) MR [2217050](#)
- [13] H. Guo, Z. Xie, and G. Yu, Quantitative K -theory, positive scalar curvature, and band width. In *Perspectives on Scalar Curvature*, edited by M. L. Gromov and H. B. Lawson, Jr., pp. 763–798, World Scientific Publishing Co. Pte. Ltd., 2023
- [14] G. Kasparov and G. Yu, The coarse geometric Novikov conjecture and uniform convexity. *Adv. Math.* **206** (2006), no. 1, 1–56 Zbl [1102.19003](#) MR [2261750](#)
- [15] G. G. Kasparov, Equivariant KK -theory and the Novikov conjecture. *Invent. Math.* **91** (1988), no. 1, 147–201 Zbl [0647.46053](#) MR [918241](#)

- [16] E. C. Lance, *Hilbert C^* -modules*. Lond. Math. Soc. Lect. Note Ser. 210, Cambridge University Press, Cambridge, 1995 MR [1325694](#)
- [17] H. B. Lawson, Jr. and M.-L. Michelsohn, *Spin geometry*. Princeton Math. Ser. 38, Princeton University Press, Princeton, NJ, 1989 Zbl [0688.57001](#) MR [1031992](#)
- [18] C. Li, A polyhedron comparison theorem for 3-manifolds with positive scalar curvature. *Invent. Math.* **219** (2020), no. 1, 1–37 Zbl [1440.53049](#) MR [4050100](#)
- [19] M. Llarull, Sharp estimates and the Dirac operator. *Math. Ann.* **310** (1998), no. 1, 55–71 Zbl [0895.53037](#) MR [1600027](#)
- [20] J. Lohkamp, Minimal smoothings of area minimizing cones. 2018, arXiv:[1810.03157](#)
- [21] J. Lott, Index theory for scalar curvature on manifolds with boundary. *Proc. Amer. Math. Soc.* **149** (2021), no. 10, 4451–4459 Zbl [1477.53067](#) MR [4305995](#)
- [22] J. Roe, Positive curvature, partial vanishing theorems and coarse indices. *Proc. Edinb. Math. Soc. (2)* **59** (2016), no. 1, 223–233 Zbl [1335.58017](#) MR [3439130](#)
- [23] J. Rosenberg and S. Stolz, Metrics of positive scalar curvature and connections with surgery. In *Surveys on surgery theory, Vol. 2*, pp. 353–386, Ann. of Math. Stud. 149, Princeton University Press, Princeton, NJ, 2001 Zbl [0971.57003](#) MR [1818778](#)
- [24] R. Schoen and S. T. Yau, Existence of incompressible minimal surfaces and the topology of three-dimensional manifolds with nonnegative scalar curvature. *Ann. of Math. (2)* **110** (1979), no. 1, 127–142 Zbl [0431.53051](#) MR [541332](#)
- [25] R. Schoen and S. T. Yau, On the structure of manifolds with positive scalar curvature. *Manuscripta Math.* **28** (1979), no. 1-3, 159–183 Zbl [0423.53032](#) MR [535700](#)
- [26] R. Schoen and S.-T. Yau, Positive scalar curvature and minimal hypersurface singularities. In *Surveys in differential geometry 2019. Differential geometry, Calabi–Yau theory, and general relativity. Part 2*, pp. 441–480, Surv. Differ. Geom. 24, International Press, Boston, MA, 2022 MR [4479726](#)
- [27] D. Slepian, Some comments on Fourier analysis, uncertainty and modeling. *SIAM Rev.* **25** (1983), no. 3, 379–393 Zbl [0571.94004](#) MR [710468](#)
- [28] M. E. Taylor, *Partial differential equations I. Basic theory*. Second edn., Appl. Math. Sci. 115, Springer, New York, 2011 Zbl [1206.35002](#) MR [2744150](#)
- [29] M. E. Taylor, *Partial differential equations II. Qualitative studies of linear equations*. Second edn., Appl. Math. Sci. 116, Springer, New York, 2011 Zbl [1206.35003](#) MR [2743652](#)
- [30] J. Wang, Z. Xie, and G. Yu, A proof of Gromov's cube inequality on scalar curvature. 2021, arXiv:[2105.12054](#)
- [31] J. Wang, Z. Xie, and G. Yu, Decay of scalar curvature on uniformly contractible manifolds with finite asymptotic dimension. 2021, arXiv:[2101.11584](#), to appear in *Commun. Pure Appl. Math.*
- [32] R. Willett and G. Yu, *Higher index theory*. Camb. Stud. Adv. Math. 189, Cambridge University Press, Cambridge, 2020 Zbl [1471.19001](#) MR [4411373](#)
- [33] Z. Xie and G. Yu, A relative higher index theorem, diffeomorphisms and positive scalar curvature. *Adv. Math.* **250** (2014), 35–73 Zbl [1306.58006](#) MR [3122162](#)
- [34] G. Yu, The coarse Baum–Connes conjecture for spaces which admit a uniform embedding into Hilbert space. *Invent. Math.* **139** (2000), no. 1, 201–240 Zbl [0956.19004](#) MR [1728880](#)
- [35] G. Yu, A characterization of the image of the Baum–Connes map. In *Quanta of maths*, pp. 649–657, Clay Math. Proc. 11, American Mathematical Society, Providence, RI, 2010 Zbl [1216.19008](#) MR [2732068](#)
- [36] R. Zeidler, Width, largeness and index theory. *SIGMA Symmetry Integrability Geom. Methods Appl.* **16** (2020), article no. 127 Zbl [1455.19005](#) MR [4181525](#)

- [37] R. Zeidler, Band width estimates via the Dirac operator. *J. Differential Geom.* **122** (2022), no. 1, 155–183 Zbl [07632811](#) MR [4507473](#)

Received 7 November 2021; revised 22 December 2021.

Zhizhang Xie

Department of Mathematics, Mailstop 3368, Texas A&M University, College Station, TX 77843-3368, USA; xie@math.tamu.edu

Jinmin Wang

Department of Mathematics, Mailstop 3368, Texas A&M University, College Station, TX 77843-3368, USA; jinmin@tamu.edu