Cycles in graphs with geometric property (T)

Jeroen Winkel

Abstract. We show that a sequence of graphs with uniformly bounded vertex degrees, number of vertices going to infinity, and with geometric property (T) has many small cycles. We also show that when a small part of such a sequence of graphs with geometric property (T) is changed, it still has geometric property (T), provided that it is still an expander. We use this to give an example of a sequence of graphs with geometric property (T) that has large cycle-free balls.

1. Introduction

In this paper, we are interested in sequences (X_n) of finite connected graphs. All graphs we consider are simple and undirected. The vertex sets will be denoted by $V(X_n)$ and the edge sets by $E(X_n)$. We will always assume that $\lim_{n\to\infty} |V(X_n)| = \infty$ and that the sequence has *uniformly bounded degree*, i.e., there is a constant d such that for all n, all vertices of X_n have degree at most d.

Let X be a finite graph. For vertices x, y, we write $x \sim y$ if (x, y) is an edge, and we write deg(x) for the degree of x. The *Laplacian* of X is the matrix Δ_X , whose rows and columns are indexed by the vertices of X, defined by

$$(\Delta_X)_{xy} = \begin{cases} \deg(x) & \text{if } x = y, \\ -1 & \text{if } x \sim y, \\ 0 & \text{else.} \end{cases}$$

It is a symmetric positive semi-definite matrix, and it has 0 as an eigenvalue. This is a simple eigenvalue if and only if X is connected.

Consider a sequence $X = (X_n)$ of finite connected graphs with uniformly bounded degree and number of vertices going to infinity. The sequence is an *expander sequence* if there is a constant h > 0 such that for all n, all positive eigenvalues of Δ_{X_n} are at least h. Equivalently, we can directly look at the operator

$$\Delta_X = \bigoplus_n \Delta_{X_n} \subseteq B(\ell^2 X).$$

²⁰²⁰ Mathematics Subject Classification. Primary 05C38; Secondary 47L55, 51F30.

Keywords. Geometric property (T), girth, cycles, Laplacian, combinatorial cost.

Here and in the following, $\ell^2 X$ denotes the Hilbert space generated by the vertices of X_1, X_2, \ldots as basis vectors. Then the sequence is an expander sequence if and only if there is h > 0 such that $\sigma(\Delta_X) \subseteq \{0\} \cup [h, \infty)$.

There is an equivalent combinatorial characterisation of expanders. For a subset of the vertices $A \subseteq V(X_n)$, let δA denote the set of edges with exactly one vertex in A. Then the sequence (X_n) is an expander if and only if it has uniformly bounded degree and there is c > 0 such that for all $A \subset V(X_n)$ with $|A| \le \frac{1}{2}|V(X)|$, we have $|\delta A| \ge c|A|$. We refer to, e.g., [7] for more details on expanders.

In [11], Willett and Yu introduced geometric property (T). It was studied in more depth by the same authors in [12]. It is a stronger property than being an expander, based on spectral gap of Δ_X in a larger algebra.

Let (X_n) be a sequence of graphs, and let T be a bounded operator in $\prod_n B(\ell^2 X_n) \subseteq B(\ell^2 X)$. For $x, y \in V(X_n)$, we denote by d(x, y) the shortest-path distance from x to y. We can view T as a matrix whose rows and columns are indexed by the vertices of X, and $T_{xy} = 0$ if $x \in X_n$, $y \in X_m$, $m \neq n$. The *propagation* of T is $\sup\{d(x, y) \mid T_{xy} \neq 0\}$, which may be infinite. The sum and product of operators with finite propagation have finite propagation again. The *algebraic uniform Roe algebra*, as introduced by Roe in [9], is defined as

$$\mathbb{C}_{\mathrm{u}}[X] = \Big\{ T \in \prod_{n} B(\ell^2 X_n) \mid T \text{ has finite propagation} \Big\}.$$

The algebraic uniform Roe algebra is a complex unital *-algebra. A representation of $\mathbb{C}_{u}[X]$ is given by a Hilbert space \mathcal{H} and a unital *-homomorphism $\pi: \mathbb{C}_{u}[X] \to B(\mathcal{H})$.

The completion of $\mathbb{C}_{u}[X]$ in the norm inherited by $B(\ell^{2}X)$ is the *reduced uniform Roe algebra* $C_{red}^{*}(X)$. In this paper, we are more interested in the completion with respect to a larger norm, namely the maximal norm $\|\cdot\|_{max}$ given by

$$\|T_{\max}\| = \sup_{(\pi,\mathcal{H})} \|\pi(T)\|,$$

where the supremum is taken over all representations (π, \mathcal{H}) . If X has uniformly bounded degree, then the maximal norm is necessarily finite (see, for example, [3, Section 3]). It is then easy to check that it is indeed a norm. Its completion is a C^* -algebra, the maximal uniform Roe algebra $C^*_{\max}(X)$ (see, for example, [10]).

For $T \in \mathbb{C}_{u}[X]$, we denote by $\sigma(T)$ the spectrum in $C^*_{red}(X)$, while $\sigma_{max}(T)$ denotes the, possibly larger, spectrum in $C^*_{max}(X)$. Recall that X is an expander sequence if and only if there is h > 0 such that $\sigma(\Delta_X) \subseteq \{0\} \cup [h, \infty)$.

Definition 1.1 ([11]). Let $X = (X_n)$ be a sequence of finite connected graphs with uniformly bounded degree and number of vertices going to infinity. Then X has geometric property (T) if there is $\gamma > 0$ such that $\sigma_{\max}(\Delta_X) \subseteq \{0\} \cup [\gamma, \infty)$.

A sequence of finite graphs (X_n) has *large girth* if for every R > 0, there is a positive integer N such that for $n \ge N$, the graphs X_n do not have any cycles of length shorter

than *R*. It was shown in [11, Corollary 7.5] that a sequence of graphs with property (T) can never have large girth, using properties of K-theory. We give a quantitative version of this result: there is some *R* such that all of the X_n have "many" *R*-cycles.

Let us make this more precise. As in [1], we define the cycle spaces of a graph.

Definition 1.2. Let *X* be a graph with edge set *E*. Let $\mathbb{C}[E]$ be the free \mathbb{R} -vector space generated by *E*. We use the convention that (x, y) = -(y, x). Let Z(X) be the subspace of $\mathbb{C}[E]$ generated by the cycles, where a cycle (x_1, x_2, \ldots, x_n) corresponds to the element $(x_1, x_2) + (x_2, x_3) + \cdots + (x_{n-1}, x_n) + (x_n, x_1)$. For any integer *R*, let $Z_R(X)$ be the subspace generated by the cycles of length at most *R*.

We can now state the first theorem of this paper.

Theorem A. Suppose $X = (X_n)$ is a sequence of finite connected graphs with uniformly bounded degree and with number of vertices going to infinity. Suppose that X has geometric property (T). Then there are constants R > 0 and $\varepsilon > 0$ such that for all large enough n, we have dim $Z_R(X_n) \ge \varepsilon |V(X_n)|$.

The *cost* of generating an equivalence relation is introduced by Levitt in [6]. Taking the supremum of all equivalence relations generated by probability measure preserving actions of a group gives the cost of this group, as introduced by Gaboriau in [2]. *Combinatorial cost* is a variant on this concept for sequences of graphs, which was defined by Elek in [1]. It measures the number of edges necessary to induce the coarse structure of a sequence of graphs.

Definition 1.3. Let (X_n) and (Y_n) be sequences of finite connected graphs such that X_n and Y_n have the same vertex set. For $x, y \in V(X_n)$, denote by $d_X(x, y)$ the path distance in X_n , and by $d_Y(x, y)$ the path distance in Y_n . We say that (X_n) and (Y_n) *induce the same coarse structure* if there is a constant L such that $d_X(x, y) \leq Ld_Y(x, y)$ and $d_Y(x, y) \leq Ld_X(x, y)$ for all n and $x, y \in V(X_n)$.

Definition 1.4 ([1]). The *combinatorial cost* of a sequence $X = (X_n)$ of graphs is defined as

$$c(X) = \inf_{Y} \liminf_{n} \frac{|E(Y_n)|}{|V(X_n)|},$$

where the infimum is taken over all sequences of graphs (Y_n) on the same vertex sets as X_n , which induce the same coarse structure as (X_n) .

In [4], it was shown that any group with property (T) has cost 1. This raises the question if a similar theorem can be proved in the combinatorial context. From Theorem A, it already follows that the infimum in Definition 1.4 is not attained for sequences of graphs with geometric property (T) (see Proposition 2.7). The question, whether sequences of graphs with geometric property (T) necessarily have cost 1, remains open.

Theorem A also raises the following question: is it true that for every sequence (X_n) with geometric property (T), there is an R such that all vertices have a cycle in their

R-neighbourhood? We show that if two graph sequences (X_n) and (Y_n) "mostly" agree, and one of them has geometric property (T) and the other one is an expander, then the other one has geometric property (T) too. This can be used to answer the above question negatively (see Corollary 3.2).

Definition 1.5. Let $X = (X_n)$ and $Y = (Y_n)$ be sequences of finite connected graphs with increasing number of vertices and uniformly bounded degree. We say that X and Y are *approximately isomorphic* if there are subgraphs $X'_n \subseteq X_n$ and $Y'_n \subseteq Y_n$ such that X'_n and Y'_n are isomorphic for all n, and

$$\lim_{n \to \infty} \frac{|V(X'_n)|}{|V(X_n)|} = \lim_{n \to \infty} \frac{|E(X'_n)|}{|E(X_n)|} = \lim_{n \to \infty} \frac{|V(Y'_n)|}{|V(Y_n)|} = \lim_{n \to \infty} \frac{|E(Y'_n)|}{|E(Y_n)|} = 1.$$

It is easy to see that we may take X'_n and Y'_n to be induced subgraphs (that is, subgraphs that contain all edges between their vertices that are present in the ambient graph).

If X and Y are approximately isomorphic and X has geometric property (T), then Y need not have geometric property (T); indeed, it does not even need to be an expander (it is only an asymptotic expander, as defined in [5]). For example, if (X_n) is a sequence of graphs with geometric property (T) with $|X_n| \ge n^2$, and we attach a path of length *n* to each X_n , then it will not be an expander anymore. However, this is the only thing that can go wrong.

Theorem B. Let $X = (X_n)$ and $Y = (Y_n)$ be approximately isomorphic sequences of finite connected graphs with uniformly bounded degree and number of vertices going to infinity. Suppose X has geometric property (T) and Y is an expander. Then Y has geometric property (T).

2. Short-cycle spaces in graphs with geometric property (T)

Let (X_n) be a sequence of graphs. The algebraic Roe algebra $\mathbb{C}_u[X]$, as recalled in Section 1, has a standard representation on $\ell^2 X$. Other representations are harder to describe explicitly. We will see how to construct a representation of $\mathbb{C}_u[X]$, starting from a representation of only part of the matrix algebra $B(\ell^2 X_n)$, for each n.

Definition 2.1. Let X be a graph, and let $B_R(\ell^2 X)$ be the set of bounded operators on $\ell^2 X$ of propagation at most R. An *R*-representation is a Hilbert space \mathcal{H} together with a linear map $\pi: B_R(\ell^2 X) \to B(\mathcal{H})$ satisfying $\pi(T^*) = \pi(T)^*$ and $\pi(TS) = \pi(T)\pi(S)$ if *T*, *S* and *TS* have propagation at most *R*.

Suppose $X = (X_n)$ is a sequence of graphs, and suppose we have a sequence of r_n representations π_n of X_n on \mathcal{H}_n . If $r_n \to \infty$, we can make a representation of $\mathbb{C}_u[X]$ as
follows: let \mathcal{U} be a free ultrafilter on \mathbb{N} , and define the ultraproduct $\mathcal{H} = \prod_n \mathcal{H}_n/\mathcal{U}$.
Define $\pi(T) = \lim_{\mathcal{U}} \pi_n(T|_{\ell^2 X_n}) \in B(\mathcal{H})$. Then $\pi: \mathbb{C}_u[X] \to B(\mathcal{H})$ is a representation.

We can use these representations to prove that a sequence of graphs with property (T) does not have large girth (which was also already proved in [11] using K-theory). Below, we first give the proof for d-regular graphs with an even integer d, to avoid technicalities. It is true in greater generality; this will follow later from Theorem A.

Proposition 2.2. Let $X = (X_n)$ be a sequence of *d*-regular finite connected graphs and an increasing number of vertices. Suppose that X has geometric property (T) and that *d* is even. Then X does not have large girth.

Proof. Suppose that X has large girth and let $t \in \mathbb{R}$. Let r_n denote the girth of X_n . There is an Eulerian cycle on X_n . This defines a direction on each edge of X_n . So we can define a function $\rho: E_n \to \{-1, 1\}$, where E_n denotes the edge set of X_n such that for each $x \in X_n$, we have $\sum_{y \sim x} \rho(x, y) = 0$. For all pairs $(x, y) \in X_n^2$, choose a shortest path $\gamma_{(x,y)} = (x = x_0, x_1, \dots, x_{d(x,y)} = y)$ from x to y. Choose it in such a way that $\gamma_{(y,x)}$ is the inverse of $\gamma_{(x,y)}$. Define $\rho: X_n^2 \to \mathbb{Z}$ by

$$\rho(x, y) = \rho(x_0, x_1) + \dots + \rho(x_{d(x, y)-1}, x_{d(x, y)})$$

Then $\rho(x, y) = -\rho(y, x)$, and if $d(x, y) + d(y, z) + d(x, z) < r_n$, we have $\rho(x, y) + \rho(y, z) = \rho(x, z)$. Now define $\pi_n: B(\ell^2 X_n) \to B(\ell^2 X_n)$ by

$$\pi_n(T)\xi(x) = \sum_y T_{xy} \exp(it\rho(x, y))\xi(y).$$

Then $\pi_n(T^*) = \pi_n(T)^*$, and if the propagations of *S* and *T* and *TS* are at most $\frac{1}{3}r_n$, then $\pi_n(TS) = \pi_n(T)\pi_n(S)$. So π_n defines a $\frac{1}{3}r_n$ -representation.

Consider the constant unit vector $\xi_n \in \ell^2 X_n$ with $\xi_n(x) = \frac{1}{\sqrt{|V(X_n)|}}$. This satisfies

$$\pi_n(\Delta_X)\xi_n = d(1 - \cos t)\xi_n.$$

Let $\pi: \mathbb{C}_u[X] \to B(\mathcal{H})$ denote the limit of the π_n . It is a representation. Let $\xi = (\xi_n) \in \mathcal{H}$. Then $\pi(\Delta_X)\xi = d(1 - \cos t)\xi$, so $d(1 - \cos t) \in \sigma_{\max}(\Delta_X)$. Since *t* can be any real number, we conclude that $[0, 2d] \subseteq \sigma_{\max}(\Delta_X)$. Then *X* does not have geometric property (T), and we have arrived at a contradiction.

If the graph is not *d*-regular for an even *d*, we do not necessarily have an Eulerian cycle. In this case, we can still prove the result. We need Lemmas 2.3 and 2.4 below to construct a cycle $v \in Z(X)$ that will play a similar role as the Eulerian cycle above. We will use it to prove something even stronger, namely that there are "many" small cycles in X (Theorem A).

Note that, if X is a finite connected, then dim($\mathbb{C}[E]$) = |E| and dim(Z(X)) = |E| - |X| + 1 (this is clear for a tree, and any time an edge is added to a graph, both sides increase by 1). We will also use the standard inner product on $\mathbb{C}[E]$, such that edges have norm 1 and are perpendicular to each other. An edge in a connected graph is called a *bridge* if the removal of this edge would render the graph disconnected.

Lemma 2.3. Let X be a finite connected graph with edge set E. There is $v \in Z(X)$ satisfying the following conditions:

- (i) For all $e \in E$, we have $v(e) \in \{-1, 0, 1\}$.
- (ii) Of all edges $e \in E$ that are not bridges, at most half satisfy v(e) = 0.

Proof. Let Z_1, \ldots, Z_K be cycles in X such that each edge that is not a bridge is contained in one of the cycles Z_i . Choose $\varepsilon_1, \ldots, \varepsilon_K \in \{0, 1\}$ uniformly and independently at random. Let $w = \sum_{i=1}^{K} \varepsilon_i Z_i$. For each $e \in E$ that is not a bridge, we have $\mathbb{P}[w(e)$ is odd] $= \frac{1}{2}$. So $\mathbb{E}[\#\{e \in E \mid w(e) \text{ is odd}\}]$ is equal to half the number of edges that are not bridges. So there is some $w \in Z(X)$ such that w(e) is odd for at least half the number of edges that are not bridges. Now let $E' \subseteq E$ be the subset of E consisting of the edges for which w(e)is odd. Then each vertex of X has an even number of adjacent edges in E'. So each component of (V(X), E') has an Eulerian cycle. Let v be the sum of these Eulerian cycles. Then v satisfies the conditions.

Lemma 2.4. Let X be a finite connected graph with edge set E and maximal degree d. Suppose there is a constant h > 0 such that for all subsets A with

$$\frac{1}{4}|V(X)| \le |A| \le \frac{1}{2}|V(X)|,$$

we have $|\delta A| \ge h|A|$. Then there is a constant c > 0, only depending on d and h, such that if |V(X)| is large enough, the number of edges that are not bridges in X is at least c|E|.

Proof. For a bridge *b*, define K_b and G_b to be the two components of the graph $(V(X), E \setminus \{b\})$, with $|K_b| \le |G_b|$. Since $\delta K_b = \{b\}$, and $h \cdot \frac{1}{4}|V(X)| > 1$ for |V(X)| large enough, we must have $|K_b| < \frac{1}{4}|V(X)|$. Now let $G = \bigcap_b G_b$ and $K = V(X) \setminus G$, where the intersection ranges over all bridges *b*. If *x*, *y* \in *G*, then all the vertices of any minimal path from *x* to *y* are also in *G*. In particular, *G* is connected.

We show that G is non-empty. Let b be a bridge such that K_b is maximal. Let x be the endpoint of b in G_b . Then $x \in G$: for if there is another bridge b' with $x \in K_{b'}$, we have either $K_b \cup \{x\} \subseteq K_{b'}$, or $G_b \subseteq K_{b'}$, and then $K_{b'}$ is too large.

Note that all bridges of X must have at least one vertex in K. Therefore, the number of edges that are not bridges is at least |G| - 1. So if we find a constant c' > 0 such that $|G| \ge c'|V(X)|$, we are done.

Let $\delta G = \{b_1, \ldots, b_N\}$. These are all bridges. For each $1 \le i \le N$, we have $|K_{b_i}| \le \frac{1}{4}|V(X)|$. The K_{b_i} are pairwise disjoint: if K_{b_i} and K_{b_j} were not disjoint for $i \ne j$, then we would find a cycle with b_i and b_j . Then we can choose some $1 \le M \le N$ such that $\min(|K|, \frac{1}{4}|V(X)|) \le \sum_{j=1}^{M} |K_{b_j}| \le \frac{1}{2}|V(X)|$. If $K < \frac{1}{4}|V(X)|$, we have $|G| \ge \frac{3}{4}|V(X)|$, and we are done. Suppose that $K \ge \frac{1}{4}|V(X)|$. Then we can apply the assumption of the lemma to $A = \bigcup_{j=1}^{M} K_{b_j}$. We have

$$|\delta A| = M \le N = |\delta G| \le d |G|,$$

therefore $|A| \leq \frac{d}{h}|G|$. So $|G| \geq \frac{h}{4d}|V(X)|$. This finishes the proof.

Corollary 2.5. Let X be a finite connected graph with edge set E and maximal degree d. Suppose there is a constant h > 0 such that for all subsets $A \subset V(X)$ with $\frac{1}{4}|V(X)| \le |A| \le \frac{1}{2}|V(X)|$, we have $|\delta A| \ge h|A|$. Then, provided that |V(X)| is large enough, there are a constant c > 0, only depending on d and h, and $v \in Z(X)$ satisfying the following conditions:

- (i) For all $e \in E$, we have $v(e) \in \{-1, 0, 1\}$.
- (ii) We have $\#\{e \in E \mid v(e) \neq 0\} \ge c|E|$.

Proof. This follows directly from Lemmas 2.3 and 2.4.

Theorem A. Suppose $X = (X_n)$ is a sequence of finite connected graphs with uniformly bounded degree and with number of vertices going to infinity. Suppose that X has geometric property (T). Then there are constants R > 0 and $\varepsilon > 0$ such that for all large enough n, we have dim $Z_R(X_n) \ge \varepsilon |V(X_n)|$.

Proof. Let $\gamma > 0$ be such that $\sigma_{\max}(\Delta_X) \subseteq \{0\} \cup [\gamma, \infty)$. Let *d* be the maximal degree of all vertices of *X*. Since *X* has geometric property (T), it is, in particular, an expander sequence by Cheeger's inequality: for each subset $A \subseteq X_n$ with $|A| \leq \frac{1}{2}|V(X_n)|$, we have $|\delta A| \geq \frac{\gamma}{2}|A|$. Only finitely many of the X_n can be trees (this follows from Lemma 2.4, for example). We can then assume without loss of generality that none of the X_n is a tree.

Define $h = \frac{\gamma}{4}$. Let c_1 be the constant from Corollary 2.5, using the constants d and h. Let $c_2 > 0$ be such that

$$c_{3} = \frac{1}{4}d^{2} - \left(1 - \frac{c_{1}}{2d}\right)\left(\frac{1}{2}d + c_{2}d\right)^{2} - \frac{c_{1}}{2d}\left(\frac{1}{2}d + c_{2}d - \frac{1}{2}\right)^{2} > 0.$$
(1)

Let t > 0 be small enough such that

$$dt^2 < \gamma \tag{2}$$

and also

$$\left| \exp(it) - 1 - it + \frac{1}{2}t^2 \right| \le c_2 t^2.$$
 (3)

Let $\varepsilon > 0$ be such that the following conditions are satisfied:

$$8\varepsilon d^2 \le \frac{1}{2}c_3 t^4,\tag{4}$$

$$2\varepsilon \le \frac{c_1}{2d},\tag{5}$$

$$4\varepsilon \le h. \tag{6}$$

Suppose for a contradiction that for each R, there is an n such that dim $Z_R(X_n) < \varepsilon |V(X_n)|$. Let E_n be the set of edges of X_n . Note that $Z(X_n) \subseteq \mathbb{C}[E_n]$ consists of the functions $\rho: E_n \to \mathbb{C}$ satisfying $\sum_{y \sim x} \rho(x, y) = 0$ for all $x \in V(X_n)$. The subset $Z_R(X_n)^{\perp} \subseteq \mathbb{C}[E_n]$ consists of all functions $\rho: E_n \to \mathbb{C}$ satisfying $\rho(x_1, x_2) + \rho(x_2, x_3) + \cdots + \rho(x_q, x_1) = 0$ for all q-cycles (x_1, x_2, \ldots, x_q) with $q \leq R$.

Let $\rho \in Z_R(X_n)^{\perp}$. We will construct a $\frac{1}{3}R$ -representation π_{ρ} of X_n . First, we extend ρ to a function on $V(X_n)^2$ as follows: for each pair $(x, y) \in V(X_n)^2$, choose a shortest path $(x = x_0, x_1, \dots, x_{d(x,y)} = y)$ from x to y and define

$$\rho(x, y) = \rho(x_0, x_1) + \rho(x_1, x_2) + \dots + \rho(x_{d(x, y)-1}, x_{d(x, y)}).$$

Since $\rho \in Z_R(X_n)^{\perp}$, this satisfies the following: if $x, y \in V(X_n)$ satisfy $d(x, y) \leq R$, then $\rho(x, y) = -\rho(y, x)$, and if $x, y, z \in V(X_n)$ satisfy $d(x, y) + d(y, z) + d(z, x) \leq R$, then $\rho(x, y) + \rho(y, z) + \rho(z, x) = 0$.

Now we define $\pi_{\rho}: B(\ell^2 X_n) \to B(\ell^2 X_n)$ by

$$\pi_{\rho}(T)\xi(x) = \sum_{y} T_{xy} \exp(i\rho(x, y))\xi(y).$$

This is a $\frac{1}{3}R$ -representation.

For a subset $B \subseteq E_n$, denote by $X_n \setminus B$ the graph $(X_n, E_n \setminus B)$. Choose edges $b_1, \ldots, b_{\dim(Z_R(X_n))}$ recursively in such a way that b_j is an edge in a cycle in $Z_R(X_n) \cap Z(X_n \setminus \{b_1, \ldots, b_{j-1}\})$. Then $Z_R(X_n) \cap Z(X_n \setminus \{b_1, \ldots, b_{j-1}, b_j\})$ has one dimension fewer than $Z_R(X_n) \cap Z(X_n \setminus \{b_1, \ldots, b_{j-1}\})$. Thus, with $B = \{b_1, \ldots, b_{\dim(Z_R(X_n))}\}$ we have $Z_R(X_n) \cap Z(X_n \setminus B) = 0$. Then we also have $Z_R(X_n) \cap \mathbb{C}[E_n \setminus B] = 0$, and counting dimensions, $Z_R(X_n) \oplus \mathbb{C}[E_n \setminus B] = \mathbb{C}[E_n]$. Since $\mathbb{C}[B] \perp \mathbb{C}(E_n \setminus B)$ and $Z_R(X_n)^{\perp} \perp Z_R(X_n)$, it follows that $\mathbb{C}[B] \cap Z_R(X_n)^{\perp} = 0$. Counting dimensions again, we get $Z_R(X_n)^{\perp} \oplus \mathbb{C}[B] = \mathbb{C}[E_n]$.

Since each b_j is an edge in a cycle in $Z(X_n \setminus \{b_1, \ldots, b_{j-1}\})$, it follows by induction that $X_n \setminus B$ is still a connected graph. Let $A \subset V(X_n)$, with $\frac{1}{4}|V(X_n)| \le |A| \le \frac{1}{2}|V(X_n)|$. Denote by $\delta_{X_n \setminus B}A$ the set of edges in $X_n \setminus B$ with exactly one vertex in A. Then we have $|\delta_{X_n \setminus B}A| \ge |\delta_{X_n}A| - |B| \ge \frac{\gamma}{2}|A| - \varepsilon |V(X_n)| \ge (\frac{\gamma}{2} - 4\varepsilon)|A| = (2h - 4\varepsilon)|A| \ge h|A|$ by inequality (6). So $X_n \setminus B$ satisfies the conditions of Corollary 2.5. We can also assume $V(X_n)$ is large enough to apply the corollary by taking R large enough.

Now let $v \in C(X_n \setminus B)$ be as in Corollary 2.5. Recall that $Z_R(X_n)^{\perp} \oplus \mathbb{C}[B] = \mathbb{C}[E_n]$. Hence there is a unique function $\rho \in Z_R(X_n)^{\perp}$ such that $\rho(e) = v(e)$ for all $e \notin B$.

Now we consider the $\frac{1}{3}R$ -representation $\pi_{t\rho}$. Let $\Delta_t = \pi_{t\rho}(\Delta_X) \in B(\ell^2 X_n)$. Let $\xi_n \in \ell^2 X_n$ be the constant function with $\xi_n(x) = 1$ for all $x \in X_n$. Then we have

$$\Delta_t \xi_n(x) = \sum_{y \sim x} (1 - \exp(it\rho(x, y)))$$

for all $x \in X_n$.

We define a partition $V(X_n) = A_1 \cup A_2 \cup A_3$. We can think of the vertices in A_1 as the 'good' vertices, the vertices in A_2 as the 'neutral' vertices, and the vertices in A_3 as the 'bad' vertices. Luckily, A_3 will be small. The sets are defined as follows:

$$A_3 = \{x \in X_n \mid (x, y) \in B \text{ for some } y \sim x\},\$$

$$A_1 = \{x \in X_n \setminus A_3 \mid v(x, y) \neq 0 \text{ for some } y \sim x\},\$$

$$A_2 = X_n \setminus (A_1 \cup A_3).$$

Since $|B| \le \varepsilon |V(X_n)|$, we have $|A_3| \le 2\varepsilon |V(X_n)|$, and since at least $c_1 |E_n|$ edges e satisfy $v(e) \ne 0$, we have $|A_1| \ge \frac{c_1}{d} |E_n| - |A_3| \ge (\frac{c_1}{d} - 2\varepsilon)|V(X_n)| \ge \frac{c_1}{2d} |V(X_n)|$ by inequality (5).

We will estimate $|\Delta_t \xi_n(x) - \frac{1}{2}dt^2|$, finding increasingly good estimates in the cases $x \in A_3, A_2, A_1$, respectively. For all $x \in X_n$, we have $|\Delta_t \xi_n(x) - \frac{1}{2}dt^2| \le 2d$. For $x \notin A_3$, we have $\rho(x, y) = v(x, y)$ for all $y \sim x$. Since $\sum_{y \sim x} v(x, y) = 0$, we have

$$\Delta_t \xi_n(x) = \sum_{y \sim x} (1 + itv(x, y) - \exp(itv(x, y))).$$

Then, using inequality (3), we have

$$\begin{aligned} \left| \Delta_t \xi_n(x) - \frac{1}{2} dt^2 \right| &\leq \sum_{y \sim x} \left| 1 + itv(x, y) - \frac{d}{2 \deg(x)} t^2 - \exp(itv(x, y)) \right| \\ &\leq \left(\sum_{y \sim x} \left| 1 + itv(x, y) - \frac{d}{2 \deg(x)} t^2 - \left(1 + itv(x, y) - \frac{1}{2} t^2 v(x, y)^2 \right) \right| \right) + c_2 dt^2 \\ &= \left| \frac{1}{2} dt^2 + c_2 dt^2 - \frac{1}{2} \sum_{y \sim x} v(x, y)^2 t^2. \end{aligned}$$

For $x \in A_2$, this is at most $(\frac{1}{2}d + c_2d)t^2$. For $x \in A_1$, there is $y \sim x$ with $v(x, y) \neq 0$, and we find that $|\Delta_t \xi_n(x) - \frac{1}{2}dt^2| \le (\frac{1}{2}d + c_2d - \frac{1}{2})t^2$.

Using inequalities (1) and (4), we now have

$$\begin{split} \left\| \Delta_t \xi_n - \frac{1}{2} dt^2 \xi_n \right\|^2 &= \sum_{x \in X_n} \left| \Delta_t \xi_n(x) - \frac{1}{2} dt^2 \right|^2 \\ &\leq \sum_{x \in A_1} \left(\frac{1}{2} d + c_2 d - \frac{1}{2} \right)^2 t^4 + \sum_{x \in A_2} \left(\frac{1}{2} d + c_2 d \right)^2 t^4 + \sum_{x \in A_3} 4 d^2 \\ &\leq \left(\left(1 - \frac{c_1}{2d} \right) \left(\frac{1}{2} d + c_2 d \right)^2 + \frac{c_1}{2d} \left(\frac{1}{2} d + c_2 d - \frac{1}{2} \right)^2 \right) |V(X_n)| t^4 \\ &+ 8\varepsilon d^2 |V(X_n)| \\ &= \left(\frac{1}{4} d^2 - c_3 \right) |V(X_n)| t^4 + 8\varepsilon d^2 |V(X_n)| \\ &\leq \left(\frac{1}{4} d^2 - \frac{1}{2} c_3 \right) |V(X_n)| t^4 = \left(1 - \frac{2c_3}{d^2} \right) \left\| \frac{1}{2} dt^2 \xi_n \right\|^2. \end{split}$$

For each *R*, we have found an n_R , a $\frac{1}{3}R$ -representation $\pi_{n_R}: B(\ell^2 X_{n_R}) \to B(\ell^2 X_{n_R})$, and a vector $\xi_{n_R} \in B(\ell^2 X_{n_R})$ satisfying

$$\left\|\pi_{n_{R}}(\Delta_{X})\xi_{n_{R}}-\frac{1}{2}dt^{2}\xi_{n_{R}}\right\|\leq\left(1-\frac{2c_{3}}{d^{2}}\right)^{\frac{1}{2}}\left\|\frac{1}{2}dt^{2}\xi_{n_{R}}\right\|.$$

Let \mathcal{U} be a free ultrafilter on \mathbb{N} , and define the ultraproduct $\mathcal{H} = \prod_n B(\ell^2 X_{n_R})/\mathcal{U}$. Let $\pi = \lim_{\mathcal{U}} \pi_{n_R}$ and $\xi = \lim_{\mathcal{U}} \frac{\xi_{n_R}}{\|\xi_{n_R}\|}$. Then π is a representation of $\mathbb{C}_u[X]$, and we have

$$\left\|\pi(\Delta_X)\xi - \frac{1}{2}dt^2\xi\right\| \le \left(1 - \frac{2c_3}{d^2}\right)^{\frac{1}{2}} \left\|\frac{1}{2}dt^2\xi\right\|.$$

It follows that

$$\sigma(\pi(\Delta_X)) \cap \left[\frac{1}{2}dt^2\left(1 - \left(1 - \frac{2c_3}{d^2}\right)^{\frac{1}{2}}\right), \frac{1}{2}dt^2\left(1 + \left(1 - \frac{2c_3}{d^2}\right)^{\frac{1}{2}}\right)\right] \neq \emptyset.$$

So $\sigma_{\max}(\Delta_X)$ contains a positive element that is at most dt^2 . This is a contradiction with inequality (2).

Remark 2.6. Note that in the above proof, ε only depends on d and γ .

We have shown that each sequence of graphs with geometric property (T) has many small cycles. It follows that we can remove a large number of cycles of the graph, while still keeping the same coarse structure. In particular, the infimum in the definition of cost is not attained (see Definition 1.4).

Proposition 2.7. Let $X = (X_n)$ be a sequence of graphs with degree at most d. Suppose there are $R, \varepsilon > 0$ such that dim $Z_R(X_n) \ge \varepsilon |V(X_n)|$ for all n. Then

$$c(X) \leq \liminf_{n} \frac{|E(X_n)|}{|V(X_n)|} - \frac{\varepsilon}{d^{R-1}}.$$

In particular, if X has geometric property (T), the infimum $c(X) = \inf_{Y} \liminf_{n \in Y_n|} \frac{|E(Y_n)|}{|V(X_n)|}$, over all sequences (Y_n) inducing the same coarse structure as X, is not attained.

Proof. Consider the set of all subgraphs of X_n , on the same vertex set, without *R*-cycles. Let Y_n be a maximal element of this set. If two vertices x, y are connected in X_n , then either they are connected in Y_n , or adding the edge (x, y) to Y_n would create an *R*-cycle in Y_n . So $d_Y(x, y) \le R - 1$.

It follows that $d_Y(x, y) \le (R-1)d_X(x, y)$ for any two vertices $x, y \in V(X_n)$. Since Y_n is a subgraph of X_n , we also have $d_Y(x, y) \ge d_X(x, y)$. This shows that X and Y induce the same coarse structure.

Since dim $Z_R(X_n) \ge \varepsilon |V(X_n)|$, there are, in particular, at least $\varepsilon |V(X_n)|$ cycles of length at most R in X_n . Of all these cycles, at least one of its edges is not in Y_n . One edge can be contained in at most d^{R-1} cycles of length at most R. So $|E(Y_n)| \le |E(X_n)| - \frac{\varepsilon}{d^{R-1}}|V(X_n)|$. This gives

$$\liminf_{n} \frac{|E(Y_n)|}{|V(X_n)|} \le \liminf_{n} \frac{|E(X_n)|}{|V(X_n)|} - \frac{\varepsilon}{d^{R-1}}$$

So $c(X) \leq \liminf_{n} \frac{|E(X_n)|}{|V(X_n)|} - \frac{\varepsilon}{d^{R-1}}$.

The last statement of the lemma follows from Theorem A and the fact that if X and Y induce the same coarse structure, also Y has geometric property (T) (see [12, Theorem 4.1]).

3. Behaviour of geometric property (T) under small changes

A natural question in light of Theorem A is the following: if a sequence of graphs (X_n) has geometric property (T), is it necessary that there is an R such that every vertex has a cycle in its R-neighbourhood? In this section, we will see that the answer is no. We will do this by proving that if we change a sequence of graphs a small amount (see Definition 1.5) while keeping expansion, we also keep geometric property (T).

Theorem B. Let $X = (X_n)$ and $Y = (Y_n)$ be approximately isomorphic sequences of finite connected graphs with uniformly bounded degree and number of vertices going to infinity. Suppose X has geometric property (T) and Y is an expander. Then Y has geometric property (T).

For the proof, we need the following proposition.

Proposition 3.1. Let $X = (X_n)$ be an expander sequence with maximum degree d, and let h > 0 such that $\sigma(\Delta_X) \subseteq \{0\} \cup [h, \infty)$. Let $\pi: \mathbb{C}_u[X] \to B(\mathcal{H})$ be a representation, and suppose that $v \in \mathcal{H}$ is a unit vector with $\Delta_X v = \eta v$ for some $\eta > 0$.

- (i) Let $F_n \subseteq V(X_n)$ be such that $\lim_{n\to\infty} \frac{|F_n|}{|V(X_n)|} = 0$, and let $P_F \in \mathbb{C}_u[X]$ denote the projection on the vertices of the F_n . Then $||P_Fv|| \le 2^{\frac{3}{4}} d^{\frac{1}{2}} h^{-\frac{1}{2}} \eta^{\frac{1}{4}}$.
- (ii) Let $G = (G_n)$ be a sequence of subgraphs of X_n such that $\lim_{n \to \infty} \frac{|V(G_n)|}{|V(X_n)|} = 0$. Then $\langle \Delta_G v, v \rangle \leq 2^{\frac{3}{2}} dh^{-1} \eta^{\frac{3}{2}}$ and $\|\Delta_G v\| \leq 2^{\frac{3}{4}} d^{\frac{1}{2}} h^{-\frac{1}{2}} \eta^{\frac{5}{4}}$.

Proof. (i) We can assume that $\eta < h$. Let $\delta > 0$ and let N be a large enough integer such that $|F_n|/|V(X_n)| < \delta$ for n > N. Let $P_{X_{\leq N}} \in \mathbb{C}_u[X]$ be the projection on the vertices of X_1 up to X_N . Then we have $\Delta_X P_{X_{\leq N}} v = \eta P_{X_{\leq N}} v$. Since η is not an eigenvalue of $\Delta_X P_{X_{\leq N}} v$, we have $P_{X_{\leq N}} v = 0$. So $P_F v = P_{F_{>N}} v$, where $P_{F_{>N}}$ is the projection on the union $F_{>N} = \bigcup_{n>N} F_n$.

We can colour the edges of X in 2d colours such that no two intersecting edges have the same colour. For each colour *i*, define the involution τ_i of X that sends each vertex in an edge with colour *i* to the other vertex of this edge, while fixing the other vertices. This defines an element in $\mathbb{C}_u[X]$ that we also denote by τ_i . As $\tau_i^2 = 1$, we have $0 \le 1 - \tau_i \le 2$. This inequality holds in $C_{\max}^*(X)$. We have $\Delta_X = \sum_{i=1}^{2d} (1 - \tau_i)$: we can check this by computing the matrix coefficients. For any $x \in X$, we have

$$\sum_{i=1}^{2d} (1 - \tau_i)_{xx} = \#\{i \mid x \text{ has an edge with colour } i\} = \deg(x)$$

and for different $x, y \in X$, we have

$$\sum_{i=1}^{2d} (1-\tau_i)_{xy} = -\#\{i \mid (x, y) \text{ is an edge with colour } i\},\$$

which is -1 if (x, y) is an edge and 0 otherwise.

For $f \in \ell^{\infty} X$, denote the corresponding multiplication operator in $B(\ell^2 X)$ by M_f . Define the positive unital linear map $\varphi: \ell^{\infty} X \to \mathbb{C}$ by $\varphi(f) = \langle M_f v, v \rangle$. For each *i*, we have $\langle (1 - \tau_i)v, v \rangle \leq \eta$, so $\|(1 - \tau_i)v\|^2 = \langle (2 - 2\tau_i)v, v \rangle \leq 2\eta$. Then for $f \in \ell^{\infty} X$, we have

$$(1 - \tau_i)\varphi(f) = \varphi(f - f \circ \tau_i)$$

= $\langle M_f v, v \rangle - \langle M_f \circ \tau_i v, v \rangle$
= $\langle M_f v, v \rangle - \langle M_f \tau_i v, \tau_i v \rangle$
= $\langle M_f v, v - \tau_i v \rangle + \langle M_f (v - \tau_i v), \tau_i v \rangle.$

We get $|(1 - \tau_i)\varphi(f)| \le 2||f||_{\infty}||v - \tau_i v|| \le 2\sqrt{2\eta}||f||_{\infty}$, so $||(1 - \tau_i)\varphi|| \le 2\sqrt{2\eta}$.

So φ is almost invariant for the involutions τ_i . Note that φ is a unit vector in the Banach space $(\ell^{\infty}X)^*$. By the Goldstine theorem, φ is in the weak closure of the set $\mathcal{P}(X) \subseteq \ell^1 X$ of probability measures on X. We will use some functional analysis to show that we can approximate φ by a probability measure $\psi \in \mathcal{P}(X)$, in such a way that it will still be almost invariant for the involutions τ_i .

Let \mathcal{B} be the real Banach space $\bigoplus_{i=1}^{2d} \ell^1(X, \mathbb{R})$ with norm $\|(\psi_i)\| = \max_i \|\psi_i\|_1$. Define the convex set $B \subseteq \mathcal{B}$ by

$$B = \{ ((1 - \tau_i)\psi) \mid \psi \in \mathcal{P}(X), |\psi(\mathbb{1}_{F>N}) - \varphi(\mathbb{1}_{F>N})| < \delta \}.$$

There is a net $\psi_{\lambda} \in \mathcal{P}(X)$ converging weakly to φ . For large λ , we have $|\psi_{\lambda}(\mathbb{1}_{F>N}) - \varphi(\mathbb{1}_{F>N})| < \delta$. Now $(1 - \tau_i)\psi_{\lambda}$ converges weakly to $(1 - \tau_i)\varphi \in (\ell^{\infty}X)^*$ for all *i*. So $((1 - \tau_i)\varphi) \in \mathcal{B}^{**}$ is in the weak closure of *B*.

Let *A* be the open ball $\{a \in \mathcal{B} \mid ||a|| < 2\sqrt{2\eta} + \delta\}$. Suppose *A* and *B* are disjoint. By the Hahn–Banach separation theorem, there are $f \in \mathcal{B}^*$ and a positive real number *s* with $f(a) < s \le f(b)$ for $a \in A$ and $b \in B$. Now $((1 - \tau_i)\varphi) \in B^{**}$ is in the weak closure of *B*, so $f(((1 - \tau_i)\varphi)) \ge s$. On the other hand, $||((1 - \tau_i)\varphi)|| \le 2\sqrt{2\eta}$, so $((1 - \tau_i)\varphi)$ is in the weak closure of $\frac{2\sqrt{2\eta}}{2\sqrt{2\eta}+\delta}A$, showing that $f(((1 - \tau_i)\varphi)) \le \frac{2\sqrt{2\eta}}{2\sqrt{2\eta}+\delta}s$, giving a contradiction.

Therefore, $A \cap B \neq \emptyset$, so there is $\psi \in \mathcal{P}(X)$ with $|\psi(\mathbb{1}_{F>N}) - \varphi(\mathbb{1}_{F>N})| < \delta$ and $||(1 - \tau_i)\psi|| < 2\sqrt{2\eta} + \delta$ for all *i*. Let $\xi = \psi^{\frac{1}{2}} \in \ell^2 X$. Then for all *i*, we have

$$\begin{aligned} \langle (1 - \tau_i)\xi, \xi \rangle &= \sum_{x \in X} (\xi(x) - \xi(\tau_i x))\xi(x) \\ &= \frac{1}{2} \sum_{x \in X} (\xi(x) - \xi(\tau_i x))^2 \\ &\leq \frac{1}{2} \sum_{x \in X} |\xi(x)^2 - \xi(\tau_i x)^2| \\ &= \frac{1}{2} \| (1 - \tau_i)\psi \|_1 \le \sqrt{2\eta} + \frac{1}{2}\delta. \end{aligned}$$

Summing over all *i* gives $\langle \Delta_X \xi, \xi \rangle \leq 2d \sqrt{2\eta} + d\delta$. Denote by ξ_c the projection of ξ on the locally constant functions in $\ell^2 X$. We get $\langle \Delta_X(\xi - \xi_c), \xi - \xi_c \rangle = \langle \Delta_X \xi, \xi \rangle \leq 2d \sqrt{2\eta} + d\delta$, but also $\langle \Delta_X(\xi - \xi_c), \xi - \xi_c \rangle \geq h \|\xi - \xi_c\|_2^2$. So

$$\|\xi - \xi_c\|_2^2 \le \frac{1}{h} (2d\sqrt{2\eta} + d\delta).$$

Since $\frac{|F_n|}{V(X_n)} < \delta$ for n > N, we have $\|\xi_{c|F_{>N}}\| < \sqrt{\delta}$. We get

$$\begin{split} \psi(\mathbb{1}_{F_{>N}}) &= \sum_{x \in F_{>N}} \xi(x)^2 \\ &= \sum_{x \in F_{>N}} (\xi - \xi_c)(x)^2 + 2 \sum_{x \in F_{>N}} \xi_c(x)(\xi - \xi_c)(x) + \sum_{x \in F_{>N}} \xi_c(x)^2 \\ &\leq \|\xi - \xi_c\|_2^2 + 2\|\xi_c|_{F_{>N}}\|_2 \cdot \|\xi - \xi_c\|_2 + \|\xi_c|_{F_{>N}}\|_2^2 \\ &\leq \frac{1}{h} (2d\sqrt{2\eta} + d\delta) + 2\sqrt{\delta} + \delta. \end{split}$$

Finally, we have

$$\begin{split} \|P_F v\|^2 &= \|P_{F>N} v\|^2 = \langle M_{\mathbb{1}_{F>N}} v, v \rangle = \varphi(\mathbb{1}_{F>N}) \le \psi(\mathbb{1}_{F>N}) + \delta \\ &\le \frac{1}{h} (2d\sqrt{2\eta} + d\delta) + 2\sqrt{\delta} + 2\delta. \end{split}$$

Letting $\delta \rightarrow 0$ gives the desired conclusion.

(ii) We recursively define $F_0 = \bigcup_n V(G_n)$ and

$$F_{k+1} = \{x \in V(X) \setminus F_{\leq k} \mid x \text{ is adjacent to a vertex in } F_k\}$$

for $k \ge 1$. Here we write $F_{\le k}$ for $F_0 \cup \cdots \cup F_k$. Let $\delta > 0$. Since there are infinitely many F_k and they are all disjoint, there is $k \ge 1$ with $\|P_{F_{k-1} \cup F_k}v\| \le \delta$. Since the graphs have uniformly bounded degree, we have $\lim_{n\to\infty} \frac{|F_{\le k-1}|}{|V(X_n)|} = 0$. By part (i), we have $\|P_{F_{\le k-1}}v\| \le 2^{\frac{3}{4}}d^{\frac{1}{2}}h^{-\frac{1}{2}}\eta^{\frac{1}{4}}$. Denote by $\Delta_{F_{\le k}}$ the Laplacian operator of the induced subgraph with vertex set $F_{< k}$. Then we have

$$P_{F_{\leq k-1}}\Delta_X = P_{F_{\leq k-1}}\Delta_{F_{\leq k}} = \Delta_{F_{\leq k}} - P_{F_k}\Delta_{F_{\leq k}} = \Delta_{F_{\leq k}} - P_{F_k}\Delta_{F_{\leq k}}P_{F_{k-1}\cup F_k}.$$

It follows that

$$\begin{split} \langle \Delta_G v, v \rangle &\leq \langle \Delta_{F_{\leq k}} v, v \rangle = \langle P_{F_{\leq k-1}} \Delta_X v, v \rangle + \langle P_{F_k} \Delta_{F_{\leq k}} P_{F_{k-1} \cup F_k} v, v \rangle \\ &\leq \eta \langle P_{F_{\leq k-1}} v, v \rangle + \| P_{F_k} \Delta_{F_{\leq k}} \| \cdot \| P_{F_{k-1} \cup F_k} v \| \cdot \| v \| \\ &< 2^{\frac{3}{2}} dh^{-1} \eta^{\frac{3}{2}} + 2d\delta. \end{split}$$

Similarly, we have

$$\begin{split} \|\Delta_G v\| &\leq \|F_{\leq k}v\| \leq \|P_{F_{\leq k-1}}\Delta_X v\| + \|P_{F_k}\Delta_{F_{\leq k}}P_{F_{k-1}\cup F_k}v\| \\ &\leq 2^{\frac{3}{4}}d^{\frac{1}{2}}h^{-\frac{1}{2}}\eta^{\frac{5}{4}} + 2d\delta. \end{split}$$

Since these inequalities hold for all $\delta > 0$, the conclusion follows.

Proof of Theorem B. Since X and Y are approximately isomorphic, we can identify the isomorphic subgraphs and assume there are induced subgraphs $Z_n \subseteq X_n \cap Y_n$ with

$$\lim_{n \to \infty} \frac{|V(Z_n)|}{|V(X_n)|} = \lim_{n \to \infty} \frac{|V(Z_n)|}{|V(Y_n)|} = 1.$$

Let *d* be the maximum degree in $X \cup Y$, let $\gamma > 0$ with $\sigma_{\max}(\Delta_X) \subseteq \{0\} \cup [\gamma, \infty)$, and let h > 0 with $\sigma(\Delta_Y) \subseteq \{0\} \cup [h, \infty)$. Suppose that *Y* does not have geometric property (T). Then the maximal spectrum of Δ_Y contains arbitrarily small positive numbers. Let $0 < \eta < h$ be in the maximal spectrum of Δ_Y . Then there are a representation $\rho: \mathbb{C}_u[Y] \to B(\mathcal{H})$ and a unit vector $v \in \mathcal{H}$ with $\Delta_Y v = \eta v$.

We will first apply Proposition 3.1 to bound v on some small subsets of Y. Then we will construct a representation of $\mathbb{C}_u[X]$ containing a vector $1 \otimes v$ that we will show is almost constant. Since X has geometric property (T), it follows that $1 \otimes v$ is close to some constant vector. This will give a contradiction because v is perpendicular to all constant vectors.

We have

$$\langle \Delta_{Y \setminus Z} v, v \rangle \le 2^{\frac{3}{2}} dh^{-1} \eta^{\frac{3}{2}}$$
 and $\|\Delta_{Y \setminus Z} v\| \le 2^{\frac{3}{4}} d^{\frac{1}{2}} h^{-\frac{1}{2}} \eta^{\frac{5}{4}}$

by Proposition 3.1 (ii).

Let $F = \{z \in Z \mid z \text{ adjacent to some } x \in X \setminus Z\}$. Since the degree of the vertices of *Y* is uniformly bounded, we have $\lim_{n\to\infty} \frac{|F \cap Y_n|}{|Y_n|} = 0$. By Proposition 3.1 (i), we have $\|P_F v\| \le 2^{\frac{3}{4}} d^{\frac{1}{2}} h^{-\frac{1}{2}} \eta^{\frac{1}{4}}$.

Now we construct a representation of $\mathbb{C}_{u}[X]$. Consider the map $E: \mathbb{C}_{u}[X] \to \mathbb{C}_{u}[Z]$ given by $E(T) = P_{Z}TP_{Z}$. This is a conditional expectation, meaning that for $T \in \mathbb{C}_{u}[X]$ and $S \in \mathbb{C}_{u}[Z]$, we have E(TS) = E(T)S and E(ST) = SE(T). We can construct the tensor product $\mathcal{H}' = \mathbb{C}_{u}[X] \otimes_{\mathbb{C}_{u}[Z]} \mathcal{H}$, as in [8, Theorem 1.8]. We repeat the construction here. First we consider the algebraic tensor product $\mathbb{C}_{u}[X] \odot \mathcal{H}$. We equip this with a conjugate symmetric form given on simple tensors by

$$\langle T_1 \odot v_1, T_2 \odot v_2 \rangle = \langle E(T_2^*T_1)v_1, v_2 \rangle.$$

It can be shown that this is positive semi-definite (see [8, Lemma 1.7] and its proof). Define the semi-norm $||w|| = \langle w, w \rangle^{\frac{1}{2}}$ for $w \in \mathbb{C}_{u}[X] \odot \mathcal{H}$. Let $\mathcal{H}' = \mathbb{C}_{u}[X] \otimes_{\mathbb{C}_{u}[Z]} \mathcal{H}$ be the Hilbert space we get by taking the quotient with respect to the kernel of $|| \cdot ||$, and then taking the completion. Let $T_1 \otimes v_1$ denote the image of $T_1 \odot v_1$ in \mathcal{H}' . It is easy to see that for $S \in \mathbb{C}_{u}[Z]$, we have $T_1S \otimes v_1 = T_1 \otimes Sv_1$. We now have a representation $\pi: \mathbb{C}_{u}[X] \to B(\mathcal{H}')$, given on simple tensors by $\pi(T_1)(T_2 \otimes v_1) = T_1T_2 \otimes v_1$.

Consider the unit vector $1 \otimes v \in \mathcal{H}'$. We have

$$\begin{split} \langle \Delta_X (1 \otimes v), 1 \otimes v \rangle &= \langle \Delta_Z v, v \rangle + \langle E(\Delta_X \setminus Z) v, v \rangle \\ &= \langle \Delta_Z v, v \rangle + \langle E(\Delta_X \setminus Z) P_F v, P_F v \rangle \\ &\leq \langle \Delta_Y v, v \rangle + \| E(\Delta_X \setminus Z) \| \cdot \| P_F v \|^2 \\ &\leq \eta + 2d \cdot 2^{\frac{3}{2}} dh^{-1} \eta^{\frac{1}{2}} \leq 8d^2 h^{-1} \eta^{\frac{1}{2}}, \end{split}$$

provided η is small enough. Let $w \in \mathcal{H}'_c$ be the projection of $1 \otimes v$ on the space of constant vectors $\mathcal{H}'_c = \ker(\rho(\Delta_X))$. Then we have

$$\begin{aligned} \langle \Delta_X(1 \otimes v), 1 \otimes v \rangle &= \langle \Delta_X(1 \otimes v - w), 1 \otimes v - w \rangle \\ &\geq \gamma \| 1 \otimes v - w \|^2. \end{aligned}$$

Combining these inequalities, we get

$$\|1 \otimes v - w\|^2 \le 8d^2h^{-1}\gamma^{-1}\eta^{\frac{1}{2}}.$$

Finally, we have

$$\begin{split} \eta &= \langle \Delta_{Y} v, v \rangle \\ &= \langle \Delta_{Z} v, v \rangle + \langle \Delta_{Y \setminus Z} v, v \rangle \\ &= \langle \Delta_{Z} (1 \otimes v), w \rangle + \langle \Delta_{Z} (1 \otimes v), 1 \otimes v - w \rangle + \langle \Delta_{Y \setminus Z} v, v \rangle \\ &= \langle \Delta_{Z} (1 \otimes v), 1 \otimes v - w \rangle + \langle \Delta_{Y \setminus Z} v, v \rangle \\ &= \langle \eta (1 \otimes v), 1 \otimes v - w \rangle - \langle 1 \otimes \Delta_{Y \setminus Z} v, 1 \otimes v - w \rangle + \langle \Delta_{Y \setminus Z} v, v \rangle \\ &\leq \eta \| 1 \otimes v - w \|^{2} + \| \Delta_{Y \setminus Z} v \| \cdot \| 1 \otimes v - w \| + 2^{\frac{3}{2}} dh^{-1} \eta^{\frac{3}{2}} \\ &\leq (8d^{2}h^{-1}\gamma^{-1} + 2^{\frac{9}{4}}d^{\frac{3}{2}}h^{-1}\gamma^{-\frac{1}{2}} + 2^{\frac{3}{2}}dh^{-1})\eta^{\frac{3}{2}}. \end{split}$$

This gives a contradiction if η is small enough. Therefore, Y must have geometric property (T).

Using the theorem, we can construct a sequence of graphs with geometric property (T) such that the graphs locally have arbitrarily large girth.

Corollary 3.2. There is a sequence of graphs (X_n) with uniformly bounded degree and number of vertices converging to infinity, satisfying geometric property (T), with a designated vertex $p_n \in V(X_n)$, such that the n-ball around p_n does not contain any cycles.

Proof. We start with a sequence of finite connected graphs (Y_n) with geometric property (T) and maximal degree d. We will construct the graphs X_n by attaching new trees to each Y_n .

Let (R_n) be an unbounded sequence of integers with $\lim_{n\to\infty} \frac{2^{R_n}}{|V(Y_n)|} = 0$. Let (T_n, p_n) be a rooted tree of depth R_n such that each vertex except for the leaves has degree 3. Connect each leaf of the tree to a different vertex in Y_n . We call the new graph X_n , and we show that (X_n) is still an expander sequence. Note that the maximal degree of X_n equals d + 1. For a subset $A \subseteq V(X_n)$, denote by $\delta_{out}A$ the outer vertex boundary, that is, the set $\{x \in V(X_n) \mid x \notin A, x \text{ adjacent to a vertex in } A\}$. Since (Y_n) is an expander sequence, there is h > 0 such that for all n and all $B \subseteq V(Y_n)$ with $|B| \le \frac{2}{3}|Y_n|$, we have $|\delta_{out}B| \ge h|B|$. Now let $C \subseteq V(X_n)$ with $|C| \le \frac{1}{2}|V(X_n)|$. Write $A = C \cap V(T_n)$ and $B = C \cap V(Y_n)$. Note that $|B| \le \frac{2}{3}|V(Y_n)|$ (provided n is large enough). We have

 $|\delta_{\text{out}}A| \ge \frac{1}{2}|A|$, and all vertices in $\delta_{\text{out}}A$ are also in $\delta_{\text{out}}C$ unless they are in B, so $|\delta_{\text{out}}C| \ge \frac{1}{2}|A| - |B|$. We also have

$$|\delta_{\text{out}}C| \ge |\delta_{\text{out}}B \cap V(Y_n)| \ge h|B|.$$

Taking a convex combination, we conclude that

$$|\delta_{\text{out}}C| \ge \left(1 + \frac{3}{2h}\right)^{-1} \left(\frac{1}{2}|A| - |B| + \frac{3}{2h}h|B|\right) = \frac{h}{2h+3}|C|.$$

Hence, (X_n) is an expander sequence.

By the condition on R_n , we see that (X_n) approximates (Y_n) . By Theorem B, the sequence (X_n) has geometric property (T). The R_n -neighbourhood of p_n is the tree T_n , so it does not contain any cycles. After taking a subsequence and renumbering the graphs, we get the sequence of graphs we wanted.

Acknowledgements. I would like to thank my advisor Tim de Laat for his support and suggestions. I would also like to thank Federico Vigolo for helpful discussions, and the referee for helpful suggestions.

Funding. The author is supported by the Deutsche Forschungsgemeinschaft under Germany's Excellence Strategy – EXC 2044-390685587, Mathematics Münster: Dynamics – Geometry – Structure.

References

- G. Elek, The combinatorial cost. *Enseign. Math.* (2) 53 (2007), no. 3–4, 225–235
 Zbl 1146.05307 MR 2455943
- [2] D. Gaboriau, Coût des relations d'équivalence et des groupes. *Invent. Math.* 139 (2000), no. 1, 41–98 Zbl 0939.28012 MR 1728876
- [3] G. Gong, Q. Wang, and G. Yu, Geometrization of the strong Novikov conjecture for residually finite groups. J. Reine Angew. Math. 621 (2008), 159–189 Zbl 1154.46042 MR 2431253
- [4] T. Hutchcroft and G. Pete, Kazhdan groups have cost 1. Invent. Math. 221 (2020), no. 3, 873– 891 Zbl 07233320 MR 4132958
- [5] A. Khukhro, K. Li, F. Vigolo, and J. Zhang, On the structure of asymptotic expanders. Adv. Math. 393 (2021), article no. 108073 Zbl 1487.46051 MR 4340225
- [6] G. Levitt, On the cost of generating an equivalence relation. Ergodic Theory Dynam. Systems 15 (1995), no. 6, 1173–1181 Zbl 0843.28010 MR 1366313
- [7] A. Lubotzky, *Discrete groups, expanding graphs and invariant measures*. Mod. Birkhäuser Class., Birkhäuser, Basel, 2010 Zbl 1183.22001 MR 2569682
- [8] M. A. Rieffel, Induced representations of C*-algebras. Adv. Math. 13 (1974), 176–257
 Zbl 0284.46040 MR 353003
- [9] J. Roe, Coarse cohomology and index theory on complete Riemannian manifolds. *Mem. Amer. Math. Soc.* 104 (1993), no. 497, x+90 pp. Zbl 0780.58043 MR 1147350

- [10] R. Willett and G. Yu, Higher index theory for certain expanders and Gromov monster groups, I. Adv. Math. 229 (2012), no. 3, 1380–1416 Zbl 1243.46060 MR 2871145
- [11] R. Willett and G. Yu, Higher index theory for certain expanders and Gromov monster groups, II. Adv. Math. 229 (2012), no. 3, 1762–1803 Zbl 1243.46061 MR 2871156
- [12] R. Willett and G. Yu, Geometric property (T). Chin. Ann. Math. Ser. B 35 (2014), no. 5, 761– 800 Zbl 1362.46073 MR 3246936

Received 11 March 2022.

Jeroen Winkel

Fachbereich Mathematik und Informatik, Westfälische Wilhelms-Universität Münster, Einsteinstrasse 62, 48149 Münster, Germany; jwinkel@uni-muenster.de