



Inkang Kim · Pierre Pansu

Local rigidity in quaternionic hyperbolic space

Received December 11, 2006 and in revised form August 17, 2007

Abstract. We study deformations of quaternionic hyperbolic lattices in larger quaternionic hyperbolic spaces and prove local rigidity results. On the other hand, surface groups are shown to be more flexible in quaternionic hyperbolic plane than in complex hyperbolic plane.

Keywords. Quaternionic hyperbolic space, rank one symmetric space, quasifuchsian representation, bending, rigidity, group cohomology

1. Introduction

1.1. 4-dimensional lattices

Lattices in $\mathrm{Sp}(n, 1)$, $n \geq 2$, when mapped to $\mathrm{Sp}(m, 1)$, cannot be deformed. This follows from K. Corlette's archimedean superrigidity theorem [5]. What about lattices in $\mathrm{Sp}(1, 1)$, i.e. in 4-dimensional hyperbolic space?

In this note we prove local rigidity of uniform lattices of $\mathrm{Sp}(1, 1)$ when mapped to $\mathrm{Sp}(2, 1)$. In complex hyperbolic geometry, such rigidity results were first discovered by D. Toledo [22]. In [8, 9], W. Goldman and J. Millson gave a local explanation of this phenomenon. Our main result is an exact quaternionic analogue of theirs.

Start with a uniform lattice Γ in $\mathrm{Sp}(1, 1)$. There is an easy manner to deform the embedding $\rho_0 : \Gamma \rightarrow \mathrm{Sp}(1, 1) \rightarrow \mathrm{Sp}(2, 1)$. Indeed, since $\mathrm{Sp}(2, 1)$ contains $\mathrm{Sp}(1, 1) \times \mathrm{Sp}(1)$, it also contains many copies of $\mathrm{Sp}(1, 1) \times U(1)$. If $H^1(\Gamma, \mathbb{R}) \neq 0$, which happens sometimes (see [17]), the trivial representation $\Gamma \rightarrow U(1)$ can be continuously deformed to a nontrivial representation ρ_1 . All such representations give rise to actions on quaternionic hyperbolic plane which stabilize a quaternionic line. Therefore, only deformations which do not stabilize any quaternionic line should be of interest.

Theorem 1.1. *Let $\Gamma \subset \mathrm{Sp}(1, 1)$ be a lattice. Embed Γ into $\mathrm{Sp}(2, 1)$ as a subgroup which stabilizes a quaternionic line.*

I. Kim: School of Mathematics, KIAS, Seoul, 130-722, Korea; e-mail: inkang@kias.re.kr

P. Pansu: Laboratoire de Mathématiques d'Orsay, UMR 8628 du CNRS, Université Paris-Sud, 91405 Orsay Cedex, France, e-mail: pierre.pansu@math.u-psud.fr

Mathematics Subject Classification (2000): 51M10, 57S25

If Γ is uniform in $\mathrm{Sp}(1, 1)$, then every small deformation of Γ in $\mathrm{Sp}(2, 1)$ again stabilizes a quaternionic line.

If Γ is not uniform in $\mathrm{Sp}(1, 1)$, then every small deformation of Γ in $\mathrm{Sp}(2, 1)$ preserving parabolics again stabilizes a quaternionic line.

Toledo's theorem inaugurated a series of global rigidity results by A. Domic and D. Toledo [6], K. Corlette [4], M. Burger, A. Iozzi and A. Wienhard [3]. By global rigidity, we mean the following: a certain characteristic number of representations, known as Toledo invariant, is maximal if and only if the representation stabilizes a totally geodesic complex hypersurface. It is highly expected that such a global rigidity should hold in quaternionic hyperbolic spaces, but we have been unable to prove it. Note that since $\mathrm{Sp}(1, 1) = \mathrm{Spin}(4, 1)^0$, there exist uniform lattices in $\mathrm{Sp}(1, 1)$ which are isomorphic to Zariski dense subgroups of $\mathrm{Sp}(4, 1)$ (see Section 7).

Question. Let $\Gamma \subset \mathrm{Sp}(1, 1)$ be a uniform lattice. Embed Γ into $\mathrm{Sp}(3, 1)$. Can one deform Γ to a Zariski dense subgroup?

1.2. 3-dimensional lattices

Uniform lattices in 3-dimensional real hyperbolic space can sometimes be deformed nontrivially in 4-dimensional real hyperbolic space (see [21, Chapter 6], or [2]). Nevertheless, when they act on quaternionic plane, all small deformations stabilize a quaternionic line, although the action on this line can be deformed nontrivially.

Theorem 1.2. *Let $\Gamma \subset \mathrm{Spin}(3, 1)^0$ be a lattice. Embed $\mathrm{Spin}(3, 1)^0$ into $\mathrm{Spin}(4, 1)^0 = \mathrm{Sp}(1, 1)$ and then into $\mathrm{Sp}(2, 1)$ in the obvious manner. This produces a discrete subgroup of $\mathrm{Sp}(2, 1)$ stabilizing a quaternionic line.*

If Γ is uniform in $\mathrm{Spin}(3, 1)^0$, then every small deformation of Γ in $\mathrm{Sp}(2, 1)$ again stabilizes a quaternionic line.

If Γ is not uniform in $\mathrm{Spin}(3, 1)^0$, then every small deformation of Γ preserving parabolics again stabilizes a quaternionic line.

If the assumption on parabolics is removed, nonuniform lattices in $\mathrm{Spin}(3, 1)^0$ can be deformed within $\mathrm{Spin}(3, 1)^0$ (see [21, Chapter 5]).

Question. Let Γ be a nonuniform lattice in $\mathrm{Spin}(3, 1)^0$. Map it to $\mathrm{Sp}(2, 1)$ via $\mathrm{Spin}(4, 1)^0 = \mathrm{Sp}(1, 1)$. Can one deform Γ to a Zariski dense subgroup?

1.3. 2-dimensional lattices

Uniform lattices in real hyperbolic plane, when mapped to $SU(2, 1)$ using the embedding $SO(2, 1) \rightarrow SU(2, 1)$, can be deformed to discrete Zariski dense subgroups of $SU(2, 1)$. On the other hand, lattices mapped via $SU(1, 1)$ and $SU(2, 1)$ are more rigid, as shown by W. Goldman and J. Millson [9]. This fact has been recently extended to higher rank groups by M. Burger, A. Iozzi and A. Wienhard [3].

It turns out that this form of rigidity of surface groups does not apply to the group $\mathrm{Sp}(2, 1)$.

Theorem 1.3. *Let Γ be the fundamental group of a closed surface of genus > 1 .*

- (1) *View Γ as a uniform lattice in $SO(2, 1)$. Map $SO(2, 1) \rightarrow \mathrm{Sp}(2, 1)$. This gives rise to a representation into $\mathrm{Sp}(2, 1)$ which can be deformed to a discrete Zariski dense representation.*
- (2) *View Γ as a uniform lattice in $SU(1, 1)$. Map $SU(1, 1) \rightarrow \mathrm{Sp}(1, 1) \rightarrow \mathrm{Sp}(2, 1)$. This gives rise to a representation into $\mathrm{Sp}(2, 1)$ fixing a quaternionic line. Then there exist small deformations which do not stabilize any quaternionic line.*

Whereas in the first case, explicit examples of deformations are provided by Thurston's bending construction, the existence of Zariski dense deformations in the second case follows from rather general principles. It would be interesting to visualize some of them.

1.4. Plan of the paper

Section 2 gives a cohomological criterion for non-Zariski dense subgroups to remain non-Zariski dense after deformation. The necessary cohomology vanishing is obtained in Section 3. Theorem 1.1 is proved in Section 4, Theorem 1.2 in Section 5. The statements for nonuniform lattices are proved in Section 6. Section 7 describes how lattices in Lie subgroups can sometimes be bent to become Zariski dense. The proof of Theorem 1.3 is completed in Section 8. We end with a remark on non-Zariski dense discrete subgroups in Section 9.

2. A relative Weil theorem

Let Γ be a finitely generated group, and G be a Lie group with Lie algebra \mathfrak{g} . The *character variety* $\chi(\Gamma, G)$ is the quotient of the space $\mathrm{Hom}(\Gamma, G)$ of homomorphisms of Γ to G by the action of G by postcomposing homomorphisms with inner automorphisms. In [23], A. Weil shows that a sufficient condition for a homomorphism $\rho : \Gamma \rightarrow G$ to define an isolated point in the character variety is that the first cohomology group $H^1(\Gamma, \mathfrak{g}_\rho)$ vanishes. In this section, we state a relative version of Weil's theorem.

Let $H \subset G$ be an algebraic subgroup of G . Let $\chi(\Gamma, H, G) \subset \chi(\Gamma, G)$ be the set of conjugacy classes of homomorphisms $\Gamma \rightarrow G$ which fall into conjugates of H . In other words, $\chi(\Gamma, H, G)$ is the set of G -orbits of elements of $\mathrm{Hom}(\Gamma, H) \subset \mathrm{Hom}(\Gamma, G)$. If $\rho \in \mathrm{Hom}(\Gamma, H)$, the representation $\mathfrak{g}_\rho = \mathrm{ad} \circ \rho$ on the Lie algebra \mathfrak{g} of G leaves the Lie algebra \mathfrak{h} of H invariant, and thus defines a quotient representation, which we shall denote by $\mathfrak{g}_\rho/\mathfrak{h}_\rho$.

Proposition 2.1. *Let $H \subset G$ be real Lie groups, with Lie algebras \mathfrak{h} and \mathfrak{g} . Let Γ be a finitely generated group. Let $\rho : \Gamma \rightarrow H$ be a homomorphism. Assume that $H^1(\Gamma, \mathfrak{g}_\rho/\mathfrak{h}_\rho) = 0$. Then $\chi(\Gamma, H, G)$ is a neighborhood of the G -conjugacy class of ρ in $\chi(\Gamma, G)$. In other words, homomorphisms $\Gamma \rightarrow G$ which are sufficiently close to ρ can be conjugated into H .*

Proof. $\text{Hom}(\Gamma, G)$ is topologized as a subset of the space G^Γ of arbitrary maps $\Gamma \rightarrow G$. Let $\Phi : G^\Gamma \rightarrow G^{\Gamma \times \Gamma}$ be the map which to a map $f : \Gamma \rightarrow G$ associates $\Phi(f) : \Gamma \times \Gamma \rightarrow G$ defined by

$$\Phi(f)(\gamma, \gamma') = f(\gamma\gamma'^{-1})f(\gamma)f(\gamma').$$

In other words, a map $f \in G^\Gamma$ is a homomorphism if and only if $\Phi(f) = 1$.

Consider the map $\Psi : G \times H^\Gamma \rightarrow G^\Gamma$ which sends $g \in G$ and $f : \Gamma \rightarrow H$ to the map $\Psi(g, f) : \Gamma \rightarrow G$ defined by

$$\Psi(g, f)(\gamma) = g^{-1}f(\gamma)g.$$

We need to prove that the image of Ψ contains a neighborhood of ρ in $\Phi^{-1}(1)$.

The cohomological assumption gives information on the differentials of Φ and Ψ . The differential $D_\rho \Phi$ is equal to $-d_1$ where d_1 denotes the coboundary $C^1(\Gamma, \mathfrak{g}_\rho) \rightarrow C^2(\Gamma, \mathfrak{g}_\rho)$. The differential of Ψ at $g = e$ and $f = \rho$ is given by

$$D_{(e,\rho)}\Psi(v, \eta) = -d_0v + \eta,$$

where d_0 denotes the coboundary $C^0(\Gamma, \mathfrak{g}_\rho) \rightarrow C^1(\Gamma, \mathfrak{g}_\rho)$. Since $\Phi(\Psi(g, f))(\gamma, \gamma') = g^{-1}\Phi(f)(\gamma, \gamma')g$ it follows that for all $f \in H^\Gamma$, $D_\rho \Phi \circ D_{(e,\rho)}\Psi = 0$. Conversely, if we assume that $H^1(\Gamma, \mathfrak{g}_\rho/\mathfrak{h}_\rho) = 0$, then any $\theta \in C^1(\Gamma, \mathfrak{g}_\rho)$ such that $D_\rho \Phi(\theta)$ takes values in the subalgebra \mathfrak{h} can be written $\theta = -d_0v + \eta$ where $v \in \mathfrak{g}$ and $\eta \in C^1(\Gamma, \mathfrak{h}_\rho)$, i.e. θ belongs to the image of $D_{(e,\rho)}\Psi$.

Clearly, $\text{Hom}(\Gamma, G)$ and $\text{Hom}(\Gamma, H)$ are real analytic varieties. To analyze a neighborhood of ρ in them, it is sufficient to analyze real analytic or even formal curves $t \mapsto \rho(t)$. In coordinates for G (in which H appears as a linear subspace), such a curve admits a Taylor expansion

$$\rho(t) = \sum_{n=0}^{\infty} a_n t^n,$$

where $a_0 = \rho$ and for $j \geq 1$, $a_j \in C^1(\Gamma, \mathfrak{g}_\rho)$ is a 1-cochain. Then $\Phi(\rho(t)) = 1$ for all t . Expanding this as a Taylor series gives

$$1 = \Phi(\rho) + D_\rho \Phi(a_1)t + (D_\rho \Phi(a_2) + D_\rho^2 \Phi(a_1, a_1))t^2 + \dots,$$

which implies that

$$D_\rho \Phi(a_1) = 0, \quad D_\rho \Phi(a_2) + D_\rho^2 \Phi(a_1, a_1) = 0, \quad \dots$$

The first equation says that a_1 is a cocycle. So is $a_1 \bmod \mathfrak{h}$, therefore there exist $v \in \mathfrak{g}$ and $b_1 \in Z^1(\Gamma, \mathfrak{h}_\rho)$ such that $a_1 = -d_0v + b_1$. Let $t \mapsto g(t)$ be an analytic curve in G with Taylor expansion $g(t) = 1 + vt + \dots$. Then the Taylor expansion of $\rho_1(t) = g(t)^{-1}\rho(t)g(t)$ takes the form $\rho_1(t) = 1 + b_1t + \dots$. In other words, up to conjugating, we arranged to bring the first term of the expansion of $\rho(t)$ into \mathfrak{h} .

The second equation now reads $D_\rho \Phi(a_2) + D_\rho^2 \Phi(b_1, b_1) = 0$. This implies that $D_\rho \Phi(a_2)$ takes its values in \mathfrak{h} . Therefore there exist $v' \in \mathfrak{g}$ and $b_2 \in Z^1(\Gamma, \mathfrak{h}_\rho)$ such

that $a_2 = -d_0v' + b_2$. Conjugating $\rho_1(t)$ by an analytic curve in G with Taylor expansion $1 + v't^2 + \dots$ kills v' and replaces a_2 with b_2 in the expansion of $\rho_1(t)$. Inductively, one can bring all terms of the expansion of $\rho(t)$ into \mathfrak{h} . The resulting curve belongs to $\text{Hom}(\Gamma, H)$. This shows that in a neighborhood of ρ , $\text{Hom}(\Gamma, G)$ coincides with $G^{-1} \text{Hom}(\Gamma, H)G$. Passing to the quotient, $\chi(\Gamma, H, G)$ coincides with $\chi(\Gamma, G)$ in a neighborhood of the conjugacy class of ρ . \square

3. A cohomology vanishing result

3.1. Preliminaries

For basic information on quaternionic hyperbolic space and surveys, see [11, 14, 19].

We regard \mathbb{H}^n as a right module over \mathbb{H} by right multiplication. Viewing $\mathbb{H} = \mathbb{C} \oplus j\mathbb{C} = \mathbb{C}^2$, left multiplication by \mathbb{H} gives \mathbb{C} -linear endomorphisms of \mathbb{C}^2 . So $\mathbb{H}^* = \text{GL}_1 \mathbb{H} \subset \text{GL}_2 \mathbb{C}$. Similarly $(x_1 + iy_1 + j(z_1 + iw_1), \dots, x_n + iy_n + j(z_n + iw_n))$ is identified with $(x_1 + iy_1, \dots, x_n + iy_n; z_1 + iw_1, \dots, z_n + iw_n)$ so that $\mathbb{H}^n = \mathbb{C}^{2n}$ and $\text{GL}_n \mathbb{H} \subset \text{GL}_{2n} \mathbb{C}$.

A \mathbb{C} -linear map $\phi : \mathbb{H}^n \rightarrow \mathbb{H}^n$ is \mathbb{H} -linear exactly when it commutes with j : $\phi(vj) = \phi(v)j$. Then it follows that

$$\text{GL}_n \mathbb{H} = \{A \in \text{GL}_{2n} \mathbb{C} : AJ = J\bar{A}\}, \quad \text{where } J = \begin{bmatrix} 0 & -I_n \\ I_n & 0 \end{bmatrix}.$$

Any element in $\text{GL}_n \mathbb{H}$ can be written as $\alpha + j\beta$ where α and β are $2n \times 2n$ complex matrices. If we write a vector in \mathbb{H}^n in the form $X + jY$ where $X, Y \in \mathbb{C}^n$, the action of $\alpha + j\beta$ on it is

$$\alpha X - \bar{\beta}Y + j(\bar{\alpha}Y + \beta X).$$

So a matrix $\alpha + j\beta$ in $\text{GL}_n \mathbb{H}$ corresponds to the matrix $\begin{bmatrix} \alpha & -\bar{\beta} \\ \beta & \bar{\alpha} \end{bmatrix}$ in $\text{GL}_{2n} \mathbb{C}$.

In this paper, we fix a quaternionic Hermitian form $\langle \cdot \rangle$ of signature $(n, 1)$ on \mathbb{H}^{n+1} as

$$\langle v, w \rangle = \sum_{i=1}^n \bar{v}_i w_i - \bar{v}_{n+1} w_{n+1}.$$

Then the Lie group $\text{Sp}(n, 1, \mathbb{H}) = \text{Sp}(n, 1)$, which is the set of matrices preserving this Hermitian form, is

$$\{A \in \text{GL}_{n+1} \mathbb{H} : A^* J' A = J'\}, \quad \text{where } J' = \begin{bmatrix} I_n & 0 \\ 0 & -1 \end{bmatrix}.$$

It is easy to see that its Lie algebra $\mathfrak{sp}(n, 1)$ is the set of matrices of the form

$$\begin{bmatrix} \text{Im } \mathbb{H} & Y \\ X & \mathfrak{sp}(n - 1, 1) \end{bmatrix},$$

where $Y + X^*J_n = 0$, $X, Y \in \mathbb{H}^n$ and $J_n = \begin{bmatrix} I_{n-1} & 0 \\ 0 & -1 \end{bmatrix}$. So we get

$$\mathfrak{sp}(n, 1) = \text{Im } \mathbb{H} \oplus \mathbb{H}^n \oplus \mathfrak{sp}(n - 1, 1).$$

Note that the adjoint action of the subgroup $\begin{bmatrix} \text{Sp}(1) & 0 \\ 0 & \text{Sp}(n-1,1) \end{bmatrix}$ preserves this decomposition. The action on the \mathbb{H}^n component is the standard action,

$$\text{Sp}(n - 1, 1)\mathbb{H}^n \text{Sp}(1)^{-1}.$$

Identifying \mathbb{H}^{n+1} with \mathbb{C}^{2n+2} as above, it is easy to see that $\text{Sp}(n, 1, \mathbb{H})$ is exactly equal to $U(2n, 2) \cap \text{Sp}(2n + 2, \mathbb{C})$, i.e. to the set of unitary matrices satisfying $AJ = J\bar{A}$. Indeed, the symplectic form with respect to the standard basis of \mathbb{C}^{2n+2} is $\begin{bmatrix} 0 & A \\ -A & 0 \end{bmatrix}$ and $A = \begin{bmatrix} I_n & 0 \\ 0 & -1 \end{bmatrix}$.

We will often complexify real Lie algebras. For any $M \in \mathfrak{gl}(2n, \mathbb{C})$, one can write

$$M = \frac{1}{2}(M - J\bar{M}J) - i\left(\frac{1}{2}(iM + iJ\bar{M}J)\right).$$

So it is easy to see that $\mathfrak{gl}(n, \mathbb{H}) = \{A \in \mathfrak{gl}(2n, \mathbb{C}) : AJ = J\bar{A}\}$ is complexified to $\mathfrak{gl}(2n, \mathbb{C})$. It is well known that $\mathfrak{u}(2n, 2) \otimes_{\mathbb{R}} \mathbb{C} = \mathfrak{gl}(2n + 2, \mathbb{C})$ and $\mathfrak{sp}(2n + 2, \mathbb{C}) \otimes_{\mathbb{R}} \mathbb{C} = \mathfrak{sp}(2n + 2, \mathbb{C}) \times \mathfrak{sp}(2n + 2, \mathbb{C})$. From these, we obtain

$$\mathfrak{sp}(n, 1) \otimes_{\mathbb{R}} \mathbb{C} = \mathfrak{sp}(2n + 2, \mathbb{C}).$$

We are particularly interested in

$$\mathfrak{sp}(1, 1) \otimes_{\mathbb{R}} \mathbb{C} = \mathfrak{sp}(4, \mathbb{C}).$$

The quaternionic hyperbolic n -space $H_{\mathbb{H}}^n$ in the unit ball model is

$$\left\{ (x_1, \dots, x_n) \mid x_i \in \mathbb{H}, \sum |x_i|^2 < 1 \right\}.$$

It can also be described as a hyperboloid model

$$\{X \in \mathbb{H}^{n+1} : \langle X, X \rangle = -1\} / \sim$$

where $X \sim Y$ iff $X = Y \text{Sp}(1)$. Then the isometry group of $H_{\mathbb{H}}^n$ is $P\text{Sp}(n, 1)$ which is a noncompact real semisimple Lie group.

A point X in the unit ball model can be mapped to $[X, 1]$ in the hyperboloid model. Then it is easy to see that the subgroup of the form

$$\begin{bmatrix} \text{Sp}(n - 1) & 0 \\ 0 & \text{Sp}(1, 1) \end{bmatrix}$$

stabilizes a quaternionic line $(0, \dots, 0, \mathbb{H})$ in the ball model. In fact, we have

Lemma 3.1. *The stabilizer of a quaternionic line $\{(0, \mathbb{H})\}$ in $\mathrm{Sp}(2, 1)$ is of the form*

$$\begin{bmatrix} \mathrm{Sp}(1) & 0 \\ 0 & \mathrm{Sp}(1, 1) \end{bmatrix}.$$

Furthermore, a parabolic element in $SO(4, 1) = \mathrm{Sp}(1, 1)$ stabilizing the quaternionic line $\{(0, \mathbb{H})\}$ is of the form in $P\mathrm{Sp}(2, 1)$

$$\begin{bmatrix} \mathrm{Sp}(1) & 0 \\ 0 & \begin{bmatrix} a & \lambda - a \\ a - \lambda & 2\lambda - a \end{bmatrix} \end{bmatrix}$$

where $a \geq 1$ is a positive real number, $\lambda \in \mathrm{Sp}(1)$ with $\mathrm{Re} \lambda = 1/a$. These elements constitute the parabolic elements in the center $\{(t, 0)\}$ of the Heisenberg group. A general parabolic element fixing a point $(0, 1)$ at infinity and not stabilizing the quaternionic line $\{(0, \mathbb{H})\}$ is of the form

$$\begin{bmatrix} * & x & -x \\ * & * & * \\ * & * & * \end{bmatrix}$$

with $x \neq 0$. These elements are parabolic elements which do not belong to the center of the Heisenberg group.

Proof. The quaternionic line $\{(0, \mathbb{H})\}$ in the hyperboloid model has coordinate $(0, \mathbb{H}, 1)$. To fix this line, it is not difficult to see that the matrix should have the form of $A = \begin{bmatrix} * & 0 & 0 \\ * & * & 0 \\ * & * & * \end{bmatrix}$. Since its inverse $J'A^*J'$ also fixes the quaternionic line, it should have the form as in the claim.

Now to prove the second claim, note that the matrix should satisfy the equation

$$\begin{bmatrix} \mathrm{Sp}(1) & 0 & 0 \\ 0 & a & b \\ 0 & c & d \end{bmatrix} (0, 1, 1) = \lambda(0, 1, 1) \quad \text{for } \lambda \in \mathrm{Sp}(1).$$

Also it should satisfy $A^*J'A = J'$. From these, we obtain

$$a + b = \lambda, \quad c + d = \lambda, \quad |a|^2 - |c|^2 = |d|^2 - |b|^2 = 1, \quad \bar{a}b - \bar{c}d = 0.$$

Then we get $\bar{a}(\lambda - a) - \bar{c}(\lambda - c) = 0$. So $(\bar{a} - \bar{c})\lambda = |a|^2 - |c|^2 = 1$, and we get $c = a - \lambda$. Now we divide A by a since a is nonzero. Note that Aa^{-1} represents the same element in $P\mathrm{Sp}(2, 1)$. Then we can assume that a is a positive real number, by conjugating A if necessary. The fact that $\mathrm{Re} \lambda = 1/a$ follows from the other two equations. So the result follows. In the Heisenberg group $\{(t, z) \mid t \in \mathrm{Im} \mathbb{H}, z \in \mathbb{H}\}$, the center $\{(t, 0)\}$ is the (ideal) boundary of the quaternionic line $\{(0, \mathbb{H})\}$. So these parabolic elements stabilizing the quaternionic line belong to the center. See [12].

To prove the last claim, we just note that $A(0, 1, 1) = \lambda(0, 1, 1)$ should be satisfied. The parabolic elements not stabilizing the quaternionic line $\{(0, \mathbb{H})\}$ should have nonzero x by the first case. □

3.2. Raghunathan's theorem

In this section we collect information concerning finite-dimensional representations of $\mathfrak{so}(5, \mathbb{C})$, which will be necessary for our main theorem. The basic theorem we will make use of is due to M. S. Raghunathan [20].

Theorem 3.2. *Let G be a connected semisimple Lie group. Let $\Gamma \subset G$ be a uniform irreducible lattice and $\rho : (\Gamma \subset G) \rightarrow \text{Aut}(E)$ a simple nontrivial linear representation. Then $H^1(\Gamma; E) = 0$ except possibly when $\mathfrak{g} = \mathfrak{so}(n+1, 1)$ (resp. $\mathfrak{g} = \mathfrak{su}(n, 1)$) and the highest weight of ρ is a multiple of the highest weight of the standard representation of $\mathfrak{so}(n+1, 1)$ (resp. of the standard representation of $\mathfrak{su}(n, 1)$ or of its contragredient representation).*

In this theorem, Raghunathan used Matsushima–Murakami's result where L^2 -cohomology is applied. We observe that as long as we use L^2 -cohomology, this theorem still holds for non-uniform lattices. This issue will be dealt with in Section 6.

3.3. Standard representation of $\mathfrak{sp}(4, \mathbb{C})$

In the previous section, we used the symplectic form with respect to the standard basis of \mathbb{C}^4

$$Q = \begin{bmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & -1 \\ 0 & 1 & 0 & 0 \end{bmatrix}.$$

Then the Lie algebra $\mathfrak{sp}(4, \mathbb{C})$ consists of the complex matrices $\begin{bmatrix} A & B \\ C & D \end{bmatrix}$ such that

$$A^t \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} D = 0,$$

$$C^t \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} C = 0,$$

$$B^t \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} + \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} B = 0.$$

An obvious choice of a Cartan subalgebra \mathfrak{h} is

$$\begin{bmatrix} x & 0 & 0 & 0 \\ 0 & y & 0 & 0 \\ 0 & -x & 0 & 0 \\ 0 & 0 & 0 & -y \end{bmatrix}.$$

Let $L_1, L_2 \in \mathfrak{h}^*$ be defined by $L_1(x, y) = x$, $L_2(x, y) = y$. Then the natural action of $\mathfrak{sp}(4, \mathbb{C})$ on \mathbb{C}^4 has the four standard basis vectors e_1, e_2, e_3, e_4 as eigenvectors with weights $L_1, L_2, -L_1, -L_2$. The highest weight is L_1 .

3.4. Representation of $\mathfrak{so}(5, \mathbb{C})$

We shall use the isomorphism of $\mathfrak{sp}(4, \mathbb{C})$ to $\mathfrak{so}(5, \mathbb{C})$. It arises from the following geometric construction.

Let $V = \mathbb{C}^4$ and ω be the symplectic form defined as before. Then

$$\bigwedge^2 V^* \otimes \bigwedge^2 V^* \rightarrow \mathbb{C}, \quad \alpha \otimes \beta \mapsto \frac{\alpha \wedge \beta}{\omega \wedge \omega},$$

is a nondegenerate quadratic form P on $\bigwedge^2 V^*$. Here since both $\alpha \wedge \beta$ and $\omega \wedge \omega$ are 4-forms, there is a constant c so that $\alpha \wedge \beta = c\omega \wedge \omega$, so the quotient should be understood as such a constant. Take the orthogonal complement W of $\mathbb{C}\omega$ with respect to this quadratic form. Any matrix A acts on 2-forms as follows: $A\alpha(v, w) = \alpha(Av, Aw)$. Then $\mathrm{Sp}(4, \mathbb{C})$ leaves W invariant and acts orthogonally on it. This gives a map from $\mathrm{Sp}(4, \mathbb{C})$ to $SO(5, \mathbb{C}) = SO(W)$, which turns out to be an isomorphism.

Next, we relate the choice of Cartan subalgebra for $\mathfrak{sp}(4, \mathbb{C})$ made in the preceding paragraph to the standard choice for $\mathfrak{so}(5, \mathbb{C})$.

We first compute the Lie algebra isomorphism derived from the group isomorphism.

Let z_1, z_2, z_3, z_4 be standard coordinates of \mathbb{C}^4 so that $dz_1 \wedge dz_3 + dz_4 \wedge dz_2 = \omega$. Let

$$\begin{aligned} \omega_6 &= \omega, & \omega_1 &= i(dz_1 \wedge dz_4 - dz_2 \wedge dz_3), \\ \omega_5 &= dz_1 \wedge dz_2 + dz_3 \wedge dz_4, & \omega_2 &= i(dz_1 \wedge dz_2 - dz_3 \wedge dz_4), \\ \omega_4 &= dz_1 \wedge dz_4 + dz_2 \wedge dz_3, & \omega_3 &= i(dz_1 \wedge dz_3 - dz_4 \wedge dz_2). \end{aligned}$$

This is an orthonormal basis of $\bigwedge^2 V^*$.

Let $A_t \in \mathrm{Sp}(4, \mathbb{C})$ so that $A_0 = I$ and $\frac{d}{dt}|_{t=0} A_t = X \in \mathfrak{sp}(4, \mathbb{C})$. Then for 1-forms α, β , one can figure out the action of X on 2-forms: $X(\alpha \otimes \beta) = \frac{d}{dt}|_{t=0} A_t(\alpha \otimes \beta) = (X\alpha) \otimes \beta + \alpha \otimes (X\beta)$. Then

$$X(\alpha \wedge \beta) = (X\alpha) \wedge \beta + \alpha \wedge (X\beta).$$

To make computation easier, we choose a basis of W as

$$v_1 = \frac{\omega_1 + i\omega_4}{\sqrt{2}}, \quad v_3 = \frac{\omega_1 - i\omega_4}{\sqrt{2}}, \quad v_2 = \frac{\omega_2 + i\omega_5}{\sqrt{2}}, \quad v_4 = \frac{\omega_2 - i\omega_5}{\sqrt{2}}, \quad v_5 = \omega_3.$$

With respect to this basis, the symmetric bilinear form P has $P(v_1, v_3) = 1 = P(v_2, v_4) = P(v_5, v_5)$ and $P(v_i, v_j) = 0$ for all other pairs. With respect to this P , one can easily see that a Cartan subalgebra of $\mathfrak{so}(5, \mathbb{C}) = \mathfrak{so}(W; P)$ can be chosen as the set of matrices of the form

$$\begin{bmatrix} x & 0 & 0 & 0 & 0 \\ 0 & y & 0 & 0 & 0 \\ 0 & 0 & -x & 0 & 0 \\ 0 & 0 & 0 & -y & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

Let (x, y, z, w) denote a diagonal matrix in $\mathfrak{sp}(4, \mathbb{C})$. Then one can easily compute that

$$(1, 0, -1, 0)v_1 = v_1, \quad (1, 0, -1, 0)v_3 = -v_3, \\ (1, 0, -1, 0)v_2 = v_2, \quad (1, 0, -1, 0)v_4 = -v_4, \quad (1, 0, -1, 0)v_5 = 0.$$

Similarly

$$(0, 1, 0, -1)v_1 = -v_1, \quad (0, 1, 0, -1)v_3 = v_3, \\ (0, 1, 0, -1)v_2 = v_2, \quad (0, 1, 0, -1)v_4 = -v_4, \quad (0, 1, 0, -1)v_5 = 0.$$

So the element $\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$ in a Cartan subalgebra of $\mathfrak{sp}(4, \mathbb{C})$ corresponds to an element in a Cartan subalgebra of $\mathfrak{so}(5, \mathbb{C})$,

$$\mathfrak{h}_1 = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

Similarly

$$\begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix} \text{ corresponds to } \mathfrak{h}_2 = \begin{bmatrix} -1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

This representation under the isomorphism to $\mathfrak{sp}(4, \mathbb{C})$ is different from the standard representation of $\mathfrak{so}(5, \mathbb{C})$ on \mathbb{C}^5 as we will see below.

Lemma 3.3. *The highest weight of the standard representation of $\mathfrak{so}(5, \mathbb{C})$ on \mathbb{C}^5 is not a multiple of the highest weight of the representation coming from $\mathfrak{sp}(4, \mathbb{C})$ on \mathbb{C}^4 .*

Proof. With respect to the symmetric bilinear form P as before, a Cartan subalgebra of $\mathfrak{so}(5, \mathbb{C})$ is the set of diagonal matrices $(x, y, -x, -y, 0)$ as noted above. Then the standard representation of $\mathfrak{so}(5, \mathbb{C})$ on \mathbb{C}^5 has eigenvectors, the standard basis e_1, e_2, e_3, e_4, e_5 , with eigenvalues $L_1, L_2, -L_1, -L_2, 0$. This has the highest weight L_1 .

The standard representation of $\mathfrak{sp}(4, \mathbb{C})$ on \mathbb{C}^4 has the highest weight L_1 as we saw in the previous section. Note that the Cartan subalgebra of $\mathfrak{sp}(4, \mathbb{C})$ is generated by the diagonal matrices $(1, 0, -1, 0)$ and $(0, 1, 0, -1)$ with dual basis L_1 and L_2 . Then under the isomorphism from $\mathfrak{sp}(4, \mathbb{C})$ to $\mathfrak{so}(\mathbb{C}\omega^\perp)$, these two diagonal matrices are mapped to the diagonal matrices $\mathfrak{h}_1 = (1, 1, -1, -1, 0)$ and $\mathfrak{h}_2 = (-1, 1, 1, -1, 0)$. Let L'_1, L'_2 be the images of L_1, L_2 under this isomorphism. Then in terms of the standard dual basis L_1, L_2 of the Cartan subalgebra of $\mathfrak{so}(5, \mathbb{C})$,

$$L'_1 = \frac{L_1 + L_2}{2}, \quad L'_2 = \frac{L_2 - L_1}{2}.$$

So the representation coming from the standard representation of $\mathfrak{sp}(4, \mathbb{C})$ on \mathbb{C}^4 has highest weight $\frac{L_1+L_2}{2}$. Actually this is the highest weight of the spin representation. \square

Corollary 3.4. *Let $\Gamma \subset \mathrm{Sp}(1, 1)$ be a uniform lattice. Then $H^1(\Gamma, \mathbb{H}^2) = 0$ where \mathbb{H}^2 denotes the standard representation of $\mathrm{Sp}(1, 1)$ restricted to Γ .*

Proof. View \mathbb{H}^2 as \mathbb{C}^4 with $\mathrm{Sp}(1, 1)$ acting on it. If we complexify the real Lie algebra $\mathfrak{sp}(1, 1)$, we get $\mathfrak{sp}(4, \mathbb{C})$. Since the standard representation of $\mathfrak{sp}(4, \mathbb{C})$ on \mathbb{C}^4 is different from the standard representation of $\mathfrak{so}(5, \mathbb{C})$ on \mathbb{C}^5 with highest weight L_1 , Theorem 3.2 (Theorem 1 of Raghunathan [20]) applies, and $H^1(\Gamma, \mathbb{H}^2) = 0$. \square

4. Proof of Theorem 1.1 (uniform case)

Let $\Gamma \subset \mathrm{Sp}(1, 1)$ be a uniform lattice. Denote by ρ the embedding $\Gamma \rightarrow \mathrm{Sp}(1, 1) \rightarrow \mathrm{Sp}(2, 1)$. Let $G = \mathrm{Sp}(2, 1)$, $H = \left[\begin{smallmatrix} \mathrm{Sp}(1) & 0 & \\ 0 & \mathrm{Sp}(1,1) & \end{smallmatrix} \right] \subset G$. As was seen in Section 3.1, the adjoint representation of G restricted to H splits as a direct sum $\mathfrak{sp}(2, 1) = \mathfrak{sp}(1) \oplus \mathbb{H}^2 \oplus \mathfrak{sp}(1, 1)$, thus $\mathfrak{g}/\mathfrak{h} = \mathbb{H}^2$, restricted to $\mathrm{Sp}(1, 1)$, is the standard representation of $\mathrm{Sp}(1, 1)$. Corollary 3.4 asserts that $H^1(\Gamma, \mathbb{H}^2)$ vanishes. Therefore $H^1(\Gamma, \mathfrak{g}_\rho/\mathfrak{h}_\rho) = 0$. According to Proposition 2.1, this implies that homomorphisms $\Gamma \rightarrow \mathrm{Sp}(2, 1)$ which are close enough to ρ can be conjugated into H , i.e. leave a quaternionic line invariant.

Since the subgroup of the form

$$\left[\begin{smallmatrix} \mathrm{Sp}(1) & 0 & 0 \\ 0 & I & 0 \\ 0 & 0 & \mathrm{Sp}(1, 1) \end{smallmatrix} \right]$$

stabilizes a quaternionic line $(0, \dots, 0, \mathbb{H})$ in the ball model, we obtain

Corollary 4.1. *Let $\Gamma \subset \mathrm{Sp}(1, 1)$ be a uniform lattice. Embed Γ into $\mathrm{Sp}(n, 1)$ as a subgroup which stabilizes a quaternionic line. Then every small deformation of Γ in $\mathrm{Sp}(n, 1)$ stabilizes a quaternionic line.*

5. 3-manifold case

In this section, we prove Theorem 1.2 for uniform 3-dimensional hyperbolic lattices. Let $\Gamma \subset \mathrm{Spin}(3, 1)^0$ be a uniform lattice. According to Proposition 2.1, local deformations of the standard representation $\rho_0 : \Gamma \rightarrow \mathrm{Spin}(3, 1)^0 \rightarrow \mathrm{Spin}(4, 1)^0 = \mathrm{Sp}(1, 1) \rightarrow \mathrm{Sp}(2, 1)$ which do not stabilize a quaternionic line are encoded in $H^1(\Gamma, \mathbb{H}^2)$. We want to show that this first cohomology is zero. The complexified Lie algebra of $SO(3, 1)$ is $\mathfrak{so}(4, \mathbb{C})$. In the notations of Section 3, the symmetric bilinear form P has a basis v_1, v_2, v_3, v_4 such that $P(v_1, v_3) = P(v_2, v_4) = 1$ and $P(v_i, v_j) = 0$ for all other pairs. The Cartan subalgebra of $\mathfrak{so}(4, \mathbb{C})$ is the set of diagonal matrices $(x, y, -x, -y)$. Then as in Lemma 3.3, the standard representation of $\mathfrak{so}(4, \mathbb{C})$ on \mathbb{C}^4 has a character which is not a multiple of the character of the representation coming from $\mathfrak{so}(4, \mathbb{C}) \subset \mathfrak{sp}(4, \mathbb{C})$. Then by Raghunathan’s Theorem 3.2, $H^1(\Gamma, \mathbb{H}^2) = 0$. Proposition 2.1 ensures that neighboring homomorphisms $\Gamma \rightarrow \mathrm{Sp}(2, 1)$ stabilize a quaternionic line.

6. Nonuniform lattices

We used Raghunathan’s theorem [20] to prove our main theorem when Γ is a uniform lattice. In this section we discuss how it generalizes, with restrictions, to nonuniform lattices.

The key point is whether Matsushima–Murakami’s vanishing theorem that Raghunathan used still holds in the nonuniform case. To apply Matsushima–Murakami’s theorem, one has to use L^2 -cohomology.

Recall that under the subgroup $\begin{bmatrix} \text{Sp}(1) & 0 \\ 0 & \text{Sp}(1,1) \end{bmatrix}$, the adjoint representation of $\text{Sp}(2, 1)$ splits as a direct sum $\mathfrak{sp}(2, 1) = \mathfrak{sp}(1) \oplus \mathbb{H}^2 \oplus \mathfrak{sp}(1, 1)$. Let ρ denote the representation of $\text{Sp}(1, 1)$ corresponding to the \mathbb{H}^2 summand. Let $M = H_{\mathbb{R}}^4 / \Gamma$ be a finite volume manifold. View Γ as a subgroup of $\text{Sp}(1, 1)$, and denote by ρ_0 the restriction of ρ to Γ . Let E be the associated flat bundle over M with fiber \mathbb{H}^2 . It is well known that

$$H^1(\Gamma, \rho_0) = H_{\text{dR}}^1(M, E)$$

where $H_{\text{dR}}^1(M, E)$ is the de Rham cohomology of smooth E -valued differential forms over M . We will denote this de Rham cohomology by $H^1(M, E)$.

In Matsushima–Murakami’s proof, specific metrics on fibers of E , depending on base points, are used. More precisely, fix a maximal compact subgroup K of $\text{Sp}(1, 1)$. Let $\mathfrak{sp}(1, 1) = \mathfrak{t} \oplus \mathfrak{p}$ be the corresponding Cartan decomposition. Fix a positive definite metric $\langle \cdot, \cdot \rangle_F$ on \mathbb{H}^2 so that $\rho(K)$ is unitary and $\rho(\mathfrak{p})$ is hermitian symmetric. Then, for two elements v, w in the fiber over a point $g \in G$, one defines

$$\langle v, w \rangle = \langle \rho(g)^{-1}v, \rho(g)^{-1}w \rangle_F.$$

Here is a concrete construction of such a metric on \mathbb{H}^2 . As before, $\mathbb{H}^{1,1} = \mathbb{H}^2$ is equipped with the signature $(1, 1)$ metric

$$Q = |q_1|^2 - |q_2|^2.$$

Then for each negative \mathbb{H} -line L in $\mathbb{H}^{1,1}$, there exists a positive definite \mathbb{H} -Hermitian metric defined by $-Q|_L \oplus Q|_{L^\perp}$ where L^\perp is the orthogonal complement of L with respect to Q .

A unit speed ray in $H_{\mathbb{R}}^4 = H_{\mathbb{H}}^1$ in terms of $\mathbb{H}^{1,1}$ coordinates can be written as $l_t = \{q_1 = \delta_t q_2\}$ where $\delta_t = (e^t - 1)/(e^t + 1)$, $0 \leq t \leq \infty$. Note that here we normalize the metric so that its sectional curvature is -1 . This can be easily computed considering a unit speed ray $r(t)$ in a ball model emanating from the origin, and $r(t)$ corresponds to the point $(r(t), 1)$ in the hyperboloid model.

Now we want to know how the metric varies along l_t as $t \rightarrow \infty$. Let $v = (v_1, v_2) \in \mathbb{H}^{1,1}$. It is easy to see that

$$b_t = \left(\frac{1}{\sqrt{1 - \delta_t^2}}, \frac{\delta_t}{\sqrt{1 - \delta_t^2}} \right), \quad a_t = \left(\frac{\delta_t}{\sqrt{1 - \delta_t^2}}, \frac{1}{\sqrt{1 - \delta_t^2}} \right)$$

are unit vectors on l_t^\perp, l_t respectively. Then the l_t component and l_t^\perp component of v are respectively

$$\left(\frac{\delta_t v_2 + \delta_t^2 v_1}{1 - \delta_t^2}, \frac{v_2 + \delta_t v_1}{1 - \delta_t^2} \right), \quad \left(\frac{\delta_t v_2 + v_1}{1 - \delta_t^2}, \frac{\delta_t^2 v_2 + \delta_t v_1}{1 - \delta_t^2} \right).$$

Hence it is easy to calculate the square of the length of v on l_t , which is

$$\frac{1 + \delta_t^2}{1 - \delta_t^2} [|v_1|^2 + |v_2|^2] + 2 \frac{\delta_t}{1 - \delta_t^2} (v_1 \bar{v}_2 + v_2 \bar{v}_1) = \frac{2\delta_t}{1 - \delta_t^2} |v_1 + v_2|^2 + \frac{1 - \delta_t}{1 + \delta_t} (|v_1|^2 + |v_2|^2).$$

In conclusion, the square of the length of v grows like $e^t |v|^2$ along the ray l_t in general. But for $v_1 + v_2 = 0$, it grows like $e^{-t} |v|^2$ along the ray. This is the case when the deformation consists in parabolic elements fixing a point $(0, 1)$ (in the ball model) and not stabilizing the quaternionic line $\{(0, \mathbb{H})\}$. See Lemma 3.1. These estimates will be used below.

Let $M = M_{\geq \epsilon} \cup M_{\leq \eta}$ be the thick-thin decomposition of M so that $\eta > \epsilon$ and $M_{\leq \eta}$ is a standard cusp part of M . Assume for simplicity that the cuspidal part is connected. It is well known that $M_{\leq \eta}$ is homeomorphic to $T \times \mathbb{R}^+$ with $ds^2 = e^{-2r} ds_T^2 + dr^2$ where T is a flat closed 3-manifold, r denotes distance from $T \times \{0\}$, and $M_{\geq \epsilon} \cap M_{\leq \eta}$ is $T \times [0, 1]$.

Let $\pi : T \times \mathbb{R}^+ \rightarrow T$ be the projection on the first factor. Since $H^k(T) = H^k(M_{\leq \eta})$ via π^* , we want to deduce that $L^2 H^k(M_{\leq \eta}) = H^k(T)$, to show that $H^k(M_{\leq \eta}) = L^2 H^k(M_{\leq \eta})$. Let α be a k -form on T . Then $|\pi^* \alpha| \sim e^{r/2} |\alpha| e^{kr}$ where r is the distance from the boundary of the thin part. Here $e^{r/2}$ comes from the fiber metric and e^{kr} comes from the base metric. Then

$$\|\pi^* \alpha\|_{L^2}^2 = \int |\alpha|^2 e^{2kr+r} e^{-3r} ds_T dr \leq \|\alpha\|_{L^2(T)}^2 \times C < \infty$$

if $2k + 1 < 3$. So the pull-back form $\pi^* \alpha$ is always an L^2 -form on $M_{\leq \eta}$ if α is a 0-form. So we have obtained

Lemma 6.1. *For a finite volume real 4-dimensional hyperbolic manifold M ,*

$$H^0(M_{\leq \eta}, E) = L^2 H^0(M_{\leq \eta}, E).$$

Proof. For any $\alpha \in H^*(T, E) = H^*(M_{\leq \eta}, E)$, its pull-back $\pi^* \alpha$ is an L^2 -form on $M_{\leq \eta}$ for $*$ = 0 as noted above. So any element in $H^0(M_{\leq \eta}, E)$ has an L^2 -representative. \square

Unfortunately, we cannot conclude that $H^1(M, E) = L^2 H^1(M, E)$. This hinders a generalization of our theorem to nonuniform lattices. Our generalization involves a restriction on the representation.

Proposition 6.2. *Let M be a finite volume hyperbolic 3-manifold such that $M = H_{\mathbb{R}}^3 / \Gamma$. Then all small deformations of $\Gamma \subset SO(3, 1) \subset Sp(1, 1)$ preserving parabolicity still stabilize a quaternionic line. The same thing holds for a finite volume hyperbolic 4-manifold.*

Proof. We give a proof only in dimension 3, since the 4-dimensional case can be obtained by the same method. Since M has finite volume, its boundary consists of tori T_i . Let $\rho_0 : \pi_1(M) \rightarrow \text{Spin}(3, 1)^0 \subset \text{Sp}(1, 1) \subset \text{Sp}(2, 1)$ be a natural representation.

If $\rho_t(\pi_1(\partial M))$ is parabolic for all small t , then by Lemma 3.1, it can contribute to the \mathbb{H}^2 summand of $\mathfrak{sp}(2, 1)$. But in this case, it can be represented by an L^2 -form. The argument goes briefly as follows.

Let $\rho_t : \pi_1(M) \rightarrow \text{Sp}(2, 1)$ be a one-parameter family of deformations so that $\rho_t(\pi_1(\partial M))$ is all parabolic. Let N be the ϵ -thick part of M . Then ∂N consists of tori and the universal cover of it in $H_{\mathbb{R}}^3$ are horospheres. Fix a component of ∂N which is a horosphere H corresponding to a component T of ∂N . Conjugating ρ_t by g_t which depends smoothly on t if necessary, we may assume that $\rho_t(\pi_1(T))$ leaves invariant a common horosphere H' in $H_{\mathbb{H}}^2$. Such a choice of g_t is possible by the following argument. Let a be an element in $\pi_1(T)$ such that all $\rho_t(a)$ are parabolic. The subset P of $\text{Sp}(2, 1)$ consisting of the parabolic elements is a smooth manifold at $\rho_0(a)$, and the map from P to $\partial H_{\mathbb{H}}^2$ associating to each element in P its unique fixed point is smooth in a neighborhood of $\rho_0(a)$.

We may assume that H' is based at $(0, 1)$ (in the ball model). Then by Lemma 3.1, the contribution of this deformation to the \mathbb{H}^2 summand is contained in the subset $\{(x, y) \mid x + y = 0\} \subset \mathbb{H}^2$. This will help us out.

Let ω be a differential form representing the infinitesimal deformation $\frac{d}{dt}\rho_t$ on this cusp. Since $\rho_t(\pi_1(T))$ fixes $(0, 1)$, ω takes its values in the subalgebra $\mathfrak{s} \subset \mathfrak{sp}(2, 1)$ of Killing fields on $H_{\mathbb{H}}^2$ which vanish at $(0, 1)$ and which are tangent to the horospheres centered at $(0, 1)$. Therefore the norm of vectors of \mathfrak{s} decays along a geodesic pointing to $(0, 1)$, with rate controlled by the maximal sectional curvature (in our case, which is the direction away from a quaternionic line, $-1/4$). In our situation, we are only concerned with the subspace $\{(v_1, v_2) \mid v_1 + v_2 = 0\} \subset \mathbb{H}^2$. So along the ray the squared norm decays like $e^{-r}|v|^2$ asymptotically.

Then integrating along a geodesic ray, we see that the 1-form ω defined on the cusp is in L^2 on the cusp. In more detail, let the cusp be $T \times [0, \infty)$ with coordinates (x, y, r) , and the metric $ds^2 = e^{-2r}ds_T^2 + dr^2$. Then the volume form on this cusp is $e^{-2r}dS_T dr$. Note that we take a metric on $H_{\mathbb{R}}^3$ whose sectional curvature is -1 . Then along $[0, \infty)$, the orthonormal basis is $\{e^r \frac{\partial}{\partial x}, e^r \frac{\partial}{\partial y}, \frac{\partial}{\partial r}\}$. Then at (x, y, r) , the norm of ω is

$$\left| \omega \left(e^r \frac{\partial}{\partial x} \right) \right|^2 + \left| \omega \left(e^r \frac{\partial}{\partial y} \right) \right|^2$$

since $\omega(\partial/\partial r) = 0$. So

$$\int_{T \times [0, \infty)} \|\omega\|^2 d\text{Vol} = \int_0^\infty e^{-r} e^{2r} e^{-2r} \int_T \|\omega_T\|^2 dS_T dr < \infty$$

where e^{-r} comes from the norm decay on $\{(v_1, v_2) \mid v_1 + v_2 = 0\}$, and e^{2r} comes from the decay of the metric on $H_{\mathbb{R}}^3$ along the ray (one should take an orthonormal basis $\{e^r \frac{\partial}{\partial x}, e^r \frac{\partial}{\partial y}, \frac{\partial}{\partial r}\}$ along the ray).

We do this for each cusp of M . Let ω_i be a 1-form which is an L^2 -representative of the deformation $\frac{d}{dt}\rho_t$ on the i -th cusp of M . Let α be a global 1-form representing the deformation $\frac{d}{dt}\rho_t$. Then

$$\omega_i = \alpha + d\phi_i$$

where ϕ_i is a function defined on the i -th cusp. Let ϕ be the union of the ϕ_i , and ξ be a smooth function so that $\xi = 1$ on cusps and 0 outside cusps. Let

$$\omega' = \alpha + d(\xi\phi) = \alpha + \phi d\xi + \xi d\phi.$$

Then on each cusp, $\omega' = \alpha + d\phi_i = \omega_i$. Thus ω' is in L^2 and $[\omega'] = [\alpha]$.

Now again we can use Matsushima–Murakami's result for this case. See [15, 16] for a similar argument in complex hyperbolic space.

So we have proved the theorem. \square

We wonder whether the theorem holds without the assumption of preserving parabolicity.

7. Bending representations

Let G be an algebraic group. The Zariski closure of a subgroup H of $G(\mathbb{R})$ is denoted by \bar{H} .

Let X be a compact orientable hyperbolic n -manifold which splits into two submanifolds with totally geodesic boundary V and W , exchanged by an involution that fixes their common boundary. Such manifolds exist in all dimensions [17]. Then $\Gamma = \pi_1(X)$ splits as an amalgamated sum $\Gamma = A \star_C B$ where $A = \pi_1(V)$, $B = \pi_1(W)$ and $C = \pi_1(\partial V)$. Here, $\bar{A} = \bar{B} = PO(n, 1)^0$ and $\bar{C} = PO(n-1, 1)^0$.

Now embed $PO(n, 1)^0$ into a larger group G . Let c belong to the centralizer $Z_G(C)$. Consider the subgroup $\Gamma_c = A \star_C cBc^{-1}$. When c is chosen along a curve in $Z_G(C)$, one obtains a special case of W. Thurston's *bending deformation* [21, Chapter 6]. In this section, we analyze the Zariski closure of Γ_c in case $G = PSp(m, 1)$ is the isometry group of m -dimensional quaternionic hyperbolic space, $m \geq n$ and $PO(n, 1)^0 \rightarrow PSp(n, 1) \rightarrow PSp(m, 1)$ in the obvious manner.

7.1. The first bending step

We find it convenient to use a geometric language, and establish a dictionary between subgroups of $G = PSp(m, 1)$ and totally geodesic subspaces of $X = H_{\mathbb{H}}^m$.

Lemma 7.1. *The subgroup of G that leaves $Y = H_{\mathbb{R}}^n \subset X$ invariant is the normalizer of $H = PO(n, 1)^0$ in G .*

Proof. If $aHa^{-1} = H$, then a maps the orbit Y of H to itself. Conversely, Y is the only orbit of H in X which is totally geodesic. If $a \in G$ normalizes H , then a maps Y to itself. \square

Second, let us determine the space of available parameters for bending, i.e. elements which commute with C .

Lemma 7.2. *Let $m \geq n \geq 2$. Let $L = PO(n-1, 1)^0 \subset PO(n, 1)^0 \subset PSp(n, 1) \subset PSp(m, 1) = G$. Let $C \subset L$ be a Zariski dense subgroup. Then the centralizer $Z_G(C)$ consists of isometries which fix $P = H_{\mathbb{R}}^{n-1}$ pointwise. As a matrix group, $Z_G(C) = Sp(m-n+1)Sp(1)$.*

Proof. Clearly, $Z_G(C) = Z_G(L)$. We know that L stabilizes the totally geodesic subspace $P = H_{\mathbb{R}}^{n-1}$ of the symmetric space $X = H_{\mathbb{H}}^m$ of G . If $a \in G$ centralizes L , then a normalizes it, thus it maps P to itself, by Lemma 7.1. Furthermore, the restriction of a to P belongs to the center of $\text{Isom}(P) = L$, thus is trivial. In other words, a fixes each point of P . Conversely, isometries of X which fix every point of P centralize L and thus C . Indeed, L is generated by geodesic symmetries with respect to points of P , and these commute with isometries fixing P . To get the matrix expression of $Z_G(C)$, view X as a subset of quaternionic projective m -space. Then for every vector $y \in \mathbb{R}^n$, extended with zero entries to give a vector in \mathbb{R}^{m+1} , there exists a quaternion $q(y)$ such that $a(y) = yq(y)$. This implies that a lifted as a matrix in $Sp(m, 1)$ is block diagonal,

$$a = \begin{bmatrix} qI_n & 0 \\ 0 & D \end{bmatrix},$$

with blocks of sizes n and $m-n+1$ respectively, $q \in Sp(1)$ and $D \in Sp(m-n+1)$. This product group maps to a subgroup of $PSp(m, 1)$ which is traditionally denoted by $Sp(m-n+1)Sp(1)$. \square

The dictionary continues with a correspondence between Zariski closures in simple groups and totally geodesic hulls in symmetric spaces.

Lemma 7.3. *Let Y_1, \dots, Y_k be totally geodesic subspaces of a symmetric space X . Then $\text{Isom}(Y_j)$ naturally embeds into $G = \text{Isom}(X)$. Furthermore, the Zariski closure of $\bigcup_j \text{Isom}(Y_j)$ equals $\text{Isom}(Z)$ where Z is the smallest totally geodesic subspace of X containing $\bigcup_j Y_j$.*

Proof. For $x \in X$, let ι_x denote the geodesic symmetry through x . Since X is symmetric, ι_x is an isometry. Such involutions generate $\text{Isom}(X)$. If $Y \subset X$ is totally geodesic, then Y is invariant under all ι_y , $y \in Y$. Therefore Y is again a symmetric space, with isometry group generated by the restrictions to Y of the ι_y . In particular, the subgroup of G generated by the ι_y , $y \in Y$, is isomorphic to $\text{Isom}(Y)$.

If γ is a geodesic joining points $x \in Y_i$ and $y \in Y_j$, then ι_x and ι_y leave γ invariant. Their restrictions to γ generate an infinite dyadic group. The Zariski closure of this group contains all ι_z where $z \in \gamma$. Therefore the Zariski closure of $\text{Isom}(Y_i) \cup \text{Isom}(Y_j)$ contains ι_z for all z belonging to the union of all geodesics intersecting both Y_i and Y_j . Since the totally geodesic closure Z is obtained by iterating this operation, one concludes that the Zariski closure of $\bigcup_j \text{Isom}(Y_j)$ contains $\text{Isom}(Z)$. Conversely, since $\text{Isom}(Z)$ is an algebraic subgroup in G , it is contained in the Zariski closure. \square

Lemma 7.4. *Let $Y = H_{\mathbb{R}}^n \subset H_{\mathbb{H}}^n = X$. Let Z be a totally geodesic subspace of X such that $Y \subsetneq Z \subsetneq X$. Assume that Z contains $a(Y)$ where $a \in G$ fixes pointwise a hyperplane P of Y but does not leave Y invariant. Then there is an isometry of X fixing Y pointwise and mapping Z to $H_{\mathbb{C}}^n$.*

Proof. View the restriction of TX to Y as a vector bundle with connection ∇ on Y . Then $TZ|_Y$ is a parallel subbundle, therefore, for $y \in Y$, T_yZ is invariant under the holonomy representation $\text{Hol}(\nabla, y)$, which we now describe.

View Y as a sheet of the hyperboloid in \mathbb{R}^{n+1} . Then a point y represents a unit vector, still denoted by y , in \mathbb{R}^{n+1} . View X as a subset of quaternionic projective space. Then the point y also represents the quaternionic line $\mathbb{H}y$ it generates. Such lines form the tautological quaternionic line bundle τ over X , a subbundle of the trivial bundle \mathbb{H}^{n+1} equipped with the orthogonally projected connection. As a connected vector bundle, $TX = \text{Hom}_{\mathbb{H}}(\tau, \tau^\perp)$. When restricted to Y , τ comes with the parallel section y . Therefore $TX|_Y = \tau^\perp = TY \otimes \mathbb{H}$. In other words, $TX|_Y$ splits as a direct sum of four parallel subbundles, each of which is isomorphic to TY . It follows that $\text{Hol}(\nabla, y)$ is the direct sum of four copies of the holonomy of the tangent connection, which is the full special orthogonal group $SO(n)$. One of these copies is T_yY , the other are its images under an orthonormal basis (I, J, K) of imaginary quaternions acting on the right.

Let us show that Z contains a copy of $H_{\mathbb{C}}^n$. Let $a \in G$ fix a hyperplane $P \subset Y$ pointwise. According to Lemma 7.2, $\text{Fix}(P) = \text{Sp}(1)\text{Sp}(1)$, so a is given by two unit quaternions q and d . Pick an origin $y \in P$. Let $u \in T_yY$ be a unit vector orthogonal to P . On $T_yX = T_yY \otimes \mathbb{H}$, a acts by the identity on T_yP and maps u to duq^{-1} . Since u is a real vector, $a(u) = udq^{-1} \in T_yY \oplus (T_yY)i$ where $i = \text{Im}(dq^{-1})$. Up to conjugating by an element of the $\text{Sp}(1)$ subgroup of G that fixes Y pointwise, one can assume that i is proportional to I , i.e. T_yZ contains uI . By assumption, $uI \notin T_yY$. By $SO(n)$ -invariance, T_yZ contains $T_yY \oplus (T_yY)I = T_yH_{\mathbb{C}}^n$, therefore Z contains $Y' = H_{\mathbb{C}}^n$.

Now $TZ|_{Y'}$ is a parallel subbundle of $TX|_{Y'}$, thus T_yZ is $U(n)$ -invariant. Under $U(n)$, T_yX splits into only two summands. Since $Z \neq X$, $T_yZ = T_yY'$, i.e. $Z = Y'$. □

Along the way, we proved the following.

Lemma 7.5. *Let $Y' = H_{\mathbb{C}}^n \subset H_{\mathbb{H}}^n = X$. Let Z be a totally geodesic subspace of X containing Y' . Then either $Z = X$ or $Z = Y'$.*

Corollary 7.6. *After bending in $P\text{Sp}(n, 1)$, a Zariski dense subgroup of $PO(n, 1)^0$ becomes Zariski dense in a conjugate of $PU(n, 1)$.*

Proof. Let $\Gamma = A \star_C B$ be Zariski dense in $PO(n, 1)^0$, with C being Zariski dense in $PO(n - 1, 1)^0$. In other words, Γ leaves $Y = H_{\mathbb{R}}^n$ invariant, and C leaves $P = H_{\mathbb{R}}^{n-1}$ invariant. Lemma 7.2 allows us to select an $a \in Z_G(C)$ which does not map Y to itself. Lemma 7.4 shows that the smallest totally geodesic subspace of $X = H_{\mathbb{H}}^n$ containing Y and $a(Y)$ is congruent to $H_{\mathbb{C}}^n$. According to Lemma 7.3, this means that the bent subgroup $A \star_C aBa^{-1}$ is Zariski dense in a conjugate of $PU(n, 1)$. □

Therefore, to obtain a Zariski dense subgroup in $P\text{Sp}(m, 1)$, $m \geq n$, one must bend several times.

7.2. Further bending steps

We shall use compact hyperbolic manifolds which contain several disjoint separating totally geodesic hypersurfaces. Again, such manifolds exist in all dimensions (see [17]). In low dimensions, a vast majority of known examples of compact hyperbolic manifolds have this property (they fall into infinitely many distinct commensurability classes, see [1]). Given such a manifold, bending can be performed several times in a row. The next lemmas show that at each step, the Zariski closure strictly increases.

Lemma 7.7. *Let $X' = H_{\mathbb{H}}^n$. Let Z be a totally geodesic subspace of $X = H_{\mathbb{H}}^m$ such that $X' \subsetneq Z \subsetneq X$. Then Z is a quaternionic subspace. Furthermore, there exists an $a \in G$ fixing X' pointwise which does not map Z into itself.*

Proof. Otherwise, Z would be $\text{Sp}(m - n)$ -invariant. In particular, for $x \in X'$, $T_x Z$ would be $\text{Sp}(m - n)$ -invariant. Since $\text{Sp}(m - n)$ acts irreducibly on $(T_x X')^\perp$, Z must be equal to X' or X , a contradiction. As Z is a negatively curved symmetric space containing $H_{\mathbb{H}}^n$, $n \geq 2$, it is a quaternionic subspace. \square

Proposition 7.8. *Let M be a compact hyperbolic n -manifold. Let $m \geq n$. Assume that M contains N disjoint separating totally geodesic hypersurfaces. Let $\Gamma = \pi_1(M) \subset \text{PO}(n, 1)^0 \rightarrow \text{P Sp}(m, 1)$. If $N \geq m - n + 2$, then Γ can be continuously deformed to a Zariski dense subgroup of $\text{P Sp}(m, 1)$.*

Proof. According to Corollary 7.6, a first bending in $\text{PU}(n, 1)$ provides us with a Zariski dense subgroup of $\text{PU}(n, 1)$.

A second bending in $\text{P Sp}(n, 1)$ gives a Zariski dense subgroup of $\text{P Sp}(n, 1)$. Indeed, the fixator of $H_{\mathbb{R}}^{n-1}$ is an $\text{Sp}(1) \text{Sp}(1)$ which contains an element a which does not map $H_{\mathbb{C}}^n$ to itself. By Lemma 7.5, no proper totally geodesic subspace of $H_{\mathbb{H}}^n$ contains both $H_{\mathbb{C}}^n$ and $a(H_{\mathbb{C}}^n)$. Lemma 7.3 implies that the bent subgroup is Zariski dense.

A third series of bendings gives a Zariski dense subgroup of $\text{P Sp}(m, 1)$. Lemma 7.7 allows us to inductively select a parameter a which strictly increases the dimension of the totally geodesic hull. After at most $m - n$ more steps, the subgroup obtained is Zariski dense, thanks to Lemma 7.3. \square

7.3. Bending along laminations

Since we need to bend surfaces of genus as low as 2, which do not admit pairs of disjoint separating closed geodesics, we describe W. Thurston’s general construction of bending along totally geodesic laminations, which does not require the leaves to be separating. We stick to the special case of totally real, totally geodesic 2-planes of $H_{\mathbb{H}}^2$.

Let $Y = H_{\mathbb{R}}^2 \subset H_{\mathbb{H}}^2 = X$. If $\ell \subset Y$ is a geodesic, the subgroup $\text{Fix}(\ell)$ of $\text{Isom}(X)$ that fixes ℓ pointwise is conjugate to $\text{Sp}(1) \text{Sp}(1)$. The Lie algebras of these subgroups form an $\text{Im } \mathbb{H} \oplus \text{Im } \mathbb{H}$ -bundle \mathcal{B} over the space \mathcal{L} of geodesics in Y . Pick once and for all an arbitrary Borel trivialization of this bundle. A lamination on Y is a closed subset of \mathcal{L}

consisting of pairwise nonintersecting geodesics. A *measured lamination* on Y is the data of a lamination λ and a transverse $\text{Im } \mathbb{H} \oplus \text{Im } \mathbb{H}$ -valued measure. By a transverse measure, we mean the data, for each continuous curve $c : [a, b] \rightarrow Y$ which crosses all geodesics of λ in the same direction, of a finite Borel $\text{Im } \mathbb{H} \oplus \text{Im } \mathbb{H}$ -valued measure μ_c on $[a, b]$, with the following compatibility: if a curve $c' : [a, b] \rightarrow Y$ can be deformed to c by sliding along λ , then $\mu_{c'} = \mu_c$. A discrete collection of geodesics, with an $\text{Im } \mathbb{H} \oplus \text{Im } \mathbb{H}$ -valued Dirac mass at each geodesic, is a simple example of a measured lamination. Since only such laminations will ultimately be used, we shall not discuss nondiscrete measured laminations further.

The Lie algebra bundle \mathcal{B} is a subbundle of the trivial bundle with fiber the Lie algebra $\mathfrak{sp}(2, 1)$. Therefore, for every transversal curve c , the measure μ_c can be pushed forward to yield an $\mathfrak{sp}(2, 1)$ -valued measure on $[a, b]$. This measure integrates into a continuous map $[a, b] \rightarrow \text{Sp}(2, 1)$ (see for example [7]). We denote the resulting element of $\text{Sp}(2, 1)$ by $\int \mu_c$. If $c = c_1 c_2$ is obtained by traversing a first curve c_1 and then a second curve c_2 , then the Chasles rule $\int \mu_{c_1 c_2} = (\int \mu_{c_1})(\int \mu_{c_2})$ holds, which allows one to extend the definition to curves which are piecewise transversal. Define a map $f : Y \rightarrow X$ as follows. Pick an origin $o \in Y$. Given $y \in Y$, join o to y with a piecewise transversal curve c_y and set $f(y) = (\int \mu_{c_y})y$. One checks that $f(y)$ does not depend on the choice of piecewise transversal curve.

For instance, in the case of a discrete lamination, f is piecewise isometric and totally geodesic away from the support of λ . At each geodesic ℓ of the lamination, f bends, i.e. the totally geodesic pieces of the surface $f(Y)$ at either side of ℓ meet at a $\text{Fix}(\ell)$ -angle equal to $\exp(\mu(\ell))$. The general case is best understood by considering limits of discrete measured laminations.

Let $\rho : \Gamma \rightarrow \text{Sp}(2, 1)$ be an isometric action of a group Γ which leaves Y and the measured lamination invariant. Then, for every piecewise transversal curve c , and $\gamma \in \Gamma$, $\int \mu_{\rho(\gamma)(c)} = \rho(\gamma)(\int \mu_c)\rho(\gamma)^{-1}$. For $\gamma \in \Gamma$, let $\rho_\lambda(\gamma) = (\int \mu_{c_\gamma})\rho(\gamma)$, where c_γ is a piecewise transversal curve joining o to $\rho(\gamma)o$. Then $\rho_\lambda : \Gamma \rightarrow \text{Sp}(2, 1)$ is a homomorphism which stabilizes $f(Y)$, and f is equivariant. Indeed, let c_1 (resp. c_2) be a piecewise transversal curve joining o to $\rho(\gamma_1)o$ (resp. to $\rho(\gamma_2)o$). Then $c_1 \rho(\gamma_1)(c_2)$ joins o to $\rho(\gamma_1 \gamma_2)o$ and

$$\begin{aligned} \rho_\lambda(\gamma_1 \gamma_2) &= \left(\int \mu_{c_1 \rho(\gamma_1)(c_2)} \right) \rho(\gamma_1 \gamma_2) = \left(\int \mu_{c_1} \right) \left(\int \mu_{\rho(\gamma_1)(c_2)} \right) \rho(\gamma_1 \gamma_2) \\ &= \left(\int \mu_{c_1} \right) \rho(\gamma_1) \left(\int \mu_{c_2} \right) \rho(\gamma_1^{-1}) \rho(\gamma_1 \gamma_2) = \rho_\lambda(\gamma_1) \rho_\lambda(\gamma_2). \end{aligned}$$

If $y \in Y$ and $\gamma \in \Gamma$, let c_y (resp. c_γ) be a piecewise transversal curve joining o to y (resp. to $\rho(\gamma)o$). Then $c_\gamma \rho(\gamma)(c_y)$ joins o to $\rho(\gamma)y$, thus

$$\begin{aligned} f(\rho(\gamma)y) &= \left(\int \mu_{c_\gamma \rho(\gamma)(c_y)} \right) \rho(\gamma)y = \left(\int \mu_{c_\gamma} \right) \left(\int \mu_{\rho(\gamma)(c_y)} \right) \rho(\gamma)y \\ &= \left(\int \mu_{c_\gamma} \right) \rho(\gamma) \left(\int \mu_{c_y} \right) \rho(\gamma)^{-1} \rho(\gamma)y = \rho_\lambda(\gamma) f(y). \end{aligned}$$

Proposition 7.9. *Let Σ be a closed hyperbolic surface with fundamental group Γ . Map $\Gamma \rightarrow SO(2, 1) \rightarrow Sp(2, 1)$. There exist measured laminations λ on Σ which make the bent group $\rho_\lambda(\Gamma)$ Zariski dense in $Sp(2, 1)$.*

Proof. As a lamination, take the lifts to $Y = \tilde{\Sigma}$ of two disjoint closed geodesics in Σ . A transversal measure in this case is simply the data of elements $a_j \in \text{Fix}(\ell_j)$ for two lifts ℓ_1, ℓ_2 . Note that the components of the complement of the two geodesics in Σ are not simply connected. In other words, each component of the complement of the support of the lifted lamination on Y is stabilized by a subgroup of Γ which is Zariski dense in $SO(2, 1)$. It follows that the Zariski closure of $\rho_\lambda(\Gamma)$ contains $SO(2, 1)$. It also contains the conjugates of $SO(2, 1)$ by the two isometries a_1 and a_2 .

According to Lemma 7.3, the Zariski closure of $\rho_\lambda(\Gamma)$ contains the isometry group of the totally geodesic hull Z of $Y \cup a_1(Y) \cup a_2(Y)$. As in the proof of Proposition 7.8, bending by a_1 gives a group which is Zariski dense in a conjugate of $PU(2, 1)$, bending by a_1 and a_2 gives a group which is Zariski dense in $P Sp(2, 1)$. \square

8. Flexibility of Fuchsian surface groups

In this section, we investigate homomorphisms of a surface group into $Sp(2, 1)$ in a neighborhood of the embedding via $SU(1, 1)$ and $Sp(1, 1)$. We shall call them *Fuchsian*, to distinguish them from the bendable homomorphisms arising from the embedding via $SO(2, 1)$.

8.1. Second order calculations

Let S be a compact Riemann surface with genus > 1 and $\rho_0 : \pi_1(S) = \Gamma \subset SU(1, 1) \rightarrow Sp(1, 1) \subset Sp(2, 1)$ be a standard representation fixing a quaternionic line in $H_{\mathbb{H}}^2$. Since $H^1(\pi_1(S), \mathbb{H}^2) \neq 0$, Proposition 2.1 does not apply. We have to investigate which infinitesimal deformations represented by elements in $H^1(\pi_1(S), \mathfrak{sp}(2, 1))$ are integrable.

The second order integrability condition for infinitesimal deformations at ϕ of representations of a group Γ in a Lie group G can be expressed in terms of the *cup-product*, a symmetric bilinear map

$$[\cdot, \cdot] : H^1(\Gamma, \mathfrak{g}_{\text{Ad } \phi}) \rightarrow H^2(\Gamma, \mathfrak{g}_{\text{Ad } \phi}).$$

For $u \in Z^1(\Gamma, \mathfrak{g}_{\text{Ad } \phi})$,

$$[u, u](\alpha, \beta) = [u(\alpha), \text{Ad } \phi(\alpha)u(\beta)].$$

It is well known [18] that for a representation ϕ from Γ to a reductive group G , if there exists a smooth path ϕ_t in $\text{Hom}(\Gamma, G)$ which is tangent to $u \in Z^1(\Gamma, \mathfrak{g}_{\text{Ad } \phi})$, then $[u, u] = 0$. According to Theorem 3 in [8], for surface groups, this necessary condition is also sufficient.

Theorem 8.1 (W. Goldman). *Let S be a closed surface, and let G be a reductive group. Let $\phi : \pi_1(S) \rightarrow G$ be a representation such that the Zariski closure of $\phi(\pi_1(S))$ is also reductive. Then for any $u \in Z^1(\pi_1(S), \mathfrak{g}_{\text{Ad}\phi})$, $[u, u] = 0$ if and only if there exists an analytic path $t \mapsto \phi_t$ in $\text{Hom}(\pi_1(S), G)$ which is tangent to u .*

8.2. Splitting of the cup-product map

The centralizer of $SU(1, 1)$ in $\text{Sp}(2, 1)$ is $\text{Sp}(1) \times U(1)$, where $\text{Sp}(1)$ is the centralizer of $\text{Sp}(1, 1)$ and $U(1) \subset \text{Sp}(1, 1)$ is the centralizer of $SU(1, 1)$ in $\text{Sp}(1, 1)$. Then by Poincaré duality,

$$H^2(\pi_1(S), \mathfrak{sp}(2, 1)) = H^0(\pi_1(S), \mathfrak{sp}(2, 1)) = \mathfrak{sp}(1) \oplus \mathfrak{u}(1).$$

Let $u \in H^1(\pi_1(S), \mathfrak{sp}(2, 1))$ split as $u = u_{\mathfrak{sp}(1)} + u_{\mathfrak{sp}(1,1)} + u_{\mathbb{H}^2}$. Since $\mathfrak{sp}(1, 1)$ and $\mathfrak{sp}(1)$ commute, $[u_{\mathfrak{sp}(1)}, u_{\mathfrak{sp}(1,1)}] = 0$. Since the subspace $\mathbb{H}^2 \subset \mathfrak{sp}(2, 1)$ is $\text{Sp}(1) \times \text{Sp}(1, 1)$ -invariant, $[u_{\mathfrak{sp}(1)}, u_{\mathbb{H}^2}]$ and $[u_{\mathfrak{sp}(1,1)}, u_{\mathbb{H}^2}]$ belong to $H^2(\pi_1(S), \mathbb{H}^2) = 0$. Therefore

$$[u, u] = [u_{\mathfrak{sp}(1)}, u_{\mathfrak{sp}(1)}] + [u_{\mathfrak{sp}(1,1)}, u_{\mathfrak{sp}(1,1)}] + [u_{\mathbb{H}^2}, u_{\mathbb{H}^2}].$$

As both $\mathfrak{sp}(1)$ and $\mathfrak{sp}(1, 1)$ are subalgebras, $[u_{\mathfrak{sp}(1)}, u_{\mathfrak{sp}(1)}]$ belongs to $H^2(\pi_1(S), \mathfrak{sp}(1)) = \mathfrak{sp}(1)$, and $[u_{\mathfrak{sp}(1,1)}, u_{\mathfrak{sp}(1,1)}]$ to $H^2(\pi_1(S), \mathfrak{sp}(1, 1)) = \mathfrak{u}(1)$. On the other hand, $[u_{\mathbb{H}^2}, u_{\mathbb{H}^2}]$ has nontrivial components $[u_{\mathbb{H}^2}, u_{\mathbb{H}^2}]_{\mathfrak{u}(1)}$ and $[u_{\mathbb{H}^2}, u_{\mathbb{H}^2}]_{\mathfrak{sp}(1)}$ on both $H^2(\pi_1(S), \mathfrak{sp}(1, 1))$ and $H^2(\pi_1(S), \mathfrak{sp}(1))$.

8.3. Homomorphisms to $\text{Sp}(1)$

In the special case of the trivial representation to $\text{Sp}(1)$, the cup-product map can be computed.

Lemma 8.2. *Let S be a closed surface. Let $\pi_1(S)$ act trivially on $\mathfrak{sp}(1)$. The quadratic map $H^1(\pi_1(S), \mathfrak{sp}(1)) \rightarrow H^2(\pi_1(S), \mathfrak{sp}(1))$, $u \mapsto [u, u]$, is onto.*

Proof. Here, $H^1(\pi_1(S), \mathfrak{sp}(1)) \simeq H^1(\pi_1(S), \mathbb{R}) \otimes \mathfrak{sp}(1)$. If $a, b \in H^1(\pi_1(S), \mathbb{R})$ and $q, q' \in \mathfrak{sp}(1)$, then

$$[a \otimes q, b \otimes q'] = a \smile b \otimes [q, q'].$$

For every $q'' \in \mathfrak{sp}(1)$, there exist $q, q' \in \mathfrak{sp}(1)$ such that $[q, q'] = q''$. Poincaré duality implies that there exist $a, b \in H^1(\pi_1(S), \mathbb{R})$ such that $a \smile b \neq 0$. Therefore the cup-product map is onto. \square

8.4. Homomorphisms to $\mathrm{Sp}(1, 1)$

A similar statement applies to $H^1(\pi_1(S), \mathfrak{sp}(1, 1))$.

Lemma 8.3. *Let S be a closed hyperbolic surface. View $\pi_1(S)$ as a subgroup of $SU(1, 1) \subset \mathrm{Sp}(1, 1)$. The quadratic map $H^1(\pi_1(S), \mathfrak{sp}(1, 1)) \rightarrow H^2(\pi_1(S), \mathfrak{sp}(1, 1)) = \mathfrak{u}(1)$, $u \mapsto [u, u]$, is onto.*

Proof. $\mathfrak{sp}(1, 1)$ consists of quaternionic 2×2 matrices $\begin{pmatrix} a & b \\ \bar{b} & d \end{pmatrix}$ with a, d imaginary quaternions. The complex matrices in $\mathfrak{sp}(1, 1)$ form the subalgebra $\mathfrak{u}(1, 1) = \mathfrak{su}(1, 1) \oplus \mathfrak{u}(1)$, where $\mathfrak{u}(1)$ consists of complex imaginary multiples of the unit matrix. As a $U(1, 1)$ -invariant projection $\mathfrak{sp}(1, 1) \rightarrow \mathfrak{u}(1) = \mathbb{R}$, we can use the linear form

$$\pi_{\mathfrak{u}(1)} \begin{pmatrix} a & b \\ \bar{b} & d \end{pmatrix} = \frac{1}{2} \mathrm{Re} e(i(a + d)).$$

Let W denote the set of matrices of the form $j \begin{pmatrix} z & w \\ -\bar{w} & \bar{z} \end{pmatrix}$, where $z, w, t \in \mathbb{C}$. Then W is a $U(1, 1)$ -invariant complement of $\mathfrak{u}(1, 1)$ in $\mathfrak{sp}(1, 1)$. Given two elements $X = j \begin{pmatrix} z & w \\ -\bar{w} & \bar{z} \end{pmatrix}$ and $X' = j \begin{pmatrix} z' & w' \\ -\bar{w}' & \bar{z}' \end{pmatrix}$ in W , one computes

$$\pi_{\mathfrak{u}(1)}([X, X']) = -\mathrm{Im}(\bar{z}z' + \bar{t}t' - 2\bar{w}w').$$

This is a symplectic structure on W (viewed as a real vector space). From Poincaré duality for local coefficient systems, it follows that the quadratic form $\pi_{\mathfrak{u}(1)}([\cdot, \cdot])$ on $H^1(\pi_1(S), W)$ is nondegenerate. In particular, it is onto. *A fortiori*, the quadratic form $[\cdot, \cdot]$ on $H^1(\pi_1(S), \mathfrak{sp}(1, 1))$ is onto. \square

8.5. Flexibility of certain Fuchsian surface groups

A surface group in $SU(n, 1)$ is *Fuchsian* if it stabilizes a complex line in complex hyperbolic space. Let us extend the terminology. Say a surface group in $\mathrm{Sp}(n, 1)$ is *Fuchsian* if it stabilizes a complex line in quaternionic hyperbolic space. Note that every complex line is contained in a unique quaternionic line.

It is well known that Fuchsian groups in $SU(2, 1)$ (or, more generally, $SU(n, 1)$) cannot be deformed to Zariski dense groups. We show that when $SU(2, 1)$ is embedded in the larger group $\mathrm{Sp}(2, 1)$, this rigidity property fails. We make essential use of the main result of [8].

Proposition 8.4. *Let S be a compact Riemann surface with genus > 1 and $\rho_0 : \pi_1(S) = \Gamma \subset SU(1, 1) \rightarrow \mathrm{Sp}(1, 1) \subset \mathrm{Sp}(2, 1)$ be a standard representation fixing a quaternionic line in $H_{\mathbb{H}}^2$. Then there exist local deformations of ρ_0 which do not stabilize any quaternionic line.*

Proof. Let $u \in H^1(\pi_1(S), \mathbb{H}^2)$ be nonzero. According to Lemmas 8.2 and 8.3, there exist $v \in H^1(\pi_1(S), \mathfrak{sp}(1))$ and $w \in H^1(\pi_1(S), \mathfrak{sp}(1, 1))$ such that $[v, v] = -[u, u]_{\mathfrak{sp}(1)}$ and $[w, w] = -[u, u]_{\mathfrak{u}(1)}$. Then $x = u + v + w \in H^1(\pi_1(S), \mathfrak{g})$ is nonzero and satisfies $[x, x] = 0$. According to Goldman's Theorem 8.1, there exists an analytic curve $t \mapsto \rho_t$ in $\text{Hom}(\pi_1(S), G)$, starting at ρ_0 , whose initial speed is a representative of the cohomology class x . Since $x \notin H^1(\pi_1(S), \mathfrak{sp}(1) \oplus \mathfrak{sp}(1, 1))$, for $t \neq 0$ small, ρ_t cannot be conjugated to the subgroup $\text{Sp}(1, 1)\text{Sp}(1)$, i.e., does not stabilize any quaternionic line. \square

Proof of Theorem 1.3. Proposition 8.4 is statement (2) of Theorem 1.3. Statement (1) of Theorem 1.3 is a consequence of the bending construction. For surfaces of sufficiently high genus, one can apply Proposition 7.8. In low genus, one has to bend along a geodesic lamination (see Proposition 7.9). \square

9. Discrete representations

Proposition 9.1. *Let Γ be a uniform lattice in $\text{Sp}(1, 1)$. Let $\rho : \Gamma \rightarrow \text{Sp}(2, 1)$ be a discrete and faithful homomorphism. Then either*

- ρ is standard, i.e. it stabilizes a quaternionic line, or
- the image is Zariski dense.

Proof. Suppose $\rho(\Gamma)$ is not Zariski dense. Then it cannot be contained in a parabolic subgroup of $\text{Sp}(2, 1)$ since Γ is not solvable. So it must stabilize a totally geodesic subspace of $H_{\mathbb{H}}^2$ (see [13]). If it stabilizes a quaternionic line, it is a standard representation, by Mostow rigidity. Suppose it stabilizes $H_{\mathbb{C}}^2$. Then $H_{\mathbb{C}}^2/\rho(\Gamma)$ is a manifold. If it is not closed, the cohomological dimension of Γ cannot be 4, which contradicts Γ being a uniform lattice in $\text{Sp}(1, 1)$. So $H_{\mathbb{C}}^2/\rho(\Gamma)$ is a closed manifold, which implies that $H_{\mathbb{C}}^2$ and $H_{\mathbb{R}}^4$ are quasi-isometric, which is impossible, again by a result of G. D. Mostow. \square

We suspect that there is no Zariski dense discrete faithful group $\rho(\Gamma)$.

Acknowledgments. We thank an anonymous referee for his valuable suggestions.

I. Kim gratefully acknowledges the partial support of KRF Grant (0409-20060066) and a warm support of IHES during his stay.

References

- [1] Allcock, D.: Infinitely many hyperbolic Coxeter groups through dimension 19. *Geom. Topology* **10**, 737–758 (2006) Zbl 1133.20025 MR 2240904
- [2] Apanasov, B. N.: Bending and stamping deformations of hyperbolic manifolds. *Ann. Global Anal. Geom.* **8**, 3–12 (1990) Zbl 0705.53013 MR 1075235
- [3] Burger, M., Iozzi, A., Wienhard, A.: Surface group representations with maximal Toledo invariant. arXiv:math/0605656.
- [4] Corlette, K.: Flat G -bundles with canonical metrics. *J. Differential Geom.* **28**, 361–382 (1988) Zbl 0676.58007 MR 0965220

- [5] Corlette, K.: Archimedean superrigidity and hyperbolic geometry. *Ann. of Math.* **135**, 165–182 (1992) Zbl 0768.53025 MR 1147961
- [6] Domic, A., Toledo, D.: The Gromov norm of the Kaehler class of symmetric domains. *Math. Ann.* **276**, 425–432 (1987) Zbl 0595.53061 MR 0875338
- [7] Estrade, A.: Exponentielle stochastique et intégrale multiplicative discontinues. *Ann. Inst. H. Poincaré Probab. Statist.* **28**, 107–129 (1992) Zbl 0760.60048 MR 1158740
- [8] Goldman, W.: Representations of fundamental groups of surfaces. In: *Geometry and Topology*, J. Alexander and J. Harer (eds.), Lecture Notes Math. 1167, Springer, 95–117 (1985) Zbl 0575.57027 MR 0827264
- [9] Goldman, W., Millson, J.: Local rigidity of discrete groups acting on complex hyperbolic space. *Invent. Math.* **88**, 495–520 (1987) Zbl 0627.22012 MR 0884798
- [10] Gromov, M., Pansu, P.: Rigidity of lattices: An introduction. In: *Geometric Topology: Recent Developments*, Lecture Notes in Math. 1504, Springer, 39–137 (1991) Zbl 0786.22015 MR 1168043
- [11] Kim, I.: Geometry on exotic hyperbolic spaces. *J. Korean Math. Soc.* **36**, 621–631 (1999) Zbl 0943.51014 MR 1736298
- [12] Kim, I.: Marked length rigidity of rank one symmetric spaces and their products. *Topology* **40**, 1295–1323 (2001) Zbl 0997.53034 MR 1867246
- [13] Kim, I.: Rigidity on symmetric spaces. *Topology* **43**, 393–405 (2004) Zbl 1049.53031 MR 2052969
- [14] Kim, I., Parker, J.: Geometry of quaternionic hyperbolic manifolds. *Math. Proc. Cambridge Philos. Soc.* **135**, 291–320 (2003) Zbl 1048.32017 MR 2006066
- [15] Koziarz, V., Maubon, J.: Harmonic maps and representations of non-uniform lattices of $PU(m, 1)$. *Ann. Inst. Fourier (Grenoble)* **58**, 507–558 (2008) Zbl 1147.22009 MR 2410381
- [16] Koziarz, V., Maubon, J.: Representations of complex hyperbolic lattices into rank 2 classical Lie groups of Hermitian type. *Geom. Dedicata* **137**, 85–111 (2008) Zbl 1159.22006 MR 2449147
- [17] Millson, J.: On the first Betti number of a constant negatively curved manifold. *Ann. of Math.* **104**, 235–247 (1976) Zbl 0364.53020 MR 0422501
- [18] Nijenhuis, A., Richardson, R. W.: Deformations of homomorphisms of Lie groups and Lie algebras. *Bull. Amer. Math. Soc.* **73**, 175–179 (1967) Zbl 0153.04402 MR 0204575
- [19] Pansu, P.: Sous-groupes discrets des groupes de Lie: rigidité, arithmétique. In: *Séminaire Bourbaki*, 1993–94, exp. 778, Astérisque **227**, 69–105 (1995) Zbl 0835.22011 MR 1321644
- [20] Raghunathan, M. S.: On the first cohomology of discrete subgroups of semi-simple Lie groups. *Amer. J. Math.* **87**, 103–139 (1965) Zbl 0132.02102 MR 0173730
- [21] Thurston, W.: *The Geometry and Topology of 3-Manifolds*. Lecture notes, Princeton (1983)
- [22] Toledo, D.: Representations of surface groups on complex hyperbolic space. *J. Differential Geom.* **29**, 125–133 (1989) Zbl 0676.57012 MR 0978081
- [23] Weil, A.: On discrete subgroups of Lie groups. *Ann. of Math.* **72**, 369–384 (1960) Zbl 0131.26602 MR 0137792