



Simon Thomas

A descriptive view of unitary group representations

Received December 30, 2012 and in revised form June 25, 2014

Abstract. In this paper, we will study the relative complexity of the unitary duals of countable groups. In particular, we will explain that if G and H are countable amenable non-type I groups, then the unitary duals of G and H are Borel isomorphic.

Keywords. Unitary dual, Borel equivalence relation, amenable group

1. Introduction

Let G be a countable group. Then the *unitary dual* \widehat{G} of G is the set of equivalence classes of irreducible unitary representations of G , equipped with its Mackey Borel structure. In more detail, for each $n \in \{1, 2, \dots, \infty\}$, let $\text{Irr}_n(G)$ be the Polish space of all irreducible unitary representations of G in some fixed separable Hilbert space \mathcal{H}_n of dimension n . Then the unitary dual is the quotient of the disjoint union $\bigsqcup \text{Irr}_n(G)$ by the unitary equivalence relation, equipped with the corresponding quotient Borel structure. Recall that the unitary equivalence relation on $\bigsqcup \text{Irr}_n(G)$ is said to be *smooth* if and only if the Mackey Borel structure on the unitary dual \widehat{G} is countably separated. Of course, this is only problematic for the restriction of the unitary equivalence relation to $\text{Irr}_\infty(G)$. The following result combines the main theorems of Glimm [22] and Thoma [47].

Theorem 1.1. *If G is a countable group, then the following conditions are equivalent:*

- (i) G is not abelian-by-finite.
- (ii) G has an infinite-dimensional irreducible representation.
- (iii) The unitary equivalence relation on the space $\text{Irr}_\infty(G)$ of infinite-dimensional irreducible unitary representations of G is not smooth.

Furthermore, by Thoma [47], a countable group G satisfies the conditions of Theorem 1.1 if and only if G is a *non-type I* group.¹ While Theorem 1.1 shows that the equivalence

S. Thomas: Mathematics Department, Rutgers University, 110 Frelinghuysen Road, Piscataway, NJ 08854-8019, USA; e-mail: sthomas@math.rutgers.edu

Mathematics Subject Classification (2010): Primary 22D10, 03E15; Secondary 43A07, 37A20

¹ The definition of a type I group can be found in Folland [17, Section 7.2]. This notion will play no role in this paper, other than to enable us to avoid repeatedly using the term “non-(abelian-by-finite)”.

classes of irreducible unitary representations of a countable non-type I group G cannot be parameterized by the points of any Polish space, this is not enough in itself to rule out a satisfactory classification. For example, the isomorphism relation on the space of rank 1 torsion-free abelian groups is not smooth; and yet Baer [3] was able to provide a perfectly satisfactory classification. However, the following more recent result of Hjorth provides a much more serious obstruction to the existence of a satisfactory classification.

Theorem 1.2 (Hjorth [25]). *If G is a countable non-type I group, then there exists a $U(\mathcal{H}_\infty)$ -invariant Borel subset $X \subseteq \text{Irr}_\infty(G)$ such that the action of the unitary group $U(\mathcal{H}_\infty)$ on X is turbulent.*

The notion of turbulence was introduced by Hjorth [26], in order to address the question of which orbit equivalence relations of continuous actions of Polish groups on Polish spaces are classifiable by countable structures. Here an analytic equivalence relation E on a Polish space X is said to admit a *classification by countable structures* if there exists a Borel map $x \mapsto \mathcal{M}_x$ from X to the space of countable structures for some countable language such that

$$x E y \Leftrightarrow \mathcal{M}_x \cong \mathcal{M}_y.$$

For example, Farah–Toms–Törnquist [15] have verified that Elliott’s classification [11] of the approximately finite-dimensional C^* -algebras by K -theoretic invariants can be implemented in a Borel manner; and this means that the isomorphism relation for approximately finite-dimensional C^* -algebras admits a classification by countable structures. On the other hand, Hjorth [26] has shown that if the continuous action of the Polish group G on the Polish space X is turbulent, then the corresponding orbit equivalence relation E_G^X does not admit a classification by countable structures. In particular, if G is a countable non-type I group, then the unitary equivalence relation \approx_G does not admit a classification by countable structures.

However, this is hopefully not the end of the story. While it does not seem feasible to find a satisfactory classification of the irreducible unitary representations of a *fixed* countable non-type I group, there remains the natural question of determining the relative complexities of the unitary duals of pairs G, H of countable non-type I groups. For example, let G be the direct sum of infinitely many copies of the symmetric group $\text{Sym}(3)$ and let \mathbb{F}_∞ be the free group on infinitely many generators. Then it seems reasonable to conjecture that the unitary dual of \mathbb{F}_∞ is strictly more complex than that of G . In order to give a precise formulation of this question, it is first necessary to recall some of the basic notions from the theory of Borel equivalence relations.

If X is a Polish space, then an equivalence relation E on X is said to be *Borel* if E is a Borel subset of $X \times X$. For example, it is well-known that unitary equivalence is an F_σ equivalence relation on $\text{Irr}_n(G)$. Suppose that E, F are Borel equivalence relations on the Polish spaces X, Y respectively. Then E is *Borel reducible* to F , written $E \leq_B F$, if there exists a Borel map $f : X \rightarrow Y$ such that for all $x, z \in X$,

$$x E z \Leftrightarrow f(x) F f(z).$$

E and F are *Borel bireducible*, written $E \sim_B F$, if both $E \leq_B F$ and $F \leq_B E$. Finally we write $E <_B F$ if both $E \leq_B F$ and $F \not\leq_B E$.

Here the idea is that Borel bireducibility captures the intuitive notion of two Borel equivalence relations having the same complexity. As we will explain next, the Borel bireducibility of the unitary equivalence relations of the countable groups G, H is equivalent to the usual notion in the literature of the Borel isomorphism of their unitary duals \widehat{G}, \widehat{H} . Recall that the unitary duals \widehat{G}, \widehat{H} are said to be *Borel isomorphic* if there exists a bijection $f : \widehat{G} \rightarrow \widehat{H}$ such that both f and f^{-1} admit Borel liftings $\varphi : \bigsqcup \text{Irr}_n(G) \rightarrow \bigsqcup \text{Irr}_n(H)$ and $\psi : \bigsqcup \text{Irr}_n(H) \rightarrow \bigsqcup \text{Irr}_n(G)$ respectively.

Remark 1.3. Some readers may be a little concerned about our apparently strong requirement that both f and f^{-1} admit Borel liftings. However, the proof of Gao [21, Corollary 5.2.4] shows that if a bijection $f : \widehat{G} \rightarrow \widehat{H}$ admits a Borel lifting, then its inverse f^{-1} also admits a Borel lifting.

Next notice, for example, that if the Borel map φ is a lifting of f , then $\pi, \pi' \in \bigsqcup \text{Irr}_n(G)$ are unitarily equivalent if and only if $\varphi(\pi), \varphi(\pi')$ are unitarily equivalent; and so φ is a Borel reduction from the unitary equivalence relation of G to the unitary equivalence relation of H . Consequently, if the unitary duals \widehat{G}, \widehat{H} are Borel isomorphic, then the unitary equivalence relations of G, H are Borel bireducible. In order to explain why the converse holds, it is useful to extend the notion of a Borel isomorphism to arbitrary quotients of Polish spaces by Borel equivalence relations.

Definition 1.4. If E, F are Borel equivalence relations on the Polish spaces X, Y then the quotients $X/E, Y/F$ are *Borel isomorphic*, written $X/E \simeq Y/F$, if there exist Borel reductions $\varphi : X \rightarrow Y$ and $\psi : Y \rightarrow X$ such that the induced maps $\widehat{\varphi} : X/E \rightarrow Y/F$ and $\widehat{\psi} : Y/F \rightarrow X/E$ are mutually inverse bijections.

There exist examples of Borel equivalence relations E, F such that E, F are Borel bireducible but not Borel isomorphic. (For example, see Friedman–Motto Ros [18].) However, the two notions coincide if we restrict our attention to the class of Borel orbit equivalence relations.

Theorem 1.5 (Motto Ros [40]). *If E, F are Borel orbit equivalence relations arising from Borel actions of Polish groups on the Polish spaces X, Y , then $E \sim_B F$ if and only if $X/E \simeq Y/F$.*

In particular, the Borel bireducibility of the unitary equivalence relations of the countable groups G, H is equivalent to the Borel isomorphism of their unitary duals \widehat{G}, \widehat{H} . Of course, if G is a countable group, then the unitary equivalence restricted to the space $\bigsqcup_{1 \leq n \in \mathbb{N}} \text{Irr}_n(G)$ of finite-dimensional irreducible unitary representations is smooth. Consequently, our focus will be on the space $\text{Irr}_\infty(G)$ of infinite-dimensional irreducible unitary representations of G .

Definition 1.6. If G is a countable non-type I group, then \approx_G denotes the unitary equivalence relation on the space $\text{Irr}_\infty(G)$.

In summary, the following result is an immediate consequence of Theorem 1.5.

Corollary 1.7. *If G, H are countable non-type I groups, then the following are equivalent:*

1. *The unitary equivalence relations \approx_G and \approx_H are Borel bireducible.*
2. *The unitary duals \widehat{G} and \widehat{H} are Borel isomorphic.*

In the remainder of this section, we will discuss the main results of this paper. We will begin by considering the countable groups with the most complex unitary duals.

Definition 1.8. The countable group G is *representation universal* if whenever H is a countable group, then \approx_H is Borel reducible to \approx_G .

For example, the free group \mathbb{F}_∞ on countably many generators is representation universal. To see this, let G be any countable group and let $p : \mathbb{F}_\infty \rightarrow G$ be a surjective homomorphism. Then the map $\pi \mapsto \pi \circ p$ is clearly a Borel reduction from \approx_G to $\approx_{\mathbb{F}_\infty}$. More generally, the same argument shows that if G, H are countable groups and G is a homomorphic image of H , then \approx_G is Borel reducible to \approx_H . On the other hand, the following basic question remains open.

Question 1.9. Suppose that G, H are countable groups such that $G \leq H$. Does it necessarily follow that \approx_G is Borel reducible to \approx_H ?

In particular, while \mathbb{F}_∞ is embeddable in $\mathrm{SL}(3, \mathbb{Z})$, it is currently not known whether or not $\mathrm{SL}(3, \mathbb{Z})$ is representation universal. On the other hand, the following result which we will prove in Section 3 implies that all countable nonabelian free groups are representation universal; and, of course, this means that very large countable groups are also representation universal. (Recall that a group G is said to be *very large* if G has a nonabelian free quotient.)


Theorem 1.10. *The free group \mathbb{F}_2 on two generators is representation universal.*

Very little is known concerning the closure properties of the class of representation universal groups. For example, it is not known whether or not a finite extension of a representation universal group is also representation universal; and, in particular, it is not known whether countable large groups are representation universal. (Recall that a group G is said to be *large* if G has a very large subgroup of finite index.)

In Sections 4 through 6, we will mainly focus on the unitary duals of countable amenable non-type I groups. Here our main result is the following theorem. (As we will explain, Theorem 1.11 is an easy consequence of results of Elliott [12] and Sutherland [46].)

Theorem 1.11. *If G is a countable non-type I group and H is a countable amenable group, then \approx_H is Borel reducible to \approx_G . In particular, if G and H are countable amenable non-type I groups, then \approx_G and \approx_H are Borel bireducible.*

The main results of this paper can be summarized graphically by the following diagram.



Here the dotted lines indicate that it is currently not known whether or not the unitary duals of *all* countable non-type I groups are Borel isomorphic. In Sections 6 and 7, we will present some evidence that suggests that countable amenable non-type I groups might not be representation universal. If this is indeed the case, then this raises the very interesting question of how many distinct unitary duals of countably infinite groups exist up to Borel isomorphism. Of course, ideally there should exist uncountably many such unitary duals. Unfortunately, the results of Section 5 suggest the possibility that perhaps the unitary duals of all countable non-amenable groups are Borel isomorphic.

Notation 1.12. If \mathcal{H} is a (possibly finite-dimensional) separable Hilbert space, then $B(\mathcal{H})$ denotes the space of bounded linear operators $T : \mathcal{H} \rightarrow \mathcal{H}$, and $U(\mathcal{H})$ denotes the corresponding unitary group.

Remark 1.13. For the convenience of the reader, the definitions of the various Borel equivalence relations that occur in this paper have been collected together in an Appendix.

2. Preliminaries

In this section, we will recall some basic notions from the theory of countable Borel equivalence relations and from the representation theory of separable C^* -algebras. As expected, C^* -algebras will enter into our study through the canonical correspondence between the irreducible unitary representations of a countable group G and the irreducible unitary representations of the associated group C^* -algebra $C^*(G)$. Countable Borel equivalence relations will enter the picture when we study the cocycles associated with the irreducible representations of semidirect products of countable groups. (It is perhaps worth mentioning that Theorem 1.2 implies that if G is a countable non-type I group, then the unitary equivalence relation \approx_G is *not* Borel reducible to any countable Borel equivalence relation.)

2.1. Countable Borel equivalence relations and ergodic theory

Let X be a standard Borel space, i.e. a Polish space equipped with its associated σ -algebra of Borel subsets. Then the Borel equivalence relation E on X is said to be *countable* if every E -equivalence class is countable. For example, suppose that G is a countable group and that X is a *standard Borel G -space*, i.e. a standard Borel space X equipped with a

Borel action $(g, x) \mapsto g \cdot x$ of G on X . Then the corresponding orbit equivalence relation E_G^X is a countable Borel equivalence relation. Conversely, by Feldman–Moore [16], if E is an arbitrary countable Borel equivalence relation on the standard Borel space X , then there exists a countable group G and a Borel action of G on X such that $E = E_G^X$.

A detailed development of the general theory of countable Borel equivalence relations can be found in Jackson–Kechris–Louveau [29]. Here we will only recall those aspects of the theory that will play an essential role in this paper.

The least complex non-smooth countable Borel equivalence relation with respect to Borel reducibility is the *Vitali equivalence relation* E_0 , which is defined on $2^{\mathbb{N}}$ by

$$x E_0 y \Leftrightarrow x(n) = y(n) \text{ for almost all } n.$$

More precisely, by Harrington–Kechris–Louveau [23], if E is any (not necessarily countable) Borel equivalence relation, then E is non-smooth if and only if $E_0 \leq_B E$. At the other extreme, let E_∞ be the orbit equivalence relation arising from the shift action of the free group \mathbb{F}_2 on two generators on $2^{\mathbb{F}_2}$. Then, by Dougherty–Jackson–Kechris [10], E_∞ is a *universal* countable Borel equivalence relation in the sense that $F \leq_B E_\infty$ for every countable Borel equivalence relation F .

If X is a Polish space, then the Polish space of all Borel probability measures on X is denoted by $\mathcal{P}(X)$. If $\mu, \nu \in \mathcal{P}(X)$, then μ and ν are said to be *equivalent*, written $\mu \sim \nu$, if for every Borel subset $A \subseteq X$,

$$\mu(A) = 0 \Leftrightarrow \nu(A) = 0.$$

If G is a countable group and X is a standard Borel G -space, then we can define a corresponding Borel action on $\mathcal{P}(X)$ by $\mu \mapsto g_*\mu$, where $g_*\mu$ is the Borel probability measure defined by

$$g_*\mu(A) = \mu(g^{-1}(A))$$

for every Borel subset $A \subseteq X$. The Borel probability measure $\mu \in \mathcal{P}(X)$ is said to be *invariant* if $g_*\mu = \mu$ for all $g \in G$, and is said to be *quasi-invariant* if $g_*\mu \sim \mu$ for all $g \in G$. If $\mu \in \mathcal{P}(X)$ is quasi-invariant, then μ is said to be *ergodic* if every G -invariant Borel subset of X is either null or conull.

Suppose that E is a countable Borel equivalence relation on the standard Borel probability space (X, μ) . Then E is said to be *measure-preserving* if Γ preserves μ for some (equivalently every) countable group Γ acting in a Borel fashion such that $E = E_\Gamma^X$. In this case, E is said to be *ergodic* if each such Γ acts ergodically on (X, μ) . If E is an ergodic measure-preserving countable Borel equivalence relation on the standard Borel probability space (X, μ) and $A \subseteq X$ is a Borel subset such that $\mu(A) > 0$, then $E \upharpoonright A$ denotes the restriction of E to the standard Borel probability space (A, μ_A) , where $\mu_A(C) = \mu(C)/\mu(A)$.

Suppose that E, F are ergodic measure-preserving countable Borel equivalence relations on the standard Borel probability spaces $(X, \mu), (Y, \nu)$ respectively. Then E and F are said to be *orbit equivalent* if, after possibly deleting measure zero subsets, there exists a Borel bijection $\theta : X \rightarrow Y$ such that $\theta_*\mu = \nu$ and $\theta(E) = F$. More generally, E and F are said to be *weakly orbit equivalent* if there exist Borel subsets $A \subseteq X$ and $B \subseteq Y$ with $\mu(A), \nu(B) > 0$ such that $E \upharpoonright A$ and $F \upharpoonright B$ are orbit equivalent.

2.2. C^* -algebras

Let \mathcal{A} be a separable C^* -algebra. For each $n \in \{1, 2, \dots, \infty\}$, let $\text{Irr}_n(\mathcal{A})$ be the Polish space of all irreducible unitary representations of \mathcal{A} in some fixed separable Hilbert space of dimension n . Then the *spectrum* $\widehat{\mathcal{A}}$ of \mathcal{A} is the quotient of the disjoint union $\bigsqcup \text{Irr}_n(\mathcal{A})$ by the unitary equivalence relation, equipped with the corresponding quotient Borel structure. By Glimm [22], the unitary equivalence relation on $\bigsqcup \text{Irr}_n(\mathcal{A})$ is smooth if and only if \mathcal{A} is a type I C^* -algebra.² If \mathcal{A} is a non-type I separable C^* -algebra, then $\approx_{\mathcal{A}}$ denotes the unitary equivalence relation on the space $\text{Irr}_{\infty}(\mathcal{A})$ of infinite-dimensional irreducible unitary representations of \mathcal{A} .

Let G be a countable group and let $\mathbb{C}[G]$ be the corresponding group algebra. Then the *group C^* -algebra* $C^*(G)$ is the completion of $\mathbb{C}[G]$ with respect to the norm

$$\|x\|_u = \sup_{\pi} \|\pi(x)\|,$$

where the supremum is taken over all cyclic $*$ -representations $\pi : \mathbb{C}[G] \rightarrow B(\mathcal{H})$. It is well-known that there is a canonical correspondence between the irreducible unitary representations of $C^*(G)$ and the irreducible unitary representations of G . (For example, see Dixmier [9, Section 13.9].) Furthermore, this canonical correspondence witnesses that $\approx_{C^*(G)}$ and \approx_G are Borel bireducible.

Remark 2.1. In 1967, Dixmier raised the question of whether the spectra of any two non-type I separable unital C^* -algebras are Borel isomorphic. In fact, this question is equivalent to asking whether countable amenable non-type I groups are representation universal. To see this, recall that if \mathcal{A} is any separable unital C^* -algebra, then \mathcal{A} is generated by a countable sequence of unitaries. It follows that there exists a surjective unital $*$ -homomorphism $f : C^*(\mathbb{F}_{\infty}) \rightarrow \mathcal{A}$ and so

$$\approx_{\mathcal{A}} \leq_B \approx_{C^*(\mathbb{F}_{\infty})} \sim_B \approx_{\mathbb{F}_{\infty}}.$$

On the other hand, combining Theorem 4.2 and Corollary 6.2 shows that if G is a countable amenable non-type I group and \mathcal{A} is any non-type I separable unital C^* -algebra, then

$$\approx_G \sim_B \approx_{C^*(G)} \leq_B \approx_{\mathcal{A}}.$$

Thus the following statements are equivalent:

- \approx_G is Borel bireducible with $\approx_{\mathbb{F}_{\infty}}$ for some (equivalently every) countable amenable non-type I group G .
- $\widehat{\mathcal{A}}$ is Borel isomorphic with $\widehat{C^*(\mathbb{F}_{\infty})}$ for every non-type I separable unital C^* -algebra \mathcal{A} .

² The definition of a type I C^* -algebra can be found in Arveson [2]. Once again, this notion will play no role in this paper.

3. The representation universality of \mathbb{F}_2

In this section, we will prove that the free group \mathbb{F}_2 on the two generators $\{a, b\}$ is representation universal. We have already noted that the free group \mathbb{F}_∞ on infinitely many generators is representation universal. Thus it is enough to show that $\approx_{\mathbb{F}_\infty}$ is Borel reducible to $\approx_{\mathbb{F}_2}$.

Notation 3.1. If $H \leq G$ are countable groups and π is a unitary representation of H , then $\text{ind}_H^G(\pi)$ denotes the corresponding induced representation of G .

We will make use of the following well-known irreducibility criterion for representations induced from normal subgroups of countable groups. (For example, see the discussion in Quigg [44, Section 1].)

Definition 3.2. Suppose that G is a countable group and that $N \trianglelefteq G$ is a normal subgroup. If π is a unitary representation of N and $g \in G$, then π^g denotes the unitary representation of N defined by $\pi^g(x) = \pi(gxg^{-1})$.

Theorem 3.3. Let G be a countable group and let $N \trianglelefteq G$ be a normal subgroup. If π is an irreducible unitary representation of N , then the following are equivalent:

- (i) $\text{ind}_N^G(\pi)$ is an irreducible unitary representation of G .
- (ii) π^g is not unitarily equivalent to π for all $g \in G \setminus N$.

Let $H = \ker h \trianglelefteq \mathbb{F}_2$, where $h : \mathbb{F}_2 \rightarrow \mathbb{Z}$ is the homomorphism such that $h(a) = 1$ and $h(b) = 0$. Then $T = \{a^n \mid n \in \mathbb{Z}\}$ is a set of coset representatives of H in \mathbb{F}_2 ; and hence, by Schreier’s Theorem, it follows that $S = \{a^n b a^{-n} \mid n \in \mathbb{Z}\}$ is a basis of H . (For example, see Serre [45, Proposition 16].) From now on, let $b_n = a^n b a^{-n}$, so that $a b_n a^{-1} = b_{n+1}$ for each $n \in \mathbb{Z}$. Let $f : \mathbb{N} \rightarrow \mathbb{N}$ be the function defined inductively by

$$f(0) = 0, \quad f(n + 1) = 1 + \sum_{i=0}^n f(i),$$

and let $K = \{f(n) \mid n \in \mathbb{N}\}$. Then it is easily checked that $|(\ell + K) \cap K| \leq 1$ for all $0 \neq \ell \in \mathbb{Z}$. Hence, if we let $c_n = b_{f(n)}$, then the infinite subset $C = \{c_n \mid n \in \mathbb{N}\} \subseteq S$ satisfies $|a^\ell C a^{-\ell} \cap C| \leq 1$ for all $0 \neq \ell \in \mathbb{Z}$. From now on, we will identify \mathbb{F}_∞ with the subgroup $\langle c_n \mid n \in \mathbb{N} \rangle$ of H and we will identify $\text{Irr}_\infty(\mathbb{F}_\infty)$ with the corresponding subspace of $\text{Irr}_\infty(H)$, i.e. we will identify each $\pi \in \text{Irr}_\infty(\mathbb{F}_\infty)$ with the corresponding representation $\pi \in \text{Irr}_\infty(H)$ obtained by setting $\pi(s) = 1$ for all $s \in S \setminus C$. Of course, with this identification, if $\pi, \sigma \in \text{Irr}_\infty(\mathbb{F}_\infty)$, then

$$\pi \approx_{\mathbb{F}_\infty} \sigma \Leftrightarrow \pi \approx_H \sigma.$$

Let $\theta : \text{Irr}_\infty(\mathbb{F}_\infty) \rightarrow \text{Rep}(\mathbb{F}_2)$ be the Borel map defined by $\pi \mapsto \theta_\pi = \text{ind}_H^{\mathbb{F}_2}(\pi)$. Then we will show that θ is a Borel reduction from $\approx_{\mathbb{F}_\infty}$ to $\approx_{\mathbb{F}_2}$.

Lemma 3.4. If $\pi \in \text{Irr}_\infty(\mathbb{F}_\infty)$, then $\theta_\pi \in \text{Irr}_\infty(\mathbb{F}_2)$.

Proof. By Theorem 3.3, it is enough to show that if $\ell \in \mathbb{Z} \setminus \{0\}$, then π^{a^ℓ} is not unitarily equivalent to π . So suppose that π^{a^ℓ} and π are unitarily equivalent. Then since

$a^\ell Sa^{-\ell} = S$ and $|a^\ell Ca^{-\ell} \cap C| \leq 1$, there exists at most one basis element $s \in S$ such that $\pi(s) \neq 1$. But this means that π is a 1-dimensional representation of H , which is a contradiction. \square

Lemma 3.5. θ is a Borel reduction from $\approx_{\mathbb{F}_\infty}$ to $\approx_{\mathbb{F}_2}$.

Proof. First note that if $\pi, \sigma \in \text{Irr}_\infty(\mathbb{F}_\infty)$ and $\pi \approx_{\mathbb{F}_\infty} \sigma$, then $\theta_\pi \approx_{\mathbb{F}_2} \theta_\sigma$. (For example, see Folland [17, Proposition 6.9].) Conversely, suppose that $\theta_\pi \approx_{\mathbb{F}_2} \theta_\sigma$. Then $\theta_\pi \upharpoonright H = \bigoplus_{\ell \in \mathbb{Z}} \pi^{a^\ell}$ and $\theta_\sigma \upharpoonright H = \bigoplus_{\ell \in \mathbb{Z}} \sigma^{a^\ell}$ are unitarily equivalent representations of H . Arguing as in the proof of Lemma 3.4 shows that if $\ell \in \mathbb{Z} \setminus \{0\}$, then π is not unitarily equivalent to σ^{a^ℓ} . Hence $\pi \approx_H \sigma$ and so $\pi \approx_{\mathbb{F}_\infty} \sigma$. \square

4. Representations of locally finite groups

In this section, we will derive the following weak form of Theorem 1.11 from a more general theorem of Elliott [12] on the Mackey Borel structures of the spectra of approximately finite-dimensional C^* -algebras.

Theorem 4.1. *If G is a countable non-type I group and H is a countable locally finite group, then \approx_H is Borel reducible to \approx_G .*

Recall that a C^* -algebra \mathcal{A} is said to be *approximately finite-dimensional* if \mathcal{A} is the closure of the union $\bigcup_{n \in \mathbb{N}} \mathcal{A}_n$ of an increasing chain $\mathcal{A}_0 \subseteq \mathcal{A}_1 \subseteq \dots$ of finite-dimensional C^* -subalgebras. For example, let H be a countable locally finite group and express $H = \bigcup_{n \in \mathbb{N}} H_n$ as the union of an increasing chain of finite subgroups H_n . Then the group C^* -algebra $C^*(H)$ is the closure of the group algebra $\mathbb{C}[H] = \bigcup_{n \in \mathbb{N}} \mathbb{C}[H_n]$ and hence $C^*(H)$ is approximately finite-dimensional. Also recall that if G is any countable group, then $\approx_{C^*(G)}$ and \approx_G are Borel bireducible. Consequently, Theorem 4.1 is an immediate consequence of the following result of Elliott [12].

Theorem 4.2 (Elliott [12]). *If \mathcal{A} is an approximately finite-dimensional C^* -algebra and \mathcal{B} is a non-type I separable C^* -algebra, then $\approx_{\mathcal{A}}$ is Borel reducible to $\approx_{\mathcal{B}}$.*

In the remainder of this section, we will point out a curious result concerning the asymptotic representation theory of finite groups, which follows easily from Thoma’s theorem [47], together with the theory of approximately finite-dimensional C^* -algebras.

Notation 4.3. If $K \leq L$ are finite groups and π, σ are irreducible representations of K, L respectively, then $\text{mult}_\pi(\sigma \upharpoonright K)$ denotes the multiplicity with which π occurs in the representation $\sigma \upharpoonright K$.

Theorem 4.4. *If $G = \bigcup_{n \in \mathbb{N}} G_n$ is the union of the strictly increasing chain of finite subgroups G_n , then the following statements are equivalent:*

- (i) G is not abelian-by-finite.
- (ii) *There exists an increasing sequence $(\ell_n \mid n \in \mathbb{N})$ and irreducible representations $\pi_n \in \text{Irr}(G_{\ell_n})$ such that for all $n \in \mathbb{N}$, $\text{mult}_{\pi_n}(\pi_{n+1} \upharpoonright G_{\ell_n}) > 1$.*

Proof. To see that (ii) implies (i), suppose that G contains an abelian subgroup A such that $[G : A] = d < \infty$. Then $A_n = G_n \cap A$ is an abelian subgroup of G_n such that $[G_n : A_n] \leq d$; and a simple application of the Frobenius reciprocity theorem shows that $\deg \pi \leq [G_n : A_n]$ for all $\pi \in \text{Irr}(G_n)$. (For example, see Remark 17.12 in Huppert [28].) Of course, if $n < m$ and $\pi \in \text{Irr}(G_n)$, $\chi \in \text{Irr}(G_m)$ satisfy $\text{mult}_\pi(\chi \upharpoonright G_n) > 1$, then $\deg \chi > \deg \pi$. Hence statement (ii) fails.

Conversely, suppose that G is not abelian-by-finite. Then, by Thoma [47], G is not of type I and so the corresponding group C^* -algebra $C^*(G)$ is a non-type I approximately finite-dimensional C^* -algebra. Notice that the Bratteli diagram associated with the increasing chain

$$\mathbb{C}[G_0] \subseteq \mathbb{C}[G_1] \subseteq \dots$$

of finite-dimensional C^* -subalgebras is precisely the branching diagram of the irreducible representations of the groups G_n in the corresponding increasing chain

$$G_0 \subseteq G_1 \subseteq \dots$$

of finite subgroups of G . Thus statement (ii) is an immediate consequence of the Lazar–Taylor characterization [38, Theorem 3.13] of type I approximately finite-dimensional C^* -algebras in terms of the multiplicities of paths through their Bratteli diagrams. \square

Question 4.5. Is there an “elementary” proof of Theorem 4.4? In other words, is Theorem 4.4 a consequence of some result concerning the branching diagrams of pairs $K \leq L$ of finite groups?

5. Cocycles and representations

Suppose that $G = A \rtimes \Gamma$ is a semidirect product of the countable groups A and Γ , where A is an infinite abelian group. Then it is well-known that the irreducible representations of G are determined by the Γ -quasi-invariant ergodic measures on the unitary dual \widehat{A} , together with the “irreducible” cocycles for the dual action of Γ on \widehat{A} . Furthermore, the unitary equivalence relation \approx_G is determined by:

- the measure equivalence relation on the space of Γ -quasi-invariant ergodic measures, together with
- the cocycle equivalence relation on the space of “irreducible” cocycles.

In his remarkable paper [46], Sutherland proved that if Γ is amenable and acts freely and ergodically on \widehat{A} , then the unitary equivalence relation \approx_H of every countable amenable group H is Borel reducible to the cocycle equivalence relation on the space of “irreducible” cocycles $\sigma : \Gamma \times \widehat{A} \rightarrow U(\mathcal{H})$, where \mathcal{H} is an infinite-dimensional separable Hilbert space. As we will see, in combination with Elliott’s work [12], this easily implies that the unitary duals of any two countable amenable non-type I groups are Borel isomorphic. In this section, we will begin our account of Sutherland’s work, slightly expanded and generalized so that it is also applicable to non-amenable groups.

If G is a countable group and \mathcal{H} is a (possibly finite-dimensional) separable Hilbert space, then $\text{Rep}(G, \mathcal{H})$ denotes the Polish space of all unitary representations of G in \mathcal{H} and $\text{Irr}(G, \mathcal{H})$ denotes the subspace of irreducible representations. For each pair $\pi, \sigma \in \text{Rep}(G, \mathcal{H})$, let

$$\mathcal{C}_G(\pi, \sigma) = \{S \in B(\mathcal{H}) \mid S \circ \pi(g) = \sigma(g) \circ S \text{ for all } g \in G\}.$$

Recall that, by Schur’s Lemma, if $\pi \in \text{Rep}(G, \mathcal{H})$, then π is irreducible if and only if $\mathcal{C}_G(\pi, \pi)$ is the vector space of scalar multiples of the identity operator. Furthermore, if $\pi, \sigma \in \text{Rep}(G, \mathcal{H})$ are both irreducible and $0 \neq T \in \mathcal{C}_G(\pi, \sigma)$, then T is a non-zero scalar multiple of a unitary operator and so π, σ are unitarily equivalent.

We are now ready to begin our discussion of cocycles and the associated unitary representations. Suppose that the countable group Γ acts freely and ergodically via measure-preserving Borel maps on the standard probability space (X, μ) . Then a Borel map $\sigma : \Gamma \times X \rightarrow U(\mathcal{H})$ is said to be a *cocycle* if for all $\gamma, \gamma' \in \Gamma$,

$$\sigma(\gamma\gamma', x) = \sigma(\gamma, \gamma' \cdot x)\sigma(\gamma', x) \quad \mu\text{-a.e. } x \in X.$$

Let $Z^1(\Gamma \times X, U(\mathcal{H}))$ be the standard Borel space of Borel cocycles, where we identify two such maps σ, σ' if for all $\gamma \in \Gamma$,

$$\sigma(\gamma, x) = \sigma'(\gamma, x) \quad \mu\text{-a.e. } x \in X.$$

Then for each cocycle $\sigma \in Z^1(\Gamma \times X, U(\mathcal{H}))$, we can define a corresponding unitary representation π_σ of Γ on the Hilbert space $L^2(X, \mathcal{H})$ by setting

$$(\pi_\sigma(\gamma) \cdot f)(x) = \sigma(\gamma^{-1}, x)^{-1} f(\gamma^{-1} \cdot x) \tag{5.1}$$

for each $\gamma \in \Gamma, x \in X$ and $f \in L^2(X, \mathcal{H})$.

Definition 5.1 (Sutherland [46]). If $\sigma, \tau \in Z^1(\Gamma \times X, U(\mathcal{H}))$, then $\text{Hom}_\Gamma(\sigma, \tau)$ consists of the bounded Borel maps $T : X \rightarrow B(\mathcal{H})$ such that for all $\gamma \in \Gamma$,

$$\sigma(\gamma, x)T(x) = T(\gamma \cdot x)\tau(\gamma, x) \quad \mu\text{-a.e. } x \in X,$$

where we again identify two such maps if they agree μ -a.e.

Here a Borel map $T : X \rightarrow B(\mathcal{H})$ is *bounded* if $\text{ess sup}_{x \in X} \|T(x)\| < \infty$. In this case, we can define a corresponding bounded linear operator $\int^\oplus T = \int^\oplus T(x) d\mu(x)$ on $L^2(X, \mathcal{H})$ by $[(\int^\oplus T)f](x) = T(x)f(x)$.

Lemma 5.2. *If $\sigma, \tau \in Z^1(\Gamma \times X, U(\mathcal{H}))$ and $T : X \rightarrow B(\mathcal{H})$ is a bounded Borel map, then $T \in \text{Hom}_\Gamma(\sigma, \tau)$ if and only if $\int^\oplus T \in \mathcal{C}_\Gamma(\pi_\sigma, \pi_\tau)$.*

Proof. A routine calculation. □

Next, by passing from Γ to a suitable semidirect product $G = A \rtimes \Gamma$ and then extending each π_σ to a corresponding unitary representation of G , we will ensure that for every $S \in \mathcal{C}_G(\pi_\sigma, \pi_\tau)$, there exists a bounded Borel map $T : X \rightarrow B(\mathcal{H})$ such that $S = \int^\oplus T$. In more detail, suppose that A is a countably infinite abelian group and that $\gamma \mapsto \varphi_\gamma$ is a homomorphism from Γ to $\text{Aut}(A)$. Then we can define an induced action of Γ on the unitary dual $X = \text{Irr}_1(A)$ of A by $\gamma \cdot x = x \circ \varphi_\gamma^{-1}$. Next suppose that μ is a Γ -invariant probability measure on X such that the action of Γ on (X, μ) is essentially free and ergodic. Let $G = A \rtimes_\varphi \Gamma$ be the corresponding semidirect product defined by

$$(a, \gamma)(a', \gamma') = (a\varphi_\gamma(a'), \gamma\gamma').$$

Then for each cocycle $\sigma \in Z^1(\Gamma \times X, U(\mathcal{H}))$, we can extend the unitary representation π_σ of Γ on $L^2(X, \mathcal{H})$ to a unitary representation of $G = A \rtimes_\varphi \Gamma$ by setting

$$(\pi_\sigma(a) \cdot f)(x) = x(a)f(x). \tag{5.2}$$

Notice that the direct integral decomposition of $\pi_0 = \pi_\sigma \upharpoonright A$ as $\pi_0 = \int^\oplus x d\mu(x)$ on $L^2(X, \mathcal{H}) = \int^\oplus \mathcal{H}_x d\mu(x)$, where each $\mathcal{H}_x = \mathcal{H}$, is the canonical direct integral decomposition of π_0 into primary representations. It follows that if $S \in \mathcal{C}_A(\pi_0, \pi_0)$, then there exists a bounded Borel map $T : X \rightarrow B(\mathcal{H})$ such that $S = \int^\oplus T$. (For example, see Nielsen [41].)

Proposition 5.3. *With the above hypotheses, if $\sigma, \tau \in Z^1(\Gamma \times X, U(\mathcal{H}))$, then the map $T \mapsto \int^\oplus T$ is a vector space isomorphism between $\text{Hom}_\Gamma(\sigma, \tau)$ and $\mathcal{C}_G(\pi_\sigma, \pi_\tau)$.*

Proof. If $S \in \mathcal{C}_G(\pi_\sigma, \pi_\tau)$, then $S \in \mathcal{C}_A(\pi_0, \pi_0)$, where $\pi_0 = \pi_\sigma \upharpoonright A = \pi_\tau$, and so there exists a bounded Borel map $T : X \rightarrow B(\mathcal{H})$ such that $S = \int^\oplus T$. Hence the result follows from Lemma 5.2. □

In particular, by Schur’s Lemma, if $\sigma \in Z^1(\Gamma \times X, U(\mathcal{H}))$, then the corresponding unitary representation π_σ of $G = A \rtimes \Gamma$ is irreducible if and only if $\text{Hom}_\Gamma(\sigma, \sigma)$ consists of the constant μ -a.e. maps $T : X \rightarrow B(\mathcal{H})$ taking values in the vector space of scalar multiples of the identity. In this case, we say that σ is an *irreducible* cocycle. Let

$$\text{Irr}(\Gamma \times X, U(\mathcal{H})) \subseteq Z^1(\Gamma \times X, U(\mathcal{H}))$$

be the set of irreducible cocycles. Since $\text{Irr}(\Gamma \times X, U(\mathcal{H}))$ is the inverse image of the set of irreducible representations under the Borel map $\sigma \mapsto \pi_\sigma$, it is a Borel subset of $Z^1(\Gamma \times X, U(\mathcal{H}))$, and hence a standard Borel space.

Next suppose that $\sigma, \tau \in \text{Irr}(\Gamma \times X, U(\mathcal{H}))$ are irreducible cocycles. Then Proposition 5.3 implies that the corresponding representations π_σ and π_τ are unitarily equivalent if and only if there exists $0 \neq T \in \text{Hom}_\Gamma(\sigma, \tau)$ such that $S = \int^\oplus T$ satisfies

$$S \circ \pi_\sigma(g) = \pi_\tau(g) \circ S \quad \text{for all } g \in G.$$

Furthermore, in this case, once again by Schur’s Lemma, S is a non-zero scalar multiple of a unitary operator. Hence, replacing T by a suitable scalar multiple, we can assume that S is a unitary operator. By Nielsen [41, Proposition 6.1(d)], it follows that $T(x) \in U(\mathcal{H})$

for μ -a.e. $x \in X$ and so we can regard T as an element of the Polish group $L(X, U(\mathcal{H}))$. Here $L(X, U(\mathcal{H}))$ is the space of Borel maps $T : X \rightarrow U(\mathcal{H})$ equipped with the group operation of pointwise multiplication, where we identify two such maps if they agree μ -a.e. (For more details, see Kechris [34, Section 19].) The group $L(X, U(\mathcal{H}))$ acts on $Z^1(\Gamma \times X, U(\mathcal{H}))$ by

$$(T \cdot \sigma)(\gamma, x) = T(\gamma \cdot x)\sigma(\gamma, x)T(x)^{-1};$$

and it is easily checked that $\text{Irr}(\Gamma \times X, U(\mathcal{H}))$ is invariant under this action. Summing up, we now see that the map $\sigma \mapsto \pi_\sigma$ is a Borel reduction from the orbit equivalence relation for the action of $L(X, U(\mathcal{H}))$ on $\text{Irr}(\Gamma \times X, U(\mathcal{H}))$ to the unitary equivalence relation on $\text{Irr}(G, L^2(X, U(\mathcal{H})))$.

Conversely, the following result will enable us to reduce the unitary equivalence relations \approx_H for various countable groups H to suitable cocycle equivalence relations.

Proposition 5.4. *Let Γ, H be countable groups and let $\rho : \Gamma \rightarrow H$ be a surjective homomorphism. Suppose that:*

- (i) Γ acts freely and ergodically via measure-preserving Borel maps on the standard probability space (X, μ) .
- (ii) $\ker \rho$ acts ergodically on (X, μ) .

Let $\pi \mapsto \sigma_\pi$ be the Borel map from $\text{Rep}(H, \mathcal{H})$ to $Z^1(\Gamma \times X, U(\mathcal{H}))$ defined by

$$\sigma_\pi(\gamma, x) = (\pi \circ \rho)(\gamma), \quad \gamma \in \Gamma, x \in X.$$

If $\sigma, \theta \in \text{Rep}(H, \mathcal{H})$, then $\text{Hom}_\Gamma(\sigma_\pi, \sigma_\theta)$ is the vector space of constant maps taking values in $\mathcal{C}_H(\pi, \theta)$.

Proof. Clearly if $T \in \mathcal{C}_H(\pi, \theta)$, then the constant map $x \mapsto T$ lies in $\text{Hom}_\Gamma(\sigma_\pi, \sigma_\theta)$. Conversely, if $T \in \text{Hom}_\Gamma(\sigma_\pi, \sigma_\theta)$, then for all $\gamma \in \Gamma$,

$$\sigma_\pi(\gamma, x)T(x) = T(\gamma \cdot x)\sigma_\theta(\gamma, x) \quad \mu\text{-a.e. } x \in X.$$

In particular, for each $\gamma \in \ker \rho$, we have

$$T(x) = T(\gamma \cdot x) \quad \mu\text{-a.e. } x \in X.$$

Since $\ker \rho$ acts ergodically on X , it follows that T is μ -a.e. constant; and since the homomorphism $\rho : \Gamma \rightarrow H$ is surjective, this constant value lies in $\mathcal{C}_H(\pi, \theta)$. \square

Following Kechris [34, Section 20(B)], we can reformulate the notion of a cocycle for an arbitrary countable ergodic measure-preserving equivalence relation E on the standard Borel probability space (X, μ) as follows. First let M be the σ -finite Borel measure defined on E by

$$M(A) = \int |A_x| d\mu(x),$$

where $A_x = \{y \in X \mid (x, y) \in A\}$. (For example, see Kechris [34, Section 6.B].) Then $Z^1(E, U(\mathcal{H}))$ denotes the standard Borel space of Borel maps $\sigma : E \rightarrow U(\mathcal{H})$ such that

$$\sigma(x, z) = \sigma(y, z)\sigma(x, y)$$

for all $x E y E z$ in some E -invariant Borel subset of X of μ -measure 1, where we identify two such maps σ, σ' if

$$\sigma(x, z) = \sigma'(x, z) \quad M\text{-a.e. } (x, z) \in E.$$

If $\sigma, \tau \in Z^1(E, U(\mathcal{H}))$, then we define $\text{Hom}_E(\sigma, \tau)$ to consist of the bounded Borel maps $T : X \rightarrow B(\mathcal{H})$ such that

$$\sigma(x, z)T(x) = T(z)\tau(x, z) \quad M\text{-a.e. } (x, z) \in E,$$

where we identify two such maps if they agree μ -a.e.

Of course, if $E = E_\Gamma^X$ is the orbit equivalence relation arising from a free ergodic measure-preserving action on (X, μ) , then for each $\sigma \in Z^1(\Gamma \times X, U(\mathcal{H}))$, we can define a corresponding cocycle $\sigma' \in Z^1(E, U(\mathcal{H}))$ by

$$\sigma'(x, z) = \sigma(g, x),$$

where $g \in \Gamma$ is the *unique* $g \in \Gamma$ such that $g \cdot x = z$; and for each $\tau \in Z^1(E, U(\mathcal{H}))$, we can define a corresponding cocycle $\tau' \in Z^1(\Gamma \times X, U(\mathcal{H}))$ by

$$\tau'(g, x) = \tau(x, g \cdot x).$$

Notice that if $\sigma_1, \sigma_2 \in Z^1(\Gamma \times X, U(\mathcal{H}))$ and $\sigma'_1, \sigma'_2 \in Z^1(E, U(\mathcal{H}))$ are the corresponding cocycles as above, then for all bounded Borel maps $T : X \rightarrow B(\mathcal{H})$,

$$T \in \text{Hom}_\Gamma(\sigma_1, \sigma_2) \Leftrightarrow T \in \text{Hom}_E(\sigma'_1, \sigma'_2).$$

Definition 5.5. A cocycle $\sigma \in Z^1(E, U(\mathcal{H}))$ is said to be *irreducible* if $\text{Hom}_E(\sigma, \sigma)$ consists of the constant μ -a.e. maps $T : X \rightarrow B(\mathcal{H})$ taking values in the vector space of scalar multiples of the identity.

For each (possibly finite-dimensional) separable Hilbert space \mathcal{H} , let $\text{Irr}(E, U(\mathcal{H}))$ be the set of irreducible cocycles.

Definition 5.6. Suppose that \mathcal{H} is an infinite-dimensional separable Hilbert space. Then \equiv_E denotes the orbit equivalence relation arising from the action of $L(X, U(\mathcal{H}))$ on $\text{Irr}(E, U(\mathcal{H}))$ defined by

$$(f \cdot \sigma)(x, z) = f(z)\sigma(x, z)f(x)^{-1}$$

for each $f \in L(X, U(\mathcal{H}))$ and $\sigma \in \text{Irr}(E, U(\mathcal{H}))$.

If $E = E_\Gamma^X$, then we will often identify $Z^1(E, U(\mathcal{H}))$ with $Z^1(\Gamma \times X, U(\mathcal{H}))$ and $\text{Irr}(E, U(\mathcal{H}))$ with $\text{Irr}(\Gamma \times X, U(\mathcal{H}))$.

It is well-known that if two ergodic measure-preserving actions of countable groups are orbit equivalent, then the associated cocycle machineries are isomorphic. (For example, see Feldman–Moore [16].) In fact, as observed by Kechris [34, Section 20(G)], the same is true when the actions are weakly orbit equivalent.

Lemma 5.7. *Suppose that E and F are countable ergodic measure-preserving equivalence relations on the standard Borel probability spaces $(X, \mu), (Y, \nu)$ respectively. If E and F are weakly orbit equivalent, then \equiv_E and \equiv_F are Borel bireducible.*

Proof. It is enough to prove the lemma for the special case where $(Y, \nu) = (A, \mu_A)$ for some Borel a.e.-complete section $A \subseteq X$, i.e. A is a Borel subset which meets μ -a.e. E -class and μ_A is the Borel probability measure defined on A by $\mu_A(Z) = \mu(Z)/\mu(A)$. For each $\sigma \in Z^1(E, U(\mathcal{H}))$, let $\sigma \upharpoonright A \in Z^1(E \upharpoonright A, U(\mathcal{H}))$ be the restriction of σ to A . By Kechris [34, Section 20(G)], if $\sigma_1, \sigma_2 \in Z^1(E, U(\mathcal{H}))$, then

$$\sigma_1 \equiv_E \sigma_2 \Leftrightarrow \sigma_1 \upharpoonright A \equiv_{E \upharpoonright A} \sigma_2 \upharpoonright A.$$

Next let $f_A : X \rightarrow A$ be a Borel map such that $f_A(x) \in E x$ for μ -a.e. $x \in X$ and such that $f_A \upharpoonright A = \text{Id}_A$. Then for each $\tau \in Z^1(E \upharpoonright A, U(\mathcal{H}))$, we can define a corresponding cocycle $\tau^A \in Z^1(E, U(\mathcal{H}))$ by $\tau^A(x, z) = \tau(f_A(x), f_A(z))$. It is easily seen that if $\tau_1, \tau_2 \in Z^1(E \upharpoonright A, U(\mathcal{H}))$, then

$$\tau_1 \equiv_{E \upharpoonright A} \tau_2 \Leftrightarrow \tau_1^A \equiv_E \tau_2^A.$$

Furthermore, by Kechris [34, Section 20(G)], for each $\sigma \in Z^1(E, U(\mathcal{H}))$,

$$\sigma \equiv_E (\sigma \upharpoonright A)^A.$$

Thus it is enough to show that if $\tau \in Z^1(E \upharpoonright A, U(\mathcal{H}))$, then

$$\tau \in \text{Irr}(E \upharpoonright A, U(\mathcal{H})) \Leftrightarrow \tau^A \in \text{Irr}(E, U(\mathcal{H})).$$

First suppose that $\tau \notin \text{Irr}(E \upharpoonright A, U(\mathcal{H}))$. Then there exists $T \in \text{Hom}_{E \upharpoonright A}(\tau, \tau)$ such that T is not a scalar multiple of the identity ν -a.e. Let $T^A : X \rightarrow B(\mathcal{H})$ be the bounded Borel map defined by $T^A(x) = T(f_A(x))$. Then $T^A \in \text{Hom}_E(\tau^A, \tau^A)$ witnesses that $\tau^A \notin \text{Irr}(E, U(\mathcal{H}))$. Conversely, suppose that $\tau \in \text{Irr}(E \upharpoonright A, U(\mathcal{H}))$ and let $T \in \text{Hom}_E(\tau^A, \tau^A)$. Then

$$\tau(x, z)T(x) = T(z)\tau(x, z) \quad M\text{-a.e. } (x, z) \in E \upharpoonright A,$$

and hence there exists $\lambda \in \mathbb{C}$ such that $T(x) = \lambda \text{Id}_{\mathcal{H}}$ for μ -a.e. $x \in A$. Since

$$\tau^A(x, f_A(x))T(x) = T(f_A(x))\tau^A(x, f_A(x)) \quad \mu\text{-a.e. } x \in X,$$

it follows that $T(x) = \lambda \text{Id}_{\mathcal{H}}$ for μ -a.e. $x \in X$. Thus $\tau^A \in \text{Irr}(E, U(\mathcal{H}))$. □

In the next section, following Sutherland [46], we will use the flexibility inherent in Lemma 5.7 to compare the unitary duals of countable amenable non-type I groups. For now, we will just record the following result.

Theorem 5.8. $\approx_{\mathbb{F}_2}$ is Borel bireducible with \equiv_{E_∞} .

Proof. In order to see that \equiv_{E_∞} is Borel reducible to $\approx_{\mathbb{F}_2}$, let μ be the usual product probability measure on $2^{\mathbb{F}_2}$ and identify $\text{Irr}(E_\infty, U(\mathcal{H}))$ with $\text{Irr}(\mathbb{F}_2 \times 2^{\mathbb{F}_2}, U(\mathcal{H}))$. Let C_2 be the cyclic group of order 2 and let

$$G = C_2 \text{ wr } \mathbb{F}_2 = A \rtimes \mathbb{F}_2,$$

where A is the base group of the wreath product. Then the induced action of \mathbb{F}_2 on the unitary dual of A is isomorphic to the shift action of \mathbb{F}_2 on $(2^{\mathbb{F}_2}, \mu)$. Consider the Borel map $\sigma \mapsto \pi_\sigma$ from $Z^1(\mathbb{F}_2 \times 2^{\mathbb{F}_2}, U(\mathcal{H}))$ to $\text{Rep}(G, L^2(2^{\mathbb{F}_2}, \mathcal{H}))$ defined by (5.1) and (5.2). Proposition 5.3 and Schur’s Lemma imply that

- (i) $\text{Irr}(\mathbb{F}_2 \times 2^{\mathbb{F}_2}, U(\mathcal{H}))$ is mapped to $\text{Irr}(G, L^2(2^{\mathbb{F}_2}, \mathcal{H}))$; and
- (ii) if $\sigma, \tau \in \text{Irr}(\mathbb{F}_2 \times 2^{\mathbb{F}_2}, U(\mathcal{H}))$, then $\sigma \equiv_{E_\infty} \tau$ if and only if $\pi_\sigma \approx_G \pi_\tau$.

Thus \equiv_{E_∞} is Borel reducible to \approx_G ; and Theorem 1.10 implies that \equiv_{E_∞} is Borel reducible to $\approx_{\mathbb{F}_2}$.

In order to see that $\approx_{\mathbb{F}_2}$ is Borel reducible to \equiv_{E_∞} , let ν be the usual product probability measure on $2^{\mathbb{F}_3}$ and consider the shift action of \mathbb{F}_3 on $(2^{\mathbb{F}_3}, \nu)$. Let $\rho : \mathbb{F}_3 \rightarrow \mathbb{F}_2$ be a surjective homomorphism. Since the shift action of \mathbb{F}_3 on $(2^{\mathbb{F}_3}, \nu)$ is strongly mixing, $\ker \rho$ acts ergodically on $(2^{\mathbb{F}_3}, \nu)$. It follows that the Borel map $\pi \mapsto \sigma_\pi$ from $\text{Rep}(\mathbb{F}_2, \mathcal{H})$ to $Z^1(\mathbb{F}_3 \times 2^{\mathbb{F}_3}, U(\mathcal{H}))$ given by Proposition 5.4 induces a Borel reduction from $\approx_{\mathbb{F}_2}$ to \equiv_E , where $E = E_{\mathbb{F}_3}^{2^{\mathbb{F}_3}}$. By Bowen [5], E is weakly orbit equivalent with $E_\infty = E_{\mathbb{F}_2}^{2^{\mathbb{F}_2}}$; and hence, by Lemma 5.7, \equiv_E is Borel bireducible with \equiv_{E_∞} . \square

Remark 5.9. As we mentioned earlier, it is currently not known whether every countable group containing a nonabelian free subgroup is representation universal. On the other hand, the methods of this section, together with the “measurable-group-theoretical solution to von Neumann’s problem” of Gaboriau–Lyons [19], suggest the possibility that perhaps every countable non-amenable group is representation universal (cf. Epstein [13]).

6. Representations of amenable groups

In the first part of this section, combining the results of Elliott [12] and Sutherland [46], we will prove that if G and H are countable amenable non-type I groups, then the unitary equivalence relations \approx_G and \approx_H are Borel bireducible. Then, in the remainder of this section, we will discuss a possible (albeit very optimistic) strategy for showing that countable amenable non-type I groups are *not* representation universal.

Theorem 6.1. *If H is a countable amenable non-type I group, then \approx_H is Borel bireducible with \equiv_{E_0} .*

Corollary 6.2. *If G and H are countable amenable non-type I groups, then \approx_G and \approx_H are Borel bireducible.* \square

Corollary 6.3. *If G is a countable non-type I group and H is any countable amenable group, then \approx_H is Borel reducible to \approx_G .*

Proof. If H is a type I group, then \approx_H is smooth and hence \approx_H is Borel reducible to \approx_G . Thus we can suppose that H is a countable amenable non-type I group. Let K be a countable locally finite non-type I group. Then by Theorem 6.1, \approx_H and \approx_K are Borel bireducible; and by Theorem 4.1, \approx_K is Borel reducible to \approx_G . \square

It should be stressed that the following is essentially just a slight reformulation of the argument in Sutherland [46]. In particular, the key idea of using the Ornstein–Weiss Theorem in this setting is due to Sutherland.

Proof of Theorem 6.1. To see that \approx_H is Borel reducible to \equiv_{E_0} , let $\Gamma = H \times \mathbb{Z}$ and let $\rho : \Gamma \rightarrow H$ be the canonical surjective homomorphism. Let ν be the usual product probability measure on $X = 2^\Gamma$ and let $E = E_\Gamma^{2^\Gamma}$. Then, arguing as in the proof of Theorem 5.8, it follows that the Borel map $\pi \mapsto \sigma_\pi$ from $\text{Rep}(H, \mathcal{H})$ to $Z^1(\Gamma \times 2^\Gamma, U(\mathcal{H}))$

given by Proposition 5.4 induces a Borel reduction from \approx_H to \equiv_E . Recall that E_0 is the orbit equivalence relation arising from the natural free ergodic action of the direct sum D of countably many copies of the cyclic group of order 2 on the Cantor space $2^{\mathbb{N}}$. By Ornstein–Weiss [42], since D and Γ are both amenable, the essentially free ergodic actions of Γ on 2^Γ and D on $2^{\mathbb{N}}$ are orbit equivalent. Applying Lemma 5.7, it follows that \equiv_E and \equiv_{E_0} are Borel bireducible, and hence \approx_H is Borel reducible to \equiv_{E_0} .

Next let $G = \bigoplus_{n \in \mathbb{N}} G_n$, where each G_n is isomorphic to $\text{Sym}(3)$. By Theorem 4.1, since G is locally finite, \approx_G is Borel reducible to \approx_H . Hence, in order to show that \equiv_{E_0} is Borel reducible to \approx_H , it is enough to show that \equiv_{E_0} is Borel reducible to \approx_G . Write $G = A \times H$, where $A = \bigoplus_{n \in \mathbb{N}} A_n$ is the direct sum of countably many copies of the cyclic group of order 3 and $H = \bigoplus_{n \in \mathbb{N}} H_n$ is the direct sum of countably many copies of the cyclic group of order 2. Then the unitary dual $Z = \text{Irr}_1(A)$ of A is the direct product of countably many copies of the cyclic group $C_3 = \{1, \xi, \xi^2\}$ of order 3. Let

$$X = \{\xi, \xi^2\}^{\mathbb{N}} \subseteq C_3^{\mathbb{N}} = Z$$

and let μ be the product probability measure on X . Then the conjugation action of H on A induces a free ergodic action of H on (X, μ) , and the corresponding orbit equivalence relation $E = E_H^X$ is clearly orbit equivalent to E_0 . Hence, by Lemma 5.7, \equiv_E and \equiv_{E_0} are Borel bireducible. Let $\sigma \xrightarrow{\pi} \pi_\sigma$ be the Borel map from $Z^1(H \times X, U(\mathcal{H}))$ to $\text{Rep}(G, L^2(X, \mathcal{H}))$ given by (5.1) and (5.2). Then, by Proposition 5.3 and Schur’s Lemma, π induces a Borel reduction from \equiv_E to \approx_G , and hence \equiv_{E_0} is Borel reducible to \approx_H . \square

In the remainder of this section, we will discuss the relationship between \equiv_{E_0} and some more commonly studied Borel equivalence relations. We will begin by considering the measure equivalence relation on a suitably restricted subspace of the Polish space $\mathbb{P}(2^{\mathbb{N}})$ of probability measures on $2^{\mathbb{N}}$. Once again, let D be the direct sum of countably many copies of the cyclic group of order 2, equipped with its natural action on the Cantor space $2^{\mathbb{N}}$. Then we can define an associated action of D on $\mathbb{P}(2^{\mathbb{N}})$ by $\mu \mapsto g_*\mu$. Recall that a probability measure $\mu \in \mathbb{P}(2^{\mathbb{N}})$ is said to be *quasi-invariant* if $g_*\mu \sim \mu$ for all $g \in D$; and if $\mu \in \mathbb{P}(2^{\mathbb{N}})$ is quasi-invariant, then μ is said to be *ergodic* if every D -invariant Borel subset of $2^{\mathbb{N}}$ is either null or conull. It is easily checked that the set $\text{QE}(2^{\mathbb{N}})$ of quasi-invariant ergodic probability measures on $2^{\mathbb{N}}$ is a Borel subset of $\mathbb{P}(2^{\mathbb{N}})$, and hence is a standard Borel space.

Definition 6.4. \sim_{qe} is the measure equivalence relation on $\text{QE}(2^{\mathbb{N}})$.

The following result is implicitly contained in Mackey [39].

Theorem 6.5. \sim_{qe} is Borel reducible to \equiv_{E_0} .

Proof. Once again, let $G = \bigoplus_{n \in \mathbb{N}} G_n$, where each G_n is isomorphic to $\text{Sym}(3)$; and write $G = A \times H$, where $A = \bigoplus_{n \in \mathbb{N}} A_n$ is the direct sum of countably many copies of the cyclic group of order 3 and $H = \bigoplus_{n \in \mathbb{N}} H_n$ is the direct sum of countably many

copies of the cyclic group of order 2. Then it is enough to show that \sim_{qe} is Borel reducible to \approx_G . As above, let $Z = \text{Irr}_1(A)$ be the unitary dual of A and let

$$X = \{\xi, \xi^2\}^{\mathbb{N}} \subseteq C_3^{\mathbb{N}} = Z.$$

Then the induced action of H on X is isomorphic to the natural action of D on $2^{\mathbb{N}}$. If the probability measure μ on X is quasi-invariant and ergodic with respect to the action of H , then we can define a corresponding irreducible unitary representation π_μ of G on $L^2(X, \mu)$ by setting

$$(\pi_\mu(g) \cdot f)(x) = \begin{cases} \sqrt{\frac{d(g_*\mu)}{d\mu}}(x) f(g^{-1}x) & \text{if } g \in H, \\ x(g)f(x) & \text{if } g \in A, \end{cases}$$

for each $f \in L^2(X, \mu)$. Suppose that ν is another probability measure on X which is quasi-invariant and ergodic with respect to the action of H . If $\mu \sim \nu$, then the unitary isomorphism $U : L^2(X, \mu) \rightarrow L^2(X, \nu)$ defined by $f \mapsto \sqrt{d\mu/d\nu} f$ satisfies

$$U^{-1}\pi_\nu(g)U = \pi_\mu(g)$$

for each $g \in G$ and hence π_ν, π_μ are unitarily equivalent. Conversely, if π_ν, π_μ are unitarily equivalent, then $\pi_\nu \upharpoonright A$ and $\pi_\mu \upharpoonright A$ are unitarily equivalent representations of A , and it is well-known that this implies that $\mu \sim \nu$. Thus \sim_{qe} is Borel reducible to \approx_G . \square

Next, following Kechris–Sofronidis [35], we will discuss the relationship between the Borel equivalence relations \sim_{qe} and E_{ℓ^2} .

Definition 6.6. E_{ℓ^2} is the Borel equivalence relation on $\mathbb{R}^{\mathbb{N}}$ defined by

$$(a_n) E_{\ell^2} (b_n) \Leftrightarrow \sum (a_n - b_n)^2 < \infty.$$

In other words, E_{ℓ^2} is the orbit equivalence relation arising from the additive action of the Banach space $\ell^2 = \{(x_n) \in \mathbb{R}^{\mathbb{N}} \mid \sum x_n^2 < \infty\}$ on $\mathbb{R}^{\mathbb{N}}$. It is well-known that E_{ℓ^2} is Borel bireducible with $E_{\ell^2} \upharpoonright (0, 1)^{\mathbb{N}}$. (For example, see the proof of Kanovei [31, Lemma 6.2.2].) As pointed out by Kechris–Sofronidis [35], the classical theorem of Kakutani [30] on equivalence of infinite product measures implies that E_{ℓ^2} is Borel reducible to \sim_{qe} . In more detail, for each sequence $\alpha = (\alpha_n) \in (0, 1)^{\mathbb{N}}$, let μ_α be the product measure defined on $2^{\mathbb{N}}$ by

$$\mu_\alpha = \prod (\alpha_n \delta_0 + (1 - \alpha_n) \delta_1),$$

where the δ_i denote the Dirac measures on $\{0, 1\}$. Then it is well-known that each μ_α is quasi-invariant and ergodic; and, by Kakutani [30], if $\alpha, \beta \in (0, 1)^{\mathbb{N}}$, then the following are equivalent:

- (i) $\mu_\alpha \sim \mu_\beta$.
- (ii) $\sum_{n=0}^{\infty} [(\sqrt{\alpha_n} - \sqrt{\beta_n})^2 + (\sqrt{1 - \alpha_n} - \sqrt{1 - \beta_n})^2] < \infty$.

Furthermore, if there exists $\varepsilon > 0$ such that $\varepsilon \leq \alpha_n, \beta_n \leq 1 - \varepsilon$ for all $n \in \mathbb{N}$, then condition (ii) is equivalent to:

- (iii) $\sum_{n=0}^{\infty} (\alpha_n - \beta_n)^2 < \infty$.

It follows easily that E_{ℓ^2} is Borel bireducible with the restriction of \sim_{qe} to the space $\{\mu_\alpha \mid \alpha \in (0, 1)^{\mathbb{N}}\}$ of quasi-invariant ergodic product measures. In particular, it is now clear that the following result holds.

Corollary 6.7. E_{ℓ^2} is Borel reducible to \equiv_{E_0} . □

Remark 6.8. Suppose that G is a countable non-type I group. Then the results in this section imply that E_{ℓ^2} is Borel reducible to \approx_G . By Kechris–Sofronidis [35], E_{ℓ^2} does not admit classification by countable structures; and hence we obtain yet another proof of Hjorth’s theorem that \approx_G does not admit classification by countable structures. (See also Farah [14] and Kerr–Li–Pichot [36].)

Question 6.9. Is \sim_{qe} Borel bireducible with E_{ℓ^2} ?

Question 6.10. Is \equiv_{E_0} Borel bireducible with E_{ℓ^2} ?

While there is currently no reason to expect a positive answer to Question 6.10, a positive answer would enable us to separate \equiv_{E_0} and \equiv_{E_∞} . In more detail, the following result will be proved in Section 7.

Theorem 6.11. E_∞ is Borel reducible to \equiv_{E_∞} .

On the other hand, as was pointed out to me by Alexander Kechris, the following result is a straightforward consequence of Popa’s Superrigidity Theorem [43].

Theorem 6.12. E_∞ is not Borel reducible to E_{ℓ^2} .

Proof. Suppose that E_∞ is Borel reducible to E_{ℓ^2} . Let Γ be a countably infinite perfect Kazhdan group (e.g. we could take $\Gamma = \text{SL}(3, \mathbb{Z})$) and consider the action of Γ on $(2^\Gamma, \mu)$, where μ is the usual product probability measure. Then the corresponding orbit equivalence relation $E = E_\Gamma^{2^\Gamma}$ is a countable Borel equivalence relation and so E is Borel reducible to E_∞ . Hence there exists a Borel reduction $f : 2^\Gamma \rightarrow \mathbb{R}^{\mathbb{N}}$ from E to E_{ℓ^2} . Let $\alpha : \Gamma \times 2^\Gamma \rightarrow \ell^2$ be the corresponding Borel cocycle defined by

$$\alpha(\gamma, x) = \text{the unique } \bar{a} \in \ell^2 \text{ such that } f(\gamma \cdot x) = f(x) + \bar{a}.$$

By Ando–Matsuzawa [1, Example 2.18], ℓ^2 is a Polish group of finite type. Hence, from Popa [43], it follows that α is equivalent to a group homomorphism, i.e. there exists a Borel map $b : 2^\Gamma \rightarrow \ell^2$ and a group homomorphism $\varphi : \Gamma \rightarrow \ell^2$ such that for all $\gamma \in \Gamma$,

$$\begin{aligned} \varphi(\gamma) &= b(\gamma \cdot x) + \alpha(\gamma, x) - b(x) \\ &= b(\gamma \cdot x) + f(\gamma \cdot x) - (b(x) + f(x)) \quad \mu\text{-a.e. } x \in 2^\Gamma. \end{aligned}$$

Since Γ is perfect, φ is the trivial homomorphism which sends every element of Γ to the identity element $\bar{0} \in \ell^2$; and so for all $\gamma \in \Gamma$,

$$b(\gamma \cdot x) + f(\gamma \cdot x) = b(x) + f(x) \quad \mu\text{-a.e. } x \in 2^\Gamma.$$

Clearly the Borel map $f' : 2^\Gamma \rightarrow \mathbb{R}^{\mathbb{N}}$ defined by $f'(x) = b(x) + f(x)$ is also a Borel reduction from E to E_{ℓ^2} . However, we have just seen that f' is Γ -invariant μ -a.e.; and since Γ acts ergodically on $(2^\Gamma, \mu)$, this implies that f' is constant μ -a.e., which is a contradiction. □

7. Quasi-regular representations of \mathbb{F}_2

In this section, we will prove that the universal countable Borel equivalence relation E_∞ is Borel reducible to the unitary equivalence relation $\approx_{\mathbb{F}_2}$ on the space of infinite-dimensional irreducible unitary representations of the free group \mathbb{F}_2 on the two generators $\{a, b\}$. Of course, combined with Theorem 5.8, this implies that E_∞ is Borel reducible to \equiv_{E_∞} . We will begin by recalling Mackey’s theorem on quasi-regular representations of countable groups.

Let G be a countable group and let $H \leq G$ be a subgroup. Then $\lambda_{G/H}$ denotes the corresponding quasi-regular representation of G on the Hilbert space $\ell^2(G/H)$. Recall that two subgroups H_0, H_1 of a group G are said to be *commensurable* if $H_0 \cap H_1$ is of finite index in both H_0 and H_1 . The *commensurator* of H_0 in G is defined to be

$$\text{Com}_G(H_0) = \{g \in G \mid H_0 \text{ and } gH_0g^{-1} \text{ are commensurable}\}.$$

The subgroup H_0 is said to be *self-commensurating* if $\text{Com}_G(H_0) = H_0$. It is easily seen that if H_0 and H_1 are self-commensurating subgroups of G , then H_0 and H_1 are conjugate in G if and only if there exists $g \in G$ such that gH_0g^{-1} and H_1 are commensurable. Thus the following result is an immediate consequence of Corollaries 1 and 2 in Mackey [39, Section 3.4]. (The formulation of Theorem 7.1 is a slight variant of Theorem 2.1 in Burger and de la Harpe [7].)

Theorem 7.1 (Mackey). *Let G be a countable group and let H_0, H_1 be subgroups of G .*

- (a) λ_{G/H_0} is irreducible if and only if H_0 is a self-commensurating subgroup.
- (b) If H_0 and H_1 are self-commensurating subgroups, then λ_{G/H_0} is unitarily equivalent to λ_{G/H_1} if and only if H_0 and H_1 are conjugate in G .

Remark 7.2. Suppose that G is a countable group and that H is a self-commensurating subgroup of G . Extending the results of Mackey [39, Section 3.4] on the induction of the 1-dimensional representations of H , Corwin [8] and Kleppner [37] proved that if π, σ are finite-dimensional irreducible representations of H , then:

- (i) $\text{ind}_H^G(\pi)$ is an irreducible representation of G ; and
- (ii) $\pi \approx_H \sigma$ if and only if $\text{ind}_H^G(\pi) \approx_G \text{ind}_H^G(\sigma)$.

For many years, it was unknown whether or not the analogous statements were also true for infinite-dimensional irreducible representations $\pi, \sigma \in \text{Irr}_\infty(H)$. However, Bekka–Curtis [4] have shown that there exists a self-commensurating subgroup H of $G = \mathbb{Z}^2 \rtimes \text{SL}(2, \mathbb{Z})$ and an infinite-dimensional irreducible representation $\pi \in \text{Irr}_\infty(H)$ such that the induced representation $\text{ind}_H^G(\pi)$ is not irreducible. It would be very interesting to find sufficient conditions on pairs $H \leq G$ of countable groups which ensure that the map $\pi \mapsto \text{ind}_H^G(\pi)$ is a Borel reduction from \approx_H to \approx_G .

Definition 7.3. If G is a countable group, then

$$\mathcal{C}(G) = \{H \leq G \mid H \text{ is a self-commensurating subgroup}\}$$

and $E_{\mathcal{C}(G)}$ denotes the conjugacy relation on $\mathcal{C}(G)$.

The following result is an immediate consequence of Theorem 7.1.

Lemma 7.4. $E_{\mathcal{C}(\mathbb{F}_2)}$ is Borel reducible to $\approx_{\mathbb{F}_2}$.

Thus, in order to show that E_∞ is Borel reducible to $\approx_{\mathbb{F}_2}$, it is enough to prove the following result.

Theorem 7.5. $E_{\mathcal{C}(\mathbb{F}_2)}$ is a universal countable Borel equivalence relation.

Let \mathbb{F}_3 be the free group on the three generators $\{a, b, c\}$. Then the following easy observation shows that it is enough to prove that $E_{\mathcal{C}(\mathbb{F}_3)}$ is a universal countable Borel equivalence relation.

Lemma 7.6. $E_{\mathcal{C}(\mathbb{F}_3)}$ is Borel reducible to $E_{\mathcal{C}(\mathbb{F}_2)}$.

Proof. By Karrass–Solitar [32, p. 950], the set $\{aba, a^2b^2a^2, a^3b^3a^3\}$ freely generates a malnormal subgroup K of \mathbb{F}_2 . (Recall that a subgroup K of a group G is said to be malnormal if $gKg^{-1} \cap K = 1$ for all $g \in G \setminus K$.) In particular, each $H \in \mathcal{C}(K)$ is also self-commensurating in \mathbb{F}_2 . Furthermore, if $H_0, H_1 \in \mathcal{C}(K)$ and $g \in \mathbb{F}_2$ satisfies $gH_0g^{-1} = H_1$, then $g \in K$. Thus the identity map is a Borel reduction from $E_{\mathcal{C}(K)}$ to $E_{\mathcal{C}(\mathbb{F}_2)}$. \square

From now on, let $\mathcal{P}_\infty(\mathbb{F}_2)$ be the standard Borel space of infinite subsets of \mathbb{F}_2 . Then it is easily seen that $(E_\infty \upharpoonright \mathcal{P}_\infty(\mathbb{F}_2)) \sim_B E_\infty$ and thus $E_\infty \upharpoonright \mathcal{P}_\infty(\mathbb{F}_2)$ is also a universal countable Borel equivalence relation. For each $A \in \mathcal{P}_\infty(\mathbb{F}_2)$, let $K(A)$ and $S(A)$ be the subgroups of \mathbb{F}_3 defined by

$$K(A) = \langle wCw^{-1} \mid w \in A \rangle, \quad S(A) = \{g \in \mathbb{F}_2 \mid gA = A\}.$$

Let $C = \langle c \rangle$. Then it is easily checked that

$$K(A) = \ast_{w \in A} wCw^{-1}$$

is the free product of the subgroups $\{wCw^{-1} \mid w \in A\}$. In particular, $K(A) \cap \mathbb{F}_2 = 1$. It is also clear that if $g \in S(A)$, then $gK(A)g^{-1} = K(A)$. Hence the subgroup $H(A) \leq \mathbb{F}_3$ generated by $K(A) \cup S(A)$ decomposes into a semidirect product:

$$H(A) = K(A) \rtimes S(A).$$

Proposition 7.7. The map $A \mapsto H(A)$ is a Borel reduction from $E_\infty \upharpoonright \mathcal{P}_\infty(\mathbb{F}_2)$ to $E_{\mathcal{C}(\mathbb{F}_3)}$.

The proof of Proposition 7.7 will be broken down into a sequence of lemmas. We will begin with the following easy observation.

Lemma 7.8. If $A \in \mathcal{P}_\infty(\mathbb{F}_2)$ and $g \in \mathbb{F}_2$, then $gH(A)g^{-1} = H(gA)$.

Proof. Clearly $gK(A)g^{-1} = K(gA)$ and $gS(A)g^{-1} = S(gA)$. \square

Most of our efforts will go into proving the following somewhat technical result.

Lemma 7.9. Suppose that $A, B \in \mathcal{P}_\infty(\mathbb{F}_2)$ and $\gamma \in \mathbb{F}_3$. If $H(A)$ and $\gamma H(B)\gamma^{-1}$ are commensurable, then there exist $g \in \mathbb{F}_2$ and $u \in K(B)$ such that $\gamma = gu$.

Before presenting the proof of Lemma 7.9, we will show how to complete the proof of Proposition 7.7.

Lemma 7.10. *If $A \in \mathcal{P}_\infty(\mathbb{F}_2)$, then $H(A) \in \mathcal{C}(\mathbb{F}_3)$.*

Proof. Suppose that $\gamma \in \text{Com}_{\mathbb{F}_3}(H(A))$. Then, by Lemma 7.9, there exist $g \in \mathbb{F}_2$ and $u \in K(A)$ such that $\gamma = gu$. Thus it is enough to show that $g \in S(A)$. To see this, first note that

$$\gamma H(A)\gamma^{-1} = gH(A)g^{-1} = H(gA).$$

Thus $H(A)$ and $H(gA)$ are commensurable; and this implies that $H(A)$ and $H(g^{-1}A)$ are also commensurable. Now suppose that $gA \neq A$. Then, after replacing g by g^{-1} if necessary, we can suppose that there exists $w_0 \in gA \setminus A$. Since $H(A)$ and $H(gA)$ are commensurable, there exists $n \geq 1$ such that

$$w_0c^nw_0^{-1} \in H(A) = K(A) \rtimes S(A).$$

Thus there exist $w_1, \dots, w_\ell \in A$, $m_1, \dots, m_\ell \in \mathbb{Z} \setminus \{0\}$ and $h \in S(A)$ such that

$$w_0c^nw_0^{-1} = w_1c^{m_1}w_1^{-1} \cdots w_\ellc^{m_\ell}w_\ell^{-1}h. \tag{7.1}$$

By considering the homomorphism $\mathbb{F}_3 \rightarrow \mathbb{F}_2$ such that $a \mapsto a$, $b \mapsto b$ and $c \mapsto 1$, we see that $h = 1$. But then (7.1) contradicts the fact that $K(\mathbb{F}_2)$ is the free product of the subgroups $\{wCw^{-1} \mid w \in \mathbb{F}_2\}$. \square

Lemma 7.11. *Suppose that $A, B \in \mathcal{P}_\infty(\mathbb{F}_2)$. If $H(A)$ and $H(B)$ are conjugate in \mathbb{F}_3 , then there exists $g \in \mathbb{F}_2$ such that $A = gB$.*

Proof. Suppose that $\gamma \in \mathbb{F}_3$ and that $H(A) = \gamma H(B)\gamma^{-1}$. Then, by Lemma 7.9, there exist $g \in \mathbb{F}_2$ and $u \in K(B)$ such that $\gamma = gu$. Thus $H(A) = H(gB)$; and arguing as in the proof of Lemma 7.10, we see that $A = gB$. \square

Proof of Proposition 7.7. By Lemma 7.10, if $A \in \mathcal{P}_\infty(\mathbb{F}_2)$, then $H(A) \in \mathcal{C}(\mathbb{F}_3)$. Suppose that $A, B \in \mathcal{P}_\infty(\mathbb{F}_2)$. By Lemma 7.8, if there exists $g \in \mathbb{F}_2$ such that $gA = B$, then $gH(A)g^{-1} = H(B)$. Conversely, by Lemma 7.11, if $H(A)$ and $H(B)$ are conjugate in \mathbb{F}_3 , then there exists $g \in \mathbb{F}_2$ such that $A = gB$. Thus the map $A \mapsto H(A)$ is a Borel reduction from $E_\infty \upharpoonright \mathcal{P}_\infty(\mathbb{F}_2)$ to $E_{\mathcal{C}(\mathbb{F}_3)}$. \square

The following argument is closely based upon the proof of Gao [20, Lemma 2].

Proof of Lemma 7.9. Suppose that $A, B \in \mathcal{P}_\infty(\mathbb{F}_2)$ and that $\gamma \in \mathbb{F}_3$ is such that $H(A)$ and $\gamma H(B)\gamma^{-1}$ are commensurable. Since

$$[K(A) : K(A) \cap \gamma H(B)\gamma^{-1}] < \infty,$$

it follows that for each $x \in A$, there exists $n_x \geq 1$ such that

$$xc^{n_x}x^{-1} = \gamma w_x g \gamma^{-1} \tag{7.2}$$

for some $w_x \in K(B)$ and $g \in S(B)$. Notice that $xc^{n_x}x^{-1}$ is a freely reduced word in a, b, c . In considering the right-hand side of (7.2), we will initially suppose that:

- (i) γ, γ^{-1} and g are freely reduced words in a, b, c ; and
- (ii) $w_x \in K(B)$ is written as a reduced word with respect to the decomposition of $K(B)$ as the free product of the subgroups $\{bcb^{-1} \mid b \in B\}$; say,

$$w_x = y_1 \cdots y_t,$$

where each $y_i = b_i c^{\ell_i} b_i^{-1}$ for some $b_i \in B$ and $\ell_i \in \mathbb{Z} \setminus \{0\}$.

Then after freely reducing the initial expression of $\gamma w_x g \gamma^{-1}$ through successive cancellations of terms of the form zz^{-1} or $z^{-1}z$ for some $z \in \{a, b, c\}$, we must eventually obtain the reduced word $x c^{n_x} x^{-1}$. From now on, for each $x \in A$, we will fix such a cancellation procedure. Note that for each $x \in A$, there exists an occurrence of c in the initial expression of $\gamma w_x g \gamma^{-1}$ which is preserved throughout the cancellation procedure and gives rise to the first occurrence of c in the reduced word $x c^{n_x} x^{-1}$. This occurrence of c in $\gamma w_x g \gamma^{-1}$ will be called the *first preserved occurrence*.

We claim that there exists at most one $x \in A$ such that the corresponding first preserved occurrence of c is a letter in γ . To see this, suppose that $x_1, x_2 \in A$ both have this property. Since γ is freely reduced, the first preserved occurrence must be the first c in γ . Thus writing $\gamma = kcu$, where $k \in \mathbb{F}_2$ and $u \in \mathbb{F}_3$, we see that

$$x_1 c^{n_{x_1}} x_1^{-1} = kcuw_{x_1} g \gamma^{-1}, \quad x_2 c^{n_{x_2}} x_2^{-1} = kcuw_{x_2} g \gamma^{-1},$$

which implies that $x_1 = k = x_2$. Similarly, we claim there exists at most one $x \in A$ such that the corresponding first preserved occurrence of c is a letter in γ^{-1} . To see this, note that if the first preserved occurrence of c is a letter in γ^{-1} , then the last preserved occurrence of c must also be a letter in γ^{-1} ; and since γ^{-1} is freely reduced, this last preserved occurrence must be the last c in γ^{-1} . Arguing as above yields the claim easily.

Thus there exists $x \in A$ such that the corresponding first preserved occurrence of c is a letter in $w_x g$. Of course, since $g \in S(B) \leq \mathbb{F}_2$, this means that the corresponding first preserved occurrence of c is actually a letter in w_x ; say, the first preserved occurrence is one of the letters in $y_i = b_i c^{\ell_i} b_i^{-1}$. Then

$$x = \gamma b_1 c^{\ell_1} b_1^{-1} \cdots b_{i-1} c^{\ell_{i-1}} b_{i-1}^{-1} b_i c^r$$

for some $r \geq 0$. Let $u = b_1 c^{\ell_1} b_1^{-1} \cdots b_{i-1} c^{\ell_{i-1}} b_{i-1}^{-1}$. Then $u \in K(B)$. If $r = 0$, then $\gamma = x b_i^{-1} u^{-1}$ has the required form, since $x b_i^{-1} \in \mathbb{F}_2$ and $u^{-1} \in K(B)$. So suppose that $r > 0$. Then we can write $x = \gamma z b_i$, where $z = u b_i c^r b_i^{-1} \in K(B)$; and, once again, γ has the required form. □

8. Unitary equivalence of arbitrary representations

In the previous sections, we have focused our attention on the unitary equivalence relation \approx_G on the space of irreducible unitary representations of the countable group G on the infinite-dimensional separable Hilbert space \mathcal{H} . In this final section, we will point out a basic open question concerning the unitary equivalence relation \approx_G^+ on the space

$\text{Rep}(G, \mathcal{H})$ of *arbitrary* unitary representations of G on \mathcal{H} . By a recent result of Hjorth–Törnquist [27], \approx_G^+ is a Borel equivalence relation; in fact, \approx_G^+ is an $F_{\sigma\delta}$ equivalence relation.

Let \sim be the measure equivalence relation on the space $\text{P}(2^{\mathbb{N}})$ of *arbitrary* probability measures on $2^{\mathbb{N}}$. (Of course, we could replace $2^{\mathbb{N}}$ with any uncountable standard Borel space X .) Then, as pointed out by Kechris [34, Section 5], the spectral theorem for unitary operators implies that the unitary equivalence relation $\approx_{\mathbb{Z}}^+$ on the space $\text{Rep}(\mathbb{Z}, \mathcal{H})$ is Borel bireducible with \sim . In particular, $\approx_{\mathbb{Z}}^+$ is not smooth and hence $\approx_{\mathbb{Z}} <_B \approx_{\mathbb{Z}}^+$. More generally, the following result is a straightforward consequence of Kanovei’s theory [31, Chapter 17] of pinned Borel equivalence relations.

Theorem 8.1 (Hjorth–Törnquist [27]). *If G is an arbitrary countably infinite group, then $\approx_G <_B \approx_G^+$.*

However, the following basic problem remains open.

Question 8.2. Is $\approx_{\mathbb{F}_\infty}^+$ Borel bireducible with $\approx_{\mathbb{Z}}^+$? Equivalently, is $\approx_{\mathbb{F}_\infty}^+$ Borel bireducible with the measure equivalence relation \sim ?

Here it is probably worth pointing out that E_∞ is Borel reducible to \sim and hence E_∞ is Borel reducible to $\approx_{\mathbb{Z}}^+$. In fact, I am currently not aware of any even vaguely plausible strategy for separating $\approx_{\mathbb{Z}}^+$ and $\approx_{\mathbb{F}_\infty}^+$.

Appendix. List of equivalence relations

In this appendix, we collect together the definitions of the various Borel equivalence relations that occur in this paper.

- \approx_G is the unitary equivalence relation on the space $\text{Irr}_\infty(G)$ of infinite-dimensional irreducible unitary representations of the countable non-type I group G .
- \approx_G^+ is the unitary equivalence relation on the space $\text{Rep}(G, \mathcal{H})$ of arbitrary unitary representations of the countable group G on the infinite-dimensional separable Hilbert space \mathcal{H} .
- $\approx_{\mathcal{A}}$ is the unitary equivalence relation on the space $\text{Irr}_\infty(\mathcal{A})$ of infinite-dimensional irreducible unitary representations of the non-type I separable C^* -algebra \mathcal{A} .
- E_G^X is the orbit equivalence relation of the countable group G acting on the standard Borel G -space X .
- E_0 is the eventual equality relation on $2^{\mathbb{N}}$.
- E_∞ is the orbit equivalence relation of the shift action of \mathbb{F}_2 on $2^{\mathbb{F}_2}$.
- \equiv_E is the orbit equivalence relation of the action of $L(X, U(\mathcal{H}))$ on the space $\text{Irr}(E, U(\mathcal{H}))$ of irreducible cocycles.
- \sim is the measure equivalence relation on the space $\text{P}(X)$ of probability measures on the uncountable Polish space X .
- \sim_{qe} is the measure equivalence relation on the space $\text{QE}(2^{\mathbb{N}})$ of quasi-invariant ergodic probability measures on $2^{\mathbb{N}}$.

- E_{ℓ^2} is the orbit equivalence relation arising from the additive action of the Banach space $\ell^2 = \{(x_n) \in \mathbb{R}^{\mathbb{N}} \mid \sum x_n^2 < \infty\}$ on $\mathbb{R}^{\mathbb{N}}$.
- $E_{\mathcal{C}(G)}$ is the conjugacy relation on the space $\mathcal{C}(G)$ of self-commensurating subgroups of the countable group G .

Acknowledgments. I would like to thank Alexander Kechris for some very helpful discussions concerning the material in this paper and for explaining that Popa's Superrigidity Theorem [43] holds for cocycles taking values in $\ell^2 = \{(x_n) \in \mathbb{R}^{\mathbb{N}} \mid \sum x_n^2 < \infty\}$. This research was partially supported by NSF Grant DMS 1101597.

References

- [1] Ando, H., Matsuzawa, Y.: On Polish groups of finite type. *Publ. RIMS Kyoto Univ.* **48**, 389–408 (2012) [Zbl 1255.46028](#) [MR 2928146](#)
- [2] Arveson, W.: *An Invitation to C^* -algebras*. *Grad. Texts in Math.* 39, Springer, New York (1976) [Zbl 0344.46123](#) [MR 0512360](#)
- [3] Baer, R.: Abelian groups without elements of finite order. *Duke Math. J.* **3**, 68–122 (1937) [Zbl 0016.20303](#) [MR 1545974](#)
- [4] Bekka, M., Curtis, R.: On Mackey's irreducibility criterion for induced representations. *Int. Math. Res. Notices* **2003**, no. 38, 2095–2101 [Zbl 1027.22010](#) [MR 1994777](#)
- [5] Bowen, L.: Stable orbit equivalence of Bernoulli shifts over free groups. *Groups Geom. Dynam.* **5**, 17–38 (2011) [Zbl 1259.37002](#) [MR 2763778](#)
- [6] Brown, N.P., Ozawa, N.: *C^* -Algebras and Finite-Dimensional Approximations*. *Grad. Stud. Math.* 88, Amer. Math. Soc., Providence, RI (2008) [Zbl 1160.46001](#) [MR 2391387](#)
- [7] Burger, M., de la Harpe, P.: Constructing irreducible representations of discrete groups. *Proc. Indian Acad. Sci. Math. Sci.* **107**, 223–235 (1997) [Zbl 0908.22005](#) [MR 1467427](#)
- [8] Corwin, L.: Induced representations of discrete groups. *Proc. Amer. Math. Soc.* **47**, 279–287 (1975) [Zbl 0271.20003](#) [MR 0387487](#)
- [9] Dixmier, J.: *C^* -algebras*. North-Holland, Amsterdam (1977) [Zbl 0372.46058](#) [MR 0458185](#)
- [10] Dougherty, R., Jackson S., Kechris, A. S.: The structure of hyperfinite Borel equivalence relations. *Trans. Amer. Math. Soc.* **341**, 193–225 (1994) [Zbl 0803.28009](#) [MR 1149121](#)
- [11] Elliott, G. A.: On the classification of inductive limits of sequences of semisimple finite-dimensional algebras. *J. Algebra* **38**, 29–44 (1976) [Zbl 0323.46063](#) [MR 0397420](#)
- [12] Elliott, G. A.: The Mackey Borel structure on the spectrum of an approximately finite-dimensional separable C^* -algebra. *Trans. Amer. Math. Soc.* **233**, 59–68 (1977) [Zbl 0369.46055](#) [MR 0454648](#)
- [13] Epstein, I.: Orbit inequivalent actions of non-amenable groups. [arXiv:0707.4215](#) (2007)
- [14] Farah, I.: A dichotomy for the Mackey Borel structure. In: *Proc. 11th Asian Logic Conference, World Sci.*, 86–93 (2011) [Zbl 1284.03228](#) [MR 2868507](#)
- [15] Farah, I., Toms, A. S., Törnquist, A.: The descriptive set theory of C^* -algebra invariants. *Int. Math. Res. Notices* **2013**, 5196–5226 [Zbl 06438759](#) [MR 3129097](#)
- [16] Feldman, J., Moore, C. C.: Ergodic equivalence relations, cohomology and von Neumann algebras I. *Trans. Amer. Math. Soc.* **234**, 289–324 (1977) [Zbl 0369.22009](#) [MR 0578656](#)
- [17] Folland, G. B.: *A Course in Abstract Harmonic Analysis*. CRC Press, Boca Raton, FL (1995) [Zbl 0857.43001](#) [MR 1397028](#)
- [18] Friedman, S.-D., Motto Ros, L.: Analytic equivalence relations and bi-embeddability. *J. Symbolic Logic* **76**, 243–266 (2011) [Zbl 1256.03050](#) [MR 2791347](#)

- [19] Gaboriau, D., Lyons, R.: A measurable-group-theoretic solution to von Neumann's problem. *Invent. Math.* **117**, 533–540 (2009) [Zbl 1182.43002](#) [MR 2534099](#)
- [20] Gao, S.: Coding subset shifts by subgroup conjugacy. *Bull. London Math. Soc.* **32**, 653–657 (2000) [Zbl 1025.03044](#) [MR 1781575](#)
- [21] Gao, S.: *Invariant Descriptive Set Theory*. CRC Press, Boca Raton, FL (2009) [Zbl 1154.03025](#) [MR 2455198](#)
- [22] Glimm, J.: Type I C^* -algebras. *Ann. of Math.* **73**, 572–612 (1961) [Zbl 0152.33002](#) [MR 0124756](#)
- [23] Harrington, L., Kechris, A. S., Louveau, A.: A Glimm–Effros dichotomy for Borel equivalence relations. *J. Amer. Math. Soc.* **3**, 903–927 (1990) [Zbl 0778.28011](#) [MR 1057041](#)
- [24] Hewitt, E., Stromberg, K.: *Real and Abstract Analysis*. Springer, New York (1965) [Zbl 0137.03202](#) [MR 0188387](#)
- [25] Hjorth, G.: Non-smooth infinite dimensional group representations. Preprint (1997)
- [26] Hjorth, G.: *Classification and Orbit Equivalence Relations*. Math. Surveys Monogr. 75, Amer. Math. Soc., Providence, RI (2000) [Zbl 0942.03056](#) [MR 1725642](#)
- [27] Hjorth, G., Törnquist, A.: The conjugacy relation on unitary representations. *Math. Res. Lett.* **19**, 525–535 (2012) [Zbl 1283.03079](#) [MR 2998137](#)
- [28] Huppert, B.: *Character Theory of Finite Groups*. de Gruyter Expositions Math. 25, de Gruyter, Berlin (1998) [Zbl 0932.20007](#) [MR 1645304](#)
- [29] Jackson, S., Kechris, A.S., Louveau, A.: Countable Borel equivalence relations. *J. Math. Logic* **2**, 1–80 (2002) [Zbl 1008.03031](#) [MR 1900547](#)
- [30] Kakutani, S.: On equivalence of infinite product measures. *Ann. of Math.* **49**, 214–224 (1948) [Zbl 0030.02303](#) [MR 0023331](#)
- [31] Kanovei, V.: *Borel Equivalence Relations: Structure and Classification*. Univ. Lecture Ser. 44, Amer. Math. Soc., Providence (2008) [Zbl 1155.03035](#) [MR 2441635](#)
- [32] Karrass, A., Solitar, D.: The free product of two groups with a malnormal amalgamated subgroup. *Canad. J. Math.* **23**, 933–959 (1971) [Zbl 0247.20028](#) [MR 0314992](#)
- [33] Kechris, A. S.: *Classical Descriptive Set Theory*. Grad. Texts in Math. 156, Springer (1995) [Zbl 0819.04002](#) [MR 1321597](#)
- [34] Kechris, A. S.: *Global Aspects of Ergodic Group Actions*. Math. Surveys Monogr. 160, Amer. Math. Soc., Providence, RI (2010) [Zbl 1189.37001](#) [MR 2583950](#)
- [35] Kechris, A. S., Sofronidis, N. E.: A strong generic ergodicity property of unitary and self-adjoint operators. *Ergodic Theory Dynam. Systems* **21**, 1459–1479 (2001) [Zbl 1062.47514](#) [MR 1855842](#)
- [36] Kerr, D., Li, H., Pichot, M.: Turbulence, representations, and trace-preserving actions. *Proc. London Math. Soc.* (3) **100**, 459–484 (2010) [Zbl 1192.46063](#) [MR 2595746](#)
- [37] Kleppner, A.: On the intertwining number theorem. *Proc. Amer. Math. Soc.* **12**, 731–733 (1961) [Zbl 0119.11101](#) [MR 0131783](#)
- [38] Lazar, A. J., Taylor, D. C.: Approximately finite-dimensional C^* -algebras and Bratteli diagrams. *Trans. Amer. Math. Soc.* **259**, 599–619 (1980) [Zbl 0437.46046](#) [MR 0567100](#)
- [39] Mackey, G. W.: *The Theory of Unitary Group Representations*. Univ. of Chicago Press (1976) [Zbl 0344.22002](#) [MR 0396826](#)
- [40] Motto Ros, L.: On the complexity of the relations of isomorphism and bi-embeddability. *Proc. Amer. Math. Soc.* **140**, 309–323 (2012) [Zbl 1245.03068](#) [MR 2833542](#)
- [41] Nielsen, O. A.: *Direct Integral Theory*. Lecture Notes in Pure Appl. Math. 61, Dekker, New York (1980) [Zbl 0482.46037](#) [MR 0591683](#)
- [42] Ornstein, D. S., Weiss, B.: Ergodic theory of amenable group actions I: The Rohlin lemma. *Bull. Amer. Math. Soc.* **2**, 161–164 (1980) [Zbl 0427.28018](#) [MR 0551753](#)

-
- [43] Popa, S.: Cocycle and orbit equivalence superrigidity for malleable actions of w-rigid groups. *Invent. Math.* **170**, 243–295 (2007) [Zbl 1131.46040](#) [MR 2342637](#)
 - [44] Quigg, J. C.: On the irreducibility of an induced representation. *Pacific J. Math.* **93**, 163–179 (1981) [Zbl 0441.22005](#) [MR 0621605](#)
 - [45] Serre, J.-P.: *Trees*. Springer (1980) [Zbl 0548.20018](#) [MR 0607504](#)
 - [46] Sutherland, C. E.: On a construction of unitary cocycles and the representation theory of amenable groups. *Ergodic Theory Dynam. Systems* **3**, 129–135 (1983) [Zbl 0522.22006](#) [MR 0743031](#)
 - [47] Thoma, E.: Über unitäre Darstellungen abzählbarer, diskreter Gruppen. *Math. Ann.* **153**, 111–138 (1964) [Zbl 0136.11603](#) [MR 0160118](#)