



Inkang Kim · Pierre Pansu

Flexibility of surface groups in classical simple Lie groups

Dedicated to Lionel Bérard-Bergery for his 65th birthday

Received October 6, 2011

Abstract. We show that a surface group of high genus contained in a classical simple Lie group can be deformed to become Zariski dense, unless the Lie group is $SU(p, q)$ (resp. $SO^*(2n)$, n odd) and the surface group is maximal in some $S(U(p, p) \times U(q - p)) \subset SU(p, q)$ (resp. $SO^*(2n - 2) \times SO(2) \subset SO^*(2n)$). This is a converse, for classical groups, to a rigidity result of S. Bradlow, O. García-Prada and P. Gothen.

Keywords. Algebraic group, symmetric space, rigidity, group cohomology, moduli space

1. Introduction

Free groups are obviously flexible. In particular, a generic free subgroup in a real algebraic group is Zariski dense. This already fails for surface groups, although they are very flexible from other points of view. The first evidence came from the following result by D. Toledo.

Theorem 1.1. (D. Toledo, 1979, 1989, [18]). *Let Γ be a discrete cocompact subgroup of $SU(1, 1)$. Map $SU(1, 1)$ as a 2×2 block in $SU(1, n)$, $n \geq 2$. Then every neighboring homomorphism $\Gamma \rightarrow SU(1, n)$ is contained in a conjugate of $S(U(1, 1) \times U(n - 1))$.*

In fact, Toledo obtained a stronger, global result: a characterization of surface subgroups of $S(U(1, 1) \times U(n - 1))$ among surface subgroups of $SU(1, n)$ by the value of a characteristic class known as Toledo's invariant, which we now define.

Let X be a Hermitian symmetric space with Kähler form Ω (the metric is normalized so that the minimal sectional curvature equals -1). Let Σ be a closed surface of negative Euler characteristic, and let $\Gamma = \pi_1(\Sigma)$ act isometrically on X . Pick a smooth equivariant map $\tilde{f} : \tilde{\Sigma} \rightarrow X$.

Definition 1.2. Define the *Toledo invariant* of the action $\rho : \Gamma \rightarrow \text{Isom}(X)$ by

$$T_\rho = \frac{1}{2\pi} \int_\Sigma \tilde{f}^* \Omega.$$

I. Kim: School of Mathematics, KIAS, Heogiro 85, Dongdaemen-gu, Seoul, 130-722, Korea; e-mail: inkang@kias.re.kr

P. Pansu: Laboratoire de Mathématiques d'Orsay, UMR 8628 du CNRS, Université Paris-Sud, 91405 Orsay Cedex, France; e-mail: pierre.pansu@math.u-psud.fr

Mathematics Subject Classification (2010): Primary 51M10; Secondary 57S25

Then

- (1) T_ρ depends continuously on ρ .
- (2) There exists $\ell_X \in \mathbb{Q}$ such that $T_\rho \in \ell_X \mathbb{Z}$.
- (3) $|T_\rho| \leq |\chi(\Sigma)| \text{rank}(X)$.

Inequality (3), known as the Milnor–Wood inequality for actions on Hermitian symmetric spaces, is due to J. Milnor [15], V. Turaev [19], A. Domic and D. Toledo [6], J.-L. Clerc and B. Ørsted [5].

Definition 1.3. Actions ρ such that $|T_\rho| = |\chi(\Sigma)| \text{rank}(X)$ are called *maximal representations*.

The following result generalizes Theorem 1.1.

Theorem 1.4 (L. Hernández Lamóneda [12], S. Bradlow, O. García-Prada, P. Gothen [1], [2]). *Maximal reductive representations of Γ to $SU(p, q)$, $p \leq q$, can be conjugated into $S(U(p, p) \times U(q - p))$. Maximal reductive representations of Γ to $SO^*(2n)$, n odd, can be conjugated into $SO^*(2n - 2) \times SO(2)$.*

In turn, Theorem 1.4 is a special case of a more general result.

Definition 1.5. Say a Hermitian symmetric space is of *tube type* if it can be realized as a domain in \mathbb{C}^n of the form $\mathbb{R}^n + iC$ where $C \subset \mathbb{R}^n$ is a proper open cone.

Example 1.6. Siegel’s upper half-spaces and Grassmannians with isometry groups $PO(2, q)$ are of tube type.

The Grassmannian $\mathcal{D}_{p,q}$, $p \leq q$, with isometry group $PU(p, q)$ is of tube type iff $p = q$.

The Grassmannian \mathcal{G}_n with isometry group $SO^*(2n)$ is of tube type iff n is even.

The exceptional Hermitian symmetric space of dimension 27 is of tube type, the other one (of dimension 16) is not.

Products of tube type spaces are of tube type, so polydisks are of tube type.

Remark 1.7. All maximal tube type subsymmetric spaces in a Hermitian symmetric space are conjugate. For instance, the maximal tube type subsymmetric space in $\mathcal{D}_{p,q}$ is $\mathcal{D}_{p,p}$. The maximal tube type subsymmetric space in \mathcal{G}_{2n+1} is \mathcal{G}_{2n} .

Theorem 1.8. (Burger, Iozzi, Wienhard [3]). *Let Γ be a closed surface group and X a Hermitian symmetric space. Every maximal representation $\Gamma \rightarrow \text{Isom}(X)$ stabilizes a maximal tube type subsymmetric space Y . Conversely, for every tube type Hermitian symmetric space X , $\text{Isom}(X)$ admits Zariski dense maximal surface subgroups.*

1.1. Results

Our main result is a converse of Theorem 1.4 (i.e. Theorem 1.8 for classical simple Lie groups).

Theorem 1. *Let G be a classical real Lie group, i.e. a real form of $SL(n, \mathbb{C})$, $O(n, \mathbb{C})$ or $Sp(n, \mathbb{C})$. Let Γ be the fundamental group of a closed surface of genus $\geq 2\dim(G)^2$. A homomorphism $\phi : \Gamma \rightarrow G$ can be approximated by Zariski dense representations, unless the symmetric space X of G is Hermitian and not of tube type, and ϕ is maximal.*

In other words, the exceptions are $G = SU(p, q)$, $q > p$, and $\phi(\Gamma)$ is contained in a conjugate of $S(U(p, p) \times U(q - p)) \subset SU(p, q)$, or $G = SO^*(2n)$, n odd, and $\phi(\Gamma)$ is contained in a conjugate of $SO^*(2n - 2) \times SO(2) \subset SO^*(2n)$.

The genus assumption is probably unnecessary, but we have been unable to remove it.

Question 1.9. Does Theorem 1 extend to exceptional simple Lie groups?

Since flexibility easily holds in compact and complex groups (see Proposition 4.3), and the rank one example F_4^{-20} is treated in [13], there remain ten cases ($E_6^6, E_6^{-14}, E_6^{-26}, E_7^7, E_7^{-5}, E_7^{-25}, E_8^8, E_8^{-24}, F_4^4, G_2^2$) with at least one rigidity case (E_6^{-14}). See [14], [16], [11] for related results.

1.2. Scheme of proof

The proof relies on

- a necessary and sufficient condition for flexibility from [13] (see Theorem 2.2 below);
- tools from Burger, Iozzi and Wienhard's theory of tight maps between Hermitian symmetric spaces, [4];
- a detailed analysis of centers of centralizers of reductive subgroups of classical simple Lie groups.

This last analysis is based on bilinear and sesquilinear algebra. This is where exceptional simple Lie groups elude us.

1.3. Plan of the paper

Section 2 recalls a result from [13]. Section 3 proves relevant consequences of the theory of tight maps. Section 4 provides a description of classical real simple Lie groups as fixed points of involutions, which helps in computing root space decompositions in the complexified Lie algebra. This is done in Section 5 for $\mathfrak{sl}(n, \mathbb{C})$ and in Section 6 for $\mathfrak{so}(n, \mathbb{C})$ and $\mathfrak{sp}(n, \mathbb{C})$. The method consists in first computing the root space decomposition in the standard representation of the complexified Lie algebra, and deducing the decomposition in the adjoint representation. The theory of tight homomorphisms allows us to exclude balancedness (with exceptions), first for real forms of $SL(n, \mathbb{C})$ in Section 7, then for real forms of $SO(n, \mathbb{C})$ and $Sp(n, \mathbb{C})$ in Section 8. Theorem 1 is proven in Section 9.

2. Flexibility criterion

As far as the flexibility of a homomorphism ϕ is concerned, a key role is played by the center of the centralizer of the image of ϕ . It splits the complexified Lie algebra of G into

root spaces \mathfrak{g}_λ . When the root λ is purely imaginary, \mathfrak{g}_λ carries a natural nondegenerate sesquilinear form defined as follows. Let $(X, X') \mapsto X \cdot X'$ denote the Killing form on $\mathfrak{g} \otimes \mathbb{C}$. Then the sesquilinear form

$$s_\lambda(X, X') = \bar{X} \cdot X'$$

is nondegenerate on \mathfrak{g}_λ . Let Ω_λ denote the imaginary part of s_λ . It is a symplectic form on \mathfrak{g}_λ viewed as a real vector space. The representation of Γ on \mathfrak{g}_λ gives rise to a homomorphism $\Gamma \rightarrow Sp(\mathfrak{g}_\lambda, \Omega_\lambda)$, an isometric action on the Siegel domain, and thus a Toledo invariant T_λ .

Definition 2.1. Let $\mathfrak{t} \subset \mathfrak{g}$ be a torus, centralized by a homomorphism $\phi : \Gamma \rightarrow G$. Among the roots of the adjoint action of \mathfrak{t} on $\mathfrak{g} \otimes \mathbb{C}$, let P be the subset of purely imaginary roots λ such that $2T_\lambda = -\chi(\Gamma) \dim_{\mathbb{C}}(\mathfrak{g}_\lambda)$, i.e. the symplectic Γ action on \mathfrak{g}_λ is maximal with positive Toledo invariant. Say \mathfrak{t} is *balanced* with respect to ϕ if 0 belongs to the interior of the sum of the convex hull of the imaginary parts of elements of P and the linear span of the real and imaginary parts of roots not in $\pm P$.

Here is a necessary and sufficient condition for flexibility, for surface groups of sufficiently large genus.

Theorem 2.2 ([13, Theorem 3]). *Let G be a semisimple real algebraic group. Let Γ be the fundamental group of a closed surface of genus $\geq 2 \dim(G)^2$. Let $\phi : \Gamma \rightarrow G$ be a homomorphism with reductive Zariski closure. Then ϕ is flexible if and only if \mathfrak{c} , the center of the centralizer of $\phi(\Gamma)$, is balanced with respect to ϕ .*

The proof of Theorem 1 will rely on this criterion: we shall describe the centers of centralizers of reductive subgroups and their roots and pile up restrictions on the set P that make nonbalancedness exceptional. The reduction from nonreductive to reductive homomorphisms will be explained in Section 9.

3. Tightness

In this section we collect properties related to maximality of representations. We start with elementary facts. Deeper results will follow from tightness theory.

3.1. Preservation of maximality

We shall need that certain embeddings of Lie groups preserve maximality.

Definition 3.1. Let $\rho : H \rightarrow G$ be a homomorphism between reductive Hermitian groups and $F : Y \rightarrow X$ denote an equivariant totally geodesic map between the corresponding Hermitian symmetric spaces. Let ω_X (resp. ω_Y) denote the Kähler form, normalized so that the minimum sectional curvature equals -1 . Say ρ (or F) is *positively maximality preserving* if

$$\frac{1}{\text{rank}(X)} F^* \omega_X = \frac{1}{\text{rank}(Y)} \omega_Y.$$

Say ρ (or F) is merely *maximality preserving* if the above equality holds up to sign.

If ρ is positively maximality preserving, then for an equivariant map $f : \tilde{S} \rightarrow Y$,

$$\int_S (F \circ f)^* \omega_X = \int_S f^* F^* \omega_X = \int_S f^* \frac{\text{rank}(X)}{\text{rank}(Y)} \omega_Y,$$

hence a homomorphism $\phi : \Gamma \rightarrow H$ is maximal if and only if $\rho \circ \phi$ is. Furthermore, positively maximality preserving maps do not change the signs of Toledo invariants.

Lemma 3.2. *Isometric and holomorphic embeddings between equal rank Hermitian symmetric spaces are positively maximality preserving.*

Example 3.3. The embedding $\mathcal{D}_{p,p} \hookrightarrow \mathcal{D}_{p,q}$ between Grassmannians, corresponding to the embedding $SU(p, p) \hookrightarrow SU(p, q)$, is isometric and holomorphic, and thus positively maximality preserving.

Example 3.4. The embedding $\mathcal{G}_n \hookrightarrow \mathcal{G}_{n+1}$ of symmetric spaces, corresponding to the embedding $SO^*(2n) \hookrightarrow SO^*(2n+2)$, is isometric and holomorphic, and thus, if n is even, positively maximality preserving.

Example 3.5. The embedding $\mathcal{S}_n \hookrightarrow \mathcal{S}_{n+1}$ of symmetric spaces, corresponding to the embedding $Sp(2n, \mathbb{R}) \hookrightarrow Sp(2n+2, \mathbb{R})$, is isometric and holomorphic, and thus never maximality preserving.

We also need to understand when maximality is preserved under linear algebraic operations. Let us start with an easy case.

Lemma 3.6. *Let Γ be a surface group. Let I_0 be a unitary 1-dimensional representation of Γ , and W a sesquilinear representation of Γ . Then $\text{Hom}(I_0, W)$ is a maximal representation if and only if W is.*

Proof. $\text{Hom}(I_0, W)$ and W are isomorphic as projective representations. Since $U(W)$ acts on the symmetric space of $SU(W)$ via its quotient $PU(W)$, maximality is a projectively invariant property. \square

But we shall need a more general case in Subsection 7.2. The following lemma is a preparation for Lemma 3.8.

Lemma 3.7. *Let Γ be a surface group. Let W, W' be sesquilinear representations of Γ . Then*

1. $T(\bar{W}) = -T(W)$.
2. $T(W \oplus W') = T(W) + T(W')$.

Proof. 1. Passing from W to \bar{W} changes the sign of the complex structure on the symmetric space X of $SU(W)$. This changes the sign of the Kähler form, and thus the sign of Toledo invariants.

2. Let Y, Y' be the symmetric spaces of $SU(W)$ and $SU(W')$ respectively. When $Y \times Y'$ is mapped to the symmetric space X of $SU(W \oplus W')$, the Kähler form ω_X of X restricts on the complex totally geodesic manifold $Y \times Y'$ to $\omega_Y + \omega_{Y'}$, so the Toledo invariants add up. \square

Lemma 3.8. *Let Γ be a surface group. Let V be a unitary representation of Γ , and W a sesquilinear representation of Γ . Then $\text{Hom}(V, W)$ is a maximal representation of Γ if and only if W is.*

Proof. Let $\phi : \Gamma \rightarrow U(\text{Hom}(V, W))$ denote the Hom of the two given representations. Changing the given action on V into the trivial representation gives rise to a representation $\psi : \Gamma \rightarrow U(\text{Hom}(V, W))$. Let us show that ϕ and ψ have equal Toledo invariants. Split $W = W^+ \oplus W^-$ into positive definite and negative definite subspaces. Then $\text{Hom}(V, W^+)$ is a maximal positive definite subspace of $\text{Hom}(V, W)$, i.e. a point in the Hermitian symmetric space X associated to $SU(\text{Hom}(V, W))$. Its $U(V) \times U(W)$ -orbit Y is totally geodesic in X , and thus contractible. Therefore, one can choose an equivariant map $\tilde{f} : \tilde{\Sigma} \rightarrow X$ whose image is contained in Y . Note that the action of $(A, B) \in U(V) \times U(W)$ on $\text{Hom}(V, W)$ is

$$(A, B, N) \mapsto B^{-1}NA,$$

hence $U(V)$ leaves invariant the subspace $\text{Hom}(V, W^+)$ (see the proof of Lemma 3.14 below). Therefore $U(V)$ fixes Y pointwise, and \tilde{f} is equivariant with respect to both ϕ and ψ . Therefore the corresponding Toledo invariants are the same. Since ψ is a direct sum of $\dim(V)$ copies of the action on W , Lemma 3.7 applies, so $T(\text{Hom}(V, W)) = \dim(V)T(W)$. Since $\text{rank}(X)$ equals $\dim(V)$ times the rank of the symmetric space associated to $SU(W)$, ψ , and thus ϕ , is maximal if and only if the original Γ action on W is. \square

3.2. Tightness

Tightness theory is a way to draw strong consequences from the existence of maximal representations.

Definition 3.9 (Burger, Iozzi, Wienhard [4]). Let G be a *reductive Hermitian* group, i.e. a connected reductive Lie group in which the center is compact and such that the symmetric spaces X_i associated to all simple noncompact factors is Hermitian. Normalize the metric on X_i so that the minimum holomorphic sectional curvature equals -1 . Let κ_G^b denote the bounded continuous cohomology class on G defined by integrating the Kähler form of $X = \prod X_i$ on triangles with geodesic sides. Let Γ be locally compact group. Say a continuous homomorphism $\phi : \Gamma \rightarrow G$ is *tight* if

$$\|\phi^* \kappa_G^b\| = \|\kappa_G^b\|.$$

Example 3.10. Maximal homomorphisms of surface groups to reductive Hermitian groups are tight.

3.3. Maximality preserving versus tight

Here is the basic mechanism which makes tightness enter our arguments: if $\phi : \Gamma \rightarrow G$ is maximal and factors through $\rho : H \rightarrow G$, then ρ is tight. There is a converse statement.

Proposition 3.11. *Let H, G be reductive Hermitian groups. Assume that the symmetric space associated to H is irreducible. Let $\rho : H \rightarrow G$ be a continuous homomorphism. Then ρ is maximality preserving if and only if ρ is tight.*

Proof. This follows from [4, Proposition 2.12]. \square

Example 3.12 ([4, Example 8.7]). The obvious embeddings $SU(n, n) \rightarrow Sp(4n, \mathbb{R})$ and $SO^*(4n) \rightarrow Sp(8n, \mathbb{R})$ are tight. It follows that $SO^*(4n) \rightarrow SU(2n, 2n)$ is tight. All three embeddings are thus maximality preserving.

Direct proofs of these facts will be given in the appendix (Lemmas 10.2 and 10.3).

3.4. Consequences of tightness

Lemma 3.13. 1. *Let $H \subset G$ be connected real algebraic groups. If G is reductive Hermitian and the embedding $H \hookrightarrow G$ is tight, then H is reductive Hermitian too.*
 2. *Let $\rho : G \rightarrow G'$ be a tight homomorphism between reductive Hermitian groups. If the kernel of ρ is compact and G' is of tube type, so is G .*

Proof. This is a combination of Theorems 7.1 and 6.2 of [4]. \square

Lemma 3.14. *Let V, V' be vector spaces equipped with nondegenerate sesquilinear forms. Assume that $\text{Hom}(V, V')$ equipped with the natural sesquilinear form*

$$(f, f') \mapsto \text{Trace}(f^* \circ f')$$

(here f^ denotes the adjoint with respect to the sesquilinear forms on V and V') has vanishing signature. Assume that the induced homomorphism $U(V) \times U(V') \rightarrow U(\text{Hom}(V, V'))$ is tight (see Definition 3.9). Then one of V and V' is definite and the other has vanishing signature.*

Proof. Since $U(\text{Hom}(V, V'))$ is Hermitian of tube type, Lemma 3.13 implies that $U(V) \times U(V')$ is of tube type, up to compact groups. This implies that each of the sesquilinear vector spaces V and V' is either of vanishing signature or definite. Clearly, if both are definite, $U(V) \times U(V')$ is compact and tightness is impossible.

Let us show that V and V' cannot both have vanishing signature, i.e. one of them must be definite. For this, we use the tightness criterion of [4, Corollary 8.2]. Let J_V denote the generator of the center of the maximal compact subgroup of $U(V)$ which defines the complex structure on the symmetric space of $U(V)$. In a splitting $V = V^+ \oplus V^-$ into a sum of positive (resp. negative) definite subspaces we have

$$J_V = \begin{pmatrix} \frac{i}{2}I_{d_V/2} & 0 \\ 0 & -\frac{i}{2}I_{d_V/2} \end{pmatrix}.$$

Use a similar splitting $V' = V'^+ \oplus V'^-$ and split accordingly $\text{Hom}(V, V')$, whose elements become block matrices $N = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$.

For a basis $\{e_i\}$ of V such that $J = (B_v(e_i, e_j))$ is diagonal with entries ± 1 , and a basis $\{d_i\}$ of V' such that $J' = (B_{v'}(d_i, d_j))$ is diagonal with entries ± 1 , we have

$$\text{Trace}(f^* \circ f') = \text{Trace}(JA^*J'B)$$

where A, B are matrices for f and f' . The action of $U \times U' \in U(V) \times U(V')$ is $(U, U', N) \mapsto U'^{-1}NU$. It is easy to see that this action preserves the natural sesquilinear form on $\text{Hom}(V, V')$.

The linearized action is $(U, U', N) \mapsto -U'N + NU$. The image of $(J_V, J_{V'})$ under the linearized action is $N \mapsto \begin{pmatrix} 0 & -iB \\ iC & 0 \end{pmatrix}$, a map whose matrix is $\begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -i & 0 \\ 0 & 0 & 0 & i \end{pmatrix}$ in a splitting of $\text{Hom}(V, V')$ into a positive subspace $\text{Hom}(V^+, V'^+) \oplus \text{Hom}(V^-, V'^-)$ and a negative subspace $\text{Hom}(V^+, V'^-) \oplus \text{Hom}(V^-, V'^+)$. Therefore the inner product with the element

$$J_{\text{Hom}(V, V')} = \begin{pmatrix} \frac{i}{2}I_d & 0 & 0 & 0 \\ 0 & \frac{i}{2}I_d & 0 & 0 \\ 0 & 0 & -\frac{i}{2}I_d & 0 \\ 0 & 0 & 0 & -\frac{i}{2}I_d \end{pmatrix},$$

where $d = d_{\text{Hom}(V, V')}/4$, vanishes. According to [4, Corollary 8.2], the homomorphism $U(V) \times U(V') \rightarrow U(\text{Hom}(V, V'))$ is not tight. We conclude that one of V and V' has vanishing signature and the other one is definite. \square

The following example of nontight embedding will be useful in the proof of Corollary 8.7.

Lemma 3.15. *The injection $\iota : O(2, 2) \hookrightarrow U(2, 2)$ is not tight.*

Proof. $O(2, 2) \subset U(2, 2)$ is the fixed point set of conjugation σ , i.e. $\sigma \circ \iota = \sigma$. Now, σ induces an orientation reversing isometry of the symmetric space X of $U(2, 2)$, which changes the sign of the Kähler form, $\sigma^*\omega_X = -\omega_X$. Moreover, ι induces a totally geodesic embedding of symmetric spaces, still denoted by $\iota : Y \rightarrow X$. Since $\iota^*\omega_X = \iota^*\sigma^*\omega_X = -\iota^*\omega_X$, we have $\iota^*\omega_X = 0$, hence ι^* is not isometric in bounded continuous cohomology, so ι is not tight. \square

3.5. Alternative definition of balancedness

One may replace symplectic structures by sesquilinear structures in the definition of balancedness.

Proposition 3.16. *Let G be a semisimple real algebraic group. Let Γ be a surface group. Let $\phi : \Gamma \rightarrow G$ be a homomorphism. Let \mathfrak{c} denote the center of the centralizer of $\phi(\Gamma)$. Let $Z_G(\mathfrak{c})$ denote its centralizer in G . Let λ be a purely imaginary root of the adjoint action of \mathfrak{c} on $\mathfrak{g} \otimes \mathbb{C}$. Let \mathfrak{g}_λ denote the corresponding root space, equipped with a sesquilinear form s_λ and its imaginary part, the symplectic form Ω_λ . Then the symplectic representation $\Gamma \rightarrow Sp(\mathfrak{g}_\lambda, \Omega_\lambda)$ is maximal with positive Toledo invariant if and only if*

- s_λ has vanishing signature;

- the sesquilinear representation $\Gamma \rightarrow U(g_\lambda, s_\lambda)$ is maximal with positive Toledo invariant.

If this is the case, then the homomorphism $Z_G(\mathfrak{c}) \rightarrow U(g_\lambda, s_\lambda)$ is tight.

Proof. Assume that $\Gamma \rightarrow Sp(g_\lambda, \Omega_\lambda)$ is maximal. It factors through $U(g_\lambda, s_\lambda)$. Thus the inclusion $U(g_\lambda, s_\lambda) \hookrightarrow Sp(g_\lambda, \Omega_\lambda)$ is tight. Lemma 3.13 implies that $U(g_\lambda, s_\lambda)$ has tube type, i.e. the signature of s_λ vanishes.

According to Example 3.12, the embedding $U(g_\lambda, s_\lambda) \hookrightarrow Sp(g_\lambda, \Omega_\lambda)$ is positively maximality preserving, so maximality and positivity of Toledo invariant do not change when passing from symplectic to unitary groups.

The maximal representation $\Gamma \rightarrow U(g_\lambda, s_\lambda)$ factors via the homomorphism $Z_G(\mathfrak{c}) \rightarrow U(g_\lambda, s_\lambda)$, which must be tight itself. \square

Now we can explain how balancedness will be analyzed in what follows. For classical simple Lie groups, the root spaces \mathfrak{g}_λ turn out to be expressible as $\text{Hom}(V, V')$ spaces, and the centralizers $Z_G(\mathfrak{c})$ are products of classical simple Lie groups. Proposition 3.16, combined with Lemmas 3.13 and 3.14, restricts the possible groups involved, as will be seen in Sections 7 and 8.

4. Classical simple Lie groups

4.1. Definition

Classical simple real Lie groups are special linear groups of division rings D with center \mathbb{R} and special unitary groups of nondegenerate binary forms over D (see [17, appendix]).

There are only three such division rings: \mathbb{R} , \mathbb{C} and \mathbb{H} , leading to three special linear groups, $SL(n, \mathbb{R})$, $SL(n, \mathbb{C})$, and $SL(n, \mathbb{H})$.

Let ι be a continuous (anti-)automorphism of D : ι can be the identity, complex conjugation or quaternionic conjugation. Let $\epsilon = \pm 1$. By a (ι, ϵ) -symmetric binary¹ form on a right D -vector space V , we mean a D -valued \mathbb{R} -bilinear map $h : V \times V \rightarrow D$ such that

- (1) for all $v, v' \in V$ and all $q \in D$, $h(v, v'q) = h(v, v')q$;
- (2) for all $v, v' \in V$, $h(v', v) = \epsilon \iota(h(v, v'))$.

Note that from (2), if ι is a conjugation, it follows that $h(vq, v') = \bar{q}h(v, v')$. Say h is nondegenerate if the only $v \in V$ such that $h(v, v') = 0$ for all $v' \in V$ is 0.

The group $U(V, h)$ consists of right D -linear self maps of V which preserve h . We are interested in the special unitary group $SU(V, h) = U(V, h) \cap SL(V, D)$.

When $D = \mathbb{R}$, the only choice for ι is the identity, leading to

- if $\epsilon = 1$, real special orthogonal groups, indexed by dimension and signature and denoted by $SO(p, q)$, where $p + q = \dim_{\mathbb{R}}(V)$;
- if $\epsilon = -1$, real symplectic groups, indexed by dimension only and denoted by $Sp(n, \mathbb{R})$, where $n = \dim_{\mathbb{R}}(V)$ is even.

¹ To avoid confusion, we keep the words Hermitian for positive definite forms, and sesquilinear for the complex case.

When $D = \mathbb{C}$ there are two choices for ι . If ι is the identity, this leads to complex special orthogonal (if $\epsilon = 1$) and symplectic (if $\epsilon = -1$) groups, indexed by dimension and denoted respectively by $SO(n, \mathbb{C})$ and $Sp(n, \mathbb{C})$. If ι is complex conjugation, $\epsilon = \pm 1$ lead to groups called special unitary groups, indexed by dimension and signature and denoted by $SU(p, q)$, where $p + q = \dim_{\mathbb{C}}(V)$.

When $D = \mathbb{H}$ there are two choices for ι . If ι is the identity, no nonzero (ι, ϵ) -symmetric forms exist. If ι is quaternionic conjugation, this leads to

- if $\epsilon = 1$, quaternionic unitary groups, indexed by dimension and signature and denoted by $Sp(p, q)$, $p + q = \dim_{\mathbb{H}}(V)$;
- if $\epsilon = -1$, quaternionic skew-unitary groups, indexed by dimension only and denoted by $SO^*(2n)$, where $n = \dim_{\mathbb{H}}(V)$ is even.

4.2. Real forms of complex groups

We shall be mainly concerned with the seven families of noncomplex groups, three attached to binary forms on real vector spaces, one on complex vector spaces and three on quaternionic vector spaces. Each of these groups is obtained as the fixed point set of an anti- \mathbb{C} -linear involutive automorphism σ of a complex Lie group, as we now explain.

Consider first the complex unitary family $U(p, q)$. Here, the data is a nondegenerate sesquilinear form s on a complex vector space V . For $f \in GL(V)$, let f^* denote the s -adjoint of f , defined by

$$\forall v, v' \in V, \quad s(f(v), v') = s(v, f^*(v')).$$

Then $\sigma(f) = (f^*)^{-1}$ is an anti- \mathbb{C} -linear involutive automorphism of $GL(V)$ and of $SL(V)$. The fixed point set of σ in $SL(V)$ is $SU(V, s)$.

The six remaining families admit a common construction.

Given a real vector space $V_{\mathbb{R}}$, let $V = V_{\mathbb{R}} \otimes \mathbb{C}$ and let $\tau =$ conjugation. Note that $\tau^{-1} = \tau$. If b is a nondegenerate quadratic or symplectic form on $V_{\mathbb{R}}$, let $B = b \otimes \mathbb{C}$.

Given a right quaternionic vector space $V_{\mathbb{H}}$, pick a basis $(1, i, j, k)$ of \mathbb{H} , and use right multiplication by i to turn $V_{\mathbb{H}}$ into a complex vector space denoted by V . Let τ be right multiplication by j . Note that $\tau^{-1} = -\tau$. For $q = a + jb \in \mathbb{H}$ with $a, b \in \mathbb{C}$, denote $\mathcal{C}(q) = a$. If h is a nondegenerate (τ, ϵ) -symmetric binary form on $V_{\mathbb{H}}$, let, for $v, v' \in V$,

$$B(v, v') = \mathcal{C}(h(vj, v')).$$

Then B is a nondegenerate $(-\epsilon)$ -symmetric \mathbb{C} -bilinear form on V .

In both cases, τ is anti- \mathbb{C} -linear, $\tau^{-1} = \eta\tau$ for some $\eta \in \pm 1$, and

$$B(\tau(v), \tau(v')) = \overline{B(v, v')}.$$

For $f \in \text{End}_{\mathbb{C}}(V)$, set

$$\sigma(f) = \tau \circ f \circ \tau^{-1}.$$

Then σ is an anti- \mathbb{C} -linear involutive automorphism of the algebra $\mathfrak{gl}(V) = \text{End}_{\mathbb{C}}(V)$ and of its subgroups $SL(V)$ and $O(V, B)$. The fixed point set of σ in $SL(V)$ is $SL(V_{\mathbb{R}})$ (resp.

$SL(V_{\mathbb{H}})$). The fixed point set of σ in $O(V, B)$, i.e., $\tau \circ f = f \circ \tau$, is $U(V_{\mathbb{R}}, b)$ (resp. $U(V_{\mathbb{H}}, h)$). This construction yields six of the seven families of noncomplex classical groups.

4.3. Consequences for roots

The special form of the involution σ for six of the seven families of noncomplex groups has the following consequence.

Lemma 4.1. *Let G belong to one of the above six families of classical simple Lie groups. Let $H \subset G$ be a reductive subgroup. Let $\mathfrak{c} \subset \mathfrak{g}$ be the center of its centralizer. Let ℓ be a root of \mathfrak{c} on V . Let I_{ℓ} denote the corresponding root space. Then $\bar{\ell}$ is a root and $I_{\bar{\ell}} = \tau(I_{\ell})$. Furthermore,*

- if $G = SL(n, \mathbb{R})$, $O(p, q)$ or $Sp(n, \mathbb{R})$, then $I_{\bar{\ell}} = \overline{I_{\ell}}$;
- if $G = SL(n, \mathbb{H})$, $SO^*(2n)$ or $Sp(p, q)$, then $I_{\ell} + I_{\bar{\ell}}$ is a quaternionic subspace;
- if $\ell \neq 0$, then I_{ℓ} and I_0 are orthogonal with respect to the $(\bar{\cdot}, \epsilon)$ -symmetric binary form h , and therefore I_0 is nondegenerate.

Proof. We let \mathbb{C} act on V on the right. Let $Z \in \mathfrak{c}$ and $v \in I_{\ell}$. Then

$$Z(\tau(v)) = \tau(Z(v)) = \tau(v\ell(Z)) = \tau(v)\overline{\ell(Z)},$$

showing that $I_{\bar{\ell}} = \tau(I_{\ell})$. For the three real families, τ is conjugation, thus $I_{\bar{\ell}} = \overline{I_{\ell}}$. For the three quaternionic families, τ is right multiplication by j , so $I_{\ell} + I_{\bar{\ell}}$ is stable by right multiplication by i and j , i.e. a quaternionic subspace.

Since h is G -invariant, for all $v, v' \in V$ and $Z \in \mathfrak{c}$ we have $h(Z(v), v') + h(v, Z(v')) = 0$. If $v \in I_{\ell}$ and $v' \in I_0$, then

$$0 = h(Z(v), v') + h(v, Z(v')) = h(v\ell(Z), v') = \overline{\ell(Z)}h(v, v').$$

This implies that $h(v, v') = 0$ if $\ell \neq 0$. This shows that I_0 is h -orthogonal to the sum of all other root spaces. Therefore the restriction of h to I_0 is nondegenerate. \square

4.4. Killing form

Here is a formula for the sesquilinear structure appearing in Theorem 2.2, valid in all cases.

Lemma 4.2. *Let $\mathfrak{g}^{\mathbb{C}} \subset \mathfrak{sl}(n, \mathbb{C})$ be a complex Lie subalgebra of $\mathfrak{sl}(n, \mathbb{C})$. Let σ be an anti- \mathbb{C} -linear involutive automorphism of $\mathfrak{g}^{\mathbb{C}}$, with fixed point set \mathfrak{g} . The natural sesquilinear form s on $\mathfrak{g} \otimes \mathbb{C}$ arising from the Killing form of \mathfrak{g} (see Section 2) is proportional to*

$$s(X, X') = \text{Trace}(\sigma(X) \circ X').$$

Proof. Since σ is anti- \mathbb{C} -linear, we have $\sigma(if) = -i\sigma(f) = -if$ for $f \in \mathfrak{g}$. It follows that the -1 -eigenspace of σ in $\mathfrak{g}^{\mathbb{C}}$ is $i\mathfrak{g}$. The map

$$\mathfrak{g}^{\mathbb{C}} \rightarrow \mathfrak{g} \otimes \mathbb{C}, \quad X \mapsto \left(\frac{X + \sigma(X)}{2}, \frac{X - \sigma(X)}{2i} \right),$$

is an isomorphism. It pulls back conjugation on $\mathfrak{g} \otimes \mathbb{C}$ to σ on $\mathfrak{g}^{\mathbb{C}}$, thus it pulls back the sesquilinear form $\bar{X} \cdot X$ of $\mathfrak{g} \otimes \mathbb{C}$ to $\sigma(X) \cdot X$ on $\mathfrak{g}^{\mathbb{C}}$. The Killing form on $\mathfrak{g}^{\mathbb{C}}$ is proportional to the restriction to $\mathfrak{g}^{\mathbb{C}}$ of the Killing form of $\mathfrak{sl}(n, \mathbb{C})$, whence the formula $\text{Trace}(\sigma(X) \circ X)$ follows. \square

4.5. Flexibility of compact and complex Lie groups

For completeness' sake, we recall here the treatment of compact and complex Lie groups from [13].

Proposition 4.3. *Let G be a compact or complex semisimple Lie group. Let $\mathfrak{c} \subset \mathfrak{g}$ be the center of the centralizer of a reductive subgroup. Then \mathfrak{c} is balanced.*

Proof. Let $X \in \mathfrak{g} \otimes \mathbb{C}$, $X = f + ig$. Then $s(X, X) = \bar{X} \cdot X = f \cdot f + g \cdot g$. If G is compact, the Killing form is negative definite, and so is s . For all roots λ of \mathfrak{c} in the adjoint representation, the sesquilinear form s_{λ} on the root space \mathfrak{g}_{λ} is negative definite. No root has vanishing signature. According to Proposition 3.16, P is empty so \mathfrak{c} is balanced.

If G is complex, then centralizers are complex Lie subgroups, \mathfrak{c} is a complex vector subspace, and roots λ are \mathbb{C} -linear maps. None of them is purely imaginary (i.e. takes purely imaginary values on \mathfrak{c}). Thus P is empty so \mathfrak{c} is balanced. \square

5. Complexified centers of centralizers in $SL(n, \mathbb{C})$

The first step is to list the possible complexified centers $\mathfrak{c} \otimes \mathbb{C}$ and describe the root structure. This depends only on the complexified Lie algebra.

Lemma 5.1. *Let \mathfrak{g} be a real Lie algebra. Let $H \subset \mathfrak{g}$ be a subset. Let $ZZ_{\mathfrak{g}}(H)$ denote the center of its centralizer. Then*

$$ZZ_{\mathfrak{g}}(H) \otimes \mathbb{C} = ZZ_{\mathfrak{g} \otimes \mathbb{C}}(H).$$

Our approach consists in using the standard complex representation V of $\mathfrak{g} \otimes \mathbb{C}$. The root space decomposition of V under $\mathfrak{c} \otimes \mathbb{C} = ZZ_{\mathfrak{g} \otimes \mathbb{C}}(H)$ is obtained from the isotypical decomposition of V under H . This easily provides us with the decomposition of $\text{End}(V)$ under $\mathfrak{c} \otimes \mathbb{C}$, and then of the invariant subspace $\mathfrak{g} \otimes \mathbb{C} \subset \text{End}(V)$.

In this section, we treat the model case of $SL(n, \mathbb{C})$, and in the next section, the more elaborate cases $O(n, \mathbb{C})$ and $Sp(n, \mathbb{C})$.

5.1. H -modules

Definition 5.2. Let H be a group. The data of a finite-dimensional complex vector space and a homomorphism of H onto a reductive real algebraic subgroup of $GL(V)$ is called an H -module.

Note that H -invariant subspaces of H -modules are again H -modules. An H -module is *irreducible* if it has no proper H -submodules.

Definition 5.3. Let $E(H)$ denote the set of equivalence classes of H -modules. Given an H -module V and $\pi \in E(H)$, let I_π , the π -isotypical component of V , be the sum of all submodules of V belonging to the equivalence class π .

The following lemma is well known, but we give a full proof since it serves as a model for orthogonal, symplectic and unitary versions of it given in the next section.

Lemma 5.4. Any H -module V splits as a direct sum of its isotypical components,

$$V = \bigoplus_{\pi \in E(H)} I_\pi.$$

Proof. Let $W \subset I_\pi$ be an irreducible invariant subspace. Since H is reductive, for each invariant subspace Z of I_π belonging to the class π there exists an H -invariant complement to Z , thus an H -invariant projector $p_Z : I_\pi \rightarrow Z$. Since such submodules Z generate I_π , for at least one such Z we have $p_Z(W) \neq 0$, thus $p_Z(W) = Z$ and W belongs to π .

More generally, if $F \subset E(H)$ is a subset, and $W \subset \sum_{\pi \in F} I_\pi$ is an irreducible invariant subspace, then W belongs to one of the classes in F . Indeed, otherwise $p_Z(W) = 0$ for all invariant subspaces Z whose class belongs to F , and these generate $\sum_{\pi \in F} I_\pi$. In particular, for every $\pi \in E(H)$, $I_\pi \cap \sum_{\pi' \neq \pi} I_{\pi'} = \{0\}$, which shows that the sum $\sum_{\pi \in E(H)} I_\pi$ is direct.

Since H is reductive, the invariant subspace $\bigoplus_{\pi \in E(H)} I_\pi$ admits an invariant complement, which contains an irreducible subspace. This is a contradiction unless $\bigoplus_{\pi \in E(H)} I_\pi = V$. \square

5.2. Centers of centralizers in $SL(n, \mathbb{C})$

Lemma 5.5. Let $H \subset SL(n, \mathbb{C})$ be a reductive subgroup. Let $\mathfrak{c} \subset \mathfrak{sl}(n, \mathbb{C})$ be the center of its centralizer. Let L denote the set of nonzero roots of \mathfrak{c} in the standard representation of $\mathfrak{sl}(n, \mathbb{C})$, and d_ℓ the dimensions of the corresponding eigenspaces. Then L has $\dim(\mathfrak{c}) + 1$ elements, which satisfy exactly one linear relation,

$$\sum_{\ell \in L} d_\ell \ell = 0.$$

Furthermore, the map

$$L \times L \setminus \text{diagonal} \rightarrow \Lambda, \quad (\ell, \ell') \mapsto \ell - \ell',$$

is one-to-one and onto the set Λ of nonzero roots of \mathfrak{c} in the adjoint representation of $\mathfrak{sl}(n, \mathbb{C})$.

Proof. Since H is reductive, the standard action of H on \mathbb{C}^n splits into irreducibles. Let us group them into isotypical components I_ℓ ; each I_ℓ is the direct sum of k_ℓ isomorphic irreducible summands. According to Schur's Lemma, the group of H -automorphisms (i.e. automorphisms which commute with H) of the representation I_ℓ is isomorphic to $GL(k_\ell, \mathbb{C})$. Then

$$Z_{GL(n, \mathbb{C})}(H) = \prod_{\ell \in L} Z_{GL(I_\ell)}(H|_{I_\ell}) \simeq \prod_{\ell \in L} GL(k_\ell, \mathbb{C}),$$

whose center is $(\mathbb{C}^*)^L$, acting on \mathbb{C}^n by multiplication by a different constant on each I_ℓ . Pick a basis of \mathbb{C}^n adapted to the splitting $\mathbb{C}^n = \bigoplus_{\ell} I_\ell$. Then the center \mathfrak{c} of $Z_{SL(n, \mathbb{C})}(H)$ consists of diagonal matrices $\text{diag}(a_1, \dots, a_n)$ whose entries corresponding to basis vectors from the same I_ℓ are equal, and which sum to 0. It follows that the elements of L generate \mathfrak{c}^* and satisfy only one linear relation, $\sum_{\ell \in L} d_\ell \ell = 0$. In particular, if (ℓ, ℓ') and (m, m') are distinct ordered pairs of distinct elements of L , then $\ell - \ell' - m + m'$ does not vanish identically on \mathfrak{c} . This shows that the map

$$L \times L \setminus \text{diagonal} \rightarrow \Lambda, \quad (\ell, \ell') \mapsto \ell - \ell',$$

which is clearly surjective onto the set of nonzero roots of the adjoint action on $\mathfrak{sl}(n, \mathbb{C})$, is injective as well. Furthermore, note that $\mathfrak{sl}(n, \mathbb{C}) \subset \text{Hom}(\mathbb{C}^n, \mathbb{C}^n) = \text{Hom}(\bigoplus_{\ell} I_\ell, \bigoplus_{\ell'} I_{\ell'})$ and the root space for $\ell - \ell'$ is $\text{Hom}(I_\ell, I_{\ell'})$. \square

6. Centralizers in orthogonal, symplectic or unitary groups

In this section, the root space decomposition of \mathfrak{g} under the center of the centralizer of a reductive subgroup is given, when $\mathfrak{g} = \mathfrak{so}(n, \mathbb{C})$ or $\mathfrak{sp}(n, \mathbb{C})$. This is a first step in handling real forms of these Lie algebras. With little extra effort, one can treat simultaneously the case of $\mathfrak{g} = \mathfrak{su}(p, q)$. This will help treating this particular real form of $\mathfrak{sl}(n, \mathbb{C})$.

Let (V, B) be a complex vector space equipped with a nondegenerate binary form B of one of the following three types:

- symmetric bilinear,
- skew-symmetric bilinear,
- symmetric sesquilinear,

which we denote by (ι, ϵ) -symmetric, $\epsilon = \pm 1$, $\iota = 1$ or $\bar{}$ (identity or conjugation). Note that the combination $(\iota, \epsilon) = (\bar{}, -1)$ makes perfect sense but does not bring anything new, since if B is a skew-symmetric sesquilinear form, then iB is symmetric sesquilinear. Let $O = O^{\iota, \epsilon}(V, B)$ denote its automorphism group (note that if B is skew-symmetric bilinear (resp. sesquilinear), this is a symplectic (resp. unitary) rather than an orthogonal group, whence the notation $O^{\iota, \epsilon}$). Let $H \subset O$ be a reductive real algebraic subgroup. In this section, we describe the center of the centralizer of H in O .

6.1. Bilinear and sesquilinear forms

Notation 6.1. Let ι be a continuous automorphism of \mathbb{C} , i.e. either conjugation or the identity. Let $\epsilon \in \{-1, 1\}$. Let V be a complex vector space. A (ι, ϵ) -symmetric form on V is a real bilinear form $B : V \times V \rightarrow \mathbb{C}$ such that for $\lambda, \lambda' \in \mathbb{C}$ and $v, v' \in V$,

- $B(\lambda v, \lambda' v') = \iota(\lambda)\lambda' B(v, v')$;
- $B(v', v) = \epsilon \iota B(v, v')$.

In other words, if $\iota = 1$ and $\epsilon = 1$, B is symmetric bilinear. If $\iota = 1$ and $\epsilon = -1$, B is skew-symmetric bilinear. If $\iota = \bar{}$ and $\epsilon = 1$, B is symmetric sesquilinear. If $\iota = \bar{}$ and $\epsilon = -1$, B is skew-symmetric sesquilinear. We shall ignore the fourth case, since if B is a skew-symmetric sesquilinear form, then iB is symmetric sesquilinear.

Notation 6.2. Let ι be a continuous automorphism of \mathbb{C} , i.e. either conjugation or the identity. Let V be a complex vector space. Then V^ι means V if $\iota = 1$, and \bar{V} if $\iota = \bar{}$. Also, $V^{\iota,*}$ means the dual vector space V^* if $\iota = 1$, and \bar{V}^* (i.e. the space of anti- \mathbb{C} -linear forms on V) if $\iota = \bar{}$.

Let B be a (ι, ϵ) -symmetric form on V . Let $\sharp_B : V \rightarrow V^{\iota,*}$ denote the \mathbb{C} -linear map which maps $v \in V$ to the anti- \mathbb{C} -linear functional

$$\sharp_B(v) : v' \mapsto B(v', v).$$

Say that B is *nondegenerate* if \sharp_B is an isomorphism. If not, its kernel is called the *nullspace* of B .

Notation 6.3. Given a linear map $L : V \rightarrow V^{\iota,*}$, there is an adjoint map $L^{\iota,\top} : V \rightarrow V^{\iota,*}$ defined by

$$\langle L^{\iota,\top}(v'), v \rangle = \langle L(v), v' \rangle,$$

where $\langle v^*, v \rangle$ denotes the evaluation of the linear or anti-linear form v^* on the vector $v \in V$.

If L is ϵ -symmetric, i.e. $L^{\iota,\top} = \epsilon \iota L$, the formula

$$B(v, v') = \langle L(v'), v \rangle$$

defines a (ι, ϵ) -symmetric form such that $\sharp_B = L$. Therefore B and \sharp_B are equivalent data.

Example 6.4. Let W be a complex vector space. Then the tautological isomorphism of $V = W \times W^{\iota,*}$ to $V^{\iota,*}$ gives rise to a tautological (ι, ϵ) -symmetric form on V ,

$$(v, v^*) \cdot (w, w^*) = \epsilon \iota(\langle v^*, w \rangle) + \langle w^*, v \rangle.$$

Any two nondegenerate $(1, \epsilon)$ -symmetric forms are isomorphic. On the other hand, $(\iota, 1)$ -symmetric forms, i.e. symmetric sesquilinear forms, take real values on the diagonal, so sign and signature issues arise: two nondegenerate $(\iota, 1)$ -symmetric forms are isomorphic if and only if they have the same signature. For instance, the tautological form of Example 6.4 has vanishing signature.

6.2. Bilinear and sesquilinear H -modules

Definition 6.5. Let H be a group. The data of a finite-dimensional complex vector space equipped with a bilinear (either symmetric or skew-symmetric) or sesquilinear form and a homomorphism of H onto a reductive real algebraic subgroup of its automorphism group will be called a (ι, ϵ) -linear H -module.

Lemma 6.6. Let W be an irreducible H -module. The space of H -invariant bilinear (resp. sesquilinear) forms on W has dimension at most 1. A nonzero H -invariant bilinear (resp. sesquilinear) form is automatically nondegenerate, and in the bilinear case, it is either symmetric or skew-symmetric.

Proof. Let b be an H -invariant bilinear form on W . Its nullspace is H -invariant. Therefore b is either zero or nondegenerate. Assume b is nonzero and denote by $\sharp_b : W \rightarrow W^{\iota,*}$ the corresponding isomorphism (in the sesquilinear case, $\iota = \text{conjugation}$, $W^{\iota,*} = \bar{W}^*$). Let b' be another H -invariant bilinear form on W . Then $L = (\sharp_b)^{-1} \circ \sharp_{b'}$ is an H -equivariant endomorphism of W , thus L is a multiple of identity (Schur's Lemma). This shows that b' is a multiple of b . Now, b has a symmetric and a skew-symmetric component. They have to be linearly dependent; in the bilinear case ($\iota = 1$), this implies that one of them vanishes. Therefore b is either symmetric or skew-symmetric. \square

Definition 6.7. Say an H -module is (ι, ϵ) -orthogonal if it admits an invariant nondegenerate (ι, ϵ) -symmetric form.

Corollary 6.8. The classification of irreducible (ι, ϵ) -linear H -modules can be deduced from the classification of irreducible H -modules: the forgetful map $E^{(\iota, \epsilon)}(H) \rightarrow E(H)$ is onto, the fiber of an equivalence class of irreducible H -modules contains one or two elements depending on whether it is (ι, ϵ) -orthogonal or not.

Lemma 6.9. Let (V, B) be a (ι, ϵ) -linear H -module. Let W and W' be distinct irreducible H -invariant subspaces. Assume that W' is not orthogonal to W . Then W' is isomorphic, as an H -module, to the (conjugate-)dual $W^{\iota,*}$ of W (in the sesquilinear case, $W^{\iota,*} = \bar{W}^*$).

Proof. The map $v \mapsto (\sharp_B(v))|_W, W' \rightarrow W^{\iota,*}$, is H -equivariant. According to Schur's Lemma, such a map is either zero or an isomorphism, and all such maps are proportional. By assumption, it does not vanish, thus W' and $W^{\iota,*}$ are isomorphic H -modules. \square

Definition 6.10. Let H be a group and V an H -module. Say V is *bi-isotypical* if there exists an irreducible H -module Z such that every irreducible invariant subspace $W \subset V$ is isomorphic either to Z or to $Z^{\iota,*}$.

Corollary 6.11. Let (V, B) be a nondegenerate (ι, ϵ) -linear H -module. Then V canonically splits as an orthogonal direct sum of its bi-isotypical components,

$$V = \bigoplus_{\pi} I_{\pi, \pi^{\iota,*}},$$

where, given an equivalence class π of irreducible H -modules, $I_{\pi, \pi^{l,*}} = I_{\pi} + I_{\pi^{l,*}}$ is the sum of all irreducible invariant subspaces of V isomorphic either to π or to $\pi^{l,*}$. Furthermore, $I_{\pi, \pi^{l,*}}$ is nondegenerate, and the centralizer $Z_{O(V,B)}(H)$ of H in the automorphism group is isomorphic to a direct product,

$$Z_{O(V,B)}(H) = \prod_{\pi} Z_{O(I_{\pi, \pi^{l,*}})}(H).$$

6.3. Examples of bi-isotypical bilinear/sesquilinear H -modules

From now on, we analyze nondegenerate bi-isotypical (l, ϵ) -linear H -modules. Here are two examples.

Example 6.12. Let $Z = \pi^{\oplus r}$ be an isotypical H -module. Set $V = Z \times Z^{l,*}$, and equip it with the canonical (l, ϵ) -symmetric form

$$(v, v^*) \cdot (w, w^*) = \epsilon l(\langle v^*, w \rangle) + \langle w^*, v \rangle.$$

Then the centralizer of H in $O^{(l, \epsilon)}(V, \cdot)$ is isomorphic to an orthogonal, symplectic or unitary group if π and $\pi^{l,*}$ are equivalent:

$$Z_{O^{(l, \epsilon)}(V, \cdot)}(H) \cong O^{(l, \epsilon)}(2r, \mathbb{C}) := \begin{cases} O(2r, \mathbb{C}) & \text{if } l = 1, \epsilon = 1, \\ Sp(2r, \mathbb{C}) & \text{if } l = 1, \epsilon = -1, \\ U(r, r) & \text{if } l = -, \epsilon = 1, \end{cases}$$

and otherwise to a general linear group,

$$Z_{O^{(l, \epsilon)}(V, \cdot)}(H) \cong GL(r, \mathbb{C}).$$

Proof. Pick a basis e_i of W and take r copies of it to form a basis of Z . Take the image of this basis under \sharp_B to get a basis of $Z^{l,*}$, i.e. for $\epsilon = 1$, choose $B(e_i, e_i) = 1$ and $e_i^* = \sharp_B(e_i)$, and then take $\{e_1, \dots, e_1^*, \dots\}$ as a basis for V . For $\epsilon = -1$, choose e_1, \dots, e_{2k} so that $B(e_i, e_{i+k}) = 1 = -B(e_{i+k}, e_i)$ and $B(e_i, e_j) = 0$ otherwise. Choose a basis for V in this case to be $\{e_1, \dots, e_k, e_{k+1}, \dots, e_{2k}, e_{k+1}^*, \dots, e_{2k}^*, e_1^*, \dots, e_k^*\}$. This gives a basis of V in which the matrix of the bilinear/sesquilinear form B equals $\begin{pmatrix} 0 & 1 \\ \epsilon & 0 \end{pmatrix}$ (blocks have size rd where $d = \dim \pi$). In this basis, the matrix of an element g of H splits into blocks of size d , with the first r diagonal blocks equal to $\pi(g)$ and the last r equal to $\pi(g^{-1})^{l, \top}$; all other blocks vanish.

If π and $\pi^{l,*}$ are equivalent, then endomorphisms of V which commute with H have matrices whose blocks of size d are scalar, i.e. proportional to the unit $d \times d$ matrix. In other words, they can be written as $A \otimes 1$ where $A \in GL(2r, \mathbb{C})$. Such a matrix preserves B if and only if

$$(A \otimes 1)^{l, \top} \left(\begin{pmatrix} 0 & 1 \\ \epsilon & 0 \end{pmatrix} \otimes 1 \right) (A \otimes 1) = \begin{pmatrix} 0 & 1 \\ \epsilon & 0 \end{pmatrix} \otimes 1,$$

i.e. A belongs to $O(2r, \mathbb{C})$ if $\epsilon = 1$, to $Sp(2r, \mathbb{C})$ if $\epsilon = -1$ and to $U(r, r)$ if $l =$ conjugation. In other words,

$$Z_{O^{(l, \epsilon)}(V, \cdot)}(H) \cong O^{(l, \epsilon)}(2r, \mathbb{C}).$$

If π and $\pi^{\iota,*}$ are not equivalent, then endomorphisms of V which commute with H preserve the splitting $V = I_\pi \oplus I_{\pi^{\iota,*}}$ and have matrices whose blocks of size d are scalar, i.e. proportional to the unit $d \times d$ matrix. In other words, they can be written as $(A \oplus A') \otimes 1$ where $A, A' \in GL(r, \mathbb{C})$. Such a matrix preserves B if and only if $A' = (A^{\iota,\top})^{-1}$. In other words,

$$Z_{O^{(\iota,\epsilon)}(V, \cdot)}(H) \cong GL(r, \mathbb{C}). \quad \square$$

Example 6.13. Let $V = \pi^{\oplus r}$ be an isotypical H -module such that π is (ι, ϵ) -orthogonal, i.e. preserves a nondegenerate (ι, ϵ) -symmetric form b . Let D be a real diagonal invertible $r \times r$ matrix. Set $B = b \otimes D$, i.e. (V, B) is an orthogonal direct sum of r real multiples of the same nondegenerate (ι, ϵ) -linear H -module. Then the centralizer of H in $O^\epsilon(V, b \otimes D)$ is isomorphic to an orthogonal/unitary group,

$$Z_{O^{\iota,\epsilon}(V, b \otimes D)}(H) \cong O^\iota(\mathbb{C}^r, D).$$

Note that in the bilinear case, $O(\mathbb{C}^r, D) = O(r, \mathbb{C})$ is a genuine orthogonal group, even when $\epsilon = -1$, i.e. when we deal with skew-symmetric forms. In the sesquilinear case, $O^\iota(\mathbb{C}^r, D) = U(p, q)$ where $p - q = \text{sign}(D)$.

Proof. Repeat the same basis of π to get a basis of V . The matrix of $B = b \otimes D$ is block diagonal with diagonal blocks of size $d = \dim \pi$, each equal to a real multiple of b . Element $g \in H$ acts by a block diagonal matrix, with $d \times d$ diagonal blocks equal to $\pi(g)$. The centralizer of H in $GL(V)$ consists of matrices with scalar $d \times d$ blocks, i.e. of the form $A \otimes 1$ for $A \in GL(r, \mathbb{C})$. (ι, ϵ) -orthogonal matrices satisfy

$$(A \otimes 1)^{\iota,\top} (b \otimes D) (A \otimes 1) = b \otimes (A^{\iota,\top} D A) = b \otimes D,$$

i.e. $A^{\iota,\top} D A = D$. Thus

$$Z_{O^\epsilon(V, B)}(H) \cong O^\iota(\mathbb{C}^r, D). \quad \square$$

6.4. Classification of bi-isotypical bi/sesquilinear H -modules

There are three cases, depending on whether π is (ι, ϵ) -orthogonal, $(\iota, -\epsilon)$ -orthogonal, or neither. In each case, we will need the following lemma.

Lemma 6.14. *Let $V = W \oplus W'$ be a (ι, ϵ) -linear H -module where W, W' are isotropic, isotypic, and the map $L : W' \rightarrow W^{\iota,*}$, $L(v') = (\sharp_B(v'))|_W$, is an isomorphism. Then V is isomorphic to Example 6.12.*

Proof. The H -map $(w + w') \mapsto (w, L(w'))$ is an isometry $W \oplus W' \rightarrow W \times W^{\iota,*}$, since

$$\begin{aligned} (v, L(v')) \cdot (w, L(w')) &= \epsilon \iota(\langle L(v'), w \rangle) + \langle L(w'), v \rangle = \epsilon \iota(\langle \sharp_B(v'), w \rangle) + \langle \sharp_B(w'), v \rangle \\ &= \epsilon \iota(B(w, v')) + B(v, w') = B(v', w) + B(v, w') \\ &= B(v + v', w + w'). \end{aligned} \quad \square$$

Proposition 6.15. *Let (V, B) be a nondegenerate bi-isotypical bilinear H -module. Assume π and $\pi^{\iota,*}$ are not isomorphic. Then (V, B) is isomorphic to Example 6.12.*

Proof. Irreducible invariant subspaces of V belong to either π or $\pi^{\iota,*}$, which do not admit nondegenerate invariant bilinear/sesquilinear forms, thus all are isotropic. Lemma 6.9 implies that any two distinct irreducible invariant subspaces of I_π are orthogonal. Thus I_π is isotropic, and so is $I_{\pi^{\iota,*}}$. Since V is nondegenerate, $\sharp_B : V \rightarrow V^{\iota,*}$ induces isomorphisms $L : I_\pi \rightarrow (I_{\pi^{\iota,*}})^{\iota,*}$ and $L' : I_{\pi^{\iota,*}} \rightarrow (I_\pi)^{\iota,*}$ related by $L^\perp = \epsilon L'$. According to Lemma 6.14, this shows that V is isomorphic, as a (ι, ϵ) -linear H -module, to Example 6.12. \square

Proposition 6.16. *Let $V = I_\pi$ be an isotypical H -module equipped with a nondegenerate ϵ -symmetric bilinear form B . Assume π is $(-\epsilon)$ -orthogonal. Then (V, B) is isomorphic to Example 6.12.*

Proof. Since π is not ϵ -orthogonal, all irreducible H -submodules of V are isotropic. Let W be one of them. An H -invariant complement to W^\perp contains an irreducible H -submodule W' , which is not orthogonal to W . The H -map $L : W' \rightarrow W^*$, $v' \mapsto (\sharp_B(v'))|_W$, is nonzero, thus an isomorphism, and Lemma 6.14 implies that $W \oplus W'$ is isomorphic to Example 6.12, in particular, it is nondegenerate. Its orthogonal is again nondegenerate, isotypic, modeled on a $(-\epsilon)$ -orthogonal H -module. By induction on dimension, V is an orthogonal direct sum of copies of Example 6.12, thus isomorphic to Example 6.12. \square

Proposition 6.17. *Let $V = I_\pi$ be an isotypical H -module equipped with a nondegenerate (ι, ϵ) -symmetric bilinear form B . Assume π is (ι, ϵ) -orthogonal. Then (V, B) is isomorphic to Example 6.13.*

Proof. Let us show that V contains at least one nondegenerate irreducible invariant subspace. Pick an irreducible H -submodule W of V . If it is nondegenerate, we are done. Otherwise, W is isotropic. An H -invariant complement to W^\perp contains an irreducible H -submodule W' , which is not orthogonal to W . If W' is nondegenerate, we are done. Otherwise, W' is isotropic too. Then $W \oplus W'$ is isomorphic to $W \times W^{\iota,*}$ equipped with the canonical (ι, ϵ) -symmetric form. Indeed, the H -map $L : W' \rightarrow W^{\iota,*}$, $v' \mapsto (\sharp_B(v'))|_W$, is nonzero, thus an isomorphism, and Lemma 6.14 applies. By assumption, there exists an ϵ -symmetric H -isomorphism $M : W \rightarrow W^{\iota,*}$. Then the graph $Z = \{(w, M(w)) \mid w \in W\}$ of M is nondegenerate. Indeed, for $v, w \in W$,

$$(v, M(v)) \cdot (w, M(w)) = \epsilon \iota(\langle M(v), w \rangle) + \langle M(w), v \rangle = 2\epsilon(\langle M(v), w \rangle)$$

cannot vanish for all w , unless $v = 0$. Thus Z is a nondegenerate irreducible (ι, ϵ) -linear H -submodule of $W \times W^{\iota,*}$, which embeds isometrically into V , so this is the required subspace.

The proof of the proposition is concluded by induction on dimension. If V is irreducible, we are done. Otherwise, we have just shown that V has at least one nondegenerate irreducible submodule, say W . Then the induction hypothesis applies to its orthogonal W^\perp . \square

Corollary 6.18. *Let (V, B) be a nondegenerate bi-isotypical (ι, ϵ) -linear H -module. Then (V, B) is isomorphic either to Example 6.12 or to Example 6.13. The centralizer of H in the automorphism group of (V, B) is isomorphic to a general linear group in the former case, an orthogonal group in the latter.*

6.5. Centers of centralizers in unitary groups

In this case, we directly get information on a real form.

Proposition 6.19. *Let (V, B) be a nondegenerate symmetric sesquilinear H -module. Let \mathfrak{c} denote the center of the centralizer of H in $U(V, B)$. Then root spaces of V under \mathfrak{c} correspond to isotypical components I_π under H . They fall into bi-isotypical components BI_π which are pairwise orthogonal. If $I_\pi = BI_\pi$, then the corresponding root of \mathfrak{c} is purely imaginary. If $I_\pi \neq BI_\pi$, then $BI_\pi = I_\pi \oplus I_{\bar{\pi}^*}$, the corresponding roots ℓ_π and $\ell_{\bar{\pi}^*}$ satisfy $\bar{\ell}_\pi + \ell_{\bar{\pi}^*} = 0$. The linear relations between roots are generated by $\bar{\ell}_\pi + \ell_{\bar{\pi}^*} = 0$, $\pi \neq \bar{\pi}^*$.*

Let N be a set which contains exactly one element of each pair $\{(\pi, \bar{\pi}^)\}$ of equivalence classes of irreducible sesquilinear H -modules occurring in V , such that $\pi \neq \bar{\pi}^*$. Then the map $(\ell_\pi)_{\pi \in N} : \mathfrak{c} \rightarrow \mathbb{C}^N$ is onto.*

Proof. If $I_\pi = BI_\pi$, then, as a sesquilinear H -module, BI_π is isomorphic to Example 6.13 and contributes a unitary factor to the centralizer of H , whose center is a purely imaginary subgroup of $SL(n, \mathbb{C})$. It acts on BI_π by multiplication by a purely imaginary number. Thus BI_π is a root space for a purely imaginary root. Otherwise, BI_π is isomorphic to Example 6.12 and contributes a general linear group factor to the centralizer. Its center is a complex subgroup of $SL(n, \mathbb{C})$, it acts on I_π (resp. $I_{\bar{\pi}^*}$) by multiplication by an unrestricted complex number (resp. the opposite of the conjugate number). This produces a subspace \mathfrak{c}' of \mathfrak{c} which admits a complex structure, the corresponding roots are \mathbb{C} -linear and half of them (to avoid the relations $\bar{\ell}_\pi + \ell_{\bar{\pi}^*} = 0$) provide complex coordinates on \mathfrak{c}' . □

Corollary 6.20. *Let L be the set of roots, and L_I the subset of roots which take only purely imaginary values. Then for every $\ell \in L \setminus L_I$, $-\bar{\ell}$ is again a root. Let L_N be a set which contains exactly one element of each pair $\{\ell, -\bar{\ell}\}$ with $\ell \in L \setminus L_I$. Then*

- I_ℓ and $I_{\ell'}$ are orthogonal unless $\ell' = -\bar{\ell}$;
- if $\ell \in L_I$, then I_ℓ is nondegenerate;
- the map $(\ell)_{\ell \in L_N} : \mathfrak{c} \rightarrow \mathbb{C}^{L_N}$ is onto.

6.6. Centers of centralizers in ϵ -orthogonal groups

We continue our convention that

$$O^\epsilon(n, \mathbb{C}) = \begin{cases} O(n, \mathbb{C}) & \text{if } \epsilon = 1, \\ Sp(n, \mathbb{C}) & \text{if } \epsilon = -1. \end{cases}$$

Note that in the latter case, n has to be even.

Proposition 6.21. *Let (V, B) be a nondegenerate bilinear H -module. Let \mathfrak{c} denote the center of the centralizer of H in $O^\epsilon(V, B)$. Under \mathfrak{c} , V splits into root spaces as follows. Each isotypical component I_π , where π is an irreducible H -module which is not equivalent to its contragredient π^* , is a root space for a nonzero root ℓ_π . It is isotropic. The sum*

of all other isotypical components constitutes the 0 root space. Bi-isotypical components $I_\pi + I_{\pi^*}$ are pairwise orthogonal. Relations among nonzero roots are generated by

$$\ell_\pi + \ell_{\pi^*} = 0.$$

Thus the number of nonzero roots is $2 \dim(\mathfrak{c})$. Let L be a set containing exactly one element of each pair (π, π^*) . Then $(\ell_\pi)_{\pi \in L} : \mathfrak{c} \rightarrow \mathbb{C}^L$ is a linear bijection.

Proof. Let L' be the set of equivalence classes of irreducible H -modules which are isomorphic to their contragredient. Then

$$Z_{O^\epsilon(V, B)}(H) = \prod_{\pi \in L} GL(r_\pi, \mathbb{C}) \times \prod_{\pi \in L'} O^\epsilon(2r_\pi, \mathbb{C}),$$

thus

$$ZZ_{O^\epsilon(V, B)}(H) = \bigoplus_{\pi \in L} \mathbb{C}(\text{id}_{I_\pi} - \text{id}_{I_{\pi^*}}),$$

i.e. $(\ell_\pi)_{\pi \in L} : \mathfrak{c} \rightarrow \mathbb{C}^L$ is a linear bijection. Furthermore, if $\pi \in L$, the bi-isotypical component $I_\pi \oplus I_{\pi^*}$ splits into two root spaces relative to roots ℓ_π and $\ell_{\pi^*} = -\ell_\pi$. \square

Corollary 6.22. *Let \tilde{L} denote the set of roots of \mathfrak{c} on \mathbb{C}^n . The roots of \mathfrak{c} in its adjoint action on $\mathfrak{so}^\epsilon(V, B)$ are exactly all differences $\ell - \ell'$ for $\ell, \ell' \in \tilde{L}$, including 2ℓ if $\epsilon = -1$ or $\dim(I_\ell) > 1$, and excluding 2ℓ if I_ℓ is 1-dimensional and $\epsilon = 1$.*

In other words, if L is a set of representatives of pairs $\{-\ell, \ell\}$ of nonzero roots, the roots of \mathfrak{c} in the adjoint representation include 0, all sums $\pm\ell \pm \ell'$ for distinct $\ell, \ell' \in L$, sometimes $\pm 2\ell$ (depending on ϵ and $\dim(I_\ell)$), and, if 0 is also a root of \mathfrak{c} on \mathbb{C}^n , all $\pm\ell$ for $\ell \in L$.

Proof. The Lie algebra $\mathfrak{so}(V, B)$ is the space of B -skew-symmetric endomorphisms of V , i.e. \mathbb{C} -linear maps $f : V \rightarrow V$ satisfying, for all $v, v' \in V$,

$$B(f(v), v') + B(v, f(v')) = 0.$$

The roots of \mathfrak{c} in the adjoint representation are differences $\lambda = \ell - \ell'$ of roots of \mathfrak{c} in \mathbb{C}^n . If $\ell' \neq \pm\ell$, the root space relative to $\ell - \ell'$ is the subspace of B -skew-symmetric elements of $\text{Hom}(I_\ell, I_{\ell'}) \oplus \text{Hom}(I_{-\ell'}, I_{-\ell})$. It never vanishes. Indeed, for every $f \in \text{Hom}(I_\ell, I_{\ell'})$, there is a unique $g \in \text{Hom}(I_{-\ell'}, I_{-\ell})$ such that (f, g) is B -skew-symmetric. The formula for g is

$$g = -\sharp_B^{-1} \circ f^\top \circ \sharp_B.$$

Here is an alternative description of $\mathfrak{so}(V, B)$: mapping $f \in \text{End}(V)$ to the bilinear form $b(v, v') = B(f(v), v')$ identifies $\text{End}(V)$ with the space $V^* \otimes V^*$ of \mathbb{C} -bilinear forms on V and $\mathfrak{so}(V, B)$ with the subspace $\Lambda^\epsilon V^*$ of $(-\epsilon)$ -symmetric \mathbb{C} -bilinear forms on V . The adjoint action of $Z \in \mathfrak{so}(V, B)$ is

$$Zb(v, v') = B([Z, f](v), v') = -b(v, Zv') - b(Zv, v').$$

If $v \in I_\ell, v' \in I_{\ell'}$ and $Z \in \mathfrak{c}$, then

$$Zb(v, v') = -(\ell + \ell')(Z)b(v, v').$$

Therefore the root space relative to 2ℓ identifies with $\Lambda^\epsilon I_\ell^*$. It vanishes if and only if $\epsilon = 1$ and $\dim(I_\ell) = 1$. \square

7. Real forms of $SL(n, \mathbb{C})$

7.1. Flexibility in $SL(n, \mathbb{R})$ and $SL(n, \mathbb{H})$

Proposition 7.1. *Let $\mathfrak{c} \subset \mathfrak{sl}(n, \mathbb{R})$ (resp. $\mathfrak{sl}(n/2, \mathbb{H})$) be the center of the centralizer of a reductive subgroup of $SL(n, \mathbb{R})$ (resp. $SL(n/2, \mathbb{H})$). Let λ be a purely imaginary root of \mathfrak{c} in its adjoint action on $\mathfrak{sl}(n, \mathbb{C})$. Then the signature of the Killing form restricted to \mathfrak{g}_λ does not vanish. It follows that \mathfrak{c} is balanced.*

Proof. Let \mathbb{C}^n denote the standard representation of $\mathfrak{sl}(n, \mathbb{C})$. As in Subsection 4.2, let $\tau(v) = \bar{v}$ in the complex case, and $\tau(v) = vj$ in the quaternionic case (here, $\mathbb{C}^n = \mathbb{H}^{n/2}$ is viewed as a right quaternionic vector space). For $f \in \mathfrak{sl}(n, \mathbb{C})$ let $\sigma(f) = \tau \circ f \circ \tau^{-1}$. Then $\text{Fix}(\sigma) = \mathfrak{g} = \mathfrak{sl}(n, \mathbb{R})$ (resp. $\mathfrak{sl}(n/2, \mathbb{H})$).

Under \mathfrak{c} , \mathbb{C}^n splits into root spaces $\mathbb{C}^n = \bigoplus_\ell I_\ell$, $\dim(I_\ell) = d_\ell$. Roots are either real or come in pairs $\{\ell, \bar{\ell}\}$ (Lemma 4.1). According to Lemma 5.5, every nonzero root λ of \mathfrak{c} in its adjoint action on $\mathfrak{sl}(n, \mathbb{C})$ can be uniquely written in the form $\ell - \ell'$. Such a root is purely imaginary if and only if $\ell' = \bar{\ell}$, i.e. $\lambda = \ell - \bar{\ell}$. The corresponding root space is

$$\mathfrak{g}_\lambda = \text{Hom}(I_\ell, I_{\bar{\ell}}).$$

Let $f \in \text{Hom}(I_\ell, I_{\bar{\ell}})$, $\sigma(f) = \bar{f} \in \text{Hom}(I_{\bar{\ell}}, I_\ell)$. Pick a basis of I_ℓ and take its image under τ as a basis of $I_{\bar{\ell}}$. Let M denote the matrix of f in the chosen basis of \mathbb{C}^n . Then the matrix of $\sigma(f)$ is $\sigma(M)$, and $\text{Trace}(\sigma(f) \circ f) = \text{Trace}(\bar{M}M)$. Write $M = S + A$ where S is symmetric and A is skew-symmetric. Then

$$\text{Trace}(\bar{M}M) = \text{Trace}(\bar{S}S) + \text{Trace}(\bar{A}A) = \text{Trace}(S^*S) - \text{Trace}(A^*A)$$

has signature $\dim(\{S\}) - \dim(\{A\}) = d_\ell(d_\ell + 1)/2 - d_\ell(d_\ell - 1)/2 = d_\ell$, which is nonzero. □

7.2. Flexibility in $SU(p, q)$

Proposition 7.2. *Let Γ be a surface group, let $\phi : \Gamma \rightarrow SU(p, q)$ a reductive homomorphism, and let \mathfrak{c} be the center of the centralizer of $\phi(\Gamma)$. Assume that \mathfrak{c} is not balanced with respect to ϕ . Then, up to conjugacy, $\phi(\Gamma)$ is contained in $S(U(p, p) \times U(q - p))$, and ϕ is maximal.*

Proof. Under \mathfrak{c} , the standard representation of $\mathfrak{sl}(n, \mathbb{C})$ splits into root spaces, $\mathbb{C}^n = \bigoplus_\ell I_\ell$. The roots of \mathfrak{c} in the adjoint representation are differences $\ell - \ell'$, and all of them indeed occur. Following Corollary 6.20, split the set of roots as $\tilde{L} = L_I \cup L_N \cup -L_N$, where L_I is the subset of purely imaginary roots. If $\ell \in L_N$, then 2ℓ is a root in the adjoint representation, and it is not purely imaginary. If $\ell \in L_I$ and $\ell' \in L_N$, then $\ell - \ell'$ is not purely imaginary. If ℓ and $\ell' \in L_N$, then $\ell - \ell'$ is not purely imaginary either, since it factors through a surjective map $\mathfrak{c} \rightarrow \mathbb{C}^{L_N}$ and a \mathbb{C} -linear form $\mathbb{C}^{L_N} \rightarrow \mathbb{C}$. So if L_N is nonempty, then the non-purely imaginary roots span \mathfrak{c}^* , and \mathfrak{c} is balanced, a contradiction. So L_N is empty, all roots on \mathbb{C}^n are purely imaginary, and the corresponding root spaces

I_ℓ are nondegenerate and pairwise orthogonal. In what follows, we shall replace roots by their imaginary parts without expressing it in the notation.

Each root λ in the adjoint representation can be expressed in a unique way as $\lambda = \ell - \ell'$, and

$$\mathfrak{g}_\lambda = \text{Hom}(I_\ell, I_{\ell'}).$$

A calculation shows that the signature of the natural Hermitian form on $\text{Hom}(I_\ell, I_{\ell'})$ equals $-\text{sign}(I_\ell) \text{sign}(I_{\ell'})$. This time, the signature is not automatically nonzero. So different arguments, based on [4], are needed.

Assume that the sesquilinear action of Γ on \mathfrak{g}_λ is maximal. The situation we are considering is as follows: since $H = \overline{\phi}(\Gamma)$ preserves I_ℓ and preserves a sesquilinear form on it, we have $\phi(\Gamma) \subset U(I_\ell)$, and similarly for $I_{\ell'}$, hence $\phi(\Gamma) \rightarrow U(I_\ell) \times U(I_{\ell'})$ and we obtain

$$\begin{array}{ccc} \phi(\Gamma) \subset Z_G(\mathfrak{c}) & \xrightarrow{\quad} & U(I_\ell) \times U(I_{\ell'}) \\ & \searrow f & \downarrow \\ & & U(\mathfrak{g}_\lambda) = U(\text{Hom}(I_\ell, I_{\ell'})) = U(I_\ell \otimes I_{\ell'}^*) \end{array}$$

where f is tight. According to Proposition 3.16, the sesquilinear space $\text{Hom}(I_\ell, I_{\ell'})$ must have vanishing signature. Also, the homomorphism $Z_G(\mathfrak{c}) \rightarrow U(\text{Hom}(I_\ell, I_{\ell'}))$ must be tight. Lemma 3.14 implies that one of I_ℓ and $I_{\ell'}$ has vanishing signature and the other one is definite. Say I_ℓ is definite, for instance. According to Lemma 3.8, maximality of $\text{Hom}(I_\ell, I_{\ell'})$ implies maximality of the Γ action on $I_{\ell'}$, with $T_\lambda = \dim(I_\ell)T(I_{\ell'})$. If instead $I_{\ell'}$ is definite, then $T_\lambda = -T(I_\ell) \dim(I_{\ell'})$.

Let D (resp. E , resp. O) denote the set of roots ℓ such that I_ℓ is definite (resp. has vanishing signature, resp. has nonvanishing signature). As in Definition 2.1, let P denote the set of roots λ such that ρ_λ is maximal with positive Toledo invariant, or equivalently, of differences $\ell - \ell'$ with $\ell \in D$ and $\ell' \in E$ such that $I_{\ell'}$ is maximal with positive Toledo invariant. Let N be the complement of $\pm P$.

Let us show that if O is nonempty, then \mathfrak{c} is balanced with respect to ϕ . Indeed, let $\ell_0 \in O$. Then for all $\ell' \neq \ell_0$, if we set $\lambda = \ell_0 - \ell'$, then ρ_λ is not maximal, thus $\ell_0 - \ell' \in N$. Since the roots ℓ span $\mathfrak{c}^* \otimes \mathbb{C}$ and satisfy the extra equation $\sum_\ell d_\ell \ell = 0$, it follows that $\text{span}_{\mathbb{C}}(\{\ell_0 - \ell' \mid \ell' \neq \ell_0\}) = \mathfrak{c}^* \otimes \mathbb{C}$. Since $\text{span}_{\mathbb{C}}(N) = \mathfrak{c}^* \otimes \mathbb{C}$, \mathfrak{c} is balanced with respect to ϕ .

From now on, we assume that O is empty. If D or E is empty, there is no room for pairs (ℓ, ℓ') for $\ell \in D$ and $\ell' \in E$, so P is empty, hence balanced. Therefore we assume that both D and E are nonempty. Let L_D (resp. L_E) denote the span of all differences $\ell - \ell'$ for $\ell \in D$ and $\ell' \in D$ (resp. for $\ell \in E$ and $\ell' \in E$). Then $\dim(L_D) = \text{card}(D) - 1$ and $\dim(L_E) = \text{card}(E) - 1$ by Lemma 5.5. Since $L_D \subset \text{span}_{\mathbb{C}}(D)$, $L_E \subset \text{span}_{\mathbb{C}}(E)$ and $\text{span}_{\mathbb{C}}(D) \cap \text{span}_{\mathbb{C}}(E)$ is the line generated by $\sum_{\ell \in D} d_\ell \ell$, we have $L_D \cap L_E = \{0\}$, thus $\dim(L_D + L_E) = \text{card}(D) + \text{card}(E) - 2 = \dim(\mathfrak{c}) - 1$. In the quotient space $\mathfrak{c}^* \otimes \mathbb{C}/L_D + L_E$, all elements of D (resp. of E) are mapped to the same vector ℓ_D (resp. ℓ_E), and $\ell_D \neq \ell_E$. Again, if one of the $\ell - \ell'$, $\ell \in D$, $\ell' \in E$, belongs to N , then $\mathfrak{c}^* \otimes \mathbb{C}/N$ vanishes, so \mathfrak{c} is balanced with respect to ϕ , a contradiction.

Therefore, all $\ell - \ell'$ with $\ell \in D$ and $\ell' \in E$ belong to $\pm P$. If there exist two pairs (ℓ, ℓ') , $\ell \in D$, $\ell' \in E$, such that $\mathfrak{S}m(\ell - \ell')$ have opposite signs in $\mathfrak{c}^*/\mathfrak{S}m(L_D + L_E)$, then \mathfrak{c} is balanced with respect to ϕ , a contradiction.

Otherwise, all $\ell - \ell'$, $\ell \in D$, $\ell' \in E$, belong to $\pm P$ and those which belong to $+P$ project to $\mathfrak{c}^*/\mathfrak{S}m(L_D + L_E)$ with equal signs. This implies that the direct sum representation $\bigoplus_{\ell \in E} I_\ell$ is maximal. In other words, $\phi(\Gamma) \subset U(p, p) \times U(q - p)$ is maximal. The symmetric spaces $\mathcal{D}_{p,p}$ and $\mathcal{D}_{p,q}$ corresponding to $SU(p, p)$ and $G = SU(p, q)$ have equal ranks and the embedding $\mathcal{D}_{p,p} \hookrightarrow \mathcal{D}_{p,q}$ is isometric and holomorphic. Therefore Example 3.3 implies that, viewed as a homomorphism $\Gamma \rightarrow G$, ϕ is maximal as well. \square

7.3. Rigidity in $SU(p, q)$

The centralizer of $SU(p, p)$ in $SU(p, q)$ with $q > p$ is $U(q - p)$ with center $\mathfrak{c} = \mathfrak{u}(1)$ generated by

$$Z = \begin{pmatrix} -\frac{2pi}{p+q} I_{q-p} & 0 \\ 0 & \frac{(q-p)i}{p+q} I_{2p} \end{pmatrix}.$$

There is only one nonzero pair of roots $\pm i$, giving rise to the root space $\text{Hom}_{\mathbb{C}}(\mathbb{C}^{q-p}, \mathbb{C}^{2p})$. The sesquilinear form s_i on $\text{Hom}_{\mathbb{C}}(\mathbb{C}^{q-p}, \mathbb{C}^{2p})$ is, in an $SU(p, p)$ -invariant manner, the direct sum of $q - p$ copies of the $U(p, p)$ -invariant Hermitian form on \mathbb{C}^{2p} . Therefore the corresponding Toledo invariant is equal to $q - p$ times the Toledo invariant obtained for $q = p + 1$. In this case, the centralizer of \mathfrak{c} is $U(1) \times SU(p, p)$, acting on the real root space via the standard complex representation of its second factor. The representation ρ_i induced on a surface subgroup $\Gamma \subset U(1) \times SU(p, p)$ is maximal if and only if the projection of Γ to $SU(p, p)$ is maximal. Thus a maximal surface subgroup of $SU(p, p)$ is not flexible in $SU(p, q)$ with $q > p$, as is well known. Such subgroups exist (Theorem 1.8) and are known to be automatically discrete [1], [3].

8. Real forms of $O(n, \mathbb{C})$ and $Sp(n, \mathbb{C})$

8.1. Non-purely imaginary roots

Lemma 8.1. *Let G be a real form of $O^\epsilon(n, \mathbb{C})$. Let $H \subset G$ be a reductive subgroup. Let $\mathfrak{c} \subset \mathfrak{g}$ be the center of its centralizer. If one of the roots of \mathfrak{c} on \mathbb{C}^n is not purely imaginary, then \mathfrak{c} is balanced.*

Proof. Assume that \mathfrak{c} is not balanced and at least one of the roots of \mathfrak{c} on \mathbb{C}^n is not purely imaginary. Consider the torus $\mathfrak{c} \otimes \mathbb{C} \subset \mathfrak{so}^\epsilon(n, \mathbb{C})$ and the set $L \cup -L$ of its nonzero roots on \mathbb{C}^n (Proposition 6.21). Write $L = L_I \cup L_N$ where $L_I \subset L$ denotes the set of roots whose restriction to \mathfrak{c} takes only purely imaginary values, and L_N its complement. By assumption, L_N is nonempty.

Assume first that L_I is nonempty too. None of the roots $\lambda = \pm \ell \pm \ell'$ with $\ell \in L_I$ and $\ell' \in L_N$ is purely imaginary, so none of them belongs to $\pm P$. They generate $\text{span}(L) = \mathfrak{c}^*$, thus \mathfrak{c} is balanced, a contradiction. Therefore, $L = L_N$.

If every root of the adjoint action is non-purely imaginary, it is balanced. Hence they do not span \mathfrak{c}^* , which implies that there exists $v \in \mathfrak{c}$ such that $\lambda(v) = 0$ for all roots λ of \mathfrak{c} in the adjoint representation which are not purely imaginary. For any distinct $\ell, \ell' \in L$, one of $\ell - \ell'$ and $\ell + \ell'$ is not purely imaginary, thus $\ell'(v) = \pm\ell(v)$.

Assume that there exists $\ell \in L$ such that ℓ or 2ℓ is a root of the adjoint action; then $\ell(v) = 0$ as well, which implies that $\ell'(v) = 0$ for all $\ell' \in L$, and $v = 0$. In other words, in that case, non-purely imaginary roots span \mathfrak{c}^* , and \mathfrak{c} is balanced, a contradiction. Therefore,

- 0 is not a root of \mathfrak{c} on \mathbb{C}^n (i.e., all root spaces correspond to elements of $\pm L$);
- for every root $\ell \in L$, 2ℓ is not a root of \mathfrak{c} in the adjoint representation.

This implies that $\epsilon = 1$ and all root spaces I_ℓ have dimension 1, i.e. $\mathfrak{c} \otimes \mathbb{C}$ is a maximal torus of $\mathfrak{so}(n, \mathbb{C})$. Its centralizer in $\mathfrak{so}(n, \mathbb{C})$ is abelian, and so are $Z_G(\mathfrak{c})$ and H , up to finite index. A homomorphism from an abelian group cannot be tight, so no symplectic action on root spaces can be maximal, P is empty, and \mathfrak{c} is balanced again, a contradiction. □

8.2. The sesquilinear structure

Proposition 8.2. *Let G be a real form of $O^\epsilon(n, \mathbb{C})$. Let $H \subset G$ be a reductive subgroup. Let $\mathfrak{c} \subset \mathfrak{g}$ be the center of its centralizer.*

1. *Let ℓ be a purely imaginary root of \mathfrak{c} on \mathbb{C}^n . Let I_ℓ denote the corresponding root space. Then I_ℓ inherits a $Z_G(\mathfrak{c})$ -invariant nondegenerate sesquilinear form s_ℓ .*
2. *Assume that all roots of \mathfrak{c} on \mathbb{C}^n are purely imaginary. Then the centralizer $Z_G(\mathfrak{c})$ is a product of the fixator of the orthogonal of the 0 root space I_0 and of unitary groups,*

$$Z_G(\mathfrak{c}) \cong G_{I_0^\perp} \times \prod_{\ell \in L} U(I_\ell, s_\ell).$$

Proof. 1. Let $s_\ell(v, v') = B(\tau(v), v')$. This is an $\eta\epsilon$ -symmetric sesquilinear form. It is nondegenerate on I_ℓ because B is and all root spaces but $I_{-\ell} = \tau(I_\ell)$ are B -orthogonal to I_ℓ . If $g \in Z_G(\mathfrak{c})$, i.e. g belongs to G and commutes with \mathfrak{c} , then g leaves all root spaces $I_{\pm\ell}$ invariant, commutes with τ , and is isometric for B , thus it is isometric for s_ℓ .

2. Let $g \in G$ commute with \mathfrak{c} . Let $g_0 \in GL(n, \mathbb{C})$ be the element which coincides with g on I_0 and fixes the sum of all I_ℓ with $\ell \in \pm L$, i.e. I_0^\perp . Then $g_0 \in G$. Indeed, g_0 commutes with τ and preserves B . Also, restrict g to each I_ℓ with $\ell \in L$. This yields an injective homomorphism $Z_G(\mathfrak{c}) \rightarrow G_{I_0^\perp} \times \prod_{\ell \in L} U(I_\ell, s_\ell)$.

Conversely, let $g_0 \in G$ fix all I_ℓ . For each $\ell \in L$, pick $g_\ell \in U(I_\ell, s_\ell)$, extend it to $I_{-\ell}$ so that $g_\ell = \tau \circ g_\ell \circ \tau^{-1}$, extend this map trivially to other $I_{\ell'}$ and to I_0 . The resulting linear map belongs to $Z_G(\mathfrak{c})$. Indeed, it preserves B , commutes with τ and preserves each I_ℓ . Multiplying g_0 with the g_ℓ 's yields the inverse isomorphism $G_{I_0^\perp} \times \prod_{\ell \in L} U(I_\ell, s_\ell) \rightarrow Z_G(\mathfrak{c})$. □

Lemma 8.3. *Let G be a real form of $O^\epsilon(n, \mathbb{C})$. Let $H \subset G$ be a reductive subgroup. Let $\mathfrak{c} \subset \mathfrak{g}$ be the center of its centralizer. Let ℓ, ℓ' be distinct purely imaginary roots of \mathfrak{c}*

on \mathbb{C}^n , and $\lambda = \ell - \ell'$ the corresponding purely imaginary root of \mathfrak{c} in its adjoint action on \mathfrak{g} . Let $I_\ell, I_{\ell'}$ and \mathfrak{g}_λ denote the corresponding root spaces.

1. If $\ell' \neq -\ell$, then, as a sesquilinear $Z_G(\mathfrak{c})$ -module, \mathfrak{g}_λ is isomorphic to $\text{Hom}(I_\ell, I_{\ell'})$ equipped with the natural sesquilinear form

$$(f, f') \mapsto \text{Trace}(f^* \circ f'),$$

where $f^* \in \text{Hom}(I_{\ell'}, I_\ell)$ is the adjoint of f with respect to the sesquilinear forms s_ℓ and $s_{\ell'}$.

2. If $\ell' = -\ell$, then, as a sesquilinear $Z_G(\mathfrak{c})$ -module, \mathfrak{g}_λ is isomorphic to the subspace of $(-\epsilon)$ -symmetric forms in the space $I_\ell^* \otimes I_\ell^*$ of \mathbb{C} -bilinear forms on I_ℓ , equipped with its natural sesquilinear form

$$(b, b') \mapsto \text{Trace}((\sharp_{s_\ell})^{-1} \circ (\overline{\sharp_b})^\top \circ (\sharp_{s_\ell})^{-1} \circ \sharp_{b'}).$$

Proof. 1. Assume first that ℓ and ℓ' are linearly independent. According to Corollary 6.22, \mathfrak{g}_λ is the space of B -skew-symmetric elements of $\text{Hom}(I_\ell, I_{\ell'}) \oplus \text{Hom}(I_{-\ell'}, I_{-\ell})$. Given $f \in \text{Hom}(I_\ell, I_{\ell'})$ and $g \in \text{Hom}(I_{-\ell'}, I_{-\ell})$, $X = (f, g)$ being B -skew-symmetric means that for all $v \in I_\ell$ and $w \in I_{-\ell'}$,

$$B(f(v), w) + B(v, g(w)) = 0.$$

Since \sharp_B identifies $I_{-\ell}$ with the dual of I_ℓ , given $f \in \text{Hom}(I_\ell, I_{\ell'})$, there exists a unique $g \in \text{Hom}(I_{-\ell'}, I_{-\ell})$ such that $X = (f, g)$ is B -skew-symmetric. This shows that, as a $Z_G(\mathfrak{c})$ -module, $\mathfrak{g}_{\ell-\ell'}$ is isomorphic to $\text{Hom}(I_\ell, I_{\ell'})$.

Since $\sigma(X) = (\tau \circ g \circ \tau^{-1}, \tau \circ f \circ \tau^{-1})$, we have

$$\sigma(X) \circ X = \eta(\tau \circ g \circ \tau \circ f, \tau \circ f \circ \tau \circ g),$$

and therefore

$$\begin{aligned} \text{Trace}(\sigma(X) \circ X) &= \eta \text{Trace}(\tau \circ g \circ \tau \circ f) + \eta \text{Trace}(\tau \circ f \circ \tau \circ g) \\ &= 2\eta \text{Trace}(\tau \circ g \circ \tau \circ f). \end{aligned}$$

If $v \in I_\ell$, then

$$\begin{aligned} s_\ell(v, \tau \circ g \circ \tau \circ f(v)) &= B(\tau(v), \tau \circ g \circ \tau \circ f(v)) = \overline{B(v, g \circ \tau \circ f(v))} \\ &= -\overline{B(f(v), \tau \circ f(v))} = -\epsilon \overline{B(\tau \circ f(v), f(v))} \\ &= -\epsilon \overline{s_\ell(f(v), f(v))} = -\epsilon \overline{s_\ell(v, f^* \circ f(v))}. \end{aligned}$$

Summing over an orthogonal basis for s_ℓ yields

$$\text{Trace}(\tau \circ g \circ \tau \circ f) = -\epsilon \overline{\text{Trace}(f^* \circ f)},$$

hence the Killing sesquilinear form on \mathfrak{g}_λ is proportional to $\text{Trace}(f^* \circ f)$.

2. Let $f \in \text{Hom}(I_\ell, I_{-\ell})$. Then f belongs to $\mathfrak{so}(V, B)$ (f is B -skew-symmetric) if and only if $f^* = -\sigma(f)$. Indeed,

$$\begin{aligned} \forall v, w \in I_\ell, B(f(v), w) + B(v, f(w)) &= 0 \\ \Leftrightarrow \forall v, w \in I_\ell, s_{-\ell}(f(v), \tau(w)) + s_\ell(v, \tau \circ f(w)) &= 0 \\ \Leftrightarrow \forall v \in I_\ell, \forall v' \in I_{-\ell}, s_{-\ell}(f(v), v') + s_\ell(v, \tau \circ f \circ \tau^{-1}(v')) &= 0 \\ \Leftrightarrow f^* = -\tau \circ f \circ \tau^{-1}. \end{aligned}$$

Thus, for B -skew-symmetric f ,

$$\text{Trace}(\sigma(f) \circ f) = \text{Trace}(\tau \circ f \circ \tau^{-1} \circ f) = -\text{Trace}(f^* \circ f).$$

Now f is B -skew-symmetric if and only if the bilinear form $b(v, v') = B(f(v), v')$ on I_ℓ is $(-\epsilon)$ -symmetric. Since $\sharp_B = f^\top \circ \sharp_B = \eta f^\top \circ \tau^\top \circ \sharp_{s_\ell}$ and $f^* = (\sharp_{s_\ell})^{-1} \circ (\overline{f})^\top \circ \sharp_{s_\ell}$, we have

$$f^* \circ f' = \eta(\sharp_{s_\ell})^{-1} \circ (\overline{b})^\top \circ (\sharp_{s_\ell})^{-1} \circ \sharp_{b'}. \quad \square$$

Lemma 8.4. *Let V be an n -dimensional complex vector space equipped with a nondegenerate sesquilinear form of signature s . Then the signatures of the induced sesquilinear forms on $V^* \otimes V^*$, $S^2 V^*$ and $\Lambda^2 V^*$ are equal to s^2 , $(s^2 + n)/2$ and $(s^2 - n)/2$ respectively.*

Proof. Fix a basis of V . If D denotes the matrix of the sesquilinear form S in this basis, and b, b' the matrices of two bilinear forms on V , the induced sesquilinear form on $V^* \otimes V^*$ is

$$\text{Trace}(\overline{b}^\top D^{-1} b' D^{-1}).$$

One can assume that S is diagonal with entries $d_m = 1$ (p times) and -1 (q times), $n = p + q, s = p - q$. Then

$$\text{Trace}(\overline{b}^\top D^{-1} b' D^{-1}) = \sum_{m, m'} d_m d_{m'} |b_{mm'}|^2$$

is diagonal again. The signature of the whole space $V^* \otimes V^*$ is $\sum_{m, m'} d_m d_{m'} = s^2$. The signature of the subspace $\Lambda^2 V^*$ of skew-symmetric forms is

$$\sum_{m < m'} d_m d_{m'} = \frac{p(p-1)}{2} + \frac{q(q-1)}{2} - pq = \frac{s^2 - n}{2}.$$

The signature of the subspace $S^2 V^*$ of symmetric forms is

$$\sum_{m \leq m'} d_m d_{m'} = n + \sum_{m < m'} d_m d_{m'} = \frac{s^2 + n}{2}. \quad \square$$

8.3. Unbalanced centers of centralizers

Recall that when $\mathfrak{c} \subset \mathfrak{g}$ is the center of the centralizer of a homomorphism of a surface group to G , P denotes the set of roots λ in the adjoint representation such that the sesquilinear action on the root space \mathfrak{g}_λ is maximal with positive Toledo invariant.

Lemma 8.5. *Let G be a real form of $O^\epsilon(n, \mathbb{C})$. Let $\phi : \Gamma \rightarrow G$ be a reductive homomorphism. Let $\mathfrak{c} \subset \mathfrak{g}$ be the center of its centralizer. Assume all the roots are purely imaginary. Let ℓ, ℓ' be distinct nonzero purely imaginary roots of \mathfrak{c} on \mathbb{C}^n , and $\lambda = \ell - \ell'$ the corresponding purely imaginary root of \mathfrak{c} in its adjoint action on \mathfrak{g} .*

1. *If $\ell' \neq -\ell$ and $\lambda \in \pm P$, then one of s_ℓ and $s_{\ell'}$ has vanishing signature and the other is definite.*
2. *If $\ell' = -\ell$, then $\lambda = 2\ell$ either is not a root or does not belong to $\pm P$.*

Proof. If $\lambda \in P$, then \mathfrak{g}_λ has vanishing signature. So does $U(\text{Hom}(I_\ell, I_{\ell'}))$ (resp. $U(\Lambda^\epsilon(I_\ell))$) with its natural sesquilinear form, according to Lemma 8.3. In particular, these groups are of tube Hermitian type. Furthermore, the $Z_G(\mathfrak{c})$ action on \mathfrak{g}_λ is tight. According to Proposition 8.2, $Z_G(\mathfrak{c})$ is a product of groups.

1. If $\ell' \neq -\ell$, then among the factors, only $U(I_\ell, s_\ell)$ and $U(I_{\ell'}, s_{\ell'})$ act nontrivially on \mathfrak{g}_λ , thus the morphism

$$U(I_\ell) \times U(I_{\ell'}) \rightarrow U(\text{Hom}(I_\ell, I_{\ell'}))$$

must be tight. Lemma 3.14 applies and one of the left hand groups is compact and the other has vanishing signature.

2. If $\ell' = -\ell$, then among the factors, only $U(I_\ell, s_\ell)$ acts nontrivially on \mathfrak{g}_λ , thus the morphism

$$U(I_\ell) \rightarrow U(\Lambda^\epsilon(I_\ell))$$

must be tight. Lemma 3.13 implies that $U(I_\ell)$ is of tube type, so s_ℓ has vanishing signature. Lemma 8.4 shows that the signature of $\Lambda^\epsilon(I_\ell)$ is $\pm \dim(I_\ell)/2$, which does not vanish, a contradiction. We conclude that if 2ℓ is a root, it does not belong to $\pm P$. \square

Proposition 8.6. *Let G be a real form of $O^\epsilon(n, \mathbb{C})$. Let $\phi : \Gamma \rightarrow G$ be a reductive homomorphism. Let $\mathfrak{c} \subset \mathfrak{g}$ be the center of its centralizer. Then \mathfrak{c} is balanced with respect to ϕ unless $\epsilon = 1$, $\dim(\mathfrak{c}) = 1$, \mathfrak{c}^* is generated by a root ℓ with a 1-dimensional root space I_ℓ , the sesquilinear form on I_0 has vanishing signature, $G_{I_0^\perp}$ is reductive Hermitian of tube type and tightly embedded in $U(I_0)$, and the homomorphism $\Gamma \rightarrow U(I_0)$ is maximal.*

Proof. Assume that \mathfrak{c} is not balanced. From Lemma 8.1, we know that roots have to be purely imaginary. We take their imaginary parts without explicit mention.

If $\epsilon = -1$ or if all $I_\ell, \ell \in L$, have dimension > 1 , then all 2ℓ are roots and do not belong to $\pm P$, so they all belong to the complement N of P (Lemma 8.5). Since they generate \mathfrak{c}^* , \mathfrak{c} is balanced, a contradiction. So $\epsilon = 1$ and the set D of roots $\ell \in L$ such that $\dim(I_\ell) = 1$ is nonempty.

Assume that D has at least two elements. According to Lemma 8.5, all combinations $\pm\ell \pm \ell'$ for $\ell, \ell' \in D$ belong to N . Since they span $\text{span}(D)$, and the multiples $2\ell''$ with

$\ell'' \notin D$ span $\text{span}(L \setminus D)$, it follows that N spans \mathfrak{c}^* and \mathfrak{c} is balanced, a contradiction. So D has exactly one element, denoted by ℓ_0 .

Assume that $L \neq D$. If $\ell \notin D$, then $2\ell \notin \pm P$, hence $2\ell \in N$. Since $\ell_0 \cup \{\ell \notin D\}$ are all roots, $\dim(\mathfrak{c}^*/\text{span}(N)) = 1$, and for all $\ell \in L \setminus D$, all combinations $\pm\ell_0 \pm \ell$ belong to $\pm P$. Since 0 does not belong to the convex hull of the image of P in $\mathfrak{c}^*/\text{span}(N)$, this convex hull contains exactly one of ℓ_0 and $-\ell_0$, say ℓ_0 . Then $\ell_0 - \ell$ and $\ell_0 + \ell$ belong to P . Thus the sesquilinear representation of Γ in $\text{Hom}(I_{\ell_0}, I_\ell \oplus I_{-\ell})$ is maximal. Lemma 3.6 allows us to replace $\text{Hom}(I_{\ell_0}, I_\ell \oplus I_{-\ell})$ with $I_\ell \oplus I_{-\ell}$. However, as a sesquilinear vector space, $I_{-\ell}$ is isomorphic to I_ℓ , so, by Lemma 3.7,

$$T(I_\ell \oplus I_{-\ell}) = T(I_\ell) + T(I_{-\ell}) = T(I_\ell) - T(I_\ell) = 0,$$

contradicting maximality. So $L = D$ and $\dim(\mathfrak{c}) = 1$.

Assume that 0 is not a root of \mathfrak{c} on \mathbb{C}^n . Then $\dim(V = I_{\ell_0} \oplus I_{-\ell_0}) = 2$ and $Z_G(\mathfrak{c}) = U(I_{\ell_0}) \cong U(1)$, which cannot have any maximality property. So 0 is a root on \mathbb{C}^n and $\pm\ell_0$ are roots of \mathfrak{c} in the adjoint representation. Since 0 does not belong to the convex hull of P in \mathfrak{c}^* , exactly one of $\pm\ell_0$ belongs to P , say $\ell_0 \in P$. Then $\text{Hom}(I_{\ell_0}, I_0)$ is a maximal sesquilinear representation of Γ , and so is I_0 , by Lemma 3.6. In particular, the signature of (I_0, s_0) vanishes, i.e. $U(I_0)$ has tube type. Also, the morphism $Z_G(\mathfrak{c}) \rightarrow PU(I_0)$ is tight. $Z_G(\mathfrak{c}) = G_{I_0^\perp} \times U(I_{\ell_0})$ acts on I_0 via the group $G_{I_0^\perp}$, so the injection $G_{I_0^\perp} \rightarrow PU(I_0)$ is tight. This implies that $G_{I_0^\perp}$ is reductive Hermitian and has tube type (Lemma 3.13). \square

Corollary 8.7. *Let G be a real form of $O^\varepsilon(n, \mathbb{C})$. Let $\phi : \Gamma \rightarrow G$ be a reductive homomorphism. Let $\mathfrak{c} \subset \mathfrak{g}$ be the center of its centralizer. Then \mathfrak{c} is balanced with respect to ϕ unless $G = SO^*(2n)$, n odd, $\phi(\Gamma) \subset SO^*(2n-2) \times SO^*(2)$ and ϕ is maximal.*

Proof. In view of Proposition 8.6, it merely remains to determine which pairs $(G, G_{I_0^\perp})$ can lead to unbalanced centralizers when G is a real form of $O(N, \mathbb{C})$, i.e. $G = O(p, q)$ or $G = SO^*(2n)$.

In the real case, I_0 is real, and $G_{I_0^\perp}$ is a real orthogonal group $O(p, q)$, which tightly injects into $U(I_0, s_0) = U(p, q)$. We also know that the signature $p - q$ of s_0 vanishes. $O(p, p)$ is reductive Hermitian only if $p = 2$. But Lemma 3.15 states that the inclusion $O(2, 2) \hookrightarrow U(2, 2)$ is not tight. So \mathfrak{c} is always balanced if $G = O(p, q)$ is a real orthogonal group.

In the quaternionic case, I_0 is quaternionic and carries a nondegenerate $(\bar{\cdot}, -1)$ -binary form (Lemma 4.1) (see also Subsection 4.2). Therefore $G_{I_0^\perp}$ is a quaternionic skew-unitary group $SO^*(2p)$ with $p = \dim_{\mathbb{H}}(I_0)$. Let ℓ denote the unique nonzero root of \mathfrak{c} on \mathbb{C}^{2n} . Then $I_0^\perp = I_\ell \oplus I_{-\ell}$ is a 2-dimensional complex vector space, thus a 1-dimensional quaternionic vector space, therefore $n = p + 1$. Since $G_{I_0^\perp}$ has tube type, p is even and n is odd. The homomorphism $\Gamma \rightarrow U(I_0) \cong U(p, p)$ is maximal. According to Lemma 10.3, the homomorphism $\Gamma \rightarrow G_{I_0^\perp} \cong SO^*(2p)$ is maximal as well. The symmetric spaces \mathcal{G}_p and \mathcal{G}_{p+1} corresponding to $G_{I_0^\perp}$ and G have equal ranks and the embedding $\mathcal{G}_p \hookrightarrow \mathcal{G}_{p+1}$ is isometric and holomorphic. Therefore Example 3.4 implies that, viewed as a homomorphism $\Gamma \rightarrow G$, ϕ is maximal as well. \square

9. Proof of Theorem 1

For homomorphisms $\Gamma \rightarrow G$ with reductive Zariski closure, the conclusion of Theorem 1 follows from Theorem 2.2, the classification of classical simple Lie groups and the case by case analysis of balancedness in Sections 7 and 8.

Here is how the problem is reduced to the case of reductive homomorphisms. Theorem 2 of [13] asserts that if $\text{genus}(\Gamma) \geq 2 \dim(G)^2$ and G is semisimple, then $\text{Hom}(\Gamma, G)$ falls into two types of connected components: in some of them, Zariski dense homomorphisms are dense; others do not contain any Zariski dense homomorphism (call them *rigid*). Let $\phi : \Gamma \rightarrow G$ be an arbitrary homomorphism. If ϕ cannot be approximated by Zariski dense homomorphisms, then it belongs to a rigid component \mathcal{C} . Proposition 8.3 of [13] asserts that \mathcal{C} contains a reductive homomorphism ψ . The reductive case of Theorem 1 implies that G is Hermitian of nontube type and ψ is maximal. Since Toledo invariants are constant on connected components of $\text{Hom}(\Gamma, G)$, ϕ is maximal as well.

10. Appendix

For the reader's convenience, we give proofs of the maximality preserving property of two embeddings between reductive Hermitian groups (Example 3.12).

We start with a preliminary observation.

Lemma 10.1. *Let $F : Y \rightarrow X$ be an equivariant totally geodesic map between Hermitian symmetric spaces. Assume Y is irreducible. Let $P \subset X$ and $Q \subset Y$ be maximal polydisks such that $F(Q) \subset P$. Then F is positively maximality preserving if and only if $F|_Q : Q \rightarrow P$ is.*

Proof. Since Y is irreducible, F is homothetic, i.e. there exists a constant c such that $F^*\omega_X = c\omega_Y$. Sectional curvature achieves its minimum along maximal polydiscs, so

$$(\omega_X)|_P = \omega_P, \quad (\omega_Y)|_Q = \omega_Q.$$

Since $\text{rank}(P) = \text{rank}(X)$ and $\text{rank}(Q) = \text{rank}(Y)$, F is positively maximality preserving if and only if

$$c = \frac{\text{rank}(X)}{\text{rank}(Y)} \Leftrightarrow c = \frac{\text{rank}(P)}{\text{rank}(Q)} \Leftrightarrow (F|_Q)^*\omega_P = \frac{\text{rank}(P)}{\text{rank}(Q)}\omega_Q$$

if and only if $F|_Q$ is positively maximality preserving. \square

Lemma 10.2. *The embedding $SU(n, n) \hookrightarrow Sp(4n, \mathbb{R})$ is positively maximality preserving.*

Proof. Let $Y = \mathcal{D}_{n,n}$ (resp. $X = \mathcal{S}_{2n}$) denote the symmetric space associated to $H = SU(n, n)$ (resp. $G = Sp(4n, \mathbb{R})$). Let $\iota : Y \rightarrow X$ denote the corresponding embedding of symmetric spaces. We must show that

$$\iota^*\omega_X = 2\omega_Y.$$

Let us first study the case when $n = 1$. Let $V_{\mathbb{C}} = \mathbb{C}^2$ be equipped with the standard symmetric sesquilinear form $v \cdot v' = \bar{v}^{\top} v'$. Let $S = \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}$. The symmetric sesquilinear form $s(v, v') = v \cdot (Sv')$ on $V_{\mathbb{C}}$ has vanishing signature. It is easy to show that $s(Av, Av') = s(v, v')$ for $v, v' \in \mathbb{C}^2$, and $A \in SL(2, \mathbb{R})$ by a direct calculation. Hence the group $H = SU(V_{\mathbb{C}}, s)$ coincides with $SL(2, \mathbb{R})$ acting on $V_{\mathbb{C}} = \mathbb{R}^2 \otimes \mathbb{C}$. Its maximal compact subgroup L is generated by $J = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$.

Let $V_{\mathbb{R}}$ denote $V_{\mathbb{C}}$ viewed as a real vector space equipped with the symplectic form $\Omega(v, v') = \Im m(s(v, v'))$. Then $H = SU(V_{\mathbb{C}}, s)$ is a subgroup of the larger symplectic group $G = Sp(V_{\mathbb{R}}, \Omega)$. Let $\rho : SU(V_{\mathbb{C}}, s) \hookrightarrow Sp(V_{\mathbb{R}}, \Omega)$ denote the inclusion homomorphism. Then $J' = \rho(J)$ is the block diagonal 4×4 matrix with 2×2 blocks $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$. As J' is a complex structure compatible with Ω and tamed by Ω , its centralizer in $Sp(V_{\mathbb{R}}, \Omega)$ is a maximal compact subgroup K of $Sp(V_{\mathbb{R}}, \Omega)$. The adjoint actions of J on $\mathfrak{h}/\mathfrak{l}$ and of J' on $\mathfrak{g}/\mathfrak{k}$ define the complex structures of the symmetric spaces Y and X associated to H and G , so the ρ -equivariant embedding $\iota : Y \hookrightarrow X$ mapping L into K is holomorphic.

Let us view $V_{\mathbb{C}}$ as $\mathbb{R}^2 \otimes \mathbb{C}$. Then $V_{\mathbb{R}} = \mathbb{R}^2 \oplus i\mathbb{R}^2$. In these coordinates, $SL(2, \mathbb{R})$ acts diagonally. The stabilizer of this decomposition in G is the standard $SL(2, \mathbb{R}) \times SL(2, \mathbb{R})$ in $Sp(4, \mathbb{R})$. By Example 3.5, the embedding of symmetric spaces $Y = \mathcal{S}_1 \hookrightarrow \mathcal{S}_2$ corresponding to each $SL(2, \mathbb{R}) \hookrightarrow G$ is isometric and holomorphic, so $SL(2, \mathbb{R}) \times SL(2, \mathbb{R}) \hookrightarrow G$ gives rise to an isometric and holomorphic map of $Y \times Y$ onto a maximal polydisk P of $X = \mathcal{S}_2$. The image $\rho(H)$ sits diagonally in the product $SL(2, \mathbb{R}) \times SL(2, \mathbb{R})$, so $\iota : Y \rightarrow P$ factors through the diagonal $\Delta : Y \rightarrow Y \times Y$. This shows that Kähler forms fit up to a factor of 2, i.e.

$$\iota^* \omega_P = \Delta^*(\text{pr}_1^* \omega_P + \text{pr}_2^* \omega_P) = 2\omega_Y.$$

In general, let $(V_{\mathbb{C}}, s)$ be the orthogonal direct sum of n copies of the $n = 1$ example just studied. Then s has vanishing signature. Each factor gives rise to a homomorphism $SU(1, 1) \hookrightarrow H = SU(V_{\mathbb{C}}, s)$ and a map $\mathcal{D}_{1,1} = \mathcal{S}_1 \hookrightarrow Y = \mathcal{D}_{n,n}$ which, according to Example 3.3, is isometric and holomorphic. The product map $\mathcal{D}_{1,1}^n \hookrightarrow Y$ is isometric and holomorphic onto a maximal polydisk Q of Y .

Let $V_{\mathbb{R}}$ be $V_{\mathbb{C}}$ viewed as a real vector space equipped with the symplectic structure $\Omega = \Im m(s)$. Each factor of $V_{\mathbb{C}}$ is the complexification of a real 2-dimensional vector space. This gives rise to commuting embeddings $SL(2, \mathbb{R}) \hookrightarrow Sp(4n, \mathbb{R})$, and the corresponding map $\mathcal{S}_1^{2n} \hookrightarrow X = \mathcal{S}_{2n}$ is a holomorphic isometry onto a maximal polydisk P of X . The restriction of $\iota : Y \rightarrow X$ to Q is the direct product of n copies of the $n = 1$ case, so again

$$\iota^* \omega_P = 2\omega_Q.$$

With Lemma 10.1, since $\text{rank}(X) = 2 \text{rank}(Y)$, this shows that ι is positively maximality preserving. \square

Lemma 10.3. *Let h be a nondegenerate (\cdot, \cdot) -skew-symmetric binary form on a $2n$ -dimensional quaternionic vector space $V_{\mathbb{H}}$ (see Subsection 4.1). Use right multiplication by i to turn $V_{\mathbb{H}}$ into a complex $4n$ -dimensional vector space $V_{\mathbb{C}}$. Let $\mathcal{C}(q) = a$ denote the complex part of a quaternion $q = a + jb$. Then*

$$s(v, v') = \mathcal{C}(h(v, v'))$$

is a nondegenerate sesquilinear form of vanishing signature on $V_{\mathbb{C}}$. Then the corresponding embedding of groups $\rho : SO^*(4n) := SU(V_{\mathbb{H}}, h) \hookrightarrow SU(2n, 2n) := SU(V_{\mathbb{C}})$ is positively maximality preserving.

Proof. Let $\iota : Y = \mathcal{G}_{2n} \rightarrow X = \mathcal{D}_{2n,2n}$ be the corresponding embedding of Hermitian symmetric spaces. Since $\text{rank}(\mathcal{G}_{2n}) = n$ and $\text{rank}(\mathcal{D}_{2n,2n}) = 2n$, we shall show that Kähler forms match up to a factor of 2, i.e. $\iota^*\omega_X = 2\omega_Y$. Following Lemma 10.1, it suffices to understand the restriction of ι to maximal polydisks.

Let $v \cdot v' = \bar{v}^\top v'$ denote the standard positive definite $\bar{\cdot}$ -symmetric binary form on \mathbb{H}^{2n} . Let $h(v, v') = v \cdot iv'$. Then h is $(\bar{\cdot}, \text{skew})$ -symmetric and nondegenerate, so we can take $V_{\mathbb{H}} = (\mathbb{H}^{2n}, h)$. The embedding $\rho : SO^*(4n) \hookrightarrow SU(2n, 2n)$ consists in taking a quaternionic matrix X , splitting it as $X = M + jM'$ where M and M' have complex entries, and letting X act on the quaternionic vector $v = a + jb$ where $(a, b) \in (\mathbb{C}^{2n})^2 = \mathbb{C}^{4n}$. Thus

$$Xv = (M + jM')(a + jb) = Ma - \bar{M}'b + j(M'a + \bar{M}b),$$

i.e.

$$\rho(X) = \begin{pmatrix} M & -\bar{M}' \\ M' & \bar{M} \end{pmatrix}. \tag{1}$$

Let $J \in GL(2n, \mathbb{H})$ denote left multiplication by i . Elements of $Sp(2n)$ which commute with J (i.e. matrices with entries in $\mathbb{C} \subset \mathbb{H}$) form a group L isomorphic to $U(2n)$. It is a maximal compact subgroup in $SO^*(4n)$. Under ρ , this subgroup is mapped to the maximal compact subgroup $K = S(U(2n) \times U(2n))$ by $M \mapsto (M, \bar{M})$. The map J belongs to the Lie algebra $\mathfrak{h} = \mathfrak{so}^*(4n)$, it generates the center of the Lie algebra \mathfrak{l} of L . Therefore the complex structure on \mathcal{G}_{2n} arises from the adjoint action of J on $\mathfrak{h}/\mathfrak{l}$. For the same reason, the complex structure on $\mathcal{D}_{2n,2n}$ arises from the adjoint action of $J' = \text{diag}(i, \dots, i, -i, \dots, -i)$ on $\mathfrak{g}/\mathfrak{k}$. Note that at the Lie algebra level we have $\rho(J) = J'$, thus ι is holomorphic.

Let us first study the case when $n = 1$. The Lie algebra

$$\mathfrak{so}^*(4) = \left\{ A \mid A^* \begin{pmatrix} i & 0 \\ 0 & i \end{pmatrix} + \begin{pmatrix} i & 0 \\ 0 & i \end{pmatrix} A = 0 \right\}$$

is isomorphic to $\mathfrak{su}(1, 1) \oplus \mathfrak{su}(2)$, it consists of matrices of the form

$$\begin{pmatrix} i\alpha & jx \\ -jx & i\alpha \end{pmatrix} + \begin{pmatrix} i\beta & y \\ -\bar{y} & -i\beta \end{pmatrix}$$

where $\alpha, \beta \in \mathbb{R}$ and $x, y \in \mathbb{C}$. The first matrix belongs to a subalgebra \mathfrak{q} isomorphic to $\mathfrak{su}(1, 1)$. A computation based on formula (1) gives

$$\rho \begin{pmatrix} i\alpha & jx \\ -jx & i\alpha \end{pmatrix} = \begin{pmatrix} i\alpha & 0 & 0 & -\bar{x} \\ 0 & i\alpha & \bar{x} & 0 \\ 0 & x & -i\alpha & 0 \\ -x & 0 & 0 & -i\alpha \end{pmatrix}.$$

We see that $\rho(\mathfrak{q})$ is contained in the subalgebra

$$\mathfrak{p} = \left\{ \begin{pmatrix} i\alpha & 0 & 0 & -\bar{x} \\ 0 & i\alpha' & -\bar{x}' & 0 \\ 0 & -x' & -i\alpha' & 0 \\ -x & 0 & 0 & -i\alpha \end{pmatrix} \mid \alpha, \alpha' \in \mathbb{R}, x, x' \in \mathbb{C} \right\},$$

which is isomorphic to $\mathfrak{su}(1, 1) \oplus \mathfrak{su}(1, 1)$ and embedded in the standard (block diagonal) manner in $\mathfrak{su}(2, 2)$. The map $\rho|_{\mathfrak{q}} : \mathfrak{q} \rightarrow \mathfrak{p}$ is the graph of an inner automorphism of $\mathfrak{su}(1, 1)$. Geometrically, this means that ι maps $Q = \mathcal{G}_2$ (a complex line of constant curvature -1) holomorphically into a maximal polydisk P of $\mathcal{D}_{2,2}$. Now, P is holomorphically isometric to $Q \times Q$ and $\iota|_Q : Q \rightarrow P$ is the graph of an isometry I of Q . It follows that

$$\iota^* \omega_P = (\text{id}, I)^*(\text{pr}_1^* \omega_Q + \text{pr}_2^* \omega_Q) = 2\omega_Q.$$

Let us map $SO^*(4)$ as a diagonal 2×2 block in $SO^*(4n)$. This yields an embedding $\mathcal{G}_2 \hookrightarrow \mathcal{G}_{2n} = Y$ which, according to Example 3.4, is isometric and holomorphic. Splitting $V_{\mathbb{H}} = \mathbb{H}^{2n}$ as an orthogonal direct sum of 2-dimensional quaternionic vector spaces yields an isometric and holomorphic embedding of \mathcal{G}_2^n onto a maximal polydisk $Q \subset \mathcal{G}_{2n}$. Then $\iota(Q)$ is contained in the standard maximal polydisk $P \subset \mathcal{D}_{2n,2n} = X$, and $\iota|_Q : Q \rightarrow P$ is a product of n copies of the map of the previous paragraph. Therefore

$$\iota^* \omega_P = \sum_{i=1}^n \iota^* \omega_{P_i} = \sum_{i=1}^n \omega_{Q_i} = 2\omega_Q.$$

Since $\text{rank}(X) = 2 \text{rank}(Y)$, this shows that ι (and thus ρ) is positively maximality preserving. \square

Acknowledgments. I. Kim gratefully acknowledges the partial support of grant (NRF-2014R1A2A2A01005574) and a warm support of IHES during his stay. The second author acknowledges support from U.S. National Science Foundation grants DMS 1107452, 1107263, 1107367 “RNMS: Geometric structures And Representation varieties” (the GEAR Network). Many thanks to Marc Burger and Jean-Louis Clerc, who explained to us tight maps and bounded symmetric domains.

References

- [1] Bradlow, S., García-Prada, O., Gothen, P.: Surface group representations and $U(p, q)$ -Higgs bundles. *J. Differential Geom.* **64**, 111–170 (2003) [Zbl 1070.53054](#) [MR 2015045](#)
- [2] Bradlow, S., García-Prada, O., Gothen, P.: Maximal surface group representations in isometry groups of classical Hermitian symmetric spaces. *Geom. Dedicata* **122**, 185–213 (2006) [Zbl 1132.14029](#) [MR 2295550](#)
- [3] Burger, M., Iozzi, A., Wienhard, A.: Surface group representations with maximal Toledo invariant. *Ann. of Math.* **172**, 517–566 (2010) [Zbl 1208.32014](#) [MR 2680425](#)
- [4] Burger, M., Iozzi, A., Wienhard, A.: Tight homomorphisms and Hermitian symmetric spaces. *Geom. Funct. Anal.* **19**, 678–721 (2009) [Zbl 1188.53050](#) [MR 2563767](#)

- [5] Clerc, J.-L., Ørsted, B.: The Gromov norm of the Kaehler class and the Maslov index. *Asian J. Math.* **7**, 269–295 (2003); Corrigendum, *ibid.* **8**, 391–393 (2004) [Zbl 1079.53120](#) [MR 2129241](#)
- [6] Domic, A., Toledo, D.: The Gromov norm of the Kaehler class of symmetric domains. *Math. Ann.* **276**, 425–432 (1987) [Zbl 0595.53061](#) [MR 0875338](#)
- [7] Goldman, W.: Discontinuous groups and the Euler class. Thesis, Univ. of California at Berkeley (1980) [MR 2630832](#)
- [8] Goldman, W.: Representations of fundamental groups of surfaces. In: *Geometry and Topology*, J. Alexander and J. Harer (eds.), Lecture Notes in Math. 1167, Springer, 95–117 (1985) [Zbl 0575.57027](#) [MR 0827264](#)
- [9] Goldman, W., Millson, J.: Local rigidity of discrete groups acting on complex hyperbolic space. *Invent. Math.* **88**, 495–520 (1987) [Zbl 0627.22012](#) [MR 0884798](#)
- [10] Goldman, W., Millson, J.: The deformation theory of representations of fundamental groups of compact Kähler manifolds. *Publ. Math. I.H.É.S.* **67**, 43–96 (1988) [Zbl 0678.53059](#) [MR 0972343](#)
- [11] Guichard, O.: Connexité et densité des représentations irréductibles des groupes de surface dans le groupe général linéaire. *Transform. Groups* **12**, 251–292 (2007) [Zbl 1181.57021](#) [MR 2323684](#)
- [12] Hernández Lamóneda, L.: Maximal representations of surface groups in bounded symmetric domains. *Trans. Amer. Math. Soc.* **324**, 405–420 (1991) [Zbl 0733.32024](#) [MR 1033234](#)
- [13] Kim, I., Pansu, P.: Density of Zariski density for surface groups. *Duke Math. J.* **163**, 1737–1794 (2014) [Zbl 1297.57070](#) [MR 3217766](#)
- [14] Li, J.: The space of surface group representations. *Manuscripta Math.* **78**, 223–243 (1993) [Zbl 0803.32020](#) [MR 1206154](#)
- [15] Milnor, J.: On the existence of a connection with curvature zero. *Comment. Math. Helv.* **32**, 215–223 (1958) [Zbl 0196.25101](#) [MR 0095518](#)
- [16] Rapinchuk, A. S., Benyash-Krivetz, V. V., Chernousov, V. I.: Representation varieties of the fundamental groups of compact orientable surfaces. *Israel J. Math.* **93**, 29–71 (1996) [Zbl 0857.14012](#) [MR 1380633](#)
- [17] Satake, I.: *Algebraic Structures of Symmetric Domains*. Kanô Memorial Lectures 4, Iwanami Shoten, Tokyo, and Princeton Univ. Press, Princeton, NJ (1980) [Zbl 0483.32017](#) [MR 0591460](#)
- [18] Toledo, D.: Representations of surface groups in complex hyperbolic space. *J. Differential Geom.* **29**, 125–133 (1989) [Zbl 0676.57012](#) [MR 0978081](#)
- [19] Turaev, B.: A cocycle of the symplectic first Chern class and Maslov indices. *Funktsional. Anal. i Prilozhen.* **18**, no. 1, 43–48 (1984) (in Russian) [Zbl 0556.55012](#) [MR 0739088](#)