

András Némethi · Baldur Sigurdsson

The geometric genus of hypersurface singularities

Received October 3, 2013 and in revised form February 20, 2014

Abstract. Using path lattice cohomology we provide a conceptual topological characterization of the geometric genus for certain complex normal surface singularities with rational homology sphere links, which is uniformly valid for all superisolated and Newton non-degenerate hypersurface singularities.

Keywords. Normal surface singularities, hypersurface singularities, links of singularities, Newton non-degenerate singularities, geometric genus, plumbing graphs, \mathbb{Q} -homology spheres, lattice cohomology, path lattice cohomology, Heegaard–Floer homology, Seiberg–Witten invariant

1. Introduction

In this introduction we present the main result of the manuscript; for detailed definitions, motivations, historical remarks and examples see the next section.

In the last years several conjectures and theorems target the topological characterization of the geometric genus p_g of complex normal surface singularities with rational homology sphere links. They are usually formulated for certain families, and any attempt to find a uniform characterization failed.

In order to have a chance for such a characterization, one has to assume two restrictions, an analytical one and a topological one. The *Casson Invariant Conjecture* (CIC) of Neumann and Wahl [46] predicts that for a complete intersection with integral homology sphere link, p_g can be determined from the Casson invariant of the link (see 2.4.1 here). This was generalized to rational homology sphere links by the first author and Nicolaescu [40]; the *Seiberg–Witten Invariant Conjecture* (SWIC) connects p_g with the Seiberg–Witten invariant of the link (associated with the canonical $spin^c$ -structure; see 2.4.2). The predicted formula was proved for several analytic families (e.g. rational, minimally elliptic, weighted homogeneous, splice quotient singularities), but it fails

A. Némethi: A. Rényi Institute of Mathematics, 1053 Budapest, Reáltanoda u. 13-15, Hungary; e-mail: nemethi.andras@renyi.mta.hu

B. Sigurdsson: Central European University, Budapest, and A. Rényi Institute of Mathematics, 1053 Budapest, Reáltanoda u. 13-15, Hungary; e-mail: sigurdsson.baldur@renyi.mta.hu

Mathematics Subject Classification (2010): Primary 32S05, 32S25, 32S50, 57M27; Secondary 14Bxx, 32Sxx, 57R57, 55N35

to hold for some sporadic hypersurfaces, namely, for certain superisolated singularities.

The present note aims to find a uniform treatment for these counterexamples, and proposes a new conceptual topological candidate for p_g , which is valid even for other important families of hypersurface singularities, e.g. for those with Newton non-degenerate principal part. The main ingredient of the topological characterization is the *path lattice cohomology* associated with the link (or, with the negative definite lattice of a fixed resolution graph).

Recall that the lattice cohomology $\{\mathbb{H}_{\text{red}}^q(M)\}_{q \geq 0}$ of the link (introduced in [35]) is a new categorification of the Seiberg–Witten invariant, that is, its ‘normalized’ Euler characteristic $eu(\mathbb{H}^*(M))$ is the Seiberg–Witten invariant. In the p_g -comparisons the main dominating term is the first module $\mathbb{H}_{\text{red}}^0(M)$, and in the superisolated case the non-vanishing of the next terms is responsible for the failure of the SWIC.

Accordingly, the proposed new invariant targets a different version of lattice cohomology, which concentrates only on the $q = 0$ part, and even optimizes it along different ‘paths’. A path is a sequence of integral cycles supported on the exceptional curve of a fixed resolution, at each step increasing only by a base element, and connecting the trivial cycle to the anticanonical cycle. For such a path γ one defines a path lattice cohomology $\mathbb{H}^0(\gamma)$, and one takes its normalized rank $eu(\mathbb{H}^0(\gamma))$. Then one shows that for any analytic type one has $p_g \leq \min_{\gamma} eu(\mathbb{H}^0(\gamma))$, hence it provides a natural topological upper bound for the geometric genus. (The authors do not know if $\min_{\gamma} eu(\mathbb{H}^0(\gamma))$ can be defined by any other construction, say, using gauge theory or low dimensional topology.)

The main result of the article is the following.

Theorem 1.0.1. *Assume that $(X, 0)$ is a normal surface singularity whose link is a rational homology sphere. Then the identity $p_g = \min_{\gamma} eu(\mathbb{H}^0(\gamma))$ is true in the following cases:*

- (a) *if $\mathbb{H}^q(M) = 0$ for $q \geq 1$ and the singular germ satisfies the SWIC Conjecture (in particular, for all weighted homogeneous and minimally elliptic singularities);*
- (b) *for superisolated singularities (with any number of cusps);*
- (c) *for singularities with non-degenerate Newton principal part.*

Moreover, since the conjecture is stable with respect to equisingular deformation of hypersurfaces, the conjecture remains valid for such deformations of any of the above cases.

The next section contains all the necessary definitions, main guiding examples, and status quo of the problem. Section 3 contains the proof for superisolated germs. In this case the link is a surgery 3-manifold. The proof has two non-trivial ingredients, already present in the recent literature: the first one provides the lattice cohomology of surgery 3-manifolds [44], the other is an application of a d -invariant vanishing result for certain L -space surgery 3-manifolds in Heegaard–Floer knot theory [5]. The next sections contain the proof of the Newton non-degenerate case; it involves deeply the very specific combinatorics of the associated toric resolution, and a lattice point counting.

2. Geometric genus formulae, conjectures, guiding examples

2.1. Preliminaries: the geometric genus

Let us fix a complex analytic normal surface singularity $(X, 0)$. Let M be its link, the oriented smooth 3-manifold which is the boundary of a convenient small representative X of the germ. Since the real cone over M is homeomorphic to X , M characterizes completely the local topology of $(X, 0)$. If we consider a resolution of X with dual resolution graph G , then M can be realized as a plumbed 3-manifold associated with G , and, in fact, it contains the same information as G itself (cf. [45]). The *topological invariants* of the germ $(X, 0)$ are read off either from the topology of M , or from the combinatorics of G .

The *analytic invariants* of $(X, 0)$ are a priori associated with the analytic structure of $(X, 0)$ read off e.g. from the local algebra $\mathcal{O}_{X,0}$, or from the analytic sheaves of a resolution $\tilde{X} \rightarrow X$ of X . The very first one, and probably the most important one, is the *geometric genus* $p_g := \dim H^1(\tilde{X}, \mathcal{O}_{\tilde{X}})$. It guides (partially) the classification of singular germs and their deformation theory [4, 19, 20, 22, 23, 31, 58, 59, 56]; it is the local analog of the global Todd index of complex manifolds. As a ‘local index’, it has several key connections with other numerical invariants (see e.g. (2.1.3)). Usually, the geometric genus cannot be determined from the link, even if we consider rather ‘simple’ singularities. For example, the hypersurface singularities $\{x^2 + y^3 + z^{18} = 0\}$ and $\{z^2 = y(x^4 + y^6)\}$ have the same link but their p_g are 3 and 2 respectively (cf. [39, 4.6]).

Nevertheless, there is a strong belief, seriously supported by the results of the last decade, that under some restrictions, p_g can be determined from M . First of all, one needs to assume that M is a *rational homology sphere*, or equivalently, the resolution exceptional divisor is a tree of rational curves (in the above example the first Betti number $b_1(M)$ of M is 2). This is still not enough. Consider for example an elliptic singularity with $p_g \geq 2$ (and even with integral homology sphere link), say the hypersurface $x^2 + y^3 + z^{13} = 0$ with $p_g = 2$. Then, by [20, 4.1], the generic (non-Gorenstein) analytic structure supported by the same topological type has $p_g = 1$ (see also [31]). For other pairs with the same integral homology sphere link, but with different geometric genus, see [26]. Hence, one needs to add some analytic restriction too to guarantee the topological characterization of p_g .

In the choice of the analytic structures one possibility would be to consider a *generic* one on each irreducible component of the moduli space of analytic structures—which, by the semicontinuity of p_g [11], would provide the smallest p_g of that moduli component. But it is equally challenging (and this is our interest here) to search for the geometric genus of special families of germs, which are the candidates providing the topological upper bound for p_g . They can be related either to special properties of $(X, 0)$ (e.g. hypersurface, ICIS, Gorenstein, \mathbb{Q} -Gorenstein), or with special constructions (see 2.1.1 below).

Example 2.1.1. The next families will play a key role in the next discussions.

(a) *Splice quotient* singularities were introduced by Neumann and Wahl [46, 47, 48]; their graph G has to satisfy some arithmetical properties, which allow one to write down from the combinatorics of G the equations of the universal abelian cover of $(X, 0)$ (up to equisingular deformation), together with the corresponding action of $H_1(M, \mathbb{Z})$ on it.

They generalize the weighted homogeneous germs, but the equations associated with different nodes might have different weights and degrees. By construction their geometric genus depends only on G ; the precise expression is given in [43] as an answer to Conjecture 2.4.2.

(b) *Superisolated* singularities were introduced by I. Luengo [25], and they played a crucial role in several testing procedures or counterexamples [2, 3, 26, 27]. If C is a projective reduced plane curve with homogeneous equation f_d of degree d , and f_{d+1} is a generic homogeneous equation of degree $d + 1$, then $f = f_d + f_{d+1} : (\mathbb{C}^3, 0) \rightarrow (\mathbb{C}, 0)$ is called superisolated. Its link and p_g are independent of the choice of f_{d+1} (see e.g. [26]); in fact, $p_g = d(d - 1)(d - 2)/6$. We will assume that C is irreducible. Then the link of f is $\mathbb{Q}HS^3$ if and only if C is a rational cuspidal curve. We write ν for the number of cusps.

(c) For *Newton non-degenerate hypersurfaces* see e.g. [16]. The principal part of such a germ is a sum of monomials situated on a fixed Newton diagram Γ with generic coefficients. The geometric genus can be recovered as the number of lattice points with all positive entries and which are ‘not above Γ ’ [29]. For more see Section 4.

2.1.2. If $(X, 0)$ is a hypersurface singularity in $(\mathbb{C}^3, 0)$ then the topological and analytic invariants are strongly related to those provided by the *embedded topological type* of $(X, 0)$, that is, the topology of the embedding $M \subset S^5$, and the numerical invariants of the Milnor fibration. Recall that the second Betti number of the Milnor fiber F is the Milnor number μ , the intersection form on $H_2(F)$ determines the Sylvester invariants μ_+ , μ_- and μ_0 , while the signature is defined by $\sigma = \mu_+ - \mu_-$. Modulo the link M , the numerical invariants p_g , μ and σ are related by two identities. Indeed, if K is the canonical class/cycle on \tilde{X} and $|\mathcal{V}|$ is the number of vertices of G , then $K^2 + |\mathcal{V}|$ is a well-defined topological invariant of M , and one has the following identities (valid in fact for any smoothing of a Gorenstein $(X, 0)$) [9, 21, 54, 57, 24]:

$$(a) \mu = 12p_g + K^2 + |\mathcal{V}| - b_1(M), \quad (b) -\sigma = 8p_g + K^2 + |\mathcal{V}|. \quad (2.1.3)$$

Hence, if any of p_g , μ or σ can be described from M then the same is true for all of them. Note that μ can be recovered from the *embedded topological type* (as the second Betti number of the universal cover of $S^5 \setminus M$), hence this fact remains true for p_g as well.

2.1.4. Notation regarding G . Regarding the link we will need the following notation and terminology. We fix a resolution $\pi : \tilde{X} \rightarrow X$ with resolution graph G as above. We assume that M is a $\mathbb{Q}HS^3$. Consider the lattice $L = H_2(\tilde{X}, \mathbb{Z})$; it is freely generated by $\{E_v\}_{v \in \mathcal{V}}$, the irreducible components of the exceptional divisor $E := \pi^{-1}(0)$ of π . It is known that G is connected and L (that is, the intersection form $\mathfrak{I} := \{(E_v, E_u)\}_{v,u}$) is negative definite. The determinant of the graph G is defined as the absolute value of $\det(\mathfrak{I})$.

Since we treat mainly hypersurface singularities, which are Gorenstein, we will assume that G is *numerically Gorenstein*. This means that the *canonical cycle* K , which satisfies the system of *adjunction relations* $(K + E_v, E_v) = -2$ for all v , is an integral cycle of L .

We define $\chi : L \rightarrow \mathbb{Z}$ by $\chi(l) = -(l, l + K)/2$. (This is the Riemann–Roch formula: for l effective $\chi(l)$ is the analytic Euler characteristic of \mathcal{O}_l .) Set $m := \min_{l \in L} \chi(l)$.

If $l'_k = \sum_v l'_{kv} E_v$ for $k = 1, 2$, then we write $\min\{l'_1, l'_2\} := \sum_v \min\{l'_{1v}, l'_{2v}\} E_v$, and $l'_1 \leq l'_2$ if $l'_{1v} \leq l'_{2v}$ for all $v \in \mathcal{V}$. The valency of the vertex $v \in \mathcal{V}$ in G is denoted by δ_v . We will write $Z_K := -K$. Furthermore, we will assume that G is a *minimal good (resolution) graph*. In that case one has the following.

Lemma 2.1.5. *Either $(X, 0)$ is rational (equivalently A-D-E, hence $Z_K = 0$) or $Z_K > E$. Moreover, in the second case, the support of $Z_K - E$ is connected.*

Proof. It is known that the only rational numerically Gorenstein graphs are of type A-D-E. Otherwise, under the assumption that the graph is a tree of rational vertices, $Z_K \geq E$ by [52, 2.8]. But $Z_K = E$ cannot happen. Indeed, $p_g = h^1(\mathcal{O}_{Z_K}) = h^1(\mathcal{O}_E) = 0$ would imply rationality, hence $Z_K = 0$. For the connectedness, see e.g. [55, 2.10] or [52, 2.6]. \square

2.1.6. In the last years there has been an intense activity to identify p_g with certain ingredients of the Seiberg–Witten (or other equivalent/similar) theories. This worked nicely for several analytic structures, but failed for some others. Though the goal of the present note is to present the parallel theory for those cases which fail the ‘Seiberg–Witten connection’, for a complete picture we need to review certain notions from this part as well.

2.2. Preliminaries: Seiberg–Witten invariant and lattice cohomology of M

Here is a short review of the lattice cohomology and path lattice cohomology of M . For more details see [32, 34, 35, 37].

2.2.1. Lattice cohomology. $\mathbb{Z}^s \otimes \mathbb{R}$ has a natural decomposition into cubes. The 0-dimensional cubes are the lattice points \mathbb{Z}^s . Any $l \in \mathbb{Z}^s$ and subset $I \subseteq \mathcal{J}$ of cardinality q define a q -dimensional cube, which has its vertices at the lattice points $(l + \sum_{j \in I'} E_j)_{I'}$, where I' runs over all subsets of I . We define the weight of any such cube \square_q by

$$w(\square_q) := \max\{\chi(v) : v \text{ is a vertex of } \square_q\}.$$

Lattice cohomology (associated with the canonical $spin^c$ structure of M) is defined as follows. For each $N \in \mathbb{Z}$, define $S_N \subset \mathbb{R}^s$ as the union of all the cubes \square_q (of any dimension) with $w(\square_q) \leq N$. Clearly, $S_N = \emptyset$ whenever $N < m$. Then for any $q \geq 0$, set

$$\mathbb{H}^q(G) := \bigoplus_{N \geq m} H^q(S_N, \mathbb{Z}), \quad \mathbb{H}_{\text{red}}^q(G) := \bigoplus_{N \geq m} \tilde{H}^q(S_N, \mathbb{Z}).$$

Then \mathbb{H}^q is $2\mathbb{Z}$ -graded, the $d = 2N$ -homogeneous elements consist of $H^q(S_N, \mathbb{Z})$. Also, \mathbb{H}^q is a $\mathbb{Z}[U]$ -module: the U -action is given by the restriction map $H^q(S_{N+1}, \mathbb{Z}) \rightarrow H^q(S_N, \mathbb{Z})$. Moreover, for $q = 0$, a base-point $l \in S_m$ provides an augmentation $H^0(S_N, \mathbb{Z}) = \mathbb{Z} \oplus \tilde{H}^0(S_N, \mathbb{Z})$, hence an augmentation of the graded $\mathbb{Z}[U]$ -modules $\mathbb{H}^0 = (\bigoplus_{N \geq m} \mathbb{Z}) \oplus \mathbb{H}_{\text{red}}^0$. The graded $\mathbb{Z}[U]$ -modules \mathbb{H}^* and $\mathbb{H}_{\text{red}}^*$ are called the *lattice cohomology* and the *reduced lattice cohomology* of G . They depend only on M , and $\mathbb{H}_{\text{red}}^*$ is a finite \mathbb{Z} -module.

2.2.2. The Seiberg–Witten invariant. Recall that the Seiberg–Witten invariants of an oriented 3-manifold M are rational numbers $\text{sw}_{\mathfrak{s}}(M)$ associated with the Spin^c -structures \mathfrak{s} of M . They can be recovered as (normalized) Euler characteristics of different cohomology theories, e.g. for their relation with Heegaard–Floer homology see [50, 51]). In the following we consider only the canonical spin^c structure, hence the symbol \mathfrak{s} will be omitted.

By [36] the normalized Euler characteristic of lattice cohomology also agrees with the Seiberg–Witten invariant:

$$-\text{sw}(M) - (K^2 + |\mathcal{V}|)/8 = eu(\mathbb{H}^*(M)), \tag{2.2.3}$$

where $eu(\mathbb{H}^*(M)) := -m + \sum_q (-1)^q \text{rank}_{\mathbb{Z}} \mathbb{H}_{\text{red}}^q(M)$. Later it will be convenient to use the following notation as well: $eu(\mathbb{H}^0(M)) := -m + \text{rank}_{\mathbb{Z}} \mathbb{H}_{\text{red}}^0(M)$.

2.3. Path lattice cohomology

The search for a topological upper bound for p_g leads to the definition of path lattice cohomology (in fact, this was the starting point of lattice cohomology as well) [32, 35].

Consider a sequence $\gamma := \{l_i\}_{i=0}^t$ with $l_i \in L$ such that $l_0 = 0$, $l_t = Z_K$, and $l_{i+1} = l_i + E_{v(i)}$ for some vertex $v(i) \in \mathcal{V}(G)$. This defines a *path* (or 1-dimensional simplicial complex) with 0-cubes $\{l_i\}_i$ and 1-cubes $[l_i, l_{i+1}]$. We can repeat the construction of lattice cohomology, but now only for those cubes which are supported by γ . Indeed, let $m_\gamma = \min_i \chi(l_i)$, and set S_N^γ to be the union of the cubes supported by γ and with weight $\leq N$. Then one defines

$$\mathbb{H}^q(\gamma) = \bigoplus_{N \geq m_\gamma} H^q(S_N^\gamma, \mathbb{Z}), \quad \mathbb{H}_{\text{red}}^q(\gamma) = \bigoplus_{N \geq m_\gamma} \tilde{H}^q(S_N^\gamma, \mathbb{Z}).$$

It turns out that $\mathbb{H}^q(\gamma) = 0$ for $q \neq 0$, $\mathbb{H}^0(\gamma) = (\bigoplus_{N \geq m_\gamma} \mathbb{Z}) \oplus \mathbb{H}_{\text{red}}^0(\gamma)$, and $\mathbb{H}_{\text{red}}^0(\gamma)$ is a finite \mathbb{Z} -module. Similarly to lattice cohomology, we set

$$eu(\mathbb{H}^0(\gamma)) := -m_\gamma + \text{rank}_{\mathbb{Z}} \mathbb{H}_{\text{red}}^0(\gamma).$$

One verifies (see [35, 3.5.2]) that

$$eu(\mathbb{H}^0(\gamma)) = \sum_{i=0}^{t-1} \max\{0, \chi(l_i) - \chi(l_{i+1})\}. \tag{2.3.1}$$

There is a natural cohomological morphism $r^* : \mathbb{H}^0(G) \rightarrow \mathbb{H}^0(\gamma)$ induced by restriction. Usually it is neither injective nor surjective. Nevertheless, r^* is onto for certain well-chosen paths. Moreover, if r^* is onto, then by [35, 3.5.4], $eu(\mathbb{H}^0(\gamma)) \leq eu(\mathbb{H}^0(G))$, hence

$$\min_{\gamma} eu(\mathbb{H}^0(\gamma)) \leq eu(\mathbb{H}^0(G)). \tag{2.3.2}$$

Intuitively, $eu(\mathbb{H}^0(G))$ depends on those lattice points $\{l_m\}_{m \in M}$ of L which realize the ‘local minima’ of χ , and also on the ‘optimal’ connecting paths of these points: for each

pair l_m and $l_{m'}$ there is a minimal $N(m, m')$ such that l_m and $l_{m'}$ can be connected by a path in $S_{N(m, m')}$. On the other hand, $eu(\mathbb{H}^0(\gamma))$ codifies similar data supported on the path γ , and $\min_\gamma eu(\mathbb{H}^0(\gamma))$ minimizes the sum $\sum_{i=0}^{t-1} \max\{0, \chi(l_i) - \chi(l_{i+1})\}$ among all the possible paths γ .

2.3.3. The analytic interpretation of $\min_\gamma eu(\mathbb{H}^0(\gamma))$. For any analytic realization, by Riemenschneider–Kodaira vanishing $h^1(\tilde{X}, \mathcal{O}_{\tilde{X}}(-Z_K)) = 0$, hence $p_g = h^1(\mathcal{O}_{Z_K})$.

Next, consider a sequence $\{l_i\}_{i=0}^t$ with $l_i \in L$ such that $l_0 = 0$, $l_t = Z_K$, and $l_{i+1} = l_i + E_{v(i)}$ as above. Recall $E_j \simeq \mathbb{P}^1$ for all j , hence $\chi(l_{i+1}) - \chi(l_i) = 1 - (E_{v(i)}, l_i)$. Then for any $0 \leq i < t$ the exact sequence $0 \rightarrow \mathcal{O}_{E_{v(i)}}(-l_i) \rightarrow \mathcal{O}_{l_{i+1}} \rightarrow \mathcal{O}_{l_i} \rightarrow 0$ induces

$$h^1(\mathcal{O}_{l_{i+1}}) - h^1(\mathcal{O}_{l_i}) \leq h^1(\mathcal{O}_{E_{v(i)}}(-l_i)) = \max\{0, \chi(l_i) - \chi(l_{i+1})\}. \tag{2.3.4}$$

Taking the sum one obtains $h^1(\mathcal{O}_{Z_K}) \leq eu(\mathbb{H}^0(\gamma))$ for any path γ , hence

$$p_g \leq \min_\gamma eu(\mathbb{H}^0(\gamma)). \tag{2.3.5}$$

Equality holds if for some γ the cohomology exact sequences split for all i .

Usually the concrete computation of $\min_\gamma eu(\mathbb{H}^0(\gamma))$ is rather difficult.

2.4. Some conjectures, results and examples

We briefly review some key steps in the topological characterization of the geometric genus via the Seiberg–Witten invariant. We start with the *Casson Invariant Conjecture* (CIC) of Neumann–Wahl:

Conjecture 2.4.1 ([46]). *Consider an isolated complete intersection singularity with signature σ , and whose link is an integral homology sphere with Casson invariant $\lambda(M)$. Then $\sigma/8 = \lambda(M)$.*

The conjecture was verified for Brieskorn–Hamm complete intersections and for those hypersurfaces which are suspensions of irreducible plane curve singularities [46] (see also [15]). Note that via (2.1.3)(b), the identity can be replaced by $p_g = -\lambda(M) - (K^2 + |\mathcal{V}|)/8$, a version independent of any smoothing. This version was verified for splice quotient singularities (without the ICIS assumption) in [42]. The original CIC is still open.

One of the difficulties of a possible proof is the lack of any characterization/description of *integral homology sphere* hypersurfaces or complete intersection links other than iterated cyclic covers (for the behavior of $\lambda(M)$ for such covers, see e.g. Collin–Saveliev [7, 8]). Having no other examples in hand, it is hard to decide whether the validity of the conjecture is guaranteed merely by the special properties of cyclic covers, or it covers a much deeper geometrical phenomenon. Moreover, integral homology sphere links appear rather rarely (e.g. among the Newton non-degenerate hypersurfaces all germs with $\mathbb{Z}HS^3$ links are of Brieskorn type, while among rational graphs there is only one, namely the E_8). Hence, it was necessary to extend the above conjecture to *rational homology sphere links*. The conjectured identity was proposed by Némethi–Nicolescu:

Conjecture 2.4.2 ([40]). *Assume that the link M of a normal surface singularity is a rational homology sphere, whose Seiberg–Witten invariant (associated with the canonical $spin^c$ -structure) is $sw(M)$. If its analytic structure is ‘nice’ then*

$$p_g = -sw(M) - (K^2 + |\mathcal{V}|)/8.$$

The conjecture is proved for splice quotient singularities (including all rational, minimal elliptic and weighted homogeneous singularities) [6, 38, 43], and for suspensions of irreducible plane curve singularities [41]. It was extended to the equivariant case (involving the Seiberg–Witten invariant of all $spin^c$ -structures and equivariant geometric genus of the universal abelian cover [6, 38]).

On the other hand, there are even hypersurface singularities which do not satisfy the conjecture. The typical counterexample are the superisolated singularities with $\nu \geq 2$ [26].

Example 2.4.3. Assume that $(X, 0)$ is a superisolated singularity with $\mathbb{Q}HS^3$ link.

(a) If $\nu = 1$, that is, C is unicuspidal, whose local cusp has local irreducible plane curve singularity knot $K \subset S^3$, then $M = S^3_{-d}(K)$. In [13] (see also [12, 44]) it is proved that the statement of Conjecture 2.4.2 is equivalent to a ‘Density Property’ of the semigroup of the local cusp. This last property was checked in [12] for ‘all known’ curves C via case-by-case verification, and it was also proved recently in [5] using the d -invariant of Heegaard–Floer theory. Hence $p_g = eu(\mathbb{H}^*(M))$.


Here we wish to emphasize an important point. The link M with $\nu = 1$ is ‘almost rational’, that is, modifying the resolution graph at only one vertex we can get a rational graph (see [33]). Consequently (see [37]), one has the vanishing $\mathbb{H}^q(M) = 0$ for $q \geq 1$. Therefore, $eu(\mathbb{H}^*(M)) = eu(\mathbb{H}^0(M))$. In particular, in this case

$$p_g \leq \min_{\gamma} eu(\mathbb{H}^0(\gamma)) \leq eu(\mathbb{H}^0(M)) = eu(\mathbb{H}^*(M)) = p_g, \tag{2.4.4}$$

hence everywhere we must have equality.

(b) Nevertheless, for $\nu \geq 2$ counterexamples to Conjecture 2.4.2 exist [26]. In this case the above vanishing has the weaker form: $\mathbb{H}^q(M) = 0$ only for $q \geq \nu$ [44, 37]. Hence, as we will see, for superisolated singularities the non-vanishing of $\mathbb{H}^q(M)$ ($1 \leq q < \nu$) obstructs the validity of Conjecture 2.4.2.

Let us consider the case C_4 of [26] (see also [35, 7.3.3]). C has degree $d = 5$ and two cusps, both with one Puiseux pair: $(3, 4)$ and $(2, 7)$. The graph G is




One shows that $m = -5$, $\text{rank}_{\mathbb{Z}}(\mathbb{H}^0) = 5$, $\text{rank}_{\mathbb{Z}}(\mathbb{H}^1) = 2$. Hence $eu(\mathbb{H}^0) = 10$, but $eu(\mathbb{H}^*) = 8$. Since for the superisolated germ with $d = 5$ one has $p_g = 10$, by equations (2.3.2) and (2.3.5) one gets $\min_{\gamma} eu(\mathbb{H}^0(\gamma)) = 10$ as well. Hence, for this superisolated germ, (2.3.5) is valid with equality, while Conjecture 2.4.2 fails.

If we take any other analytic structure supported by the above graph, $p_g \leq 10$ still holds by (2.3.5), hence superisolated germs realize the optimal upper bound.

Note also that this topological type supports another natural analytic structure, namely a splice quotient analytic type: it is the \mathbb{Z}_5 -factor of the complete intersection $\{z_1^3 + z_2^4 + z_3^5 z_4 = z_3^7 + z_4^2 + z_1^4 z_2 = 0\} \subset (\mathbb{C}^4, 0)$ by the diagonal action $(\alpha^2, \alpha^4, \alpha, \alpha)$ ($\alpha^5 = 1$). By [43] it satisfies the SWIC Conjecture 2.4.2, hence $p_g = 8$.

In particular, in their choices of the topological characterization of their p_g , some analytic structures prefer $eu(\mathbb{H}^*)$, some of them the extremal $\min_\gamma eu(\mathbb{H}^0(\gamma))$ (and there might even exist other choices).

(c) We can ask whether the choice between $eu(\mathbb{H}^*)$ and $\min_\gamma eu(\mathbb{H}^0(\gamma))$ is uniform in the case of hypersurface singularities, that is, if all hypersurfaces choose $\min_\gamma eu(\mathbb{H}^0(\gamma))$, as superisolated germs do. We expect that this is not the case; here is a possible candidate which makes a different choice. We consider the suspension $f(x, y, z) := (x^3 + y^2)^2 + yx^5 + z^{19}$ of an irreducible plane curve singularity with two Puiseux pairs. Note that in this case the link is even an integral homology sphere. The graph of $\{f = 0\}$ is



The following facts were checked by Helge M. Pedersen (via a computer program based on the ‘Reduction Theorem’ of [17]): $m = -18$, $\text{rank } \mathbb{H}_{\text{red}}^0 = 26$, hence $eu(\mathbb{H}^0) = 44$, and $\text{rank } \mathbb{H}^1 = 8$. Since the graph has only two nodes, one has $\mathbb{H}^q = 0$ for $q \geq 2$ (cf. [37]), hence $eu(\mathbb{H}^*) = 44 - 8 = 36$. On the other hand, the Milnor number of the plane curve singularity is 16, hence the Milnor number of f is $\mu = 16 \cdot 18 = 288$. Then, by 2.1.3, one gets $p_g = 36$ (as expected, since this germ satisfies both Conjectures 2.4.1 and 2.4.2, cf. [41]).

Therefore (cf. (2.4.4)), $36 \leq \min_\gamma eu(\mathbb{H}^0(\gamma)) \leq 44$. Computer search in the ‘reduced lattice’ of [17] gives 44, however at this moment we do not know if the ‘Reduction Theorem’ of [17] works for $\min_\gamma eu(\mathbb{H}^0(\gamma))$ too. Still, we expect that $p_g = eu(\mathbb{H}^*) < \min_\gamma eu(\mathbb{H}^0(\gamma))$.

2.5. The new proposed identity

Having in mind the conclusion of Example 2.4.3(b)–(c), we can ask how accidental the superisolated example is; or, what are the choices of other important families of hypersurfaces, e.g. of Newton non-degenerate ones. Here we wish to recall that in [6] it is shown that for such germs one can recover from M the Newton diagram of the equation, hence the equisingularity type of the germ too. Hence, in principle, p_g can be recovered from M ; however this statement does not indicate any topological candidate for p_g . (For comparison, a similar statement regarding the possibility to recover the equisingularity type of suspensions of irreducible curves from their link is provided in [28]. In that case the choice is $p_g = eu(\mathbb{H}^*)$.)

The next theorem says that the extremal choice $\min_\gamma eu(\mathbb{H}^0(\gamma))$ is not accidental at all: in fact, all superisolated and Newton non-degenerate germs prefer uniformly exactly this one.

Theorem 2.5.1. *Assume that $(X, 0)$ is a normal surface singularity with rational homology sphere link. Then the identity $p_g = \min_{\gamma} eu(\mathbb{H}^0(\gamma))$ is true in the following cases:*

- (a) *if $p_g = eu(\mathbb{H}^0(M))$ [this happens e.g. whenever $\mathbb{H}^q(M) = 0$ for $q \geq 1$ and $(X, 0)$ satisfies the SWIC (Conjecture 2.4.2); in particular, the conjecture is true for all weighted homogeneous and minimally elliptic singularities];*
- (b) *for superisolated singularities with any number of cusps (in this case, in fact, $p_g = \min_{\gamma} eu(\mathbb{H}^0(\gamma)) = eu(\mathbb{H}^0(M))$ too);*
- (c) *for singularities with non-degenerate Newton principal part.*

Since the conjecture is stable with respect to equisingular deformation of hypersurfaces, it remains valid for such deformations of any of the above cases.

Corollary 2.5.2. *A superisolated singularity, where C is an irreducible rational unicuspidal curve, satisfies the Seiberg–Witten Invariant Conjecture 2.4.2 (predicted in [12, 13] too).*


Indeed, in this case the graph is ‘almost rational’ (cf. [33]), hence $\mathbb{H}^q(M) = 0$ for $q \geq 1$ by [37]. In particular, $eu(\mathbb{H}^*(M)) = eu(\mathbb{H}^0(M))$. Hence part (b) of the theorem suffices.

2.5.3. Let us repeat the meaning of the identity in Theorem 2.5.1. For any sequence $\gamma := \{l_i\}_{i=0}^t$ with $l_i \in L$ where $l_0 = 0$, $l_t = Z_K$, and $l_{i+1} = l_i + E_{v(i)}$ ($v(i) \in \mathcal{V}$) we set $eu(\mathbb{H}^0(\gamma)) := \sum_{i=0}^{t-1} \max\{0, \chi(l_i) - \chi(l_{i+1})\} = \sum_{i=0}^{t-1} \max\{0, -1 + (l_i, E_{v(i)})\}$. The statement is that $p_g = eu(\mathbb{H}^0(\gamma))$ for a well-chosen path γ .

2.5.4. Part (a) follows easily via the inequalities (2.3.2) and (2.3.5). By isolating (a) we wish to emphasize that for several ‘simple cases’, one has all the equalities $p_g = \min_{\gamma} eu(\mathbb{H}^0(\gamma)) = eu(\mathbb{H}^0(M)) = eu(\mathbb{H}^*(M))$. (This explains why earlier, when we had complete information only about these simple cases, it was difficult to predict the general behavior.)

Regarding (a), note also that the condition $p_g = \min_{\gamma} eu(\mathbb{H}^0(\gamma))$ is more restrictive: there exist examples with $p_g = \min_{\gamma} eu(\mathbb{H}^0(\gamma))$ but $p_g < eu(\mathbb{H}^0(M))$ (see next example).

Example 2.5.5 ([17]). Consider the germ $(X, 0) = \{x^{13} + y^{13} + x^2y^2 + z^3 = 0\}$ with non-degenerate Newton principal part with $p_g = 5$. The graph is



Then $m = -1$, $eu(\mathbb{H}^0) = 6$, $\text{rank } \mathbb{H}^1 = 1$, and $\min_{\gamma} eu(\mathbb{H}^0(\gamma)) = 5$.

2.5.6. The proof of part (b) of Theorem 2.5.1 will be given in Section 3. In fact, we will show that for superisolated singularities we have $p_g = eu(\mathbb{H}^0(M))$, hence (a) applies.

The Newton non-degenerate case is treated in the remaining sections.

3. The proof of Theorem 2.5.1(b) for superisolated singularities

For the proof of Theorem 2.5.1(b) we have to combine two results from the literature. In order to provide a more complete picture, we give some additional details as well.

In the last years we have witnessed a focused effort to prove the (SWI) Conjecture 2.4.2 for superisolated singularities with $\nu = 1$ cusp, or to understand the main cause of its failure in particular cases with $\nu \geq 2$. This materialized in several results [12, 13, 14, 26, 33, 34, 44] and a new conjecture regarding the algebraic realization of cuspidal rational curves of fixed degree and given local singularities, the ‘semigroup distribution property’ [12, 13]. These results and their reformulations in different languages of Heegaard–Floer or Seiberg–Witten theory (although they were originally motivated and guided by the SWIC and the distribution property), reorganized and put together in a new puzzle, provide the proof of Theorem 2.5.1. This also shows that, in fact, for *certain analytic structures* the statement of 2.5.1 is the *right guiding statement* and not the one in the SWIC.

Let $f = f_d + f_{d+1} : (\mathbb{C}^3, 0) \rightarrow (\mathbb{C}, 0)$ be a superisolated singularity as in Example 2.1.1(b), where $C = \{f_d = 0\} \subset \mathbb{C}\mathbb{P}^2$ is an irreducible rational cuspidal curve with ν cusps. Let $(C, p_i) \subset (\mathbb{C}\mathbb{P}^2, p_i)$ be the local singularities of C ; denote their Milnor numbers by μ_i , and their local links by $K_i \subset S^3$. Then the link of $(X, 0) = \{f = 0\}$ is $S^3_{-d}(K)$, where K is the connected sum $K_1 \# \dots \# K_\nu$. For the plumbing graph constructed from the embedded resolution graphs of (C, p_i) and the integer d , see [44, 2.3]. The degree d and the local singularity types (C, p_i) are related: the rationality of C implies $\sum_i \mu_i = (d - 1)(d - 2)$. In particular, any statement like the SWIC (or our main theorem) makes a bridge between the local topological types (C, p_i) and the global degree d . Computations show that for general $S^3_{-d}(K)$ (when the pair (K, d) has no algebraic realization as above) such identities cannot be expected. Hence such identities provide criterions for the algebraic realizations of a degree d curve with given local singularities.

3.0.1. The proof splits into two parts. The first part, done in [44], is valid for *any* surgery 3-manifold $S^3_{-d}(K)$. It provides $\mathbb{H}^*(S^3_{-d}(K))$, where K is a connected sum of algebraic knots in S^3 , and d is an arbitrary positive integer (and we do not assume the existence of a degree d curve with local types $K_i \subset S^3$, nor even the identity $\sum_i \mu_i = (d - 1)(d - 2)$). The cohomology is described completely in terms of the semigroups $\{S_i\}_{i=1}^\nu$ of the algebraic knots $K_i \subset S^3$. The construction uses the ‘reduction theorem’ of [17], which describes the lattice cohomology (defined a priori in the lattice L) in the first quadrant of a lattice of rank ν . The formula for $eu(\mathbb{H}^0)$ is the following. For any $n \in \mathbb{Z}_{\geq 0}$ set

$$\Sigma_n := \left\{ (\beta_1, \dots, \beta_\nu) \in (\mathbb{Z}_{\geq 0})^\nu : \sum_i \beta_i = n + 1 \right\},$$

and for any $(\beta_1, \dots, \beta_\nu) \in (\mathbb{Z}_{\geq 0})^\nu$ define the weight

$$\overline{W}(\beta_1, \dots, \beta_\nu) := \sum_{i=1}^\nu |\{k_i \notin S_i : k_i \geq \beta_i\}|.$$

Then, by [44, 6.1.15] (see also Remark 6.1.19, loc. cit.),

$$eu(\mathbb{H}^0(S^3_{-d}(K))) = \sum_{j \geq 0} \min \{ \overline{W} \text{ restricted to } \Sigma_{jd} \}. \tag{3.0.2}$$

The sum in (3.0.2) is finite: since $\{k_i \notin S_i : k_i \geq \mu_i\} = \emptyset$, if $jd + 1 \geq \sum_i \mu_i$ we get $\min(\{\overline{W}|_{\Sigma_{jd}}\}) = 0$. For example, if $\sum_i \mu_i = (d - 1)(d - 2)$, then for $j \geq d - 2$ we get a zero contribution.

Remark 3.0.3. Though $\mathbb{H}^0(\gamma)$ is not mentioned in [44], in its Section 6 one can see clearly that $\min_{\gamma} eu(\mathbb{H}^0(\gamma)) = eu(\mathbb{H}^0)$ even in this general topological context; nevertheless, we do not need this—in our case it will follow automatically from the estimate of the second part.

3.0.4. The second ingredient is the main result of [5] (as a possible answer to the conjectures of [12]), and it is valid only in the presence of the algebraic realization of $S^3_{-d}(K)$. Its proof uses surgery properties of the d -invariant of Heegaard–Floer theory.

[5, Theorem 5.4] reads as follows. If for any $(\beta_1, \dots, \beta_v) \in \mathbb{Z}^v$ one writes

$$\overline{W}^*(\beta_1, \dots, \beta_v) := \sum_{i=1}^v |\{k_i \in \mathbb{Z} \setminus S_i : k_i \geq \beta_i\}|,$$

then

$$\min \{ \overline{W}^* \text{ restricted to } \Sigma_{jd} \} = (j - d + 1)(j - d + 2)/2. \tag{3.0.5}$$

Since $S_i \subset \mathbb{Z}_{\geq 0}$, the minimum in (3.0.5) is realized for $(\beta_1, \dots, \beta_v) \in (\mathbb{Z}_{\geq 0})^v$, hence the minima in (3.0.5) and (3.0.2) agree. (Indeed, if $\beta_1 < 0$ and $\beta_v > 0$, then $\overline{W}^*(\beta_1, \dots, \beta_v) \geq \overline{W}^*(\beta_1 + 1, \dots, \beta_v - 1)$.) Hence (3.0.5) and (3.0.2) combined give

$$eu(\mathbb{H}^0(S^3_{-d}(K))) = \sum_{j=0}^{d-2} j(j + 1)/2 = d(d - 1)(d - 2)/6 = p_g.$$

This together with inequalities (2.3.2) and (2.3.5) ends the proof (cf. part (a) of the theorem).

For us, in fact, the main geometric meaning of the ‘arithmetical result’ of [5] is exactly the statement of Theorem 2.5.1(b).

4. Preliminaries on Newton non-degenerate singularities

4.1. The Newton boundary [16]

For any set $S \subset \mathbb{N}^3$ denote by $\Gamma_+(S) \subset \mathbb{R}^3$ the convex closure of $\bigcup_{p \in S} (p + \mathbb{R}_+^3)$. The collection of all boundary faces of $\Gamma_+(S)$ is denoted by \mathcal{F} , while the set of compact faces of $\Gamma_+(S)$ by \mathcal{F}_c . By definition, the Newton boundary (or diagram) $\Gamma(S)$ associated with S is the union of compact boundary faces of $\Gamma_+(S)$. Let $\Gamma_-(S)$ denote the cone with base $\Gamma(S)$ and vertex 0.

Let $f: (\mathbb{C}^3, 0) \rightarrow (\mathbb{C}, 0)$ be an analytic function germ defined by a convergent power series $\sum_p a_p z^p$, where $p = (p_1, p_2, p_3)$ and $z^p := z_1^{p_1} z_2^{p_2} z_3^{p_3}$. By definition, the *Newton boundary* $\Gamma(f)$ of f is $\Gamma(\text{supp}(f))$, where $\text{supp}(f) = \{p : a_p \neq 0\}$, and we write $\Gamma_{\pm}(f)$ for $\Gamma_{\pm}(\text{supp}(f))$. The *Newton principal part* of f is $\sum_{p \in \Gamma(f)} a_p z^p$. Similarly, for any q -dimensional face Δ of $\Gamma(f)$, set $f_{\Delta}(z) := \sum_{p \in \Delta} a_p z^p$. We say that f is *non-degenerate* on Δ if the system of equations $\partial f_{\Delta} / \partial z_1 = \partial f_{\Delta} / \partial z_2 = \partial f_{\Delta} / \partial z_3 = 0$ has no solution in $(\mathbb{C}^*)^3$. When f is non-degenerate on every q -face of $\Gamma(f)$, we say (after Kouchnirenko [16]) that f has a *non-degenerate Newton principal part*. In what follows we assume that f has this property. Moreover, we also assume that f defines an isolated singularity at the origin. This can be characterized by $\Gamma(f)$ as follows (cf. [16, 1.13(ii)] or [6, 2.1]):

$$\Gamma(f) \text{ has a vertex on every coordinate plane, and it has a vertex at most 1 away from any chosen coordinate axis.} \tag{4.1.1}$$

Nevertheless, we will not assume that $\Gamma(f)$ is ‘convenient’ (recall that $\Gamma(f)$ is *convenient* if it intersects all the coordinate axes). In fact, if f is not convenient, there are several ways to complete the diagram to a convenient one without modifying the equisingularity type. Hence, in general, several diagrams might produce the same equisingularity type. From all possible diagrams we choose a *minimal one* (with respect to inclusion; for its existence and ‘almost unicity’ see [6, §3]). This minimal diagram, in general, is not convenient.

Furthermore, as always in this note, we assume that the link M of $(X, 0) = \{f = 0\}$ is a *rational homology sphere*. This, in terms of the Newton boundary $\Gamma(f)$, reads as follows (cf. [53]):

$$M \text{ is a rational homology sphere} \Leftrightarrow \Gamma(f) \cap \mathbb{N}_{>0}^3 = \emptyset. \tag{4.1.2}$$

Lemma 4.1.3 ([6, §2.3]). *Under the above assumptions, if a face of $\Gamma(f)$ is not a triangle then it is a trapezoid. Up to permuting coordinates, its vertices are: $(p, 0, n)$, $(0, q, n)$, $(r_1, r_2 + tq, 0)$ and $(r_1 + tp, r_2, 0)$, where $p, q > 0$, $\text{gcd}(p, q) = 1$, $t \geq 1$ and $r_1, r_2 \geq 0$.*

For any face of the diagram at most one edge might have inner lattice points, and if an edge has inner lattice points then that edge sits in a coordinate plane.

We will denote by $\langle p, q \rangle = \sum_i p_i q_i$ the standard scalar product on \mathbb{R}^3 . Let N_{Δ} be the primitive normal vector of a 2-face $\Delta \in \mathcal{F}$ oriented so that $\langle N_{\Delta}, (1, 1, 1) \rangle > 0$. In the case of two adjacent faces Δ and ∇ from \mathcal{F} , we write $t_{\Delta, \nabla} - 1$ for the number of interior lattice points of the common edge. If both faces are compact then $t_{\Delta, \nabla} = 1$ by (4.1.2). Furthermore, let $n_{\Delta, \nabla}$ be the greatest common divisor of the 2-minors of the vectors N_{Δ} and N_{∇} .

4.2. Oka’s algorithm for G

Let $f: (\mathbb{C}^3, 0) \rightarrow (\mathbb{C}, 0)$ be a germ as in 4.1. We recall the combinatorial algorithm of M. Oka (based on toric resolution), which provides a dual resolution graph G of $(X, 0)$ from $\Gamma(f)$ (cf. [49, Theorem 6.1]).

4.2.1. The algorithm. The graph G is a subgraph of a larger graph \tilde{G} , whose construction is the following. We start with a set of vertices, each of them corresponding to a face from \mathcal{F} (we will call them *face vertices*). Consider two adjacent faces Δ and ∇ from \mathcal{F} . Then we connect the corresponding vertices by $t_{\Delta,\nabla}$ copies of the following chain.


If $n_{\Delta,\nabla} > 1$ then let $0 < c_{\Delta,\nabla} < n_{\Delta,\nabla}$ be the unique integer for which

$$N_{\Delta,\nabla} := (N_{\nabla} + c_{\Delta,\nabla}N_{\Delta})/n_{\Delta,\nabla} \tag{4.2.2}$$

is an integral vector. Let us write $n_{\Delta,\nabla}/c_{\Delta,\nabla}$ as a (negative) continued fraction:

$$\frac{n_{\Delta,\nabla}}{c_{\Delta,\nabla}} = e_1 - \frac{1}{e_2 - \frac{1}{\dots - \frac{1}{e_k}}} \tag{4.2.3}$$

with each $e_i \geq 2$. Then the chain with the corresponding self-intersection numbers is



The left ends of all the $t_{\Delta,\nabla}$ copies of the chain (marked by Δ) are identified with the face vertex corresponding to Δ , and similarly for the right ends marked by ∇ .

If $n_{\Delta,\nabla} = 1$ then the chain consists of an edge connecting the vertices Δ and ∇ (we put $t_{\Delta,\nabla}$ such edges). Also, in this case we set $c_{\Delta,\nabla} := 0$ and $N_{\Delta,\nabla} := N_{\nabla}$.

Next, we compute the decoration b_{Δ} of any face vertex $\Delta \in \mathcal{F}_c$ from the equation

$$b_{\Delta}N_{\Delta} + \sum_{\nabla \in \mathcal{F}_{\Delta}} t_{\Delta,\nabla}N_{\Delta,\nabla} = 0, \tag{4.2.4}$$

where \mathcal{F}_{Δ} is the collection of all 2-faces of $\Gamma_+(f)$ adjacent to Δ .

In this way we obtain the graph \tilde{G} . Notice that the face vertices corresponding to *non-compact faces* are not decorated.

Proposition 4.2.5. (a) [49] *If we delete all the vertices corresponding to non-compact faces (and all the edges adjacent to them) we get a dual resolution graph G .*

(b) [6, 4.2.5] *Under the above choice of the ‘minimal’ Newton diagram (cf. 4.1), the graph G is the minimal good resolution graph. In particular, the nodes (vertices with valency $\delta_v \geq 3$) are exactly the face vertices associated with compact faces of the diagram.*

(c) [6, 3.3.11] *If Δ is a compact face and ∇ is an adjacent non-compact face, then $n_{\Delta,\nabla} > 1$. Hence, such an edge of Δ produces a non-trivial chain (leg) of G .*

We denote by $\{b_v\}_{v \in \mathcal{V}}$ the decorations (self-intersections) of the corresponding vertices.

4.2.6. The ‘extended’ graph G^e . Replace each edge of \tilde{G} connecting a face vertex \tilde{v} of $\tilde{G} \setminus G$ and another vertex w of G by an arrow with arrowhead a and supporting vertex w . Then G^e consists of G equipped with this type of arrowheads.

The set of vertices of G is denoted by \mathcal{V} , the set of arrowheads of G^e by \mathcal{A} , and we refer to $\mathcal{V}^e := \mathcal{V} \cup \mathcal{A}$ as the set of (arrowhead and non-arrowhead) vertices of G^e . Let \mathcal{N} be the set of (non-arrowhead) face vertices of G^e ; these are the nodes in both graphs G and G^e .

To each $v \in \mathcal{V}^e$ we associate a vector N_v in $\mathbb{Z}_{\geq 0}^3$. If v is a face vertex of G corresponding to Δ then $N_v = N_\Delta$. If $a \in \mathcal{A}$ corresponds to $\Delta \in \mathcal{F} \setminus \mathcal{F}_c$ then $N_a = N_\Delta$. All the other vectors are determined in a unique way by the following identities (see [6]):

Fix $v \in \mathcal{V}$, and set $\mathcal{V}_v^e := \{w \in \mathcal{V}^e : w \text{ adjacent to } v \text{ in } G^e\}$ and $\mathcal{V}_v := \mathcal{V}_v^e \cap \mathcal{V}$. Then

$$b_v N_v + \sum_{w \in \mathcal{V}_v^e} N_w = 0. \tag{4.2.7}$$

(In this procedure it helps to know that the vector associated with a neighbor of a face vertex v corresponding to Δ on the chain in the direction w corresponding to ∇ is $N_{\Delta, \nabla}$.)

If v is on a chain connecting the face vertices w_1 and w_2 , then $N_v = r_1 N_{w_1} + r_2 N_{w_2}$ with $r_1, r_2 \in \mathbb{Q}_{>0}$. Since for any $v \in \mathcal{N}$ all the entries of N_v are strictly positive, the same property remains true for all $v \in \mathcal{V}$. By (4.2.7) (which typically characterizes the multiplicities of the function encoded in an embedded resolution graph), or by [49],

$$\text{the multiplicity of } z_i \text{ along } E_v \text{ is the } i\text{-th coordinate of } N_v \text{ (} v \in \mathcal{V}\text{)}. \tag{4.2.8}$$

In fact, if we decorate the (arrowhead and non-arrowhead) vertices of G^e by the i -th coordinate of the vectors N_v , then we get the embedded resolution graph of the germ z_i on $(X, 0)$, where the arrowheads with zero decorations can be deleted, while the arrowheads with positive decorations represent strict transforms of $z_i = 0$ which are not necessarily transversal to their supporting curves (the corresponding decoration is the intersection multiplicity of the strict transform with the supporting curve). This decoration can be larger than 1 if the corresponding non-compact face is not a coordinate plane.

- Definition 4.2.9.** (i) For each $v \in \mathcal{V}^e$ we denote by ℓ_v the linear function $\langle N_v, \cdot \rangle$.
 (ii) If $l = \sum_{v \in \mathcal{V}} l_v E_v \in L$, then we write $m_v(l) = l_v$. We extend this by setting $m_v(l) := -1$ for any $v \in \mathcal{V}^e \setminus \mathcal{V}$. (For a motivation of the value -1 see the proof of Lemma 4.3.4.)
 (iii) For any $l \in L$ and $v \in \mathcal{V}^e$ we define the half-space $H_v^\geq(l) := \{p \in \mathbb{R}^3 \mid \ell_v(p) \geq m_v(l)\}$, and write $H_v^=(l)$ for its boundary plane. Moreover, we define the *rational polytope*

$$\Gamma_+^e(l) := \bigcap_{v \in \mathcal{V}^e} H_v^\geq(l).$$

4.3. Divisorial valuations, weights and the canonical cycle

Let f be as in (4.1) and L the lattice of rank $|\mathcal{V}|$ associated with G , with E_v as its basis. The resolution with exceptional curve $E = \bigcup_v E_v$ is denoted by \tilde{X} (as in 2.1.4).

- Definition 4.3.1.** (a) For a non-empty finite set S in $\mathbb{Z}_{\geq 0}^3$ and $v \in \mathcal{V}$ define $\text{wt}_v(S) = \min_{p \in S} \ell_v(p)$ and $\text{wt}(S) := \sum_{v \in \mathcal{V}} \text{wt}_v(S) E_v \in L$. For $0 \neq h \in \mathbb{C}\{z\}$ set $\text{wt}_v(h) = \text{wt}_v(\text{supp}(h))$ and $\text{wt}(h) = \text{wt}(\text{supp}(h))$.

(b) For any $\bar{h} \in \mathcal{O}_{X,0}$, denote by $\text{div}_v(\bar{h})$ the order of vanishing on E_v of the pullback of \bar{h} to \tilde{X} , and set $\text{div}(\bar{h}) = \sum_v \text{div}_v(\bar{h})E_v$. If $h \in \mathbb{C}\{z\}$, let $\text{div}(h) = \text{div}(h|_X)$.

Remark 4.3.2. The functions wt_v are order functions. For any $p \in \mathbb{N}^3$, we have $\text{wt}(z^p) = \text{div}(z^p)$ (cf. 4.2.8). Since for all $h_1, h_2 \in \mathbb{C}\{z\}$ with $\text{supp}(h_1) \cap \text{supp}(h_2) = \emptyset$ one has $\text{wt}(h_1 + h_2) = \min\{\text{wt}(h_1), \text{wt}(h_2)\}$, and the div_v are valuations, we get $\text{wt}(h) \leq \text{div}(h)$ for any $h \in \mathbb{C}\{z\}$.

Lemma 4.3.3 ([49, Theorem (9.1)]). *The (anti)canonical divisor $Z_K = -K \in L$ satisfies*

$$Z_K - E = \text{wt}(f) - \text{wt}(z_1 z_2 z_3).$$

Lemma 4.3.4. $\Gamma_+^e(Z_K - E) = \Gamma_+(f) - (1, 1, 1)$.

Proof. Let us analyse the non-compact faces of $\Gamma_+(f)$. Up to a permutation of coordinates they have the form $z_1 + az_2 = a$ for some $a \in \mathbb{Z}_{\geq 0}$. This follows from (4.1.1) or from [6, 3.1.2]. This plane shifted by $(1, 1, 1)$ gives the plane $z_1 + az_2 = -1$, a fact which explains Definition 4.2.9 for $v \notin \mathcal{V}$ too. Otherwise the statement follows from Lemma 4.3.3. □

5. The proof of Theorem 2.5.1(c)

Since part (a) covers the rational hypersurfaces, we assume that $(X, 0)$ is not rational. Let \tilde{X} be a minimal good resolution (provided by Prop. 4.2.5); we write \mathcal{O} for $\mathcal{O}_{\tilde{X}}$. Let G be the corresponding resolution graph. By Lemma 2.1.5 we get $Z_K > E$.

5.1. The sequence $\{z_i\}_i$ (preliminaries)

5.1.1. From the definition of path cohomology, by paragraphs 2.3.3 and 2.5.3, we have to construct a sequence $\{l_i\}_{i=0}^t$ with $l_i \in L$ such that $l_0 = 0, l_t = Z_K, l_{i+1} = l_i + E_{v(i)}$ with

$$p_g = \sum_{i=0}^{t-1} (h^1(\mathcal{O}_{l_{i+1}}) - h^1(\mathcal{O}_{l_i})) = \sum_{i=0}^{t-1} h^1(\mathcal{O}_{E_{v(i)}}(-l_i)).$$

5.1.2. We need the dual picture. Take a sequence $\{z_i\}_{i=0}^t$ with $z_i \in L$ where $z_0 = 0, z_t = Z_K, z_{i+1} = z_i + E_{v(i)}$. Then the pair of sheaves $\mathcal{O}(-z_{i+1}) \hookrightarrow \mathcal{O}(-z_i)$ gives an exact sequence

$$0 \rightarrow H^0(\tilde{X}, \mathcal{O}(-z_{i+1})) \rightarrow H^0(\tilde{X}, \mathcal{O}(-z_i)) \rightarrow H^0(\mathcal{O}_{E_{v(i)}}(-z_i)) \rightarrow \dots \quad (5.1.3)$$

Since $H^0(\mathcal{O})/H^0(\mathcal{O}(-Z_K)) \simeq H^0(\mathcal{O}_{Z_K}) = \mathbb{C}^{p_g}$, and since we have $h^0(\mathcal{O}_{E_{v(i)}}(-z_i)) = \max\{0, 1 - (E_{v(i)}, z_i)\}$, it follows that

$$p_g = \sum_{i=0}^{t-1} \dim \frac{H^0(\tilde{X}, \mathcal{O}(-z_i))}{H^0(\tilde{X}, \mathcal{O}(-z_{i+1}))} \leq \sum_{i=0}^{t-1} \max\{0, 1 - (E_{v(i)}, z_i)\}. \quad (5.1.4)$$

The dual statement of 5.1.1 requires the existence of $\{z_i\}_i$ with equality in (5.1.4) for all i . The duality is realized by $z_i := Z_K - l_{t-i}$ and Serre duality.

[Indeed, if $z_{i+1} = z_i + E_{v(i)}$, $l_{j+1} = l_j + E_{u(j)}$, $z_i = Z_K - l_{t-i}$, then with $j := t - 1 - i$ one has $u(j) = v(i)$ and $h^1(\mathcal{O}_{E_{u(j)}}(-l_j)) = h^0(\mathcal{O}_{E_{v(i)}}(-z_i))$ by Serre duality.]

Since $H^1(\mathcal{O}_{E_j}) = H^1(\mathcal{O}_E) = 0$, in fact, in 5.1.1 we need a sequence $\{l_i\}_i$ which starts with $l_0 = E$ and ends with $l_t = Z_K$ (and otherwise is as in 5.1.1).

By dual considerations (since $H^0(\mathcal{O}(-Z_K)) \hookrightarrow H^0(\mathcal{O}(-Z_K + E))$ is an isomorphism) it is enough to construct a sequence $\{z_i\}_{i=0}^t$ which starts with $z_0 = 0$ and ends with $z_t = Z_K - E$ (and otherwise is as above). This is what we will do.

5.1.5. The plan of the proof. To a sequence $\{z_i\}_i$ of elements of L (with $z_0 = 0$, $z_t = Z_K - E$, $z_{i+1} = z_i + E_{v(i)}$) we associate the sequence $\{\Gamma_+^e(z_i)\}_i$ of polytopes. Note that $\Gamma_+^e(z_t) = \Gamma_+(f) - (1, 1, 1) \subset \Gamma_+^e(z_0) \subset (\mathbb{R}_{\geq -1})^3$. Clearly $\Gamma_+^e(z_{i+1}) \subset \Gamma_+^e(z_i)$. This filtration realizes a partition of the lattice points $(\Gamma_-(f) - (1, 1, 1)) \cap \mathbb{Z}_{\geq 0}^3$ by

$$P_i := (\Gamma_+^e(z_i) \setminus \Gamma_+^e(z_{i+1})) \cap \mathbb{Z}_{\geq 0}^3. \tag{5.1.6}$$

Then, for the sequence $\{z_i\}_i$ provided by the algorithm 5.3.1 we show that for $0 \leq i < t$,

$$\max\{0, 1 - (E_{v(i)}, z_i)\} \leq |P_i|, \tag{5.1.7}$$

$$|P_i| \leq \dim \frac{H^0(\tilde{X}, \mathcal{O}(-z_i))}{H^0(\tilde{X}, \mathcal{O}(-z_{i+1}))}. \tag{5.1.8}$$

These two facts together with (5.1.4) show that in (5.1.4) we must have equality.

Note that based on (5.1.7) and (5.1.4) together with the result of Merle–Teissier [29], which says that $|\Gamma_-(f) \cap \mathbb{Z}_{\geq 0}^3| = p_g$, or in the above terms $\sum_i |P_i| = p_g$, we could already conclude our proof. Nevertheless, by providing an independent argument for the additional (5.1.8), besides the proof of our theorem we reprove the Merle–Teissier result as well.

Some of the steps can be analysed easily.

Lemma 5.1.9. *Assume that along the sequence one has $(E_{v(i)}, z_i) > 0$ for some i . Then $P_i = \emptyset$, hence (5.1.7) is valid. Moreover, in (5.1.8) one has equality (with both sides zero).*

Proof. Assume that $P_i \neq \emptyset$, set $p \in P_i$ and $v := v(i)$. Then $\ell_v(p) = m_v(z_i)$ and $\ell_w(p) \geq m_w(z_i)$ for every $w \in \mathcal{V}$. Moreover, since all the entries of p are non-negative, $\ell_w(p) \geq 0$ for every $w \in \mathcal{V}^e \setminus \mathcal{V}$ as well. Therefore, by (4.2.7), $0 = (b_v \ell_v + \sum_{w \in \mathcal{V}_v^e} \ell_w)(p) \geq b_v m_v(z_i) + \sum_{w \in \mathcal{V}_v} m_w(z_i) = (E_v, z_i) > 0$, a contradiction. For the second statement use (5.1.3). □

5.2. The operation $l \mapsto c(l)$ and the ratio test

In the construction of the sequence $\{z_i\}_i$ a part of the ‘easy’ steps are provided by the next ‘completion operation’.

If $l = \sum_v m_v E_v \in L$, then we define the support of l as $|l| := \{v \in \mathcal{V} : m_v \neq 0\}$.

In the next proposition the meaning of the condition involved is motivated by the adjunction formula $(E_v, Z_K - E) = 2 - \delta_v$. Note that $\delta_v \leq 2$ if $v \notin \mathcal{N}$.

Proposition 5.2.1. (A) Fix $l \in L$. Then there exists a unique element $c(l) \in L$ with

- (1) $m_n(c(l)) = m_n(l)$ for all $n \in \mathcal{N}$;
- (2) $(c(l), E_v) \leq 2 - \delta_v$ for every $v \notin \mathcal{N}$;
- (3) $c(l)$ is minimal with properties (1) and (2).

(Clearly, by (1), $c(l)$ depends only on $\{m_n(l)\}_{n \in \mathcal{N}}$.)

- (B) Let l be as in (A), and $z \in L$ another cycle such that $z \leq c(l)$ and $m_n(z) = m_n(l)$ for all $n \in \mathcal{N}$. We construct a ‘computation sequence’ $\{x_j\}_{j=0}^T$ starting with $x_0 = z$ as follows. First, one takes $x_0 = z$. Then, if x_j is already constructed, and it does not satisfy (A)(2), that is, there exists v with $(x_j, E_v) > 2 - \delta_v$, then $x_{j+1} = x_j + E_v$ for such v . If x_j satisfies (A)(2) then we stop and $j = T$. Then for any such computation sequence, $x_T = c(l)$.
- (C) If $l_i \in L$ are such that $m_n(l_1) \leq m_n(l_2)$ for all $n \in \mathcal{N}$, then $c(l_1) \leq c(l_2)$.
- (D) If G is not an A_n -graph, then $c(0) = 0$. (For A_n one has $c(0) = -E$.)
- (E) $c(Z_K - E) = Z_K - E$.

Proof. The proof of (A)–(B) is an alteration of Artin’s proof of the existence of the fundamental cycle [4] and of the existence of the Laufer computation sequence providing this cycle [19]. The major steps are the following. First, note that by the negative definiteness of the intersection form there exists a cycle with properties (A)(1–2). Then one shows (as in [4]) that if $c'(l)$ and $c''(l)$ satisfy (A)(1–2) then $\min\{c'(l), c''(l)\}$ also satisfies them, hence there exists a unique minimal element $c(l)$ satisfying (A)(1–2). For (B), one checks by induction that $x_j \leq c(l)$ for every j . Hence the sequence must stop and $x_T \leq c(l)$. Since x_T satisfies (A)(1–2), and $c(l)$ is minimal with this property, $x_T = c(l)$.

(C) $c(l_2) - (l_2 - l_1)$ satisfies (A)(1–2) and restricted on \mathcal{N} is l_1 , hence $c(l_1) \leq c(l_2) - (l_2 - l_1)$.

(D) It is an elementary arithmetical verification on the chains of $G \setminus \mathcal{N}$.

(E) Since $Z_K - E$ satisfies the conditions (A)(1–2), $l := Z_K - E - c(Z_K - E) \geq 0$ by (A)(3). But $(l, E_v) \geq 0$ for all v in the support of l , hence by the negative definiteness of G we have $l \leq 0$ too. Hence $l = 0$. \square

Definition 5.2.2 (The ratio test). We fix $l \in L$, and we consider the ratio $R(v) := m_v(l)/m_v(Z_K - E)$ for every $v \in \mathcal{V}$. We say that the ratio test for l chooses the vertex $v \in \mathcal{V}$ if $R(v) = \min\{R(w) : w \in \mathcal{V}\}$ and $R(v) < R(w) < \infty$ for at least one adjacent vertex w of v . (If $m_v(Z_K - E) = 0$ then $R(v) = \infty$, which is ‘larger than any real number’.)

Lemma 5.2.3. Assume that for some effective l supported on \mathcal{N} one has $0 < c(l) < Z_K - E$. Then the ratio test for $c(l)$ always makes a choice, that is, it is not possible that $v \mapsto R(v)$ is a constant on $|Z_K - E|$. Moreover, the chosen v is in the support of $Z_K - E - c(l)$.

Proof. Since the support of $Z_K - E$ is connected (cf. 2.1.5), if $R(v)$ is constant on this support then $c(l) = r(Z_K - E)$ for some $r \in (0, 1)$ (in the complement of this support both $c(l)$ and $Z_K - E$ are zero). Then for any w with $\delta_w = 1$ we would get $r = r \cdot (Z_K - E, E_w) = (c(l), E_w) \in \mathbb{Z}$, a contradiction. Hence, a choice v exists. Moreover, $m_v(c(l)) = m_v(Z_K - E)$ would imply $1 = R(v) = \min_w\{R(w)\}$, or $c(l) \geq Z_K - E$. \square

5.3. The algorithm for $\{z_i\}_i$

Definition 5.3.1. The sequence $\{z_i\}_{i=0}^t$ is constructed in the following steps.

S1. $\bar{z}_0 = 0$.

S2. Assume that \bar{z}_i is already constructed, and $\bar{z}_i < Z_K - E$. If $i = 0$ then we choose an arbitrary E_v in the support of $Z_K - E$ and we write $E_{v(0)} := E_v$. If $i > 0$, then $\bar{z}_i > 0$, and the ratio test for \bar{z}_i makes a choice (cf. Lemma 5.2.3), say $v(i)$. Then (in both cases) set $\bar{z}_{i+1} = c(\bar{z}_i + E_{v(i)})$. Note that by Lemma 5.2.3 and Proposition 5.2.1 one has $\bar{z}_i + E_{v(i)} \leq \bar{z}_{i+1} \leq Z_K - E$.

S3. If $\bar{z}_{i+1} < Z_K - E$ then run Step S2 again. If $\bar{z}_i = Z_K - E$ then stop and set $\bar{t} := i$.

Note that different choices might produce different sequences $\{\bar{z}_i\}_{i=1}^{\bar{t}}$.

For any $i \geq 0$ one has $c(\bar{z}_i + E_{v(i)}) \geq \bar{z}_i + E_{v(i)}$. For $i = 0$ this follows from Proposition 5.2.1(D), while for $i > 0$ from part (C). Hence by 5.2.1(B) there exists an ‘intermediate’ sequence between $\bar{z}_i + E_{v(i)}$ and \bar{z}_{i+1} . Completing the above sequence with these intermediate steps we get the desired sequence $\{z_i\}_{i=0}^t$ for which $z_{i+1} - z_i$ is always a base element. Along these intermediate steps (cf. Lemma 5.1.9), the inequalities (5.1.7)–(5.1.8) hold with equality, since $(E_{v(i)}, x_j) > 2 - \delta_{v(i)} \geq 0$.

The very first step $0 = z_0 \rightsquigarrow z_1 = E_{v(0)}$ is also easy: $(E_{v(0)}, z_0) = 0$, the factor space $H^0(\mathcal{O})/H^0(\mathcal{O}(-E_{v(0)}))$ is 1-dimensional, and since for any $v \in \mathcal{V}$ all the entries of N_v are positive, $(N_v, p) > 0$ for all $\mathbb{Z}_{\geq 0}^3 \setminus (0, 0, 0)$. Hence $P_0 = \{(0, 0, 0)\}$, and it has cardinality 1.

The next lemma analyzes the step $\bar{z}_i \rightsquigarrow \bar{z}_i + E_{v(i)}$ ($i > 0$).

Lemma 5.3.2. $\delta_{v(i)} = 2$ cannot happen. If $\delta_{v(i)} = 1$ then $(\bar{z}_i, E_{v(i)}) = 1$.

Proof. The choice of $v(i)$ implies

$$\frac{(\bar{z}_i, E_{v(i)})}{\sum m_w(\bar{z}_i)} > \frac{(Z_K - E, E_{v(i)})}{\sum m_w(Z_K - E)} = \frac{2 - \delta_{v(i)}}{\sum m_w(Z_K - E)},$$

where the sums run over $w \in \mathcal{V}_{v(i)}$. Then use the fact that \bar{z}_i is c of an element. □

5.3.3. This shows (using again Lemma 5.1.9) that what remains to verify is the validity of (5.1.7)–(5.1.8) in the step $\bar{z}_i \rightsquigarrow \bar{z}_i + E_{v(i)}$ whenever $i > 0$, $v(i)$ is a face vertex (that is, node with $\delta_{v(i)} \geq 3$) and $(\bar{z}_i, E_{v(i)}) \leq 0$. In what follows, we assume all these facts.

It is also convenient to define, for any $n \in \mathcal{N}$ and cycle $l > 0$,

$$F_n(l) := \left(\bigcap_{w \in \mathcal{V} \setminus n} H_w^{\geq}(l) \right) \cap H_n^{\leq}(l) \quad \text{and} \quad F_n^{nb}(l) := \left(\bigcap_{w \in \mathcal{V}_n} H_w^{\geq}(l) \right) \cap H_n^{\leq}(l).$$

Let C_n (resp. C_n^{nb}) be the real cone over $F_n(Z_K - E)$ (resp. $F_n^{nb}(Z_K - E)$) with vertex 0.

The definition of $F_n(l)$ should be compared with the definition of the face $F_n^e(l)$ of $\Gamma_+^e(l)$ which sits in $H_n^{\leq}(l)$, defined by $(\bigcap_{w \in \mathcal{V}^e \setminus n} H_w^{\geq}(l)) \cap H_n^{\leq}(l)$ (which, having extra equations indexed by $\mathcal{V}^e \setminus \mathcal{V}$, is a subset of $F_n(l)$). Also, $F_n(l) \subset F_n^{nb}(l)$ since $\mathcal{V}_n \subset \mathcal{V} \setminus n$.

Lemma 5.3.4. Fix $n \in \mathcal{N}$. Then $F_n^{nb}(Z_K - E) = F_n(Z_K - E) = F_n^e(Z_K - E)$. In particular, $C_n^{nb} = C_n$.

Proof. With simplified notation we have $F_n^{nb} \supset F_n \supset F_n^e$. Next note that $\mathcal{V}_n^e = \mathcal{V}_n$ (cf. 4.2.5(c)), that is, if $n' \in \mathcal{V}^e \setminus \mathcal{V}$ is connected to n by a chain to n , then this chain is non-empty.

If n' is another face vertex of \mathcal{V} connected to n by a chain with vertices v_1, \dots, v_k , then all the intersections $H_n^-(Z_K - E) \cap H_{v_j}^-(Z_K - E)$ agree with the intersection $H_n^-(Z_K - E) \cap H_{n'}^-(Z_K - E)$. This follows from the adjunction relation and identities (4.2.7). The same is true if $n' \in \mathcal{V}^e \setminus \mathcal{V}$, and $H_{n'}^-(Z_K - E)$ is $\{p : \ell_{n'}(p) = -1\}$ (compatibly with the convention $m_{n'} = -1$, cf. Definition 4.2.9(ii)).

Therefore, F_n^{nb} equals the polygon of H_n^- cut out by the intersections of type $H_{n'}^{\geq}$, where n' are as in the previous paragraph (neighbor extended face-vertices). But, by Lemma 4.3.4, this equals the corresponding face of $\Gamma_+(f) - (1, 1, 1)$, and also F_n^e , since the n -face of $\Gamma_+(f)$ is also cut out by the equations given by neighbor face-vertices n' . □

5.3.5. Let us consider again the step $\bar{z}_i \rightsquigarrow \bar{z}_i + E_{v(i)}$ of the algorithm with restrictions as in 5.3.3.

We still denote $P_i := (\Gamma_+^e(\bar{z}_i) \setminus \Gamma_+^e(\bar{z}_i + E_{v(i)})) \cap \mathbb{Z}_{\geq 0}^3$.

In the affine plane $H_i := H_{v(i)}^-(\bar{z}_i)$ there are several polygons:

- $F_i := F_{v(i)}(\bar{z}_i)$ cut out by the half-spaces indexed by $\mathcal{V} \setminus v(i)$;
- $F_i^{nb} := F_{v(i)}^{nb}(\bar{z}_i)$ cut out by the half-spaces indexed by the ‘neighbors’ $\mathcal{V}_{v(i)}$;
- $F_i^{cn} := C_{v(i)} \cap H_i$ cut out by the cone over $F_{v(i)}(Z_K - E)$.

By Lemma 5.3.4, the cone $C_{v(i)}$ is given by the inequalities

$$\frac{\ell_w(p)}{m_w(Z_K - E)} \geq \frac{\ell_{v(i)}(p)}{m_{v(i)}(Z_K - E)} \quad \text{for all } w \in \mathcal{V}_{v(i)}.$$

This restricted to $H_i = \{l_{v(i)}(p) = m_{v(i)}(\bar{z}_i)\}$ transforms into

$$\ell_w(p) \geq \frac{m_{v(i)}(\bar{z}_i)}{m_{v(i)}(Z_K - E)} \cdot \frac{m_w(Z_K - E)}{m_w(\bar{z}_i)} \cdot m_w(\bar{z}_i) = r_{i,w} \cdot m_w(\bar{z}_i) \quad (w \in \mathcal{V}_{v(i)}). \quad (5.3.6)$$

By the ratio test $r_{i,w} \leq 1$. The inequalities which cut out F_i^{nb} in H_i are $\ell_w(p) \geq m_w(\bar{z}_i)$ ($w \in \mathcal{V}_{v(i)}$), hence they are more restrictive than those from (5.3.6). Hence $F_i^{nb} \subset F_i^{cn}$.

For technical purposes we need another polygon F_i^{cn-} satisfying $F_i^{nb} \subset F_i^{cn-} \subset F_i^{cn}$.

First we associate to any $w \in \mathcal{V}_{v(i)}$ an integer $\epsilon_j \in \{0, 1\}$. Fix w and let I_w be the edge of F_i^{cn} determined by $\ell_w(p) = r_{i,w} \cdot m_w(\bar{z}_i)$. Then $\epsilon_w = 0$ except if the following happens: I_w contains at least one lattice point (hence $r_{i,w} \cdot m_w(\bar{z}_i) \in \mathbb{Z}$), and $r_{i,w} \cdot m_w(\bar{z}_i) < m_w(\bar{z}_i)$. In this case $\epsilon_w = 1$. We define $F_i^{cn-} \subset H_i$ cut out by the inequalities $\ell_w(p) \geq r_{i,w} \cdot m_w(\bar{z}_i) + \epsilon_w$ for all $w \in \mathcal{V}_{v(i)}$. Hence

$$F_i \subset F_i^{nb} \subset F_i^{cn-} \subset F_i^{cn}. \quad (5.3.7)$$

The next lemma compares the lattice points of these polygons.

Lemma 5.3.8. $P_i = F_i \cap \mathbb{Z}_{\geq 0}^3 = F_i^{nb} \cap \mathbb{Z}_{\geq 0}^3 = F_i^{cn-} \cap \mathbb{Z}_{\geq 0}^3$.

Proof. The first equality follows from the fact that $\ell_{v(i)}$ takes integral values on the lattice points, hence all the lattice points of $\Gamma_+^e(\bar{z}_i) \setminus \Gamma_+^e(\bar{z}_i + E_{v(i)})$ are on the boundary face F_i .

For the other ones, via (5.3.7), it is enough to show that $F_i^{cn-} \cap \mathbb{Z}_{\geq 0}^3 \subset P_i$. Take $p_0 \in F_i^{cn-} \cap \mathbb{Z}_{\geq 0}^3$; we wish to show that $p_0 \in P_i$.

Run the algorithm from $z_0 = 0$ to $z_t = Z_K - E$. Replacing $\Gamma_+^e(z_k)$ by $\Gamma_+^e(z_{k+1})$, we say that we pass the lattice points in their difference. Along the algorithm we have to pass all the lattice points of $(\Gamma_+(f) - (1, 1, 1)) \cap \mathbb{Z}_{\geq 0}^3$, each of them exactly once. By the above discussion (see also 5.1.9), we pass a lattice point only at steps of type $\bar{z}_j \rightsquigarrow \bar{z}_j + E_{v(j)}$, where $v(j) \in \mathcal{N}$. Along this step we pass exactly the lattice points P_j . Moreover, by (5.3.7) applied for $v(j)$, these lattice points are situated in the cone $C_{v(j)}$.

Hence, if our chosen lattice point p_0 is situated only in the cone $C_{v(i)}$, then we can pass it only at step $v(i)$ by the plane H_i , hence it is in P_i .

Assume next that p_0 is situated at the intersection of two cones $C_{v(i)}$ and $C_{n'}$, hence on an edge I_w of F_i^{cn} (w being on the chain connecting $v(i)$ and n').

Then along the algorithm we pass p_0 either at step $v(i)$ by H_i , or before that step. In the first case $p_0 \in P_i$. We show that the second case cannot happen.

Indeed, if we pass p_0 at step $j < i$ with $v(j) = n'$, then $\ell_{v(j)}(p_0) = m_{v(j)}(\bar{z}_j)$. Since $\bar{z}_i \geq \bar{z}_j + E_{v(j)}$, one gets $\ell_{v(j)}(p_0) < m_{v(j)}(\bar{z}_i)$.

Claim. $\ell_{v(i)}(p_0) = m_{v(i)}(\bar{z}_i)$ and $\ell_{v(j)}(p_0) < m_{v(j)}(\bar{z}_i)$ implies $\ell_w(p_0) < m_w(\bar{z}_i)$.

Proof of Claim. Write $n = v(i)$. The Claim follows from the existence of positive rational numbers a and b with $(\dagger) \ell_w = a\ell_n + b\ell_{n'}$ and $(\ddagger) m_w \geq am_n + bm_{n'}$.

The numbers a and b are the following. Let (w, v_1, \dots, v_k) be the chain connecting n and n' . Let d and d' be the determinant of the chains (w, v_1, \dots, v_k) and (v_1, \dots, v_k) (where the determinant of the empty graph is 1). Then $a = d'/d$ and $b = 1/d$. This follows from weighted summation of identities (4.2.7) for (\dagger) and of inequalities 5.2.1(2) for (\ddagger) .

Hence, we have $\ell_w(p_0) < m_w(\bar{z}_i)$, which implies that $\epsilon_w = 1$, hence by the construction of F_i^{cn-} we get $p_0 \notin F_i^{cn-}$, a contradiction. \square

Remark 5.3.9. By Oka's algorithm 4.2.1, the greatest common divisor of the 2-minors of the 2×3 -matrix formed by N_v and N_w of two neighbor vertices (v, w) is one. Hence ℓ_w is a primitive linear function on H_i , and $F_i^{cn-} \cap \mathbb{Z}_{\geq 0}^3$ is obtained from $F_i^{cn} \cap \mathbb{Z}_{\geq 0}^3$ by eliminating $I_w \cap \mathbb{Z}_{\geq 0}^3$ with $\epsilon_w = 1$ (and no other points).

5.4. Polygons in an affine space

In this section we establish some combinatorial facts about triangles and trapezoids in affine planes of \mathbb{R}^3 . (For motivation see Lemma 4.1.3.)

Definition 5.4.1. A triangle or trapezoid Δ_{st} in an affine plane A_{st} is *standard* if its vertices are integral points, it contains no integral points in its interior and at most one of its edges contains integral points in its interior. If a standard polygon has an edge with integral points in its interior, call that side the *long* edge, and the others the *short* edges.

Otherwise, any edge may be referred to as long or short. In this way, all standard polygons Δ_{st} have a long edge. Let $t(\Delta_{st}) - 1$ be the number of interior lattice points of the long edge. (In the case of a trapezoid the long edge is automatically parallel to another edge; cf. Lemma 4.1.3.)

Let Δ_{st} be a standard polygon with edges $\{I_w\}_w$. To each edge I_w we associate an integral affine function $l_{st,w}$, which is constant along I_w , $l_{st,w}|_{\Delta_{st}} \geq l_{st,w}|_{I_w}$, and its restriction to A_{st} is primitive. ($l_{st,w}$ is *primitive* if its restriction $l_{st,w} : A_{st} \cap \mathbb{Z}^3 \rightarrow \mathbb{Z}$ is onto.) The restriction of $l_{st,w}$ to A_{st} is unique up to an additive integral constant.

5.4.2. Assuming that the link of the singularity defined by f is a rational homology sphere (hence (4.1.2) is valid), all faces of $\Gamma_+(f)$ are either standard triangles or standard trapezoids; cf. [6, 2.3] or 4.1 here. (There is only one exception to this, namely the Newton polygon of $z_1^{2a} + z_2^{2b} + z_3^{2c}$, a, b, c pairwise relative primes, in which case three edges contain interior lattice points. Since this is a weighted homogeneous Newton diagram, which case is covered by part (a) of Theorem 2.5.1 whose proof was already given, we can assume that our Newton diagram is not of this type. Alternatively, the interested reader might run the argument below slightly modified to check this case too.) Moreover, the edge of such a face which contains integral interior points is necessarily contained in a coordinate plane.

Accordingly, in our application and proof, the standard polygons are $\Delta_{st} = F_{v(i)}^{nb}(Z_K - E)$, which by Lemma 5.3.4 are the faces of $\Gamma_+(f)$ shifted by $-(1, 1, 1)$.

Lemma 5.4.3. *Consider $\Delta_{st} = F_{v(i)}^{nb}(Z_K - E)$ with long edge I_1 , and an adjacent short edge I_2 . Then the map $(l_{st,1}, l_{st,2}) : A_{st} \cap \mathbb{Z}^3 \rightarrow \mathbb{Z}^2$ is a \mathbb{Z} -affine isomorphism.*

Proof. Since Δ_{st} contains no integral points in its interior, neither does the parallelogram spanned by the two primitive vectors supported on its edges I_1 and I_2 . □

5.4.4. A polygon Δ in the affine plane A is a *small scalar translate* of a standard polygon Δ_{st} if

$$\text{there is a vector } V \in \mathbb{R}^3 \text{ and } r \in (0, 1) \text{ such that } \Delta = V + r \cdot \Delta_{st}. \tag{5.4.5}$$

Let Δ be a small scalar translate of Δ_{st} ; we denote its edges by the same symbols $\{I_w\}_w$ (I_1 corresponding to the long edge). Then for each w there exists a unique integral primitive affine function l'_w on A such that $l'_w|_{I_w}$ is constant with value in $(-1, 0]$, and $l'_w|_{\Delta} \geq l'_w|_{I_w}$. Additionally, we select $\epsilon_w \in \{0, 1\}$ (for motivation see 5.3.5) so that if a *short edge* contains a lattice point then ϵ_w might be 0 or 1, otherwise it is zero. Finally, set $l_w := l'_w - \epsilon_w$.

In our applications, F_i^{cn} is a small scalar translate whose equations $l'_w(p) \geq l'_w|_{I_w}$ are replaced by $l'_w(p) \geq l'_w|_{I_w} + \epsilon_w$ in order to obtain F_i^{cn-} .

Lemma 5.4.6. *With the above notation one has:*

- (a) *All the lattice points of Δ are on a line parallel to I_1 .*
- (b) *The function $t(\Delta_{st})l_1 + \sum_{w>1} l_w$ is constant (say c) on A . Moreover, $\max\{0, c+1\} = |(\Delta \setminus \bigcup_{\epsilon_w=1} I_w) \cap \mathbb{Z}^3|$.*

Proof. The proof splits into the cases when Δ_{st} is a triangle or when it is a trapezoid. First we fix the equations $l_{st,w}$ of the edges of Δ_{st} . We abbreviate $t := t(\Delta_{st})$. By Lemma 5.4.3, in the case of a triangle, we can take integral linear coordinates (x, y) on A_{st} so that $l_{st,1} = y, l_{st,2} = x$ and $l_{st,3} = -x - ty$. In the case of trapezoid (see Lemma 4.1.3 too) we can take $l_{st,1} = y, l_{st,2} = x, l_{st,3} = 1 - y,$ and $l_{st,4} = -x - (t - 1)y + t$. In both cases the sum $t l_{st,1} + \sum_{w>1} l_{st,w}$ is constant on A_{st} . Then note that the corresponding equations of Δ , or their modifications by $\{\epsilon_w\}_w$, are obtained from the above ones by adding constants, hence $t l_1 + \sum_{w>1} l_w$ is constant too. Evaluating on a lattice point we find that this constant is an integer c .

In fact, in A too, we can take an affine coordinate such that $l_1 = y$ and $l_2 = x$.

In Δ_{st} any line parallel to I_2 contains at most two lattice points (two is realized by the endpoints of that edge). Therefore, a parallel line to I_2 in Δ contains at most one lattice point (by (5.4.5)). Hence all the lattice points of Δ are on the line $y = 0$. This shows (a).

In order to prove the second part of (b), first take $\epsilon_w = 0$ for all w .

In the case of a triangle, $l_3 = -x - ty + c$. If $c \geq 0$ then the lattice points of Δ are exactly $\{(0, 0), \dots, (c, 0)\}$. If $c < 0$ then Δ has no lattice point.

Next assume that Δ is a trapezoid. By the choice of $l_1 = y$ and $l_2 = x$, the vertex $(a, b) := I_1 \cap I_2$ is in $(-1, 0]^2$. If $r \in (0, 1)$ is the homothety factor of Δ then the other vertices are $(a, b + r), (a + r, b + r)$ and $(a + tr, b)$. Since $a + r \in (-1, 1)$, we have two cases.

If $a + r \geq 0$, then $l_3 = -y$. Therefore, $l_4 = -x - (t - 1)y + c$ for some c , which is the sum $t l_1 + \sum_{w>1} l_w$. Again, if $c \geq 0$ then the lattice points of Δ are exactly $\{(0, 0), \dots, (c, 0)\}$, while if $c < 0$ then Δ has no lattice point.

If $a + r < 0$, then $l_3 = -y - 1$. Clearly $\Delta \cap \mathbb{Z}^3 = \emptyset$, and by a computation $c < 0$ too.

Finally notice that each $\epsilon_w = 1$ decreases both sides of (b) by 1 till either we use all the edges with $\epsilon_w = 1$, or c becomes negative (hence both sides of (b) become zero). \square

Corollary 5.4.7 (The proof of (5.1.7)). *Consider the step $\bar{z}_i \rightsquigarrow \bar{z}_i + E_{v(i)}$ where $v(i)$ is a face vertex and $(E_{v(i)}, \bar{z}_i) \leq 0$, as in 5.3.3. Then (5.1.7) holds*

$$1 - (E_{v(i)}, \bar{z}_i) \leq |P_i|.$$

Proof. Write $m_u = m_u(\bar{z}_i)$. Recall that $\Delta = F_i^{cn}$ is given by $\ell_w(p) \geq r_{i,w} \cdot m_w$, and F_i^{cn-} is given by $\ell_w(p) \geq r_{i,w} \cdot m_w + \epsilon_w$ ($w \in \mathcal{V}_{v(i)}$). Hence $l_w = \ell_w - \lceil r_{i,w} \cdot m_w \rceil - \epsilon_w$, and $m_w \geq \lceil r_{i,w} \cdot m_w \rceil + \epsilon_w$ for all w . (For the definition of $r_{i,w}$ see (5.3.6).)

Then $\mathcal{V}_{v(i)}^e = \mathcal{V}_{v(i)}$ (cf. Proposition 4.2.5(c) and (4.2.7) imply, for any $p \in H_{v(i)}^{\bar{z}_i}$,

$$\begin{aligned} -(E_{v(i)}, \bar{z}_i) &= -b_{v(i)} m_{v(i)} - \sum_{w \in \mathcal{V}_{v(i)}} m_w = -b_{v(i)} \ell_{v(i)}(p) - \sum_{w \in \mathcal{V}_{v(i)}} m_w \\ &= \sum_{w \in \mathcal{V}_{v(i)}} (\ell_w(p) - m_w) \leq \sum_{w \in \mathcal{V}_{v(i)}} l_w(p). \end{aligned}$$

Since by assumption $(E_{v(i)}, \bar{z}_i) \leq 0$, the sum $c := \sum_{w \in \mathcal{V}_{v(i)}} l_w$ is non-negative. Hence, by Lemma 5.4.6, $c + 1$ is the number of lattice points in F_i^{cn-} , which is $|P_i|$ by Lemma 5.3.8. \square

5.5. The proof of (5.1.8)

The proof is based on the comparison of the divisorial valuation/filtration with the Newton filtration. For notation see Subsection 4.3.

Lemma 5.5.1 ([10]). *For any $0 \neq g \in \mathbb{C}\{z\}$ and $v \in \mathcal{V}$ we denote by g_v the principal part of g with respect to wt_v , the weight corresponding to v . Then for any non-zero $h \in \mathbb{C}\{z\}$, $\text{wt}_v(h) < \text{div}_v(h)$ if and only if h_v is divisible by f_v over the ring $\mathbb{C}\{z\}[z_1^{-1}, z_2^{-1}, z_3^{-1}]$.*

As an immediate consequence we have

Proposition 5.5.2. *At each step $\bar{z}_i \rightsquigarrow \bar{z}_i + E_{v(i)}$ we have*

$$|P_i| \leq \dim \frac{H^0(\tilde{X}, \mathcal{O}(-\bar{z}_i))}{H^0(\tilde{X}, \mathcal{O}(-\bar{z}_i - E_{v(i)}))}.$$

Proof. Consider the functions z^p for $p \in P_i$. Since $\text{wt}(z^p) = \text{div}(z^p)$ (cf. 4.3.2), we have $z^p \in H^0(\tilde{X}, \mathcal{O}(-\bar{z}_i)) \setminus H^0(\tilde{X}, \mathcal{O}(-\bar{z}_i - E_{v(i)}))$. Therefore, it is enough to show that the family $(z^p)_{p \in P_i}$ is linearly independent modulo $H^0(\tilde{X}, \mathcal{O}(-\bar{z}_i - E_{v(i)}))$. This can be verified as follows. Assume that there exist $a_p \in \mathbb{C}$ (not all zero) such that

$$g := \sum_{p \in P_i} a_p z^p \in H^0(\tilde{X}, \mathcal{O}(-\bar{z}_i - E_{v(i)})). \quad (5.5.3)$$

Then g is its own principal part with respect to the weight corresponding to $v(i)$. We have $\text{wt}_{v(i)}(g) = m_{v(i)}(\bar{z}_i)$, but by (5.5.3), $\text{div}_{v(i)}(g) \geq m_{v(i)}(\bar{z}_i + E_{v(i)}) = m_{v(i)}(\bar{z}_i) + 1$. By Lemma 5.5.1 this means that for some $h \in \mathbb{C}\{z_1, z_2, z_3\}[z_1^{-1}, z_2^{-1}, z_3^{-1}]$ one has $g = f_{v(i)} \cdot h$. But this is impossible, since the support of g lies on a segment (cf. Lemma 5.4.6(b)) whereas the same cannot be said about $f_{v(i)}$ (since $v(i)$ is a face vertex), hence about $f_{v(i)} \cdot h$ neither. \square

Acknowledgments. The first author is partially supported by OTKA Grant 100796. The second author is supported by the PhD program of the CEU, Budapest and by the ‘Lendület’ and ERC program ‘LTDBud’ at Rényi Institute.

References

- [1] Arnold, V. I., Gusein-Zade, S. M., Varchenko, A. N.: Singularities of Differentiable Maps. Vols. 1 and 2, Monogr. Math. 82–83, Birkhäuser, Boston (1988) [Zbl 0691.58002](#)(Vol. I) [Zbl 0659.58002](#)(Vol. II) [MR 1088739](#)(Vol. I) [MR 1039615](#)(Vol. II)
- [2] Artal-Bartolo, E.: Forme de Jordan de la monodromie des singularités superisolées de surfaces. Mem. Amer. Math. Soc. **109**, no. 525, x + 84 pp. (1994) [Zbl 0918.32017](#) [MR 1204839](#)
- [3] Artal-Bartolo, E., Cassou-Noguès, P., Luengo, I., Melle Hernández, A.: Monodromy conjecture for some surface singularities. Ann. Sci. École Norm. Sup. (4) **35**, 605–640 (2002) [Zbl 1020.32025](#) [MR 1981174](#)

- [4] Artin, M.: On isolated rational singularities of surfaces. *Amer. J. Math.* **88**, 129–136 (1966) [Zbl 0142.18602](#) [MR 0199191](#)
- [5] Borodzik, M., Livingston, Ch.: Heegaard–Floer homology and rational cuspidal curves. *Forum Math. Sigma* **2**, e28, 23 pp. (2014) [Zbl 1325.14047](#) [MR 3347955](#)
- [6] Braun, G., Némethi, A.: Invariants of Newton non-degenerate surface singularities. *Compos. Math.* **143**, 1003–1036 (2007) [Zbl 1124.14033](#) [MR 2339837](#)
- [7] Collin, O., Saveliev, N.: A geometric proof of the Fintushel–Stern formula. *Adv. Math.* **147**, 304–314 (1999) [Zbl 0957.57024](#) [MR 1734525](#)
- [8] Collin, O., Saveliev, N.: Equivariant Casson invariants via gauge theory. *J. Reine Angew. Math.* **541**, 143–169 (2001) [Zbl 0989.57013](#) [MR 1876288](#)
- [9] Durfee, A.: The signature of smoothings of complex surface singularities. *Math. Ann.* **232**, 85–98 (1978) [Zbl 0346.32016](#) [MR 0466620](#)
- [10] Ebeling, W., Gusein-Zade, S.: On divisorial filtrations associated with a Newton diagram. *J. Singularities* **3**, 1–7 (2011) [Zbl 1292.32008](#) [MR 2785856](#)
- [11] Elkik, R.: Singularités rationnelles et déformations. *Invent. Math.* **47**, 139–147 (1978) [Zbl 0363.14002](#) [MR 0501926](#)
- [12] Fernández de Bobadilla, J., Luengo, I., Melle-Hernández, A., Némethi, A.: On rational cuspidal projective plane curves. *Proc. London Math. Soc.* **92**, 99–138 (2006) [Zbl 1115.14021](#) [MR 2192386](#)
- [13] Fernández de Bobadilla, J., Luengo, I., Melle-Hernández, A., Némethi, A.: On rational cuspidal curves, open surfaces and local singularities. In: *Singularity Theory*, World Sci., Hackensack, NJ, 411–442 (2007) [Zbl 1124.14026](#) [MR 2342921](#)
- [14] Fernández de Bobadilla, J., Luengo, I., Melle-Hernández, A., Némethi, A.: Classification of rational unicuspidal projective curves whose singularities have one Puiseux pair. In: *Real and Complex Singularities*, Trends Math., Birkhäuser, 31–45 (2007) [Zbl 1120.14019](#) [MR 2280130](#)
- [15] Fintushel, R., Stern, R. J.: Instanton homology of Seifert fibered homology 3-spheres. *Proc. London Math. Soc.* (3) **61**, 109–137 (1991) [Zbl 0705.57009](#) [MR 1051101](#)
- [16] Kouchnirenko, A. G.: Polyèdres de Newton et nombre de Milnor. *Invent. Math.* **32**, 1–31 (1976) [Zbl 0328.32007](#) [MR 0419433](#)
- [17] László, T., Némethi, A.: Reduction theorem for lattice cohomology. *Int. Math. Res. Notices* **2015**, 2938–2985 [Zbl 06447941](#) [MR 3373041](#)
- [18] Laufer, H. B.: *Normal Two-Dimensional Singularities*. Ann. of Math. Stud. 71, Princeton Univ. Press (1971) [Zbl 0245.32005](#) [MR 0320365](#)
- [19] Laufer, H. B.: On rational singularities. *Amer. J. Math.* **94**, 597–608 (1972) [Zbl 0251.32002](#) [MR 0330500](#)
- [20] Laufer, H. B.: On minimally elliptic singularities. *Amer. J. Math.* **99**, 1257–1295 (1977) [Zbl 0384.32003](#) [MR 0568898](#)
- [21] Laufer, H. B.: On μ for surface singularities. In: *Several Complex Variables* (Williamstown, MA, 1975), Proc. Sympos. Pure Math. 30, Amer. Math. Soc., 45–49 (1977) [Zbl 0363.32009](#) [MR 0450287](#)
- [22] Laufer, H. B.: Weak simultaneous resolution for deformations of Gorenstein surface singularities. In: *Singularities* (Arcata, CA, 1981), Proc. Sympos. Pure Math. 40, Part 2, 1–29 (1983) [Zbl 0568.14008](#) [MR 0713236](#)
- [23] Laufer, H. B.: Strong simultaneous resolution for surface singularities. In: *Complex Analytic Singularities*, Adv. Stud. Pure Math. 8, North-Holland, 207–214 (1986) [Zbl 0627.14014](#) [MR 0894294](#)
- [24] Looijenga, E., Wahl, J.: Quadratic functions and smoothing surface singularities. *Topology* **25**, 261–291 (1986) [Zbl 0615.32014](#) [MR 0842425](#)

- [25] Luengo, I.: The μ -constant stratum is not smooth. *Invent. Math.* **90**, 139–152 (1987) [Zbl 0627.32018](#) [MR 0906582](#)
- [26] Luengo, I., Melle-Hernández, A., Némethi, A.: Links and analytic invariants of superisolated singularities. *J. Algebraic Geom.* **14**, 543–565 (2005) [Zbl 1084.32022](#) [MR 2129010](#)
- [27] Melle-Hernández, A.: Milnor numbers for surface singularities. *Israel J. Math.* **115**, 29–50 (2000) [Zbl 0956.32027](#) [MR 1749672](#)
- [28] Mendris, R., Némethi, A.: The link of $\{f(x, y) + z^n = 0\}$ and Zariski's Conjecture. *Compos. Math.* **141**, 502–524 (2005) [Zbl 1077.32016](#) [MR 2134278](#)
- [29] Merle, M., Teissier, B.: Conditions d'adjonction, d'après Du Val. In: *Séminaire sur les singularités des surfaces (Polytechnique)*, 1976–1977, exp. 14, 16 pp.
- [30] Milnor, J.: *Singular Points of Complex Hypersurfaces*. Ann. of Math. Stud. 61, Princeton Univ. Press (1968) [Zbl 0184.48405](#) [MR 0239612](#)
- [31] Némethi, A.: “Weakly” elliptic Gorenstein singularities of surfaces. *Invent. Math.* **137**, 145–167 (1999) [Zbl 0934.32018](#) [MR 1703331](#)
- [32] Némethi, A.: On the Ozsváth–Szabó invariant of negative definite plumbed 3-manifolds. *Geom. Topol.* **9**, 991–1042 (2005) [Zbl 1138.57301](#) [MR 1703331](#)
- [33] Némethi, A.: On the Heegaard–Floer homology of $S^3_{-d}(K)$ and unicuspidal rational plane curves. In: *Geometry and Topology of Manifolds*, Fields Inst. Comm. 47, 219–234 (2005) [Zbl 1101.57005](#) [MR 2189934](#)
- [34] Némethi, A.: Graded roots and singularities. In: *Singularities in Geometry and Topology (Trieste, 2005)*, World Sci., Hackensack, NJ, 394–463 (2007) [Zbl 1125.14019](#) [MR 2311495](#)
- [35] Némethi, A.: Lattice cohomology of normal surface singularities. *Publ. RIMS Kyoto Univ.* **44**, 507–543 (2008) [Zbl 1149.14029](#) [MR 2426357](#)
- [36] Némethi, A.: The Seiberg–Witten invariants of negative definite plumbed 3-manifolds. *J. Eur. Math. Soc.* **13**, 959–974 (2011) [Zbl 1231.32020](#) [MR 2800481](#)
- [37] Némethi, A.: Two exact sequences for lattice cohomology. In: *Noncommutative Geometry and Global Analysis*, Contemp. Math. 546, Amer. Math. Soc., 249–269 (2011) [Zbl 1244.32016](#) [MR 2815139](#)
- [38] Némethi, A.: The cohomology of line bundles of splice-quotient singularities. *Adv. Math.* **229**, 2503–2524 (2012) [Zbl 1241.32024](#) [MR 2880230](#)
- [39] Némethi, A.: Invariants of normal surface singularities. In: *Real and Complex Singularities (San Carlos, 2002)*, Contemp. Math. 354, Amer. Math. Soc., 161–208 (2004) [Zbl 1076.14007](#) [MR 2087811](#)
- [40] Némethi, A., Nicolaescu, L. I.: Seiberg–Witten invariants and surface singularities. *Geom. Topol.* **6**, 269–328 (2002) [Zbl 1031.32023](#) [MR 1914570](#)
- [41] Némethi, A., Nicolaescu, L. I.: Seiberg–Witten invariants and surface singularities: splittings and cyclic covers. *Selecta Math. (N.S.)* **11**, 399–451 (2005) [Zbl 1110.14006](#) [MR 2215260](#)
- [42] Némethi, A., Okuma, T.: On the Casson Invariant Conjecture of Neumann–Wahl. *J. Algebraic Geom.* **18**, 135–149 (2009) [Zbl 1154.14025](#) [MR 2868183](#)
- [43] Némethi, A., Okuma, T.: The Seiberg–Witten invariant conjecture for splice-quotients. *J. London Math. Soc.* **28**, 143–154 (2008) [Zbl 1149.14030](#)
- [44] Némethi, A., Román, F.: The lattice cohomology of $S^3_{-d}(K)$. In: *Zeta Functions in Algebra and Geometry*, Contemp. Math. 566, Amer. Math. Soc., 261–292 (2012) [Zbl 1270.57046](#) [MR 2427056](#)
- [45] Neumann, W. D.: A calculus for plumbing applied to the topology of complex surface singularities and degenerating complex curves. *Trans. Amer. Math. Soc.* **268**, 299–344 (1981) [Zbl 0546.57002](#) [MR 0632532](#)

- [46] Neumann, W., Wahl, J.: Casson invariant of links of singularities. *Comment. Math. Helv.* **65**, 58–78 (1991) [Zbl 0704.57007](#) [MR 1036128](#)
- [47] Neumann, W., Wahl, J.: Universal abelian covers of surface singularities. In: *Trends in Singularities*, Trends Math., Birkhäuser, Basel, 181–190 (2002) [Zbl 1072.14502](#) [MR 1900786](#)
- [48] Neumann, W., Wahl, J.: Universal abelian covers of quotient-cusps. *Math. Ann.* **326**, 75–93 (2003) [Zbl 1032.14010](#) [MR 1981612](#)
- [49] Oka, M.: On the resolution of hypersurface singularities. In: *Complex Analytic Singularities*, Adv. Stud. Pure Math. 8, North-Holland, 405–436 (1987) [Zbl 0622.14012](#) [MR 0894303](#)
- [50] Ozsváth, P. S., Szabó, Z.: Holomorphic disks and topological invariants for closed three-manifolds. *Ann. of Math.* **159**, 1027–1158 (2004) [Zbl 1073.57009](#) [MR 2113019](#)
- [51] Ozsváth, P. S., Szabó, Z.: Holomorphic discs and three-manifold invariants: properties and applications. *Ann. of Math.* **159**, 1159–1245 (2004) [Zbl 1081.57013](#) [MR 2113020](#)
- [52] Popescu-Pampu, P.: Numerically Gorenstein surface singularities are homeomorphic to Gorenstein ones. *Duke Math. J.* **159**, 539–559 (2011) [Zbl 1227.14010](#) [MR 2831877](#)
- [53] Saito, M.: Exponents and Newton polyhedra of isolated hypersurface singularities. *Math. Ann.* **281**, 411–417 (1988) [Zbl 0628.32038](#) [MR 0954149](#)
- [54] Steenbrink, J. H. M.: Mixed Hodge structures associated with isolated singularities. In: *Singularities (Arcata, CA, 1981)*, Proc. Sympos. Pure Math. 40, Part 2, Amer. Math. Soc., 513–536 (1983) [Zbl 0515.14003](#) [MR 0713277](#)
- [55] Veys, W.: Stringy invariants of normal surfaces. *J. Algebraic Geom.* **13**, 115–141 (2004) [Zbl 1060.14021](#) [MR 2008717](#)
- [56] Wahl, M. J.: Equisingular deformations of normal surface singularities. I. *Ann. of Math.* **104**, 325–356 (1976) [Zbl 0358.14007](#) [MR 0422270](#)
- [57] Wahl, J.: Smoothings of normal surface singularities. *Topology* **20**, 219–246 (1981) [Zbl 0484.14012](#) [MR 0608599](#)
- [58] Yau, S. S.-T.: Gorenstein singularities with geometric genus equal to one. *Amer. J. Math.* **101**, 813–854 (1979) [Zbl 0415.14005](#) [MR 0536042](#)
- [59] Yau, S. S.-T.: On maximally elliptic singularities. *Trans. Amer. Math. Soc.* **257**, 269–329 (1980) [Zbl 0343.32009](#) [MR 0552260](#)