



Francesco Caravenna · Rongfeng Sun · Nikos Zygouras

Polynomial chaos and scaling limits of disordered systems

Received December 20, 2013 and in revised form July 1, 2014

Abstract. Inspired by recent work of Alberts, Khanin and Quastel [AKQ14a], we formulate general conditions ensuring that a sequence of multi-linear polynomials of independent random variables (called *polynomial chaos* expansions) converges to a limiting random variable, given by a *Wiener chaos* expansion over the d -dimensional white noise. A key ingredient in our approach is a *Lindeberg principle* for polynomial chaos expansions, which extends earlier work of Mossel, O’Donnell and Oleszkiewicz [MOO10]. These results provide a unified framework to study the *continuum and weak disorder scaling limits* of statistical mechanics systems that are *disorder relevant*, including the disordered pinning model, the (long-range) directed polymer model in dimension $1 + 1$, and the two-dimensional random field Ising model. This gives a new perspective in the study of disorder relevance, and leads to interesting new continuum models that deserve further studies.

Keywords. Continuum limit, finite size scaling, Lindeberg principle, polynomial chaos, Wiener chaos, disordered pinning model, directed polymer model, random field Ising model

Contents

1. Introduction	2
1.1. Continuum limits of disordered systems	2
1.2. General strategy and results	4
1.3. Discussion and perspectives	7
2. From polynomial to Wiener chaos via Lindeberg	11
2.1. White noise in a nutshell	11
2.2. Polynomial chaos	12
2.3. Convergence of polynomial chaos to Wiener chaos	13
2.4. Lindeberg principle for polynomial chaos	16
3. Scaling limits of disordered systems	19
3.1. Disordered pinning model	19
3.2. Directed polymer model	23
3.3. Random field Ising model	27

F. Caravenna: Dipartimento di Matematica e Applicazioni, Università degli Studi di Milano-Bicocca, via Cozzi 55, 20125 Milano, Italy; e-mail: francesco.caravenna@unimib.it

R. Sun: Department of Mathematics, National University of Singapore, 10 Lower Kent Ridge Road, 119076 Singapore; e-mail: matsr@nus.edu.sg

N. Zygouras: Department of Statistics, University of Warwick, Coventry CV4 7AL, UK; e-mail: N.Zygouras@warwick.ac.uk

Mathematics Subject Classification (2010): Primary 82B44; Secondary 82D60, 60K35

4. Proof of the Lindeberg principle	31
5. Proof of the convergence to Wiener chaos	37
6. Proof for the disordered pinning model	42
7. Proof for the directed polymer model	46
8. Proof for the random field Ising model	49
Appendix A. The Cameron–Martin shift	56
Appendix B. Technical lemmas	57
References	63

1. Introduction

In this paper, we consider statistical mechanics models defined on a lattice, in which disorder acts as an external “random field”. We focus on models that are *disorder relevant*, in the sense that arbitrarily weak disorder changes the qualitative properties of the model. We will show that when the homogeneous model (without disorder) has a non-trivial continuum limit, disorder relevance manifests itself via the convergence of the disordered model to a *disordered continuum limit* if the disorder strength and lattice mesh are suitably rescaled.

Our approach is inspired by recent work of Alberts, Khanin and Quastel [AKQ14a] on the directed polymer model in dimension $1+1$. Here we follow a different path, establishing a general convergence result for polynomial chaos expansions based on a Lindeberg principle. This extends earlier work of Mossel, O’Donnell and Oleszkiewicz [MOO10] to optimal (second) moment assumptions, and is of independent interest.

In this section, we present somewhat informally the main ideas of our approach in a unified framework, emphasizing the natural heuristic considerations. The precise formulation of our results is given in Sections 2 and 3, which can be read independently (both of each other and of the present one). The proofs are contained in Sections 4 to 8, while some technical parts have been deferred to the Appendices. Throughout the paper, we use the conventions $\mathbb{N} := \{1, 2, 3, \dots\}$ and $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$, and we denote by *Leb* the Lebesgue measure on \mathbb{R}^d .

1.1. Continuum limits of disordered systems

Consider an open set $\Omega \subseteq \mathbb{R}^d$ and define the lattice $\Omega_\delta := (\delta\mathbb{Z})^d \cap \Omega$ for $\delta > 0$. Suppose that a *reference probability measure* $\mathbb{P}_{\Omega_\delta}^{\text{ref}}$ is given on $\mathbb{R}^{\Omega_\delta}$, which describes a real-valued field $\sigma = (\sigma_x)_{x \in \Omega_\delta}$. We focus on the case when each σ_x takes two possible values (typically $\sigma_x \in \{0, 1\}$ or $\sigma_x \in \{-1, 1\}$).

Let $\omega := (\omega_x)_{x \in \Omega_\delta}$ be i.i.d. random variables (also independent of σ) with zero mean, unit variance, and locally finite exponential moments, which represent the *disorder*. Probability and expectation for ω will be denoted respectively by \mathbb{P} and \mathbb{E} .

Given $\lambda > 0$, $h \in \mathbb{R}$ and a \mathbb{P} -typical realization of the disorder ω , we define the *disordered model* as the following probability measure $\mathbb{P}_{\Omega_\delta; \lambda, h}^\omega$ for the field $\sigma = (\sigma_x)_{x \in \Omega_\delta}$:

$$\mathbb{P}_{\Omega_\delta; \lambda, h}^\omega(d\sigma) := \frac{e^{\sum_{x \in \Omega_\delta} (\lambda\omega_x + h)\sigma_x}}{Z_{\Omega_\delta; \lambda, h}^\omega} \mathbb{P}_{\Omega_\delta}^{\text{ref}}(d\sigma), \quad (1.1)$$

where the normalizing constant, called the *partition function*, is defined by

$$Z_{\Omega_\delta; \lambda, h}^\omega := \mathbb{E}_{\Omega_\delta}^{\text{ref}} \left[e^{\sum_{x \in \Omega_\delta} (\lambda \omega_x + h) \sigma_x} \right]. \quad (1.2)$$

The *quenched free energy* $F(\lambda, h)$ of the model is defined as the rate of exponential growth of $Z_{\Omega_\delta; \lambda, h}^\omega$ as $\Omega \uparrow \mathbb{R}^d$ for fixed δ (or equivalently¹ as $\delta \downarrow 0$ for fixed Ω):

$$F(\lambda, h) := \limsup_{\Omega \uparrow \mathbb{R}^d} \frac{1}{|\Omega_\delta|} \mathbb{E}[\log Z_{\Omega_\delta; \lambda, h}^\omega] = \limsup_{\delta \downarrow 0} \frac{1}{|\Omega_\delta|} \mathbb{E}[\log Z_{\Omega_\delta; \lambda, h}^\omega]. \quad (1.3)$$

Discontinuities in the derivatives of the free energy correspond to phase transitions. A fundamental question is: does arbitrary disorder (i.e., $\lambda > 0$) radically change the behavior of the homogeneous model (i.e., $\lambda = 0$), such as the qualitative properties of the law of the field σ and/or the smoothness of the free energy in h ? When the answer is affirmative, the model is called *disorder relevant*. In such cases, we will show that the disordered model typically admits a non-trivial scaling limit as $\delta \downarrow 0$, provided $\lambda, h \rightarrow 0$ at appropriate rates.

Informally speaking, our key assumption is that the discrete field $\sigma = (\sigma_x)_{x \in \Omega_\delta}$, under the reference law $\mathbf{P}_{\Omega_\delta}^{\text{ref}}$ and after a suitable rescaling, converges as $\delta \downarrow 0$ to a “continuum field” $\sigma = (\sigma_x)_{x \in \Omega}$, possibly distribution-valued, with law $\mathbf{P}_\Omega^{\text{ref}}$. (Our precise assumptions will be about the convergence of correlation functions, see (1.11).) Although the approach we follow is very general, we describe three specific models, to be discussed extensively in what follows.

1. The *disordered pinning model* ($d = 1$). Let $\tau = (\tau_k)_{k \geq 0}$ be a *renewal process* on \mathbb{N} with $\mathbb{P}(\tau_1 = n) = n^{-(1+\alpha)+o(1)}$, with $\alpha \in (1/2, 1)$. Take $\Omega = (0, 1)$, $\delta = 1/N$ for $N \in \mathbb{N}$, and define $\mathbf{P}_{\Omega_\delta}^{\text{ref}}$ as the law of $(\sigma_x := \mathbb{1}_{\delta\tau}(x))_{x \in \Omega_\delta}$, where $\delta\tau = \{N^{-1}\tau_n\}_{n \geq 0}$ is viewed as a random subset of Ω . The continuum field $\mathbf{P}_\Omega^{\text{ref}}$ is $(\sigma_x = \mathbb{1}_\tau(x))_{x \in (0, 1)}$ where τ denotes the α -stable regenerative set (the zero level set of a Bessel($2(1 - \alpha)$) process).
2. The (long-range) *directed polymer model*. Let $(S_n)_{n \geq 0}$ be a random walk on \mathbb{Z} with i.i.d. increments, in the domain of attraction of an α -stable law, with $1 < \alpha \leq 2$. Take $\Omega = (0, 1) \times \mathbb{R}$, $\delta = 1/N$ for $N \in \mathbb{N}$ and, abusing notation, set $\Omega_\delta := ((\delta\mathbb{Z}) \times (\delta^{1/\alpha}\mathbb{Z})) \cap \Omega$. The “effective dimension” for this model is therefore $d_{\text{eff}} := 1 + 1/\alpha$. Define $\mathbf{P}_{\Omega_\delta}^{\text{ref}}$ as the law of the field $(\sigma_x := \mathbb{1}_{A_\delta}(x))_{x \in \Omega_\delta}$, where $A_\delta := \{(n/N, S_n/N^{1/\alpha})\}_{n \geq 0}$ is viewed as a random subset of Ω . The continuum field $\mathbf{P}_\Omega^{\text{ref}}$ is $(\sigma_x = \mathbb{1}_A(x))_{x \in (0, 1) \times \mathbb{R}}$ where $A = \{(t, X_t)\}_{t \geq 0}$ and $(X_t)_{t \geq 0}$ is an α -stable Lévy process (Brownian motion for $\alpha = 2$).
3. The *random field Ising model* ($d = 2$). Take any bounded and connected set $\Omega \subseteq \mathbb{R}^2$ with smooth boundary and define $\mathbf{P}_{\Omega_\delta}^{\text{ref}}$ to be the *critical* Ising model on Ω_δ with inverse temperature $\beta = \beta_c = \frac{1}{2} \log(1 + \sqrt{2})$ and + boundary condition. The (distribution-valued) continuum field $\mathbf{P}_\Omega^{\text{ref}}$ has been recently constructed in [CGN12a, CGN13], using breakthrough results on the scaling limit of correlation functions of the critical two-dimensional Ising model, determined in [CHI12].

¹ We assume the natural consistency condition that $\mathbf{P}_{(c\Omega)_{c\delta}}^{\text{ref}}$ coincides with $\mathbf{P}_{\Omega_\delta}^{\text{ref}}$ for any $c > 0$.

The restrictions on the dimensions and parameters of these models are linked precisely to the disorder relevance issue, as will be explained later.

Since the reference law $\mathbf{P}_{\Omega_\delta}^{\text{ref}}$ has a weak limit $\mathbf{P}_\Omega^{\text{ref}}$ as $\delta \downarrow 0$, a natural question emerges: can one obtain a limit also for the disordered model $\mathbf{P}_{\Omega_\delta; \lambda, h}^\omega$, under an appropriate scaling of the coupling constants λ, h ? (We mean, of course, a *non-trivial* limit, which keeps track of λ, h ; otherwise, it suffices to let $\lambda, h \rightarrow 0$ very fast to recover the “free case” $\mathbf{P}_\Omega^{\text{ref}}$.)

A natural strategy is to look at the exponential weight in (1.1). As $\delta \downarrow 0$, the discrete disorder $\omega = (\omega_x)_{x \in \Omega_\delta}$ approximates the white noise $W(dx)$, which is a sort of random signed measure on Ω (see Subsection 2.1 for more details). Then one might hope to define the candidate continuum disordered model $\mathbf{P}_{\Omega; \hat{\lambda}, \hat{h}}^W$ by

$$\frac{d\mathbf{P}_{\Omega; \hat{\lambda}, \hat{h}}^W}{d\mathbf{P}_\Omega^{\text{ref}}}(\boldsymbol{\sigma}) := \frac{e^{\int_\Omega \sigma_x (\hat{\lambda} W(dx) + \hat{h} dx)}}{\mathbf{Z}_{\Omega; \hat{\lambda}, \hat{h}}^W}, \quad (1.4)$$

in analogy with (1.1), with $\mathbf{Z}_{\Omega; \hat{\lambda}, \hat{h}}^W$ defined accordingly, as in (1.2).

Unfortunately, formula (1.4) typically makes no sense for $\hat{\lambda} \neq 0$, because the configurations of the continuum field $\boldsymbol{\sigma} = (\sigma_x)_{x \in \Omega}$ under $\mathbf{P}_\Omega^{\text{ref}}$ are too rough or “thin” for the integral over $W(dx)$ to be meaningful (cf. the three motivating models listed above). We stress here that the difficulty is substantial and not just technical: for pinning and directed polymer models, one can show [AKQ14b, CSZ14] that the scaling limit $\mathbf{P}_{\Omega; \hat{\lambda}, \hat{h}}^W$ of $\mathbf{P}_{\Omega_\delta; \lambda, h}^\omega$ exists, but for $\hat{\lambda} \neq 0$ it is *not absolutely continuous with respect to* $\mathbf{P}_\Omega^{\text{ref}}$. In particular, it is hopeless to define the continuum disordered model through a Radon–Nikodym density, as in (1.4).

1.2. General strategy and results

In this paper we focus on the disordered partition function $Z_{\Omega_\delta; \lambda, h}^\omega$. We show that when $\delta \downarrow 0$ and λ, h are scaled appropriately, the partition function admits a non-trivial limit in distribution, which is explicit and *universal* (i.e., it does not depend on the fine details of the model).

Switching from the random probability law $\mathbf{P}_{\Omega_\delta; \lambda, h}^\omega$ to the random number $Z_{\Omega_\delta; \lambda, h}^\omega$ is of course a simplification, whose relevance may not be evident. It turns out that the partition function *contains essential information on the model*. In fact, the scaling limit of $Z_{\Omega_\delta; \lambda, h}^\omega$, for sufficiently many domains Ω and “boundary conditions”, allows one to reconstruct *the full scaling limit of* $\mathbf{P}_{\Omega_\delta; \lambda, h}^\omega$. This task has been achieved in [AKQ14b] for the directed polymer model based on simple random walk, and in [CSZ14] for the disordered pinning model. We discuss the case of the long-range directed polymer model in Remark 3.9 below.

The scaling limit of $Z_{\Omega_\delta; \lambda, h}^\omega$ can also describe the *universal behavior* of the free energy $F(\lambda, h)$ as $\lambda, h \rightarrow 0$ (cf. Subsection 1.3). This explains the key role of the partition function and is a strong motivation for focusing on it in the first place.

We now describe our approach. The idea is to consider the so-called “high-temperature expansion” ($|\lambda|, |h| \ll 1$) of the partition function $Z_{\Omega_\delta; \lambda, h}^\omega$. When the field

takes two values (say $\sigma_x \in \{0, 1\}$, for simplicity), we can factorize and “linearize” the exponential in (1.2):

$$Z_{\Omega_\delta}^\omega; \lambda, h = \mathbb{E}_{\Omega_\delta}^{\text{ref}} \left[\prod_{x \in \Omega_\delta} (1 + \varepsilon_x \sigma_x) \right], \quad \text{where} \quad \varepsilon_x := e^{\lambda \omega_x + h} - 1. \quad (1.5)$$

Let us introduce the k -point correlation function $\psi_{\Omega_\delta}^{(k)}(x_1, \dots, x_k)$ of the field under the reference law, defined for $k \in \mathbb{N}$ and distinct $x_1, \dots, x_k \in \Omega$ by

$$\psi_{\Omega_\delta}^{(k)}(x_1, \dots, x_k) := \mathbb{E}_{\Omega_\delta}^{\text{ref}} [\sigma_{x_1} \cdots \sigma_{x_k}], \quad (1.6)$$

where we set $\sigma_x := \sigma_{x_\delta}$ with x_δ being the point in Ω_δ closest to $x \in \Omega$. (We define the correlation function on all points of Ω for later convenience, and set it to be zero whenever $(x_i)_\delta = (x_j)_\delta$ for some $i \neq j$.) A binomial expansion of the product in (1.5) then yields

$$Z_{\Omega_\delta}^\omega; \lambda, h = 1 + \sum_{k=1}^{|\Omega_\delta|} \frac{1}{k!} \sum_{(x_1, \dots, x_k) \in (\Omega_\delta)^k} \psi_{\Omega_\delta}^{(k)}(x_1, \dots, x_k) \prod_{i=1}^k \varepsilon_{x_i}, \quad (1.7)$$

where the $k!$ accounts for the fact that we sum over *ordered* k -uples (x_1, \dots, x_k) . We have rewritten the partition function as a multi-linear polynomial of the independent random variables $(\varepsilon_x)_{x \in \Omega_\delta}$ (what is called a *polynomial chaos* expansion), with coefficients given by the k -point correlation function of the reference field. Note that, by Taylor expansion,

$$\mathbb{E}[\varepsilon_x] \simeq h + \frac{1}{2} \lambda^2 =: h', \quad \text{Var}[\varepsilon_x] \simeq \lambda^2. \quad (1.8)$$

The crucial fact is that, for $|\lambda|, |h| \ll 1$, the distribution of a polynomial chaos expansion, like the right hand side of (1.7), is insensitive the marginal distribution of the random variables $(\varepsilon_x)_{x \in \Omega_\delta}$, as long as mean and variance are kept fixed. *A precise formulation of this loosely stated invariance principle is given in Section 2 (Theorems 2.6 and 2.8) in the form of a Lindeberg principle.* Denoting by $(\tilde{\omega}_x)_{x \in \Omega_\delta}$ a family of i.i.d. standard Gaussians, by (1.8) we can then approximate

$$Z_{\Omega_\delta}^\omega; \lambda, h \simeq 1 + \sum_{k=1}^{|\Omega_\delta|} \frac{1}{k!} \sum_{(x_1, \dots, x_k) \in (\Omega_\delta)^k} \psi_{\Omega_\delta}^{(k)}(x_1, \dots, x_k) \prod_{i=1}^k (\lambda \tilde{\omega}_{x_i} + h'). \quad (1.9)$$

Let us now introduce a white noise $W(\cdot)$ on \mathbb{R}^d (see Subsection 2.1): setting $\Delta := (-\delta/2, \delta/2)^d$, we replace each $\tilde{\omega}_x$ by $\delta^{-d/2} W(x + \Delta)$. Since $h' = h' \delta^{-d} \text{Leb}(x + \Delta)$, the inner sum in (1.9) coincides, up to boundary terms, with the following (deterministic + stochastic) integral:

$$\int \cdots \int_{\Omega^k} \psi_{\Omega_\delta}^{(k)}(x_1, \dots, x_k) \prod_{i=1}^k (\lambda \delta^{-d/2} W(dx_i) + h' \delta^{-d} dx_i). \quad (1.10)$$

(We recall that $\psi_{\Omega_\delta}^{(k)}(x_1, \dots, x_k)$ is a piecewise constant function.)

We can now state our crucial assumption: we suppose that, for every $k \in \mathbb{N}$, there exist a symmetric function $\psi_{\Omega}^{(k)} : (\mathbb{R}^d)^k \rightarrow \mathbb{R}$ and an exponent $\gamma \in [0, \infty)$ such that

$$(\delta^{-\gamma})^k \psi_{\Omega_{\delta}}^{(k)}(x_1, \dots, x_k) \xrightarrow{\delta \downarrow 0} \psi_{\Omega}^{(k)}(x_1, \dots, x_k) \quad \text{in } L^2(\Omega^k). \quad (1.11)$$

By (1.10), if we fix $\hat{\lambda} \geq 0$, $\hat{h} \in \mathbb{R}$ and rescale the coupling constants as follows:

$$\lambda = \hat{\lambda} \delta^{d/2-\gamma}, \quad h' = \hat{h} \delta^{d-\gamma} \quad (\text{where } h' := h + \lambda^2/2), \quad (1.12)$$

equations (1.9)–(1.10) suggest that $Z_{\Omega_{\delta}; \lambda, h}^{\omega}$ converges in distribution as $\delta \downarrow 0$ to a random variable which admits a *Wiener chaos expansion* with respect to the white noise $W(\cdot)$:

$$Z_{\Omega_{\delta}; \lambda, h}^{\omega} \xrightarrow{\delta \downarrow 0} Z_{\Omega; \hat{\lambda}, \hat{h}}^W := 1 + \sum_{k=1}^{\infty} \frac{1}{k!} \int \cdots \int_{\Omega^k} \psi_{\Omega}^{(k)}(x_1, \dots, x_k) \prod_{i=1}^k (\hat{\lambda} W(dx_i) + \hat{h} dx_i). \quad (1.13)$$

This is precisely what happens, by our main convergence results in Section 2 (Theorems 2.3 and 2.5). It is natural to call the random variable $Z_{\Omega; \hat{\lambda}, \hat{h}}^W$ in (1.13) the *continuum partition function*, because it is the scaling limit of $Z_{\Omega_{\delta}; \lambda, h}^{\omega}$.

Remark 1.1. The L^2 convergence in (1.11) typically requires $\gamma < d/2$ (cf. (1.16) below), which means that the disorder coupling constants λ, h vanish as $\delta \downarrow 0$, by (1.12). The fact that the continuum partition function $Z_{\Omega; \hat{\lambda}, \hat{h}}^W$ in (1.13) is nevertheless a *random object* (for $\hat{\lambda} > 0$) is a manifestation of *disorder relevance*. We elaborate more on this issue in Subsection 1.3.

Let us finally take a quick look at the three motivating models. *The complete results are described in Section 3 (Theorems 3.1, 3.8 and 3.14).* Note that the scaling exponents in (1.12) are determined by the dimension d and by the exponent γ appearing in (1.11).

1. For the *disordered pinning model* ($d = 1$), one has $\gamma = 1 - \alpha$ by renewal theory [D97]. Relation (1.12) (for $\delta = 1/N$) yields

$$\lambda = \hat{\lambda}/N^{\alpha-1/2}, \quad h' = \hat{h}/N^{\alpha}.$$

Notice that $h' = (\text{const})\lambda^{2\alpha/(2\alpha-1)}$ is precisely the scaling of the critical curve [G10].

2. For the (long-range) *directed polymer model* ($d_{\text{eff}} = 1 + 1/\alpha$), one has $\gamma = 1/\alpha$ by Gnedenko's local limit theorem [BGT87]. Therefore (1.12) (for $\delta = 1/N$ and d replaced by d_{eff}) gives

$$\lambda = \hat{\lambda}/N^{(\alpha-1)/(2\alpha)}, \quad h' = \hat{h}/N.$$

(The parameter h' is actually irrelevant for this model, and one usually sets $h' \equiv 0$, i.e., $h = -\frac{1}{2}\lambda^2$.) In the case $\alpha = 2$, when the underlying random walk has zero mean and finite variance, one recovers the scaling $\lambda \approx N^{-1/4}$ determined in [AKQ14a].

3. For the *random field Ising model* ($d = 2$) one has $\gamma = 1/8$ [CHI12], hence by (1.12),

$$\lambda = \hat{\lambda}\delta^{7/8}, \quad h = \hat{h}\delta^{15/8}. \quad (1.14)$$

(Note that h instead of h' appears in this relation; moreover one should look at the normalized partition function $\exp(-\frac{1}{2}\lambda^2|\Omega_\delta|)Z_{\Omega_\delta;\lambda,h}^\omega$. This is because $\sigma_x \in \{-1, 1\}$ instead of $\{0, 1\}$, hence the starting relation (1.5) requires a correction.)

1.3. Discussion and perspectives

We now collect some comments and observations and point out some further directions of research.

1. (Disorder relevance). The main motivation of our approach is to understand the issue of *disorder relevance*, i.e. whether the addition of a small amount of disorder modifies the nature of the phase transition of the underlying homogeneous model. Remarkably, the key condition (1.11), which determines the class of models to which our approach applies, is consistent with the Harris criterion for disorder relevance, as we now discuss.

First we note that when $\gamma > 0$, which is the most interesting case, condition (1.11) indicates that the reference law has polynomially decaying correlations, which signifies that *we are at the critical point of a continuous phase transition*. In our context, this means that the order parameter $m_h := \lim_{\delta \downarrow 0} |\Omega_\delta|^{-1} \mathbb{E}_{\Omega_\delta;0,h}[\sum_{x \in \Omega_\delta} \sigma_x]$ in the homogeneous model ($\lambda = 0$) vanishes continuously, but non-analytically, as $h \rightarrow 0$ (cf. (1.1) and (1.2)).

When (1.11) holds *pointwise*, with $\gamma > 0$, the limiting correlation function typically diverges polynomially on diagonals, with the same exponent γ :

$$\psi_\Omega^{(k)}(x_1, \dots, x_k) \approx \|x_i - x_j\|^{-\gamma} \quad \text{as } x_i \rightarrow x_j. \quad (1.15)$$

To have finite L^2 norm (which is necessary for L^2 convergence in (1.11)), such a local divergence must be locally square-integrable in \mathbb{R}^d : this means that $(d-1) - 2\gamma > -1$, i.e.

$$\gamma < d/2. \quad (1.16)$$

(Note that $d/2 - \gamma$ is precisely the scaling exponent of the coupling constant λ in (1.12).)

Relation (1.16) matches the *Harris criterion* for disorder relevance [H74]. This was originally introduced in the context of the Ising model with *bond disorder*, but it can be naturally rephrased for general disordered system (cf. [G10], [CCFS86]): denoting by ν the *correlation length exponent* of the homogeneous system ($\lambda = 0$), it asserts that a d -dimensional system is disorder-relevant when $\nu < 2/d$ and irrelevant when $\nu > 2/d$ (the remaining case $\nu = 2/d$ being dubbed *marginal*). The exponent ν is usually defined in terms of field correlations:

$$|\mathbb{E}_{\Omega_\delta;0,h}(\sigma_x \sigma_y) - \mathbb{E}_{\Omega_\delta;0,h}(\sigma_x)\mathbb{E}_{\Omega_\delta;0,h}(\sigma_y)| \underset{\delta \downarrow 0}{\approx} e^{-\frac{|x-y|}{\delta \xi(h)}} \quad \text{with } \xi(h) \underset{h \downarrow 0}{\approx} h^{-\nu}. \quad (1.17)$$

Since such an exponent can be difficult to compute, it is typical to consider alternative notions of correlation length $\xi(h)$, linked to finite size scaling [CCFS86, CCFS89]. In our context, it is natural to define

$$\xi(h)^{-1} := \max\{\delta > 0 : Z_{\Omega_\delta; 0, h} > A\}, \quad (1.18)$$

where $A > 0$ is a fixed large constant, whose precise value is immaterial. Recalling (1.12), we can rewrite (1.13) for $\lambda = 0$ as

$$\lim_{h \downarrow 0} Z_{\Omega_\delta; 0, h} = Z_{\Omega; 0, \hat{h}} \quad \text{with} \quad \delta = \delta_{h, \hat{h}} := (h/\hat{h})^{1/(d-\gamma)}.$$

If $\hat{h} \mapsto Z_{\Omega; 0, \hat{h}}$ is increasing (e.g., when $\psi_\Omega^{(k)}(x_1, \dots, x_k) \geq 0$, by (1.13)), denoting by \hat{h}_A the unique solution to $Z_{\Omega; 0, \hat{h}} = A$, under some natural regularity assumptions we have

$$\xi(h) \underset{h \downarrow 0}{\sim} (\delta_{h, \hat{h}_A})^{-1} = \left(\frac{\hat{h}_A}{h}\right)^{1/(d-\gamma)} \approx h^{-\nu}, \quad \text{with} \quad \nu = \frac{1}{d-\gamma}.$$

This shows that *Harris' condition* $\nu < 2/d$ coincides with the key condition $\gamma < d/2$ of our approach (cf. (1.16)), ensuring the square-integrability of the limiting correlations.

The correlation length (1.18) is expected to be equivalent to the classical one (1.17), in the sense that it should have the same critical exponent (cf. [G07] for disordered pinning models) when the phase transition is continuous, that is, when $\gamma > 0$ in (1.15). If $\gamma = 0$, which is the signature of a discontinuous (first order) phase transition, our approach still applies and gives the scaling limit of the disorder partition function, but there is no direct link with disorder relevance (cf. (1.24) below and the following discussion).

In summary, our approach suggests an *alternative view on disorder relevance*, in which the randomness survives in the continuum limit with vanishing coupling constants. In fact, relation (1.13) can be seen as a rigorous *finite size scaling* relation [Car88] for disordered systems (the special case of non-disordered pinning models is treated in [Soh09]).

Remark 1.2. We can now explain the parameter restrictions in the motivating models: condition (1.16) is fulfilled by the *disordered pinning model* ($d = 1$, $\gamma = 1 - \alpha$) when $\alpha > 1/2$, by the (long-range) *directed polymer model* ($d_{\text{eff}} = 1 + 1/\alpha$, $\gamma = 1/\alpha$) when $\alpha > 1$, and by the critical *random field Ising model* ($d = 2$, $\gamma = 1/8$).

2. (Universality). The convergence in distribution of the discrete partition function $Z_{\Omega_\delta; \lambda, h}^\omega$ toward its continuum counterpart $Z_{\Omega; \hat{\lambda}, \hat{h}}^W$ (cf. (1.13)) is an instance of *universality*. In fact:

- the details of the disorder distribution are irrelevant: any family $(\omega_x)_{x \in \Omega_\delta}$ of i.i.d. random variables with zero mean, finite variance and locally finite exponential moments scales in the limit to the same continuum object, namely white noise $W(\cdot)$;
- also the fine details of the reference law $\mathbb{P}_{\Omega_\delta}^{\text{ref}}$ disappear in the limit: any family $\psi_{\Omega_\delta}^{(k)}$ of discrete k -point correlation functions converging to the same limit (1.11) yields the same continuum partition function $Z_{\Omega; \hat{\lambda}, \hat{h}}^W$ in (1.13).

At a deeper level, the continuum partition function sheds light on the discrete free energy $F(\lambda, h)$ (cf. (1.3)) in the weak disorder regime $\lambda, h \rightarrow 0$. Defining the *continuum free energy*

$$\mathbf{F}(\hat{\lambda}, \hat{h}) := \limsup_{\Omega \uparrow \mathbb{R}^d} \frac{1}{\text{Leb}(\Omega)} \mathbb{E}[\log Z_{\Omega; \hat{\lambda}, \hat{h}}^W], \quad (1.19)$$

and setting $\lambda_\delta := \delta^{d/2-\gamma} \hat{\lambda}$ and $h_\delta := \delta^{d-\gamma} \hat{h} - \frac{1}{2}(\lambda_\delta)^2$ (cf. (1.12)), one is led to the following

$$\text{Conjecture: } \lim_{\delta \downarrow 0} \frac{F(\lambda_\delta, h_\delta)}{\delta^d} = \mathbf{F}(\hat{\lambda}, \hat{h}).^2 \quad (1.20)$$

The heuristics goes as follows: by (1.3) we can write (replacing lim sup by lim for simplicity)

$$\lim_{\delta \downarrow 0} \frac{F(\lambda_\delta, h_\delta)}{\delta^d} = \lim_{\delta \downarrow 0} \lim_{\Omega \uparrow \mathbb{R}^d} \frac{1}{\delta^d} \frac{1}{|\Omega_\delta|} \mathbb{E}[\log Z_{\Omega_\delta; \lambda_\delta, h_\delta}^\omega]; \quad (1.21)$$

on the other hand, applying (1.13) in (1.19) (assuming uniform integrability) and noting that $\text{Leb}(\Omega) = \lim_{\delta \downarrow 0} \delta^d |\Omega_\delta|$, one gets

$$\mathbf{F}(\hat{\lambda}, \hat{h}) = \lim_{\Omega \uparrow \mathbb{R}^d} \lim_{\delta \downarrow 0} \frac{1}{\delta^d} \frac{1}{|\Omega_\delta|} \mathbb{E}[\log Z_{\Omega_\delta; \lambda_\delta, h_\delta}^\omega]. \quad (1.22)$$

Therefore proving (1.20) amounts to *interchanging the infinite volume* ($\Omega \uparrow \mathbb{R}^d$) and *continuum and weak disorder* ($\delta \downarrow 0$) limits. This is in principle a delicate issue, but we expect relation (1.20) to hold in many interesting cases, such as the three motivating models in the specified parameter range (and, more generally, when the continuum correlations are “non-trivial”; see the next point). This is an interesting open problem.

Relation (1.20) implies that the discrete free energy $F(\lambda, h)$ has a *universal shape for weak disorder* $\lambda, h \rightarrow 0$. This leads to sharp predictions on the asymptotic behavior of free energy-related quantities, such as critical curves and order parameters. Consider, e.g., the average magnetization $\langle \sigma_0 \rangle_{\beta_c, h}$ in the critical Ising model on \mathbb{Z}^2 with a homogeneous external field $h > 0$. If relation (1.20) holds (with $d = 2$, $\lambda_\delta = \hat{\lambda} \delta^{7/8}$, $h_\delta = \hat{h} \delta^{15/8}$, cf. (1.14), and we look at the case $\hat{\lambda} = 0$), differentiating both sides with respect to \hat{h} suggests that

$$\lim_{h \downarrow 0} \frac{\langle \sigma_0 \rangle_{\beta_c, h}}{h^{1/15}} = \frac{\partial \mathbf{F}}{\partial \hat{h}}(0, 1), \quad (1.23)$$

which would sharpen the results in [CGN12b]. Analogous predictions can be formulated for disorder pinning and directed polymer models (see Section 3). Of course, proving such precise estimates is likely to require substantial additional work, but having a candidate for the limiting constants, as in (1.23), can be of great help.

² A notational remark: for the directed polymer model, the denominator in (1.20) should be δ instead of δ^d , due to a different normalization of the discrete free energy (cf. Subsection 3.2); for the random field Ising model, one should set $h_\delta := \delta^{d-\gamma} \hat{h}$ without the “ $-\frac{1}{2}(\lambda_\delta)^2$ correction”, as already discussed after (1.14). These notational details are discussed in Section 3 for each model, while here we keep a unified approach.

Remark 1.3. Relation (1.20) (in a stronger form) has been proved in [BdH97, CG10] for the so-called *disordered copolymer model*, by means of a subtle coarse-graining procedure. We mention that our approach can also be applied to the copolymer model, yielding a Wiener chaos expansion as in (1.13) for the continuum partition function.

3. (First order phase transitions). Relation (1.11) can hold with $\gamma = 0$ (i.e., the k -point correlation function converges without rescaling) and with a “trivial” factorized limit:

$$\psi_{\Omega_\delta}^{(k)}(x_1, \dots, x_k) \xrightarrow{\delta \downarrow 0} \psi_\Omega^{(k)}(x_1, \dots, x_k) =: \varrho^k \quad \text{with } \varrho \in (0, \infty). \quad (1.24)$$

This is typical for a system at the critical point of a *first order phase transition* (i.e., the order parameter $m_h := \lim_{\delta \downarrow 0} |\Omega_\delta|^{-1} \mathbb{E}_{\Omega_\delta; 0, h}[\sum_{x \in \Omega_\delta} \sigma_x]$, as a function of h , has a jump discontinuity at $h = 0$). Examples include the *pinning model* for $\alpha > 1$ and the *Ising model* for $\beta > \beta_c$. Plugging (1.24) into (1.13) and performing integration (cf. [J97, §3.2]), one gets

$$\mathbf{Z}_{\Omega; \hat{\lambda}, \hat{h}}^W = \exp\{\varrho \hat{\lambda} W(\Omega) + (\varrho \hat{h} - \frac{1}{2}(\varrho \hat{\lambda})^2) \text{Leb}(\Omega)\}. \quad (1.25)$$

This explicit formula allows exact asymptotic computations on the discrete model: e.g., relation (1.13) yields for suitable values of $\zeta \in \mathbb{R}$ (when uniform integrability holds)

$$\mathbb{E}[(\mathbf{Z}_{\Omega_\delta; \lambda, h}^\omega)^\zeta] \xrightarrow{\delta \downarrow 0} \mathbb{E}[(\mathbf{Z}_{\Omega; \hat{\lambda}, \hat{h}}^W)^\zeta] = \exp\{\varrho \zeta (\hat{h} - \frac{1}{2} \varrho \hat{\lambda}^2 (1 - \zeta)) \text{Leb}(\Omega)\}. \quad (1.26)$$

Incidentally, for disordered pinning models with $\alpha > 1$, this estimate clarifies the strategy for the sharp asymptotic behavior of the critical curve $h_c(\lambda)$ as $\lambda \downarrow 0$, determined in [BC⁺14] (even though the proof therein is carried out with different techniques).

Unfortunately, the continuum partition function (1.25) can fail to capture some key properties of the discrete model, because it is shared by many “too different” models: relation (1.24) requires that the field variables under the reference law \mathbb{P}^{ref} become uncorrelated as $\delta \downarrow 0$, but is insensitive to the *correlation decay*, which could be polynomial, exponential, or even finite-range (as in the extreme case of a “trivial” reference law \mathbb{P}^{ref} , under which $(\sigma_x)_{x \in \Omega_\delta}$ are i.i.d. with $\mathbb{E}^{\text{ref}}(\sigma_x) = \varrho$). Since the correlation decay can affect the discrete free energy substantially, conjecture (1.20) usually *fails* under (1.24).

For example, for disordered pinning models with $\alpha > 1$, one always has $F(\lambda, h) \geq 0$ (there is only a polynomial cost for the underlying renewal process not to return before time N , and the energy of such a renewal configuration is 0). On the other hand, if the renewal jump distribution has finite exponential moments, then there is an exponential cost for the renewal not to return before time N , and $F(\lambda, h) < 0$ if h is sufficiently negative. Both models satisfy (1.24) with $\varrho = 1/\mathbb{E}[\tau_1]$ and thus their continuum partition functions coincide, but their free energies depend on finer detail of the renewal distribution (beyond the value of $\mathbb{E}[\tau_1]$) and are therefore radically different, causing (1.20) to fail. The continuum free energy is $F(\hat{\lambda}, \hat{h}) = \varrho \hat{h} - \frac{1}{2}(\varrho \hat{\lambda})^2$ (cf. (1.25) and (1.19)), which can attain negative values.

4. (Moment assumptions). In our convergence results, Theorems 3.1, 3.8 and 3.14, we assume that the disorder variables $(\omega_x)_{x \in \Omega_\delta}$ have finite exponential moments, which guarantees that the expectation and variance in (1.8) are well-defined. However, this assumption can be relaxed to finite moments. The necessary number of moments depends on the model and can be determined by the requirement that the typical maximum value of the variables ω_x “sampled” by the field does not exceed the reciprocal of the disorder strength λ (so that a truncation of ω_x at level λ^{-1} provides a good approximation).

For example, in the long-range directed polymer model, one expects that the path will be confined (at weak disorder) to a box of size $N \times N^{1/\alpha}$. If the disorder variables have a polynomial tail $\mathbb{P}(\omega_x > y) \approx y^{-\eta}$ as $y \uparrow \infty$, their maximum in such a box is of the order $N^{\frac{1}{\eta} \frac{1+\alpha}{\alpha}}$. Since $\lambda \approx N^{-\frac{\alpha-1}{2\alpha}}$ for this model (cf. Theorem 3.8), one expects the validity of the convergence result as long as $N^{\frac{1}{\eta} \frac{1+\alpha}{\alpha}} \ll N^{\frac{\alpha-1}{2\alpha}}$, i.e. for $\eta \geq 2(\alpha + 1)/(\alpha - 1)$. For $\alpha = 2$, this gives $\eta \geq 6$, which was conjectured in [AKQ14a] and recently proved in [DZ].

Similarly, for the pinning model the number of relevant variables is of order N and $\lambda \approx N^{-(\alpha-1/2)}$ (cf. Theorem 3.1), leading to a conjectured value $\eta \geq 2/(2\alpha - 1)$; for the RFIM, the number of relevant variables is of order N^2 and $\lambda \approx N^{-7/8}$ (cf. Theorem 3.14), leading to a conjectured value $\eta \geq 16/7$.

2. From polynomial to Wiener chaos via Lindeberg

In this section, which can be read independently of the previous one, we first recall the main properties of white noise on \mathbb{R}^d (Subsection 2.1) and define polynomial chaos expansions (Subsection 2.2). We then formulate our main general theorem (Subsection 2.3), ensuring convergence of polynomial chaos toward Wiener chaos expansions. This is based on a Lindeberg principle (Subsection 2.4) which extends results in [MOO10] to optimal second moment assumptions. Subsections 2.3 and 2.4 can be read independently.

The space of Lebesgue square-integrable functions $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is denoted by $L^2(\mathbb{R}^d)$, and we set $\|f\|_{L^2(\mathbb{R}^d)}^2 = \int_{\mathbb{R}^d} f(x)^2 dx$. For more details on the white noise, we refer to [J97, PT10].

2.1. White noise in a nutshell

By *white noise on \mathbb{R}^d* we mean a Gaussian process $W = (W(f))_{f \in L^2(\mathbb{R}^d)}$ with $\mathbb{E}[W(f)] = 0$ and $\text{Cov}(W(f), W(g)) = \int_{\mathbb{R}^d} f(x)g(x) dx$, defined on some probability space $(\Omega_W, \mathcal{A}, \mathbb{P})$. Since the specified covariance is a symmetric and positive definite function, such a process exists (and is unique in law).

If A_1, A_2, \dots are disjoint Borel sets with finite Lebesgue measure, it follows that the random variables $(W(A_i) := W(\mathbb{1}_{A_i}))_{i=1,2,\dots}$ are independent $\mathcal{N}(0, \text{Leb}(A_i))$ and the relation $W(\bigcup_{i \geq 1} A_i) = \sum_{i \geq 1} W(A_i)$ holds a.s. Consequently, it is suggestive to use the notation

$$\int_{\mathbb{R}^d} f(x) W(dx) := W(f), \quad (2.1)$$

even though $W(\cdot)$ is a.s. *not* a signed measure on \mathbb{R}^d . For $d = 1$, $W(f)$ coincides with the usual Wiener integral $\int f(t) dW_t$ with respect to the Brownian motion $W_t := W(\mathbb{1}_{[0,t]})$.

One can define a multi-dimensional stochastic integral $W^{\otimes k}(f)$, for $k \in \mathbb{N}$ and suitable $f : (\mathbb{R}^d)^k \rightarrow \mathbb{R}$, as follows. For “special indicator functions” $f = \mathbb{1}_{A_1 \times \dots \times A_k}$ built over *disjoint* bounded Borel sets $A_1, \dots, A_k \subseteq \mathbb{R}^d$, one sets $W^{\otimes k}(f) := W(\mathbb{1}_{A_1}) \cdots W(\mathbb{1}_{A_k})$. This definition is extended, by linearity, to the space \mathcal{S}_k of “special simple functions”, i.e. finite linear combinations of special indicator functions. Since a permutation of the arguments of f leaves $W^{\otimes k}(f)$ invariant, it is sufficient to consider *symmetric* functions f , which we do henceforth. One then observes that $\mathbb{E}[W^{\otimes k}(f)] = 0$ and the crucial *Itô isometry* is satisfied:³

$$\mathbb{Cov}(W^{\otimes k}(f), W^{\otimes l}(g)) = k! \mathbb{1}_{\{k=l\}} \int_{(\mathbb{R}^d)^k} f(x_1, \dots, x_k) g(x_1, \dots, x_k) dx_1 \cdots dx_k. \quad (2.2)$$

Since \mathcal{S}_k is dense in $L^2((\mathbb{R}^d)^k)$, one can finally extend the definition of $W^{\otimes k}(f)$ to every symmetric $f \in L^2((\mathbb{R}^d)^k)$ in such a way that (2.2) still holds. As in (2.1), we will write

$$\int \cdots \int_{(\mathbb{R}^d)^k} f(x_1, \dots, x_k) W(dx_1) \cdots W(dx_k) := W^{\otimes k}(f). \quad (2.3)$$

We also set $W^{\otimes 0}(c) := c$ for $c \in L^2((\mathbb{R}^d)^0) := \mathbb{R}$.

Note that $W^{\otimes k}(f)$ is a random variable defined on $(\Omega_W, \mathcal{A}, \mathbb{P})$, with zero mean (for $k \geq 1$) and finite variance, which is measurable with respect to the σ -algebra $\sigma(W)$ generated by the white noise W . (One can show that $W^{\otimes k}(f)$ is non-Gaussian for $k > 1$ and $f \neq 0$.) Remarkably, *every* square-integrable random variable X defined on Ω_W , which is measurable with respect to $\sigma(W)$, can be written as the L^2 convergent series

$$X = \sum_{k=0}^{\infty} \frac{1}{k!} W^{\otimes k}(f_k), \quad (2.4)$$

called a *Wiener chaos expansion*, for a unique choice of symmetric functions $f_k \in L^2((\mathbb{R}^d)^k)$ satisfying $\sum_{k=0}^{\infty} \frac{1}{k!} \|f_k\|_{L^2(\mathbb{R}^d)}^2 < \infty$, by (2.2). In other terms, the multiple stochastic integrals $W^{\otimes k}(f)$ span the whole Hilbert space $L^2(\Omega_W, \sigma(W), \mathbb{P})$.

2.2. Polynomial chaos

Let \mathbb{T} be a finite or countable index set (e.g., $\mathbb{T} = \{1, \dots, N\}$, $\mathbb{T} = \mathbb{N}$, $\mathbb{T} = \mathbb{Z}^d$). We set

$$\mathcal{P}^{\text{fin}}(\mathbb{T}) := \{I \subseteq \mathbb{T} : |I| < \infty\}.$$

Any function $\psi : \mathcal{P}^{\text{fin}}(\mathbb{T}) \rightarrow \mathbb{R}$ determines a (formal if $|\mathbb{T}| = \infty$) *multi-linear polynomial* Ψ :

$$\Psi(x) = \sum_{I \in \mathcal{P}^{\text{fin}}(\mathbb{T})} \psi(I) x^I, \quad \text{where } x^I := \prod_{i \in I} x_i \quad \text{with } x^\emptyset := 1. \quad (2.5)$$

We say that $\psi : \mathcal{P}^{\text{fin}}(\mathbb{T}) \rightarrow \mathbb{R}$ is the *kernel* of Ψ .

³ For the isometry (2.2) it is essential to “avoid diagonals”: this is the reason for taking special indicator functions, corresponding to products of *disjoint* Borel sets.

Let now $\zeta := (\zeta_i)_{i \in \mathbb{T}}$ be a family of independent (but not necessarily identically distributed) random variables. We say that a random variable X admits a *polynomial chaos expansion* with respect to ζ if it can be expressed as $X = \Psi(\zeta) = \Psi((\zeta_i)_{i \in \mathbb{T}})$ for some multi-linear polynomial Ψ . Of course, when $|\mathbb{T}| = \infty$ some care is needed: by $X = \Psi(\zeta)$ we mean that for any sequence $\Lambda_N \subset \mathbb{T}$ with $|\Lambda_N| < \infty$ and $\Lambda_N \uparrow \mathbb{T}$ one has

$$X = \lim_{N \rightarrow \infty} \sum_{I \subseteq \Lambda_N} \psi(I) \zeta^I \quad \text{in probability.} \quad (2.6)$$

Remark 2.1. When $\text{Var}(\zeta_i) \neq 0$ for all $i \in \mathbb{T}$, we can assume that all the variances are equal with no loss of generality: it suffices to redefine $\psi(I) \rightarrow \psi(I) (\prod_{i \in I} \text{Var}(\zeta_i))^{-1/2}$.

Remark 2.2. When the independent random variables $\zeta := (\zeta_i)_{i \in \mathbb{T}}$ have zero mean and the same variance σ^2 , an easy sufficient condition for (2.6), with L^2 convergence, is

$$\sum_{I \in \mathcal{P}^{\text{fin}}(\mathbb{T})} (\sigma^2)^{|I|} \psi(I)^2 < \infty, \quad (2.7)$$

because $\mathbb{E}[\zeta^I \zeta^J] = 0$ for $I \neq J$. For variables with non-zero mean $\mu := (\mu_i)_{i \in \mathbb{T}}$ (always with the same variance σ^2), sharp conditions for L^2 convergence in (2.6) involve μ and the kernel ψ jointly. As we show below, practical sufficient ‘‘factorized’’ conditions are

$$\sum_{i \in \mathbb{T}} \mu_i^2 < \infty, \quad \exists \varepsilon > 0 : \sum_{I \in \mathcal{P}^{\text{fin}}(\mathbb{T})} (1 + \varepsilon)^{|I|} (\sigma^2)^{|I|} \psi(I)^2 < \infty. \quad (2.8)$$

2.3. Convergence of polynomial chaos to Wiener chaos

Consider for $\delta \in (0, 1)$ an index set $\mathbb{T}_\delta \subset \mathbb{R}^d$ and a family of polynomial chaos expansions $(\Psi_\delta(\zeta_\delta))_{\delta \in (0, 1)}$, defined from kernels $\psi_\delta : \mathcal{P}^{\text{fin}}(\mathbb{T}_\delta) \rightarrow \mathbb{R}$ and from independent random variables $\zeta_\delta := (\zeta_{\delta, x})_{x \in \mathbb{T}_\delta}$. If \mathbb{T}_δ converges to the continuum \mathbb{R}^d as $\delta \downarrow 0$ (e.g., $\mathbb{T}_\delta := (\delta \mathbb{Z})^d$), then after suitable scaling, the random variables $(\zeta_{\delta, x})_{x \in \mathbb{T}_\delta}$ approximate the white noise $W(dx)$ on \mathbb{R}^d . If the kernel ψ_δ , suitably rescaled, converges as $\delta \downarrow 0$ to a continuum kernel $\psi_0 : \mathcal{P}^{\text{fin}}(\mathbb{R}^d) \rightarrow \mathbb{R}$, it is plausible that the polynomial chaos expansion $\Psi_\delta(\zeta_\delta)$ approximates a *Wiener chaos expansion* Ψ_0 (cf. (2.3)–(2.4)) with kernel ψ_0 . This is precisely what we are going to show.

First we introduce some notation. Each random variable ζ_x indexed by a point x in an index set $\mathbb{T} \subset \mathbb{R}^d$ will be associated with a cell in \mathbb{R}^d containing x , and functions defined on \mathbb{T}^k will be extended to functions defined on $(\mathbb{R}^d)^k$.

- Let $\mathcal{B}(\mathbb{R}^d)$ denote the Borel subsets of \mathbb{R}^d . Given a locally finite set $\mathbb{T} \subset \mathbb{R}^d$, we call $\mathcal{C} : \mathbb{T} \rightarrow \mathcal{B}(\mathbb{R}^d)$ a *tessellation* of \mathbb{R}^d indexed by \mathbb{T} if $(\mathcal{C}(x))_{x \in \mathbb{T}}$ form a disjoint partition of \mathbb{R}^d such that $x \in \mathcal{C}(x)$ for each $x \in \mathbb{T}$. We call $\mathcal{C}(x)$ the *cell* associated with $x \in \mathbb{T}$. In most cases, $(\mathcal{C}(x))_{x \in \mathbb{T}}$ will be the cells of a cubic lattice. However, there are natural examples where this is not the case, such as the directed polymer model defined from a simple symmetric random walk, or the Ising model defined on non-cubic lattices.

- Once a tessellation \mathcal{C} is fixed, any function $f : \mathbb{T} \rightarrow \mathbb{R}$ is automatically extended to $f : \mathbb{R}^d \rightarrow \mathbb{R}$ by setting $f(y) := f(x)$ for all $y \in \mathcal{C}(x)$, for each $x \in \mathbb{T}$. Note that for such extensions, $\|f\|_{L^2(\mathbb{R}^d)}^2 = \sum_{x \in \mathbb{T}} f(x)^2 \text{Leb}(\mathcal{C}(x))$.
- Analogously, for any $\psi : \mathcal{P}^{\text{fin}}(\mathbb{T}) \rightarrow \mathbb{R}$, we first extend it to $\psi : \bigcup_{k=0}^{\infty} \mathbb{T}^k \rightarrow \mathbb{R}$ by setting $\psi(x_1, \dots, x_k) := \psi(\{x_1, \dots, x_k\})$ if the x_i are distinct, and $\psi(x_1, \dots, x_k) := 0$ otherwise. We then extend it to $\psi : \bigcup_{k=0}^{\infty} (\mathbb{R}^d)^k \rightarrow \mathbb{R}$ by assigning value $\psi(x_1, \dots, x_k)$ to all points in $\mathcal{C}(x_1) \times \dots \times \mathcal{C}(x_k)$, for each $k \in \mathbb{N}$ and $x_1, \dots, x_k \in \mathbb{T}$.
- Given $\psi : \mathcal{P}^{\text{fin}}(\mathbb{R}^d) \rightarrow \mathbb{R}$, its extension to $\psi : \bigcup_{k=0}^{\infty} (\mathbb{R}^d)^k \rightarrow \mathbb{R}$ is defined similarly (no cells involved). It will be clear from the context which version of ψ is being used.
- Finally, given a measurable function $\psi : \bigcup_{k=0}^{\infty} (\mathbb{R}^d)^k \rightarrow \mathbb{R}$, we denote by $\|\psi\|_{L^2((\mathbb{R}^d)^k)}$ the L^2 norm of the restriction of ψ to $(\mathbb{R}^d)^k$, i.e.

$$\|\psi\|_{L^2((\mathbb{R}^d)^k)}^2 = \int \dots \int_{(\mathbb{R}^d)^k} \psi(x_1, \dots, x_k)^2 dx_1 \dots dx_k.$$

We are now ready to state our main convergence result, proved in Section 5.

Theorem 2.3 (Convergence of polynomial chaos to Wiener chaos, L^2 case). *Assume that for $\delta \in (0, 1)$ the following ingredients are given:*

- Let \mathbb{T}_δ be a locally finite subset of \mathbb{R}^d .
- Let $\zeta_\delta := (\zeta_{\delta,x})_{x \in \mathbb{T}_\delta}$ be independent random variables in L^2 with the same variance,

$$\mathbb{E}[\zeta_{\delta,x}] = \mu_\delta(x) \quad \text{and} \quad \text{Var}(\zeta_{\delta,x}) = \sigma_\delta^2,$$

such that $((\zeta_{\delta,x} - \mathbb{E}[\zeta_{\delta,x}])^2)_{\delta \in (0,1), x \in \mathbb{T}_\delta}$ are uniformly integrable.

- Let $\Psi_\delta(z)$ be a formal multi-linear polynomial with kernel $\psi_\delta : \mathcal{P}^{\text{fin}}(\mathbb{T}_\delta) \rightarrow \mathbb{R}$ (cf. (2.5)).
- Let \mathcal{C}_δ be a tessellation of \mathbb{R}^d indexed by \mathbb{T}_δ , where every cell $\mathcal{C}_\delta(x)$ has the same volume $v_\delta := \text{Leb}(\mathcal{C}_\delta(x))$.

Assume that $v_\delta \rightarrow 0$ as $\delta \downarrow 0$, and that the following conditions are satisfied:

- (i) There exist $\sigma_0 \in (0, \infty)$ and $\mu_0 \in L^2(\mathbb{R}^d)$ such that

$$\lim_{\delta \downarrow 0} \sigma_\delta = \sigma_0, \quad \lim_{\delta \downarrow 0} \|\bar{\mu}_\delta - \mu_0\|_{L^2(\mathbb{R}^d)} = 0, \quad \text{where} \quad \bar{\mu}_\delta(x) := v_\delta^{-1/2} \mu_\delta(x). \quad (2.9)$$

- (ii) There exists $\psi_0 : \mathcal{P}^{\text{fin}}(\mathbb{R}^d) \rightarrow \mathbb{R}$, with $\|\psi_0\|_{L^2((\mathbb{R}^d)^k)} < \infty$ for every $k \in \mathbb{N}_0$ such that

$$\lim_{\delta \downarrow 0} \|\bar{\psi}_\delta - \psi_0\|_{L^2((\mathbb{R}^d)^k)}^2 = 0, \quad \text{where} \quad \bar{\psi}_\delta(I) := v_\delta^{-|I|/2} \psi_\delta(I). \quad (2.10)$$

- (iii) For some $\varepsilon > 0$ (or even $\varepsilon = 0$ if $\mu_\delta(x) \equiv 0$)

$$\lim_{\ell \rightarrow \infty} \limsup_{\delta \downarrow 0} \sum_{I \in \mathcal{P}^{\text{fin}}(\mathbb{T}_\delta), |I| > \ell} (1 + \varepsilon)^{|I|} (\sigma_\delta^2)^{|I|} \psi_\delta(I)^2 = 0. \quad (2.11)$$

Then the polynomial chaos expansion $\Psi_\delta(\zeta_\delta)$ is well-defined and converges in distribution

as $\delta \downarrow 0$ to a random variable Ψ_0 with an explicit Wiener chaos expansion:

$$\Psi_\delta(\zeta_\delta) \xrightarrow[\delta \downarrow 0]{d} \Psi_0 := \sum_{k=0}^{\infty} \frac{1}{k!} \int \cdots \int_{(\mathbb{R}^d)^k} \psi_0(y_1, \dots, y_k) \prod_{i=1}^k (\sigma_0 W(dy_i) + \mu_0(y_i) dy_i), \quad (2.12)$$

where $W(\cdot)$ denotes a white noise on \mathbb{R}^d .

The series in (2.12) converges in L^2 , and $\mathbb{E}[\Psi_\delta(\zeta_\delta)^2] \rightarrow \mathbb{E}[\Psi_0^2]$. Consequently, for any coupling of $\Psi_\delta(\zeta_\delta)$ and Ψ_0 such that $\Psi_\delta(\zeta_\delta) \rightarrow \Psi_0$ a.s., one has $\mathbb{E}[|\Psi_\delta(\zeta_\delta) - \Psi_0|^2] \rightarrow 0$.

The convergence (2.12) extends to the joint distribution of a finite collection of polynomial chaos expansions $(\Psi_{i,\delta}(\zeta_\delta))_{1 \leq i \leq M}$ provided $(\Psi_{i,\delta})_{\delta \in (0,1)}$ satisfies (ii)–(iii) above for each i .

Remark 2.4. Let us be more precise about the random variable Ψ_0 in (2.12). Setting $\nu(x) := \mu_0(x)/\sigma_0$, we can rewrite it as

$$\Psi_0 = \sum_{k=0}^{\infty} \frac{1}{k!} \int \cdots \int_{(\mathbb{R}^d)^k} \psi_0(y_1, \dots, y_k) \sigma_0^k \prod_{i=1}^k (W(dy_i) + \nu(y_i) dy_i), \quad (2.13)$$

which can be viewed as a “Wiener chaos expansion with respect to the *biased* white noise $W_\nu(dx) := W(dx) + \nu(x)dx$ ”. The rigorous definition of such an expansion goes as follows. For every fixed $k \in \mathbb{N}$, the integral over $(\mathbb{R}^d)^k$ in (2.13) can be defined by expanding the product and integrating out the “deterministic coordinates” (those corresponding to $\nu(y_i)dy_i$), obtaining a finite sum of well-defined (lower-dimensional) ordinary stochastic integrals, as in (2.3). After regrouping terms, the series in (2.13) becomes an ordinary Wiener chaos expansion, as in (2.4). In analogy with the polynomial case (2.8), we show in Section 5 that the L^2 convergence of the series is ensured by the conditions that $\mu_0 \in L^2(\mathbb{R}^d)$ and that

$$\exists \varepsilon > 0 : \sum_{k=0}^{\infty} \frac{1}{k!} (1 + \varepsilon)^k (\sigma_0^2)^k \|\psi_0\|_{L^2((\mathbb{R}^d)^k)}^2 < \infty, \quad (2.14)$$

which follow from assumptions (i)–(iii) of Theorem 2.3.

2.3.1. Beyond the L^2 case. There is a useful alternative interpretation of (2.12)–(2.13). If $(\Omega_W, \mathcal{A}, \mathbb{P})$ is the probability space on which the white noise $W = (W(f))_{f \in L^2(\mathbb{R}^d)}$ is defined, for every $\nu \in L^2(\mathbb{R}^d)$ we introduce a new probability \mathbb{P}_ν on Ω_W by

$$\frac{d\mathbb{P}_\nu}{d\mathbb{P}} := e^{W(\nu) - \frac{1}{2}\mathbb{E}[W(\nu)^2]} = e^{\int_{\mathbb{R}^d} \nu(x) W(dx) - \frac{1}{2} \int_{\mathbb{R}^d} \nu(x)^2 dx}. \quad (2.15)$$

It turns out that the “biased stochastic integrals” in (2.13) have the same joint distribution as the ordinary stochastic integrals (with ν replaced by 0) under the probability \mathbb{P}_ν , by the *Cameron–Martin theorem* (cf. Appendix A). As a consequence, the random variable Ψ_0 in (2.12) enjoys the following equality in distribution, with $\nu(x) := \mu_0(x)/\sigma_0$:

$$\Psi_0 \stackrel{d}{=} \sum_{k=0}^{\infty} \frac{1}{k!} \int \cdots \int_{(\mathbb{R}^d)^k} \psi_0(y_1, \dots, y_k) \sigma_0^k W(dy_1) \cdots W(dy_k) \quad \text{under } \mathbb{P}_\nu \quad (2.16)$$

provided the series (either in (2.12) or (2.16), equivalently) converges in probability.

Let us now assume the weaker version of relation (2.14) for $\varepsilon = 0$ holds, i.e.

$$\sum_{k=0}^{\infty} \frac{1}{k!} (\sigma_0^2)^k \|\Psi_0\|_{L^2((\mathbb{R}^d)^k)} < \infty. \quad (2.17)$$

Under this condition, the series in (2.16) converges in L^2 under the original probability \mathbb{P} , by the Itô isometry (2.2). Since the Radon–Nikodym density (2.15) has finite moments of all orders, it follows by (2.16) and an application of the Hölder inequality (see (5.11) for the details) that the series in (2.12) defining Ψ_0 converges in L^p for every $p \in (0, 2)$ when (2.17) holds (even though it might not converge in L^2 if (2.14) fails).

As a consequence, by performing an L^p analysis for $p < 2$, we can weaken condition (iii) in Theorem 2.3, setting $\varepsilon = 0$ in (2.11), under mild restrictions on the disorder distribution (due to the implementation of a change of measure as in (2.16) for polynomial chaos).

Theorem 2.5 (Convergence of polynomial chaos to Wiener chaos, L^{2-} case). *Let the assumptions of Theorem 2.3 hold, with condition (iii) weakened by setting $\varepsilon = 0$ in (2.11). Assume further that $\lim_{\delta \downarrow 0} \|\mu_\delta\|_\infty = 0$, and that either of the following two conditions is satisfied:*

- (a) $\inf_{\delta \in (0,1), x \in \mathbb{T}_\delta} \min\{\mathbb{P}(\zeta_{\delta,x} > 0), \mathbb{P}(\zeta_{\delta,x} < 0), \text{Var}(\zeta_{\delta,x} \mid \zeta_{\delta,x} > 0), \text{Var}(\zeta_{\delta,x} \mid \zeta_{\delta,x} < 0)\} > 0;$
- (b) $\forall C > 0: \lim_{\delta \downarrow 0} \sum_{I \in \mathcal{P}^{\text{fin}}(\mathbb{T}_\delta), |I| > \|\mu_\delta\|_\infty^{-1}} e^{C\|\mu_\delta\|_\infty |I|} (\sigma_\delta^2)^{|I|} \psi_\delta(I)^2 = 0.$ (2.18)

Then the polynomial chaos expansion $\Psi_\delta(\zeta_\delta)$ is well-defined and converges in distribution as $\delta \downarrow 0$ to the random variable Ψ_0 defined by (2.12), or equivalently (2.16). For all $p \in (0, 2)$, the series therein converges in L^p , and furthermore $\mathbb{E}[|\Psi_\delta|^p] \rightarrow \mathbb{E}[|\Psi_0|^p]$. The conclusion extends to finite collections $(\Psi_{i,\delta}(\zeta_\delta))_{1 \leq i \leq M}$.

2.4. Lindeberg principle for polynomial chaos

The key ingredients in our proof of Theorem 2.3 are two Lindeberg principles for polynomial chaos. As we discuss in Remark 2.7, they extend [MOO10, Theorem 3.18] in two ways: firstly, we relax the finite third moment assumption of [MOO10] to an optimal condition of uniform integrability of the square of the random variables; secondly, we allow random variables with non-zero mean.

We point out that the first extension is actually not needed for our applications to disordered systems, due to the assumption of finite exponential moments for the disorder random variables. However, it is an extension of general interest and will be useful if one attempts to weaken the moment assumptions on the disorder random variables, as discussed at the end of Section 1.3. We remark that Lindeberg principles have also played crucial roles in recent breakthrough results on random matrices [C06, TV11].

Given a polynomial chaos expansion $\Psi(\zeta)$ with respect to a family $\zeta := (\zeta_i)_{i \in \mathbb{T}}$ of independent random variables (cf. Subsection 2.2), we will control how the distribution of $\Psi(\zeta)$ changes when we replace ζ by independent Gaussian random variables $\xi := (\xi_i)_{i \in \mathbb{T}}$ with the same mean and variance as ζ .

Given a multi-linear polynomial $\Psi(x) = \Psi((x_i)_{i \in \mathbb{T}})$ as in (2.5), with kernel ψ , we set

$$C_\Psi := \sum_{I \in \mathcal{P}^{\text{fin}}(\mathbb{T}), I \neq \emptyset} \psi(I)^2, \quad (2.19)$$

and define the *influence* of the i -th variable x_i on Ψ by

$$\text{Inf}_i[\Psi] := \sum_{I \in \mathcal{P}^{\text{fin}}(\mathbb{T}), I \ni i} \psi(I)^2. \quad (2.20)$$

Note that if $\zeta = (\zeta_i)_{i \in \mathbb{T}}$ are independent random variables with zero mean and unit variance,

$$C_\Psi = \text{Var}[\Psi(\zeta)], \quad \text{Inf}_i[\Psi] = \mathbb{E}[\text{Var}[\Psi(\zeta) \mid (\zeta_j)_{j \in \mathbb{T} \setminus \{i\}}]],$$

which is just the influence of the random variable ζ_j on $\Psi(\zeta)$ introduced in [MOO10] (for more on the notion of influence, see e.g. [KKL88, BK92] and the references in [MOO10]). We also define the degree ℓ truncations $\Psi^{\leq \ell}$ and $\Psi^{> \ell}$ of the multi-linear polynomial Ψ by

$$\Psi^{\leq \ell}(x) := \sum_{I \in \mathcal{P}^{\text{fin}}(\mathbb{T}), |I| \leq \ell} \psi(I)x^I, \quad \Psi^{> \ell}(x) := \sum_{I \in \mathcal{P}^{\text{fin}}(\mathbb{T}), |I| > \ell} \psi(I)x^I, \quad (2.21)$$

whose kernels will be denoted by $\psi^{\leq \ell}(I) = \psi(I)\mathbb{1}_{\{|I| \leq \ell\}}$ and $\psi^{> \ell}(I) = \psi(I)\mathbb{1}_{\{|I| > \ell\}}$.

We are now ready to state and comment on our Lindeberg principles, which will be proved in Section 4.

Theorem 2.6 (Lindeberg principle, zero mean case). *Let $\zeta = (\zeta_i)_{i \in \mathbb{T}}$ and $\xi = (\xi_i)_{i \in \mathbb{T}}$ be families of independent random variables with zero mean and unit variance. Let $\Psi(x)$ be a multi-linear polynomial as in (2.5) with $C_\Psi = \sum_{I \in \mathcal{P}^{\text{fin}}(\mathbb{T})} \psi(I)^2 < \infty$. Then the polynomial chaos expansions $\Psi(\zeta)$, $\Psi(\xi)$ are well-defined L^2 random variables. For $M \in [0, \infty]$ defining the maximal truncated moments*

$$m_2^{>M} := \sup_{X \in \bigcup_{i \in \mathbb{T}} \{\zeta_i, \xi_i\}} \mathbb{E}[X^2 \mathbb{1}_{|X| > M}], \quad m_3^{\leq M} := \sup_{X \in \bigcup_{i \in \mathbb{T}} \{\zeta_i, \xi_i\}} \mathbb{E}[|X|^3 \mathbb{1}_{|X| \leq M}]. \quad (2.22)$$

Then for every $f : \mathbb{R} \rightarrow \mathbb{R}$ of class \mathcal{C}^3 with

$$C_f := \max\{\|f'\|_\infty, \|f''\|_\infty, \|f'''\|_\infty\} < \infty, \quad (2.23)$$

every $\ell \in \mathbb{N}$, and every $M \in (0, \infty]$ large enough such that $m_2^{>M} \leq 1/4$, one has

$$\begin{aligned} |\mathbb{E}[f(\Psi(\zeta))] - \mathbb{E}[f(\Psi(\xi))]| &\leq C_f \left\{ 2\sqrt{C_{\Psi^{> \ell}}} + C_{\Psi^{\leq \ell}} 16\ell^2 m_2^{>M} \right. \\ &\quad \left. + C_{\Psi^{\leq \ell}} 70^{\ell+1} (m_3^{\leq M})^\ell \sqrt{\max_{i \in \mathbb{T}} \text{Inf}_i[\Psi^{\leq \ell}]} \right\}, \quad (2.24) \end{aligned}$$

where C_\cdot , $\text{Inf}_i[\cdot]$ and $\Psi^{\leq \ell}$, $\Psi^{> \ell}$ are defined in (2.19)–(2.21).

Intuitively, this theorem shows that $\Psi(\zeta)$ and $\Psi(\xi)$ are close in distribution when the right hand side of (2.24) is small. Despite their technical appearance, each of the three terms inside the brackets can be easily controlled:

- The first term is controlled by $C_{\Psi^{>\ell}} = \sum_{|I|>\ell} \psi(I)^2$, which is small for ℓ large.
- The second term is controlled by $m_2^{>M}$, which is small for M large if the random variables $(\zeta_i)_{i \in \mathbb{T}}$ and $(\xi_i)_{i \in \mathbb{T}}$ have *uniformly integrable squares* (e.g., if they are i.i.d.).
- The third term is controlled by the *maximal influence* $\max_{i \in \mathbb{T}} \text{Inf}_i[\Psi^{\leq \ell}]$, which is small if the multi-linear polynomial $\Psi^{\leq \ell}$ is sufficiently “spread out”.

In particular, we shall see that the conditions of Theorem 2.3 allow us to exploit (2.24).

Remark 2.7. When the polynomial $\Psi = \Psi^{\leq \ell}$ has degree ℓ and the random variables ζ_i, ξ_i have third absolute moments bounded by $m_3 < \infty$, relation (2.24) for $M = \infty$ reduces to

$$|\mathbb{E}[f(\Psi(\zeta))] - \mathbb{E}[f(\Psi(\xi))]| \leq C_f C_\Psi 70^{\ell+1} (m_3)^\ell \max_{i \in \mathbb{T}} \sqrt{\text{Inf}_i[\Psi]}.$$

This is the key estimate proved by Mossel, O’Donnell and Oleszkiewicz [MOO10, Theorem 3.18 under hypothesis **H2**], with the prefactor $70^{\ell+1}$ instead of 30^ℓ . Our Theorem 2.6 thus provides an extension of [MOO10, Theorem 3.18] to finite second-moment assumptions.

Some of the results in [MOO10] are formulated in the more general setting of multi-linear polynomials over *orthonormal ensembles*. Although we stick for simplicity to the case of independent random variables, our approach can also be adapted to deal with orthonormal ensembles. In fact, we follow the same line of proof of [MOO10], which is based on Lindeberg’s original approach, with two refinements: a sharper approximation of the remainder in Taylor’s expansion and a fine truncation on the random variables (cf. Section 4 for details).

As a corollary to Theorem 2.6, we can treat the case where we add non-zero mean to the random variables $(\zeta_i)_{i \in \mathbb{T}}$ and $(\xi_i)_{i \in \mathbb{T}}$. The following result is also proved in Section 4.

Theorem 2.8 (Lindeberg principle, non-zero mean case). *Let $\zeta = (\zeta_i)_{i \in \mathbb{T}}$ and $\xi = (\xi_i)_{i \in \mathbb{T}}$ be as in Theorem 2.6, and define the maximal truncated moments $m_2^{>M}, m_3^{\leq M}$ by (2.22). Let $\mu := (\mu_i)_{i \in \mathbb{T}}$ be a family of real numbers with*

$$c_\mu := \sum_{i \in \mathbb{T}} \mu_i^2 < \infty, \quad (2.25)$$

and define the μ -biased families $\tilde{\zeta} := \zeta + \mu = (\zeta_i + \mu_i)_{i \in \mathbb{T}}$ and $\tilde{\xi} := \xi + \mu = (\xi_i + \mu_i)_{i \in \mathbb{T}}$. Let $\Psi(x)$ be a multi-linear polynomial as in (2.5). Setting for $\varepsilon > 0$,

$$\Psi^{(\varepsilon)}(x) = \sum_{I \in \mathcal{P}^{\text{fin}}(\mathbb{T})} (1 + \varepsilon)^{|I|/2} \psi(I) x^I, \quad (2.26)$$

assume that $C_{\Psi^{(\varepsilon)}} = \sum_{I \in \mathcal{P}^{\text{fin}}(\mathbb{T})} (1 + \varepsilon)^{|I|} \psi(I)^2 < \infty$ for some $\varepsilon > 0$. Then the polynomial chaos expansions $\Psi(\tilde{\zeta})$ and $\Psi(\tilde{\xi})$ are well-defined L^2 random variables.

For every $f : \mathbb{R} \rightarrow \mathbb{R}$ of class \mathcal{C}^3 with $C_f < \infty$ (cf. (2.23)), every $\ell \in \mathbb{N}$ and every $M \in [0, \infty]$ large enough such that $m_2^{>M} \leq 1/4$, one has

$$\begin{aligned} |\mathbb{E}[f(\Psi(\tilde{\zeta}))] - \mathbb{E}[f(\Psi(\tilde{\xi}))]| &\leq e^{2c_\mu/\varepsilon} C_f \left\{ 2\sqrt{C_{\Psi^{(\varepsilon), >\ell}}} + C_{\Psi^{(\varepsilon), \leq \ell}} 16\ell^2 m_2^{>M} \right. \\ &\quad \left. + C_{\Psi^{(\varepsilon), \leq \ell}} 70^{\ell+1} (m_3^{\leq M})^\ell \sqrt{\max_{i \in \mathbb{T}} \text{Inf}_i[\Psi^{(\varepsilon), \leq \ell}]} \right\}, \end{aligned} \quad (2.27)$$

where C_\cdot , $\text{Inf}_i[\cdot]$ are defined in (2.19), (2.20) and $\Psi^{(\varepsilon), >\ell}$, $\Psi^{(\varepsilon), \leq \ell}$ are defined as in (2.21).

3. Scaling limits of disordered systems

In this section, which can be read independently of Sections 1 and 2, we consider three much-studied statistical mechanics models: the *disordered pinning model* (Subsection 3.1), the (long-range) *directed polymer model* in dimension $1 + 1$ (Subsection 3.2), and the two-dimensional *random field Ising model* (Subsection 3.3). For each model, we show that the partition function has a non-trivial limit in distribution, in the continuum and weak disorder regime, given by an explicit Wiener chaos expansion with respect to the white noise on \mathbb{R}^d (see Subsection 2.1 for some reminders). The proofs, given in Sections 6–8, are based on the general convergence results of Section 2 (cf. Theorem 2.3).

For each model, the disorder will be given by a countable family of i.i.d. random variables ω_i with zero mean, finite variance and locally finite exponential moments:

$$\mathbb{E}[\omega_i] = 0, \quad \text{Var}(\omega_i) = 1, \quad \exists t_0 > 0 : \Lambda(t) := \log \mathbb{E}[e^{t\omega_i}] < \infty \text{ for } |t| < t_0. \quad (3.1)$$

Our approach actually works in the much more general setting when the disorder is given by a triangular array of independent (but not necessarily identically distributed) random variables, in the spirit of Theorem 2.3, but we stick to the i.i.d. case for the sake of simplicity.

3.1. Disordered pinning model

Consider a discrete renewal process $\tau := \{\tau_n\}_{n \geq 0}$, that is, $\tau_0 = 0$ and the increments $\{\tau_n - \tau_{n-1}\}_{n \geq 1}$ are i.i.d. \mathbb{N} -valued random variables. We assume that τ is *non-terminating*, that is, $\mathbb{P}(\tau_1 < \infty) = 1$, and that

$$\mathbb{P}(\tau_1 = n) = L(n)/n^{1+\alpha}, \quad \forall n \in \mathbb{N}, \quad (3.2)$$

where $\alpha \in [0, \infty)$ and $L : (0, \infty) \rightarrow (0, \infty)$ is a slowly varying function [BGT87]. One could also consider the periodic case, when (3.2) holds for $n \in p\mathbb{N}$ and $\mathbb{P}(\tau_1 = n) = 0$ if $n \notin p\mathbb{N}$, for some period $p \in \mathbb{N}$. For simplicity, we focus on the aperiodic case $p = 1$.

Let $\omega = (\omega_n)_{n \in \mathbb{N}_0}$ be a sequence of i.i.d. random variables, independent of τ , satisfying (3.1). The *disordered pinning model* is the random probability law $\mathbb{P}_{N, \beta, h}^\omega$ on subsets on \mathbb{N}_0 , indexed by ω and by $N \in \mathbb{N}$, $\beta \geq 0$ and $h \in \mathbb{R}$, defined by

$$d\mathbb{P}_{N, \beta, h}^\omega(\tau) := \frac{1}{Z_{N, \beta, h}^\omega} e^{\sum_{n=1}^N (\beta \omega_n - \Lambda(\beta) + h)} \mathbb{1}_{\{n \in \tau\}} d\mathbb{P}(\tau),$$

where we recall that $\Lambda(\beta) := \log \mathbb{E}[e^{\beta\omega_1}]$, and the *partition function* $Z_{N,\beta,h}^\omega$ is defined by

$$Z_{N,\beta,h}^\omega := \mathbb{E}\left[e^{\sum_{n=1}^N (\beta\omega_n - \Lambda(\beta) + h)\mathbb{1}_{\{n \in \tau\}}}\right]. \quad (3.3)$$

We also consider the *conditioned partition function*

$$Z_{N,\beta,h}^{\omega,c} := \mathbb{E}\left[e^{\sum_{n=1}^N (\beta\omega_n - \Lambda(\beta) + h)\mathbb{1}_{\{n \in \tau\}}} \mid N \in \tau\right]. \quad (3.4)$$

The disordered pinning model exhibits an interesting localization/delocalization phase transition. This can be quantified via the (quenched) free energy, which is defined as

$$F(\beta, h) := \lim_{N \rightarrow \infty} \frac{1}{N} \log Z_{N,\beta,h}^\omega = \lim_{N \rightarrow \infty} \frac{1}{N} \mathbb{E}\left[\log Z_{N,\beta,h}^\omega\right] \quad \mathbb{P}(d\omega)\text{-a.s.} \quad (3.5)$$

By restricting the partition function to configurations such that $\tau \cap [1, N] = \emptyset$, it is easily seen that $F(\beta, h) \geq 0$. The localized and delocalized regimes (\mathcal{L} , \mathcal{D} respectively) can be defined as

$$\mathcal{L} := \{(\beta, h) : F(\beta, h) > 0\}, \quad \mathcal{D} := \{(\beta, h) : F(\beta, h) = 0\}.$$

We refer to [G10] for more information on the structure of the phase transition, and in particular for quantitative estimates on the *critical curve*

$$h_c(\beta) := \sup\{h \in \mathbb{R} : F(\beta, h) = 0\} = \inf\{h \in \mathbb{R} : F(\beta, h) > 0\}. \quad (3.6)$$

We can now state our main result on the disordered pinning model, to be proved in Section 6. To lighten notation, we write $Z_{Nt,\beta,h}^\omega$ to mean $Z_{\lfloor Nt \rfloor, \beta, h}^\omega$.

Theorem 3.1 (Scaling limit of disordered pinning models). *Let the aperiodic renewal process τ either satisfy (3.2) for some $\alpha \in (1/2, 1)$, or have finite mean $\mathbb{E}[\tau_1] < \infty$ (which happens in particular when (3.2) holds with $\alpha > 1$). For $N \in \mathbb{N}$, $\hat{\beta} > 0$ and $\hat{h} \in \mathbb{R}$, set*

$$\beta_N = \begin{cases} \hat{\beta} \frac{L(N)}{N^{\alpha-1/2}} & \text{if } 1/2 < \alpha < 1, \\ \hat{\beta} \frac{1}{\sqrt{N}} & \text{if } \mathbb{E}[\tau_1] < \infty, \end{cases} \quad h_N = \begin{cases} \hat{h} \frac{L(N)}{N^\alpha} & \text{if } 1/2 < \alpha < 1, \\ \hat{h} \frac{1}{N} & \text{if } \mathbb{E}[\tau_1] < \infty. \end{cases} \quad (3.7)$$

Then, for every $t \geq 0$, the conditioned partition function $Z_{Nt,\beta_N,h_N}^{\omega,c}$ of the disordered pinning model converges in distribution as $N \rightarrow \infty$ to the random variable $\mathbf{Z}_{t,\hat{\beta},\hat{h}}^{W,c}$ given by

$$\mathbf{Z}_{t,\hat{\beta},\hat{h}}^{W,c} := 1 + \sum_{k=1}^{\infty} \frac{1}{k!} \int \cdots \int_{[0,t]^k} \psi_t^c(t_1, \dots, t_k) \prod_{i=1}^k (\hat{\beta} W(dt_i) + \hat{h} dt_i), \quad (3.8)$$

where $W(\cdot)$ denotes white noise on \mathbb{R} and $\psi_t^c(t_1, \dots, t_k)$ is a symmetric function, defined for $0 < t_1 < \dots < t_k < t$ by

$$\psi_t^c(t_1, \dots, t_k) = \begin{cases} \frac{C_\alpha^k t^{1-\alpha}}{t_1^{1-\alpha} (t_2 - t_1)^{1-\alpha} \dots (t_k - t_{k-1})^{1-\alpha} (t - t_k)^{1-\alpha}} & \text{if } 1/2 < \alpha < 1, \\ \frac{1}{\mathbb{E}[\tau_1]^k} & \text{if } \mathbb{E}[\tau_1] < \infty, \end{cases} \quad (3.9)$$

where $C_\alpha := \alpha \sin(\pi\alpha)/\pi$. The series in (3.8) converges in L^2 , and one has the convergence of the corresponding second moments: $\mathbb{E}[(Z_{Nt, \beta_N, h_N}^{\omega, c})^2] \rightarrow \mathbb{E}[(Z_{t, \hat{\beta}, \hat{h}}^{W, c})^2]$ as $N \rightarrow \infty$.

An analogous statement holds for the free (unconditioned) partition function $Z_{Nt, \beta_N, h_N}^\omega$, where the limiting random variable $Z_{t, \hat{\beta}, \hat{h}}^W$ is defined as in (3.8), with kernel

$$\psi(t_1, \dots, t_k) := \begin{cases} \frac{C_\alpha^k}{t_1^{1-\alpha} (t_2 - t_1)^{1-\alpha} \dots (t_k - t_{k-1})^{1-\alpha}} & \text{if } 1/2 < \alpha < 1, \\ \frac{1}{\mathbb{E}[\tau_1]^k} & \text{if } \mathbb{E}[\tau_1] < \infty. \end{cases} \quad (3.10)$$

When $\mathbb{E}[\tau_1] < \infty$, for both the free and conditioned case, the continuum partition function has an explicit distribution: for every $t \geq 0$

$$Z_{t, \hat{\beta}, \hat{h}}^{W, c} \stackrel{d}{=} Z_{t, \hat{\beta}, \hat{h}}^W \stackrel{d}{=} \exp\left\{ \frac{\hat{\beta}}{\mathbb{E}[\tau_1]} W_t + \left(\frac{\hat{h}}{\mathbb{E}[\tau_1]} - \frac{\hat{\beta}^2}{2\mathbb{E}[\tau_1]^2} \right) t \right\}, \quad (3.11)$$

where $W = (W_t)_{t \geq 0}$ denotes a standard Brownian motion.

Remark 3.2. The stochastic integrals in (3.8) can be rewritten more directly as follows: denoting by $W = (W_t)_{t \geq 0}$ a standard Brownian motion, we have

$$Z_{t, \hat{\beta}, \hat{h}}^W := 1 + \sum_{k=1}^{\infty} \int_{0 < t_1 < \dots < t_k < t} \dots \int \psi_t(t_1, \dots, t_k) \prod_{i=1}^k (\hat{\beta} dW_{t_i} + \hat{h} dt_i), \quad (3.12)$$

where the integrals can be viewed as ordinary Ito integrals: it suffices to first integrate over $(\hat{\beta} dW_{t_1} + \hat{h} dt_1)$ for $t_1 \in (0, t_2)$, then over $(\hat{\beta} dW_{t_2} + \hat{h} dt_2)$ for $t_2 \in (0, t_3)$, etc.

Remark 3.3. Theorem 3.1 extends readily to the convergence of the joint distribution of a finite collection of partition functions (conditioned or free). Analogously, the two parameter family of partition functions

$$Z_{\beta_N, h_N}^{\omega, c}(Ns, Nt) := \mathbb{E}\left[e^{\sum_{n=Ns+1}^{Nt-1} (\beta \omega_n - \Lambda(\beta) + h)} \mathbb{1}_{\{n \in \tau\}} \mid Ns, Nt \in \tau \right] \quad \text{for } 0 < s < t, \quad (3.13)$$

converges in finite-dimensional distributions to a 2-parameter process $(Z_{\hat{\beta}, \hat{h}}^{W, c}(s, t))_{0 < s < t}$. In [CSZ14] we upgrade this result to a convergence in distribution in the space of continuous functions, equipped with the uniform topology. This allows us to construct the continuum limit of the disordered pinning measure $P_{N, \beta, h}^\omega$.

Remark 3.4. It is natural to call the random variable $Z_{t,\hat{\beta},\hat{h}}^W$ in Theorem 3.1 the *continuum partition function* and to define the corresponding *continuum free energy*

$$\mathbf{F}^{(\alpha)}(\hat{\beta}, \hat{h}) = \lim_{t \rightarrow \infty} \frac{1}{t} \mathbb{E}[\log Z_{t,\hat{\beta},\hat{h}}^W], \quad (3.14)$$

where \mathbb{E} denotes expectation with respect to the white noise W , provided the limit exists. For $1/2 < \alpha < 1$, we expect that the continuum and discrete free energies are related via

$$\mathbf{F}^{(\alpha)}(\hat{\beta}, \hat{h}) = \lim_{\delta \downarrow 0} \frac{F(\delta^{\alpha-1/2} L(1/\delta) \hat{\beta}, \delta^\alpha L(1/\delta) \hat{h})}{\delta}. \quad (3.15)$$

This would follow if one could interchange the limits in the formal computation

$$\mathbf{F}^{(\alpha)}(\hat{\beta}, \hat{h}) = \lim_{t \rightarrow \infty} \frac{1}{t} \log Z_{t,\hat{\beta},\hat{h}}^W \stackrel{d}{=} \lim_{t \rightarrow \infty} \frac{1}{t} \lim_{N \rightarrow \infty} \log Z_{tN,\beta_N,h_N}^\omega, \quad (3.16)$$

where β_N, h_N scale as in (3.7). Such an interchange of limits has been made possible in the related copolymer model, cf. [BdH97, CG10]. Proving (3.15) is a very interesting open problem. Even the existence of the continuum free energy in (3.14)—possibly also in the \mathbb{P} -a.s. sense, as for the discrete case (3.5)—is a non-trivial issue.

Relation (3.15) is appealing because of its implication of universality: it states that the discrete free energy $F(\beta, h)$ has a *universal shape* in the weak disorder regime $\beta, h \rightarrow 0$, given by the continuum free energy, which depends only on the parameter α and not on finer details of the renewal distribution. Inverting the relation $\beta = \delta^{\alpha-1/2} L(1/\delta)$, one can rewrite (3.15) for $\hat{\beta} = 1$ as

$$\mathbf{F}^{(\alpha)}(1, \hat{h}) = \lim_{\beta \downarrow 0} \frac{F(\beta, \tilde{L}(1/\beta) \beta^{2\alpha/(2\alpha-1)} \hat{h})}{\tilde{L}(1/\beta) \beta^{2/(2\alpha-1)}}, \quad (3.17)$$

where $\tilde{L}(\cdot)$ and $\hat{L}(\cdot)$ are suitable slowly varying functions determined by $L(\cdot)$.

Indeed, given a slowly varying function ϕ and $\gamma > 0$, we define the slowly varying functions $\bar{\phi}_\gamma$ and ϕ^* by

$$\bar{\phi}_\gamma(x) := \frac{1}{\phi(x^{1/\gamma})}, \quad \phi^*(x\phi(x)) \sim \frac{1}{\phi(x)} \quad \text{as } x \rightarrow \infty,$$

where the existence of ϕ^* is guaranteed by [BGT87, Theorem 1.5.13]. Then as $\beta \downarrow 0$,

$$\delta^{\alpha-1/2} L(1/\delta) = \beta \Rightarrow \frac{1}{\delta^{\alpha-1/2}} \bar{L}_{\alpha-1/2} \left(\frac{1}{\delta^{\alpha-1/2}} \right) \sim \frac{1}{\beta} \Rightarrow L \left(\frac{1}{\delta} \right) \sim (\bar{L}_{\alpha-1/2})^* \left(\frac{1}{\beta} \right),$$

hence from $\delta^{\alpha-1/2} L(1/\delta) = \beta$ we deduce that $\delta \sim \beta^{2/(2\alpha-1)} \hat{L}(1/\beta)$ and $\delta^\alpha L(1/\delta) \sim \beta^{2\alpha/(2\alpha-1)} \tilde{L}(1/\beta)$, where

$$\hat{L}(x) := [(\bar{L}_{\alpha-1/2})^*(x)]^{-2/(2\alpha-1)}, \quad \tilde{L}(x) := [(\bar{L}_{\alpha-1/2})^*(x)]^{-1/(2\alpha-1)}. \quad (3.18)$$

Plugging this into (3.15) (with $\hat{\beta} = 1$), we get (3.17).

Define the critical curve of the continuum free energy in analogy with (3.6), i.e.

$$\mathbf{h}_c^{(\alpha)}(\hat{\beta}) := \sup\{\hat{h} \in \mathbb{R} : \mathbf{F}^{(\alpha)}(\hat{\beta}, \hat{h}) = 0\} = \inf\{\hat{h} \in \mathbb{R} : \mathbf{F}^{(\alpha)}(\hat{\beta}, \hat{h}) > 0\}.$$

Relation (3.17) leads us to the following

Conjecture 3.5. For any disordered pinning model satisfying (3.2) with $\alpha \in (1/2, 1)$, the critical curve $h_c(\beta)$ has the following universal asymptotic behavior (with $L(\cdot)$ defined by (3.18)):

$$\lim_{\beta \downarrow 0} \frac{h_c(\beta)}{\tilde{L}(1/\beta)\beta^{2\alpha/(2\alpha-1)}} = h_c^{(\alpha)}(1).$$

Further support to this conjecture is provided by the fact that (non-matching) upper and lower bounds for $h_c(\beta)$ of the order $\tilde{L}(1/\beta)\beta^{2\alpha/(2\alpha-1)}$ were proved in [A08, AZ09].

Remark 3.6. The case when (3.2) holds with $\alpha = 1$ and $E[\tau_1] = \infty$, i.e. $\sum_{n \in \mathbb{N}} L(n)/n = \infty$, can also be included in Theorem 3.1 (we have omitted it for notational lightness), by setting

$$\beta_N = \hat{\beta} \frac{\ell(N)}{\sqrt{N}}, \quad h_N = \hat{h} \frac{\ell(N)}{N}, \quad \text{where} \quad \ell(N) := \sum_{n=1}^N \frac{L(n)}{n}$$

is a slowly varying function, and with $\psi_t^c(t_1, \dots, t_k) = \psi(t_1, \dots, t_k) \equiv 1$. This is easily checked from the proof in Section 6, because $P(n \in \tau) \sim 1/\ell(n)$ [BGT87, Theorem 8.7.5].

On the other hand, the case $\alpha = 1/2$ appears to be fundamentally different, because the continuum kernels ψ_t^c, ψ are no longer L^2 integrable, and therefore the stochastic integrals are not properly defined. When $\alpha = 1/2$ and $\sum_{n \in \mathbb{N}} 1/(nL(n)^2) = \infty$ —in particular, when $L(n) \sim \text{const}$ as $n \rightarrow \infty$, as for the simple random walk on \mathbb{Z} —we expect that a non-trivial continuum limit should exist. This appears to be a challenging open problem.

3.2. Directed polymer model

Consider a random walk $S = (S_n)_{n \in \mathbb{N}_0}$ on \mathbb{Z} with law P . Let $\omega = (\omega(n, x))_{n \in \mathbb{N}, x \in \mathbb{Z}}$ be a family of i.i.d. random variables, independent of S , with zero mean, unit variance and locally finite exponential moments (cf. (3.1)). The $(1 + 1)$ -dimensional *directed polymer model* is the random probability law $P_{N, \beta}^\omega$ for the walk S defined for $N \in \mathbb{N}$ and $\beta \geq 0$ by

$$dP_{N, \beta}^\omega(S) := \frac{1}{Z_{N, \beta}^\omega} e^{\sum_{n=1}^N (\beta \omega(n, S_n) - \Lambda(\beta))} dP(S), \quad (3.19)$$

where we recall that $\Lambda(\beta) := \log E[e^{\beta \omega_{n, x}}]$ and the *partition function* $Z_{N, \beta}^\omega$ is defined by

$$Z_{N, \beta}^\omega = E[e^{\beta \sum_{n=1}^N \omega(n, S_n)}] e^{-\Lambda(\beta)N}. \quad (3.20)$$

For $y \in \mathbb{Z}$, we also define the *constrained* point-to-point partition function $Z_{N, \beta}^\omega(y)$ and the *conditioned* point-to-point partition function $Z_{N, \beta}^{\omega, c}(y)$, setting

$$\begin{aligned} Z_{N, \beta}^\omega(y) &= E[e^{\beta \sum_{n=1}^N \omega(n, S_n)} \mathbb{1}_{\{S_N = y\}}] e^{-\Lambda(\beta)N}, \\ Z_{N, \beta}^{\omega, c}(y) &= E[e^{\beta \sum_{n=1}^N \omega(n, S_n)} \mid S_N = y] e^{-\Lambda(\beta)N}. \end{aligned} \quad (3.21)$$

Plainly, $Z_{N, \beta}^\omega = \sum_{y \in \mathbb{Z}} Z_{N, \beta}^\omega(y) = \sum_{y \in \mathbb{Z}} Z_{N, \beta}^{\omega, c}(y) P(S_N = y)$.

When $(S_n)_{n \geq 1}$ is the simple symmetric random walk on \mathbb{Z} , we have the much-studied *directed polymer in random medium*. First introduced in the physics literature in [HH85], this model has received particular attention due to its connection to the Kardar–Parisi–Zhang (KPZ) equation and its universality class (see [CSY04], [C12] for a review). In particular, the point-to-point partition function can be thought of as an approximation of the solution of the stochastic heat equation (SHE), whose logarithm is the so-called Hopf–Cole solution of the KPZ equation. This was made rigorous in [AKQ14a], by showing that when $\beta = \beta_N$ is scaled as $N^{-1/4}$ (the so-called intermediate disorder regime) and y is scaled as $N^{-1/2}$, the point-to-point partition function $Z_{Nt,\beta}^\omega(y)$ converges in distribution to a continuum process, which solves the SHE.

Our approach allows us to extend the results in [AKQ14a]. Not only can we deal with general zero mean, finite variance random walks, which are the natural “universality class” of the simple symmetric random walk on \mathbb{Z} ; we can also consider random walks attracted to stable laws with index $\alpha \in (1, 2)$, exploring new universality classes. When allowing “big jumps”, it is natural to call $P_{N,\beta}^\omega$ the *long-range* directed polymer model.

Let us now state precisely our assumptions on the random walk.

Assumption 3.7. Let $S = (S_n)_{n \geq 0}$ be a random walk on \mathbb{Z} , with $S_0 = 0$ and with i.i.d. increments $(S_n - S_{n-1})_{n \geq 1}$, such that for some $\alpha \in (1, 2]$ the following holds:

- (Case $\alpha = 2$): $E[S_1] = 0$ and $\sigma^2 := \text{Var}(S_1) < \infty$;
- (Case $1 < \alpha < 2$): $E[S_1] = 0$ and there exist $\gamma \in [-1, 1]$ and $C \in (0, \infty)$ such that

$$P(S_1 > n) \sim \left(C \frac{1+\gamma}{2}\right) \frac{1}{n^\alpha}, \quad P(S_1 < -n) \sim \left(C \frac{1-\gamma}{2}\right) \frac{1}{n^\alpha}, \quad \text{as } n \rightarrow \infty. \quad (3.22)$$

This means that the random walk S is in the *domain of normal attraction* of a stable law of index $\alpha \in (1, 2]$ and (for $\alpha < 2$) skewness parameter γ . (The adjective “normal” refers to the absence of slowly varying functions.) In other words, $S_n/n^{1/\alpha}$ converges in distribution as $n \rightarrow \infty$ to a random variable Y , which has law $\mathcal{N}(0, \sigma^2)$ if $\alpha = 2$, while for $1 < \alpha < 2$,

$$E[e^{itY}] = e^{-c_\alpha C |t|^\alpha (1 - i\gamma (\text{sign } t) \tan \frac{\pi\alpha}{2})} \quad \text{for a suitable } c_\alpha > 0. \quad (3.23)$$

We remark that Y satisfies the same conditions as S_1 in Assumption 3.7 and has an absolutely continuous law, with a bounded and continuous density $g(\cdot)$. For $t > 0$ we set

$$g_t(x) := \frac{1}{t^{1/\alpha}} g\left(\frac{x}{t^{1/\alpha}}\right). \quad (3.24)$$

We stress that $g(\cdot)$ depends only on the parameters (α, σ^2) or (α, γ, C) in Assumption 3.7.

The *period* of a random walk S on \mathbb{Z} is the largest $p \in \mathbb{N}$ such that $P(S_1 \in p\mathbb{Z} + r) = 1$ for some $r \in \{0, \dots, p-1\}$. For instance, the simple symmetric random walk on \mathbb{Z} has period $p = 2$, because $P(S_1 \in 2\mathbb{Z} + 1) = 1$. To lighten notation, given $(s, y) \in \mathbb{R}^+ \times \mathbb{R}$, we write $Z_{s,\beta}^{\omega,c}(y)$ to mean $Z_{\tilde{s},\beta}^{\omega,c}(\tilde{y})$, where $\tilde{s} := \lfloor s \rfloor$ and $\tilde{y} := \max\{z \in p\mathbb{Z} + r\tilde{s} : z \leq y\}$.

We are ready to state our main result for this model, to be proved in Section 7.

Theorem 3.8 (Scaling limits of directed polymer models). *Let S be a random walk on \mathbb{Z} satisfying Assumption 3.7 for some $\alpha \in (1, 2]$. For $N \in \mathbb{N}$ and $\hat{\beta} > 0$, set*

$$\beta_N := \hat{\beta}/N^{(\alpha-1)/(2\alpha)}.$$

For every $t \geq 0$ and $x \in \mathbb{R}$, the conditioned point-to-point partition function $Z_{Nt, \beta_N}^{\omega, c}(N^{1/\alpha}x)$ converges in distribution as $N \rightarrow \infty$ to the random variable $Z_{t, \hat{\beta}}^{W, c}(x)$ given by

$$Z_{t, \hat{\beta}}^{W, c}(x) := 1 + \sum_{k=1}^{\infty} \frac{\hat{\beta}^k}{k!} \int \cdots \int_{(\{0, t\} \times \mathbb{R})^k} \psi_{t, x}^c((t_1, x_1), \dots, (t_k, x_k)) \prod_{i=1}^k W(dt_i dx_i), \quad (3.25)$$

where $W(\cdot)$ denotes a space-time white noise (i.e., white noise on \mathbb{R}^2) and the symmetric function $\psi_{t, x}^c((t_1, x_1), \dots, (t_k, x_k))$ is defined for $0 < t_1 < \dots < t_k < t$ by

$$\psi_{t, x}^c((t_1, x_1), \dots, (t_k, x_k)) := \left(\prod_{i=1}^k \sqrt{p} g_{t_i - t_{i-1}}(x_i - x_{i-1}) \right) \frac{g_{t-t_k}(x - x_k)}{g_t(x)}, \quad (3.26)$$

where $x_0 := 0$ and $p \in \mathbb{N}$ is the period of the random walk. The series in (3.25) converges in L^2 and furthermore $\mathbb{E}[(Z_{Nt, \beta_N}^{\omega, c}(N^{1/\alpha}x))^2] \rightarrow \mathbb{E}[(Z_{t, \hat{\beta}}^{W, c}(x))^2]$ as $N \rightarrow \infty$.

An analogous statement holds for the free partition function Z_{Nt, β_N}^{ω} , where the limiting random variable $Z_{t, \hat{\beta}}^W$ is defined as in (3.25), with kernel

$$\psi_t((t_1, x_1), \dots, (t_k, x_k)) := \prod_{i=1}^k \sqrt{p} g_{t_i - t_{i-1}}(x_i - x_{i-1}).$$

Remark 3.9. Theorem 3.8 extends to the convergence of the joint distribution of a finite collection of partition functions (conditioned or free). In particular, the four-parameter family $Z_{Ns, Nt, \beta_N}^{\omega, c}(N^{1/\alpha}x, N^{1/\alpha}y)$, defined for (s, x) and (t, y) in $\mathbb{R}^+ \times \mathbb{R}$ by

$$Z_{Ns, Nt, \beta_N}^{\omega, c}(N^{1/\alpha}x, N^{1/\alpha}y) := \mathbb{E}\left[e^{\sum_{n=N_s+1}^{N_t-1} (\beta\omega(n, S_n) - \Lambda(\beta))} \mid S_{N_s} = N^{1/\alpha}x, S_{N_t} = N^{1/\alpha}y \right],$$

converges in finite-dimensional distributions to a four-parameter family of continuum conditioned partition functions $\{Z_{s, t, \hat{\beta}}^{W, c}(x, y)\}_{(s, x), (t, y) \in \mathbb{R}^+ \times \mathbb{R}}$.

Similar to the case $\alpha = 2$ studied in [AKQ14b], we expect that this convergence can be upgraded to a convergence in distribution in the space of continuous functions, equipped with the uniform topology. We can then use these partition functions to define a *continuum long-range directed polymer model* (which corresponds intuitively to an “ α -stable Lévy process in a white noise random medium”), by specifying its finite-dimensional distributions as done in [AKQ14b] for the Brownian case $\alpha = 2$.

Remark 3.10. The free energy of the (discrete) directed polymer model is defined by

$$F(\beta) := \lim_{N \rightarrow \infty} \frac{1}{N} \mathbb{E}[\log Z_{N,\beta}^\omega],$$

where we expect that the limit exists (also \mathbb{P} -a.s. and in $L^1(d\mathbb{P})$), as in the usual setting. It is natural to define the free energy of the continuum model analogously, i.e.

$$F(\hat{\beta}) := \lim_{t \rightarrow \infty} \frac{1}{t} \mathbb{E}[\log \mathbf{Z}_{t,\hat{\beta}}^W],$$

assuming of course that the limit exists. We stress that $F(\cdot)$ is a *universal* quantity, which depends only on the parameters (α, σ^2) or (α, γ, C) in Assumption 3.7 (furthermore, the parameters σ^2, C enter as simple scale factors). We also note that $F(\hat{\beta}) = F(1)\hat{\beta}^{(\alpha-1)/(2\alpha)}$, by an easy scaling argument. In analogy with Remark 3.4, we are led to the following

Conjecture 3.11. For any directed polymer model satisfying Assumption 3.7, the free energy $F(\beta)$ has the following universal asymptotic behavior for weak disorder:

$$\lim_{\beta \downarrow 0} \frac{F(\beta)}{\beta^{2\alpha/(\alpha-1)}} = \lim_{\delta \downarrow 0} \frac{F(\delta^{(\alpha-1)/(2\alpha)})}{\delta} = F(1).$$

When $\alpha = 2$ we would then get $F(\beta) \sim F(1)\beta^4$, which is supported by the (non-matching) upper and lower bounds on $F(\beta)$ obtained in [L10].

Remark 3.12. For $1 < \alpha < 2$, the function $g_t(\cdot)$ in (3.24) is the marginal density of the α -stable Lévy process $(X_t)_{t \in \mathbb{R}^+}$ whose infinitesimal generator is given by a multiple of

$$\Delta^{\alpha/2, \gamma} f(x) := \int_{-\infty}^{\infty} (f(x+y) - f(x) - yf'(x)) \left(\frac{1+\gamma}{|y|^{1+\alpha}} \mathbb{1}_{\{y>0\}} + \frac{1-\gamma}{|y|^{1+\alpha}} \mathbb{1}_{\{y<0\}} \right) dy. \quad (3.27)$$

In the symmetric case $\gamma = 0$, this reduces to the much studied *fractional Laplacian*

$$\Delta^{\alpha/2} f(x) := \int_{-\infty}^{\infty} (f(x+y) - f(x) - yf'(x)) \frac{1}{|y|^{1+\alpha}} dy. \quad (3.28)$$

Let us stick for simplicity to the symmetric case $\gamma = 0$. It is natural to call $\mathbf{Z}_{t,\hat{\beta}}^{W,c}(x)$ in (3.25) the continuum *conditioned* point-to-point partition function. Introducing the continuum *constrained* point-to-point partition function by

$$\mathbf{Z}_{t,\hat{\beta}}^W(x) := \mathbf{Z}_{t,\hat{\beta}}^{W,c}(x) g_t(x),$$

one can check that the process $u(t, x) = \mathbf{Z}_{t,\hat{\beta}}^W(x)$ has a version that is continuous in (t, x) and, up to a scaling factor, it is a mild solution to the one-dimensional stochastic PDE

$$\begin{cases} \partial_t u = \Delta^{\alpha/2} u + \sqrt{p} \hat{\beta} \dot{W} u, \\ u(0, \cdot) = \delta_0(\cdot), \end{cases} \quad (3.29)$$

which we can call the *stochastic fractional heat equation* (SFHE), generalizing the usual SHE (which corresponds to $\alpha = 2$). Uniqueness of mild solutions for the SFHE follows from standard techniques (see discussions in [CJKS14] and references therein).

Let us also consider the process $A_{\alpha, \hat{\beta}}(\cdot) := \log \mathbf{Z}_{1, \hat{\beta}}^W(\cdot)$. When $\alpha = 2$, this is the *cross-over process* studied in [ACQ11, SS10], which owes its name to the fact that its one-point marginals interpolate between the Gaussian distribution (in the limit $\hat{\beta} \rightarrow 0$) and the Tracy–Widom GUE distribution (in the limit $\hat{\beta} \rightarrow \infty$). When $\alpha < 2$, it is easy to see that $A_{\alpha, \hat{\beta}}(\cdot)$ is again asymptotically Gaussian for $\hat{\beta} \rightarrow 0$ (the contribution from the first stochastic integral in its Wiener chaos expansion is dominant over the iterated integrals, which are multiplied by higher powers of $\hat{\beta}$). However, it is far from obvious whether $A_{\alpha, \hat{\beta}}(\cdot)$ converges to some asymptotic process $A_{\alpha, \infty}(\cdot)$ as $\hat{\beta} \rightarrow \infty$ and whether such a process describes some universality class for long-range random polymers, last passage percolation and growth models, generalizing the so-called Airy process obtained for $\alpha = 2$. Besides a very recent work on the limit shapes of long-range first-passage percolation model [CD13], long-range polymer type models do not appear to have been studied systematically before.

3.3. Random field Ising model

Given a bounded $\Omega \subseteq \mathbb{Z}^2$, we set

$$\partial\Omega := \{x \in \mathbb{Z}^2 \setminus \Omega : \|x - y\| = 1 \text{ for some } y \in \Omega\}.$$

For a fixed parameter $\beta \geq 0$, representing the “inverse temperature”, the *Ising model on Ω with + boundary condition (and zero external field)* is the probability measure \mathbb{P}_Ω^+ on the set of spin configurations $\{\pm 1\}^\Omega$, where each $\sigma := (\sigma_x)_{x \in \Omega} \in \{\pm 1\}^\Omega$ has probability

$$\mathbb{P}_\Omega^+(\sigma) := \frac{1}{Z_\Omega^+} \exp\left\{\beta \sum_{x \sim y \in \Omega \cup \partial\Omega} \sigma_x \sigma_y\right\} \prod_{x \in \partial\Omega} \mathbb{1}_{\{\sigma_x = +1\}}. \quad (3.30)$$

Here $x \sim y$ denotes an unordered nearest-neighbor pair in \mathbb{Z}^2 , and Z_Ω^+ is the normalizing constant. The value of β will soon be fixed, which is why we do not indicate it in \mathbb{P}_Ω^+ .

Remark 3.13. It is well-known that as $\Omega \uparrow \mathbb{Z}^2$, the sequence of probability measures \mathbb{P}_Ω^+ has a unique infinite volume limit $\mathbb{P}_{\mathbb{Z}^2}^+$, which of course depends on β , such that

$$E_{\mathbb{Z}^2}^+(\sigma_0) > 0 \quad \text{if and only if} \quad \beta > \beta_c := \frac{1}{2} \log(1 + \sqrt{2}).$$

By an obvious symmetry, for $\beta > \beta_c$ there also exists an infinite volume Gibbs measure $\mathbb{P}_{\mathbb{Z}^2}^-$ satisfying $E_{\mathbb{Z}^2}^-(\sigma_0) < 0$. Given the coexistence of multiple infinite volume Gibbs measures, the Ising model is said to have a first order phase transition for $\beta > \beta_c$. The same result holds in higher dimensions, with different values of β_c .

A question that attracted significant interest was whether this picture will be altered by the addition of a small random external field. After long debates, this question was settled by Bricmont and Kupiainen [BK88], who showed that the first order phase transition

persists for the random field Ising model in dimensions $d \geq 3$ at low temperatures (i.e., for large β), and by Aizenman and Wehr [AW90] who showed the absence of first order phase transition in dimension 2 at any temperature. See [B06, Chap. 7] for an overview.

Henceforth we fix $\beta = \beta_c := \frac{1}{2} \log(1 + \sqrt{2})$, so that P_Ω^+ denotes the two-dimensional critical Ising model. Let $\omega := (\omega_x)_{x \in \mathbb{Z}^2}$ be a family of i.i.d. random variables satisfying (3.1), representing the disorder. Given $\lambda := (\lambda_x)_{x \in \mathbb{Z}^2} \geq 0$ and $h := (h_x)_{x \in \mathbb{Z}^2} \in \mathbb{R}$, representing the disorder strength and bias respectively, the random field Ising model (RFIM) is the probability measure $P_{\Omega, \lambda, h}^{+, \omega}$ on $\{\pm 1\}^\Omega$ with

$$P_{\Omega, \lambda, h}^{+, \omega}(\sigma) = \frac{1}{Z_{\Omega, \lambda, h}^{+, \omega}} \exp\left\{ \sum_{x \in \Omega} (\lambda_x \omega_x + h_x) \sigma_x \right\} P_\Omega^+(\sigma), \quad (3.31)$$

where the normalizing constant, called the *partition function*, is given by

$$Z_{\Omega, \lambda, h}^{+, \omega} = E_\Omega^+ \left[\exp\left\{ \sum_{x \in \Omega} (\lambda_x \omega_x + h_x) \sigma_x \right\} \right]. \quad (3.32)$$

Note that we allow the disorder strength λ and bias h to vary from site to site. Also observe that choosing P_Ω^+ as a ‘‘reference law’’ means that $Z_{\Omega, \lambda, h}^{+, \omega} = 1$ for $\lambda, h \equiv 0$ (with $\beta = \beta_c$).

Fix now a bounded open set $\Omega \subseteq \mathbb{R}^2$ with piecewise smooth boundary, and define the rescaled lattice $\Omega_\delta := \Omega \cap (\delta\mathbb{Z})^2$, for $\delta > 0$. We are going to obtain a non-trivial limit in distribution for the partition function $Z_{\Omega_\delta, \lambda, h}^{+, \omega}$, in the continuum and weak disorder regime $\delta, \lambda, h \rightarrow 0$. We build on recent results of Chelkak, Hongler and Izyurov [CHI12, Theorem 1.3] on the continuum limit of the spin correlations under $P_{\Omega_\delta}^+$ (the two-dimensional critical Ising model with + boundary condition): for all $n \in \mathbb{N}$ and distinct $x_1, \dots, x_n \in \Omega$,

$$\lim_{\delta \downarrow 0} \delta^{-n/8} E_{\Omega_\delta}^+ [\sigma_{x_1} \cdots \sigma_{x_n}] = \mathcal{C}^n \phi_\Omega^+(x_1, \dots, x_n), \quad (3.33)$$

where $\phi_\Omega^+ : \bigcup_{n \in \mathbb{N}} \Omega^n \rightarrow \mathbb{R}$ is a symmetric function and $\mathcal{C} := 2^{5/48} e^{-\frac{3}{2} \zeta'(-1)}$, with ζ' denoting the derivative of Riemann’s zeta function.⁴ (For simplicity, in (3.33) we have set $\sigma_x := \sigma_{x_\delta}$, where x_δ denotes the point in Ω_δ closest to $x \in \Omega$.)

We can complement (3.33) with a uniform bound (see Lemma 8.1 below): there exists $C \in (0, \infty)$ such that for all $n \in \mathbb{N}$, $x_1, \dots, x_n \in \Omega$ and $\delta \in (0, 1)$,

$$0 \leq \delta^{-n/8} E_{\Omega_\delta}^+ [\sigma_{x_1} \cdots \sigma_{x_n}] \leq \prod_{i=1}^n \frac{C}{d(x_i, \partial\Omega \cup \{x_1, \dots, x_n\} \setminus \{x_i\})^{1/8}}, \quad (3.34)$$

where $d(x, A) := \inf_{y \in A} \|x - y\|$ and the RHS is shown to be in L^2 in Lemma 8.3 below. Therefore the convergence (3.33) also holds in $L^2(\Omega)$, and $\|\phi_\Omega^+\|_{L^2(\Omega^n)} < \infty$ for every $n \in \mathbb{N}$.

We can now state our main result for the RFIM, to be proved in Section 8.

⁴ In [CHI12], the mesh size of Ω_δ is $\sqrt{2} \delta$ instead of δ , which accounts for the difference between our formula for \mathcal{C} and that in [CHI12, (1.3)]. Exact formulas for $\phi_\Omega^+(x_1, \dots, x_n)$ are available for $n = 1, 2$ when Ω is the upper half-plane \mathbb{H} [CHI12, (1.4)]. The general case ϕ_Ω^+ can be obtained from $\phi_{\mathbb{H}}^+$ via a conformal map.

Theorem 3.14 (Scaling limits of RFIM). *Let $\Omega \subseteq \mathbb{R}^2$ be bounded and simply connected open set with piecewise smooth boundary, and define $\Omega_\delta := \Omega \cap (\delta\mathbb{Z})^2$ for $\delta > 0$.*

Let $\omega := (\omega_x)_{x \in \mathbb{Z}^2}$ be i.i.d. random variables satisfying (3.1) and define $\omega_\delta = (\omega_{\delta,x})_{x \in \Omega_\delta}$ by $\omega_{\delta,x} := \omega_{x/\delta}$. Given continuous functions $\hat{\lambda} : \bar{\Omega} \rightarrow (0, \infty)$ and $\hat{h} : \bar{\Omega} \rightarrow \mathbb{R}$, define

$$\lambda_{\delta,x} := \hat{\lambda}(x)\delta^{7/8}, \quad h_{\delta,x} := \hat{h}(x)\delta^{15/8}, \quad (3.35)$$

and set $\lambda_\delta = (\lambda_{\delta,x})_{x \in \Omega_\delta}$, $h_\delta = (h_{\delta,x})_{x \in \Omega_\delta}$. Then the rescaled partition function

$$e^{-\frac{1}{2}\|\hat{\lambda}\|_{L^2(\Omega)}^2} \delta^{-1/4} \mathbf{Z}_{\Omega_\delta, \lambda_\delta, h_\delta}^{+, \omega_\delta} \quad (3.36)$$

converges in distribution as $\delta \downarrow 0$ to a random variable $\mathbf{Z}_{\Omega, \hat{\lambda}, \hat{h}}^{+, W}$ with Wiener chaos expansion

$$\mathbf{Z}_{\Omega, \hat{\lambda}, \hat{h}}^{+, W} = 1 + \sum_{n=1}^{\infty} \frac{C^n}{n!} \int \cdots \int_{\Omega^n} \phi_{\Omega}^+(x_1, \dots, x_n) \prod_{i=1}^n [\hat{\lambda}(x_i) W(dx_i) + \hat{h}(x_i) dx_i], \quad (3.37)$$

where $W(\cdot)$ denotes a white noise on \mathbb{R}^2 and $\phi_{\Omega}^+(\cdot)$, C are as in (3.33).

Remark 3.15. We require the continuity of $\hat{\lambda}$ and \hat{h} and the strict positivity of $\hat{\lambda}$ for technical simplicity: these conditions can be relaxed with a more careful analysis.

We call the random variable $\mathbf{Z}_{\Omega, \hat{\lambda}, \hat{h}}^{+, W}$ in (3.37) the *continuum RFIM partition function*. The fact that the continuum correlation function ϕ_{Ω}^+ is *conformally covariant*, proved in [CHI12, Theorem 1.3], allows one to deduce the conformal covariance of $\mathbf{Z}_{\Omega, \hat{\lambda}, \hat{h}}^{+, W}$.

Corollary 3.16 (Conformal covariance). *Let $\Omega, \tilde{\Omega} \subseteq \mathbb{R}^2$ be bounded and simply connected open sets with piecewise smooth boundaries, and let $\varphi : \tilde{\Omega} \rightarrow \Omega$ be conformal. Let $\hat{\lambda} : \bar{\Omega} \rightarrow (0, \infty)$ and $\hat{h} : \bar{\Omega} \rightarrow \mathbb{R}$ be continuous functions. Then*

$$\mathbf{Z}_{\Omega, \hat{\lambda}, \hat{h}}^{+, W} \stackrel{d}{=} \mathbf{Z}_{\tilde{\Omega}, \tilde{\lambda}, \tilde{h}}^{+, W},$$

where we set $\tilde{\lambda}(z) := |\varphi'(z)|^{7/8} \hat{\lambda}(\varphi(z))$ and $\tilde{h}(z) := |\varphi'(z)|^{15/8} \hat{h}(\varphi(z))$ for all $z \in \tilde{\Omega}$.

Remark 3.17. Recently, Camia, Garban and Newman [CGN12a, CGN13] showed that for a *deterministic* external field, more precisely $\lambda_{\delta,x} \equiv 0$ and $h_{\delta,x} \equiv \hat{h}\delta^{15/8}$ for fixed $\hat{h} \in \mathbb{R}$, the Ising measure $\mathbb{P}_{\Omega_\delta, \lambda_\delta, h_\delta}^+$ converges as $\delta \downarrow 0$ (in a suitable sense) to a limiting distribution-valued process $\Phi_{\Omega}^{\infty, \hat{h}}$. Theorem 3.14 can be regarded as a first step towards the extension of this convergence to the case of a *random* external field, where $\mathbf{Z}_{\Omega, \hat{\lambda}, \hat{h}}^{+, W}$ plays the role of the partition function of a *continuum Ising model with random external field* ($\hat{\lambda}(x)W(dx) + \hat{h}(x)dx$) and *+ boundary condition*. The next step towards the construction of such a continuum model would be to identify the joint distribution of the partition functions $(\mathbf{Z}_{\Gamma, \hat{\lambda}, \hat{h}}^{b, W})_{\Gamma, b}$ (note that they are random variables which are functions of the disorder W) for a large enough family of subdomains $\Gamma \subseteq \Omega$ and “boundary conditions” b .

Remark 3.18. Consider the case when the disorder strength and bias are constant, i.e. $\lambda_x \equiv \lambda \geq 0$ and $h_x \equiv h \in \mathbb{R}$. The free energy $F(\lambda, h)$ of the critical random field Ising model can be defined as follows: setting $\Omega := (-1/2, 1/2)^2$ and $\Lambda_N := (N\Omega) \cap \mathbb{Z}^2 = \{[-N/2], \dots, [N/2]\}^2$, we define

$$F(\lambda, h) := \lim_{N \rightarrow \infty} \frac{1}{N^2 |\Omega|} \mathbb{E}[\log Z_{\Lambda_N, \lambda, h}^{+, \omega}] = \lim_{\delta \downarrow 0} \frac{\delta^2}{|\Omega|} \mathbb{E}[\log Z_{\Omega_\delta, \lambda, h}^{+, \omega}], \quad (3.38)$$

where the limit exists by standard superadditivity arguments and is independent of the choice of Ω . Note that $F(0, 0) = 0$, that is $F(\lambda, h)$ represents the *excess* free energy with respect to the critical Ising model (cf. (3.32)).

It is natural to define a continuum free energy $\mathbf{F}(\hat{\lambda}, \hat{h})$ for $\hat{\lambda} \geq 0, \hat{h} \in \mathbb{R}$ by

$$\mathbf{F}(\hat{\lambda}, \hat{h}) := \lim_{\Omega \uparrow \mathbb{R}^2} \frac{1}{\text{Leb}(\Omega)} \mathbb{E}[\log Z_{\Omega, \hat{\lambda}, \hat{h}}^{+, W}],$$

provided the limit exists (at least along sufficiently nice domains $\Omega \uparrow \mathbb{R}^2$), where \mathbb{E} denotes expectation with respect to the white noise $W(\cdot)$. In analogy with Remark 3.4, Theorem 3.14 suggests the following conjecture on the universal behavior of the free energy $F(\lambda, h)$ in the weak disorder regime $\lambda, h \rightarrow 0$.

Conjecture 3.19. The following asymptotic relation holds:

$$\lim_{\delta \downarrow 0} \frac{F(\hat{\lambda} \delta^{7/8}, \hat{h} \delta^{15/8})}{\delta^2} = \mathbf{F}(\hat{\lambda}, \hat{h}), \quad \forall \hat{\lambda} \geq 0, \hat{h} \in \mathbb{R}. \quad (3.39)$$

One can go one step further: differentiating both sides of (3.39) with respect to \hat{h} and setting $\delta = h^{8/15}$, for $\hat{h} = 1$ one would obtain

$$\lim_{h \rightarrow 0} \frac{\langle \sigma_0 \rangle_{\hat{\lambda} h^{7/15}, h}}{h^{1/15}} = \frac{\partial \mathbf{F}}{\partial \hat{h}}(\hat{\lambda}, 1), \quad (3.40)$$

where $\langle \sigma_0 \rangle_{\lambda, h} := \frac{\partial F}{\partial h}(\lambda, h) = \lim_{N \rightarrow \infty} \mathbb{E}[E_{\Lambda_N, \lambda, h}^{+, \omega}[\sigma_0]]$ represents the average magnetization in the infinite-volume random field critical Ising model (cf. (3.38) and (3.31)). Relation (3.40) is supported by (non-matching) upper and lower bounds of the order $h^{1/15}$ for $\langle \sigma_0 \rangle_{\lambda, h}$ in the non-disordered case $\lambda = 0$, recently proved by Camia, Garban and Newman [CGN12b].

In light of Theorem 3.14, and the fact that for the two-dimensional Ising model below the critical temperature a disordered external field smoothens the phase transition [AW90], it is natural to conjecture that a similar smoothing effect occurs at the critical temperature:

Conjecture 3.20. For any fixed $\lambda > 0$, the average magnetization in the infinite-volume random field critical Ising model on \mathbb{Z}^2 satisfies

$$\langle \sigma_0 \rangle_{\lambda, h} \sim Ch^\gamma \quad \text{as } h \downarrow 0 \quad \text{for some } \gamma > \frac{1}{15}. \quad (3.41)$$

4. Proof of the Lindeberg principle

In this section, we prove Theorems 2.6 and 2.8 on the Lindeberg principle for polynomial chaos expansions. We first deduce Theorem 2.8 from Theorem 2.6, starting with the following lemma which controls how a multi-linear polynomial $\Psi(x) = \Psi((x_i)_{i \in \mathbb{T}})$ is affected by a shift $x \mapsto x + \mu$, where $\mu := (\mu_i)_{i \in \mathbb{T}}$.

Lemma 4.1 (Effect of adding a mean). *Let $\Psi(x) = \Psi((x_i)_{i \in \mathbb{T}})$ be a multi-linear polynomial as in (2.5), with kernel ψ , and let $\mu := (\mu_i)_{i \in \mathbb{T}}$ be a family of real numbers. Then $\tilde{\Psi}(x) := \Psi(x + \mu)$ is a multi-linear polynomial, i.e. $\tilde{\Psi}(x) = \sum_{J \in \mathcal{P}^{\text{fin}}(\mathbb{T})} \tilde{\psi}(J) x^J$, with kernel*

$$\tilde{\psi}(J) = \sum_{I \in \mathcal{P}^{\text{fin}}(\mathbb{T}), I \supseteq J} \psi(I) \mu^{I \setminus J}. \quad (4.1)$$

For $\varepsilon > 0$, let $\Psi^{(\varepsilon)}(x)$ be defined as in (2.26), and recall the definitions of C_Ψ from (2.19) and $\text{Inf}_j[\Psi]$ from (2.20). Then, setting $c_\mu := \sum_{i \in \mathbb{T}} \mu_i^2$, for any $\varepsilon > 0$ we have

$$C_{\tilde{\Psi}} \leq e^{c_\mu/\varepsilon} C_{\Psi^{(\varepsilon)}}, \quad \text{Inf}_j[\tilde{\Psi}] \leq e^{c_\mu/\varepsilon} \text{Inf}_j[\Psi^{(\varepsilon)}]. \quad (4.2)$$

Proof. Note that (4.1) follows from the expansion

$$\begin{aligned} \tilde{\Psi}(x) &= \Psi(x + \mu) = \sum_I \psi(I) (x + \mu)^I = \sum_I \psi(I) \sum_{J \subseteq I} \mu^{I \setminus J} x^J \\ &= \sum_J \left(\sum_{I \supseteq J} \psi(I) \mu^{I \setminus J} \right) x^J. \end{aligned}$$

For any $\varepsilon > 0$, we can apply Cauchy–Schwarz to write

$$\begin{aligned} \tilde{\psi}(J)^2 &\leq \left(\sum_{I \supseteq J} (\varepsilon^{-1} \mu^2)^{I \setminus J} \right) \left(\sum_{I \supseteq J} \varepsilon^{|I \setminus J|} \psi(I)^2 \right) \\ &= (1 + \varepsilon^{-1} \mu^2)^{\mathbb{T} \setminus J} \left(\sum_{I \supseteq J} \varepsilon^{|I \setminus J|} \psi(I)^2 \right), \end{aligned} \quad (4.3)$$

and note that

$$(1 + \varepsilon^{-1} \mu^2)^{\mathbb{T} \setminus J} \leq (1 + \varepsilon^{-1} \mu^2)^{\mathbb{T}} \leq e^{c_\mu/\varepsilon}.$$

Therefore

$$\begin{aligned} C_{\tilde{\Psi}} &:= \sum_J \tilde{\psi}(J)^2 \leq e^{c_\mu/\varepsilon} \sum_I \left(\sum_{J \subseteq I} \varepsilon^{|I \setminus J|} \right) \psi(I)^2 \\ &= e^{c_\mu/\varepsilon} \sum_I (1 + \varepsilon)^{|I|} \psi(I)^2 =: e^{c_\mu/\varepsilon} C_{\Psi^{(\varepsilon)}}, \end{aligned}$$

proving the first relation in (4.2). The second relation is obtained similarly:

$$\begin{aligned} \text{Inf}_j[\tilde{\Psi}] &:= \sum_{J \ni j} \tilde{\psi}(J)^2 \leq e^{c_\mu/\varepsilon} \sum_{I \ni j} \left(\sum_{J, j \in J \subseteq I} \varepsilon^{|I \setminus J|} \right) \psi(I)^2 \\ &\leq e^{c_\mu/\varepsilon} \sum_{I \ni j} (1 + \varepsilon)^{|I|} \psi(I)^2 =: e^{c_\mu/\varepsilon} \text{Inf}_j[\Psi^{(\varepsilon)}]. \end{aligned} \quad (4.4)$$

This concludes the proof of the lemma. \square

Proof of Theorem 2.8. Since $\Psi(\tilde{\zeta}) = \Psi(\zeta + \mu) =: \tilde{\Psi}(\zeta)$, by Lemma 4.1 the conditions $c_\mu < \infty$ and $C_{\Psi^{(\varepsilon)}} < \infty$ ensure that $C_{\tilde{\Psi}} < \infty$. The polynomial chaos expansion $\Psi(\tilde{\zeta})$ is then a well-defined L^2 random variable, by Remark 2.2, and the same holds for $\Psi(\tilde{\xi})$.

To prove (2.27), we are first going to truncate Ψ to degree ℓ , i.e. we consider $\Psi^{\leq \ell}$ and $\Psi^{> \ell}$, defined in (2.21). Note that

$$\begin{aligned} |\mathbb{E}[f(\Psi(\tilde{\zeta})) - f(\Psi^{\leq \ell}(\tilde{\zeta}))]| &\leq \|f'\|_\infty \mathbb{E}[|\Psi(\tilde{\zeta}) - \Psi^{\leq \ell}(\tilde{\zeta})|] \\ &\leq \|f'\|_\infty \mathbb{E}[|\Psi^{> \ell}(\tilde{\zeta})|^2]^{1/2}, \end{aligned} \quad (4.5)$$

and the same bound holds when $\tilde{\zeta}$ is replaced by $\tilde{\xi}$, therefore

$$\begin{aligned} |\mathbb{E}[f(\Psi(\tilde{\zeta}))] - \mathbb{E}[f(\Psi(\tilde{\xi}))]| &\leq 2\|f'\|_\infty \mathbb{E}[|\Psi^{> \ell}(\tilde{\zeta})|^2]^{1/2} \\ &\quad + |\mathbb{E}[f(\Psi^{\leq \ell}(\tilde{\zeta}))] - \mathbb{E}[f(\Psi^{\leq \ell}(\tilde{\xi}))]|, \end{aligned} \quad (4.6)$$

where we use the fact that, since ζ and ξ are independent with zero mean,

$$\mathbb{E}[|\Psi^{> \ell}(\zeta)|^2] = \mathbb{E}[|\Psi^{> \ell}(\xi)|^2] = \sum_{|I| > \ell} \psi(I)^2 = C_{\Psi^{> \ell}}.$$

To bound the first term in (4.6), we write $\tilde{\Psi}^{> \ell}(x) := \Psi^{> \ell}(x + \mu)$, which has kernel $\tilde{\psi}^{> \ell}$ (note that first we truncate the kernel and then we shift $x \mapsto x + \mu$). Since ζ and ξ are independent with zero mean and variance one, by Lemma 4.1 we have

$$\mathbb{E}[|\Psi^{> \ell}(\tilde{\zeta})|^2] = \mathbb{E}[|\tilde{\Psi}^{> \ell}(\zeta)|^2] = \sum_J \tilde{\psi}^{> \ell}(J)^2 =: C_{\tilde{\Psi}^{> \ell}} \leq e^{c\mu/\varepsilon} C_{\Psi^{(\varepsilon), > \ell}}.$$

Substituting this bound into (4.6) then leads to the first term in (2.27).

To bound the second term in (4.6), we write $\tilde{\Psi}^{\leq \ell}(x) := \Psi^{\leq \ell}(x + \mu)$, and then apply Theorem 2.6 to obtain

$$\begin{aligned} |\mathbb{E}[f(\Psi^{\leq \ell}(\tilde{\zeta}))] - \mathbb{E}[f(\Psi^{\leq \ell}(\tilde{\xi}))]| &= |\mathbb{E}[f(\tilde{\Psi}^{\leq \ell}(\zeta))] - \mathbb{E}[f(\tilde{\Psi}^{\leq \ell}(\xi))]| \\ &\leq C_f C_{\tilde{\Psi}^{\leq \ell}} \left(16\ell^2 m_2^{> M} + 70^{\ell+1} (m_3^{\leq M})^\ell \max_{i \in \mathbb{T}} \sqrt{\text{Inf}_i[\tilde{\Psi}^{\leq \ell}]} \right). \end{aligned}$$

Applying the bounds in Lemma 4.1 to $\Psi^{\leq \ell}$ then gives the remaining terms in (2.27), where we have combined and upper bounded factors of $e^{c\mu/\varepsilon}$. \square

Proof of Theorem 2.6. We note that it is sufficient to prove the theorem in the case $|\mathbb{T}| < \infty$, because the general case follows by considering finite $\Lambda_N \uparrow \mathbb{T}$. For notational simplicity, we assume that $\mathbb{T} = [N] := \{1, \dots, N\}$.

Step 1. We first truncate Ψ to a degree ℓ polynomial $\Psi^{\leq \ell}$. By the same calculations as in (4.5) and (4.6), we have

$$\begin{aligned} |\mathbb{E}[f(\Psi(\zeta))] - \mathbb{E}[f(\Psi(\xi))]| &\leq 2\|f'\|_\infty \mathbb{E}[|\Psi^{> \ell}(\zeta)|^2]^{1/2} \\ &\quad + |\mathbb{E}[f(\Psi^{\leq \ell}(\zeta))] - \mathbb{E}[f(\Psi^{\leq \ell}(\xi))]|. \end{aligned} \quad (4.7)$$

This leads to the first term on the right hand side of (2.24).

Step 2. For a fixed $f \in C_b^3(\mathbb{R})$, we denote

$$f(\Psi^{\leq \ell}(x)) =: g(x_1, \dots, x_N).$$

For a vector $x \in \mathbb{R}^N$ and $y \in \mathbb{R}$, we also set

$$h_j^x(y) := g(x_1, \dots, x_{j-1}, y, x_{j+1}, \dots, x_N).$$

Using this notation, we can write

$$\begin{aligned} f(\Psi^{\leq \ell}(\zeta)) - f(\Psi^{\leq \ell}(\xi)) &= \sum_{j=1}^N (g(\zeta_1, \dots, \zeta_j, \xi_{j+1}, \dots, \xi_N) - g(\zeta_1, \dots, \zeta_{j-1}, \xi_j, \dots, \xi_N)) \\ &= \sum_{j=1}^N (h_j^{X^j}(\zeta_j) - h_j^{X^j}(\xi_j)), \end{aligned} \quad (4.8)$$

where we have used the notation

$$X^j := (X_1^j, \dots, X_N^j) := (\zeta_1, \dots, \zeta_j, \xi_{j+1}, \dots, \xi_N). \quad (4.9)$$

Next, we will be Taylor expanding each term in (4.8). For this we note that for $y \in \mathbb{R}$,

$$h_j^x(y) = h_j^x(0) + y \frac{dh_j^x}{dy}(0) + \frac{y^2}{2} \frac{d^2 h_j^x}{dy^2}(0) + R_j^x(y),$$

where the remainder term has the form

$$R_j^x(y) = \frac{1}{2} \int_0^y \frac{d^3 h_j^x}{dy^3}(t)(y-t)^2 dt = \frac{1}{2} \frac{d^2 h_j^x}{dy^2}(0)y^2 + \int_0^y \frac{d^2 h_j^x}{dy^2}(t)(y-t) dt,$$

from which it follows that

$$|R_j^x(y)| \leq \min \left\{ \frac{|y|^3}{6} \left\| \frac{d^3 h_j^x}{dy^3} \right\|_{\infty}, y^2 \left\| \frac{d^2 h_j^x}{dy^2} \right\|_{\infty} \right\}. \quad (4.10)$$

Inserting the Taylor expansion into (4.8) we obtain

$$\begin{aligned} f(\Psi^{\leq \ell}(\zeta)) - f(\Psi^{\leq \ell}(\xi)) &= \sum_{j=1}^N \left\{ h_j^{X^j}(0) + \zeta_j \frac{dh_j^{X^j}}{dy}(0) + \frac{\zeta_j^2}{2} \frac{d^2 h_j^{X^j}}{dy^2}(0) + R_j^{X^j}(\zeta_j) \right. \\ &\quad \left. - h_j^{X^j}(0) - \xi_j \frac{dh_j^{X^j}}{dy}(0) - \frac{\xi_j^2}{2} \frac{d^2 h_j^{X^j}}{dy^2}(0) - R_j^{X^j}(\xi_j) \right\}. \end{aligned}$$

Since ζ_i and ξ_i both have zero mean and unit variance, taking expectation, we get

$$\begin{aligned} |\mathbb{E}[f(\Psi^{\leq \ell}(\zeta))] - \mathbb{E}[f(\Psi^{\leq \ell}(\xi))]| &= \left| \sum_{j=1}^N \mathbb{E}[R_j^{X^j}(\zeta_j) - R_j^{X^j}(\xi_j)] \right| \\ &\leq \sum_{j=1}^N \left[\mathbb{E}[|R_j^{X^j}(\zeta_j)|] + \sum_{j=1}^N \mathbb{E}[|R_j^{X^j}(\xi_j)|] \right]. \end{aligned} \quad (4.11)$$

Since the estimates for both sums are identical, we will focus on the first one.

Step 3. The derivatives of $h_j^x(\cdot)$ are easily computed:

$$\begin{aligned} &\frac{d^m h_j^x}{dy^m}(y) \\ &= f^{(m)}(\Psi^{\leq \ell}(x_1, \dots, x_{j-1}, y, x_{j+1}, \dots, x_N)) \left(\frac{\partial \Psi^{\leq \ell}(x_1, \dots, x_{j-1}, y, x_{j+1}, \dots, x_N)}{\partial y} \right)^m \\ &= f^{(m)}(\Psi^{\leq \ell}(x_1, \dots, x_{j-1}, y, x_{j+1}, \dots, x_N)) \left(\sum_{I \ni j, |I| \leq \ell} \psi(I) x^{I \setminus \{j\}} \right)^m. \end{aligned}$$

Then, setting $C_f = \max\{\|f'\|_\infty, \|f^{(2)}\|_\infty, \|f^{(3)}\|_\infty\}$, we can apply (4.10) to bound

$$\sum_{j=1}^N \mathbb{E}[|R_j^{X^j}(\zeta_j)|] \leq C_f \sum_{j=1}^N \mathbb{E}[\varphi(L_j(X^j))], \quad (4.12)$$

where X^j was introduced in (4.9), and we define

$$\varphi(x) := \min\{|x|^3/6, |x|^2\}, \quad L_j(x) := \sum_{I \ni j, |I| \leq \ell} \psi(I) x^I. \quad (4.13)$$

Notice that $L_j(x)$ includes the variable x_j in the product as a result of absorbing the powers of y in (4.10). Also note that $\varphi(x) = \varphi(|x|)$ and

$$\varphi(a+b) \leq \varphi(2 \max\{|a|, |b|\}) \leq \varphi(2|a|) + \varphi(2|b|) \leq 4|a|^2 + \frac{4}{3}|b|^3.$$

Writing $L_j(X^j) = (L_j(X^j) - L_j(X^{j-})) + L_j(X^{j-})$, where $X^{j-} := (X_1^{j-}, \dots, X_N^{j-})$ is a suitably truncated version of X^j , we then obtain

$$\mathbb{E}[\varphi(L_j(X^j))] \leq 4\mathbb{E}[(L_j(X^j) - L_j(X^{j-}))^2] + \frac{4}{3}\mathbb{E}[|L_j(X^{j-})|^3]. \quad (4.14)$$

The two terms on the right hand side will give rise respectively to the two terms on the right hand side of (2.24).

Step 4. We now describe the truncation procedure. This new ingredient, with respect to [MOO10], is tailored to control random variables with finite second moments and

uniformly integrable squares. Fix $M \in (0, \infty)$. We want to decompose any real-valued random variable Y with *zero mean* and finite variance in the following way:

$$Y = Y^- + Y^+, \quad (4.15)$$

where Y^- , Y^+ are functions of Y and possibly of some extra randomness, such that

$$\begin{aligned} \mathbb{E}[Y^-] = \mathbb{E}[Y^+] = 0, \quad Y^- Y^+ = 0, \\ |Y^-| \leq |Y| \mathbb{1}_{\{|Y| \leq M\}}, \quad \mathbb{E}[(Y^+)^2] \leq 2\mathbb{E}[Y^2 \mathbb{1}_{\{|Y| > M\}}]. \end{aligned} \quad (4.16)$$

If $\mathbb{E}[Y \mathbb{1}_{\{-M \leq Y \leq M\}}] = 0$ we are done: just choose $Y^- := Y \mathbb{1}_{\{-M \leq Y \leq M\}}$ and $Y^+ := Y - Y^-$. If, on the other hand, $\mathbb{E}[Y \mathbb{1}_{\{-M \leq Y \leq M\}}] > 0$ (the strictly negative case is analogous), we set

$$\bar{T} := \sup\{T \in [0, M] : \mathbb{E}[Y \mathbb{1}_{\{-M \leq Y \leq T\}}] \leq 0\} \in [0, M].$$

Note that $\mathbb{E}[Y \mathbb{1}_{\{-M \leq Y \leq \bar{T}\}}] \geq 0$, because $T \mapsto \mathbb{E}[Y \mathbb{1}_{\{-M \leq Y \leq T\}}]$ is right-continuous. If $\mathbb{E}[Y \mathbb{1}_{\{-M \leq Y \leq \bar{T}\}}] = 0$, and we define $Y^- := Y \mathbb{1}_{\{-M \leq Y \leq \bar{T}\}}$ and $Y^+ := Y - Y^-$, then all the properties in (4.16) are clearly satisfied, except the last one that will be checked below. Finally, we consider the case $\mathbb{E}[Y \mathbb{1}_{\{-M \leq Y \leq \bar{T}\}}] > 0$ (then necessarily $\bar{T} > 0$). Since $\mathbb{E}[Y \mathbb{1}_{\{-M \leq Y < \bar{T}\}}] \leq 0$ by definition of \bar{T} , we must have $\mathbb{P}(Y = \bar{T}) > 0$. Then take a random variable U uniformly distributed in $(0, 1)$ and independent of Y , and define

$$Y^- := Y(\mathbb{1}_{\{-M \leq Y < \bar{T}\}} + \mathbb{1}_{\{Y = \bar{T}, U \leq \varrho\}}), \quad \text{where} \quad \varrho := \frac{-\mathbb{E}[Y \mathbb{1}_{\{-M \leq Y < \bar{T}\}}]}{\bar{T} \mathbb{P}(Y = \bar{T})} \in (0, 1).$$

Set $Y^+ := Y - Y^-$. Then all the properties (4.16) but the last one are clearly satisfied.

For the last property, we write

$$\begin{aligned} \mathbb{E}[(Y^+)^2] &= \mathbb{E}[(Y^+)^2 \mathbb{1}_{\{|Y| > M\}}] + \mathbb{E}[(Y^+)^2 \mathbb{1}_{\{|Y| \leq M\}}] \\ &= \mathbb{E}[Y^2 \mathbb{1}_{\{|Y| > M\}}] + \mathbb{E}[(Y^+)^2 \mathbb{1}_{\{|Y| \leq M\}}], \end{aligned}$$

because $Y^+ = Y$ on the event $\{|Y| > M\}$. For the second term, since $0 \leq Y^+ \leq M$ on the event $\{|Y| \leq M\}$, we can write $(Y^+)^2 \leq M Y^+$ (no absolute value needed). Since $Y^- = Y^- \mathbb{1}_{\{|Y| \leq M\}}$ has zero mean by (4.16), we obtain

$$\begin{aligned} \mathbb{E}[(Y^+)^2 \mathbb{1}_{\{|Y| \leq M\}}] &\leq M \mathbb{E}[Y^+ \mathbb{1}_{\{|Y| \leq M\}}] = M \mathbb{E}[(Y^+ + Y^-) \mathbb{1}_{\{|Y| \leq M\}}] \\ &= M \mathbb{E}[Y \mathbb{1}_{\{|Y| \leq M\}}] = M(-\mathbb{E}[Y \mathbb{1}_{\{|Y| > M\}}]) \leq \mathbb{E}[Y^2 \mathbb{1}_{\{|Y| > M\}}], \end{aligned}$$

where we have used the fact that $\mathbb{E}[Y] = 0$ by assumption, and Markov's inequality. The last relation in (4.16) is proved.

Step 5. We apply the decomposition (4.15) to the random variables $(X_i^j)_{i \in [N]}$, defined in (4.9), where the extra randomness used in the construction is taken independently for each variable; then we write

$$X_i^j = X_i^{j-} + X_i^{j+},$$

and the properties in (4.16) are satisfied. Note that by (4.13) we can write

$$L_j(X^j) - L_j(X^{j-}) = \sum_{I \ni j, |I| \leq \ell} \psi(I) \sum_{\Gamma \subseteq I, |\Gamma| \geq 1} (X^{j+})^\Gamma (X^{j-})^{I \setminus \Gamma}.$$

Since the random variables $X_1^{j-}, X_1^{j+}, X_2^{j-}, X_2^{j+}, \dots$ are orthogonal in L^2 by construction, setting $\sigma_{\pm, i}^2 := \mathbb{E}[(X_i^{j\pm})^2]$ and observing that $\sigma_{-, i}^2 + \sigma_{+, i}^2 = \mathbb{V}\text{ar}(X_i^j) = 1$ we obtain

$$\begin{aligned} \mathbb{E}[(L_j(X^j) - L_j(X^{j-}))^2] &= \sum_{I \ni j, |I| \leq \ell} \psi(I)^2 \sum_{\Gamma \subseteq I, |\Gamma| \geq 1} (\sigma_+^2)^\Gamma (\sigma_-^2)^{I \setminus \Gamma} \\ &= \sum_{I \ni j, |I| \leq \ell} \psi(I)^2 (1 - (\sigma_-^2)^I) \leq \sum_{I \ni j, |I| \leq \ell} \psi(I)^2 (1 - (1 - \bar{\sigma}_+^2)^{|I|}), \end{aligned}$$

where

$$\bar{\sigma}_+^2 := \max_{i=1, \dots, N} \sigma_{+, i}^2 = \max_{i=1, \dots, N} \mathbb{E}[(X_i^{j+})^2] \leq 2 \max_{i=1, \dots, N} \mathbb{E}[(X_i^j)^2 \mathbb{1}_{\{|X_i^j| > M\}}] \leq 2m_2^{\leq M},$$

having used (4.16) and the definition of $m_2^{\leq M}$ in (2.22) (recall (4.9)). Since $1 - (1 - p)^n \leq np$, we obtain

$$\begin{aligned} \sum_{j=1}^N \mathbb{E}[(L_j(X^j) - L_j(X^{j-}))^2] &\leq 2m_2^{\leq M} \sum_{j=1}^N \left(\sum_{I \ni j, |I| \leq \ell} |I| \psi(I)^2 \right) \\ &\leq 2m_2^{\leq M} \ell^2 \sum_{|I| \leq \ell} \psi(I)^2. \end{aligned}$$

Tracing back through (4.7), (4.11), (4.12) and (4.14), we note that this gives the first term on the right hand side of (2.24).

Step 6. We finally consider the contribution of the second term in (4.14). We apply the hypercontractivity results in [MOO10]: by Propositions 3.16 and 3.12 therein, denoting by $\|Y\|_q := \mathbb{E}[|Y|^q]^{1/q}$ the usual L^q norm, for every $q > 2$ we have

$$\|L_j(X^{j-})\|_q \leq (B_q)^\ell \|L_j(X^{j-})\|_2, \quad (4.17)$$

where

$$B_q := 2\sqrt{q-1} \max_{i \in [N]} \frac{\|X_i^{j-}\|_q}{\|X_i^{j-}\|_2}.$$

Let us set $Y := X_i^j$ for short and choose $q = 3$. Since $|Y^-| \leq M$, by (4.16) we have

$$\|Y^-\|_3 \leq \mathbb{E}[|Y|^3 \mathbb{1}_{\{|Y| \leq M\}}]^{1/3} \leq (m_3^{\leq M})^{1/3},$$

where we recall (4.9) and the definition of $m_3^{\leq M}$ from (2.22). On the other hand, again

by (4.16),

$$\begin{aligned}\|Y^-\|_2^2 &= \|Y\|_2^2 - \|Y^+\|_2^2 = \mathbb{E}[Y^2] - \mathbb{E}[(Y^+)^2] \geq \mathbb{E}[Y^2] - 2\mathbb{E}[Y^2 \mathbf{1}_{\{|Y|>M\}}] \\ &= 1 - 2\mathbb{E}[(X_i^j)^2 \mathbf{1}_{\{|X_i^j|>M\}}] \geq 1 - 2m_2^{>M},\end{aligned}$$

hence

$$B_3 \leq 2\sqrt{2} \frac{(m_3^{\leq M})^{1/3}}{\sqrt{1 - 2m_2^{>M}}} \leq 4(m_3^{\leq M})^{1/3}$$

provided $m_{>M}^{(2)} \leq 1/4$, as in the assumptions of Theorem 2.6. Therefore, (4.17) for $q = 3$ yields

$$\mathbb{E}[|L_j(X^{j-})|^3] \leq 64^\ell (m_3^{\leq M})^\ell \mathbb{E}[L_j(X^{j-})^2]^{3/2}.$$

Note that, since $\mathbb{E}[(X_i^{j-})^2] \leq \mathbb{E}[(X_i^j)^2] = 1$, we have

$$\mathbb{E}[L_j(X^{j-})^2] = \sum_{I \ni j, |I| \leq \ell} \psi(I)^2 \prod_{i \in I} \mathbb{E}[(X_i^{j-})^2] \leq \sum_{I \ni j, |I| \leq \ell} \psi(I)^2 = \text{Inf}_j[\Psi^{\leq \ell}].$$

Therefore

$$\begin{aligned}\sum_{j=1}^N \mathbb{E}[|L_j(X^{j-})|^3] &\leq 64^\ell (m_3^{\leq M})^\ell \left(\max_{i \in [N]} \sqrt{\text{Inf}_i[\Psi^{\leq \ell}]} \right) \sum_{j=1}^N \sum_{I \ni j, |I| \leq \ell} \psi(I)^2 \\ &\leq \ell 64^\ell (m_3^{\leq M})^\ell \left(\max_{i \in [N]} \sqrt{\text{Inf}_i[\Psi^{\leq \ell}]} \right) \sum_{|I| \leq \ell} \psi(I)^2.\end{aligned}\tag{4.18}$$

This gives the third term on the right hand side of (2.24) (because $\frac{8}{3} \ell 64^\ell \leq 70^{\ell+1}$), which completes the proof. \square

5. Proof of the convergence to Wiener chaos

In this section, we prove Theorems 2.3 and 2.5 on the convergence of polynomial chaos expansions to Wiener chaos expansions.

Proof of Theorem 2.3. Let $W(\cdot)$ be the d -dimensional white noise used to define the Wiener chaos expansion for Ψ_0 in Theorem 2.3. Given the tessellation \mathcal{C}_δ indexed by \mathbb{T}_δ , where the cell $\mathcal{C}_\delta(x)$ has the same volume v_δ for each $x \in \mathbb{T}_\delta$, we define

$$\xi_{\delta,x} := \mu_\delta(x) + v_\delta^{-1/2} \int_{\mathcal{C}_\delta(x)} \sigma_\delta W(dy) = v_\delta^{-1/2} \int_{\mathcal{C}_\delta(x)} (\sigma_\delta W(dy) + \bar{\mu}_\delta(y) dy), \tag{5.1}$$

where we recall that $\bar{\mu}_\delta := v_\delta^{-1/2} \mu_\delta$ by (2.9). Note that $\xi_\delta := (\xi_{\delta,x})_{x \in \mathbb{T}}$ is a family of independent Gaussian random variables with the same mean and variance as $\zeta_\delta = (\zeta_{\delta,x})_{x \in \mathbb{T}_\delta}$.

We recall that our goal is to show that $\Psi_\delta(\zeta_\delta) \rightarrow \Psi_0$ in distribution. The strategy is first to focus on $\Psi_\delta(\xi_\delta)$ instead of $\Psi_\delta(\zeta_\delta)$: we can write $\Psi_\delta(\xi_\delta)$ as a Wiener chaos expansion with respect to $W(\cdot)$, like Ψ_0 , and show that $\mathbb{E}[|\Psi_\delta(\xi_\delta) - \Psi_0|^2] \rightarrow 0$ as $\delta \downarrow 0$; then we use the Lindeberg principle, Theorem 2.8, to replace $\Psi_\delta(\xi_\delta)$ by $\Psi_\delta(\zeta_\delta)$.

Step 1. We first show that for each degree $k \in \mathbb{N}_0$,

$$\lim_{\delta \downarrow 0} \mathbb{E}[|\Psi_\delta^{(k)}(\xi_\delta) - \Psi_0^{(k)}|^2] = 0, \quad (5.2)$$

where $\Psi_\delta^{(k)}$ is a polynomial with kernel

$$\psi_\delta^{(k)}(I) := \psi_\delta(I) \mathbb{1}_{\{|I|=k\}},$$

and similarly $\Psi_0^{(k)}$ is defined as Ψ_0 in (2.12), except the kernel ψ_0 therein is replaced by $\psi_0^{(k)}(I) := \psi_0(I) \mathbb{1}_{\{|I|=k\}}$ (that is, we take the k -th term in the sum). Recalling from (2.10) that

$$\bar{\psi}_\delta(I) := v_\delta^{-|I|/2} \psi_\delta(I),$$

and extending $\bar{\psi}_\delta$ to a function defined on \mathbb{R}^k , as discussed before Theorem 2.3, we can write

$$\begin{aligned} \Psi_\delta^{(k)}(\xi_\delta) &= \sum_{I \in \mathcal{P}^{\text{fin}}(\mathbb{T}_\delta), |I|=k} \psi_\delta(I) \xi_\delta^I \\ &= \frac{1}{k!} \int \cdots \int_{(\mathbb{R}^d)^k} \bar{\psi}_\delta(y_1, \dots, y_k) \prod_{i=1}^k (\sigma_\delta W(dy_i) + \bar{\mu}_\delta(y_i) dy_i) = \\ &= \frac{1}{k!} \sum_{I \subset [k]: \{1, \dots, k\}} \int_{(\mathbb{R}^d)^{k-|I|}} \left(\int_{(\mathbb{R}^d)^{|I|} } \sigma_\delta^{k-|I|} \bar{\psi}_\delta(y_1, \dots, y_k) \prod_{i \in I} \bar{\mu}_\delta(y_i) dy_i \right) \prod_{j \in [k] \setminus I} W(dy_j). \end{aligned}$$

A similar expansion holds for $\Psi_0^{(k)}$ with $\bar{\psi}_\delta, \bar{\mu}_\delta, \sigma_\delta$ replaced respectively by ψ_0, μ_0, σ_0 . Comparing the two expansions term by term for each $I \subset [k]$, we obtain, by the triangle inequality and the Ito isometry (2.2),

$$\begin{aligned} &\mathbb{E}[|\Psi_\delta^{(k)}(\xi_\delta) - \Psi_0^{(k)}|^2]^{1/2} \\ &\leq \frac{1}{k!} \sum_{I \subset [k]} \left(\int_{(\mathbb{R}^d)^{k-|I|} } \left\{ \int_{(\mathbb{R}^d)^{|I|} } \left(\sigma_\delta^{k-|I|} \bar{\psi}_\delta \prod_{i \in I} \bar{\mu}_\delta(y_i) - \sigma_0^{k-|I|} \psi_0 \prod_{i \in I} \mu_0(y_i) \right) \prod_{i \in I} dy_i \right\}^2 \right. \\ &\quad \left. \cdot \prod_{j \in [k] \setminus I} dy_j \right)^{1/2}. \quad (5.3) \end{aligned}$$

To see that each term above tends to 0 as $\delta \downarrow 0$, assume without loss of generality that either $I = \emptyset$ or $I = [n]$ for some $1 \leq n \leq k$. We then write in a telescopic sum

$$\begin{aligned} &\sigma_\delta^{k-n} \bar{\psi}_\delta \prod_{i=1}^n \bar{\mu}_\delta(y_i) - \sigma_0^{k-n} \psi_0 \prod_{i=1}^n \mu_0(y_i) \\ &= \Delta_0 \prod_{i=1}^n \bar{\mu}_\delta(y_i) + \sigma_0^{k-n} \psi_0 \Delta_1 \prod_{i=2}^n \bar{\mu}_\delta(y_i) + \cdots + \sigma_0^{k-n} \psi_0 \prod_{i=1}^{n-1} \mu_0(y_i) \Delta_n, \quad (5.4) \end{aligned}$$

where $\Delta_0 := \sigma_\delta^{k-n} \bar{\psi}_\delta - \sigma_0^{k-n} \psi_0$ and $\Delta_i := \bar{\mu}_\delta(y_i) - \mu_0(y_i)$ for $i = 1, \dots, n$. The contribution of each term from (5.4) to the integrals in (5.3) can be bounded by applying

Cauchy–Schwarz to the inner integral, in such a way that the ψ term is separated from the product of the μ 's. It is then easily seen that all terms tend to 0 as $\delta \downarrow 0$ by the assumptions in Theorem 2.3 that $\bar{\mu}_\delta \rightarrow \mu_0$ and $\bar{\psi}_\delta \rightarrow \psi_0$ in L^2 , together with $\sigma_\delta \rightarrow \sigma_0 \in (0, \infty)$. This implies (5.2).

Step 2. We next give a uniform L^2 bound on the tail of the series for $\Psi_\delta(\zeta_\delta)$ and Ψ_0 . More precisely, for any $\ell < N$, let $\Psi_\delta^{(\ell, N)} := \sum_{\ell < k < N} \Psi_\delta^{(k)}$ and $\Psi_0^{(\ell, N)} := \sum_{\ell < k < N} \Psi_0^{(k)}$. Write $\Psi_\delta^{>\ell}$ and $\Psi_0^{>\ell}$ for the case $N = \infty$. We will show that

$$\lim_{\ell \rightarrow \infty} \limsup_{\delta \downarrow 0} \mathbb{E}[|\Psi_\delta^{>\ell}(\zeta_\delta)|^2] = 0, \quad \text{and} \quad \Psi_0 = \sum_{k=0}^{\infty} \Psi_0^{(k)} \quad \text{converges in } L^2. \quad (5.5)$$

Together with (5.2) and the fact that $\mathbb{E}[\Psi_\delta^{(\ell, N)}(\xi_\delta)^2] = \mathbb{E}[\Psi_\delta^{(\ell, N)}(\zeta_\delta)^2]$ for all $0 \leq \ell < N \leq \infty$, it follows that

$$\mathbb{E}[\Psi_\delta(\zeta_\delta)^2] = \mathbb{E}[\Psi_\delta(\xi_\delta)^2] \rightarrow \mathbb{E}[\Psi_0^2] \quad \text{as } \delta \downarrow 0,$$

which is one of the claims in Theorem 2.3.

If we denote $\xi_{\delta, x} = \mu_\delta(x) + \tilde{\xi}_{\delta, x}$ and let $\tilde{\Psi}_\delta^{(\ell, N)}(x) := \Psi_\delta^{(\ell, N)}(x + \mu_\delta)$, then for $\varepsilon > 0$ as specified in Theorem 2.3(iii) we can apply Lemma 4.1 (actually a modification of it, where we take into account that the random variables do not have normalized variance) to obtain

$$\begin{aligned} \mathbb{E}[|\Psi_\delta^{(\ell, N)}(\xi_\delta)|^2] &= \mathbb{E}[|\tilde{\Psi}_\delta^{(\ell, N)}(\tilde{\xi}_\delta)|^2] \\ &\leq e^{\frac{1}{\varepsilon \sigma_\delta^2} \sum_{x \in \mathbb{T}_\delta} \mu_\delta(x)^2} \sum_{\substack{I \in \mathcal{P}^{\text{fin}}(\mathbb{T}_\delta) \\ \ell < |I| < N}} (1 + \varepsilon)^{|I|} (\sigma_\delta^2)^{|I|} \psi_\delta(I)^2. \end{aligned} \quad (5.6)$$

A similar relation to (5.6) holds with ζ_δ replacing ξ_δ . Since $\sum_{x \in \mathbb{T}_\delta} \mu_\delta(x)^2 = \|\bar{\mu}_\delta\|_{L^2(\mathbb{R}^d)}^2$ (recall the extension of $f : \mathbb{T}_\delta \rightarrow \mathbb{R}$ to $f : \mathbb{R}^d \rightarrow \mathbb{R}$ as specified before Theorem 2.3), the assumptions in Theorem 2.3(i)&(iii) immediately imply the first limit in (5.5) if we let $N \uparrow \infty$ in (5.6). It also shows that $\Psi_\delta(\xi_\delta) = \sum_{k=0}^{\infty} \Psi_\delta^{(k)}(\xi_\delta)$, as well as $\Psi_\delta(\zeta_\delta)$, are L^2 convergent series.

By (5.2), we can let $\delta \downarrow 0$ in (5.6) to obtain

$$\begin{aligned} \mathbb{E}[|\Psi_0^{(\ell, N)}|^2] &= \lim_{\delta \downarrow 0} \mathbb{E}[|\Psi_\delta^{(\ell, N)}(\xi_\delta)|^2] \\ &\leq e^{\frac{1}{\varepsilon \sigma_0^2} \|\mu_0\|_{L^2(\mathbb{R}^d)}^2} \limsup_{\delta \downarrow 0} \sum_{\substack{I \in \mathcal{P}^{\text{fin}}(\mathbb{T}_\delta) \\ |I| > \ell}} (1 + \varepsilon)^{|I|} (\sigma_\delta^2)^{|I|} \psi_\delta(I)^2. \end{aligned}$$

By assumptions (i) and (iii) of Theorem 2.3, this implies that $\Psi_0 = \sum_{k=0}^{\infty} \Psi_0^{(k)}$ is an L^2 convergent series, proving the second relation in (5.5).

We also observe that if $\mu_\delta(x) \equiv 0$ there is no need to apply Lemma 4.1: relation (5.6) holds without the exponential prefactor and with $\varepsilon = 0$, because $\xi_\delta(x)$, $x \in \mathbb{T}_\delta$ are random variables with zero mean and finite variance σ_δ (cf. Remark 5.5).

Step 3. We now use the Lindeberg principle, Theorem 2.8, to show that for each $\ell \in \mathbb{N}_0$, $\Psi_\delta^{\leq \ell}(\zeta_\delta) := \sum_{k=0}^{\ell} \Psi_\delta^{(k)}(\zeta_\delta)$ has the same limiting distribution as $\Psi_\delta^{\leq \ell}(\xi_\delta)$ as $\delta \downarrow 0$. Together with the L^2 convergence of $\Psi_\delta^{\leq \ell}(\xi_\delta)$ to $\Psi_0^{\leq \ell} := \sum_{k=0}^{\ell} \Psi_0^{(k)}$ proved in Step 1 (cf. (5.2)), as well as the uniform L^2 bound on $\Psi_\delta^{\leq \ell}(\zeta_\delta)$ and $\Psi_0^{\leq \ell}$ shown in Step 2 (cf. (5.5)), this implies that $\Psi_\delta(\zeta_\delta)$ converges in distribution to Ψ_0 as $\delta \downarrow 0$.

It suffices to show that for all $f \in \mathcal{C}^3$ with $C_f := \max\{\|f'\|_\infty, \|f''\|_\infty, \|f'''\|_\infty\} < \infty$,

$$\lim_{\delta \downarrow 0} |\mathbb{E}[f(\Psi_\delta^{\leq \ell}(\zeta_\delta)) - f(\Psi_\delta^{\leq \ell}(\xi_\delta))]| = 0. \quad (5.7)$$

With ε as specified in Theorem 2.3(iii), we can apply Theorem 2.8 (actually a slight modification of it, taking into account the non-normalized variance σ_δ of the variables used here): the absolute value on the left hand side of (5.7) is bounded by

$$e^{\frac{2}{\varepsilon \sigma_\delta^2} \|\bar{\mu}_\delta\|_{L^2(\mathbb{R}^d)}^2} C_f \widehat{\mathcal{C}}_{\Psi_\delta^{(\varepsilon), \leq \ell}} \left(16\ell^2 \frac{m_2^{\geq M}}{\sigma_\delta^2} + 70^{\ell+1} \left(\frac{m_3^{\leq M}}{\sigma_\delta^3} \right)^\ell \max_{i \in \mathbb{T}_\delta} \sqrt{\widehat{\text{Inf}}_i[\Psi_\delta^{(\varepsilon), \leq \ell}]} \right), \quad (5.8)$$

where

$$\begin{aligned} \widehat{\mathcal{C}}_{\Psi_\delta^{(\varepsilon), \leq \ell}} &:= \sum_{|I| \leq \ell} (1 + \varepsilon)^{|I|} (\sigma_\delta^2)^{|I|} \psi_\delta(I)^2 = \sum_{k=0}^{\ell} (1 + \varepsilon)^k \frac{1}{k!} \|\sigma_\delta \bar{\psi}_\delta\|_{L^2((\mathbb{R}^d)^k)}^2 \\ \widehat{\text{Inf}}_i[\Psi_\delta^{(\varepsilon), \leq \ell}] &:= \sum_{I \ni i, |I| \leq \ell} (1 + \varepsilon)^{|I|} (\sigma_\delta^2)^{|I|} \psi_\delta(I)^2 \\ &= \sum_{k=1}^{\ell} \frac{(1 + \varepsilon)^k}{(k-1)!} (\sigma_\delta^2)^k \|\bar{\psi}_\delta \mathbb{1}_{\{x_1 \in \mathcal{C}_\delta(i)\}}\|_{L^2((\mathbb{R}^d)^k)}^2, \end{aligned}$$

and we recall that $\mathcal{C}_\delta(i) \subset \mathbb{R}^d$ is the cell indexed by $i \in \mathbb{T}_\delta$ in the tessellation \mathcal{C}_δ . We are left with showing that (5.8) vanishes as $\delta \downarrow 0$. Note that

- $\widehat{\mathcal{C}}_{\Psi_\delta^{(\varepsilon), \leq \ell}}$ is uniformly bounded by Theorem 2.3(ii)–(iii);
- $\|\bar{\mu}_\delta\|_{L^2(\mathbb{R}^d)}^2 \rightarrow \|\mu_0\|_{L^2(\mathbb{R}^d)}^2$ and $\sigma_\delta \rightarrow \sigma_0 > 0$ by Theorem 2.3(i);
- $m_3^{\leq M} \leq M^3$ and $m_2^{\geq M}$ can be made arbitrarily small by choosing M large, by the definition (2.22) (to be applied to the centered variables $\zeta_{\delta,x} - \mathbb{E}[\zeta_{\delta,x}]$) and the fact that $((\zeta_{\delta,x} - \mathbb{E}[\zeta_{\delta,x}])^2)_{\delta \in (0,1), x \in \mathbb{T}_\delta}$ are uniformly integrable by assumption.

It only remains to verify that $\widehat{\text{Inf}}_i[\Psi_\delta^{(\varepsilon), \leq \ell}]$ vanishes as $\delta \downarrow 0$, uniformly in $i \in \mathbb{T}_\delta$. Since

$$\widehat{\text{Inf}}_i[\Psi_\delta^{(\varepsilon), \leq \ell}] \leq 2 \sum_{k=1}^{\ell} \frac{(1 + \varepsilon)^k}{(k-1)!} (\sigma_\delta^2)^k (\|\bar{\psi}_\delta - \psi_0\|_{L^2((\mathbb{R}^d)^k)}^2 + \|\psi_0 \mathbb{1}_{\{x_1 \in \mathcal{C}_\delta(i)\}}\|_{L^2((\mathbb{R}^d)^k)}^2),$$

one has $\|\bar{\psi}_\delta - \psi_0\|_{L^2((\mathbb{R}^d)^k)}^2 \rightarrow 0$ by Theorem 2.3(ii), and $\|\psi_0 \mathbb{1}_{\{x_1 \in \mathcal{C}_\delta(i)\}}\|_{L^2((\mathbb{R}^d)^k)}^2 \rightarrow 0$ uniformly in $i \in \mathbb{T}_\delta$ because $\text{Leb}(\mathcal{C}_\delta(i)) = v_\delta \downarrow 0$ uniformly in $i \in \mathbb{T}_\delta$. This completes Step 3 and establishes the convergence of $\Psi_\delta(\zeta_\delta)$ to Ψ_0 in distribution.

Step 4. Lastly, to show that the convergence of $\Psi_\delta(\zeta_\delta)$ to Ψ_0 in distribution extends to the joint distribution of a finite collection of polynomial chaos expansions $(\Psi_{i,\delta}(\zeta_\delta))_{1 \leq i \leq M}$, we note that by the Cramér–Wold device, it suffices to show the convergence of $\sum_{i=1}^M c_i \Psi_{i,\delta}(\zeta_\delta)$ to $\sum_{i=1}^M c_i \Psi_{i,0}$ for any $c_1, \dots, c_M \in \mathbb{R}$. This follows from what we have proved so far, since $\sum_{i=1}^M c_i \Psi_{i,\delta}(x)$ is also a polynomial that satisfies all the required conditions. \square

Proof of Theorem 2.5. Instead of performing an L^2 estimate on the tail series $\Psi_\delta^{>\ell}(\zeta_\delta)$ as in Step 2 in the proof of Theorem 2.3, we shall give an L^p estimate for any $p \in (0, 2)$. More precisely, we replace relation (5.5) by the following one: for any $p \in (0, 2)$,

$$\Psi_0 = \sum_{k=0}^{\infty} \Psi_0^{(k)} \quad \text{converges in } L^p, \quad \text{and} \quad \lim_{\ell \rightarrow \infty} \limsup_{\delta \downarrow 0} \mathbb{E}[|\Psi_\delta^{>\ell}(\zeta_\delta)|^p] = 0, \quad (5.9)$$

and we show that this holds under either condition (a) or (b) in Theorem 2.5. The rest of the proof of Theorem 2.3 then carries over without change.

The key to proving (5.9) is a change of measure argument. For $\ell < N$, let $\Psi_0^{(\ell, N)}$ and $\Psi_\delta^{(\ell, N)}$ be defined as in Step 2 in the proof of Theorem 2.3. Note that $\Psi_0^{(\ell, N)}$ is a finite sum of stochastic integrals with respect to the *biased* white noise $\sigma_0 W(dx) + \mu_0(x)dx$. By the discussion in Subsection 2.3.1, cf. (2.15), the joint distribution of these stochastic integrals is absolutely continuous with respect to the unbiased case $\mu_0(x) \equiv 0$, with Radon–Nikodym derivative

$$f(W) := \exp \left\{ \frac{1}{\sigma_0} \int \mu_0(y) W(dy) - \frac{1}{2\sigma_0^2} \int \mu_0(y)^2 dy \right\}. \quad (5.10)$$

Therefore, using $\Psi_{0, \mu_0 \equiv 0}^{(\ell, N)}$ to denote $\Psi_0^{(\ell, N)}$ with $\mu_0(x) \equiv 0$, for any $p \in (0, 2)$ we have

$$\begin{aligned} \mathbb{E}[|\Psi_0^{(\ell, N)}|^p] &= \mathbb{E}[f(W)|\Psi_{0, \mu_0 \equiv 0}^{(\ell, N)}|^p] \leq \mathbb{E}[f(W)^{\frac{2}{2-p}}]^{\frac{2-p}{2}} \mathbb{E}[|\Psi_{0, \mu_0 \equiv 0}^{(\ell, N)}|^2]^{p/2} \\ &= e^{\frac{p}{2(2-p)} \|\mu_0/\sigma_0\|_{L^2(\mathbb{R}^d)}^2} \mathbb{E}[|\Psi_{0, \mu_0 \equiv 0}^{(\ell, N)}|^2]^{p/2}, \end{aligned} \quad (5.11)$$

by Hölder’s inequality. By Theorem 2.3, when $\mu_\delta = \mu_0 \equiv 0$ it is enough to assume that condition (iii) therein holds with $\varepsilon = 0$ to guarantee that $\Psi_{0, \mu_0 \equiv 0} = \sum_{k=0}^{\infty} \Psi_{0, \mu_0 \equiv 0}^{(k)}$ is an L^2 convergent series. Therefore $\Psi_0 = \sum_{k=0}^{\infty} \Psi_0^{(k)}$ is convergent in L^p , by (5.11).

To control $\mathbb{E}[|\Psi_\delta^{>\ell}(\zeta_\delta)|^p]$ via a change of measure for ζ_δ is more subtle, since $(\zeta_{\delta, i})_{i \in \mathbb{T}_\delta}$ are not assumed to have finite exponential moments. We will instead perform an exponential change of measure on a bounded subset of the support of $\zeta_{\delta, i}$. Since by assumption $\|\mu_\delta\|_\infty \rightarrow 0$ and $((\zeta_{\delta, i} - \mu_\delta)^2)_{i \in \mathbb{T}_\delta}$ are uniformly integrable, also $(\zeta_{\delta, i}^2)_{i \in \mathbb{T}_\delta}$ are uniformly integrable. We can then apply Lemma B.1 of Appendix B: there exist independent random variables $\tilde{\zeta}_{\delta, i}$, whose law is absolutely continuous with respect to the

law of $\zeta_{\delta,i}$, with density $f_{\delta,i}(x)$, which satisfy (B.1)–(B.3). We can then write

$$\begin{aligned} \mathbb{E}[|\Psi_{\delta}^{(\ell,N)}(\zeta_{\delta})|^p] &= \mathbb{E}\left[\prod_{i \in \mathbb{T}_{\delta}} f_{\delta,i}(\zeta_{\delta,i})^{p/2} |\Psi_{\delta}^{(\ell,N)}(\zeta_{\delta})|^p \prod_{i \in \mathbb{T}_{\delta}} f_{\delta,i}(\zeta_{\delta,i})^{-p/2}\right] \\ &\leq \mathbb{E}\left[\prod_{i \in \mathbb{T}_{\delta}} f_{\delta,i}(\zeta_{\delta,i}) |\Psi_{\delta}^{(\ell,N)}(\zeta_{\delta})|^2\right]^{p/2} \mathbb{E}\left[\prod_{i \in \mathbb{T}_{\delta}} f_{\delta,i}(\zeta_{\delta,i})^{-\frac{p}{2-p}}\right]^{\frac{2-p}{2}} \\ &= \mathbb{E}[|\Psi_{\delta}^{(\ell,N)}(\tilde{\zeta}_{\delta})|^2]^{p/2} \prod_{i \in \mathbb{T}_{\delta}} \mathbb{E}[f_{\delta,i}(\zeta_{\delta,i})^{-\frac{p}{2-p}}]^{\frac{2-p}{2}}. \end{aligned} \quad (5.12)$$

Applying (B.1), we have

$$\prod_{i \in \mathbb{T}_{\delta}} \mathbb{E}[f_{\delta,i}(\zeta_{\delta,i})^{-\frac{p}{2-p}}]^{\frac{2-p}{2}} \leq e^{C_p \frac{2-p}{2} \sum_{i \in \mathbb{T}_{\delta}} \mu_{\delta}(i)^2} = e^{C_p \frac{2-p}{2} \|\bar{\mu}_{\delta}\|_{L^2(\mathbb{R}^d)}^2},$$

which is uniformly bounded for δ close to 0 (recall that $\bar{\mu}_{\delta} \rightarrow \mu_0$ in L^2). To bound the first factor in (5.12), we use the fact that $(\tilde{\zeta}_{\delta,i})_{i \in \mathbb{T}_{\delta}}$ are independent with zero mean to obtain

$$\mathbb{E}[|\Psi_{\delta}^{(\ell,N)}(\tilde{\zeta}_{\delta})|^2] = \sum_{\ell < |I| < N} \left(\prod_{i \in I} \mathbb{E}[\tilde{\zeta}_{\delta,i}^2] \right) \psi_{\delta}(I)^2 \leq \sum_{\ell < |I| < N} e^{C' \sum_{i \in \mathbb{T}_{\delta}} \mu_{\delta}(i)^2} (\sigma_{\delta}^2)^{|I|} \psi_{\delta}(I)^2, \quad (5.13)$$

where in the inequality we have applied (B.3), provided ζ satisfies condition (a) in Theorem 2.5. Combined with the assumption in Theorem 2.5 that (2.11) holds with $\varepsilon = 0$, and tracing back to (5.12), we then obtain the desired L^p bound on $\Psi_{\delta}^{>\ell}(\zeta_{\delta})$ in (5.9).

If we assume instead that condition (b) in Theorem 2.5 holds, then we can modify the calculation in (5.13) by applying the bound $\mathbb{E}[\tilde{\zeta}_{\delta,i}^2] \leq \sigma_{\delta}^2(1 + C\mu_{\delta}(i))$, stated in (B.1), to obtain

$$\mathbb{E}[|\Psi_{\delta}^{(\ell,N)}(\tilde{\zeta}_{\delta})|^2] \leq \sum_{\ell < |I| < N} e^{C\|\mu_{\delta}\|_{\infty}|I|} (\sigma_{\delta}^2)^{|I|} \psi_{\delta}(I)^2. \quad (5.14)$$

Theorem 2.5(b) and condition (2.11) with $\varepsilon = 0$ then give the desired bound in (5.9). \square

6. Proof for the disordered pinning model

In this section we prove Theorem 3.1. We recall that $\tau = (\tau_k)_{k \geq 0}$ is an aperiodic renewal process such that either $\mathbb{E}[\tau_1] < \infty$, or relation (3.2) holds for some $\alpha \in (1/2, 1)$. Note that

$$u(n) := \mathbb{P}(n \in \tau) \sim \begin{cases} \frac{1}{\mathbb{E}[\tau_1]} & \text{if } \mathbb{E}[\tau_1] < \infty, \\ \frac{C_{\alpha}}{L(n)n^{1-\alpha}} & \text{if } 1/2 < \alpha < 1 \text{ (where } C_{\alpha} := \alpha \sin(\pi\alpha)/\pi), \end{cases} \quad (6.1)$$

where the first asymptotic relation is the classical renewal theorem, while the second one is due to Doney [D97, Thm. B] (see also [G07, §A.5]). We also recall that $\omega = (\omega_n)_{n \in \mathbb{N}}$, representing the disorder, is an i.i.d. sequence of random variables satisfying (3.1).

Proof of Theorem 3.1. It suffices to rewrite the partition function as a polynomial chaos expansion and then to check that all the conditions of Theorem 2.3 are satisfied. We only consider the conditioned partition function $Z_{Nt, \beta_N, h_N}^{\omega, c}$, as the proof for the free one follows the same lines. We also set $t = 1$, to lighten notation.

Step 1. Consider the lattice $\mathbb{T}_N := \frac{1}{N}\mathbb{N}$ for $N \in \mathbb{N}$. Note that in Section 2 we used the notation \mathbb{T}_δ (where δ would equal $1/N$): here we prefer \mathbb{T}_N , as it indicates the size of the polymer. Consequently, all the quantities in this section will be indexed by N instead of δ . For each $t \in \mathbb{T}_N$ we define the cell $\mathcal{C}_N(t) := (t - 1/N, t]$, which has volume $v_N = 1/N$.

Step 2. We now rewrite the conditioned partition function $Z_{N, \beta, h}^{\omega, c}$ (cf. (3.4)), as a polynomial chaos expansion. This was already done in (1.5)–(1.7), in terms of the random variables $\varepsilon_i = e^{\beta\omega_i - \Lambda(\beta) + h} - 1$. It is actually convenient to rescale the ε_i so that their variance is of order one, in order to apply Theorem 2.3. Since $\text{Var}(\varepsilon_i) \sim \beta^2$ as $\beta \downarrow 0$, recalling (3.7) we set

$$a_N = \begin{cases} \frac{1}{\sqrt{N}} & \text{if } \mathbb{E}[\tau_1] < \infty, \\ \frac{L(N)}{N^{\alpha-1/2}} & \text{if } \frac{1}{2} < \alpha < 1, \end{cases} \quad (6.2)$$

so that $\beta_N = \hat{\beta}a_N$, and we define the random variables $\zeta_N = (\zeta_{N,t})_{t \in \mathbb{T}_N}$ by

$$\zeta_{N,t} := \frac{1}{a_N} (e^{\beta_N \omega_{Nt} - \Lambda(\beta_N) + h_N} - 1). \quad (6.3)$$

In this way, arguing as in (1.5)–(1.7), we can write

$$Z_{N, \beta_N, h_N}^{\omega, c} = \Psi_N(\zeta_N) := 1 + \sum_{k=1}^N \frac{1}{k!} \sum_{(t_1, \dots, t_k) \in (\mathbb{T}_N)^k} \psi_N^c(t_1, \dots, t_k) \prod_{i=1}^k \zeta_{N, t_i}, \quad (6.4)$$

where the kernel $\psi_N^c(t_1, \dots, t_k)$ is a symmetric function, which vanishes when $t_i = t_j$ for some $i \neq j$ or when some $t_i \notin (0, 1]$, and for $0 =: t_0 < t_1 < \dots < t_k \leq 1$ is defined by

$$\begin{aligned} \psi_N^c(t_1, \dots, t_k) &:= a_N^k \mathbb{P}(\{Nt_1, \dots, Nt_k\} \subseteq \tau \mid N \in \tau) \\ &= \frac{u(N(1-t_k))}{u(N)} \prod_{i=1}^k a_N u(N(t_i - t_{i-1})) \end{aligned} \quad (6.5)$$

(recall (6.1)). We extend $\psi_N^c(t_1, \dots, t_k)$ from $(\mathbb{T}_N)^k$ to \mathbb{R}^k in the usual way, as a piecewise constant function on products of cells. The same is done for $(s, t) \mapsto u(N(t-s))$.

Step 3. We now verify that the conditions of Theorem 2.3 are satisfied. By our assumptions (3.1) on the disorder, for every fixed $N \in \mathbb{N}$ the random variables $(\zeta_{N,t})_{t \in \mathbb{T}_N}$ are i.i.d. with mean and variance given by

$$\begin{aligned} \mu_N &:= \mathbb{E}(\zeta_{N,t}) = \frac{1}{\sigma_N} (e^{h_N} - 1) \sim \frac{\hat{h}}{\sqrt{N}}, \\ \sigma_N^2 &:= \text{Var}(\zeta_{N,t}) = \frac{1}{a_N^2} (e^{\Lambda(2\beta_N) - 2\Lambda(\beta_N)} - 1) e^{2h_N} \sim \frac{\beta_N^2}{a_N^2} \rightarrow \hat{\beta}^2, \quad \text{as } N \rightarrow \infty. \end{aligned} \quad (6.6)$$

Since $v_N = 1/N$, condition (i) of Theorem 2.3 is satisfied with $\sigma_0 = \hat{\beta}$ and $\mu_0(t) = \hat{h}\mathbb{1}_{(0,1]}(t)$. (More precisely, redefining $\zeta_{N,t}$ as $(\zeta_{N,t} - \mathbb{E}[\zeta_{N,t}])$ when $t \notin (0, 1]$ —which is harmless, since such values of t do not contribute to (6.4)—one has $\mu_N(t) = \mu_N\mathbb{1}_{(0,1]}(t) \rightarrow \mu_0(t)$ in $L^2(\mathbb{R})$.)

To prove that the random variables $((\zeta_{N,t} - \mu_N)^2)_{N \in \mathbb{N}, t \in \mathbb{T}_N}$ are uniformly integrable, we show that the moments $\mathbb{E}[(\zeta_{N,t} - \mu_N)^4]$ are uniformly bounded. Since $\Lambda(\beta) = O(\beta)$ as $\beta \downarrow 0$, by (3.1), for every $k \in \mathbb{N}$ we can estimate

$$\begin{aligned} & \mathbb{E}[(e^{\beta\omega - \Lambda(\beta)} - 1)^{2k}] \\ & \leq 2^{2k} e^{-2k\Lambda(\beta)} \mathbb{E}[(e^{\beta\omega} - 1)^{2k}] + 2^{2k} (1 - e^{-\Lambda(\beta)})^{2k} \\ & = 2^{2k} e^{-2k\Lambda(\beta)} \mathbb{E}\left[\left(\int_0^\beta \omega e^{t\omega} dt\right)^{2k}\right] + 2^{2k} (1 - e^{-\Lambda(\beta)})^{2k} \\ & \leq 2^{2k} e^{-2k\Lambda(\beta)} \beta^{2k} \int_0^\beta \mathbb{E}[\omega^{2k} e^{2kt\omega}] \frac{dt}{\beta} + 2^{2k} (1 - e^{-\Lambda(\beta)})^{2k} = O(\beta^{2k}). \end{aligned} \quad (6.7)$$

Recalling that $\beta_N = \hat{\beta}a_N$ and $h_N = o(1)$, we obtain the desired bound:

$$\mathbb{E}[(\zeta_{N,t} - \mu_N)^4] \leq \frac{e^{4h_N}}{a_N^4} O(\beta_N^4) = O(1).$$

Let us check condition (ii) of Theorem 2.3. The renewal estimates in (6.1) imply that, for fixed $0 < s < t$,

$$\lim_{N \rightarrow \infty} \sqrt{N} a_N u(N(t-s)) = \begin{cases} \frac{1}{\mathbb{E}[\tau_1]} & \text{if } \mathbb{E}[\tau_1] < \infty, \\ \frac{C_\alpha}{(t-s)^{1-\alpha}} & \text{if } 1/2 < \alpha < 1. \end{cases} \quad (6.8)$$

Recalling the definitions (6.5), (3.9) of the discrete and continuum kernels ψ_N^c, ψ_t^c (for $t = 1$), as well as the fact that $v_N = 1/N$, it follows that for every fixed $k \in \mathbb{N}$ the convergence

$$v_N^{-k/2} \psi_N^c(t_1, \dots, t_k) \xrightarrow{N \rightarrow \infty} \psi_1^c(t_1, \dots, t_k) \quad (6.9)$$

holds *pointwise*, for distinct points t_1, \dots, t_k . To obtain the required L^2 convergence, it suffices to exhibit an L^2 domination. The case $\mathbb{E}[\tau_1] < \infty$ is easy: by (6.1) there exists $A \in (0, \infty)$ such that $1/A \leq u(n) \leq A$ for every $n \in \mathbb{N}$, and since $v_N^{-1/2} a_N = 1$ it follows that

$$v_N^{-k/2} \psi_N^c(t_1, \dots, t_k) \leq A^{k+2} \mathbb{1}_{(0,1]^k}(t_1, \dots, t_k). \quad (6.10)$$

We now focus on the case $1/2 < \alpha < 1$. By Karamata's representation theorem for slowly varying functions [BGT87, Theorem 1.3.1], we can write $L(n) = c(n) \exp(\int_1^n \frac{\varepsilon(u)}{u} du)$ for some functions $c(x) \rightarrow c > 0$ and $\varepsilon(x) \rightarrow 0$ as $x \rightarrow \infty$. It follows that for any $\eta > 0$ there exists a constant $A' = A'_\eta \in (0, \infty)$ such that

$$\frac{1}{A'} \left(\frac{n}{m}\right)^{-\eta} \leq \frac{L(n)}{L(m)} \leq A' \left(\frac{n}{m}\right)^\eta, \quad \forall n, m \in \mathbb{N} \text{ with } m \leq n.$$

Recalling (6.1) and (6.2), for possibly a larger constant $A \in (0, \infty)$ we have

$$\frac{C_\alpha}{A(t-s)^{1-\alpha-\eta}} \leq \sqrt{N} a_N u(N(t-s)) \leq \frac{AC_\alpha}{(t-s)^{1-\alpha+\eta}}, \quad (6.11)$$

which plugged into (6.5) implies that for $0 < t_1 < \dots < t_k \leq 1$,

$$v_N^{-k/2} \psi_N^c(t_1, \dots, t_k) \leq \frac{A^{k+2} C_\alpha^k}{t_1^{1-\alpha'} \dots (t_k - t_{k-1})^{1-\alpha'} (1 - t_k)^{1-\alpha'}}, \quad (6.12)$$

where we set $\alpha' := \alpha - \eta$ for short. If we choose $\eta > 0$ sufficiently small, so that $\alpha' > 1/2$ (recall that $\alpha > 1/2$), we have obtained the required L^2 domination.

We finally check condition (iii) of Theorem 2.3, that is, relation (2.11). Since σ_N^2 is bounded (cf. (6.6)), we let $B \in (0, \infty)$ denote a constant such that $(1 + \varepsilon)\sigma_N^2 \leq B$, so that

$$\sum_{I \subseteq \mathbb{T}_N, |I| > \ell} (1 + \varepsilon)^{|I|} (\sigma_N^2)^{|I|} \psi_N^c(I)^2 \leq \sum_{k > \ell} B^k \sum_{\substack{(t_1, \dots, t_k) \in (\mathbb{T}_N)^k \\ 0 < t_1 < \dots < t_k \leq 1}} \psi_N^c(t_1, \dots, t_k)^2. \quad (6.13)$$

If $E[\tau_1] < \infty$, applying (6.10) and recalling that $v_N = 1/N$ we see that this expression is bounded by

$$\sum_{k > \ell} B^k A^{2(k+2)} (v_N)^k \frac{N^k}{k!} = \sum_{k > \ell} \frac{A^{2(k+2)} B^k}{k!},$$

which is arbitrarily small for ℓ large, proving (2.11). If $\alpha \in (1/2, 1)$ we apply (6.12): setting $\chi := 2(1 - \alpha') < 1$ for short and bounding sums by integrals, we estimate (6.13) by

$$\begin{aligned} \sum_{k > \ell} B^k A^{2(k+2)} C_\alpha^{2k} \int \dots \int_{0 < t_1 < \dots < t_k < 1} \frac{dt_1 \dots dt_k}{t_1^\chi \dots (t_k - t_{k-1})^\chi (1 - t_k)^\chi} \\ \leq \sum_{k > \ell} B^k A^{2(k+2)} C_\alpha^{2k} c_1 e^{-c_2 k \log k}, \end{aligned}$$

where for the last inequality we have applied Lemma B.3 below (we recall that $\chi < 1$). Again, the sum can be made arbitrarily small by choosing ℓ large, proving (2.11).

Step 4. Lastly, we prove formula (3.11) when $E[\tau_1] < \infty$. Since $\psi_t^c(t_1, \dots, t_k) = \left(\frac{1}{E[\tau_1]}\right)^k$ in this case, formula (3.8) for $\hat{h} = 0$ yields

$$\begin{aligned} \mathbf{Z}_{t, \hat{\beta}, 0}^{W, c} &= 1 + \sum_{k=1}^{\infty} \frac{1}{k!} \left(\frac{\hat{\beta}^k}{E[\tau_1]} \right)^k \int \dots \int_{[0, t]^k} W(dt_1) \dots W(dt_k) \\ &= e^{\frac{\hat{\beta}}{E[\tau_1]} W([0, t]) - \frac{1}{2} \left(\frac{\hat{\beta}}{E[\tau_1]} \right)^2 t}, \end{aligned} \quad (6.14)$$

where the second equality follows by [J97, Theorem 3.33 and Example 7.12]. This shows that (3.11) holds for $\hat{h} = 0$, because $W([0, t]) \sim \mathcal{N}(0, t)$. In the general case, we introduce the tilted law $\tilde{\mathbb{P}}$ defined by $d\tilde{\mathbb{P}}/d\mathbb{P} = \exp\{(\hat{h}/\hat{\beta})W([0, t]) - \frac{1}{2}(\hat{h}/\hat{\beta})^2 t\}$ and note that $\mathbf{Z}_{t, \hat{\beta}, \hat{h}}^{W, c}$ under \mathbb{P} has the same law as $\mathbf{Z}_{t, \hat{\beta}, 0}^{W, c}$ under $\tilde{\mathbb{P}}$ (cf. (2.15) and (2.16)). Since $W([0, t]) \sim \mathcal{N}((\hat{h}/\hat{\beta})t, t)$ under $\tilde{\mathbb{P}}$, formula (3.11) is proved also when $\hat{h} \neq 0$. \square

7. Proof for the directed polymer model

In this section we prove Theorem 3.8. We recall that $S = (S_n)_{n \geq 0}$ is a random walk on \mathbb{Z} satisfying Assumption 3.7. We denote by $p \in \mathbb{N}$ the period of the random walk, so that $P(S_1 \in p\mathbb{Z} + r) = 1$ for some $r \in \{0, \dots, p-1\}$. Introducing the lattice

$$\mathbb{T} := \{(n, k) \in \mathbb{Z}^2 : k \in p\mathbb{Z} + rn\}, \quad (7.1)$$

we have $P(S = (S_n)_{n \geq 0} \in \mathbb{T}) = 1$. Defining

$$q_n(k) := P(S_n = k), \quad \forall n \geq 0, k \in \mathbb{Z}, \quad (7.2)$$

Gnedenko's local limit theorem [BGT87, Theorem 8.4.1] yields

$$\sup_{k \in \mathbb{Z}, (n,k) \in \mathbb{T}} |n^{1/\alpha} q_n(k) - pg(k/n^{1/\alpha})| \xrightarrow{n \rightarrow \infty} 0, \quad (7.3)$$

where $g(\cdot)$ denotes the density of the stable law to which S is attracted. We also recall that $\omega = (\omega(n, k))_{n \in \mathbb{N}, k \in \mathbb{Z}}$ is an i.i.d. sequence of random variables satisfying (3.1).

Proof of Theorem 3.8. As in the proof of Theorem 3.1, the strategy is to rewrite the partition function as a polynomial chaos expansion and then to apply Theorem 2.3. We focus on the conditioned point-to-point partition function $Z_{Nt, \beta_N}^{\omega, c}(N^{1/\alpha}x)$, as the free one follows the same lines. For notational simplicity, we set $t = 1$.

Step 1. We introduce for $N \in \mathbb{N}$ the rescaled lattice

$$\mathbb{T}_N := \{(N^{-1}n, N^{-1/\alpha}k) : (n, k) \in \mathbb{T}\} \subseteq \mathbb{R}^2$$

(cf. (7.1)). Note that we use N instead of $\delta := 1/N$ as a subscript, as it indicates the ‘‘length’’ of the polymer. For each $(t, x) \in \mathbb{T}_N$, we define the cell $\mathcal{C}_N((t, x)) := (t - 1/N, t] \times (x - p/N^{1/\alpha}, x]$, which has volume $v_N = pN^{-(1+1/\alpha)}$.

Step 2. We rewrite the conditioned partition function $Z_{N, \beta_N}^{\omega, c}(N^{1/\alpha}x)$, defined in (3.21), as a polynomial chaos expansion, using the random variables $\zeta_N = (\zeta_N(s, y))_{(s,y) \in \mathbb{T}_N}$ given by

$$\zeta_N(s, y) := N^{(\alpha-1)/(2\alpha)} (e^{\beta_N \omega(Ns, N^{1/\alpha}y) - \Lambda(\beta_N)} - 1), \quad (7.4)$$

where the prefactor has been chosen so that $\text{Var}(\zeta_N(s, y)) = O(1)$ (see below). Arguing as in (1.5)–(1.7), we can write

$$Z_{N, \beta_N}^{\omega, c}(N^{1/\alpha}x) = \Psi_N(\zeta_N) := 1 + \sum_{k=1}^N \frac{1}{k!} \sum_{(z_1, \dots, z_k) \in (\mathbb{T}_N)^k} \psi_{N, (1, x)}^c(z_1, \dots, z_k) \prod_{i=1}^k \zeta_N(z_i),$$

where $\psi_{N, (1, x)}^c$ is a symmetric function of $(z_1, \dots, z_k) \in (\mathbb{T}_N)^k$, which vanishes when

$z_i = z_j$ for some $i \neq j$ and when some $z_i \notin (0, 1] \times \mathbb{R}$, and for distinct $z_1 = (t_1, x_1), \dots, z_k = (t_k, x_k)$, say with $0 < t_1 < \dots < t_k \leq 1$, it is defined by (recall (7.2))

$$\begin{aligned} \psi_{N,(1,x)}^c((t_1, x_1), \dots, (t_k, x_k)) &:= \frac{\mathbb{P}(S_{Nt_1} = N^{1/\alpha}x_1, \dots, S_{Nt_k} = N^{1/\alpha}x_k \mid S_N = x)}{(N^{(\alpha-1)/(2\alpha)})^k} \\ &= \frac{q_{N(1-t_k)}(N^{1/\alpha}(x - x_k))}{q_N(N^{1/\alpha}x)} \prod_{i=1}^k (N^{-(\alpha-1)/(2\alpha)} q_{N(t_i-t_{i-1})}(N^{1/\alpha}(x_i - x_{i-1}))), \end{aligned} \quad (7.5)$$

where $(t_0, x_0) := (0, 0)$. The kernel $\psi_{N,(1,x)}^c(z_1, \dots, z_k)$ is extended from $(\mathbb{T}_N)^k$ to $(\mathbb{R}^2)^k$ in the usual way, as a piecewise constant function which is constant on every product of cells. The function $((s, x), (t, y)) \mapsto q_{N(t-s)}(N^{1/\alpha}(y - x))$ is extended in the same way.

Step 3. We now check the assumptions of Theorem 2.3. Recalling (3.1) and the fact that $\beta_N = \hat{\beta}N^{-(\alpha-1)/(2\alpha)}$ (cf. Theorem 3.8), for every fixed $N \in \mathbb{N}$ the random variables $(\zeta_N(z))_{z \in \mathbb{T}_N}$ are i.i.d. with zero mean $\mu_N(z) \equiv 0$ and variance given by

$$\sigma_N^2 = \mathbb{V}\text{ar}(\zeta_N(z)) = N^{(\alpha-1)/\alpha} e^{\Lambda(2\beta_N) - 2\Lambda(\beta_N)} \sim N^{(\alpha-1)/\alpha} \beta_N^2 \xrightarrow{N \rightarrow \infty} \hat{\beta}^2. \quad (7.6)$$

Condition (i) of Theorem 2.3 is thus satisfied with $\mu_0(z) \equiv 0$ and $\sigma_0 = \hat{\beta}$. The uniform integrability of $(\zeta_N(z)^2)_{N \in \mathbb{N}, z \in \mathbb{T}_N}$ is easily checked as for the pinning model (cf. (6.7)).

Let us check condition (ii). Recalling the definition (3.24) of the function $g_t(\cdot)$, we observe that by (7.3), for fixed $0 < s < t$ and $x, y \in \mathbb{R}$,

$$\lim_{N \rightarrow \infty} N^{1/\alpha} q_{N(t-s)}(N^{1/\alpha}(y - x)) = pg_{t-s}(y - x).$$

Recalling the definition (3.26) of the continuum kernel $\psi_{t,x}^c$ (for $t = 1$), since $v_N = pN^{-1-1/\alpha}$, it follows by (7.5) that for every $k \in \mathbb{N}$ one has the *pointwise* convergence

$$\lim_{N \rightarrow \infty} v_N^{-k/2} \psi_{N,(1,x)}^c((t_1, x_1), \dots, (t_k, x_k)) = \psi_{1,x}^c((t_1, x_1), \dots, (t_k, x_k)). \quad (7.7)$$

We need to show that this convergence also holds in L^2 . Since the density $g(\cdot)$ is bounded, relation (7.3) implies that for some constant $A \in (0, \infty)$,

$$q_n(k) \leq An^{-1/\alpha}, \quad \forall n \in \mathbb{N}_0, k \in \mathbb{Z}. \quad (7.8)$$

For fixed $x \in \mathbb{R}$, one has $q_N(N^{1/\alpha}x) \geq \frac{p}{2}g(x)/N^{1/\alpha}$ for large N , again by (7.3), hence the prefactor in (7.5) is upper bounded as

$$\frac{q_{N(1-t_k)}(N^{1/\alpha}(x - x_k))}{q_N(N^{1/\alpha}x)} \leq \frac{C_x}{(1-t_k)^{1/\alpha}}, \quad \text{where } C_x := \frac{2A}{pg(x)}.$$

Applying (7.8) to each term in the product in (7.5) and recalling that $v_N = pN^{-1-1/\alpha}$, for $0 < t_1 < \dots < t_k \leq 1$ we get

$$\begin{aligned} [v_N^{-k/2} \psi_{N,(1,x)}^c((t_1, x_1), \dots, (t_k, x_k))]^2 &\leq \frac{C_x}{(1-t_k)^{1/\alpha}} \frac{q_{N(1-t_k)}(N^{1/\alpha}(x-x_k))}{q_N(N^{1/\alpha}x)} \\ &\quad \times \prod_{i=1}^k (v_N^{-1/2} N^{-(\alpha-1)/(2\alpha)})^2 \left(\frac{A}{N^{1/\alpha}(t_i-t_{i-1})^{1/\alpha}} \right) q_{N(t_i-t_{i-1})}(N^{1/\alpha}(x_i-x_{i-1})) \\ &= C'_{k,x} (N^{1/\alpha})^k \frac{\mathbb{P}(S_{Nt_1} = N^{1/\alpha}x_1, \dots, S_{Nt_k} = N^{1/\alpha}x_k \mid S_N = N^{1/\alpha}x)}{t_1^{1/\alpha} (t_2-t_1)^{1/\alpha} \dots (t_k-t_{k-1})^{1/\alpha} (1-t_k)^{1/\alpha}}, \end{aligned} \quad (7.9)$$

where we set $C'_{k,x} := A^k p^{-k} C_x$. A further application of (7.8) also yields

$$[v_N^{-k/2} \psi_{N,(1,x)}^c((t_1, x_1), \dots, (t_k, x_k))]^2 \leq \frac{A^k C'_{k,x}}{t_1^{2/\alpha} (t_2-t_1)^{2/\alpha} \dots (t_k-t_{k-1})^{2/\alpha} (1-t_k)^{2/\alpha}}. \quad (7.10)$$

We now decompose the domain $\{0 < t_1 < \dots < t_k < 1\} \times \mathbb{R}^k$ as $D_1 \cup D_2 \cup D_3$, where

$$\begin{aligned} D_1 &:= \bigcap_{i=1}^k \{t_i - t_{i-1} > \eta, |x_i| < M\} \cap \{1 - t_k > \eta\}, \\ D_2 &:= \bigcup_{i=1}^k \{t_i - t_{i-1} \leq \eta\} \cup \{1 - t_k \leq \eta\}, \quad D_3 := \bigcup_{i=1}^k \{|x_i| \geq M\}, \end{aligned}$$

for fixed $\eta, M \in (0, \infty)$. Inequality (7.10) shows that the rescaled kernel $v_N^{-k/2} \psi_{N,(1,x)}^c(\cdot)$ is uniformly bounded on the (bounded) set D_1 , hence the convergence (7.7) holds in L^2 on D_1 . If we show that the integrals of $[v_N^{-k/2} \psi_{N,(1,x)}^c(\cdot)]^2$ and of $[\psi_{1,x}^c(\cdot)]^2$ over the sets D_2 and D_3 can be made arbitrarily small for suitable η, M , we are done. Since $\psi_{1,x} \in L^2([0, 1]^k \times \mathbb{R}^k)$, by (3.24) there is nothing to prove for $\psi_{1,x}^c(\cdot)$ and we may focus on $\psi_{N,(1,x)}^c(\cdot)$. By (7.9),

$$\begin{aligned} &\int_{D_2} |v_N^{-k/2} \psi_{N,(1,x)}^c(\cdot)|^2 \\ &= \sum_{(t_1, x_1), \dots, (t_k, x_k) \in (\mathbb{T}_N)^k} v_N^k [v_N^{-k/2} \psi_{N,(1,x)}^c((t_1, x_1), \dots, (t_k, x_k))]^2 \\ &\leq C'_{k,x} p^k \frac{1}{N^k} \sum_{\substack{(t_1, \dots, t_k) \in \frac{1}{N} \mathbb{N} \\ 0 < t_1 < \dots < t_k \leq 1 \\ t_i - t_{i-1} \leq \eta \text{ for some } i}} \frac{1}{t_1^{1/\alpha} \dots (t_k - t_{k-1})^{1/\alpha} (1 - t_k)^{1/\alpha}}, \\ &\leq C'_{k,x} p^k \int_{\substack{0 < t_1 < \dots < t_k < 1 \\ t_i - t_{i-1} \leq \eta \text{ for some } i}} \frac{dt_1 \dots dt_k}{t_1^{1/\alpha} \dots (t_k - t_{k-1})^{1/\alpha} (1 - t_k)^{1/\alpha}}, \end{aligned} \quad (7.11)$$

which vanishes as $\eta \downarrow 0$ (recall that $\alpha > 1$). Analogously, using again (7.9), we get

$$\begin{aligned} \int_{D_3} |v_N^{-k/2} \psi_{N,(1,x)}^c(\cdot)|^2 &\leq C'_{k,x} p^k \left(\int \cdots \int_{0 < t_1 < \cdots < t_k < 1} \frac{dt_1 \cdots dt_k}{t_1^{1/\alpha} \cdots (t_k - t_{k-1})^{1/\alpha} (1 - t_k)^{1/\alpha}} \right) \\ &\times \mathbb{P} \left(\max_{0 \leq n \leq N} |S_n| \geq N^{1/\alpha} M \mid S_N = N^{1/\alpha} x \right). \end{aligned} \quad (7.12)$$

As $N \rightarrow \infty$, the last probability converges to $\mathbb{P}(\sup_{0 \leq t \leq 1} |X_t| \geq M \mid X_1 = x)$, where $X = (X_t)_{t \geq 0}$ is the stable Lévy process to which the random walk is attracted [Lig68], hence it can be made as small as one wishes, uniformly in $N \in \mathbb{N}$, by choosing M large. This completes the verification of condition (ii) in Theorem 2.3.

Finally, we check (iii). Since σ_N^2 is bounded (cf. (7.6)), we have $(1 + \varepsilon)\sigma_N^2 \leq B$ for some $B \in (0, \infty)$ (we can even set $\varepsilon = 0$, because $\mu_N \equiv 0$ in this case). Applying (7.11) for $\eta = 1$, i.e. with no restriction on $t_i - t_{i-1}$ (equivalently, (7.12) with $M = 0$), we obtain

$$\begin{aligned} \sum_{I \subset \mathbb{T}_N, |I| > \ell} (1 + \varepsilon)^{|I|} (\sigma_N^2)^{|I|} \psi_{N,(1,x)}^c(I)^2 &\leq \sum_{k > \ell} B^k \frac{1}{k!} \sum_{(z_1, \dots, z_k) \in (\mathbb{T}_N)^k} \psi_{N,(1,x)}^c(z_1, \dots, z_k)^2 \\ &= \sum_{k > \ell} B^k \frac{1}{k!} \|v_N^{-k/2} \psi_{N,(1,x)}^c\|_{L^2((\mathbb{R}^2)^k)}^2 \\ &\leq \sum_{k > \ell} \tilde{B}_x^k \int \cdots \int_{0 < t_1 < \cdots < t_k < 1} \frac{dt_1 \cdots dt_k}{t_1^{1/\alpha} \cdots (t_k - t_{k-1})^{1/\alpha} (1 - t_k)^{1/\alpha}} \leq \sum_{k > \ell} \tilde{B}_x^k c_1 e^{-c_2 k \log k}, \end{aligned}$$

where, recalling that $C'_{k,x} = A^k p^{-k} C_x$, we have set $\tilde{B}_x := BA\sqrt{p} \max\{1, C_x\}^2$, which is a finite constant for every fixed $x \in \mathbb{R}$, and we have applied Lemma B.3 for the last inequality. This shows that (2.11) holds, hence condition (iii) in Theorem 2.3 is verified. \square

8. Proof for the random field Ising model

In this section we prove Theorem 3.14 and Corollary 3.16. We recall that the disordered partition function $Z_{\Omega_\delta, \lambda_\delta, h_\delta}^{+, \omega_\delta}$ is defined as in (3.32), where $\Omega_\delta := \Omega \cap (\delta\mathbb{Z})^2$ (with $\Omega \subseteq \mathbb{R}^2$ being a fixed bounded, simply connected open set with piecewise smooth boundary) and where:

- $\mathbb{P}_{\Omega_\delta}^+$ (with expectation $\mathbb{E}_{\Omega_\delta}^+$) denotes the critical Ising model on \mathbb{Z}^2 , defined as in (3.30);
- $\omega_\delta = (\omega_{\delta,x})_{x \in \Omega_\delta}$ is an i.i.d. family of random variables satisfying (3.1);
- $\lambda_\delta = (\lambda_{\delta,x})_{x \in \Omega_\delta}$, $h_\delta = (h_{\delta,x})_{x \in \Omega_\delta}$ are defined by

$$\lambda_{\delta,x} := \hat{\lambda}(x)\delta^{7/8}, \quad h_{\delta,x} := \hat{h}(x)\delta^{15/8} \quad (8.1)$$

(cf. (3.35)), where $\hat{\lambda} : \bar{\Omega} \rightarrow (0, \infty)$ and $\hat{h} : \bar{\Omega} \rightarrow \mathbb{R}$ are fixed continuous functions.

The heart of our proof are pointwise and L^2 estimates for the critical Ising correlation functions, in particular near the diagonals (see Lemmas 8.1–8.3 below). Complementary L^1 estimates have recently been established in [CGN12a, Prop. 3.9].

Proof of Theorem 3.14. We are going to apply Theorem 2.3 with $v_\delta = \delta^2$, rewriting the partition function in terms of a polynomial chaos expansion.

Step 1. By (3.32), we can write

$$Z_{\Omega_\delta, \lambda_\delta, h_\delta}^{+, \omega_\delta} = \mathbb{E}_{\Omega_\delta}^+ \left[\prod_{x \in \Omega_\delta} (\cosh(\xi_{\delta, x}) + \sigma_x \sinh(\xi_{\delta, x})) \right], \quad \text{where} \quad (8.2)$$

$$\xi_{\delta, x} := \lambda_{\delta, x} \omega_{\delta, x} + h_{\delta, x}.$$

Recall the notation $\alpha^I := \prod_{x \in I} \alpha_x$. A binomial expansion of the product yields

$$\begin{aligned} e^{-\frac{1}{2} \|\hat{\lambda}\|_{L^2(\Omega)}^2} Z_{\Omega_\delta, \lambda_\delta, h_\delta}^{+, \omega_\delta} &= e^{-\frac{1}{2} \|\hat{\lambda}\|_{L^2(\Omega)}^2} \delta^{-1/4} \sum_{I \subseteq \Omega_\delta} \cosh(\xi_{\delta, \cdot})^{\Omega_\delta \setminus I} \mathbb{E}_{\Omega_\delta}^+ [\sigma^I] \sinh(\xi_{\delta, \cdot})^I \\ &= e^{-\frac{1}{2} \|\hat{\lambda}\|_{L^2(\Omega)}^2} \delta^{-1/4} \cosh(\xi_{\delta, \cdot})^{\Omega_\delta} \sum_{I \subseteq \Omega_\delta} \mathbb{E}_{\Omega_\delta}^+ [\sigma^I] \tanh(\xi_{\delta, \cdot})^I. \end{aligned} \quad (8.3)$$

We first show that the prefactor before the sum converges to 1 in probability as $\delta \downarrow 0$.

Recalling the definition (8.2) of $\xi_{\delta, x}$ and the fact that $\omega_{\delta, x}$ have zero mean, unit variance and locally finite exponential moments (cf. (3.1)), a Taylor expansion yields

$$\mathbb{E}[\log \cosh(\xi_{\delta, x})] = \frac{\lambda_{\delta, x}^2}{2} + O(h_{\delta, x}^2 + \lambda_{\delta, x}^4) = \frac{\hat{\lambda}(x)^2}{2} \delta^{7/4} + O(\delta^{7/2}), \quad (8.4)$$

where the term $O(\delta^{7/2})$ is uniform over $x \in \Omega_\delta$, by the continuity of $\hat{\lambda}, \hat{h}$. Therefore, as $\delta \downarrow 0$,

$$\sum_{x \in \Omega_\delta} \mathbb{E}[\log \cosh(\xi_{\delta, x})] = \frac{1}{2} \|\hat{\lambda}\|_{L^2(\Omega)}^2 \delta^{-1/4} + o(1), \quad (8.5)$$

and the prefactor in (8.3) can be rewritten as

$$\exp \left\{ \sum_{x \in \Omega_\delta} (\log \cosh(\xi_{\delta, x}) - \mathbb{E}[\log \cosh(\xi_{\delta, x})]) \right\} (1 + o(1)).$$

The sum is over $|\Omega_\delta| = O(\delta^{-2})$ i.i.d. centered random variables, hence it converges to zero in probability provided $\text{Var}[\log \cosh(\xi_{\delta, x})] = o(\delta^2)$. This is checked by a Taylor expansion:

$$\text{Var}(\log \cosh(\xi_{\delta, x})) \leq \mathbb{E}[(\log \cosh(\xi_{\delta, x}))^2] = O(\lambda_{\delta, x}^4) = O(\delta^{7/2}) = o(\delta^2).$$

Step 2. It remains to verify that the sum in (8.3) converges to the desired Wiener chaos expansion, namely (3.37). Defining the family $\zeta_\delta = (\zeta_{\delta, x})_{x \in \Omega_\delta}$ by

$$\zeta_{\delta, x} := \frac{\tanh(\xi_{\delta, x})}{\text{Var}(\tanh(\xi_{\delta, x}))^{1/2}},$$

the sum in (8.3) can be written as a polynomial chaos expansion $\Psi_\delta(\zeta_\delta) := \sum_{I \subseteq \Omega_\delta} \psi_\delta(I) \zeta_{\delta, \cdot}^I$, where

$$\psi_\delta(I) := \text{Var}(\tanh(\xi_{\delta, \cdot}))^{|I|/2} \mathbb{E}_{\Omega_\delta}^+ [\sigma^I], \quad I \subseteq \Omega_\delta. \quad (8.6)$$

We are thus left with checking that the conditions in Theorem 2.3 are satisfied.

By a Taylor expansion, as $\delta \downarrow 0$ one has

$$\begin{aligned}\mathbb{E}[\tanh(\xi_{\delta,x})] &= h_{\delta,x} + O(h_{\delta,x}^3 + \lambda_{\delta,x}^3) = \hat{h}(x)\delta^{15/8} + O(\delta^{21/8}), \\ \mathbb{E}[\tanh(\xi_{\delta,x})^2] &= \lambda_{\delta,x}^2 + O(h_{\delta,x}^2 + \lambda_{\delta,x}^4) = \hat{\lambda}(x)^2\delta^{7/4} + O(\delta^{7/2}),\end{aligned}\tag{8.7}$$

where the $O(\cdot)$ terms are uniform in x , by the continuity of $\hat{\lambda}, \hat{h}$. Therefore, uniformly in x ,

$$\mu_\delta(x) := \mathbb{E}[\zeta_{\delta,x}] = \frac{\hat{h}(x)}{\hat{\lambda}(x)}\delta + o(\delta), \quad \sigma_\delta^2 := \mathbb{V}\text{ar}[\zeta_{\delta,x}] = 1.$$

Recalling that $v_\delta = \delta^2$ and $\hat{\lambda} : \overline{\Omega} \rightarrow (0, \infty)$ is continuous (hence uniformly bounded away from zero), condition (i) of Theorem 2.3 is satisfied with $\sigma_0 = 1$ and $\mu_0(x) := \hat{h}(x)/\hat{\lambda}(x)$.

The uniform integrability of $((\zeta_{\delta,x} - \mu_\delta(x))^2)_{\delta \in (0,1), x \in \Omega_\delta}$ holds because the moments $\mathbb{E}[(\zeta_{\delta,x} - \mu_\delta(x))^4]$ are uniformly bounded, as for the disordered pinning model (cf. (6.7)).

Step 3. It remains to check conditions (ii) and (iii) of Theorem 2.3. By (8.7),

$$\mathbb{V}\text{ar}(\tanh(\xi_{\delta,x})) = \hat{\lambda}(x)^2\delta^{7/4} + O(\delta^{7/2}),$$

hence for fixed $I = \{x_1, \dots, x_n\} \subseteq \Omega$, by (8.6),

$$\psi_\delta(I) = \hat{\lambda}(\cdot)^{|I|}\delta^{7|I|/8}\mathbb{E}_{\Omega_\delta}^+[\sigma^I] = \hat{\lambda}(x_1) \cdots \hat{\lambda}(x_n)\delta^{7n/8}\mathbb{E}_{\Omega_\delta}^+[\sigma_{x_1} \cdots \sigma_{x_n}].$$

As $v_\delta = \delta^2$, relation (3.33), recently proved by Chelkak, Hongler and Izyurov [CHI12], shows immediately that for every $n \in \mathbb{N}$ and distinct $x_1, \dots, x_n \in \Omega$,

$$\begin{aligned}\lim_{\delta \downarrow 0} v_\delta^{-n/2} \psi_\delta(\{x_1, \dots, x_n\}) &= \hat{\lambda}(x_1) \cdots \hat{\lambda}(x_n) \mathcal{C}^n \phi_\Omega^+(x_1, \dots, x_n) \\ &=: \psi_0(\{x_1, \dots, x_n\}).\end{aligned}\tag{8.8}$$

(Incidentally, since $\sigma_0 = 1$ and $\mu_0(x) := \hat{h}(x)/\hat{\lambda}(x)$, the Wiener chaos expansion (2.12) of Theorem 2.3 matches the expansion (3.37) of Theorem 3.14.)

To extend the pointwise convergence (8.8) to L^2 convergence, we need uniform bounds on $\psi_\delta(I)$. By Lemma 8.1 below, we have the following bound uniformly in $\delta \in (0, 1)$:

$$\begin{aligned}v_\delta^{-|I|/2} \psi_\delta(I) &\leq (C \|\hat{\lambda}\|_\infty)^{|I|} \prod_{i=1}^{|I|} \frac{1}{d(x_i, \partial\Omega \cup I \setminus \{x_i\})^{1/8}} \\ &=: (C \|\hat{\lambda}\|_\infty)^{|I|} f_\Omega(x_1, \dots, x_{|I|}),\end{aligned}\tag{8.9}$$

where by Lemma 8.3 below, given $|I| = n$ for any $n \in \mathbb{N}$,

$$\frac{1}{n!} \|f_\Omega\|_{L^2(\Omega^n)}^2 \leq C^n (n!)^{-3/4}.\tag{8.10}$$

Combined with (8.8), this shows that conditions (ii) and (iii) of Theorem 2.3 are satisfied, which completes the proof of Theorem 3.14. \square

We next state and prove the lemmas needed to establish (8.9) and (8.10).

Lemma 8.1. *Let Ω , Ω_δ for $\delta > 0$, and $E_{\Omega_\delta}^+$ be as introduced at the beginning of this section. Then there exists $C = C(\Omega) \in (0, \infty)$ such that for any $I = \{x_1, \dots, x_n\} \subseteq \Omega$ with $|I| = n$,*

$$0 \leq E_{\Omega_\delta}^+[\sigma_{x_1} \cdots \sigma_{x_n}] \leq C^n \delta^{n/8} \prod_{i=1}^n \frac{1}{d(x_i, \partial\Omega \cup I \setminus \{x_i\})^{1/8}}, \quad (8.11)$$

where for any $x \in \Omega$ we define $\sigma_x := \sigma_{x_\delta}$ with x_δ being the point in Ω_δ closest to x , and we set $d(x, A) := \inf_{y \in A} \|x - y\|$.

Proof. If $B(x; r)$ denotes the ball of radius r centered at x , and $\mathbb{B}(x; r)_\delta := B(x; r) \cap (\delta\mathbb{Z})^2$, then (3.33) with $\Omega = B(0; 1)$, $n = 1$ and $x_1 = 0$ implies that for some $C \in (0, \infty)$,

$$E_{\mathbb{B}(0;1)_\delta}^+[\sigma_0] \leq C \delta^{1/8} \quad \text{for all } \delta \in (0, 1). \quad (8.12)$$

Then, for any $x \in \Omega$, by imposing + boundary condition on the ball $B(x; r)$ with radius $r := d(x, \partial\Omega)$ and applying the FKG inequality [G06, Chapter 2], we obtain

$$E_{\Omega_\delta}^+[\sigma_x] \leq E_{\mathbb{B}(x;r)_\delta}^+[\sigma_x] = E_{\mathbb{B}(0;r)_\delta}^+[\sigma_0] = E_{\mathbb{B}(0;1)_{\delta/r}}^+[\sigma_0] \leq C \frac{\delta^{1/8}}{r^{1/8}} = \frac{C \delta^{1/8}}{d(x, \partial\Omega)^{1/8}}, \quad (8.13)$$

where in the last inequality we applied (8.12).

Inequality (8.11) follows by applying (8.13) and Lemma 8.2 below, choosing Ω_i therein to be $\Omega_i \cap (\delta\mathbb{Z})^2$, where Ω_i is the ball centered at x_i with radius $\frac{1}{4}d(x_i, \partial\Omega \cup I \setminus \{x_i\})$. \square

Lemma 8.2. *Let $x_1, \dots, x_n \in \Omega \subseteq \mathbb{Z}^2$ and suppose $x_i \in \Omega_i \subseteq \Omega$ with $\Omega_i \cap (\Omega_j \cup \partial\Omega_j) = \emptyset$ for all $i \neq j$. Then*

$$0 \leq E_{\Omega}^+[\sigma_{x_1} \cdots \sigma_{x_n}] \leq \prod_{i=1}^n E_{\Omega_i}^+[\sigma_i]. \quad (8.14)$$

Proof. Relation (8.14) is a consequence of the Griffiths–Kelly–Sherman (GKS) inequalities (see e.g. [E06, Chapter V.3]). We recall that P_{Ω}^+ denotes the Ising measure on $\{\pm 1\}^{\Omega}$ with inverse temperature $\beta \in (0, \infty)$ and zero external field (cf. (3.30)). (The fact that $\beta = \beta_c$ is immaterial for this proof, and we could even include a positive external field in P_{Ω}^+ .) Given $h = (h_x)_{x \in \Omega}$, let $P_{\Omega, h}^{\text{free}}$ denote the Ising measure with external (site-dependent) field h , i.e.

$$P_{\Omega, h}^{\text{free}}(\sigma) = \frac{1}{Z_{\Omega, \beta, h}^{\text{free}}} \exp\left\{ \sum_{x \sim y \in \Omega} \beta \sigma_x \sigma_y + \sum_{x \in \Omega} h_x \sigma_x \right\}.$$

Since $P_{\Omega}^+ = P_{\Omega, h^+}^{\text{free}}$ with the choice $h_x^+ := \beta|\{y \in \partial\Omega : y \sim x\}|$, we may focus on $P_{\Omega, h}^{\text{free}}$.

Let $I := \{x_1, \dots, x_n\}$ and $\sigma_I := \sigma_{x_1} \cdots \sigma_{x_n}$. If $h \geq 0$ (that is, $h_x \geq 0$ for all $x \in \Omega$), then

$$E_{\Omega, h}^{\text{free}}[\sigma_I] \geq 0,$$

by the first GKS inequality, proving the first bound in (8.14). Always for $h \geq 0$,

$$\frac{\partial E_{\Omega, h}^{\text{free}}[\sigma_I]}{\partial h_y} = E_{\Omega, h}^{\text{free}}[\sigma_I \sigma_y] - E_{\Omega, h}^{\text{free}}[\sigma_I] E_{\Omega, h}^{\text{free}}[\sigma_y] \geq 0, \quad \forall y \in \Omega,$$

by the second GKS inequality. Therefore $E_{\Omega, h}^{\text{free}}[\sigma_I]$ is increasing in h_y for every $y \in \Omega$. If we start with $h = h^+$ and increase h_y to ∞ for each $y \in \bigcup_{i=1}^n \partial\Omega_i$, the resulting Ising measure is equivalent to imposing $+$ boundary condition on $\bigcup_{i=1}^n \partial\Omega_i$. Under this limiting measure, the distribution of the spin configurations on the disjoint subdomains $(\Omega_i)_{1 \leq i \leq n}$ factorizes, leading to the second bound in (8.14). \square

Lemma 8.3. *For any $n \in \mathbb{N}$ and distinct $x_1, \dots, x_n \in \Omega$, set $I := \{x_1, \dots, x_n\}$ and define*

$$f_{\Omega}(x_1, \dots, x_n) := \prod_{i=1}^n \frac{1}{d(x_i, \partial\Omega \cup I \setminus \{x_i\})^{1/8}}. \quad (8.15)$$

Then there exists $C = C(\Omega) < \infty$ such that for all $n \in \mathbb{N}$,

$$\|f_{\Omega}\|_{L^2(\Omega^n)}^2 \leq C^n (n!)^{1/4}. \quad (8.16)$$

Proof. To prove (8.16), it suffices to show that for all $n \in \mathbb{N}$,

$$\|f_{\Omega}\|_{L^2(\Omega^n)}^2 \leq C n^{1/4} \|f_{\Omega}\|_{L^2(\Omega^{n-1})}^2, \quad (8.17)$$

where $\|f_{\Omega}\|_{L^2(\Omega^0)}^2 := 1$.

Note that in

$$\|f_{\Omega}\|_{L^2(\Omega^n)}^2 = \int \cdots \int_{\Omega^n} \prod_{i=1}^n \frac{1}{d(x_i, \partial\Omega \cup I \setminus \{x_i\})^{1/4}} dx_1 \cdots dx_n, \quad (8.18)$$

we can divide the domain of integration Ω for x_n into disjoint open sets $\Omega_0, \dots, \Omega_{n-1}$ (modulo a set of measure 0), such that

$$\begin{aligned} x_n \in \Omega_0 & \quad \text{if and only if} \quad d(x_n, \partial\Omega \cup I \setminus \{x_n\}) = d(x_n, \partial\Omega), \\ x_n \in \Omega_i & \quad \text{if and only if} \quad d(x_n, \partial\Omega \cup I \setminus \{x_n\}) = d(x_n, x_i), \quad 1 \leq i \leq n-1. \end{aligned} \quad (8.19)$$

We next bound $f_{\Omega}(x_1, \dots, x_n)$ in terms of $f_{\Omega}(x_1, \dots, x_{n-1})$. First consider the case $x_n \in \Omega_0$. For each $1 \leq i \leq n-1$, either

$$d(x_i, \partial\Omega \cup I \setminus \{x_i\}) = d(x_i, \partial\Omega \cup I' \setminus \{x_i\}), \quad (8.20)$$

where $I' := \{x_1, \dots, x_{n-1}\}$, or

$$d(x_i, \partial\Omega \cup I \setminus \{x_i\}) = d(x_i, x_n). \quad (8.21)$$

In the latter case, by the triangle inequality and the assumption $x_n \in \Omega_0$, we find that

$$d(x_i, \partial\Omega) \leq d(x_i, x_n) + d(x_n, \partial\Omega) \leq 2d(x_i, x_n), \quad (8.22)$$

and hence

$$\frac{1}{d(x_i, \partial\Omega \cup I \setminus \{x_i\})} = \frac{1}{d(x_i, x_n)} \leq \frac{2}{d(x_i, \partial\Omega)} \leq \frac{2}{d(x_i, \partial\Omega \cup I' \setminus \{x_i\})}. \quad (8.23)$$

If N_0 denotes the number of points among $\{x_1, \dots, x_{n-1}\}$ such that (8.21) holds, then

$$f_\Omega(x_1, \dots, x_n)^2 \leq 2^{N_0/4} f_\Omega(x_1, \dots, x_{n-1})^2 \frac{1}{d(x_n, \partial\Omega)^{1/4}}. \quad (8.24)$$

We claim that $N_0 \leq 6$, which would then imply

$$\int_{\Omega^{n-1} \times \Omega_0} \dots \int f_\Omega(x_1, \dots, x_n)^2 dx_1 \dots dx_n \leq 2^{3/2} \|f_\Omega\|_{L^2(\Omega^{n-1})}^2 \int_\Omega \frac{dx_n}{d(x_n, \partial\Omega)^{1/4}}, \quad (8.25)$$

where the last integral is bounded by some constant $C_3(\Omega) < \infty$, because Ω is assumed to be a bounded simply connected domain with a piecewise smooth boundary.

To verify the claim that $N_0 \leq 6$, assume without loss of generality that x_1, \dots, x_k are the points which satisfy (8.21). In particular, $d(x_i, x_n) \leq d(x_i, x_j)$ for all $1 \leq i \neq j \leq k$. We may shift the origin to x_n and assume without loss of generality that $x_i \in \mathbb{R}^2$ has polar coordinates (r_i, ϑ_i) , and the directional vectors $e^{i\vartheta_1}, \dots, e^{i\vartheta_k}$ are ordered counter clockwise on the unit circle. For any two adjacent $e^{i\vartheta_j}$ and $e^{i\vartheta_{j+1}}$ on the unit circle, in order to satisfy

$$\max\{d(x_j, 0), d(x_{j+1}, 0)\} \leq d(x_j, x_{j+1}),$$

it is necessary that the angle between $e^{i\vartheta_j}$ and $e^{i\vartheta_{j+1}}$ be at least $\pi/3$. It then follows that there can be at most six such points.

We now consider the case $x_n \in \Omega_i$ for $1 \leq i \leq n-1$ (recall (8.19)). Without loss of generality, assume that $x_n \in \Omega_1$. By the same reasoning as above, for each $1 \leq i \leq n-1$, either (8.20) or (8.21) holds; in the latter case we can replace (8.22) by

$$d(x_i, x_1) \leq d(x_i, x_n) + d(x_n, x_1) \leq 2d(x_i, x_n),$$

because $x_n \in \Omega_1$. Thus for $i \geq 2$ (8.23) still holds if we replace $d(x_i, \partial\Omega)$ by $d(x_i, x_1)$ therein. The case $i = 1$ needs to be dealt with separately: for this we simply bound

$$\begin{aligned} \frac{1}{d(x_1, \partial\Omega \cup I \setminus \{x_1\})^{1/4}} &\leq \frac{1}{d(x_1, \partial\Omega \cup I' \setminus \{x_1\})^{1/4}} + \frac{1}{d(x_1, x_n)^{1/4}} \\ &= \frac{1}{d(x_1, \partial\Omega \cup I' \setminus \{x_1\})^{1/4}} \left(1 + \frac{d(x_1, \partial\Omega \cup I' \setminus \{x_1\})^{1/4}}{d(x_1, x_n)^{1/4}} \right). \end{aligned}$$

We thus obtain the following analogue of (8.24) (with $N_0 \leq 6$) when $x_n \in \Omega_1$:

$$\begin{aligned} f_\Omega(x_1, \dots, x_n)^2 &\leq 2^{3/2} f_\Omega(x_1, \dots, x_{n-1})^2 \left\{ \left(1 + \frac{d(x_1, \partial\Omega \cup I' \setminus \{x_1\})^{1/4}}{d(x_1, x_n)^{1/4}} \right) \frac{1}{d(x_n, x_1)^{1/4}} \right\}. \end{aligned}$$

Bounding the term in brackets by $C_4(\Omega)/d(x_1, x_n)^{1/2}$ for some $C_4(\Omega) < \infty$ (recall that Ω is a bounded set), we obtain

$$\begin{aligned} \int_{\Omega^{n-1} \times \Omega_1} \cdots \int f_{\Omega}(x_1, \dots, x_n)^2 dx_1 \cdots dx_n \\ \leq 2^{3/2} C_4(\Omega) \int_{\Omega^{n-1} \times \Omega_1} \cdots \int \frac{f_{\Omega}(x_1, \dots, x_{n-1})^2}{d(x_n, x_1)^{1/2}} dx_1 \cdots dx_n \\ \leq C_5(\Omega) |\Omega_1|^{3/4} \|f_{\Omega}\|_{L^2(\Omega^{n-1})}^2 \end{aligned} \quad (8.26)$$

for some $C_5(\Omega) < \infty$, where we have applied the Hardy–Littlewood rearrangement inequality (see e.g. [LL01, Theorem 3.4]) to bound

$$\int_{\Omega_1} \frac{dx_n}{d(x_n, x_1)^{1/2}} \leq \int_{\Omega_1^*} \frac{dx_n}{d(x_n, 0)^{1/2}} = \int_0^{r^*} \frac{2\pi r}{r^{1/2}} dr = \frac{4}{3} \pi^{1/4} |\Omega_1|^{3/4},$$

where Ω_1^* is the ball centered at the origin with the same area $|\Omega_1^*| = \pi r_*^2$ as Ω_1 .

Combining (8.25) and (8.26), and the analogue for $x_n \in \Omega_i$ with $2 \leq i \leq n-1$, we obtain

$$\begin{aligned} \|f_{\Omega}\|_{L^2(\Omega^n)}^2 &\leq C_6(\Omega) (1 + |\Omega_1|^{3/4} + \cdots + |\Omega_{n-1}|^{3/4}) \|f_{\Omega}\|_{L^2(\Omega^{n-1})}^2 \\ &\leq C_6(\Omega) \left(1 + (n-1) \left(\frac{|\Omega_1| + \cdots + |\Omega_{n-1}|}{n-1} \right)^{3/4} \right) \|f_{\Omega}\|_{L^2(\Omega^{n-1})}^2 \\ &\leq C_2(\Omega) n^{1/4} \|f_{\Omega}\|_{L^2(\Omega^{n-1})}^2, \end{aligned} \quad (8.27)$$

where we have applied Jensen's inequality to the function $g(x) = x^{3/4}$. This establishes (8.17) and concludes the proof of Theorem 3.14. \square

Proof of Corollary 3.16. Let $\varphi'(\cdot)$ denote the complex derivative of the conformal map $\varphi : \tilde{\Omega} \rightarrow \Omega$. Since $|\varphi'(z)|^2$ equals the Jacobian determinant of φ , for all $f, g \in L^2(\Omega)$ we have

$$\int_{\Omega} f(x)g(x) dx = \int_{\tilde{\Omega}} f(\varphi(z))g(\varphi(z))|\varphi'(z)|^2 dz, \quad (8.28)$$

by the change of variables formula. As a consequence, if $W(\cdot)$ denotes a white noise on \mathbb{R}^2 , the processes $(\int_{\Omega} f(x) W(dx))_{f \in L^2(\Omega)}$ and $(\int_{\tilde{\Omega}} f(\varphi(z))|\varphi'(z)| W(dz))_{f \in L^2(\Omega)}$ have the same distribution: they are both centered Gaussian processes with the same covariance (8.28). This extends to an equality in distribution for multiple integrals (recall Subsection 2.1):

$$\begin{aligned} \int \cdots \int_{\Omega^n} f(x_1, \dots, x_n) \prod_{i=1}^n W(dx_i) \\ \stackrel{d}{=} \int \cdots \int_{\tilde{\Omega}^n} f(\varphi(z_1), \dots, \varphi(z_n)) \prod_{i=1}^n [|\varphi'(z_i)| W(dz_i)], \end{aligned}$$

jointly for $n \in \mathbb{N}$ and symmetric $f \in L^2(\Omega^n)$. Informally, we have $W(d\varphi(z)) \stackrel{d}{=} |\varphi'(z)|W(dz)$, which is the stochastic analogue of $d\varphi(z) = |\varphi'(z)|^2 dz$. Recalling (3.37), it follows that

$$\mathbf{Z}_{\Omega, \hat{\lambda}, \hat{h}}^{+, W} \stackrel{d}{=} 1 + \sum_{n=1}^{\infty} \frac{C^n}{n!} \int \cdots \int_{\tilde{\Omega}^n} \phi_{\Omega}^+(\varphi(z_1), \dots, \varphi(z_n)) \prod_{i=1}^n [\hat{\lambda}(\varphi(z_i)) |\varphi'(z_i)| W(dz_i) + \hat{h}(\varphi(z_i)) |\varphi'(z_i)|^2 dz_i]. \quad (8.29)$$

By [CHI12, Theorem 1.3], the function ϕ_{Ω}^+ is conformally covariant with

$$\phi_{\Omega}^+(\varphi(z_1), \dots, \varphi(z_n)) = \phi_{\tilde{\Omega}}^+(z_1, \dots, z_n) \prod_{i=1}^n |\varphi'(z_i)|^{-1/8}, \quad (8.30)$$

hence $\mathbf{Z}_{\Omega, \hat{\lambda}, \hat{h}}^{+, W} \stackrel{d}{=} \mathbf{Z}_{\tilde{\Omega}, \tilde{\lambda}, \tilde{h}}^{+, W}$ with $\tilde{\lambda}(z) := |\varphi'(z)|^{7/8} \hat{\lambda}(\varphi(z))$ and $\tilde{h}(z) := |\varphi'(z)|^{15/8} \hat{h}(\varphi(z))$. \square

Appendix A. The Cameron–Martin shift

Recalling Subsection 2.1, let $W = (W(f))_{f \in L^2(\mathbb{R}^d)}$ be a white noise on \mathbb{R}^d defined on the probability space $(\Omega_W, \mathcal{A}, \mathbb{P})$. We denote by $L^0 := L^0(\Omega_W, \sigma(W), \mathbb{P})$ the space of (equivalence classes of) a.s. finite random variables that are measurable with respect to the σ -algebra generated by W , equipped with the topology of convergence in probability. Note that all the multi-dimensional stochastic integrals $W^{\otimes k}(f)$ belong to L^0 .

Let us now fix $\mathbf{v} \in L^2(\mathbb{R}^d)$, representing the *bias*. Given $k \in \mathbb{N}$ and a symmetric square-integrable function $f : (\mathbb{R}^d)^k \rightarrow \mathbb{R}$, the ‘‘biased stochastic integral’’

$$W_{\mathbf{v}}^{\otimes k}(f) = \int \cdots \int_{(\mathbb{R}^d)^k} f(x_1, \dots, x_k) \prod_{i=1}^k (W(dx_i) + \mathbf{v}(x_i) dx_i) \quad (A.1)$$

was defined in Remark 2.4 by expanding the product and integrating out the deterministic variables corresponding to $\mathbf{v}(x_i) dx_i$, thus reducing to a sum of lower-dimensional ordinary (unbiased) stochastic integrals. In particular, for $k = 1$ we can write $W_{\mathbf{v}}(f) = W_{\mathbf{v}}^{\otimes 1}(f)$ as

$$W_{\mathbf{v}}(f) := W(f) + \int_{\mathbb{R}^d} f(x) \mathbf{v}(x) dx = W(f) + \mathbb{E}[W(f) \xi] \quad \text{with } \xi := W(\mathbf{v}), \quad (A.2)$$

by the Itô isometry (2.2).

Thus $W_{\mathbf{v}}(f) = \varrho_{\xi}(W(f))$, where we define the map $\varrho_{\xi}(X) := X + \mathbb{E}[X \xi]$ for every one-dimensional stochastic integral X . By [J97, Theorem 14.1], such a map admits a unique extension $\varrho_{\xi} : L^0 \rightarrow L^0$, called the *Cameron–Martin shift*, which is continuous, linear and satisfies

$$\varrho_{\xi}(1) = 1, \quad \varrho_{\xi}(XY) = \varrho_{\xi}(X) \varrho_{\xi}(Y) \quad \forall X, Y \in L^0. \quad (A.3)$$

As a consequence, the multi-dimensional biased stochastic integrals (A.1) correspond to

$$W_{\mathbf{v}}^{\otimes k}(f) = \varrho_{\xi}(W^{\otimes k}(f)). \quad (A.4)$$

This is easily checked for “special simple functions” f (recall Subsection 2.1) using (A.2)–(A.3), and is extended to general symmetric $f \in L^2((\mathbb{R}^d)^k)$ using the continuity of ϱ_ξ .

For any $X \in L^0$, the random variable $\varrho_\xi(X)$ has the same distribution as X under the probability \mathbb{P}_ν defined in (2.15), by [J97, Theorem 14.1(iii)–(iv)]. In particular, choosing for X the series in (2.13) whenever it converges in probability, one obtains relation (2.16).

Appendix B. Technical lemmas

The following lemma is used in the proof of Theorem 2.5.

Lemma B.1 (Exponential tilting). *Let $(\zeta_{\delta,i})_{\delta \in (0,1), i \in \mathbb{T}_\delta}$ be a family of independent random variables in L^2 with mean $\mu_\delta(i)$ and variance σ_δ^2 , with $\sigma_\delta \rightarrow \sigma_0 \in (0, \infty)$ and $\|\mu_\delta\|_\infty := \sup_{i \in \mathbb{T}_\delta} |\mu_\delta(i)| \rightarrow 0$ as $\delta \downarrow 0$. Assume further that $(\zeta_{\delta,i}^2)_{\delta \in (0,1), i \in \mathbb{T}_\delta}$ are uniformly integrable. Then one can construct independent random variables $(\tilde{\zeta}_{\delta,i})_{\delta \in (0,1), i \in \mathbb{T}_\delta}$ such that*

$$\mathbb{P}(\tilde{\zeta}_{\delta,i} \in dx) = f_{\delta,i}(x) \mathbb{P}(\zeta_{\delta,i} \in dx),$$

and there exist $\delta_0, C \in (0, \infty)$ and $C_p \in (0, \infty)$ for every $p \in \mathbb{R}$ such that

$$\mathbb{E}[\tilde{\zeta}_{\delta,i}] = 0, \quad \mathbb{E}[\tilde{\zeta}_{\delta,i}^2] \leq \sigma_\delta^2(1 + C|\mu_\delta(i)|), \quad \mathbb{E}[f_{\delta,i}(\zeta_{\delta,i})^p] \leq 1 + C_p \mu_\delta(i)^2, \quad (\text{B.1})$$

for all $\delta \in (0, \delta_0)$ and $i \in \mathbb{T}_\delta$. Furthermore, if

$$\inf_{\delta \in (0,1), i \in \mathbb{T}_\delta} \min \{ \mathbb{P}(\zeta_{\delta,i} > 0), \mathbb{P}(\zeta_{\delta,i} < 0), \mathbb{V}\text{ar}(\zeta_{\delta,i} \mid \zeta_{\delta,i} > 0), \mathbb{V}\text{ar}(\zeta_{\delta,i} \mid \zeta_{\delta,i} < 0) \} > 0, \quad (\text{B.2})$$

then there exists $C' \in (0, \infty)$ such that the following improved bound holds:

$$\mathbb{E}[\tilde{\zeta}_{\delta,i}^2] \leq \sigma_\delta^2(1 + C' \mu_\delta(i)^2). \quad (\text{B.3})$$

The proof of Lemma B.1 is an easy corollary of the following general result, which concerns exponential tilting of a single random variable in order to shift its mean to zero (since the random variables are not assumed to have finite exponential moments, the tilting is performed on a bounded subset). The assumptions in Lemma B.1 guarantee that conditions (B.5) and (B.9) are fulfilled, and the constants in (2.15), (B.7), (B.10) are uniformly bounded.

Theorem B.2. *Let X be a square-integrable random variable and let $A > 0$ be such that*

$$\mathbb{E}[X^2 \mathbb{1}_{\{|X| > A\}}] \leq \frac{1}{4} \mathbb{E}[X^2]. \quad (\text{B.4})$$

Assume that $\mathbb{E}[X]$ is sufficiently small, more precisely,

$$|\mathbb{E}[X]| \leq \varepsilon := \frac{\mathbb{E}[X^2]^2}{144A^3}. \quad (\text{B.5})$$

Then one can define a random variable \tilde{X} such that $\mathbb{P}(\tilde{X} \in dx) = f(x)\mathbb{P}(X \in dx)$, satisfying

$$\mathbb{E}[f(X)^p] \leq 1 + C_p \mathbb{E}[X]^2 \quad \forall p \in \mathbb{R} \quad \text{with } C_p := \frac{4e^{|p|}}{A\varepsilon}, \quad (\text{B.6})$$

$$\mathbb{E}[\tilde{X}] = 0, \quad \mathbb{E}[\tilde{X}^2] \leq \mathbb{E}[X^2] + C|\mathbb{E}[X]| \quad \text{with } C := \frac{A^{3/2}}{\sqrt{\varepsilon}}. \quad (\text{B.7})$$

If $\mathbb{E}[X] \geq 0$, and A is chosen such that

$$\mathbb{E}[X^2 \mathbb{1}_{\{X > A\}}] \leq \frac{1}{4} \mathbb{E}[X^2 \mathbb{1}_{\{X \geq 0\}}], \quad (\text{B.8})$$

(replace X by $-X$ if $\mathbb{E}[X] \leq 0$), and further assume that

$$|\mathbb{E}[X \mid X \geq 0]| \leq \varepsilon' := \frac{\mathbb{E}[X^2 \mid X \geq 0]^2}{144A^3}, \quad (\text{B.9})$$

then we can define \tilde{X} such that (B.6) holds with ε replaced by ε' , while (B.7) is improved to

$$\mathbb{E}[\tilde{X}] = 0, \quad \mathbb{E}[\tilde{X}^2] \leq \mathbb{E}[X^2] + C' \mathbb{E}[X]^2 \quad \text{with } C' := \frac{A}{2\mathbb{P}(X \geq 0)\varepsilon'}. \quad (\text{B.10})$$

Proof. Without loss of generality, we assume that $\mathbb{E}[X] \geq 0$ (otherwise consider $-X$).

Step 1 (Strategy). We will fix $I \subseteq \mathbb{R}$, which is either $[-A, A]$ (assuming (B.4)–(B.5)), or $[0, A]$ (assuming (B.8)–(B.9)), and we define random variables Y, Z with laws

$$\mathbb{P}(Y \in \cdot) := \mathbb{P}(X \in \cdot \mid X \in I), \quad \mathbb{P}(Z \in \cdot) := \mathbb{P}(X \in \cdot \mid X \notin I). \quad (\text{B.11})$$

Taking independent copies of X, Y, Z , we have the following equality in distribution:

$$X \stackrel{\text{dist}}{=} \mathbb{1}_{\{X \in I\}} Y + \mathbb{1}_{\{X \notin I\}} Z \quad (\text{note that } |Y| \leq A). \quad (\text{B.12})$$

We then exponentially tilt Y , defining for $\lambda \in \mathbb{R}$ a random variable Y_λ with law

$$\mathbb{P}(Y_\lambda \in dx) := e^{\lambda x - F(\lambda)} \mathbb{P}(Y \in dx), \quad \text{where } F(\lambda) := \log \mathbb{E}[e^{\lambda Y}]. \quad (\text{B.13})$$

As we will show at the end of the proof, we can choose $\lambda = \tilde{\lambda} \in \mathbb{R}$ such that

$$\mathbb{E}[Y_{\tilde{\lambda}}] = -\frac{\mathbb{P}(X \notin I)}{\mathbb{P}(X \in I)} \mathbb{E}[Z]. \quad (\text{B.14})$$

If we define \tilde{X} replacing Y by $Y_{\tilde{\lambda}}$ in the definition (B.12), that is,

$$\tilde{X} := \mathbb{1}_{\{X \in I\}} Y_{\tilde{\lambda}} + \mathbb{1}_{\{X \notin I\}} Z, \quad (\text{B.15})$$

then $\mathbb{E}[\tilde{X}] = 0$ by construction. Moreover $\mathbb{P}(\tilde{X} \in dx) = f(x)\mathbb{P}(X \in dx)$ with density

$$f(x) = e^{\tilde{\lambda}x - F(\tilde{\lambda})} \mathbb{1}_{\{x \in I\}} + \mathbb{1}_{\{x \in \mathbb{R} \setminus I\}} = 1 + (e^{\tilde{\lambda}x - F(\tilde{\lambda})} - 1) \mathbb{1}_{\{x \in I\}}. \quad (\text{B.16})$$

The rest of the proof is devoted to estimating $\mathbb{E}[\tilde{X}^2]$ and $\mathbb{E}[f(X)^p]$. We are going to use the following bounds on $\tilde{\lambda}$, which will be proved at the very end:

$$I := [-A, A], \text{ assuming (B.4)–(B.5): } |\tilde{\lambda}| \leq \frac{|\mathbb{E}[X]|}{2A^{3/2}\sqrt{\varepsilon}}; \quad (\text{B.17})$$

$$I := [0, A], \text{ assuming (B.8)–(B.9): } |\tilde{\lambda}| \leq \frac{|\mathbb{E}[X]|}{\sqrt{6}A^{3/2}\sqrt{\varepsilon'}\sqrt{\mathbb{P}(X \in I)\mathbb{P}(X \geq 0)}}; \quad (\text{B.18})$$

$$\text{In either case: } |\tilde{\lambda}| \leq \frac{1}{27A}. \quad (\text{B.19})$$

Step 2 (Bounds on $\mathbb{E}[\tilde{X}^2]$). Denote $G(\lambda) := \mathbb{E}[Y_\lambda^2] = \mathbb{E}[Y^2 e^{\lambda Y}] / \mathbb{E}[e^{\lambda Y}]$. Recalling (B.15) and (B.12), we can write

$$\begin{aligned} \mathbb{E}[\tilde{X}^2] &= \mathbb{E}[X^2] + \mathbb{P}(X \in I)(\mathbb{E}[Y_\lambda^2] - \mathbb{E}[Y^2]) \\ &= \mathbb{E}[X^2] + \mathbb{P}(X \in I) \int_0^{\tilde{\lambda}} G'(\lambda) d\lambda. \end{aligned} \quad (\text{B.20})$$

Since $G'(\lambda) = \mathbb{E}[(Y_\lambda)^3] - \mathbb{E}[(Y_\lambda)^2]\mathbb{E}[Y_\lambda]$ and $|Y_\lambda| \leq A$, we have $|G'(\lambda)| \leq 2A^3$, and hence

$$\mathbb{E}[\tilde{X}^2] \leq \mathbb{E}[X^2] + 2A^3\mathbb{P}(X \in I)|\tilde{\lambda}|. \quad (\text{B.21})$$

Applying (B.17), we obtain precisely the second bound in (B.7).

To prove (B.10), assume (B.8)–(B.9) and set $I := [0, A]$. By (B.12)–(B.14),

$$\mathbb{E}[Y_\lambda] = \mathbb{E}[Y] - \frac{\mathbb{E}[X]}{\mathbb{P}(X \in I)}, \quad (\text{B.22})$$

and hence $\mathbb{E}[Y_\lambda] \leq \mathbb{E}[Y]$. Since $\mathbb{E}[Y_\lambda] = F'(\lambda)$ is increasing in λ (because $F''(\lambda) = \text{Var}[Y_\lambda] \geq 0$), it follows that $\tilde{\lambda} \leq 0$. We then refine (B.20) as follows:

$$\mathbb{E}[\tilde{X}^2] = \mathbb{E}[X^2] + \mathbb{P}(X \in I)\tilde{\lambda}G'(0) + \mathbb{P}(X \in I) \int_0^{\tilde{\lambda}} \left(\int_0^\lambda G''(s) ds \right) d\lambda. \quad (\text{B.23})$$

Note that $G'(0) = \mathbb{E}[Y^3] - \mathbb{E}[Y^2]\mathbb{E}[Y] \geq 0$, because $Y \in [0, A]$ and hence Y^2 and Y are positively correlated. Therefore the second term in (B.23) is bounded by 0. Also note that

$$G''(\lambda) = \mathbb{E}[(Y_\lambda)^4] - 2\mathbb{E}[(Y_\lambda)^3]\mathbb{E}[Y_\lambda] + 2\mathbb{E}[(Y_\lambda)^2]\mathbb{E}[Y_\lambda]^2 - \mathbb{E}[(Y_\lambda)^2]^2,$$

and hence $|G''(\lambda)| \leq 6A^4$. Substituting into (B.23) then yields

$$\mathbb{E}[\tilde{X}^2] \leq \mathbb{E}[X^2] + 3A^4\mathbb{P}(X \in I)\tilde{\lambda}^2. \quad (\text{B.24})$$

Applying (B.18), we obtain precisely (B.10).

Step 3 (Bounds on $\mathbb{E}[f(X)^p]$). Recall f from (B.16) and $F(\tilde{\lambda})$ from (B.13). Since $F(0) = 0$ and $|F'(\tilde{\lambda})| = |\mathbb{E}[Y_{\tilde{\lambda}}]| \leq A$ (cf. (B.13)), we have $|F(\tilde{\lambda})| \leq A|\tilde{\lambda}|$ and hence $|\tilde{\lambda}x - F(\tilde{\lambda})| \leq 2A|\tilde{\lambda}|$ for every $x \in I \subseteq [-A, A]$. Applying (B.19), we obtain

$$e^{-2/27} \leq f(x) \leq e^{2/27}, \quad \forall x \in \mathbb{R}. \quad (\text{B.25})$$

For any $p \in \mathbb{R}$, Taylor expansion gives $y^p \leq 1 + p(y - 1) + C'_p(y - 1)^2$ for all $y \in [e^{-2/27}, e^{2/27}]$ with

$$C'_p := \max_{y \in [e^{-2/27}, e^{2/27}]} |p(p-1)y^{p-2}| = |p(p-1)|e^{\frac{2}{27}|p-2|} \leq 2e^{|p|}. \quad (\text{B.26})$$

Therefore

$$\begin{aligned} f(x)^p &\leq 1 + p(f(x) - 1) + C'_p(f(x) - 1)^2 \\ &= 1 + (p - 2C'_p)(f(x) - 1) + C'_p(e^{2\tilde{\lambda}x - 2F(\tilde{\lambda})} - 1)\mathbb{1}_{\{x \in I\}}. \end{aligned}$$

Since f is a probability density, recalling the definition of F from (B.13), we obtain

$$\mathbb{E}[f(X)^p] \leq 1 + C'_p \mathbb{P}(X \in I)(e^{F(2\tilde{\lambda}) - 2F(\tilde{\lambda})} - 1). \quad (\text{B.27})$$

Since $|F''(\lambda)| = |\mathbb{E}[Y_{\lambda}^2] - \mathbb{E}[Y_{\lambda}]^2| \leq 2A^2$, the Mean Value Theorem and (B.19) yield

$$0 \leq F(2\tilde{\lambda}) - 2F(\tilde{\lambda}) = (F(2\tilde{\lambda}) - F(\tilde{\lambda})) - (F(\tilde{\lambda}) - F(0)) \leq 4A^2\tilde{\lambda}^2 \leq \frac{4}{27^2} \leq 1, \quad (\text{B.28})$$

where the first inequality holds by convexity of F (note that $F''(\lambda) = \text{Var}[Y_{\lambda}] \geq 0$).

Consider first the case $I = [-A, A]$, assuming (B.4)–(B.5): since $e^x - 1 \leq 2x$ for $0 \leq x \leq 1$, applying (B.28), (B.17) and (B.26) we obtain

$$\mathbb{E}[f(X)^p] \leq 1 + 2C'_p(F(2\tilde{\lambda}) - 2F(\tilde{\lambda})) \leq 1 + C'_p 8A^2\tilde{\lambda}^2 \leq 1 + \frac{4e^{|p|}}{A\varepsilon} \mathbb{E}[X]^2, \quad (\text{B.29})$$

proving (B.6).

The case $I = [0, A]$, assuming (B.8)–(B.9), is similar: we keep the term $\mathbb{P}(X \in I)$ in (B.27) when writing (B.29), so that applying (B.18) gives

$$\mathbb{E}[f(X)^p] \leq 1 + C'_p \mathbb{P}(X \in I) 8A^2\tilde{\lambda}^2 \leq 1 + \frac{16e^{|p|}}{6A\varepsilon'} \mathbb{E}[X]^2 \leq 1 + \frac{4e^{|p|}}{A\varepsilon'} \mathbb{E}[X]^2,$$

which coincides with (B.6), where ε is replaced by ε' .

Step 4 (Bounds on $\tilde{\lambda}$). We finally show that one can choose $\lambda = \tilde{\lambda}$ so that (B.14) holds and the bounds (B.17)–(B.19) are satisfied.

Since $F'(\lambda) = \mathbb{E}[Y_{\lambda}]$ (cf. (B.13)), we can rewrite (B.14) (cf. (B.22)), as

$$F'(\tilde{\lambda}) - F'(0) = \tilde{x}, \quad \text{where} \quad \tilde{x} := -\frac{\mathbb{E}[X]}{\mathbb{P}(X \in I)}. \quad (\text{B.30})$$

Since $F'''(\lambda) = \mathbb{E}[Y_\lambda^3] - 3\mathbb{E}[Y_\lambda]\mathbb{E}[Y_\lambda^2] + 2\mathbb{E}[Y_\lambda]^3$ and $|Y_\lambda| \leq A$, we have $|F'''(\lambda)| \leq 6A^3$. Therefore

$$F''(\lambda) \geq F''(0) - 6A^3|\lambda| \geq \frac{F''(0)}{2} \quad \text{for } |\lambda| \leq c := \frac{F''(0)}{12A^3} = \frac{\text{Var}(Y)}{12A^3}.$$

In particular, equation (B.30) has exactly one solution $\tilde{\lambda} \in [-c, c]$ provided

$$|\tilde{\lambda}| \leq \frac{F''(0)}{2}c, \quad \text{that is,} \quad |\mathbb{E}[X]| \leq \frac{\mathbb{P}(X \in I) \text{Var}(Y)^2}{24A^3}, \quad (\text{B.31})$$

in which case $\tilde{\lambda}$ satisfies

$$|\tilde{\lambda}| \leq \frac{|\tilde{\lambda}|}{\frac{1}{2}F''(0)} = \frac{2|\mathbb{E}[X]|}{\mathbb{P}(X \in I) \text{Var}(Y)}. \quad (\text{B.32})$$

It only remains to check that condition (B.31) is indeed satisfied, under either assumptions (B.4)–(B.5) or (B.8)–(B.9), and to show that (B.32) yields the bounds (B.17)–(B.19). For this we need to estimate $\mathbb{P}(X \in I)$ and $\text{Var}(Y)$.

In order to avoid repetitions, let \mathbb{P}^* denote the original law \mathbb{P} when we assume (B.4)–(B.5) or the conditional law $\mathbb{P}(\cdot | X \geq 0)$ when we assume (B.8)–(B.9). In either case $I = [-A, A]$ or $I = [0, A]$ we can write $\mathbb{P}(Y \in \cdot) := \mathbb{P}(X \in \cdot | X \in I) = \mathbb{P}^*(X \in \cdot | |X| \leq A)$, therefore

$$\text{Var}(Y) = \text{Var}^*(X | |X| \leq A). \quad (\text{B.33})$$

Note that assumptions (B.4)–(B.5) and (B.8)–(B.9) can be written as follows:

$$\mathbb{E}^*[X^2 \mathbb{1}_{\{|X| > A\}}] \leq \frac{1}{4}\mathbb{E}^*[X^2], \quad |\mathbb{E}^*(X)| \leq \frac{\mathbb{E}^*[X^2]^2}{144A^3}. \quad (\text{B.34})$$

Since $\mathbb{E}^*[X^2 \mathbb{1}_{\{|X| \leq A\}}] \leq A^2 \mathbb{P}^*(|X| \leq A)$, it follows that

$$A^2 \geq \frac{3}{4} \frac{\mathbb{E}^*[X^2]}{\mathbb{P}^*(|X| \leq A)} \geq \frac{3}{4} \mathbb{E}^*[X^2]. \quad (\text{B.35})$$

We thus get

$$\begin{aligned} \mathbb{P}^*(|X| \leq A) &= 1 - \mathbb{P}^*(|X| > A) \geq 1 - \frac{\mathbb{E}^*[X^2 \mathbb{1}_{\{|X| > A\}}]}{A^2} \geq 1 - \frac{\mathbb{E}^*[X^2]}{4A^2} \geq \frac{2}{3}, \\ |\mathbb{E}^*[X \mathbb{1}_{\{|X| > A\}}]| &\leq \frac{\mathbb{E}^*[X^2 \mathbb{1}_{\{|X| > A\}}]}{A} \leq \frac{\mathbb{E}^*[X^2]}{4A} \leq \frac{\sqrt{\mathbb{E}^*[X^2]}}{2\sqrt{3}}. \end{aligned} \quad (\text{B.36})$$

Together with (B.34) and (B.35), this gives

$$\begin{aligned} |\mathbb{E}^*[X \mathbb{1}_{\{|X| \leq A\}}]| &\leq |\mathbb{E}^*[X]| + |\mathbb{E}^*[X \mathbb{1}_{\{|X| > A\}}]| \leq \left(\frac{(4/3)^{3/2}}{144} + \frac{1}{2\sqrt{3}} \right) \sqrt{\mathbb{E}^*[X^2]} \\ &\leq \frac{1}{3} \sqrt{\mathbb{E}^*[X^2]}, \end{aligned}$$

which yields

$$|\mathbb{E}^*[X \mid |X| \leq A]| = \frac{|\mathbb{E}^*[X \mathbb{1}_{\{|X| \leq A\}}]|}{\mathbb{P}^*(|X| \leq A)} \leq \frac{1}{2} \sqrt{\mathbb{E}^*[X^2]}.$$

Applying one more time (B.34) we get

$$\mathbb{E}^*[X^2 \mid |X| \leq A] = \frac{|\mathbb{E}^*[X^2 \mathbb{1}_{\{|X| \leq A\}}]|}{\mathbb{P}^*(|X| \leq A)} \geq \frac{3}{4} \mathbb{E}^*[X^2],$$

which finally yields (cf. (B.33))

$$\text{Var}(Y) = \mathbb{E}^*[X^2 \mid |X| \leq A] - \mathbb{E}^*[X \mid |X| \leq A]^2 \geq \frac{1}{2} \mathbb{E}^*[X^2]. \quad (\text{B.37})$$

By (B.5) and (B.9), and applying (B.37) and (B.36), we obtain

$$|\mathbb{E}^*[X]| \leq \frac{\mathbb{E}^*[X^2]^2}{144A^3} \leq \frac{\text{Var}(Y)^2}{36A^3} \leq \frac{\mathbb{P}^*(|X| \leq A) \text{Var}(Y)^2}{24A^3}. \quad (\text{B.38})$$

Consider first the case $I = [-A, A]$, assuming (B.4)–(B.5): since $\mathbb{P}^*(|X| \leq A) = \mathbb{P}(X \in I)$, inequality (B.38) coincides precisely with the condition (B.31) to be checked. Next we consider the case $I = [0, A]$, assuming (B.8)–(B.9), where we recall that $\mathbb{P}^*(\cdot) = \mathbb{P}(\cdot \mid X \geq 0)$. By assumption $\mathbb{E}[X] \geq 0$, we have $|\mathbb{E}[X]| \leq |\mathbb{E}[X \mathbb{1}_{\{X \geq 0\}}]| = \mathbb{P}(X \geq 0) |\mathbb{E}^*[X]|$. Since we can write $\mathbb{P}^*(|X| \leq A) = \mathbb{P}(X \in I) / \mathbb{P}(X \geq 0)$, (B.38) again yields (B.31).

To conclude, for $I = [-A, A]$, applying (B.36) and (B.37) to (B.32) and recalling the definition of ε in (B.5) gives (B.17). For $I = [0, A]$, the bound (B.18) follows similarly, recalling the definition of ε' in (B.9) and observing that $\mathbb{P}(X \in I) \geq \frac{2}{3} \mathbb{P}(X \geq 0)$ by (B.36). Finally, to obtain (B.19) from (B.17)–(B.18), apply (B.35) and the assumptions (B.4), (B.8). \square

Finally, we prove the following bound on iterated integrals.

Lemma B.3. *Let $\chi \in [0, 1)$. Then there exist $c_1, c_2 > 0$ such that for all $k \in \mathbb{N}$,*

$$\int \cdots \int_{0 < t_1 < \cdots < t_k < 1} \frac{dt_1 \cdots dt_k}{t_1^\chi \cdots (t_k - t_{k-1})^\chi} \leq c_1 e^{-c_2 k \log k}, \quad (\text{B.39})$$

$$\int \cdots \int_{0 < t_1 < \cdots < t_k < 1} \frac{dt_1 \cdots dt_k}{t_1^\chi \cdots (t_k - t_{k-1})^\chi (1 - t_k)^\chi} \leq c_1 e^{-c_2 k \log k}. \quad (\text{B.40})$$

Proof. It is enough to prove (B.40), since the integral therein bounds (B.39). Recognizing the density of the Dirichlet distribution (with parameters $k + 1$ and $1 - \chi$) allows us to evaluate

$$\int \cdots \int_{0 < t_1 < \cdots < t_k < 1} \frac{dt_1 \cdots dt_k}{t_1^\chi \cdots (t_k - t_{k-1})^\chi (1 - t_k)^\chi} = \frac{\Gamma(1 - \chi)^{k+1}}{\Gamma((k + 1)(1 - \chi))},$$

and (B.40) follows by standard properties of the gamma function $\Gamma(\cdot)$. \square

Acknowledgments. We are grateful to Tom Spencer for fruitful discussions and we thank the referee for helpful comments. We also thank Niccolò Torri and Ran Wei for spotting some mistakes in the draft. We acknowledge support of NUS grant R-146-000-148-112 (R.S.), a Marie Curie International Reintegration Grant within the 7th European Community Framework Programme IRG-246809 and EPSRC grant EP/L012154/1 (N.Z.), and ERC Advanced Grant 267356 VARIS (F.C.). We also thank for the kind hospitality of the Institute of Mathematics at the University of Leiden, of Academia Sinica in Taipei and of the Department of Mathematics at Hokkaido University in Sapporo, where parts of this work were completed.

References

- [AW90] Aizenman, M., Wehr, J.: Rounding effects of quenched randomness on first-order phase transitions. *Comm. Math. Phys.* **130**, 489–528 (1990) [Zbl 0714.60090](#) [MR 1060388](#)
- [AKQ14a] Alberts, T., Khanin, K., Quastel, J.: The intermediate disorder regime for directed polymers in dimension $1 + 1$. *Ann. Probab.* **42**, 1212–1256 (2014) [Zbl 1292.82014](#) [MR 3189070](#)
- [AKQ14b] Alberts, T., Khanin, K., Quastel, J.: The continuum directed random polymer. *J. Statist. Phys.* **154**, 305–326 (2014) [Zbl 1291.82143](#) [MR 3162542](#)
- [A08] Alexander, K. S.: The effect of disorder on polymer depinning transitions. *Comm. Math. Phys.* **279**, 117–146 (2008) [Zbl 1175.82034](#) [MR 2377630](#)
- [AZ09] Alexander, K. S., Zygouras, N.: Quenched and annealed critical points in polymer pinning models. *Comm. Math. Phys.* **291**, 659–689 (2009) [Zbl 1188.82154](#) [MR 2534789](#)
- [ACQ11] Amir, G., Corwin, I., Quastel, J.: Probability distribution of the free energy of the continuum directed random polymer in $1+1$ dimensions. *Comm. Pure Appl. Math.* **64**, 466–537 (2011) [Zbl 1222.82070](#) [MR 2796514](#)
- [BC⁺14] Berger, Q., Caravenna, F., Poisat, J., Sun, R., Zygouras, N.: The critical curve of the random pinning and copolymer models at weak coupling. *Comm. Math. Phys.* **326**, 507–530 (2014) [MR 3165465](#)
- [BGT87] Bingham, N. H., Goldie, C. M., Teugels, J. L.: *Regular Variation*. *Encyclopedia Math. Appl.* **27**, Cambridge Univ. Press, Cambridge (1987) [Zbl 0667.26003](#) [MR 1015093](#)
- [BdH97] Bolthausen, E., den Hollander, F.: Localization transition for a polymer near an interface. *Ann. Probab.* **25**, 1334–1366 (1997) [Zbl 0885.60022](#) [MR 1457622](#)
- [BK92] Bourgain, J., Kahn, J., Kalai, G., Katznelson, Y., Linial, N.: The influence of variables in product spaces. *Israel J. Math.* **77**, 55–64 (1992) [Zbl 0771.60002](#) [MR 1194785](#)
- [B06] Bovier, A.: *Statistical Mechanics of Disordered Systems: a Mathematical Perspective*. Cambridge Univ. Press (2006) [Zbl 1108.82002](#) [MR 2252929](#)
- [BK88] Bricmont, J., Kupiainen, A.: Phase transition in the 3d random field Ising model. *Comm. Math. Phys.* **116**, 539–572 (1988) [MR 0943702](#)
- [CGN12a] Camia, F., Garban, C., Newman, C. M.: Planar Ising magnetization field I. Uniqueness of the critical scaling limit. *Ann. Probab.* **43**, 528–571 (2015) [Zbl 1332.82012](#) [MR 3305999](#)
- [CGN12b] Camia, F., Garban, C., Newman, C. M.: The Ising magnetization exponent on \mathbb{Z}^2 is $1/15$. *Probab. Theory Related Fields* **160**, 175–187 (2014) [Zbl 1307.82004](#) [MR 3256812](#)
- [CGN13] Camia, F., Garban, C., Newman, C. M.: Planar Ising magnetization field II. Properties of the critical and near-critical scaling limits. *Ann. Inst. H. Poincaré Probab. Statist.* **52**, 146–161 (2016) [Zbl 1338.82009](#) [MR 3449298](#)

- [CG10] Caravenna, F., Giacomini, G.: The weak coupling limit of disordered copolymer models. *Ann. Probab.* **38**, 2322–2378 (2010) [Zbl 1242.82022](#) [MR 2683632](#)
- [CSZ14] Caravenna, F., Sun, R., Zygouras, N.: The continuum disordered pinning model. *Probab. Theory Related Fields* **164**, 17–59 (2016) [Zbl 1341.82040](#) [MR 3449385](#)
- [Car88] Cardy, J. (ed.): *Finite-Size Scaling*. North-Holland (1988) [MR 1103125](#)
- [C06] Chatterjee, S.: A generalization of the Lindeberg principle. *Ann. Probab.* **34**, 2061–2076 (2006) [Zbl 1117.60034](#) [MR 2294976](#)
- [CD13] Chatterjee, S., Dey, P. S.: Multiple phase transitions in long-range first-passage percolation on square lattices. *Comm. Pure Appl. Math.* **69**, 203–256 (2016) [Zbl 1332.60133](#) [MR 3434612](#)
- [CCFS86] Chayes, J. T., Chayes, L., Fischer, D. S., Spencer, T.: Finite-size scaling and correlation lengths for disordered systems. *Phys. Rev. Lett.* **57**, 2999–3002 (1986) [MR 0925751](#)
- [CCFS89] Chayes, J. T., Chayes, L., Fischer, D. S., Spencer, T.: Correlation length bounds for disordered Ising ferromagnets. *Comm. Math. Phys.* **120**, 501–523 (1989) [Zbl 0658.60137](#) [MR 0981216](#)
- [CHI12] Chelkak, D., Hongler, C., Izyurov, C.: Conformal invariance of spin correlations in the planar Ising model. *Ann. of Math.* **181**, 1087–1138 (2015) [Zbl 1318.82006](#) [MR 3296821](#)
- [CSY04] Comets, F., Shiga, T., Yoshida, N.: Probabilistic analysis of directed polymers in a random environment: a review. In: *Stochastic Analysis on Large Scale Interacting Systems*, *Adv. Stud. Pure Math.* 39, Math. Soc. Japan, 115–142 (2004) [Zbl 1114.82017](#) [MR 2073332](#)
- [CJKS14] Conus, D., Joseph, M., Khoshnevisan, D., Shiu, S-Y.: Initial measures for the stochastic heat equation. *Ann. Inst. H. Poincaré Probab. Statist.* **50**, 136–153 (2014) [Zbl 1288.60077](#) [MR 3161526](#)
- [C12] Corwin, I.: The Kardar–Parisi–Zhang equation and universality class. *Random Matrices Theory Appl.* **1**, 1130001, 76 pp. (2012) [Zbl 1247.82040](#) [MR 2930377](#)
- [DZ] Dey, P., Zygouras, N.: High temperature limits of directed polymers with heavy tail disorder. *Ann. Probab.* (to appear)
- [D97] Doney, R. A.: One-sided local large deviation and renewal theorems in the case of infinite mean. *Probab. Theory Related Fields* **107**, 451–465 (1997) [Zbl 0883.60022](#) [MR 1440141](#)
- [E06] Ellis, R.: *Entropy, Large Deviations, and Statistical Mechanics*. Springer, Berlin (2006) [Zbl 1102.60087](#) [MR 2189669](#)
- [G07] Giacomini, G.: *Random Polymer Models*. Imperial College Press, London (2007) [Zbl 1125.82001](#) [MR 2380992](#)
- [G10] Giacomini, G.: *Disorder and Critical Phenomena Through Basic Probability Models*. *Lecture Notes in Math.* 2025, Springer (2011) [Zbl 1230.82004](#) [MR 2816225](#)
- [G06] Grimmett, G.: *The Random-Cluster Model*. *Grundlehren Math. Wiss.* 333, Springer, Berlin (2006) [Zbl 1122.60087](#) [MR 2243761](#)
- [H74] Harris, A. B.: Effect of random defects on the critical behaviour of Ising models. *J. Phys. C* **7**, 1671–1692 (1974)
- [HH85] Huse, D. A., Henley, C. L.: Pinning and roughening of domain wall in Ising systems due to random impurities. *Phys. Rev. Lett.* **54**, 270–2711 (1985)
- [J97] Janson, S.: *Gaussian Hilbert Spaces*. *Cambridge Tracts in Math.* 129, Cambridge Univ. Press, Cambridge (1997) [Zbl 0887.60009](#) [MR 1474726](#)
- [KKL88] Kahn, J., Kalai, G., Linial, N.: The influence of variables on boolean functions. In: *Proceedings of the 29th Annual Symposium on Foundations of Computer Science*, 68–80 (1988)

- [L10] Lacoïn, H.: New bounds for the free energy of directed polymer in dimension 1+1 and 1+2. *Comm. Math. Phys.* **294**, 471–503 (2010) [Zbl 1227.82098](#) [MR 2579463](#)
- [LL01] Lieb, E. H., Loss, M.: *Analysis*. 2nd ed., *Grad. Stud. Math.* 14, Amer. Math. Soc., Providence, RI (2001) [Zbl 0966.26002](#) [MR 1817225](#)
- [Lig68] Liggett, T. L.: An invariance principle for conditioned sums of independent random variables. *J. Math. Mech.* **18**, 559–570 (1968) [Zbl 0181.20502](#) [MR 0238373](#)
- [MOO10] Mossel, E., O’Donnell, R., Oleszkiewicz, K.: Noise stability of functions with low influences: Invariance and optimality. *Ann. of Math.* **171**, 295–341 (2010) [Zbl 1201.60031](#) [MR 2630040](#)
- [PT10] Peccati, G., Taqqu, M. S.: *Wiener Chaos: Moments, Cumulants and Diagrams*. Springer (2010) [Zbl 1231.60003](#) [MR 2791919](#)
- [SS10] Sasamoto, T., Spohn, H.: Exact height distributions for the KPZ equation with narrow wedge initial condition. *Nuclear Phys. B* **834**, 523–542 (2010) [Zbl 1204.35137](#) [MR 2628936](#)
- [Soh09] Sohler, J.: Finite size scaling for homogeneous pinning models. *Alea* **6**, 163–177 (2009) [Zbl 1172.60336](#) [MR 2506863](#)
- [TV11] Tao, T., Vu, V.: Random matrices: Universality of the local eigenvalue statistics. *Acta Math.* **206**, 127–204 (2011) [Zbl 1217.15043](#) [MR 2784665](#)