



Izzet Coskun · Jack Huizenga · Matthew Woolf

## The effective cone of the moduli space of sheaves on the plane

Received July 15, 2014

**Abstract.** Let  $\xi$  be the Chern character of a stable coherent sheaf on  $\mathbb{P}^2$ . We compute the cone of effective divisors on the moduli space  $M(\xi)$  of semistable sheaves on  $\mathbb{P}^2$  with Chern character  $\xi$ . The computation hinges on finding a good resolution of the general sheaf in  $M(\xi)$ . This resolution is determined by Bridgeland stability and arises from a well-chosen Beilinson spectral sequence. The existence of a good choice of spectral sequence depends on remarkable number-theoretic properties of the slopes of exceptional bundles.

**Keywords.** Moduli spaces of sheaves, Bridgeland stability, effective cone, Brill–Noether divisors

### Contents

1. Introduction . . . . .	1421
2. Preliminaries . . . . .	1425
3. The corresponding exceptional bundle . . . . .	1433
4. Transcendentality of the generalized Cantor set . . . . .	1436
5. The Beilinson spectral sequence . . . . .	1445
6. The Kronecker fibration . . . . .	1454
7. The effective cone of the moduli space . . . . .	1460
References . . . . .	1465

### 1. Introduction

In this paper, we determine the cone of effective divisors on any moduli space of semi-stable sheaves on  $\mathbb{P}^2$ .

The effective and movable cones are among the most important invariants of a variety  $X$  (see [La]). They dictate both the geometry and arithmetic of  $X$ . The *movable cone*  $\text{Mov}(X)$ , which is spanned by divisor classes whose stable base locus does not contain

---

I. Coskun, M. Woolf: Department of Mathematics, Statistics and CS,  
University of Illinois at Chicago, Chicago, IL 60607, USA;  
e-mail: [coskun@math.uic.edu](mailto:coskun@math.uic.edu), [mwoolf@uic.edu](mailto:mwoolf@uic.edu)

J. Huizenga: Department of Mathematics, The Pennsylvania State University,  
University Park, PA 16802, USA; e-mail: [huizenga@math.psu.edu](mailto:huizenga@math.psu.edu)

*Mathematics Subject Classification (2010):* Primary 14J60; Secondary 14E30, 14D20, 13D02

any divisorial components, controls the rational maps from  $X$  to other projective varieties. The *effective cone*  $\text{Eff}(X)$ , which is spanned by classes of effective divisors, plays a prominent role in determining the basic birational invariants of  $X$  such as the Kodaira dimension and is expected to dictate the asymptotics of rational points on  $X$  via Manin's conjecture (see [T]).

Let  $\xi$  be the Chern character of a stable coherent sheaf on  $\mathbb{P}^2$ . Then the Gieseker moduli space  $M(\xi)$ , which parameterizes  $S$ -equivalence classes of semistable sheaves on  $\mathbb{P}^2$  with Chern character  $\xi$ , is an irreducible, normal, factorial projective variety (see [LP]).

When  $\xi$  is the Chern character of an *exceptional* or *height zero* bundle (see §2),  $M(\xi)$  is a point or has Picard rank one, respectively. In these cases there is nothing further to discuss. Hence, we will always assume that  $M(\xi)$  is not an exceptional or height zero moduli space. There is a natural pairing on the Grothendieck group  $K(\mathbb{P}^2)$  that associates to two Chern characters  $\xi$  and  $\zeta$  the Euler characteristic  $\chi(\xi^*, \zeta)$ . The Picard group of  $M(\xi)$  is a free abelian group of rank 2, naturally identified with  $\xi^\perp$  in  $K(\mathbb{P}^2)$  with respect to this pairing [DLP]. Correspondingly, the Néron–Severi space  $\text{NS}(M(\xi))$  is a two-dimensional vector space equal to  $\text{Pic}(M(\xi)) \otimes \mathbb{R}$ . Therefore, in order to determine the effective and movable cones, it suffices to specify their two extremal rays.

We now describe our results on the effective cone of  $M(\xi)$  in greater detail. Since  $M(\xi)$  is factorial, in order to construct effective Cartier divisors, it suffices to impose codimension 1 geometric conditions on sheaves. Brill–Noether conditions provide a systematic method for constructing effective divisors. Let  $\zeta \in \xi^\perp$  be the Chern character of a vector bundle  $V$ . There are  $h^0$ - and  $h^2$ -Brill–Noether loci on  $M(\xi)$  depending on whether the slope  $\mu(\xi \otimes \zeta)$  is positive or negative. These are related by duality and are responsible for the two different extremal rays of the effective cone. For simplicity, we will assume that the slope is positive and describe the  $h^0$ -Brill–Noether loci first and later explain the necessary changes needed for  $h^2$ -Brill–Noether loci. Define

$$D_V := \{U \in M(\xi) : h^0(U \otimes V) \neq 0\}$$

and endow it with the natural determinantal scheme structure. The locus  $D_V$  is either an effective divisor or  $D_V = M(\xi)$ . Consequently, if there exists  $U \in M(\xi)$  such that  $h^0(U \otimes V) = 0$ , then  $D_V$  is an effective Cartier divisor called the *Brill–Noether divisor associated to  $V$* .

**Definition 1.1.** A sheaf  $U \in M(\xi)$  is *cohomologically orthogonal* to a vector bundle  $V$  if  $h^i(U \otimes V) = 0$  for every  $i$ .

If  $U$  is cohomologically orthogonal to  $V$ , then  $\chi(U \otimes V) = 0$  and, by the Riemann–Roch Theorem, the discriminant of  $V$  is determined by the slope of  $V$  (see §2). Furthermore, the Brill–Noether divisor  $D_V$  is an effective divisor that does not contain  $U$  in its base locus. Hence, the following problem naturally arises in studying stable base loci of linear systems on  $M(\xi)$ .

**Problem 1.2** (Higher rank interpolation). Given a sheaf  $U \in M(\xi)$ , determine the minimum slope  $\mu^+ \in \mathbb{Q}$  such that

- (1)  $\mu(\xi) + \mu^+ \geq 0$ , and
- (2)  $U$  is cohomologically orthogonal to a vector bundle  $V$  of slope  $\mu^+$ .

The name of the problem is a holdover from the case where  $M(\xi)$  is a Hilbert scheme of points  $\mathbb{P}^{2[m]}$ . In that case the problem amounts to determining when a zero-dimensional scheme  $Z$  can impose independent conditions on sections of a vector bundle. This problem was previously solved in [H] for a general scheme  $Z$  and in [CHI] for monomial schemes. In this paper, we will solve the problem for general sheaves in any moduli space  $M(\xi)$ . This will determine the effective cone of  $M(\xi)$ .

Our answer to the interpolation problem is in terms of the classification of stable vector bundles on  $\mathbb{P}^2$  due to Drézet and Le Potier [DLP, LP]. The classification, which we will recall in §2, can best be phrased in terms of the slope  $\mu$  and the discriminant  $\Delta$ . There is a fractal-like curve  $\delta(\mu)$  in the  $(\mu, \Delta)$ -plane such that the moduli space of vector bundles with invariants  $(r, \mu, \Delta)$  is positive-dimensional if and only if  $\Delta \geq \delta(\mu)$  and  $r$  is a sufficiently divisible integer so that the Chern class  $c_1$  and the Euler characteristic  $\chi$  are integers. In addition, there are exceptional bundles whose moduli spaces are isolated points. The slopes of exceptional bundles are in one-to-one correspondence with dyadic integers (see §2). For each exceptional slope  $\alpha$ , there is an interval  $I_\alpha = (\alpha - x_\alpha, \alpha + x_\alpha)$  such that the curve  $\delta(\mu)$  is defined on  $(\alpha - x_\alpha, \alpha]$  by a parabola  $P_1^\alpha$  and on  $[\alpha, \alpha + x_\alpha)$  by a parabola  $P_2^\alpha$  (see §2 for explicit equations).

Each Chern character  $\xi$  determines a (possibly degenerate, if  $r(\xi) = 0$ ) parabola  $Q_\xi$  in the  $(\mu, \Delta)$ -plane parameterizing pairs  $(\mu, \Delta)$  that are orthogonal to  $\xi$ . Assume that the rightmost intersection of  $Q_\xi$  with the line  $\Delta = 1/2$  occurs along the interval  $I_\alpha$  containing the exceptional slope  $\alpha$ . Our answer depends on three cases.

First, in special cases, the invariants of the exceptional bundle  $E_\alpha$  of slope  $\alpha$  may lie on the parabola  $Q_\xi$ . In this case, let  $(\mu^+, \Delta^+)$  be the slope and discriminant of  $E_\alpha$ . Second, if the invariants of  $E_\alpha$  are below the parabola  $Q_\xi$ , let  $(\mu^+, \Delta^+)$  be the intersection point of  $Q_\xi$  with the parabola  $P_1^\alpha$ . Finally, if the invariants of  $E_\alpha$  are above the parabola  $Q_\xi$ , let  $(\mu^+, \Delta^+)$  be the intersection point of  $Q_\xi$  with the parabola  $P_2^\alpha$  (see §3). Let  $\xi^+$  be the Chern character  $(r^+, \mu^+, \Delta^+)$ , where  $r^+$  is sufficiently high and divisible. The character  $\xi^+$  is always a stable Chern character. Let  $V$  denote a general point of the moduli space  $M(\xi^+)$ .

With this notation, the main theorem of this paper is the following.

**Theorem 1.3.** *Let  $\xi$  be the Chern character of a stable sheaf such that  $M(\xi)$  has Picard rank 2. Let  $U$  be a general point of  $M(\xi)$ .*

- (1) *The slope  $\mu^+$  is the solution to the higher rank interpolation problem for  $U$ .*
- (2) *The Brill–Noether divisor  $D_V$  spans an extremal ray of  $\text{Eff}(M(\xi))$ .*
- (3) *If the invariants of  $E_\alpha$  do not lie on  $Q_\xi$ , then  $D_V$  also spans an extremal ray of  $\text{Mov}(M(\xi))$ .*

The proof of Theorem 1.3 relies on finding an appropriate resolution of  $U$ . In §5, we will give a resolution of  $U$  in terms of a triad of exceptional bundles. The choice of optimal triad is determined by the intersection of  $Q_\xi$  with the line  $\Delta = 1/2$ , and the resolution naturally arises from the Beilinson spectral sequence. This resolution allows one to compute the cohomology of  $U \otimes V$  effectively.

We will show that  $M(\xi)$  always admits a dominant rational map to a moduli space of appropriately chosen Kronecker modules (see §6). As long as the invariants of the

exceptional bundle  $E_\alpha$  do not lie on  $Q_\xi$ , this map has positive-dimensional fibers. Consequently, the pullback of the ample generator from the moduli space of Kronecker modules gives an edge of both the effective and movable cones. When the invariants of  $E_\alpha$  lie on  $Q_\xi$ , the corresponding rational map is birational and contracts a divisor. The exceptional locus gives an extremal edge of the effective cone, and the pullback of the ample generator on the moduli space of Kronecker modules defines a movable class.

The curve  $\delta(\mu)$  intersects the line  $\Delta = 1/2$  in a Cantor set  $C$  (see §4), which consists precisely of the real numbers lying in no interval  $I_\alpha$ . One of the main technical points that we have to address is to show that the parabola  $Q_\xi$  does not intersect the line  $\Delta = 1/2$  along  $C$  so that there is always a well-defined interval  $I_\alpha$ . In §4, we will prove that every point of the Cantor set which is not an endpoint of an interval  $I_\alpha$  is transcendental (see Theorem 4.1). We will further show that  $Q_\xi$  cannot pass through an endpoint of an interval  $I_\alpha$ . Since the intersection of  $Q_\xi$  with  $\Delta = 1/2$  is either a rational number or a quadratic irrational, it follows that  $Q_\xi$  does not intersect the Cantor set. These results rely on remarkable properties of the continued fraction expansions of points of  $C$  and deep Diophantine approximation results such as Roth's Theorem [R] and the Schmidt Subspace Theorem [Sc].

We now discuss the other edge of the effective cone. When the rank of  $\xi$  is at least three, consider the Serre dual Chern character  $\xi^D$ . Duality gives a birational map  $M(\xi) \dashrightarrow M(\xi^D)$  which is an isomorphism in codimension 1. It therefore induces an isomorphism of Picard groups which preserves the effective cones. The edge of  $\text{Eff}(M(\xi^D))$  constructed in Theorem 1.3 is carried to the other edge of  $\text{Eff}(M(\xi))$ . More precisely, if  $D_V$  is an  $h^0$ -Brill–Noether divisor giving an edge of  $\text{Eff}(M(\xi^D))$ , then the  $h^2$ -Brill–Noether divisor

$$\{U \in M(\xi) : h^2(U \otimes V^*) \neq 0\}$$

corresponding to  $V^*$  gives the other edge of  $\text{Eff}(M(\xi))$ .

When the rank of  $\xi$  is 1 or 2, there is a morphism  $j : M(\xi) \rightarrow M^{\text{DUY}}(\xi)$  to the Donaldson–Uhlenbeck–Yau moduli space constructed by Jun Li (see [Li], [HuL]). The exceptional locus of  $j$  is an extremal effective divisor spanning the other edge of the effective cone. The morphism  $j$  is defined by a nef divisor giving the other edge of the movable cone. The exceptional locus of  $j$  has a simple geometric description.

- When  $\xi$  has rank 2, the exceptional locus of  $j$  is the locus of singular sheaves.
- When  $\xi$  has rank 1, the space  $M(\xi)$  is isomorphic to a Hilbert scheme of points  $\mathbb{P}^{2[m]}$ , and  $M^{\text{DUY}}(\xi)$  is isomorphic to the symmetric product  $\mathbb{P}^{2(n)}$ . The morphism  $j$  is the Hilbert–Chow morphism, and its exceptional locus is the locus of nonreduced schemes.

When  $\xi$  has rank 0 and  $M(\xi)$  has Picard rank 2, then we must have  $d := c_1(\xi) \geq 3$ , and  $M(\xi)$  parameterizes  $S$ -equivalence classes of pure one-dimensional sheaves with Chern character  $\xi$ . Let  $s : M(\xi) \rightarrow |\mathcal{O}_{\mathbb{P}^2}(d)| = \mathbb{P}^N$  be the morphism sending a sheaf to its support. Then  $s^*\mathcal{O}_{\mathbb{P}^N}(1)$  spans the other extremal ray of both the movable and the effective cones [W].

Bridgeland stability conditions were effectively used to clarify the birational geometry of the Hilbert scheme of points on  $\mathbb{P}^2$  in [ABCH], [LZ] and [Oh]. In [H], the second author determined the collapsing wall without carrying out the entire minimal model program.

The current paper is a natural continuation of this line of work, where we show that the collapsing wall is given by an appropriately chosen exceptional bundle. The ample cone of  $M(\xi)$  is computed in [CH2]. There have been parallel developments using Bridgeland stability in the study of moduli spaces of sheaves on other surfaces such as K3 surfaces [BM1], [BM2], [MY1], [MY2], abelian surfaces [MM], [YY], [Y], Enriques surfaces [Ne], and Hirzebruch and del Pezzo surfaces [BC].

### *Organization of the paper*

In §2, we will recall basic facts concerning coherent sheaves, exceptional bundles, the classification of stable bundles, the Picard group of the moduli space, and Brill–Noether divisors. In §3, we will show how to associate an exceptional bundle to a given Chern character  $\xi$  and state our theorem on the effective cone precisely. In §4, we will study the number-theoretic properties of points in the Cantor set  $C$  and show that every point of  $C$  which is not an endpoint of an interval is transcendental. In §5, we will use the corresponding exceptional bundle to choose an appropriate triad of exceptional bundles. We then use the Beilinson spectral sequence to determine an important resolution of the general sheaf  $U \in M(\xi)$ . In §6, we define a rational map from the moduli space of sheaves on  $\mathbb{P}^2$  to a moduli space of Kronecker modules. This map arises from the special resolution of the general sheaf. In §7, we will determine the effective cone of the moduli space of sheaves. We close the paper with a sample explicit computation of the effective cone of a moduli space.

## 2. Preliminaries

In this section, we collect basic facts concerning the classification of stable bundles on  $\mathbb{P}^2$  and the geometry of the Gieseker moduli spaces of semistable sheaves on  $\mathbb{P}^2$ . We refer the reader to books by Le Potier [LP] and Huybrechts and Lehn [HuL] for more details.

### *2.1. Stability; slope and discriminant; Euler pairing*

In this paper all sheaves will be coherent. Let  $E$  be a coherent sheaf. The dimension  $\dim E$  of  $E$  is the dimension of the support  $\text{Supp}(E)$ . We say  $E$  is *pure of dimension  $d$*  if  $\dim E = d$  and every nontrivial coherent subsheaf  $F \subset E$  has  $\dim F = d$ . If  $E$  has dimension  $d$ , the Hilbert polynomial  $P_E(m) = \chi(E(m))$  of  $E$  is of the form

$$P_E(m) = \alpha_d \frac{m^d}{d!} + O(m^{d-1}).$$

The *reduced Hilbert polynomial*  $p_E$  is defined by

$$p_E = P_E/\alpha_d.$$

Then  $E$  is (*Gieseker*) *semistable* (resp. *stable*) if  $E$  is pure and for every nontrivial  $F \subset E$  we have  $p_F \leq p_E$  (resp.  $<$ ), where polynomials are compared for all sufficiently large  $m$ .

Fix a Chern character  $\xi = (r, \text{ch}_1, \text{ch}_2)$ . If the rank  $r$  is positive, then the *slope*  $\mu$  and the *discriminant*  $\Delta$  are defined by

$$\mu = \text{ch}_1 / r \quad \text{and} \quad \Delta = \frac{1}{2}\mu^2 - \text{ch}_2 / r.$$

The rank, slope and discriminant determine the Chern character. Hence, we can equivalently record the Chern character by  $\xi = (r, \mu, \Delta)$ . The classification of stable vector bundles on  $\mathbb{P}^2$  is most conveniently expressed in terms of these invariants, so we will primarily use them instead of  $\text{ch}_1$  and  $\text{ch}_2$ . If  $r > 0$ , the Riemann–Roch formula reads

$$\chi(E) = r(P(\mu) - \Delta),$$

where  $P(m)$  is the Hilbert polynomial of  $\mathcal{O}_{\mathbb{P}^2}$  given by

$$P(m) = \frac{1}{2}(m^2 + 3m + 2).$$

Furthermore,

$$\mu(E \otimes F) = \mu(E) + \mu(F) \quad \text{and} \quad \Delta(E \otimes F) = \Delta(E) + \Delta(F).$$

These definitions easily extend to Chern characters in  $K(\mathbb{P}^2) \otimes \mathbb{R}$ , and we observe that if  $0 \neq c \in \mathbb{R}$  then  $\xi$  and  $c\xi$  have the same slope and discriminant. Note that semistability of a sheaf  $E$  is equivalent to requiring that for every proper subsheaf  $F \subset E$  we have  $\mu(F) \leq \mu(E)$ , with  $\Delta(F) \geq \Delta(E)$  in case of equality. A character  $\xi$  is (*semi*)*stable* if it is the Chern character of a (semi)stable sheaf.

Given two sheaves  $E$  and  $F$ , the Euler characteristic of the pair  $(E, F)$  is defined by the formula

$$\chi(E, F) = \sum_{i=0}^2 (-1)^i \text{ext}^i(E, F).$$

When both sheaves have nonzero rank, the Euler characteristic is computed by the Riemann–Roch formula

$$\chi(E, F) = r(E)r(F)(P(\mu(F) - \mu(E)) - \Delta(E) - \Delta(F)).$$

The derived dual induces a homomorphism  $K(\mathbb{P}^2) \rightarrow K(\mathbb{P}^2)$ . We will write  $\xi^*$  for the dual Chern character. The Euler characteristic depends only on Chern characters, so it induces a bilinear pairing  $(\xi, \zeta) = \chi(\xi^*, \zeta)$  on  $K(\mathbb{P}^2) \otimes \mathbb{R}$ . Correspondingly,  $\xi^\perp$  denotes the orthogonal complement of  $\xi$  in  $K(\mathbb{P}^2) \otimes \mathbb{R}$  with respect to this pairing.

## 2.2. Exceptional bundles

A stable vector bundle  $E$  on  $\mathbb{P}^2$  is called an *exceptional bundle* if  $\text{Ext}^1(E, E) = 0$ . The line bundles  $\mathcal{O}_{\mathbb{P}^2}(n)$  and the tangent bundle  $T_{\mathbb{P}^2}$  are examples of exceptional bundles. An *exceptional slope*  $\alpha$  is the slope of an exceptional bundle. If  $\alpha$  is an exceptional slope,

there is a unique exceptional bundle  $E_\alpha$  of slope  $\alpha$ . The rank of  $E_\alpha$  is the smallest positive integer  $r_\alpha$  such that  $r_\alpha\alpha$  is an integer. The discriminant  $\Delta_\alpha$  is then given by

$$\Delta_\alpha = \frac{1}{2}(1 - 1/r_\alpha^2).$$

We also write  $\xi_\alpha = \text{ch}(E_\alpha)$ . All exceptional bundles can be constructed starting with line bundles via the process of mutation [D2]. Exceptional bundles are precisely the stable bundles  $E$  on  $\mathbb{P}^2$  with  $\Delta(E) < 1/2$  [LP, Proposition 16.1.1]. They are rigid and their moduli spaces consist of a single reduced point [LP, Corollary 16.1.5].

There is a one-to-one correspondence between the set  $\mathcal{E}$  of exceptional slopes and dyadic integers,  $\varepsilon : \mathbb{Z}[1/2] \rightarrow \mathcal{E}$ , defined inductively by  $\varepsilon(n) = n$  for  $n \in \mathbb{Z}$  and

$$\varepsilon\left(\frac{2p+1}{2^{q+1}}\right) = \varepsilon\left(\frac{p}{2^q}\right) \cdot \varepsilon\left(\frac{p+1}{2^q}\right),$$

where  $\alpha \cdot \beta$  is defined by

$$\alpha \cdot \beta = \frac{\alpha + \beta}{2} + \frac{\Delta_\beta - \Delta_\alpha}{3 + \alpha - \beta}.$$

Define the *order* of an exceptional slope  $\alpha \in \mathcal{E}$  to be the smallest natural number  $q$  such that  $\alpha = \varepsilon(p/2^q)$ .

**Example 2.1.** We record the first several exceptional slopes in the interval  $[0, 1/2]$  together with their orders.

$p/2^q$	0	$\frac{1}{16}$	$\frac{1}{8}$	$\frac{3}{16}$	$\frac{1}{4}$	$\frac{5}{16}$	$\frac{3}{8}$	$\frac{7}{16}$	$\frac{1}{2}$
$\varepsilon(p/2^q)$	0	$\frac{13}{34}$	$\frac{5}{13}$	$\frac{75}{194}$	$\frac{2}{5}$	$\frac{179}{433}$	$\frac{12}{29}$	$\frac{70}{169}$	$\frac{1}{2}$
$\text{ord}(\varepsilon(p/2^q))$	0	4	3	4	2	4	3	4	1

### 2.3. The classification of stable bundles

The classification of positive-dimensional moduli spaces of stable vector bundles on  $\mathbb{P}^2$  is expressed in terms of a fractal-like curve  $\delta$  in the  $(\mu, \Delta)$ -plane. Let

$$\delta(\mu) = \sup_{\{\alpha \in \mathcal{E} : |\mu - \alpha| < 3\}} (P(-|\mu - \alpha|) - \Delta_\alpha).$$

For each exceptional slope  $\alpha \in \mathcal{E}$ , there is an interval  $I_\alpha = (\alpha - x_\alpha, \alpha + x_\alpha)$  such that the function  $\delta(\mu)$  is defined on  $I_\alpha$  by

$$\delta(\mu) = P(-|\mu - \alpha|) - \Delta_\alpha \quad \text{if } \mu \in I_\alpha,$$

where

$$x_\alpha = \frac{3 - \sqrt{5 + 8\Delta_\alpha}}{2}.$$

The graph of  $\delta(\mu)$  is an increasing concave up parabola on the interval  $[\alpha - x_\alpha, \alpha]$  and a decreasing concave up parabola on the interval  $[\alpha, \alpha + x_\alpha]$ . The graph of  $\delta$  over  $I_\alpha$  is

symmetric across the vertical line  $\mu = \alpha$  and is invariant under translation by integers. Furthermore,  $\delta(\alpha \pm x_\alpha) = 1/2$  for every  $\alpha \in \mathcal{E}$ . The graph of  $\delta$  intersects the line  $\Delta = 1/2$  in a Cantor set

$$C := \mathbb{R} \setminus \bigcup_{\alpha \in \mathcal{E}} I_\alpha.$$

The following result can be viewed as the first structure theorem for  $C$ ; we will strengthen this theorem in §4.

**Theorem 2.2** ([D3, Theorem 1]). *Every rational number  $q \in \mathbb{Q}$  lies in some interval  $I_\alpha$ ; equivalently, the Cantor set  $C$  consists entirely of irrational numbers.*

The fundamental theorem on the existence of moduli spaces of semistable sheaves due to Drézet and Le Potier is the following.

**Theorem 2.3** ([DLP], [LP]). *Let  $\xi = (r, \mu, \Delta)$  be a Chern character of positive integer rank. There exists a positive-dimensional moduli space of semistable sheaves  $M(\xi)$  with Chern character  $\xi$  if and only if  $c_1 = r\mu \in \mathbb{Z}$ ,  $\chi = r(P(\mu) - \Delta) \in \mathbb{Z}$  and  $\Delta \geq \delta(\mu)$ . In this case,  $M(\xi)$  is a normal, irreducible, factorial projective variety of dimension  $r^2(2\Delta - 1) + 1$ .*

Height zero characters  $\xi$  are characters with invariants lying on the curve  $\Delta = \delta(\mu)$ .

Moreover,  $M(\xi)$  is a Mori dream space. In particular, the effective cone is closed. This has been proved in [W, §3], but only explicitly stated for the rank 0 case. We briefly sketch the proof in the higher rank case and refer to [W] for additional details.

**Theorem 2.4.** *Let  $\xi$  be the Chern character of a stable coherent sheaf. Then the moduli space  $M(\xi)$  is a Mori dream space.*


*Proof.* See [W] for the rank 0 case. We assume  $r(\xi) > 0$ . By [BCHM], a log Fano variety is a Mori dream space. Consequently, it suffices to show that  $M(\xi)$  is a log Fano variety. By [HuL, Theorems 8.2.8 and 8.3.3] the anti-canonical bundle of  $M(\xi)$  is nef. By the proof of [W, Proposition 3.3],  $M(\xi)$  has canonical singularities. Briefly,  $M(\xi)$  is the good quotient of a smooth variety [LP, §14.5]. Consequently,  $M(\xi)$  has rational singularities [B]. Since  $M(\xi)$  is Gorenstein and every rational Gorenstein singularity is canonical [K, Theorem 11.1], we conclude that  $M(\xi)$  has canonical singularities. Hence, the pair  $(M(\xi), \epsilon E)$  is a klt-pair for every effective divisor  $E$  and sufficiently small  $\epsilon > 0$ . It will become clear in §7 that there are effective divisors  $E$  such that  $-K_{M(\xi)} - \epsilon E$  is ample for all sufficiently small  $\epsilon > 0$ . We conclude that  $M(\xi)$  is a log Fano variety, hence a Mori dream space.  $\square$

#### 2.4. Picard group of the moduli space

A theorem of Drézet determines the Picard group of  $M(\xi)$ .

**Theorem 2.5** (Drézet). *If  $\Delta > \delta(\mu)$ , then the Picard group of  $M(\xi)$  is a free abelian group on two generators. Furthermore, in this case,  $\text{Pic}(M(\xi)) \otimes \mathbb{R}$  is naturally identified with  $\xi^\perp$ . If  $\Delta = \delta(\mu)$ , then the Picard group of  $M(\xi)$  is an infinite cyclic group. (See Figure 1.)*





**Fig. 1.** The curve  $\delta(\mu) = \Delta$  occurring in the classification of stable bundles. The invariants of the first several exceptional bundles are also displayed. A Chern character  $\xi = (r, \mu, \Delta)$  with  $c_1(\xi) \in \mathbb{Z}$  and  $\chi(\xi) \in \mathbb{Z}$  has a moduli space  $M(\xi)$  of Picard rank 2 if and only if the point  $(\mu, \Delta)$  lies in the shaded region above the curve. If  $(\mu, \Delta)$  lies on the curve, the moduli space has Picard rank 1.

In  $M(\xi)$ , linear equivalence and numerical equivalence coincide and the Néron–Severi space  $\text{NS}(M(\xi)) = \text{Pic}(M(\xi)) \otimes \mathbb{R}$ . When the Picard rank of a projective variety is 1, then the ample, effective and movable cones coincide and are equal to the half-space spanned by any effective divisor. Therefore, for the rest of the paper, we will always assume that  $\xi$  is a Chern character satisfying  $\Delta > \delta(\mu)$ . In this case, the Néron–Severi space is a two-dimensional vector space. Hence, the cones  $\text{Mov}(M(\xi))$  and  $\text{Eff}(M(\xi))$  are determined by specifying their two extremal rays.

At times it will be necessary to consider the open subvariety  $M^s(\xi)$  of  $M(\xi)$  parameterizing stable sheaves. The space  $M^s(\xi)$  is a coarse moduli space for flat families of stable sheaves with Chern character  $\xi$ . On the other hand, the identification of  $S$ -equivalence classes of sheaves prevents  $M(\xi)$  from being a coarse moduli space for isomorphism classes of sheaves unless  $\xi$  is a primitive character. When the Picard rank of  $M(\xi)$  is at least 2, the complement of  $M^s(\xi)$  has codimension at least 2 by [LP, Lemma 18.3.1]. Thus when studying questions about divisors, there is typically no harm in working on  $M^s(\xi)$  instead of  $M(\xi)$  as necessary.

Most of the effective divisors considered in this paper will be certain Brill–Noether divisors. It will be important to understand the class of a Brill–Noether divisor in terms of the isomorphism  $\xi^\perp \cong \text{NS}(M(\xi))$ . We first recall the construction of this isomorphism more explicitly. We summarize the discussion in [LP, Chapter 18].

Let  $\mathcal{U}/S$  be a flat family of semistable sheaves of Chern character  $\xi$  on  $\mathbb{P}^2$  over a variety  $S$ , and let  $p : S \times \mathbb{P}^2 \rightarrow S$  and  $q : S \times \mathbb{P}^2 \rightarrow \mathbb{P}^2$  be the two projections. A homomorphism  $\lambda_{\mathcal{U}} : K(\mathbb{P}^2) \rightarrow \text{Pic}(S)$  is described as the composition

$$K(\mathbb{P}^2) \xrightarrow{q^*} K^0(S \times \mathbb{P}^2) \xrightarrow{[\mathcal{U}]} K^0(S \times \mathbb{P}^2) \xrightarrow{-p_*} K^0(S) \xrightarrow{\det} \text{Pic } S,$$

where  $p_i = \sum(-1)^i R^i p_*$ . If  $\mathcal{U}$  is replaced by a family  $\mathcal{U} \otimes p^*L$  for some line bundle  $L$  on  $S$ , then the moduli map  $S \rightarrow M(\xi)$  does not change, but

$$\lambda_{\mathcal{U} \otimes p^*L}(\zeta) = \lambda_{\mathcal{U}}(\zeta) \otimes L^{\otimes -(\xi, \zeta)}.$$

For an integral class  $\zeta \in \xi^\perp$ , there is a line bundle denoted by  $\lambda_M(\zeta)$  on  $M(\xi)$  with the property that if  $\mathcal{U}/S$  is a flat family of semistable sheaves with Chern character  $\xi$  and  $f : S \rightarrow M(\xi)$  is the moduli map, then

$$\lambda_{\mathcal{U}}(\zeta) = f^* \lambda_M(\zeta).$$

This induces a linear map  $\lambda_M : \xi^\perp \rightarrow \text{NS}(M(\xi))$  which is an isomorphism when  $\Delta > \delta(\mu)$ .

**Warning 2.6.** We have chosen to normalize  $\lambda_M$  by using  $-p_i$  instead of  $p_i$ , as is considered in [HuL] and [LP]. We will discuss the reason for this change below in Remark 2.10.

When they exist, Brill–Noether divisors provide explicit sections of some of these line bundles.

**Proposition 2.7.** *Let  $\xi$  be a positive-rank Chern character such that  $M(\xi)$  is of Picard rank 2. Suppose  $V$  is a stable vector bundle of Chern character  $\zeta$  such that the general  $U \in M(\xi)$  is cohomologically orthogonal to  $V$ . Endow the locus*

$$D_V = \overline{\{U \in M^s(\xi) : H^1(U \otimes V) \neq 0\}} \subset M(\xi)$$

with the natural determinantal scheme structure.

- (1)  $D_V$  is an effective divisor;
- (2) If  $\mu(U \otimes V) > -3$ , then  $\mathcal{O}_{M(\xi)}(D_V) \cong \lambda_M(\zeta)$ .
- (3) If  $\mu(U \otimes V) < 0$ , then  $\mathcal{O}_{M(\xi)}(D_V) \cong \lambda_M(\zeta)^* = \lambda_M(-\zeta)$ .

*Proof.* We first recall how to give  $D_V$  a determinantal scheme structure; the fact that it is a divisor will also follow. Let  $\mathcal{U}/S$  be a complete flat family of stable sheaves with Chern character  $\xi$ . We will describe a determinantal scheme structure on the locus  $D_{V,S}$  of  $s \in S$  such that  $U_s \in D_V$ . If  $M^s(\xi)$  admits a universal family, this gives the correct scheme structure on  $D_V$ . Otherwise, one can work with the universal family on the stack  $\mathcal{M}^s(\xi)$  and then take the image in  $M^s(\xi)$  (see [A, Theorem 4.16 and Example 8.7]).

Suppose  $\mu(U \otimes V) > -3$ . Let  $d \gg 0$ , and fix some  $s \in S$ . We first describe  $D_{V,S}$  locally near  $s$ . Let  $C \subset \mathbb{P}^2$  be a general curve of degree  $d$ , so that  $C$  avoids any singularities of  $U_s$ . Replacing  $S$  by a smaller open subset  $S'$ , we may assume  $C$  avoids the singularities of  $U_s$  for all  $s \in S'$ . Consider the exact sequence

$$0 \rightarrow \mathcal{U} \otimes q^*V \rightarrow \mathcal{U} \otimes q^*(V(d)) \rightarrow \mathcal{U} \otimes q^*(V(d)|_C) \rightarrow 0$$

of sheaves on  $S' \times \mathbb{P}^2$ . For a fixed  $s \in S'$  we have a sequence

$$H^1(U_s \otimes V(d)) \rightarrow H^1(U_s \otimes V(d)|_C) \rightarrow H^2(U_s \otimes V).$$

The first cohomology group here vanishes since  $d \gg 0$  and the moduli space  $M(\xi)$  is bounded. The last group vanishes by stability since  $\mu(U \otimes V) > -3$ . Thus the middle group vanishes, and we conclude  $R^1 p_*(\mathcal{U} \otimes q^*(V(d)|_C)) = 0$ . We also have  $p_*(\mathcal{U} \otimes q^*V) = 0$  since the general  $U_S$  is cohomologically orthogonal to  $V$ . We then obtain an exact sequence

$$0 \rightarrow p_*(\mathcal{U} \otimes q^*(V(d))) \xrightarrow{\phi} p_*(\mathcal{U} \otimes q^*(V(d)|_C)) \rightarrow R^1 p_*(\mathcal{U} \otimes q^*V) \rightarrow 0$$

on  $S'$ . The sheaves  $A = p_*(\mathcal{U} \otimes q^*(V(d)))$  and  $B = p_*(\mathcal{U} \otimes q^*(V(d)|_C))$  are vector bundles of the same rank since  $d$  is sufficiently large and  $(\xi, \zeta) = 0$ . The divisor  $D_{V,S'}$  is the zero locus of the section  $\det \phi$  of the line bundle  $\mathcal{H}om(\det A, \det B) \cong \det R^1 p_*(\mathcal{U} \otimes q^*V)$ , so it has a scheme structure.

In fact, the scheme structure on  $D_{V,S'}$  described locally in the previous paragraph is independent of the choice of  $d$  and  $C$ , and the construction globalizes to give a scheme structure on the divisor  $D_{V,S}$ . Precisely, over an open set  $S'$  which is sufficiently small that  $A$  and  $B$  are isomorphic trivial bundles, the Fitting ideal sheaf  $\text{Fitt}_0(R^1 p_*(\mathcal{U} \otimes q^*V)) \subset \mathcal{O}_{S'}$  is generated by  $\det \phi$ , viewed as a section of  $\mathcal{O}_{S'}$  via the trivializations of  $A$  and  $B$  (see [E, Chapter 20]). This Fitting ideal is therefore the ideal sheaf of  $D_{V,S}$ , and the scheme structure on  $D_{V,S}$  is determined.

We can further identify

$$\text{Fitt}_0(R^1 p_*(\mathcal{U} \otimes q^*V))^* \cong \det R^1 p_*(\mathcal{U} \otimes q^*V).$$

Indeed, on an open set  $S'$  where  $A$  and  $B$  are locally trivial, there is an induced trivialization of the line bundle  $\det R^1 p_*(\mathcal{U} \otimes q^*V)$  and a local section  $\det \phi$ , viewed as a section of  $\mathcal{O}_{S'}$ . The collection of all such local sections forms an effective Cartier divisor by the existence of the Fitting ideal. This induces a global section of  $\det R^1 p_*(\mathcal{U} \otimes q^*V)$  that yields an isomorphism  $\det R^1 p_*(\mathcal{U} \otimes q^*V) \cong \mathcal{O}_S(D_{V,S})$ .

Finally, using  $V$  to compute  $\lambda_{\mathcal{U}}(\xi')$ , we find

$$\lambda_{\mathcal{U}}(\zeta) = \det(-p_1[\mathcal{U} \otimes q^*V]) = \det([R^1 p_*(\mathcal{U} \otimes q^*V)]) = \mathcal{O}_S(D_{V,S}),$$

with  $R^2 p_*(\mathcal{U} \otimes q^*V)$  vanishing by stability and the slope assumption. We conclude the corresponding result holds for the moduli space as well.

When  $\mu(U \otimes V) < 0$ , we instead consider an exact sequence

$$0 \rightarrow \mathcal{U} \otimes q^*(V(-d)) \rightarrow \mathcal{U} \otimes q^*V \rightarrow \mathcal{U} \otimes q^*V|_C \rightarrow 0$$

for  $d \gg 0$  and perform a similar calculation. We find  $\mathcal{O}_S(D_{V,S}) \cong \det R^2 p_*(\mathcal{U} \otimes q^*V)$  and compute  $\lambda_{\mathcal{U}}(\zeta)$  as before. □

**Remark 2.8.** In the proposition, we described the divisor  $D_V$  as the locus of all  $U$  where  $H^1(U \otimes V)$  jumps in rank. As  $\chi(U \otimes V) = 0$ , if  $\mu(U \otimes V) \geq 0$  then this locus can instead be described as the locus where  $H^0(U \otimes V)$  jumps rank. Similarly, if  $\mu(U \otimes V) \leq -3$ , it can be described as the locus where  $H^2(U \otimes V)$  jumps rank. These descriptions are perhaps preferable, in light of the fact that the divisor class of  $D_V$  exhibits qualitatively different behavior in each case despite the uniform description in terms of  $H^1$ .

It is also worth pointing out that  $U$  and  $V$  are never cohomologically orthogonal if  $-3 < \mu(U \otimes V) < 0$ . For if they are, then  $\lambda_M(\zeta) = \lambda_M(-\zeta)$ . As  $\lambda_M$  is an isomorphism, this is impossible. The hypothesis that the Picard rank of  $M(\xi)$  is 2 is crucial here. If  $\xi$  is a stable Chern character such that  $\chi(\xi) = 0$  and  $-3 < \mu(\xi) < 0$ , then any  $U \in M(\xi)$  is cohomologically orthogonal to  $\mathcal{O}_{\mathbb{P}^2}$ . It follows that  $M(\xi)$  has Picard rank at most 1. (This is also clear from Theorem 2.5.)

2.5. Halves of Néron–Severi space

There are two distinguished Chern characters  $\zeta_0, \zeta_1$  in  $\xi^\perp$ . They are uniquely determined by the requirements

$$\begin{aligned} r(\zeta_0) &= r(\xi), & r(\zeta_1) &= 0, \\ c_1(\zeta_0) &= 0, & c_1(\zeta_1) &= r(\xi). \end{aligned}$$

Line bundles  $\mathcal{L}_i$  are defined by  $\mathcal{L}_i = \lambda_M(\zeta_i)$ . The line bundle  $\mathcal{L}_1 := \lambda_M(\zeta_1)$  gives the Jun Li morphism  $j : M(\xi) \rightarrow M^{\text{DUY}}(\xi)$  to the Donaldson–Uhlenbeck–Yau compactification by slope-semistable sheaves, hence is nef. For  $n \gg 0$ , the line bundle  $\mathcal{L}_0 \otimes \mathcal{L}_1^n$  is ample [HuL].

We view the Chern character  $\zeta_1$  as spanning a line in  $\xi^\perp$  which splits this plane into two open halves, depending on the rank of a Chern character. We call the open half-plane of characters with positive rank the *primary* half, and the open half-plane of characters with negative rank the *secondary* half. We also split  $\text{NS}(M(\xi))$  into primary and secondary halves according to the isomorphism  $\text{NS}(M(\xi)) \cong \xi^\perp$ .

Consider a closed convex cone  $\Lambda \subset \text{NS}(M(\xi))$  which contains the nef cone and is not a half-plane. The effective and movable cones both have these properties. Since  $\Lambda$  contains both  $\mathcal{L}_1$  and  $\mathcal{L}_0 \otimes \mathcal{L}_1^n$  for some  $n \gg 0$ , exactly one of the edges of  $\Lambda$  lies in the primary half of  $\text{NS}(M(\xi))$ ; we call this edge the *primary edge* of  $\Lambda$ . We call the other edge of  $\Lambda$  the *secondary edge*. Describing  $\Lambda$  then amounts to computing its primary and secondary edges.

When the edge of a cone is given by a Brill–Noether divisor, it is simple to determine whether the edge is primary or secondary. The next result follows immediately from Proposition 2.7.

**Corollary 2.9.** *Keep the notation and hypotheses from Proposition 2.7.*

- (1) *If  $\mu(U \otimes V) > -3$ , then  $[D_V]$  lies in the primary half of  $\text{NS}(M(\xi))$ .*
- (2) *If  $\mu(U \otimes V) < 0$ , then  $[D_V]$  lies in the secondary half of  $\text{NS}(M(\xi))$ .*

**Remark 2.10.** Without our sign convention on  $\lambda_M$ , the  $h^0$ -Brill–Noether divisor associated to a bundle  $V$  of character  $\zeta$  with  $\mu(U \otimes V) > -3$  corresponds to the Chern character  $-\zeta$  of negative rank. The primary half of  $\xi^\perp$  must be taken to consist of the negative rank characters. Furthermore, the classes  $\zeta_0, \zeta_1$  would have negative rank and  $c_1$ , respectively. As we will be thinking of  $\text{NS}(M(\xi))$  primarily in terms of  $h^0$ -Brill–Noether divisors, our sign convention is advantageous.

### 3. The corresponding exceptional bundle

For the rest of the paper,  $\xi$  will denote a Chern character of positive rank such that the moduli space  $M(\xi)$  of semistable sheaves with Chern character  $\xi$  is of Picard rank 2. We will only lift this restriction in §7.2, where we discuss the case of torsion sheaves.

In this section, we introduce an exceptional bundle which controls the effective cone of divisors on  $M(\xi)$ . We also discuss how the effective cone is determined by this bundle. We focus on describing the primary edge of the effective cone, as defined in Section 2.5.

Recall that  $\xi^\perp$  is a linear space which is naturally identified with  $\text{NS}(M(\xi))$ . The rank of an orthogonal class is relatively unimportant, so it is convenient to work with slopes and discriminants instead of Chern characters. Hence, for any  $\zeta \in K(\mathbb{P}^2)$  we introduce the parabola

$$Q_\zeta = \{(\mu, \Delta) : (1, \mu, \Delta) \text{ lies in } \zeta^\perp\} \subset \mathbb{R}^2$$

as a subset of the  $(\mu, \Delta)$ -plane.

The Riemann–Roch formula shows that  $Q_\xi$  is the concave up parabola with equation

$$Q_\xi : P(\mu(\xi) + \mu) - \Delta(\xi) = \Delta.$$

where  $P(x) = \frac{1}{2}(x^2 + 3x + 2)$  is the Hilbert polynomial of  $\mathcal{O}_{\mathbb{P}^2}$ . Consider the *reference parabola*  $Q_{\xi_0}$  corresponding to the Chern character  $\xi_0$  of  $\mathcal{O}_{\mathbb{P}^2}$ . It is defined by the equation

$$Q_{\xi_0} : P(\mu) = \Delta,$$

so  $Q_\xi$  is obtained from  $Q_{\xi_0}$  by translation by the vector  $(-\mu(\xi), -\Delta(\xi))$ . The hypothesis that  $M(\xi)$  is positive-dimensional gives  $\Delta(\xi) > 1/2$ . The minimum value of  $P(x)$  occurs at  $P(-3/2) = -1/8$ , so  $Q_\xi$  passes below the line  $\Delta = 0$ .

Observe that every parabolic arc in the curve  $\delta(\mu) = \Delta$  is a portion of some parabola  $Q_{\xi_\alpha}$ , where  $\xi_\alpha$  denotes the Chern character of an exceptional bundle  $E_\alpha$ . If  $\xi, \zeta$  are any two linearly independent (rational) Chern characters of positive rank, then the intersection  $Q_\xi \cap Q_\zeta$  consists of a single point  $\{(\mu, \Delta)\}$  with rational coordinates, corresponding to the intersection of planes  $\xi^\perp \cap \zeta^\perp$ .

**Theorem 3.1.** *The parabola  $Q_\xi$  intersects the line  $\Delta = 1/2$  in two points. If  $\mu_0 \in \mathbb{R}$  is the larger of the two slopes such that  $(\mu_0, 1/2) \in Q_\xi$ , then there is a (unique) exceptional slope  $\alpha \in \mathcal{E}$  such that  $\mu_0 \in I_\alpha$ .*

*Proof.* We first explicitly calculate

$$\mu_0 = \frac{-3 - 2\mu(\xi) + \sqrt{5 + 8\Delta(\xi)}}{2}$$

from the formula for  $Q_\xi$ . Then either  $\mu_0$  is rational (in case  $5 + 8\Delta(\xi) \in \mathbb{Q}$  is a square), or it is a quadratic irrational. If  $\mu_0$  is rational, then  $\mu_0 \in I_\alpha$  for some  $\alpha$  by Theorem 2.2.

Suppose  $\mu_0$  is a quadratic irrational and that  $\mu_0$  is not in any  $I_\alpha$ , so that it lies in the generalized Cantor set  $C = \mathbb{R} \setminus \bigcup_\alpha I_\alpha$ . We show in the next section that any  $x \in C$  which is *not* an endpoint of some interval  $I_\alpha$  is transcendental. Hence it suffices to show

that  $\mu_0$  cannot be an endpoint. Without loss of generality, assume that  $\mu_0 = \alpha + x_\alpha$  for some  $\alpha \in \mathcal{E}$ . The point  $(\alpha + x_\alpha, 1/2)$  is the endpoint of one of the parabolic arcs in the curve  $\delta(\mu) = \Delta$ , so it lies on a parabola  $Q_{\xi_\beta}$  for some exceptional slope  $\beta$ . Then  $Q_\xi \cap Q_{\xi_\beta} = \{(\mu_0, 1/2)\}$  is a single point with rational coordinates. Therefore,  $\mu_0$  is rational, a contradiction.  $\square$

**Definition 3.2.** The exceptional slope  $\alpha$  in Theorem 3.1 is the (primary) corresponding exceptional slope to  $\xi$ .

It is now easy to describe the ray spanning the primary edge of the effective cone. The behavior of the effective cone depends on the sign of the pairing  $(\xi, \xi_\alpha) = \chi(\xi \otimes \xi_\alpha)$ . We keep the identification  $\text{NS}(M(\xi)) \cong \xi^\perp$  in mind, and recall that the primary half of this space corresponds to Chern characters of positive rank.

- (1) If  $(\xi, \xi_\alpha) > 0$ , the primary edge of the effective cone is spanned by a positive rank Chern character in  $\xi^\perp \cap (\xi_{-\alpha})^\perp$ .
- (2) If  $(\xi, \xi_\alpha) = 0$ , the primary edge of the effective cone is spanned by the orthogonal Chern character  $\xi_\alpha$ .
- (3) If  $(\xi, \xi_\alpha) < 0$ , the primary edge of the effective cone is spanned by a positive rank Chern character in  $\xi^\perp \cap (\xi_{-\alpha-3})^\perp$ .

Note that the previous intersections are also easily computed in the  $(\mu, \Delta)$ -plane as intersections of parabolas. For example, in the first case, the primary edge of the effective cone corresponds to the point  $Q_\xi \cap Q_{\xi_{-\alpha}}$ .

**Definition 3.3.** The (primary) corresponding orthogonal invariants  $(\mu^+, \Delta^+)$  to  $\xi$  are defined by

- (1)  $\{(\mu^+, \Delta^+)\} = Q_\xi \cap Q_{\xi_{-\alpha}}$  if  $(\xi, \xi_\alpha) > 0$ ,
- (2)  $(\mu^+, \Delta^+) = (\alpha, \Delta_\alpha)$  if  $(\xi, \xi_\alpha) = 0$ , and
- (3)  $\{(\mu^+, \Delta^+)\} = Q_\xi \cap Q_{\xi_{-\alpha-3}}$  if  $(\xi, \xi_\alpha) < 0$ .


A (primary) orthogonal Chern character  $\xi^+$  to  $\xi$  is defined to be any character  $\xi^+ = (r^+, \mu^+, \Delta^+)$  where  $r^+$  is sufficiently large and divisible.

We will usually drop the parenthesized word “primary” when discussing the previous constructions, as we will only briefly discuss the secondary edge of the effective cone in §7.3. With this notation, we summarize the preceding discussion with the most concise statement of the theorem we will eventually prove.


**Theorem 3.4.** In terms of the isomorphism  $\text{NS}(M(\xi)) \cong \xi^\perp$ , the primary edge of the effective cone of  $M(\xi)$  is spanned by any primary orthogonal character  $\xi^+$ .

**Remark 3.5.** It is useful to sketch the intersections defining the corresponding orthogonal invariants in the  $(\mu, \Delta)$ -plane in the three separate cases and to compare them with the classification of stable vector bundles. First, in Figure 2 we display the important features of the portion of the curve  $\delta(\mu) = \Delta$  lying over the interval  $I_\alpha$  of slopes. The curve consists of parabolic arcs  $Q_{\xi_{-\alpha}}$  and  $Q_{\xi_{-\alpha-3}}$  meeting at the vertical line  $\mu = \alpha$ .

Next, in Figure 3 we overlay the sketch of the curve  $\delta(\mu) = \Delta$  with the parabola  $Q_\xi$  in several possible arrangements.



**Fig. 2.** Schematic diagram of the portion of the curve  $\delta(\mu) = \Delta$  over the interval  $I_\alpha$  in the  $(\mu, \Delta)$ -plane.



**Fig. 3.** Possible relative positions of the curve  $\delta(\mu) = \Delta$  and the parabola  $Q_\xi$ .

- (1) We have  $(\xi, \xi_\alpha) > 0$  and  $(\mu^+, \Delta^+)$  lies on the  $\delta$ -curve.
- (1') Here we still have  $(\xi, \xi_\alpha) > 0$ , but  $(\mu^+, \Delta^+)$  does *not* lie on the  $\delta$ -curve. Correspondingly,  $\mu^+ > \alpha$ .
- (2) In this case  $Q_\xi$  passes through the Chern character  $\xi_\alpha$ , and  $(\xi, \xi_\alpha) = 0$ .
- (3) Finally, when  $(\xi, \xi_\alpha) < 0$ , the point  $(\mu^+, \Delta^+)$  always lies on the  $\delta$ -curve.

When we are in case (1), (2), or (3), we observe that  $\mu^+$  is the minimum possible slope of a *stable* Chern character  $\zeta$  orthogonal to  $\xi$  such that  $\mu(\xi \otimes \zeta) \geq 0$ . Conversely, in case (1'), there exist stable Chern characters  $\zeta$  orthogonal to  $\xi$  with  $\mu(\zeta) < \mu^+$  such that  $\mu(\xi \otimes \zeta) \geq 0$ , and such Chern characters span noneffective rays in  $\text{Pic}(M(\xi))$ .

When  $(\xi, \xi_\alpha) < 0$ , it is clear by the Intermediate Value Theorem that  $Q_\xi$  intersects the parabola  $Q_{\xi-\alpha-3}$  as in case (3) above. To verify that the schematic diagram is correct when  $(\xi, \xi_\alpha) > 0$ , we must perform a small check.

**Lemma 3.6.** *Suppose  $(\xi, \xi_\alpha) > 0$ . Then  $\mu^+ > \alpha - x_\alpha$ .*

*Proof.* Let  $\mu_0 \in \mathbb{R}$  be the larger of the two slopes such that  $(\mu_0, 1/2) \in Q_\xi$ . Then  $\mu_0 \in I_\alpha$ . Let  $(\mu_1, \Delta_1)$  and  $(\mu_2, \Delta_2)$  be the vertices of  $Q_\xi$  and  $Q_{\xi-\alpha}$ , respectively. We

compute

$$\mu_1 = -3/2 - \mu(\xi), \quad \Delta_1 = -1/8 - \Delta(\xi), \quad \mu_2 = -3/2 + \alpha, \quad \Delta_2 = -1/8 - \Delta_\alpha.$$

Let  $U$  be a slope-stable bundle of character  $\xi$ . Since  $(\xi, \xi_\alpha) > 0$ , at least one of  $h^0(E_\alpha \otimes U) > 0$  or  $h^2(E_\alpha \otimes U) > 0$  holds. However, we will see in Lemma 5.2(1) that  $\mu(E_\alpha \otimes U) > -3/2$ , and by Serre duality and stability this guarantees  $H^2(E_\alpha \otimes U) = 0$ . Thus  $h^0(E_\alpha \otimes U) > 0$ , which by stability implies  $\mu(\xi) + \alpha > 0$ , from which we conclude  $\mu_2 > \mu_1$ . Since  $\Delta(\xi) > 1/2$ , we also have  $\Delta_2 > \Delta_1$ .

The point  $p = (\alpha - x_\alpha, 1/2)$  lies on  $Q_{\xi-\alpha}$  and above  $Q_\xi$ . The vertex of  $Q_{\xi-\alpha}$  lies to the left of  $p$ . Since  $Q_\xi$  and  $Q_{\xi-\alpha}$  are translates of one another, they intersect in a single point, and the Intermedial Value Theorem shows the vertex of  $Q_{\xi-\alpha}$  lies above  $Q_\xi$ . An elementary calculation implies that  $\mu^+ > \alpha - x_\alpha$ , as in Figure 3.  $\square$

The following result is now immediate from Theorem 2.3.

**Proposition 3.7.** *The Chern character  $\xi^+$  is semistable.*

**Remark 3.8.** Case (1') of the previous remark never arises for Chern characters  $\xi$  of rank 1 (or rank 0, once the appropriate definitions have been made—see §7.2). In particular, for the Hilbert scheme of points, the primary edge of the effective cone always corresponds to the stable orthogonal Chern character of smallest positive slope [H]. It follows from results in [H] and [CH1] that if  $\xi$  is the Chern character of an ideal sheaf of  $n$  points and  $\zeta$  is a sufficiently divisible stable Chern character with  $\mu(\zeta) > 0$  such that  $(\xi, \zeta) = 0$ , then the image of the rational map

$$\mathbb{P}^{2[n]} \times M(\zeta) \dashrightarrow M(\xi \otimes \zeta), \quad (I_Z, V) \mapsto I_Z \otimes V,$$

is not contained entirely in the  $\Theta$ -divisor of sheaves with a section.

The natural analog of this statement is false in higher rank. For an example, consider the moduli space  $M(\xi) = M(r, \mu, \Delta) = M(2, 0, 11/2)$ . The associated exceptional slope is  $\alpha = 2$ , and  $(\xi, \xi_\alpha) = 1$ . Furthermore,  $\mu^+ = 9/4 > \alpha$ , so we are in case (1') of the previous remark. The point of intersection of  $Q_\xi$  with the  $\delta$ -curve has  $\mu$ -coordinate  $21/10$ . If  $21/10 \leq \mu < 9/4$  is rational, then for sufficiently divisible  $r$  we can find Chern characters  $\zeta = (r, \mu, \Delta)$  orthogonal to  $\xi$  that are stable. However, these characters correspond to noneffective divisor classes on  $M(\xi)$ . Thus, assuming the rational map  $M(\xi) \times M(\zeta) \dashrightarrow M(\xi \otimes \zeta)$  is defined, its image must land in the  $\Theta$ -divisor.

#### 4. Transcendentalty of the generalized Cantor set

Let  $C = \mathbb{R} \setminus \bigcup_{\alpha \in \mathcal{E}} I_\alpha$  be the generalized Cantor set, where  $\mathcal{E}$  is the set of exceptional slopes. In this section, we prove the following number-theoretic fact about points of  $C$ .

**Theorem 4.1.** *Let  $x$  be a point of  $C$ . Then either  $x$  is an endpoint of an interval  $I_\alpha$  and is a quadratic irrational number, or  $x$  is transcendental.*



Theorem 4.1 completes the proof of Theorem 3.1, establishing the existence of the corresponding exceptional slope. Before giving the details, we will outline the roadmap of the proof. The reader interested solely in geometry can safely skip to the next section, but we have written the argument for readers with minimal number theory background in mind.

The proof of the theorem relies on remarkable number-theoretic properties of exceptional slopes. Since the set  $C$  is stable under translation by integers  $x \mapsto x + k$  and inversion  $x \mapsto -x$ , we will restrict ourselves to  $C \cap (0, 1/2)$ . Every rational number  $q$  has an even length  $q^e$  and an odd length  $q^o$  continued fraction expansion. The even continued fraction expansion  $\beta^e$  of an exceptional slope  $0 < \beta < 1/2$  is a palindrome made of a string of ones and twos beginning and ending with a two. Furthermore, given two exceptional slopes

$$\alpha = \varepsilon\left(\frac{p-1}{2^q}\right) \quad \text{and} \quad \beta = \varepsilon\left(\frac{p+1}{2^q}\right),$$

the even partial fraction expansion of  $\gamma = \alpha.\beta = \varepsilon(p/2^q)$  is obtained by concatenating the odd partial fraction expansion of  $\beta$  with the even partial fraction expansion of  $\alpha$  with a two in between (see §4.3), i.e.

$$\gamma^e = \beta^o 2 \alpha^e.$$

Using this fact, we give an explicit algorithm for computing the continued fraction expansion of any point of the Cantor set  $C$  to arbitrary length. The crux of the matter is to show that given  $x \in C$  which is not an endpoint of  $I_\alpha$  for some  $\alpha$ , the continued fraction expansion of  $x$  is not eventually periodic (Theorem 4.16). A celebrated theorem of Euler and Lagrange [Da] asserts that a number is a quadratic irrational if and only if its continued fraction expansion is eventually periodic. This suffices to prove that  $x$  is not a quadratic irrational. To prove the stronger fact that  $x$  is transcendental, we need a theorem of Adamczewski and Bugeaud.

**Theorem 4.2** (Adamczewski–Bugeaud, [AB, Theorem 2.1]). *If the continued fraction expansion of a number  $x$  begins in arbitrarily long palindromes, then  $x$  is either quadratic irrational or transcendental.*

If  $x \in C$  is not an endpoint of some interval  $I_\alpha$ , we will show that its continued fraction expansion is not eventually periodic and begins in arbitrarily long palindromes. We conclude that  $x$  is transcendental.

We now turn to the details. At times, it will be convenient to consider certain portions of the Cantor set. Let  $C' = C \cap (-1, 1)$  and  $C'' = C \cap (0, 1/2)$  for brevity. By symmetry, it suffices to prove the theorem for  $C''$ . Similarly, we define restricted intervals of exceptional slopes  $\mathcal{E}' = \mathcal{E} \cap (-1, 1)$  and  $\mathcal{E}'' = \mathcal{E} \cap (0, 1/2)$ .

#### 4.1. The left-right sequences of points in $C'$

Recall that the order of an exceptional slope  $\alpha \in \mathcal{E}$  is the smallest natural number  $q$  such that  $\alpha = \varepsilon(p/2^q)$ . Using the order, we rewrite

$$C' = C \cap (-1, 1) = (-1, 1) \setminus \bigcup_{\alpha \in \mathcal{E}} I_\alpha = \bigcap_{q=0}^{\infty} \left( (-1, 1) \setminus \bigcup_{\substack{\alpha \in \mathcal{E} \\ \text{ord } \alpha = q}} I_\alpha \right).$$

Define

$$C_n = \bigcap_{q=0}^n \left( (-1, 1) \setminus \bigcup_{\substack{\alpha \in \mathcal{E} \\ \text{ord } \alpha = q}} I_\alpha \right),$$

setting  $C_{-1} = (-1, 1)$  for convenience. Then  $C_0$  is a union of two intervals,

$$C_0 = [-1 + x_{-1}, 0 - x_0] \cup [0 + x_0, 1 - x_1].$$

If  $x \in C$ , then  $x$  is either in the left or in the right interval of  $C_0$ . To form  $C_1$ , one removes an interior open subinterval from each component of  $C_0$ , which yields a left and right subinterval of each component of  $C_0$ . At each stage,  $C_n$  comprises  $2^{n+1}$  disjoint intervals, with each component of  $C_{n-1}$  containing two of the intervals of  $C_n$ , a “left” and a “right” subinterval. Associated to an  $x \in C'$ , we build an infinite word  $\sigma_x$  in the alphabet  $\Sigma = \{L, R\}$ ; the  $n$ th letter of  $\sigma_x$  corresponds to whether  $x$  lies in the left or right part of the component  $I \subset C_{n-1}$  containing  $x$ .

Conversely, given an infinite word  $\sigma \in \Sigma^{\mathbb{N}}$ , there is a unique  $x_\sigma \in C'$  with  $\sigma_{x_\sigma} = \sigma$ . Thus points in  $C'$  determine and are determined by left-right sequences.

#### 4.2. Action of the alphabet $\Sigma = \{L, R\}$ on $\mathcal{E}'$

Let us develop a more concrete description of the points in the set  $C'$ .

An exceptional slope  $\gamma \in \mathcal{E}$  can be uniquely written as  $\gamma = \varepsilon(p/2^q)$ , where  $q \geq 0$  and  $p$  is odd unless  $q = 0$ . Then setting

$$\alpha = \varepsilon\left(\frac{p-1}{2^q}\right) \quad \text{and} \quad \beta = \varepsilon\left(\frac{p+1}{2^q}\right)$$

we have  $\gamma = \alpha.\beta$ . We call  $\alpha$  and  $\beta$  the *left* and *right parent slopes* of  $\gamma$ , and write

$$\alpha = \text{par}_L(\gamma) \quad \text{and} \quad \beta = \text{par}_R(\gamma).$$

We define

$$\gamma \cdot L = \text{par}_L(\gamma).\gamma = \alpha.\gamma \quad \text{and} \quad \gamma \cdot R = \gamma.\text{par}_R(\gamma) = \gamma.\beta.$$

Denoting by  $\Sigma^*$  the monoid of finite strings in the alphabet  $\Sigma$ , we induce a right action of  $\Sigma^*$  on  $\mathcal{E}$ . The next easy lemma establishes that exceptional slopes can be understood either in terms of dyadic rationals via the map  $\varepsilon$  or in terms of finite left-right sequences.

**Lemma 4.3.** *There is a bijection  $\Sigma^* \rightarrow \mathcal{E}'$  given by*

$$\sigma \mapsto 0 \cdot \sigma.$$

*That is, every  $\gamma \in \mathcal{E}'$  admits a unique finite left-right sequence  $\sigma_\gamma \in \Sigma^*$  such that*

$$\gamma = 0 \cdot \sigma_\gamma.$$

In light of this correspondence, we extend the notion of parents to strings in  $\Sigma^*$ . Given  $\sigma \in \Sigma^*$ , we write  $\text{par}_L(\sigma)$  for the sequence corresponding to the left parent of  $0 \cdot \sigma$ ; we similarly define  $\text{par}_R(\sigma)$ .

The action of  $\Sigma^*$  on  $\mathcal{E}'$  can be naturally extended to a function  $\mathcal{E}' \times \Sigma^{\mathbb{N}} \rightarrow C'$ . Specifically, for  $\gamma \in \mathcal{E}'$  and  $\sigma \in \Sigma^{\mathbb{N}}$  we define  $\gamma \cdot \sigma$  to be the point in the Cantor set associated to the concatenated left-right sequence  $\sigma_\gamma \sigma$ ; in symbols,

$$\gamma \cdot \sigma = x_{\sigma_\gamma \sigma}.$$

As a special case, we note the identity

$$0 \cdot \sigma = x_\sigma.$$

This action has a natural continuity property: if  $\sigma|_n$  denotes the length  $n$  initial segment of  $\sigma$ , then

$$\gamma \cdot \sigma = \lim_{n \rightarrow \infty} \gamma \cdot \sigma|_n$$

in the real numbers. The key point is that the exceptional slope  $0 \cdot \sigma|_n$  lies in the same component of  $C_{n-1}$  as  $x_\sigma$  does for each  $n$ .

**Example 4.4.** Consider the right endpoint  $\alpha + x_\alpha$  of an interval  $I_\alpha$ , where  $\alpha \in \mathcal{E}'$ . Using a bar to denote an infinite repeating block, we have

$$\alpha + x_\alpha = \alpha \cdot R\bar{L}.$$

Similarly,

$$\alpha - x_\alpha = \alpha \cdot L\bar{R}.$$

It follows that the endpoints of the intervals  $I_\alpha$  correspond precisely to the left-right sequences  $\sigma \in \Sigma^{\mathbb{N}}$  which are *eventually constant*.

To prove the theorem, we will show that if the left-right sequence  $\sigma_x$  of an  $x \in C''$  is not eventually constant, then its continued fraction expansion is not eventually repeating.

**Remark 4.5.** If  $x \in C'' = C \cap (0, 1/2)$ , then the word  $\sigma_x \in \Sigma^{\mathbb{N}}$  begins with  $RL$ . Conversely, if  $\sigma \in \Sigma^{\mathbb{N}}$  is a word beginning with  $RL$ , then  $0 \cdot \sigma$  lies in  $C''$ . Similarly, if  $\gamma \in \mathcal{E}'' = \mathcal{E} \cap (0, 1/2)$ , then the word  $\sigma_\gamma \in \Sigma^*$  begins with  $RL$ , and any  $\sigma \in \Sigma^*$  beginning with  $RL$  has  $0 \cdot \sigma \in \mathcal{E}''$ .

**Example 4.6.** Let  $\gamma \in \mathcal{E}''$ . It is easy to describe  $\text{par}_L(\sigma_\gamma)$  and  $\text{par}_R(\sigma_\gamma)$  in terms of  $\sigma_\gamma$ . Since  $\gamma \in \mathcal{E}''$ , the sequence  $\sigma_\gamma$  begins with  $RL$ , and it is not a constant sequence. We can write either

$$\sigma_\gamma = \sigma'RL^n \quad \text{or} \quad \sigma_\gamma = \sigma'LR^n$$

for some word  $\sigma' \in \Sigma^*$  and some  $n \geq 1$ . In the first case, where  $\sigma_\gamma$  ends in  $L$ , we have

$$\text{par}_L(\sigma_\gamma) = \sigma' \quad \text{and} \quad \text{par}_R(\sigma_\gamma) = \sigma'RL^{n-1}.$$

Analogously, if  $\sigma_\gamma$  ends in  $R$ , then

$$\text{par}_L(\sigma_\gamma) = \sigma'LR^{n-1} \quad \text{and} \quad \text{par}_R(\sigma_\gamma) = \sigma'.$$

One qualitative consequence of the example is important enough to single out.

**Corollary 4.7.** *If  $\gamma \in \mathcal{E}''$ , then the parent slopes of  $\gamma$  have associated left-right sequences which are initial segments of  $\sigma_\gamma$ .*

### 4.3. Continued fractions

The continued fraction expansions of exceptional slopes have a remarkable inductive structure, which we recall here.

Given real numbers  $a_0, a_1, \dots, a_k$ , we write  $[a_0; a_1, \dots, a_k]$  for the continued fraction

$$[a_0; a_1, \dots, a_k] := a_0 + \frac{1}{a_1 + \frac{1}{\ddots + \frac{1}{a_k}}}$$

assuming this expression is defined. Any rational number  $\alpha \in \mathbb{Q}$  has two finite continued fraction expansions as above where  $a_0 \in \mathbb{Z}$  and all the  $a_i$  with  $i > 0$  are positive integers. One of them  $\alpha = [a_0; a_1, \dots, a_k]$  has  $a_k = 1$ , and the other is given by  $\alpha = [a_0; a_1, \dots, a_{k-2}, a_{k-1} + 1]$ .

For convenience, we work with the set  $\mathcal{E}''$  of exceptional slopes in the interval  $(0, 1/2)$ . If  $\alpha \in \mathcal{E}''$  and  $\alpha = [a_0; a_1, \dots, a_k]$ , then clearly  $a_0 = 0$ . It is useful to consider the string of integers  $a_1 \dots a_k \in (\mathbb{N}_{>0})^*$ . We write  $\alpha^e \in (\mathbb{N}_{>0})^*$  for the *even* length string  $a_1 \dots a_k$  of natural numbers such that  $\alpha = [0; a_1, \dots, a_k]$ . Similarly, we write  $\alpha^o \in (\mathbb{N}_{>0})^*$  for the *odd* length string with this property.

**Theorem 4.8** ([H, Theorem 3.2]). *Let  $\gamma \in \mathcal{E}'' = \mathcal{E} \cap (0, 1/2)$  be an exceptional slope, and set  $\alpha = \text{par}_L(\gamma)$  and  $\beta = \text{par}_R(\gamma)$ . Then*

$$\gamma^e = \beta^o 2 \alpha^e.$$

*That is, the even length continued fraction expansion of  $\gamma$  is obtained from the odd length expansion of  $\beta$  by concatenating a 2 and the even length expansion of  $\alpha$ .*

*Furthermore,*

- (1) *every number in  $\gamma^e$  and  $\gamma^o$  is a 1 or a 2, so that these words actually are in  $\{1, 2\}^*$ , and*
- (2) *the word  $\gamma^e$  is a palindrome.*

By Corollary 4.7, the left-right sequences of the parents of  $\gamma$  are initial segments of the left-right sequence of  $\gamma$ . This fact makes it very easy to compute the continued fraction of an exceptional slope from its left-right sequence. As a basis for the inductive algorithm, one uses that  $0^e$  is the empty string and  $(1/2)^o$  is the string 2.


**Example 4.9.** Consider the word  $RLLLLR$  and the corresponding exceptional slope  $\gamma = 0 \cdot RLLLLR$ . We compute the continued fraction expansion of  $\gamma$  by computing the continued fraction expansions of the exceptional slopes corresponding to initial segments.

$\sigma$	$\text{par}_L(\sigma)$	$\text{par}_R(\sigma)$	$(0 \cdot \sigma)^e$	$(0 \cdot \sigma)^o$
$\emptyset$	N/A	N/A	$\emptyset$	N/A
$R$	$\emptyset$	N/A	11	2
$RL$	$\emptyset$	$R$	22	211
$RLL$	$\emptyset$	$RL$	2112	21111
$RLLL$	$\emptyset$	$RLL$	211112	2111111
$RLLLR$	$RLLL$	$RLL$	211112211112	2111122111111
$RLLRR$	$RLLLR$	$RLL$	211112211112211112	2111122111122111111

We summarize our various points of view on  $\gamma$  by concluding

$$\begin{aligned} \gamma &= 0 \cdot RLLRR \\ &= \varepsilon \left( \frac{1}{2} - \frac{1}{4} - \frac{1}{8} - \frac{1}{16} + \frac{1}{32} + \frac{1}{64} \right) \\ &= [0; 2, 1, 1, 1, 1, 2, 2, 1, 1, 1, 1, 2, 2, 1, 1, 1, 2] \\ &= \frac{19760}{51641}. \end{aligned}$$

**Example 4.10.** It is also easy to systematically compute the continued fraction expansions of all exceptional slopes with order at most  $k$  for any fixed  $k$ . Figure 4 shows the computation of the slopes in  $[0, 1/2]$  up to order 5. In the diagram, the order increases from left to right. Except for those of order 5, we have recorded both the even and odd length continued fraction expansions of the exceptional slopes. We always write the odd



**Fig. 4.** The continued fraction expansions of exceptional slopes of order at most 5. Orders are listed at the top of the diagram.

length expansion above the even length expansion. To avoid cluttering the diagram, we have only recorded the even length expansions for slopes of order 5. To find the continued fraction of an exceptional slope  $\alpha$ , trace its left-right sequence along the solid lines, starting from the expansion  $\emptyset$  of 0.

#### 4.4. Repeating structures

We say a word  $\sigma = a_1 \dots a_k$  in some alphabet  $S$  has *period*  $p > 0$  if  $a_i = a_{i+p}$  for all  $1 \leq i \leq k - p$ . In this case,  $\sigma$  is the length  $k$  initial segment of the infinite word  $\overline{a_1 \dots a_p}$ .

**Lemma 4.11.** *Let  $\sigma \in S^*$  have length  $k$ , and suppose  $p$  is the smallest period of  $\sigma$ . If  $\sigma$  has another period  $p'$  and  $p + p' \leq k$ , then  $p$  divides  $p'$ .*

*Proof.* Suppose  $p$  does not divide  $p'$ , and write  $p' = pq + r$  with  $0 < r < p$ . We claim  $\sigma$  has period  $r$ , violating the minimality of  $p$ . Write  $\sigma = a_1 \dots a_k$ , and let  $i$  be an index with  $1 \leq i \leq k - r$ . There is some  $m \geq 0$  such that  $1 \leq i - mp \leq p$ . Then using the two periods of  $\sigma$ , we have

$$a_i = a_{i-mp} = a_{i-mp+p'} = a_{i-mp+p'-(q-m)p} = a_{i+r},$$

where the hypothesis  $p + p' \leq k$  ensures  $i - mp + p' \leq k$ . □

Many exceptional slopes have continued fraction expansions exhibiting a simple repeating structure.

**Proposition 4.12.** *Let  $\gamma \in \mathcal{E}''$  be an exceptional slope such that  $\sigma_\gamma$  ends in an  $R$ . Write*

$$\sigma_\gamma = \sigma'LR^n.$$

*If  $\beta = \text{par}_R(\gamma)$  and  $\alpha = \text{par}_L(\beta)$  (caution:  $\alpha$  is the parent of  $\beta$ , not of  $\gamma$ ), then*

$$\gamma^e = (\beta^o 2)^{n+1} \alpha^e.$$

*The word  $\alpha^e$  is an initial segment of  $\beta^o$ . Thus the length of  $\beta^o 2$  is a period of  $\gamma^e$ .*

*Furthermore, unless  $\beta = 1/2$ , the length of  $\beta^o 2$  is the smallest period of  $\gamma^e$ . If  $\beta = 1/2$ , then  $\gamma^e = 2^{2n+2}$ .*

*Proof.* The proof is an analysis of the continued fraction algorithm. We have  $\sigma_\beta = \sigma'$  as in Example 4.6. Observe that  $\text{par}_L(\sigma_\beta L) = \sigma_\alpha$  and  $\text{par}_R(\sigma_\beta L) = \sigma_\beta$ , while for  $n \geq 1$  we have

$$\text{par}_L(\sigma_\beta LR^n) = \sigma_\beta LR^{n-1} \quad \text{and} \quad \text{par}_R(\sigma_\beta LR^n) = \sigma_\beta.$$

Using this data one computes the continued fraction expansion of  $\gamma$  using Theorem 4.8 as in the following table.

$\sigma$	$\text{par}_L(\sigma)$	$\text{par}_R(\sigma)$	$(0 \cdot \sigma)^e$	$(0 \cdot \sigma)^o$
$\sigma_\beta$	$\sigma_\alpha$		$\beta^e$	$\beta^o$
$\sigma_\beta L$	$\sigma_\alpha$	$\sigma_\beta$	$\beta^o 2 \alpha^e$	
$\sigma_\beta LR$	$\sigma_\beta L$	$\sigma_\beta$	$(\beta^o 2)^2 \alpha^e$	
$\sigma_\beta LRR$	$\sigma_\beta LR$	$\sigma_\beta$	$(\beta^o 2)^3 \alpha^e$	
$\vdots$	$\vdots$	$\vdots$	$\vdots$	
$\sigma_\beta LR^n$	$\sigma_\beta LR^{n-1}$	$\sigma_\beta$	$(\beta^o 2)^{n+1} \alpha^e$	

Let us show that the tail  $\alpha^e$  of the continued fraction expansion of  $\gamma$  is an initial segment of  $\beta^o$ . If  $\beta = 1/2$ , then  $\alpha = 0$  and there is nothing to prove, so assume  $0 < \alpha < \beta < 1/2$ . In this case, we recall  $\beta = \alpha \cdot \text{par}_R(\beta)$ . Therefore,

$$\beta^e = (\text{par}_R(\beta))^o 2 \alpha^e.$$

The words  $\alpha^e$  and  $\beta^e$  are both palindromes (Theorem 4.8), so  $\alpha^e$  is a *proper* initial segment of  $\beta^e$ . Since  $0 < \beta < 1/2$ , the first digit in  $\beta^e$  is a 2, and thus its last digit is also a 2. Then  $\beta^o$  is obtained from  $\beta^e$  by replacing its final digit with 11, and we conclude  $\alpha^e$  is also an initial segment of  $\beta^o$ .

Suppose  $p$  is the length of the smallest period of  $\gamma^e$ . By Lemma 4.11,  $p$  divides the length of  $\beta^o 2$ . If  $p$  is smaller than this length, we can write  $\beta^o 2$  as a concatenation of  $k \geq 2$  copies of some word  $\delta$ :

$$\beta^o 2 = \delta^k.$$

Under the assumption  $0 < \beta < 1/2$ , the expansion  $\beta^e$  is a palindrome beginning and ending in 2; write

$$\beta^e = 2b_2b_3 \dots b_{l-1}2.$$

Then

$$\beta^o 2 = 2b_2b_3 \dots b_{l-1}112.$$

The word  $\delta$  must then begin with 2 and end in 112, so be of the form

$$\delta = 2d_2 \dots d_{m-3}112.$$

Defining

$$\delta' = 2d_2 \dots d_{m-3}2,$$

we have  $\beta^e = \delta^{k-1}\delta'$ . Expanding gives

$$\beta^e = (2d_2 \dots d_{m-3}\underline{112})\delta^{k-2}(2d_2 \dots d_{m-3}2).$$

Recalling that  $\beta^e$  is a palindrome, we obtain a contradiction by comparing the underlined digits in the previous expression. □

4.5. Continued fractions of points in  $C''$

Irrational numbers have unique continued fraction expansions. For an  $x \in C''$ , the expansion takes the form  $[0; a_1, a_2, \dots]$ , with all  $a_i$  in  $\{1, 2\}$  since  $x$  is a limit of exceptional slopes. We write  $x^f$  for the word  $a_1 a_2 \dots \in \{1, 2\}^{\mathbb{N}}$ .

**Lemma 4.13.** *Let  $\gamma \in \mathcal{E}''$ , and let  $\sigma \in \Sigma^*$  be nonempty.*

- (1) *If the first letter of  $\sigma$  is  $R$ , then  $\gamma^e$  is a proper initial segment of  $(\gamma \cdot \sigma)^e$ .*
- (2) *If the first letter of  $\sigma$  is  $L$ , then  $\gamma^o$  is a proper initial segment of  $(\gamma \cdot \sigma)^e$ .*

*Proof.* Induct on the length  $k$  of  $\sigma$ . If  $\sigma = R$ , then

$$(\gamma \cdot R)^e = (\gamma \cdot \text{par}_R(\gamma))^e = (\text{par}_R(\gamma))^o 2\gamma^e.$$

Since  $(\gamma \cdot R)^e$  is a palindrome,  $\gamma^e$  is a proper initial segment of  $(\gamma \cdot R)^e$ . Similarly, if  $\sigma = L$ , then  $\gamma^o$  is a proper initial segment of  $(\gamma \cdot L)^e$ .

For the inductive step, without loss of generality assume  $\sigma$  begins with  $R$  and has length  $k + 1$ . Let  $\sigma' = \sigma|_k$  be the length  $k$  initial segment of  $\sigma$ . By induction,  $\gamma^e$  is a proper initial segment of  $(\gamma \cdot \sigma')^e$ . Furthermore, either  $(\gamma \cdot \sigma')^e$  or  $(\gamma \cdot \sigma')^o$  is a proper initial segment of  $(\gamma \cdot \sigma)^e$  by the previous paragraph. As  $\gamma^e$  is a proper initial segment of  $(\gamma \cdot \sigma')^e$ , it is also a proper initial segment of  $(\gamma \cdot \sigma')^o$ . Thus in either case,  $\gamma^e$  is a proper initial segment of  $(\gamma \cdot \sigma)^e$ . □

We then obtain the following result by continuity. The continued fraction expansion of  $x \in C''$  can be understood in terms of the fractions of exceptional slopes approximating it.

**Corollary 4.14.** *Let  $x \in C''$ , and let  $\sigma_x$  be the left-right sequence corresponding to  $x$ . Let  $\sigma'$  be an initial segment of  $\sigma_x$ , and let  $\gamma = 0 \cdot \sigma'$ .*

- (1) *If  $\sigma'$  precedes an  $R$  in  $\sigma$ , then  $\gamma^e$  is an initial segment of  $x^f$ .*
- (2) *If  $\sigma'$  precedes an  $L$  in  $\sigma$ , then  $\gamma^o$  is an initial segment of  $x^f$ .*

Since even length expansions of exceptional slopes are palindromes, we deduce the next corollary.

**Corollary 4.15.** *If  $x \in C''$  is not an endpoint of an interval  $I_\alpha$ , then  $x^f$  begins in arbitrarily long palindromes.*

Having studied the continued fractions of exceptional slopes in great detail, the next result is now easy.

**Theorem 4.16.** *If  $x \in C''$  has a left-right sequence  $\sigma_x$  which is not eventually constant, then  $x^f$  is not eventually repeating. Thus  $x$  is not a quadratic irrational.*

*Proof.* Assume  $x^f$  is eventually repeating with period  $p$ , beginning from the  $c$ th digit of  $x^f$ . Fix  $N$  very large relative to  $p$  and  $c$ . Since  $\sigma_x$  is not eventually constant, we may find an initial segment  $\sigma'$  of  $\sigma_x$  which ends in  $R$  such that  $\sigma'$  and the parents of  $\sigma'$  have



length at least  $N$ . Correspondingly, let  $\alpha$  and  $\beta$  be the parents of  $\gamma = 0 \cdot \sigma'$ , so that  $\gamma = \alpha \cdot \beta$ . As in Proposition 4.12, we have

$$\gamma^e = (\beta^o 2)^n \alpha^e$$

for some  $n \geq 2$ . Then  $\beta^o$  and  $\alpha^e$  have length at least  $N$ . Either  $\gamma^e$  or  $\gamma^o$  is an initial segment of  $x^f$  by Corollary 4.14, so  $x^f$  begins like

$$x^f = \beta^o 2 \beta^o 2 \dots$$

Let  $p' > N$  be the length of  $\beta^o 2$ . Writing  $x^f = a_1 a_2 \dots$ , consider the word

$$\sigma = a_c a_{c+1} \dots a_{2p'}$$

By construction,  $\sigma$  has periods  $p$  and  $p'$ , and  $\sigma$  has length more than  $p + p'$ . Lemma 4.11 shows that  $p$  is in fact the shortest period of  $\sigma$ . Again by Lemma 4.11,  $p$  divides  $p'$  and the word  $\beta^o 2$  is a power of a shorter word. As in the proof of Proposition 4.12, this is a contradiction.  $\square$

Theorem 4.1 follows at once from Corollary 4.15, Theorem 4.16 and Theorem 4.2.

### 5. The Beilinson spectral sequence

In this section, we use the generalized Beilinson spectral sequence to describe a canonical resolution of the general sheaf in a moduli space  $M(\xi)$ . This resolution illuminates the structure of the effective cone.

#### 5.1. Generalities on the Beilinson spectral sequence

Following [D2, D3], a *triad* is a triple  $(E, G, F)$  of exceptional bundles where the slopes are of the form  $(\alpha, \alpha \cdot \beta, \beta)$ ,  $(\beta - 3, \alpha, \alpha \cdot \beta)$ , or  $(\alpha \cdot \beta, \beta, \beta + 3)$  for some exceptional slopes  $\alpha, \beta$  of the form

$$\alpha = \varepsilon \left( \frac{p}{2q} \right), \quad \beta = \varepsilon \left( \frac{p+1}{2q} \right). \tag{*}$$

Corresponding to the triad  $(E, G, F)$  is a fourth exceptional bundle  $M$  defined as the cokernel of the canonical map  $\text{ev}^* : G \rightarrow F \otimes \text{Hom}(G, F)^*$ . The collection  $(E^*(-3), M^*, F^*)$  is another triad.

The bundles of a triad form a *strong exceptional collection* for the bounded derived category  $D^b(\mathbb{P}^2)$ . In other words, if  $A, B$  are two bundles in a triad with  $A$  listed before  $B$ , then  $\text{Ext}^i(A, B) = 0$  for  $i > 0$  and  $\text{Ext}^i(B, A) = 0$  for all  $i$ . Furthermore, a triad is a full collection, in the sense that it generates the derived category. This fact is a consequence of the generalized Beilinson spectral sequence, which we now recall.

**Theorem 5.1** ([D2]). *Let  $U$  be a coherent sheaf, and let  $(E, G, F)$  be a triad. Write*

$$\begin{aligned} G_{-2} &= E, & F_{-2} &= E^*(-3), \\ G_{-1} &= G, & F_{-1} &= M^*, \\ G_0 &= F, & F_0 &= F^*, \end{aligned}$$

and set  $G_i = F_i = 0$  if  $i \notin \{-2, -1, 0\}$ . There is a spectral sequence with  $E_1$  page

$$E_1^{p,q} = G_p \otimes H^q(U \otimes F_p)$$

which converges to  $U$  in degree 0 and to 0 in all other degrees.

In fact, every full strong exceptional collection for  $D^b(\mathbb{P}^2)$  is a triad. Gorodentsev and Rudakov [GR] studied exceptional bundles on projective space contemporaneously with Drézet, and in their language every *helix* on  $\mathbb{P}^2$  is *constructive*; i.e. any full strong exceptional collection is obtained from  $(\mathcal{O}, \mathcal{O}(1), \mathcal{O}(2))$  by a sequence of *mutations*.

Drézet [D1, Question 4] remarks that by choosing a triad carefully it is possible to obtain interesting information about a sheaf  $U$ . For example, to solve the existence problem for  $M(\xi)$ , one chooses a triad related to the exceptional bundle  $E_\alpha$  such that  $\mu(\xi) \in I_\alpha$ . He also asks if there is a “best choice” for the Beilinson spectral sequence. Our study fits this theme: to understand  $\text{Eff}(M(\xi))$ , we will choose a triad related to the corresponding exceptional bundle.

### 5.2. Choosing the triad

Since the map  $\varepsilon : \mathbb{Z}[1/2] \rightarrow \mathcal{E}$  is a bijection, the corresponding exceptional slope to  $\xi$  (see Definition 3.2) can be written uniquely in the form  $\alpha.\beta$ , where  $\alpha, \beta$  are as in equation (\*). Denote by  $U$  a general sheaf in  $M(\xi)$ . As discussed in §3, the qualitative behavior of the effective cone depends on the sign of  $(\xi, \xi_{\alpha.\beta})$ . Which triad we use to decompose  $U$  depends on the sign of this pairing as well.

- (1) If  $(\xi, \xi_{\alpha.\beta}) > 0$ , we decompose  $U$  according to the triad  $(E_{-\alpha-3}, E_{-\beta}, E_{-(\alpha.\beta)})$ .
- (2) If  $(\xi, \xi_{\alpha.\beta}) < 0$ , we use the triad  $(E_{-(\alpha.\beta)-3}, E_{-\alpha-3}, E_{-\beta})$ .
- (3) When  $(\xi, \xi_{\alpha.\beta}) = 0$ , the spectral sequence from either of the previous two triads degenerates and gives the same result.

We will primarily focus on the first case in this section, briefly summarizing the results in the other two cases at the end.

### 5.3. The spectral sequence when $(\xi, \xi_{\alpha.\beta}) > 0$

Assuming  $(\xi, \xi_{\alpha.\beta}) > 0$ , we aim to describe the Beilinson spectral sequence of the general  $U \in M(\xi)$  with respect to the triad  $(E_{-\alpha-3}, E_{-\beta}, E_{-(\alpha.\beta)})$ . In the notation of Theorem 5.1, we have

$$\begin{aligned} G_{-2} &= E_{-\alpha-3}, & F_{-2} &= E_\alpha, \\ G_{-1} &= E_{-\beta} & F_{-1} &= E_{\alpha.(\alpha.\beta)}, \\ G_0 &= E_{-(\alpha.\beta)} & F_0 &= E_{\alpha.\beta}. \end{aligned}$$

(To compute  $F_{-1}$ , notice that the only exceptional bundle  $F_{-1}$  such that  $(E_\alpha, F_{-1}, E_{\alpha,\beta})$  is a triad is  $F_{-1} = E_{\alpha,(\alpha,\beta)}$ .) Then the associated spectral sequence has the following first page:

$$\begin{aligned} E_{-\alpha-3} \otimes H^2(U \otimes E_\alpha) &\rightarrow E_{-\beta} \otimes H^2(U \otimes E_{\alpha,(\alpha,\beta)}) \rightarrow E_{-(\alpha,\beta)} \otimes H^2(U \otimes E_{\alpha,\beta}), \\ E_{-\alpha-3} \otimes H^1(U \otimes E_\alpha) &\rightarrow E_{-\beta} \otimes H^1(U \otimes E_{\alpha,(\alpha,\beta)}) \rightarrow E_{-(\alpha,\beta)} \otimes H^1(U \otimes E_{\alpha,\beta}), \\ E_{-\alpha-3} \otimes H^0(U \otimes E_\alpha) &\rightarrow E_{-\beta} \otimes H^0(U \otimes E_{\alpha,(\alpha,\beta)}) \rightarrow E_{-(\alpha,\beta)} \otimes H^0(U \otimes E_{\alpha,\beta}). \end{aligned}$$

Many of these terms typically vanish for a given  $U$ . To see which vanishings we can expect, we perform a numerical calculation.

**Lemma 5.2.** *With the notation of this subsection, the following inequalities hold:*

- (1)  $\mu(U \otimes E_\alpha) > -3/2$ ,
- (2)  $\chi(U \otimes E_\alpha) < 0$ ,
- (3)  $\chi(U \otimes E_{\alpha,(\alpha,\beta)}) < 0$ .

*Proof.* (1) The reference parabola  $Q_{\xi_0}$  corresponding to  $\mathcal{O}_{\mathbb{P}^2}$  passes through the points  $(0, 1)$  and  $(-3, 1)$  in the  $(\mu, \Delta)$ -plane. It follows that any parabola obtained by translating  $Q_{\xi_0}$  downward by more than  $1/2$  intersects the line  $\Delta = 1/2$  in two points more than three units apart.

Let  $\mu_0 \in \mathbb{R}$  be the larger of the two slopes such that  $(\mu_0, 1/2) \in Q_\xi$ , and let  $(\mu_1, \Delta_1) \in \mathbb{Q}^2$  be the vertex of  $Q_\xi$ . The parabola  $Q_\xi$  is obtained from  $Q_{\xi_0}$  by translation by  $(-\mu(\xi), -\Delta(\xi))$ . Since  $\Delta(\xi) > 1/2$ , the previous paragraph shows that

$$\mu_0 - \mu_1 > 3/2.$$

The vertex of  $Q_{\xi_0}$  occurs at  $(-3/2, -1/8)$ , so  $\mu_1 = -\mu(\xi) - 3/2$ . By construction of the corresponding exceptional slope,  $\mu_0 \in I_{\alpha,\beta}$ , so  $\mu_0 < \alpha.\beta + x_{\alpha,\beta}$ . Then

$$\begin{aligned} \mu(U \otimes E_\alpha) &= \mu(\xi) + \alpha = -3/2 - \mu_1 + \alpha > \alpha - \mu_0 > \alpha - (\alpha.\beta + x_{\alpha,\beta}) \\ &\geq (\sqrt{5} - 5)/2 \approx -1.38 > -3/2, \end{aligned}$$

with the worst case of the “ $\geq$ ” occurring when  $\alpha$  and  $\alpha.\beta$  are consecutive integers.

(2 & 3) Consider the parabola  $Q_{\xi-(\alpha,\beta)}$  giving the arc of the curve  $\delta(\mu) = \Delta$  over the interval  $[\alpha.\beta - x_{\alpha,\beta}, \alpha.\beta]$ . The parabola  $Q_\xi$  intersects  $Q_{\xi-(\alpha,\beta)}$  at some point  $(\mu_2, \Delta_2)$ , and  $\mu_2 > \alpha.\beta - x_{\alpha,\beta}$  by Lemma 3.6. We have  $\mu_0 < \mu_2$ , so to the left of  $\mu_2$  the parabola  $Q_{\xi-(\alpha,\beta)}$  lies above the parabola  $Q_\xi$ . Therefore, if  $\zeta$  is any Chern character with invariants on  $Q_{\xi-(\alpha,\beta)}$  such that  $\mu(\zeta) < \mu_2$ , we observe  $(\xi, \zeta) < 0$ .

The invariants of  $E_\alpha$  and  $E_{\alpha,(\alpha,\beta)}$  both lie on  $Q_{\xi-(\alpha,\beta)}$  since  $\chi(E_{\alpha,\beta}, E_\alpha) = \chi(E_{\alpha,(\alpha,\beta)}, E_\alpha) = 0$ . The required inequalities then follow from the previous paragraph.  $\square$

As a consequence of Lemma 5.2(1) and semistability, the  $H^2$ -terms in the spectral sequence vanish for any  $U \in M(\xi)$ . We will show that for the general  $U \in M(\xi)$  the

$H^0$ - and  $H^1$ -terms are governed by the Euler characteristic. That is, the bundles  $U \otimes E_\alpha$ ,  $U \otimes E_{\alpha,(\alpha,\beta)}$ , and  $U \otimes E_{\alpha,\beta}$  are all nonspecial, so that the spectral sequence takes the form

$$\begin{aligned} E_{-\alpha-3} \otimes \mathbb{C}^{m_1} &\rightarrow E_{-\beta} \otimes \mathbb{C}^{m_2} \rightarrow 0, \\ 0 &\rightarrow 0 \rightarrow E_{-(\alpha,\beta)} \otimes \mathbb{C}^{m_3}, \end{aligned}$$

where

$$m_1 = -\chi(U \otimes E_\alpha), \quad m_2 = -\chi(U \otimes E_{\alpha,(\alpha,\beta)}), \quad m_3 = \chi(U \otimes E_{\alpha,\beta}).$$

**Proposition 5.3.** *With the notation of this subsection, if  $U \in M(\xi)$  is general, then the Beilinson spectral sequence degenerates as above.*

*More precisely, the general  $U \in M(\xi)$  admits a resolution of the form*

$$0 \rightarrow E_{-\alpha-3}^{m_1} \xrightarrow{\psi} E_{-\beta}^{m_2} \oplus E_{-(\alpha,\beta)}^{m_3} \rightarrow U \rightarrow 0.$$

*Conversely, a general sheaf with this resolution is semistable.*

*Proof.* Consider a sheaf  $U$  defined as the cokernel of a map

$$E_{-\alpha-3}^{m_1} \xrightarrow{\psi} E_{-\beta}^{m_2} \oplus E_{-(\alpha,\beta)}^{m_3} \rightarrow U \rightarrow 0,$$

with  $\psi \in \text{Hom}(E_{-\alpha-3}^{m_1}, E_{-\beta}^{m_2} \oplus E_{-(\alpha,\beta)}^{m_3})$  general.

*Step 1:*  $\text{ch}(U) = \xi$  and  $\psi$  is injective. Since we do not know that  $\psi$  is injective, we first compute the Chern character of the mapping cone of  $\psi$  in the derived category. The convergence of the Beilinson spectral sequence provides the necessary equality

$$\xi = -m_1 \text{ch}(E_{-\alpha-3}) + m_2 \text{ch}(E_{-\beta}) + m_3 \text{ch}(E_{-\alpha,\beta}).$$

To see that  $\psi$  is injective, we first recall a Bertini-type theorem [H, Proposition 2.6]: if  $E$  and  $F$  are vector bundles of ranks  $m, n$  on a smooth variety  $X$  such that  $\mathcal{H}om(E, F)$  is globally generated, then for a general  $\psi \in \text{Hom}(E, F)$  the degeneracy locus  $D_k(\psi) = \{x \in X : \text{rk } \psi_x \leq k\}$  is either empty or has the expected codimension  $(m - k)(n - k)$ .

In our present circumstances, the sheaves

$$\mathcal{H}om(E_{-\alpha-3}, E_{-\beta}) \quad \text{and} \quad \mathcal{H}om(E_{-\alpha-3}, E_{-(\alpha,\beta)})$$

are both globally generated [H, Lemma 5.4], from which we deduce that

$$\mathcal{H}om(E_{-\alpha-3}^{m_1}, E_{-\beta}^{m_2} \oplus E_{-(\alpha,\beta)}^{m_3})$$

is globally generated as well. Since  $\text{rk } \xi \geq 1$ , the Bertini-type theorem implies that  $\psi_x$  is injective for general  $x \in \mathbb{P}^2$ . Thus  $\psi$  is injective as a map of sheaves.

*Step 2:*  $U$  has the expected zeroes in its Beilinson spectral sequence. It is enough to check

$$H^0(U \otimes E_\alpha) = H^0(U \otimes E_{\alpha,(\alpha,\beta)}) = H^1(U \otimes E_{\alpha,\beta}) = 0,$$

and these vanishings follow immediately from orthogonality properties of exceptional bundles. For instance,  $\mathcal{H}om(E_\beta, E_\alpha)$  and  $\mathcal{H}om(E_{\alpha,\beta}, E_\alpha)$  both have no cohomology, while  $\text{Ext}^1(E_{\alpha+3}, E_\alpha) \cong \text{Ext}^1(E_\alpha, E_\alpha)^* = 0$  since exceptional bundles are rigid. This gives  $H^0(U \otimes E_\alpha) = 0$ ; the other vanishings are proved similarly.

*Step 3:  $U$  is stable.* For brevity, write  $A = E_{-\alpha-3}^{m_1}$  and  $B = E_{-\beta}^{m_2} \oplus E_{-(\alpha,\beta)}^{m_3}$ . Let

$$S \subset \text{Hom}(A, B)$$

be the open subset parameterizing injective sheaf maps with torsion-free cokernels.

The set  $S$  is in fact nonempty; to see this we show the general cokernel  $U$  is torsion-free. The Bertini-type theorem shows the map  $\psi_x$  on fibers is an injection outside of codimension 2 in case  $r(\xi) = 1$  and an injection everywhere if  $r(\xi) > 1$ . Hence, any torsion in  $U$  occurs in codimension at least 2. No sheaf with torsion in codimension 2 can be a cokernel of an injective map of locally free sheaves, so in fact  $U$  must be torsion-free in this case as well.

Consider the family  $\mathcal{U}/S$  of quotients parameterized by  $S$ . We claim that  $\mathcal{U}$  is a complete family of *prioritary* sheaves. A *prioritary* sheaf is a torsion-free sheaf  $U$  such that

$$\text{Ext}^2(U, U(-1)) = 0.$$

Drézet [D2] shows that if  $\alpha < \beta$  are two exceptional slopes then  $\text{Ext}^i(E_\alpha, E_\beta) = 0$  for  $i > 0$ . From this vanishing theorem, it follows easily that the sheaves parameterized by  $S$  are all *prioritary*.

To see  $\mathcal{U}/S$  is a complete family, fix an  $s \in S$ , let  $U_s$  be the corresponding *prioritary* sheaf, and consider the Kodaira–Spencer map

$$\text{Hom}(A, B) = T_s S \rightarrow \text{Ext}^1(U_s, U_s);$$

we must show this map is surjective. The Kodaira–Spencer map factors as the composition

$$\text{Hom}(A, B) \xrightarrow{i} \text{Hom}(A, U_s) \xrightarrow{j} \text{Ext}^1(U_s, U_s),$$

where the maps come from the natural long exact sequences

$$\begin{aligned} \text{Hom}(A, B) &\xrightarrow{i} \text{Hom}(A, U_s) \rightarrow \text{Ext}^1(A, A), \\ \text{Hom}(A, U_s) &\xrightarrow{j} \text{Ext}^1(U_s, U_s) \rightarrow \text{Ext}^1(B, U_s). \end{aligned}$$

From the exact sequence

$$\text{Ext}^1(B, B) \rightarrow \text{Ext}^1(B, U_s) \rightarrow \text{Ext}^2(B, A)$$

and the orthogonality properties of exceptional bundles, we conclude  $\text{Ext}^1(A, A) = \text{Ext}^1(B, U_s) = 0$ , so  $i, j$ , and the Kodaira–Spencer map are all surjective.

The Artin stack  $\mathcal{P}(\xi)$  of *prioritary* sheaves with Chern character  $\xi$  is an irreducible stack containing the stack  $\mathcal{M}(\xi)$  of *semistable* sheaves with Chern character  $\xi$  as a dense open subset [HiL]. It follows that the general sheaf in  $\mathcal{M}(\xi)$  is parameterized by a point of  $S$ . □

From the perspective of Bridgeland stability, it is better to interpret the resolution of a general  $U \in M(\xi)$  discussed in the previous proposition from the viewpoint of the derived category. We first record a well-known lemma which will be important throughout the rest of the paper.

**Lemma 5.4.** *Consider a pair of two-term complexes*

$$\begin{aligned} W &= E_{-\alpha-3} \otimes \mathbb{C}^{m_1} \xrightarrow{d} E_{-\beta} \otimes \mathbb{C}^{m_2}, \\ W' &= E_{-\alpha-3} \otimes \mathbb{C}^{m'_1} \xrightarrow{d'} E_{-\beta} \otimes \mathbb{C}^{m'_2}, \end{aligned}$$

each sitting in degrees  $-1$  and  $0$ . Every homomorphism  $W \rightarrow W'$  in the derived category  $D^b(\mathbb{P}^2)$  is realized by a homomorphism of complexes, so

$$\mathrm{Hom}_{D^b(\mathbb{P}^2)}(W, W') = \mathrm{Hom}_{\mathrm{Kom}(\mathbb{P}^2)}(W, W').$$

*Proof.* We consider the second hyperext spectral sequence which has  $E_1$  page

$$E_1^{i,j} = \prod_p \mathrm{Ext}^j(\mathcal{K}_1^p, \mathcal{K}_2^{p+i})$$

with

$$E_1^{i,j} \Rightarrow \mathrm{Ext}^{i+j}(\mathcal{K}_1^\bullet, \mathcal{K}_2^\bullet)$$

which comes from taking an injective resolution of  $\mathcal{K}_2^\bullet$  and using the two spectral sequences coming from a double complex. Applying this to the complexes  $W$  and  $W'$ , we get

$$\begin{aligned} E_1^{0,0} &= \mathrm{Hom}(E_{-\alpha-3} \otimes \mathbb{C}^{m_1}, E_{-\alpha-3} \otimes \mathbb{C}^{m'_1}) \oplus \mathrm{Hom}(E_{-\beta} \otimes \mathbb{C}^{m_2}, E_{-\beta} \otimes \mathbb{C}^{m'_2}), \\ E_1^{1,0} &= \mathrm{Hom}(E_{-\alpha-3} \otimes \mathbb{C}^{m_1}, E_{-\beta} \otimes \mathbb{C}^{m'_2}). \end{aligned}$$

The map  $E_1^{0,0} \rightarrow E_1^{1,0}$  is  $(f, g) \mapsto -d' \circ f + g \circ d$ , so we can identify the kernel with  $\mathrm{Hom}_{\mathrm{Kom}(\mathbb{P}^2)}(W, W')$ . On the other hand, the spectral sequence degenerates at the  $E_2$  page, so we can also identify the kernel with  $\mathrm{Hom}_{D^b(\mathbb{P}^2)}(W_1, W_2)$ .  $\square$

**Theorem 5.5.** *With the notation of this subsection, let  $U \in M(\xi)$  be general. Let  $W \in D^b(\mathbb{P}^2)$  be the mapping cone of the canonical evaluation map*

$$E_{-(\alpha,\beta)} \otimes \mathrm{Hom}(E_{-(\alpha,\beta)}, U) \rightarrow U,$$

so that there is a distinguished triangle

$$E_{-(\alpha,\beta)} \otimes \mathrm{Hom}(E_{-(\alpha,\beta)}, U) \rightarrow U \rightarrow W \rightarrow \cdot.$$

Then  $W$  is isomorphic to a complex of the form

$$E_{-\alpha-3} \otimes \mathbb{C}^{m_1} \rightarrow E_{-\beta} \otimes \mathbb{C}^{m_2}$$

sitting in degrees  $-1$  and  $0$ . Any two complexes of this form which are isomorphic to  $W$  are in the same orbit under the natural action of  $\mathrm{GL}(m_1) \times \mathrm{GL}(m_2)$  on the space of such complexes.

Furthermore,  $W$  is also isomorphic to the complex  $E_{-\alpha-3} \otimes \mathbb{C}^{m_1} \rightarrow E_{-\beta} \otimes \mathbb{C}^{m_2}$  appearing in the Beilinson spectral sequence for  $U$ .

*Proof.* It is easy to show that if

$$0 \rightarrow A \rightarrow B \oplus C \rightarrow D \rightarrow 0$$

is an exact sequence of sheaves, then the mapping cone of  $C \rightarrow D$  is isomorphic to the complex  $A \rightarrow B$  sitting in degrees  $-1$  and  $0$ . Suppose we have a resolution

$$0 \rightarrow E_{-\alpha-3}^{m_1} \xrightarrow{\psi} E_{-\beta}^{m_2} \oplus E_{-(\alpha,\beta)}^{m_3} \rightarrow U \rightarrow 0$$

of a general  $U$  as in Proposition 5.3. Then  $m_3 = \text{hom}(E_{-(\alpha,\beta)}, U)$ , so since  $U$  is semistable the map  $E_{-(\alpha,\beta)}^{m_3} \rightarrow U$  can be identified with the canonical evaluation  $E_{-(\alpha,\beta)} \otimes \text{Hom}(E_{-(\alpha,\beta)}, U) \rightarrow U$ . Thus the mapping cone of this evaluation is the complex given by the first component of  $\psi$ . By Lemma 5.4, any two complexes of the form  $E_{-\alpha-3}^{m_1} \rightarrow E_{-\beta}^{m_2}$  which are isomorphic in the derived category are in the same orbit of the  $\text{GL}(m_1) \times \text{GL}(m_2)$  action.

Finally, we show that  $W$  is isomorphic to the complex which appears in the Beilinson spectral sequence for  $U$ . We recall how to compute the  $E_1$  page of the spectral sequence. Let  $p_i : \mathbb{P}^2 \times \mathbb{P}^2 \rightarrow \mathbb{P}^2$  be the projections, and let  $\Delta \subset \mathbb{P}^2 \times \mathbb{P}^2$  be the diagonal. There is a resolution of the diagonal

$$0 \rightarrow E_{-\alpha-3} \boxtimes E_\alpha \rightarrow E_{-\beta} \boxtimes E_{\alpha,(\alpha,\beta)} \rightarrow E_{-(\alpha,\beta)} \boxtimes E_{\alpha,\beta} \rightarrow \mathcal{O}_\Delta \rightarrow 0.$$

The complex appearing in the Beilinson spectral sequence is then the natural map

$$R^1 p_{1*}(E_{-\alpha-3} \boxtimes (E_\alpha \otimes U)) \rightarrow R^1 p_{1*}(E_{-\beta} \boxtimes (E_{\alpha,(\alpha,\beta)} \otimes U)),$$

which, by the projection formula, is a map

$$E_{-\alpha-3} \otimes H^1(E_\alpha \otimes U) \rightarrow E_{-\beta} \otimes H^1(E_{\alpha,(\alpha,\beta)} \otimes U).$$

We split the resolution of the diagonal into two short exact sequences

$$\begin{aligned} 0 \rightarrow E_{-\alpha-3} \boxtimes E_\alpha \rightarrow E_{-\beta} \boxtimes E_{\alpha,(\alpha,\beta)} \rightarrow M \rightarrow 0, \\ 0 \rightarrow M \rightarrow E_{-(\alpha,\beta)} \boxtimes E_{\alpha,\beta} \rightarrow \mathcal{O}_\Delta \rightarrow 0. \end{aligned}$$

Tensoring with  $p_2^*(U)$  and applying  $Rp_{1*}$ , we get triangles

$$\begin{aligned} \Phi_{E_{-\alpha-3} \boxtimes E_\alpha}(U) \rightarrow \Phi_{E_{-\beta} \boxtimes E_{\alpha,(\alpha,\beta)}}(U) \rightarrow \Phi_M(U) \rightarrow \cdot, \\ \Phi_M(U) \rightarrow \Phi_{E_{-(\alpha,\beta)} \boxtimes E_{\alpha,\beta}}(U) \rightarrow \Phi_{\mathcal{O}_\Delta}(U) \rightarrow \cdot, \end{aligned}$$

where  $\Phi_F : D^b(\mathbb{P}^2) \rightarrow D^b(\mathbb{P}^2)$  is the Fourier–Mukai transform with kernel  $F$ . Computing these transforms using Proposition 5.3, we obtain two different triangles

$$\begin{aligned} E_{-\alpha-3} \otimes H^1(E_\alpha \otimes U) \rightarrow E_{-\beta} \otimes H^1(E_{\alpha,(\alpha,\beta)} \otimes U) \rightarrow \Phi_M(U)[1] \rightarrow \cdot, \\ \Phi_M(U) \rightarrow E_{-\alpha,\beta} \otimes \text{Hom}(E_{-\alpha,\beta}, U) \rightarrow U \rightarrow \cdot \end{aligned}$$

involving  $\Phi_M(U)$ ; notice that the map  $E_{-\alpha,\beta} \otimes \text{Hom}(E_{-\alpha,\beta}, U) \rightarrow U$  is the canonical one since the map  $E_{-(\alpha,\beta)} \boxtimes E_{\alpha,\beta} \rightarrow \mathcal{O}_\Delta$  is the trace map. Therefore  $\Phi_M(U)[1]$  is isomorphic to  $W$  by the second triangle. On the other hand,  $\Phi_M(U)[1]$  is also isomorphic to the complex in the Beilinson spectral sequence by the first triangle.  $\square$

**Remark 5.6.** For experts on Bridgeland stability, we remark that the resolution we have found arises from Bridgeland stability. We refer the reader to [H, §8 and §9] for the necessary definitions and results. We use the notation in that paper. Briefly, for every  $s \in \mathbb{R}$ , let  $\mathcal{F}_s$  (resp.,  $\mathcal{Q}_s$ ) denote the full subcategory of sheaves whose Harder–Narasimhan filtration with respect to the Mumford slope has maximal (resp., minimal) slope at most (resp., greater than)  $s$ . Then, for every  $s, t$  with  $s \in \mathbb{R}$  and  $t > 0$ , there exists a Bridgeland stability condition  $\sigma_{s,t} = (\mathcal{A}_s, Z_{s,t})$  on  $\mathbb{P}^2$ , where

$$\mathcal{A}_s := \{E^\bullet \in \mathcal{D}^b(\mathbb{P}^2) : H^{-1}(E^\bullet) \in \mathcal{F}_s, H^0(E^\bullet) \in \mathcal{Q}_s, H^i(E^\bullet) = 0 \text{ if } i \neq 0, -1\},$$

$$Z_{s,t}(E) := - \int_{\mathbb{P}^2} e^{-(s+it)L} \text{ch}(E),$$

and  $L$  denotes the hyperplane class. Given a fixed Chern character  $\xi$ , the  $st$ -plane has a wall and chamber decomposition, where the Bridgeland walls in  $s < \mu(\xi)$  are finitely many nested semicircles [ABCH] with centers along the  $s$ -axis. With this notation, the precise theorem is as follows.

**Theorem 5.7.** *Let  $U \in M(\xi)$  be general. Let  $\Lambda = W(U, E_{-(\alpha,\beta)})$  be the wall in the  $st$ -plane of stability conditions where  $U$  and  $E_{-(\alpha,\beta)}$  have the same  $\mu_{s,t}$ -slope. There is an exact sequence*

$$0 \rightarrow E_{-(\alpha,\beta)} \otimes \text{Hom}(E_{-(\alpha,\beta)}, U) \rightarrow U \rightarrow W \rightarrow 0$$

in the corresponding categories  $\mathcal{A}_s$ . The kernel and cokernel objects are  $\sigma_{s,t}$ -semistable outside of walls nested inside  $\Lambda$ , so  $U$  is  $\sigma_{s,t}$ -semistable along and outside  $\Lambda$ , but not stable inside  $\Lambda$ . The wall  $\Lambda$  has center  $(s_0, 0)$  satisfying  $s_0 = -\mu^+ - 3/2$ , where  $\mu^+$  is the corresponding orthogonal slope to  $\xi$ .

*Proof.* Since  $U$  is Gieseker stable, it is  $\sigma_{s,t}$ -stable whenever  $s < \mu(\xi)$  and  $t \gg 0$ . If we show that  $U$  is  $\sigma_{s,t}$ -semistable along  $\Lambda$ , then it follows that  $U$  is  $\sigma_{s,t}$ -semistable everywhere outside  $\Lambda$ . Indeed, if  $0 \rightarrow F \rightarrow U$  destabilizes  $U$  along some wall outside  $\Lambda$ , then this would violate either the Gieseker semistability of  $U$  or the  $\sigma_{s,t}$ -semistability of  $U$  along  $\Lambda$ .

As in [H, §9], one shows that  $E_{-(\alpha,\beta)}$  and  $W$  are  $\sigma_{s,t}$ -semistable for  $(s, t)$  outside semicircular walls of radius at most  $\sqrt{5}/2$ . Observe that if  $E_{-(\alpha,\beta)}$  is destabilized along some wall  $\Lambda'$  then either  $\Lambda$  is nested inside  $\Lambda'$  or  $\Lambda'$  is nested inside  $\Lambda$ , since both  $\Lambda$  and  $\Lambda'$  belong to a common foliation of the  $st$ -plane by semicircles. The same argument applies to  $W$ . We will prove that  $\Lambda$  has radius at least  $\sqrt{5}/2$ , and therefore  $E_{-(\alpha,\beta)}$  and  $W$  are  $\sigma_{s,t}$ -semistable along  $\Lambda$ .

Since  $(\xi, \xi_{\alpha,\beta}) > 0$ , the orthogonal Chern character  $\xi^+$  satisfies

$$(\xi^+, \xi) = (\xi^+, \xi_{-(\alpha,\beta)}) = 0$$

(see Remark 3.5). Then by [CH1, Proposition 4.1], the wall  $\Lambda$  is the semicircular wall with center  $(s_0, 0)$  and radius  $\rho > 0$  given by

$$\mu^+ = -s_0 - 3/2 \quad \text{and} \quad 2\Delta^+ = \rho^2 - 1/4.$$

The inequality  $\rho > \sqrt{5}/2$  follows at once from  $\Delta^+ > 1/2$ . □



5.4. *The spectral sequence when  $(\xi, \xi_{\alpha,\beta}) < 0$ .*

As mentioned earlier, in this case we use the spectral sequence for  $U$  associated to the triad  $(E_{-(\alpha,\beta)-3}, E_{-\alpha-3}, E_{-\beta})$ . This time,

$$\begin{aligned} G_{-2} &= E_{-(\alpha,\beta)-3}, & F_{-2} &= E_{\alpha,\beta}, \\ G_{-1} &= E_{-\alpha-3}, & F_{-1} &= E_{(\alpha,\beta),\beta}, \\ G_0 &= E_{-\beta}, & F_0 &= E_{\beta}. \end{aligned}$$

It is easily shown that the general  $U$  has a spectral sequence of the form

$$\begin{aligned} E_{-(\alpha,\beta)-3} \otimes \mathbb{C}^{m_3} &\rightarrow 0 \rightarrow 0 \\ 0 \rightarrow E_{-\alpha-3} \otimes \mathbb{C}^{m_1} &\rightarrow E_{-\beta} \otimes \mathbb{C}^{m_2}, \end{aligned}$$

with

$$m_1 = \chi(U \otimes E_{(\alpha,\beta),\beta}), \quad m_2 = \chi(U \otimes E_{\beta}), \quad m_3 = -\chi(U \otimes E_{\alpha,\beta}).$$

The analog of Theorem 5.5 is substantially nicer in this case, since the analog of the object  $W$  of the derived category in Theorem 5.5 is a sheaf.

**Theorem 5.8.** *With the notation of this subsection, let  $U \in M(\xi)$  be general, and let*

$$E_{-\alpha-3} \otimes \mathbb{C}^{m_1} \rightarrow E_{-\beta} \otimes \mathbb{C}^{m_2}$$

*be the complex appearing in the generalized Beilinson spectral sequence for  $U$ , sitting in degrees  $-1$  and  $0$ . The map in this complex is injective, so the complex is isomorphic to a sheaf  $W$  with resolution*

$$0 \rightarrow E_{-\alpha-3} \otimes \mathbb{C}^{m_1} \rightarrow E_{-\beta} \otimes \mathbb{C}^{m_2} \rightarrow W \rightarrow 0.$$

*Any two resolutions of  $W$  of this form are conjugate under the  $GL(m_1) \times GL(m_2)$  action. Finally, the sheaf  $U$  fits in a triangle*

$$W \rightarrow U \rightarrow E_{-(\alpha,\beta)-3}[1] \otimes \text{Hom}(U, E_{-(\alpha,\beta)-3}[1])^* \rightarrow \cdot,$$

*where the second map is the canonical one.*

The obvious analog of Theorem 5.7 holds here as well.

5.5. *The spectral sequence when  $(\xi, \xi_{\alpha,\beta}) = 0$*

In this case, either of the previous spectral sequences degenerates to an exact sequence

$$0 \rightarrow E_{-\alpha-3} \otimes \mathbb{C}^{m_1} \rightarrow E_{-\beta} \otimes \mathbb{C}^{m_2} \rightarrow U \rightarrow 0,$$

where  $m_1 = -\chi(U \otimes E_\alpha)$  and  $m_2 = \chi(U \otimes E_\beta)$ . Comparing the two different spectral sequences shows we have equalities

$$-\chi(U \otimes E_\alpha) = \chi(U \otimes E_{(\alpha,\beta),\beta}), \quad -\chi(U \otimes E_{\alpha,(\alpha,\beta)}) = \chi(U \otimes E_\beta).$$

We have chosen to represent each  $m_i$  by its simplest expression. Furthermore, the next result follows easily from analyzing the first spectral sequence for  $U$ .

**Proposition 5.9.** *With the notation of this subsection, a sheaf  $U \in M^s(\xi)$  lies in the Brill–Noether locus  $D_{E_{\alpha,\beta}}$  if and only if it does not have a resolution of the form*

$$0 \rightarrow E_{-\alpha-3} \otimes \mathbb{C}^{m_1} \rightarrow E_{-\beta} \otimes \mathbb{C}^{m_2} \rightarrow U \rightarrow 0.$$

Furthermore,  $D_{E_{\alpha,\beta}}$  is a divisor.

*Proof.* By orthogonality properties of exceptional bundles, any stable sheaf  $U$  with a resolution of the above form has  $H^0(U \otimes E_{\alpha,\beta}) = 0$ , and thus does not lie in  $D_{E_{\alpha,\beta}}$ . This also shows  $D_{E_{\alpha,\beta}}$  is a divisor. Conversely, if  $H^0(U \otimes E_{\alpha,\beta}) = 0$ , then the first spectral sequence for  $U$  takes the form

$$\begin{aligned} E_{-\alpha-3} \otimes \mathbb{C}^{m_1+d_1} &\rightarrow E_{-\beta} \otimes \mathbb{C}^{m_2+d_2} \rightarrow 0, \\ E_{-\alpha-3} \otimes \mathbb{C}^{d_1} &\rightarrow E_{-\beta} \otimes \mathbb{C}^{d_2} \rightarrow 0, \end{aligned}$$


for some nonnegative integers  $d_1, d_2$ . If either  $d_1$  or  $d_2$  is positive, the sequence cannot converge to a sheaf in degree 0, so we conclude  $U$  has a resolution of the expected form.  $\square$

### 6. The Kronecker fibration

The generalized Beilinson spectral sequence described in the previous section allows us to describe a natural rational map from a moduli space  $M(\xi)$  to a moduli space of Kronecker modules. These moduli spaces have Picard rank 1, and analyzing this fibration will allow us to determine the effective cone of  $M(\xi)$ .

#### 6.1. Moduli spaces of Kronecker modules

Let  $V$  be an  $N$ -dimensional vector space with  $N \geq 3$ . A linear map  $e : B \otimes V \rightarrow A$  for  $b$ - and  $a$ -dimensional vector spaces  $B, A$ , respectively, can be viewed as a representation of the Kronecker quiver  $Q$  consisting of two vertices and  $N$  arrows from the first vertex to the second:



We call such a representation a *Kronecker  $V$ -module*. The *dimension vector*  $\underline{\dim} e \in \mathbb{N}^2$  of such a Kronecker module  $e$  is the vector  $(b, a)$ . If  $f : B' \otimes V \rightarrow A'$  is another Kronecker

$V$ -module, a homomorphism  $f \rightarrow e$  is a pair of maps  $\beta : B' \rightarrow B$  and  $\alpha : A' \rightarrow A$  making the square

$$\begin{array}{ccc} B \otimes V & \longrightarrow & A \\ \beta \otimes \text{id} \uparrow & & \uparrow \alpha \\ B' \otimes V & \longrightarrow & A' \end{array}$$

commute. The space of Kronecker  $V$ -modules forms an abelian category.

Fix a dimension vector  $(b, a)$  and vector spaces  $B, A$  of dimensions  $b, a$ . Every Kronecker  $V$ -module  $e$  with  $\underline{\dim} e = (b, a)$  is isomorphic to an element of  $\text{Hom}_{\mathbb{C}}(B \otimes V, A)$ . The natural action of  $\mathbb{C}^*$  on this space preserves the isomorphism type of the corresponding Kronecker modules, and there is an induced action of  $\text{SL}(B) \times \text{SL}(A)$  on  $\mathbb{P}\text{Hom}_{\mathbb{C}}(B \otimes V, A)$ . A Kronecker  $V$ -module  $e$  is *(GIT)-semistable* if any (equivalently, all) of the corresponding points of  $\mathbb{P}\text{Hom}_{\mathbb{C}}(B \otimes V, A)$  are semistable points of this action. We denote by  $Kr_N(\underline{\dim} e)$  the quotient space of semistable modules.

The *expected dimension* of  $Kr_N(b, a)$  is

$$\text{edim } Kr_N(b, a) = abN - a^2 - b^2 + 1.$$

If the expected dimension of  $Kr_N(b, a)$  is positive then stable modules with vector  $(b, a)$  exist, and thus  $Kr_N(b, a)$  has the expected dimension. (We will only need this result in case  $N = \chi(E, G)$  for two exceptional bundles  $E, G$  appearing in a triad  $(E, G, F)$  on  $\mathbb{P}^2$ ; in this case  $Kr_N(b, a)$  is isomorphic to a height 0 moduli space  $M(\xi)$  of sheaves on  $\mathbb{P}^2$  by [D3, Theorem 2]. Under this correspondence, stable modules correspond to stable sheaves, and the open set  $M^s(\xi)$  is nonempty by [LP, Corollary 15.4.6].)

For our present work, it is important to note that semistability of Kronecker modules can be detected by the existence of orthogonal modules. To explain this, we define the *Euler characteristic* of a pair  $(f, e)$  of modules by

$$\chi(f, e) = \text{hom}_Q(f, e) - \text{ext}_Q^1(f, e);$$

all the higher Ext terms vanish. The Euler characteristic of a pair can be computed in terms of the numerical invariants of the module and the quiver. Precisely, if  $\underline{\dim} f = (b', a')$  and  $\underline{\dim} e = (b, a)$ , then

$$\chi(f, e) = b'b + a'a - Nb'a.$$

It is useful to note that

$$\text{edim } Kr_N(\underline{\dim} e) = 1 - \chi(e, e).$$

**Theorem 6.1** ([S, SvdB]). *A Kronecker  $V$ -module  $e$  is semistable if and only if it admits a semistable left orthogonal module, i.e. there is some semistable Kronecker  $V$ -module  $f$  such that*

$$\text{Hom}_Q(f, e) = \text{Ext}_Q^1(f, e) = 0.$$

*Semistability is also equivalent to the existence of a semistable right orthogonal module.*

Suppose  $f$  and  $e$  are Kronecker modules with  $\chi(f, e) = 0$ . If  $\underline{\dim} e = (b, a)$  then

$$\underline{\dim} f = k(a, Na - b)$$

for some  $k \in \mathbb{Q}_{>0}$ , so that the dimension vector of  $f$  is determined up to a scalar multiple by  $\underline{\dim} e$ . We also observe that  $\text{edim } Kr_N(\underline{\dim} f)$  is positive if and only if  $\text{edim } Kr_N(\underline{\dim} e)$  is positive, since

$$\text{edim } Kr_N(\underline{\dim} f) - 1 = k^2(\text{edim } Kr_N(\underline{\dim} e) - 1).$$

All the Kronecker spaces we will need to consider going forward are isomorphic to height zero moduli spaces  $M(\xi)$  of sheaves on  $\mathbb{P}^2$ , and so have Picard group  $\mathbb{Z}$ . If  $f$  is a left orthogonal module to  $e$ , then the locus

$$D_f = \{e' : \text{Hom}_Q(f, e') \neq 0\} \subset Kr_N(\underline{\dim} e)$$

is an effective generator of  $\text{Pic } Kr_N(\underline{\dim} e) \otimes \mathbb{Q}$  such that  $e$  is not in the base locus of  $D_f$ .

### 6.2. Kronecker modules coming from complexes

Suppose  $(E, G, F)$  is a triad of exceptional bundles. Consider the category of complexes of the form

$$E^{n_1} \rightarrow G^{n_2} \quad (n_i \in \mathbb{N}),$$

where the homomorphisms are maps of complexes. Corresponding to such a complex, we can form a Kronecker  $\text{Hom}(E, G)^*$ -module

$$\mathbb{C}^{n_1} \otimes \text{Hom}(E, G)^* \rightarrow \mathbb{C}^{n_2}.$$

Since exceptional bundles are simple, homomorphisms of complexes are the same as homomorphisms of the corresponding Kronecker modules. In fact, this correspondence is an equivalence of categories, and it preserves the Euler characteristic of pairs.

### 6.3. Moduli spaces of sheaves birational to Kronecker spaces

In this subsection we study some simple moduli spaces of sheaves on the plane. The orthogonal invariants  $(\mu^+, \Delta^+)$  of an arbitrary moduli space  $M(\xi)$  are always of this form unless they are the invariants of an exceptional bundle, so studying these spaces is particularly important. These moduli spaces also include the height 0 moduli spaces of Picard rank 1.

**Proposition 6.2.** *Fix an exceptional slope  $\alpha, \beta$ , and suppose  $\xi$  is a stable Chern character with  $(\xi, \xi_{-(\alpha, \beta)}) = 0$  and  $\Delta(\xi) > 1/2$ . We have either  $\mu(\xi) > \alpha, \beta - x_{\alpha, \beta}$  or  $\mu(\xi) < \alpha, \beta - 3 + x_{\alpha, \beta}$ . Let  $N = \chi(E_{\beta-3}, E_\alpha)$ .*

(1) *If  $\mu(\xi) > \alpha, \beta - x_{\alpha, \beta}$ , the general  $V \in M(\xi)$  admits a resolution of the form*

$$0 \rightarrow E_{\beta-3}^{n_1} \rightarrow E_\alpha^{n_2} \rightarrow V \rightarrow 0$$

*unique up to the action of  $\text{GL}(n_1) \times \text{GL}(n_2)$ , and  $M(\xi)$  is birational to  $Kr_N(n_1, n_2)$ .*

(2) If  $\mu(\xi) < \alpha.\beta - 3 + x_{\alpha,\beta}$  and  $r(\xi) \geq 2$ , the general  $V \in M(\xi)$  admits a resolution

$$0 \rightarrow V \rightarrow E_{\beta-3}^{n_1} \rightarrow E_{\alpha}^{n_2} \rightarrow 0$$

unique up to the action of  $GL(n_1) \times GL(n_2)$ , and  $M(\xi)$  is birational to  $Kr_N(n_1, n_2)$ .

*Proof.* The invariants of  $\xi$  lie on the parabola  $Q_{\xi-(\alpha,\beta)}$ , which intersects the line  $\Delta = 1/2$  at the points  $\alpha.\beta - 3 + x_{\alpha,\beta}$  and  $\alpha.\beta - x_{\alpha,\beta}$ . Since  $\Delta(\xi) > 1/2$ , it follows that either  $\mu(\xi) > \alpha.\beta - x_{\alpha,\beta}$  or  $\mu(\xi) < \alpha.\beta - 3 + x_{\alpha,\beta}$ .

(1) For  $\mu(\xi) > \alpha.\beta - x_{\alpha,\beta}$ , it is easy to show that the associated exceptional slope to  $\xi$  is  $-(\alpha.\beta) = (-\beta).(-\alpha)$ . The results of §5.5 then give that there is a resolution of the general  $V \in M(\xi)$  of the form

$$0 \rightarrow E_{\beta-3}^{n_1} \rightarrow E_{\alpha}^{n_2} \rightarrow V \rightarrow 0,$$

where  $n_1 = -\chi(E_{\beta}, V)$  and  $n_2 = \chi(E_{\alpha}, V)$ . We also know that the map in a general complex

$$E_{\beta-3}^{n_1} \rightarrow E_{\alpha}^{n_2}$$

is injective with semistable cokernel of Chern character  $\xi$  (Proposition 5.3), and that if two general semistable bundles with Chern character  $\xi$  are isomorphic then the isomorphism is induced by an isomorphism of their resolutions.

We now associate to a two-term complex as above a Kronecker  $\text{Hom}(E_{\beta-3}, E_{\alpha})^*$ -module  $e$  with dimension vector  $(n_1, n_2)$  as in Section 6.2. Then

$$\dim M(\xi) = 1 - \chi(V, V) = 1 - \chi(e, e) = \text{edim } Kr_N(n_1, n_2),$$

so since  $\dim M(\xi) > 0$  we conclude  $Kr_N(n_1, n_2)$  has positive expected dimension. Thus the general such module is stable. Furthermore, isomorphism of Kronecker modules corresponds to isomorphism of complexes, so we obtain a birational map  $M(\xi) \dashrightarrow Kr_N(n_1, n_2)$ .

(2) Since the rank is at least 2, the general  $V \in M(\xi)$  is locally free and taking duals gives a birational map  $M(\xi) \dashrightarrow M(\xi^*)$ . This case then follows by applying case (1) to the Chern character  $\xi^*$  satisfying  $(\xi^*, \xi_{\alpha,\beta-3}) = 0$ .  $\square$

#### 6.4. The Kronecker fibration

Let  $\xi$  be a stable Chern character corresponding to a moduli space of semistable sheaves with Picard rank 2. Let  $\alpha.\beta$  be the corresponding exceptional slope, and let  $\xi^+$  be the corresponding orthogonal character.

Let  $U$  be a general sheaf in  $M(\xi)$ . We saw in §5 that the Beilinson spectral sequence gives a complex  $W$  of the form

$$E_{-\alpha-3}^{m_1} \rightarrow E_{-\beta}^{m_2},$$

and that the isomorphism class of  $W$  depends only on  $U$ . Equivalently,  $U$  determines an isomorphism class of  $\text{Hom}(E_{-\alpha-3}, E_{-\beta})^*$ -modules. Let  $N = \chi(E_{-\alpha-3}, E_{-\beta})$ . We will show that since  $U$  is general, this complex is semistable; furthermore, the set-theoretic association  $U \mapsto W$  is in fact a rational map  $M(\xi) \dashrightarrow Kr_N(m_1, m_2)$ . We call this map the *Kronecker fibration*.

**Proposition 6.3.** *The Kronecker fibration  $\pi : M(\xi) \dashrightarrow Kr_N(m_1, m_2)$  is a dominant rational map, and  $Kr_N(m_1, m_2)$  is positive-dimensional.*

*Proof.* We first show  $Kr_N(m_1, m_2)$  is positive-dimensional. By Proposition 3.7, the Chern character  $\xi^+$  is semistable. Let  $V$  be a general semistable vector bundle with character  $\xi^+$ . First suppose  $(\xi, \xi_{\alpha,\beta}) > 0$ . By definition,  $\{(\mu^+, \Delta^+)\} = Q_{\xi_{-\alpha,\beta}} \cap Q_\xi$ , and  $\mu^+ > \alpha.\beta - x_{\alpha,\beta}$  by Lemma 3.6. According to Proposition 6.2, the bundle  $V$  has a resolution

$$0 \rightarrow E_{\beta-3}^{n_1} \rightarrow E_\alpha^{n_2} \rightarrow V \rightarrow 0.$$

By contrast, if  $U \in M(\xi)$  is general, we have a distinguished triangle

$$E_{-(\alpha,\beta)}^{m_3} \rightarrow U \rightarrow W \rightarrow \cdot,$$

where  $W$  is a complex

$$W : E_{-\alpha-3}^{m_1} \rightarrow E_{-\beta}^{m_2}.$$

As  $(\mu^+, \Delta^+) \in Q_\xi$ , we have  $\chi(V^*, U) = 0$ . Orthogonality properties of exceptional bundles show  $\chi(V^*, E_{-(\alpha,\beta)}) = 0$  as well, and therefore  $\chi(V^*, W) = 0$ . By Serre duality, this is equivalent to  $\chi(W, V^*(-3)) = 0$ , and we further obtain  $\chi(W, V^*(-3)[1]) = 0$ . As  $V^*(-3)[1]$  is isomorphic in the derived category to a complex

$$V^*(-3)[1] : E_{-\alpha-3}^{n_1} \rightarrow E_{-\beta}^{n_2}$$

sitting in degrees  $-1$  and  $0$ , both  $W$  and  $V^*(-3)[1]$  correspond to Kronecker  $\text{Hom}(E_{-\alpha-3}, E_{-\beta})^*$ -modules, say  $f$  and  $e$ , respectively. We conclude that  $\chi(f, e) = 0$ , so  $\underline{\dim} f$  is a left-orthogonal dimension vector to  $\underline{\dim} e$ . The space  $M(\text{ch } V)$  is positive-dimensional, so Proposition 6.2 shows that  $Kr_N(n_1, n_2)$  is positive-dimensional. By the discussion at the end of §6.1,  $Kr_N(m_1, m_2)$  is positive-dimensional as well. In case  $(\xi, \xi_{\alpha,\beta}) < 0$ , a similar argument works. When  $(\xi, \xi_{\alpha,\beta}) = 0$ , the positive-dimensionality was proved in Proposition 6.2.

Next we show that  $U \mapsto W$  gives a dominant rational map  $M(\xi) \dashrightarrow Kr_N(m_1, m_2)$ . Again we focus on the case  $(\xi, \xi_{\alpha,\beta}) > 0$ . Since  $Kr_N(m_1, m_2)$  is nonempty, a general complex  $W_0$  of the form  $E_{-\alpha-3}^{m_1} \rightarrow E_{-\beta}^{m_2}$  is semistable when regarded as a Kronecker module. If  $W_0$  is sufficiently general, then by Proposition 5.3 and Theorem 5.5 there is a stable sheaf  $U_0 \in M^s(\xi)$  such that the corresponding complex in the Beilinson spectral sequence is  $W_0$ .

Let  $\mathcal{U}$  on  $\mathbb{P}^2 \times S$  be a complete family of stable sheaves on  $\mathbb{P}^2$  of Chern character  $\xi$ , parameterized by  $S$ , and assume  $U_0$  is a sheaf parameterized by a point  $0 \in S$ . We consider a relative version of the map giving  $W$  in the Beilinson spectral sequence. Let  $q_i$  be the projections from  $\mathbb{P}^2 \times S$ , and let  $p_i$  and  $p_{ij}$  be the projections from  $\mathbb{P}^2 \times \mathbb{P}^2 \times S$  to its factors. The relative version of the map giving  $W$  is obtained by applying  $R^1 p_{13*}$  to the map

$$\psi : p_1^*(E_{-\alpha-3}) \otimes p_2^*(E_\alpha) \otimes p_{23}^*(\mathcal{U}) \rightarrow p_1^*(E_{-\beta}) \otimes p_2^*(E_{\alpha.(\alpha,\beta)}) \otimes p_{23}^*(\mathcal{U})$$

of sheaves on  $\mathbb{P}^2 \times \mathbb{P}^2 \times S$ ; after applying the projection formula, this is the map

$$R^1 p_{13*}(\psi) : q_1^*(E_{-\alpha-3}) \otimes q_2^* R^1 q_{2*}(q_1^*(E_\alpha) \otimes \mathcal{U}) \rightarrow q_1^*(E_{-\beta}) \otimes q_2^* R^1 q_{2*}(q_1^*(E_{\alpha.(\alpha.\beta)}) \otimes \mathcal{U}).$$

By cohomology and base change and Proposition 5.3, over an open dense subset of  $S$  the sheaves

$$R^1 q_{2*}(q_1^*(E_\alpha) \otimes \mathcal{U}) \quad \text{and} \quad R^1 q_{2*}(q_1^*(E_{\alpha.(\alpha.\beta)}) \otimes \mathcal{U})$$

are vector bundles of ranks  $m_1$  and  $m_2$ , respectively. We can thus view  $R^1 p_{13*}(\psi)$  as a flat family of Kronecker modules over this smaller subset. Since  $W_0$  is semistable, these modules are semistable over a smaller open set. The moduli property of  $Kr_N(m_1, m_2)$  gives us a rational map  $S \dashrightarrow Kr_N(m_1, m_2)$  which coincides with the set-theoretic association  $U \mapsto W$ .

In particular, suppose we realize  $M^s(\xi)$  as the GIT quotient  $R^s // G$  of a subvariety  $R^s$  of a Quot scheme as in [HuL, §4.3]. As in the previous paragraph, we obtain a  $G$ -invariant rational map  $R^s \dashrightarrow Kr_N(m_1, m_2)$ ; furthermore, there is a  $G$ -invariant open subset in the domain of definition of this map. Since the quotient  $R^s // G$  is a uniform categorical quotient, this map descends to a dominant rational map  $M(\xi) \dashrightarrow Kr_N(m_1, m_2)$ .  $\square$

Our final result of this section characterizes when the general fiber of the Kronecker fibration is positive-dimensional.

**Theorem 6.4.** *Let  $\pi : M(\xi) \dashrightarrow Kr_N(m_1, m_2)$  be the Kronecker fibration.*

- (1) *If  $(\xi, \xi_{\alpha.\beta}) \neq 0$ , then the general fiber of  $\pi$  is positive-dimensional.*
- (2) *If  $(\xi, \xi_{\alpha.\beta}) = 0$ , then  $\pi$  is birational.*

*Proof.* Part (2) was covered in Proposition 6.2. Let us show that if  $(\xi, \xi_{\alpha.\beta}) \neq 0$  then the general fiber of  $\pi$  is positive-dimensional. We focus on the case  $(\xi, \xi_{\alpha.\beta}) > 0$ , the other case being similar. Since  $\pi$  is dominant, we know  $\dim M(\xi) \geq \dim Kr_N(m_1, m_2)$ , and we must show the inequality is strict. By computing dimensions, this inequality is equivalent to

$$\chi(U, U) \leq \chi(W, W).$$

Expanding each side by additivity of the Euler characteristic, we have the equivalent inequality

$$m_3(\xi, \xi_{\alpha.\beta}) + \chi(W, U) \leq \chi(W, U) - m_3 \chi(W, E_{-(\alpha.\beta)}).$$

Since  $m_3 = (\xi, \xi_{\alpha.\beta}) > 0$ , this is the same as

$$(\xi, \xi_{\alpha.\beta}) \leq -\chi(W, E_{-(\alpha.\beta)}) = m_1 \chi(E_{-\alpha-3}, E_{-(\alpha.\beta)}) - m_2 \chi(E_{-\beta}, E_{-(\alpha.\beta)}).$$

Recalling  $m_1 = -(\xi, \xi_\alpha)$ ,  $m_2 = -(\xi, \xi_{\alpha.(\alpha.\beta)})$ , we see that there are nonzero constants  $C_1, C_2$  (independent of  $\xi$ ) such that the inequality is equivalent to

$$(\xi, \xi_{\alpha.\beta}) \leq (\xi, C_1 \xi_\alpha + C_2 \xi_{\alpha.(\alpha.\beta)}).$$

Noting that  $\{\xi_\alpha, \xi_{\alpha.\beta}, \xi_{\alpha.\beta}\}$  form a basis for  $K(\mathbb{P}^2) \otimes \mathbb{R}$ , we finally conclude there is some nonzero  $\zeta \in K(\mathbb{P}^2) \otimes \mathbb{R}$  such that the inequality  $\dim M(\xi) \geq \dim Kr_N(n_1, n_2)$  is equivalent to

$$(\xi, \zeta) \geq 0.$$

Here  $\zeta$  depends only on the corresponding exceptional slope  $\alpha.\beta$ , and not on the particular Chern character  $\xi$ . We must verify that the inequality is strict.

Consider the set

$$\Omega = \left\{ \xi' : \begin{array}{l} \Delta(\xi') > \delta(\mu(\xi')), \\ \alpha.\beta \text{ is the corresponding exceptional slope to } \xi', \\ \text{and } (\xi', \xi_{\alpha.\beta}) > 0 \end{array} \right\} \subset K(\mathbb{P}^2) \otimes \mathbb{Q}.$$

Clearly  $\Omega$  is an open set. If it intersects the plane  $\zeta^\perp$ , then  $\Omega$  contains some rational Chern character  $\xi'$  such that  $(\xi', \zeta) < 0$ , a contradiction. Thus  $\Omega$  and  $\zeta^\perp$  must be disjoint, and we conclude  $(\xi, \zeta) > 0$ . This implies  $\dim M(\xi) > \dim Kr_N(n_1, n_2)$ .  $\square$

### 7. The effective cone of the moduli space

In this section we prove our main theorem on the effective cone of divisors on  $M(\xi)$  by making use of the Kronecker fibration.

#### 7.1. The primary edge of the effective cone

Let  $\xi$  be a stable Chern character of positive rank such that  $M(\xi)$  has Picard rank 2. Let  $\alpha.\beta$  be the associated exceptional slope. We first consider the nonexceptional case.

**Theorem 7.1.** *Suppose  $(\xi, \xi_{\alpha.\beta}) \neq 0$ . Let  $\xi^+$  be a primary corresponding orthogonal character to  $\xi$ . If  $V \in M(\xi^+)$  is general, then the Brill–Noether locus*

$$D_V = \{U \in M(\xi) : h^0(U \otimes V) \neq 0\}$$

*is a divisor spanning the primary edge of the effective cone of  $M(\xi)$ . Under the isomorphism  $NS(M(\xi)) \cong \xi^\perp$ , it corresponds to  $\xi^+$ .*

*Proof.* We consider the case  $(\xi, \xi_{\alpha.\beta}) > 0$ , the other case being similar. A general  $U$  in  $M(\xi)$  fits into a triangle

$$E_{-\alpha.\beta}^{m_3} \rightarrow U \rightarrow W \rightarrow \cdot,$$

while  $V$  has a resolution

$$0 \rightarrow E_{\beta-3}^{n_1} \rightarrow E_\alpha^{n_2} \rightarrow V \rightarrow 0.$$

We wish to show  $V \otimes U$  has no cohomology. We find  $Hom(E_{\alpha.\beta}, V)$  has no cohomology. As in the first part of the proof of Theorem 6.4, we reduce to showing  $Hom(W, V^*(-3)[1])$  has no cohomology. But if  $f, e$  are the Kronecker modules corresponding to  $W$  and  $V^*(-3)[1]$ , respectively, this is equivalent to the vanishing of



$\text{Hom}(f, e)$ , which is true by Theorem 6.1. Then  $D_V$  is just the pullback of the ample divisor  $D_f \subset \text{Kr}(\underline{\dim} e)$  under the map  $\pi : M(\xi) \dashrightarrow \text{Kr}(\underline{\dim} e)$ . It follows that  $D_V$  spans an edge of the effective cone. The computation of the class of  $D_V$  and the fact that it spans the primary edge of the effective cone follows from Proposition 2.7 and Corollary 2.9.  $\square$

**Corollary 7.2.** *If  $(\xi, \xi_{\alpha,\beta}) \neq 0$ , then the primary edges of the effective and movable cones of divisors of  $M(\xi)$  coincide.*

*Proof.* Since  $D_V$  is the pullback of an ample divisor from the space of Kronecker modules, its stable base locus is contained in the indeterminacy locus of  $\pi$ . As  $M(\xi)$  is normal, the indeterminacy locus of  $\pi$  has codimension at least 2. Thus  $D_V$  spans a movable class.  $\square$

The exceptional case requires a slightly different argument.

**Theorem 7.3.** *Suppose  $(\xi, \xi_{\alpha,\beta}) = 0$ . The Brill–Noether divisor  $D_{E_{\alpha,\beta}}$  spans the primary edge of the effective cone of  $M(\xi)$ . Furthermore,  $D_{E_{\alpha,\beta}}$  is reduced and irreducible. Under the isomorphism  $\text{NS}(M(\xi)) \cong \xi^\perp$ , the divisor corresponds to the exceptional Chern character  $\xi_{\alpha,\beta}$ .*

*Proof.* By Proposition 5.9,  $D_{E_{\alpha,\beta}}$  is an effective divisor, and it is the complement of the locus of sheaves  $U$  with a resolution of the expected form

$$0 \rightarrow E_{-\alpha-3}^{m_1} \rightarrow E_{-\beta}^{m_2} \rightarrow U \rightarrow 0.$$

To see that it is extremal, we produce a dual moving curve. Let

$$S = \mathbb{P} \text{Hom}(E_{-\alpha-3}^{m_1}, E_{-\beta}^{m_2}),$$

and let  $\mathcal{U}/S$  be the universal cokernel sheaf. It is a complete family of prioritary sheaves with  $\Delta > \delta(\mu)$ , so by [LP, Lemma 18.3.1] the open set  $S' \subset S$  consisting of those  $s$  such that  $U_s$  is stable has complement of codimension at least 2. Thus for a general sheaf  $U \in M(\xi)$  we can find a complete curve in  $S'$  passing through a point  $s$  corresponding to  $U$ . Any such curve is disjoint from  $D_{E_{\alpha,\beta}}$ , so this divisor is extremal.

The divisor  $D_{E_{\alpha,\beta}}$  is reduced and irreducible because it is extremal and  $\xi_{\alpha,\beta}$  is a generator of  $K(\mathbb{P}^2) \cap \mathbb{Q}\xi_{\alpha,\beta}$ . Indeed,  $c_1(\xi_{\alpha,\beta})$  and  $r_{\alpha,\beta}$  are coprime, so no integral class  $\zeta$  of rank smaller than  $r_{\alpha,\beta}$  has slope  $\alpha/\beta$ .  $\square$

In the exceptional case, the primary edge of the movable cone of divisors is considerably more challenging to describe.

### 7.2. The rank 0 case

In this subsection, we discuss the primary edge of the effective cone of a moduli space  $M(\xi)$  of semistable rank 0 sheaves. We normalize our Chern character  $\xi = (0, d, \chi)$  in terms of the first Chern class  $d$  and Euler characteristic  $\chi$ . Assuming the Picard rank is 2, we have  $d \geq 3$ . The general point  $U \in M(\xi)$  corresponds to a line bundle of Euler characteristic  $\chi$  supported on a smooth plane curve of degree  $d$ .


The primary difference with the positive rank case is that the parabolas  $Q_\xi$  from the positive rank case degenerate to vertical lines in the  $(\mu, \Delta)$ -plane. As before, we define a locus

$$Q_\xi = \{(\mu, \Delta) : (1, \mu, \Delta) \text{ lies in } \xi^\perp\} \subset \mathbb{R}^2$$

as a subset of the  $(\mu, \Delta)$ -plane. A Riemann–Roch calculation shows  $Q_\xi$  is the vertical line

$$Q_\xi : \mu = -\chi/d.$$

It therefore intersects the line  $\Delta = 1/2$  in the rational point  $(-\chi/d, 1/2)$ . The corresponding exceptional slope to  $\xi$  is the slope  $\alpha \in \mathcal{E}$  with  $-\chi/d \in I_\alpha$ . The orthogonal invariants  $(\mu^+, \Delta^+)$  are defined in the same way as in the general case. In Figure 5 we display the possible arrangements of the curves involved.



**Fig. 5.** Graphical depiction of the corresponding orthogonal invariants in the rank 0 case, (1) when  $(\xi, \xi_\alpha) > 0$ , (2) when  $(\xi, \xi_\alpha) = 0$ , and (3) when  $(\xi, \xi_\alpha) < 0$ .

Once the orthogonal invariants have been defined, there is no major difficulty in carrying out arguments similar to those given for the general case. The necessary minor technical changes were already considered in [W]. All the results of §7.1 hold verbatim.

### 7.3. The secondary edge of the effective cone

We now discuss the behavior of the secondary edge of the effective cone.

When  $\xi$  has small rank, the secondary edge of the effective cone exhibits special behavior, which we now recall.

- If  $r(\xi) = 0$  and  $d = c_1(\xi) \geq 3$ , there is a map  $M(\xi) \rightarrow |\mathcal{O}_{\mathbb{P}^2}(d)|$  sending a sheaf to its support. This map has positive-dimensional fibers, and the pullback of  $\mathcal{O}(1)$  spans the secondary edge of the nef, movable, and effective cones. See [W] for more information about this class.
- If  $r(\xi) = 1$ , then  $M(\xi)$  is isomorphic to a Hilbert scheme of points in the plane. The secondary edge of the effective cone is spanned by the exceptional divisor  $B$  of the Hilbert–Chow morphism  $\mathbb{P}^{2[n]} \rightarrow \mathbb{P}^{2(n)}$ . The secondary edge of the movable cone is spanned by the divisor  $H$  given as the locus of schemes meeting a fixed line in the plane [ABCH].

- If  $r(\xi) = 2$ , the exceptional locus  $E$  of the morphism  $M(\xi) \rightarrow M^{\text{DUY}}(\xi)$  is divisorial, and it spans the secondary edge of the effective cone. Its support is the locus of singular sheaves. The secondary edge of the movable cone coincides with the ray giving this morphism. This case was considered in detail in [St].

We note that the classes of the previous extremal effective divisors can be easily determined in terms of the isomorphism  $\lambda_M : \xi^\perp \rightarrow \text{NS}(M(\xi))$ . We concentrate on the rank 2 case, as the lower rank cases are already well-understood [W, ABCH].

**Proposition 7.4.** *When  $r(\xi) = 2$ , the secondary edge of the effective cone is the ray spanned by  $\lambda_M(\zeta)$ , where  $\zeta$  is a negative rank orthogonal Chern character with  $\mu(\xi \otimes \zeta) = -3/2$ .*

*Proof.* Strømme [St, Proposition 4.6] describes how to give a scheme structure to the locus  $E$  of singular sheaves, at least when  $M(\xi)$  admits a universal family. When  $M(\xi)$  does not carry a universal family, the scheme structure can still first be described locally by similar methods to the proof of Proposition 2.7. If  $\mathcal{U}/S$  is a complete family of sheaves on  $\mathbb{P}^2$  with Chern character  $\xi$ , the divisor  $E_S$  on  $S$  of singular sheaves is cut out by the 0th Fitting ideal of the sheaf  $\mathcal{F} = p_* \mathcal{E}xt^1(\mathcal{U}, q^* \mathcal{O}_{\mathbb{P}^2}(-1))$  on  $S$ , and  $\mathcal{O}_S(E_S) \cong \det \mathcal{F}$ . Strømme describes a canonical locally free resolution of  $\mathcal{F}$  of the form

$$0 \rightarrow A \rightarrow B \rightarrow \mathcal{F} \rightarrow 0,$$

where  $A$  and  $B$  are sheaves whose determinants are of the form  $\det A = \lambda_{\mathcal{U}}(\zeta_1)$  and  $\det B = \lambda_{\mathcal{U}}(\zeta_2)$  for some  $\zeta_i \in K(\mathbb{P}^2)$ . An easy computation shows that  $\det \mathcal{F} = \lambda_{\mathcal{U}}(\zeta)$ , where  $\zeta = \zeta_2 - \zeta_1$  is an orthogonal Chern character to  $\xi$  with  $\mathcal{O}_{M(\xi)}(E) = \lambda_M(\zeta)$ , and  $\mu(\xi \otimes \zeta) = -3/2$ . □

Suppose  $r(\xi) \geq 3$ . The complement of the open set in  $M(\xi)$  consisting of stable locally-free sheaves has codimension at least 2 by [LP, Corollary 17.1.5 and Lemma 18.3.1]. Thus the functor  $\text{Hom}(-, \mathcal{O}_{\mathbb{P}^2}(-3))$  gives a birational map  $M(\xi) \dashrightarrow M(\xi^D)$ , where  $\xi^D$  denotes the Serre dual Chern character, and this map is an isomorphism between the loci of stable locally free sheaves. It therefore induces an isomorphism of Picard groups  $\text{Pic}(M(\xi)) \cong \text{Pic}(M(\xi^D))$  which preserves effective cones.

However, this isomorphism does *not* preserve the primary edge of the effective cone. The  $h^2$ -Brill–Noether divisor  $D_V$  on  $M(\xi)$  is carried to the  $h^0$ -Brill–Noether divisor  $D_{V^*}$  on  $M(\xi^D)$ , and vice versa. Thus, by Corollary 2.9, the isomorphism takes the primary edge of the effective cone of  $M(\xi^D)$  to the secondary edge of the effective cone of  $M(\xi)$ . We can therefore determine the secondary edge of the effective cone of  $M(\xi)$  in terms of the primary edge of the effective cone of the Serre dual moduli space.

**Definition 7.5.** Suppose  $r(\xi) \geq 3$ . Let  $(\mu_D^+, \Delta_D^+)$  be the corresponding orthogonal invariants to  $\xi^D$ . The *secondary corresponding orthogonal invariants* to  $\xi$  are  $(\mu^-, \Delta^-)$ , given by

$$\mu^- = -\mu_D^+ \quad \text{and} \quad \Delta^- = \Delta_D^+.$$

The *secondary corresponding exceptional slope* to  $\xi$  is given by  $-\alpha_D$ , where  $\alpha_D$  is the primary corresponding exceptional slope to  $\xi^D$ .

The following theorem is immediate.

**Theorem 7.6.** *If  $r(\xi) \geq 3$ , let  $\xi^- = (r^-, \mu^-, \Delta^-)$  where  $r^-$  is sufficiently large and divisible. There exist stable vector bundles  $V$  of Chern character  $\xi^-$  cohomologically orthogonal to the general  $U \in M(\xi)$ . The  $h^2$ -Brill–Noether divisor  $D_V$  spans the secondary edge of the effective cone of  $M(\xi)$ .*


Corollary 7.2 and Theorem 7.3 also have immediate generalizations for the secondary edges of the effective and movable cones.

**Example 7.7.** We close the paper with a typical computation of the effective cone of a moduli space  $M(\xi)$ .

Consider the Chern character  $\xi = (r, \mu, \Delta) = (3, 2/3, 17/9)$ . The moduli space  $M(\xi)$  has dimension  $r^2(2\Delta - 1) + 1 = 26$ . The parabola  $Q_\xi$  intersects the line  $\Delta = 1/2$  at the two points  $(\mu_0^\pm, 1/2)$  given by

$$\mu_0^\pm = \frac{1}{6}(-13 \pm \sqrt{181}), \quad \text{so } \mu_0^+ \approx 0.076 \quad \text{and} \quad \mu_0^- \approx -4.409.$$

Then  $\mu_0^+ \in I_0$  and  $\mu_0^- \in I_{-22/5}$ , so the primary and secondary corresponding exceptional slopes are 0 and  $-22/5$ , respectively (see Figure 6).



**Fig. 6.** Computing the corresponding orthogonal invariants for the Chern character  $\xi$  of rank 3, slope  $2/3$ , and discriminant  $17/9$ .

Consider the primary edge of the effective cone. We have  $(\xi, \xi_0) = 1$ , so the primary orthogonal invariants  $(\mu^+, \Delta^+)$  are given by the intersection  $Q_\xi \cap Q_{\xi_0}$ . We find  $(\mu^+, \Delta^+) = (1, 3)$ . A primary orthogonal Chern character is any character  $\xi^+ = (r^+, \mu^+, \Delta^+)$  with sufficiently large and divisible rank. There are stable vector bundles  $V$

with Chern character  $\xi^+$  cohomologically orthogonal to the general  $U \in M(\xi)$ , and the Brill–Noether divisors  $D_V$  span the primary edge of the effective cone.

To make the proof of the theorems explicit in this case, we observe that the general  $U \in M(\xi)$  has a resolution of the form

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^2}(-2)^4 \rightarrow \mathcal{O}_{\mathbb{P}^2}(-1)^6 \oplus \mathcal{O}_{\mathbb{P}^2} \rightarrow U \rightarrow 0.$$

The Kronecker fibration is a rational map  $\pi : M(\xi) \dashrightarrow Kr_3(4, 6)$ , and this space of Kronecker modules has dimension 21 and Picard rank 1. The divisor  $D_V$  is the pullback of a multiple of the ample generator on  $Kr_3(4, 6)$ , so it is an extremal effective divisor.

For the secondary edge of the effective cone, we observe that  $Q_\xi$  passes through the invariants  $\xi_{-22/5}$  of the exceptional bundle of slope  $-22/5$ . The primary corresponding exceptional slope to the Serre dual character  $\xi^D$  is  $22/5$ , and we find  $(\mu^-, \Delta^-) = (-22/5, 12/25)$ . Furthermore, the general  $U_D \in M(\xi^D)$  admits a resolution of the form

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^2}(-7) \rightarrow T_{\mathbb{P}^2}(-6)^2 \rightarrow U_D \rightarrow 0.$$

We have  $\text{hom}(\mathcal{O}_{\mathbb{P}^2}(-7), T_{\mathbb{P}^2}(-6)) = 15$ , so the Kronecker map is a birational map  $M(\xi^D) \dashrightarrow Kr_{15}(1, 2)$ . The Brill–Noether divisor  $D_{E_{22/5}}$  spans the primary edge of the effective cone of  $M(\xi^D)$ . Correspondingly, the  $h^2$ -Brill–Noether divisor  $D_{E_{-22/5}}$  on  $M(\xi)$  spans the secondary edge of the effective cone of  $M(\xi)$ .

*Acknowledgments.* We are grateful to Daniele Arcara, Arend Bayer, Aaron Bertram, Lawrence Ein, Joe Harris, Brendan Hassett, Emanuele Macrì and Yuri Tschinkel for many useful conversations about the geometry of the moduli spaces of sheaves. We would also like to thank the Vietnam Institute for Advanced Study in Mathematics (VIASM) for providing a wonderful working environment while this work was completed. Finally, we would like to thank the referees for providing many useful comments which improved the paper.

During the preparation of this article the first author was partially supported by the NSF CAREER grant DMS-0950951535, and an Alfred P. Sloan Foundation Fellowship and the second author was partially supported by a National Science Foundation Mathematical Sciences Postdoctoral Research Fellowship

## References

- [AB] Adamczewski, B., Bugeaud, Y.: Palindromic continued fractions. *Ann. Inst. Fourier (Grenoble)* **57**, 1557–1574 (2007) [Zbl 1126.11036](#) [MR 2364142](#)
- [A] Alper, J.: Good moduli spaces for Artin stacks. *Ann. Inst. Fourier* **63**, 2349–2402 (2013) [Zbl 1314.14095](#) [MR 3237451](#)
- [ABCH] Arcara, D., Bertram, A., Coskun, I., Huizenga, J.: The minimal model program for the Hilbert scheme of points on  $\mathbb{P}^2$  and Bridgeland stability. *Adv. Math.* **235**, 580–626 (2013) [Zbl 1267.14023](#) [MR 3010070](#)
- [BM1] Bayer, A., Macrì, E.: Projectivity and birational geometry of Bridgeland moduli spaces. *J. Amer. Math. Soc.* **27**, 707–752 (2014) [Zbl 1314.14020](#) [MR 3194493](#)
- [BM2] Bayer, A., Macrì, E.: MMP for moduli of sheaves on K3s via wall-crossing: nef and movable cones, Lagrangian fibrations. *Invent. Math.* **198**, 505–590 (2014) [Zbl 1308.14011](#) [MR 3279532](#)

- [BC] Bertram, A., Coskun, I.: The birational geometry of the Hilbert scheme of points on surfaces. In: *Birational Geometry, Rational Curves, and Arithmetic*, Springer, New York, 15–55 (2013) [Zbl 1273.14032](#) [MR 3114922](#)
- [BCHM] Birkar, C., Cascini, P., Hacon, C., McKernan, J.: Existence of minimal models for varieties of log general type. *J. Amer. Math. Soc.* **23**, 405–468 (2010) [Zbl 1210.14019](#) [MR 2601039](#)
- [B] Boutot, J.-F.: Singularités rationnelles et quotients par les groupes réductifs. *Invent. Math.* **88**, 65–68 (1987) [Zbl 0619.14029](#) [MR 0877006](#)
- [CH1] Coskun, I., Huizenga, J.: Interpolation, Bridgeland stability and monomial schemes in the plane. *J. Math. Pures Appl.* **102**, 930–971 (2014) [Zbl 06361756](#) [MR 3271294](#)
- [CH2] Coskun, I., Huizenga, J.: The ample cone of moduli spaces of sheaves on the plane. *Algebraic Geom.* **3**, 106–136 (2016) [Zbl 06609387](#) [MR 3455422](#)
- [Da] Davenport, H.: *The Higher Arithmetic*. 8th ed., Cambridge Univ. Press, Cambridge (2008) [Zbl 1204.11002](#) [MR 2462408](#)
- [D1] Drézet, J.-M.: Exceptional bundles and moduli spaces of stable sheaves on  $\mathbf{P}_n$ . In: *Vector Bundles in Algebraic Geometry* (Durham, 1993), London Math. Soc. Lecture Note Ser. 208, Cambridge Univ. Press, Cambridge, 101–117 (1995) [Zbl 0860.14018](#) [MR 1338414](#)
- [D2] Drézet, J.-M.: Fibrés exceptionnels et suite spectrale de Beilinson généralisée sur  $\mathbf{P}_2(\mathbf{C})$ . *Math. Ann.* **275**, 25–48 (1986) [Zbl 0578.14013](#) [MR 0849052](#)
- [D3] Drézet, J.-M.: Fibrés exceptionnels et variétés de modules de faisceaux semi-stables sur  $\mathbf{P}_2(\mathbf{C})$ . *J. Reine Angew. Math.* **380**, 14–58 (1987) [Zbl 0613.14013](#) [MR 0916199](#)
- [DLP] Drézet, J.-M., Le Potier, J.: Fibrés stables et fibrés exceptionnels sur  $\mathbf{P}_2$ . *Ann. Sci. École Norm. Sup. (4)* **18**, 193–243 (1985) [Zbl 0586.14007](#) [MR 0816365](#)
- [GR] Gorodentsev, A. L., Rudakov, A. N.: Exceptional vector bundles on projective spaces. *Duke Math. J.* **54**, 115–130 (1987) [Zbl 0646.14014](#) [MR 0885779](#)
- [E] Eisenbud, D.: *Commutative Algebra*. Grad. Texts in Math. 150, Springer, New York (1995) [Zbl 0819.13001](#) [MR 1322960](#)
- [HiL] Hirschowitz, A., Laszlo, Y.: Fibrés génériques sur le plan projectif. *Math. Ann.* **297**, 85–102 (1993) [Zbl 0808.14007](#) [MR 1238408](#)
- [HK] Hu, Y., Keel, S.: Mori dream spaces and GIT. *Michigan Math. J.* **48**, 331–348 (2000) [Zbl 1077.14554](#) [MR 1786494](#)
- [H] Huizenga, J.: Effective divisors on the Hilbert scheme of points in the plane and interpolation for stable bundles. *J. Algebraic Geom.* **25**, 19–75 (2016) [Zbl 06543158](#) [MR 3419956](#)
- [HuL] Huybrechts, D., Lehn, M.: *The Geometry of Moduli Spaces of Sheaves*, 2nd ed., Cambridge Math. Library, Cambridge Univ. Press, Cambridge (2010) [Zbl 1206.14027](#) [MR 2665168](#)
- [K] Kollár, J.: Singularities of pairs. In: *Algebraic Geometry—Santa Cruz 1995*, Proc. Sympos. Pure Math. 62, Part 1, Amer. Math. Soc., Providence, RI, 221–287 (1997) [Zbl 0905.14002](#) [MR 1492525](#)
- [La] Lazarsfeld, R.: *Positivity in Algebraic Geometry. I*. *Ergeb. Math. Grenzgeb.* 48, Springer, Berlin (2004) [Zbl 1093.14501](#) [MR 2095471](#)
- [LP] Le Potier, J.: *Lectures on Vector Bundles*. Cambridge Stud. Adv. Math. 54, Cambridge Univ. Press, Cambridge (1997) [Zbl 0872.14003](#) [MR 1428426](#)
- [Li] Li, J.: Algebraic geometric interpretation of Donaldson’s polynomial invariants. *J. Differential Geom.* **37**, 417–466 (1993) [Zbl 0809.14006](#) [MR 1205451](#)
- [LZ] Li, C., Zhao, X.: The MMP for deformations of Hilbert schemes of points on the projective plane. [arXiv:1312.1748](#) (2013)

- [MM] Maciocia, A., Meachan, C.: Rank 1 Bridgeland stable moduli spaces on a principally polarized abelian surface. *Int. Math. Res. Notices* **2013**, 2054–2077 [Zbl 1312.14036](#) [MR 3053413](#)
- [MYY1] Minamide, H., Yanagida, S., Yoshioka, K.: Fourier–Mukai transforms and the wall-crossing behavior for Bridgeland’s stability conditions. [arXiv:1106.521v2](#) (2012)
- [MYY2] Minamide, H., Yanagida, S., Yoshioka, K.: Some moduli spaces of Bridgeland’s stability conditions, *Int. Math. Res. Notices* **2014**, 5264–5327 [Zbl 1327.14087](#) [MR 3267372](#)
- [Ne] Nuer, H.: Projectivity and birational geometry of moduli spaces of Bridgeland stable objects on an Enriques surface. *Proc. London Math. Soc.* **113**, 345–386 (2016) [Zbl 06647887](#) [MR 3551850](#)
- [Oh] Ohkawa, R.: Moduli of Bridgeland semistable objects on  $\mathbf{P}^2$ . *Kodai Math. J.* **33**, 329–366 (2010) [Zbl 1198.14040](#) [MR 2681544](#)
- [R] Roth, K. F.: Rational approximations to algebraic numbers. *Mathematika* **2**, 1–20 (1955); corrigendum, 168. [Zbl 0064.28501](#) [MR 0072182](#)
- [Sc] Schmidt, W. M.: Norm form equations. *Ann. of Math. (2)* **96**, 526–551 (1972) [Zbl 0226.10024](#) [MR 0314761](#)
- [S] Schofield, A.: Semi-invariants of quivers. *J. London Math. Soc. (2)* **43**, 385–395 (1991) [Zbl 0779.16005](#) [MR 1113382](#)
- [SvdB] Schofield, A., van den Bergh, M.: Semi-invariants of quivers for arbitrary dimension vectors. *Indag. Math. (N.S.)* **12**, 125–138 (2001) [Zbl 1004.16012](#) [MR 1908144](#)
- [St] Strømme, S. A.: Ample divisors on fine moduli spaces on the projective plane. *Math. Z.* **187**, 405–423 (1984) [Zbl 0533.14006](#) [MR 0757480](#)
- [T] Tschinkel, Y.: Fujita’s program and rational points. In: *Higher Dimensional Varieties and Rational Points* (Budapest, 2001), *Bolyai Soc. Math. Stud.* 12, Springer, Berlin, 283–310 (2003) [Zbl 1112.14021](#) [MR 2011749](#)
- [W] Woolf, M.: Nef and effective cones on the moduli space of torsion sheaves on the projective plane. [arXiv:1305.1465](#) (2013)
- [YY] Yanagida, S., Yoshioka, K.: Bridgeland’s stabilities on abelian surfaces. *Math. Z.* **276**, 571–610 (2014) [Zbl 1292.14012](#) [MR 3150219](#)
- [Y] Yoshioka, K.: Bridgeland’s stability and the positive cone of the moduli spaces of stable objects on an abelian surface. [arXiv:1206.4838v2](#) (2014)