

Peter Keevash

Counting designs

Received April 18, 2015

Abstract. We give estimates on the number of combinatorial designs, which prove (and generalise) a conjecture of Wilson from 1974 on the number of Steiner Triple Systems. This paper also serves as an expository treatment of our recently developed method of Randomised Algebraic Construction: we give a simpler proof of a special case of our result on clique decompositions of hypergraphs, namely triangle decompositions of quasirandom graphs.

Keywords. Design theory, graph theory

1. Introduction

When does a graph G have a triangle decomposition? (By this we mean a partition of its edge set into triangles.) There are two obvious necessary ‘divisibility conditions’: the number of edges must be divisible by three, and the degree of any vertex must be even. We say that G is *tridivisible* if it satisfies these divisibility conditions. In 1847 Kirkman proved that any tridivisible complete graph has a triangle decomposition; equivalently, there is a Steiner Triple System on n vertices if n is 1 or 3 mod 6. In [5] we showed more generally that a tridivisible graph has a triangle decomposition if we assume a certain pseudorandomness condition. In fact, we proved a more general result on clique decompositions of simplicial complexes, which in particular proved the Existence Conjecture for combinatorial designs.

One purpose of the current paper is to illustrate the new technique (Randomised Algebraic Construction) of [5] in the simplified setting of triangle decompositions; we will also prove a conjecture of Wilson [12] on the number of Steiner Triple Systems. These results are proved in the next three sections, roughly following the method of [5], but introducing some novelties in technique that lead to considerable simplifications in the case of triangle decompositions; the material here closely follows a lecture series that the author recently gave at the Israel Institute for Advanced Studies. In Section 5 we sketch an argument of Bennett and Bohman [1] on the random greedy matching process and adapt the calculations to the version needed in this paper. We generalise from Steiner Triple

P. Keevash: Mathematical Institute, University of Oxford,
Oxford, UK; e-mail: keevash@maths.ox.ac.uk

Mathematics Subject Classification (2010): 05B05, 05C51

Systems to designs in Section 6. We conclude by noting that it remains an open problem to obtain an asymptotic formula for the number of designs, or even just for the number of regular graphs.

2. Triangle decompositions

We start by stating our result that tridivisible pseudorandom graphs have triangle decompositions. The pseudorandomness condition is as follows. Let G be a graph on n vertices. The *density* of G is $d(G) = |E(G)|/\binom{n}{2}$. We say that G is c -*typical* if every vertex has $(1 \pm c)d(G)n$ neighbours and every pair of vertices have $(1 \pm c)d(G)^2n$ common neighbours. (We write $b \pm c$ for any real between $b - c$ and $b + c$.)

Theorem 2.1. *There exist $0 < c_0 < 1$ and $n_0 \in \mathbb{N}$ such that if $n \geq n_0$ and G is a c -typical tridivisible graph on n vertices with $d(G) > n^{-10^{-7}}$ and $c < c_0 d(G)^{10^6}$ then G has a triangle decomposition.*

Note that in Theorem 2.1 we allow the density to decay polynomially with n ; this will be important for the application in the next subsection, but in many cases of interest one can consider $d(G)$ and c to be fixed constants independent of n . One such consequence of Theorem 2.1 noted in [5] is that the standard random graph model $G(n, 1/2)$ with high probability (whp) has a partial triangle decomposition that covers all but $(1 + o(1))n/4$ edges. Indeed, deleting a perfect matching on the set of vertices of odd degree and then at most two 4-cycles gives a graph satisfying the hypotheses of the theorem. This is asymptotically best possible, as whp there are $(1 + o(1))n/2$ vertices of odd degree, and any set of edge-disjoint triangles must leave at least one edge uncovered at each vertex of odd degree.

We remark that our definition of typicality here is weaker than that used in [5]. In fact, for most of the paper we will assume the stronger version, then explain at the end how the proof can be modified to work with the current definition. We also make the (well-known) remark that typicality implies the standard regularity property (for appropriate constants) that appears in Szemerédi's Regularity Lemma, but the converse is not true, as regularity allows individual vertices to behave badly, even to be isolated.

2.1. The number of Steiner Triple Systems

Another purpose of our paper is to prove the following conjecture of Wilson [12] on the number of *Steiner Triple Systems* on n vertices, i.e. triangle decompositions of the complete graph K_n ; denote this by $STS(n)$.

Theorem 2.2. *If n is 1 or 3 mod 6, then $STS(n) = (n/e^2 + o(n))^{n^2/6}$.*

Note that K_n is tridivisible if and only if n is 1 or 3 mod 6, so $STS(n) = 0$ for all other n . The upper bound in Theorem 2.2 was recently proved by Linial and Luria [8], who showed that $STS(n) \leq (n/e^2 + O(\sqrt{n}))^{n^2/6}$. Our lower bound will be $STS(n) \geq (n/e^2 + O(n^{1-a}))^{n^2/6}$ for some small $a > 0$.

Theorem 2.2 will follow quite easily from Theorem 2.1 and the semirandom method (nibble). It will be most convenient for us to apply the results of Bohman, Frieze and Lubetzky [2] on the triangle removal process (although we could make do with a simpler nibble argument, or the argument of Bennett and Bohman [1] sketched in Section 5). We say that an event E holds *with high probability* (whp) if $\mathbb{P}(E) = 1 - e^{-\Omega(n^c)}$ for some $c > 0$ as $n \rightarrow \infty$; note that when n is sufficiently large, by union bounds we can assume that any specified polynomial number of such events all occur.

In the triangle removal process, we start with the complete graph K_n , and at each step we delete the edges of a uniformly random triangle in the current graph. It is shown in [2] that whp the process persists until only $O(n^{3/2+o(1)})$ edges remain, but we will stop at $n^{2-10^{-7}}$ edges (i.e. at the nearest multiple of 3 to this number) so that we can apply Theorem 2.1. We need the following additional facts from [2] about this stopped process: whp the final graph is $n^{-1/3}$ -typical, and when $pn^2/2$ edges remain the number of choices for the deleted triangle is $(1 \pm n^{-2/3})(pn)^3/6$.

Proof of Theorem 2.2. Consider the following procedure for constructing a Steiner Triple System on n vertices: run the triangle removal process until $n^{2-10^{-7}}$ edges remain, then apply Theorem 2.1 (if its hypotheses are satisfied, which occurs in $1 - o(1)$ proportion of all instances of the process). Writing m for the number of steps and $p(i) = 1 - 6i/n^2$, we see that the logarithm of the number of choices in this procedure is

$$L_1 = \sum_{i=1}^m (\log(p(i)^3 n^3/6) \pm 2n^{-2/3}) = (n^2/6)(\log(n^3/6) - 3 \pm n^{-10^{-8}}),$$

since $\sum_{i=1}^m \log p(i) = (1 + O(n^{-10^{-7}} \log n))(n^2/6) \int_0^1 \log p \, dp$ and $\int_0^1 \log p \, dp = -1$. Also, for any fixed Steiner Triple System, the logarithm of the number of times it is counted by this procedure is at most

$$L_2 = \sum_{i=1}^m \log(p(i)n^2/6) = (n^2/6)(\log(n^2/6) - 1 \pm n^{-10^{-8}}).$$

Therefore $\log(STS(n)) \geq L_1 - L_2 = (n^2/6)(\log(n) - 2 \pm 2n^{-10^{-8}})$, which implies the stated bound on $STS(n)$. □

2.2. Strategy

The strategy of the proof of Theorem 2.1 is encapsulated by the following setup (we give motivation and discussion below). We say that $J \subseteq G$ is c -bounded if $|J(v)| < c|V(G)|$ for every $v \in V(G)$, where $J(v) = \{u \in V(G) : uv \in J\}$ is the *neighbourhood* of v in J .

Setup 2.3. Suppose we have $G^* \subseteq G$ with a ‘template’ triangle decomposition T such that

Nibble: $G \setminus G^*$ contains a set N of edge-disjoint triangles with ‘leave’ $L := (G \setminus G^*) \setminus \bigcup N$ that is c_1 -bounded.

Cover: For any $L \subseteq G \setminus G^*$ that is c_1 -bounded, there is a set M^c of edge-disjoint triangles such that $L = (G \setminus G^*) \cap \bigcup M^c$ and the ‘spill’ $S := G^* \cap \bigcup M^c$ is c_2 -bounded.

Hole: For any tridivisible $S \subseteq G^*$ that is c_2 -bounded, there are ‘outer’ and ‘inner’ sets M^o, M^i of edge-disjoint triangles in G^* such that $\bigcup M^o$ is c_3 -bounded and $(S, \bigcup M^i)$ is a partition of $\bigcup M^o$.

Completion Given L, M^c, M^o and M^i as above, there are sets M_1, M_2, M_3, M_4 of edge-disjoint triangles in G such that $(L, \bigcup M_2)$ is a partition of $\bigcup M_1, \bigcup M_3 = \bigcup M_4, M_3 \subseteq T$ and $M_2 \subseteq M_4$.

The key step is choosing T (which determines G^*). We will use our method of Randomised Algebraic Construction, which takes a particularly simple form for triangle decompositions. To motivate the construction, suppose that $V(G)$ is an abelian group, and consider the set Σ of triples xyz such that $x + y + z = 0$. We note that Σ is a good ‘model’ for a triangle decomposition, as for any xy there is a unique z such that $x + y + z = 0$. However, we cannot simply take Σ , as not all such xyz are triangles of G ; moreover, x, y, z may not even be pairwise distinct.

The idea of the construction is that a suitable random subset of Σ can act as a template, which covers a constant fraction of G . Next we find an approximate decomposition of the rest of G by random greedy algorithms: this is accomplished by steps **Nibble** and **Cover** of Setup 2.3. After these steps, every edge of G has been covered once or twice, and the spill S is the set of edges that have been covered twice. Finally, we use local modifications built into the template to turn the approximate decomposition into an exact decomposition: this is accomplished by steps **Hole** and **Completion** of Setup 2.3.

To motivate **Completion**, we imagine first that we have **Hole** and also $M^o \subseteq T$. Then we could delete M^o and take M^i instead, thus reducing by one the multiplicity of every edge in S , so that we have a triangle decomposition of G . However, specifying a triangle of T is very restrictive, as there are only order(n^2) such triangles out of a total of order(n^3) triangles in G . If we had chosen T uniformly at random it would be hopeless to obtain any useful configuration formed by triangles of T . However, the algebraic structure implies that certain configurations of triangles are dense within a sparse configuration space (described by linear constraints). This forms the basis of a modification procedure that replaces M^c, M^o and M^i by other sets of triangles with the same properties, where M_1 plays the role of $M^c \cup M^i$, M_2 of M^o , and each triangle f of M_2 can be embedded in a small subgraph that has one triangle decomposition (part of M_4) using f and another triangle decomposition (part of M_3) contained in T .

It is not hard to see that G contains a triangle decomposition in Setup 2.3. Indeed, we start by taking the sets N provided by **Nibble** and then the sets M^c and S provided by **Cover**. Now we note that $S = \bigcup T + \bigcup N + \bigcup M^c - G$ is tridivisible, as any integer linear combination of tridivisible graphs is tridivisible. So we can apply **Hole** to obtain M^o and M^i . Then we can apply **Completion** to obtain M_1, M_2, M_3, M_4 . Finally, $M = N \cup M_1 \cup (M_4 \setminus M_2) \cup (T \setminus M_3)$ is a triangle decomposition of G . Thus the remainder of the proof will be to show that we can achieve Setup 2.3.

2.3. Template

We choose the template as follows.

Construction 2.4. Let $a \in \mathbb{N}$ be such that $2^{a-2} < |V(G)| \leq 2^{a-1}$. Let $\pi : V(G) \rightarrow \mathbb{F}_{2^a} \setminus \{0\}$ be a uniformly random injection. Let

$$T = \{xyz \in K_3(G) : \pi(x) + \pi(y) + \pi(z) = 0\} \quad \text{and} \quad G^* = \bigcup T.$$

To avoid cumbersome notation, we use xyz to denote either the vertex set $\{x, y, z\}$ or the edge set $\{xy, xz, yz\}$ of a triangle. The context determines which interpretation is intended, e.g. in Construction 2.4 the graph G^* is the (disjoint) union of the edge-sets of the triangles in T .

In this subsection we will show that whp the pair (G, G^*) is ‘typical’ (in a precise sense defined below); this will allow us to implement the approximate decomposition in steps **Nibble** and **Cover**. Moreover, we will show in Section 4 that G^* is ‘linearly typical’ (roughly speaking: we can count subgraph extensions with linear constraints on the vertices); this will imply the existence of the local modifications used in steps **Hole** and **Completion**.

We start with some notation and preliminary observations. Throughout the paper we write $n = |V(G)|$. We identify G with its edge set $E(G)$, so that $|G|$ denotes the number of edges of G (rather than the number of vertices, as is used by some authors). We write $[n] = \{1, \dots, n\}$. We define

$$\gamma = 2^{-a}n,$$

and note that $1/4 < \gamma \leq 1/2$. We observe that if $x, y, z \in \mathbb{F}_{2^a} \setminus \{0\}$ and $x + y + z = 0$ then x, y, z are pairwise distinct. We note that $+1 = -1$ in \mathbb{F}_{2^a} , so we can use $+$ and $-$ interchangeably in \mathbb{F}_{2^a} -arithmetic. We consider \mathbb{F}_{2^a} as a vector space over \mathbb{F}_2 , and observe that any two nonzero elements span a subspace of dimension two.

Next we introduce the stronger typicality assumption used in [5]. We say that G is (c, h) -typical if

$$\left| \bigcap_{x \in S} G(x) \right| = (1 \pm |S|c)d(G)^{|S|}n \quad \text{for any } S \subseteq V(G) \text{ with } |S| \leq h.$$

Note that being c -typical is essentially the same as being $(c, 2)$ -typical (up to a factor of 2 in c). For most of the paper we will assume that G is $(c, 16)$ -typical; at the end we will explain how the proof can be modified to work with the weaker assumption that G is c -typical.

Now we define the typicality condition for (G, G^*) and show that it holds whp. Let G^* be a subgraph of G . We say that (G, G^*) is (c, h) -typical if

$$\left| \bigcap_{x \in S^*} G^*(x) \cap \bigcap_{x \in S \setminus S^*} G(x) \right| = (1 \pm |S|c)d(G^*)^{|S^*|}d(G)^{|S|-|S^*|}n$$

for any $S^* \subseteq S \subseteq V(G)$ with $|S| \leq h$. For a convenient statement of the following lemma we will assume that c is not too small; since the typicality conditions become stronger as c decreases, without loss of generality we can assume $c > c_0d(G)^{10^6}/2$.

Lemma 2.5. *Whp $d(G^*) = (1 \pm 3c)\gamma d(G)^3$ and (G, G^*) is $(6c, 16)$ -typical.*

The proof uses the following consequence of Azuma’s inequality.

Definition 2.6. Let S_n be the symmetric group, $f : S_n \rightarrow \mathbb{R}$ and $b \geq 0$. We say that f is b -Lipschitz if for any $\sigma, \sigma' \in S_n$ such that $\sigma = \tau \circ \sigma'$ for some transposition $\tau \in S_n$ we have $|f(\sigma) - f(\sigma')| \leq b$.

Lemma 2.7 (see e.g. [9]). Suppose $f : S_n \rightarrow \mathbb{R}$ is b -Lipschitz, $\sigma \in S_n$ is uniformly random and $X = f(\sigma)$. Then

$$\mathbb{P}(|X - \mathbb{E}X| > t) \leq 2e^{-t^2/2nb^2}.$$

Proof of Lemma 2.5. We start by estimating $\mathbb{E}|G^*| = \sum_{e \in G} \mathbb{P}(e \in G^*)$. For any $e = xy$, given $\pi(x)$ and $\pi(y)$, we have $e \in G^*$ if and only if $\pi(z) = \pi(x) + \pi(y)$ for some z such that $xyz \in K_3(G)$. Since G is $(c, 16)$ -typical, there are $(1 \pm 2c)d(G)^2n$ choices for z . Each satisfies $\pi(z) = \pi(x) + \pi(y)$ with probability $(2^a - 3)^{-1}$, so $\mathbb{E}|G^*| = |G|(1 \pm 2c)d(G)^2n(2^a - 3)^{-1}$. We can view π as $\sigma \circ \pi_0$, where $\pi_0 : V(G) \rightarrow \mathbb{F}_{2^a} \setminus \{0\}$ is any fixed injection and σ is a random permutation of $\mathbb{F}_{2^a} \setminus \{0\}$. Any transposition of σ affects $|G^*|$ by $O(n)$, so by Lemma 2.7, whp we have $d(G^*) = (1 \pm 2.1c)\gamma d(G)^3$.

Similarly, we consider any $S^* \subseteq S \subseteq V(G)$ with $|S| \leq 16$, write $Y = \bigcap_{x \in S^*} G^*(x) \cap \bigcap_{x \in S \setminus S^*} G(x)$, and estimate $\mathbb{E}|Y| = \sum_{y \in V(G)} \mathbb{P}(y \in Y)$. For any $y \in \bigcap_{x \in S} G(x)$, given $\pi(y)$ and $\pi(x)$ for all $x \in S$, we have $y \in Y$ if and only if for all $x \in S^*$ there is $xyz_x \in K_3(G)$ such that $\pi(z_x) = \pi(x) + \pi(y)$. Since G is $(c, 16)$ -typical, there are $(1 \pm |S|c)d(G)^{|S|}n$ choices for y . By excluding $O(1)$ choices of y we can assume $\pi(x) + \pi(y) \neq \pi(x')$ for all $x, x' \in S$. Then for each $x \in S^*$ there are $(1 \pm 2c)d(G)^2n$ choices for z_x , and for any set of choices, with probability $(1 + O(1/n))2^{-a|S^*|}$ they all satisfy $\pi(z_x) = \pi(x) + \pi(y)$. This gives

$$\mathbb{E}|Y| = O(1) + (1 \pm |S|c)d(G)^{|S|}n \cdot ((1 \pm 2c)d(G)^2n)^{|S^*|} \cdot (1 + O(1/n))2^{-a|S^*|}.$$

Any transposition of σ affects $|Y|$ by $O(1)$, so by Lemma 2.7, whp we have $|Y| = (1 \pm (3|S| + 1)c)d(G)^{|S|}(\gamma d(G)^2)^{|S^*|}n = (1 \pm 6|S|c)d(G^*)^{|S|}d(G)^{|S| - |S^*|}n$. \square

Since $d(G^*) = (1 \pm 3c)\gamma d(G)^3$ and $1/4 < \gamma < 1/2$, we have $0.24d(G)^3 < d(G^*) < 0.51d(G)$ for small c . Also, as (G, G^*) is $(6c, 16)$ -typical, we can deduce that $G \setminus G^*$ is $50c$ -typical. Indeed, for any $v \in V(G)$ we have

$$\begin{aligned} |(G \setminus G^*)(v)| &= (1 \pm c)d(G)n - (1 \pm 6c)d(G^*)n \\ &= (d(G) - d(G^*))n \pm 6c(d(G) + d(G^*))n = (1 \pm 20c)d(G \setminus G^*)n. \end{aligned}$$

Furthermore, for any $u, v \in V(G)$ we estimate $|(G \setminus G^*)(u) \cap (G \setminus G^*)(v)|$ as

$$\begin{aligned} |G(u) \cap G(v)| - |G^*(u) \cap G^*(v)| &= |G(u) \cap G^*(v)| + |G^*(u) \cap G^*(v)| \\ &= (1 \pm 2c)d(G)^2n - 2(1 \pm 12c)d(G)d(G^*)n + (1 \pm 12c)d(G^*)^2n \\ &= (d(G) - d(G^*))^2n \pm 12c(d(G) + d(G^*))^2n = (1 \pm 50c)d(G \setminus G^*)^2n. \end{aligned}$$

Applying the following theorem, we deduce **Nibble** with $c_1 = (50c)^{1/4}$.

Theorem 2.8. *There are $b_0 > 0$ and $n_0 \in \mathbb{N}$ such that if $n > n_0$, $n^{-0.1} < b < b_0$ and G is a b -typical graph on n vertices with $d(G) > b$, then there is a set N of edge-disjoint triangles in G such that $L = G \setminus \bigcup N$ is $b^{1/4}$ -bounded.*

We remark that the parameters in Theorem 2.8 are not very sharp: we have just fixed some convenient values that suffice for our purposes. Similar results are well-known, but we are not aware of any reference that implies the theorem as stated, so we will sketch a proof in Section 5.

For convenient reference, we give here the values of some other parameters that will be used below:

$$c_2 = 10^2 c_1 d(G)^{-6}, \quad c_3 = 10^{20} c_2 d(G)^{-50}$$

$$c_4 = 10^{20} c_3 d(G)^{-100}, \quad c_5 = 10^{10} c_4 d(G)^{-180}.$$

The tightest constraint on c that will be required in our calculations is

$$100c_5 = 10^{54} (50c)^{1/4} d(G)^{-336} < 10^{-6} d(G)^{180};$$

this holds for small c_0 if $c < c_0 d(G)^{3000}$. (This is the bound we need if G is $(c, 16)$ -typical, but if G is c -typical we need the stronger bound in Theorem 2.1.)

2.4. Cover

Consider the following random greedy algorithm. Let $L = \{e_i : i \in [t]\}$ (with edges ordered arbitrarily). Let $M^c = \{T_i : i \in [t]\}$ be triangles such that T_i consists of e_i and two edges of G^* , and is chosen uniformly at random from all such triangles that are edge-disjoint from all previous choices; if there is no available choice for T_i then the algorithm aborts.

To analyse the algorithm we require a concentration inequality. We say that a random variable Y is (μ, C) -dominated if there are constants μ_1, \dots, μ_m with $\sum_{i=1}^m \mu_i < \mu$, and we can write $Y = \sum_{i=1}^m Y_i$ with $|Y_i| \leq C$ for all i , and conditional on any given values of Y_j for $j < i$ we have $\mathbb{E}|Y_i| < \mu_i$. The following lemma follows easily from Freedman's inequality [3] (see [5, Lemma 2.7]).

Lemma 2.9. *If Y is (μ, C) -dominated then $\mathbb{P}(|Y| > 2\mu) < 2e^{-\mu/6C}$.*

Sometimes we will use a modified inequality with 2 replaced by $1 + c$. We also note that if the Y_i are independent (not necessarily identically distributed) indicator variables, we recover a version of the Chernoff bound for (pseudo)binomial variables (where better concentration is known). For the following lemma, we recall that L is c_1 -bounded, where $c_1 = (50c)^{1/4}$, and that $c_2 = 10^2 c_1 d(G)^{-6}$.

Lemma 2.10. *Whp the algorithm to choose M^c does not abort, and $S := G^* \cap \bigcup M^c$ is c_2 -bounded.*

Proof. For $i \in [t]$ we let \mathcal{B}_i be the bad event that $S_i := G^* \cap \bigcup_{j < i} T_j$ is not c_2 -bounded. We define a stopping time τ be the smallest i for which \mathcal{B}_i holds or the algorithm aborts, or ∞ if there is no such i . It suffices to show whp $\tau = \infty$.

We fix $t_0 \in [t]$ and bound $\mathbb{P}(\tau = t_0)$ as follows. For any $i < t_0$, since \mathcal{B}_i does not hold, S_i is c_2 -bounded. Writing $e_i = v_i v'_i$, we can bound the number of excluded choices for T_i by $c_2 n < |G^*(v_i) \cap G^*(v'_i)|/2$, so at most one half of the triangles on e_i are excluded.

Next we fix $e = vv' \in G^*$, and estimate $r_e := \sum_{i \leq t_0} \mathbb{P}'(e \subseteq T_i)$, where \mathbb{P}' denotes the conditional probability given the choices made before step i . We compare r_e to the expected number of times that e would be covered if we chose all triangles independently. To be precise, we let

$$E_e := \sum_{i \leq t_0} \mathbb{P}(e \subseteq T'_i),$$

where each T'_i is a uniform random triangle consisting of e_i and two edges of G^* , and $(T'_i : i \in [t])$ are independent. By the bound on excluded choices, $\mathbb{P}'(e \subseteq T_i) < 2\mathbb{P}(e \subseteq T'_i)$, so $r_e < 2E_e$.

The i th summand in E_e is only nonzero when $e_i \cap e \neq \emptyset$. As L is c_1 -bounded, the number of such i is at most $|L(v)| + |L(v')| < 2c_1 n$. Also, for each i such that $e_i \cup e$ spans a triangle, we have

$$\mathbb{P}(e \subseteq T'_i) = |G^*(v_i) \cap G^*(v'_i)|^{-1} < 2d(G^*)^{-2} n^{-1}.$$

Therefore $E_e < 4c_1 d(G^*)^{-2} < c_2/4$.

Finally, fix $v \in V(G)$ and consider

$$X = |S_{t_0}(v)| = \sum_{i \leq t_0} X_i, \quad \text{where } X_i = \sum_{v \in e \in G^*} 1_{e \subseteq T_i}.$$

We have $|X_i| \leq 2$ and

$$\sum_{i \leq t_0} \mathbb{E}'(X_i) = \sum_{i \leq t_0} \sum_{v \in e \in G^*} \mathbb{P}'(e \subseteq T_i) = \sum_{v \in e \in G^*} r_e < c_2 n/2.$$

By Lemma 2.9 we have $\mathbb{P}(X \geq c_2 n) < 2e^{-c_2 n/2^4}$. Taking a union bound over $i \leq t_0 \leq t$, we see that whp $|S(v)| < c_2 n$, i.e. S is c_2 -bounded and $\tau = \infty$. \square

Below we will require several more random greedy algorithms similar to that above. One could formulate an abstract general lemma to cover all cases [5, Lemma 4.11], but here we will prefer the more intuitive approach of identifying the key principles of the proof, so that it will be clear how it may be adapted to future instances. For a general random greedy algorithm, we identify some desired boundedness conclusion, then at each step of the algorithm, assuming that boundedness has not failed, we show that at most one half (say) of the choices of the required configuration have been excluded. Then for each edge e in the underlying graph H we estimate the expected number E_e of times that e would be covered if we chose all configurations independently. If $E_e < b/4$ and the configurations have constant size (not depending on n) then the graph of all covered edges is whp b -bounded.

We record some estimates that are useful for such arguments. Suppose H is a small fixed graph ($|H| \leq 500$ say), $F \subseteq V(H)$ and ϕ is an embedding of $H[F]$ in G^* . We call $E = (\phi, F, H)$ an *extension*. Let $X_E(G^*)$ be the number of embeddings ϕ^* of H in G^* that restrict to ϕ on F . We suppose that E is *16-degenerate*, meaning that we can construct the embedding one vertex at a time, so that at each step we add a vertex adjacent to at most 16 existing vertices. As (G, G^*) is $(6c, 16)$ -typical, when we add a vertex adjacent to $t \leq 16$ existing vertices, there are $(1 \pm 6tc)d(G^*)^t n$ choices. Multiplying these estimates, we obtain the following estimate for $X_E(G^*)$.

Lemma 2.11. *Suppose $E = (\phi, F, H)$ is a 16-degenerate extension with $|H| \leq 500$. Then*

$$X_E(G^*) = (1 \pm 7|H|c)d(G^*)^{|H \setminus H[F]|} n^{|V(H)| - |F|}.$$

Now suppose that we wish to exclude embeddings ϕ^* that use some edge in J which is c -bounded. Fix $e \in H \setminus H[F]$ and consider the embeddings ϕ^* with $\phi^*(e) \in J$. If $e \cap F \neq \emptyset$ there are at most cn choices for the embedding of e , and then at most $n^{|V(H)| - |F| - 1}$ choices for the remainder of ϕ^* . If $e \cap F = \emptyset$ there are at most cn^2 choices for the embedding of e , and then at most $n^{|V(H)| - |F| - 2}$ choices for the remainder of ϕ^* . Thus at most $|H|cn^{|V(H)| - |F|}$ choices of ϕ^* are excluded, which is a negligible fraction of $X_E(G^*)$.

3. Integral relaxations

In this section we establish **Hole**. Our first step is to consider an integral relaxation, in the following sense. Instead of thinking of $(S, \bigcup M^i)$ as a partition of $\bigcup M^o$, we think of S as a weighted sum of edge sets of triangles, where triangles in M^o have weight 1 and triangles in M^i have weight -1 . We can express this by the equation $\Phi A = S$, where Φ is the corresponding ± 1 -vector indexed by triangles, and A is the inclusion matrix of triangles against edges, i.e. $A_{fe} = 1_{e \subseteq f}$ for any edge e and triangle f . It is straightforward to show that this equation has a solution if we allow Φ to have any integer weights on triangles (see [4, 13, 14] for more general results).

It will be more convenient to work with linear maps rather than matrices. For any graph H we define \mathbb{Z} -linear boundary/shadow maps $\partial_j : \mathbb{Z}^{K_i(H)} \rightarrow \mathbb{Z}^{K_j(H)}$ for $i \geq j \geq 0$ by $\partial_j(e) = \sum \binom{e}{j}$ for $e \in K_i(H)$, i.e. for $J \in \mathbb{Z}^{K_i(H)}$ and $f \in K_j(H)$ we define $\partial_j(J)_f = \sum_{f \subseteq e \in K_i(H)} J_e$. For example, if $J \in \mathbb{Z}^H$ then $\partial_1(J) \in \mathbb{Z}^{V(H)}$ is defined by $\partial_1(J)_v = \sum_{v \in e \in H} J_e$.

It will also be notationally convenient to identify vectors with (generalised) sets. It is standard to identify $v \in \{0, 1\}^X$ with the set $\{x \in X : v_x = 1\}$. Similarly, we can identify $v \in \mathbb{N}^X$ with the multiset in X in which x has multiplicity v_x (for our purposes $0 \in \mathbb{N}$). We also apply similar notation and terminology to that for multisets to vectors $v \in \mathbb{Z}^X$ ('intssets'). Here our convention is that 'for each $x \in v$ ' means that x is considered $|v_x|$ times in any statement or algorithm, and has a sign attached to it (the same as that of v_x); we also refer to x as a 'signed element' of v . For $v \in \mathbb{Z}^X$ we write $v = v^+ - v^-$, where $v_x^+ = \max\{v_x, 0\}$ and $v_x^- = \max\{-v_x, 0\}$ for $x \in X$. Given $J \in \mathbb{N}^G$ and $v \in V(G)$,

we define $J(v) \in \mathbb{N}^{V(G)}$ by $J(v)_u = 1_{uv \in G} J_{uv}$. Then we can extend the definition of boundedness to multigraphs: J is c -bounded if $|J(v)| < cn$ for every $v \in V(G)$.

With this notation, our integral relaxation of **Hole** is expressed by the following lemma (in which K_n denotes the complete graph on $V(G)$); for **Hole** we will need the additional properties that $\Phi(f) = 0$ for any $f \in K_3(K_n) \setminus K_3(G^*)$, and $\Phi(f) \in \{0, 1, -1\}$ for all $f \in K_3(G^*)$, as then we can write $\Phi = M^o - M^i$.

Lemma 3.1. *There is $\Phi \in \mathbb{Z}^{K_3(K_n)}$ with $\partial_2 \Phi = S$ such that $\partial_2 \Phi^+$ is $100c_2$ -bounded.*

Proof. We will construct $\Phi = \Phi_0 + \Phi_1 + \Phi_2$ such that $J^0 = S - \partial_2 \Phi_0$, $J^1 = J^0 - \partial_2 \Phi_1$, $J^2 = J^1 - \partial_2 \Phi_2$ satisfy $\partial_i J^i = 0$ for $i = 0, 1, 2$. Recalling that S is tridivisible, each J^i will be tridivisible, in the ‘intgraph’ sense: i.e. $\sum_e J_e^i$ is divisible by 3 and $\sum_u J_{uv}^i$ is divisible by 2 for all v .

Step 0: For Φ_0 , we choose $|S|/3$ independent uniformly random triangles in K_n ; then $J^0 = S - \partial_2 \Phi_0$ satisfies $\partial_0 J^0 = 0$. For each vertex v , the number of these triangles containing v is binomial with mean $|S|/n < c_2 n/2$, so by the Chernoff bound whp $\partial_2 \Phi_0$ is $1.1c_2$ -bounded.

Step 1: We let $J^* = \partial_1 J^0$, so $\partial_0 J^* = 2\partial_0 J^0 = 0$, i.e. $|J^{*+}| = |J^{*-}|$. Note for all $x \in V(G)$ that J_x^* is even, as J^0 is tridivisible, and $|J_x^*| < 1.1c_2 n$. We fix an arbitrary sequence $((x_i^+, x_i^-) : i \in [|J^{*+}|/2])$ so that each $x \in V(G)$ occurs $J_x^{*+}/2$ times as some x_i^+ and $J_x^{*-}/2$ times as some x_i^- . For each i we choose $a_i b_i \subseteq V(G) \setminus \{x_i^+, x_i^-\}$ independently uniformly at random, and let $\Phi_1 = \sum_{i \in [|J^{*+}|/2]} (\{x_i^+ a_i b_i\} - \{x_i^- a_i b_i\})$; then $J^1 = J^0 - \partial_2 \Phi_1$ satisfies $\partial_1 J^1 = 0$.

We claim that whp $\partial_2 \Phi_1^\pm$ are $8c_2$ -bounded. To see this, we first fix any $e \in K_n$ and estimate the expected contributions to e from each step i , according to whether e contains x_i^+ , x_i^- , or neither. Each endpoint of e occurs at most $0.6c_2 n$ times as x_i^\pm , and for such i we cover e with probability $2/(n-2)$, so the expected contribution to $(\partial_2 \Phi_1^\pm)_e$ from all such i is at most $2.5c_2$. At any other step, we cover e with probability $\binom{n-2}{2}^{-1}$, so the total expected contribution to $(\partial_2 \Phi_1^\pm)_e$ from these steps is at most $1.1c_2$. Now, for each vertex v , summing over its incident edges, we see that $|\partial_2 \Phi_1^\pm(v)|$ are both $(4c_2 n, 1)$ -dominated, so the claim holds by Lemma 2.9.

Step 2: We start by fixing an arbitrary expression $J^1 = \sum_{C \in \mathcal{C}_0} C$, where each C is a closed walk in K_n with edge weights alternating between 1 and -1 , and there are no cancellations, i.e. every edge appears in the sum only with weight 1 or only with weight -1 . As is well-known, such an expression may be found by a greedy algorithm: each C can be obtained by following an arbitrary alternating walk on the signed elements of J^1 until we return to our starting point using an edge with the opposite sign to that of the first edge, whereupon we add $-C$ to J^1 and repeat the procedure. (We note that this argument leads to a convenient shortcut for triangle decompositions, but does not generalise to hypergraph decompositions.)

Next we express each $C \in \mathcal{C}_0$ as a sum of signed four-cycles in the complete graph K_n on $V(G)$, where we write each closed walk of length $2m$ as a chain of $m-1$ signed four-




Fig. 1. Decomposing even signed cycles.

cycles, using the identity (see Figure 1)

$$\begin{aligned} & \sum_{i=1}^{m-1} (-1)^i (\{x_i x_{i+1}\} - \{x_{i+1} y_{i+1}\} + \{y_{i+1} y_i\} - \{y_i x_i\}) \\ &= \{x_1 y_1\} + (-1)^m \{x_m y_m\} + \sum_{i=1}^{m-1} (-1)^i \{x_i x_{i+1}\} + \sum_{i=1}^{m-1} (-1)^i \{y_i y_{i+1}\}. \end{aligned}$$

This identity can be used as is if $x_i \neq y_i$ for $i \in [m]$. For each i such that $x_i = y_i$, we note that $1 < i < m$, $x_{i-1} \neq y_{i-1}$, $x_{i+1} \neq y_{i+1}$, and $x_{i+1} \neq y_{i-1}$, so we can replace the four-cycles for summands $i - 1$ and i by

$$\begin{aligned} & (-1)^{i-1} (\{x_{i-1} x_i\} - \{x_i x_{i+1}\} + \{x_{i+1} y_{i-1}\} - \{y_{i-1} x_{i-1}\}), \quad \text{and} \\ & (-1)^i (\{x_{i+1} y_{i-1}\} - \{y_{i-1} y_i\} + \{y_i y_{i+1}\} - \{y_{i+1} x_{i+1}\}). \end{aligned}$$

Thus we can write $J^1 = \sum_{C \in \mathcal{C}} C$, where each summand is a signed four-cycle in K_n . Furthermore, the above construction has the property that for each $v \in V(G)$ and $w \in \{-1, 1\}$ we use at most $3|J^{1+}(v)| < 24c_2n$ edges at v with weight w .

For each $C = \{ab\} - \{bc\} + \{cd\} - \{da\} \in \mathcal{C}$ we choose $x \in V(G) \setminus \{a, b, c, d\}$ independently uniformly at random, and add $\{xab\} - \{xbc\} + \{xcd\} - \{xda\}$ to Φ_2 ; then $\partial_2 \Phi_2 = \sum_{C \in \mathcal{C}} C = J^1$. Let Γ denote the multigraph formed by summing $\{xa, xb, xc, xd\}$ over all such C . For any $e \in K_n$, at most $48c_2n$ elements of \mathcal{C} can contribute to Γ_e , so $\mathbb{E}\Gamma_e < 49c_2n$. Then for any v , summing over its incident edges, we see that $|\Gamma(v)|$ is $(49c_2n, 4)$ -dominated, so by Lemma 2.9 (modified), whp Γ is $50c_2$ -bounded. Defining $\Phi = \Phi_0 + \Phi_1 + \Phi_2$, we have $\partial_2 \Phi = S$ and $\partial_2 \Phi^+$ is $100c_2$ -bounded. \square

To obtain **Hole**, we will modify Φ using the following ‘octahedral’ configurations (see Figure 2). Consider a copy of $K_{2,2,2}$, the complete tripartite graph with two points in each part, with parts $\{(j, 0), (j, 1)\}$ for $j \in [3]$. Its triangles are $\{f_x : x \in \{0, 1\}^3\}$, where $f_x = \{(j, x_j) : j \in [3]\}$. The sign of f_x is $s(f_x) = (-1)^{\sum x}$. Thus each edge is in one triangle of each sign. Defining $\Omega = \sum_{x \in \{0,1\}^3} s(f_x) \{f_x\} \in \mathbb{Z}^{K_3(K_{2,2,2})}$, we see that $\partial_2 \Omega = 0$. This gives a method to eliminate any signed triangle f from Φ without altering $\partial_2 \Phi$: we add some copy of Ω with the opposite sign to f in which (say) $f_{000} = f$, thus replacing f by seven other signed triangles that have the same total 2-shadow. Similarly (and more importantly), we can eliminate any pair of triangles f, f' that have opposite sign and share an edge e , replacing f, f' by six other signed triangles that have the same total 2-shadow and do not use e . We apply this method in the following two-phase algorithm.




Fig. 2. An octahedron with signed triangles.

Octahedral Elimination Algorithm (Phase I). We eliminate all triangles in Φ , according to a random greedy algorithm, where in each step we consider some original signed element f of Φ , and choose an octahedral configuration Ω_f to replace f . We refer to edges of Ω_f not in f as *new edges*, and choose Ω_f uniformly at random subject to the new edges belonging to G^* and being disjoint from $\partial_2\Phi^+$ and all new edges from previous steps.

Let Φ' denote the result of Phase I (if it does not abort). Then $\partial_2\Phi' = \partial_2\Phi = S$, and we can write $\partial_2\Phi'^+ = \partial_2\Phi^+ + \Gamma$, where Γ is the graph of new edges, and every signed element of Φ' contains at most one edge of $\partial_2\Phi^+$.

Octahedral Elimination Algorithm (Phase II). We replace all signed edges apart from those in S and Γ . To do this, we fix a sequence \mathcal{S} of pairs of signed elements of Φ' , so that (i) for each $ff' \in \mathcal{S}$, there is some $e \in \partial_2\Phi^+$ such that f and f' both contain e , and f and f' have opposite signs, and (ii) the multiset consisting of all e as in (i) is $\partial_2\Phi^-$. Now we eliminate each $ff' \in \mathcal{S}$, according to a random greedy algorithm, by subtracting some copy $\Omega_{ff'}$ of Ω with $f_{000} = f$ and $f_{001} = f'$, or vice versa, depending on the signs. We refer to edges of $\Omega_{ff'}$ not in f or f' as *new edges*, and choose $\Omega_{ff'}$ uniformly at random subject to the new edges belonging to G^* and being distinct from $\partial_2\Phi^+ \cup \Gamma$ and all new edges from previous steps.

Let Ψ denote the result of this algorithm (if it does not abort) and Γ' the graph of new edges for Phase II. Then $\partial_2\Psi = S$ and $\partial_2\Psi^- = \Gamma \cup \Gamma' \subseteq G^*$. This implies $\Psi(f) = 0$ for any $f \in K_3(K_n) \setminus K_3(G^*)$, and $\Psi(f) \in \{0, 1, -1\}$ for all $f \in K_3(G^*)$, so $\Psi = M^o - M^i$, where M^o and M^i are as in **Hole**, once we have verified the boundedness condition.

Lemma 3.2. *Whp the Octahedral Elimination Algorithm produces M^o and M^i as in **Hole**.*

Proof. We first show that whp Γ is c'_2 -bounded, where $c'_2 = 10^5 c_2 d(G^*)^{-9}$. The proof follows the discussion after the proof of Lemma 2.10, where a configuration for f consists of the new edges of some Ω_f . By Lemma 2.11, at each step, the number of choices of Ω_f with all new edges belonging to G^* (with no excluded configurations) is $(1 \pm 60c)d(G^*)^9 n^3$. Assuming that the graph of previous new edges is c'_2 -bounded, as $\partial_2\Phi^+$ is $100c_2$ -bounded, the number of excluded configurations is at most $10c'_2 n^3$, which

is less than half of the total. Next, for each $e \in G^*$, we consider separately the contributions to E_e , according to whether e intersects f in 0 or 1 vertex (there is no contribution to new edges from triangles containing e). There are at most $600c_2n$ signed elements of Φ that intersect e in one vertex. For each of these, a random configuration covers e with probability at most $3n^2/(1 - 60c)d(G^*)^9n^3$, so the total contribution to E_e from such elements is at most $2000c_2d(G^*)^{-9}$. Also, Φ has at most $100c_2n^2$ signed elements, and for each one that is disjoint from e the contribution to E_e is at most $6n/(1 - 60c)d(G^*)^9n^3$, so the total contribution from such elements is at most $1000c_2d(G^*)^{-9}$. We obtain $E_e < 3000c_2d(G^*)^{-9}$, which implies the claimed bound on Γ .

Next we claim that whp Γ' is c_2'' -bounded, where $c_2'' = 20c_2'd(G^*)^{-7}$. The argument is very similar to that given for Γ . Now a configuration for ff' consists of the new edges of some $\Omega_{ff'}$. By Lemma 2.11, at each step, the number of choices of $\Omega_{ff'}$ with all new edges belonging to G^* (with no excluded configurations) is $(1 \pm 50c)d(G^*)^7n^2$. Assuming that the graph of previous new edges is c_2'' -bounded, as $\partial_2\Phi^+ \cup \Gamma$ is $2c_2'$ -bounded, the number of excluded configurations is at most $10c_2''n^2$, which is less than half of the total. Next, for each $e \in G^*$, we consider separately the contributions to E_e according to whether e intersects $f \cup f'$ in 0 or 1 vertex (there is no contribution to new edges if $e \subseteq f \cup f'$).

First we consider those $ff' \in \mathcal{S}$ that intersect e in one vertex x . There are two choices for $x \in e$. If $x \in f \cap f'$ then there are at most $200c_2n$ choices for $f \cap f'$ in $\partial_2\Phi^+ \cup \partial_2\Phi^-$, which determines f and f' . If $\{x\} = f \setminus f'$ then there are at most $|\Gamma(x)| < c_2'n$ choices for f , and so f' . The same bound applies if $\{x\} = f' \setminus f$, so there are at most $5c_2'n$ such ff' . Each contributes at most $2n/(1 - 50c)d(G^*)^7n^2$ to E_e , so the total contribution from such ff' is at most $11c_2'd(G^*)^{-7}$. Also, $|\mathcal{S}| = |\partial_2\Phi^-| < 100c_2n^2$, and for each $ff' \in \mathcal{S}$ with $e \cap (f \cup f') = \emptyset$ the contribution to E_e is at most $2/(1 - 50c)d(G^*)^7n^2$, so the total contribution from such elements is at most $300c_2d(G^*)^{-7}$. We obtain $E_e < 12c_2'd(G^*)^{-7}$, which implies the claimed bound on Γ' . Recalling that $d(G^*) > 0.24d(G)^3$ and $c_3 = 10^{20}c_2d(G)^{-50}$ we see that $\bigcup M^o = \partial_2\Psi^+ = \mathcal{S} \cup \Gamma \cup \Gamma'$ is c_3 -bounded, so we have the required properties for **Hole**. \square

4. Completion

For **Completion**, we divide the analysis into two parts. Firstly, we will determine what conditions on M_1 and M_2 enable us to find M_3 and M_4 . Secondly, we will show that the sets M^c , M^o and M^i from **Cover** and **Hole** can be modified to give M_1 and M_2 satisfying the required conditions. For convenient notation we suppress the embedding $\pi : V(G) \rightarrow \mathbb{F}_{2^a}$ whenever we do not need to refer to it, instead thinking of $V(G)$ as a subset of \mathbb{F}_{2^a} .

4.1. Shuffles

Suppose we have a set M_2 of edge-disjoint triangles in G^* , and we want to find sets M_3 and M_4 of edge-disjoint triangles in G^* such that $\bigcup M_3 = \bigcup M_4$, $M_3 \subseteq T$ and

$M_2 \subseteq M_4$. Our basic building blocks (‘shuffles’) will be edge-disjoint subgraphs of G^* , each having two different triangle decompositions, one only using triangles in T , and the other including any specified triangle of M_2 . Then the unions over all blocks of the two triangle decompositions will give M_3 and M_4 as required.

We define the shuffles as follows. Fix $x = (x_1, x_2, x_3) \in \mathbb{F}_{2^a}^3$ and $t = (t_1, t_2) \in \mathbb{F}_{2^a}^2$ such that $\{x_1, x_2, x_3, t_1, t_2\}$ is linearly independent over \mathbb{F}_2 . Let $\langle x \rangle$ be the subspace of \mathbb{F}_{2^a} generated by $\{x_1, x_2, x_3\}$. The xt -shuffle S_{xt} is the complete tripartite graph with parts $t_i + \langle x \rangle = \{t_i + y : y \in \langle x \rangle\}$, $i \in [3]$, where $t_3 := t_1 + t_2$. If $S_{xt} \subseteq G^*$ then it has a triangle decomposition M_{3xt} only using triangles in T : take all triangles $y_1y_2y_3$ where each $y_i \in t_i + \langle x \rangle$ and $y_1 + y_2 + y_3 = 0$. We define another triangle decomposition M_{4xt} of S_{xt} by translating each triangle of M_{3xt} by (x_1, x_2, x_3) , i.e. M_{4xt} consists of all triangles $y_1y_2y_3$ where $y_i \in t_i + \langle x \rangle$ for each i and $x_1 + x_2 + x_3 + y_1 + y_2 + y_3 = 0$.

To construct M_3 and M_4 , we choose shuffles according to a random greedy algorithm, where in each step we consider some $z_1z_2z_3 \in M_2$, and choose some shuffle $S_{xt} \subseteq G^*$ such that $z_i = t_i + x_i$ for all $i \in [3]$. We will see in Lemma 4.2 that the Randomised Algebraic Construction is whp such that there are many choices for such a shuffle. This is the most important property of the construction, and it would not hold if we had chosen the template to be a uniformly random set of edge-disjoint triangles; in fact the expected number of shuffles (or any ‘shuffle-like’ configuration) would be $o(1)$. First we identify a property that we need for triangles in M_2 so that the required shuffles exist and can be chosen to be edge-disjoint. We say that $z_1z_2z_3$ is *octahedral* if $z_1 + z_2 + z_3 \neq 0$ and there is a copy K' of $K_{2,2,2}$ in G such that $\pi(K')$ has parts $\{z_1, z_2 + z_3\}$, $\{z_2, z_1 + z_3\}$ and $\{z_3, z_1 + z_2\}$; we call K' the *associated octahedron* of $z_1z_2z_3$. We assume

(P1) all triangles in M_2 are octahedral, with edge-disjoint associated octahedra.

Remark 4.1. The associated octahedron has all the properties that we require for the construction of M_3 and M_4 , so we could implement our algorithm without using shuffles. This remark was communicated to the author by Yang, and independently by Glebov and Luria. We have opted to keep the shuffle argument in this paper, as it indicates how to treat general (hyper)graphs (we only see how to dispense with it for triangles), and also illustrates the arguments needed for Subsection 4.4.

Lemma 4.2. *Under the random choice of π used in the definition of T , whp for any octahedral $z_1z_2z_3$ there are $(1 \pm 200c)d(G)^{180}\gamma^{18}2^{2a}$ shuffles $S_{xt} \subseteq G^*$ such that $t_i + x_i = z_i$ for $i \in [3]$.*

Proof. We can write the number of such shuffles as a sum of indicator variables, $X = \sum 1_{E(K, \ell, x, t)}$, where the sum ranges over all (K, ℓ, x, t) such that K is a copy of $K_{8,8,8}$ in G containing the associated octahedron K' of $z_1z_2z_3$, ℓ is a bijective labelling of each part of K by \mathbb{F}_2^3 , we let $E(K, \ell, x, t)$ be the event that $\pi(w) = t_i + \ell(w) \cdot x$ for all $i \in [3]$ and w in the i th part of K , and we assume ℓ is consistent with K' , in that $\ell(\pi^{-1}(z_i)) = e_i$ and $\ell(\pi^{-1}(z_i + z_j)) = e_i + e_j$ for $\{i, j\} \subseteq [3]$.

As G is $(c, 16)$ -typical, there are $(1 \pm 181c)d(G)^{180}n^{18}$ choices of (K, ℓ) . There are $2^{2a} - O(n)$ choices of t , which determines x given z , as only $O(n)$ choices of t are excluded by the condition that $\{x_1, x_2, x_3, t_1, t_2\}$ is linearly independent over \mathbb{F}_2 : there are

$O(1)$ possible linear relations between them, and each such relation is linearly independent or contradictory to the system $t_i + x_i = z_i$ for $i \in [3]$ (as $z_1 + z_2 + z_3 \neq 0$), so is satisfied by at most 2^a choices of t . Given (K, ℓ, x, t) , conditional on $\pi|_{K'}$, we have $\mathbb{P}(E(K, \ell, x, t)) = (1 + O(1/n))2^{-18a}$. Therefore $\mathbb{E}X = (1 \pm 182c)d(G)^{180}\gamma^{18}2^{2a}$.

Also, any transposition τ of π affects X by at most $100 \cdot 2^a$. To see this, we estimate the number of shuffles containing $z_1z_2z_3$ and any fixed $v \in \mathbb{F}_{2^a} \setminus \{z_1, z_2, z_3, z_1 + z_2, z_1 + z_3, z_2 + z_3\}$. Consider any $j \in [3]$, $b \in \mathbb{F}_2^3 \setminus \{e_j, (1, 1, 1) - e_j\}$, and the equations $t_i + b \cdot x = v$ and $t_i + x_i = z_i$ for $i \in [3]$ in (t, x) . We have four linearly independent constraints, so there are at most 2^a solutions. Including multiplicative factors for i, b and τ gives the required bound. Now by Lemma 2.7 whp $X = (1 \pm 200c)d(G)^{180}\gamma^{18}2^{2a}$. \square

4.2. Linear extensions

We digress to note a more general estimate for future reference. Suppose H is a graph, $y = (y_i : i \in [g])$ are variables, and for all $v \in V(H)$ we have distinct linear forms $L_v(y) = c_v + \sum_{i \in S_v} y_i$ for some $c_v \in \mathbb{F}_{2^a}$ and $S_v \subseteq [g]$. We call $E = (L, H)$ a linear extension with base $F = \{v \in V(H) : S_v = \emptyset\}$. Let $X_E(G^*)$ be the number of L -embeddings of H , i.e. embeddings ϕ of H in G^* such that for some $y \in \mathbb{F}_{2^a}^g$ we have $\phi(v) = L_v(y)$ for all $v \in V(H)$. The above argument (see also [5, Lemma 5.15]) gives the following formula analogous to that obtained for shuffles.

Lemma 4.3. *Let $E = (L, H)$ be a 16-degenerate linear extension with $|H| \leq 500$. Suppose*

- H has a triangle decomposition M such that $L_x + L_y = L_z$ for each $xyz \in M$,
- the incidence matrix of $\{S_v : v \in V(H)\}$ has full column rank $g \geq 1$.

Then

$$X_E(G^*) = (1 \pm 1.1|H|c)d(G)^{|H \setminus H[F]|}\gamma^{|V(H) \setminus F|}2^{ga}.$$

4.3. Shuffle algorithm

Recalling our general framework for random greedy algorithms, we want to show that, of the potential shuffles S_{xt} with $t_i + x_i = z_i$ for $i \in [3]$, at most half are excluded due to sharing an edge with a previous shuffle, assuming some boundedness condition on the graph Γ of new edges from previous shuffles. We classify the potential restrictions according to the label of the shuffle edge involved, which is specified by some $\{j, k\} \subseteq [3]$ and $b_j, b_k \in \mathbb{F}_2^3$ such that $b_j \notin \{e_j, (1, 1, 1) - e_j\}$ or $b_k \notin \{e_k, (1, 1, 1) - e_k\}$ (here we do not consider edges of the associated octahedra: these are already determined, and edge-disjoint by (P1)). For any $v_jv_k \in G^*$, the shuffles excluded because of mapping the given labelled shuffle edge to v_jv_k are given by the (x, t) -solutions of the system \mathcal{S} of equations $t_j + b_j \cdot x = v_j, t_k + b_k \cdot x = v_k$ and $t_i + x_i = z_i$ for $i \in [3]$. There may be 0, 1 or 2^a solutions. We can ignore the case of no solutions, as it does not exclude anything. For the cases with one solution, we can bound the number of excluded choices by the number of edges covered by all shuffles, which is $192|M_2|$.

It remains to consider the case that \mathcal{S} has 2^a solutions, which occurs when one of the equations is redundant, due to being a linear combination of the other equations. There are a constant number of linear combinations, and each constrains (v_j, v_k) to lie on a line, as may be seen from general considerations of linear algebra, or simply by enumerating the possibilities: without loss of generality $t_k + b_k \cdot x = v_k$ is redundant, due to

- (i) $b_k = e_k$ and $v_k = z_k$,
- (ii) $b_k = (1, 1, 1) - e_k$ and $v_k = z_1 + z_2 + z_3 - z_k$,
- (iii) $b_j + b_k = e_j + e_k$ and $v_j + v_k = z_j + z_k$,
- (iv) $b_j + b_k = e_i$ and $v_j + v_k = z_i$, where $[3] = \{i, j, k\}$.

In cases (i) and (ii) where v_k is fixed, assuming that Γ is c_5 -bounded, there are at most $c_5 n$ choices for v_j such that $v_j v_k \in \Gamma$. In cases (iii) and (iv) we need an additional boundedness condition:

We say that Γ is *linearly c_5 -bounded* if Γ is c_5 -bounded and also contains at most $c_5 2^a$ edges from any line of the form $\{(x_1 + \mu, x_2 + \mu) : \mu \in \mathbb{F}_{2^a}\}$.

We also need similar conditions so that we can avoid the associated octahedra; writing Δ for the union of all associated octahedra of triangles in M_2 , we will ensure that

(P2) Δ is linearly c_4 -bounded.

Then the total number of excluded shuffles is at most $192(|M_2| + (c_4 + c_5)2^{2a}) < 200c_5 2^{2a}$, which is less than half of the total.

Next we fix $e \in G^*$ and estimate E_e . To do so, we fix b_j, b_k as above, write $e = v_j v_k$ and estimate the sum over $z_1 z_2 z_3 \in M_2$ of the probability p that a random shuffle S_{xt} with $t_i + x_i = z_i$ for $i \in [3]$ satisfies $t_j + b_j \cdot x = v_j$ and $t_k + b_k \cdot x = v_k$. For fixed $z_1 z_2 z_3$, if the system \mathcal{S} as above has N solutions then $p = N / (1 \pm 200c)d(G)^{180} \gamma^{18} 2^{2a}$. When $N = 1$ the total contribution is at most $|M_2| / (1 - 200c)d(G)^{180} \gamma^{18} 2^{2a} < 1.1c_4 d(G)^{-180} \gamma^{-18}$. If $N = 2^a$ then (z_1, z_2, z_3) is constrained to lie in a certain plane (this can be seen by linear algebra, or by considering each possibility as above: e.g. in case (iii) the plane is $v_j + v_k = z_j + z_k$). Thus we see the final property that we need from M_2 :

(P3) M_2 contains at most $c_4 2^a$ elements $z_1 z_2 z_3$ from any *basic plane* of the form $b \cdot z = v$ where $b \in \mathbb{F}_2^3 \setminus \{0\}$.

(Note that by (P1) we can assume $v \neq 0$ in (P3).) Then the total contribution is at most $c_4 2^a \cdot 2^a / (1 - 200c)d(G)^{180} \gamma^{18} 2^{2a}$. Summing over $\{j, k\}$, b_j and b_k , we can estimate $E_e < 250c_4 d(G)^{-180} \gamma^{-18} = c_5/4$. Applying Lemma 2.9 as in the proof of Lemma 2.10, we deduce that whp the boundedness assumptions on Γ used above do not fail (linear boundedness follows in the same way as boundedness), and so the algorithm does not abort. This completes the analysis of the first part of **Completion**: given M_1 and M_2 as in **Completion**, under the conditions (P1)–(P3) on M_2 , we can find M_3 and M_4 as in **Completion**.

4.4. Octahedral Elimination Algorithm

To complete the proof of **Completion**, and so of the theorems, it remains to show that we can find M_1 and M_2 satisfying the conditions (P1)–(P3). We apply a similar two-phase algorithm to that used in **Hole**.

Phase I. We start with $\Phi = M^c + M^i - M^o$, so $\partial_2\Phi = L$, $\partial_2\Phi^+ = \bigcup(M^c \cup M^i)$, $\partial_2\Phi^- = \bigcup M^o$. Next we eliminate all triangles in Φ according to a random greedy algorithm, where in each step we consider some original signed element f of Φ , and choose an octahedral configuration Ω_f to replace f . We say that a triangle f' of Ω_f is *far* if $|f' \cap f| \leq 1$, and that Ω_f is *valid* if (i) none of its triangles are template triangles, with the possible exception of f , and (ii) all of its far triangles are octahedral, and their associated octahedra share edges only in Ω_f , in which case we denote their union by the extended configuration Ω_f^+ . We say that an edge of Ω_f^+ not in f is *new*, and choose a valid Ω_f uniformly at random subject to the new edges being distinct from all new edges from previous steps.

Let Φ' denote the result of Phase I (if it does not abort). We have $\partial_2\Phi' = \partial_2\Phi = L$, and writing Γ for the graph of new edges, we see that every signed element of Φ' is either a far triangle consisting of three edges of Γ , or is not far and consists of two edges of Γ and one edge of $\partial_2\Phi^+$.

Phase II. Now we will eliminate all triangles of Φ' apart from those that contain an edge of L or were far in the previous modification procedure. We partition all such triangles into a sequence \mathcal{S} of pairs of signed elements of Φ' , so that for each $ff' \in \mathcal{S}$, there is some $e \in \partial_2\Phi^+$ such that f and f' both contain e , and f and f' have opposite signs. We eliminate each $ff' \in \mathcal{S}$, according to a random greedy algorithm, by subtracting some copy $\Omega_{ff'}$ of Ω with $f_{000} = f$ and $f_{001} = f'$, or vice versa, depending on the signs. Now we say that $\Omega_{ff'}$ is *valid* if all of its triangles apart from f and f' are octahedral, and their associated octahedra share edges only in $\Omega_{ff'}$, in which case we denote their union by the extended configuration $\Omega_{ff'}^+$. We refer to edges of $\Omega_{ff'}^+$ not in f or f' as *new* edges, and choose a valid $\Omega_{ff'}$ uniformly at random subject to the new edges being distinct from Γ and all new edges from previous steps.

Let Ψ denote the result of this algorithm (if it does not abort) and Γ' the graph of new edges for Phase II. Since $\partial_2\Psi = \partial_2\Phi = L$, defining $M_1 = \Psi^+$ and $M_2 = \Psi^-$, we see that $\bigcup M_2 = \Gamma \cup \Gamma'$ and $\bigcup M_1 = L \cup \Gamma \cup \Gamma'$, so $(L, \bigcup M_2)$ is a partition of $\bigcup M_1$. The following lemma completes the proof of **Completion**, and so of the theorems, under the assumption that G is $(c, 16)$ -typical.

Lemma 4.4. *Whp M_2 satisfies (P1)–(P3).*

Proof. To analyse Phase I, we first estimate the number of choices for an extended configuration on a triangle f . This can be described by the linear extension (Ω_f^+, L) , where Ω_f^+ is as above, we have variables $z = (z_1, z_2, z_3)$, which we also use to label the vertices of $\Omega_f \setminus f$, we define $L_x = x$ for all $x \in \Omega_f$, and define L_x for all other x as required for the far triangles in Ω_f to be octahedral, i.e. in the associated octahedron for a triangle abc , the linear forms on the two vertices in each of the three parts are $\{L_a, L_b + L_c\}$, $\{L_b, L_c + L_a\}$ and $\{L_c, L_a + L_b\}$. By Lemma 4.3 whp G^* is such that for any triangle f in Φ , there are $(1 \pm 60c)d(G)^{45}\gamma^{15}2^{3a}$ valid choices of Ω_f . Here we also use the fact that for any triangle abc of Ω_f other than f there are only 2^{2a} solutions to $L_a(z) + L_b(z) + L_c(z) = 0$. The precise exponents of $d(G)$ and γ (which are not important for the argument) may be easily calculated from the observation that adding an octahedron to a triangle adds three new vertices and nine new edges, and Ω_f^+ is the composition of five such extensions.

Next we claim that whp the graph Γ of new edges is linearly c'_3 -bounded, where $c'_3 = 400c_3d(G)^{-45}\gamma^{-15}$. We assume this bound on the current graph of new edges and estimate how many configurations are excluded. Consider any edge uu' of the extended configuration. Suppose first that $uu' \cap f = \emptyset$. If $L_u(y) + L_{u'}(y)$ is not constant, then for any $vv' \in G^*$ the number of L -embeddings with $L_u(y) = v$ and $L_{u'}(y) = v'$ is at most 2^a . There are at most $45(|M^c| + |M^i| + |M^o|) < 100c_3n^2$ choices for a previous new edge vv' , so this excludes at most $100c_3n^22^a$ configurations. On the other hand, if $L_u(y) + L_{u'}(y)$ is constant, then $L_u(y)$ and $L_{u'}(y)$ are constrained to lie on a basic line; there are at most c'_32^a choices for vv' by linear boundedness, and each such vv' excludes at most 2^{2a} configurations. The latter estimate also applies to the case when one of u or u' is in f . Summing these bounds over all uu' , we see that fewer than half of the total configurations are excluded.

Next we fix any edge uu' of the extended configuration and any $vv' \in G^*$, and estimate the sum over $f \in \Phi$ of the probability p that a random configuration satisfies $L_u(y) = v$ and $L_{u'}(y) = v'$. If $uu' \cap f = \emptyset$ and $L_u(y) + L_{u'}(y)$ is not constant, then $p < 2^a/(1 - 60c)d(G)^{45}\gamma^{15}2^{3a}$ for any $f \in \Phi$. There are at most c_3n^2 choices for f , so the total contribution is at most $2c_3d(G)^{-45}\gamma^{-15}$. Otherwise, if $L_u(y) + L_{u'}(y)$ is constant or one of u or u' is in f , then one vertex of f is specified by $(L_u(y), L_{u'}(y))$. For example, writing $f = abc$, in the associated octahedron for az_2z_3 , if $u = z_2$ and $u' = a + z_2$ then a is specified by $(L_u(y), L_{u'}(y))$. Then there are at most $2c_3n$ choices for f (as $\bigcup M^o$ is c_3 -bounded). For each such f we have a contribution of at most $2^{2a}/(1 - 60c)d(G)^{45}\gamma^{15}2^{3a}$, so again the total contribution is at most $2c_3d(G)^{-45}\gamma^{-15}$. Summing these bounds over all uu' we can estimate $E_{vv'} < 100c_3d(G)^{-45}\gamma^{-15} = c'_3/4$. Applying Lemma 2.9 as in the proof of Lemma 2.10, we deduce the claimed bound on Γ .

We also claim that whp there are at most $2c'_32^a$ far triangles in any basic plane $\Pi = \{z : b \cdot z = v\}$. To see this, we first consider the contribution from the template triangles $\Pi^* = \Pi \cap T$. Since $z_1 + z_2 + z_3 = 0$ is linearly independent or contradictory to the defining equation of Π , we have $|\Pi^*| \leq 2^a$. Summing $E_{vv'} < c'_3/4$ over an edge vv' in each triangle of Π^* , by Lemma 2.9 we see that whp Π contains at most c'_32^a template triangles. Now fix any far non-template triangle f' of the extended configuration, any $g \in K_3(G^*)$, and estimate the sum over $f \in \Phi$ of the probability p that a random configuration satisfies $L_{f'}(y) = g$. If $f' \cap f = \emptyset$ then as f' is non-template it determines the configuration, so $p < 1/(1 - 60c)d(G)^{45}\gamma^{15}2^{3a}$, giving a total contribution of at most $2c_3d(G)^{-45}\gamma^{-15}n^{-1}$. Otherwise, f' determines one of the associated octahedra, so specifies one vertex of f , for example, writing $f = abc$, we see that if $f' = \{z_2, a + z_2, a + z_3\}$ then a is specified. Then there are at most $2c_3n$ choices for f ; for each such f we have $p < 2^a/(1 - 60c)d(G)^{45}\gamma^{15}2^{3a}$, so again the total contribution is at most $2c_3d(G)^{-45}\gamma^{-15}n^{-1}$. Summing over f' and applying Lemma 2.9 as in the proof of Lemma 2.10, we deduce the claimed bound on Π . This completes the analysis of Phase I.

To analyse Phase II, we first estimate the number of choices for an extended configuration on a pair ff' . This can be described by the linear extension $(\Omega_{ff'}^+, L)$, where $\Omega_{ff'}^+$ is as above, we have variables $z = (z_1, z_2)$, which we also use to label the ver-

tices of $\Omega_{ff'} \setminus (f \cup f')$, we define $L_x = x$ for all $x \in \Omega_{ff'}$, and define L_x for all other x as required for the triangles in $\Omega_{ff'}$ other than f and f' to be octahedral. The linear forms are distinct, as f and f' are not template triangles. By Lemma 4.3 there are $(1 \pm 60c)d(G)^{53}\gamma^{20}2^{2a}$ valid $\Omega_{ff'}$. Again, the precise exponents of $d(G)$ and γ are not important for the argument, but are straightforward to calculate: e.g. γ appears with exponent 20, as $\Omega_{ff'}$ and each of the seven associated octahedra adds three new vertices to the extension, but four vertices in the associated octahedra belong to the base of the extension, being the third vertex of the template triangle containing an edge in f or f' other than e .

We claim that whp Γ' is linearly $c_4/2$ -bounded. The argument is very similar to that given above for Γ . Assuming this bound on the current graph of new edges, one can show similarly to before that fewer than half of the total configurations are excluded. We also need to estimate the sum over $f \in \Phi$ of the probability that a random configuration satisfies $L_u(y) = v$ and $L_{u'}(y) = v'$, for any uu' in the extended configuration and $vv' \in G^*$. For most choices for uu' the required bound follows as before, but there is an additional case, namely when $uu' \cap (f \cup f') = \emptyset$ and $L_u(y) + L_{u'}(y)$ is constant, it may be that no vertex of $f \cup f'$ is specified by $(L_u(y), L_{u'}(y))$, but instead some pair (not e) is constrained to lie on a basic line. For example, writing $f = abc$ and $f' = abc'$, if $u = b + z_1$ and $u' = c + z_1$ then $(L_u(y), L_{u'}(y))$ specifies $b + c$, but not b or c . In this case, we use the fact that Γ is linearly c'_3 -bounded to see that there are at most $c'_3 2^a$ choices for ff' . Each such ff' contributes at most $2^a / (1 - 60c)d(G)^{53}\gamma^{20}2^{2a}$, giving a total contribution of at most $2c'_3 d(G)^{-53}\gamma^{-20}$. Summing over all uu' we estimate $E_{vv'} < 100c'_3 d(G)^{-53}\gamma^{-20}$, so the claim follows from Lemma 2.9.

Finally, M_2 satisfies the conditions (P1)–(P3): indeed, (P1) holds by definition of the extended configurations and random greedy algorithms, (P2) holds as $\Delta \subseteq \Gamma \cup \Gamma'$, and (P3) holds as whp Ψ has at most $c_4 n$ triangles in any basic plane: this holds for the new triangles in this algorithm by the same argument as for Φ' , and we may include the far triangles from the previous algorithm in this estimate. \square

This completes the proof of **Completion**, and so of Theorem 2.1, under the assumption that G is $(c, 16)$ -typical. The following modification proves the theorem under the assumption that G is c -typical. It is well-known that G is $c^{1/50}$ -regular (say) in the ‘Szemerédi sense’ (see e.g. [6, Theorem 2.2]). For $S \subseteq V(G)$ with $|S| \leq 16$, say that S is *good* if $|\bigcap_{x \in S} G(x)| = (1 \pm |S|c^{1/100})d(G)^{|S|}n$, otherwise *bad*. As G is c -typical, there are no bad sets of size 1 or 2. By regularity, for $k \leq 15$, any good k -set is contained in at most $c^{1/100}n$ bad $(k + 1)$ -sets. The proof of Lemma 2.5 still applies if we assume that all subsets of S are good. Thus we can avoid using bad sets with negligible changes to the calculations.

5. Random triangle removal

In this section we sketch a proof of Theorem 2.8, by describing how to apply the analysis of random greedy hypergraph matching by Bennett and Bohman [1] (we choose this for simplicity, but there are several other alternative approaches).

Consider the random triangle removal algorithm starting with G rather than K_n . The intuition is that after i steps the remaining graph G^i should look like a random subgraph of G where edges are retained independently with probability $p = 1 - 3t$, where $t = i/|G|$. For any $e \in G$ let $T_e(i)$ denote the number of triangles of G^i containing e , and $Q(i)$ denote the total number of triangles of G^i . By assumption, $T_e(0) = (1 \pm b)D$ for all $e \in G$, where $D = d(G)^2n$. Note also that $Q(0) = (1 \pm b)|G|D/3$.

We will show that for $0 \leq i \leq (1 - b^{1/4})|G|/3$ whp

$$Q(i) = |G|Dp^3/3 \pm e_q \quad \text{and} \quad T_e(i) = Dp^2 \pm e_d \quad \text{for all } e \in G^i,$$

where $e_q = 2(1 - 3 \log p)^2 b|G|D$ and $e_d = 2(1 - 3 \log p)b^{2/3}D$. We restrict attention to the upper bounds, as the lower bounds are similar. A convenient reformulation is to show negativity of the shifted variables

$$Q^+(i) = Q(i) - |G|Dp^3/3 - e_q \quad \text{and} \quad T_e^+(i) = T_e(i) - Dp^2 - e_d.$$

This follows whp from martingale concentration inequalities (e.g. [3]) after we verify the following ‘trend’ and ‘boundedness’ hypotheses, supposing that the required estimates for Q and T_e hold at previous steps (i.e. $i < \tau$, where the stopping time τ is the first step where any of the required estimates fails, or ∞); we use primes to denote conditional expectation given the history of the process.

Trend hypothesis: If $Q^+(i) \geq -b|G|D$ then $\mathbb{E}'Q^+(i+1) \leq Q^+(i)$; if $T_e^+(i) \geq -b^{2/3}D$ then $\mathbb{E}'T_e^+(i+1) \leq T_e^+(i)$.

Boundedness hypothesis: $(Q^+(i+1) - Q^+(i))^2|G|p \log n < (b|G|D)^2$, and $-\Theta < T_e^+(i+1) - T_e^+(i) < \theta$ with $\theta < \Theta/10$ and $\theta\Theta|G|p \log n < (b^{2/3}D)^2$.

We start with the boundedness hypothesis, which holds with room to spare. Indeed, $|Q^+(i+1) - Q^+(i)| = O(n)$, so $(Q^+(i+1) - Q^+(i))^2|G|p \log n = O(n^4 \log n)$, whereas $(b|G|D)^2 > b^2 d(G)^3 n^6/4 > n^{5.5}/4$. Also, for any $e \in G^i$ we have $-2 \leq T_e(i+1) - T_e(i) \leq 0$. The change in Dp^2 is $O(Dp/|G|)$ and in e_d is $O(b^{2/3}D/p|G|)$, so we can take $\Theta = 3$ and $\theta = O(p + b^{2/3}/p)D/|G|$. Then $\theta\Theta|G|p \log n = O(p^2 + b^{2/3})D \log n = O(n \log n)$, whereas $(b^{2/3}D)^2 = b^{4/3}d(G)^4 n^2 > n^{1.4}$.

Next consider the trend hypothesis for Q . Conditional on the required estimates at step i , if $Q^+(i) \geq -b|G|D$ we have

$$\begin{aligned} \mathbb{E}'[Q(i+1) - Q(i)] &= -Q(i)^{-1} \sum_{e \in G^i} T_e(i)^2 + O(1) \\ &= -9Q(i)/|G|p \pm 2|G|pe_d^2/Q(i) + O(1) \\ &\leq -3Dp^2 - 9e_q/|G|p + 9bD/p + (6 + o(1))e_d^2/Dp^2. \end{aligned}$$

Also, since e_q is increasing, the one-step change in $-|G|Dp^3/3 - e_q$ is at most $(1 + O(1/|G|))3Dp^2$. As $p \geq b^{1/4}$, we deduce

$$\begin{aligned} \mathbb{E}'[Q^+(i+1) - Q^+(i)] &\leq -9e_q/|G|p + 9bD/p + 7e_d^2/Dp^2 \\ &\leq -18(1 - 3 \log p)^2 bD/p + 9bD/p + 28(1 - 3 \log p)^2 b^{4/3}D/p^2 \leq 0. \end{aligned}$$

Now consider the trend hypothesis for T_e . Conditional on the required estimates at step i , if $T_e^+(i) \geq -b^{2/3}D$ we have

$$\begin{aligned} \mathbb{E}'[T_e(i+1) - T_e(i)] &= -Q(i)^{-1}T_e(i)2(Dp^2 \pm e_d + O(1)) \\ &\leq -2Q(i)^{-1}(Dp^2 + e_d - b^{2/3}D)(Dp^2 - e_d - O(1)) \\ &\leq -6(Dp^2)^2/|G|Dp^3 + (6 + o(1))(b^{2/3}D)(Dp^2)/|G|Dp^3 + O(e_d/|G|^2p^2). \end{aligned}$$

Also, the one-step change in $-Dp^2$ is at most $(1 + O(1/|G|))6Dp/|G|$, and in $-e_d$ is at most $-(1 + O(1/|G|p))18b^{2/3}D/|G|p$. We deduce

$$\mathbb{E}'[T_e^+(i+1) - T_e^+(i)] \leq -10b^{2/3}D/|G|p + O(\log^2 p)bD/|G|p^2 \leq 0.$$

Thus the required estimates hold, i.e. for $0 \leq i \leq (1 - b^{1/4})|G|/3$, whp we have $Q(i) = |G|Dp^3/3 \pm e_q$ and $T_e(i) = Dp^2 \pm e_d$ for all $e \in G^i$.

Now we apply the same method to deduce the boundedness conclusion of Theorem 2.8. For $0 \leq i \leq (1 - b^{1/4})|G|/3$ we show whp

$$|G^i(v)| = p|G(v)| \pm e_v \quad \text{for any vertex } v,$$

where $e_v = 2b^{1/3}d(G)n$. The upper bound $G_v^+(i) := |G^i(v)| - p|G(v)| - e_v \leq 0$ will follow whp after we verify the following two conditions.

Trend hypothesis: If $G_v^+(i) \geq -b^{1/3}d(G)n$ then $\mathbb{E}'G_v^+(i+1) \leq G_v^+(i)$.

Boundedness hypothesis: $-\Theta < G_v^+(i+1) - G_v^+(i) < \theta$ with $\theta < \Theta/10$ and $\theta\Theta|G|p \log n < (b^{1/3}d(G)n)^2$.

For the boundedness hypothesis, we can take $\Theta = 2$ and $\theta = 3|G(v)|/|G|$, so that $\theta\Theta|G|p \log n = 6p|G(v)| \log n = O(n \log n)$, whereas $(b^{1/3}d(G)n)^2 > b^3n^2 \geq n^{1.7}$. For the trend hypothesis, if $G_v^+(i) \geq -b^{1/3}d(G)n$ we have

$$\begin{aligned} \mathbb{E}'[|G^{i+1}(v)| - |G^i(v)|] &= - \sum_{e:v \in e \in G^i} T_e(i)/Q(i) \\ &\leq -(p|G(v)| + e_v - b^{1/3}d(G)n)(Dp^2 - e_d)/(|G|Dp^3/3 + e_q) \\ &\leq -3|G(v)|/|G| - 3(e_v - b^{1/3}d(G)n)/|G|p + O(e_d)\frac{|G(v)|}{|G|Dp^2} + O(e_q)\frac{|G(v)|}{|G|^2Dp^3}. \end{aligned}$$

The one-step change in $-p|G(v)| - e_v$ is at most $3|G(v)|/|G|$, so

$$pn \mathbb{E}'[G_v^+(i+1) - G_v^+(i)] \leq -6b^{1/3} + O(e_d/Dp) + O(e_q/|G|Dp^2) \leq 0,$$

as $e_d/Dp = O(b^{2/3}p^{-1} \log p)$, $e_q/|G|Dp^2 = O(bp^{-2} \log^2 p)$ and $p \geq b^{1/4}$.

Thus, letting N be the set of triangles removed during the process, we see that whp $L = G \setminus \bigcup N$ is $b^{1/4}$ -bounded.

6. The number of designs

In this section we generalise Theorem 2.2 to estimate the number of designs. We start by describing the results of [5] on the existence of designs. Let D be a q -graph (i.e. a set of subsets of size q) of a set X of size n . We say that D is a *design* with parameters (n, q, r, λ) if every subset of X of size r belongs to exactly λ elements of S . Note that if $q = 3, r = 2, \lambda = 1$ then D is a Steiner Triple System. The necessary divisibility conditions generalise in a straightforward way: if D exists then $\binom{q-i}{r-i}$ must divide $\lambda \binom{n-i}{r-i}$ for every $0 \leq i \leq r - 1$; to see this, fix any i -subset I of X and consider the sets in D that contain I . In [5] we proved the ‘Existence Conjecture’, which states that these divisibility conditions are also sufficient for the existence of a design with parameters (n, q, r, λ) , assuming q, r, λ are fixed and $n > n_0(q, r, \lambda)$ is large. We will generalise Theorem 2.2 as follows.

Theorem 6.1. *For any q, r, λ there is n_0 such that if $n > n_0$ and $\binom{q-i}{r-i} \mid \lambda \binom{n-i}{r-i}$ for all $0 \leq i \leq r - 1$, and if we write $Q = \binom{q}{r}$ and $N = \binom{n-r}{q-r}$, then the number $D(n, q, r, \lambda)$ of designs with parameters (n, q, r, λ) satisfies*

$$D(n, q, r, \lambda) = \lambda! \binom{n}{r} ((\lambda/e)^{Q-1} N + o(N))^{\lambda Q^{-1}} \binom{n}{r}.$$

The proof of Theorem 6.1 follows that of Theorem 2.2: the lower bound generalises the argument given earlier in this paper, and the upper bound generalises that of Linial and Luria [8].

We start with the lower bound. In the same way as a Steiner Triple System can be viewed as a triangle decomposition of K_n , we can view a design with parameters (n, q, r, λ) as a K_q^r -decomposition of λK_n^r , where K_q^r denotes the complete r -graph on q vertices and λK_n^r denotes the multi(hyper)graph in K_n^r in which each edge has multiplicity λ . To generalise Theorem 2.1, we first need to define the divisibility and typicality conditions for general r -graphs (we will omit multiplicities in the definitions, as we do not need them for our application here).

For $S \subseteq V(G)$, the *neighbourhood* $G(S)$ is the $(r - |S|)$ -graph $\{f \subseteq V(G) \setminus S : f \cup S \in G\}$. We say that G is K_q^r -divisible if $\binom{q-i}{r-i}$ divides $|G(e)|$ for any i -set $e \subseteq V(G)$, for all $0 \leq i \leq r$. We say that G is (c, h) -typical if there is some $p > 0$ such that for any set A of $(r - 1)$ -subsets of $V(G)$ with $|A| \leq h$ we have $|\bigcap_{S \in A} G(S)| = (1 \pm c)p^{|A|}n$.

Now we can state the r -graph generalisation of Theorem 2.1. When $d(G)$ is at least a constant independent of n this follows from [5, Theorem 1.4]; the same proof shows that $d(G)$ can decay polynomially in n .

Theorem 6.2. *For any $q > r \geq 1$ there are $c_0, a \in (0, 1)$ and $h, \ell, n_0 \in \mathbb{N}$ such that if $n \geq n_0$ and G is a K_q^r -divisible (c, h) -typical r -graph on n vertices with $d(G) > n^{-a}$ and $c < c_0 d(G)^\ell$ then G has a K_q^r -decomposition.*

In the proof of Theorem 6.1 it is more convenient to count designs together with a choice for each $e \in K_n^r$ of a bijection between the λ copies of e in λK_n^r and the λ sets of the design containing e ; we will refer to such a structure as an *edge-labelled design* with

parameters (n, q, r, λ) and denote their number by $D^*(n, q, r, \lambda)$. As $D^*(n, q, r, \lambda) = \lambda! \binom{n}{r} D(n, q, r, \lambda)$, it suffices to show

$$D^*(n, q, r, \lambda) = ((\lambda/e)^{Q-1} N + o(N))^{\lambda Q^{-1} \binom{n}{r}}.$$

Proof of Theorem 6.1. For the lower bound we start by setting aside a random subgraph R of K_n^r in which each edge is chosen with probability $n^{-a/Q}$, where we can apply Theorem 6.2 with a and ℓ , and we suppose without loss of generality that $Q \ll h \ll \ell \ll 1/a$ (i.e. parameters are chosen from left to right to satisfy various inequalities below).

Next we will consider the random greedy matching process in the following auxiliary hypergraph A . We let $V(A)$ consist of λ copies of each edge of $K_n^r \setminus R$ and $\lambda - 1$ copies of each edge of R . We let $E(A)$ consist of all $\binom{q}{r}$ -sets in $V(A)$ that are edge-sets of a copy of K_q^r in K_n^r .

In the random greedy matching process, we start with A , and at each step we select a uniformly random edge e of the current hypergraph, then delete all vertices of e (and all incident edges) to obtain the hypergraph for the next step. We stop the process when fewer than $n^{r-3\ell a}$ vertices of A remain and let L denote the multigraph in K_n^r consisting of the remaining vertices of A . Similarly to the previous section, one can adapt the analysis of Bennett and Bohman [1] to show that whp (i) when $p\lambda \binom{n}{r}$ edges remain the number of choices for the next edge of the process is $(1 \pm n^{-1/Q})(p\lambda)^Q \binom{n}{q}$, and (ii) $|L(e)| < 2n^{1-3\ell a}$ for any $(r - 1)$ -set $e \subseteq [n]$.

Next we apply a random greedy algorithm to sequentially cover each edge of L by a copy of K_q^r in which all other edges are in R ; as usual, at each step we make a uniformly random choice subject to not using any previously covered edge. The proof of Lemma 2.10 generalises to show that whp the algorithm does not abort, and writing S for the subgraph of R covered by the algorithm, we see that whp $|S(e)| < n^{1-2\ell a}$ for any $(r - 1)$ -set $e \subseteq [n]$. By Chernoff bounds whp R is $(n^{-2\ell a}, h)$ -typical. Also whp $|R| > \frac{1}{2}n^{r-a/Q}$, so $R' := R \setminus S$ is $(c_0n^{-\ell a}, h)$ -typical with $|R'| > n^{r-a}$. Furthermore, R' was obtained from λK_n^r by deleting edge-disjoint copies of K_q^r , and λK_n^r is K_q^r -divisible by assumption, so R' is K_q^r -divisible. Therefore R' has a K_q^r -decomposition by Theorem 6.2. Combining this with the previous choices of K_q^r 's we have constructed an edge-labelled design with parameters (n, q, r, λ) (the vertices of A specify the edge-labelling).

Now we may calculate similarly to the proof of Theorem 2.2. Writing m for the number of steps in the random greedy matching process, and $p(i) = 1 - n^{-a/Q} - i Q \binom{n}{r}^{-1}$ for the approximate density at the i th step, we deduce that the logarithm of the number of choices is

$$L_1 = \sum_{i=1}^m (\log(p(i)^Q \lambda^Q \binom{n}{q}) \pm 2n^{-1/Q}) = \lambda Q^{-1} \binom{n}{r} (\log(\lambda^Q \binom{n}{q}) - Q \pm n^{-a/2Q}).$$

Also, for any fixed design, the logarithm of the number of times it is counted is at most

$$L_2 = \sum_{i=1}^m \log(p(i)\lambda Q^{-1} \binom{n}{r}) = \lambda Q^{-1} \binom{n}{r} (\log(\lambda Q^{-1} \binom{n}{r}) - 1 \pm n^{-a/2Q}).$$

As $\binom{n}{q} Q \binom{n}{r}^{-1} = N + o(N)$ we deduce

$$\log D^*(n, q, r, \lambda) \geq \lambda Q^{-1} \binom{n}{r} \log((\lambda/e)^{Q-1} N + o(N)).$$

For the upper bound in Theorem 6.1 we apply the Entropy Method, following the argument of Linial and Luria [8] for Steiner Triple Systems (see their paper for motivation and exposition of the method). We let X be a uniformly random edge-labelled design with parameters (n, q, r, λ) , and consider the entropy $H(X) = -\sum_D \mathbb{P}(X = D) \log \mathbb{P}(X = D)$ (using natural logarithms). As $D^*(n, q, r, \lambda) = e^{H(X)}$, it suffices to estimate $H(X)$.

We consider the labelled edges of λK_n^r in a uniformly random order: it is convenient to select $\mu = (\mu_e) \in [0, 1]^{\lambda K_n^r}$ uniformly at random, and to proceed by decreasing order of μ_e . At each step, when we consider e , we reveal the block X_e of X that contains e and is assigned to e according to the edge-labelling. Conditional on μ , we have

$$H(X) = \sum_{e \in \lambda K_n^r} H(X_e | (X_{e'} : \mu(e') > \mu(e))).$$

We estimate $H(X_e | (X_{e'} : \mu(e') > \mu(e))) \leq \log N_e^\mu$, where N_e^μ is the size of the support of the random variable $X_e | (X_{e'} : \mu(e') > \mu(e))$, i.e. $N_e^\mu = 1$ if e is a labelled edge of $X_{e'}$ for some e' that precedes e , otherwise N_e^μ is the number of choices of a labelled q -set f containing e (i.e. we fix labellings of the other $Q - 1$ edges in f) such that for each such labelled edge e' in f , no labelled edge of the block $X_{e'}$ precedes e .

Next we condition on X , fix e , and write F_e for the event that $\mu(e') \leq \mu(e)$ for all $e' \in X_e$. We estimate $\mathbb{E} \log N_e^\mu$, where the expectation is with respect to μ , and we suppress the X -conditioning in our notation. We have

$$\mathbb{E} \log N_e^\mu = \mathbb{E}(\mathbb{E}[\log N_e^\mu | \mu_e]) = \mathbb{E}(\mu_e^{Q-1} \mathbb{E}[\log N_e^\mu | \mu_e, F_e])$$

and by Jensen's inequality $\mathbb{E}[\log N_e^\mu | \mu_e, F_e] \leq \log \mathbb{E}[N_e^\mu | \mu_e, F_e]$.

Now we write $\mathbb{E}[N_e^\mu | \mu_e, F_e] = 1 + \sum_f \mathbb{P}[E_f | \mu_e, F_e]$, where the sum is over all labelled q -sets $f \neq X_e$ containing e , and E_f is the event that for each labelled edge e' in f , no labelled edge of the block $X_{e'}$ precedes e . Note that there are only $O(N/n)$ such f with $|f \cap f'| > r$ for some block f' of X . For any other such f we have $\mathbb{P}[E_f | \mu_e, F_e] = \mu_e^{Q(Q-1)}$. We deduce $\mathbb{E}[N_e^\mu | \mu_e, F_e] = \mu_e^{Q(Q-1)} \lambda^{Q-1} N + O(N/n)$.

Finally,

$$\begin{aligned} \log D^*(n, q, r, \lambda) = H(X) &\leq \sum_{e \in \lambda K_n^r} \mathbb{E} \mu_e^{Q-1} \log \mathbb{E}[N_e^\mu | \mu_e, F_e] \\ &= \lambda \binom{n}{r} \int_0^1 t^{Q-1} [\log(t^{Q(Q-1)} \lambda^{Q-1} N) + O(1/n)] dt. \end{aligned}$$

For any $A, B, C > 0$ we have $\int_0^1 t^{A-1} \log(Ct^B) dt = A^{-1} \log C - A^{-2} B$. Setting $A = Q$, $B = Q(Q - 1)$, $C = \lambda^{Q-1} N$ we deduce

$$\log D^*(n, q, r, \lambda) \leq \lambda Q^{-1} \binom{n}{r} \log((\lambda/e)^{Q-1} N + o(N)). \quad \square$$

7. Concluding remarks

Although we have proved (and generalised) Wilson's conjecture, one may still ask for more precise estimates (even an asymptotic formula) for the number of Steiner Triple Systems, and more generally designs. Such results have been obtained by Kuperberg, Lovett and Peled [7], using very different methods to ours, but only for designs within a certain range of parameters. One open case of particular interest (recently drawn to my attention by Ron Peled) is the problem of estimating the number $G(n, d)$ of d -regular graphs on n vertices. These may be viewed as designs with parameters $(n, 2, 1, d)$, for which our methods give $G(n, d) = d!^{-n} (dn/e + o(dn))^{dn/2}$. Much more precise results have been obtained by McKay and Wormald, including asymptotic enumeration for $d = \omega(n/\log n)$ (see [10]) and $d = o(\sqrt{n})$ (see [11]); their conjecture in [10] regarding a general asymptotic formula remains open.

Acknowledgments. We thank an anonymous referee for some helpful remarks.

Research supported in part by ERC grant 239696.

References

- [1] Bennett, P., Bohman, T.: A natural barrier in random greedy hypergraph matching. [arXiv:1210.3581](#) (2012)
- [2] Bohman, T., Frieze, A., Lubetzky, E.: Random triangle removal. *Adv. Math.* **280**, 379–438 (2015) [Zbl 1312.05123](#) [MR 3350225](#)
- [3] Freedman, D. A.: On tail probabilities for martingales. *Ann. Probab.* **3**, 100–118 (1975) [Zbl 0313.60037](#) [MR 0380971](#)
- [4] Graver, J. E., Jurkat, W. B.: The module structure of integral designs. *J. Combin. Theory Ser. A* **15**, 75–90 (1973) [Zbl 0259.05020](#) [MR 0329930](#)
- [5] Keevash, P.: The existence of designs. [arXiv:1401.3665](#) (2014)
- [6] Keevash, P.: A hypergraph regularity method for generalised Turán problems. *Random Structures Algorithms* **34**, 123–164 (2009) [Zbl 1182.05125](#) [MR 2478542](#)
- [7] Kuperberg, G., Lovett, S., Peled, R.: Probabilistic existence of regular combinatorial objects. *Geom. Funct. Anal.* **27**, 919–972 (2017) [Zbl 1369.05024](#) [MR 3678505](#)
- [8] Linial, N., Luria, Z.: An upper bound on the number of Steiner triple systems. *Random Structures Algorithms* **43**, 399–406 (2013) [Zbl 1278.05030](#) [MR 3124689](#)
- [9] McDiarmid, C.: Concentration. In: *Probabilistic Methods for Algorithmic Discrete Mathematics*, *Algorithms Combin.* **16**, 195–248 (1998) [Zbl 0927.60027](#) [MR 1678578](#)
- [10] McKay, B. D., Wormald, N. C.: Asymptotic enumeration by degree sequence of graphs of high degree. *Eur. J. Combin.* **11**, 565–580 (1990) [Zbl 0739.05043](#) [MR 1078713](#)
- [11] McKay, B. D., Wormald, N. C.: Asymptotic enumeration by degree sequence of graphs with degrees $o(n^{1/2})$. *Combinatorica* **4**, 369–382 (1991) [Zbl 0742.05047](#) [MR 1137769](#)
- [12] Wilson, R. M.: Nonisomorphic Steiner triple systems. *Math. Z.* **135**, 303–313 (1974) [Zbl 0264.05009](#) [MR 0340046](#)
- [13] Wilson, R. M.: The necessary conditions for t -designs are sufficient for something. *Utilitas Math.* **4**, 207–215 (1973) [Zbl 0286.05005](#) [MR 0325415](#)
- [14] Wilson, R. M.: Signed hypergraph designs and diagonal forms for some incidence matrices. *Designs Codes Cryptography* **17**, 289–297 (1999) [Zbl 0938.05047](#) [MR 1715268](#)